The PPE2 protein of *Mycobacterium tuberculosis* is responsible for the development of
hyperglycemia and insulin resistance during tuberculosis

Manoj Kumar Bisht[1], Vandana Maurya[1,2], Priyanka Dahiya[1] Vijaya Lakshmi Valluri[3], Sudip

Ghosh[4], and Sangita Mukhopadhyay[1,¶,]

[1]Laboratory of Molecular Cell Biology, BRIC-Centre for DNA Fingerprinting and

Diagnostics, Inner Ring Road, Uppal, Hyderabad - 500039, Telangana, India

[2]Graduate Studies, Regional Centre for Biotechnology, Faridabad, Haryana, India

[3]Immunology & Molecular Biology Department, Bhagwan Mahavir Medical Research Centre

(BMMRC), AC Guards, Masab Tank, Hyderabad – 500004, Telangana, India

[4]ICMR-National Institute of Nutrition, Hyderabad - 500007, Telangana, India

¶To whom correspondence should be addressed: Laboratory of Molecular Cell Biology,

BRIC-Centre for DNA Fingerprinting and Diagnostics (CDFD), Inner Ring Road, Uppal,

Hyderabad - 500039, Telangana, India. Tel: +91-40-27216134; E-mail: sangita@cdfd.org.in

Running Title: PPE2 protein causes insulin resistance

Keywords: PPE2 protein; Mice; *Mycobacterium tuberculosis*; Adipose tissue; Insulin

resistance; Lipolysis.

Abstract

Diabetes is a known risk factor for tuberculosis (TB), but clinical evidences suggest that TB itself can induce hyperglycaemia and insulin resistance, though the underlying mycobacterial factors are not known. Herein, we implicate PPE2, a secretory PE/PPE family protein of *Mycobacterium tuberculosis* (Mtb), as a key modulator of adipose tissue physiology that contributes to the development of insulin resistance. In mice, PPE2 caused fat loss, adipocyte hypertrophy, immune cell infiltration, impaired glucose tolerance, reduced expression of PPAR-γ, C/EBP-α, adiponectin and higher insulin resistance. Transcriptomic analysis revealed PPE2 altered expression of genes associated with chemokine/cytokine, ribosomal biogenesis and lipase signaling. PPE2 induced lipolysis by activating cAMP–PKA–HSL axis, increased circulating free fatty acids, a feature also observed in TB patient sera. Interestingly, PPE2-immunization mitigated these effects, suggesting its potential as a subunit vaccine. Overall, this study identifies PPE2 as a key link between Mtb-infection, adipose tissue dysfunction and insulin resistance.

Introduction

Mycobacterium tuberculosis (Mtb) remains one of the deadliest bacterial pathogens, causing approximately 1.5 million deaths annually. Each year, over 10 million people are infected, but about 5-10% of them develop clinical symptoms of active tuberculosis (TB) disease (WHO 2023). In the majority of cases, the bacilli enter a dormant phase during which individuals test positive on immunologic tests but do not exhibit clinical symptoms, a condition referred as latent tuberculosis (Lin and Flynn, 2010). Latent tuberculosis can reactivate into active tuberculosis when the immune system becomes compromised due to factors such as undernutrition, smoking, HIV co-infection, use of immunosuppressive drugs and diabetes (Ai et al., 2016). It has been found that people with diabetes are 1.5-2.4 times higher risk of developing TB (Franco et al., 2024) and diabetes is responsible for 10.6% of the total TB deaths (Kyu et al., 2018). The co-occurrence of tuberculosis and diabetes (TB-DM) complicates TB treatment and drastically worsens disease outcomes. Several studies have reported that TB can exacerbate the development and severity of diabetes-associated complications (Magee et al., 2018; Mugusi et al., 1990). The TB-DM co-occurrence is a growing global health concern with an established bidirectional relationship between the two conditions (Khattak et al., 2024; Gautam et al., 2020). Diabetes is a well-known risk factor for tuberculosis with an estimated global prevalence of 15.3% among TB patients, varying by country of origin, age, sex, region, level of country income and development (Noubiap et al., 2019). The prevalence of diabetes in Nigeria (15%), Tanzania (11%), and Ethiopia (10%) (Alebel et al., 2019) is notable while in India it is as high as 25.3% (Viswanathan et al., 2012), which has the second largest number of diabetic patients in the world (Atlas IDFD, 2025).

Several clinical studies indicate that TB patients are prone to develop insulin resistance and

hyperglycemia. In a Nigerian study, it has been found that about 37% of the patients with

active pulmonary TB had impaired glucose tolerance, which could be reversed during anti-

tuberculosis treatment (Oluboyo and Erasmust 1990). More recently many studies have

shown that TB patients with no known history of diabetes can develop hyperglycemia

(Magee et al., 2018; Lin et al., 2017). Though transient hyperglycemia is commonly observed

in many TB patients, 8-87% of TB patients are newly diagnosed with dysglycemia without a

prior history (Mave et al., 2017; Magee et al., 2018).

Development of hyperglycemia, glucose intolerance and insulin resistance can generally be

attributed to metabolic changes, induction of pro-inflammatory conditions in the host, and

dysregulation of adipose tissue functions (Guilherme et al., 2008). Interestingly, Mtb is

known to persist in the adipose tissue and to regulate pulmonary pathology and has been

implicated in tuberculosis and changes in adipose tissue behavior (Erol, 2008; Neyrolles et

al., 2006; Beigier-bompadre et al., 2017). Mtb persistence in adipose tissue is known to cause

adipocyte hypertrophy, lipolysis, changes in physiology and infiltration of immune cells in

adipose tissue (Beigier-bompadre et al., 2017; Bisht et al., 2023a). Monocyte chemoattractant

protein-1 (MCP-1) expression was found to be higher in mycobacterium-infected visceral

adipose, which has been linked to increased immune cell infiltration, insulin resistance and

hepatic steatosis (Kanda et al., 2006). Adipose tissue-resident Mtb is known to alter its

physiology which may cause a shift in the metabolism of fatty acids (Neyrolles, 2006;

Beigier-bompadre et al., 2017; Erol, 2008).

Though direct interaction between Mtb and host adipose tissue exists, the nature and outcome

of such interaction are not well understood. Therefore, we speculated that certain Mtb

factors/proteins, which are either exposed on the cell wall or secreted into the host circulation

or directly released into the adipose tissue milieu by resident Mtb, are likely to be involved in

modulating adipose tissue physiology and functions. In this study, we investigated the role of

a secretory mycobacterial protein, PPE2, in the modulation of adipose tissue physiology and

functions. PPE2 belongs to the PE/PPE family and possesses a nuclear localization signal

sequence and a leucine zipper DNA-binding motif, and is secreted during Mtb infection in

mice (Bhat et al., 2013; Bhat et al, 2017; Abraham et al., 2014; Bisht et al., 2023b). PPE2

appears to be a pleiotropic protein and plays an important role in mycobacterial virulence

(Srivastava et al., 2019; Pal et al., 2021). It inhibits nitric oxide (NO) production by binding

to the GATA-1-like elements in the iNOS promoter, inhibits the production of reactive

oxygen species (ROS) by destabilizing NADPH-oxidase complex, as well as inhibits

myelopoiesis to reduce the production of myeloid cells like leukocytes, mast cells, platelets

and red blood cells (RBCs) (Bhat et al., 2017; Srivastava et al., 2019; Pal et al., 2021; Pal and

Mukhopadhyay, 2021).

To the best of our knowledge, for the first time, our study provides evidence that PPE2 can

modulate gene expression and adipose tissue functions leading to glucose intolerance and

hyperglycemia. Using recombinantly purified protein, a surrogate non-pathogenic

mycobacterium expressing PPE2 and a *ppe2*-null mutant of a clinical *M. tuberculosis* strain

(CDC1551)*,* we have shown that PPE2 interferes with adipogenesis by inhibiting adipogenic

transcription factors, inhibiting adipokines and promoting adipocyte hypertrophy and

lipolysis, which can lead to the development of impaired glucose intolerance and insulin

resistance.

Results

1. Recombinant PPE2 protein (rPPE2) inhibits differentiation of adipocytes

Mycobacterium tuberculosis has been previously reported to inhibit adipogenic transcription factors like PPAR-γ and adipocyte secreted proteins like adiponectin, in the adipose tissue of infected mice (Ayyappan et al., 2019). PPE2 has been identified to be a secretory protein (Bhat et al., 2013; Pal et al., 2022), and its presence has been demonstrated in the bloodstream of mice infected with *M. smegmatis* and *M. tuberculosis* (Bisht et al., 2023b). Therefore, it is speculated that PPE2 may exert a systemic effect on the regulation of adipose tissue functions and thereby may contribute to diabetes-related complications. Therefore, to investigate the impact of PPE2 on adipogenesis, differentiation of 3T3-L1 pre-adipocyte was carried out in the presence of recombinantly purified PPE2 protein (rPPE2). Accordingly, 3T3-L1 cells were treated with a differentiation cocktail medium (Sadowski et al., 1992), in the absence or presence of various concentrations of rPPE2 and triglyceride accumulation was measured using Oil Red O staining at day 10 post-treatment. In the presence of rPPE2, the intensity of Oil Red O staining was found to be decreased in a concentration-dependent manner (Figure 1A, B), suggesting that PPE2 inhibits adipocyte differentiation. Significant inhibition was observed at rPPE2 concentration of 3 µg/ml, which was used for subsequent experiments. No cytotoxic effect of rPPE2 was observed when the 3T3-L1 preadipocytes were treated for 48 hours (Figure S1), suggesting that the observed inhibition of adipogenesis was due to physiological effect of rPPE2. To further confirm the specificity of inhibition, 3 µg/ml and 5 µg/ml of anti-PPE2 antibody (in house raised in Balb/c mice (Bisht et al., 2023b)) was used to neutralize the effect of rPPE2 during differentiation. It was found that, in

the presence of anti-PPE2 Ab, the inhibitory effect of rPPE2 on adipogenesis was neutralised

in a concentration-dependent manner (Figure 1C).

Next, the effect of PPE2 on adipocyte differentiation was examined in the context of

mycobacterial infection. Accordingly, 3T3-L1 pre-adipocytes were infected with either *M.*

smegmatis harbouring vector control (Msmeg-pVV) or *M. smegmatis* expressing PPE2

(Msmeg-PPE2), followed by induction of differentiation and Oil Red O staining. Cells

infected with Msmeg-PPE2 exhibited a significant reduction in adipocyte differentiation

compared to those infected with Msmeg-pVV (Figure S2), indicating that PPE2 could inhibit

adipocyte differentiation when present in the context of bacilli. However, infection with

Msmeg-pVV also caused a noticeable reduction in adipogenesis compared to uninfected

control cells, indicating that other factors intrinsic to *M. smegmatis* or infection-triggered

signaling may also contribute to the inhibition of adipogenesis. Next, we examined the effect

of PPE2 on critical adipogenic factors like PPAR-γ and C/EBP-α and fatty acid synthase

(FAS) during differentiation of 3T3-L1 cells. In the presence of rPPE2, there was a marked

decrease in the expression of PPAR-γ, C/EBP-α and FAS during differentiation (Figure 1D).

These findings suggest that PPE2 inhibits 3T3-L1 differentiation by downregulating key

adipogenic genes involved in adipocyte differentiation, highlighting its potential role in

modulating adipose tissue physiology and function.

2. PPE2 protein alters adipose tissue physiology during mice infection.

Mtb can persist within adipose tissue and alter its physiology, potentially leading to a shift in

the metabolism of fatty acids (Neyrolles et al., 2006, Beigier-Bompadre et al., 2017, Erol,

2008). While Mtb is known to cause inhibition of adipogenesis and promote lipolysis in

adipose tissue (Ayyappan et al., 2019), the mechanisms responsible for this effect remained unclear. Since we found that PPE2 inhibits adipogenesis in 3T3-L1 cells, we investigated its impact on adipose tissue physiology and function during infection in mice. Balb/c mice of 9-11 weeks old were infected with either Msmeg-pVV or Msmeg-PPE2 and sacrificed on day 2 and day 7 post-infection. The two major fat depots, the perigonadal (PG) and the visceral (VS) fat tissues, were analysed to assess fat mass, tissue pathology, gene expression pattern and bacterial survival. Mice infected with Msmeg-PPE2 showed significant loss of adipose tissue mass (Figure S3) compared to both the uninfected control and Msmeg-pVV infected groups, indicating that PPE2 inhibits adipogenesis in both perigonadal and visceral fat depots (Figure 2A, B). Notably, loss of adipose tissue was also observed in Msmeg-pVV-infected mice, implying that either increased energy demands to combat infection or some intrinsic factors from *M. smegmatis* may also contribute to adipose tissue atrophy.

Next, we examined the expression patterns of PPAR-γ, C/EBP-α, as well as adiponectin in these infected mice. PPAR-γ is a key transcription factor involved in regulating adipocyte differentiation, lipid metabolism and insulin sensitivity (Maréchal et al., 2018), while C/EBP-α is also required for adipogenesis as well as normal adipocyte function, while adiponectin is known to play important roles in variety of physiological functions including lipid metabolism, energy homeostasis, inflammation, and insulin sensitivity (Khoramipour et al., 2021). Infection with Msmeg-PPE2 resulted in a marked decrease in the expression of PPAR-γ, C/EBP-α and adiponectin in both PG and VS adipose tissues compared to mice infected with Msmeg-pVV at both 2^{nd} and 7^{th} day post-infection (dpi) (Figure 2C- F). These findings suggest that PPE2 during mycobacterial infection leads to altered gene expression, which potentially contributes to the modulation of the physiology and function of adipose tissue.

Adipose tissue is composed of various cell types including preadipocytes, adipocytes, endothelial cells, blood cells, fibroblasts, macrophages and several types of other immune cells (Saetang and Sangkhathat, 2018; Richard et al., 2020). During mycobacterial infection, immune cells infiltrate into adipose tissue and play a role in regulation of inflammation (Beigier-Bompadre et al., 2017). Histological analyses of adipose tissues stained with H&E revealed that mice infected with Msmeg-PPE2 exhibited increased immune cell infiltration and adipocyte hypertrophy by 7th dpi (Figure 2G and H). Higher infiltration of immune cells as well as adipose tissue hypertrophy was found to be more pronounced at 7th dpi in both the fat depots. Notably, at 7th dpi, an increased cell surface area was observed in perigonadal and visceral fat tissues of Msmeg-PPE2-infected mice in comparison to uninfected or Msmeg-pVV-infected mice (Figure 2I and J). These findings collectively indicate that Msmeg-PPE2 infection leads to enhanced immune cell infiltration and adipose tissue hypertrophy, influencing the cellular composition and morphology of adipocytes.

We also assessed mycobacterial survival within perigonadal and visceral fat tissues. Adipose tissues from mice infected with Msmeg-PPE2 had a significantly higher colony forming unit (CFU) count as compared to those infected with Msmeg-pVV at both 2nd and 7th dpi (Figure 2K and L). This suggests that PPE2 contributes to enhanced survival of mycobacteria within PG and VS adipose tissues.

These data suggest that PPE2 can significantly influence the adipose tissue at the morphological and molecular levels. Since adipose tissue plays a crucial role in regulating immune metabolism and glucose homeostasis, an intraperitoneal glucose tolerance test

(IPGTT) was performed in mice at 7^{th} dpi. After overnight fast, all mice received an

intraperitoneal injection of dextrose (2 g/kg body mass), and blood glucose levels were

monitored over 120 minutes. Interestingly, mice infected with Msmeg-PPE2 exhibited

increased glucose intolerance in comparison to both uninfected and Msmeg-pVV-infected

mice at 7^{th} dpi (Figure 2M). Analyses of the area under the curve (AUC) indicate that

Msmeg-PPE2-infected mice had higher glucose intolerance compared to other groups (Figure

2N). These findings indicate that PPE2 may impair glucose homeostasis through its effects on

adipose tissue physiology.

3. Administration of recombinantly purified PPE2 protein causes metabolic

perturbations in mice.

In the previous section, it was found that PPE2, when presented in the context of bacterium

can modulate adipose tissue physiology in mice. To further confirm that the observed

changes can be directly attributed to the PPE2 protein, mice were injected intraperitoneally

with 3 mg/kg of either recombinantly purified PPE2 protein (rPPE2) or bovine serum

albumin (BSA) as a control on alternate days for a total duration of 12 days and sacrificed on

day 14. The morphological changes along with histopathology and gene expression pattern

were analysed to assess the direct impact of PPE2 on adipose tissue physiology. As expected,

on morphological examination of PG and VS fat depots, significant reduction in both size and

mass was observed in rPPE2-administered mice compared to those administered with BSA

(Figure 3A, B, C). Furthermore, rPPE2-administration led to significant decrease in the

expressions of PPAR-γ, C/EBP-α and adiponectin in the PG fat tissue in comparison to those

treated with BSA (Figure 3D). These data indicate that PPE2 can specifically affect the

expression of key genes involved in adipogenesis and immune-metabolic functions.

Histopathological studies of PG and VS fat tissues in rPPE2-treated mice revealed

pronounced adipocyte hypertrophy and increased infiltration of immune cells into the adipose

tissue marked by intense staining of the nucleus of the infiltrated cells (Figure 3E).

To assess the impact of rPPE2 on glucose homeostasis, an IPGT test was performed. The

results indicate a distinctive metabolic effect, with rPPE2-treated mice displaying a lower rate

of glucose clearance as compared to the control mice treated with BSA (Figure 3F and G).

This observation indicates that PPE2 is a potential mycobacterial factor for development of

hyperglycaemia during tuberculosis.

238   4. Immunization of mice with rPPE2 ameliorates PPE2-induced adipose tissue

239   pathophysiology and functions.

Since PPE2 is a secretory protein and detectable in the circulation of Mycobacterium-infected

mice (Bisht et al., 2023b) and capable of promoting glucose intolerance, we next sought to

determine whether neutralization of circulating PPE2 could ameliorate PPE2-induced glucose

intolerance during mycobacterial infection. Accordingly, mice were immunized with rPPE2

protein using Freund's incomplete adjuvant, followed by three booster doses at 15-day

intervals. All the immunized mice showed a higher titre of anti-PPE2 antibody in the sera

(Figure S4). Subsequently, these mice were infected with either Msmeg-pVV or Msmeg-

PPE2, while mice injected with PBS served as unimmunized control group. As expected, loss

of PG and VS fat tissues was effectively prevented in Msmeg-PPE2 infected mice which

were previously immunized with rPPE2 on 7[th] dpi (Figure 4A, B and C). This is reflected in

significant improvement in the total body fat of the Msmeg-PPE2-infected mice which were

immunized previously. In contrast, unimmunized mice showed significant loss of total body

fat content (Figure 4D). Histopathological examination of both PG and VS adipose tissues showed marked improvement in the pathology in mice immunized with rPPE2 when compared to unimmunized mice (Figure 4E). Moreover, reduced infiltration of cells was observed in PPE2-immunized mice compared to unimmunized mice infected with Msmeg-PPE2 (Figure 4E). The rPPE2-immunized mice also exhibited significantly improved glucose tolerance as revealed by IPGT test (Figure 4F and G). In summary, rPPE2-immunized mice showed marked improvement in adipose tissue content, total body fat content, tissue pathology and glucose tolerance when challenged with infection, underscoring the role of PPE2 in modulating adipose tissue physiology and function.

5. PPE2 alters adipose tissue physiology and induces glucose intolerance in mice during infection with *M. tuberculosis*

The role of adipose tissue in regulating pulmonary pathology, inflammation and *M. tuberculosis* load in murine TB model is now well established (Ayyappan et al., 2019). Disruptions in the metabolic equilibrium within the adipose tissue and/or dysfunction of adipocytes during Mtb infection can contribute to the development of insulin resistance (Oswal et al., 2022). In several clinical studies, stress-induced hyperglycemia has been reported in tuberculosis patients which tend to normalize with proper management (Magee et al., 2018). Moreover, increased prevalence of impaired glucose tolerance and diabetes appears to be common among the TB patients (Oluboyo and Erasmus, 1990; Jeon et al., 2010; Mugusi et al., 1990; Raghuraman et al., 2014). However, the exact etiological mechanisms responsible for the development of glucose intolerance and insulin resistance in TB patients are not yet clearly understood.

In the previous sections, we demonstrated that the PPE2 protein of *M. tuberculosis* can

significantly remodel the adipose tissue and can cause significant impaired glucose tolerance

in mice. To further confirm its pathophysiological effects, we used a *ppe2*-null mutant of a

clinical Mtb strain CDC1551 (CDC1551-ΔPPE2) and compared its effect with the wild-type

strain (CDC1551) and a complemented strain where the genomic deletion of *ppe2* was

complemented with a functional copy of the *ppe2* gene (CDC1551-ΔPPE2::PPE2).

Accordingly, 9-10 weeks Balb/c mice were aerosol-infected either with CDC1551 (wild-

type) or CDC1551-ΔPPE2 (*ppe2*KO) or CDC1551-ΔPPE2::PPE2 and the infected mice were

sacrificed after 30- and 60-dpi. Expectedly, a significant loss of adipose tissue mass was

observed in both PG (Figure 5A and B) and VS (Figure 5C and D) fat depot in mice infected

with the wild-type CDC1551 and the *ppe2*-complemented strains as compared to uninfected

controls and mice infected with *ppe2*-null mutants (Figure 5A-D, Figure S5), suggesting

impairment of normal adipogenesis. Histological analyses revealed enhanced infiltration of

immune cells into both PG and VS adipose tissues of the mice infected with wild-type Mtb as

well as *ppe2*-complemented strain as compared to uninfected or *ppe2*KO strain-infected mice

at both time points (Figure 5E). While adipocyte hypertrophy was notably conspicuous in the

fat tissues of both the wild-type and *ppe2*-complmented infected mice, *ppe2*KO-infected mice

also exhibited some degree of hypertrophy in adipocytes as compared to those of uninfected

controls (Figure 5F-I).

Since increased infiltration of immune cells like macrophages, T-cells (Wu et al., 2007) into

the adipose tissue, have been implicated in adipocyte dysfunction and development of insulin

resistance (Apovian et al., 2008; Guilherme et al., 2008) and Mtb is known to promote

infiltration of these cells (Beigier-Bompadre et al., 2017), we next examined the degree of T-

cell infiltration in the adipose tissue in these infected mice using immunofluorescence

microscopy (Figure S6). Significantly increased CD3[+] T-cell infiltration was observed in PG

and VS adipose tissues of the mice infected with wild-type and *ppe2*-complemented Mtb

strains as compared to those observed in *ppe2*-KO strain-infected mice at both 30 and 60 dpi

(Figure 5J-M).

We also examined the changes in the expression of key genes associated with the

maintenance and regulation of adipose tissue, such as PPAR-γ and CD36 (a scavenger

receptor that is regulated by PPAR-γ and is involved in the uptake of long-chain fatty acids

and other lipids) (Maréchal et al., 2018; Chen et al., 2022) in the perigonadal fat tissue of

these infected mice. It was found that expression of both PPAR-γ and CD36 was significantly

downregulated in mice infected with wild-type and *ppe2*-complemented strains, in

comparison to mice infected with *ppe2*KO or uninfected control (Figure 5N and O). These

results reiterate our earlier observations and underscore that PPE2 interferes with

adipogenesis by suppressing the expression of adipogenic genes like PPAR-γ.

Since, adipose tissue was previously speculated to be as an important dormancy niche for

Mtb (Neyrolles et al., 2006; Ayyappan et al., 2019), we assessed PPE2's impact on the

survival of Mtb in adipose tissue particularly in perigonadal adipose tissue, for being a major

reservoir of fat in mice. It was found that mice infected with wild-type as well as *ppe2*-

complementated strains exhibited significantly higher bacterial load compared to *ppe2*KO

strain-infected mice (Figure 5P). These data suggest that PPE2 promotes persistence of Mtb

in adipose tissue.

We also assessed the impact of PPE2 on the rate of glucose clearance using IPGTT assay and

degree of insulin resistance using homeostatic model assessment of insulin resistance

(HOMA-IR) (Matthews et al. 1985), where the conversion factor was adapted from Knopp et

al. (2019). As expected, mice infected with the wild-type and *ppe2*-complemented strains

exhibited markedly higher glucose intolerance compared to uninfected control mice and those

infected with the *ppe2*KO strain at both 30 and 60 dpi, as indicated by blood glucose levels

and AUC at 30-minute intervals (Figure 5Q and R). At day 60 post infection, groups infected

with wild-type Mtb and *ppe2*KO strain complemented with a functional *ppe2* gene, showed

higher fasting blood glucose and HOMA-IR compared to the groups uninfected or infected

with *ppe2*KO strain (Figure 5S and T). Therefore, these data indicate that mycobacterial

infection can influence glucose homeostasis leading to the development of early

hyperglycaemia which can later lead to development of insulin resistance and PPE2 is one of

the potential etiological factors responsible for such metabolic perturbations.

6. PPE2 modulates the transcriptome of perigonadal adipose tissue

Results obtained from the experiments carried out in the earlier sections indicate that PPE2

can inhibit adipogenesis by modulating expressing of key adipogenic genes required for

adipogenesis. To further assess its impact on the global gene expression profile in the adipose

tissue, we performed RNA-seq in perigonadal adipose tissue collected from uninfected, wild-

type and *ppe2*KO-infected mice at 60 dpi. Compared to the uninfected, 3569 and 2441 genes

were found to be differentially expressed in wild-type and *ppe2*KO-infected mice adipose

tissue, respectively, while 1261 genes were differentially expressed in *ppe2*KO and wild-type

adipose tissue (Figure 6A). However, with ±1.3 log2 fold change cut-off, we observed 807

and 661 genes differentially expressed in wild-type and *ppe2*KO adipose tissue, respectively,

and 191 genes between *ppe2*KO and wild-type adipose tissues (Figure 6B and C). In

significantly modulated DEGs among *ppe2*KO and wild-type infected mice, 96 genes

overlapped with mice infected with wild-type Mtb, and 76 were unique DEGs in *ppe2*KO and

wild-type infected mice adipose tissues. To identify key signaling pathways influenced by

PPE2 in adipose tissue, we performed gene set enrichment analysis (GSEA) using gene

ontology, KEGG, and REACTOME pathway databases (Subramanian et al., 2005) using the

DEGs between wild-type- and *ppe2*KO-infected mice (Figure 6D and E). As expected, gene

sets associated with tissue-specific immune responses (chemokine and cytokine signaling)

were highly upregulated in mice infected with wild-type Mtb (Figure 6F and Figure S7),

while gene sets related to oxidative phosphorylation were found to be downregulated in these

mice (Figure 6F). Phospholipase C activity was found to be highly enriched in the adipose

tissue of wild-type CDC1551 compared to uninfected and *ppe2*KO-infected mice, as

represented by the genes *TXK, CD86, GPR55*, *Angiotensin*, and *LPAR1* (Figure 6G). Both

gene ontology and Reactome pathway analyses showed enrichment of genes related to

cytosolic ribosomal protein synthesis and ribosomal biogenesis in the adipose tissue of wild-

type Mtb-infected mice compared to that of *ppe2*KO-infected mice (Figure 6D, E and Figure

6H). Genes related to cytoplasmic translation of both small and large ribosomal subunits were

found to be enriched in the adipose tissue of CDC155-infected mice. In addition, KEGG

pathways analysis revealed enrichment of pathways related to prion disease, Alzheimer,

Parkinson's disease and other neurodegenerative pathways suggesting an increase in the

unfolded proteins in the endoplasmic reticulum (ER). ER stress is known to trigger

inflammation and lipolysis in adipocytes leading to development of dyslipidemia and insulin

resistance (Khan and Wang, 2014; Foley et al., 2021; Deng et al., 2012). To validate the

occurrence of ER stress, we performed real-time PCR analysis to assess the expression of ER

stress markers like ATF4, a transcription factor that regulates stress response and GRP78, a

chaperone which facilitates protein folding during unfolded protein response (UPR). It was

found that the expression of both ATF4 and GRP78 was significantly upregulated in the

perigonadal adipose tissue of wild-type Mtb-infected mice as compared to uninfected and

*ppe2*KO mice groups, confirming induction of ER stress in the adipose tissue (Figure 6I and

J).

8. PPE2 induce lipolysis during Mtb infection in adipocyte cell lines and in *vivo*.

Since, ER stress in adipose tissue is known to trigger lipolysis through activation of

cAMP/PKA pathway (Song et al., 2017; Deng et al., 2012), we next examined whether PPE2

can trigger lipolysis in mature 3T3-L1 adipocytes *in vitro* by measuring free glycerol released

in the medium due to hydrolysis of triglycerides (Schweiger et al., 2014). Treatment of

mature 3T3-L1 adipocytes with rPPE2 protein for 24 hours resulted in a dose-dependent

increase of free glycerol in the medium (Figure 7A). Infiltration of immune cells and

hypertrophy are known to stimulate production of pro-inflammatory cytokines such as TNF-

α, which in turn is also known to trigger lipolysis (Samad et al., 1999). To determine whether

PPE2-induced lipolysis occurs independently of TNF-α production, we treated mature 3T3-

L1 adipocytes with a TNF-α inhibitor in the presence of rPPE2. While the inhibitor was able

to effectively suppress TNF-α-induced free glycerol release, it did not significantly affect

lipolysis induced by rPPE2 (Figure 7B). Since insulin possess antilipolytic properties (Arner

et al., 1981), we next examined whether PPE2 can override antilipolytic effect of insulin and

it was found that insulin failed to suppress rPPE2-induced lipolysis in 3T3-L1 adipocytes

(Figure S8). Therefore, we speculated that PPE2 interferes with insulin's ability to activate

phosphodiesterase activity in adipocytes. To confirm this, we analysed the expression of

phosphodiesterase 3B (PDE3B), a critical enzyme that converts cyclic adenosine

monophosphate (cAMP) into 5'AMP and serves as a primary mediator of insulin's

antilipolytic effects (Kitamura et al., 1999; DiPilato et al., 2015). Semi-quantitative PCR

analysis revealed that PPE2 treatment could significantly downregulate PDE3B expression in

mature adipocytes (Figure 7C). Reduction in PDE3B is associated with elevated cAMP

levels, which activate cAMP-dependent protein kinase (PKA) and activated PKA

phosphorylates hormone-sensitive lipase (HSL), a key enzyme involved in lipolysis

(Degerman et al., 1998). To confirm the involvement of this pathway, we assessed HSL

phosphorylation levels in rPPE2-treated adipocytes at 6- and 24-hour post-treatment. The

results showed a significant increase in HSL phosphorylation, indicating enhanced enzymatic

activity (Figure 7D). Furthermore, pretreatment with H89-dihydrochloride, a PKA-specific

inhibitor, significantly reduced free glycerol release in rPPE2-treated adipocytes (Figure 7E).

Collectively, these data suggest that PPE2 promotes lipolysis in adipocytes by reducing

PDE3B expression, and activating PKA-HSL pathway through elevated cAMP levels.

Since PPE2 was found to promote lipolysis, we next measured serum non-esterified free fatty

acids (NEFA) levels in these mice. It was found that the mice infected with wild-type Mtb

and *ppe2*-complemented *ppe2*KO strains had higher NEFA content compared to those

infected with *ppe2*KO strain and uninfected control mice at 60 dpi (Figure 7F). From these

data, it appears that PPE2 contributes to sustain lipolytic activity and elevated NEFA levels in

circulation during mycobacterial infection. To further validate this hypothesis, we measured

the plasma NEFA content of TB patients (n = 51) and compared it those of healthy controls

(n = 14). The TB patients were found to have significantly higher NEFA content compared to

heathy controls (Figure 7G), suggesting presence of dyslipidaemia, an established risk factor

for development of insulin resistance.

Discussion

Association of tuberculosis with diabetes has been suspected for almost a century ago (Root, 1934). It is now well established that diabetic patients are at increased risk of developing tuberculosis. It has been reported that diabetic patients are at 3-fold increased risk of developing tuberculosis compared to those with normoglycemia (Jeon and Murray, 2008). TB-patients with diabetic conditions were found to have higher treatment failures as well as mortality compared to non-diabetic TB-patients (Dooley et al., 2009; Viswanathan et al., 2014). On the other hand, many TB patients with no family history of diabetes were found to develop insulin resistance (Magee et al., 2018; Lin et al., 2017). It has been found that about 8-87% of TB patients are newly diagnosed with diabetes without prior conditions (Mave et al., 2017; Magee et al., 2018). However, the exact etiological connections from Mtb that trigger development of diabetes are not well understood.

Although the adipose tissue was once thought to function as a passive fat storage site, it is now recognized as a dynamic endocrine organ that also plays a crucial role in regulating glucose homeostasis and lipid metabolism (Coelho et al., 2013). Dysregulation of adipose tissue functions and inflammatory conditions under certain pathophysiological disorders like obesity are known to promote insulin resistance and subsequently to type 2 diabetes. Interestingly, it has been shown that in active TB patients, Mtb can persist in the host adipose tissue (Kashyap et al., 2012; Neyrolles et al., 2006; Agarwal et al., 2014) and this persistence can modulate adipose tissue behavior (Erol, 2008; Neyrolles et al 2006; Beigier-Bompadre et al., 2017). Mycobacterial persistence in adipose tissue is known to cause adipocyte hypertrophy, lipolysis, change in physiology and infiltration of immune cells (Beigier-bompadre et al., 2017; Martinez et al., 2019; Bisht et al., 2023a). Therefore, occurrence of

hyperglycemia, insulin resistance and metabolic alterations during tuberculosis suggests

possible involvement of etiologic factors/proteins from Mtb in the development of pre-

diabetic/diabetic conditions.

Earlier, we found that PPE2, a member of the PE/PPE family, unique to pathogenic

Mycobacteria sp. is secreted into the circulation (Bisht et al., 2023b) which can translocate to

the nucleus, have pleiotropic functions and as a virulent factor promotes survival of Mtb

inside the host (Bisht et al., 2023b; Pal et al., 2021). Therefore, it can have a systemic effect

on the host and access various tissues including the adipose tissue during Mtb infection.

Moreover, adipose tissue is now recognized as a significant reservoir for Mtb (Ayyappan et

al., 2019; Neyrolles et al., 2006; Agarwal et al., 2014), it is possible that mycobacterial

secreted proteins can directly modulate adipocyte physiology and functions. Given its role in

mycobacterial survival and a pleiotropic protein, we speculated that PPE2 is one of the

potential etiological factors from Mtb which can interfere with adipocyte functions including

those governing lipid metabolism, insulin sensitivity, and inflammatory responses, ultimately

contributing to metabolic dysregulation during infection.

To test our hypothesis, we first examined whether PPE2 inhibits differentiation of adipocytes

in vitro using 3T3-L1 fibroblast cell line, since impaired adipogenesis is linked to insulin

resistance and reduced abdominal fat content in mice with high-fat diets (Gustafson et al.,

2015; Ji et al., 2014). It was found that recombinantly purified PPE2 protein could

significantly inhibit adipogenesis by suppressing expression of adipogenic transcription

factors PPAR-γ and C/EBP-α (Figure 1). PPAR-γ is regarded as the "master regulator" of

adipogenesis (Farmer, 2006; Park et al., 2016) and together with C/EBP-α, it regulates early

adipogenesis events, fatty acid transport and insulin signalling/sensitization (Wu et al., 2020;

LeBlanc et al., 2016; Gerstner et al., 2024). When a surrogate non-pathogenic bacterium *M.*

smegmatis expressing Mtb PPE2 was used to infect mice, significant reduction in the adipose

tissue mass was observed (Figure 2A, 2B, Figure S3) which was also accompanied by

marked decrease in expression of PPAR-γ and C/EBP-α (Figure 2C, 2D, 2E, 2F). Similar

results were also obtained when the mice were administered with recombinant PPE2 protein

(Figure 3A, 3B, 3C) which could be reversed by immunizing mice with rPPE2 (Figure 4A,

4B, 4C, 4D). In addition, downstream targets of PPAR-γ, such as fatty acid synthase (FAS)

and adiponectin, were similarly downregulated upon exposure to PPE2 both *in vivo* and *in*

vitro. These data suggest a broad suppression of the PPAR-γ signaling axis implicating its

possible involvement in tuberculosis-induced metabolic perturbations (Oswal et al., 2022).

PPAR-γ and C/EBP-α also regulate the expression of several makers associated with mature

adipocytes such as adiponectin, fatty acid-binding protein 4 (FABP4), glucose transporter

type 4 (GLUT4) etc. (Gustafson et al., 2015). Among the several isoforms of glucose

transporters, GLUT4 is primarily responsible for insulin-stimulated glucose uptake in major

tissues involved in glucose homeostasis such as adipocytes, muscles, heart and brain (Navale

and Paranjape, 2016). Thus PPE2-mediated inhibition of inhibition of adipogenesis and

repression of PPAR-γ axis, thereby contributes to limiting GLUT4-expressing mature

adipocytes and thereby impair insulin responsiveness and promote glucose intolerance as

observed in mice infected with Msmeg-PPE2 or wild-type Mtb or mice administered with

rPPE2 (Figures 2M, 5Q-R and 3F). While the Mtb aerosol model can be questioned for

bacterial load effects, it provides crucial *in vivo* validation that PPE2 function is relevant in

the context of mycobacterial infection.

Profound weight loss or wasting is a significant clinical feature associated with TB patients

which greatly influences disease severity and prognosis. It has been found that TB patients

undergo significant changes in the whole-body compositions with substantial depletion of fat

mass which could due to combination of several factors like less energy intake due to loss of

appetite, altered metabolism due to inflammatory responses (Paton and Ng, 2006). The

decrease in body fat is also correlated with TB-associated complications and has been linked

to exacerbated pulmonary pathology (Ayyappan et al., 2019). In this study, both recombinant

protein treatment and mice infection models showed significant loss in the adipose tissue

mass in both PG and VS fat depots in the presence of PPE2 (Figures 2, 3 and 5). In contrast,

mice immunized with rPPE2 protein did not show severe fat loss in both tissues (Figure 4).

Dual-energy X-ray absorptiometry (DEXA) analysis also confirmed that total body fat loss

was resisted in PPE2-immunized mice (Figure 4D). Although food intake was not

quantitatively assessed, we speculate that neutralization of PPE2 by immunization played a

key role in preserving the adipose tissue. Fat loss during TB is likely driven by chronic

inflammation in the adipose tissue and production of TNF-α, IL-2, IL-1β and IFN-γ during

mycobacterial infection (Kim et al., 2011; Beigier-Bompadre et al., 2017) some of which

were also found to be upregulated in the adipose tissue transcriptome of the mice infected

with wild-type Mtb strain (Figure 6).

Adipocyte dysfunction is often characterized by adipocyte hypertrophy, increased oxidative

stress, chronic low-grade inflammation and reduced ability to generate new adipocytes from

its precursor pre-adipocytes (Klöting and Blüher, 2014; Reyes-Farias et al., 2021). Under

normal physiological conditions, the major roles of adipose tissue include insulin-stimulated

glucose uptake and storage of excess energy in the form of triglyceride, in addition to its

endocrine functions like secretion of hormones like leptin, resistin and various other

cytokines (Scheja and Heeren, 2019). When adipogenesis is impaired, it often leads to

development of insulin resistance (Hammarstedt et al., 2018). This is because the

differentiating preadipocytes buffer excess circulating free fatty acids by uptaking and

converting them into triglycerides. However, when adipogenesis is restricted, the buffering

capacity is hindered, the excess free fatty acids are ectopically deposited in other tissues like

muscle and liver, the major sites for insulin-stimulated glucose homeostasis and also get

deposited in kidney and heart (Hocking et al., 2013; Morigny et al., 2016), leading to

development of insulin resistance in peripheral tissues. Adipocyte hypertrophy often leads to

immune cell infiltration into the adipose tissue, leading to induction of oxidative stress,

NLRP3 inflammasome activation and subsequent production of pro-inflammatory cytokines

which further trigger the development of local and systemic inflammation and insulin

resistance (Kursawe et al., 2016; Susca et al., 2024). We found that PPE2 can promote

immune cell infiltration in adipocytes, including T-cells. Immune cell infiltration into the

adipose tissue is widely implicated as an important link in the development of insulin

resistance during obesity by causing adipose tissue dysfunction (Guzik et al., 2017).

Infiltration of Mtb-specific T-cells and NK cells into perigonadal adipose tissue was found to

be associated with distinct gene expression profiles and elevated interferon-gamma signaling,

indicating localized tissue inflammation (Beigier-Bompadre et al., 2017). Infiltration of

immune cells is known to promote lipolysis, leading to elevated circulating free fatty acids,

and thereby contributing to peripheral insulin resistance (Beigier-Bompadre et al., 2017;

Guilherme et al., 2008). We found that PPE2 also inhibit the expression of adiponectin in

mice adipose tissue, either when administered as a pure protein or infected with *M.*

smegmatis expressing PPE2. Adiponectin, a hormone secreted by the adipose tissue, plays a

crucial role in various physiological functions including lipid metabolism, energy

homeostasis, inflammation and insulin sensitivity (Khoramipour et al., 2021). Adiponectin is known to inhibit inflammation in various tissues including adipose tissue and found to be protective against diabetes (Ouchi and Walsh, 2007; Siitonen et al., 2011; Lindsay et al., 2002; Ziemke and Mantzoros, 2010). Adiponectin also enhances glucose tolerance in a diabetic rat model where a single dose of adiponectin resulted in significant reduction in the blood glucose levels (Tishinsky et al., 2012). Adiponectin also improves insulin sensitivity through several direct and indirect mechanisms like decreasing triglyceride content of the adipose tissue, activation of AMPK cascade to stimulate beta-oxidation of fatty acids and PPAR-α phosphorylation (Katira and Tan, 2016). Therefore, inhibition of adiponectin expression by PPE2 appears to be another mechanism by which it can promote glucose intolerance and subsequent insulin resistance.

A role of PPE2 in modulation of adipose tissue physiology was further confirmed by using a *ppe2*-null mutant (*ppe2*KO) of a clinical strain of *M. tuberculosis* (CDC1551). The observed suppression of expression of adipogenic genes like PPAR-γ and CD36 gene in the adipose tissue of mice infected with wild-type *M. tuberculosis* (CDC1551) was rescued in mice infected with *ppe2*KO. Functional complementation of *ppe2*KO with *ppe2* gene reverted this phenotype is similar to that of mice infected with wild-type strain. Also, these mice showed exacerbated adipose tissue pathology compared to the *ppe2*-null mutant which had almost normal adipose tissue histology similar to the uninfected control mice. In addition to impaired glucose tolerance, mice infected with wild-type as well as *ppe2*KO complemented with *ppe2* gene showed pronounced insulin resistance at day 60 dpi as determined HOMA-IR. All the experiments like *in vitro* cell line model with recombinantly purified PPE2 protein (rPPE2) Figure 1); *in vivo* mice infection model using *M. smegmatis* expressing Mtb PPE2 (Figure 2), *in vivo* mice treatment with rPPE2 (Figure 3); rescuing of adverse effects of PPE2 in mice

pre-immunized with rPPE2 (Figure 4) and infection of mice with wild-type *versus ppe2* null-

mutant (*ppe2* KO) and complemented Mtb strains (Figure 5) clearly demonstrated that

observed effects of PPE2 on adipose tissue physiology (which include impaired glucose

homeostasis, lipolysis as well as changes in gene expression) are due to the pathological

effects of PPE2 but not merely due to differences in bacterial burden.

To the best of our knowledge for the first time our data indicate that Mtb PPE2 is one of the

etiological factors which is involved in triggering adipocyte dysfunction, hyperglycemia as

well as insulin resistance observed in TB patients. PPE2 appears to exert its effects through

multiple mechanisms which are possibly interconnected. PPE2 interferes with adipogenesis

by inhibiting PPAR-γ and C/EBP-α, promotes adipocyte hypertrophy, inhibits expression of

adiponectin and promotes infiltration of immune cells and lipolysis which are orchestrated

together to induce glucose intolerance and eventually insulin resistance. In addition, PPE2

was also found to modulate expression of a large number of genes in the adipose tissue of

infected mice belonging to several classes of genes performing wide ranges of physiological

functions which include tissue specific immune responses, phospholipase C activity,

ribosomal biogenesis, as well as proteins involved in neurodegenerative disorders. The ability

of PPE2 to modulate such large number of genes is not so surprising, as earlier we have

shown that PPE2 mimics several features of a eukaryotic transcription factor. PPE2 was

found to possess a functional nuclear localization signal (NLS) and a DNA-binding domain,

allowing it to translocate to the nucleus and bind to GATA-1 like elements (Bhat et al., 2017;

Pal et al., 2021). In addition, PPE2 was also found to contain a SRC homology 3 motif (SH3)

(Srivastava et al., 2019), a protein-protein interaction domain, characteristic of several

kinases involved in cellular signaling (Kurochkina and Guha, 2013). Therefore, PPE2 acting

as a eukaryotic transcription factor is likely to have a profound influence on the transcriptome

as well as the protein-protein interactome of the adipose tissue.

Although we did not explore the exact molecular mechanisms by which PPE2 modulate the

expressions of adipogenic transcription factors and adiponectin, we delineated the possible

mechanisms by which PPE2 promote lipolysis which can trigger development of insulin

resistance by promoting adipose tissue inflammation (Morigny et al., 2016). PPE2 was found

to trigger ER stress in the adipose tissue, which in turn activate cAMP/PKA leading to

increased activation of HSL leading to increased lipolysis of adipocyte triglycerides.

Increased lipolysis is likely to elevate circulating NEFA, and we found that TB patients had

significantly higher NEFA levels in their serum. This confirms that PPE2-mediated

dyslipidaemia is a clinical hallmark of TB-associated metabolic perturbance. Interestingly,

pre-immunization of mice with rPPE2 was able to counter the adverse effects of PPE2 in

mouse models of infection used in this study. Therefore, the development of PPE2 subunit

vaccine may constitute an effective strategy to clinically manage TB-associated insulin

resistance and exacerbated diabetic complications in TB patients.

Materials and methods

Ethical statement and sample collection

All animal experimental protocols were approved by the Institutional Animal Ethics

Committee of Centre for DNA Fingerprinting and Diagnostics (CDFD), Hyderabad, India.

Balb/c mice were maintained at the animal house facility of CDFD, Hyderabad, under

standard conditions. The study involving human sample collection was approved by the

Institutional Ethical Committee for Biomedical Research, at Bhagwan Mahavir Medical

Research Centre (BMMRC) (No 941/BMMRC/2020/IEC). The study included 51 active

pulmonary TB patients and 14 BCG-vaccinated healthy controls with no prior TB history.

About 5 ml peripheral blood was collected, plasma was separated and stored at –80 °C until

further use. The control group (8 males, 6 females; age 16–40) and TB group (39 males, 12

females; age 18–75) were all HIV-negative. All the samples were collected after obtaining

written informed consents from the study participants. Active TB was diagnosed using smear

microscopy, chest radiography, molecular tests, and clinical evaluation, following NTEP

guidelines (India).

Reagents

Antibodies and reagents used in this study are anti-phospho-HSL (Cell Signaling

Technology, USA), anti-HSL antibody (St Jhon Laboratory, USA), anti-β-actin antibody

(Cell Signaling Technology, USA), TNA-α inhibitor (654256-M, Millipore, USA), H89

dihydrochloride (H89, Santa Cruz, USA), NEFA Kit (FUJIFILM, Japan). Free glycerol

reagent was purchased from Sigma-Aldrich, USA (F6428-40mL).

Purification of recombinant PPE2 protein

PPE2 protein was purified as described by us earlier (Bhat et al., 2017). Sequence

information is provided in Table 1. For affinity purification of the recombinant PPE2 protein,

the CDS was amplified from BAC-Rv329 (A5) using a forward primer (5′-

CGAGAGCTCATGACCGCCCCGATCTGGATGG-3′) and a reverse primer (5′-

GCAGAATTCTCACTCCACCCGGGTCGCTGAGT-3′) and cloned into the *Sac*I and

*Eco*R1 site of a T7 polymerase driven *Escherichia coli* expression vector pRSET A

(Invitrogen, Carlsbad, USA) in frame with an N-terminal histidine tag to facilitate

purification using metal affinity purification resins. The pRSET A vector containing the

PPE2 gene was then transformed into *Escherichia coli* BL21 (DE3) pLysS cells and a

primary inoculum was prepared from a single transformed colony. The primary culture was

inoculated into 800 ml terrific broth in the presence of chloramphenicol (35 μg/ml) and

ampicillin (100 μg/ml). The culture was grown in a shaker incubator at 37°C till the

absorbance reached to about 0.5. At this stage, expression of the recombinant protein was

induced by adding IPTG (Isopropyl β- d-1-thiogalactopyranoside, Sigma-Aldrich, USA) to a

final concentration of 1 mM and incubation was continued further for 16 hours at 18°C in a

shaker incubator to favor soluble expression of the protein. Cells were then harvested by

centrifugation and the pellets were suspended in lysis buffer (PBS containing 5% glycerol,

0.3% sodium lauroyl sarcosine). The cell suspension was then sonicated and the lysate was

centrifuged at 12000 rpm for 30 minutes. The soluble recombinant protein present in the

supernatant was allowed to bind with the TALON resin (Clontech, USA) for an hour, loaded

into a column and washed with washing buffer (PBS containing 5% glycerol and 20 mM

Imidazole). The resin-bound protein was eluted using elution buffer (PBS containing 5%

glycerol and 200 mM imidazole). Eluted samples were loaded onto an SDS-PAGE gel to

confirm the presence of the recombinant PPE2 protein. The PPE2-positive fractions were

pooled together and dialyzed against PBS containing 5% glycerol to remove imidazole from

the protein samples. The concentration of the purified protein was estimated using

MicroBCA^{TM} Protein Assay Kit from Thermo Scientific (Rockford, USA) following

manufacturer's instructions. To remove LPS contamination, the recombinant proteins were

treated with 10% (v/v) polymyxin B-agarose (Sigma-Aldrich, USA; binding capacity 200 –

500 μg of LPS/ml) for 1 hour at 4°C as described by us earlier (Bhat et al., 2012).

Cloning and expression of *ppe2* gene of *M. tuberculosis* in *M. smegmatis* bacteria

The non-pathogenic *M. smegmatis* mc^{2}155 bacteria were received from ATCC and were

grown in Middlebrook 7H9 medium (HiMedia Laboratories, India) supplemented with 0.05%

Tween-80 (Sigma-Aldrich, USA) and 10% Albumin-dextrose-catalase (ADC) (HiMedia

Laboratories, India), For generating *M. smegmatis* expressing PPE2 protein of *M.*

tuberculosis (Msmeg-PPE2), *ppe2* was cloned in a mycobacterial shuttle vector pVV16 into

Nde I – *Pst* I sites for expression in *M. smegmatis* as described earlier (Bisht et al., 2023b).

Transformation of pVV16-*ppe2* construct into *M. smegmatis* (Msmeg-PPE2) was carried out

as described earlier (Bhat et al., 2017). *M. smegmatis* harboring the empty vector (Msmeg-

pVV16) alone was used as control.

Culture of bacterial strains

The DH5α strain of *E. coli*, used for cloning was grown in Luria-Bertani broth (Becton

Dickinson, Difco Laboratories). Wild-type *M. tuberculosis* strain (CDC1551) and *ppe2*

deleted mutant *M. tuberculosis* strain (CDC1551-ΔPPE2) and complemented strain

(CDC1551-ΔPPE2::PPE2) were maintained as described earlier (Bisht et al., 2023b) in

ABSL3/BSL3 facility of the University of Delhi South Campus (UDSC), Delhi, India.

Mycobacterial strains were grown in Middlebrook 7H9 Broth supplemented with 10%

albumin, dextrose, catalase, and NaCl (HiMedia Laboratories) along with 0.2% glycerol

(Sigma-Aldrich, USA) and 0.05% Tween-80 (Sigma-Aldrich, USA).

Differentiation of 3T3-L1 preadipocytes into mature adipocyte cells

3T3-L1 fibroblast cells were differentiated into adipocytes using a differentiation cocktail

containing dexamethasone, insulin and IBMX. In brief, 3T3-L1 cells (obtained from ATCC)

were cultured in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10%

fetal bovine serum and antibiotics (all from Invitrogen USA) and were maintained in an

incubator at 37°C and 5% of CO_2. At confluence, preadipocytes were induced to differentiate

into adipocytes by culturing them in differentiation cocktail (DMI) comprising of 1 μM

dexamethasone (Sigma-Aldrich, USA), 0.5 mM isobutylmethylxantine (Sigma-Aldrich,

USA) and 10 μg/ml bovine insulin (Sigma-Aldrich, USA). On day 2, the medium was

replaced with DMEM with 10% fetal bovine serum and insulin (10 μg/ml). From day 4

onwards, medium was replaced with 10% FBS in DMEM and antibiotics but without insulin,

and this medium was changed every 2 days up to day 8 after confluence, when ~95% of the

cells were differentiated to adipocytes.

Measurement of degree of differentiation of 3T3-L1 cells in the presence of rPPE2

The degree of differentiation was measured by staining with Oil Red O as described by Wu et

al. (2007). 3T3-L1 cells were differentiated either in the absence or presence of different

concentrations of recombinantly purified PPE2 protein of *M. tuberculosis* (rPPE2). On day

10, the cells were fixed with 10% paraformaldehyde for one hour and subsequently washed

with distilled water. The fixed cells were stained with 0.7% solution of Oil Red O stain

(Sigma-Aldrich, USA) in isopropanol, followed by multiple washes with distilled water. To

quantify Oil Red O stain retained by the lipid droplets present in matured 3T3-L1 cells, the

dye was extracted using isopropanol and the absorbance was measured at 490 nm using a

spectrophotometer. Also, total RNA was isolated from differentiating 3T3-L1 adipocyte

culture medium containing rPPE2 (3 μg/ml) at various time points and adipogenic markers

like PPAR-γ, C/EBP-α, fatty acid synthase (FAS) expression was measured by reverse

transcription polymerase chain reaction (RT-PCR) using gene-specific primers.

.

Effect of PPE2 on adipose tissue physiology *in vivo*

Balb/c mice of 9-11 weeks of age were infected with 100 million of either *M. smegmatis*

carrying empty vector pVV16 (Msmeg-pVV) or *M. smegmatis* expressing PPE2 (Msmeg-

PPE2) through intravenous route and were sacrificed on 7[th] day post-infection. The

perigonadal and visceral fat tissue were harvested for measuring the weight and also used for

histopathology and extraction of total RNA. Histopathology was carried out by hematoxylin

and eosin (H&E) staining.

RNA isolation and RT-PCR and semi-quantitative reverse transcriptase PCR

Total RNA was extracted from 3T3-L1 or perigonadal and visceral fat tissue using trizol

following the manufacturer's protocol (Invitrogen, USA). After converting the total RNA

into cDNA, the levels of expression of PPAR-γ, C/EBP-α, FAS, CD36 and Adiponectin was

measured by qPCR. GAPDH was used as an internal control. The primers used are as

follows; GAPDH (F 5'ACAACTTTGTCAAGCTCATTTCCT3' and R

5'GATAGGGCCTCTCTTGCTCA3'), PPAR-γ (F 5'ATTCTCAGTGGAGACCGC3' and

5'AGCAGGGGGTGAAGGCTC3'), C/EBP-α (F 5'GAGAATGGGGGCACCACC3' and R

5'GCAACAGCGGCTGGCGAC3'), FAS (F 5'ATCCCCGGCTGCCCCATG3' and R

5'CCGAAGCCAGTGCTCGCT3'), CD36 (F 5' TGGCCTTACTTGGGATTGG3' and R

5'CCAGTGTATATGTAGGCTCATCCA3') and Adiponectin (F

5'GCCTGTCCCCATGAGTAC3' and R 5'TCTTCGGCATGACTGGGC3'). Also, the levels

of expression of ATF4 (F 5'CATGCCAGATGAGCTCTTGA3' and R

(5'CAAGAATGTAAAGGGGGCA3') and GRP78 (F 5'AGATCTTCTCCACGGCTTCC3'

and R (5'GGAGCAGGAGGAATTCCAGT3') in perigonadal fat tissue were checked where

β-actin (F 5'3GGCTGTATTCCCCTCCATCG' and R

5'CCAGTTGGTAACAATGCCATGT3') was used as internal control. PDE3B level in 3T3-

L1 was measured by semi-quantitative reverse transcriptase PCR using the following primer;

PDE3B (F 5' ATTGAGTGGCAGAACCAGTTTCC3' and R 5'

CTGCTGCGATCCCACCTTGAACA3').

Protein treatment in mice

Balb/c mice of 9-11 weeks old mice were administered with either bovine serum albumin

(BSA) or rPPE2 intravenously (3 mg/kg) on alternate days for 12 days. On day 14, glucose

tolerance test was performed using standard protocol as described in the following section.

Mice were sacrificed on day 15th and perigonadal and visceral adipose tissue weights were

measured. Also, perigonadal and visceral tissues were harvested for histopathology analyses

using H&E staining and RNA isolation was carried out for measuring the expression of

PPAR-γ, C/EBP-α and Adiponectin.

Histopathology

For histopathology analyses, tissue samples were fixed with 4% paraformaldehyde and

embedded in paraffin wax, and then sectioned (4~5 µm) for hematoxylin and eosin (H&E)

staining. For this, the sections were deparaffinized and hydrated in distilled water and stained

with Hematoxylin (30 seconds), washed in distilled water, and counterstained with Eosin (1%

in distilled water for 30 seconds). Slides were washed, dehydrated, and mounted on slides for

imaging. Tissue section images were acquired using Nikon ECLIPSE Ni-U light upright

microscope.

Intraperitoneal glucose tolerance test (IPGTT)

Balb/c mice were starved overnight and an intraperitoneal injection of glucose was given at a

dose of 2 g/kg as 25% glucose solution in sterile saline. Blood glucose levels were monitored

from the tail tip vein using a glucometer at 30, 60, 90 and 120 minutes post-injection. The

data plotted as percent of initial glucose concentration versus time (Veloso et al., 2019;

Martinez et al., 2019). The area under the curve was calculated using the trapezoid method

using Graph Pad Prism (GraphPad Software, USA).

Immunization of mice with rPPE2 and analysis of fat tissue distribution

Balb/c mice were immunized subcutaneously with 100 μg of rPPE2 protein in incomplete

Freund's adjuvant. Subsequently, three booster doses (100 μg rPPE2/mouse) were

administered subcutaneously in incomplete Freund's adjuvant at 15 days intervals. Sera were

tested for the presence of PPE2-specific antibody by ELISA. These immunized Balb/c mice

were infected with 100 million Msmeg-pVV/Msmeg-PPE2 through intravenous route and

were sacrificed at 7[th] day post-infection. Total body fat percentage was measured by a Dual-

energy X-ray Absorptiometry machine (Discovery QDR series, Hologic, USA). After

sacrificing the mice, the perigonadal and visceral fat content was measured by weighing the

tissues and harvested for histopathology by H&E staining.

M. tuberculosis **infection**

Balb/c mice of 9-11 weeks of age were aerosol-infected with approximately 200-300 cfu per

mice of either CDC1551 (wild-type *M. tuberculosis)* or CDC1551-ΔPPE2 *(ppe2*KO*) or*

CDC1551-ΔPPE2::PPE2 *(ppe2*-complemented *ppe2*KO strain) and sacrificed at 30 days post-

infection (dpi) and 60 dpi respectively. The perigonadal and visceral fat tissue were harvested

and the weight of these tissues was measured. The histopathology of perigonadal and visceral

fat tissue was performed using hematoxylin and eosin (H&E) staining. Additionally, RNA

was extracted from perigonadal and visceral fat tissue using Trizol following the

manufacturer's protocol (Invitrogen, USA) and the levels of expression of PPAR-γ and CD36

were measured by qPCR. GAPDH was used as control**.** Perigonadal adipose tissue from mice

sacrificed at day 60 post-infection was used for mRNA sequencing study.

Transcriptome analysis

Total RNA was extracted from the perigonadal fat tissues from uninfected, wild-type- and

*ppe2*KO CDC1551-infected mice using Direct-ZOL Mini Prep Kit (Zymo Research) as per

the manufacturer's protocol. RNA quality and quantity were assessed using a NanoDrop

spectrophotometer and Agilent Tape station with high-sensitivity RNA ScreenTape. Paired-

end RNA-sequencing libraries were prepared using NEBNext® Ultra™ II Directional RNA

Library Prep Kit for Illumina (NEB) as per the manufacturer's protocol. The libraries were

purified using AMPureXP beads and analyzed on 4200 Tape Station system (Agilent

Technologies) using high-sensitivity D1000 Screen tape as per the manufacturer's protocol.

The sequencing was performed on an Illumina MiSeq platform. The raw reads were

processed using Trimmomatic v0.39 to remove adaptor sequences, ambiguous reads (reads

with unknown nucleotide 'N' larger than 5%), and low-quality sequences (reads with more

than 10% quality threshold (QV) < 25phred score). High-quality reads with a minimum

length of 100 nucleotides (nt) after trimming were retained. These high-quality reads

(QV>25) were mapped on the mouse reference genome (GRCm39.110) obtained from

ENSEMBLE using STAR (v2.7.10a) using default parameters. FeatureCounts (version 2.0.3)

was used to count the number of reads mapped to each gene. Differential gene expression

analysis was performed using the DESeq2 R package between the groups. Log2 Fold Change

values greater than zero were considered upregulated, while those less than zero were

downregulated, along with a P-value threshold of 0.05 for statistically significant results.

Finally, the gene set enrichment analysis (GSEA) was performed using software GSEA

(4.3.3) from the UC San Diego and Broad Institute website following the standard data

processing pipeline (GSEA (gsea-msigdb.org)) (Subramanian et al., 2005; Mootha et al.,

2003)

Colony forming unit (CFU) count

For CFU count, perigonadal adipose tissue from *M. tuberculosis*-infected mice was

homogenized in saline solution and plated on 7H11 plates supplemented with 10% OADC

(HIMEDIA, India). Plates were next incubated at 37°C and the colonies were counted after

30-35 days post-plating.

NEFA estimation

Non-esterified free fatty acids (NEFA) were analyzed in human and mouse sera samples

using LabAssay™ NEFA (FFA) kit from FUJIFILM, Japan, following the manufacturer's

protocols.

Western blotting

3T3-L1 cells were lysed using RIPA buffer (50 mM Tris-HCl [pH 8], 150 mM NaCl, 2 mM

EDTA, 1% NP-40, 0.5% Sodiumdeoxycholate, 0.1% SDS and protease inhibitors) and the

proteins were resolved on SDS-PAGE followed by Western blotting using specific primary

antibody and appropriate combinations of HRP (horseradish peroxidase) conjugated

secondary Ab (Sigma-Aldrich, USA). Bound HRP enzyme was detected by

chemiluminescence following the manufacturer's protocol (GE Healthcare, UK).

Measurement of lipolysis

Lipolysis in mature 3T3-L1 adipocytes was measured by estimating free glycerol released in

the medium using Free Glycerol Reagent Kit (F6428-40mL, Sigma-Aldrich, USA) following

the manufacturer's protocol. Briefly, 10 µl of culture medium was added to 800 µl of free

glycerol reagent and kept at 37□ for 5 minutes. The reaction results in production of a

quinoneimine dye which was measured at 540 nm using a spectrophotometer. The results

were expressed as µg of glycerol released per mg protein.

Statistical analysis

846 GraphPad Prism software, version 8.0 was used for determining the significance of the

847 difference in the mean between the samples. Student t-test was performed to calculate

848 significance between two samples. $p < 0.05$ was considered to be significant.

Acknowledgement

We thank UDSC-ABSL3/BSL3 facility for providing access to the BSL3 facility at the University of Delhi South Campus for the work related to the use of the pathogenic strain of *Mycobacterium tuberculosis*. We thank Dr Garima Khare from UDSC and her team members, Dr Prachi Nagpal and Dr Nupur Angrish, for their assistance during the mouse aerosol infection. We also thank Dr Sandeep Kumar Kotturu and Dr. N.V. Sanjeev Kumar for kind help. The authors gratefully acknowledge the financial support by the Science and Engineering Research Board (SERB), Department of Science and Technology (DST), Government of India (JCB/2021/000035), Council of Scientific and Industrial Research (CSIR), Govt. of India (37WS(0020)/2023-24/EMR-II/ASPIRE), Department of Biotechnology (DBT), Govt of India (BT/PR51149/MED/29/1660/2023) to SM and SG, Indian Council of Medical Research (ICMR), Govt. of India (2021-10087/GTGE/ADHOC-BMS and IIRPSG-2024-01-01453) and a core grant from CDFD by DBT. VM was supported by the DST-INSPIRE fellowship, Govt. of India.

Declaration of interests

A patent application based on the results described in this paper is being filed by CDFD, Hyderabad, in which S.M. M.K.B and S.G. are listed as inventors.

Author contributions

M.K.B., S.G. and S.M. conceived, designed the experiments, and analyzed the data. M.K.B. performed all the experiments. V.M. and P.D. helped in 3T3-L1 and mouse infection

872 experiments. V.L.V. provided healthy control and TB patients' plasma samples. M.K.B., S.G.

873 and S.M. wrote the manuscript. S.G. and S.M. corrected and edited the manuscript.

References

Abraham, P.R., Latha, G.S., Valluri, V.L., and Mukhopadhyay, S. (2014). *Mycobacterium*
tuberculosis PPE protein Rv0256c induces strong B cell response in tuberculosis patients.
Journal of molecular epidemiology and evolutionary genetics in infectious diseases. **22**, 244-
249. 10.1016/j.meegid.2013.06.023.
Agarwal, P., Khan, S. R., Verma, S. C., Beg, M., Singh, K., Mitra, K., Gaikwad, A. N., Akhtar, M. S.,
and Krishnan, M. Y. (2014). *Mycobacterium tuberculosis* persistence in various adipose depots
of infected mice and the effect of anti-tubercular therapy. *Microbes and Infection* **16**, 571-80.
10.1016/j.micinf.2014.04.006.
Ai, J.W., Ruan, Q.L., Liu, Q.H., and Zhang, W.H. (2016). Updates on the risk factors for latent
tuberculosis reactivation and their managements. *Emerging Microbes & infections* **5**, e10.
10.1038/emi.2016.10
Alebel, A., Wondemagegn, A.T., Tesema, C., Kibret, G.D., Wagnew, F., Petrucka, P., Arora, A.,
Ayele, A.D., Alemayehu, M., and Eshetie, S. (2019). Prevalence of diabetes mellitus among
tuberculosis patients in Sub-Saharan Africa: a systematic review and meta-analysis of
observational studies. *BMC Infectious Diseases*. **19**, 254. 10.1186/s12879-019-3892-8.
Apovian, C.M., Bigornia, S., Mott, M., Meyers, M.R., Ulloor, J., Gagua, M., McDonnell, M., Hess,
D., Joseph, L., and Gokce, N. (2008). Adipose macrophage infiltration is associated with
insulin resistance and vascular endothelial dysfunction in obese subjects. *Arteriosclerosis,*
Thrombosis, and Vascular Biology. **28**, 1654–1659. 10.1161/ATVBAHA.108.170316.
Arner, P., Bolinder, J., Engfeldt, P., and Ostman, J. (1981). The antilipolytic effect of insulin in
human adipose tissue in obesity, diabetes mellitus, hyperinsulinemia and starvation.
Metabolism **30**, 753–760. 10.1016/0026-0495(81)90020-2.
Ayyappan, J.P., Ganapathi, U., Lizardo, K., Vinnard, C., Subbian, S., Perlin, D.S., and Nagajyothi,
J.F. (2019). Adipose tissue regulates pulmonary pathology during TB infection. *mBio* **10**,
e02771-18. 10.1128/mBio.02771-18.
Beigier-Bompadre, M., Montagna, G.N., Kühl, A.A., Lozza, L., Weiner, J., Kupz, A., Vogelzang, A.,
Mollenkopf, H.J., Löwe, D., Bandermann, S., Dorhoi, A., Brinkmann, V., Matuschewski, K.,
and Kaufmann, S.H.E. (2017). *Mycobacterium tuberculosis* infection modulates adipose tissue
biology. *PLoS Pathogens*. **13**, e1006676. 10.1371/journal.ppat.1006676.
Bhat, K.H., Das, A., Srikantam, A., and Mukhopadhyay, S. (2013). PPE2 protein of *Mycobacterium*
tuberculosis may inhibit nitric oxide in activated macrophages. *Annals of the New York*
Academy of Sciences. **1283**, 97–101. 10.1111/nyas.12070.

Bhat, K.H., Srivastava, S., Kotturu, S.K., Ghosh, S., and Mukhopadhyay, S. (2017). The PPE2 protein of *Mycobacterium tuberculosis* translocates to the host nucleus and inhibits nitric oxide production. *Scientific Reports.* **7**, 39706. 10.1038/srep39706.

Bisht, M.K., Dahiya, P., Ghosh, S., and Mukhopadhyay, S. (2023a). The cause-effect relation of tuberculosis on incidence of diabetes mellitus. *Frontiers in Cellular and Infection Microbiology.* **26**, 13:1134036. 10.3389/fcimb.2023.1134036.

Bisht, M.K., Pal, R., Dahiya, P., Naz, S., Sanyal, P., Nandicoori, V.K., Ghosh, S., and Mukhopadhyay, S. (2023b). The PPE2 protein of *Mycobacterium tuberculosis* is secreted during infection and facilitates mycobacterial survival inside the host. *Tuberculosis (Edinb.)* **143**, 102421. 10.1016/j.tube.2023.102421.

Chen, Y., Zhang, J., Cui, W., and Silverstein, R.L. (2022). CD36, a signaling receptor and fatty acid transporter that regulates immune cell metabolism and fate. *The Journal of Experimental Medicine.* **219**, e20211314. 10.1084/jem.20211314.

Coelho, M., Oliveira, T., and Fernandes, R. (2013). Biochemistry of adipose tissue: an endocrine organ. *Archives of medical science.* **9**, 191–200. 10.5114/aoms.2013.33181

Degerman, E., Landström, T.R., Wijkander, J., Holst, L.S., Ahmad, F., Belfrage, P., and Manganiello, V. (1998). Phosphorylation and activation of hormone-sensitive adipocyte phosphodiesterase type 3B. *Methods* **14**, 43–53. 10.1006/meth.1997.0564.

Deng, J., Liu, S., Zou, L., Xu, C., Geng, B., and Xu, G. (2012). Lipolysis response to endoplasmic reticulum stress in adipose cells. *The Journal of Biological Chemistry.* **287**, 6240–6249. 10.1074/jbc.M111.299115.

DiPilato, L.M., Ahmad, F., Harms, M., Seale, P., Manganiello, V., and Birnbaum, M.J. (2015). The role of PDE3B phosphorylation in the inhibition of lipolysis by insulin. *Molecular and Cellular Biology.* **35**, 2752–2760. 10.1128/MCB.00422-15.

Dooley, K.E., Tang, T., Golub, J.E., Dorman, S.E., and Cronin, W. (2009). Impact of diabetes mellitus on treatment outcomes of patients with active tuberculosis. *The American Journal of Tropical Medicine and Hygiene.* **80**, 634–639.

Erol, A. (2008). Visceral adipose tissue specific persistence of *Mycobacterium tuberculosis* may be reason for the metabolic syndrome. *Medical Hypotheses.* **71**, 222–228. 10.1016/j.mehy.2008.03.028.

Farmer, S.R. (2006). Transcriptional control of adipocyte formation. *Cell Metabolism.* **4**, 263–273. 10.1016/j.cmet.2006.07.001.

Foley, K.P., Chen, Y., Barra, N.G., Heal, M., Kwok, K., Tamrakar, A.K., Chi, W., Duggan, B.M., Henriksbo, B.D., Liu, Y., and Schertzer, J.D. (2021). Inflammation promotes adipocyte lipolysis via IRE1 kinase. *The Journal of Biological Chemistry.* **296**, 100440. 10.1016/j.jbc.2021.100440.

Franco, J. V., Bongaerts, B., Metzendorf, M. I., Risso, A., Guo, Y., Peña Silva, L., Boeckmann, M., Schlesinger, S., Damen, J. A., Richter, B., Baddeley, A., Bastard, M., Carlqvist, A., Garcia-Casal, M. N., Hemmingsen, B., Mavhunga, F., Manne-Goehler, J., and Viney, K. (2024). Diabetes as a risk factor for tuberculosis disease. *The Cochrane Database of Systematic Reviews.* **8**, CD016013. 10.1002/14651858.CD016013

Gautam, S., Shrestha, N., Mahato, S., Nguyen, T.P.A., and Mishra, S.R. (2021). Diabetes among tuberculosis patients and its impact on tuberculosis treatment in South Asia: a systematic review and meta-analysis. *Scientific Reports.* **11**, 2113. 10.1038/s41598-021-81057-2.

Gerstner, M., Heller, V., Fechner, J., et al. (2024). Prmt6 represses the pro-adipogenic Ppar-gamma–C/ebp-alpha transcription factor loop. *Scientific Reports.* **14**, 6656. 10.1038/s41598-024-57310-9.

Guilherme, A., Virbasius, J.V., Puri, V., and Czech, M.P. (2008). Adipocyte dysfunctions linking obesity to insulin resistance and type 2 diabetes. *Nature reviews. Molecular Cell Biology.* **9**, 367–377. 10.1038/nrm2391.

Gustafson, B., Hedjazifar, S., Gogg, S., Hammarstedt, A., and Smith, U. (2015). Insulin resistance and impaired adipogenesis. *Trends in Endocrinology and Metabolism: TEM.* **26**, 193–200. 10.1016/j.tem.2015.01.006.

Guzik, T.J., Skiba, D.S., Touyz, R.M., and Harrison, D.G. (2017). The role of infiltrating immune
cells in dysfunctional adipose tissue. *Cardiovascular Research* **113**, 1009–1023.
10.1093/cvr/cvx108.
Hammarstedt, A., Gogg, S., Hedjazifar, S., Nerstedt, A., and Smith, U. (2018). Impaired adipogenesis
and dysfunctional adipose tissue in human hypertrophic obesity. *Physiological Reviews*. **98**,
1911–1941. 10.1152/physrev.00034.2017.
Hocking, S., Samocha-Bonet, D., Milner, K. L., Greenfield, J. R., and Chisholm, D. J. (2013).
Adiposity and insulin resistance in humans: the role of the different tissue and cellular lipid
depots. *Endocrine Reviews* **34**, 463–500. 10.1210/er.2012-1041.
International Diabetes Federation (IDF) (2025). *Diabetes Atlas*, 11th Edition.
https://diabetesatlas.org/atlas/11th-edition/.
Jeon, C. Y., and Murray, M. B. (2008). Diabetes mellitus increases the risk of active tuberculosis: a
systematic review of 13 observational studies. *PLoS Medicine* **5**, e152.
Jeon, C. Y., Harries, A. D., Baker, M. A., Hart, J. E., Kapur, A., Lönnroth, K., Ottmani, S. E.,
Goonesekera, S., and Murray, M. B. (2010). Bi-directional screening for tuberculosis and
diabetes: a systematic review. *Tropical Medicine & International Health* **15**, 1300–1314.
10.1111/j.1365-3156.2010.02632.x.
Ji, E., Jung, M., Park, J., Lee, J., Kim, J., Han, S., Lee, H., and Kim, Y. (2014). Inhibition of
adipogenesis in 3T3-L1 cells and suppression of abdominal fat accumulation in high-fat diet-
feeding C57BL/6J mice after downregulation of hyaluronic acid. *International Journal of*
Obesity, **38**, 1035–1043. 10.1038/ijo.2013.202
Kanda, H., Tateya, S., Tamori, Y., Kotani, K., Hiasa, K., Kitazawa, R., Kitazawa, S., Miyachi, H.,
Maeda, S., Egashira, K., and Kasuga, M. (2006). MCP-1 contributes to macrophage infiltration
into adipose tissue, insulin resistance, and hepatic steatosis in obesity. *The Journal of Clinical*
Investigation **116**, 1494–1505. 10.1172/JCI26498.
Kashyap, R. S., Rajan, A. N., Ramteke, S. S., Agrawal, V. S., Kelkar, S. S., Purohit, H. J., Taori, G.
M., and Daginawala, H. F. (2012). Presence of multiple strains of *Mycobacterium tuberculosis*
in adipose tissue samples from tuberculosis patients. *PLoS One* **7**, e30626.
Katira, A., and Tan, P. H. (2016). Evolving role of adiponectin in cancer–controversies and update.
Cancer Biology & Medicine. **13**, 101–119. 10.28092/j.issn.2095-3941.2015.0092.
Khan, S., and Wang, C. H. (2014). ER stress in adipocytes and insulin resistance: mechanisms and
significance (Review). *Molecular Medicine Reports*. **10**, 2234–2240. 10.3892/mmr.2014.2532.
Khattak, M., Rehman, A. U., Muqaddas, T., Hussain, R., Rasool, M. F., Saleem, Z., Almalki, M. S.,
Alturkistani, S. A., Firash, S. Z., Alzahrani, O. M., Bahauddin, A. A., Abuhussain, S. A.,
Najjar, M. F., Elsabaa, H. M. A., and Haseeb, A. (2024). Tuberculosis (TB) treatment
challenges in TB-diabetes comorbid patients: a systematic review and meta-analysis. *Annals of*
Medicine. **56**, 2313683. 10.1080/07853890.2024.2313683.
Khoramipour, K., Chamari, K., Hekmatikar, A. A., Ziyaiyan, A., Taherkhani, S., Elguindy, N. M.,
and Bragazzi, N. L. (2021). Adiponectin: structure, physiological functions, role in diseases,
and effects of nutrition. *Nutrients*. **13**, 1180. 10.3390/nu13041180.
Kim, J. S., Ryu, M. J., Byun, E. H., Kim, W. S., Whang, J., Min, K. N., et al. (2011). Differential
immune response of adipocytes to virulent and attenuated *Mycobacterium tuberculosis*.
Microbes and Infection. **13**, 1242–1251. 10.1016/j.micinf.2011.07.002.
Kitamura, T., Kitamura, Y., Kuroda, S., Hino, Y., Ando, M., Kotani, K., Konishi, H., Matsuzaki, H.,
Kikkawa, U., Ogawa, W., and Kasuga, M. (1999). Insulin-induced phosphorylation and
activation of cyclic nucleotide phosphodiesterase 3B by the serine-threonine kinase Akt.
Molecular and Cellular Biology. **19**, 6286–6296. 10.1128/MCB.19.9.6286.
Klöting, N., and Blüher, M. (2014). Adipocyte dysfunction, inflammation and metabolic syndrome.
Reviews in Endocrine and Metabolic Disorders. **15**, 277–287. 10.1007/s11154-014-9301-0.
Knopp, J. L., Holder-Pearson, L., and Chase, J. G. (2019). Insulin units and conversion factors: a story
of truth, boots, and faster half-truths. *Journal of Diabetes Science and Technology*. **13**, 597–
600. 10.1177/1932296818805074.
Kurochkina, N., and Guha, U. (2013). SH3 domains: modules of protein-protein interactions.
Biophysical Reviews. **5**, 29–39. 10.1007/s12551-012-0081-z.

Kursawe, R., Dixit, V. D., Scherer, P. E., Santoro, N., Narayan, D., Gordillo, R., Giannini, C., Lopez, X., Pierpont, B., Nouws, J., Shulman, G. I., and Caprio, S. (2016). A role of the inflammasome in the low storage capacity of the abdominal subcutaneous adipose tissue in obese adolescents. *Diabetes.* **65**, 610–618. 10.2337/db15-1478.

Kyu, H. H., et al. (2018). The global burden of tuberculosis: results from the Global Burden of Disease Study 2015. *Lancet Infectious Diseases.* **18**, 261–284. 10.1016/S1473-3099(17)30703-X.

LeBlanc, S. E., Wu, Q., Lamba, P., Sif, S., and Imbalzano, A. N. (2016). Promoter-enhancer looping at the PPARγ2 locus during adipogenic differentiation requires the Prmt5 methyltransferase. *Nucleic Acids Research.* **44**, 5133–5147. 10.1093/nar/gkw129.

Lin, P. L., and Flynn, J. L. (2010). Understanding latent tuberculosis: a moving target. *Journal of Immunology.* **185**, 15–22. 10.4049/jimmunol.0903704.

Lin, Y., Yuan, Y., Zhao, X., Liu, J., Qiu, L., He, X., Bai, Y., Li, B., Zeng, Z., Yang, G., Meng, X., Yuan, C., Zheng, H., Kang, W., and Harries, A. D. (2017). The change in blood glucose levels in tuberculosis patients before and during anti-tuberculosis treatment in China. *Global Health Action.* **10**, 1289737. 10.1080/16549716.2017.1289737.

Lindsay, R. S., Funahashi, T., Hanson, R. L., Matsuzawa, Y., Tanaka, S., Tataranni, P. A., Knowler, W. C., and Krakoff, J. (2002). Adiponectin and development of type 2 diabetes in the Pima Indian population. *Lancet.* **360**, 57–58. 10.1016/S0140-6736(02)09335-2.

Magee, M. J., Salindri, A. D., Kyaw, N. T. T., Auld, S. C., Haw, J. S., and Umpierrez, G. E. (2018). Stress hyperglycemia in patients with tuberculosis disease: epidemiology and clinical implications. *Current Diabetes Reports.* **18**, 71. 10.1007/s11892-018-1036-y.

Maréchal, L., Laviolette, M., Rodrigue-Way, A., Sow, B., Brochu, M., Caron, V., and Tremblay, A. (2018). The CD36-PPARγ pathway in metabolic disorders. *International Journal of Molecular Sciences.* **19**, 1529. 10.3390/ijms19051529.

Martinez, N., Cheng, C. Y., Ketheesan, N., Cullen, A., Tang, Y., Lum, J., West, K., Poidinger, M., Guertin, D. A., Singhal, A., and Kornfeld, H. (2019). mTORC2/Akt activation in adipocytes is required for adipose tissue inflammation in tuberculosis. *EBioMedicine.* **45**, 314-327. 10.1016/j.ebiom.2019.06.052.

Matthews, D. R., Hosker, J. P., Rudenski, A. S., Naylor, B. A., Treacher, D. F., and Turner, R. C. (1985). Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia.* **28**, 412–419. 10.1007/BF00280883.

Mave, V., Meshram, S., Lokhande, R., Kadam, D., Dharmshale, S., Bharadwaj, R., and Golub, J. E. (2017). Prevalence of dysglycemia and clinical presentation of pulmonary tuberculosis in Western India. *International Journal of Tuberculosis and Lung Disease.* **21**, 1280–1287.

Mootha, V. K., Lindgren, C. M., Eriksson, K.F., Subramanian. A., Sihag, S., Lehar, J., Puigserver, P., Carlsson, E., Ridderstråle, M., Laurila, E., Houstis, N., Daly, M. J., Patterson, N., Mesirov, J. P., Golub, T. R., Tamayo, P., Spiegelman, B., Lander, E. S., Hirschhorn, J. N., Altshuler, D., and Groop, L. C. (2003). PGC-1α-responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. *Nature Genetics.* **34**, 267-273. 10.1038/ng1180.

Morigny, P., Houssier, M., Mouisel, E., and Langin, D. (2016). Adipocyte lipolysis and insulin resistance. *Biochimie.* **125**, 259-66. 10.1016/j.biochi.2015.10.024.

Mugusi, F., Swai, A. B., Alberti, K. G., and McLarty, D. G. (1990). Increased prevalence of diabetes mellitus in patients with pulmonary tuberculosis in Tanzania. *Tubercle.* **71**, 271–276. 10.1016/0041-3879(90)90040-f

Navale, A. M., and Paranjape, A. N. (2016). Glucose transporters: physiological and pathological roles. *Biophysical Reviews.* **8**, 5–9. 10.1007/s12551-015-0186-2

Neyrolles, O., Hernández-Pando, R., Pietri-Rouxel, F., Fornès, P., Tailleux, L., Barrios Payán, J. A., Pivert, E., Bordat, Y., Aguilar, D., Prévost, M. C., Petit, C., and Gicquel, B. (2006). Is adipose tissue a place for *Mycobacterium tuberculosis* persistence? *PLoS One.* **1**, e43. 10.1371/journal.pone.0000043

Noubiap, J. J., Nansseu, J. R., Nyaga, U. F., Nkeck, J. R., Endomba, F. T., Kaze, A. D., Agbor, V. N., and Bigna, J. J. (2019). Global prevalence of diabetes in active tuberculosis: a systematic

review and meta-analysis of data from 2.3 million patients with tuberculosis. *Lancet Global Health*. **7**, e448–e460. 10.1016/S2214-109X(18)30487-X

Oluboyo, P. O., and Erasmus, R. T. (1990). The significance of glucose intolerance in pulmonary tuberculosis. *Tuberculosis*. **71**, 135–138. 10.1016/0041-3879(90)90010-6

Oswal, N., Lizardo, K., Dhanyalayam, D., Ayyappan, J. P., Thangavel, H., Heysell, S. K., and Nagajyothi, J. F. (2022). Host metabolic changes during *Mycobacterium tuberculosis* infection cause insulin resistance in adult mice. *Journal of Clinical Medicine*. **11**, 1646. 10.3390/jcm11061646

Ouchi, N., and Walsh, K. (2007). Adiponectin as an anti-inflammatory factor. *Clinica Chimica Acta;* **380**, 24–30.

Pal, R., Ghosh, S., and Mukhopadhyay, S. (2021). Moonlighting by PPE2 protein: Focus on mycobacterial virulence. *Journal of Immunology*. **207**, 2393–2397. 10.4049/jimmunol.2100212

Pal, R., and Mukhopadhyay, S. (2021). PPE2 protein of *Mycobacterium tuberculosis* affects myeloid hematopoiesis in mice. *Immunobiology*. **226**, 152051. 10.1016/j.imbio.2020.152051

Pal, R., Bisht, M. K., and Mukhopadhyay, S. (2022). Secretory proteins of *Mycobacterium tuberculosis* and their roles in modulation of host immune responses: focus on therapeutic targets. *The FEBS Journal*. **289**, 4146–4171. 10.1111/febs.16369

Park, H. S., Ju, U. I., Park, J. W., et al. (2016). PPARγ neddylation essential for adipogenesis is a potential target for treating obesity. *Cell Death & Differentiation*. **23**, 1296–1311. 10.1038/cdd.2016.6

Paton, N. I., and Ng, Y. M. (2006). Body composition studies in patients with wasting associated with tuberculosis. *Nutrition*. **22**, 245–251. 10.1016/j.nut.2005.06.009

Raghuraman, S., Vasudevan, K. P., Govindarajan, S., Chinnakali, P., and Panigrahi, K. C. (2014). Prevalence of diabetes mellitus among tuberculosis patients in urban Puducherry. *North American Journal of Medical Sciences*. **6**, 30–34. 10.4103/1947-2714.125863

Reyes-Farias, M., Fos-Domenech, J., Serra, D., Herrero, L., and Sánchez-Infantes, D. (2021). White adipose tissue dysfunction in obesity and aging. *Biochemical Pharmacology*. **192**, 114723. 10.1016/j.bcp.2021.114723

Richard, A. J., White, U., Elks, C. M., and Stephens, J. M. (2020). Adipose tissue: Physiology to metabolic dysfunction. In K. R. Feingold et al. (Eds.), *Endotext*. MDText.com, Inc.

Root, H. F. (1934). The association of diabetes and tuberculosis. *New England Journal of Medicine*. **210**, 1–13.

Sadowski, H. B., Wheeler, T. T., and Young, D. A. (1992). Gene expression during 3T3-L1 adipocyte differentiation. Characterization of initial responses to the inducing agents and changes during commitment to differentiation. *Journal of Biological Chemistry*. **267**, 4722–4731.

Saetang, J., and Sangkhathat, S. (2018). Role of innate lymphoid cells in obesity and metabolic disease (Review). *Molecular Medicine Reports*. **17**, 1403–1412. 10.3892/mmr.2017.8038

Samad, F., Uysal, K. T., Wiesbrock, S. M., Pandey, M., Hotamisligil, G. S., and Loskutoff, D. J. (1999). Tumor necrosis factor α is a key component in the obesity-linked elevation of plasminogen activator inhibitor 1. *Proceedings of the National Academy of Sciences of the United States of America*. **96**, 6902–6907. 10.1073/pnas.96.12.6902

Scheja, L., and Heeren, J. (2019). The endocrine function of adipose tissues in health and cardiometabolic disease. *Nature Reviews Endocrinology*. **15**, 507–524. 10.1038/s41574-019-0230-6

Schweiger, M., Eichmann, T. O., Taschler, U., Zimmermann, R., Zechner, R., and Lass, A. (2014). Measurement of lipolysis. *Methods in Enzymology*. **538**, 171–193. 10.1016/B978-0-12-800280-3.00010-4

Siitonen, N., Pulkkinen, L., Lindström, J., Kolehmainen, M., Eriksson, J. G., Venojärvi, M., Ilanne-Parikka, P., Keinänen-Kiukaanniemi, S., Tuomilehto, J., and Uusitupa, M. (2011). Association of ADIPOQ gene variants with body weight, type 2 diabetes and serum adiponectin concentrations: the Finnish Diabetes Prevention Study. *BMC Medical Genetics*. **12**, 5. 10.1186/1471-2350-12-5

Song, Y. F., Hogstrand, C., Wei, C. C., Wu, K., Pan, Y. X., and Luo, Z. (2017). Endoplasmic reticulum (ER) stress and cAMP/PKA pathway mediated Zn-induced hepatic lipolysis. *Environmental Pollution*. **228**, 256–264. 10.1016/j.envpol.2017.05.046

Srivastava, S., Battu, M. B., Khan, M. Z., Nandicoori, V. K., and Mukhopadhyay, S. (2019).
Mycobacterium tuberculosis PPE2 protein interacts with p67[phox] and inhibits reactive oxygen
species production. *Journal of Immunology*. **203**, 1218–1229. 10.4049/jimmunol.1801143

Subramanian, A., Tamayo, P., Mootha, V. K., Mukherjee, S., Ebert, B. L., Gillette, M. A., Paulovich,
A., Pomeroy, S. L., Golub, T. R., Lander, E. S., and Mesirov, J. P. (2005). Gene set enrichment
analysis: a knowledge-based approach for interpreting genome-wide expression profiles.
Proceedings of the National Academy of Sciences of the United States of America. **102**, 15545–
15550. 10.1073/pnas.0506580102

Susca, N., Leone, P., Prete, M., Cozzio, S., and Racanelli, V. (2024). Adipose failure through
adipocyte overload and autoimmunity. *Autoimmunity Reviews*. 23, 103502.
10.1016/j.autrev.2023.103502

Tishinsky, J. M., Robinson, L. E., and Dyck, D. J. (2012). Insulin-sensitizing properties of
adiponectin. *Biochimie*. **94**, 2131–2136. 10.1016/j.biochi.2012.01.017

Veloso, C. C., Oliveira, M. C., Rodrigues, V. G., Oliveira, C. C., Duarte, L. P., Teixeira, M. M.,
Ferreira, A. V. M., and Perez, A. C. (2019). Evaluation of the effects of extracts of *Maytenus*
imbricata (Celastraceae) on the treatment of inflammatory and metabolic dysfunction induced
by high-refined carbohydrate diet. *Inflammopharmacology*. **27**, 539-548. 10.1007/s10787-018-
0496-7.

Viswanathan, V., Kumpatla, S., Aravindalochanan, V., Rajan, R., Chinnasamy, C., Srinivasan, R., et
al. (2012). Prevalence of diabetes and pre-diabetes and associated risk factors among
tuberculosis patients in India. *PLoS One*. **7**, e41367. 10.1371/journal.pone.0041367.

Viswanathan, V., Vigneswari, A., Selvan, K., Satyavani, K., Rajeswari, R., and Kapur, A. (2014).
Effect of diabetes on treatment outcome of smear-positive pulmonary tuberculosis–a report
from south India. *Journal of Diabetes and Its Complications*. **28**, 162–165.
10.1016/j.jdiacomp.2013.12.003

World Health Organization. (2023). *Global Tuberculosis Report 2023*. Geneva: World Health
Organization. License: CC BY-NC-SA 3.0 IGO.

Wu, H., Li, X., and Shen, C. (2020). Peroxisome proliferator-activated receptor γ in white and brown
adipocyte regulation and differentiation. *Physiological Research*. **69**, 759–773.
10.33549/physiolres.934411

Wu, H., Ghosh, S., Perrard, X. D., Feng, L., Garcia, G. E., Perrard, J. L., Sweeney, J. F., Peterson, L.
E., Chan, L., Smith, C. W., and Ballantyne, C. M. (2007). T-cell accumulation and regulated on
activation, normal T cell expressed and secreted upregulation in adipose tissue in obesity.
Circulation. **115**, 1029-38. 10.1161/CIRCULATIONAHA.106.638379

Ziemke, F., and Mantzoros, C. S. (2010). Adiponectin in insulin resistance: lessons from translational
research. *The American Journal of Clinical Nutrition*. **91**, 258S–261S.
10.3945/ajcn.2009.28449C

1165

1166

Figure Legends

Figure 1. Recombinantly purified PPE2 protein of *M. tuberculosis* (rPPE2) inhibits differentiation of 3T3-L1 cells into adipocytes. (A) 3T3-L1 cells were treated with different concentrations of rPPE2 and induced to differentiate using DMI cocktail (dexamethasone, insulin and IBMX). At day 10 post-induction of differentiation, the accumulated triglycerides in adipocytes were stained with Oil Red O and image of differentiated adipocytes were taken. **(B)** For quantitative measurement of lipid accumulation, lipid-bound Oil Red O stain was extracted with isopropanol and absorbance of the extracted solution was measured at 490 nm using isopropanol as blank. **(C)** Differentiation of rPPE2-treated 3T3-L1 cells in the absence or presence of anti-PPE2 Ab. **(D)** 3T3-L1 cells were differentiated in the absence or presence of 3 µg/ml rPPE2 and changes in the expressions of PPAR-γ, C/EBP-α and FAS genes at various days post-differentiation were measured by qPCR. GAPDH transcript level was used as an internal control. The results shown are mean ± SEM of 3 different experiments. ns = non-significant.

Figure 2. Infection of mice with *M. smegmatis* expressing PPE2 protein of *M. tuberculosis* alters adipose tissue physiology. (A, B) Balb/c mice were infected with 100 million of either Msmeg-pVV or Msmeg-PPE2 bacilli and sacrificed at 2^{nd} and 7^{th} day post-infection (dpi) to measure the weight of perigonadal (**A**) and visceral (**B**) fat tissue. **(C, D, E, F)** Expression levels of PPAR-γ, C/EBP-α and adiponectin in perigonadal tissue and in visceral tissue were quantified by qPCR at day 2 (**C** and **E**) and day 7 (**D** and **F**) post-infection (dpi). GAPDH transcript level was used as an internal control. **(G and H)** Histological analyses of perigonadal (**G**) and visceral (**H**) adipose tissue sections were performed at day 2 and day 7 post-infection by staining with hematoxylin and eosin (H&E)

and photographs of representative sections are presented at 40X magnification (scale bar = 100 μm). Black arrows indicate adipocyte hypertrophy (enlarged adipocyte) and the red arrows point to the infiltrated immune cells into the adipose tissue. **(I and J)** Adipocyte cell area for perigonadal (**I**) and visceral (**J**) fat tissue was measured using ImageJ software. **(K and L)** Bacterial load in perigonadal (**K**) and visceral (**L**) fat tissue was measured at day 2 and day 7 post-infection. For **(M and N)** For glucose tolerance test at day 7 post-infection, mice were fasted overnight followed by intraperitoneal injection of glucose (2 mg/kg body weight) and blood glucose levels were monitored for 120 minutes (**M**) and the Area under the curve (AUC) was calculated (**N**). The results shown are mean ± SEM of 4 mice.

Figure 3. Administration of rPPE2 protein leads to adipose tissue loss and impaired glucose tolerance in mice. Balb/c mice were intraperitoneally administered with 3 mg/kg of either BSA or recombinant PPE2 protein (rPPE2) on alternate days for 12 days and mice were sacrificed at day 14 post-treatment. **(A)** Representative photographic images of perigonadal and visceral fat tissues are shown. (**B and C**) Weight of the total perigonadal fat mass (**B**) and visceral fat mass (**C**) are recorded. **(D)** Transcript levels of PPAR-γ, C/EBP-α and adiponectin genes are quantified in perigonadal fat tissue using qPCR where GAPDH transcript level was used as internal control. **(E)** Histopathology analyses of perigonadal and visceral fat tissue (representative photographs of hematoxylin and eosin (H&E)-stained sections) is shown at 40X magnification (scale bar = 100 μm). Black arrows indicate adipocyte hypertrophy (enlarged adipocyte) and the red arrows indicate infiltrated immune cells into the adipose tissue. (**F and G**) Mice were starved overnight and glucose tolerance test was performed for 120 minutes at day 14 post-treatment (**F**) and the mean Area under the curve (AUC) was calculated **(G).** The results shown are mean ± SEM of 4 mice.

1215

Figure 4. **Pre-immunization with rPPE2 protects the mice from detrimental effects of**

PPE2 on adipose tissue. Mice were immunized with 3 doses of recombinant PPE2 protein

with Freund's incomplete adjuvant at 15 days interval. Next, non-immunized and rPPE2-

immunized mice were infected with Msmeg-pVV or Msmeg-PPE2 and a change in fat

content was observed after day 7. (**A**) Photographs of perigonadal and visceral fat tissues are

shown**.** Weight of perigonadal (**B**) and visceral (**C**) fat tissues of various groups is recorded.

(**D**) Percentage change in total body fat as determined by Dual-energy X-ray Absorptiometry

(DEXA) is shown. (**E**) At day 7, sections of perigonadal and visceral fat tissues were

prepared and stained with hematoxylin and eosin and photographs of the representative

sections were visualized at 40X magnification (scale bar = 100 µm). (**F**) A glucose tolerance

test was performed on 7th dpi and (**G**) Area under the curve (AUC) of glucose concentration

is shown. The results shown are mean ± SEM of 4 mice.

1228

Figure 5. PPE2 alters adipose tissue physiology during *M. tuberculosis* infection in mice.

Balb/c mice were infected through aerosol route either with *M. tuberculosis* clinical strain

CDC1551 (wild-type) or CDC1551-ΔPPE2 (*ppe2*KO) or CDC1551-ΔPPE2::PPE2

(complemented) strains and were sacrificed at day 30 (**A** and **C**) and day 60 (**B** and **D**) post-

infection (dpi). Perigonadal (PG) and visceral (VS) fat tissues were harvested and weight of

PG fat tissue (**A and B**) and VS fat tissue (**C and D**) was measured. (**E**) Also, representative

photographs of hematoxylin and eosin (H&E)-stained sections of PG at both day 30 and day

60 post-infection were visualized at 40X magnification (scale bar = 100 µM). Black arrow

shows adipocyte hypertrophy (enlarged adipocyte) and red arrow represent infiltrated

immune cells into the adipose tissue. (**F-I**) Graphs showing cell area of adipocytes in PG (**F**

and H) and VS (**G and I**) fat tissues at 30 dpi (**F and G**) and 60 dpi (**H and I**) respectively.

(**J-M**) Number of infiltrated CD3[+] T-cells were counted using ImageJ software and plotted as

cells per field of view for PG (**J and L**) and VS (**K and M**) adipose tissue sampled at 30 dpi

(**J and K**) and 60 dpi (**L and M**) respectively. (**N and O**) Changes of expression of PPAR-γ

and CD36 in perigonadal tissue at day 30 (**Ni, Nii**) and day 60 (**Oi, Oii**) post-infection were

measured by qPCR and GAPDH transcript level was used as an internal control. (**P**) Number

of bacilli in PG fat tissue at 30 (**Pi**) and 60 (**Pii**) dpi respectively. (**Q and R**) IPGTT

performed on mice along with corresponding AUC data obtained at day 30 (**Qi, Qii**) and day

60 (**Ri**, **Rii**) post-infection respectively. (**S**) Fasting blood glucose levels of mice at 60 dpi

and (**T**) HOMA-IR scores in each mice group were calculated at day 60 post-infection. The

results shown are mean ± SEM of 5 mice.

1250

Figure 6. PPE2 modulates the adipose tissue transcriptome during infection. Perigonadal

fat tissues were harvested from mice (n = 3) infected with wild-type, *ppe2*KO and

complemented strains at 60 dpi and subjected to RNA sequencing using Illumina platform.

(**A**) Venn diagram showing all the differentially expressed genes (DEGs) in all subsets (**B**)

Venn diagram showing significant DEGs after applying ±1.3-fold change cutoff. (**C**) Volcano

plot showing significant DEGs in *ppe2*KO versus wild-type-infected mice in PG adipose

tissue. (**D**) Gene set enrichment analysis (GSEA) for gene ontology analysis (**E**) KEGG

pathway analysis in *ppe2*KO versus CDC1551 adipose tissues. (**F**) Heat map of DEGs for

chemokine, metabolic pathways, cytokines and TB susceptibility genes in PG adipose tissues.

(**G**) Heat map showing differentially expressed ribosomal biogenesis genes (**H**) Heat map

showing differential expression of phospholipase C activity genes. (**I**) Perigonadal fat tissue

were harvested from mice either left uninfected or infected with various strains of *M.*

tuberculosis at day 60 post-infection and used for cDNA synthesis. Transcript levels of ATF4

and GRP78 as measured by qPCR with β-actin as an internal control. The results shown are

mean ± SEM of 4 mice.

1266

Figure 7. PPE2 induces lipolysis in adipocytes by targeting the cAMP-PKA-HSL

signaling axis. (**A**) 3T3-L1 cells were differentiated into mature adipocytes and then treated

with various concentration of rPPE2 and after 24 hours medium was removed from cells and

amount of glycerol released was quantified. (**B**) 3T3-L1 mature adipocytes were incubated

with 3 µg/ml rPPE2 or TNF-α cytokine in the absence or presence of TNF-α inhibitor and the

free glycerol released was quantified. (**C**) Mature 3T3-L1 adipocytes were treated with 3

µg/ml of rPPE2 and after 24 hours, cells were harvested for semi-quantitative reverse

transcriptase PCR analysis (**Ci**) and densitometric quantification was carried out using

ImageJ software (**Cii**). (**D**) In a separate experiment, rPPE2-treated 3T3-L1 mature

adipocytes were harvested after 6 hours and 24 hours, and the levels of phosphorylated

hormone-sensitive lipase (p-HSL) and total HSL were assessed by Western blotting using

anti-p-HSL Ab and anti-HSL Ab respectively (**Di**). Densitometric quantification of the

Western blots was carried out using ImageJ software (**Dii**). (**E**) Glycerol release was

quantified in rPPE2-treated (3 µg/ml) mature 3T3-L1 adipocytes at 24 hours post-treatment

in the absence or presence of H89, a PKA-specific inhibitor. (**A-E**) All the data represent

mean ± SEM of 3 independent experiments. (**F**) Levels of non-esterified free fatty acid

(NEFA) measured in sera of mice either left uninfected or infected with various strains of *M.*

tuberculosis at day 60 post-infection. The results shown are mean ± SEM of 5 mice. (**G**)

NEFA levels in the plasma of healthy controls and TB patients.

1286

1287

Supplementary Figures

Figure S1. rPPE2 does not cause 3T3-L1 cell toxicity. 3T3-L1 cells (2×10^5/ml) were treated with different concentrations of rPPE2 for 48 hours and cell cytotoxicity was measured by MTT assay. In brief, MTT (3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide, Sigma-Aldrich, USA) was added at 1 mg/ml and incubated for 4 hours. The cells were lysed overnight using 100 μl of lysis buffer (20% SDS and 50% dimethyl formamide) and the absorbance was measured at 550 nm. ns = non-significant

Figure S2. PPE2 inhibits 3T3-L1 adipocyte differentiation during infection. Undifferentiated 3T3-L1 pre-adipocytes were infected with *M. smegmatis* harboring pVV16 empty vector (Msmeg-pVV) or *M. smegmatis* expressing PPE2 (Msmeg-PPE2) at 1:10 MOI (multiplicity of infection) for 24 hours. Following infection, the cells were washed with gentamicin to a final concentration of 50 μg/ml to remove extracellular bacteria and these infected cells were induced to differentiate into matured adipocytes. At day 10 post-differentiation, the cells were fixed with 10% paraformaldehyde and stained with Oil Red O. For quantification of Oil Red O stain, the dye was extracted using isopropanol and the absorbance was measured at 490 nm using a spectrophotometer.

Figure S3. Photographs of perigonadal and visceral fat tissues of mice infected with either Msmeg-pVV or Msmeg-PPE2. Balb/c mice were infected with 100 million of either Msmeg-pVV or Msmeg-PPE2 through intravenous route and were sacrificed on 2nd and 7th day post-infection and the perigonadal and visceral fat tissue were harvested and photographs of the perigonadal and visceral fat tissues from infected mice are shown.

1311

Figure S4. Levels of anti-PPE2 Ab in mice immunized with rPPE2. Balb/c mice (n = 8) were immunized with 100 µg of rPPE2 using incomplete Freund's adjuvant. Two booster doses were given at 15 days intervals and sera were collected on the 45[th] day of immunization. Pre-bleed sera were collected at day 0. Sera were diluted and the levels of anti-PPE2 Ab were checked by ELISA. For ELISA, a 96-well microtiter plate was coated with rPPE2 protein at 1 µg/well diluted in carbonate buffer and incubated overnight at 4^{o}C. After blocking the wells with 2% BSA in PBS, mice sera at various dilutions were added and plate was incubated at 37^{o}C for 1 hour. After washing with PBS-T (0.05% Tween 20 in PBS), the plate was incubated with anti-mouse Ab bound to horseradish peroxidase (HRP) (1:10000) followed by washing with PBS-T. The plate was then washed and developed with TMB (3, 3', 5, 5'-Tetramethylbenzidine) substrate (BD Biosciences, USA). The reaction was stopped with 1N H_2SO_4 and the absorbance was measured at 450/550 nm in an ELISA microplate reader.

1325

Figure S5. Photographs of perigonadal and visceral fat tissues of mice infected with various strains of *M. tuberculosis*. Balb/c mice were infected *via* aerosol route with wild-type CDC1551 or CDC1551-ΔPPE2 or CDC1551-ΔPPE2::PPE2 and were sacrificed at day 30 or day 60 post-infection. Perigonadal and visceral fat tissues were harvested and photographs of perigonadal and visceral fat tissues are shown.

1331

Figure S6. Infiltration of CD3^{+} T-cells in perigonadal and visceral fat tissues. Balb/c mice were infected *via* aerosol route with wild-type *M. tuberculosis* strain CDC1551 or

CDC1551-ΔPPE2 or CDC1551-ΔPPE2::PPE2 and were sacrificed at day 30 or day 60 post-

infection. Perigonadal and visceral fat tissues were harvested. The tissues were fixed in 4%

paraformaldehyde and embedded in paraffin wax and then sectioned using a microtome. The

tissue sections were then deparaffinized with 2 changes of xylene followed by sequential

hydration with two changes of 100% ethanol for 3 minutes each followed by 95%, 75% and

50% ethanol for 1 minute each. Antigens were retrieved by heating the sections in sodium

citrate buffer (10 mM, pH 6) at 95-100°C for 30 minutes, followed by cooling at room

temperature. The sections were then premetallized by washing twice with PBS for 2 min

each, followed by incubation with 0.1% Triton-X 100 in PBS. Blocking was carried out in

PBS containing 5% bovine serum albumin (BSA) for 30 minutes at room temperature. After

3 washes with PBS, the sections were incubated with Pacific Blue™-conjugated anti-mouse

CD3 antibody (BioLegend, USA) overnight at 4°C, followed by washing thrice with PBS for

3 minutes each and imaged using a confocal microscope.

1347

Figure S7. Cystoscape map showing upregulated chemokine map in perigonadal adipose

tissue of mice infected with wild-type CDC1551 when compared with adipose tissue of mice

infected with CDC1551-ΔPPE2.

.

Figure S8. Insulin does not inhibit rPPE2-induced lipolysis. The mature 3T3-L1

adipocytes were treated with rPPE2 protein (3 µg/ml) and after 30 minutes, insulin (1 µg/ml)

was added to the medium and lipolysis was measured after 24 hours by estimating free

glycerol released in the medium using Free Glycerol Reagent Kit (F6428-40mL, Sigma-

Aldrich, USA). Ns = non-significant

1357

1358

Figure 1

Figure 2

M Glucose tolerance test

Glucose (% of initial)

Legend:
- Uninfected
- Msmeg-pVV
- Msmeg-PPE2

N

p < 0.01

p < 0.01

Figure 2

Figure 3

Figure 4

Figure 5

Figure 5

Figure 6

Figure 6

Figure 7