Autosomal Allelic Inactivation: Variable Replication and Dosage Sensitivity

Michael B. Heskett[1], Athanasios E. Vouzas[2], Brian Johnstone[3], Krister P. Freese[4], Phillip A. Yates[5], Philip F. Copenhaver[6], Paul T. Spellman[1], David M. Gilbert[2], Mathew J. Thayer[5],*.

[1]Division of Hematology Oncology
Departments of Medicine and Human Genetics
David Geffen School of Medicine
University of California, Los Angeles

[2]San Diego Biomedical Research Institute
3525 John Hopkins Court
San Diego, CA 92121

[4]Shriners Hospitals for Children—Portland
3101 SW Sam Jackson Park Rd,
Portland, Oregon, 97239

[3]Department of Orthopedics and Rehabilitation
[5]Department of Chemical Physiology and Biochemistry
[6]Department of Cell, Developmental and Cancer Biology
Oregon Health & Science University
3181 S. W. Sam Jackson Park Road
Portland, Oregon 97239

*Corresponding author.

ORCID: 0000-0001-6483-1661

TEL: (503) 494-2447

FAX: (503) 494-7368

Email:thayerm@ohsu.edu

Abstract

Autosomal monoallelic gene expression and asynchronous replication between alleles are established features of imprinted genes and genes regulated by allelic exclusion. Inactivation/Stability Centers (I/SCs) are recently described autosomal loci that exhibit epigenetic regulation of allelic expression and replication timing, with differences that can be comparable to those observed between the active and inactive X chromosomes [1]. Here we characterize >100 autosomal loci with allele-specific epigenetic regulation of replication timing and gene expression, defining them as I/SCs. I/SCs are approximately 1 megabase in size and can contain both protein-coding and noncoding genes. In different single cell derived clones, these genes may be expressed from a single allele, the opposite allele, both alleles, or not expressed at all. This stochastic, yet mitotically stable, pattern indicates that the choice of which allele is expressed is independent of parent of origin and independent of the expression status of the other allele. Similarly, alleles within I/SCs show varying replication timing, either earlier or later, that is also independent of the other allele. Additionally, we identify syntenic loci in the mouse genome that display epigenetic regulation of allelic replication timing, highlighting the genomic organization and conservation of I/SC-associated regulation between human and mouse genomes. The allele-restricted regulation described here creates extensive cellular mosaicism through a stable epigenetic mechanism. This mosaicism impacts numerous dosage-sensitive genes associated with human diseases such as Alzheimer, Parkinson, epilepsy, deafness, and impaired intellectual development.

Introduction:

The relationship between monoallelic expression and asynchronous replication between alleles of autosomal genes, exemplified by genes subject to genomic imprinting or allelic exclusion, has been the focus of extensive research for over three decades [2-4]. In recent work, we employed haplotype-phased RNA expression and replication timing assays on multiple single-cell-derived Lymphoblastoid Cell Line (LCL) clones from two unrelated individuals to identify loci exhibiting allele-restricted expression and replication timing [1]. This clonal, allele-specific analysis revealed an unexpected genomic feature affecting megabase-sized loci across every autosome, characterized by allele-restricted gene expression (including random monoallelic expression) and variable replication timing (including asynchronous replication between alleles). The magnitude of allelic differences in expression and replication timing observed in some of the clones was comparable to the large differences detected between the active and inactive X chromosomes [1]. We have named these autosomal loci Inactivation/Stability Centers (I/SCs) [1,5-7].

I/SCs can encompass both coding and noncoding genes that may be expressed from a single allele in some clones, expressed from the opposite allele in other clones, biallelically expressed in other clones, or silent on both alleles in yet other clones. This pattern of allele-restricted expression indicates that each allele independently adopts either an expressed or silent state. Importantly, because these expression states are mitotically stable, allele-autonomous, and independent of parental origin, we refer to the choice of the expressed allele as stochastic [1]. To describe this phenomenon, we and others have used the term Allelic Expression Imbalance (AEI) to refer to the stochastic

nature of autosomal random monoallelic expression [1,8-11]. Additionally, the replication

timing profiles on autosomes are highly variable and clone-specific, exhibiting both

synchronous and asynchronous patterns in different clones. We refer to this as Variable

Epigenetic Replication Timing (VERT) [1].

Random monoallelic expression in the olfactory, immune, and central nervous

systems is well established and plays a critical role in generating cell-specific identities

[12-14]. However, increasing evidence indicates that autosomal genes outside these

systems also exhibit allele-specific expression, suggesting that AEI plays a broader role

across diverse cell types [4,13]. The functional diversity of these genes implies that AEI

contributes to a broad array of cellular processes and is not limited to specialized

systems.

In the present study, we aimed to determine whether I/SCs could be detected in

the genomes of clonal populations of human primary cells derived from a tissue distinct

from those previously examined. To this end, we employed articular cartilage progenitor

(ACP) cells as a second model system [15]. ACP cells are non-transformed yet are

particularly well-suited for clonal analysis due to several advantages: 1) they can be

readily isolated from amputated tissue of patients with polydactyly, 2) they exhibit high

proliferative capacity, and 3) they can be efficiently expanded from single cells [16]. We

identify an additional 163 VERT regions in the human genome, increasing the total

number of potential I/SCs to 363, which collectively cover approximately 12% of the

human genome. Of these, 112 loci display both AEI and VERT at the same genomic

position and therefore represent high-confidence I/SCs.

Notably, we found that many previously reported "random monoallelic" genes,

including those encoding olfactory and vomeronasal receptors[3,17], antigen receptors[18],

and homophilic cell adhesion proteins[12], are located within VERT regions. This confirms

that these well-established monoallelically expressed genes reside within genomic loci

that display the defining characteristics of I/SCs: allele-restricted expression and variable

replication timing.

The relationship between genotype and phenotype is central to understanding

human genetic disease. While most loss-of-function mutations are recessive,

indicating that a single normal allele is sufficient for typical function, dominantly inherited

disorders often result from gain-of-function mutations. However, a significant fraction of

dominant diseases arises through haploinsufficiency (i.e. heterozygous for a loss of

function mutation), where a single functional allele fails to maintain normal physiology.

Although haploinsufficiency may affect up to 40% of human genes [19-21], the underlying

molecular mechanisms remain poorly understood. Current models suggest that

insufficient gene dosage may arise from a lack of redundancy, critical expression

thresholds, stoichiometric imbalances, or toxicity associated with a lack of

overexpression buffering [22-25]. In contrast, insights from X-linked disease inheritance

illustrate how cellular mosaicism and cellular selection can shape phenotypic outcomes:

in XX individuals, differences in the survival or expansion of cells expressing either the

wild-type or mutant alleles can influence whether a disease manifests as recessive or

dominant [21,26,27].

The magnitude of AEI is often quantified using allelic expression ratios, which

represent the proportion of transcripts derived from each allele. Importantly, recent

findings have demonstrated that even modest allelic expression biases can

have functional consequences, particularly in disease contexts. One recent study

directly linked small but stable AEI to variable clinical outcomes among individuals

carrying the same pathogenic mutation, underscoring AEI as a potential modulator of

disease expressivity [28]. These observations highlight the importance of considering AEI

as a disease modifier that may shape phenotypic diversity and influence the penetrance

of genetic disorders.

Here, we demonstrate that numerous genes implicated in autosomal human

genetic diseases, such as Parkinson (DNAJC6, LRRK2 and SNCA), epilepsy (SCN1A,

GABR1, and SAMD12), deafness (ATP11A, EPS8, and MYO6), and

neurodevelopmental disorders (KCNA1, RETREG1, ROBO1, and TIAM1), are located

within VERT regions and display AEI. Furthermore, mapping I/SCs across all

autosomes reveals 87 known dominantly inherited single-gene disorders that map

within VERT regions, where loss of function mutations in most of these genes result in

haploinsufficiency [29]. We propose that epigenetically regulated stochastic AEI can

contribute to pathogenic phenotypes in heterozygous individuals by either reducing the

number of functional cells below a critical disease threshold, akin to X linked dominant

disorders in females, or by disrupting the normal mosaicism of gene expression within

disease-relevant tissues.

Results

Here, we employed articular cartilage progenitor (ACP) cells[15] as a distinct primary cell type

from those used in our previous studies, serving as a second model system to identify

I/SCs. ACP cells were isolated from amputated digits of patients with polydactyly, and

genomic DNA from their parents was sequenced to generate phased genomes for allele-specific analyses. We examined two independent sets of ACP clones derived from unrelated individuals, one female (ACP7) and one male (ACP6), using haplotype-phased Repli-seq and RNA-seq (Figure 1a and 1b).

To benchmark the magnitude of allele-specific epigenetic differences in replication timing and gene expression, we incorporated two internal controls. First, analysis of female ACP7 clones enabled direct comparison of autosomal allelic differences with those observed between the active and inactive X chromosomes within the same clones, thereby establishing statistical thresholds for VERT and AEI assignment (Figure 1c; Supplementary Tables 1–3). Second, we examined allelic replication timing at known imprinted loci and identified 12 imprinted regions displaying allelic replication imbalance that is comparable in magnitude to the VERT regions detected at non-imprinted loci. Imprinted regions with differences in allelic replication timing are highlighted in Figure 1e, Supplementary Figure 1, as well as in subsequent figures (Figures 2d, 3c, 6g, 7e, and 7g).

To identify genomic regions with epigenetic variation in replication timing between alleles, we calculated the standard deviation (SD) of Repli-seq data across each allele for each 250 kb genomic window. Windows with an SD value ≥ 2.5 standard deviations above the genome-wide median were classified as outliers. This analysis identified 163 additional VERT regions (Supplementary Table 1). Figure 1e illustrates this analysis for both ACP (ACP6 and ACP7) and LCL (EB3-2 and GM12878) clones on chromosome 1, where windows are indicated as gray circles and outliers indicated in different colors as shown.

Next, to detect coding genes with AEI in ACP cells, we employed RNA-seq on

each clone and analyzed the number of reads at heterozygous SNPs. We defined AEI

as differential expression between alleles using recently established criteria [1,28]. Genes

were considered AEI-positive if they exhibited $\geq$80% allelic imbalance (AEI $\geq$ 0.80 or $\leq$

0.20) with FDR $\leq$ 0.05 in at least one clone. We selected the $\geq$80% allelic imbalance

threshold as our criterion for significant AEI because a recent study demonstrated that

allelic imbalance, as low as a 65%/35%, is enough to affect disease penetrance in

humans [28]. Additional analyses using more stringent thresholds (AEI $\geq$ 0.90 or $\geq$ 0.95)

are provided in Supplementary Table 2. In addition, to exclude imprinted genes and

effects due to genetic polymorphisms (i.e. expression Quantitative Trait Loci, eQTL[30])

we required that each AEI gene also be biallelically expressed (AEI < 0.70) in at least

one clone from the same individual with $\geq$ 20 informative reads. Using these criteria, we

identified 256 (2.4% of expressed genes) autosomal coding genes exhibiting AEI across

the two ACP clone sets (Supplementary Table 2).

In addition, we also sought to identify AEI of extremely long noncoding RNAs (i.e.

ASAR lincRNAs [1,5,7,31]) expressed in the ACP clones. For this analysis we refer to

regions of the genome with >50 kb of contiguous transcription of noncoding DNA as

Transcribed Loci (TL) [1]. Using the same criteria used for the coding genes described

above, we identified 23 TLs exhibiting AEI across the two ACP clone sets

(Supplementary Table 3).

Allelic Expression Imbalance and Variable Epigenetic Replication Timing define

Inactivation/Stability Centers.

By integrating the locations of genes exhibiting AEI with genomic regions displaying

VERT, we identified overlapping loci that define I/SCs. A schematic representation of a

hypothetical I/SC, showing a shaded VERT region in the replication profile and allele-

specific expression of coding and noncoding genes, is provided in Figure 1d. Figures 1f

and 1g illustrate a representative I/SC located on chromosome 1 at approximately 240 Mb

(marked by an arrow in Figure 1e), where VERT was detected across both sets of ACP

clones. Figure 1h shows the corresponding UCSC Genome Browser view of this locus,

with genes displaying AEI indicated by asterisks (see Supplementary Table 2).

Additional examples of I/SCs located on chromosomes 18, 6, and 9 are presented in

Figure 2. In total, we identified 112 loci where both VERT and AEI were detected at the

same genomic position, representing high-confidence I/SCs (Supplementary Table 4).

One of the objectives of this study was to assess whether different tissues utilize

distinct sets of I/SCs, as might be expected given that each cell type expresses a unique

repertoire of genes potentially subject to AEI. To explore this, we compared the genomic

locations of VERT regions identified in the two ACP clonal sets and the two LCL clonal

sets. As shown in Figure 1i, Venn diagrams reveal a modest degree of tissue-specific

overlap: ACP6 clones share more VERT regions with ACP7 than with either LCL clone

set, and the reciprocal relationship holds for the LCL clones. These findings suggest

that while a subset of I/SCs are shared across tissues, a modest component of tissue-

specific epigenetic regulation of allelic replication timing also exists.

At the molecular level, stochastic monoallelic expression contributes to cellular

individuality and introduces functional variability among individual cells within complex

biological systems [12-14]. This mechanism is particularly well characterized in the

olfactory, immune, and central nervous systems, where the expression of diverse

receptor-type molecules through random monoallelic expression confers cell-specific

functional identities. To further investigate the functional characteristics of protein-

coding genes located within VERT regions, we annotated the genes residing within

the VERT regions identified across all four clonal datasets (ACP clones: ACP6 and

ACP7; and LCL clones: EB3-2, and GM12878; see Supplementary Table 5). To assess

whether these genes are functionally related, we performed a Gene Ontology (GO)

enrichment analysis. This analysis revealed a significant enrichment for genes involved

in plasma membrane localization, cell-cell adhesion and neurodevelopmental

processes, the most significant hits for each set of clones are presented in Figure 1j

(see full details in Supplementary Table 6). This analysis suggests that I/SC-associated

epigenetic regulation may play a broader role in the establishment of cellular identity

and tissue organization particularly in the nervous system.

Gene clusters with AEI map to VERT regions.

In the olfactory, immune, and central nervous systems, random monoallelic

expression of gene family members, often organized in genomic clusters, plays a key

role in establishing cell-specific identities [12-14]. Consistent with these observations, we

found that VERT regions frequently coincide with gene clusters that contain known AEI

genes (Table 1). For example, Figure 3a and 3b highlight a cluster of 11 olfactory

receptor (OR) genes located on human chromosome 19 at approximately 15 Mb. This

region also harbors a cluster of Cytochrome P450 (CYP4F family, n = 6) genes.

The interspersed genomic arrangement of the CYP4F genes among the OR genes,

indicates that both gene families reside within the same I/SC and therefore both may be

subject to AEI.

A second example of a gene cluster is shown in Figure 3c and 3d, where

the major histocompatibility complex (MHC), comprising 16 HLA genes (which are

known to be subject to AEI [28,32]), lies within a VERT region. Notably, this locus also

contains a cluster of Lymphocyte Antigen 6 genes (LY6G family, n = 6), which are

interspersed among the HLA genes. This genome organization indicates that the LY6G

genes, like their neighboring HLA genes, are within the same I/SC and suggests that

they too may be subject to stochastic allelic expression imbalance**.**

Mapping mouse VERTs identifies known AEI gene clusters and syntenic I/SCs.

Much of our current understanding of stochastic autosomal allelic expression is derived

from studies conducted in the mouse [4,9-11,17,18,33-40]. To assess whether VERT

regions could be identified at expected loci in the mouse genome, specifically at sites of

known AEI, we analyzed previously published Repli-seq data from mouse pre-B cell

clones reported by Blumenfeld et al [41]. This analysis revealed 117 loci in the mouse

genome that display VERT (Supplemental Table 7).

Consistent with our findings in the human genome, the VERT regions detected in

the mouse genome frequently overlap with genes that are known to display AEI,

including 5 olfactory receptor gene clusters located on four different autosomes (Table

1). Figure 4 presents three examples: two olfactory receptor (Or) gene clusters on

mouse chromosome 2 (Figure 4a–4c) and the protocadherin (Pcdh) gene cluster, which

are known to generate cellular individuality in the central nervous system through

stochastic expression of different Pcdh family members [12,42], located on mouse

chromosome 18 (Figure 4d–4e). In addition to these gene clusters with VERT, we also

detected immunoglobulin and T cell receptor gene loci with VERT. As detailed in

Supplemental Figure 2, we detected VERT at the T cell receptor alpha (Tra) locus on

mouse chromosome 14 at approximately 54 Mb. One striking feature of this locus is that

it also contains an olfactory receptor cluster, containing 37 Or genes, and a

vomeronasal receptor gene (Vmn2r88). All of these multigene families are well

documented sites of AEI [3,12,17,42,43], highlighting the genomic organization and

conservation of I/SC-associated regulation at known AEI genes in the mouse. Gene

Ontology analysis of the mouse genes located within VERT regions showed significant

enrichment for biological processes related to the plasma membrane and cell-cell

adhesion (Figure 1j), which is consistent with the human VERT gene dataset and

reinforces the conservation of this regulatory mechanism across species.

Previous studies analyzed a relatively small number of mouse AEI genes to

model their human counterparts with respect to autosomal-dominant disorders [38,44]. To

further investigate the conservation of I/SC regulation between the human and mouse

genomes, we identified VERT regions that map to syntenic loci across the two species.

This analysis revealed 21 loci where VERT is detected in both genomes, as detailed in

Table 2. Figure 5 presents three representative examples of syntenic VERT regions: two

loci on the short arm of human chromosome 5 with corresponding regions on mouse

chromosome 15 (Figs. 5a–5g), and a third locus on human chromosome 16 with its syntenic

region on mouse chromosome 16 (Figs. 5h–5k). Additional examples of syntenic loci

showing AEI and/or VERT, including mouse App (see Gimelbrant et al. [45] for AEI of

human APP), Mocos and Fhod3 (see Figure 2a for VERT and AEI of human MOCOS

and FHOD3), and a cluster of 5 sodium channel genes, Scn1a, Scn2a, Scn3a, Scn7a

and Scn9a (see Figure 6a-c below for VERT and AEI of the syntenic human SCN gene

cluster) are shown in Supplemental Figure 3.

We point out that many of the mouse genes that are located within VERT regions

are associated with various human genetic diseases (Supplemental Table 8). One of

the loci highlighted in Figure 5a-5d labeled "hi", exhibits both VERT and AEI (Figure 5c

and [1]), affecting the protein-coding gene SEMA5A, a gene implicated in autism and

neurodevelopmental disorders [46-48]. Notably, this locus also contains a syntenic pair of

taste receptor genes: TAS2R1 in humans and Tas2r119 in mice. These genes are part

of the taste receptor family, which are known to harbor chromatin marks characteristic

of AEI-regulated loci [49]. Consistent with this observation, we also detected VERT at a

taste receptor gene cluster (Tas2r) on mouse chromosome 6 (Table 1).

A second example of synteny is shown in Figure 5f-5g, where human RETREG1

and mouse Retreg1 are located within VERT regions. Human RETREG1 is a known

AEI gene [45], and mutations in RETREG1 cause sensory and autonomic neuropathy [29].

A third example is shown in Figure 5h-5k, where both human GRIN2A and mouse

Grin2a are located within VERT regions, and loss of function mutations in GRIN2A in

humans and Grin2a in mice, are known to cause autosomal dominant (i.e.

haploinsufficiency) epilepsy [29,50,51]. These findings indicate that I/SC-associated

epigenetic regulation is conserved across species, affects functionally related genes

and gene families, and may influence both human and mouse genetic inheritance

patterns (see below).

Neurodevelopmental disease genes map to I/SCs.

To further investigate the relationship between I/SC regulation and human disease

genes, we annotated the human protein-coding genes located within VERT regions (see

Supplemental Table 5). This analysis identified 979 genes overlapping VERT regions, of

which 272 are associated with single-gene disorders according to the OMIM database

[29]. Among these genes, 155 are linked to autosomal recessive

diseases, 87 to autosomal dominant diseases, and 30 exhibit both dominant and

recessive inheritance patterns (Supplemental Table 9). Consistent with our GO

analysis, which revealed an enrichment for neurodevelopmental processes (see Figure

1j), we found that various neurodevelopmental disease genes were prominently

represented among the disease-associated genes (Table 3). Moreover, it was not

unexpected to observe gene clusters associated with neurodevelopmental disorders

residing within VERT regions. For example, Figure 6a–6c depicts the location, I/SC

profile, and genomic organization of a cluster of sodium channel genes (SCN family, n =

5). Mutations in SCN1A, SCN2A, SCN3A, and SCN9A genes are known to

cause autosomal dominant developmental and epileptic encephalopathies, with

haploinsufficiency as a primary mechanism (Table 3 and [29,52,53]). We also note that the

syntenic locus in the mouse displays VERT, with the 5 syntenic mouse Scn genes

residing within a VERT region on mouse chromosome 2 (Supplementary Figure 3e and

3f).

A second example of a cluster of epilepsy disease genes is shown in Figure 6d–

6e, where four members of the gamma-aminobutyric acid type A receptor gene family

(GABR family, n = 4) reside adjacent to a VERT region; notably, GABRA1 displays AEI

(Figure 6e, and Supplementary Table 2), and mutations in GABRA1, GABRB2, and

GABRG2 cause autosomal dominant developmental and epileptic encephalopathies

[29,54,55]. We also note that a second GABR gene cluster (n = 4), where GABRA2 and

GABRB1 are also associated with autosomal dominant developmental and epileptic

encephalopathies, is adjacent to a VERT region on chromosome 4 (see Table 1). Two

additional examples are shown in Figure 6g–6j, where OXR1 and SAMD12 are located

in different VERT regions on chromosome 8, and mutations in each result in autosomal

dominant epilepsy [29].

Another prominent group of human disease genes located within VERT

regions are those associated with Parkinson disease (Table 3). As shown in Figure 7,

four well-characterized Parkinson disease genes, SNCA**,** LRRK2**,** DNAJC6**,**

and PARKN, are located within or adjacent to VERT regions**.** Importantly,

both SNCA and LRRK2 were among the first genes identified as displaying AEI in the

study by Gimelbrant et al. in 2007 [45]. In our own dataset, we found

that DNAJC6 exhibits AEI in LCL clones [1]. We also highlight that JAK1, a gene involved

in immune signaling, lies within the same I/SC as DNAJC6 (Figure 7f). JAK1 is a known

AEI gene [56,57], and mutations in JAK1 cause autosomal dominant autoinflammation,

immune dysregulation, and eosinophilia (AIIDE). Importantly, a recent study indicated

that AEI of wild-type and mutant JAK1 alleles correlates with variable disease

penetrance [28].

Taken together, these findings underscore the role of I/SCs in regulating disease-

relevant gene clusters and emphasizes their conservation, synteny**,** and functional

relevance in both neurological and immune-related disorders. These examples highlight

how stochastic epigenetic regulation within I/SCs can contribute to phenotypic variability

and disease risk.

Discussion:

The co-occurrence of random monoallelic expression and asynchronous replication

between alleles of autosomal genes is well documented [2-4], yet its developmental

potential and contribution to human disease have not been systematically investigated.

In this study, we expand on the discovery and characterization of Inactivation/Stability

Centers (I/SCs), by identifying an additional 163 VERT regions using clonal lines of

primary ACP cells. This brings the total number of VERT regions to 363, covering

approximately 12% of the human genome. In addition, we identified 112 loci where we

detected both VERT and AEI, which represent high confidence I/SCs (Supplementary

Table 4).

To assess the evolutionary conservation of I/SC regulation, we analyzed

replication timing data from mouse pre-B cell clones, identifying 117 mouse VERT loci.

Many of these regions overlap with known mouse AEI genes and gene clusters

including olfactory, vomeronasal, taste and antigen receptors as well

as protocadherin gene families. Notably, we identified 21 syntenic VERT regions shared

between human and mouse genomes. These findings reinforce and extend previous

observations indicating a high degree of conservation of AEI genes [4] and suggest

that I/SCs are a fundamental and widespread feature of autosomal epigenetic regulation

in mammals.

I/SC regulation imparts tissue-relevant variation

Our clonal analysis of ACP and LCL cells, combined with haplotype-phased replication-

timing and expression profiling, allowed us to identify I/SCs that appear tissue-enriched

as well as a subset that is shared across tissues and conserved between species. Although

the number of clones and individuals examined here is limited, these observations are

consistent with a model in which some I/SCs serve broader or more ubiquitous

functions, while others are preferentially engaged in a tissue-dependent manner,

reflecting underlying differences in gene expression programs. Consistent with this

interpretation, comparisons of VERT regions revealed greater overlap within tissue type

(ACP6 vs. ACP7; EB3-2 vs. GM12878) than between tissues. However, given the

modest sample size and potential batch effects, these findings should be interpreted as

preliminary evidence suggesting, but not definitively establishing, a link between I/SC

activity and cellular identity. Expanded analyses across additional tissues, individuals,

and clonal lines will be required to rigorously define the extent of tissue specificity in

I/SC regulation.

Importantly, we identified 94 VERT regions shared across two or more clonal

sets, representing I/SCs that likely reflect core regulatory regions active across

individuals and cell types (Figure 1i). As detailed in Supplemental Table 1, we

identified VERT regions detected exclusively in single clone sets. These private VERT

regions may reflect inter-individual variability in epigenetic regulation, context-dependent

I/SC usage, but more likely reflect a limitation of sampling—i.e., regions that would be

detected in additional clones if a larger number were analyzed. Thus, while our findings

provide strong support for both shared and tissue specific I/SCs, they also underscore

the importance of clone number and diversity in defining the full landscape of autosomal

epigenetic regulation.

In the context of development, AEI introduces stochastic variability in gene

dosage across clonal populations, contributing to cellular heterogeneity during lineage

specification and tissue formation. Such heterogeneity could serve as a substrate

for cellular selection, enabling the developing tissue to fine-tune function or respond

adaptively to environmental cues. This model aligns with observations that AEI often

affects genes involved in cell adhesion, signal transduction, and neurodevelopment,

where even small variations in expression levels could significantly impact cell fate and

connectivity.

At the molecular level, I/SCs contribute to cellular individuality by

enabling stochastic monoallelic expression of specific genes. This mechanism is

particularly critical in biological systems that require diversity at the single-cell level,

such as the olfactory, vomeronasal, immune, and nervous systems. Consistent with

this, many I/SCs coincide with gene clusters, including olfactory, vomeronasal, and

antigen receptor genes, as well as HLA and protocadherin genes, all of which are

known to undergo stochastic monoallelic expression. We identified numerous additional

gene clusters, e.g. taste receptors, residing within I/SCs, suggesting that these other

gene families may also participate in tissue mosaicism and/or cellular individuality. Our

current model proposes that I/SC regulation is integral to tissue development,

where stochastic, epigenetically driven allelic regulation introduces phenotypic

variability among cells. This variability may promote functional diversity and selective

advantage during development. Under this model, each tissue would engage a distinct

but partially overlapping set of I/SC-regulated genes, aligned with its

unique transcriptional and functional requirements. This framework positions I/SCs as

key players in promoting cellular individuality, tissue mosaicism, and developmental

plasticity.

Many of the genes that reside within I/SCs are regulated by allelic exclusion.

Allelic exclusion is a regulatory mechanism by which only one allele of a gene is

expressed while the other is actively silenced, ensuring monoallelic expression in each

cell. This process is well characterized in the immune system, where it plays a critical

role in B and T cell receptor gene rearrangements, ensuring that each lymphocyte

produces a single antigen receptor [18]. Furthermore, functional monoallelic expression is

achieved via a secondary mechanism that involves product inhibition. For example, in

developing B cells, expression of a productive immunoglobulin heavy chain allele

suppresses rearrangement and expression of the second allele, maintaining clonal

specificity [18]. Allelic exclusion has also been observed in olfactory receptors, where

product feedback ensures expression of a single allele of only one member of a large

gene family, ensuring that each olfactory receptor neuron responds to a single odorant

[2,33,58-61].

I/SC regulation acts through mechanisms distinct from X inactivation.

Numerous studies in both human and mouse cells support the notion that allele-specific

epigenetic regulation of autosomal protein-coding genes is widespread

[4,11,13,37,38,40,58,62,63]. Although stochastic monoallelic expression on autosomes may

appear similar to X chromosome inactivation (XCI), the mechanisms behind them are

distinct: First, XCI is initiated in preimplantation embryos, as early as the 4-8 cell stage

in the mouse and 8-16 cell stage in humans [64-69], and later becomes established during

differentiation of the inner cell mass of the blastocyst in both species [68,70]. In contrast,

autosomal stochastic monoallelic expression appears later in development, with allelic

biases that can be erased and reestablished during differentiation [10,38]. For example, a

recent study found that allele-specific autosomal inactivation is absent during early

blood lineage specification but detectable in hematopoietic stem cells and their

differentiated progeny [9]. Second, in XCI, there is strong coordination along the

chromosomes: genes on the active X chromosome share both expression and early

replication profiles, whereas genes on the inactive X show silencing and late replication.

In contrast, autosomal inactivation lacks such coordination: allelic expression and

replication timing are unlinked within a chromosome and independent across the

genome [1,7,71]. Taken together, these observations indicate that autosomal allelic

inactivation is established through a mechanism that is fundamentally distinct from XCI

and arises at a different developmental stage, underscoring that the two processes

represent separate epigenetic systems.

An alternative model for haploinsufficiency

The relationship between genotype and phenotype is fundamental to understanding

human genetic disease. At the scale of the whole genome, loss of function alleles are

generally not deleterious when heterozygous and are therefore haplosufficient [21,24]. Not

surprisingly, most of the loss of function mutations in single genes in humans are

recessive [21,29]. Therefore, for most human disease genes a single normal allele is

sufficient to achieve the typical phenotype. Dominantly inherited disorders comprise a

diverse group of conditions that are often severe and poorly understood, except in

cases of gain-of-function mutations, where dominant negative or interfering activities

induce the abnormal phenotype. However, many dominantly inherited diseases result

from copy number changes, either gain or loss of wild-type alleles [29,72].

Haploinsufficiency, predicted to apply to as much as 40% of the human genome [19,20],

refers to a heterozygous deletion or complete loss-of-function mutation that results in an

altered phenotype, indicating that a single normal allele is insufficient. The

underlying molecular mechanisms responsible for most haploinsufficient human disease

genes remain largely unknown.

The existence of two distinct populations of cells in XX individuals, one

expressing genes from the maternally derived X chromosome and the other from the

paternally derived X, has profound implications for both development and

the manifestation of X-linked diseases, where both dominant and recessive inheritance

patterns are observed in females [29]. A key factor influencing these inheritance patterns

is the presence or absence of cellular selection during development. In certain X-linked

disorders, strong selection favors the survival or expansion of cells in which the mutant

allele is on the inactive X chromosome, resulting in a predominance of wild-type cells

and a clinical presentation consistent with X-linked recessive inheritance [21,26]. In

contrast, when cellular selection is absent, heterozygous females typically retain

a ~50:50 mosaic of wild-type and mutant-expressing cells. The presence of mutant-

expressing cells, essentially null cells, is thought to underlie X-linked dominant

phenotypes [27,73]. In such cases, heterozygous loss of function mutations leads to a

condition of haploinsufficiency, in which the presence of a single normal allele is

insufficient to maintain normal function. Interestingly, although single X-linked alleles are

sufficient in males, the mosaic expression pattern in females unmasks phenotypic

consequences that are otherwise hidden. This highlights an underappreciated

mechanism of haploinsufficiency: the cellular mosaicism inherent to X inactivation can

reduce effective gene dosage below functional thresholds in individuals with

heterozygous mutations, particularly in tissues where cells expressing the mutant (null)

allele persist.

One of the most striking implications of I/SC regulation is its relevance to human

genetic disease, particularly those involving haploinsufficiency. Among the 979

genes mapped within human VERT regions, 272 are associated with monogenic

diseases, including 87 with autosomal dominant inheritance, many of which are

classified as haploinsufficient [29]. Our findings offer a new mechanistic

model for autosomal haploinsufficiency: in the absence of compensatory upregulation

and/or cellular selection mechanisms, stochastic allelic inactivation results in or expands

a functionally null cell population, even when one allele is wild type. This mirrors the

situation observed in X-linked dominant disorders with loss of function mutations, where

the absence of cellular selection results in the persistence of null cells. This model

expands upon prior theories of haploinsufficiency that assume strict biallelic expression,

which attribute dominant phenotypes to factors such as threshold-dependent

expression, stoichiometric imbalance in multiprotein complexes, or inability to buffer

expression noise [22-25]. In addition, our model fits nicely with a previous study that

modeled stochastic gene expression of disease genes and proposed "haploinsufficiency

syndromes might result from an increased susceptibility to stochastic delays of gene

initiation or interruptions of gene expression" [74]. Our data suggest that epigenetically

regulated stochastic AEI can contribute to pathogenic phenotypes in heterozygous

individuals by either reducing the number of functional cells below a critical disease

threshold or by disrupting the normal mosaicism of gene expression within disease-

relevant tissues. This expands the conceptual framework of haploinsufficiency to include epigenetic, rather than purely genetic, sources of dosage reduction.

Limitations of our study.

While our findings provide substantial evidence for the existence and functional relevance of I/SCs, several important limitations remain to be addressed. First, the number of clones analyzed in our studies, while sufficient to identify robust patterns of AEI and VERT, limits our ability to capture the full spectrum of stochastic epigenetic states across the genome. For example, we frequently observe AEI of genes located adjacent to VERT regions, indicating that the boundaries of the epigenetic mechanisms functioning at I/SCs remain undefined. This raises important questions about how far the influence of I/SC-associated regulation extends beyond the VERT region. In addition, the specific epigenetic marks, including DNA methylation, histone modifications, and chromatin accessibility, at key regulatory elements such as replication origins, enhancers, and promoters within I/SCs have yet to be defined. Furthermore, the higher-order chromatin organization of I/SCs, including their relationship to topologically associating domains (TADs) and A/B compartments [49,75], remains largely unexplored and may hold critical insights into their regulatory architecture and functional constraints. Second, our analysis was conducted in a restricted number of cell types, namely LCLs and ACPs, and therefore does not represent the full diversity of I/SC usage across human tissues. Third, our study was limited to a small number of individuals, which may constrain the identification of inter-individual variability in I/SC regulation. Fourth, the developmental timing of I/SC establishment remains unresolved; it is unclear when during lineage commitment or differentiation these regions acquire their allele-specific states. A related and equally

important question is whether I/SCs can be reset during reprogramming into induced

pluripotent stem cells (iPSCs) and subsequently reestablished upon differentiation. This

issue is particularly relevant given the widespread use of iPSCs in human disease

modeling and regenerative medicine [76-78]. Understanding how I/SCs behave during

cellular reprogramming and lineage specification will be critical for interpreting AEI-

related phenotypes in iPSC-derived systems.

An additional open question is how the magnitude of stochastic

AEI influences disease states, particularly in the context of variable

penetrance and expressivity. For example, a recent study employed allelic expression

ratios to identify functionally relevant AEI in genes associated with inborn errors of

immunity, showing that individuals with the same pathogenic mutation can

exhibit divergent clinical outcomes depending on the degree of allelic imbalance [28].

These findings suggest that AEI can modulate disease severity, acting as an epigenetic

layer of regulation that determines whether a given mutation manifests clinically. We

anticipate that the biological significance of AEI will vary depending on the function of

the gene, its dosage sensitivity, and the cellular context in which it is expressed.

Addressing these questions will be essential for achieving a comprehensive

understanding of the formation, function, and disease relevance of I/SCs in human

biology.

Conclusion

In conclusion, our work extends the catalog of I/SCs in the human genome and

supports a model in which stochastic, allele-specific epigenetic regulation at autosomal

loci contributes to cellular diversity and disease susceptibility. Moreover, by

identifying syntenic I/SCs in the mouse genome, our findings greatly enhance the potential utility of mouse models for mechanistic studies of disease-causing mutations at conserved loci. Our findings present a potential mechanism for incomplete penetrance, variable expressivity, and tissue-specific vulnerability, particularly in disorders caused by haploinsufficiency. In such cases, stochastic silencing of the wild-type allele in a subset of cells may lead to functional null states, even in heterozygous individuals, reducing the effective number of functional cells below a critical threshold. This may be especially relevant in the nervous system, where precise gene dosage is often essential for maintaining neuronal function. Moreover, I/SC regulation at disease genes raises the possibility that stochastic epigenetic regulation could act as an epigenetic modifier of disease risk, helping to explain why some individuals carrying pathogenic mutations remain asymptomatic. Understanding the full scope of I/SC regulation in both normal development and disease pathogenesis will be essential for identifying novel regulatory mechanisms and potential therapeutic targets. As tools for single-cell and allele-resolved epigenomics continue to advance, I/SCs may emerge as key elements in both basic biology and precision medicine.

RESOURCE AVAILABILITY

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Mathew Thayer (thayerm@ohsu.edu).

Materials availability statement

All data generated or analyzed during this study are included in the manuscript and supporting files. The Repli-seq and RNA-seq raw sequencing data generated in this study have been deposited in the European Nucleotide Archive database under accession code PRJEB110291 and are available without restriction.

Acknowledgments:

596   M.J.T. was supported by NIH (R01GM130703, and RF1NS142814).
597   M.B.H was supported by NIH NCI 4K00CA245677-03
598   DMG was supported by NIH R01GM083337
599

Author Contribution Statement

MBH, MJT, and DMG conceived of the experimental design for the RNA expression and
replication timing assays. KF performed the surgeries to obtain the amputed digits. BJ
and PAY performed the ACP cultures and clone generation. MBH and AEV carried out
the RNA-seq assays, including data analysis. MBH and AEV carried out the Repli-seq
assays, including data analysis. PFC and PTS provided advice during preparation of the
manuscript. MBH, DMG and MJT wrote the manuscript.
607

Declaration of interests

609   The authors declare no competing interests.

610
611

Literature Cited:

1. Heskett, M.B., Vouzas, A.E., Smith, L.G., Yates, P.A., Boniface, C., Bouhassira, E.E., Spellman, P.T., Gilbert, D.M., and Thayer, M.J. (2022). Epigenetic control of chromosome-associated lncRNA genes essential for replication and stability. Nature communications *13*, 6301. 10.1038/s41467-022-34099-7.

2. Goldmit, M., and Bergman, Y. (2004). Monoallelic gene expression: a repertoire of recurrent themes. Immunol Rev *200*, 197-214.

3. Chess, A., Simon, I., Cedar, H., and Axel, R. (1994). Allelic inactivation regulates olfactory receptor gene expression. Cell *78*, 823-834. 10.1016/s0092-8674(94)90562-2.

4. Chess, A. (2016). Monoallelic Gene Expression in Mammals. Annu Rev Genet *50*, 317-327. 10.1146/annurev-genet-120215-035120.

5. Stoffregen, E.P., Donley, N., Stauffer, D., Smith, L., and Thayer, M.J. (2011). An autosomal locus that controls chromosome-wide replication timing and mono-allelic expression. Hum Mol Genet *20*, 2366-2378.

6. Thayer, M.J. (2012). Mammalian chromosomes contain cis-acting elements that control replication timing, mitotic condensation, and stability of entire chromosomes. Bioessays *34*, 760-770.

7. Donley, N., Smith, L., and Thayer, M.J. (2015). ASAR15, A cis-Acting Locus that Controls Chromosome-Wide Replication Timing and Stability of Human Chromosome 15. PLoS Genet *11*, e1004923. 10.1371/journal.pgen.1004923.

8. Crowley, J.J., Zhabotynsky, V., Sun, W., Huang, S., Pakatci, I.K., Kim, Y., Wang, J.R., Morgan, A.P., Calaway, J.D., Aylor, D.L., et al. (2015). Analyses of allele-specific gene expression in highly divergent mouse crosses identifies pervasive allelic imbalance. Nat Genet *47*, 353-360. 10.1038/ng.3222.

9. Kubasova, N., Alves-Pereira, C.F., Gupta, S., Vinogradova, S., Gimelbrant, A., and Barreto, V.M. (2022). In Vivo Clonal Analysis Reveals Random Monoallelic Expression in Lymphocytes That Traces Back to Hematopoietic Stem Cells. Frontiers in cell and developmental biology *10*. 10.3389/fcell.2022.827774.

10. Marion-Poll, L., Forêt, B., Zielinski, D., Massip, F., Attia, M., Carter, A.C., Syx, L., Chang, H.Y., Gendrel, A.V., and Heard, E. (2021). Locus specific epigenetic modalities of random allelic expression imbalance. Nature communications *12*, 5330. 10.1038/s41467-021-25630-3.

11. Zwemer, L.M., Zak, A., Thompson, B.R., Kirby, A., Daly, M.J., Chess, A., and Gimelbrant, A.A. (2012). Autosomal monoallelic expression in the mouse. Genome Biol *13*, R10. 10.1186/gb-2012-13-2-r10.

12. Yagi, T. (2013). Genetic basis of neuronal individuality in the mammalian brain. Journal of neurogenetics *27*, 97-105. 10.3109/01677063.2013.801969.

13. Savova, V., Chun, S., Sohail, M., McCole, R.B., Witwicki, R., Gai, L., Lenz, T.L., Wu, C.T., Sunyaev, S.R., and Gimelbrant, A.A. (2016). Genes with monoallelic expression contribute disproportionately to genetic diversity in humans. Nat Genet *48*, 231-237. 10.1038/ng.3493.

14. Savova, V., Vigneau, S., and Gimelbrant, A.A. (2013). Autosomal monoallelic expression: genetics of epigenetic diversity? Curr Opin Genet Dev *23*, 642-648. 10.1016/j.gde.2013.09.001.

15. Johnstone, B., Stoddart, M.J., and Im, G.I. (2020). Multi-Disciplinary Approaches for Cell-Based Cartilage Regeneration. J Orthop Res *38*, 463-472. 10.1002/jor.24458.

660   16.   Pattappa, G., Reischl, F., Jahns, J., Schewior, R., Lang, S., Zellner, J.,
661   Johnstone, B., Docheva, D., and Angele, P. (2021). Fibronectin Adherent Cell
662   Populations Derived From Avascular and Vascular Regions of the Meniscus
663   Have Enhanced Clonogenicity and Differentiation Potential Under Physioxia.
664   Front Bioeng Biotechnol *9*, 789621. 10.3389/fbioe.2021.789621.
665   17.   Rodriguez, I., Feinstein, P., and Mombaerts, P. (1999). Variable patterns of
666   axonal projections of sensory neurons in the mouse vomeronasal system. Cell
667   *97*, 199-208. 10.1016/s0092-8674(00)80730-8.
668   18.   Mostoslavsky, R., Singh, N., Tenzen, T., Goldmit, M., Gabay, C., Elizur, S., Qi,
669   P., Reubinoff, B.E., Chess, A., Cedar, H., and Bergman, Y. (2001).
670   Asynchronous replication and allelic exclusion in the immune system. Nature
671   *414*, 221-225.
672   19.   Backman, J.D., Li, A.H., Marcketta, A., Sun, D., Mbatchou, J., Kessler, M.D.,
673   Benner, C., Liu, D., Locke, A.E., Balasubramanian, S., et al. (2021). Exome
674   sequencing and analysis of 454,787 UK Biobank participants. Nature *599*, 628-
675   634. 10.1038/s41586-021-04103-z.
676   20.   Jurgens, S.J., Wang, X., Choi, S.H., Weng, L.C., Koyama, S., Pirruccello, J.P.,
677   Nguyen, T., Smadbeck, P., Jang, D., Chaffin, M., et al. (2024). Rare coding
678   variant analysis for human diseases across biobanks and ancestries. Nat Genet
679   *56*, 1811-1820. 10.1038/s41588-024-01894-5.
680   21.   Zschocke, J., Byers, P.H., and Wilkie, A.O.M. (2023). Mendelian inheritance
681   revisited: dominance and recessiveness in medical genetics. Nat Rev Genet *24*,
682   442-463. 10.1038/s41576-023-00574-0.
683   22.   Huang, N., Lee, I., Marcotte, E.M., and Hurles, M.E. (2010). Characterising and
684   predicting haploinsufficiency in the human genome. PLoS Genet *6*, e1001154.
685   10.1371/journal.pgen.1001154.
686   23.   Morrill, S.A., and Amon, A. (2019). Why haploinsufficiency persists. Proc Natl
687   Acad Sci U S A *116*, 11866-11871. 10.1073/pnas.1900437116.
688   24.   Wilkie, A.O. (1994). The molecular basis of genetic dominance. J Med Genet *31*,
689   89-98. 10.1136/jmg.31.2.89.
690   25.   Veitia, R.A., Zschocke, J., and Birchler, J.A. (2025). Gene Dosage Sensitivity and
691   Human Genetic Diseases. J Inherit Metab Dis *48*, e70058. 10.1002/jimd.70058.
692   26.   Minks, J., Robinson, W.P., and Brown, C.J. (2008). A skewed view of X
693   chromosome inactivation. J Clin Invest *118*, 20-23. 10.1172/jci34470.
694   27.   Boone, P.M., Buenaventura, T., King, J.W.D., and Merkenschlager, M. (2025). X-
695   linked competition - implications for human development and disease. Nat Rev
696   Genet. 10.1038/s41576-025-00840-3.
697   28.   Stewart, O., Gruber, C., Randolph, H.E., Patel, R., Ramba, M., Calzoni, E.,
698   Huang, L.H., Levy, J., Buta, S., Lee, A., et al. (2025). Monoallelic expression can
699   govern penetrance of inborn errors of immunity. Nature *637*, 1186-1197.
700   10.1038/s41586-024-08346-4.
701   29.   McKusick, V. (2025). Online Mendelian Inheritance in Man, OMIM®.
702   30.   Gilad, Y., Rifkin, S.A., and Pritchard, J.K. (2008). Revealing the architecture of
703   gene regulation: the promise of eQTL studies. Trends Genet *24*, 408-415.
704   10.1016/j.tig.2008.06.001.
705   31.   Heskett, M.B., Smith, L.G., Spellman, P., and Thayer, M.J. (2020). Reciprocal
706   monoallelic expression of ASAR lncRNA genes controls replication timing of

human chromosome 6. RNA (New York, N.Y.) *26*, 724-738.
10.1261/rna.073114.119.

32.   Fu, R., Qin, P., Zou, X., Hu, Z., Hong, N., Wang, Y., and Jin, W. (2021). A
Comprehensive Characterization of Monoallelic Expression During
Hematopoiesis and Leukemogenesis via Single-Cell RNA-Sequencing. Frontiers
in cell and developmental biology *9*, 702897. 10.3389/fcell.2021.702897.

33.   Zakharova, I.S., Shevchenko, A.I., and Zakian, S.M. (2009). Monoallelic gene
expression in mammals. Chromosoma *118*, 279-290.

34.   Xu, J., Carter, A.C., Gendrel, A.V., Attia, M., Loftus, J., Greenleaf, W.J.,
Tibshirani, R., Heard, E., and Chang, H.Y. (2017). Landscape of monoallelic
DNA accessibility in mouse embryonic stem cells and neural progenitor cells. Nat
Genet *49*, 377-386. 10.1038/ng.3769.

35.   Nag, A., Savova, V., Fung, H.L., Miron, A., Yuan, G.C., Zhang, K., and
Gimelbrant, A.A. (2013). Chromatin signature of widespread monoallelic
expression. eLife *2*, e01256. 10.7554/eLife.01256.

36.   Li, S.M., Valo, Z., Wang, J., Gao, H., Bowers, C.W., and Singer-Sam, J. (2012).
Transcriptome-wide survey of mouse CNS-derived cells reveals monoallelic
expression within novel gene families. PLoS One *7*, e31751.

37.   Gendrel, A.V., Marion-Poll, L., Katoh, K., and Heard, E. (2016). Random
monoallelic expression of genes on autosomes: Parallels with X-chromosome
inactivation. Semin Cell Dev Biol. 10.1016/j.semcdb.2016.04.007.

38.   Gendrel, A.V., Attia, M., Chen, C.J., Diabangouaya, P., Servant, N., Barillot, E.,
and Heard, E. (2014). Developmental dynamics and disease potential of random
monoallelic gene expression. Dev Cell *28*, 366-380.
10.1016/j.devcel.2014.01.016.

39.   Eckersley-Maslin, M.A., Thybert, D., Bergmann, J.H., Marioni, J.C., Flicek, P.,
and Spector, D.L. (2014). Random monoallelic gene expression increases upon
embryonic stem cell differentiation. Dev Cell *28*, 351-365.
10.1016/j.devcel.2014.01.017.

40.   Barreto, V.M., Kubasova, N., Alves-Pereira, C.F., and Gendrel, A.V. (2021). X-
Chromosome Inactivation and Autosomal Random Monoallelic Expression as
"Faux Amis". Frontiers in cell and developmental biology *9*, 740937.
10.3389/fcell.2021.740937.

41.   Blumenfeld, B., Masika, H., Farago, M., Yehuda, Y., Halaseh, L., Vardi, O.,
Rapoport, R., Levin-Klein, R., Cedar, H., Bergman, Y., and Simon, I. (2021).
Chromosomal coordination and differential structure of asynchronous replicating
regions. Nature communications *12*, 1035. 10.1038/s41467-021-21348-4.

42.   Mountoufaris, G., Chen, W.V., Hirabayashi, Y., O'Keeffe, S., Chevee, M.,
Nwakeze, C.L., Polleux, F., and Maniatis, T. (2017). Multicluster Pcdh diversity is
required for mouse olfactory neural circuit assembly. Science *356*, 411-414.
10.1126/science.aai8801.

43.   Williams, D.L., Sikora, V.M., Hammer, M.A., Amin, S., Brinjikji, T., Brumley, E.K.,
Burrows, C.J., Carrillo, P.M., Cromer, K., Edwards, S.J., et al. (2021). May the
Odds Be Ever in Your Favor: Non-deterministic Mechanisms Diversifying Cell
Surface Molecule Expression. Frontiers in cell and developmental biology *9*,
720798. 10.3389/fcell.2021.720798.

44.   Savova, V., Vinogradova, S., Pruss, D., Gimelbrant, A.A., and Weiss, L.A.
(2017). Risk alleles of genes with monoallelic expression are enriched in gain-of-

function variants and depleted in loss-of-function variants for
neurodevelopmental disorders. Mol Psychiatry *22*, 1785-1794.
10.1038/mp.2017.13.
45. Gimelbrant, A., Hutchinson, J.N., Thompson, B.R., and Chess, A. (2007).
Widespread monoallelic expression on human autosomes. Science *318*, 1136-
1140.
46. Ito, H., Morishita, R., and Nagata, K.I. (2017). Autism spectrum disorder-
associated genes and the development of dentate granule cells. Med Mol
Morphol *50*, 123-129. 10.1007/s00795-017-0161-z.
47. Lin, L., Lesnick, T.G., Maraganore, D.M., and Isacson, O. (2009). Axon guidance
and synaptic maintenance: preclinical markers for neurodegenerative disease
and therapeutics. Trends Neurosci *32*, 142-149. 10.1016/j.tins.2008.11.006.
48. Mosca-Boidron, A.L., Gueneau, L., Huguet, G., Goldenberg, A., Henry, C., Gigot,
N., Pallesi-Pocachard, E., Falace, A., Duplomb, L., Thevenon, J., et al. (2016). A
de novo microdeletion of SEMA5A in a boy with autism spectrum disorder and
intellectual disability. European journal of human genetics : EJHG *24*, 838-843.
10.1038/ejhg.2015.211.
49. Richer, S., Tian, Y., Schoenfelder, S., Hurst, L., Murrell, A., and Pisignano, G.
(2023). Widespread allele-specific topological domains in the human genome are
not confined to imprinted gene clusters. Genome Biol *24*, 40. 10.1186/s13059-
023-02876-2.
50. Reutlinger, C., Helbig, I., Gawelczyk, B., Subero, J.I., Tönnies, H., Muhle, H.,
Finsterwalder, K., Vermeer, S., Pfundt, R., Sperner, J., et al. (2010). Deletions in
16p13 including GRIN2A in patients with intellectual disability, various
dysmorphic features, and seizure disorders of the rolandic region. Epilepsia *51*,
1870-1873. 10.1111/j.1528-1167.2010.02555.x.
51. Camp, C.R., Vlachos, A., Klöckner, C., Krey, I., Banke, T.G., Shariatzadeh, N.,
Ruggiero, S.M., Galer, P., Park, K.L., Caccavano, A., et al. (2023). Loss of
Grin2a causes a transient delay in the electrophysiological maturation of
hippocampal parvalbumin interneurons. Commun Biol *6*, 952. 10.1038/s42003-
023-05298-9.
52. Ding, J., Li, X., Tian, H., Wang, L., Guo, B., Wang, Y., Li, W., Wang, F., and Sun,
T. (2021). SCN1A Mutation-Beyond Dravet Syndrome: A Systematic Review and
Narrative Synthesis. Front Neurol *12*, 743726. 10.3389/fneur.2021.743726.
53. Wolff, M., Johannesen, K.M., Hedrich, U.B.S., Masnada, S., Rubboli, G.,
Gardella, E., Lesca, G., Ville, D., Milh, M., Villard, L., et al. (2017). Genetic and
phenotypic heterogeneity suggest therapeutic implications in SCN2A-related
disorders. Brain *140*, 1316-1336. 10.1093/brain/awx054.
54. Hernandez, C.C., Hu, N., Shen, W., and Macdonald, R.L. (2023). Epileptic
Encephalopathy GABRB Structural Variants Share Common Gating and
Trafficking Defects. Biomolecules *13*. 10.3390/biom13121790.
55. Hamdan, F.F., Myers, C.T., Cossette, P., Lemay, P., Spiegelman, D., Laporte,
A.D., Nassif, C., Diallo, O., Monlong, J., Cadieux-Dion, M., et al. (2017). High
Rate of Recurrent De Novo Mutations in Developmental and Epileptic
Encephalopathies. Am J Hum Genet *101*, 664-685. 10.1016/j.ajhg.2017.09.008.
56. Gruber, C.N., Calis, J.J.A., Buta, S., Evrony, G., Martin, J.C., Uhl, S.A., Caron,
R., Jarchin, L., Dunkin, D., Phelps, R., et al. (2020). Complex Autoinflammatory
Syndrome Unveils Fundamental Principles of JAK1 Kinase Transcriptional and

Biochemical Function. Immunity *53*, 672-684.e611.
10.1016/j.immuni.2020.07.006.

57. Ross, S.H., and Cantrell, D.A. (2020). Of Mosaicism and Mechanisms: How
JAK1 Goes Awry. Immunity *53*, 481-484. 10.1016/j.immuni.2020.08.010.

58. Khamlichi, A.A., and Feil, R. (2018). Parallels between Mammalian Mechanisms
of Monoallelic Gene Expression. Trends Genet *34*, 954-971.
10.1016/j.tig.2018.08.005.

59. Dalton, R.P., Lyons, D.B., and Lomvardas, S. (2013). Co-opting the unfolded
protein response to elicit olfactory receptor feedback. Cell *155*, 321-332.
10.1016/j.cell.2013.09.033.

60. Lyons, D.B., Allen, W.E., Goh, T., Tsai, L., Barnea, G., and Lomvardas, S.
(2013). An epigenetic trap stabilizes singular olfactory receptor expression. Cell
154, 325-336. 10.1016/j.cell.2013.06.039.

61. Pourmorady, A.D., Bashkirova, E.V., Chiariello, A.M., Belagzhal, H., Kodra, A.,
Duffié, R., Kahiapo, J., Monahan, K., Pulupa, J., Schieren, I., et al. (2024). RNA-
mediated symmetry breaking enables singular olfactory receptor choice. Nature
625, 181-188. 10.1038/s41586-023-06845-4.

62. Chess, A. (2012). Mechanisms and consequences of widespread random
monoallelic expression. Nat Rev Genet *13*, 421-428.

63. Reinius, B., and Sandberg, R. (2015). Random monoallelic expression of
autosomal genes: stochastic transcription and allele-level regulation. Nat Rev
Genet *16*, 653-664. 10.1038/nrg3888.

64. Brockdorff, N., and Turner, B.M. (2015). Dosage compensation in mammals.
Cold Spring Harb Perspect Biol *7*, a019406. 10.1101/cshperspect.a019406.

65. Dossin, F., and Heard, E. (2022). The Molecular and Nuclear Dynamics of X-
Chromosome Inactivation. Cold Spring Harb Perspect Biol *14*.
10.1101/cshperspect.a040196.

66. Galupa, R., and Heard, E. (2018). X-Chromosome Inactivation: A Crossroads
Between Chromosome Architecture and Gene Regulation. Annu Rev Genet *52*,
535-566. 10.1146/annurev-genet-120116-024611.

67. Payer, B., and Lee, J.T. (2008). X chromosome dosage compensation: how
mammals keep the balance. Annu Rev Genet *42*, 733-772.
10.1146/annurev.genet.42.110807.091711 [doi].

68. van den Berg, I.M., Galjaard, R.J., Laven, J.S., and van Doorninck, J.H. (2011).
XCI in preimplantation mouse and human embryos: first there is remodelling….
Hum Genet *130*, 203-215. 10.1007/s00439-011-1014-9.

69. Werner, J.M., Hover, J., and Gillis, J. (2024). Population variability in X-
chromosome inactivation across 10 mammalian species. Nature communications
15, 8991. 10.1038/s41467-024-53449-1.

70. van den Berg, I.M., Laven, J.S., Stevens, M., Jonkers, I., Galjaard, R.J., Gribnau,
J., and van Doorninck, J.H. (2009). X chromosome inactivation is initiated in
human preimplantation embryos. Am J Hum Genet *84*, 771-779.
10.1016/j.ajhg.2009.05.003.

71. Donley, N., Stoffregen, E.P., Smith, L., Montagna, C., and Thayer, M.J. (2013).
Asynchronous Replication, Mono-Allelic Expression, and Long Range Cis-Effects
of ASAR6. PLoS Genet *9*, e1003423.

72. Rehm, H.L., Berg, J.S., Brooks, L.D., Bustamante, C.D., Evans, J.P., Landrum,
M.J., Ledbetter, D.H., Maglott, D.R., Martin, C.L., Nussbaum, R.L., et al. (2015).

ClinGen--the Clinical Genome Resource. N Engl J Med *372*, 2235-2242.
10.1056/NEJMsr1406261.

73.   Giovenino, C., Trajkova, S., Pavinato, L., Cardaropoli, S., Pullano, V., Ferrero, E., Sukarova-Angelovska, E., Carestiato, S., Salmin, P., Rinninella, A., et al. (2023). Skewed X-chromosome inactivation in unsolved neurodevelopmental disease cases can guide re-evaluation For X-linked genes. European journal of human genetics : EJHG *31*, 1228-1236. 10.1038/s41431-023-01324-w.

74.   Cook, D.L., Gerber, A.N., and Tapscott, S.J. (1998). Modeling stochastic gene expression: implications for haploinsufficiency. Proc Natl Acad Sci U S A *95*, 15641-15646. 10.1073/pnas.95.26.15641.

75.   Pope, B.D., Ryba, T., Dileep, V., Yue, F., Wu, W., Denas, O., Vera, D.L., Wang, Y., Hansen, R.S., Canfield, T.K., et al. (2014). Topologically associating domains are stable units of replication-timing regulation. Nature *515*, 402-405. 10.1038/nature13986.

76.   Li, Y., and Darabi, R. (2022). Role of epigenetics in cellular reprogramming; from iPSCs to disease modeling and cell therapy. J Cell Biochem *123*, 147-154. 10.1002/jcb.30164.

77.   Janiszewski, A., Talon, I., Chappell, J., Collombet, S., Song, J., De Geest, N., To, S.K., Bervoets, G., Marin-Bejar, O., Provenzano, C., et al. (2019). Dynamic reversal of random X-Chromosome inactivation during iPSC reprogramming. Genome Res *29*, 1659-1672. 10.1101/gr.249706.119.

78.   Tcw, J. (2019). Human iPSC application in Alzheimer's disease and Tau-related neurodegenerative diseases. Neurosci Lett *699*, 31-40. 10.1016/j.neulet.2019.01.043.

79.   Pattappa, G., Markway, B.D., Docheva, D., and Johnstone, B. (2023). Physioxic Culture of Chondrogenic Cells. Methods Mol Biol *2598*, 45-63. 10.1007/978-1-0716-2839-3_5.

80.   Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics (Oxford, England) *25*, 1754-1760. 10.1093/bioinformatics/btp324.

81.   Garrison, E.a.M., G (2024). Haplotype-based variant detection from short-read sequencing. arXiv *1207.3907v2*.

82.   Martin, M., Ebert, P., and Marschall, T. (2023). Read-Based Phasing and Analysis of Phased Variants with WhatsHap. Methods Mol Biol *2590*, 127-138. 10.1007/978-1-0716-2819-5_8.

83.   Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., and Durbin, R. (2009). The Sequence Alignment/Map format and SAMtools. Bioinformatics (Oxford, England) *25*, 2078-2079. 10.1093/bioinformatics/btp352.

84.   Marchal, C., Sasaki, T., Vera, D., Wilson, K., Sima, J., Rivera-Mulia, J.C., Trevilla-García, C., Nogues, C., Nafie, E., and Gilbert, D.M. (2018). Genome-wide analysis of replication timing by next-generation sequencing with E/L Repli-seq. Nat Protoc *13*, 819-839. 10.1038/nprot.2017.148.

85.   Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics (Oxford, England) *29*, 15-21. 10.1093/bioinformatics/bts635.

86. Depienne, C., van den Maagdenberg, A., Kühnel, T., Ishiura, H., Corbett, M.A.,
and Tsuji, S. (2023). Insights into familial adult myoclonus epilepsy pathogenesis:
How the same repeat expansion in six unrelated genes may lead to cortical
excitability. Epilepsia *64 Suppl 1*, S31-s38. 10.1111/epi.17504.
87. Edvardson, S., Nicolae, C.M., Noh, G.J., Burton, J.E., Punzi, G., Shaag, A.,
Bischetsrieder, J., De Grassi, A., Pierri, C.L., Elpeleg, O., and Moldovan, G.L.
(2019). Heterozygous RNF13 Gain-of-Function Variants Are Associated with
Congenital Microcephaly, Epileptic Encephalopathy, Blindness, and Failure to
Thrive. Am J Hum Genet *104*, 179-185. 10.1016/j.ajhg.2018.11.018.
88. Uehara, T., Tsuchihashi, T., Yamada, M., Suzuki, H., Takenouchi, T., and
Kosaki, K. (2019). CNOT2 haploinsufficiency causes a neurodevelopmental
disorder with characteristic facial features. Am J Med Genet A *179*, 2506-2509.
10.1002/ajmg.a.61356.
89. Liao, C., Fu, F., Li, R., Yang, W.Q., Liao, H.Y., Yan, J.R., Li, J., Li, S.Y., Yang, X.,
and Li, D.Z. (2013). Loss-of-function variation in the DPP6 gene is associated
with autosomal dominant microcephaly and mental retardation. Eur J Med Genet
56, 484-489. 10.1016/j.ejmg.2013.06.008.
90. Ismail, V., Zachariassen, L.G., Godwin, A., Sahakian, M., Ellard, S., Stals, K.L.,
Baple, E., Brown, K.T., Foulds, N., Wheway, G., et al. (2022). Identification and
functional evaluation of GRIA1 missense and truncation variants in individuals
with ID: An emerging neurodevelopmental syndrome. Am J Hum Genet *109*,
1217-1241. 10.1016/j.ajhg.2022.05.009.
91. Blanchet, P., Bebin, M., Bruet, S., Cooper, G.M., Thompson, M.L., Duban-Bedu,
B., Gerard, B., Piton, A., Suckno, S., Deshpande, C., et al. (2017). MYT1L
mutations cause intellectual disability and variable obesity by dysregulating gene
expression and development of the neuroendocrine hypothalamus. PLoS Genet
13, e1006957. 10.1371/journal.pgen.1006957.
92. Bertoli-Avella, A.M., Kandaswamy, K.K., Khan, S., Ordonez-Herrera, N.,
Tripolszki, K., Beetz, C., Rocha, M.E., Urzi, A., Hotakainen, R., Leubauer, A., et
al. (2021). Combining exome/genome sequencing with data repository analysis
reveals novel gene-disease associations for a wide range of genetic disorders.
Genet Med *23*, 1551-1568. 10.1038/s41436-021-01159-0.
93. Cohen, J.S., Srivastava, S., Farwell Hagman, K.D., Shinde, D.N., Huether, R.,
Darcy, D., Wallerstein, R., Houge, G., Berland, S., Monaghan, K.G., et al. (2017).
Further evidence that de novo missense and truncating variants in ZBTB18
cause intellectual disability with variable features. Clin Genet *91*, 697-707.
10.1111/cge.12861.
94. Sakaguchi, Y., Uehara, T., Suzuki, H., Kosaki, K., and Takenouchi, T. (2017).
Truncating mutation in CSNK2B and myoclonic epilepsy. Hum Mutat *38*, 1611-
1612. 10.1002/humu.23307.
95. Paulhus, K., Ammerman, L., and Glasscock, E. (2020). Clinical Spectrum of
KCNA1 Mutations: New Insights into Episodic Ataxia and Epilepsy Comorbidity.
Int J Mol Sci *21*. 10.3390/ijms21082802.
96. Müller, P., Takacs, D.S., Hedrich, U.B.S., Coorg, R., Masters, L., Glinton, K.E.,
Dai, H., Cokley, J.A., Riviello, J.J., Lerche, H., and Cooper, E.C. (2023). KCNA1
gain-of-function epileptic encephalopathy treated with 4-aminopyridine. Ann Clin
Transl Neurol *10*, 656-663. 10.1002/acn3.51742.

97. Boulin, T., Itani, O., El Mouridi, S., Leclercq-Blondel, A., Gendrel, M., Macnamara, E., Soldatos, A., Murphy, J.L., Gorman, M.P., Lindsey, A., et al. (2021). Functional analysis of a de novo variant in the neurodevelopment and generalized epilepsy disease gene NBEA. Mol Genet Metab *134*, 195-202. 10.1016/j.ymgme.2021.07.013.

98. Zhao, S., Yudin, Y., and Rohacs, T. (2020). Disease-associated mutations in the human TRPM3 render the channel overactive via two distinct mechanisms. eLife *9*. 10.7554/eLife.55634.

99. Palmer, E.E., Kumar, R., Gordon, C.T., Shaw, M., Hubert, L., Carroll, R., Rio, M., Murray, L., Leffler, M., Dudding-Byth, T., et al. (2017). A Recurrent De Novo Nonsense Variant in ZSWIM6 Results in Severe Intellectual Disability without Frontonasal or Limb Malformations. Am J Hum Genet *101*, 995-1005. 10.1016/j.ajhg.2017.10.009.

100. Jaillard, S., Andrieux, J., Plessis, G., Krepischi, A.C., Lucas, J., David, V., Le Brun, M., Bertola, D.R., David, A., Belaud-Rotureau, M.A., et al. (2011). 5q12.1 deletion: delineation of a phenotype including mental retardation and ocular defects. Am J Med Genet A *155a*, 725-731. 10.1002/ajmg.a.33758.

101. Ohta, E., Kawakami, F., Kubo, M., and Obata, F. (2013). Dominant-negative effects of LRRK2 heterodimers: a possible mechanism of neurodegeneration in Parkinson's disease caused by LRRK2 I2020T mutation. Biochem Biophys Res Commun *430*, 560-566. 10.1016/j.bbrc.2012.11.113.

102. Taymans, J.M., Fell, M., Greenamyre, T., Hirst, W.D., Mamais, A., Padmanabhan, S., Peter, I., Rideout, H., and Thaler, A. (2023). Perspective on the current state of the LRRK2 field. NPJ Parkinsons Dis *9*, 104. 10.1038/s41531-023-00544-7.

103. Kobayashi, H., Krüger, R., Markopoulou, K., Wszolek, Z., Chase, B., Taka, H., Mineki, R., Murayama, K., Riess, O., Mizuno, Y., and Hattori, N. (2003). Haploinsufficiency at the alpha-synuclein gene underlies phenotypic severity in familial Parkinson's disease. Brain *126*, 32-42. 10.1093/brain/awg010.

104. Konno, T., Ross, O.A., Puschmann, A., Dickson, D.W., and Wszolek, Z.K. (2016). Autosomal dominant Parkinson's disease caused by SNCA duplications. Parkinsonism Relat Disord *22 Suppl 1*, S1-6. 10.1016/j.parkreldis.2015.09.007.

105. Blauwendraat, C., Makarious, M.B., Leonard, H.L., Bandres-Ciga, S., Iwaki, H., Nalls, M.A., Noyce, A.J., and Singleton, A.B. (2021). A population scale analysis of rare SNCA variation in the UK Biobank. Neurobiol Dis *148*, 105182. 10.1016/j.nbd.2020.105182.

Methods

Cell culture

All human tissue samples were obtained in accordance with protocols approved by the Oregon Health & Science University Institutional Review Board (OHSU IRB# 00019018) and the Western Institutional Review Board (WIRB# 20193303). Informed consent was obtained from all donors or their legal guardians. ACP cells were grown in Dulbecco's Modified Eagle's Medium low glucose (Life Technologies) supplemented with 10% fetal bovine serum (Hyclone). Single cell clones were isolated by plating individual cells in 6 well dishes, and were expanded for >20 population doublings [79]. All cells were grown in a humidified incubator at 37°C in a 5% carbon dioxide atmosphere.

Haplotype Phasing of ACP cells

Short-read whole genome sequencing was obtained from two parent/child trios with a range of ~7-14x genome coverage. BWA [80] was used for alignment, and FreeBayes [81] used to call putative germline variants jointly for each trio. Variants were filtered to remove sites with Genotype Quality score of less than 30, and sites with missing genotypes for any member of the trio were removed. WhatsHap [82] was then used to phase variants in the child utilizing mendelian inheritance patterns and reads containing multiple variants. 1,679,207 high-confidence sites were phased in ACP7, and 837,323 sites were phased in ACP6.

Allele-specific Repli-Seq

Short read sequence data was aligned with BWA with standard parameters. BAM files were then deduplicated and sorted with SAMtools [83]. BCFtools mpileup [83] was then

used to produce allele-specific read counts at phased heterozygous loci. Library size

normalization was performed to produce counts per million informative reads. Read

counts were then grouped into 250kb sliding genomic windows with no overlap. E/L

Repli-Seq values were generated by taking the Log base 2 of early versus late values

[84]. Quantile normalization was performed on each group of subclones within each

individual sample to allow for comparison among subclones. For each sample,

variability of allele-specific RT among subclones was defined as the Standard Deviation

(degrees of freedom = 1) of repli-seq at each 250kb window. Genome-wide std. dev.

values generated a log-normal distribution, and we set a threshold of $Z>=2.25$ on the

log-normal to identify regions with highly variable allele-specific RT among subclones.

1017

Allele-specific RNA-seq

Nuclei were isolated by centrifugation for 0.5 minutes from ACP clones following lysis in

0.5% NP40, 140 mM NaCl, 10 mM Tris-HCl (pH 7.4), and 1.5 mM $MgCl_2$. Nuclear RNA

was isolated using Trizol reagent using the manufacturer's instructions, followed by

DNase treatment to remove possible genomic DNA contamination. Briefly, ribosomal

RNAs were removed using the Ribo-Zero kit (Illumina), RNA was fragmented into 250-

300bp fragments, and cDNA libraries were prepared using the Directional RNA Library

Prep Kit (NEB). Paired end sequencing was done on a NovaSeq 6000 at the OHSU

MPSSR core facility.

Strand specific short read sequence data was aligned with STAR [85]. SAMtools [83]

was used to remove duplicates and sort BAM files. Strand-specific, allele-specific reads

at heterozygous phased SNPs were counted using BCFtools mpileup [83]. For gene-

centric allele-specific analyses, the GENCODE v44 Gene annotation GTF file was used

to define gene start and stop coordinates. The informative strand-specific, allele-specific

read counts for each heterozygous SNP were then pooled together for all SNPs within

each gene. Allelic expression imbalance was calculated as the fraction of reads derived

from the maternal vs the paternal allele.

$$AEI = \frac{|\text{Paternal allele reads} - \text{Maternal allele reads}|}{\text{Paternal allele reads} + \text{Maternal allele reads}} \times 100$$

1035

For example, completely balanced allelic expression is 50% AEI, and completely

imbalanced AEI is 100%. A threshold of AEI>= 70%, and FDR corrected binomial p-

value<=0.05 was used to classify AEI in each individual subclone. The standard

deviation (degrees of freedom=1) was used for each gene to generate a metric of AEI

variability among subclones. The genome-wide distribution of std. dev. was generated

across all genes, and the top 2.5% of variable genes were classified as highly variable

among subclones.

Figure Legends:

Figure 1: Genome wide variable replication-timing and expression imbalance. a) Single cell derived ACP clones (color coded) were isolated, expanded and processed for Repli-seq and RNA-seq. The maternal and paternal haplotypes were used to analyze allele-specific replication timing and expression. b) The top panel illustrates the expected replication timing profile in multiple clones across an autosomal region. The regions that display VERT are highlighted by shading. The bottom panel illustrates the expected expression variability in different clones from the same individual. c) We define VERT regions in Repli-seq data across each allele for each 250 kb genomic window. Windows with an SD value ≥ 2.5 standard deviations above the genome-wide median were classified as outliers (top panels). We define AEI as an allelic bias that is ≥80% allelic imbalance (AEI ≥ 0.80 or ≤ 0.20). The bottom panel shows both autosomal and X linked expression data. d) Illustration of an Inactivation/Stability Center showing the clonal and allele-restricted variability in replication timing and expression of both protein coding and noncoding genes. e) The standard deviation (Std. Dev.) in 250 kb windows (circles) across human chromosome 1. Outlier windows from all four sets of clones (ACP6, ACP7, EB3-2, and GM12878) are highlighted in different colors as shown. The top panel illustrates the G-banding (blue shading) pattern for human chromosome 1, with 23 VERT regions highlighted in red. The location of an imprinted region with asynchronous replication, containing the maternally expressed TP73 gene (geneimprint.com), is marked with an +. The arrow marks the VERT region shown in panels f-h below. f-h) I/SC located between 239 and 242 mb of chromosome 1 (see panel e). f) VERT (shaded region) and AEI of genes (rectangles) detected in the ACP6 clones, each clone was color coded as shown. For replication, the solid lines represent the paternal (Hap1) allele and the dotted line represents the maternal (Hap2) allele. g) VERT (shaded region) and AEI of genes (rectangles) detected in the ACP7 clones, each clone was color coded as shown. h) UCSC Genome Browser view of the genomic region in panels f and g. The shaded area highlights the VERT region. The asterisks mark the genes highlighted in panels f and g. i) Venn diagram illustrating the number of VERT regions that overlap between the different clonal populations. j) Gene Ontology (GO) enrichment analysis of the genes located within the VERT regions, the most significant hits for each set of clones, organized by cell type, are shown. GO analysis of the mouse genes located within the pre-B cell clone VERT regions are also shown.

Figure 2. Examples of I/SCs. a, d and g) VERT regions on human chromosomes 18, 6 and 9, respectively. The top panels illustrate the G-banding (blue shading) pattern for each human chromosome, with VERT regions highlighted in red. The standard deviation (Std. Dev.) in 250 kb windows (circles) across each human chromosome is shown. Outlier windows from all four sets of clones (ACP6, ACP7, EB3-2, and GM12878) are highlighted in different colors as shown. The arrows and red boxes mark the VERT regions shown in panels b, e, and h. b, e and h) I/SCs highlighted in panels a, d, and g. For replication, the solid lines represent the paternal (Hap1) allele and the dotted lines represent the maternal (Hap2) allele. The regions with VERT are shaded and the AEI is shown for each clone in different colors. The location of an imprinted

region with asynchronous replication, containing the paternally expressed LIN28B gene (geneimprint.com), is marked with an + in panel d. c, f and i) UCSC Genome Browser views of the genomic regions in panels b, e and h. The shaded areas highlight the VERT regions. Genes with AEI are marked with +, *, or [#]. b and c) The + marks MOCOS, which shows AEI in ACP cells (Supplemental Table 2) and ++ marks FHOD3, which shows AEI in LCLs [1]. e and f) The * marks from DNAH8, ** marks transcripts from KCNK5, and *** marks non-coding transcripts from ENSG00000293066 (also named: TL:6-39.9_1 [1]). All three of these genes show AEI in LCL clones [1]. h and i) The [#] marks transcripts from DAPK1, [##] marks transcripts from SPIN1. Both genes show AEI in LCL clones [1].

Figure 3. Human gene clusters with AEI map to VERT regions. a and c) VERT regions on human chromosome 19 and 6, respectively. The top panels illustrate the G-banding (blue shading) pattern for each human chromosome, with VERT regions highlighted in red. The standard deviation (Std. Dev.) in 250 kb windows (circles) across each human chromosome is shown. Outlier windows from all four sets of clones (ACP6, ACP7, EB3-2, and GM12878) are highlighted in different colors as shown. The red boxes and arrows identify the I/SCs in panel b and d. The location of an imprinted region with asynchronous replication, containing the paternally expressed LIN28B gene (geneimprint.com), is marked with an + in panel c. b) UCSC genome browser view illustrating the genomic location highlighted in panel a. The shaded area highlights the VERT region. The red underlines mark 11 olfactory receptor genes, and the green underlines mark 6 cytochrome P450 family (CYP4F) genes. d) UCSC Genome Browser view illustrating the genomic location highlighted in panel c. The shaded area highlights the VERT region. The red underlines mark 16 HLA genes, and the green underlines mark 6 Lymphocyte antigen 6 family (LY6G) genes.

Figure 4. Mouse gene clusters with AEI map to VERT regions. a and d) VERT regions on mouse chromosome 2 and 18, respectively. The standard deviation (Std. Dev.) in 50 kb windows (circles) across each mouse chromosome are shown. Outlier windows from pre-B cell clones are highlighted in orange. The red boxes and arrows identify the VERT regions in panels b (i) and c (ii). b and c) UCSC genome browser views illustrating the genomic locations represented by i and ii in panel a. b) Genome Browser view of the VERT region showing the location of 29 olfactory receptor genes (Or genes). The shaded area represents the VERT region. c) Genome Browser view of the VERT region in ii above, showing the location of 242 olfactory receptor genes (Or genes). The shaded area represents the VERT region. e) Genome Browser view of the VERT region in panel d, showing the location of clustered Protocaderhin genes (15 Pcdha, 23 Pcdhb, and 22 Pcdhg genes). The shaded area represents the VERT region.

Figure 5. Human and Mouse syntenic regions display VERT. a and b) The short arm of human chromosome 5, with 11 VERT regions (panel a) and the centromeric region of mouse chromosome 15, with 4 VERT regions (panel b) are shown as orange circles. a) Two human VERT regions are highlighted, hi and hii, and are syntenic with locations on mouse chromosome 15, mi and mii, that also show VERT (panel b). Note that the gene order in both syntenic regions are inverted with respect to the telomeres when

comparing the human and mouse loci. c) Illustration of the I/SC on human chromosome 5 between 7.75 and 10.5 mb (hi). The shaded area marks the VERT region, and AEI of the coding gene SEMA5A in different clones is highlighted. For replication, the paternal (Hap1) and maternal (Hap2) alleles are indicated. d) UCSC Genome Browser view of the human VERT region (shaded) in hi, showing the location of SEMA5A and a taste receptor gene (TAS2R1). e) UCSC Genome Browser view of the mouse VERT region (shaded) in mi, showing the location of Sema5a and a taste receptor gene (Tas2r119). f) UCSC Genome Browser view of the human VERT region (shaded) in hii above, highlighting the location of the AEI gene RETREG1 [45]. g) UCSC Genome Browser view of the mouse VERT region (shaded) in mii, showing the synteny between human and mouse for the protein coding genes in panel f, including Retreg1. h) VERT regions on human chromosome 16. The standard deviation (Std. Dev.) in 250 kb windows (circles) across human chromosome 16 is shown. Outlier windows are highlighted in different colors as shown. The red box and arrow mark the genomic location in panel i. i) UCSC Genome Browser view illustrating the genomic location represented in panel h, showing the location of GRIN2A (red box). j) VERT regions on mouse chromosome 16. The standard deviation in 50 kb windows (circles) is shown. Outlier windows from pre-B cell clones are highlighted in orange. k) UCSC Genome Browser view illustrating the genomic location of the VERT region (shaded area) represented in panel j above, with the location of Grin2a highlighted with a red box.

Figure 6. Epilepsy genes map to I/SCs. a, d, and g) VERT regions on human chromosomes 2, 5, and 8, respectively. The top panels illustrate the G-banding (blue shading) pattern for each human chromosome, with VERT regions highlighted in red. The standard deviation (Std. Dev.) in 250 kb windows (circles) across each human chromosome is shown. Outlier windows from all four sets of clones (ACP6, ACP7, EB3-2, and GM12878) are highlighted in different colors as shown. The location of an imprinted region with asynchronous replication, containing the paternally expressed DLGAP2 gene (geneimprint.com), is marked with an + in panel g. The arrows and red boxes mark the VERT regions shown in panels b, c, e, f, h, i, j and k. b, e, h, j) Illustrations of I/SCs on chromosomes 2, 5, and 8, respectively. The shaded areas mark the VERT regions. Each clone was color coded to illustrate the VERT and AEI in different clones. The paternal (Hap1) and maternal (Hap2) alleles are indicated for replication and expression. c) UCSC Genome Browser view illustrating the genomic location of the VERT region (shaded area) represented in panels a and b, with the location of 5 sodium channel genes (SCN3A, SCN2A, SCN1A, SCN9A, and SCN7A) highlighted with red underlines. e) Illustration of an I/SC on chromosome 5. The shaded area marks the VERT region. AEI of GABRA1 is shown. f) UCSC Genome Browser view illustrating the genomic location of the VERT region (shaded area) represented in panels d and e above, with the location of 4 GABA receptor genes (GABRB2, GABRBA6, GABRA1, and GABRG2) highlighted with red underlines. h) Illustration of an I/SC (A and red box) on chromosome 8. Each ACP6 clone was color coded to illustrate the VERT and AEI in different clones. The paternal (Hap1) and maternal (Hap2) alleles are indicated for replication and expression. The shaded area marks the VERT region. The location and expression of OXR1 is indicated. i) UCSC Genome Browser view illustrating the genomic location of the VERT region (shaded area) represented in panels g (A) and h above, with the location of the OXR1 gene highlighted with a red underline. We note that ANGPT1 (marked with *), located within this I/SC

displays AEI (see Supplementary Table 2). j) Illustration of an I/SC (B and red box) on chromosome 8. AEI of SAMD12 is indicated. k) UCSC Genome Browser view illustrating the genomic location of the VERT region (shaded area) represented in panels g (B) and j above, with the location of the SAMD12 highlighted with a red underline.

Figure 7. Parkinson disease genes map to I/SCs. a, c, e, and g) VERT regions on human chromosomes 4, 12, 1, and 6, respectively. The top panels illustrate the G-banding (blue shading) pattern for each human chromosome, with VERT regions highlighted in red. The standard deviation (Std. Dev.) in 250 kb windows (circles) across each human chromosome is shown. Outlier windows from all four sets of clones (ACP6, ACP7, EB3-2, and GM12878) are highlighted in different colors as shown. The location of an imprinted region with asynchronous replication, containing the paternally expressed LIN28B gene (geneimprint.com), is marked with an + in panel g. The location of an imprinted region with asynchronous replication, containing the maternally expressed TP73 gene (geneimprint.com), is marked with an + in panel e. The arrows and red boxes mark the VERT regions shown in panels b, d, f, and h, respectively. b) UCSC Genome Browser view illustrating the genomic location of the VERT region (shaded area) represented in panel a, with the location of the SNCA gene highlighted with a red underline. d) UCSC Genome Browser view illustrating the genomic location of the VERT region (shaded area) highlighted in panel c, with the location of the LRRK2 gene highlighted with a red underline. f) UCSC Genome Browser view illustrating the genomic location of the VERT region (shaded area) represented in panel e, with the location of the DNAJC6 gene highlighted with a red underline; also highlighted with a green underline is the JAK1 gene. h) UCSC Genome Browser view illustrating the genomic location of the VERT region (shaded area) represented in panel g, with the location of the PRKN gene highlighted with a red underline.

Supplementary Figure 1. Imprinted regions display asynchronous replication. a and d) VERT regions on human chromosome 14 and 20. The standard deviation in 250 kb windows (circles) are shown. Outlier windows from all four sets of clones (ACP6, ACP7, EB3-2, and GM12878) are highlighted in different colors as shown. The regions highlighted by the red boxes mark known imprinted loci. b and e) For replication timing, solid lines represent the paternal allele (Hap1) and the dotted line represents the maternal allele (Hap2). The regions with asynchrony are shaded and the gene expression is shown for each clone in a different color. c) UCSC Genome Browser view of the asynchronous region in panel a. c and b) The single asterisk marks the paternally expressed DLK gene, and the double asterisks mark the maternally expressed MEG8 gene. f) UCSC Genome Browser view of the VERT region in panel d. f and e) The # marks the paternally expressed L3MBTL1 gene.

Supplementary Figure 2. VERT detected at the mouse T cell receptor alpha locus. UCSC Genome Browser view of the mouse VERT region (shaded) detected at the T cell receptor alpha (Tra) locus on mouse chromosome 14. The location of 37 Or genes, marked by red underlines, and the vomeronasal receptor gene Vmn2r88, marked by a green underline.

Supplementary Figure 3. Mouse syntenic regions display VERT. a, c and e) VERT
regions on mouse chromosome 16, 2, and 18. The standard deviation in 50 kb windows
(circles) is shown. Outlier windows from pre-B cell clones are highlighted in orange. The
VERT regions highlighted by red boxes are expanded in panels b, d, and f. b) UCSC
Genome Browser view of the VERT region in a above, highlighting the location of the
mouse App gene (red box). The shaded area represents the VERT region. d) UCSC
Genome Browser view of the VERT region in panel c, highlighting the location of the
mouse Mocos and Fhod3 genes (red boxes). The shaded area represents the VERT
region. f) UCSC Genome Browser view of the VERT region in panel c, highlighting the
location of five mouse Scn genes (red boxes). The shaded area represents the VERT
region.
1242
1243

RESOURCE AVAILABILITY

Source data are provided with this paper. Further information and requests for resources
and reagents should be directed to and will be fulfilled by the lead contact, Mathew
Thayer (thayerm@ohsu.edu).

Materials availability statement

All reagents generated in this study will be made freely available upon request.

1250
1251 # Data Availability Statement
1252
1253 The Repli-seq and RNA-seq raw sequencing data generated in this study have been
1254 deposited in the European Nucleotide Archive database under accession code
1255 PRJEB110291 and are available without restriction.
1256

Human VERT Region			Gene Cluster			
Chr	Start	Stop	Genes	# of Genes	Start	stop
1	13,250,000	15,000,000	PRAMEF	23	12,768,939	13,422,488
2	101,633,538	102,383,540	IL1R	5	101,987,899	102,401,443
2	165,250,000	166,000,000	SCN	5	165,081,204	166,499,088
2	176,250,000	176,500,000	HOXD	9	176,091,062	176,191,493
4	46,750,000	47,500,000	GABR	4	46,035,583	47,431,569
4	68,384,282	68,634,282	UGT	9	68,525,283	69,660,413
5	651,700	750,000	SLC	5	468,888	1,448,927
5	148,000,000	148,250,000	SPINK	7	147,822,992	148,343,066
5	162,250,000	162,750,000	GABR	4	161,282,809	162,159,566
6	31,250,000	32,750,000	HLA	15	30,489,509	33,093,613
			LY6	6	31,669,259	31,722,476
8	11,250,000	12,392,491	DEFB	6	11,970,747	12,320,078
8	80,587,765	83,000,000	FABP	4	81,277,942	81,536,065
8	142,500,000	142,750,000	LY6	8	142,695,707	143,162,219
11	104,750,000	105,000,000	CASP	5	104,883,181	105,141,441
15	23,357,400	23,500,000	GOLGA	5	22,451,995	23,451,545
15	28,750,000	29,000,000	GOLGA	6	28,344,604	28,855,059
17	9,750,000	10,500,000	MYH	6	10,297,552	10,658,865
19	15,000,000	15,250,000	OR	11	14,792,476	15,952,060
			CYP	6	15,507,105	15,936,041
19	20,750,000	21,000,000	ZNF	37	19,667,773	24,119,420
19	36,313,700	36,500,000	ZNF	29	36,171,632	37,791,237
19	43,000,000	43,250,000	PSG	10	42,713,521	43,272,861
20	24,250,000	24,750,000	CST	11	23,429,902	24,957,423
21	30,627,681	30,877,681	KRTAP	33	30,276,201	31,041,124

1257

Mouse VERT Region			Gene Cluster			
Chr	Start	Stop	Genes	# of Genes	Start	stop
2	36,322,857	36,622,857	Or	28	36,225,624	37,193,679
2	64,622,866	66,522,868	Scn	5	65,287,042	66,637,709
2	88,950,000	89,950,000	Or	239	85,160,859	90,147,389
4	59,850,000	59,950,000	Mup	22	59,950,000	61,824,646
5	91,150,000	91,350,000	Cxcl	5	91,186,938	91,335,955
5	95,950,000	96,250,000	Pramel	23	93,891,822	96,243,866
6	40,650,000	41,150,000	Tas2r	8	40,386,430	43,196,843
			Or	24	40,386,430	43,196,843
6	66,650,000	66,850,000	Vmn1	8	66,498,911	66,758,688
7	113,809,945	113,909,945	Or	24	113,790,926	114,306,041
11	73,450,000	74,050,000	Or	31	73,160,403	74,180,092
18	37,350,000	37,750,000	Pcdh	23	37,298,052	37,684,088

1258   **Table 1. VERT regions at gene clusters in the human and mouse genomes**

Mouse VERT			Synteny (%)	Human Syntenic Region			Human VERT		
Chr	Start	stop		Chr	start	Stop	Chr	start	Stop
1	73,042,585	73,542,585	54.0	2	216,836,864	217,400,721	2	217,000,000	217,250,000
1	88,681,147	89,081,147	33.0	2	234,565,154	235,053,857	2	234,000,000	235,000,000
1	189558318	189,758,318	49.8	1	214,144,079	214,410,956	1	214,326,657	216,000,000
2	64,842,287	66,742,287	46.5	2	164,610,983	166,685,353	2	165,250,000	166,000,000
3	18,204,164	18,404,164	33.7	8	64,684,742	64,876,026	8	64,337,443	65,087,765
3	57,253,499	58,553,499	32.6	3	149,407,811	150,704,641	3	149,282,213	150,032,213
3	78,253,385	78,653,385	26.1	4	159,655,594	159,952,325	4	159,328,848	159,828,848
3	82,453,385	82,853,385	32.6	4	154,688,934	155,210,940	4	154,500,000	154,750,000
4	105034592	105,234,592	62.8	1	56,331,724	56,556,339	1	55,500,000	56,750,000
4	142286667	142,686,667	40.6	1	14,022,915	14,523,879	1	13,250,000	15,000,000
8	10,500,000	11,200,000	43.2	13	108,978,718	110,065,357	13	109,250,000	109,500,000
8	24,029,544	24,929,544	37.3	8	40,152,844	41,090,451	8	40,250,000	40,750,000
8	39,817,687	40,717,687	28.6	8	15,936,545	16,964,053	8	15,750,000	17,000,000
9	116969950	117,469,950	42.3	3	28,837,647	29,402,683	3	29,000,000	29,500,000
11	46,927,325	48,127,325	35.2	5	155,465,678	156,698,430	5	155,500,000	156,250,000
12	33,715,134	33,815,134	36.1	7	19,330,767	19,416,599	7	19,250,000	19,500,000
12	105278049	105,578,049	41.6	14	95,853,630	96,269,387	14	96,000,000	96,250,000
13	11,095,964	11,672,619	29.9	1	237,711,340	238,470,922	1	237,604,200	238,750,000
15	25,120,331	25,920,331	42.9	5	16,534,412	17,724,462	5	16,250,000	17,000,000
15	32,220,391	32,420,391	60.7	5	9,377,059	9,575,503	5	8,749,888	9,499,888
16	9,067,771	9,667,771	41.6	16	9,476,924	10,028,307	16	9,750,000	10,000,000

1259 **Table 2. VERT regions showing synteny between human and mouse genomes.**

VERT Gene	Epilepsy Disease Genes (Inheritance: Autosomal Recessive-AR; Autosomal Dominant-AD; Loss of Function-LoF; Gain of Function-GoF)	Ref.
ALG14	Myasthenic syndrome (AR); Intellectual developmental disorder with epilepsy (AR); Myopathy, epilepsy, and progressive cerebral atrophy (AR)	29
CACNA2D1	Developmental and epileptic encephalopathy (AR)	29
GABRB1	Developmental and epileptic encephalopathy (AD, LoF & GoF)	54
GRIN2A	Epilepsy (AD, LoF)	50
NTRK2	Developmental and epileptic encephalopathy (AD); Obesity (AD)	55
OXR1	Cerebellar hypoplasia/atrophy, epilepsy, and global developmental delay (AR)	29
RAPGEF2	Epilepsy, familial adult myoclonic (AD, TTTCA repeat expansion)	86
RNF13	Developmental and epileptic encephalopathy (AD, GoF)	87
SAMD12	Epilepsy (AD, TTTCA repeat expansion)	86
SCN1A	Developmental and epileptic encephalopathy (AD, LoF); Dravet syndrome (AD, LoF); Generalized epilepsy with febrile seizures (AD, LoF)	52
SCN2A	Developmental and epileptic encephalopathy (AD, LoF & GoF)	53

1260

VERT Gene	Intellectual Developmental Genes (Inheritance: Autosomal Recessive-AR; Autosomal Dominant-AD; Loss of Function-LoF; Gain of Function-GoF)	Ref.
ALG14	Intellectual developmental disorder with epilepsy (AR); Myasthenic syndrome (AR); Myopathy, epilepsy, and progressive cerebral atrophy (AR)	29
CEP104	Intellectual developmental disorder (AR); Joubert syndrome (AR)	29
CNOT2	Intellectual developmental disorder (AD, LoF)	88
DPP6	Intellectual developmental disorder (AD, LoF)	89
ELP2	Intellectual developmental disorder (AR)	29
FMN2	Intellectual developmental disorder (AR)	29
GRIA1	Intellectual developmental disorder (AD, GoF; AR, LoF)	90
IMPA1	Intellectual developmental disorder (AR)	29
IQSEC1	Intellectual developmental disorder (AR)	29
MYT1L	Intellectual developmental disorder (AD, LoF)	91
PDZD8	Intellectual developmental disorder (AR)	29
RSRC1	Intellectual developmental disorder (AR)	29
TAF4	Intellectual developmental disorder (AD, LoF)	92
TNIK	Intellectual developmental disorder (AR)	29
TUSC3	Intellectual developmental disorder (AR)	29
ZBTB18	Intellectual developmental disorder (AD, LoF)	93

1261

VERT Gene	Neurodevelopment Disease Genes (Inheritance: Autosomal Recessive-AR; Autosomal Dominant-AD; Loss of Function-LoF; Gain of Function-GoF)	Ref.
ATP8A2	Cerebellar ataxia (AR)	29
CSNK2B	Poirier-Bienvenu neurodevelopmental syndrome (AD, LoF)	94
DPH5	Neurodevelopmental disorder (AR)	29
KCNA1	Episodic ataxia/myokymia syndrome (AD, LoF & GoF)	95,96
LMBRD2	Developmental delay with variable neurologic and brain abnormalities (AD)	29
NBEA	Neurodevelopmental disorder (AD, LoF)	97
RETREG1	Neuropathy (AR)	29
ROBO1	Nystagmus (AR); Neurooculorenal syndrome (AR); Pituitary hormone deficiency (AD)	29
TIAM1	Neurodevelopmental disorder (AR)	29
TRPM3	Neurodevelopmental disorder (AD, GoF)	98
VARS1	Neurodevelopmental disorder with microcephaly (AR)	29
ZSWIM6	Acromelic frontonasal dysostosis, (AD); Neurodevelopmental disorder (AD, GoF)	99,100

1262

VERT Gene	Parkinson Disease Gene (Inheritance: Autosomal Recessive-AR; Autosomal Dominant-AD; Loss of Function-LoF; Gain of Function-GoF)	Ref.
DNAJC6	Parkinson disease 19 (AR)	29
LRRK2	Parkinson disease (AD, GoF)	101,102
PRKN	Parkinson disease (AR)	29
SLC18A2	Parkinsonism-dystonia (AR)	29
SNCA	Parkinson disease 1 & 4 (AD, LoF/GoF)	103-105

Table 3. Neurodevelopmental genes within VERT regions.

a Human Chr18: 14 VERT regions

ACP6 ACP7 EB3-2 GM12878

b

c Chr18: 34.5-37.5 mb

+ MOCOS; ++ FHOD3

d Human Chr6: 20 VERT Regions

ACP6 ACP7 EB3-2 GM12878

e

f Chr6: 38.3-41.5 mb

*DNAH8; **KCNK5; ***ENSG00000293066

g Human Chr9: 18 VERT Regions

ACP6 ACP7 EB3-2 GM12878

h

i Chr9: 86.3-89.0 mb

#DAPK1; ##SPIN1

a Human Chr19: 5 VERT Regions

ACP6 ACP7 EB3-2 GM12878

b Chr19: 14.8-16.0 mb. Olfactory Receptors (11 OR genes)

Cytochrome P450 cluster (6 CYP4F genes)

c Human Chr6: 20 VERT Regions

ACP6 ACP7 EB3-2 GM12878

d Chr6:30.9-33.5 mb. Major Histocompatibility (16 HLA genes)

Lymphocyte antigen 6 family cluster (6 LY6G genes)

a Mouse Chr2: 9 VERT Regions

Replication / Std. Dev.

Chromosome position

b i. Chr2: 36.2-37.25 mb. Olfactory Receptors (29 Or genes)

Scale 500 kb mm39
chr2: 36,300,000 36,500,000 36,700,000 36,900,000 37,100,000
GENCODE VM37

Or1j8 Or1j11 Or1j14 Or1j16 Or1j18 Or1j1 Or1n2 Or12k5 Or1b1 Or1af1 Or5c1
Or1j10 Or1j13 Or1j15 Or1j17 Or1j4 Or1ak2 Or12k8 Or1l4
Or1j12 Or1j21 Or1n1b Or1q1 Or1l4b
Or1j19 Or1l8 Or12k7
Or1j20
Or1n1

c ii. Chr2: 85-90 mb: Olfactory Receptors (243 Or genes)

Scale 2 Mb mm39
chr2: 85,500,000 86,500,000 87,500,000 88,500,000 89,500,000
GENCODE VM37

d Mouse Chr18: 8 VERT Regions

Replication / Std. Dev.

Chromosome position

e Chr18: 37-38 mb: Protocadherins (15 Pcdha, 23 Pcdhb, and 22 Pcdhg genes)

Scale 500 kb mm39
chr18: 37,100,000 37,300,000 37,500,000 37,700,000 37,900,000
GENCODE VM37

Pcdha | Pcdhb | Pcdhg

a Human Chr4: 18 VERT Regions

b Chr4: 88.7-90 mb

SNCA (PARK1 and PARK4: Autosomal Dominant)

c Human Chr12: 18 VERT Regions

d Chr12: 39.5-41.6 mb

LRRK2 (PARK8: Autosomal Dominant)

e Human Chr1: 19 VERT Regions

f Chr1: 63-66.5 mb

JAK1 (AIIDE: Autosomal Dominant)   DNAJC6 (PARK19: Autosomal Recessive)

g Human Chr6: 20 VERT Regions

h Chr6: 160.8-162.9 mb

PRKN (PARK2: Autosomal Recessive)