NIPBL dosage shapes genome folding by tuning the rate of cohesin loop extrusion

Rini Shah[1,*], Maxime M. C. Tortora[2,*], Nessim Louafi[3], Hadi Rahmaninejad[2], Karissa L. Hansen[1,4], Erika C. Anderson[1], David Wen[1,4], Luca Giorgetti[3], Geoffrey Fudenberg[2,$], Elphège P. Nora[1,5,6$]

1. Cardiovascular Research Institute, University of California, San Francisco, San Francisco, CA, USA
2. Department of Quantitative and Computational Biology, University of Southern California, Los Angeles, CA, USA
3. Friedrich Miescher Institute for Biomedical Research, Basel, Switzerland
4. Developmental and Stem Cell Biology Graduate Program, University of California, San Francisco, CA, USA
5. Department of Biochemistry and Biophysics, University of California, San Francisco, San Francisco, CA, USA
6. Chan-Zuckerberg Biohub San Francisco, CA, USA

* These authors contributed equally
$ Corresponding authors: elphege.nora@ucsf.edu and fudenber@usc.edu

Summary

Cohesin loop extrusion is a major driver of chromosome folding, but how its dynamics are controlled to shape the genome remains elusive. Here we disentangle the contributions of the cohesin cofactors NIPBL, WAPL and PDS5 to loop extrusion kinetics using titrated perturbations and calibrated biophysical models. We demonstrate that beyond contributing to cohesin loading, cellular NIPBL dosage quantitatively controls the rate of loop extrusion. This role explains how lowering NIPBL abundance compensates for elevated cohesin lifetime caused by inactivation of WAPL, the cohesin unloader, and how the NIPBL:PDS5 balance in cells sets the net rate of extrusion. We validate the predicted consequences of altered extrusion rate on chromosome morphology and cohesin dynamics in single cells and demonstrate its impact on compartmentalization and transcription. These findings provide a mechanistic basis for the genetic interactions between cohesin cofactors and the molecular origin of haploinsufficiency in cohesinopathies, such as Cornelia de Lange syndrome.

Highlights

- The abundance, extrusion rate and lifetime of cohesin differentially regulate genome folding.
- The cofactors NIPBL, PDS5, and WAPL each modulate multiple aspects of extrusion kinetics.
- The cellular dosage of NIPBL and its titration by PDS5 sets the net rate of extrusion.
- Loop extrusion rates tune compartmentalization and transcription patterns.

Keywords

Chromosome folding, cohesin loop extrusion, genomic compartmentalization, NIPBL/SCC2, Hi-C, polymer simulations

Introduction

The physical organization of chromosomes influences countless genomic processes, including transcription, DNA repair, replication, recombination and segregation [1]. Throughout the domains of life, Structural Maintenance of Chromosome (SMC)-family protein complexes act as key architects of chromosome folding, through their ability to extrude DNA loops[2,3]. Cohesin has emerged as the main loop extrusion machinery during interphase in vertebrates [4,5], with multiple cofactors enabling cycles of loading, extrusion, and release from chromatin[6].

Mutations in cohesin cofactors, such as WAPL, PDS5 and NIPBL (Figure S1A), can cause chromosome morphology defects and disrupt crucial downstream genomic processes. WAPL mis-expression disrupts heterochromatin[7], compacts chromosomes[8], derails VDJ recombination[9,10] and skews protocadherin promoter choice[11,12]. Alterations in PDS5 across organisms misfold chromosomes[13], compromise DNA condensation, replication and repair[14–16], disrupt meiotic chromosome axis formation and impair interhomolog recombination[17,18]. NIPBL depletion disrupts genome organization[19–22], long-range transcriptional regulation[23–25], and heterozygous mutations in humans cause the severe developmental disorder Cornelia de Lange Syndrome[26,27]. Importantly, mutations in cohesin often produce distinct phenotypic and clinical outcomes from those in its cofactors [28,29], suggesting they perturb different molecular functions. We still do not understand why mutations that reduce overall cohesin dosage are rarely observed in Cornelia de Lange syndrome, whereas heterozygous NIPBL mutations, which typically only reduce overall NIPBL expression by 30-40%, nonetheless cause profound developmental abnormalities[30]. To link chromosome misfolding to transcriptional dysregulation and congenital malformations, we must mechanistically understand how the interplay between individual cohesin cofactors modulates the dynamics of loop extrusion.

Defining specific roles of cohesin cofactors poses biological and technical challenges. Biologically, each cofactor influences multiple facets of DNA loop extrusion. For example NIPBL, initially characterized as a cohesin loader[31–33], is also required for loop extrusion activity *in vitro* [34,35], and its functions in loop extrusion *in vivo* appear to extend beyond acting as a transient loader[36,37]. In contrast, WAPL promotes cohesin unloading from DNA, and its depletion from cells results in the extension of DNA loops [13,38]. PDS5 facilitates WAPL-mediated unloading[39] but also competes with NIPBL for cohesin binding *in vitro*[40,41]. PDS5 depletion from cells extends DNA loops[42–47], resembling the effects of WAPL inactivation, but with differences whose mechanistic basis remains unclear[13,42,48].

Given the biological complexity of the interplay between cohesin cofactors, disentangling their molecular functions in genome folding also poses technical challenges. First, cofactor essentiality limits genetic perturbations in living cells, and changes in their stoichiometry can trigger compensatory effects that obscure mechanistic interpretation. While such limitations may be somewhat alleviated by the use of *in vitro* assays[34,35], these assays are highly sensitive to reconstitution conditions, and do not fully recapitulate native chromatin context or nucleoplasmic availabilities. Second, properties of loop extrusion are not direct readouts from cellular assays, including Hi-C and microscopy. A promising approach to circumvent this limitation, and link the underlying properties of loop extrusion to genome folding features observed *in vivo*, involves the use of *in silico* models[1,4]. Such models, however, require fitting multiple parameters which have been underconstrained by available experimental measurements. New strategies for modeling and inference are thus required to obtain

an unambiguous mapping between cofactor perturbations and individual biophysical parameters of loop
extrusion and gain molecular insight into clinically relevant aspects of cohesin biology.

Here we determined how individual cohesin cofactors influence distinct biophysical parameters of loop extrusion:
the number of loaded cohesins (*separation* between extruders), their residence time on chromatin (*lifetime*), and
how fast they extrude (*rate*). We established these molecular mechanisms with: (1) an unprecedented suite of
degron tags on cohesin and cofactors that enable precisely titrated perturbations in mouse embryonic stem cells
(ESCs); (2) calibrated biophysical models capable of using Hi-C readouts after cellular perturbations to infer
alterations to parameters of loop extrusion. Leveraging this approach, we found that while NIPBL depletion
lowers the fraction of chromatin-bound cohesin, the resulting Hi-C defects are best explained by a reduction in
extrusion rate rather than by decreased loading alone. Furthermore, titrating NIPBL abundance to emulate
altered dosage in Cornelia de Lange Syndrome revealed that NIPBL expression levels quantitatively tune the
rate of loop extrusion in cells. Since extruded loop length is the product of cohesin extrusion rate and lifetime,
the lower extrusion rates caused by NIPBL depletion explain how co-depletion of WAPL, which increases
lifetime, restores normal genome folding patterns, even though cohesin dynamics are not restored[38,49].
Furthermore, depleting PDS5 both accelerates extrusion rates, by competing with NIPBL, and increases cohesin
lifetime, likely by supporting WAPL unloading[50]. We validated model inferences via microscopy measurements
of chromosome morphology and cohesin dynamics in single cells. In sum, our observations reveal a modular
control system for cohesin extrusion that links 3D genome organization to transcriptional states, with direct
relevance to understanding the molecular etiology of cohesinopathies.

Results

Reduced cohesin loading does not explain chromosome folding defects upon NIPBL depletion

Since NIPBL is needed for cohesin loading, we first investigated whether defects caused by loss of NIPBL can
be explained solely by loss of cohesin loading. To acutely deplete NIPBL from mouse embryonic stem cells
(ESCs), we used a cell line[23] where the endogenous protein can be degraded with the $FKBP12^{F36V}$ degron upon
induction by the small molecule dTAG-13[51] - hereafter abbreviated FKBP and dTAG. In these cells, levels of
NIPBL (and its binding partner MAU2) drop below 10% within hours without causing cell cycle defects[23]. Given
the established role of NIPBL in loading cohesin, one might expect that NIPBL-depleted cells would closely
resemble cohesin-depleted cells. We therefore compared the effect of depleting NIPBL to degrading the core
cohesin subunit RAD21, using cells we previously generated[52–54] with the auxin-inducible degron (AID)[55]. Using
Hi-C, we observed marked defects in chromosome folding within hours of depleting NIPBL, resembling short
term RAD21 degradation (Figure 1A and S1B)[23]. These defects arise from impaired cohesin loop extrusion, not
sister chromatid cohesion, as contacts between sisters only marginally contribute to bulk Hi-C patterns in
mammalian cells - even in G2[56].

To precisely measure the amount of cohesin left on chromatin after NIPBL depletion, we tagged the endogenous
core cohesin subunit RAD21 with a Halotag (which had no effect on chromosome folding, Figure S1B) in the
NIPBL-FKBP cells. We then pre-extracted soluble proteins from the cells and precisely quantified the remaining

chromatin-bound cohesin by flow-cytometry[57] (Figure S1C, Methods). With this approach, we estimated that ~60-70% of RAD21 molecules are bound to chromatin in WT ESCs (Figure 1B), in line with previous orthogonal estimates from live imaging and Western blot across a variety of mammalian cell types[36,58–61]. NIPBL depletion only reduced the abundance of chromatin-bound cohesin two-fold (Figure 1B) even after 24h (two full cell cycles, figure S1D), consistent with previous RNAi experiments[13]. Depleting NIPBL did not measurably change cohesin residence time as measured in inverse fluorescence recovery after photobleaching (iFRAP) (Figure 1C), indicating the reduction in chromatin-bound cohesin does not result from increased unloading.

To assess whether the two-fold reduction in chromatin-bound cohesin accounts for the chromosome folding defects observed upon NIPBL depletion, we engineered cells in which cohesin abundance can be directly reduced. We created RAD21-FKBP-Halotag cells and titrated dTAG concentrations to achieve precise intermediate levels of bound RAD21 that we quantified by flow cytometry (Figure S1E). The dramatic Hi-C defects caused by NIPBL depletion were not recapitulated by reducing chromatin-bound cohesin levels two- or even three-fold (Figure 1D and 1E, Figure 2B conditions that do not induce confounding cell cycle defects, Figure S1F).

Acute NIPBL depletion therefore disrupts chromosome folding more severely than expected from the remaining amount of chromatin-bound cohesin. Thus, NIPBL must control chromosome folding through functions beyond simply loading cohesin onto chromatin.

Calibration of biophysical simulations with experimental measurements

Biochemical reconstitution approaches demonstrated that NIPBL is required for cohesin loop extrusion *in vitro* [34,35]. NIPBL only binds cohesin intermittently, rapidly binding and unbinding while cohesin is on DNA - both *in vitro*[62] and *in vivo*[36]. We therefore reasoned that reducing the abundance of NIPBL in cells would increase the amount of time a cohesin molecule spends between two NIPBL binding events, thereby reducing the average rate of loop extrusion. To explore this idea, we first needed to understand how slower extrusion rates would manifest in Hi-C data, and how this would compare to lowering cohesin loading.

To connect changes in cohesin dynamics with genome-wide consequences, we built polymer models tailored to ESCs to predict how variations in loop extrusion parameters would affect genome folding (Figure 2A). Specifically, we determined how simulated contact frequency ($P(s)$) scales with genomic separation (s), which can be directly compared with similar curves computed from experimental Hi-C data. Building on previous approaches[63–65], we coupled a 1D lattice model of loop extruder dynamics with a 3D bead-spring polymer model of the chromatin fiber (Figure S2A). We parameterized the unperturbed basal state based on available data (similar to previous studies[64,66], Figure S2B and Methods), simulating 50 Mb of chromatin as a fiber of 50 nm monomers, each representing 2.5 kilobases (kb) of DNA. We set the separation between extruders in kb based on cohesin abundance previously measured by Fluorescence Correlation Spectroscopy (FCS)[67] (Figure S2B). We set the extruder lifetime based on FRAP data[58], and the correspondence between simulated and experimental time based on locus tracking data[68,69]. Finally, A/B compartmentalization was modeled as a block copolymer that mirrors the arrangement of active and inactive regions along an endogenous chromosome

(Figure S2C, S2D and Methods). To estimate the rate of extrusion, for which direct *in vivo* measurements are currently lacking, we varied the number of simulated 3D molecular dynamics steps of the polymer per update of extruder positions along the 1D polymer lattice and fit to the experimental Hi-C *P(s)* in untagged wild-type cells (Figure S2E, S2F and Methods). The resulting model for WT untagged cells corresponds to an average separation of 185 kb between extruders, a lifetime of 22 minutes, and average extrusion rate of 0.25 kb/sec.

Polymer simulations support a prominent role for NIPBL beyond cohesin loading in controlling loop extrusion rate

Using simulations, we investigated what changes in extrusion parameters could account for the chromosome folding defects we observed after NIPBL depletion. Experimental *P(s)* curves from NIPBL-depleted (with NIPBL-FKBP) or partially RAD21-depleted cells (with RAD21-FKBP-Halotag) both had reduced contact frequencies between ~20 kb-2 Mb (Figure 2B). With polymer simulations, we then assessed how *P(s)* curves change with decreasing numbers of extruders, equivalent to increasing the genomic separation between extruders (Figure 2C). Clear, though subtle, differences were observed in simulations across the range of extruder separations considered. We thus considered the goodness-of-fit between simulated and experimental *P(s)* curves to determine the best-fitting simulation parameters. Remarkably, the best fit to the experimental *P(s)* profile from cells with chromatin-bound RAD21 reduced to 50% was achieved in simulations with 2-fold increased extruder separation (Figure 2D). Similarly, the best fit to the experimental *P(s)* profiles from cells with chromatin-bound RAD21 reduced to 30% was achieved with simulations with 3-fold increased extruder separation. Therefore, this fitting approach between experimental and simulated data can account for subtle changes in *P(s)* by correctly incriminating the loop extrusion parameter corresponding to the number of extruders (compare experiments in Figure 2B to best fitting simulations in 2E).

We next explored the effect of decreasing extrusion rates on *P(s)* in simulations (Figure 2F). We observed that lowering extrusion rates had much greater effects than increasing cohesin separation (compare to Figure 2C). Although NIPBL-depleted cells have 2-fold less bound cohesin (Figure 1B), increasing extruder separation alone yielded poor fits to the experimental Hi-C data (Figure 2G). In contrast, lowering extrusion rates in the context of 50% chromatin-bound cohesin provided an excellent fit to the experimental NIPBL Hi-C, with best agreement obtained at 40-50% of initial extrusion rates (Figure 2H). Further supporting our inference, varying extrusion rate alone offered poor fits to the partially depleted RAD21 datasets (Figure S2G). Simulations thus provide quantitative support that the Hi-C defects observed after NIPBL depletion cannot be explained by reduced cohesin loading alone, and instead suggest that NIPBL abundance could shape chromosome folding by modulating the rate of loop extrusion.

Qualitative support for the differential impact of NIPBL versus RAD21 depletion comes from considering how the inflection point in *P(s)* shifts in these two conditions (Figure S2H). This feature of *P(s)*, termed s[P'max], relates to the average length of loops when they are sufficiently separated[70]. For NIPBL depletion, s[P'max] decreased (from ~90 kb to ~55 kb). In contrast, after partial cohesin depletion, s[P'max] increased (up to ~150 kb for 50% remaining bound cohesin, or ~210 kb for 30%) - Figure S2H. These distinct effects of NIPBL vs. partial cohesin depletions on s[P'max] were reproduced in simulations (Fig 2E). By monitoring the location of extruders on simulated polymers we observed that, in the baseline regime, nearly 70% of extruders have collided on at least one side (Figure 2I). At higher extruder separations, collisions are reduced and average loop lengths

increased (Figure S2I), explaining how depleting RAD21 led to a rightward shift in the *P(s)* inflection point (Figure 2B and 2E).

Altogether, simulations and experiments indicate that cohesin collisions limit loop growth in the normal regime (Figure 2J). Lowering cohesin abundance with partial RAD21 depletion increases the separation between extruders and enables loops to grow slightly larger. Therefore, the *P(s)* measured after NIPBL depletion cannot be explained by reduced cohesin loading. These observations further support the idea that, in addition to the reduction of bound cohesin levels, NIPBL-depleted cells experience a lower rate of extrusion.

Reduced extrusion rate after NIPBL depletion, not reduced cohesin loading, can counteract the increase in cohesin lifetime caused by WAPL inactivation

A role for NIPBL in setting the rate of loop extrusion yields testable predictions about how it interacts genetically with cofactors controlling other steps of the extrusion process. Several studies have reported that impairing NIPBL functions can prevent the 3D genome defects observed upon WAPL inactivation[38,49,71]. WAPL unloads cohesin and thereby controls lifetime[32,72,73]. Yet, NIPBL co-depletion does not restore cohesin lifetime(Haarhuis et al. 2017). How, then, do NIPBL and WAPL depletions offset each other in chromosome folding?

To distinguish whether compensation arises from reduced extrusion rate or decreased cohesin loading, we engineered WAPL-AID ESCs[52,74] to either co-deplete NIPBL or partially co-deplete RAD21. As previously reported in other cell types, Hi-C patterns and *P(s)* curves from cells acutely co-depleted for WAPL and NIPBL closely resembled that of undepleted cells (Figure 3A and 3B). In contrast, simultaneously depleting WAPL and reducing RAD21 levels did not cause Hi-C patterns to resemble those of undepleted cells, even when titrated to precisely match the levels obtained after NIPBL co-depletion (Figure 3A, 3B and S3A). This indicates that NIPBL co-depletion mitigates WAPL inactivation through means other than cohesin loading.

Viewing NIPBL as controlling the rate of loop extrusion immediately clarifies how its co-depletion counteracts the genome folding defects caused by WAPL inactivation, in spite of cohesin lifetime remaining aberrantly high. Indeed, given that processivity, the average distance traveled by an unimpeded extruder along the genome, is the product of lifetime and rate (Figure 3C), the decrease in loop extrusion rate from NIPBL depletion offsets the increase in lifetime from WAPL depletion (Figure 3D).

Beyond ascertaining the role of NIPBL in controlling the rate of loop extrusion, these results demonstrate that opposing changes to distinct extrusion parameters can counterbalance one another to buffer chromosome architecture despite altered cohesin dynamics, and provide a long-sought mechanistic explanation for the genetic interaction between NIPBL and WAPL.

Hi-C based evidence that PDS5 depletion accelerates loop extrusion rates

We next considered the role of cellular PDS5 in loop extrusion. PDS5 is a protein that binds cohesin and promotes WAPL-mediated cohesin unloading [75–77]. Yet PDS5-depleted cells do not exhibit Hi-C patterns identical to WAPL-depleted cells, indicating that PDS5 has functions in genome folding beyond mediating unloading[13,42,48]. *In vitro* assays revealed that PDS5 docks to cohesin on the same site as NIPBL, and their binding is mutually exclusive: PDS5-bound cohesin has no ATPase activity, and cannot extrude loops [34,40,41]. We therefore tested whether the differences in genome folding between WAPL- and PDS5-depleted cells arise from how PDS5 competes with NIPBL for cohesin binding: if PDS5 depletion enables NIPBL to bind cohesin more efficiently or frequently, one would expect extrusion rates to increase.

We performed Hi-C in cells depleted for both PDS5A and PDS5B (hereafter PDS5AB) for 6h using the AID system, as we observed that both paralogs contribute to genome folding in ESCs (Figure S4A-D). In line with previous reports in other cell types[13,42], PDS5AB depletion resulted in more altered *P(S)*(Figure 4B), and less sharply defined Hi-C dots compared to WAPL depletion (Figure 4A). Consistent with unloading defects, both WAPL and PDS5AB depletion increase the amount of chromatin-bound cohesin (Figure S4E). These results are not solely driven by cell cycle defects, as short-term depletion of WAPL in ESCs does not affect the cell cycle[52], nor does PDS5AB depletion (Figure S4F). Longer PDS5AB depletion caused S-phase stress, as previously reported[15,78], but only beyond 24h.

To elucidate how WAPL and PDS5AB depletion differentially impact genome folding, we used simulations and inferred which kinetic parameters of extrusion are altered in each situation. Starting from the benchmarked model of untagged wild-type ESCs (Figure 2), we increased cohesin lifetime and extrusion rate either in isolation or in combination, and compared *P(s)* from each simulation to the experimental Hi-C after WAPL or PDS5AB depletion. To limit our sweep to lifetime and extrusion rate, we calibrated extruder separation to match the amount of chromatin-bound cohesin measured experimentally after each perturbation (by flow cytometry, Figures S4E and S4G). Instead of converging on single values, this analysis highlighted a large number of lifetime and rate combinations that could fit the experimental Hi-C *P(s)*, indicating that this metric alone was unable to distinguish the contribution of higher cohesin lifetime from that of higher extrusion rate (Figure 4D). This can be explained by simulated *P(s)* being largely determined by cohesin processivity, which depends equally on lifetime and rate (processivity = lifetime * rate). We therefore sought additional metrics beyond *P(s)* to disambiguate these two components.

We noticed that, in simulations, increasing extrusion rate reduced the segregation of A and B compartments more effectively than increasing extruder lifetimes (Figure 4E). We therefore used compartmentalization scores to disambiguate the contribution of cohesin lifetime versus extrusion rate to processivity. Experimentally, PDS5AB depletion lowered compartment scores further than WAPL depletion (Figure 4F). Requiring a joint best-fit to both experimental *P(s)* and compartment scores after WAPL and PDSAB depletion (Figure S4G) led to a unique set of best fitting parameters for each condition. PDS5AB depletion only increases cohesin lifetime three-fold (about half of WAPL depletion), but increases extrusion rates three times (Figure 4G). These observations indicate that even though cohesin lifetime is less perturbed by lowering PDS5AB dosage than WAPL dosage, the increase in extrusion rate drives more profound changes in genome folding.

Microscopy-based evidence that PDS5 depletion does not fully disable unloading but accelerates loop extrusion

We next tested predictions of the PDS5AB and WAPL simulations by considering orthogonal manifestations of loop extrusion that were not used to inform the selection of model parameters. Disabling unloading causes extruding cohesins to collide and form axial structures known as vermicelli[8,79], visible as spatial clusters of cohesins in the nucleus. We therefore analyzed the location of extruders in individual simulated conformations, performing *in silico* microscopy, using the parameters for lifetime and extrusion rate that we previously identified as best fitting the Hi-C data. While simulations of WAPL depletion exhibited some clustering of extruders, simulations of PDS5AB depletion displayed more pronounced axial structures (Figure 5A). We also observed this pattern experimentally using fixed-cell microscopy: PDS5AB depleted cells displayed stronger cohesin clustering than WAPL depleted cells (Figure 5B). Microscopy thus argues that cells depleted for PDS5AB experience faster extrusion rates than WAPL depleted cells, supporting the distinct roles of these cofactors we deduced from fitting simulations to Hi-C.

To more directly ascertain that PDS5AB depletion increases the rate of cohesin extrusion, we assayed the speed of vermicelli formation in synchronized cells. Cohesin is naturally removed from chromosomes in mitosis, only initiating extrusion at the onset of G1[80]. If chromosome extrusion is faster after PDS5AB than WAPL depletion, then compaction into vermicelli structures in early G1 should also occur faster after PDS5AB depletion. To test this, we used nocodazole to synchronize ESCs in M phase (Figure 5C, S5A and S5B), at which point most cohesin is removed from chromosome arms through the normal prophase pathway. We then depleted either WAPL or PDS5AB for three hours by adding auxin, collected mitotic cells by mitotic shake-off, and released cells into G1 by washing off nocodazole (keeping auxin in the medium to suppress either WAPL or PDS5AB expression). A microscopy time-course over six hours after release into G1 revealed that vermicelli form faster and end up more pronounced in the absence of PDS5AB than in the absence of WAPL (Figure 5C). These experiments therefore ascertain that PDS5AB depletion accelerates the rate of loop extrusion, and rule out possible technical confounders in asynchronous cells, such as eventual differences in cell cycle length or degron leakiness between cell lines.

We then tested the model prediction that WAPL depletion leads to a greater increase in cohesin lifetime than PDS5AB depletion using iFRAP for RAD21-Halotag in live ESCs. We observed that PDS5AB depletion increased the residence time of cohesin, but not as much as WAPL depletion (Figure 5D, S5C and S5D), in line with RNAi experiments in HeLa cells[13]. iFRAP thus quantitatively supports the parameters inferred from our calibrated polymer simulations and Hi-C fitting: while cohesin lifetime increases upon PDS5AB depletion, it does not reach the levels seen after WAPL depletion.

Altogether, these observations ascertain that PDS5 normally limits the rate of loop extrusion, accounting for its WAPL-independent role in genome folding. Higher extrusion rates readily explain the Hi-C *P(s)* observed upon PDS5AB depletion (Figure 5E), as well as the stronger nuclear clustering of cohesin seen by microscopy (Figure 5E). These results also demonstrate how polymer simulations, although only instructed by Hi-C data and cohesin abundance from flow-cytometry data, can both quantitatively predict meso-scale chromosome organization observable by microscopy, and assign specific molecular roles in extrusion kinetics to cohesin cofactors.

The NIPBL:PDS5 balance shapes chromosome folding by tuning the rate of cohesin extrusion

Given that NIPBL and PDS5 can compete for cohesin, we wondered if beyond controlling cohesin loading and unloading, their relative abundance sets the net rate of extrusion in cells. We therefore performed Hi-C after acute co-depletion of NIPBL and PDS5AB (Figure 6A and B). If PDS5 and NIPBL have opposing effects on extrusion rate, their co-depletion should restore normal extrusion dynamics. However, since PDS5AB depletion also increases cohesin lifetime, PDS5AB+NIPBL co-depleted cells should still be partly deficient in unloading. Therefore, the expectation is that Hi-C in PDS5AB+NIPBL co-depleted cells would resemble WAPL-depleted cells, which have increased lifetime. This is indeed what we observed experimentally (Figure 6A-B).

We next tested whether the resemblance in Hi-C patterns between PDS5AB+NIPBL co-depleted cells and WAPL depleted cells arises from balancing of extrusion rates or balancing of loading and unloading. We generated RAD21-FKBP-Halogtag subclones from the PDS5AB-AID ESCs and, in the context of PDS5AB depletion, titrated RAD21 using limiting amounts of dTAG to reduce bound cohesin to levels that are comparable (80%) or lower (45%) than those in PDS5AB+NIPBL co-depleted cells (Figure S6A). Unlike co-depleting NIPBL, lowering cohesin abundance on top of PDS5AB depletion did not create Hi-C patterns reminiscent of WAPL depletion (Figure 6A and B). On the contrary, reducing RAD21 abundance shifted the $P(s)$ inflection point to substantially higher genomic distances, consistent with a decrease in cohesin-cohesin collisions. As shown in Figure 2, such collisions normally constrain loop growth, and hence fewer collisions would enable loops to extend further upon PDS5AB depletion.

To understand the complex chromosome-folding defects that result from disrupting two cohesin cofactors, we simulated each co-depletion. We assigned cohesin separation multipliers using flow cytometry measurements of chromatin-bound RAD21 (Figures S6A, S4B). We set lifetime and extrusion rate by multiplying the parameters from each of the two single depletions (from Figures 2 and 4). The resulting simulated $P(s)$ for co-depletions mirrored those for experimental data, supporting the mechanistic explanations we provide for individual co-factors: in PDS5AB+NIPBL co-depletion, the effect of NIPBL depletion lowering extrusion rate is counteracted by PDS5AB depletion raising extrusion rate and lifetime. As a result, after PDS5AB+NIPBL co-depletion rate is re-balanced but lifetime remains elevated, explaining the close resemblance to WAPL depletion (Figure 6B, 6C and 6D). In contrast, reducing cohesin levels by partial depletion of RAD21 does not compensate for the increase of rate and lifetime caused by PDS5AB depletion.

We reached similar insight modeling co-depletions with WAPL: simulations of WAPL+NIPBL co-depletion (Figure S6B) also matched experimental data (Figure 3B). Increasing lifetime (due to WAPL depletion) while reducing extrusion rate (due to NIPBL depletion) results in normal processivity and therefore a $P(s)$ curve similar to WT untagged cells. In contrast, RAD21 partial depletion does not compensate for the longer lifetime in WAPL depletion (Figure S6B).

Altogether, our observations demonstrate that the cellular balance between NIPBL and PDS5 abundance sets the overall rate of cohesin extrusion (Figure 6E), in line with a molecular competition scenario[40,41] combined with short-lived NIPBL binding to cohesin[36,62]. In this context, a single cohesin molecule would undergo extrusion for as long as NIPBL remains bound to it. When NIPBL unbinds, extrusion pauses until the next binding event. PDS5 would shield cohesin from NIPBL, thereby extending the interval between extrusion bursts.

NIPBL expression levels determine the rate of cohesin extrusion

A key prediction of a molecular competition scenario between NIPBL and PDS5 is that increasing NIPBL expression in cells with normal PDS5 levels should accelerate extrusion. Remarkably, we observed that simply overexpressing NIPBL induces vermicelli formation (Figure 6F), consistent with faster extrusion rates.

Given the haploinsufficient nature of NIPBL for human development, we sought to understand how the rate of cohesin loop extrusion scales with NIPBL expression levels. For this we generated a mouse ESC line expressing NIPBL-FKBP fused to GFP, alongside RAD21-Halotag. This allowed precise monitoring of intermediate NIPBL levels through flow cytometry, after titrating its expression level to generate a range of partial dosage reductions that include and extend beyond the expression levels observed in Cornelia de Lange syndrome. Consistent with the original NIPBL-FKBP line[23] (lacking GFP), basal NIPBL levels were slightly reduced (~80% of WT untagged cells) as estimated by Western blot, likely reflecting slight leakiness of the degron system (Figure S6C). Using limiting amounts of dTAG we generated NIPBL-FKBP-GFP cells with ~50% and ~22% NIPBL levels (Figure S6D) and profiled them by Hi-C, and analyzed them side-by side with previously described NIPBL-FKBP undepleted (80% NIPBL levels) and depleted to ~7%. We measured bound RAD21-Halotag levels by flow cytometry (Figure S6E), enabling calibration of simulated extruder separation. We performed a series of polymer simulations sweeping across a range of extrusion rate multipliers to identify which rate values best fit experimental Hi-C *P(s),* for each NIPBL dose (Figure 6G). We found that inferred rates increased with NIPBL levels across the five samples (Figure 6G). These analyses indicate that NIPBL dosage quantitatively tunes the rate of DNA loop extrusion in cells, thereby modulating genome folding, with immediate implications for clinical contexts where dosage is only partially reduced, such as Cornelia de Lange syndrome.

Cohesin regulators differentially modulate genomic compartmentalization and transcriptomes

To evaluate the generalizability of our models, we examined their predictions for compartmentalization (Figure S7A-B). To simulate WAPL+NIPBL and PDS5AB+NIPBL co-depletions, lifetime and extrusion rate were estimated by multiplying inferred parameters from single perturbations. All simulations used experimentally-determined cohesin separation, obtained as above by flow cytometry of chromatin-bound RAD21. Remarkably, the ranking of compartment scores across experimental depletions (Figure 7A) was identical to the ranking of simulations (Figure 7B and S7A-C), aside from partial co-depletions of RAD21 on WAPL and PDS5AB backgrounds. Indeed, PDS5AB+NIPBL codepletion had higher compartmentalization than WAPL alone, and WAPL+NIPBL co-depletion had higher compartmentalization than even wild type cells, both in simulations and experiments.

Importantly, reducing the extrusion rate in PDS5AB mutants, via NIPBL co-depletion, greatly restored compartment strength across genomic distances in *cis* and in *trans* (Figure S7D). While partial RAD21 depletion increased compartmentalization in simulations on the background of PDS5AB or WAPL depletion, experimental compartmentalization was enhanced more than predicted for both of these scenarios. We speculate that this stems from the greater relative availability of these residual subunits when cohesin is simultaneously downregulated. Indeed, the higher PDS5AB:RAD21 and WAPL:RAD21 ratios in these co-depleted cell lines

could help rescue cohesin lifetimes and extrusion rates towards their basal values. This effect is neglected by the current assumption that the net changes in extrusion parameters induced by simultaneous co-depletions are strictly multiplicative (Figure S7C). Together, our findings illustrate the broad applicability of our approach to infer extrusion lifetime and rate, and showcase how varying cohesin parameters – in particular, of extrusion rate – can dramatically alter compartmentalization patterns (Figure 7C).

Finally, we tested whether Hi-C defects tracked with overall transcriptional changes by performing RNA-seq across six depletion conditions, including our previously reported NIPBL degradation[23]. We profiled transcriptomes after 1 day of depletion to capture enough dysregulated genes to compare cellular states between perturbations, refraining from correlating transcriptional dysregulations with local genomic features, as the secondary defects would confound such analyses. Samples with similar contacts versus distance curves (as reflected by their relative *P(s)* inflection points, or Hi-C s[P'max] values, see Figure S2G) displayed similar transcriptomes (as estimated by PC1 from principal component analysis, Figure 7D and 7E). Notably, NIPBL+WAPL co-depletion not only produced Hi-C patterns resembling those of untagged cells, but also led to clustering of RNA-seq profiles with untagged controls. Similarly, NIPBL+PDS5AB depletion and WAPL depletion, which caused similar Hi-C patterns, led to similar transcriptomic profiles. PDS5AB depletion, which caused the most significant change in processivity, also induced the greatest overall transcriptomic alterations. Altogether, these findings illustrate that changes in loop extrusion kinetics not only affect 3D genome folding but can also have a profound impact on overall transcriptional states, providing molecular insight into how genetic interactions between cohesin cofactors can correct transcriptomes disrupted by alterations in specific loop extrusion parameters.

Discussion

Here we combined experimental and computational approaches to derive a quantitative understanding of how cohesin cofactors contribute to loop extrusion and shape genome architecture. We uncovered a prominent role for NIPBL beyond cohesin loading: higher NIPBL expression levels increase the rate of extrusion through a mechanism antagonized by PDS5. These findings provide a long-sought mechanistic explanation for how modest reductions in NIPBL expression, such as those found in Cornelia de Lange syndrome, can disrupt genome organization, even when overall cohesin levels on chromatin remain high[81,82]. By integrating genetic perturbations and polymer simulations in a unified framework, our approach quantitatively defines how the relative abundance of NIPBL and PDS5 sets the rate of extrusion and shapes chromosome folding patterns in living cells. These findings align with previous studies that demonstrated NIPBL and PDS5 can compete for cohesin *in vitro*[40,41] and that PDS5 inhibits loop extension[42,45]. Furthermore, we reveal that extrusion rate and cohesin lifetime exert distinct yet overlapping effects on chromosome folding, and uncover a previously unrecognized, yet critical, role for the extrusion rate in modulating compartmentalization between active and inactive chromatin. Finally, we provide new mechanistic insight into how cofactors that modulate distinct aspects of loop extrusion can nonetheless produce compensatory effects on genome folding and transcriptional states.

An experimentally calibrated biophysical framework connects dynamics of individual cohesins to chromosome-scale organization

We employed biophysically calibrated simulations of loop extrusion to infer the role of individual cohesin cofactors from a diverse array of perturbation conditions. We approached this challenge as a system of three unknowns - loaded cohesin number, extrusion rate, and lifetime - that we solved by quantitatively estimating three metrics experimentally: flow cytometry for chromatin-bound cohesin, and *P(s)* and compartment scores from Hi-C. After setting the loaded number with the flow cytometry estimate, this then required sweeping lifetime and extrusion rate in simulations and determining their respective values by fitting to the combination of Hi-C metrics. Prior approaches largely used *P(s)* and its derivatives[63,83] to fit polymer models to Hi-C data and infer loop sizes and parameters of extrusion. However, as we demonstrate here, *P(s)* curves alone are insufficient to distinguish between altered extrusion rate or lifetime (Figure 4). Additional consideration of compartment strengths allowed us to disambiguate these two parameters and specifically understand the role of extrusion rate in genome folding. This provided mechanistic insight into the distinct contributions of WAPL and PDS5 to loop extrusion, where PDS5 depletion moderately increases both extrusion rate and lifetime, whereas WAPL depletion does not increase rate but elevates lifetime to greater extent .

We first validated our parameter inference approach with partial RAD21 depletions that had known levels of residual cohesin. To our knowledge, this was the first demonstration that simulations can quantitatively reproduce the effects of titrated cohesin dosage. Second, as orthogonal validation of our predictions, we turned to imaging measurements: we measured cohesin lifetime in live cells with iFRAP; we also compared simulation coordinates to experimental vermicelli images in fixed cells, via a quantitative numerical microscopy protocol. Altogether these approaches underscore the value of simulations for integrating multimodal data, along with their ability to consistently capture genome folding structure and kinetics at both the population (Hi-C) and single-cell level.

An important consideration of our simulation approach was to explicitly model extrusion rate for comparison with experimental datasets. This enabled us to infer a prominent role of NIPBL abundance on extrusion rate, beyond its role in loading. Explicitly modeling extrusion rate contrasts with previous investigations of the more technical parameter of 3D-to-1D timesteps inside of simulations, which controls the relative kinetics of cohesin versus the relaxation rate of the polymer [64,84]. Importantly, our estimated average rate of ~0.25 kb/s in wild-type ESCs reflects the average overall chromatin-bound cohesin molecules, and only a subset are expected to be associated with NIPBL and actively engaged in extrusion at any given time[36]. Indeed, high-resolution live-cell tracking of DNA loci has reported rare bursts reaching ~2.7 kb/s, which might specifically correspond to the rate of NIPBL-bound, actively extruding cohesin[85]. We speculate that variations in the abundance of NIPBL and chromatin-bound cohesin across cell types could account for differences in estimated rates across studies (e.g., 0.85 kb/s in HeLa[86]).

Our simulations and experimental titration of RAD21 in the background of various other perturbations supports the prevalence of cohesin-cohesin collisions, even in the basal regime in WT untagged ESCs. Indeed, accounting for reduced collisions was essential for accurately predicting *P(s)* changes after partial RAD21 depletion. Still, our observations call for caution when interpreting derivatives of *P(s),* as the quantitative relationship with loop size is incompletely determined in biologically relevant scenarios[70]. Indeed, in simulations, the inflection point in *P(s)* was substantially less than loop size, which in turn was substantially less than the processivity of an unimpeded extruder (Figure S2I). While frequent cohesin collisions can lead to stacking of loops, higher-order rosettes are however rare in cells[53] and likely transient because of WAPL-mediated turnover.

Finally, our observations for interphase cohesin parallels recent observations in mitosis where condensin-condensin encounters mainly result in collisions rather than bypass [87].

NIPBL for loop extrusion: loader, processivity factor or both?

A substantial body of literature has established NIPBL as a cohesin loader [31–33]. However, these investigations primarily addressed the role of NIPBL in establishing sister-chromatid cohesion, not in the context of cohesin extrusion. Extruding and cohesive forms of cohesin differ significantly - for example cohesion involves topological entrapment and does not require NIPBL for maintenance[56,88]. In contrast, reconstitution studies demonstrated that the association of NIPBL with cohesin is continuously required for extrusion *in vitro*[34,35,62], which has led to the proposal that NIPBL functions as a processivity factor during cohesin extrusion[5], and that there may be no *bona fide* loading complex for extrusive cohesin [34]. Our *in vivo* data demonstrate that the contribution of NIPBL to processivity plays a prominent role at all levels of 3D genome folding examined.

The observation that NIPBL depletion reduces cohesin levels on chromatin does not necessarily imply that NIPBL performs a dedicated loading reaction for extrusion. Indeed, NIPBL may license extrusion after cohesin associates with DNA, potentially via the STAG1/2 subunit[20,89–91]. In this context NIPBL would still effectively act as a loader, by enabling cohesin to engage in the extrusion cycles that prevents its dissociation from chromatin until WAPL-mediated release[86]. Further investigation using cell types that can be efficiently synchronized, along with inactivation strategies that completely disable NIPBL activity, will be crucial to resolve these questions.

Dual roles of cellular PDS5: lowering extrusion rates and promoting unloading

We disentangle contributions of PDS5 on extrusion rate from unloading. While genetic data supports the antagonism between PDS5 and NIPBL, the molecular basis of their competition remains incompletely understood. In particular, it is unclear whether PDS5 actively displaces NIPBL in cells, or whether it only binds cohesin after NIPBL spontaneously dissociates. It will be of particular interest to compare the dissociation kinetics of NIPBL and PDS5 *in vivo*, and to examine their interplay with the dynamics of cohesin acetylation, which is known to alter the affinity of PDS5 for cohesin[42] and interfere with WAPL unloading[92]. The exact role of PDS5 during unloading remains to be clarified. Possible requirements for release include: simultaneous PDS5 and WAPL co-binding, sequential binding[50,86,93], or independent pathways operated by PDS5 and WAPL [13].

The interplay of cohesin-cofactors imparts exquisite tunability to interphase extrusion

Our findings demonstrate that altering cohesin cofactor levels can selectively influence certain extrusion parameters while buffering others. For example, concomitant downregulation of PDS5 and NIPBL preserves extrusion rates while increasing cohesin lifetime, and downregulation of both WAPL and NIPBL preserves overall processivity despite elevated cohesin lifetime. Notably, while increases in extrusion rate and cohesin lifetime both enhance processivity equally, our observations highlight that they lead to different outcomes in cohesin dynamics and compartmentalization. We hypothesize that this balancing act enables cells to finely tune specific aspects of genome organization while preserving others, allowing flexible yet precise control over long-range chromosomal interactions. For example, such modularity may be key for orchestrating communication between

cis-regulatory elements while independently shaping patterns of genome compartmentalization in ways that contribute to long-term epigenetic memory[94].

Developmental relevance

The expression levels of cohesin and its regulators vary across cell types, providing regulatory opportunities to fine-tune specific aspects of genome folding and achieve distinct genomic functions. For example, WAPL, PDS5A, and PDS5B levels are especially high in ESCs, where extrusion processivity appears lower[95], and their expression is crucial for the maintenance of pluripotency and proper silencing of facultative heterochromatin[43,52,96,97]. In contrast, other cell types naturally express very low levels of WAPL and experience higher extrusion processivity, as seen in pro-B cells, where high cohesin processivity facilitates long-range VDJ recombination of the *Igh* locus[9,10], and in olfactory sensory neurons, where it allows long-range enhancer activation across the large *Pcdh* loci[11,12]. By modulating the chromosomal interactions that underlie Topologically Associating Domains[98], cell type-specific changes in extrusion kinetics offer a dynamic mechanism to reconfigure regulatory landscapes across cell types.

Loop extrusion is a major player in the morphogenesis of meiotic chromosomes across species[99,100]. Consistent with the role of PDS5 in lowering extrusion rates, disabling PDS5 in germ cells shortens the meiotic chromosome axis and affects crossover frequency in yeast and mice [18,101]. It is tempting to speculate that, akin to our observations in interphase, the NIPBL:PDS5 ratio directly regulates extrusion rates in meiotic cells, with immediate relevance for the evolution of genetic variation.

Of note, some organisms possess several paralogs for PDS5 (e.g., five in *Arabidopsis thaliana*)[102], the specific functions of which remain to be investigated. While meiosis-specific paralogs exist for all core cohesin subunits in many organisms, NIPBL or PDS5 are typically expressed in both somatic and germ cells. The distinct evolutionary constraints exerted on cohesin subunits versus their dynamically associating subunits highlight a division of labor with developmental implications that remain to be explored.

Insight for cohesinopathies

Our work underscores that NIPBL and cohesin abundance differentially influence chromosome folding, bringing new perspectives to the understanding of cohesinopathies. Downregulation of cohesin itself results in fewer but longer DNA loops, as extruders collide less often. In contrast, reduced NIPBL abundance slows the growth of DNA loops, and can cause stronger increases to compartmentalization than cohesin downregulation alone. These mechanistic insights shed light on how altered cohesin or NIPBL expression lead to distinct genome folding defects and are associated with divergent disease outcomes[103]. Recognizing that NIPBL expression quantitatively tunes extrusion rates is especially relevant to our understanding of the molecular mechanisms underlying Cornelia de Lange syndrome: lower extrusion rates not only shorten DNA loops but also alter their dynamics, dysregulating transcription of gene loci in cell types where they rely on distal enhancers, ultimately derailing developmental processes[23]. A lower rate of extrusion can account for the impaired distribution of cohesin along chromosomes reported in models of Cornelia de Lange syndrome[25,82]. Beyond Cornelia de Lange

syndrome mutations that reduce NIPBL dosage, these considerations are also relevant to mutations that lower the extrusion activity of cohesin without necessarily altering NIPBL abundance directly[104].

Transcriptional and cell differentiation defects caused by NIPBL impairment can be prevented or even reversed through inhibition of WAPL[49,71,105,106], highlighting significant therapeutic potential. By revealing that this compensation hinges on the rate of loop extrusion rather than cohesin loading, our findings will inform rational approaches to therapeutic intervention in Cornelia de Lange syndrome and other cohesinopathies.

Summary

Altogether, this work illuminates how distinct cohesin cofactors regulate different steps of the extrusion process, and how their tunable expression can modulate multiple aspects of chromosome organization. This loop extrusion control system, while operating at the molecular level, enables the emergence of higher-order structural and functional properties of the genome. Our findings open new avenues for investigating the molecular processes contributing to the dynamic organization of chromatin, with broad implications for understanding genome function and stability in health and disease.

Limitations of the study

As our models are calibrated for extruder separation using experimental data, the accuracy of our extrusion parameter estimates is constrained by how precisely extrusive chromatin-bound cohesin can be quantified. Future developments in quantification methodologies will sharpen the estimates of loop extrusion parameters derived with our framework. Additionally, cohesin extrusion parameters may vary across the genome, which genome-wide $P(s)$-based analyses do not capture. Our baseline model was calibrated to published estimates of absolute cohesin abundance (from FCS and mass spectrometry) and cohesin lifetime (from FRAP), which might be imprecise. However, such uncertainty in baseline parameters is not expected to affect their inferred fold changes after cofactor perturbations. Our simulations do not model extrusion barriers such as CTCF, the transcription machinery or MCM complexes. As these factors have minor effects on genome-wide Hi-C $P(s)$ [13,107–109], they are not expected to significantly alter our parameter inferences. Another limitation is the use of cycling cells, due to the technical challenge of arresting large numbers of ESCs and releasing them while maintaining viability, which precluded bulk assays in precisely staged populations. Finally, while degron-mediated strategies are useful to circumvent the essentiality of cohesin cofactors, the depletions are not complete, and basal expression is often lower than in untagged cells (especially for AID tags).

Resource availability

Lead contact

For further information and requests, please contact the lead contact.

Materials availability

Stable material generated in this study, such as cell lines and plasmids, will be made available by the lead contact upon request upon completion of a Material Transfer Agreement (MTA).

Data and code availability

Hi-C and RNA-seq data have been deposited to the Gene Expression Omnibus (GEO) database. Requests related to polymer simulations will be fulfilled by co-corresponding author Geoffrey Fudenberg. Microscopy data reported in this paper will be shared by the lead contact upon request. Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work the authors used chatGPT as a writing and coding assistant. After using this tool the authors reviewed and further edited, and take full responsibility for the content of the publication.

Acknowledgements

We thank Elzo de Wit for helpful discussions and sharing WAPL-AID ESCs. We are grateful to Daniele Canzio, Benoit Bruneau and the Nora lab for critical reading of the manuscript. We are deeply appreciative of the Luke Lavis lab and the Open Chemistry team (Janelia) for the gift of JF and JFX dyes. We are thankful to: Kevin So in the lab of Benoit Bruneau for providing technical support isolating PDS5-AID cells at the early phase of this project, Pia Mach for advice optimizing microscopy analyses of ESCs, Laurent Gelman and the Facility for Advanced Imaging and Microscopy (FAIM) at the FMI. We thank the Nora and Fudenberg labs for stimulating discussions.

Funding

Grant NIH 1R35GM142792-01 to EN, the Chan-Zuckerberg Biohub San Francisco Investigator program to EN, the Hellman Foundation UCSF Fellows program to EN. Research in GF's lab is supported by NIH R35GM143116 to GF. Research in LG's lab is supported from the Novartis Research Foundation, the Swiss National Science Foundation (grant no. 310030_192642). KH and DW were supported by National Institute of Health training grant 5T32HD007470 for the UCSF Developmental and Stem Cell Biology Graduate Program and the UCSF Discovery Fellows Program, and KH received support from the California Institute for Regenerative Medicine Scholars Training Program education grant EDUC4-12812 for UCSF. DW was supported by the National Science Foundation GRFP and NICHD Predoctoral Training in Developmental Biology Grant 2T32HD007470. EA was supported by postdoctoral fellowships from the Helen Hay Whitney Foundation

and the Regenerative Medicine Scholars Training Program education grant EDUC4-12812 for UCSF. Part of the sequencing was performed at the UCSF CAT, supported by UCSF PBBR, RRP IMIA, and NIH 1S10OD028511-01 grants.

Author contributions

RS performed tissue culture, genome editing, fractionation-flow-cytometry, Hi-C, RNA-seq, Western blot and contributed to RNA-seq analyses. MMCT performed polymer simulations and analysis of experimental Hi-C data. NL performed iFRAP experiments and analyses under the supervision of LG. HR contributed to polymer simulations. EA contributed to genome editing experiments. KH and DW contributed to the RNA-seq analyses. GF developed preliminary polymer simulations and supervised polymer simulations. EPN performed most genome editing experiments, fixed cell microscopy, cell synchronization experiments, supervised all wet lab experiments except iFRAP, analyzed experimental Hi-C, RNA-seq and fixed cell microscopy. EPN and GF jointly designed the project. EPN and GF wrote the manuscript and assembled the figures with input from other authors.

Declaration of interests

The authors declare no competing interests.

Methods

Plasmid Construction

Plasmids were assembled using Gibson assembly (SBI MC010B-1) or restriction-ligation (NEB Quick Ligase M2200). The list of plasmids generated in this study can be found in the supplementary table and their annotated sequence maps as supplementary information.

Culturing mouse embryonic stem cells (ESCs)

ESCs were cultured in DMEM+Glutamax (ThermoFisher 10569044) supplemented with 15% Fetal Bovine Serum (ThermoFisher 10437028), 550μM b-mercaptoethanol (ThermoFisher 21985-023), 1X non-essential amino-acids (ThermoFisher 11140050), 10^4U of Leukemia inhibitory factor (Millipore ESG1107), PD0325901 (APExBIO A3013), CHIR-99021 (APExBIO A3011). Cells were maintained at a density of 0.2-1.5x10^5 cells/cm2 by passaging using TrypLE (ThermoFisher 12605010) every 24-48h on 0.1% gelatin-coated dishes at 37°C and 5% CO_2. 0.1% gelatin was prepared in 1X PBS using Gelatin powder (Sigma Aldrich G1890). Medium was changed daily when cells were not passaged. Cells were checked for mycoplasma infection and tested negative.

Degron inductions

Degradation of proteins tagged with the Auxin-Inducible Degron (44 amino-acid AID* truncation with residues 71–114 (Nora et al., 2017)) was induced by adding the auxin analog Indole-3-acetic acid sodium (IAA, Sigma Cat #I5148-2G) to a final concentration of 500μM (from a 1000x filter-sterilized stock diluted in water). Degradation of proteins tagged with the FKBP degron (FKBP12^{F36V} (Nabet et al., 2018)) was induced by adding dTAG-13 (abbreviated dTAG throughout the manuscript - Sigma Cat, SML2601) to a final concentration of 500nM (from a 1000x stock diluted in DMSO). We noticed equivalent degradation efficiency in ESCs compared to using the dTAG-v1 inducer at 500nM.

Genome editing of ESCs

For transfection, plasmids were prepared using the Nucleobond Midi kit (Macherey Nagel 740410.5) followed by isopropanol precipitation. Constructs were not linearized. Transfections were typically done using the Neon system (Thermo Fisher) with a 100 μL tip and 1 million cells at 1400 V, 10 ms, 3 pulses. To create knock-in cells, ESCs were co-transfected with 5 μg of Cas9-sgRNA expressing vector and 10-15 μg of targeting construct.

sgRNAs were cloned either into a Cas9 vector (pX330, Addgene #42230) or Cas9-2A-puro vector (pEN243, identical to pX459 Addgene 62988). Homology arms of targeting vectors ranged from 800 base pairs (bp) - 1 kilobase (kb) in length.

After electroporation, cells were seeded in a 9 cm^2 well and left to recover for 48 hours. Cells were plated at limited dilution and grown for around 8 days changing medium every other day until single colonies could be picked. Individual colonies were genotyped by polymerase chain reaction (PCR) and validated with Sanger sequencing. Homozygous or heterozygous clones were identified, expanded, validated by western blot, and cryopreserved. For serial editing, resulting validated clonal lines were used to repeat the transfection process with another targeting construct.

When the targeting construct contained an FRT flanked resistance cassette, we adopted the following strategy to avoid having to subclone the cells again after removal of the resistance cassette: after transfection of the targeting construct containing the FRT flanked resistance cassette, we added antibiotic in the medium 48h after transfection and kept the cells under selection for at least 5 days. Final antibiotic concentrations: Puromycin: 1 µg/mL; Blasticidin: 10 µg/mL; Hygromycin: 200 µg/mL; Geneticin: 200 µg/mL. We then transfected this heterogenous pool with pCAGGS-FlpO using the Neon electroporation system with a 10 µL tip and 0.1 million cells at 1400 V, 10 ms, 3 pulses, using 0.5µg DNA. Cells were seeded in a 9 cm^2 well and subcloned without antibiotic after 48h. PCR genotyping and antibiotic resistance tests identified homozygous knockin clones with the FRT-flanked selection cassette removed.

The list of cell lines, genotypes, order of transfection and corresponding vectors is provided as a supplementary table.

Western blotting

ESCs were dissociated using TrypLE, resuspended in culture medium, pelleted at 100g for 5 minutes at urrp # whp shudwxh. Cell pellets were washed in PBS, pelleted again, snap frozen in dry ice and stored in −70°C freezer. In total, 10-20 million cells were used to prepare nuclear extracts. Cell pellets stored at −70°C were taken out on ice, resuspended in 10 mM HEPES (pH 7.9), 2.5 mM MgCl$_2$, 0.25 M sucrose, 0.1% NP40, 1 mM DTT, and 1X HALT protease inhibitors (ThermoFisher 78437) and kept on ice for 10 minutes to allow swelling. After centrifugation at 500g at 4°C for 5 minutes, the supernatant was discarded and the pellets were resuspended on ice in 25 mM HEPES (pH 7.9), 1.5 mM MgCl$_2$, 700 mM NaCl, 0.5 mM DTT, 0.1 mM EDTA, 20% glycerol, 1 mM DTT, and 250 U benzonase (Sigma-Aldrich E1014), and incubated on ice for 30-60 minutes. The nuclear lysates were centrifuged at 18,000g, 4°C, 10 minutes, to eliminate the cellular debris. Protein concentration from the supernatants was measured using Pierce BCA Kit (ThermoFisher 23225).

Equal amounts of protein lysates were heated in 1X Laemmli buffer (Biorad 1610747) at 95°C for 10 minutes and subjected to electrophoresis on a 4-15% polyacrylamide gel (Biorad 17000927). For NIPBL westerns, 60 µg protein was loaded. The separated proteins were transferred onto a polyvinylidene difluoride (PVDF) membrane (Millipore IPFL00010) using phosphate-based transfer buffer (10 mM sodium phosphate monobasic, 10 mM sodium phosphate dibasic) at 4°C, 600 mA, 2 hours. After the completion of transfer, membranes were blocked in the 5% blocking buffer (5% skimmed milk prepared in 1X TBS-T) and incubated overnight at 4°C with the primary antibodies prepared in the 5% blocking buffer. Next day, the membranes were washed four times with 1X TBS-T buffer (20 mM Tris-Cl buffer-pH 7.4, 500 mM NaCl and 0.1% Tween 20) and incubated with the appropriate secondary antibodies conjugated with horseradish peroxidase (HRP) for an hour at RT (typically 1:5000). Following this, the membranes were again washed four times with 1X TBS-T buffer. The blots were developed using Pierce ECL western blotting substrate (ThermoScientific 32209) and detected using Chemidoc XRS+ imager (Biorad).

For validating homozygous knockin and degron depletion efficiency we used the following antibodies:
PDS5A: Bethyl Labs A300-089A 1:1000

687 PDS5B: Bethyl Labs IHC-00381 1:1000
688 TBP (loading control): Cell Signaling technology 8515S 1:2000
689 RAD21: Millipore 05-908 1:1000
690 NIPBL: sc-374625 1:1000
691 MAU2: Abcam ab183033 1:1000
692
693
Flow cytometry to measure total and chromatin bound RAD21
ESCs were seeded at a density of 0.5 x 10^6 per well of a gelatinized 6-well plate. Next day, the degron containing
ESC lines were treated with either IAA or dTAG-13 or both for 6 hours. At the end of 6 hours, media was aspirated
out and replaced with medium containing 100nM or 500nM of Halo-ligand JFX549 or JFX646 (kind gift of Luke
Lavis, HMMI Janelia Research Campus) and specified concentration of degron inducers (IAA and/or dTAG-13).
The cells were incubated at 37°C, 5% CO_2 for 30 minutes to allow JFX549 to bind the Halo-tag on RAD21. After
30 minutes, media was aspirated, cells were washed twice with 1X PBS and fresh media containing the specified
concentration of degron inducers (IAA and/or dTAG-13) was added to the well. Cells were incubated at 37°C,
5% CO_2 for 30 minutes to allow for unbound JFX549 or JFX646 to diffuse out of the cells. Following 30 minutes
incubation, cells were rinsed with 1X PBS, dissociated using TrypLE, resuspended in fresh culture medium
containing appropriate amounts of IAA or dTAG-13 and centrifuged at 100g for 5 minutes at RT. Supernatant
was aspirated and the cell pellets were resuspended in 1 ml of 1X PBS. Resuspended cells were equally divided
into two tubes to measure total RAD21 and bound RAD21 and spun at 100g for 5 minutes at RT. Supernatant
was discarded.
708
For measuring total RAD21, cell pellets were resuspended in 100ul of 4% formaldehyde (prepared in 1X PBS,
diluted from 16% methanol-free formaldehyde - ThermoFisher 28908) and incubated at RT for 15 minutes. After
incubation, 1 ml of ice-cold 1% BSA (diluted from 3% BSA in 1X PBS) was added and cells were spun at 300g
for 5 minutes. Supernatant was discarded. Cell pellets were resuspended in 100ul of ice-cold CSK-T buffer
(25mM HEPES pH7.4, 150mM NaCl, 1mM EDTA, 3mM MgCl2, 300mM Sucrose, 0.5% TritonX-100) on ice and
incubated on ice for 10 minutes. After incubation, 1 ml of ice-cold 1% BSA (diluted from 3% BSA in 1X PBS) was
added and cells were spun at 300g for 5 minutes. Supernatant was discarded. Cell pellets were resuspended in
300ul of 1mg/ml DAPI solution prepared in 1X PBS. Incubated for 5-10 minutes and acquired using BD Versa
flow cytometer using PE-A (for JFX549), APC-A (for JFX646) and V450 (for DAPI) channels. 100,000 events
were acquired.
719
For measuring chromatin-bound RAD21, cell pellets were resuspended in 100ul of ice-cold CSK-T buffer (25mM
HEPES pH7.4, 150mM NaCl, 1mM EDTA, 3mM MgCl2, 300mM Sucrose, 0.5% TritonX-100) on ice and
incubated on ice for 10 minutes. After incubation, 1 ml of ice-cold 1% BSA (diluted from 3% BSA in 1X PBS) was
added and cells were spun at 300g for 5 minutes. Supernatant was discarded, pellets were resuspended in 100ul
of 4% formaldehyde (prepared in 1X PBS, diluted from 16% methanol-free formaldehyde) and incubated at RT
for 15 minutes. After incubation, 1 ml of ice-cold 1% BSA (diluted from 3% BSA in 1X PBS) was added and cells
were spun at 300g for 5 minutes. Supernatant was discarded and pellets were resuspended in 300ul of 1mg/ml
DAPI solution prepared in 1X PBS . Incubated for 5-10 minutes and acquired using BD Versa flow cytometer
using PE-A (for JFX549) and V450 (for DAPI) channels. 100,000 events were acquired.

For complete depletion of PDS5A/B or WAPL, respective degron lines were treated with 500 µM of IAA for 6
hours. For complete depletion of NIPBL, the NIPBL-FKBP degron line was treated with 500nM of dTAG-13 for 6
hours. For co-depletion of PDS5A/B and NIPBL, WAPL and NIPBL, respective degron lines were treated with
500 µMof IAA and 500nM of dTAG-13 for 6 hours. To achieve partial RAD21 depletion in the background of
complete PDS5A/B-depletion, cells were treated with 500 µMof IAA (to completely deplete PDS5A/B) and 10nM
or 25nM dTAG-13 (to partially deplete RAD21) for 6 hours. To achieve partial RAD21 depletion in the background
of complete WAPL-depletion, cells were treated with 500 µMof IAA (to completely deplete WAPL) and 8nM or
25nM dTAG-13 (to partially deplete RAD21) for 6 hours. To achieve partial depletion of RAD21, cells were treated
with 20nM or 45nM of dTAG-13 for 6 hours.

ESC line devoid of any degron-tag and without Halo-tag on RAD21 served as a negative control. ESC line without
any degron-tag and with Halo-tag on RAD21 served as a positive control (no degron). The median fluorescence
for PE-A channel from the single cell gated for total RAD21 and bound RAD21 was subtracted from the
respective values coming from the cells devoid of Halo-tag. This eliminated any non-specific background from
the Halo ligand. Mean fluorescence for RAD21 was plotted for both total RAD21 and bound RAD21 for No degron
(RAD21^{Halo}) and different degron lines.

Percent bound RAD21 was calculated using following formula, taking the mean fluorescence across each cell
population:

Step 1: For all samples, remove aspecific signal by subtracting fluorescence levels of Halo-labeled untagged
cells (not expressing the halotag) from the fluorescence levels of Halo-labeled RAD21-Halotag cells

Step 2: Calculate % Bound RAD21 as follows:
(i) %Bound RAD21 = (RAD21 after pre-extraction)*100/(Total RAD21)
(ii) Percent bound RAD21 relative to No degron was derived as follows:
%Bound RAD21 relative to No degron = (Bound RAD21 in degron line)*100/(Bound RAD21 in No degron)

Cell cycle analysis by EdU incorporation and DAPI staining
The Click-iT Plus EdU Alexa Fluor 647 Flow Cytometry Assay Kit (Thermo Fisher Scientific C10634) was used
following the manufacturer's instructions. We used 20 µM EdU, pulsed for 1 hour and 30 minutes, and proceeded
with 1 million cells, incorporating DAPI after EdU staining.

Mitotic synchronization and release, cell cycle analysis by DAPI and RAD21 vermicelli microscopy

Three T75 flasks per genotype were incubated overnight (9h) with 100 ng/mL Nocodazole (from a 1/40 dilution
followed by a 1/500 dilution from a 2mg/ml stock, Sigma-Alrdich M1404). IAA (auxin analog) was then added at
500 µM final to the culture medium to induce WAPL or PDS5AB depletion for 3h. Mitotic shake-off was then

performed by firmly tapping the flaks, and the supernatant culture medium containing the dislodged mitotic cells was collected, pooling the three T75 flasks in a 50mL conical tube. Cells were spun, resuspended in 25mL 1XPBS containing IAA, spun again, resuspended in 25mL 1XPBS containing IAA again counted, spun once more and resuspended at 1 million cells per mL in ES culture medium with nocodazole but with 500 µM IAA. This procedure yielded around 10M cells across the three T75 flasks. The '0h release' timepoint was immediately collected by aliquoting 1M cells. For the other timepoints 1M cells were seeded in a gelatinized tissue culture 9 cm^2 well for later hourly collection, in ESC medium with 500 µM IAA. For the first two hours cells typically only loosely attached and could be singularized by pipetting. For later timepoints cells were detached and singularized as during regular passaging using TrypIE. For each time-point we collected three fractions. The synchronized/released cell pellet was resuspended in 1XPBS with 500 µM IAA and split in two fractions:

- Microscopy fraction (to quantify RAD21 vermicelli scores): 0.1M to 0.2M cells were seeded in a well of a glass-bottom 96-well plate (Greiner 655892). To ensure cells adhere the glass had been pre-coated with Poly-L Lysine. To coat the glass, we incubated 50 µL per well of 0.01 % (w/v) Poly-L-lysine solution (P8920-100ML) diluted in water for 10 minutes at room temperature and dried for at least 30min (up to days). After 10 minutes of incubating the cells onto the coated glass excess volume was removed by pipetting (adhered cells stayed on the coated glass a monolayer of single-cells) and cells were incubated with 100uL of 3% formaldehyde diluted in 1XPBS (from a 37% solution Electron Microscopy Sciences 15686). After 10 minutes wells were rinsed twice with PBS and left in 100uL 1xPBS until all the timepoints were collected. Fixed cells were kept at least overnight at 4°C once all timepoints were collected. For Halotag ligand reaction and H3S10Ph immunostaining, cells were first permeabilized by incubating in 0.5% Triton in 1xPBS for 5 minutes, and rinsed three times with 1xPBS. Then cells were blocked by incubating with 1%BSA in 1xPBS 15 minutes. Incubation was changed for a 1/5000 dilution of anti-H3S10Ph antibody (Millipore 06-570) in 1%BSA 1xPBS for 30min, followed by three 5-minute washes in 1xPBS. Cells were then incubated with a 1/10000 of goat anti-rabbit Alexa-647 (ThermoFisher 21245) and 100nM JFX549 for 30 minutes at room temperature. After three 5-minute washes in 1xPBS, cells were incubated with 1µg/mL DAPI in 1xPBS 10 minutes, rinsed twice with 1xPBS and left in 1xPBS until imaging, within 24h. Imaging was performed on a spinning disk microscope (Zeiss) with a 60x 1.4NA oil objective equipped with 405nm, 488nm, 561nm, 640nm excitation lasers at the Histology and Light Microscopy Core of the Gladstone Institutes. Cells were imaged in 3D using a step size of 0.3µm.

- Flow cytometry fraction (to assess overall synchronization and release efficiency): 400 µL of 1XPBS was added to a 100µL aliquot of cells and transferred into a FACS tube. 70% ethanol was then added dropwise while gently vortexing, and tubes were stored at least overnight at -20. Flow-cytometry fractions for each time-point were transferred to a 1.5mL tube, spun, resuspended in 1mL 1XPBS, spun again, resuspended in 1mL 1XPBS with 0.5% Triton X-100 1µg/mL DAPI and incubated at room temperature 30min. Cells were then processed on a BD Versa flow cytometer using V450 (for DAPI) channels.

RAD21 vermicelli microscopy from asynchronous ESCs

For RAD21-Halotag imaging in asynchronous cells, adherent cultures were obtained by seeding 100,000 ESCs per well of a laminin-coated glass-bottom 8-well slides (Ibidi 80807) for overnight growth, adding IAA for the

time indicated in figure panels. To coat the slides, laminin (L2020-1MG) was diluted 1:100 in 1XPBS (10-
20μg/ml working solution), and 100-150μL was added per well for at least 2h up to overnight at 37°C (in tissue
culture incubator). Laminin was then aspirated from the well prior to dispensing cells, without rinsing. We found
laminin coating minimized the size of the ESC colonies, helping with high-magnification imaging.

When ready for staining, adherent cells were washed with 150μL 1XPBS per well, followed by 3%
formaldehyde diluted in 1XPBS (from a 37% solution Electron Microscopy Sciences 15686) for 10 minutes.
Cells were then rinsed three times with 150μL 1XPBS, permeabilized by incubating in 0.5% Triton in 1xPBS for
5 minutes, and rinsed three times with 1xPBS. Cells were incubated with 100nM JFX549 1μg/mL DAPI in
1xPBS for 30 minutes, washed twice with 1xPBS for 10 minutes and left in 1xPBS until imaging, within 24h. As
most asynchronous ESCs are in S/G2 phases H3S10Ph staining was used in this analysis.

Slides were imaged on a spinning disk microscope (Nikon) with a 60x 1.4NA oil objective equipped with
405nm, 491nm, 561nm, 640nm excitation lasers. at the USCF Center for Advance Light Microscopy. Cells
were imaged in 3D using a step size of 0.3μm.

Quantification of RAD21 vermicelli scores from microscopy images
Proprietary file formats (.czi for Zeiss instrument, mitotic release timecourse; .nd2 for Nikon instrument for
asynchronous cells) were open in imageJ using Bio-Formats importer[110]. A series of custom scripts were used
first to extract and sort stacks. A mask for nuclei was then created by creating a Z projection and applying a
Gaussian Blur (sigma = 2) followed by a Mean filter (radius=5). Filtered nuclei were then manually thresholded
for each image and binarized. Touching nuclei were separated using Watershed processing (tolerance = 1.5)
and resulting particles were filtered for size (100-450) and circularity (0.6-1) the resulting collection of region or
interests was then used to analyze the other channels. The red channel (JFX549 staining RAD21-Halotag) was
quantified by extracting the mean and the standard deviation for each nucleus. For the analysis of synchronized
cells, the paired infra-red channel (H3S10Ph staining) was quantified by extracting the mean signal. After
processing all nuclei across several image stacks the RAD21-Halotag vermicelli score, or coefficient of variation
(StdDev/Mean) was calculated in R. For the analysis of synchronized cells, unreleased mitotic cells were
eliminated from the analysis by removing the nuclei with H3S10Ph by thresholding the infra-red signal in R.

Inverse Fluorescence Recovery after Photobleaching (iFRAP)
The cells were imaged on an inverted motorized stand Zeiss AxioObserver7 equipped with CSU-W1 Confocal
Scanner Unit (Yokogawa) with Dual T2, a 50um pinhole disk unit, 2 sCMOS cameras (Prime 95B,
photometrics),a Visitron VS-Homogenizer ,an ASI MS2000 X,Y, ZPiezo drive (300um travel range) and
controlled with VisiView (version 6.0.0.35) imaging software (Visitron Systems GmbH). Illumination was achieved
with 561 Cobolt Jive (Cobolt, 200mW). For FRAP, illumination was done using a 473 nm FRAP laser.
A plan Apochromat 40X/1.3 oil objective was used, resulting in a pixel size of 0.275 μm in x,y. A 405/488/561
dichroic was used. The emission filter in the scan head was a 575 nm long pass. Imaging was done in a
humidified incubation chamber at 37° C supplied with 8% CO_2.
A ROI was defined on the field of view and few cells (~10) were selected for FRAP. A rectangular ROI was
defined for each nuclei encompassing around 80% of the nuclei area at the focal plane. 2D timelapse images

were acquired every 30 s for 250 frames, further referred to as movies. Images were taken at 30% max laser power (~ 3.528mW) to limit bleaching. The focus was kept using a Zeiss definiteFocus 2 system between every frame. After 4 frames, the FRAP sequence was initiated on the selected nuclei at 100% laser power. For FRAP, a laser of 22 um was focused on the focal plane and was shined on each pixel for 300ms. The focusing of the FRAP laser to the imaging focal plane and calibration of the x,y direction of the laser was done before every acquisition using a dedicated well not used for the imaging. The calibration was done by moving the beam to specific locations on the field of view and adjusting its focus to obtain the smallest beam at the imaging focal plane.

The cells were imaged on a μ-Slide 8 Well glass bottom (Ibidi, 80807). Briefly, a day before imaging ~200.000 cells were split in Gibco™ FluoroBrite™ DMEM (Gibco™) onto the dish. On the day of the imaging 100nM of Halo-JF-549 ligand (Janelia Fluor® NHS ester Tocris lab, 6147) was added for 30 minutes to the imaging medium. Then cells were washed three times with 1X PBS followed by addition of fresh imaging media.

Image analysis

The movies were saved as ome.tf2 format in the VisiView software. The images were analyzed using a custom GUI python script. We selected 2 circular ROI for both bleached and unbleached regions and updated their position every 5 frames (2.5 minutes). Missing positions were linearly interpolated. The mean intensity in the unbleached (I_unbleached(t)) and bleached (I_bleached(t) regions at every frame are then extracted.

To account for photobleaching we defined for every movie 10 unbleached nuclei and segmented them using Otsu's intensity thresholding method. To account for cell motion nuclei are linked using maximum overlap as a linking score, nuclei tracks are then built using the most overlapping nuclei. The mean intensity over time (I_photobleaching(t)) is then extracted. Furthermore, to account for the background intensity, another circular ROI was defined in a place on the image without any cells and (I_background) was extracted and averaged over all frames.

For every movie I_photobleaching(t) - I_background is computed and normalized to 1 as the initial difference in order to obtain a photobleaching curve.

Raw I_unbleached(t) and I_bleached(t) are averaged over all the nuclei in a given movie and the difference I_unbleached(t)-I_bleached(t) is computed and normalized by the bleaching to create the iFRAP signal,

iFRAP = I_unbleached(t)-I_bleached(t)/I_photobleaching(t) -I_background

Individual nuclei traces were visually checked and traces corresponding to nuclei leaving the FOV were manually removed. To account for initial cohesin concentration and equilibration of the system after bleaching, the iFRAP signal was further normalized to its level at frame 6 (1.5 minutes after bleaching).

More detailed explanation and raw code is available on
https://nesslfy.github.io/cohesin_residence_time/index.html

RNA-seq

0.5 million ESCs were seeded in a 9 cm^2 tissue culture well (-/+ dTAG) and collected after 24 hours for RNA extraction. ESCs were dissociated using TrypLE, resuspended in culture medium, spun down at 100g for 5 minutes at RT. Supernatant was discarded and the cell pellets were washed once with 1X PBS. Cell pellets were

resuspended in 1 mL of Trizol (ThermoFisher 15596-018), incubated at RT for 5 minutes. 250 µL of chloroform was added, vortexed vigorously, incubated at RT for 10 minutes and spun at 12,000g for 5 minutes at 4°C. Upper aqueous layer was transferred to a fresh tube, mixed with an equal volume of isopropanol, vortexed briefly, incubated at RT for 10 minutes and centrifuged at 12,000g at 4°C for 15 minutes to pellet the RNA. Supernatant was discarded and the pellet was washed twice with 75% ethanol and allowed to air dry for 5 minutes. RNA was resuspended in 50 µL of Nuclease-free water. RNA concentration was measured using Nanodrop and quality was assessed using Bioanalyzer (Total RNA Pico kit- Agilent 5067-1513).

1.5 µg of RNA was treated with the Turbo DNase kit (Invitrogen AM1907), incubated at 37°C for 30 minutes. DNase was inactivated using 2 µL of inactivation reagent by incubating at RT for 5 minutes followed by centrifugation at 13500 rpm for 90 seconds at RT. Supernatant was transferred to a fresh tube. 1000 ng of total RNA was used as input for poly(A) capture and library preparation using the Illumina Stranded mRNA kit (Illumina 20040532) per manufacturer's instructions. Quality and quantity of libraries were assessed using QuBit and Bioanalyzer (High sensitivity DNA Chip - Agilent 5067-4626). Libraries were sequenced on HiSeq 4000 using 100 bp paired end or NovaSeq 6000 using 150 bp paired end.

Analysis of RNA-seq data
Reads were trimmed to 100bp paired-end. Alignment of raw RNA-seq data was produced using STAR version 2.7.7a with default parameters and basic two-pass mapping in mm10. GENCODE VM23 (GRCm38.p6) was used as the reference gene set. Size factor normalization and differential expression analyses were done using DESeq2 (version 1.36.0) after filtering out genes with less than 10 raw reads. Principal Component Analysis was conducted in DESeq2 on auxin/dTAG-treated samples with the three replicates separate. Replicates were then collapsed by averaging their PC1 coordinates after PCA, resulting in a single point per condition.

Hi-C
Hi-C was performed using the kit from Arima Genomics as per manufacturer's instructions (202103-1577). Briefly, ESCs were dissociated using TrypLE, resuspended in culture medium and counted. 5-10 million cells were spun down at 100g for 5 minutes at RT. Supernatant was discarded, and the cell pellets were resuspended in 5 ml of 1X PBS. Cells were crosslinked using 2% formaldehyde (diluted from 37% formaldehyde - Electron Microscopy Sciences 15686), mixed by inverting the tubes 10 times and incubated at RT for 10 minutes. Quenching was done using the Stop solution 1 provided with the kit, mixed by inverting the tubes 10 times and incubated at RT for 5 minutes. Samples were then incubated on ice for 15 minutes and spun down at 100g for 5 minutes at 4°C. After discarding the supernatant, cells were resuspended in cold 1X PBS, counted, aliquoted at a density of $1x10^6$ cells/tube and spun down at 100g for 5 minutes at 4°C. Supernatant was discarded, cell pellets were snap frozen on dry ice and stored at -70°C.

Fixed cells were thawed and Hi-C was performed as per manufacturer's guidelines. Libraries were prepared using TruSeq DNA Nano LP kit from Illumina (20015964). Quality and quantity of libraries were assessed using Qubit and Bioanalyzer (High sensitivity DNA Chip- Agilent 5067-4626). Libraries were sequenced on NextSeq 500 using 75 paired end or NextSeq 2000 using 101 bp paired (and then trimmed to 75 before analysis).

Analysis of Hi-C data

Reads were trimmed to 75bp (if sequenced beyond that) using *cutadapt*(Martin, 2011). Data was mapped to mm10 using the *distiller-nf* pipeline (https://github.com/open2c/distiller-nf), iteratively corrected and converted into .cool format with *cooler*[111], only considering read pairs with both sides mapping with high confidence (mapq ≥ 30). After ascertaining the concordance of each replicate using either *cooltools* (Open2C et al., 2024a) or *Genova*(van der Weide et al., 2021) (P(s) curves and aggregate peak analysis), the independent Hi-C replicates (up to two) were merged for each condition. Multiresolution coolers (.mcool) were then generated with *cooler* and balanced using iterative correction. P(s) plots and Hi-C contact maps were generated using *cooltools* (Open2C et al., 2024a). Compartment scores were obtained from the saddle plots using *cooltools* and were normalized to 1 in the limit of full A/B segregation, with a score of 0 denoting ideal A/B mixing. For computations of distance-dependent compartment scores (Fig. S7D), experimental observed-over-expected maps were aggregated across all chromosomes in which compartment patterns were found to visually align with the first Hi-C eigenvector (ev1) (chromosomes 1, 2, 4, 5, 6, 8, 9, 11, and 13-19). Hi-C counts were binned against ev1 and partitioned as a function of distance from the main diagonal in 500 kb increments in order to create ensembles of distance-dependent saddle plots, from which compartment scores were similarly calculated.

Polymer model of ESC chromatin with loop extrusion and compartmentalization

We modeled chr2:66,000,000-116,000,000 using Polychrom-HooMD (https://github.com/open2c/polychrom-hoomd). As previously[64], we modeled chromatin as a fiber with monomers of individual diameter 50 nm, each representing 2.5 kb of DNA at a density of 0.2 monomers per unit volume, amounting to a physiological DNA concentration of $0.01\ bp/nm^3$. Molecular dynamics (MD) simulations were conducted using periodic boundary conditions, and were initiated from a compact conformation which was allowed to expand until the radius of gyration equilibrated over $O(10^7)$ MD steps. Based on the comparison of the computed single-locus mean-squared displacements with in vivo measurements in yeast and mammals[86], each MD step was estimated to amount to 5 ms of physical time. Genomic positions of A and B compartments were determined in a standard fashion from the first principal eigenvector of the experimental Hi-C correlation matrix obtained in WT embryos at 25 kb resolution, and computed using *cooltools*. While this resolution was sufficient to reproduce the main features of the experimental maps, coarse graining compartment positions to 25 kb contributes to the greater dynamic range of simulated compartment scores. A self-affinity interaction of magnitude 0.05 kT was applied between B-type loci, which was found by iterative optimization to yield a maximum Spearman rank correlation coefficient between the simulated and experimental observed-over-expected maps in wild-type conditions (Fig. S2D). To optimize statistical sampling, as well as mitigate potential finite-size effects, each simulation was set up to include 10 identical replicates of the chromosomal region of interest, thus amounting to a total system size of 200,000 monomers (or, equivalently, 500 Mb).

As previously[64], we coupled this 3D bead-spring heteropolymer model with a 1D lattice model of loop extruder dynamics. Extruder leg positions from the 1D lattice at each timestep were used to specify positions of additional harmonic bonds between monomers in the 3D simulation, with identical stepping dynamics in A and B compartments. As originally implemented for cohesin extrusion and as recently hypothesized for condensin-

condensin encounters[87], we assumed that extruders collided without a chance to bypass each-other. Under wild-type conditions, an extruder separation of 185 kb was used to match the loaded cohesin abundance reported in ESCs[67]. Similarly, the mean extruder residence time on chromatin was set to 1320 s based on FRAP measurements of cohesin dynamics *in vivo* [58]. During each lattice update in a wild-type 1D simulation, 4000 MD steps were executed. This value was found from a systematic sweep to yield optimal agreement with the experimental contact frequency vs. distance curve in untagged cells, and amounts to an average extrusion rate of 0.25kb/s.

Simulated Hi-C and contact-vs-distance profiles

In silico contact maps were generated from MD simulation trajectories using the monomerResolutionContactMapSubchains function from the contact_maps module as implemented in the polykit library, using the default capture radius (2.3 monomers, 115 nm). Maps were averaged across 10^4 equilibrated conformations obtained from 10 independent runs. We then determined contact frequencies as a function of genomic distance using the expected_cis function of *cooltools*. For comparisons with experimental data, the obtained *in silico* P(s) curves were linearly interpolated using the interp function of the numpy library to synchronize bin positions to those used for experimental data. *In silico* curves were then rescaled such that the value at 50 kb matches that measured in the *in vivo* sample. Accordingly, the goodness-of-fit parameter (R^2) was defined as

$$R^2 = 1 - \frac{\sum_s \ [P_{exp}(s) - P(s)]^2}{\sum_s \ [P_{exp}(s) - \underline{P}_{exp}]^2},$$

where the summation and average ($\underline{P}_{exp}$) are both performed over the genomic range $[50\ kb, 5\ Mb]$. For visualization, derivatives of the contact frequency curve (P'(S)) were computed with the NGrsh¤>rIwd>w¤N>dGDIB function from srGTFDw¢srGTphu¤dIdGTNhN¢

In silico microscopy

Simulated microscopy images were obtained from individual chromosome conformations by binning computed cohesin spatial positions into discrete voxels of size 100 nm x 100 nm x 100 nm. Optical diffraction effects in the (x, y) plane were mimicked by running the resulting 3D raster through a Gaussian convolution filter of width 250 nm, following which a maximum intensity projection was performed along the z direction. To account for the presence of unloaded cohesins, additional unbound cohesins were placed randomly throughout the simulated volume, with a number determined from FACS estimates for the bound fraction. Vermicelli scores were subsequently defined as in experiments from the ratio of the standard deviation and the mean of the normalized cohesin fluorescence signal.

References

1. Dekker, J., and Mirny, L.A. (2024). The chromosome folding problem and how cells solve it. Cell *187*, 6424–6450. https://doi.org/10.1016/j.cell.2024.10.026.

2. Yatskevich, S., Rhodes, J., and Nasmyth, K. (2019). Organization of Chromosomal DNA by SMC Complexes. Annu. Rev. Genet. *53*, 445–482. https://doi.org/10.1146/annurev-genet-112618-043633.

3. Hoencamp, C., and Rowland, B.D. (2023). Genome control by SMC complexes. Nat. Rev. Mol. Cell Biol. *24*, 633–650. https://doi.org/10.1038/s41580-023-00609-8.

4. Fudenberg, G., Abdennur, N., Imakaev, M., Goloborodko, A., and Mirny, L.A. (2017). Emerging Evidence of Chromosome Folding by Loop Extrusion. Cold Spring Harb. Symp. Quant. Biol. *82*, 45–55. https://doi.org/10.1101/sqb.2017.82.034710.

5. Davidson, I.F., and Peters, J.-M. (2021). Genome folding through loop extrusion by SMC complexes. Nat. Rev. Mol. Cell Biol. *22*, 445–464. https://doi.org/10.1038/s41580-021-00349-7.

6. de Wit, E., and Nora, E.P. (2022). New insights into genome folding by loop extrusion from inducible degron technologies. Nat. Rev. Genet., 1–13. https://doi.org/10.1038/s41576-022-00530-4.

7. Vernì, F., Gandhi, R., Goldberg, M.L., and Gatti, M. (2000). Genetic and Molecular Analysis of wings apart-like (wapl), a Gene Controlling Heterochromatin Organization in Drosophila melanogaster. Genetics *154*, 1693–1710. https://doi.org/10.1093/genetics/154.4.1693.

8. Tedeschi, A., Wutz, G., Huet, S., Jaritz, M., Wuensche, A., Schirghuber, E., Davidson, I.F., Tang, W., Cisneros, D.A., Bhaskara, V., et al. (2013). Wapl is an essential regulator of chromatin structure and chromosome segregation. Nature *501*, 564–568. https://doi.org/10.1038/nature12471.

9. Hill, L., Ebert, A., Jaritz, M., Wutz, G., Nagasaka, K., Tagoh, H., Kostanova-Poliakova, D., Schindler, K., Sun, Q., Bönelt, P., et al. (2020). Wapl repression by Pax5 promotes V gene recombination by Igh loop extrusion. Nature *584*, 142–147. https://doi.org/10.1038/s41586-020-2454-y.

10. Dai, H.-Q., Hu, H., Lou, J., Ye, A.Y., Ba, Z., Zhang, X., Zhang, Y., Zhao, L., Yoon, H.S., Chapdelaine-Williams, A.M., et al. (2021). Loop extrusion mediates physiological Igh locus contraction for RAG scanning. Nature *590*, 338–343. https://doi.org/10.1038/s41586-020-03121-7.

11. Kiefer, L., Gaudin, S., Rajkumar, S.M., Servito, G.I.F., Langen, J., Mui, M.H., Nawsheen, S., and Canzio, D. (2024). Tuning cohesin trajectories enables differential readout of the Pcdhα cluster across neurons. Science *385*, eadm9802. https://doi.org/10.1126/science.adm9802.

12. Kiefer, L., Chiosso, A., Langen, J., Buckley, A., Gaudin, S., Rajkumar, S.M., Servito, G.I.F., Cha, E.S., Vijay, A., Yeung, A., et al. (2023). WAPL functions as a rheostat of Protocadherin isoform diversity that controls neural wiring. Science *380*, eadf8440. https://doi.org/10.1126/science.adf8440.

13. Wutz, G., Várnai, C., Nagasaka, K., Cisneros, D.A., Stocsits, R.R., Tang, W., Schoenfelder, S., Jessberger, G., Muhar, M., Hossain, M.J., et al. (2017). Topologically associating domains and chromatin loops depend on cohesin and are regulated by CTCF, WAPL, and PDS5 proteins. EMBO J. *36*, 3573–

3599. https://doi.org/10.15252/embj.201798004.

14. Denison, S.H., Kafer, E., and May, G.S. (1993). Mutation in the Bimd Gene of Aspergillus Nidulans Confers a Conditional Mitotic Block and Sensitivity to DNA Damaging Agents. Genetics *134*, 1085–1096.

15. Morales, C., Ruiz-Torres, M., Rodríguez-Acebes, S., Lafarga, V., Rodríguez-Corsino, M., Megías, D., Cisneros, D.A., Peters, J.-M., Méndez, J., and Losada, A. (2020). PDS5 proteins are required for proper cohesin dynamics and participate in replication fork protection. J. Biol. Chem. *295*, 146–157. https://doi.org/10.1074/jbc.RA119.011099.

16. Hartman, T., Stead, K., Koshland, D., and Guacci, V. (2000). Pds5p Is an Essential Chromosomal Protein Required for Both Sister Chromatid Cohesion and Condensation in Saccharomyces cerevisiae. J. Cell Biol. *151*, 613–626.

17. van Heemst, D., James, F., Pöggeler, S., Berteaux-Lecellier, V., and Zickler, D. (1999). Spo76p is a conserved chromosome morphogenesis protein that links the mitotic and meiotic programs. Cell *98*, 261–271. https://doi.org/10.1016/s0092-8674(00)81020-x.

18. Viera, A., Berenguer, I., Ruiz-Torres, M., Gómez, R., Guajardo, A., Barbero, J.L., Losada, A., and Suja, J.A. (2020). PDS5 proteins regulate the length of axial elements and telomere integrity during male mouse meiosis. EMBO Rep. *21*, e49273. https://doi.org/10.15252/embr.201949273.

19. Schwarzer, W., Abdennur, N., Goloborodko, A., Pekowska, A., Fudenberg, G., Loe-Mie, Y., Fonseca, N.A., Huber, W., H Haering, C., Mirny, L., et al. (2017). Two independent modes of chromatin organization revealed by cohesin removal. Nature *551*, 51–56. https://doi.org/10.1038/nature24281.

20. Alonso-Gil, D., Cuadrado, A., Giménez-Llorente, D., Rodríguez-Corsino, M., and Losada, A. (2023). Different NIPBL requirements of cohesin-STAG1 and cohesin-STAG2. Nat. Commun. *14*, 1326. https://doi.org/10.1038/s41467-023-36900-7.

21. Aboreden, N.G., Zhao, H., Shan, F., Liu, F., Zhang, H., and Blobel, G.A. (2025). Cis-regulatory chromatin contacts form de novo in the absence of loop extrusion. Preprint at bioRxiv, https://doi.org/10.1101/2025.01.12.632634 https://doi.org/10.1101/2025.01.12.632634.

22. Popay, T.M., Pant, A., Munting, F., Black, M.E., Haghani, N., and Dixon, J.R. (2024). Genome-wide in vivo dynamics of cohesin-mediated loop extrusion and its role in transcription activation. Preprint at bioRxiv, https://doi.org/10.1101/2024.10.02.616323 https://doi.org/10.1101/2024.10.02.616323.

23. Hansen, K.L., Adachi, A.S., Braccioli, L., Kadvani, S., Boileau, R.M., Pokorny, B., Shah, R., Anderson, E.C., Martinovic, M., Zhang, K., et al. (2024). Synergy between cis-regulatory elements can render cohesin dispensable for distal enhancer function. Preprint at bioRxiv, https://doi.org/10.1101/2024.10.04.615095 https://doi.org/10.1101/2024.10.04.615095.

24. Rollins, R.A., Morcillo, P., and Dorsett, D. (1999). Nipped-B, a Drosophila homologue of chromosomal adherins, participates in activation by remote enhancers in the cut and Ultrabithorax genes. Genetics *152*, 577–593. https://doi.org/10.1093/genetics/152.2.577.

25. Garcia, P., Fernandez-Hernandez, R., Cuadrado, A., Coca, I., Gomez, A., Maqueda, M., Latorre-Pellicer, A., Puisac, B., Ramos, F.J., Sandoval, J., et al. (2021). Disruption of NIPBL/Scc2 in Cornelia de Lange

Syndrome provokes cohesin genome-wide redistribution with an impact in the transcriptome. Nat. Commun. *12*, 4551. https://doi.org/10.1038/s41467-021-24808-z.

26. Tonkin, E.T., Wang, T.-J., Lisgo, S., Bamshad, M.J., and Strachan, T. (2004). NIPBL, encoding a homolog of fungal Scc2-type sister chromatid cohesion proteins and fly Nipped-B, is mutated in Cornelia de Lange syndrome. Nat. Genet. *36*, 636–641. https://doi.org/10.1038/ng1363.

27. Krantz, I.D., McCallum, J., DeScipio, C., Kaur, M., Gillis, L.A., Yaeger, D., Jukofsky, L., Wasserman, N., Bottani, A., Morris, C.A., et al. (2004). Cornelia de Lange syndrome is caused by mutations in NIPBL, the human homolog of Drosophila melanogaster Nipped-B. Nat. Genet. *36*, 631–635. https://doi.org/10.1038/ng1364.

28. Antony, J., Chin, C.V., and Horsfield, J.A. (2021). Cohesin Mutations in Cancer: Emerging Therapeutic Targets. Int. J. Mol. Sci. *22*, 6788. https://doi.org/10.3390/ijms22136788.

29. Piché, J., Van Vliet ,Patrick Piet, Pucéat ,Michel, and and Andelfinger, G. (2019). The expanding phenotypes of cohesinopathies: one ring to rule them all! Cell Cycle *18*, 2828–2848. https://doi.org/10.1080/15384101.2019.1658476.

30. Mannini, L., Cucco, F., Quarantotti, V., Krantz, I.D., and Musio, A. (2013). Mutation Spectrum and Genotype–Phenotype Correlation in Cornelia de Lange Syndrome. Hum. Mutat. *34*, 10.1002/humu.22430. https://doi.org/10.1002/humu.22430.

31. Ciosk, R., Shirayama, M., Shevchenko, A., Tanaka, T., Toth, A., Shevchenko, A., and Nasmyth, K. (2000). Cohesin's binding to chromosomes depends on a separate complex consisting of Scc2 and Scc4 proteins. Mol. Cell *5*, 243–254. https://doi.org/10.1016/s1097-2765(00)80420-7.

32. Gillespie, P.J., and Hirano, T. (2004). Scc2 couples replication licensing to sister chromatid cohesion in Xenopus egg extracts. Curr. Biol. CB *14*, 1598–1603. https://doi.org/10.1016/j.cub.2004.07.053.

33. Takahashi, T.S., Yiu, P., Chou, M.F., Gygi, S., and Walter, J.C. (2004). Recruitment of Xenopus Scc2 and cohesin to chromatin requires the pre-replication complex. Nat. Cell Biol. *6*, 991–996. https://doi.org/10.1038/ncb1177.

34. Davidson, I.F., Bauer, B., Goetz, D., Tang, W., Wutz, G., and Peters, J.-M. (2019). DNA loop extrusion by human cohesin. Science *366*, 1338–1345. https://doi.org/10.1126/science.aaz3418.

35. Kim, Y., Shi, Z., Zhang, H., Finkelstein, I.J., and Yu, H. (2019). Human cohesin compacts DNA by loop extrusion. Science *366*, 1345–1349. https://doi.org/10.1126/science.aaz4475.

36. Rhodes, J., Mazza, D., Nasmyth, K., and Uphoff, S. (2017). Scc2/Nipbl hops between chromosomal cohesin rings after loading. eLife *6*, e30000. https://doi.org/10.7554/eLife.30000.

37. Alonso-Gil, D., and Losada, A. (2023). NIPBL and cohesin: new take on a classic tale. Trends Cell Biol. *33*, 860–871. https://doi.org/10.1016/j.tcb.2023.03.006.

38. Haarhuis, J.H.I., van der Weide, R.H., Blomen, V.A., Yáñez-Cuna, J.O., Amendola, M., van Ruiten, M.S., Krijger, P.H.L., Teunissen, H., Medema, R.H., van Steensel, B., et al. (2017). The Cohesin Release Factor WAPL Restricts Chromatin Loop Extension. Cell *169*, 693-707.e14.

1118   https://doi.org/10.1016/j.cell.2017.04.013.

39. Chan, K.-L., Roig, M.B., Hu, B., Beckouët, F., Metson, J., and Nasmyth, K. (2012). Cohesin's DNA exit gate is distinct from its entrance gate and is regulated by acetylation. Cell *150*, 961–974. https://doi.org/10.1016/j.cell.2012.07.028.

40. Kikuchi, S., Borek, D.M., Otwinowski, Z., Tomchick, D.R., and Yu, H. (2016). Crystal structure of the cohesin loader Scc2 and insight into cohesinopathy. Proc. Natl. Acad. Sci. *113*, 12444–12449. https://doi.org/10.1073/pnas.1611333113.

41. Petela, N.J., Gligoris, T.G., Metson, J., Lee, B.-G., Voulgaris, M., Hu, B., Kikuchi, S., Chapard, C., Chen, W., Rajendra, E., et al. (2018). Scc2 Is a Potent Activator of Cohesin's ATPase that Promotes Loading by Binding Scc1 without Pds5. Mol. Cell *70*, 1134-1148.e7. https://doi.org/10.1016/j.molcel.2018.05.022.

42. van Ruiten, M.S., van Gent, D., Sedeño Cacciatore, Á., Fauster, A., Willems, L., Hekkelman, M.L., Hoekman, L., Altelaar, M., Haarhuis, J.H.I., Brummelkamp, T.R., et al. (2022). The cohesin acetylation cycle controls chromatin loop length through a PDS5A brake mechanism. Nat. Struct. Mol. Biol. *29*, 586–591. https://doi.org/10.1038/s41594-022-00773-z.

43. Bsteh, D., Moussa, H.F., Michlits, G., Yelagandula, R., Wang, J., Elling, U., and Bell, O. (2023). Loss of cohesin regulator PDS5A reveals repressive role of Polycomb loops. Nat. Commun. *14*, 8160. https://doi.org/10.1038/s41467-023-43869-w.

44. Yu, D., Chen, G., Wang, Y., Wang, Y., Lin, R., Liu, N., Zhu, P., Liu, H., Hu, T., Feng, R., et al. (2022). Regulation of cohesin-mediated chromosome folding by PDS5 in mammals. EMBO Rep. *23*, e54853. https://doi.org/10.15252/embr.202254853.

45. Bastié, N., Chapard, C., Dauban, L., Gadal, O., Beckouët, F., and Koszul, R. (2022). Smc3 acetylation, Pds5 and Scc2 control the translocase activity that establishes cohesin-dependent chromatin loops. Nat. Struct. Mol. Biol. *29*, 575–585. https://doi.org/10.1038/s41594-022-00780-0.

46. Costantino, L., Hsieh, T.-H.S., Lamothe, R., Darzacq, X., and Koshland, D. (2020). Cohesin residency determines chromatin loop patterns. eLife *9*, e59889. https://doi.org/10.7554/eLife.59889.

47. Göbel, A.-M., Zhou, S., Wang, Z., Tzourtzou, S., Himmelbach, A., Zheng, S., Pradillo, M., Liu, C., and Jiang, H. (2024). Mutations of PDS5 genes enhance TAD-like domain formation Arabidopsis thaliana. Nat. Commun. *15*, 9308. https://doi.org/10.1038/s41467-024-53760-x.

48. Dauban, L., Montagne, R., Thierry, A., Lazar-Stefanita, L., Bastié, N., Gadal, O., Cournac, A., Koszul, R., and Beckouët, F. (2020). Regulation of Cohesin-Mediated Chromosome Folding by Eco1 and Other Partners. Mol. Cell *77*, 1279-1293.e4. https://doi.org/10.1016/j.molcel.2020.01.019.

49. Luppino, J.M., Field, A., Nguyen, S.C., Park, D.S., Shah, P.P., Abdill, R.J., Lan, Y., Yunker, R., Jain, R., Adelman, K., et al. (2022). Co-depletion of NIPBL and WAPL balance cohesin activity to correct gene misexpression. PLoS Genet. *18*, e1010528. https://doi.org/10.1371/journal.pgen.1010528.

50. Murayama, Y., and Uhlmann, F. (2015). DNA Entry into and Exit out of the Cohesin Ring by an Interlocking Gate Mechanism. Cell *163*, 1628–1640. https://doi.org/10.1016/j.cell.2015.11.030.

51. Nabet, B., Roberts, J.M., Buckley, D.L., Paulk, J., Dastjerdi, S., Yang, A., Leggett, A.L., Erb, M.A., Lawlor, M.A., Souza, A., et al. (2018). The dTAG system for immediate and target-specific protein degradation. Nat. Chem. Biol. *14*, 431–441. https://doi.org/10.1038/s41589-018-0021-8.

52. Liu, N.Q., Maresca, M., van den Brand, T., Braccioli, L., Schijns, M.M.G.A., Teunissen, H., Bruneau, B.G., Nora, E.P., and de Wit, E. (2021). WAPL maintains a cohesin loading cycle to preserve cell-type-specific distal gene regulation. Nat. Genet. *53*, 100–109. https://doi.org/10.1038/s41588-020-00744-4.

53. Hafner, A., Park, M., Berger, S.E., Murphy, S.E., Nora, E.P., and Boettiger, A.N. (2023). Loop stacking organizes genome folding from TADs to chromosomes. Mol. Cell *83*, 1377-1392.e6. https://doi.org/10.1016/j.molcel.2023.04.008.

54. Szabo, Q., Donjon, A., Jerković, I., Papadopoulos, G.L., Cheutin, T., Bonev, B., Nora, E.P., Bruneau, B.G., Bantignies, F., and Cavalli, G. (2020). Regulation of single-cell genome organization into TADs and chromatin nanodomains. Nat. Genet. *52*, 1151–1157. https://doi.org/10.1038/s41588-020-00716-8.

55. Nishimura, K., Fukagawa, T., Takisawa, H., Kakimoto, T., and Kanemaki, M. (2009). An auxin-based degron system for the rapid depletion of proteins in nonplant cells. Nat. Methods *6*, 917–922. https://doi.org/10.1038/nmeth.1401.

56. Mitter, M., Gasser, C., Takacs, Z., Langer, C.C.H., Tang, W., Jessberger, G., Beales, C.T., Neuner, E., Ameres, S.L., Peters, J.-M., et al. (2020). Conformation of sister chromatids in the replicated human genome. Nature *586*, 139–144. https://doi.org/10.1038/s41586-020-2744-4.

57. Forment, J.V., and Jackson, S.P. (2015). A flow cytometry–based method to simplify the analysis and quantification of protein association to chromatin in mammalian cells. Nat. Protoc. *10*, 1297–1307. https://doi.org/10.1038/nprot.2015.066.

58. Hansen, A.S., Pustova, I., Cattoglio, C., Tjian, R., and Darzacq, X. (2017). CTCF and cohesin regulate chromatin loop stability with distinct dynamics. eLife *6*, e25776. https://doi.org/10.7554/eLife.25776.

59. Holzmann, J., Politi, A.Z., Nagasaka, K., Hantsche-Grininger, M., Walther, N., Koch, B., Fuchs, J., Dürnberger, G., Tang, W., Ladurner, R., et al. (2019). Absolute quantification of cohesin, CTCF and their regulators in human cells. eLife *8*, e46269. https://doi.org/10.7554/eLife.46269.

60. Cuadrado, A., Giménez-Llorente, D., De Koninck, M., Ruiz-Torres, M., Kojic, A., Rodríguez-Corsino, M., and Losada, A. (2022). Contribution of variant subunits and associated factors to genome-wide distribution and dynamics of cohesin. Epigenetics Chromatin *15*, 37. https://doi.org/10.1186/s13072-022-00469-0.

61. Gerlich, D., Koch, B., Dupeux, F., Peters, J.-M., and Ellenberg, J. (2006). Live-Cell Imaging Reveals a Stable Cohesin-Chromatin Interaction after but Not before DNA Replication. Curr. Biol. *16*, 1571–1578. https://doi.org/10.1016/j.cub.2006.06.068.

62. Barth, R., Davidson, I.F., Torre, J. van der, Taschner, M., Gruber, S., Peters, J.-M., and Dekker, C. (2025). SMC motor proteins extrude DNA asymmetrically and can switch directions. Cell *188*, 749-763.e21. https://doi.org/10.1016/j.cell.2024.12.020.

63. Fudenberg, G., Imakaev, M., Lu, C., Goloborodko, A., Abdennur, N., and Mirny, L.A. (2016). Formation of

Chromosomal Domains by Loop Extrusion. Cell Rep. *15*, 2038–2049.
https://doi.org/10.1016/j.celrep.2016.04.085.

64. Nuebler, J., Fudenberg, G., Imakaev, M., Abdennur, N., and Mirny, L.A. (2018). Chromatin organization by an interplay of loop extrusion and compartmental segregation. Proc. Natl. Acad. Sci. U. S. A. *115*, E6697–E6706. https://doi.org/10.1073/pnas.1717730115.

65. Goloborodko, A., Imakaev, M.V., Marko, J.F., and Mirny, L. (2016). Compaction and segregation of sister chromatids via active loop extrusion. eLife *5*, e14864. https://doi.org/10.7554/eLife.14864.

66. Gabriele, M., Brandão, H.B., Grosse-Holz, S., Jha, A., Dailey, G.M., Cattoglio, C., Hsieh, T.-H.S., Mirny, L., Zechner, C., and Hansen, A.S. (2022). Dynamics of CTCF- and cohesin-mediated chromatin looping revealed by live-cell imaging. Science *376*, 496–501. https://doi.org/10.1126/science.abn6583.

67. Cattoglio, C., Pustova, I., Walther, N., Ho, J.J., Hantsche-Grininger, M., Inouye, C.J., Hossain, M.J., Dailey, G.M., Ellenberg, J., Darzacq, X., et al. (2019). Determining cellular CTCF and cohesin abundances to constrain 3D genome models. eLife *8*, e40164. https://doi.org/10.7554/eLife.40164.

68. Mach, P., Kos, P.I., Zhan, Y., Cramard, J., Gaudin, S., Tünnermann, J., Marchi, E., Eglinger, J., Zuin, J., Kryzhanovska, M., et al. (2022). Cohesin and CTCF control the dynamics of chromosome folding. Nat. Genet. *54*, 1907–1918. https://doi.org/10.1038/s41588-022-01232-7.

69. Hajjoul, H., Mathon, J., Ranchon, H., Goiffon, I., Mozziconacci, J., Albert, B., Carrivain, P., Victor, J.-M., Gadal, O., Bystricky, K., et al. (2013). High-throughput chromatin motion tracking in living yeast reveals the flexibility of the fiber throughout the genome. Genome Res. *23*, 1829–1838. https://doi.org/10.1101/gr.157008.113.

70. Polovnikov, K.E., Brandão, H.B., Belan, S., Slavov, B., Imakaev, M., and Mirny, L.A. (2023). Crumpled Polymer with Loops Recapitulates Key Features of Chromosome Organization. Phys. Rev. X *13*, 041029. https://doi.org/10.1103/PhysRevX.13.041029.

71. Park, D.S., Nguyen, S.C., Isenhart, R., Shah, P.P., Kim, W., Barnett, R.J., Chandra, A., Luppino, J.M., Harke, J., Wai, M., et al. (2023). High-throughput Oligopaint screen identifies druggable 3D genome regulators. Nature *620*, 209–217. https://doi.org/10.1038/s41586-023-06340-w.

72. Kueng, S., Hegemann, B., Peters, B.H., Lipp, J.J., Schleiffer, A., Mechtler, K., and Peters, J.-M. (2006). Wapl controls the dynamic association of cohesin with chromatin. Cell *127*, 955–967. https://doi.org/10.1016/j.cell.2006.09.040.

73. Gandhi, R., Gillespie, P., and Hirano, T. (2006). Human Wapl Is a Cohesin-binding Protein that Promotes Sister Chromatid Resolution in Mitotic Prophase. Curr. Biol. CB *16*, 2406–2417. https://doi.org/10.1016/j.cub.2006.10.061.

74. Liu, N.Q., Magnitov, M., Schijns, M.M.G.A., van Schaik, T., Teunissen, H., van Steensel, B., and de Wit, E. (2025). Extrusion fountains are restricted by WAPL-dependent cohesin release and CTCF barriers. Nucleic Acids Res. *53*, gkaf549. https://doi.org/10.1093/nar/gkaf549.

75. Shintomi, K., and Hirano, T. (2009). Releasing cohesin from chromosome arms in early mitosis: opposing actions of Wapl–Pds5 and Sgo1. Genes Dev. *23*, 2224–2236. https://doi.org/10.1101/gad.1844309.

76. Ouyang, Z., Zheng, G., Tomchick, D.R., Luo, X., and Yu, H. (2016). Structural Basis and IP 6 Requirement for Pds5-Dependent Cohesin Dynamics. Mol. Cell *62*, 248–259. https://doi.org/10.1016/j.molcel.2016.02.033.

77. Nasmyth, K.A., Lee, B.-G., Roig, M.B., and Löwe, J. (2023). What AlphaFold tells us about cohesin's retention on and release from chromosomes. eLife *12*, RP88656. https://doi.org/10.7554/eLife.88656.

78. Carvajal-Maldonado, D., Byrum, A.K., Jackson, J., Wessel, S., Lemaçon, D., Guitton-Sert, L., Quinet, A., Tirman, S., Graziano, S., Masson, J.-Y., et al. (2019). Perturbing cohesin dynamics drives MRE11 nuclease-dependent replication fork slowing. Nucleic Acids Res. *47*, 1294–1310. https://doi.org/10.1093/nar/gky519.

79. Gassler, J., Brandão, H.B., Imakaev, M., Flyamer, I.M., Ladstätter, S., Bickmore, W.A., Peters, J.-M., Mirny, L.A., and Tachibana, K. (2017). A mechanism of cohesin-dependent loop extrusion organizes zygotic genome architecture. EMBO J. *36*, 3600–3618. https://doi.org/10.15252/embj.201798083.

80. Zhang, H., Emerson, D.J., Gilgenast, T.G., Titus, K.R., Lan, Y., Huang, P., Zhang, D., Wang, H., Keller, C.A., Giardine, B., et al. (2019). Chromatin Structure Dynamics During the Mitosis to G1-Phase Transition. Nature *576*, 158–162. https://doi.org/10.1038/s41586-019-1778-y.

81. Sakata, T., Tei, S., Izumi, K., Krantz, I.D., Bando, M., and Shirahige, K. (2025). A common molecular mechanism underlying Cornelia de Lange and CHOPS syndromes. Curr. Biol. *35*, 1353-1363.e5. https://doi.org/10.1016/j.cub.2025.01.044.

82. Remeseiro, S., Cuadrado, A., Kawauchi, S., Calof, A.L., Lander, A.D., and Losada, A. (2013). Reduction of Nipbl impairs cohesin loading locally and affects transcription but not cohesion-dependent functions in a mouse model of Cornelia de Lange Syndrome. Biochim. Biophys. Acta *1832*, 10.1016/j.bbadis.2013.07.020. https://doi.org/10.1016/j.bbadis.2013.07.020.

83. Gibcus, J.H., Samejima, K., Goloborodko, A., Samejima, I., Naumova, N., Nuebler, J., Kanemaki, M.T., Xie, L., Paulson, J.R., Earnshaw, W.C., et al. (2018). A pathway for mitotic chromosome formation. Science *359*, eaao6135. https://doi.org/10.1126/science.aao6135.

84. Kadam, S., Sahoo, S., Kumar, P.B.S., and Padinhateeri, R. (2025). Dynamical properties of chromatin provide insights into key folding principles. Preprint at bioRxiv, https://doi.org/10.1101/2024.05.07.592992 https://doi.org/10.1101/2024.05.07.592992.

85. Lee, J., Chen, L.-F., Gaudin, S., Gupta, K., Spakowitz, A., and Boettiger, A.N. (2025). Kinetic organization of the genome revealed by ultra-resolution, multiscale live imaging. Preprint at bioRxiv, https://doi.org/10.1101/2025.03.27.645817 https://doi.org/10.1101/2025.03.27.645817.

86. Tortora, M.M.C., and Fudenberg, G. (2024). The physical chemistry of interphase loop extrusion. Preprint at bioRxiv, https://doi.org/10.1101/2024.08.23.609419 https://doi.org/10.1101/2024.08.23.609419.

87. Samejima, K., Gibcus, J.H., Abraham, S., Cisneros-Soberanis, F., Samejima, I., Beckett, A.J., Puáčeková, N., Abad, M.A., Spanos, C., Medina-Pritchard, B., et al. (2025). Rules of engagement for condensins and cohesins guide mitotic chromosome formation. Science *388*, eadq1709. https://doi.org/10.1126/science.adq1709.

88. Kim, E., Barth, R., and Dekker, C. (2023). Looping the Genome with SMC Complexes. Annu. Rev. Biochem. *92*, 15–41. https://doi.org/10.1146/annurev-biochem-032620-110506.

89. Porter, H., Li, Y., Neguembor, M.V., Beltran, M., Varsally, W., Martin, L., Cornejo, M.T., Pezić, D., Bhamra, A., Surinova, S., et al. (2023). Cohesin-independent STAG proteins interact with RNA and R-loops and promote complex loading. eLife *12*, e79386. https://doi.org/10.7554/eLife.79386.

90. Shi, Z., Gao, H., Bai, X., and Yu, H. (2020). Cryo-EM structure of the human cohesin-NIPBL-DNA complex. Science *368*, 1454–1459. https://doi.org/10.1126/science.abb0981.

91. Li, Y., Muir, K.W., Bowler, M.W., Metz, J., Haering, C.H., and Panne, D. (2018). Structural basis for Scc3-dependent cohesin recruitment to chromatin. eLife *7*, e38356. https://doi.org/10.7554/eLife.38356.

92. Wutz, G., Ladurner, R., St Hilaire, B.G., Stocsits, R.R., Nagasaka, K., Pignard, B., Sanborn, A., Tang, W., Várnai, C., Ivanov, M.P., et al. (2020). ESCO1 and CTCF enable formation of long chromatin loops by protecting cohesinSTAG1 from WAPL. eLife *9*, e52091. https://doi.org/10.7554/eLife.52091.

93. Ouyang, Z., and Yu, H. (2017). Releasing the cohesin ring: A rigid scaffold model for opening the DNA exit gate by Pds5 and Wapl. BioEssays *39*, 1600207. https://doi.org/10.1002/bies.201600207.

94. Owen, J.A., Osmanović, D., and Mirny, L. (2023). Design principles of 3D epigenetic memory systems. Science *382*, eadg3053. https://doi.org/10.1126/science.adg3053.

95. Bonev, B., Mendelson Cohen, N., Szabo, Q., Fritsch, L., Papadopoulos, G.L., Lubling, Y., Xu, X., Lv, X., Hugnot, J.-P., Tanay, A., et al. (2017). Multiscale 3D Genome Rewiring during Mouse Neural Development. Cell *171*, 557-572.e24. https://doi.org/10.1016/j.cell.2017.09.043.

96. Stelloh, C., Reimer, M.H., Pulakanti, K., Blinka, S., Peterson, J., Pinello, L., Jia, S., Roumiantsev, S., Hessner, M.J., Milanovich, S., et al. (2016). The cohesin-associated protein Wapal is required for proper Polycomb-mediated gene silencing. Epigenetics Chromatin *9*, 14. https://doi.org/10.1186/s13072-016-0063-7.

97. Arruda, N.L., Bryan, A.F., and Dowen, J.M. (2022). PDS5A and PDS5B differentially affect gene expression without altering cohesin localization across the genome. Epigenetics Chromatin *15*, 30. https://doi.org/10.1186/s13072-022-00463-6.

98. Nora, E.P., Lajoie, B.R., Schulz, E.G., Giorgetti, L., Okamoto, I., Servant, N., Piolot, T., van Berkum, N.L., Meisig, J., Sedat, J., et al. (2012). Spatial partitioning of the regulatory landscape of the X-inactivation centre. Nature *485*, 381–385. https://doi.org/10.1038/nature11049.

99. Schalbetter, S.A., Fudenberg, G., Baxter, J., Pollard, K.S., and Neale, M.J. (2019). Principles of meiotic chromosome assembly revealed in S. cerevisiae. Nat. Commun. *10*, 4795. https://doi.org/10.1038/s41467-019-12629-0.

100. Zuo, W., Chen, G., Gao, Z., Li, S., Chen, Y., Huang, C., Chen, J., Chen, Z., Lei, M., and Bian, Q. (2021). Stage-resolved Hi-C analyses reveal meiotic chromosome organizational features influencing homolog alignment. Nat. Commun. *12*, 5827. https://doi.org/10.1038/s41467-021-26033-0.

101. Song, M., Zhai, B., Yang, X., Tan, T., Wang, Y., Yang, X., Tan, Y., Chu, T., Cao, Y., Song, Y., et al.

(2021). Interplay between Pds5 and Rec8 in regulating chromosome axis length and crossover frequency. Sci. Adv. *7*, eabe7920. https://doi.org/10.1126/sciadv.abe7920.

102. Quiroz, D., Pierce, A., and Monroe, G. (2025). Evolutionary trajectories of cohesin gatekeepers across the Tree of Life. bioRxiv, 2025.05.08.652912. https://doi.org/10.1101/2025.05.08.652912.

103. Horsfield, J., Print, C.G., and Mönnich, M. (2012). Diverse Developmental Disorders from The One Ring: Distinct Molecular Pathways Underlie the Cohesinopathies. Front. Genet. *3*. https://doi.org/10.3389/fgene.2012.00171.

104. Panarotto, M., Davidson, I.F., Litos, G., Schleiffer, A., and Peters, J.-M. (2022). Cornelia de Lange syndrome mutations in NIPBL can impair cohesin-mediated DNA loop extrusion. Proc. Natl. Acad. Sci. *119*, e2201029119. https://doi.org/10.1073/pnas.2201029119.

105. Kean, C.M., Tracy, C.J., Mitra, A., Rahat, B., Van Winkle, M.T., Gebert, C.M., Noeker, J.A., Calof, A.L., Lander, A.D., Kassis, J.A., et al. (2022). Decreasing Wapl dosage partially corrects embryonic growth and brain transcriptome phenotypes in Nipbl+/− embryos. Sci. Adv. *8*, eadd4136. https://doi.org/10.1126/sciadv.add4136.

106. Linares-Saldana, R., Kim, W., Bolar, N.A., Zhang, H., Koch-Bojalad, B.A., Yoon, S., Shah, P.P., Karnay, A., Park, D.S., Luppino, J.M., et al. (2021). BRD4 orchestrates genome folding to promote neural crest differentiation. Nat. Genet. *53*, 1480–1492. https://doi.org/10.1038/s41588-021-00934-8.

107. Nora, E.P., Goloborodko, A., Valton, A.-L., Gibcus, J.H., Uebersohn, A., Abdennur, N., Dekker, J., Mirny, L.A., and Bruneau, B.G. (2017). Targeted Degradation of CTCF Decouples Local Insulation of Chromosome Domains from Genomic Compartmentalization. Cell *169*, 930-944.e22. https://doi.org/10.1016/j.cell.2017.05.004.

108. Barutcu, A.R., Blencowe, B.J., and Rinn, J.L. (2019). Differential contribution of steady-state RNA and active transcription in chromatin organization. EMBO Rep. *20*.

109. Dequeker, B.J.H., Scherr, M.J., Brandão, H.B., Gassler, J., Powell, S., Gaspar, I., Flyamer, I.M., Lalic, A., Tang, W., Stocsits, R., et al. (2022). MCM complexes are barriers that restrict cohesin-mediated loop extrusion. Nature, 1–7. https://doi.org/10.1038/s41586-022-04730-0.

110. Linkert, M., Rueden, C.T., Allan, C., Burel, J.-M., Moore, W., Patterson, A., Loranger, B., Moore, J., Neves, C., Macdonald, D., et al. (2010). Metadata matters: access to image data in the real world. J. Cell Biol. *189*, 777–782. https://doi.org/10.1083/jcb.201004104.

111. Abdennur, N., and Mirny, L.A. (2020). Cooler: scalable storage for Hi-C data and other genomically labeled arrays. Bioinforma. Oxf. Engl. *36*, 311–316. https://doi.org/10.1093/bioinformatics/btz540.

Figure 1

Figure 1: Acute NIPBL depletion causes profound defects in genome folding despite leaving substantial amounts of cohesin on chromatin

(A) Hi-C contact maps (20kb bins) after acute RAD21 or NIPBL depletion with inducible degrons in mouse embryonic stem cells.

(B) Flow cytometry of Halo-tagged RAD21 cells (see Methods). Acute NIPBL depletion only reduces chromatin-bound cohesin 2 fold, even after 1-2 cell divisions (see Figure S1D).

(C) iFRAP analyses indicate cohesin residence time (lifetime) remains similar after NIPBL depletion. n=16-18 cells. Shown are medians +/- standard deviation.

(D) Reducing cohesin dosage causes distinct Hi-C patterns from NIPBL depletion.

(E) Flow cytometry quantification of Halo-tagged RAD21 levels after partial induction of the FKBP degron with limiting dTAG amounts, using RAD21-FKBP-Halotag cells.

Figure S1

Figure S1: Additional characterization of degron cells, related to Figure 1

(A) Schematic of cohesin core subunits and cofactors.

(B) *P(s)* plots after near-complete degradation of RAD21 or NIPBL as in[23] and Halo-tagging of RAD21.

(C) Procedure to measure total and chromatin-bound cohesin RAD21 with flow cytometry.

(D) Depleting NIPBL for longer than 6h does not further decrease the amount of bound cohesin, even after 24h (1-2 cell divisions).

(E) Partial depletions of cohesin levels by dTAG titration in RAD21-FKBP degron cells.

(F) Partial cohesin depletion conditions used for Hi-C do not interfere with cell cycle.

Figure 2

1356

Figure 2: Polymer models support a role for NIPBL beyond cohesin loading in controlling the rate of loop extrusion

(A) Overview of the polymer modeling approach.

(B) *P(s)* curves from experimental Hi-C after NIPBL or partial RAD21 depletion.

(C) *in silico* exploration of the effect of cohesin separation on *P(s)* curves from polymer simulations (the basal separation modeling WT untagged cells was set to 185 kb).

(D) Fit scores between simulated $P(s)$ curves with varying cohesin separation and experimental Hi-C from cells with measured 50% (top) or 30% bound cohesin (bottom). Note the best model fits closely match the levels of cohesin measured experimentally by flow cytometry.

(E) Simulated $P(s)$ curves corresponding to the parameters that best fit the experiments.

(F) *in silico* exploration of the effect of extrusion rates on $P(s)$ curves from polymer simulations (the basal rate modeling WT untagged cells was set to 0.25 kb/sec).

(G) Fit scores between simulated $P(s)$ curves with varying cohesin separation and experimental Hi-C from NIPBL depleted cells.

(H) Fit scores between simulated $P(s)$ curves with varying extrusion rates and experimental Hi-C from NIPBL depleted cells. Note models better fit reduced extrusion rates than increased cohesin separation .

(I) The simulated fraction of extruders with at least one collided side diminishes as the separation between them increases. Collisions between cohesins therefore limit loop growth, explaining why reducing cohesin levels to 50% or 30% increases loop size.

(J) Summary cartoons.

Figure S2

A Tuning extrusion rates in simulations

1D extruder updates → 3D polymer updates

1D lattice (genomic position)
extruder 'legs'

3D bead spring monomers

B Calibrating basal simulation parameters using experimental data with normal cohesin

parameter	experimental data type
Extruder number	Fluorescence correlation spectroscopy Cattoglio et al. eLife 2019
Extruder lifetime	RAD21 FRAP Hansen et al. eLife 2017
Net extrusion rate	Deduced from sweeps (Fig S2E-F)
Correspondence between simulated and physical time	Locus tracking Mach et al. Nature Genetics 2022 Hajjoul et al. Genome Research 2013
Self affinity between B-type monomers (compartmentalization)	Hi-C in WT untagged cells (this study)

C Assigning compartment types in polymer simulations

chr2 70Mb 85Mb 100Mb 115Mb

Experimental Hi-C

log10 (contacts): -1.5, -2.0, -2.5, -3.0, -3.5, -4.0, -4.5

Experimental Eigenvector 1 — A-type / B-type

compartment type assignment to model monomers (not drawn to scale)

D Calibrating compartmentalization strength in silmulations

A-type B-type

ε

ø

Spearman R coefficient: 0.76, 0.72, 0.68, 0.64, 0.60

ε (k_BT): 0.02, 0.04, 0.06, 0.08, 0.10

Best fit

ε = 0.02 k_BT ε = 0.03 k_BT ε = 0.04 k_BT ε = 0.05 k_BT (Best fit)
ε = 0.06 k_BT ε = 0.07 k_BT ε = 0.08 k_BT ε = 0.09 k_BT

log10 (observed/expected): +1.0, +0.5, 0.0, -0.5, -1.0, -1.5

E Deducing basal extrusion rate from fitting simulated 3D-to-1D steps to experimental data

Experimental vs. simulated *P(s)* for various rates

- - - Experimental Hi-C (untagged cells)

(Simulated) Hi-C counts (P)

genomic separation (s): 100kb, 1Mb

3D-to-1D steps: 7000, 6500, 6000, 5500, 5000, 4500, 4000, 3500, 3000, 2500

Fit to experimental *P(s)* - untagged cells

R^2: 0.995, 0.990, 0.985, 0.980, 0.975, 0.970, 0.965

3D-to-1D steps: 3000, 4000, 5000, 6000, 7000

Best fit

F Correspondence of 3D-to-1D steps to extrusion rates

extrusion rate (bp/s): 400, 350, 300, 250, 200, 150

3D-to-1D steps: 3000, 4000, 5000, 6000, 7000

G Simulation fits to experiments

Fits to partial RAD21 depletion

R^2: 1.00, 0.98, 0.96, 0.94, 0.92, 0.90

rate multiplier: 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1.0

50% RAD21 left (measured)
30% RAD21 left (measured)

H

depletion:
untagged control
RAD21 partial (50% left)
RAD21 partial (30% left)
NIPBL

Hi-C counts (P): 10^{-3}, 10^{-4}, 10^{-5}, 10^{-6}

Experiments

Slope: -0.5, -1.0, -1.5, -2.0, -2.5

genomic separation (s): 100kb, 1Mb, 10Mb

s[P'max]

s[P'max]
NIPBL depletion: **55kb**
untagged: **90kb**
RAD21 partial (50% left): **150kb**
RAD21 partial (30% left): **210kb**

I

Genomic distance (kb): 325, 300, 275, 250, 225, 200, 175, 150, 125

processivity
mean loop size
s[P'max]

Separation multiplier: 1.0, 1.5, 2.0, 2.5, 3.0, 3.5, 4.0, 4.5, 5.0

Figure S2: Additional information about modeling approaches, related to Figure 2

(A) Extrusion simulations alternate between updating positions of polymer beads in 3D and extruder leg positions in 1D (along the chromosome). Tuning extrusion rates in simulations requires modification of the distance traveled and the relative relaxation rate of the polymer fiber.

(B) Experimental data used to calibrate simulations.

(C) Illustration of how compartment identity is assigned in simulations.

(D) The self-affinity (interaction energy) between B-type monomers in simulation was determined by fitting to experimental Hi-C from untagged control cells (see methods). Heatmaps highlight experimental (lower left corner) v.s simulated (top right corner) over a range of self affinities. Note that with this layout for compartments, compartmentalization had minimal impact on simulated *P(S)* for the range of B-B attraction strengths considered.

(E) Sweep of polymer model versus lattice model timescales (i.e., 3D-to-1D steps), where each Molecular Dynamics step was estimated to amount to 5 ms of physical time (methods). 4000 3D-to-1D steps best fit the experimental Hi-C *P(s)* of WT untagged cells.

(F) Effective average extrusion rate as a function of 3D-to-1D steps, showing that 4000 3D-to-1D steps in WT untagged cells correspond to an extrusion rate of 250 bp/s.

(G) Fit scores between simulated *P(s)* curves with varying extrusion rate and experimental Hi-C from cells with partial cohesin depletions. Varying extrusion rate produces worse fit to experimental partial cohesin depletion than lower extruder separation (as shown in Figure 2D).

(H) s[P'max] is defined as the inflection point of the Hi-C *P(s)*, or the first maximum of its slope.

(I) Scaling of mean loop size and s[P'max] according to extruder separation in simulations.

Figure 3

Figure 3: Reduced extrusion rate after NIPBL co-depletion, not reduced loading, mitigates genome misfolding caused by increased lifetime upon WAPL inactivation

(A and B) Experimental Hi-C (A) and corresponding *P(s)* curves (B) showing that chromosome folding after acute co-depletion of WAPL + NIPBL resembles that of normal cells, but is distinct from patterns observed with WAPL + partial RAD21 co-depletion at equivalent amount of bound cohesin (see Figure S6A).

(C) Considering that processivity is the product of lifetime and rate provides an immediate explanation for how NIPBL depletion (which reduces extrusion rate, see Figures 1 and 2) can counteract the effect of WAPL depletion (which increases lifetime).

(D) Summary cartoons.

Figure S3

1412
1413

Figure S3: RAD21 quantifications, related to Figure 3

(A) Flow cytometry quantification of RAD21, highlighting the conditions used for Hi-C.

1416

Figure 4

Figure 4: Compartment scores disambiguate the contribution of cohesin lifetime and extrusion rate to genome misfolding upon WAPL or PDS5AB depletion

(A) Hi-C after acute depletion of WAPL or combined PDS5A and PDS5B (hereafter abbreviated PDS5AB).

(B) Experimental *P(s)* curves highlighting the greater impact of depleting PDS5AB than WAPL.

(C) Goodness-of-fit between experimental and simulated *P(s)* as a function of lifetime and extrusion rate after WAPL depletion, or (D) PDS5AB depletion. Red boxes highlight that many combinations of these parameters produce good fits ($R^2 > 0.95$, red boxes) to the experimental *P(s)*. The basal lifetime and rate modeling WT untagged cells was set to 22 minutes and 0.25 kb/sec respectively).

(E) Simulations reveal that compartment scores decrease more steeply upon increasing extrusion rate than increasing cohesin lifetime. Lower scores indicate weaker separation of A and B compartments. Here base extruder separation was used for all simulations.

(F) Experimental Hi-C contact maps (chromosome-wide, 1 Mb bins) and corresponding compartment scores (for the same chr2:66,000,000-116,000,000 mm10 region as used in simulations) are more altered upon depleting PDS5AB than WAPL.

(G) Simulated *P(s)* with multipliers chosen from fitting both *P(s)* and compartment scores (note 0.8x cohesin separation comes from calibration to the experimentally-measured amount of chromatin-bound RAD21, see Figure S4E).

Figure S4

Figure S4: Additional characterization of PDS5 degron cells, related to Figure 4

(A) Western blot quantification of PDS5A and (B) PDS5B depletion.

(C and D) Experimental Hi-C (C) and corresponding *P(s)* curves (D) after depleting either PDS5A and PDS5B, highlighting that each paralog contributes to overall chromosome folding in ESCs.

(E) Flow cytometry quantification of RAD21-Halotag after PDS5AB or WAPL depletion. Degron leakiness of the AID system, as seen in (A), accounts for the elevated chromatin-bound fraction prior to adding auxin in PDS5AB-AID or WAPL-AID cells.

(F) PDS5AB depletion does not cause cell cycle defects within the first day, but longer depletions result in replication stress.

(G) Cartoon summarizing the experimentally calibrated biophysical framework to measure separation and infer lifetime and rate parameters.

Figure 5

1449
1450

Figure 5: PDS5AB depletion accelerates the rate of cohesin loop extrusion

(A) Example of simulated chromosomes (left) and *in silico* microscopy images obtained from analyzing the location of extruders (right). Simulations were conducted using extrusion parameters that best fit Hi-C

 experiments (see Figure 4). The models predict that cohesin clusters more strongly into axial structures
 (vermicelli) upon PDS5AB depletion compared to WAPL depletion.
 (B) Experimental staining of RAD21 in Halo-tagged subclones (JFX646 dye) highlighting stronger clustering
 into vermicelli after PDS5AB depletion than WAPL depletion. Vermicelli scores correspond to the coefficient of
 variation in (simulated) microscopy images (see Methods).
 (C) Cohesin clustering into vermicelli occurs more rapidly after G1 entry following PDS5AB depletion than
 WAPL depletion, consistent with PDS5AB depletion accelerating the rate of loop extrusion.
 (D) Inverse Fluorescence Recovery After Photobleaching (iFRAP) of RAD21-Halotag after either WAPL or
 PDS5AB depletion, supporting the model predictions that depleting PDS5AB does not increase cohesin
 lifetime as much as depleting WAPL. n=12-18 cells. Shown are medians +/- standard deviation.
 (E) Summary cartoons.

Figure S5

Figure S5: Additional characterizations of degron cells, related to Figure 5

(A) Synchronization controls of DNA content (DAPI) measured by flow cytometry after nocodazole-induced mitotic block. Note that only a fraction of cells re-enter G1 after nocodazole washoff, precluding any bulk assays with this setup.

(B) Microscopy examples of unreleased cells blocked in M-phase (left), or after 2h of nocodazole washoff (right). Cells that have not re-entered G1 are eliminated from downstream vermicelli analyses based on their high H3S10Ph immunostaining.

(C-D) Inverse fluorescence recovery after photobleaching (iFRAP) of RAD21-Halotag after WAPL of PDS5AB depletion with untreated samples. Lifetime increase in PDS5AB-AID cells prior to auxin treatment can be attributed to leakiness of the degron. n=12-18 cells. Shown are medians +/- standard deviation.

Figure 6

Figure 6: The dosage of NIPBL and its balance with PDS5 shapes chromosome folding by tuning the rate of cohesin extrusion

(A and B) Experimental Hi-C (A) and corresponding *P(s)* curves (B) after acute co-depletion of PDS5AB + NIPBL closely resemble WAPL depletion, and are distinct from PDS5AB + partial RAD21 co-depletion.

(C) Simulated *P(s)* curves and corresponding parameters of loop extrusion. Rate multipliers for the single NIPBL depletion were obtained as the best fit to experimental *P(s)*, in Figure 2. Lifetime and rate multipliers for PDS5AB-only depletion were obtained as the joint best fit to experimental *P(s)* and compartment scores, in Figure 4. Multipliers for co-depletions were computed by multiplying loop extrusion parameters of the single mutants. Separation multipliers in simulations were calibrated to match experimental measurements of chromatin-bound RAD21 by flow cytometry in each of the single and co-depletions. Simulations indicate that reducing rate (NIPBL depletion) can offset the increase in rate caused by PDS5AB depletion, but not the concomitant increase in lifetime - explaining why NIPBL+PDS5AB *P(s)* closely match that of WAPL

(D) Depleting PDS5AB increases processivity by increasing both cohesin lifetime and extrusion rates. Lowering extrusion rate by co-depleting NIPBL and PDS5AB offsets the extrusion rate but does not restore lifetime. Considering the separate effect of PDS5AB depletion on cohesin lifetime and extrusion rate therefore explains why co-depleting NIPBL + PDS5AB leads to similar genome folding patterns as depleting WAPL alone.

(E) Cartoon model illustrating how the NIPBL:PDS5 balance sets the net rate of cohesin loop extrusion

(F) Transient NIPBL over-expression in RAD21-Halotag ESCs is sufficient to cause chromosome compaction into vermicelli. Inset shows a representative cell with high NIPBL expression.

(G) Approach to determine how extrusion rates scale with NIPBL dosage: (top) Hi-C maps were collected at four levels of experimentally-titrated NIPBL dosages; (bottom left) best fits to simulations were computed to infer relative extrusion rates (as a multiplier relative to baseline untagged cells - 0.25kb/s). (bottom right) inferred extrusion rates were plotted as a function of NIPBL expression levels (circles: data points obtained with NIPBL-FKBP cells; squares: NIPBL-FKBP-GFP cells; triangle: untagged control cells). Bound cohesin levels were measured with RAD21-Halotag subclones (see Figure S6E).

Figure S6

1507

Figure S6: Additional characterization of degron cells, related to Figure 6

(A) Flow cytometry quantification of RAD21-Halotag, highlighting the conditions used for Hi-C in the indicated PDS5AB-AID backgrounds.

(B) Simulated *P(s)* curves and corresponding parameters of loop extrusion as in Figure 6C, but in the WAPL-depletion background. These simulations contrast the effect of co-depleting NIPBL or RAD21 together with WAPL, highlighting how an increase in lifetime (WAPL depletion) can be offset by a decrease in rate (NIPBL depletion) - yielding *P(s)* that resemble the basal situation in untagged cells. Lifetime increase cannot be offset by reducing the amount of extruders (partial RAD21 depletion).

(C) Western blot quantification of basal NIPBL levels in NIPBL-FKBP and NIPBL-FKBP-GFP ESCs, and confirmation of degron potency with 500mM dTAG (maximum depletion).

1518 (D) NIPBL-FKBP-GFP levels after dTAG titration, normalized to basal levels in untagged cells as estimated by
1519 Western-blots. Doses used for Hi-C are indicated.
1520 (E) Flow cytometry quantification of RAD21-Halotag, highlighting the conditions used for Hi-C and
1521 corresponding simulations in the indicated NIBPL-degron cells.
1522

Figure 7

1523

Figure 7: Effect of altering combinations of cohesin cofactors on genomic compartmentalization and transcriptomes

(A) Ordering of experimental depletions by compartment scores/

(B) Ordering of simulated depletions by compartment scores. Separation values correspond to the experimentally measured chromatin-bound RAD21; the lifetime and rate multipliers for co-depletions (underlined) are the product of the best fits for single depletions, as in Figure 3 and 6. Note the identical ordering of compartment scores between simulations and experiments in (A).

(C) Illustration of how extrusion rate alters genomic compartmentalization.

(D) RNA-seq PCA in the indicated genotypes treated with auxin or dTAG or 1d, showing PC1 and PC2 and all three replicates separately. Tir1 are control cells expressing the auxin receptor transgene but without any degron knockin. Transcriptomes of NIPBL-FKBP cells were obtained from Hansen et al.[23].

(E) *top:* RNA-seq Principal component analysis PC1 values separate depletion conditions according to their overall transcriptomic similarity, averaging replicates. *Bottom:* Samples ordered by genomic distance of their inflection points (s[P'max]) derived from Hi-C *P(s)* curves (see Figure S2H). Samples with similar Hi-C s[P'max] (which relates to loop size Figure S2H and S2I) have similar transcriptomes.

Figure S7

A Simulated conformation examples

lifetime x0.25 rate x0.25   *lifetime x1 rate x1*   *lifetime x4 rate x4*

A-type segment
B-type segment
Extruders

1μm

B Simulated compartment scores

C Hi-C saddle plots and compartment scores

untagged — *Sim: 0.21* / *Exp: 0.2*
WAPL-AID 6h — *Sim: 0.16* / *Exp: 0.16*
PDS5AB-AID 6h — *Sim: 0.12* / *Exp: 0.06*

NIPBL-FKBP 6h — *Sim: 0.3* / *Exp: 0.3*
WAPL-AID + NIPBL-FKBP 6h — *Sim: 0.23* / *Exp: 0.26*
PDS5AB-AID + NIPBL-FKBP 6h — *Sim: 0.15* / *Exp: 0.18*

RAD21-FKBP (50%left) 6h — *Sim: 0.24* / *Exp: 0.28*
WAPL-AID + RAD21-FKBP(80%left) 6h — *Sim: 0.17* / *Exp: 0.22*
PDS5AB-AID + RAD21-FKBP(80%left) 6h — *Sim: 0.12* / *Exp: 0.12*

RAD21-FKBP(50% left) 6h — *Sim: 0.24* / *Exp: 0.28*
WAPL-AID + RAD21-FKBP(45%left) 6h — *Sim: 0.18* / *Exp: 0.27*
PDS5AB-AID + RAD21-FKBP(45%left) 6h — *Sim: 0.12* / *Exp: 0.16*

D Compartment scores vs. genomic distance

cis *trans* untagged
PDS5AB-AID + NIPBL-FKBP
PDS5AB-AID + RAD21-FKBP(45%left) 6h
PDS5AB-AID + RAD21-FKBP(80%left) 6h
PDS5AB-AID

1541
1542

Figure S7:, related to Figure 7

(A) Example of simulated conformations highlighting the spatial positions of compartment types and extruders.

(B) Simulated compartment scores obtained with a range of lifetime and rate multipliers, at basal extruder
separation.

(C) Comparison of Hi-C saddle plots obtained from simulations and experiments with corresponding
compartment scores.

(D) Experimental compartment scores from Hi-C across genomic distances in *cis* and in *trans* (see methods).

Supplementary material

Video S1: vidS1.mp4 - Differential effects of chromatin lifetime and extrusion rate on single-cohesin behavior, related to Figure 7.

Representative simulation time-lapse (sped up x50 relative to physical time) of extrusion by isolated cohesin at baseline rate and increased lifetime (left), or baseline lifetime and increased rate (right). The chromatin fiber is artificially stretched by an external force of magnitude ~0.5 pN applied at the chain extremities to facilitate visualization.

Video S2: vidS2.mp4 - Simulated impacts of extrusion parameters on local chromatin structure and dynamics, related to Figure 7.

Same as Video S1 for a stretched 2.5 Mb chromatin segment subjected to extrusion at either baseline (top left) or altered cohesin separations, lifetimes and rates, as respectively inferred for NIPBL- (top right), WAPL- (bottom left) and PDS5-depleted mutants (bottom right). See Figure 7B for the full list of parameter values used.

Table 1: Cell-Lines-Vectors-sgRNAs_v1.xlsx
Table 2: Supp_Table_2_HiC_mapping_statistics.xlsx
Archive 1: Annotated plasmid maps.zip