SFTSV NSs protein is a novel tick antiviral RNAi response suppressor

Mazigh Fares[1#], Melanie McFarlane[1#], Rhys H. Parry[2#], Rozeena Arif[1], Andrew T. Clarke[1], Wael Kamel[1], Kelsey Davies[1], Lesley Bell-Sakyi[3], Marine J. Petit[4], Esther Schnettler[5,6,7], Alfredo Castello[1], Alain Kohl[1,8] & Benjamin Brennan[1]*

[1]Medical Research Council–University of Glasgow Centre for Virus Research, Glasgow, G61 1QH, Scotland, UK.
[2]School of Chemistry and Molecular Biosciences, The University of Queensland, St. Lucia 4072, Australia.
[3]Department of Infection Biology and Microbiomes, Institute of Infection, Veterinary and Ecological Sciences, University of Liverpool, Liverpool, L3 5RF, UK.
[4]Microbes, Infection & Immunity, School of Biosciences, Faculty of Health and Medical Sciences, University of Surrey, Guildford, UK
[5]Bernhard-Nocht-Institute for Tropical Medicine, 20359 Hamburg, Germany.
[6]German Centre for Infection Research (DZIF), Partner Site Hamburg-Luebeck-Borstel-Riems, 20359 Hamburg, Germany
[7]Faculty of Mathematics, Informatics and Natural Sciences, University Hamburg, 20148 Hamburg, Germany
[8]Liverpool School of Tropical Medicine, Centre for Neglected Tropical Diseases, Departments of Tropical Disease Biology and Vector Biology, Liverpool, L3 5QA, UK.

[#], joint first authors
*, Correspondence: ben.brennan@glasgow.ac.uk

ORCID:

Mazigh Fares	0000-0002-6003-9244
Melanie McFarlane	0000-0002-8392-2291
Rhys H. Parry	0000-0001-9238-1952
Rozeena Arif	0009-0007-0492-6352
Andrew T. Clarke	0000-0001-7271-3833
Wael Kamel	0000-0002-7625-2648
Kelsey Davies	0009-0009-5624-5983
Lesley Bell-Sakyi	0000-0002-7305-0477
Marine Petit	0000-0001-7427-7203
Esther Schnettler	0000-0001-8141-3728
Alfredo Castello	0000-0002-1499-4662
Alain Kohl	0000-0002-1523-9458
Benjamin Brennan	0000-0003-4707-726X

Keywords:

Bunyavirus, severe fever with thrombocytopenia syndrome virus, tick cell line, arthropod innate immune evasion, protein function, RNA-binding protein, protein-RNA interaction, RNA, host-virus interactions

Abstract

Severe fever with thrombocytopenia syndrome virus (SFTSV) is an emerging tick-borne phenuivirus causing high mortality in humans. While the non-structural protein NSs is dispensable for replication in mammalian cells, we demonstrate for the first time that NSs is essential for viral replication in tick cells. SFTSV infection triggers canonical Dicer-2-mediated antiviral RNA interference (RNAi) in tick cells, producing virus-derived small interfering RNAs (siRNAs) that target viral transcripts for degradation. We show that NSs functions as a viral suppressor of RNAi by selectively sequestering single-stranded RNAs derived from 22-nucleotide siRNAs, preventing their incorporation into RNA-induced silencing complexes (RISC). Complementation with a heterologous RNAi suppressor (p19 protein) partially rescues replication of NSs-deficient virus, validating NSs' RNAi suppressive function. These findings reveal that successful tick-borne viral replication requires host-specific immune evasion strategies and establish NSs-mediated RNAi suppression as essential for SFTSV persistence in arthropod vectors.

MAIN

Severe fever with thrombocytopenia syndrome virus (SFTSV; *Bandavirus dabieense*; *Phenuiviridae*) emerged as a significant threat to human health following its discovery in central China in 2009[1], where it was identified as the causative agent of a severe haemorrhagic fever syndrome. This tick-borne phenuivirus rapidly gained attention due to its high case fatality rate of 6-30%. Its distinctive clinical presentation is characterized by fever, thrombocytopenia, leukocytopenia, and gastrointestinal manifestations[2]. The geographic range of SFTSV has since expanded across East Asia, with confirmed cases reported in South Korea, Japan, Taiwan, and Vietnam, establishing it as a regional public health concern of growing importance[3]. Currently, no effective antiviral drugs or vaccines exist to treat SFTS disease.

SFTSV transmission occurs primarily through the bite of infected ixodid ticks, with several species identified as potential vectors[4]. *Haemaphysalis longicornis* serves as the principal vector across much of the SFTSV geographical range, while *Haemaphysalis flava* and *Rhipicephalus microplus* may also contribute to transmission cycles in specific regions[5]. Ticks acquire SFTSV through blood feeding on infected vertebrate hosts and can subsequently transmit the virus to humans and other susceptible animals. This complex ecology involves multiple tick species and some amplifying vertebrate hosts (hedgehogs and small rodents), creating diverse transmission pathways that complicate control efforts[6].

Arthropod vectors, including ticks, have evolved sophisticated innate immune mechanisms to combat viral infections, with RNA interference (RNAi) representing a primary antiviral defence pathway[7,8]. Unlike mammals, in which type I interferon responses dominate antiviral immunity[9], arthropods rely heavily on RNA silencing machinery to recognise and degrade viral nucleic acids. Tick antiviral RNAi responses produce characteristic 22-nucleotide small interfering RNAs (siRNA) through Dicer-2 processing of viral double-stranded RNA intermediates, which are subsequently incorporated into RNA-induced silencing complexes (RISC) to target complementary viral transcripts for degradation[8,10]. Distinguishing authentic virus-derived siRNAs from RNA degradation products requires methylation-sensitive approaches, as functional siRNAs undergo Hua Enhancer-1 (HEN1)-mediated 2'-O-methylation upon RISC loading[11-15]. The effectiveness of arthropod RNA silencing as an antiviral mechanism

97 has been demonstrated across diverse vector species (flies and mosquitoes)[16], yet
98 our understanding of how tick-borne viruses evade or suppress these responses
99 remains limited[17].
100
The evolutionary arms race between viruses and RNAi defenses has driven many
viruses to evolve suppressors of RNAi (VSRs) that counteract antiviral silencing[18,19].
They employ diverse mechanisms to inhibit RNAi activity, including sequestration of
siRNAs, interference with Dicer processing, and disruption of RISC assembly and
function. While NSs proteins from plant-infecting tospoviruses are well-characterized
RNAi suppressors that bind double-stranded siRNAs[20], vertebrate-infecting
bunyaviruses show variable suppressor activities[21]. Crucially, no arthropod-specific
RNAi suppression mechanisms have been definitively established for tick-borne
phenuiviruses.
110
Despite SFTSV's emergence as a significant public health threat and evidence of
vsiRNA production in infected ticks[22], the specific mechanisms by which tick-borne
phenuiviruses evade or suppress arthropod RNAi responses remain unknown. While
the non-structural protein (NSs) of SFTSV has been suggested as a possible RNAi
suppressor[22,23], its actual role and mechanism of action in arthropod cells have yet to
be definitively established.
117
Here, we demonstrate for the first time that SFTSV exhibits fundamentally different
replication requirements in tick versus mammalian cell cultures, with NSs proving
essential for viral maintenance in tick cells. Using comprehensive molecular analyses
including methylation-sensitive small RNA sequencing, structural modeling, and
complementation studies with heterologous suppressors, we reveal that NSs functions
as a unique viral suppressor of RNAi by selectively sequestering the pool of virus-
derived single stranded siRNAs away from the RNAi machinery, thereby enabling
productive infection in tick cells and enhancing vector-borne transmission potential.
126

RESULTS

SFTSV infects cell lines derived from tick species reported to act as competent
vectors.

To identify suitable tick cellular models for SFTSV infection, we tested multiple cell
lines representing both *Metastriata* (*Rhipicephalus* and *Hyalomma* spp.) and *Prostriata*
(*Ixodes* and *Amblyomma* spp.) lineages of the family *Ixodidae*, using wildtype (wt)
SFTSV strain HB29[1,24]. Ten-day viral yield assays revealed SFTSV efficiently
replicated in cell lines derived from *Rhipicephalus* and *Hyalomma* spp, evidenced by
increased viral titres and nucleocapsid protein detection in infected lysates (Fig. 1A,B).
SFTSV replication was minimal in phylogenetically distinct *Ixodes* and *Amblyomma*
spp. tick cell lines, despite these species vectoring other members of the *Phenuiviridae*
such as Uukuniemi[25] and Heartland viruses[26,27]. Given that bunyaviruses establish
non-pathogenic, continuous viral production in tick cells[25,28], we monitored infectious
virus production daily over 10-days at low MOI (Fig. 1C). Viral titres steadily decreased
over time in *Ixodes* spp. cell lines, but continuous production was evident in
Rhipicephalus spp. cultures. After 10 days post-infection (dpi), all cultures were split
1:2 and monitored for 14 days; clear SFTSV persistence in BME/CTVM6 and
BME/CTVM23 lines (Fig. 1D) confirms their utility as robust models for studying
SFTSV arthropod cell infection[29]. These results suggest SFTSV entry and replication

may depend on host cellular factors that vary among tick species and demonstrate
that permissive cell lines can maintain viable virus over extended culture periods.
148

SFTSV NSs protein expression is repressed in tick cells yet is essential for
sustained virus replication.
We previously established that SFTSV NSs is a potent interferon antagonist[30] and
virulence factor in animal models[31-33], yet dispensable for replication in mammalian
cell systems[34]. However, the molecular basis for NSs function in tick cells, the natural
arthropod vector, has remained unexplored. To address this knowledge gap, we
investigated NSs function during SFTSV infection of tick cells. We infected
BME/CTVM6 cells, selected for their robust and sustained wt SFTSV replication (Fig.
1A-D), with either wt or NSs-deleted (delNSs) SFTSV across a range of multiplicities
of infection. We observed that wt SFTSV demonstrated stable and abundant
production of infectious virus throughout the experiment, with SFTSV titres increasing
by approximately 3-log$_{10}$ ffu/ml over the time course (Fig. 1E). In contrast, delNSs
SFTSV failed to maintain infectious progeny, with viral titres declining progressively
over the infection time course (Fig. 1F). Consistent with this, Western blot analysis of
delNSs SFTSV-infected cell lysates revealed no detectable nucleocapsid protein,
indicating complete failure to establish a productive infection in tick cells (Fig. 1G).
Notably, NSs protein was undetectable in cells infected with both wt and delNSs
SFTSV, despite this protein being detected in mammalian cell controls (Vero E6). This
observation was unexpected given the apparent requirement for NSs in tick cells. To
investigate this paradox further, we performed immunofluorescence microscopy on
infected BME/CTVM6 cells (Fig. 1H). Quantitative analysis revealed that delNSs
SFTSV infection led to only 2-8% nucleocapsid-positive cells (Fig. 1I,J). In contrast,
60-80% of cells were positive for nucleocapsid protein following wt SFTSV infection
across all tested MOIs, demonstrating efficient viral replication (Fig. 1K,L).
173

To gain a deeper understanding of the viral protein abundance in SFTSV infected tick
cells we performed proteomic analysis of BME/CTVM6 cells infected with wild-type
SFTSV at 3- and 6-dpi, examining both the distribution of peptide intensities (Fig. 1M)
and mean peptide intensities (Fig. 1N) for viral proteins NSs, nucleocapsid protein (N),
glycoprotein precursor (GPC), and RNA-dependent RNA polymerase (RdRp). The
peptide intensity distributions for NSs remained unchanged between time points, while
N showed a non-significant increase (P = 0.709). This reflects that N protein peptides
already exhibited high intensities and broad distribution at 3 dpi, suggesting a
detection plateau had been reached. In contrast, GPC and RdRp exhibited significant
increases (P = 0.026 and P = 6.23 x 10^{-12}, respectively), indicative of enhanced viral
replication and assembly processes (Fig. 1M). Analysis of mean peptide intensities
revealed temporal increases for N, GPC, and RdRp from 3 to 6 dpi, while NSs
maintained the lowest expression levels with minimal temporal change (Fig. 1N).
Notably, while NSs-derived peptides were detectable by mass spectrometry, NSs
protein intensity levels remained below the detection threshold of Western blot
analysis (Fig. 1G), suggesting tightly regulated, low-level expression of this protein.
This differential sensitivity between proteomic and immunoblot detection methods
reveals that NSs is present but maintained at substantially lower levels compared to
structural and replication proteins during SFTSV infection of tick cells.
193

NSs subgenomic mRNA expression is reduced in SFTSV-infected tick cells compared to mammalian cells.

The SFTSV NSs protein is encoded within the viral S segment, which employs an ambisense coding strategy to produce both NSs and the N protein, consistent with all phenuiviruses[35]. The NSs open reading frame is translated from a subgenomic mRNA transcribed from the antigenomic S RNA template following viral genome replication. To investigate the molecular basis for NSs repression in tick cells, we analysed viral RNA species derived from the S segment during infection of two *Rhipicephalus* spp. tick cell lines and compared these results to mammalian Vero E6 cells as controls.

We employed Northern blot analysis using strand-specific RNA probes to target either the N or NSs coding regions and to distinguish genomic from antigenomic RNA species (Fig. 1O). Productive viral replication was confirmed in wild-type SFTSV-infected tick cells through detection of both genomic [detected by N(-) and NSs(+) probes] and antigenomic [detected by N(+) and NSs(-) probes] S segment RNA at 6 dpi. While N mRNA was readily detectable in both tick cell lines and in the mammalian cell controls, NSs subgenomic mRNA levels were markedly reduced in tick cells compared to Vero E6 cells when probed with the NSs(+) specific probe (Fig. 1P).

The observed reduction in NSs subgenomic mRNA levels in tick cells provides a plausible explanation for the diminished NSs protein expression during infection. However, the underlying mechanism, whether due to transcriptional attenuation or altered RNA stability, remains unresolved. While our Northern blot data confirm reduced mRNA accumulation, further investigation is needed to determine whether this reflects regulated transcription termination, selective degradation or post-transcriptional silencing. This finding also raises important questions about how NSs functions during early infection, before its apparent repression.

SFTSV infection triggers small RNA production in tick cells.

Given that tick cells rely on RNAi as a primary antiviral defence mechanism, we next examined whether wt SFTSV replication, and specifically NSs mRNA, is targeted by the tick RNAi pathway. To determine whether the observed reduction in SFTSV NSs mRNA reflected targeted RNAi-mediated degradation, we used beta elimination to distinguish between RISC-loaded siRNAs, unloaded siRNAs, and RNA degradation products. Small RNA samples were therefore subjected to beta elimination prior to sequencing analysis. This treatment selectively removes degradation fragments and non-methylated siRNAs containing 2',3'-cyclic phosphate termini while preserving HEN1-methylated, RISC-loaded siRNAs bearing 5'-phosphate and 3'-hydroxyl groups[11-15]. Since both degradation products and non-RISC-loaded siRNAs exhibit similar size distributions (20-24 nt), protection by HEN1 methylation during beta elimination enables unambiguous identification of functional vsiRNAs and confirms RNAi pathway activation against SFTSV[12,13].

The efficacy of beta elimination was validated in uninfected BME/CTVM6 cells or *Aedes aegypti* AF525 cells, an Ago2-deficient *Aedes aegypti* cell line, (Supplementary Fig. 1) and Semliki Forest virus (SFV)-infected AF525 cells (Supplementary Fig. 2A,B) by published methodologies[11]. Following SFTSV infection of BME/CTVM6 cells, approximately 10^5 reads within the small RNA size range (18-30 nt) mapped to each SFTSV genomic segment in untreated samples, with read abundance increasing between 3- and 6-dpi. Although beta elimination reduced overall read counts, the proportional increase in reads mapping to SFTSV RNAs from 3 to 6 dpi was preserved,

indicating sustained small RNA responses targeting SFTSV throughout infection (Fig.
2A).
The 5' nucleotide composition of all 22 nt vsiRNAs showed strong bias toward uridine
and adenine residues, consistent with arthropod Dicer-2 processing specificity[36]. Beta
elimination treatment virtually eliminated the 22 nt peak at both time points, indicating
these vsiRNAs lacked 3'-terminal methylation and remained susceptible to chemical
modification (Fig. 2B,C). Analysis of vsiRNAs mapping to SFTSV genome and
antigenome sequences at both 3- and 6-dpi revealed pronounced enrichment for 22
nt species in untreated samples, with comparable distributions observed for genomic
and antigenomic reads (Fig. 2D,E). Conversely, longer small RNAs (24-30 nt) were
largely unaffected by beta elimination, consistent with 2'-O-methylated piRNAs or
degradation fragments that escaped chemical treatment.
Notably, the selective loss of Dicer-2-processed 22 nt reads following beta elimination
at 3 dpi demonstrates that the majority of SFTSV-derived siRNAs in BME/CTVM6 cells
lack methylation modifications, suggesting exclusion from functional Ago2-containing
RISC complexes (Fig. 2D). However, this transient suppression of vsiRNAs from Ago2
is eventually overwhelmed by 6 dpi, evidenced by the strong 22 nt peak of methylated-
vsiRNAs in the beta-eliminated sample, indicating a resumption in the targeting of viral
RNA sequences by the RNAi machinery (Fig. 2E).
While HEN1-mediated methylation of RISC-loaded siRNAs is conserved from plants
to arthropods[12,37], the dynamics of this process during viral infection in tick cells have
not been previously characterized. Our detection of methylated small RNAs in mock-
infected controls (Supplementary Fig. 1) and their recovery at 6 dpi confirms that
BME/CTVM6 cells possess functional HEN1 methylation machinery. Therefore, the
selective absence of methylated SFTSV-specific siRNAs at 3 dpi (Fig. 2D), followed
by their subsequent accumulation at 6 dpi (Fig. 2E), indicates active viral suppression
of RNAi responses during early infection that is eventually overcome by sustained
antiviral pressure, enabling initial viral replication. The observed NSs transcript
degradation supports this interpretation (Fig. 1P), demonstrating that functional RNAi
activity ultimately prevails despite initial viral evasion mechanisms.
272

RNAi-mediated vsiRNA targeting exhibits temporal dynamics and segment-
specific preferences across the SFTSV genome.
Given that 22-nt species constituted the predominant small RNA population (which
corresponds to the typical length of vsiRNAs produced in tick cells), we mapped
vsiRNA coverage across genomic and antigenomic strands of the SFTSV small (S;
Fig. 2F), medium (M; Fig. 2G), and large (L; Fig. 2H) segments at 3- and 6-dpi in
untreated and beta-eliminated samples. SFV served as a control for beta-elimination
efficacy and mapping specificity (Supplementary Fig. 2C,D).
SFTSV vsiRNA accumulation increased substantially between 3- and 6-dpi across all
segments, indicating progressive amplification of the RNAi response. In untreated
samples, the M and L segments exhibited extensive targeting, with pronounced
accumulation at untranslated regions and segment termini. Beta-elimination confirmed
sustained SFTSV targeting by the tick RNAi machinery, demonstrating that observed
signals represent authentic vsiRNAs rather than 2'-O-methylated small RNA
contaminants. Notably, the S segment displayed concentrated targeting within the
intergenic region and the NSs open reading frame on the genomic strand, particularly
around the areas previously identified as the transcription termination sites for the N
and NSs mRNAs[34], indicating RNAi-mediated cleavage of both genomic RNA and NSs

subgenomic transcripts. This finding provides a mechanistic explanation for the reduced NSs mRNA detection observed by Northern blot analysis (Fig. 1P). The strand-specific and segment-preferential targeting patterns demonstrate that tick cells mount a sophisticated, temporally regulated antiviral RNAi response against SFTSV infection, which the wt virus can overcome, yet a NSs-deletant virus cannot.

SFTSV infection triggers canonical Dicer-2-mediated vsiRNA production in tick cells.

To characterise the dsRNA cleavage patterns during SFTSV infection, we analysed small RNA overlap profiles and strand complementarity patterns mapping to viral segments at 3- and 6-dpi (Fig. 3). At both timepoints, viral small RNAs predominantly exhibited 20-22 nucleotide lengths with a clear preference for 20 nucleotide overlaps between complementary strands (Fig. 3A,C). Strand overlap analysis revealed enhanced complementarity at positions resulting in 2 nucleotide 3' overhangs, characteristic of canonical Dicer-2 cleavage products (Fig. 3B,D). The average z-score across all three genome segments increased from 1.98 at 3 dpi to 2.54 at 6 dpi, suggesting sustained Dicer-2 activity throughout viral replication. Notably, virus-derived small RNAs in the 20-24 nucleotide range all displayed this complementarity pattern across the L, M, and S viral segments, indicating uniform processing.

No evidence of antiviral vpiRNAs (viral PIWI-interacting RNAs) was detected in the expected 24-30 nt range in infected tick cells at either timepoint. As a methodological control, SFV-infected AF525 cells displayed the characteristic sharp 19 nucleotide overlap signature resulting from impaired passenger strand processing in the absence of functional Ago2, along with piRNA signatures in the 24-30 nucleotide range with 10 nucleotide overlaps, as previously documented[11] (Supplementary Fig. 3). This contrasts with the broader overlap patterns observed in tick cells, confirming the host species-specific nature of antiviral RNA processing mechanisms.

Viral RNA silencing suppressor partially rescues replication-defective SFTSV in tick cells.

The findings above suggested SFTSV NSs functions as an RNA silencing suppressor in tick cells. To test this hypothesis, we engineered a recombinant delNSs SFTSV expressing the tomato bushy stunt virus (TBSV) p19 protein, a well-characterized RNA silencing suppressor that sequesters siRNAs to counteract antiviral RNA interference[38,39]. Both delNSs-p19V5 and delNSs-eGFP control viruses replicated efficiently in mammalian Vero cells, confirming viral viability and demonstrating again that NSs is dispensable for replication in mammalian systems (Supplementary Fig. 4A,B). In contrast, these viruses showed markedly different phenotypes in tick cells. Wild-type SFTSV established robust infection with abundant viral N protein foci in BME/CTVM6 cells (Fig. 4A), while delNSs-eGFP showed no detectable infection by immunofluorescence (Fig. 4B) and no viral proteins by Western blot (Fig. 4C). Critically, functional complementation with p19 partially rescued the replication-defective phenotype, enabling viral persistence and limited local spread as evidenced by discrete N protein foci at 6 dpi (Fig. 4D,E). Western blot analysis confirmed viral N protein expression and V5-tagged p19 detection in delNSs-p19V5 infected cells (Fig. 4C). Growth kinetics revealed that delNSs-p19V5 achieved only intermediate replication levels with declining viral titers over time, indicating that viral replication can be recovered in tick cells (Fig. 4F). These complementation studies support a role for

339 NSs in counteracting tick antiviral responses. To elucidate the underlying mechanism,
340 we next investigated the biochemical properties of NSs.
341

SFTSV NSs exhibits structural features consistent with RNA-binding activity.

Given that RNA silencing suppression typically requires direct interaction with RNA
substrates, we investigated whether NSs functions as an RNA-binding protein. We
generated a high-confidence structural model of the NSs protein using AlphaFold2[40,41]
(Fig. 5A and Supplementary Fig. 5). The predicted structure revealed a compact
globular protein with multiple α-helical regions and β-sheet domains. Electrostatic
surface analysis demonstrated several distinct positively charged patches distributed
across the protein surface (Fig. 5B), a characteristic feature of nucleic acid-binding
proteins that facilitates interaction with the negatively charged phosphate backbone of
RNA. We next employed computational approaches to identify potential RNA-binding
domains (RBDs) within the NSs sequence. RBDetect analysis, which identifies RNA-
binding sites based on sequence homology to known cellular RNA-binding
sequences[42,43], revealed two discrete regions with significant similarity to established
RBDs: residues SER13-ARG22 and TRP133-THR141 (Fig. 5C; Supplementary Table
1). The probability scores for RNA-binding activity peaked at approximately 0.9 for the
first region and 0.6 for the second region, indicating strong predicted binding potential.
To validate these predictions using an independent approach, we applied BindUP
analysis, which predicts RNA-binding sites based on surface physicochemical
properties[44] (Supplementary Table 2). This analysis identified a region within α-helix 9
as having high probability for nucleic acid interaction. Notably, when the outputs from
both RBDetect and BindUP were mapped onto the AlphaFold2 structural model, the
predicted binding sites showed substantial overlap with spatial continuity (Fig. 5D),
strengthening confidence in these predictions. Importantly, all identified putative RBDs
were located within regions of high structural confidence in the AlphaFold2 model
(Supplementary Fig. 5E).
367

SFTSV NSs binds 22 nt vsiRNAs in infected cells.

To validate NSs as an RNA-binding protein, we performed immunoprecipitation of V5-
tagged p19 and NSs from recombinant virus-infected BME/CTVM6 cells at 2 dpi.
Although protein expression levels in whole cell lysates varied between replicates,
both proteins were detected in immunoprecipitated samples (Fig. 6A). Small RNA-seq
analysis of co-immunoprecipitated RNAs revealed that p19 lysates were enriched for
19-22 nt RNAs as expected[45], whereas NSs immunoprecipitates showed enrichment
for 22 nt RNAs (the characteristic products of Dicer-2 processing) which comprised
50-55% of all detected reads across three independent replicates (Fig. 6B).
Analysis of small RNA composition demonstrated that 22 nt reads accounted for 75%
of mapped reads derived primarily from SFTSV RNA in NSs immunoprecipitates, with
approximately 45% from antigenomic sense and 30% from genomic sense sequences.
In contrast, p19-associated SFTSV siRNAs were approximately 20nt in length, though
this varied between replicates (Fig. 6C; Supplementary Fig. 6). When comparing
normalized read counts across viral genome segments, both proteins showed similar
binding levels, although siRNAs derived from the SFTSV M segment were the most
reproducibly immunoprecipitated by NSs across replicates (Fig. 6D).
Genome mapping analysis revealed distinct siRNA mapping patterns between the two
proteins. The p19-associated 22 nt siRNAs mapped to limited viral genome regions

(Fig. 6E, Supplementary Fig. 7), whereas NSs-bound 22 nt RNAs mapped to multiple
reproducible hotspots distributed across all three viral segments (Fig. 6F,
Supplementary Fig. 8). Notably, hotspots derived from antigenomic sense S segment
RNA corresponded to regions within the genome with strongly predicted secondary
structures, such as the siRNA hotspot identified mapping to nt position 1210-1238 of
the antigenomic S RNA, which corresponds to the head region of a hairpin structure
(Fig. 6F; Supplementary Fig. 9).
Crucially, NSs immunoprecipitation revealed a statistically significant reduction (**p ≤
0.01) in the proportion of 22 nt siRNA duplex pairs (those with 20 nt overlaps),
decreasing from approximately 45% in total infected cell lysates to less than 10% in
NSs-bound samples. This reduction was specific to vsiRNAs, as *R. microplus*
genome-derived siRNAs showed no difference in duplex abundance (Fig. 6G). Size-
specificity analysis confirmed similar single-strand enrichment for 21 nt and 23 nt
siRNAs but not for 24 nt or 28 nt species, demonstrating that NSs selectively binds
vsiRNA single strands within a defined size range (Supplementary Fig. 10). This
sequestration likely occurs post-Dicer processing, as correctly processed vsiRNA
duplexes were robustly and reproducibly detected in infected tick cells (Fig. 3). These
findings support a model in which NSs suppresses RNAi by selectively sequestering
single-stranded 22-nt vsiRNAs, likely at a step preceding their incorporation into RISC.
The observed reduction in duplex vsiRNA abundance, combined with structural
predictions, provides strong support for this mechanism. However, direct evidence of
impaired Argonaute loading, such as through Ago2 immunoprecipitation or silencing
reporter assays, has yet to be demonstrated. Whether NSs binding fully blocks RISC
assembly or transiently delays vsiRNA incorporation to permit productive viral
replication remains to be determined. The molecular details and stoichiometric
requirements of NSs–vsiRNA interactions warrant further investigation.
413
To validate our computational predictions, we performed molecular modelling using
authentic 22-nucleotide vsiRNAs identified from NSs immunoprecipitation
(Supplementary Table 3). Docking analyses revealed multiple RNA-binding interfaces
within 4Å contact distance for vsiRNAs derived from all three SFTSV genomic
segments: S (Fig. 6H), M (Fig. 6I), and L (Fig. 6J).
Notably, the predicted RNA contact residues clustered within the same protein region
identified by both RBdetect and BindUP algorithms (Fig. 5D), providing highly
convergent evidence for the location of functional RNA-binding sites from orthogonal
computational approaches. This structural validation strongly supports the siRNA-
binding capacity of SFTSV NSs protein in tick cells and establishes a molecular
framework for understanding NSs-mediated RNAi suppression.
425
DISCUSSION
This study reveals a fundamental dichotomy in SFTSV replication strategies between
mammalian and arthropod hosts, establishing NSs as a critical RNA silencing
suppressor essential for tick cell infection. The striking observation that NSs protein is
virtually undetectable in productively-infected tick cells, despite being absolutely
required for viral replication, challenges conventional paradigms of viral protein
function and highlights the sophisticated evolutionary adaptations that enable arboviral
transmission.
The species-specific permissivity patterns observed across tick cell lines (Fig. 1A,B)
provide crucial insights into possible SFTSV vector competence. The successful
replication in *R. microplus* cells (BME/CTVM6 and BME/CTVM23) mirrors the known

epidemiological distribution of SFTSV-positive ticks, suggesting that cellular factors
governing viral entry and replication are phylogenetically conserved traits that
determine vector competence in nature. Cell lines from two other species,
Rhipicephalus decoloratus (BDE/CTVM16) and *Hyalomma anatolicum* (HAE/CTVM9)
originating from East Africa and India respectively, also supported SFTSV replication,
although the virus has not been reported from these tick species or geographic
regions, suggesting the potential for spread beyond its currently known distribution.
The sustained viral production over extended culture periods (Fig. 1C,D) confirms the
establishment of infections expected of arthropod-borne bunyaviruses[25,28,46,47]. The
failure of SFTSV to replicate in *Ixodes* spp. cell lines, despite these ticks vectoring
related phenuiviruses[48], indicates that host range determinants may operate at the
cellular level.
Here we show that SFTSV NSs protein expression is repressed in tick cells yet is
unexpectedly essential for sustained virus replication. This contrasts with mosquito-
borne bunyaviruses such as Rift Valley fever virus, where NSs repression does not
impact virus replication in mosquito cells and instead facilitates viral persistence[47,49-
51], suggesting distinct host-specific requirements for NSs in different arthropod
vectors. The paradoxical requirement for NSs in tick cells represents a remarkable
evolutionary balancing act that extends beyond recent findings in mammalian
systems[23]. Our demonstration that delNSs SFTSV fails to sustain infectious progeny
in tick cells (Fig. 1F) and showed no detectable nucleocapsid protein (Fig. 1G),
contrasts sharply with dispensable NSs function in mammalian cells[34]. Proteomic
analysis revealed differential temporal expression of SFTSV proteins in tick cells, with
structural proteins (N, GPC) and the RNA-dependent RNA polymerase (RdRp)
showing significant increases from 3 to 6 dpi, while the non-structural protein NSs
remained at consistently low levels throughout infection. The detection of NSs by mass
spectrometry despite its absence in Western blots suggests tight regulation of this
protein in infected tick cells, perhaps compounded by antbody sensitivity (Fig. 1G,H).
The paradox of NSs essentiality despite undetectable protein levels may be resolved
through several non-exclusive mechanisms. Firstly, based on established proteomics
intensity-to-concentration relationships[52,53], the observed mean peptide intensity for
SFTSV NSs (Log2 = 15) suggests protein concentrations in the low nanomolar range
(1-10 nM) range, which, while below Western blot detection limits, align with the
nanomolar binding affinities characteristic of other viral RNA silencing suppressors[54].
Nevertheless, future direct quantification of NSs:vsiRNA binding stoichiometry, and
assessment of competitive Ago2 loading inhibition *in vivo*, will be essential to
experimentally validate this proposed depletion mechanism; Second, NSs may exhibit
exceptionally high RNA-binding affinity or cooperative binding that amplifies its
suppressive capacity at low concentrations; Third, temporal or subcellular
compartmentalization may concentrate NSs activity during critical early infection
phases, when RNAi activation would otherwise prevent productive replication. The
profound replication defect observed in NSs-deficient infections (Fig. 1I,J) reinforces
the notion that even low-abundance NSs is functionally essential in tick cells. These
mechanisms may collectively explain how limited quantities of NSs achieve effective
RNAi suppression in the tick host.
482
483 The selective reduction of NSs subgenomic mRNA in tick cells provides a mechanistic
484 explanation for the protein's low abundance while maintaining its essential function.
485 Our Northern blot analysis (Fig. 1P) demonstrates that while both genomic and
486 antigenomic viral RNAs accumulate to similar levels in tick and mammalian cells, NSs

mRNA shows markedly reduced levels specifically in tick cells, suggesting a post-transcriptional regulation mechanisms unique to arthropod hosts. This regulatory strategy represents an evolutionary compromise that maximizes NSs activity during critical early infection phases while minimizing detection by host surveillance mechanisms.

The demonstration that SFTSV infection triggers robust small RNA production in tick cells (Fig. 2A) establishes RNAi as a primary antiviral defence mechanism. Our beta elimination analyses definitively prove that the observed 22-nucleotide small RNAs represent authentic virus-derived siRNAs rather than degradation products (Fig. 2B-E). This finding significantly extends previous work by Xu *et al.*, who demonstrated vsiRNA production during SFTSV infection of *Haemaphysalis longicornis* through blood feeding[22]. Our study advances this understanding by providing detailed mechanistic insights into vsiRNA processing, demonstrating temporal dynamics of the RNAi response, and establishing the functional consequences for viral replication in cell culture systems.

The emergence of methylated vsiRNAs at 6 dpi (Fig. 2E), which are absent at 3 dpi, supports a model wherein the host RNAi machinery eventually overcomes transient viral suppression. Given that NSs is expressed at low levels in infected tick cells (undetectable by Western blot; Fig. 1G), complete sequestration of all vsiRNAs would present a significant stoichiometric challenge. This constraint is further amplified by the temporal dynamics of antiviral RNAi responses, as vsiRNA accumulation increases substantially between 3 and 6 dpi. The appearance of RISC-loaded vsiRNAs at 6 dpi therefore indicates that sustained RNAi pathway activation can ultimately saturate the suppressive capacity of this low-abundance viral suppressor, enabling functional antiviral silencing despite initial evasion mechanisms.

Our observations support a 'depletion strategy' whereby viral suppressors of RNAi achieve functional suppression without requiring abundant expression. To address potential stoichiometric limitations, we propose that NSs operates through high-affinity binding kinetics rather than simple mass action. Assuming dissociation constants comparable to characterized VSRs (Kd ~0.1-1 nM[54]) and estimated cellular vsiRNA concentrations of 10-100 nM during peak infection, even low-abundance NSs could effectively compete for substrate binding. Additionally, preferential sequestration of newly processed single-stranded guide RNAs before RISC loading would be more efficient than competing with mature RISC complexes, explaining how limited NSs quantities can achieve functional suppression despite abundant total vsiRNA pools. Such high-affinity interactions are characteristic of other viral RNAi suppressors, including tombusvirus p19, which functions at substoichiometric concentrations (Kd ~ 0.17 nM) and achieves >90% siRNA depletion despite limited protein abundance[45].

The progressive amplification of vsiRNA accumulation across all viral segments between 3- and 6-dpi (Fig. 2F-H) indicates that tick cells mount increasingly robust antiviral responses over time. The concentrated targeting within the unencapsidated S segment intergenic region[55] and NSs open reading frame provides direct mechanistic evidence for RNAi-mediated NSs mRNA degradation, explaining the reduced NSs protein levels observed. This segment-specific targeting pattern suggests that viral RNA secondary structures or transcript abundance influence RNAi machinery accessibility, with implications for understanding viral evolutionary constraints.

The demonstration of canonical Dicer-2-mediated vsiRNA production through strand
overlap analysis (Fig. 3) confirms that tick antiviral RNAi pathways follow established
arthropod paradigms. The predominance of 20-22 nucleotide products with
characteristic 2-nucleotide 3' overhangs indicates uniform processing across all viral
segments[8]. Notably, the absence of detectable vpiRNA responses contrasts with
mosquito antiviral systems[46,56-59], suggesting lineage-specific adaptations in arthropod
small RNA-mediated immunity. However, the complete absence of vpiRNA production
warrants further mechanistic investigation to determine whether this reflects genuine
pathway absence or differential regulation.
546
The successful complementation of NSs-deficient virus with the heterologous p19
suppressor (Fig. 4) provides strong evidence that NSs functions as an RNAi
suppressor in tick cells. The partially restored viral protein expression and intermediate
replication kinetics in delNSs-p19V5 infected cells demonstrate functional
complementation while revealing that NSs possesses unique properties beyond
simple siRNA sequestration. This incomplete complementation suggests NSs may: (i)
exhibit specific binding preferences for SFTSV-derived vsiRNAs through evolved
sequence or structural recognition motifs, (ii) interact with additional host factors
beyond the core RNAi machinery, or (iii) function through temporal coordination with
other viral proteins during the replication cycle. The heterologous p19 suppressor,
while functionally competent, lacks these virus-specific adaptations that have co-
evolved to optimise NSs performance within the SFTSV life cycle.
559
Our integrated structural and functional analyses establish NSs as a novel siRNA-
binding protein that selectively sequesters 22 nt vsiRNAs through size-specific
molecular recognition. This finding extends the established paradigm of bunyaviral
RNA silencing suppression beyond the well-characterized tospoviral NSs proteins,
which bind both siRNAs and miRNAs with varying specificity depending on
phylogenetic origin; American clade tospoviruses exhibit broad-spectrum binding to
both long and short dsRNA molecules, whereas Eurasian clade NSs proteins
demonstrate restricted binding to short dsRNA substrates[20,60,61]. Notably, while RNA
silencing suppressor activity has not been clearly demonstrated for NSs proteins from
vertebrate-infecting bunyaviruses in either vertebrate hosts or arthropod vectors[20], our
results provide definitive mechanistic evidence for this function. Javaid et al.[23] recently
demonstrated SFTSV NSs interacts with AGO2 to regulate RNAi in mammalian cells.
However, our work provides the first direct evidence that NSs functions as an essential
viral suppressor of RNAi specifically in arthropod cells, revealing distinct mechanistic
requirements across host systems.
While our immunoprecipitation and computational modeling provide strong evidence
for NSs-vsiRNA interactions, quantitative binding parameters remain to be
determined. The inherent challenges associated with NSs biochemical
characterization, including insolubility during bacterial expression[24,62] and extensive
C-terminal disorder that complicates *in vitro* folding, have limited direct binding studies.
Despite these technical constraints, our cellular interaction data and structural
predictions provide compelling evidence for the RNA-binding capacity and size-
specific vsiRNA recognition observed in infected tick cells.
583
584 The convergence of computational predictions, AlphaFold2 structural modeling (Fig.
585 5), and experimental validation reveals that NSs functions through single-strand siRNA
586 sequestration, with discrete binding hotspots that suggest recognition of specific viral

RNA secondary structural motifs (Fig. 6, Supplementary Fig. 9). This size-selective mechanism operates at a critical temporal junction, post-Dicer processing, but preceding RISC loading, effectively intercepting single-stranded guide RNAs before Ago2 incorporation. This represents a strategically optimal intervention point that preemptively neutralizes host antiviral responses by depleting the cellular pool of functional guide RNAs. While our data strongly support NSs-mediated sequestration of siRNAs upstream of RISC incorporation, future studies using Ago2 immunoprecipitation or RNAi reporter assays will be needed to directly confirm impaired Argonaute loading.

Importantly, our overlap analysis coupled with molecular docking experiments unambiguously confirms the single-stranded nature of the bound RNA substrates, identifying a previously unrecognized bunyaviral mechanism for RNAi suppression that diverges from canonical dsRNA-binding suppressors. The molecular docking validation using authentic vsiRNA sequences derived from viral infections provides direct structural confirmation and establishes a comprehensive molecular framework for understanding NSs-mediated RNAi evasion. This mechanistic insight represents a substantial functional advancement beyond the previous protein-protein interaction studies (NSs/Ago2) conducted in mammalian cell systems.

Our findings highlight a key evolutionary adaptation in tick-borne phenuiviruses: the requirement for RNAi suppression in arthropod vectors, but not mammalian hosts. This host-specific dependency likely constrains viral evolution and presents an opportunity for vector-targeted antiviral strategies. The identification of NSs as a siRNA-sequestering protein, rather than an upstream pathway inhibitor, reveals a novel mechanism of immune evasion and offers tractable targets for intervention. Disrupting NSs/vsiRNA interactions or enhancing guide RNA stability could block viral replication in ticks without affecting mammalian infection. While our findings are derived from established tick cell line models, *in vivo* validation in whole *H. logicornis* or *R. microplus* spp. ticks will be essential to confirm the physiological relevance of NSs-mediated RNAi suppression during natural infection. These insights establish NSs as a critical determinant of SFTSV vector competence and uncover a previously unrecognised function for this viral protein.

 METHODS

 Experimental model and subject detail

Cell culture
Vero E6 cells[63] were cultured in DMEM (Gibco) supplemented with 10% foetal bovine
serum (FBS) (Gibco). HuH7-Lunet-T7 cells[64], which stably express T7 RNA
polymerase were obtained from R. Bartenschlager, Heidelberg University and were
grown in DMEM (Gibco) supplemented with 2 mM L-glutamine, nonessential amino
acids, and 10% FBS (Gibco). All mammalian cells were cultured at 37 °C with 5% CO_2.
The tick cell lines used in this study were *Amblyomma americanum* AAE2[65] and
AAE12[66], *Amblyomma variegatum* AVL/CTVM17[67], *R. microplus* BME/CTVM2[67],
BME/CTVM6[67], and BME/CTVM23[68], *R. decoloratus* BDE/CTVM16[67], *H. anatolicum*
HAE/CTVM9[69], *Ixodes scapularis* ISE6[70],; *Ixodes ricinus* IRE/CTVM19 and
IRE/CTVM20[71] and *Rhipicephalus sanguineus* RML-RSE[72], sourced from the Tick Cell
Biobank at the University of Liverpool. The tick cell lines were grown at either 28 °C or
32 °C in sealed flat-sided culture tubes (Nunc) containing 2 ml of appropriate complete
culture medium (L-15, L-15B, L-15B300, L-15/MEM, L-15/H-Lac, L-15/L-15B or L-
15/H-lac/L-15B)[67,73,74] as described previously for maintenance, and transferred to
sealed T25 or T80 non-vented flasks to bulk for experimentation. Aag2-AF525 cells,
an Ago2 knockout cell line derived from Aag2-AF5 (*Aedes aegypti* mosquito; European
Collection of Cell Cultures (ECACC): 19022601) cells by CRISPR/Cas9[11,75] were
maintained in Leibovitz's L-15 medium supplemented with 10% FBS (Gibco), 10%
tryptose phosphate broth (Gibco), and penicillin-streptomycin (final concentration 100
units/ml, 100 µg/ml; Gibco) and maintained at 28 °C.

Plasmids
Plasmids for the recovery of SFTSV have been described previously[24,34]. The
expression plasmids pTM1-HB29L and pTM1-HB29N contain the SFTSV HB29 L and
N ORFs under control of the T7 promoter and encephalomyocarditis virus internal
ribosome entry site sequence. The rescue plasmids pTVT7-HB29S, pTVT7-HB29M,
and pTVT7-HB29L contain full-length cDNAs of the SFTSV strain HB29 antigenome
segments flanked by T7 promoter and hepatitis delta ribozyme sequences. Plasmid
pTVT7-HB29SNSsV5 encodes an S segment expressing NSs with a C-terminal V5
epitope tag (GKPIPNPLLGLDST). Plasmid pTVT7-HB29SdelNSs:TBSVp19V5
encodes an S segment in which the NSs ORF is replaced with the tomato bushy stunt
virus p19 protein, also with a C-terminal V5 epitope tag (TBSV; Accession No.
M21958.1).

Generation of recombinant viruses from cDNA
Recombinant SFTSV were generated by transfecting HuH7-Lunet-T7 cells with 0.1 µg
pTM1-HB29L, 0.5 µg pTM1-HB29N, and 1 µg each of pTVT7-based plasmids
expressing wt or recombinant viral antigenomic segments. Transfections used 3 µl
TransIT-LT1 (Mirus Bio LLC) per µg of DNA as the transfection reagent. After 5 days,
transfection supernatants were collected, clarified by low-speed centrifugation, and
stored at -80°C. Recombinant virus stocks were amplified in Vero E6 cells at 37 °C by
infection at an MOI of 0.01, with culture medium harvested at 7 dpi. The genome
segments of recovered viruses were amplified by RT-PCR and sequenced to confirm
that no mutations had occurred during the rescue process.

Virus Production

SFTSV strains (rHB29pp [wt SFTSV][24] or recombinant viruses rHB2912aaNSs [delNSs][34], rHB29delNSsGFP [delNSs eGFP][34], rHB29NSsV5 [NSsV5] and rHB29delNSsp19V5 [p19V5]) were generated using reverse genetics technologies. The Hubei 29 (HB29) strain is based on a plaque-purified stock called Hubei 29pp (HB29pp) provided by Amy Lambert (CDC Arbovirus Diseases Branch, Fort Collins, CO). Working stocks of SFTSV were generated in Vero E6 cells by infecting at a low multiplicity of infection and harvesting the cell culture medium 7 dpi. Recombinant virus stocks were confirmed by Sanger sequencing to ensure no mutations had occurred during propagation. Previous data have demonstrated that recombinant viruses have the same replication kinetics and viral protein expression profiles as the wt/parental viruses in time course of infection studies[24,34].

Virus titration by immunofocus assays

Virus titres were determined by focus-forming assays in Vero E6 cells. Confluent Vero E6 cells were infected with 10-fold serial dilutions of virus prepared in phosphate-buffered saline containing 2% FBS and incubated for 1 h at 37 °C. Following infection, cells were overlaid with MEM supplemented with 2% FBS and 0.6% Avicel (FMC Biopolymer). Cells were incubated for 6 days before fixation and immunostaining as described previously[24].

RNA extraction from cell cultures

Total RNA was extracted from cultured cells using 1 ml TRIzol reagent (Thermo Fisher Scientific) according to the manufacturer's protocol. RNA concentrations were measured using a NanoDrop spectrophotometer, and RNA was stored at -80 °C until use.

Western blotting

Cell lysates were prepared at indicated timepoints by adding 300 μl lysis buffer (100 mM Tris-HCl pH 6.8, 4% SDS, 20% glycerol, 200 mM DTT, 0.2% bromophenol blue, and 25 U/ml Pierce Universal Nuclease [Thermo Fisher]). Proteins were separated on 4-12% SDS-polyacrylamide gradient gels (Invitrogen) and transferred to Hybond-C Extra membranes (Amersham). Membranes were blocked in saturation buffer (PBS containing 5% dry milk and 0.1% Tween 20) for 1 h at room temperature. Primary antibodies included SFTSV anti-N or anti-NSs polyclonal rabbit antibodies[24], anti-tubulin monoclonal antibody (Sigma), or anti-V5 antibody (Abcam). After washing, membranes were incubated with goat anti-rabbit DyLight 680 or goat anti-mouse DyLight 800 secondary antibodies (Thermo Fisher). Proteins were visualized using a Li-Cor Odyssey CLx infrared imaging system. Data shown in the manuscript are representative gels from 2-3 independent replicates.

Immunofluorescence microscopy

BME/CTVM6 cells previously seeded onto 13mm coverslips for 24 hours were infected with wt or recombinant SFTS viruses at indicated MOIs. At defined timepoints post-infection, cells were fixed with 8% paraformaldehyde in PBS for 15 min at room temperature, permeabilized with 0.5% Triton X-100 for 10 min and blocked with 4% milk in PBS for 1 h. Cells were incubated with primary antibodies overnight at 4 °C, washed 3x with PBS, followed by staining with Alexa Fluor-conjugated secondary

717　antibodies for 1 h at room temperature. Nuclei were counterstained with DAPI (1
718　µg/ml). Images were acquired using a Zeiss LSM 880 confocal microscope and
719　number of nuclei and infection rates quantified using ImageJ (version 1.52p).
720

Northern blotting

BME/CTVM6 or Vero E6 cells were infected with wt SFTSV, and total cellular RNA was
extracted at 72 h post-infection using TRIzol reagent (Invitrogen). RNA samples (1 µg)
were electrophoresed through 1.2% agarose gels in TAE buffer[76] and transferred via
capillary action to positively charged nylon membranes (Roche). Membranes were
hybridized with digoxigenin-labelled RNA probes (150 ng each) complementary to
negative-sense genomic RNA, positive-sense antigenomic RNA, or subgenomic
mRNA sequences encoding N or NSs proteins. Detection was performed using a DIG
Northern Starter Kit (Roche) according to the manufacturer's instructions. Data shown
in the manuscript are representative gels from 2-3 independent replicates.
731

Small RNA sequencing and library preparation

Total RNA (1 µg per sample) was extracted and processed using the BGI Genomics
Small RNA Library Construction Protocol for DNBSEQ platform (DNBSeq, UMI small
library, SE50). Following sequential 3' and 5' adaptor ligation, reverse transcription,
and PCR amplification (16–17 cycles), libraries were PAGE-purified, circularized to
single-stranded DNA, and quality-assessed using an Agilent Bioanalyzer. Sequencing
was performed on a DNBSEQ-G400 instrument with 50 bp single-end reads.
739

Small RNA methylation analysis

Methylation status of small RNAs was determined using a beta-elimination assay as
previously described[37]. BME/CTVM6 cells (2.5×10^5 cells/cm^2) were infected with wt
SFTSV (MOI: 0.1) or mock-infected for 3 or 6 days. AF525 cells infected with SFV
(MOI: 10, 24 h) served as positive controls for 2'-O-methylation. Following TRIzol LS
extraction, RNA samples (200 µl) were treated with 5 µl 20x borate buffer (pH 8.6) and
either 12.5 µl 200 mM sodium periodate or nuclease-free water (controls), incubated
for 15 min at room temperature, then treated with 10 µl glycerol for an additional 15
min. RNA was precipitated overnight at -20 °C using GlycoBlue coprecipitant, 3 M
sodium acetate (pH 5.2), and absolute ethanol, recovered by centrifugation (16,000 x
g, 15 min, 4 °C), washed with 70% ethanol, and resuspended in 100 µL sodium borate
buffer (55 mM, pH 9.5). RNA was purified using the Monarch RNA Cleanup Kit and
sequenced by BGI Tech Solutions using DNBSEQ UMI technology (minimum 500 ng
input, 20 million clean reads per sample).
754

Small RNA immunoprecipitation assay and RNA extraction

BME/CTVM6 cells were infected with either NSsV5- or p19V5- expressing SFTS
viruses (MOI: 1) and cell lysates prepared at 2 dpi to assess viral protein-RNA
interactions. Immunoprecipitation was performed using Protein G Dynabeads (Thermo
Fisher). Lysis buffer contained 20 mM Tris-HCl pH 7.5, 150 mM NaCl, 5 mM MgCl$_2$,
0.5% NP-40, and protease/phosphatase inhibitors (Roche). Wash buffer consisted of
50 mM Tris-HCl pH 7.5, 200 mM NaCl, 1 mM EDTA, and 1% NP-40. Dynabeads (20
µL) were washed three times, conjugated with 5 µg anti-V5 antibody (Abcam) for 1 h
at 4°C, then washed again. Cells were lysed in 100 µL lysis buffer per 10^6 cells for 20

min on ice, clarified by centrifugation (16,000 x g, 20 min), and immunoprecipitated
overnight at 4 °C with rotation. After three washes, proteins were eluted in LDS buffer
at 90 °C for 10 min. For RNA extraction, eluates were treated with proteinase K (1 µg,
1 h, 50 °C; Sigma) then processed with TRIzol following standard phenol-chloroform
extraction. RNA was precipitated with isopropanol, washed with 75% ethanol, and
resuspended in nuclease-free water.
770

Small RNA computational analysis

Small RNA bioinformatics pipeline
Small RNA libraries (Supplementary Table 4 and 5) were sequenced and processed
using a comprehensive bioinformatics pipeline described on the Zenodo repository.
Raw sequencing reads were quality-filtered and adapter-trimmed using fastp v0.23[77,78]
(1_small_RNA_fastp_processor.sh) with small RNA-optimized parameters: length
filtering (18-32 nt), Phred quality score threshold Q20, unqualified base percentage
limit 40%, maximum 2 N bases per read, tail trimming with window size 1, poly-G and
poly-X trimming enabled, and overrepresentation analysis activated. Adapter
sequences were auto detected. FASTA files were standardized and formatted for
optimal analysis (2_fasta_formatter.sh).
782

Read alignment and quality control
Reference genome indices were constructed using Bowtie v1.3.1[79,80]
(3_bowtie_index_builder.sh), and trimmed reads were aligned to viral (SFTSV HB29
strain) and host (*R. microplus* GCA_002176555.1) reference genomes
(4_recursive_bowtie_alignment_pipeline_outputs_standspecificBAM.sh,
4a_bowtie_stringent_smallRNA_aligner.sh) with stringent parameters: zero
mismatches allowed (-v 0), maximum 5 mapping locations per read (-m 5), best
alignment reporting (--best --strata), and exhaustive search (--tryhard). Post-alignment
filtering retained only uniquely mapping reads with MAPQ ≥20, excluding unmapped
(SAM flag 4), secondary (flag 256), and supplementary (flag 2048) alignments. BAM
files were strand-separated based on SAM flag 16 (reverse complement) using
SAMtools v1.15[81]. Quality filtering statistics and mapping efficiencies were tracked
throughout the pipeline to ensure data integrity.
796

Small RNA sequence analysis and characterization
Read length distributions and nucleotide composition analysis were performed using
custom scripts (5_Length_Distribution_Analysis.sh) and
(5a_strand_specific_Length_Distribution_Analysis.sh) calculating first- and last-base
frequencies (A, T/U, G, C percentages) across read lengths 1-50 nt individually, with
reads >50 nt grouped. Data transformation for visualization was conducted using R
scripts (5_for_generating_length_vs_nt_pc_graph_data.R,
5a_for_generating_strand_specific_length_vs_nt_pc_graph_data.R). Chromosome-
specific read counting was performed across all BAM files
(6_bam_chromosome_read_counter.sh).
Coverage analysis employed a custom Python pipeline
(7_rpm_normalized_coverage_tool.py) utilizing pysam v0.19.1[82] with CIGAR-aware
position counting (equivalent to bedtools -split behaviour) and RPM normalization
based on total input reads from original FASTQ files rather than mapped reads. The
most abundant small RNA sequences were identified and characterized
(9_top_mapping_small_rna_finder.py).

813
Ping-pong amplification analysis

Ping-pong amplification signatures (for vpiRNAs) were detected using a custom algorithm (8_pingpong_smallRNA_analyzer.py) analysing overlapping antisense small RNA pairs within a 1-19 nucleotide window, requiring complementary reads of 23-29 nt length with ≥20 nt overlap on opposite strands. The algorithm calculated Z-scores for overlap frequencies and identified statistically significant ping-pong pairs using both raw counts and probability-based H-signatures. Data consolidation across samples was performed (8a_ping_pong_data_consolidator.R) and formatted for analysis (8b_ping_pong_matrix_formatter.R).

Small RNA classification

Comprehensive small RNA characterization was conducted using an integrated pipeline (10_small_rna_analysis.py) with configuration management (10a_small_rna_config.yaml, 10b_setup_and_run.sh) that classified small RNA populations as miRNA-like (20-24 nt), siRNA-like (20-25 nt), or piRNA-like (24-32 nt) based on size distribution profiles. All analyses maintained strand-specific resolution with forward strand values reported as positive and reverse strand as negative for visualization.

AlphaFold prediction of SFTSV NSs

The sequence of SFTSV NSs (Hubei-29; GenBank ID: AJD86041) was used to query ColabFold's AlphaFold2 structure prediction algorithm using the advanced beta notebook[40,41]. Structure prediction was performed using MMseqs2 for multiple sequence alignment generation, with a maximum MSA depth of 512:1024 sequences and MSA subsampling enabled. Five models were generated with two random seeds each, using 3 recycle iterations and pTM scoring enabled. The ten resulting models were ranked by pLDDT scores, with the highest-ranking model (rank_1_model_5_ptm_seed_0; pLDDT: 72.77, pTMscore: 0.7104) selected for further structural analysis (Supplementary Fig. 5). Subsequent structural analysis was performed using UCSF ChimeraX[83].

SFTSV NSs RNA binding prediction

RNA binding prediction for the SFTSV NSs protein sequence was performed using RBDetect[42], a machine learning model trained by Shrinkage Discriminant Analysis (SDA) with a dataset of 8,891 experimentally identified polypeptides from RBDmap experiments, using positive examples (RNA-bound polypeptides) and negative examples (RNA-released polypeptides)[43]. For each amino acid position in the NSs protein sequence, RBDetect assigns a probability value for RNA binding based on the fragment centred at that position. A Hidden Markov model was then used to visualize the probabilities sequentially, determining the most probable RNA binding regions across the protein structure. Predicted RNA binding sites (Supplementary Table 1) were mapped onto the AlphaFold-predicted NSs structure to identify potential functional domains. We also utilised BindUP[84], a computational tool that predicts RNA-binding regions in protein sequences using physicochemical and structural features rather than sequence homology. The tool exploits characteristic surface properties of RNA-binding proteins (RBPs), including electrostatic potential, hydrophobicity, and shape complementarity, to distinguish them from non-RBPs. BindUP employs a

machine-learning framework trained on curated RBP and non-RBP datasets. The software computes surface patches on protein structures (experimental or predicted) and scores their likelihood of mediating RNA interactions[84] (Supplementary Table 2).

RNA secondary structure prediction

RNA secondary structures were predicted using the forna web server (http://rna.tbi.univie.ac.at/forna/, accessed May 2025) powered by the ViennaRNA package[85]. The 76-nucleotide RNA sequence derived from nucleotides 1191-1266 of the antigenomic HB29 S RNA (GenBank: KP202165) was subjected to minimum free energy folding calculations using Turner energy parameters at 37°C. The predicted secondary structure with the lowest free energy was selected for analysis and visualization. No folding constraints were applied during the prediction process.

Modelling of RNA binding to SFTSV NSs

Structural modelling of SFTSV NSs-vsiRNA interactions was performed using AlphaFold3. The SFTSV NSs protein sequence (Hubei-29; GenBank ID: AJD86041) was submitted to the AlphaFold3 webserver (version 2.0) with representative 22-nucleotide vsiRNA sequences identified from NSs immunoprecipitation experiments in SFTSV-infected BME/CTVM6 tick cells. vsiRNA sequences were selected based on abundance in NSs-IP samples from identified hotspots within each viral RNA segment (Supplementary Table 3). Five independent structural predictions were generated for each NSs-vsiRNA complex using default parameters. Models were selected based on per-residue confidence scores (pLDDT >70 for NSs functional domains) and interface confidence scores (>0.5 for protein-RNA contacts). Structural analysis was performed using UCSF ChimeraX (version 1.6)[83], focusing on RNA-binding interfaces, contact residues within 4 Å of vsiRNA substrates, and structural determinants of vsiRNA recognition (Supplementary Table 6).

Proteomic analysis

To estimate relative abundance of SFTSV proteins, we analyzed proteomic data from project PXD054068, which comprises whole-cell proteomes of BME/CTVM6 tick cells infected with SFTSV (MOI 1) at 3- and 6-dpi with four biological replicates. Corresponding peptides for each viral protein [non-structural protein (NSs; AJD86041), nucleocapsid protein (N; AJD86040), glycoprotein precursor (GPC; AJD86039) or RNA-dependent RNA polymerase (RdRp; AJD86038)] were pooled across replicates at each time point, and peptide intensity distributions were analyzed. Scripts used to generate plots and statistics are available from the Zenodo online repository (https://doi.org/10.5281/zenodo.15877198). Statistical significance of changes in peptide intensities between time points was assessed using Welch's t-test.

Biosafety

All experiments with infectious SFTSV were performed in the Richard Elliott Biosafety Laboratories (REBL) at the MRC-University of Glasgow Centre for Virus Research, under containment level 3 (CL-3) conditions, approved by the UK Health and Safety Executive (GM223/20.2a).

Quantification and Statistical Analysis

Experiments were performed with a minimum of n=2 biological replicates unless otherwise stated, with each biological sample tested in technical replicates where applicable. Technical replicates were averaged before statistical analysis. Data are presented as mean ± SD e.g. for virus titres and cell counts. For immunofluorescence quantification, n=3 randomised fields of view were analysed per condition. Small RNA sequencing experiments used n=3 biologically independent samples. Statistical significance for multiple group comparisons was assessed using Brown-Forsythe and Welch's ANOVA with Dunnett's T3 multiple comparison test. Statistical significance was set at $p \leq 0.05$, with * $p \leq 0.05$, ** $p \leq 0.01$; 'ns' indicates not significant. Individual sample sizes are indicated in figure legends. Statistical analyses were performed using GraphPad Prism Version 10.5.0 (673).

ACKNOWLEDGEMENTS

We thank Prof. Ulrike Munderloh, University of Minnesota, for permission to use the ISE6, AAE2 and AAE12 cell lines. We also thank Dr Matthew Arnold (Royal Veterinary College) for support with the initial AlphaFold modelling.

FUNDING

This research was funded by a Wellcome Trust/Royal Society Sir Henry Dale Fellowship (210462/Z/18/Z) awarded to B.B. (supporting M.F., M.McF., K.D., A.T.C and B.B.). The facilities used in this study were funded by the UK Medical Research Council (MC_UU_00034/4, MC_UU_00034/8). R.H.P. was supported by a Medical Research Future Fund, Early to Mid-Career Researcher Grant MRF2022950. L.B.S. was supported by the UK BBSRC grant no. BB/P024270/1 and the Wellcome Trust grant no. 223743/Z/21/Z. M.J.P. was funded by European Union's Horizon 2020 Research and Innovation Program under the Marie Skłodowska-Curie grant agreement No 890970. E.S. was funded by German Centre for Infection Research (DZIF) (TTU 01.708). A.C. is funded by the European Research Council (ERC) Consolidator Grant 'vRNP-capture' 101001634 and the MRC grants MR/R021562/1 and MC_UU_00034/2 (supporting R.A & W.K.). A.K., was funded by the UK Medical Research Council (MC_UU_12014/8, MC_UU_00034/4). This research was funded in whole or in part by the Wellcome Trust.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

AUTHORS' CONTRIBUTIONS

Conceptualization: M.F., M.McF., E.S., A.C., A.K., B.B., Data Curation: M.F., M.McF., R.H.P., R.A., A.T.C., M.P., B.B., Formal Analysis: M.F., M.McF., R.H.P., R.A., A.T.C., W.K., M.P., A.C., A.K., B.B., Funding Acquisition: B.B., Investigation: M.F., M.McF., R.H.P., R.A., A.T.C., W.K., K.D., A.C., A.K., B.B., Methodology: M.F., M.McF., R.H.P., M.P., E.S., B.B., Project administration: B.B., Resources: L.B.S., B.B., Software: R.H.P., B.B., Supervision: A.C., A.K., B.B., Validation: M.F., M.McF., R.H.P., R.A., A.T.C., W.K., K.D., M.P., E.S., A.C., A.K., B.B., Visualization: M.F., M.McF., R.H.P.,

955 R.A., A.T.C., W.K., B.B., Writing – original draft: M.F., M.McF., A.K., B.B., Writing –
956 review & editing: M.F., M.McF., R.H.P., R.A., A.T.C., W.K., K.D., L.B.S., M.P., E.S.,
957 A.C., A.K., B.B.
958
959 **DECLARATION OF INTERESTS**
960
961 The authors declare no competing interests.
962
963 **DATA AVAILABILITY**
964
All data that support the findings of this study are openly available from Enlighten
Research Data (https://doi.org/10.5525/gla.researchdata.2028). All sequencing data
generated in this study have been deposited in the NCBI Sequence Read Archive
under BioProject accession number PRJNA1283909. The bioinformatic scripts used
for analysis can be found in the Zenodo online repository
(https://doi.org/10.5281/zenodo.15877198).
971

REFERENCES

Yu, X. J. *et al.* Fever with thrombocytopenia associated with a novel bunyavirus in China. *N Engl J Med* **364**, 1523-1532 (2011). https://doi.org:10.1056/NEJMoa1010095

Casel, M. A., Park, S. J. & Choi, Y. K. Severe fever with thrombocytopenia syndrome virus: emerging novel phlebovirus and their control strategy. *Exp Mol Med* **53**, 713-722 (2021). https://doi.org:10.1038/s12276-021-00610-1

Li, J. C., Zhao, J., Li, H., Fang, L. Q. & Liu, W. Epidemiology, clinical characteristics, and treatment of severe fever with thrombocytopenia syndrome. *Infect Med (Beijing)* **1**, 40-49 (2022). https://doi.org:10.1016/j.imj.2021.10.001

Jang, H. *et al.* Seasonal dynamics of Haemaphysalis tick species as SFTSV vectors in South Korea. *Microbiol Spectr* **12**, e0048924 (2024). https://doi.org:10.1128/spectrum.00489-24

Hu, Y. Y. *et al.* Role of three tick species in the maintenance and transmission of Severe Fever with Thrombocytopenia Syndrome Virus. *PLoS Negl Trop Dis* **14**, e0008368 (2020). https://doi.org:10.1371/journal.pntd.0008368

Pérez, L. J., Baele, G., Hong, S. L., Cloherty, G. A. & Berg, M. G. Ecological Changes Exacerbating the Spread of Invasive Ticks has Driven the Dispersal of Severe Fever with Thrombocytopenia Syndrome Virus Throughout Southeast Asia. *Mol Biol Evol* **41** (2024). https://doi.org:10.1093/molbev/msae173

Barry, G. *et al.* Gene silencing in tick cell lines using small interfering or long double-stranded RNA. *Exp Appl Acarol* **59**, 319-338 (2013). https://doi.org:10.1007/s10493-012-9598-x

Schnettler, E. *et al.* Induction and suppression of tick cell antiviral RNAi responses by tick-borne flaviviruses. *Nucleic Acids Res* **42**, 9436-9446 (2014). https://doi.org:10.1093/nar/gku657

Hoffmann, H. H., Schneider, W. M. & Rice, C. M. Interferons and viruses: an evolutionary arms race of molecular interactions. *Trends Immunol* **36**, 124-138 (2015). https://doi.org:10.1016/j.it.2015.01.004

Gammon, D. B. & Mello, C. C. RNA interference-mediated antiviral defense in insects. *Curr Opin Insect Sci* **8**, 111-120 (2015). https://doi.org:10.1016/j.cois.2015.01.006

Scherer, C. *et al.* An Aedes aegypti-Derived Ago2 Knockout Cell Line to Investigate Arbovirus Infections. *Viruses* **13** (2021). https://doi.org:10.3390/v13061066

Horwich, M. D. *et al.* The Drosophila RNA methyltransferase, DmHen1, modifies germline piRNAs and single-stranded siRNAs in RISC. *Curr Biol* **17**, 1265-1272 (2007). https://doi.org:10.1016/j.cub.2007.06.030

Yang, Z., Vilkaitis, G., Yu, B., Klimasauskas, S. & Chen, X. Approaches for studying microRNA and small interfering RNA methylation in vitro and in vivo. *Methods Enzymol* **427**, 139-154 (2007). https://doi.org:10.1016/S0076-6879(07)27008-9

Alefelder, S., Patel, B. K. & Eckstein, F. Incorporation of terminal phosphorothioates into oligonucleotides. *Nucleic Acids Res* **26**, 4983-4988 (1998). https://doi.org:10.1093/nar/26.21.4983

15 Elmer, K., Helfer, S., Mirkovic-Hosle, M. & Forstemann, K. Analysis of endo-
siRNAs in Drosophila. *Methods Mol Biol* **1173**, 33-49 (2014).
https://doi.org/10.1007/978-1-4939-0931-5_4
16 Blair, C. D. & Olson, K. E. The role of RNA interference (RNAi) in arbovirus-
vector interactions. *Viruses* **7**, 820-843 (2015).
https://doi.org/10.3390/v7020820
17 Wilson, A. *et al.* Uukuniemi virus infection causes a pervasive remodelling of
the RNA-binding proteome in tick cells. *PLoS Pathog* **21**, e1013393 (2025).
https://doi.org/10.1371/journal.ppat.1013393
18 Song, L. *et al.* Silencing suppressors: viral weapons for countering host cell
defenses. *Protein Cell* **2**, 273-281 (2011). https://doi.org/10.1007/s13238-011-
1037-y
19 van Mierlo, J. T. *et al.* Novel Drosophila viruses encode host-specific
suppressors of RNAi. *PLoS Pathog* **10**, e1004256 (2014).
https://doi.org/10.1371/journal.ppat.1004256
20 Hedil, M. & Kormelink, R. Viral RNA Silencing Suppression: The Enigma of
Bunyavirus NSs Proteins. *Viruses* **8** (2016). https://doi.org/10.3390/v8070208
21 Blakqori, G. *et al.* La Crosse bunyavirus nonstructural protein NSs serves to
suppress the type I interferon system of mammalian hosts. *J Virol* **81**, 4991-
4999 (2007). https://doi.org/10.1128/jvi.01933-06
22 Xu, Y. *et al.* Antiviral RNA interference in disease vector (Asian longhorned)
ticks. *PLoS Pathog* **17**, e1010119 (2021).
https://doi.org/10.1371/journal.ppat.1010119
23 Javaid, N., Jang, T. W., Fu, Y. & Choi, Y. SFTSV NSs interacts with AGO2 to
regulate the RNAi pathway for viral replication. *J Virol* **99**, e0220524 (2025).
https://doi.org/10.1128/jvi.02205-24
24 Brennan, B. *et al.* Reverse genetics system for severe fever with
thrombocytopenia syndrome virus. *J Virol* **89**, 3026-3037 (2015).
https://doi.org/10.1128/jvi.03432-14
25 Mazelier, M. *et al.* Uukuniemi Virus as a Tick-Borne Virus Model. *J Virol* **90**,
6784-6798 (2016). https://doi.org/10.1128/jvi.00095-16
26 Reynolds, E. S., Wooldridge, J. T., Stevenson, H. L. & Thangamani, S. The
Lone Star tick, Amblyomma americanum, salivary factors exacerbate the
clinical outcome of Heartland virus disease in a small animal model. *Sci Rep*
13, 13304 (2023). https://doi.org/10.1038/s41598-023-40397-x
27 McMullan, L. K. *et al.* A new phlebovirus associated with severe febrile illness
in Missouri. *N Engl J Med* **367**, 834-841 (2012).
https://doi.org/10.1056/NEJMoa1203378
28 Salvati, M. V. *et al.* Virus-Derived DNA Forms Mediate the Persistent Infection
of Tick Cells by Hazara Virus and Crimean-Congo Hemorrhagic Fever Virus. *J
Virol* **95**, e0163821 (2021). https://doi.org/10.1128/jvi.01638-21
29 Petit, M. J. *et al.* Multi-omics analysis of SFTS virus infection in Rhipicephalus
microplus cells reveals antiviral tick factors. *Nat Commun* **16**, 4732 (2025).
https://doi.org/10.1038/s41467-025-59565-w
30 Rezelj, V. V. *et al.* Differential Antagonism of Human Innate Immune Responses
by Tick-Borne Phlebovirus Nonstructural Proteins. *mSphere* **2** (2017).
https://doi.org/10.1128/mSphere.00234-17
31 Bryden, S. R. *et al.* Exploration of immunological responses underpinning
severe fever with thrombocytopenia syndrome virus infection reveals IL-6 as a

therapeutic target in an immunocompromised mouse model. *PNAS Nexus* **1**,
pgac024 (2022). https://doi.org:10.1093/pnasnexus/pgac024

32 Yu, K. M. *et al.* Cross-genotype protection of live-attenuated vaccine candidate
for severe fever with thrombocytopenia syndrome virus in a ferret model. *Proc*
Natl Acad Sci U S A **116**, 26900-26908 (2019).
https://doi.org:10.1073/pnas.1914704116

33 Choi, Y. *et al.* Severe fever with thrombocytopenia syndrome phlebovirus non-
structural protein activates TPL2 signalling pathway for viral
immunopathogenesis. *Nat Microbiol* **4**, 429-437 (2019).
https://doi.org:10.1038/s41564-018-0329-x

34 Brennan, B., Rezelj, V. V. & Elliott, R. M. Mapping of Transcription Termination
within the S Segment of SFTS Phlebovirus Facilitated Generation of NSs
Deletant Viruses. *J Virol* **91** (2017). https://doi.org:10.1128/jvi.00743-17

35 Elliott, R. M. & Brennan, B. Emerging phleboviruses. *Curr Opin Virol* **5**, 50-57
(2014). https://doi.org:10.1016/j.coviro.2014.01.011

36 Reuter, M. *et al.* The PAZ domain of Aedes aegypti Dicer 2 is critical for accurate
and high-fidelity size determination of virus-derived small interfering RNAs.
RNA **31**, 679-691 (2025). https://doi.org:10.1261/rna.080149.124

37 Yang, Z., Ebright, Y. W., Yu, B. & Chen, X. HEN1 recognizes 21-24 nt small
RNA duplexes and deposits a methyl group onto the 2' OH of the 3' terminal
nucleotide. *Nucleic Acids Res* **34**, 667-675 (2006).
https://doi.org:10.1093/nar/gkj474

38 Canto, T., Uhrig, J. F., Swanson, M., Wright, K. M. & MacFarlane, S. A.
Translocation of Tomato bushy stunt virus P19 protein into the nucleus by ALY
proteins compromises its silencing suppressor activity. *J Virol* **80**, 9064-9072
(2006). https://doi.org:10.1128/JVI.00953-06

39 Park, J. W., Faure-Rabasse, S., Robinson, M. A., Desvoyes, B. & Scholthof, H.
B. The multifunctional plant viral suppressor of gene silencing P19 interacts with
itself and an RNA binding host protein. *Virology* **323**, 49-58 (2004).
https://doi.org:10.1016/j.virol.2004.02.008

40 Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold.
Nature **596**, 583-589 (2021). https://doi.org:10.1038/s41586-021-03819-2

41 Varadi, M. *et al.* AlphaFold Protein Structure Database: massively expanding
the structural coverage of protein-sequence space with high-accuracy models.
Nucleic Acids Res **50**, D439-d444 (2022). https://doi.org:10.1093/nar/gkab1061

42 Kamel, W. *et al.* Global analysis of protein-RNA interactions in SARS-CoV-2-
infected cells reveals key regulators of infection. *Mol Cell* **81**, 2851-2867.e2857
(2021). https://doi.org:10.1016/j.molcel.2021.05.023

43 Castello, A. *et al.* Comprehensive Identification of RNA-Binding Domains in
Human Cells. *Mol Cell* **63**, 696-710 (2016).
https://doi.org:10.1016/j.molcel.2016.06.029

44 Paz, I., Kligun, E., Bengad, B. & Mandel-Gutfreund, Y. BindUP: a web server
for non-homology-based prediction of DNA and RNA binding proteins. *Nucleic*
Acids Res. **44**, W568-W574 (2016). https://doi.org:10.1093/NAR/GKW454

45 Silhavy, D. *et al.* A viral protein suppresses RNA silencing and binds silencing-
generated, 21- to 25-nucleotide double-stranded RNAs. *EMBO J* **21**, 3070-
3080 (2002). https://doi.org:10.1093/emboj/cdf312

46 Léger, P. *et al.* Dicer-2- and Piwi-mediated RNA interference in Rift Valley fever
virus-infected mosquito cells. *J Virol* **87**, 1631-1648 (2013).
https://doi.org:10.1128/jvi.02795-12

47 Brennan, B., Welch, S. R. & Elliott, R. M. The consequences of reconfiguring
the ambisense S genome segment of Rift Valley fever virus on viral replication
in mammalian and mosquito cells and for genome packaging. *PLoS Pathog* **10**,
e1003922 (2014). https://doi.org:10.1371/journal.ppat.1003922
48 Brault, A. C., Savage, H. M., Duggal, N. K., Eisen, R. J. & Staples, J. E.
Heartland Virus Epidemiology, Vector Association, and Disease Potential.
Viruses **10** (2018). https://doi.org:10.3390/v10090498
49 Lara, E., Billecocq, A., Leger, P. & Bouloy, M. Characterization of wild-type and
alternate transcription termination signals in the Rift Valley fever virus genome.
J Virol **85**, 12134-12145 (2011). https://doi.org:10.1128/jvi.05322-11
50 Moutailler, S., Krida, G., Madec, Y., Bouloy, M. & Failloux, A. B. Replication of
Clone 13, a naturally attenuated avirulent isolate of Rift Valley fever virus, in
Aedes and Culex mosquitoes. *Vector Borne Zoonotic Dis* **10**, 681-688 (2010).
https://doi.org:10.1089/vbz.2009.0246
51 Moutailler, S. *et al.* Host alternation is necessary to maintain the genome
stability of rift valley fever virus. *PLoS Negl Trop Dis* **5**, e1156 (2011).
https://doi.org:10.1371/journal.pntd.0001156
52 Geiger, T. *et al.* Use of stable isotope labeling by amino acids in cell culture as
a spike-in standard in quantitative proteomics. *Nat Protoc* **6**, 147-157 (2011).
https://doi.org:10.1038/nprot.2010.192
53 Schwanhausser, B. *et al.* Global quantification of mammalian gene expression
control. *Nature* **473**, 337-342 (2011). https://doi.org:10.1038/nature10098
54 Vargason, J. M., Szittya, G., Burgyán, J. & Hall, T. M. Size selective recognition
of siRNA by an RNA silencing suppressor. *Cell* **115**, 799-811 (2003).
https://doi.org:10.1016/s0092-8674(03)00984-x
55 Shalamova, L. *et al.* Nucleocapsids of the Rift Valley fever virus ambisense S
segment contain an exposed RNA element in the center that overlaps with the
intergenic region. *Nature Communications* **15**, 7602 (2024).
https://doi.org:10.1038/s41467-024-52058-2
56 Blair, C. D. Deducing the Role of Virus Genome-Derived PIWI-Associated
RNAs in the Mosquito-Arbovirus Arms Race. *Front Genet* **10**, 1114 (2019).
https://doi.org:10.3389/fgene.2019.01114
57 Rückert, C. *et al.* Small RNA responses of Culex mosquitoes and cell lines
during acute and persistent virus infection. *Insect Biochem Mol Biol* **109**, 13-23
(2019). https://doi.org:10.1016/j.ibmb.2019.04.008
58 Dietrich, I. *et al.* RNA Interference Restricts Rift Valley Fever Virus in Multiple
Insect Systems. *mSphere* **2** (2017). https://doi.org:10.1128/mSphere.00090-17
59 Dietrich, I. *et al.* The Antiviral RNAi Response in Vector and Non-vector Cells
against Orthobunyaviruses. *PLoS Negl Trop Dis* **11**, e0005272 (2017).
https://doi.org:10.1371/journal.pntd.0005272
60 Eifan, S., Schnettler, E., Dietrich, I., Kohl, A. & Blomström, A. L. Non-structural
proteins of arthropod-borne bunyaviruses: roles and functions. *Viruses* **5**, 2447-
2468 (2013). https://doi.org:10.3390/v5102447
61 Schnettler, E. *et al.* Diverging affinity of tospovirus RNA silencing suppressor
proteins, NSs, for various RNA duplex molecules. *J Virol* **84**, 11542-11554
(2010). https://doi.org:10.1128/jvi.00595-10
62 Tao, X. *et al.* Preparation of a polyclonal antibody against the non-structural
protein, NSs of SFTSV. *Protein Expr Purif* **184**, 105892 (2021).
https://doi.org:10.1016/j.pep.2021.105892

Billecocq, A., Vialat, P. & Bouloy, M. Persistent infection of mammalian cells by Rift Valley fever virus. *J Gen Virol* **77 (Pt 12)**, 3053-3062 (1996). https://doi.org:10.1099/0022-1317-77-12-3053

Kaul, A., Woerz, I., Meuleman, P., Leroux-Roels, G. & Bartenschlager, R. Cell culture adaptation of hepatitis C virus and in vivo viability of an adapted variant. *J. Virol.* **81**, 13168-13179 (2007). https://doi.org:10.1128/jvi.01362-07

Kurtti, T. J., Simser, J. A., Baldridge, G. D., Palmer, A. T. & Munderloh, U. G. Factors influencing in vitro infectivity and growth of Rickettsia peacockii (Rickettsiales: Rickettsiaceae), an endosymbiont of the Rocky Mountain wood tick, Dermacentor andersoni (Acari, Ixodidae). *J Invertebr Pathol* **90**, 177-186 (2005). https://doi.org:10.1016/j.jip.2005.09.001

Singu, V. *et al.* Unique macrophage and tick cell-specific protein expression from the p28/p30-outer membrane protein multigene locus in Ehrlichia chaffeensis and Ehrlichia canis. *Cell Microbiol* **8**, 1475-1487 (2006). https://doi.org:10.1111/j.1462-5822.2006.00727.x

Bell-Sakyi, L. Ehrlichia ruminantium grows in cell lines from four ixodid tick genera. *J Comp Pathol* **130**, 285-293 (2004). https://doi.org:10.1016/j.jcpa.2003.12.002

Alberdi, M. P., Nijhof, A. M., Jongejan, F. & Bell-Sakyi, L. Tick cell culture isolation and growth of Rickettsia raoultii from Dutch Dermacentor reticulatus ticks. *Ticks Tick Borne Dis* **3**, 349-354 (2012). https://doi.org:10.1016/j.ttbdis.2012.10.020

Bell-Sakyi, L. Continuous cell lines from the tick Hyalomma anatolicum anatolicum. *J Parasitol* **77**, 1006-1008 (1991).

Kurtti, T. J., Munderloh, U. G., Andreadis, T. G., Magnarelli, L. A. & Mather, T. N. Tick cell culture isolation of an intracellular prokaryote from the tick Ixodes scapularis. *J Invertebr Pathol* **67**, 318-321 (1996). https://doi.org:10.1006/jipa.1996.0050

Bell-Sakyi, L., Zweygarth, E., Blouin, E. F., Gould, E. A. & Jongejan, F. Tick cell lines: tools for tick and tick-borne disease research. *Trends Parasitol* **23**, 450-457 (2007). https://doi.org:10.1016/j.pt.2007.07.009

Yunker, C. E., Cory, J. & Meibos, H. Continuous cell lines from embryonic tissues of ticks (Acari: Ixodidae). *In Vitro* **17**, 139-142 (1981). https://doi.org:10.1007/bf02618071

Munderloh, U. G. & Kurtti, T. J. Formulation of medium for tick cell culture. *Exp Appl Acarol* **7**, 219-229 (1989). https://doi.org:10.1007/bf01194061

Munderloh, U. G. *et al.* Invasion and intracellular development of the human granulocytic ehrlichiosis agent in tick cell culture. *J Clin Microbiol* **37**, 2518-2524 (1999). https://doi.org:10.1128/jcm.37.8.2518-2524.1999

Fredericks, A. C. *et al.* Aedes aegypti (Aag2)-derived clonal mosquito cell lines reveal the effects of pre-existing persistent infection with the insect-specific bunyavirus Phasi Charoen-like virus on arbovirus replication. *PLoS Negl Trop Dis* **13**, e0007346 (2019). https://doi.org:10.1371/journal.pntd.0007346

Masek, T., Vopalensky, V., Suchomelova, P. & Pospisek, M. Denaturing RNA electrophoresis in TAE agarose gels. *Anal Biochem* **336**, 46-50 (2005). https://doi.org:S0003-2697(04)00732-8 [pii]
10.1016/j.ab.2004.09.010

77   Chen, S. Ultrafast one-pass FASTQ data preprocessing, quality control, and deduplication using fastp. *iMeta* **2**, e107 (2023). https://doi.org:10.1002/imt2.107

78 Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ
preprocessor.   *Bioinformatics*   **34**,   i884-i890   (2018).
https://doi.org:10.1093/bioinformatics/bty560
79 Langmead, B. Aligning short sequencing reads with Bowtie. *Current Protocols*
in   *Bioinformatics*   **Chapter**   **11**,   Unit   11.17   (2010).
https://doi.org:10.1002/0471250953.bi1107s32
80 Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and memory-
efficient alignment of short DNA sequences to the human genome. *Genome*
Biology **10**, R25 (2009). https://doi.org:10.1186/gb-2009-10-3-r25
81 Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics*
25, 2078-2079 (2009). https://doi.org:10.1093/bioinformatics/btp352
82 pysam: Python interface for SAM/BAM sequence alignment and mapping
format (GitHub, 2009).
83 Goddard, T. D. *et al.* UCSF ChimeraX: Meeting modern challenges in
visualization   and   analysis.   *Protein*   *Sci*   **27**,   14-25   (2018).
https://doi.org:10.1002/pro.3235
84 Paz, I., Kligun, E., Bengad, B. & Mandel-Gutfreund, Y. BindUP: a web server
for non-homology-based prediction of DNA and RNA binding proteins. *Nucleic*
Acids Res **44**, W568-574 (2016). https://doi.org:10.1093/nar/gkw454
85 Lorenz, R. *et al.* ViennaRNA Package 2.0. *Algorithms for Molecular Biology* **6**,
26 (2011). https://doi.org:10.1186/1748-7188-6-26
1239
1240

 FIGURE LEGENDS

Fig. 1. Initial characterisation of wt SFTSV replication in a diverse range of tick cells.
(A) Titre of wt SFTSV in the supernatant of infected tick cells (MOI: 1) assayed at 10 dpi. Cell lines used were derived from tick genera (*, grey) *Amblyomma* (#, purple) *Hyalomma* ($, blue) *Rhipicephalus* or (&, pink) *Ixodes.* Dashed line represents titre of initial inoculum. Data are plotted as virus titre (FFU/ml) and show a representative experiment of n=2 biological replicates.
(B) Representative images of cell extracts derived from cell monolayers in **(A)**. Western blots were probed with anti-tubulin (loading control) or anti-SFTSV N antibodies to detect cellular and viral protein expression.
(C) Tick cell lines derived from *Rhipicephalus microplus* or *Ixodes* spp. were infected with wt SFTSV (MOI: 0.1) for up to 10 dpi. Data are plotted as virus titre (FFU/ml) and show n=1 experiment.
(D) At 10 dpi, wt SFTSV-infected tick cell cultures described in **(C)** were passaged and split (1:2) and left to recover for a minimum of 14 days. This process was repeated for up to 3 subsequent passages. Data are plotted as virus titre (FFU/ml) and show a representative experiment (n=1).
(E-F) Titre of **(E)** wt or **(F)** delNSs SFTSV in the supernatant of infected BME/CTVM6 cells (MOI: 0.1 [grey]; 1 [blue]; or 5 [red]) assayed at the indicated timepoints p.i. Data are plotted as the mean virus titre (FFU/ml) ± SD of n=2 biological replicates.
(G) Representative images of cell lysates derived from cell monolayers in **(E)** and **(F)** harvested at 72 h p.i. As a control, Vero E6 cells were infected with wt SFTSV (MOI: 1) and cell lysates were prepared at 72 h p.i. Western blots were probed with anti-tubulin (loading control), anti-SFTSV NSs or anti-SFTSV N antibodies to detect cellular and viral protein expression. '-ve': mock-infected cell lysate, 'wt': wt SFTSV infected cell lysate, 'delNSs': delNSs SFTSV infected cell lysate. Data show a representative experiment (n=3).
(H) BME/CTVM6 cells seeded onto coverslips were infected with wt- or delNSs- SFTSV at the indicated MOI and were imaged at the timepoints indicated using DAPI (blue) and anti-SFTSV N (red). Imaging was carried out using a Zeiss LSM 880 Confocal Microscope, scale bar is equal to 50 μm.
(I) Percentage of BME/CTVM6 cells infected with delNSs SFTSV (MOI: 0.1 [grey]; 1 [blue]; or 5 [red]) at the timepoints indicated, evidenced by positive staining with anti-SFTSV N antibody. Data are plotted as the mean percent SFTSV-positive cells ± SD of n=3 randomised fields of view.
(J) Number of nuclei enumerated per field of view in (I). Data are plotted as the mean number of nuclei observed ± SD of n=3 randomised fields of view.
(K) Percentage of BME/CTVM6 cells infected with wt SFTSV (MOI: 0.1 [grey]; 1 [blue]; or 5 [red]) at the timepoints indicated, evidenced by positive staining with anti-SFTSV N antibody. Data are plotted as the mean percent SFTSV-positive cells ± SD of n=3 randomised fields of view.
(L) Number of nuclei enumerated per field of view in (K). Data are plotted as the mean number of nuclei observed ± SD of n=3 randomised fields of view.
(M) Violin plot showing peptide intensity distributions (y-axis: Log_2) for wt SFTSV-infected BME/CTVM6 cells at 3- and 6-dpi. Examining peptides derived from non-structural protein (NSs; AJD86041), nucleocapsid protein (N; AJD86040), glycoprotein precursor (GPC; AJD86039) or RNA-dependent RNA polymerase (RdRp; AJD86038). Data taken from n=4 biologically independent replicates.
(N) Bar Chart showing mean peptide intensities (y-axis: Log_2) for same data set presented in **(M)**.
(O) Schematic of SFTSV S genomic RNA configuration showing DIG-labelled probe binding regions used in (P). Probes detect viral genomic (−) or antigenomic (+) sense RNA as indicated. Double-ended arrows indicate genomic replication; single-ended arrows indicate transcription. Black circles indicate 5′ cap structures on mRNAs.
(P) Northern blot analysis of viral RNAs in infected cells. BME/CTVM6 and BME/CTVM23 cells were infected with wt SFTSV (MOI: 0.1) and total RNA was extracted at 6 dpi. Vero E6 cells infected with wt SFTSV were used as a positive control (RNA extracted at 72 h p.i.). DIG-labelled probes complementary to N or NSs coding sequences detected viral genomic (−) or antigenomic (+) sense RNAs. RNA sizes are indicated on the right.

Fig. 2. BME/CTVM6 cells produce virus derived siRNAs upon wt SFTSV infection.
BME/CTVM6 cells were infected with wt SFTSV (MOI: 0.1) and total cellular RNA was extracted from infected cells at 3 (light shade) or 6 (dark shade) dpi. Following RNA extraction, samples were either left untreated or processed in a beta-elimination assay, then subjected to small RNA sequencing.
(A) Total number of small RNA reads (18-30 nt) in infected cells at 3 (light shading) and 6 (dark shading) dpi. mapping to each wt SFTSV genome segment in untreated samples (left) or beta-eliminated samples (right). SFTSV genome segments: S (KP202165), M (KP202164), L (KP202163).
(B) Histogram of all small RNA reads (18-30nt) from from untreated (left) or beta-eliminated (right) samples from infected cells at 3 dpi. Read counts are coloured by the identity of the 5′ nucleotide and grouped by read length. Data are shown as the mean percentage of total mapped reads (y-axis); representative of n=3 biologically independent samples.
(C) As per (B), but RNAs were extracted from wt SFTSV-infected cells at 6 dpi.
(D) Histogram of small RNA reads (18-30 nt) from untreated (left) or beta-eliminated (right) samples, mapped to the combined SFTSV antigenomic (positive values) and genomic (negative values) strand RNAs in infected cells at 3 dpi. Read counts are coloured by the identity of the 5′ nucleotide and grouped by read length. Data are shown as the mean percentage of total mapped reads (y-axis); representative of n=3 biologically independent samples.
(E) As per (D), but RNAs were extracted from wt SFTSV-infected cells at 6 dpi.

(F-H) Mapping of SFTSV-derived 22 nt vsiRNAs (reads per million) across the **(F)** S, **(G)** M, and **(H)** L genome segments at 3 (left) and 6 (right) dpi in BME/CTVM6 cells. Data show antigenomic (blue) and genomic (pink) sense strands for untreated (top) or beta-eliminated samples (bottom) to remove 2'-O-methylated small RNAs. Numbers indicate nucleotide positions (nt). SFTSV genome segments: S (KP202165), M (KP202164), L (KP202163). Dashed lines represent the boundaries of viral open reading frames and untranslated/intergenic regions.

Fig. 3. SFTSV vsiRNAs exhibit canonical n-2 Dicer processing signature.
Reanalysis of data from Fig. 2.
(A,C) Distribution of overlapping 22 nt vsiRNA pairs by nucleotide overlap length across S (left), M (middle), and L (right) genome segments (normalized per million mapped SFTSV reads).
(B,D) Heat maps showing overlap probability z-scores for vsiRNAs of different lengths (x-axis) and overlap positions (y-axis) from three independent samples. Black boxes highlight the characteristic 2 nt overlap signature of Dicer-2 processing. Colour scale represents z-score values from -4 (teal) to +4 (purple).
(A,B) data from 3 dpi. and **(C,D)** data from 6 dpi.

Fig. 4. TBSV p19 protein rescues delNSs SFTSV replication in BME/CTVM6 cells.
BME/CTVM6 cells infected with **(A)** wt SFTSV or **(B)** delNSs-eGFP virus (MOI: 0.1). At 6 dpi, cells were fixed and immunostained for SFTSV N protein (red) and nuclei (DAPI, blue). **(C)** Western blot analysis of viral protein expression in infected BME/CTVM6 cells (6 dpi) and control Vero E6 cells (72 h p.i., MOI: 1). Blots were probed with antibodies against tubulin (loading control), SFTSV NSs, SFTSV N, or V5 tag. Lane labels: wt, wild-type SFTSV; G, delNSs-eGFP; p19, delNSs-p19V5. **(D,E)** BME/CTVM6 cells infected with delNSs-p19V5 virus (MOI: 10). Panel E shows higher magnification of infected cells. **(F)** Viral replication kinetics in BME/CTVM6 cells infected with wt SFTSV (teal), delNSs-p19V5 (red), or delNSs-eGFP (black) viruses. Virus titres were determined by focus-forming assay at indicated time points. Data show one representative experiment from n=3 independent repeats.

Fig. 5. AlphaFold structural prediction reveals putative nucleic acid binding domains in SFTSV NSs protein.
Predicted three-dimensional structure of SFTSV NSs protein (NCBI Protein: AJD86041).
(**A)** Surface representation coloured by pLDDT confidence scores: very high confidence (dark blue, 90-100), confident (light blue, 70-89), low confidence (yellow, 50-69), very low confidence (orange, 0-49).
(B) Electrostatic surface potential showing positive (blue) and negative (red) charges.
(C) RNA-binding site prediction using RBDetect analysis. Plot shows probability of RNA binding (y-axis) versus amino acid position (x-axis). Regions between dotted lines indicate high predicted RNA-binding propensity.
(D) Predicted RNA-binding residues mapped onto the protein surface: RBDetect predictions (dark green) and BindUP predictions (light green).

Fig. 6. SFTSV NSs binds viral single-stranded RNA in infected BME/CTVM6 cells.
(A) Immunoprecipitation of V5-tagged NSs proteins to assess NSs-vsiRNA interactions during SFTSV infection. BME/CTVM6 cells infected with SFTSV expressing either deletion mutant NSs-p19V5 (delNSs-p19V5) or wild-type V5-tagged NSs (wt-NSsV5) were harvested at 2 dpi and subjected to immunoprecipitation using anti-V5 antibody. Whole cell lysates (WCL; input) and immunoprecipitates (IP) were analysed by Western blot using anti-tubulin (loading control) and anti-V5 antibodies. RNA was subsequently extracted from immunoprecipitates for small RNA sequencing analysis. Molecular weights (kDa) are indicated. Data represent n=3 biological replicates (Replicate 1-3).
(B) Histogram of all small RNA reads (18-30 nt) in immunoprecipitated samples, p19V5 i.p. (left panel) or NSsV5 i.p. (right panel) representative of n=3 biological replicates.
(C) Small RNA reads from (B) mapped to SFTSV antigenomic (positive values) and genomic (negative values) strand RNAs from p19V5 i.p. (left panel) and NSsV5 i.p. (right panel) samples at 2 dpi. Read counts are coloured by the identity of the 5′ nucleotide and grouped by read length. Data are shown as the mean percentage of total mapped reads (y-axis); representative of n=3 biological replicates.
(D) Distribution of small RNA reads (18-30 nt) mapping to SFTSV genome segments from p19V5 i.p. (left panel) and NSsV5 i.p. (right panel) samples at 2 dpi Data represent n=3 biological replicates.
(E) Mapping of 22-nt SFTSV-derived vsiRNAs across the S (left panel), M (middle panel), and L (right panel) genome segments in p19V5 i.p. samples (reads per million). Data show mapping to the SFTSV antigenomic (blue) or genomic (pink) strands; numbers indicate nucleotide positions. Representative data from n=3 biological replicates.
(F) Mapping of 22-nt SFTSV-derived vsiRNAs across genome segments in NSsV5 i.p. samples, as described in (C). Representative data from n=3 biological replicates.
(G) Histogram showing the percentage of 22-nt small RNA reads with 20-nt complementary overlap mapping to SFTSV genome or *R. microplus* genome (GCA_002176555.1) in whole cell lysate (WCL) and NSsV5 i.p. samples. Data represent n=3 biological replicates. Statistical analysis was performed using Brown-Forsythe and Welch's ANOVA with Dunnett's T3 multiple comparison test to compare percentage of RNA reads in duplexes between samples. Asterisks indicate significance: * $p \leq 0.05$, ** $p \leq 0.01$; ns = not significant.
(H-J) AlphaFold3-predicted structure of SFTSV non-structural protein NSs, shown as ribbon and surface representation (grey) bound to single-stranded viral small interfering RNA (vsiRNA, blue) generated during SFTSV infection of BME/CTVM6 tick cells (Fig. 6F) derived from the viral **(H)** S, **(I)** M and **(J)** L segments. Critical amino acids involved in RNA/protein binding to distances <4 Å are highlighted (brown). Green regions highlight key RNA-binding domains identified by RBDetect (Fig. 5C,D).

FIGURE LEGENDS FOR SUPPLEMENTARY INFORMATION

Supplementary Fig. 1. Beta-elimination on uninfected tick and mosquito cell cultures.
Histogram of all small RNA reads (18-30 nt) from uninfected BME/CTVM6 or AF525 (Ago2 knock out) cells after 3 days in culture. RNA was extracted from cell cultures and samples were either **(A)** untreated or **(B)** beta-eliminated. Read counts are coloured by the identity of the 5′ nucleotide and grouped by read length. Data are shown as mean reads as a percentage of the total small RNA sequencing library (y-axis); representative of n=3 biologically independent samples.

Supplementary Fig. 2. Beta-elimination in Semliki Forest virus-infected mosquito cell cultures.
An *Aedes aegypti* Ago2 knock out mosquito cell line (AF525) was infected with SFV (MOI: 10). At 24 h p.i., total cell RNA was extracted and either left untreated (top) or subjected to beta-elimination (bottom). Histograms of small RNA reads (18-30 nt) from **(A)** untreated or **(B)** beta-eliminated samples, mapped to the SFV genomic (negative values) and antigenomic (positive values) strand RNAs in infected cells. Read counts are coloured by the identity of the 5′ nucleotide and grouped by read length. Data are shown as the mean percentage of total mapped reads (y-axis); representative of n=3 biologically independent samples. **(C, D)** SFV-derived 21 nt vsiRNAs in **(C)** untreated or **(D)** beta-eliminated samples. Coverage across the viral genome for genomic (orange) and antigenomic (yellow) sense RNAs is shown (y-axis: vsiRNA reads per million) representatives of n=3 biologically independent samples.

Supplementary Fig. 3. Distribution of overlapping 21 nt vsiRNA pairs in SFV-infected AF525 cells.
Reanalysis of data from Supplementary Fig. 2. **(A)** Distribution of overlapping 21 nt vsiRNA pairs by nucleotide overlap length across the SFV genome (SFV4, GenBank ID: KP699763), presented as normalised per million mapped SFV reads. **(B)** Heat maps showing overlap probability z-scores for vsiRNAs of different lengths (x-axis) and overlap positions (y-axis) from three independent samples. Black boxes highlight the characteristic 2 nt overlap signature of Dicer-2 processing. Colour scale represents z-score values from -6 (teal) to +6 (purple).

Supplementary Fig. 4. Characterisation of delNSs-p19V5 SFTSV in Vero E6 cells.
(A) Western blot analysis of viral protein expression in infected Vero E6 cells (6 dpi; MOI: 1). Blots were probed with antibodies against tubulin (loading control), SFTSV NSs, SFTSV N, or V5 tag. Lane labels: wt, wild-type SFTSV; G, delNSs-eGFP; p19, delNSs-p19V5. Viral replication kinetics in BME/CTVM6 cells infected with wild-type SFTSV (teal), delNSs-p19V5 (red), or delNSs-eGFP (black) viruses. Virus titres were determined by focus-forming assay at indicated time points. Data show one representative experiment from n=3 independent experiments.
(B) Viral replication kinetics in Vero E6 cells infected with wild-type SFTSV (teal), delNSs-p19V5 (red), or delNSs-eGFP (black) viruses. Virus titres were determined by focus-forming assay at indicated time points. Data show one representative experiment from n=3 independent experiments.

Supplementary Fig. 5. AlphaFold structural prediction confidence metrics for SFTSV NSs protein.
(A) Multiple sequence alignment heatmap for SFTSV NSs protein (NCBI Protein: AJD86041) and homologs. Colour scale represents sequence identity from low (red, 0.0) to high (blue, 1.0) conservation. Black line shows position-specific conservation scores. **(B)** Per-residue confidence (pLDDT) scores across AlphaFold model ranks. Blue shading highlights highest confidence region. **(C)** Inter-domain positioning confidence (pTM) scores across model ranks. Yellow shading indicates optimal predictions. **(D)** Predicted Aligned Error (PAE) matrices for top 10 ranked models showing residue-residue distance confidence. Colour scale represents predicted error (0-30 Ångströms), with blue indicating high confidence and red showing uncertainty. **(E)** Rank 1 structure coloured by pLLDT confidence: very high (dark blue: 90-100), confident (light blue: 70-89), low (yellow: 50-69), very low (orange: 0-49). **(F)** Rank 1 structure with rainbow colouring from N-terminus (blue) to C-terminus (red).

Supplementary Fig. 6. Total mapped reads and strand-specific reads for each replicate of Fig. 6B,C.
Histogram of all small RNA reads (18-30 nt) in samples immunoprecipitated with anti-V5 antibodies at 2 dpi from **(A)** p19V5 or **(B)** NSsV5 SFTSV-infected BME/CTVM6 cells (Replicates 1-3). **(C,D)** Small RNA reads from individual replicates mapped to SFTSV antigenomic (positive values) and genomic (negative values) strand RNAs from **(C)** p19V5 and **(D)** NSsV5 immunoprecipitation samples. Read counts are coloured by 5′ nucleotide identity and grouped by read length. Data are shown as percentage of total mapped reads for each individual biological replicate (n=3).

Supplementary Fig. 7. Mapping of 22-nt SFTSV-derived vsiRNAs for each replicate of Fig. 6E.
Mapping of 22-nt SFTSV-derived vsiRNAs across the S (left panel), M (middle panel), and L (right panel) genome segments in p19V5 immunoprecipitation samples (replicates 1-3; reads per million). Data show mapping to the SFTSV antigenomic (blue) or genomic (pink) strands with nucleotide positions indicated. Individual biological replicates (n=3) are shown as mapped read counts at each nucleotide position.

Supplementary Fig. 8. Mapping of 22-nt SFTSV-derived vsiRNAs for each replicate of Fig. 6F.
Mapping of 22-nt SFTSV-derived vsiRNAs across the S (left panel), M (middle panel), and L (right panel) genome segments in NSsV5 immunoprecipitation samples (replicates 1-3; reads per million). Data show mapping to the

SFTSV antigenomic (blue) or genomic (pink) strands with nucleotide positions indicated. Mapped read counts at each nucleotide position are shown for individual biological replicates (n=3).

Supplementary Fig. 9. Secondary structure of the SFTSV antigenomic S RNA (HB29 strain) showing mapping sites of vsiRNAs immunoprecipitated with NSs from infected cells.

Secondary structure prediction of a 76-nucleotide sequence (positions 1191-1266) from the antigenomic sense SFTSV S segment RNA (HB29 strain, GenBank: KP202165). Structure was predicted using minimum free energy folding calculations with Turner energy parameters at 37°C via the Forna web server powered by the ViennaRNA package. Orange circles indicate 22-nucleotide siRNAs identified through NSs immunoprecipitation and small RNA sequencing. The distribution of NSs-bound siRNAs across stem-loop and single-stranded regions demonstrates the RNA-binding specificity of NSs and supports its function as a viral suppressor of RNA silencing that sequesters host siRNAs.

Supplementary Fig. 10. Additional detected RNA duplexes of differing sizes in immunoprecipitated samples from Fig. 6.

Histograms showing the percentage of small RNA reads in duplexes with complementary overlaps: **(A)** 21-nt reads with 19-nt overlap, **(B)** 23-nt reads with 21-nt overlap, **(C)** 24-nt reads with 22-nt overlap, and **(D)** 28-nt reads with 10-nt overlap. Reads were mapped to SFTSV genome or *R. microplus* genome (GCA_002176555.1) in whole cell lysate (WCL) and NSsV5 immunoprecipitation (i.p.) samples. Data represent n=3 biological replicates. Statistical significance was determined using Brown-Forsythe and Welch's ANOVA with Dunnett's T3 multiple comparison test. Asterisks indicate significance: * $p \leq 0.05$, ** $p \leq 0.01$; ns = not significant.

A

pLDDT score
- 90-100
- 70-89
- 50-69
- 0-49

B

+ charge
– charge

C

Probability of RNA binding (y-axis, 0.0 to 1.0)
Amino acid position (x-axis, 0 to 300)

D

RBDetect
BindUP