1   **Elucidating design principles for Ribozyme-Enabled Tissue Specificity (RETS) to allow precise**
2   **expression without specialized promoters**
3

4   Max M. Combest[1], Josh Conlin[1], Vivia Van De Mark[2], Morgan Boisen[3], Hadley Colwell[1], and Arjun
5   Khakhar[1*]
6   [1]Colorado State University Department of Biology
7   [2]Colorado State University Walter Scott Jr. College of Engineering
8   [3]Colorado State University Department of Biochemistry and Molecular Biology
9

10 **Corresponding Author:**
11 Arjun Khakhar
12 **Email:** khakhar@colostate.edu

Abstract

Tissue- or cell type- specific expression of transgenes is often essential for interrogation of biological phenomenon or predictable engineering of multicellular organisms but can be stymied by cryptic enhancers that make identification of promoters that generate desired expression profiles challenging. In plants the months-to-years long timeline associated with prototyping putative tissue-specific promoters in transgenic lines deepens this challenge. We have developed a novel strategy called *Ribozyme Enabled Tissue Specificity* (RETS) that leverages the knowledge of where and when genes are expressed derived from transcriptomic studies to enable tissue-specific expression without needing characterized promoters. It uses a split self-splicing ribozyme based on a group I intron from *Tetrahymena thermophila* to enable conditional reconstitution of a transgene mRNA in the presence of a secondary tissue-specific mRNA of choice. We elucidate the design features that enable flexible swapping of transgenes and targets, enhancing transgene expression, and circumventing host RNA interference responses. We then show that these innovations enable tissue-specific and dose-dependent expression of transgenes in *Arabidopsis thaliana*. Finally, we demonstrate the utility of RETS both for creating genetically encoded biosensors to study the spatiotemporal patterns of gene expression *in planta* as well as for engineering tissue-specific changes in organ size. RETS provides a novel avenue to study expression patterns of native loci with non-destructive imaging, complementing the weakness of existing approaches. Additionally, the spatiotemporal control of transgene expression afforded by RETS enables precision engineering of plant phenotypes which will facilitate enhancing crops without the trade-offs associated with constitutive expression.

Keywords

Tissue-specific expression, Ribozymes, plant development, expression biosensor, plant synthetic biology

53 **Main Text**
54
Introduction
Precision spatiotemporal control of transgene expression is often critical for both interrogating
gene function as well as engineering novel traits in plants (Bull & Khakhar, 2023; Hong et al., 2003;
Maizel & Weigel, 2004) . The expression of transgenes in inappropriate cell and tissue types can cause
embryonic lethality or generate negative pleiotropic phenotypes that confound experiments or engineering
goals (Bandopadhyay et al., 2010) . Modern transcriptomics has generated temporally resolved maps of
native gene expression at tissue and single-cell resolution across a range of plant species (He et al.,
2023; Lim et al., 2022) . These data have been leveraged to identify genes with desired expression
profiles and their promoters have been used to enable tissue-specific transgene expression (Tian et al.,
2023) . However, identifying the sequence necessary to recapitulate a transcripts expression pattern is
often stymied by the presence of cryptic enhancers that can be embedded in introns or present hundreds
of kilobases away from the gene (Hummel et al., 2023; Tan et al., 2023; Tian et al., 2022; Ullah et al.,
2018; Yasmeen et al., 2023) . Additionally, the limited existing understanding of regulatory elements (Tian
et al., 2022, 2023; Yasmeen et al., 2023) makes it challenging to reliably predict if these promoters will
generate consistent expression patterns across different genomic insertion points, genetic backgrounds,
and environmental conditions. This often necessitates iterative prototyping in stable transgenic lines to
identify novel tissue-specific promoters, which severely limits the utility of this approach thanks to the
months- to years-long timeline associated with plant transgenesis (Bandopadhyay et al., 2010). Thus,
there is a pressing need for tools that can leverage transcriptomic data to enable spatiotemporal control
of transgene expression without the need to find and characterize new promoters. Recent advances in
engineering catalytic RNA motifs, called ribozymes, provide a promising avenue to achieve this goal.
Since the discovery of self-splicing group I introns, groups have been working on ways to
engineer them to facilitate targeted *trans*-splicing (Gambill et al., 2023; Hasegawa et al., 2004, 2006;
Hasegawa & Rao, 2006; Kruger et al., 1982; Sullenger & Cech, 1994; Watanabe & Sullenger, 2000; Zaug
et al., 1986) . A group I intron from the rRNA of *Tetrahymena thermophila* functions as a ribozyme (TtRz),
catalyzing its own excision and the ligation of its bordering exons without the need for any protein
partners (Zaug & Cech, 1986) . Recently, Gambill et al. developed an RNA detection platform based on
these motifs in bacteria (Gambill et al., 2023) wherein it was embedded in the coding sequence of a
reporter. This transcript was then split into two at the first loop of the ribozyme, producing two separate
RNA molecules, which were extended with guide sequences that are complementary to a chosen target
RNA. When the guides bind to their target, they scaffold the two portions of the TtRz close enough to
reform a functional structure and allow for the splicing reaction to occur, ligating the neighboring exons
and thereby reconstituting the reporter coding sequence to enable expression (Figure 1). This mechanism
represents an ideal starting point to enable tissue-specific expression without tissue-specific promoters.
89

90
91 **Figure 1. Graphical abstract.** (Left) Tissue-specific expression can be complicated by cryptic enhancer
92 sequences that make identifying tissue-specific promoters challenging. (Right) Ribozyme Enabled Tissue
93 Specificity, RETS, allows tissue-specific expression of transgenes without the need for characterizing
94 tissue-specific promoters.
95

In this work we have developed a promoter-independent tissue-specific expression system in
plants called *Ribozyme Enabled Tissue Specificity* (RETS). Here, a chosen cargo is divided into two
inactive transcripts that bear split ribozyme sequences fused to guide sequences that are complementary
to a tissue-specific transcript of choice. The two transcripts are expressed constitutively, but the protein
cargo is directed to be expressed only in those tissues where the target RNA is available to scaffold the
ribozyme and promote splicing-based reconstitution of the translatable mRNA, thereby producing tissue-
specific expression without the need for tissue-specific promoters. We use transient expression assays in
Nicotiana benthamiana to elucidate design features that enable flexible swapping of transgenes and
targets, enhance transgene expression, and circumvent host RNA interference (RNAi) responses. We go
on to demonstrate that the system can drive tissue-specific and dose-dependent expression of
transgenes in stable transgenic lines of the model plant *Arabidopsis thaliana*. Finally, we demonstrate the
utility of RETS both for creating genetically encoded biosensors to study the spatiotemporal patterns of
gene expression *in planta* as well as for engineering tissue-specific changes in organ size.

Results

The RETS system enables modular RNA-gated gene expression *in planta*

We first aimed to test whether the RNA-templated reconstitution of a split self-splicing ribozyme
that was previously demonstrated in bacteria (Gambill et al., 2023) could be modified to be functional *in*
planta. We constructed a transfer DNA (T-DNA) plasmid encoding constitutive expression cassettes for
the two halves of a split fluorescent reporter with the split ribozymes targeted to a test transcript, the
coding sequence of an *Arabidopsis*-codon-optimized Cas9, as well as a second fluorescent protein,
Venus, to enable ratiometric normalization to control for differential delivery. Here the split reporter, a red
fluorescent protein (mScarlet), was split at the U-base of the start codon, AUG (Figure 2A). We reasoned
that this modular split site within the start codon would enable the reporter to be easily swapped with
other proteins in the future. To enable transcript dependent reconstitution, the two halves of the ribozymes
were fused to 200-base (bp) long guide arms that had sequence complementarity with two adjacent
regions on the Cas9 transcript. A detailed description of the construct design can be found in the methods
section and supplementary materials (Supplemental RETS construct design). If this system functioned
correctly, we would expect mScarlet signal only in the presence of the Cas9 transcript. From here
forward, RETS constructs will be described with the notation, *Target*:RETS:Cargo, so this first construct
that expresses mScarlet in response to Cas9 is referred to as "*Cas9*:RETS:mScarlet".

Figure 2. RETS conditions cargo expression on presence of target RNA. A) Schematic describing
the RETS mechanism. Here the type I intron ribozyme (orange) is split with each section fused to either
the AU or G of the start codon of an mScarlet reported (red). Additionally, the other side of both sections
are fused to guide sequences (light yellow) with sequence complementarity with a trigger mRNA (dark
yellow). Annealing to this trigger brings the two halves of RETS together enabling splicing dependent
reconstitution and expression of mScarlet. B) Schematic showing how the TMV Omega enhancer (blue)
can be incorporated 5' of the split start codon to enhance mScarlet output. C) Boxplots summarizing
mScarlet signal normalized by Venus signal from transient fluorescent assays of the constructs show in A
(No enhancer) and B (Omega enhancer) with (gray) or without (yellow) co-infiltration with the trigger
transcript. Each dot represents an independent biological replicate and ** represents p ≤ 0.005.
141
142

To test *Cas9*:RETS:mScarlet, we used *Agrobacterium* infiltration (Kapila et al., 1997)   to
transiently express this construct, either with or without co-infiltration of a second T-DNA expressing the
Cas9 transcript, in the leaves of *Nicotiana benthamiana*. At 96hrs post infiltration, we observed that the
ratiometrically normalized mScarlet output was significantly higher when the Cas9 transcript was present
than when it was absent, with an average fold change of 1.97x (p=0.004) (Figure 2C). We further
hypothesized that the fold change of RETs could be enhanced via the inclusion of a translational
enhancer to generate more output signal from each correctly spliced transcript (Figure 2B). To test this,
we included a strong translational enhancer, the omega leader sequence from tobacco mosaic
virus(Gallie, 2002) , upstream of the start codon and tested it in tandem with the original construct using
the assay described above. At 96hrs post infiltration, the ratiometrically normalized mScarlet fold change
of the system with the omega sequence in the presence of Cas9, at 2.78x (p=0.0003) (Figure2C), was
significantly greater than the one lacking it. Based on these results, all future constructs incorporated this
enhancer. Taken together, these results demonstrate that RETS is functional *in planta* and that its output
can be enhanced via the incorporation of translational enhancers.

RETS enables tissue-specific expression of transgenes without needing tissue-specific promoters

Building on the transient validation of the RETS mechanism, we set out to test the utility of the
RETS system to enable tissue-specific transgene expression, by mimicking the expression pattern of
native genes, in stable transgenic lines. As a proof-of-concept we chose to use the *Arabidopsis* sucrose
transporter gene, *SUCROSE-PROTON SYMPORTER 1* (*SUC1),* which has been demonstrated by prior
studies to be expressed in trichomes (Lasin et al., 2020; Sivitz et al., 2006, 2008) . These studies showed
that *SUC1*'s expression pattern is dependent on the presence of enhancer elements in its introns and so
cannot be duplicated by reporters using only its promoter (Lasin et al., 2020) . This makes it a good
demonstration of the promoter-independent functionality of the RETS system. We constructed a new T-
DNA expressing a modified version of the previously described RETS construct, with guide sequences
that target the first exon of At*SUC1*, and used them to generate transgenic *SUC1:*RETS:mScarlet *A.
thaliana* lines. The reporter expression across several independent lines was subsequently characterized
using fluorescence microscopy. We observed that mScarlet fluorescence was localized to the bases and
cell bodies of young trichomes of rosette leaves (Figure 3B-D, See Figure S1A to compare to
untransformed Col-0) as expected (Lasin et al., 2020; Sivitz et al., 2006) . These results demonstrate that
the RETS system can program transgene expression to mimic the expression pattern of native genes
without needing tissue-specific promoters. It also shows how the RETS system can be flexibly
reprogramed to target a transcript of choice by swapping guide sequences.

 Figure 3. RETS enables tissue-specific expression of cargo with some caveats for 200nt guide
sequences. A) Cartoon describing how RETS would restrict cargo expression to trichomes due to
trichome specific expression of SUC1 (blue). B-D) Representative fluorescence microscopy images of
mScarlet signal from adult leaves of SUC1:RETS:mScarlet expressing plants with the trichomes
highlighted by dashed white lines. E) Cartoon describing how unexpected mScarlet fluorescence in
vascular tissues could be driven by cross-reactivity to vascularly localized members of the sucrose
transporter family, SUC2 and SUC9 (teal). F) Representative fluorescence microscopy images of
mScarlet signal from adult leaves of SUC1:RETS:mScarlet expressing plants with mScarlet signal in
vasculature highlighted with white arrows. G) Cartoon depicting how the production of dsRNA
intermediates during the process of target recognition could trigger RNA silencing directed at the target
and/or the RETS cassettes. H) Box plots summarizing RT-PCR expression data for AtSUC1 relative to a
housekeeping gene in wildtype (gray) and SUC1:RETS:mScarlet expressing plants (orange). Each dot
represents a biological replicate and * corresponds to p ≤ 0.05.

Unexpectedly, while characterizing these lines we also observed expression of mScarlet
throughout the vasculature of adult leaves where *SUC1* is not expressed (Figure 3F). We hypothesized
that this might be due to non-specific binding of the guide sequences to non-target transcripts leading to
reconstitution of mScarlet. An alignment of the 400 bases of *SUC1* targeted by the guide sequences with
other members of the SUC transporter family revealed a significant (~75%) sequence similarity to
vascularly localized(Sivitz et al., 2006) members, namely *SUC2* and *SUC9* (Figure S2A). This incomplete
homology is may be sufficient to trigger the RETS mechanism and highlights the need for transcript
specific guide sequences. Additionally, we hypothesized that the transient formation of duplex RNA during
the association of the RETS transcripts with the target could act as a trigger for RNAi, potentially
downregulating the target and both RETS transcripts, and leading to a reduction in mScarlet signal as
well as unwanted pleiotropies (Figure 3G). To assess whether this was occurring we measured the
expression of the target, *SUC1*, as well as a housekeeping gene to enable normalization in both
SUC1:RETS:mScarlet lines and wildtype *A. thaliana* seedlings via RT-qPCR. We observed that the RETS
lines had consistently lower levels of *SUC1* expression (mean decrease of 32.5%, p=0.034) (Figure 3H)
than wildtype seedlings indicating a silencing effect. Taken together, these results show that a lack of
guide sequence specificity and silencing both limit the functionality of RETS for enabling tissue-specific
expression.

Reducing guide length improves fold-change of RETS and reduces silencing

Seeking to resolve the design issues identified by characterization of the *SUC1*:RETS:mScarlet,
we chose to modify the length and target of the guide sequences (Figure 4A). To reduce the chances of
inducing an RNAi response from RNA duplex formation we redesigned new RETS constructs that have
shorter guides of either 30, 25 or 20 bp each. These lengths were chosen as prior work has shown that
RNA duplexes under 37 bp in length reduce the efficiency of DICER based generation of siRNA
(Kakiyama et al., 2019) . The functionality of these shorter guide sequences was tested via transient
expression experiments in *N. benthamiana*, either with or without co-infiltration with a constitutively
expressed copy of *AtSUC1* as the trigger, using the ratiometrically normalized reporter assay described
above. We observed that all three guide lengths generated significantly higher normalized mScarlet signal
in the presence of the trigger compared to when it was absent, demonstrating all three guide lengths were
functional. Additionally, we observed an increased fold change in mScarlet output for the shortened
guides compared to the previously test long (200bp) guides, ranging from 2.3-4.5x for each of the
shortened guide sequences compared to 1.8 for 200 base guides (Figure S3). Interestingly we observed
that one of the constructs with the shortest guide length (20bp), generated the largest fold change in
reporter signal (4.5x, p = 3.4×10^{-8}, Figure S3). We next made versions of the shortened-guide construct
that targeted a more unique section of the At*SUC1* transcript to reduce off-target activity. We used the
NCBI BLAST search tool (Sayers et al., 2024) to identify a region that has no significant similarity to any
other endogenous transcript. These were also tested using the same transient expression assay in *N.*
benthamina described above. Here again we observed that they retained the capacity to reconstitute the
mScarlet transcript in the presence of the *SUC1* transcript and that they had a higher fold change in
response to the target than the original RETS construct with 200 bp long guides. Overall, we observed
that the RETS constructs with shorter guides outperformed the 200bp version both in terms of absolute
activation by the target and fold change. For the shortened guides targeting the unique sequence, fold
increase in mScarlet output for each construct varied between 1.7 and 2.0 (Figure S3). Among these, the
best performing was the 30-base guide construct targeting the unique region of SUC1 (Figure 4B).

Reducing RETS guide sequence length eliminates host RNAi and off-target splicing while preserving activity

Figure 4. Reducing RETS guide length eliminates host RNAi and increases specificity. A) Cartoon
depicting how RETS guide sequences were shortened to reduce host RNAi while preserving tissue-
specific expression. B) Box plots summarizing normalized mScarlet signal from transient assays
comparing SUC1:RETS:mScarlet constructs with either the original 200 bp guides (red) or new 30 bp
guides directed at a new, unique, region of SUC1 (orange) either with (blue) or without (gray) the SUC1
trigger. Each dot is an independent biological replicate and * or *** correspond to p ≤ 0.05 or p ≤ 0.0005
respectively. C) Representative fluorescence microscopy images of mScarlet signal in adult leaves of a
line expressing SUC1:RETS:mScarlet with shortened guides. Trichomes are highlighted with dashed
white lines. D) Representative fluorescence microscopy images of mScarlet signal in the vasculature of
adult leaves of plants expressing SUC1:RETS:mScarlet with either 200 bp, 30bp, 25bp, or 20bp long
guide sequences. Vasculature is highlighted with dashed white lines. E) Boxplots summarizing the
quantification of mScarlet signal from images of the vasculature in adult leaves of plants expressing
SUC1:RETS:mScarlet with either 200 bp (red) or 30bp (orange) long guide sequences. Every dot is an
independent measurement and *** corresponds to p ≤ 0.0005. F) Box plots summarizing expression
AtSUC1 relative to a housekeeping gene in wildtype (gray) and SUC1:RETS:mScarlet with 30bp guide
expressing plants (orange). Each dot represents a biological replicate and n.s. stands for non-significant
difference.
255

Having determined that the shortened and more specific guide sequences were functional, we
used these T-DNAs to generate stable *Arabidopsis* lines to test if these modifications reduced the RNAi
and off target effects associated with the 200 bp guides. We first tested the impact of shortened guides on
silencing by measuring the expression of *SUC1* in rosette leaves of the T1 progeny of these lines along
with wildtype *A. thaliana* grown in parallel as described above. We observed that all the lines with the
shorter guides directed to the new target sequence had no significant difference in SUC1 expression
compared to the wildtype control, demonstrating that reducing the guide length eliminated the RNAi
response (Figure 4F, S4 for other guide lengths).

We further characterized the expression of the mScarlet reporter in the leaves of these lines to
determine whether these modified guides were able to drive trichome expression without the off-target
vascular expression seen in the 200 bp long guides. We observed expression of the reporter in trichomes
across the lines with shortened guides which demonstrates that this modification did not impair tissue-
specific expression (Figure 4C, S5). We also observed that all the lines with RETS constructs that
targeted the more specific region of *SUC1* had no visible signal in the vasculature compared to those
targeting the non-specific region that did, irrespective of guide length (Figure 4D). We further quantified
the difference in reporter signal from the vasculature in lines with either the original non-specific 200 bp
guide sequences or the specific 30 bp guides and observed that the more specific guides reduced the off-
target signal to background levels (Figure 4E). Taken together these results show that shortening RETS
guide sequences can reduce RNAi as well as enhance transcript accumulation in response to the trigger,
and that targeting unique regions of genes is necessary to ensure on-target tissue-specific expression.

RETS enables studying expression patterns of genes *in planta*

Our characterization of *SUC1*:RETS:mScarlet plant lines demonstrates how this tool could be
used to study the expression patterns of native genes *in planta*. To further demonstrate its utility and the
ease of target programmability we designed two new RETS constructs which drive mScarlet expression in
the presence of different genes (Figure 5A and 5C). We chose *PRODUCTION OF ANTHOCYANIN*
PIGMENT 1 (*PAP1*), a MYB transcription factor which has an intron encoded enhancer that drives its
sucrose inducibility and expression in floral sepals (Broeckling et al., 2016) , and *STERILE APETALA*
(*SAP*), a transcription factor involved in floral identity[18] that previous antisense hybridization studies
(Byzova et al., 1999) determined is expressed in the flanks of young flower buds. Like the *SUC1* RETS
constructs, the split reporter was an Omega-enhanced mScarlet, but the splice site was moved to the
middle of the gene at the second U base of codon F66. This choice was made to facilitate more efficient
qPCR verification of splicing and to demonstrate how the splice site could be reconfigured for future

cargoes (Supplemental RETS construct design). In these constructs the guides were 50 bp long as we
aimed to limit the RNAi observed with 200 bp guides but had not validated that 20 bp guides were
functional yet. T-DNAs encoding expression cassettes for these split reporters as well as a constitutively
expressed Venus were constructed and used to generate stable *A. thaliana* transgenic lines. The
expression of the mScarlet reporter in these lines across target and non-target tissues was subsequently
characterized using fluorescence microscopy.

Figure 5. RETS can be re-targeted to other genes for dose-dependent tissue-specific expression.
299
A) Cartoon depicting the expected expression of the mScarlet reporter in the flanks of young buds in
SAP:RETS:mScarlet expressing lines, based on the previously validated expression of SAP mRNA (teal)
in this tissue. B) Representative fluorescence microscopy images mScarlet signal in dissected buds of
independent SAP:RETS:mScarlet lines. Dashed white lines highlight where individual buds are in the
image. C) Cartoon depicting the expected expression of the mScarlet reporter in the base of sepals in
PAP1:RETS:mScarlet expressing lines, based on the previously validated expression of PAP1 mRNA
(purple) in this tissue. D) Representative fluorescence microscopy images mScarlet signal in dissected
flower bases of independent PAP1:RETS:mScarlet lines. Dashed white lines highlight where the flower
base and pedicle are in the image. E) Cartoon depicting how the intron encoded enhancer in PAP1
triggers sucrose induced expression of PAP1, thereby altering the concentration of spliced mScarlet
mRNA. F) Scatter plot depicting paired measurements of spliced mScarlet mRNA and PAP1 expression
relative to a housekeeping gene in PAP1:RETS:mScarlet seedlings treated with 0%, 2%, 4%, or 5%
sucrose for 24hrs. Each dot is an independent biological replicate, and darker red colors represent higher
sucrose concentrations.

We dissected and imaged the apexes of developing inflorescences and observed mScarlet signal
only in the flanks of young developing flower buds (Figure 5B) in the *SAP*:RETS:mScarlet lines, and not

in the adjacent tissues where *SAP* is not known to be expressed, demonstrating the functionality of this
reporter. In the *PAP1*:RETS:mScarlet lines we imaged adult flowers and observed mScarlet signal
localized in the sepals as expected and absent from adjacent floral pedicel tissues (Figure 5D). Critically,
across both these constructs the two halves of the RETS system and the Venus normalization control
were all driven by constitutive promoters (UBQ1 and p35s respectively) and while Venus fluorescence
was observed in all cells of the plant the RETS-gated mScarlet signal was only observed in cells where
the target transcript is expressed (Figure S6). This, coupled with the distinct distribution of mScarlet signal
across the *SUC1*:RETS:mScarlet, *SAP*:RETS:mScarlet, and *PAP1*:RETS:mScarlet lines demonstrates
that RETS can drive tissue-specific expression by piggybacking off the expression profile of a gene,
independent of whether it has regulatory elements outside the canonical promoter.

The utility of RETS-based reporters as a tool to study expression patterns necessitates that the
reporter signal match native gene expression in both localization and strength. To test this, we measured
whether the RETS-based reconstitution of the mScarlet reporter had a dose-dependent relationship to the
abundance of the target transcript by leveraging the sucrose inducibility of *PAP1* (Broeckling et al., 2016) .
We treated 4-day old *PAP1*:RETS:mScarlet seedlings for 24hrs with either 0%, 2%, 4%, or 5% sucrose
and subsequently harvested RNA from them to enable quantification of both *PAP1* expression as well as
splicing based reconstitution of mScarlet using RT-qPCR. We detected significantly higher PAP1
expression as well as correctly spliced mScarlet transcript in the sucrose treated samples as compared to
the mock treated control, and the levels of RETS-spliced mScarlet correlated well with the PAP1
expression (R^2 = 0.866) (Figure 5F). This validates the target dose dependency of the RETS-based
reporters. Taken together, these results demonstrate the utility of RETS-based reporters as genetically
encoded biosensors to study the expression patterns of native genes.

RETS enables precision engineering of plant morphology

Beyond studying native gene expression patterns, the RETS system's capacity to enable tissue-
specific expression of transgenes without needing tissue-specific promoters also opens frontiers for
precision engineering of plant phenotypes. As a proof-of-concept of this approach we designed a RETS
construct that targets a gene with a tissue-specific expression profile and reconstitutes a developmental
regulator that generates predictable changes to cell expansion and thereby organ size. We chose to
target the gene, *KNOTTED*-like from *Arabidopsis thaliana* 2 (KNAT2), which is known to be expressed
specifically in the floral meristem and root elongation zone based on tissue-specific RNA-seq (Fucile et
al., 2011; Waese et al., 2017) (Figure 6A) so we could restrict the phenotypic alterations to these tissues.
For the reconstituted developmental regulator, we chose a stabilized mutant from a family of
transcriptional regulators called DELLA proteins (Davière & Achard, 2016) . Prior studies have
demonstrated that this mutated DELLA, called GAIrht, when expressed constitutively or from its native
promoter, restricts cell expansion and hence leads to a reduction in organ size across all vegetative
tissues in the plant (Willige et al., 2007) . It has also been shown to inhibit the transition from vegetative to
floral growth and thereby delay flowering (Li et al., 2016; Willige et al., 2007; Zhang et al., 2023) . Thus,
we hypothesize that this *KNAT2*:RETS:GAIrht construct would result in plants that have a reduction in
root length, as the expression of GAIrht in the root elongation zone would prevent cell expansion, but no
change in the size of other vegetative tissues (Figure 6A). We would also predict that plants expressing
this construct would have delayed inflorescence bolting due to expression of GAIrht in floral meristems
(Figure 6A).

Figure 6. RETS enables tissue-specific engineering of plant morphology and flowering. A) A
cartoon depicting that KNAT2 is expressed in the elongation zone and shoot apical meristem (blue). The
middle boxes and plant image on the right show the KNAT2 dependent reconstitution of GAIrht should
inhibit root cell expansion and the transition to flowering but not reduce leaf area due to the lack of KNAT2
expression in leaves. B) Box plots summarizing the flowering time, quantified as days to bolting post light
shock, in either wildtype (gray) or in four independent KNAT2:RETS:GAIrht lines (reds). C) Box plots
summarizing area of the same age leaf in either wildtype (gray) or in four independent
KNAT2:RETS:GAIrht lines (reds). D) Box plots summarizing primary root length of seedlings either
wildtype (gray) or in four independent KNAT2:RETS:GAIrht lines (reds). Each dot is an independent
biological replicate and *, **, and *** correspond to p ≤ 0.05, p ≤ 0.005, and p ≤ 0.0005 respective.

        A T-DNA encoding expression cassettes for each of the two halves of the *KNAT2*:RETS:GAIrht
construct as well as a constitutively expressed Venus marker were constructed and used to generate
transgenic *A. thaliana* lines. These lines were phenotyped in tandem with wildtype *A. thaliana* controls to
assess if we observed tissue-specific changes in organ size. We observed that 3 days post germination
on ½ MS agar plates *KNAT2*:RETS:GAIrht seedlings across all lines showed significantly reduced root
lengths, between 0.56 and 0.78 fold, compared wildtype controls (Figure 6D). These seedlings were
subsequently moved to soil and then phenotyped 7 days after the start of bolting. No significant
differences were present in the *KNAT2*:RETS:GAIrht lines compared to the wildtype control in terms of
developmentally equivalent leaf area (Figure 6C). This is consistent with what we would expect if RETS
was successfully restricting the expression of GAIrht to tissues expressing KNAT2. We also observed that
KNAT2:RETS:GAIrht plants began bolting a mean of 2.4 days later than the wildtype plants (20.9 days vs.
18.5 days) which is consistent with RETS driven expression of GAIrht in floral meristems (Figure 6B).
Taken together, these results demonstrate that RETS can enable tissue-specific modulation of plant
phenotypes without defined tissue-specific promoters.

Discussion

The challenge of precise spatiotemporal control of transgene expression is often dependent on
the complexity of native transcriptional regulation (Bandopadhyay et al., 2010). The canonical proximal
promoter region 1-2Kb upstream of a gene's transcription start site is often assumed to contain most of
the relevant regulatory elements that drive the expression profile of a cognate gene. This assumption can
provide a convenient way for synthetic biologists to choose promoter regions that express desired
transgenes in specific tissues(Tian et al., 2022, 2023; Yasmeen et al., 2023) , but it is becoming clear that
this model is not generally applicable. Genes such as SUC1 (Lasin et al., 2020), PAP1 (Broeckling et al.,
2016), and AGAMOUS (Hong et al., 2003) in *Arabidopsis thaliana* have been shown to have native
expression profiles that depend on the presence of intronic enhancer elements, and more such genes are
continually being discovered (Back & Walther, 2021; Meng et al., 2021; Ullah et al., 2018) . Other
situations such as when a crucial enhancer element is genomically distant from the affected gene
(Hummel et al., 2023) or when a gene is controlled by long noncoding RNAs (Csorba et al., 2014) or
microRNAs (Dong et al., 2022) that require native-sequence specificity complicate a synthetic biologist's
ability to engineer transcriptional control mechanisms that mimic native expression patterns.
Growing transcriptomic resources for model and crop plant species provide a wealth of
information on the precise patterns of expression of native genes (Fucile et al., 2011; Waese et al.,
2017) . Our work shows that the RETS platform can leverage this information to enable spatiotemporally
precise transgene expression without being confounded by cryptic regulatory motifs. The RETS platform
builds on the work of previous groups that aimed to engineer the *Tetrahymena* group I intron to enable
targeted and/or conditional *trans*-splicing (Alexander et al., 2005; Gambill et al., 2023; Hasegawa et al.,
2004, 2006; Hasegawa & Rao, 2006; Kruger et al., 1982; Sullenger & Cech, 1994; Watanabe & Sullenger,
2000; Zaug et al., 1986). Whereas past work focused on *in vitro*, mammalian, yeast, and bacterial cell
lines, our work is focused on how it can be adapted to solve a grand challenge in multicellular synthetic
biology, namely spatiotemporally precise transgene expression. Our principal innovations are the
elucidation of design principles for RETS to enable flexible coupling to transgenes, overcome low
expression or off-target activity, and avoid host RNAi responses. We also demonstrate how these
innovations make the RETS platform a useful tool for studying gene expression patterns and engineering
plants.
The inclusion of the Omega leader sequence as the 5'UTR of the RETS splice product provided
approximately 41% greater induction above background compared to the same construct lacking it. The
likely mechanism for this increase is the Omega leader functioning to enhance the rate at which
ribosomes associate with it by recruiting elongation factors (Gallie, 2002) , increasing amount of protein
produced. We hypothesize that the mRNA splice product created by the RETS system is relatively low
abundance, so the translational enhancement is helpful to produce a robust signal. In the future the RETS
system's output strength could be improved by using stronger translational enhancers or alternative signal
amplification mechanisms. A welcome byproduct of the inclusion of the omega enhancer was the creation
of a generalized splice site at the U-base of the start codon 'AUG'. Because the internal guide sequence
of the ribozyme must be changed for every new splice site (Köhler et al., 1999) , and each splice site has
the potential to have variable efficiency (Hasegawa et al., 2006), the establishment of a consistent and
functional site at the start codon allows for the modular swapping of cargoes. This contrasts with the
recent work on a similar *in planta* ribozyme-based system which depended on a split GFP to provide a
consistent splice site and a P2A sequence to introduce modular cargos (Y. Liu et al., 2025) . In addition to
reducing cargo size requirements, we hypothesize our method for enabling modular coupling would
circumvent the significantly reduced expression that results from placing a cargo downstream of a P2A
sequence (Z. Liu et al., 2017) .
Additionally, our in-depth characterization of RETS functionality in stable transgenic lines
revealed two major issues with the use of split TtRz-based systems in plants, and likely other multicellular
eukaryotes, that were not apparent from the largely transient studies carried out in this recent study (Y.
Liu et al., 2025) . Firstly, even though longer guides have been shown to be more effective in other hosts
(Ayre et al., 1999; Gambill et al., 2023; Hasegawa et al., 2006) , the formation of dsRNA intermediates
prior to splicing triggered an RNAi response leading to the formation of siRNAs that knockdown the RETS
transcripts and/or the target sequence, drastically impacting the system's effectiveness. This problem was
evident in RETS lines with 200bp guide sequences where the expression of the target, At*SUC1*, was
observed to be suppressed compared to wildtype plants. The second issue was that the association of
the guides to a target did not require a 100% sequence match, the free energy of association of the two
guides only needed to be sufficient to bring the two pieces together. This meant that imperfect matches,

such as matches to genes closely related to the chosen target, could facilitate the splicing reaction. This
issue was demonstrated in stable lines with the observation that RETS-spliced mScarlet signal was
detected in the vasculature of rosette leaves, a tissue expected to have expression of the SUC1 relatives
SUC2 and SUC9. Our results demonstrated that reducing guide sequence length below minimum duplex
requirements for DICER proteins to efficiently generate siRNAs (Kakiyama et al., 2019) eliminated host
RNAi responses and generally outperformed longer guides in terms of both fold change and absolute on-
states. Additionally, this shortening enabled selection of a highly specific target region which eliminated
off-target activity associated with cross targeting. However, there remain many aspects of guide sequence
design that remain to be elucidated in future work, such as the precise lower limit of length that preserves
functionality, as well as the impact of guide secondary structure and target binding energy on splicing
efficiency.
A powerful application of RETS is the construction genetically encoded biosensors that enable
studying spatiotemporal patterns of gene expression, without disrupting endogenous loci, by post-
transcriptionally coupling reporter expression to transcript abundance. The tissue-specific and dose
dependent expression of reporter signal in the *SUC1*:RETS:mScarlet, *PAP1*:RETS:mScarlet, and
SAP:RETS:mScarlet lines demonstrate that RETS-reporters can be used to determine both the location
and strength of gene expression. This system has several advantages over existing strategies to study
gene expression. They enable non-destructive, cellular-resolution visualization of transcript abundance
over developmental time in contrast to RNA-seq, RT-qPCR, or FISH, all of which necessitate sacrificing
the tissue being studied, making them well suited for studying transcriptional dynamics. They can also be
easily multiplexed, thanks to the ease of swapping targets and coupled transgenes, which could enable
coupling different targets to spectrally separable reporters in future studies. However, these tools also
have limitations, such as relatively low signal and the need to express two transcripts per target.
Overcoming these challenges will be necessary to improve the utility of RETS as a tool for studying gene
expression.
Beyond studying expression patterns, the tissue-specific transgene expression enabled by RETS
opens the door to sophisticated plant engineering. Localizing the impact of transgene expression to
specific tissues has the potential to circumvent the negative pleiotropies sometimes associated with
constitutive overexpression. Our results showing that RETS enables targeted dwarfing of roots and
delayed flowering without associated dwarfing in leaves, through coupling expression of a stabilized
DELLA mutant (GAIrht) to the cell-types where KNAT2 is expressed, serves as a proof-of-concept of this
approach. These stabilized DELLA mutants were responsible for generating cereal morphologies with
semi-dwarfed shoots that drove the green revolution but also resulted in significant negative pleiotropies
such as reduced nitrogen uptake by roots (Wu et al., 2021). This highlights how, while our work focuses
on a model plant *A. thaliana*, the underlying technology has major translational relevance for agriculture.
The broad conservation of the stabilized DELLA mutant's impact on plant physiology (Willige et al.,
2007) , coupled with the host independence of TtRzs (Ayre et al., 1999; Gambill et al., 2023; Hasegawa
et al., 2006; Kalvapalle et al., 2025; Y. Liu et al., 2025; Sullenger & Cech, 1994; Watanabe & Sullenger,
2000) and its easy programmability, means this strategy could be translated to crop plants in the future to
facilitate trade-off free enhancement of morphology. A key limitation of this approach is that the
expression of the transgene is inherently limited by the expression of the tissue-specific target. In the
future, the utility of RETS for phenotype engineering could be enhanced by coupling it with secondary
mechanisms that can generate a desired level of transgene expression in response to a tissue-specific
trigger.
Taken together, this work demonstrates that RETS can facilitate the expression of diverse protein
cargos in specific plant tissues without the need for a defined tissue-specific promoter. The key benefit of
this method being the straightforward way it allows researchers to take easily accessible tissue- and
single-cell level transcriptomic data and translate it into a tissue-specific expression pattern for their
chosen transgene. These capabilities will make RETS will be valuable for plant synthetic biology projects
that require precision timing or targeting of expression. In the future these tools could also be expanded
to respond externally derived transcripts, such as viral genomes or trans-kingdom RNAs, to enable the
detection and management of plant pathogens.
501
502 **Materials and Methods**
503
General RETS and cloning Methods

RETS constructs were designed based on vectors described in Gambill et al. (2023) (Gambill et
al., 2023) . The 419 base *Tetrahymena thermophila* rRNA intron ribozyme (TtRz) was split at the 15th
residue to produce two fragments: (1) An upstream fragment wherein the first 6 bases are the final 6
bases of the upstream exon (also referred to as the internal guide sequence binding site, IGSBS) with the
6th base being a required U that defines the exon splice junction. The following 9 bases are the P1 loop of
the TtRz. (2) A downstream fragment that consists of the next 11 bases of the P1 loop, a 6-base internal
guide sequence (IGS) that binds complementarily to the IGSBS perfectly aside from a G that creates a
wobble pairing with the splice-site U, the remaining 387 bases of the TtRz, then the downstream exon. To
the 3' end of fragment (1) a first guide sequence is added that is complementary to a target RNA of
choice, preceded by helper stem sequence (GGATCA). To the 5' end of fragment (2) a second guide
sequence is added, also complementary to the target RNA, and this guide is followed by the second
strand of the helper stem (TGATCC). The first and second guide sequences are chosen such that their
binding sites are separated by 5 bases of target transcript to accommodate the helper stem and assist
with the necessary association of the two RETS transcripts when complexed with the target RNA. Guide
RNA length varied from 200 and 20 bases between experiments as noted.

Most RETS constructs in this work were designed to output the fluorescent protein, mScarlet
except for the final experiments that output an autoactive rht variant of the Arabidopsis DELLA protein,
GAI. Exon splice site arrangements were either designed to split the AUG start codon (utilizing the start
codon's 'U' base as the splice site) or a codon in the middle of the output protein's coding sequence
(UU/U of F66 in mScarlet, or CU/C of L56 in GAIrht). For each splice site, the IGS needed to be modified
to match the bases in the IGSBS of the upstream exon (accounting for the required G/U wobble pairing).
In the cases where the start codon served as the splice site, the final 4 bases of the omega enhancer
(UUAC) completed the IGSBS. After the first experiment, all output transcripts included an omega
enhancer prior to their coding sequences, whether the splice site was at the start codon or not. The two
cassettes expressing the first and second RETS transcripts were driven by the Arabidopsis UBQ1
(pUBQ1) and CaMV 35s (p35s) promoters respectively, and both were terminated with the Arabidopsis
HSP18.2 terminator (tHSP). In every case, a third cassette with a nuclear localized yellow fluorescent
protein, venus (driven by pUBQ1, terminated by tHSP), was included for ratiometric normalization and to
simplify transformant selection. These three cassettes were cloned into binary vectors using a 2-level
modular golden gate cloning system (Čermák et al., 2017) to create the final plasmids for transformation
into agrobacterium and subsequent use in plants.

Agrobacterium transient Infiltration

Constructs were transformed into agrobacterium strain GV3101 using the freeze-thaw method
and positive transformants were selected on LB plates with 50ug/ml kanamycin. Single isolated colonies
were picked after 3 days growing at 28C. Selected colonies were used to inoculate liquid LB cultures that
were grown shaking at 28C overnight (12-18 hours). Cultures were centrifuged at 2500g for 5 minutes to
pellet bacteria, and the pellets were washed once with infiltration buffer (10 mM MgCl2, 10 mM 2-(N-
morpholino) ethanesufonic acid (MES) pH 5.6), then resuspended in the same buffer supplemented with
100uM of acetosyringone. These bacterial suspensions were adjusted to OD600 of 1.0 and combined 1:1
with each other to facilitate codelivery of RETS_Omega+Cas9, RETS_NO-OMEGA+Cas9 and the RETS
suspensions were also diluted 1:1 with acetosyringone-supplemented infiltration buffer to provide Cas9 (-)
conditions with equivalent effective optical densities as the Cas9 (+) treatments (0.5 effective per
construct). Diluted cultures were incubated for 2-4 hours at room temperature and then gently infiltrated
into fully expanded leaves of 4-6 week old *Nicotiana benthamiana* plants using needleless syringes. After
72-96 hours, 7mm leaf disks were taken with a core borer from the infiltrated zones of each leaf and their
fluorescence readings were analyzed by a Tecan Spark plate reader. Background autofluorescence was
determined by analyzing buffer and Cas9-only infiltrated zones, and the mScarlet and Venus readings
from corresponding RETS-infiltrated zones were assessed based on a fold change above the average
background reading. Final normalized readings (Figure 2C) were determined based on a ratiometric
comparison (mScarlet/Venus) between the mScarlet output (proportional to RETS-mediated splicing) and
the Venus reading (constitutive, proportional to the number of cells in the leaf disk). All p values reported
were calculated using a student's T Test.

Plant growth

When grown for seed or tissue for microscopy, *Arabidopsis thaliana* ecotype Columbia (Col-0)
plants are grown in soil (3:1 ratio of Promix and turface) in a growth room or growth chamber at ~22-24C
at 60% humidity and 120-150 µmol/m²/s light with a 16/8hr day-night cycle. For transient assays,
Nicotiana benthamiana plants were grown in Promix for 3-5 weeks in a growth chamber set to 25C, the
same humidity and light conditions as *Arabidopsis*. Stable *Arabidopsis* lines were generated in the
Columbia background (Col-0) using the agrobacterium floral dip method. Single isolated colonies of
Agrobacterium tumifaciens bearing the desired binary plasmids were used to start 100mL LB cultures
(supplemented with kanamycin and gentamicin) that were grown to confluence shaking overnight at 28C.
Cultures were centrifuged for 10 minutes at 4000g to aspirate the growth medium and pellets were
resuspended in 200mL of transformation buffer (ddH$_2$0 with 5% sucrose, 0.03% Silwet L-77). Densely
planted Col-0 plants that have bolted with the first flowers beginning to open are then dipped into the
bacterial solution and swirled for ~10s and allowed to dry in darkness for 24hrs. The plants are then
returned to light and allowed to set T0 seed. Transgenic plants are germinated under selection by
inducing light shock on antibiotic-supplemented medium. T0 seed is surface sterilized with 70% ethanol,
then seeds are spread onto ½ Murashige-Skoog medium plates (0.8% w/v phytoagar) with 50mg/mL
kanamycin and allowed to stratify in darkness at 4C for at least 48hrs. Plates are transferred to light for
6hrs to induce germination and then covered again for 48-72hrs at 22-25C to induce etiolation. Plates are
then uncovered and over the next 3-5 days, the seedlings that survive the etiolation and produce adult
leaves are considered to have passed selection and move on to further analysis. Selected lines are
transferred to soil, used for experiments and/or allowed to set T1 seed. T1 plants are then either used for
experiments or put through another round of selection to produce a T2 generation and so on.

Microscope imaging

All microscope images were taken using a Leica DM5500 B upright fluorescence microscope
using a CTR5500 controller system, a mercury halide lamp, and the LAS X software package.
Fluorescence was resolved using mScarlet (570±20nm excitation, 609±34nm emission) and GFP
(470±40nm excitation, 525±50nm emission) filter cubes for mScarlet and Venus visualization respectively.
Magnification was set to 10x for every image using an HCX PL Fluotar 5x objective and a 2x ocular lens
prior to image acquisition. All images were 1 second exposures, in FIM light mode with the FIM and
transmission light field set to the maximum settings of 100% and 6 respectively. Images were exported as
raw TIFF files and prepared for publication in imageJ. Plant material for imaging was prepared as either
excised leaves (SUC1), mature flowers (PAP1), or developing flower buds (SAP), that were set on a glass
slide either with a drop of 0.5% Tween-20 solution and a glass coverslip (SUC1) or imaged dry and
uncovered (PAP1/SAP1).

Plant phenotyping and imageJ analysis.

For Figure 4E, microscope images (mScarlet channel, all with the same capture settings and
lookup table thresholds) of leaves of SUC1:RETS:mScarlet lines with either 200 base (old target) or 30
base guides (new target) were analyzed with the ROI tool in imageJ. A background ROI was selected on
a region of the leaf without observable vasculature and 5 ROIs were taken along the vascular features
that were visible. The mean mScarlet signal from the background measurement was subtracted from the
other 5 ROIs to produce the "Vascular relative mScarlet signal" reported in Figure 4E. For synchronized
growth experiments (Figure 6), T1 KNAT2:RETS:GAIrht and Col-0 seeds were arranged in rows on ½ MS
plates without antibiotics, stratified and light shocked as previously described, and placed covered,
vertically in the growth chamber. After 72hrs etiolating in darkness, the plates were scanned with an office
copier alongside a ruler to provide scale. The plates were then examined under a fluorescent microscope
to identify which plants were true transgenics – plants with no identifiable Venus fluorescence signal in
the roots were excluded from the analyses. After measurement, the seedlings were allowed to grow
vertically on plates for ~2 weeks before a subset of each genotype were transferred to soil in a growth
chamber. Over the following days, each individual plant was monitored for signs of bolting. Seven days
after the first sign of bolting, each plant was removed and fully dissected, with each leaf and the
inflorescence being arranged in developmental order and scanned. All p values reported were calculated
using a student's T Test. All code for analyzing and plotting data, as well as all raw data, can be found at
https://github.com/arjunkhakhar/RETS_paper.

Sucrose treatments

To examine sucrose dependent induction of PAP1 and correlated induction of RETS-spliced
mScarlet (Figure 5F), seedlings of Col-0 and PAP1:RETS:mScarlet were germinated on solid ½ MS
media and light shocked as normal. Plants were transferred to light after 72hrs in darkness and allowed to
grow for 24 more hours. Seedlings were then transferred to liquid ½ MS media supplemented with either
0%, 2%, 4%, or 5% sucrose (w/v). Seedlings in liquid media were returned to the growth chamber and
allowed to incubate for 24hrs. Seedlings were then pooled, 5-8 per replicate, to create 4 replicates per
treatment for RNA extraction and subsequent RT-qPCR analysis. All code for analyzing and plotting data,
as well as all raw data, can be found at https://github.com/arjunkhakhar/RETS_paper.

Expression Analysis

For relative gene expression analysis, sampling was done by harvesting at least 5 whole
seedlings either 10 days post germination (for SUC1 silencing analysis: Figure 3H, 4H, and S4) or 4 days
post germination (for PAP1 sucrose tests: Figure 5F). RNA was extracted using the Trizol method
following manufacturer's protocols. RNA was then treated with TurboDNase according to the
manufacturer's 'Rigorous DNase treatment' protocol. Synthesis of cDNA was performed with LunaScript
reverse transcriptase mastermix using either the 'Supermix' of oligo-dT and random hexamer primers
(SUC1 silencing experiment), or gene specific reverse primers to recognize PAP1, SUC1, mScarlet and
the housekeeping gene PP2aa3. Fold changes in gene expression were calculated using the ddCT
method. All p values reported were calculated using a student's T Test. All code for analyzing and plotting
data, as well as all raw data, can be found at https://github.com/arjunkhakhar/RETS_paper.

630
631
632
633   **Abbreviations**

634
RETS – Ribozyme Enabled Tissue Specificity
TtRz – *Tetrahymena thermophila* Ribozyme
SUC1 – Sucrose-proton symporter 1
RNAi – RNA Interference
SAP – Sterile Apetala
PAP1 – Production of Anthocyanin Pigment 1
KNAT2 – *KNOTTED*-like from *Arabidopsis thaliana* 2

642
643

Author information

Research was performed at Colorado State University:

Biology Building

251 W. Pitkin Street

Fort Collins, CO 80521

United States of America

Author Contributions

MC designed the research, performed research, analyzed data, and wrote the paper; JC performed
research; VV performed research; MB performed research; HC performed research; AK acquired funding,
designed the research, and revised the paper.

Conflict of Interest

The authors declare no conflict of interest

Supporting Information

Supplementary Materials: Additional methods, supporting data including figures S1-S6, and tables S1-S3 listing constructs, primers, and genes used in this study (DOC).

Acknowledgments

AK's time and the reagents costs for this work were funded by a grant from NIGMS (5R35GM155313). MC was partially funded by a grant from the Institute of Cannabis Research at Colorado State University in Pueblo (ICR-FY23-Nachappa).

References

Alexander, R. C., Baum, D. A., & Testa, S. M. (2005). 5' Transcript Replacement in Vitro Catalyzed by a Group I Intron-Derived Ribozyme †. *Biochemistry*, *44*(21), 7796–7804. https://doi.org/10.1021/bi047284a

Ayre, B. G., Köhler, U., Goodman, H. M., & Haseloff, J. (1999). Design of highly specific cytotoxins by using trans-splicing ribozymes. *Proceedings of the National Academy of Sciences*, *96*(7), 3507–3512. https://doi.org/10.1073/pnas.96.7.3507

Back, G., & Walther, D. (2021). Identification of cis-regulatory motifs in first introns and the prediction of intron-mediated enhancement of gene expression in Arabidopsis thaliana. *BMC Genomics*, *22*(1), 390. https://doi.org/10.1186/s12864-021-07711-1

Bandopadhyay, R., Haque, I., Singh, D., & Mukhopadhyay, K. (2010). Transgenic Crop Plants, Principles and Development. *Transgenic Crop Plants*, 145–186. https://doi.org/10.1007/978-3-642-04809-8_5

Broeckling, B. E., Watson, R. A., Steinwand, B., & Bush, D. R. (2016). Intronic Sequence Regulates Sugar-Dependent Expression of Arabidopsis thaliana Production of Anthocyanin Pigment-1/MYB75. *PLoS ONE*, *11*(6), e0156673. https://doi.org/10.1371/journal.pone.0156673

Bull, T., & Khakhar, A. (2023). Design principles for synthetic control systems to engineer plants. *Plant Cell Reports*, *42*(12), 1875–1889. https://doi.org/10.1007/s00299-023-03072-z

Byzova, M. V., Franken, J., Aarts, M. G. M., Almeida-Engler, J. de, Engler, G., Mariani, C., Campagne, M. M. V. L., & Angenent, G. C. (1999). Arabidopsis STERILE APETALA, a multifunctional gene regulating inflorescence, flower, and ovule development. *Genes & Development*, *13*(8), 1002–1014. https://doi.org/10.1101/gad.13.8.1002

Čermák, T., Curtin, S. J., Gil-Humanes, J., Čegan, R., Kono, T. J. Y., Konečná, E., Belanto, J. J., Starker, C. G., Mathre, J. W., Greenstein, R. L., & Voytas, D. F. (2017). A Multipurpose Toolkit to Enable Advanced Genome Engineering in Plants. *The Plant Cell*, *29*(6), 1196–1217. https://doi.org/10.1105/tpc.16.00922

Csorba, T., Questa, J. I., Sun, Q., & Dean, C. (2014). Antisense COOLAIR mediates the coordinated switching of chromatin states at FLC during vernalization. *Proceedings of the National Academy of Sciences*, *111*(45), 16160–16165. https://doi.org/10.1073/pnas.1419030111

Davière, J.-M., & Achard, P. (2016). A Pivotal Role of DELLAs in Regulating Multiple Hormone Signals. *Molecular Plant*, *9*(1), 10–20. https://doi.org/10.1016/j.molp.2015.09.011

Dong, Q., Hu, B., & Zhang, C. (2022). microRNAs and Their Roles in Plant Development. *Frontiers in Plant Science*, *13*, 824240. https://doi.org/10.3389/fpls.2022.824240

Fucile, G., Biase, D. D., Nahal, H., La, G., Khodabandeh, S., Chen, Y., Easley, K., Christendat, D., Kelley, L., & Provart, N. J. (2011). ePlant and the 3D Data Display Initiative: Integrative Systems Biology on the World Wide Web. *PLoS ONE*, *6*(1), e15237. https://doi.org/10.1371/journal.pone.0015237

Gallie, D. R. (2002). The 5'-leader of tobacco mosaic virus promotes translation through enhanced recruitment of eIF4F. *Nucleic Acids Research*, *30*(15), 3401–3411. https://doi.org/10.1093/nar/gkf457

Gambill, L., Staubus, A., Mo, K. W., Ameruoso, A., & Chappell, J. (2023). A split ribozyme that links detection of a native RNA to orthogonal protein outputs. *Nature Communications*, *14*(1), 543. https://doi.org/10.1038/s41467-023-36073-3

Hasegawa, S., Choi, J. W., & Rao, J. (2004). Single-Cell Detection of Trans-Splicing Ribozyme In Vivo Activity. *Journal of the American Chemical Society*, *126*(23), 7158–7159. https://doi.org/10.1021/ja049144u

Hasegawa, S., Gowrishankar, G., & Rao, J. (2006). Detection of mRNA in Mammalian Cells with a Split Ribozyme Reporter. *ChemBioChem*, *7*(6), 925–928. https://doi.org/10.1002/cbic.200600061

Hasegawa, S., & Rao, J. (2006). Modulating the splicing activity of Tetrahymena ribozyme via RNA self-assembly. *FEBS Letters*, *580*(6), 1592–1596. https://doi.org/10.1016/j.febslet.2006.01.090

He, Z., Luo, Y., Zhou, X., Zhu, T., Lan, Y., & Chen, D. (2023). scPlantDB: a comprehensive database for exploring cell types and markers of plant cell atlases. *Nucleic Acids Research*, *52*(D1), D1629–D1638. https://doi.org/10.1093/nar/gkad706

Hong, R. L., Hamaguchi, L., Busch, M. A., & Weigel, D. (2003). Regulatory Elements of the
Floral Homeotic Gene AGAMOUS Identified by Phylogenetic Footprinting and
Shadowing[W]. *The Plant Cell*, *15*(6), 1296–1309. https://doi.org/10.1105/tpc.009548

Hummel, N. F. C., Zhou, A., Li, B., Markel, K., Ornelas, I. J., & Shih, P. M. (2023). The trans-
regulatory landscape of gene networks in plants. *Cell Systems*, *14*(6), 501-511.e4.
https://doi.org/10.1016/j.cels.2023.05.002

Kakiyama, S., Tabara, M., Nishibori, Y., Moriyama, H., & Fukuhara, T. (2019). Long DCL4-
substrate dsRNAs efficiently induce RNA interference in plant cells. *Scientific Reports*, *9*(1),
6920. https://doi.org/10.1038/s41598-019-43443-9

Kalvapalle, P. B., Staubus, A., Dysart, M. J., Gambill, L., Gamas, K. R., Lu, L. C., Silberg, J. J.,
Stadler, L. B., & Chappell, J. (2025). Information storage across a microbial community using
universal RNA barcoding. *Nature Biotechnology*, 1–8. https://doi.org/10.1038/s41587-025-
02593-0

Kapila, J., Rycke, R. D., Montagu, M. V., & Angenon, G. (1997). An Agrobacterium-mediated
transient gene expression system for intact leaves. *Plant Science*, *122*(1), 101–108.
https://doi.org/10.1016/s0168-9452(96)04541-4

Köhler, U., Ayre, B. G., Goodman, H. M., & Haseloff, J. (1999). Trans-splicing ribozymes for
targeted gene delivery. *Journal of Molecular Biology*, *285*(5), 1935–1950.
https://doi.org/10.1006/jmbi.1998.2447

Kruger, K., Grabowski, P. J., Zaug, A. J., Sands, J., Gottschling, D. E., & Cech, T. R. (1982).
Self-splicing RNA: Autoexcision and autocyclization of the ribosomal RNA intervening
sequence of tetrahymena. *Cell*, *31*(1), 147–157. https://doi.org/10.1016/0092-8674(82)90414-
7

Lasin, P., Weise, A., Reinders, A., & Ward, J. M. (2020). Arabidopsis Sucrose Transporter
AtSuc1 introns act as strong enhancers of expression. *Plant and Cell Physiology*, *61*(6),
1054–1063. https://doi.org/10.1093/pcp/pcaa029

Li, M., An, F., Li, W., Ma, M., Feng, Y., Zhang, X., & Guo, H. (2016). DELLA proteins interact
with FLC to repress flowering transition. *Journal of Integrative Plant Biology*, *58*(7), 642–
655. https://doi.org/10.1111/jipb.12451

Lim, P. K., Zheng, X., Goh, J. C., & Mutwil, M. (2022). Exploiting plant transcriptomic
databases: Resources, tools, and approaches. *Plant Communications*, *3*(4), 100323.
https://doi.org/10.1016/j.xplc.2022.100323

Liu, Y., Rajput, R., Islam, M. T., Valle, I. D., Yao, T., Agrawal, R., Boone, B. A., Eckert, C. A.,
Abraham, P. E., Chen, J., Tuskan, G. A., & Yang, X. (2025). A split ribozyme system for in
vivo plant RNA imaging and genetic engineering. *Plant Biotechnology Journal*.
https://doi.org/10.1111/pbi.14612

Liu, Z., Chen, O., Wall, J. B. J., Zheng, M., Zhou, Y., Wang, L., Vaseghi, H. R., Qian, L., & Liu,
J. (2017). Systematic comparison of 2A peptides for cloning multi-genes in a polycistronic
vector. *Scientific Reports*, *7*(1), 2193. https://doi.org/10.1038/s41598-017-02460-2

Maizel, A., & Weigel, D. (2004). Temporally and spatially controlled induction of gene
expression in Arabidopsis thaliana. *The Plant Journal*, *38*(1), 164–171.
https://doi.org/10.1111/j.1365-313x.2004.02027.x

Meng, F., Zhao, H., Zhu, B., Zhang, T., Yang, M., Li, Y., Han, Y., & Jiang, J. (2021). Genomic
editing of intronic enhancers unveils their role in fine-tuning tissue-specific gene expression
in Arabidopsis thaliana. *The Plant Cell*, *33*(6), 1997–2014.
https://doi.org/10.1093/plcell/koab093

Sayers, E. W., Beck, J., Bolton, E. E., Brister, J. R., Chan, J., Connor, R., Feldgarden, M., Fine,
A. M., Funk, K., Hoffman, J., Kannan, S., Kelly, C., Klimke, W., Kim, S., Lathrop, S.,
Marchler-Bauer, A., Murphy, T. D., O'Sullivan, C., Schmieder, E., … Pruitt, K. D. (2024).
Database resources of the National Center for Biotechnology Information in 2025. *Nucleic
Acids Research*, *53*(D1), D20–D29. https://doi.org/10.1093/nar/gkae979

Sivitz, A. B., Reinders, A., Johnson, M. E., Krentz, A. D., Grof, C. P. L., Perroux, J. M., &
Ward, J. M. (2006). Arabidopsis Sucrose Transporter AtSUC9. High-Affinity Transport
Activity, Intragenic Control of Expression, and Early Flowering Mutant Phenotype. *Plant
Physiology*, *143*(1), 188–198. https://doi.org/10.1104/pp.106.089003

Sivitz, A. B., Reinders, A., & Ward, J. M. (2008). Arabidopsis Sucrose Transporter AtSUC1 Is
Important for Pollen Germination and Sucrose-Induced Anthocyanin Accumulation. *Plant
Physiology*, *147*(1), 92–100. https://doi.org/10.1104/pp.108.118992

Sullenger, B. A., & Cech, T. R. (1994). Ribozyme-mediated repair of defective mRNA by
targeted trans-splicing. *Nature*, *371*(6498), 619–622. https://doi.org/10.1038/371619a0

Tan, Y., Yan, X., Sun, J., Wan, J., Li, X., Huang, Y., Li, L., Niu, L., & Hou, C. (2023). Genome-
wide enhancer identification by massively parallel reporter assay in Arabidopsis. *The Plant
Journal*, *116*(1), 234–250. https://doi.org/10.1111/tpj.16373

Tian, C., Li, J., & Wang, Y. (2023). From qualitative to quantitative: the state of the art and
challenges for plant synthetic biology. *Quantitative Biology*, *11*(3), 214–230.
https://doi.org/10.15302/j-qb-022-0326

Tian, C., Zhang, Y., Li, J., & Wang, Y. (2022). Benchmarking Intrinsic Promoters and
Terminators for Plant Synthetic Biology Research. *BioDesign Research*, *2022*, 1–12.
https://doi.org/10.34133/2022/9834989

Ullah, F., Hamilton, M., Reddy, A. S. N., & Ben-Hur, A. (2018). Exploring the relationship
between intron retention and chromatin accessibility in plants. *BMC Genomics*, *19*(1), 21.
https://doi.org/10.1186/s12864-017-4393-z

Waese, J., Fan, J., Pasha, A., Yu, H., Fucile, G., Shi, R., Cumming, M., Kelley, L. A., Sternberg,
M. J., Krishnakumar, V., Ferlanti, E., Miller, J., Town, C., Stuerzlinger, W., & Provart, N. J.
(2017). ePlant: Visualizing and Exploring Multiple Levels of Data for Hypothesis Generation
in Plant Biology. *The Plant Cell*, *29*(8), 1806–1821. https://doi.org/10.1105/tpc.17.00073

Watanabe, T., & Sullenger, B. A. (2000). Induction of wild-type p53 activity in human cancer
cells by ribozymes that repair mutant p53 transcripts. *Proceedings of the National Academy of*
Sciences, *97*(15), 8490–8494. https://doi.org/10.1073/pnas.150104097

Willige, B. C., Ghosh, S., Nill, C., Zourelidou, M., Dohmann, E. M. N., Maier, A., &
Schwechheimer, C. (2007). The DELLA Domain of GA INSENSITIVE Mediates the
Interaction with the GA INSENSITIVE DWARF1A Gibberellin Receptor of Arabidopsis.
The Plant Cell, *19*(4), 1209–1220. https://doi.org/10.1105/tpc.107.051441

Wu, K., Xu, H., Gao, X., & Fu, X. (2021). New insights into gibberellin signaling in regulating
plant growth–metabolic coordination. *Current Opinion in Plant Biology*, *63*, 102074.
https://doi.org/10.1016/j.pbi.2021.102074

Yasmeen, E., Wang, J., Riaz, M., Zhang, L., & Zuo, K. (2023). Designing artificial synthetic
promoters for accurate, smart, and versatile gene expression in plants. *Plant Communications*,
4(4), 100558. https://doi.org/10.1016/j.xplc.2023.100558

Zaug, A. J., Been, M. D., & Cech, T. R. (1986). The Tetrahymena ribozyme acts like an RNA
restriction endonuclease. *Nature*, *324*(6096), 429–433. https://doi.org/10.1038/324429a0

Zaug, A. J., & Cech, T. R. (1986). The Intervening Sequence RNA of Tetrahymena Is an
Enzyme. *Science*, *231*(4737), 470–475. https://doi.org/10.1126/science.3941911

Zhang, C., Jian, M., Li, W., Yao, X., Tan, C., Qian, Q., Hu, Y., Liu, X., & Hou, X. (2023).
Gibberellin signaling modulates flowering via the DELLA–BRAHMA–NF-YC module in
Arabidopsis. *The Plant Cell*, *35*(9), 3470–3484. https://doi.org/10.1093/plcell/koad166

Supplementary Materials

Supplemental RETS construct design

For figure 2, RETS constructs were designed to detect the coding sequence of an *Arabidopsis*-codon-
optimized Cas9 transcript. Guide RNA sequences were 200 bases in length, and the mScarlet cargo
transcript was split at the start codon. Upstream of the start codon was either the complete 67-base
omega enhancer (RETS_Omega) or just the final 4 bases of the enhancer to provide a complete IGSBS
(RETS_NO-Omega). A third, target construct was prepared in parallel which expressed Cas9 under
control of the *Arabidopsis* UBQ10 promoter, terminated by tHSP.

For figure 3, the RETS construct was re-designed to target exon 1 of *Arabidopsis* Sucrose transporter 1,
AtSUC1. For figure 4, to reduce the likelihood of inducing dsRNA-dependent RNA silencing, RETS
constructs with shorter guides were produced. Targeting the same region of AtSuc1, 3 constructs were
made that utilize pairs of 30nt, 25nt, and 20nt RNA guide arms instead of the 200nt guide arms of the
initial construct. In addition, 3 more constructs were designed that had 30, 25, and 20nt guide arms re-
targeted to a region of AtSuc1 that is more divergent from the other members of the Suc gene family (see
figure S2).

To facilitate more rapid prototyping, a constitutive copy of the AtSuc1 cDNA was cloned into a binary
vector under control of the CaMV 35s promoter. This construct allowed for characterization of the new
Suc1:RETS constructs in *N. benthamiana* leaves with transient expression via agroinfiltration.

For the PAP1 and SAP-detecting constructs in figure 5, the backbone of the cassettes was modified
slightly: The guides were all made to be 50nt in length, and the split-site of the cargo was designed to be
U197 of mScarlet, the second base of F66. This change altered the IGSBS, and thus the IGS of the TtRz
was modified to be 5'-GACUGC-3' to reflect that change, allowing for successful reconstitution of the TtRz
P1 helix, facilitating the splicing reaction.

For figure 6, Autoactive DELLA protein mutant GAI(*rht*) was constructed by amplifying *Arabidopsis* GAI1
(AT1G14920) from *Col-0* without the first 55 amino acids (using codon for M56 as the new start),
analogous to what was described by Willige et al (2007). The RETS construct designed to detect KNAT2
was redesigned to reconstitute GAI(*rht*), splitting the transcript at U167, and having an IGSBS and IGS of
5'-GAUUCU-3' and 5'-GGAAUC-3' respectively. The constitutive Venus cassette was included to facilitate
selection-free identification of transformants.

874 **Supplemental Figure S1. Representative *A. thaliana* Col-0 wildtype fluorescent microscope**
875 **images using the mScarlet channel.** A) Shows wildtype trichomes, compare to Figure 3B-D, 3F and 4C.
876 B) wildtype vasculature, compare to Figure 4D. C) Shows developing wildtype flower buds, compare to
877 Figure 5B. D) Shows wildtype adult flower sepals, compare to Figure 5D. Displayed lookup table
878 thresholds are calibrated to the respective experimental images they are compared to.

A

	cov	pid	
200nt RETS guide target	100.0%	100.0%	1 [
AtSUC1	100.0%	98.5%	
AtSUC2	100.0%	73.7%	
AtSUC9	99.2%	73.8%	

Supplemental Figure S2. Multiple sequence alignment of RETS target sequences and SUC family genes. A) Multiple sequence alignment of the original RETS construct target sequence with 200-base guides targeting a region of AtSUC1. Target sequence is aligned to the homologous region of AtSUC1 and the major vascularly-localized members of the SUC family, SUC2 and SUC9. B) Alignment of a re-designed target site for a RETS construct with 30-base guides and SUC1, 2, and 9. Note that in both alignments, there is an intentional 5-base gap in the intended target (SUC1) not covered by the RETS guides, this allows for steric space between the two 200 or 30-base guide arms. Alignments were done using Clustal Omega and visualized with MView.

898
899
Supplemental Figure S3. Transient expression analysis of different sized RETS guides targeting
AtSUC1. Experiments were done by co-agroinfiltrating the various constructs into *Nicotiana benthamiana*
leaves with or without an exogenous copy of p35S:AtSUC1 (Trigger). Blue boxplots represent data from
leaf discs co-infiltrated with the trigger plasmid, grey boxplots represent leaf discs that did not have the
trigger co-infiltrated with the RETS construct. Y-axis represents the normalized mScarlet fluorescence
signal, i.e. the SUC1:RETS:mScarlet construct mScarlet fluorescence divided by the constitutive Venus
fluorescence that was measured simultaneously. Each point represents the data from an individual leaf
disc. The letter after the underscore on the x-axis label represents whether the guide sequence targets
the old site (o) -the same site as the 200-base construct- or the new site (n), the site that more specifically
binds to SUC1 with less cross-reactivity to SUC2/9. Statistical significance determined by Students' t-test:
* = p ≤ 0.05; ** = p ≤ 0.005; *** = p ≤ 0.0005.

911
912

914
915

Supplemental Figure S4. Quantitative RT-PCR measurements of AtSUC1 for seedlings of Col-0
and SUC1:RETS:mScarlet lines with different guide lengths. The grey boxplot represents the
distribution of SUC1 expression in wildtype Col-0 seedlings, the orange boxplots represent the SUC1
expression from SUC1:RETS:mScarlet lines. Each point represents a biological replicate of 5-8 pooled
seedlings. The letter after the underscore on the x-axis label represents whether the guide sequence
targets the old site (o) -the same site as the 200-base construct- or the new site (n), the site that more
specifically binds to SUC1 with less cross-reactivity to SUC2/9. Statistically significant differences to Col-0
determined by Students' t-test: * = $p \leq 0.05$; ** = $p \leq 0.005$; *** = $p \leq 0.0005$.

925
926

Supplemental Figure S5. Additional microscope images of SUC1:RETS:mScarlet lines with shorter guides directed to a more unique site in AtSUC1. Images were taken of adult rosette leaves with the mScarlet channel, focusing on the trichomes. A) Images of a line with 25-base guides. B) Images of a line with 20-base guides.

Supplemental Figure S6. Images of SAP and PAP1:RETS:mScarlet lines in the Venus channel to show uniform distribution of the T-DNA containing the RETS cassettes. A) Developing flower buds of 3 independent SAP:RETS:mScarlet lines with constitutive Venus expression. Compare to the tissue-specific distribution of mScarlet in Figure 5B. B) Adult flower sepals of 3 independent PAP1:RETS:mScarlet lines with constitutive Venus expression. Compare to the tissue-specific distribution of mScarlet in Figure 5D.

941 **Supplemental Table S1. Plasmid constructs used in this work.**

Plasmid code (P#)	Plasmid name	Plasmid use	Plasmid map link
P354	pTRANS_230d-pUBQ10:NLS-AtCas9:tHSP - pGmUbi:NLS-MCP-DREB2A-tUBQ1 - p35s:NLS-PCP-TPLN300:tNos	Cas9 Expression	https://benchling.com/s/seq-G4OTRmGF2rlvtg1XD3n8?m=slm-YpMR0Gk0KZGiCgmfjprl
P1699	Cas9_RETS-fixedIGS/IGSBS_Omega	Cas9:RETS:mScarlet with Omega enhancer	https://benchling.com/s/seq-yFpTyqR8Lbqq0ENl2P0L?m=slm-a1ATqcjVP8cdO6y94aKL
P1700	Cas9_RETS-fixedIGS/IGSBS_NO-Omega	Cas9:RETS:mScarlet NO Omega enhancer	https://benchling.com/s/seq-EjioHrHC792M8dNK78ps?m=slm-rBe20gX1IW9HGKFsIuKX
P1574	TtRz-AtSuc1-sensing(FIXED-IGS_and_IGSBS)_mScarlet-output_Venus-norm	SUC1:RETS:mScarlet 200nt guides	https://benchling.com/s/seq-hRlp6xrNZB73xSatXqlY?m=slm-XLVQzwZY1GL198ZUiWrr
P1735	AtSuc1-RETS_mScarlet-output_30baseGuides_oldTarg	SUC1:RETS:mScarlet 30nt guides	https://benchling.com/s/seq-gmR1CuYqzcwmf75gDk5E?m=slm-QP5bxh5DTAYFraO1FKHN
P1736	AtSuc1-RETS_mScarlet-output_25baseGuides_oldTarg	SUC1:RETS:mScarlet 25nt guides	https://benchling.com/s/seq-pYdtpCcHJk0hnL8tiEyV?m=slm-DGTS8KwQcXewqRwwTs7U
P1737	AtSuc1-RETS_mScarlet-output_20baseGuides_oldTarg	SUC1:RETS:mScarlet 20nt guides	https://benchling.com/s/seq-LzkQTRKXtgF89wDO0L3G?m=slm-vLITlLNva4I2QbFxbSv5
P1738	AtSuc1-RETS_mScarlet-output_30baseGuides_NEWTarg	SUC1:RETS:mScarlet 30nt guides NEW TARGET	https://benchling.com/s/seq-a7OpThgtvFLvTbqXsvBs?m=slm-1KtrrGL8oP34IAqiRBi3
P1739	AtSuc1-RETS_mScarlet-output_25baseGuides_NEWTarg	SUC1:RETS:mScarlet 25nt guides NEW TARGET	https://benchling.com/s/seq-1BeQ0mDyoOWyN3EJvnvE?m=slm-jY3uwxDvKNdPgGFiFluZ
P1740	AtSuc1-RETS_mScarlet-output_20baseGuides_NEWTarg	SUC1:RETS:mScarlet 20nt guides NEW TARGET	https://benchling.com/s/seq-8ayahgGo17DGo9LFdbRK?m=slm-6QFkbTVMqcGZVc0ZSfwH
P1800	p35s_AtSuc1(No-Introns)_tHSP	Exogenous trigger for SUC1:RETS:mScarlet agroinfiltration	https://benchling.com/s/seq-Q37qKbA9Vghg6uels8pm?m=slm-iRgBCT0Gz0DlP91046v1
P1696	AtPAP1-RETS_Split-mScarlet-output	PAP1:RETS:mScarlet 50nt guides	https://benchling.com/s/seq-pXjzRAe2eMAFp13touu8?m=slm-9Vn6hEtIVxOs4JflzNNu

P1697	AtSterileApetala-RETS_Split-mScarlet-output	SAP:RETS:mScarlet 50nt guides	https://benchling.com/s/seq-XgQMqqNkkeOoKySGQAlu?m=slm-g2Aenj7zfU4nSxCsK3Xe
P1781	Knat2-targeting_RETS-GAIrht-output	KNAT2:RETS:GAIrht 50nt guides	https://benchling.com/s/seq-b6doB1c35mnhimuNpxWr?m=slm-YbSZo7q1F5DN5tXN5mUH

951 **Supplemental Table S2. Primers used in this work.**

Primer code (O#)	Primer name	Primer use	Primer sequence
O2058	qPCR-AtSuc1(exon1)-f1	AtSUC1 qPCR	TCCCATGTGGATGCTTCTAATC
O2059	qPCR-AtSuc1(exon1)-r1	AtSUC1 qPCR	CACGACCCATCCAATCAGTATC
o2800	qPCR_PAP1-f	AtPAP1 qPCR	CCTACAACACCGGCACTAAA
o2801	qPCR_PAP1-r	AtPAP1 qPCR, gene specific cDNA synthesis	TCCAAGGCATGGAGGATTAAC
o843	TMV-5UTR_mScarlet-qPCR-f	mScarlet qPCR	CCAACAACAACAAACAACAAACAAC
o844	TMV-5UTR_mScarlet-qPCR-r	mScarlet qPCR	GATCCTTCCATGTGGACCTTAAA
o273	[PP2A]_q-f	Arabidopsis qPCR housekeeping gene, PP2aa3 (At1G13320)	AACGTGGCCAAAATGATGC
o274	[PP2A]_q-r	Arabidopsis qPCR housekeeping gene, PP2aa3 (At1G13320)	AACCGCTTGGTCGACTATCG
o275	[PP2A]-r	PP2aa3 gene specific cDNA synthesis	CTAGACATCATCACATTGTCAATAG
o2805	qPCR_mScarlet-acrossF66split-r2	mScarlet gene specific cDNA synthesis (across F66 RETS split)	TTATAAAGGCACGACTGCCATAC
o2318	BsaI-CATG_AtGAI(dFirst55aa)-f	Amplifying the front section of GAIrht from Col-0	aaggtctcaCATGATGTCTAATGTTCAAGAAGACGATCTTTC
o2319	AtGAI(dFirst55aa)_BsaI-TTCT-r	Amplifying the front section of GAIrht from Col-0	aaggtctcaAGAATCGCGTCACCGG
o2323	BsaI-CTCG_BackHalf-GAI(rht)-f	Amplifying the back section of GAIrht from Col-0	aaggtctcaCTCGCAATCAGTTCGCTATCGATTCGG
o2324	BackHalf-GAI(rht)_BsaI-ATAT-r	Amplifying the back section of GAIrht from Col-0	aaggtctcaATATCTAATTGGTGGAGAGTTTCCAAGC

952
953
954

955 **Supplemental Table S3. Arabidopsis target genes used in this work.**

Gene name	TAIR accession #
SUC1	AT1G71880
SUC2	AT1G22710
SUC9	AT5G06170
SAP	AT5G35770
PAP1	AT1G56650
KNAT2	AT1G70510
GAI1	AT1G14920