High-salt diet modulates endocrine regulation between
cortisol and FGF23

Matthias B. Moor[1,2], Klaudia Kopper[2,3], Antoinette Pechère-Bertschi[4,5], Michael S. Sagmeister[6],

Rowan S Hardy[7,8], Eric Feraille[2,4,5], Daniel G. Fuster[1,2], Johannes Loffing[2,3], Ganesh Pathare[2,3,9,10]

[1]Department of Nephrology and Hypertension, Inselspital, Bern University Hospital, University of

Bern, Bern, Switzerland.

[2]Swiss National Centre of Competence in Research "Kidney Control of Homeostasis",

Switzerland

[3]Institute of Anatomy, University of Zurich, Zurich, Switzerland

[4]Service of Nephrology and Hypertension, University Hospital Geneva, Geneva, Switzerland.

[5]Department of Cell Physiology and Metabolism, University of Geneva, Geneva, Switzerland.

[6]Institute of Applied Health Research, College of Medicine and Health, University of Birmingham,

Birmingham B15 2TT, UK.

[7]MRC Arthritis Research UK Centre for Musculoskeletal Ageing Research, University of

Birmingham, Birmingham, B15 2TT, UK.

[8]Department of Biomedical Sciences, School of Infection, Inflammation & Immunology, College of

Medicine and Health, University of Birmingham, Birmingham B15 2TT, UK.

[9]Institute of Veterinary Anatomy, University of Zurich, Zurich, Switzerland

[10]Department of Infectious Diseases and Public Health, City University of Hong Kong, Hong

Kong SAR

*Corresponding author:

Ganesh Pathare, BVSc, PhD

Department of Infectious Diseases and Public Health

Jockey Club College of Veterinary Medicine and Life Sciences

City University of Hong Kong, Hong Kong SAR

Email: gpathare@cityu.edu.hk

Keywords: High-salt-diet, cortisol (corticosterone), FGF23, bone and kidney

Competing Interests

The authors declare no potential conflicts of interest.

ABSTRACT

Excessive dietary salt intake is a global health concern, affecting cardiovascular, renal, and bone health. While the renin-angiotensin-aldosterone system (RAAS) is a known regulator of dietary salt-induced hormonal responses, the impact of adrenal cortisol remains unclear. Here, we performed a retrospective analysis in individuals (n=292) consuming a random diet. Dietary salt intake positively correlated with urinary cortisol and inversely correlated with plasma fibroblast growth factor 23 (FGF23), a bone-derived hormone regulating phosphate and vitamin D homeostasis. Controlled salt diets in healthy individuals confirmed a dose-dependent increase in urinary cortisol and suppression of plasma FGF23. In mice, oral corticosterone, a cortisol analogue, reduced circulating FGF23 levels. RNA-seq analysis of corticosterone-treated MC3T3 osteoblasts identified suppression of FGF23 via glucocorticoid receptor activation, anti-inflammatory pathways, and reduced osteoblast activity. Our findings reveal a novel endocrine cascade where high salt intake elevates cortisol and suppresses FGF23, with potential implications for bone, kidney, and cardiovascular health.

INTRODUCTION

Excessive dietary salt intake, a hallmark of the Western diet, is a global health concern associated with cardiovascular disease, bone loss, and autoimmune disorders, contributing significantly to morbidity and mortality[1–3]. Beyond its well-established effects on blood pressure, high salt intake disrupts metabolic signaling and hormonal responses[3]. The renin-angiotensin-aldosterone system (RAAS) is the primary hormonal pathway influenced by dietary salt[4]. Notably, adrenal-derived cortisol is known for its role in stress response and immunosuppression, but it may also contribute to salt homeostasis[5,6]. However, the mechanisms underlying this interaction and its impact on the pathophysiology of high salt intake are not yet fully understood.

Fibroblast growth factor 23 (FGF23), a bone-derived hormone, is central to phosphate and vitamin D homeostasis[7]. Elevated FGF23 is implicated in cardiovascular and kidney diseases, while its deficiency causes aging-like phenotypes in animal models[8,9]. Recent studies suggest a link between sodium homeostasis and FGF23 levels, but the mechanisms remain unclear[10–14.]. Given cortisol's emerging role in salt homeostasis, and its established effects on bone remodeling through glucocorticoid receptors (GR), it is crucial to investigate the relationship between dietary salt, cortisol, and FGF23.

Here, we hypothesize that dietary salt intake modulates endocrine regulation between cortisol and FGF23. To test this, we analyzed human cohort data and data from healthy individuals with and without dietary salt intervention, and performed mechanistic experiments in mice and osteoblast cell lines. Our findings reveal a previously unrecognized endocrine cascade, suggesting that dietary salt elevates cortisol signaling, which may contribute to FGF23 suppression.

RESULTS

1. High salt intake in humans activates cortisol signaling and suppresses FGF23 formation.

To investigate the association between dietary sodium intake and cortisol signaling, we reanalyzed data from the thoroughly phenotyped population of the Bern Kidney Stone Registry (BKSR), a single-center observational cohort of kidney stone formers[15]. We found that 24-hour urinary sodium, a reliable surrogate marker of dietary sodium intake, was associated with 24-h urinary cortisol (Fig. 1A). Further, urinary sodium was negatively associated with plasma FGF23 levels (Fig. 1B). Notably, urinary cortisone and cortisol tended to show an inverse relationship with plasma FGF23; however, only the association with urinary cortisone but not cortisol reached statistical significance (Fig. 1C, D). After adjusting the models for renal function, age, sex and body mass index in multivariable regression, the association with sodium remained whereas the association with cortisone was attenuated and did not remain significant (p=0.12) (Figures 1E-F).

Next, we analyzed serum and urine samples from a prospective crossover-controlled study involving healthy male volunteers[16]. These participants underwent three 7-day dietary interventions: a low-salt diet (LSD, 3 g NaCl/day), a normal-salt diet (NSD, 6 g NaCl/day), and a

high-salt diet (HSD, 15 g NaCl/day), with constant potassium intake (Fig. 1G). Dietary sodium

intake dose-dependently increased both 24-hour urinary sodium excretion (Fig. 1H) and cortisol

levels (Fig. 1G). Interestingly, dietary sodium also dose-dependently suppressed both intact and

cFGF23 levels, with a more pronounced effect observed in assays measuring cFGF23 (Fig. 1I, J).

Importantly, no significant differences in serum or urinary phosphate levels were observed across

the dietary interventions (Fig. 1K, L).

2. Corticosterone suppresses FGF23 formation by GR activation, anti-inflammation, and

repression of osteoblast activity

C57Bl6 wild-type mice were administered corticosterone orally for 2 weeks. At the end of the

treatment period, blood samples were collected, and intact FGF23 (iFGF23) and C-terminal FGF23

(cFGF23) levels were measured. We found that corticosterone treatment significantly suppressed

both iFGF23 and cFGF23 levels in mice (Fig. 2A, B).

To investigate the mechanisms for FGF23 suppression by corticosterone, *in vitro* experiments were

conducted using MC3T3 cells. Treatment with 100 nM corticosterone for 24 hours significantly

reduced both cFGF23 levels in the cell supernatant and *Fgf23* mRNA expression in cell lysates

(Fig. 2C, D). *Fgf23* mRNA expression was dose-dependently suppressed when cells were treated

with 0–200 nM corticosterone (Fig. 2E). Treatment with hydrocortisone (100 nM), a synthetic

cortisol analogue, also resulted in a reduction in *Fgf23* mRNA expression (Fig. 2F). Pretreatment

with RU-486 (GR blocker) completely rescued the corticosterone-induced suppression of *Fgf23*

mRNA, suggesting that the effects of corticosterone on *Fgf23* expression are mediated through

GR activation (Fig. 2G).

RNA-seq analysis was performed on MC3T3 cells treated with 100 nM corticosterone for 24 hours.

Differential gene expression analysis revealed 269 upregulated and 245 downregulated genes

(padj < 0.05; $\log_2$FoldChange > 1), visualized in a volcano plot (Fig. 2H). Gene ontology (GO)

analysis of biological processes revealed key pathways, including "ossification," "inflammatory

response," "osteoblast differentiation," and "bone mineralization." (Fig. 2I). Further information is

provided in the supplementary datasets 1 and 2 for DEGs and in the complete list of GO pathways.

Genes associated with direct corticosteroid signaling, including *Sgk1*, *Fkbp5*, and *Per1*, were

significantly upregulated, highlighting GR activation[17] (Fig. 2H/I). Anti-inflammatory pathways were

prominently modulated, with a strong upregulation of classical corticosteroid-responsive genes

such as *Hp* (Haptoglobin), *Orm1/2*, and *Serpine1* and a downregulation of *Ednra*[18] (Fig. 2H/I).

Simultaneously, genes involved in osteoblast differentiation, such as *Runx2*, *Pth1r*, and *Ibsp*, were

downregulated, indicating reduced osteoblast activity and extracellular matrix remodeling[19].

Suppression of *Bglap* and *Tnfrsf11b* suggest enhanced bone mineralization and bone resorption

(Fig. 2H/I). These findings provide evidence that corticosteroids suppress FGF23 through

combined GR signaling, anti-inflammatory effects, and suppression of osteoblast activity (Fig.

2H/I).

DISCUSSION

Our study identifies a potential endocrine cascade by which high salt intake elevates cortisol signaling and suppresses FGF23 levels, revealing non-classical pathophysiological consequences of high-salt diets on systemic health.

We measured 24-hour urinary cortisol for its accuracy in reflecting systemic cortisol secretion, overcoming the limitations of diurnal variation and inter-individual variability in serum levels[20]. Prior interventional studies with smaller sample sizes suggested that dietary salt intake modulates urinary cortisol levels[5,6]. While real-world dietary sodium variations are less extreme than those in interventional settings, our retrospective analysis showed that even among individuals on random diets, urinary cortisol positively correlates with dietary salt intake. Two possible mechanisms may explain the cortisol rise with high salt intake. First, both aldosterone and cortisol are synthesized in the adrenal cortex, share initial biosynthetic steps, and their levels are affected due to competition for shared precursors and enzymes in the context of salt intake. Suppression of RAAS activity by high salt reduces the demand for aldosterone synthesis, which depends on the enzyme *CYP11B2*. This reduction may divert precursors such as 11-deoxycorticosterone toward cortisol synthesis, potentially increasing cortisol levels[1,21]. Second, a recent study in mice suggest high-salt intake elevates corticosterone levels due to activated hypothalamus-pituitary (HPA) axis and stress response[22].

We found that high salt-mediated cortisol signaling was negatively associated with plasma FGF23. In the retrospective analysis of individuals consuming random diets, urinary cortisone correlated more strongly with FGF23 than cortisol, likely due to its higher stability. While the association did not remain statistically significant after adjustment, potential residual confounding by renal function cannot be excluded. Notably, in a controlled dietary salt intervention in young adults, high salt intake consistently increased urinary cortisol and suppressed plasma FGF23, supporting this relationship under experimental conditions. FGF23 is primarily recognized for its role in phosphate and vitamin D homeostasis, but its connection to salt metabolism is of emerging interest[10–14]. Aldosterone, a key hormone of the RAAS, directly enhances FGF23 production in bone via mineralocorticoid receptors[10,13]. This was linked to a potential mechanism for FGF23 regulation through dietary salt intervention in previous studies[14]. Our findings suggest that elevated cortisol and reduced aldosterone may collectively contribute to FGF23 suppression in response to high salt intake. However, our study provides limited direct evidence for high salt-induced cortisol suppressing FGF23 levels and highlights the need for further investigation. Furthermore, elaborate *in vivo* experiments are warranted to determine the relative contributions of cortisol and aldosterone to FGF23 regulation under a high-salt diet. Previous studies with potent glucocorticoids dexamethasone and prednisolone have shown both stimulatory and inhibitory effects on FGF23 levels in mice[23,24]. In contrast, our results indicate that the naturally occurring hormone corticosterone suppresses FGF23 production in mice and osteoblast cells via GR

188 activation, anti-inflammatory signaling pathways, and transcriptional repression of osteoblast
189 activity.
190
The interplay between dietary salt, cortisol signaling, and FGF23 may have important systemic
implications. Both high salt intake and elevated cortisol levels contribute to hypertension,
autoimmune diseases, inflammation and metabolic syndrome, bone loss and fracture risks,
therefore they may create a vicious cycle and amplify the burden of chronic diseases[1,3]. Future
studies are warranted to explore pharmacological strategies to counteract glucocorticoid-mediated
effects of high-salt diets on bone and cardiovascular health. Furthermore, both cortisol and FGF23
are used in many diagnostic tests[20,25]. Our results suggest that variations in dietary salt intake may
influence the interpretations of these measurements.
199
200 In conclusion, our study uncovers an endocrine cascade wherein dietary salt intake elevates
201 cortisol signaling, suppressing FGF23 levels. These findings suggest an unconventional role of
202 dietary salt in systemic health and highlight the need for further study.
203
METHODS
Human cohort analysis
We retrospectively analyzed data from the BKSR[15], a single-center observational cohort consisting
of individuals. Plasma C-terminal FGF23 (cFGF23) levels were quantified using ELISA. Steroids
in urinary samples were quantified using mass spectrometric analysis on a gas chromatograph
7890A coupled to a mass-selective Hewlett-Packard 5975C detector. We analyzed data regarding
urinary and serum parameters from 292 individuals. Associations between urinary sodium,
cortisol/cortisone, and plasma FGF23 levels were assessed using generalized linear models, after
square-root or log transformation as appropriate based on residuals evaluation. Adjustments were
made for age, sex, estimated glomerular filtration rate (creatinine CKD-EPI) and body mass index.
214
Dietary salt intervention study
Urine and serum samples were employed from previous perspective, crossover-controlled study
comprising healthy male volunteers[16]. Participants underwent three 7-day dietary interventions:
low-salt diet (LSD, 3 g NaCl/day), normal-salt diet (NSD, 6 g NaCl/day), and high-salt diet (HSD,
15 g NaCl/day), with constant potassium intake. Serum and 24-h urine samples were collected
after each dietary phase. Concentrations of urinary free cortisol were measured using an ELISA
(Diametra), whereas iFGF23 and cFGF23 levels were measured using an ELISA (Quidel).
Changes in plasma hormone levels and urinary parameters across the dietary interventions were
compared using paired t-tests (n=12).
224
225 **Mouse experiments:**
226 C57Bl6 male mice were housed in standard conditions with ad lib access to standard chow and
227 water. All animal experiments were performed in accordance with the guidelines and regulations

approved by the University of Birmingham. Drinking water was supplemented with either corticosterone (100 µg/mL, 0.6% ethanol), or vehicle (0.6% ethanol), with an average consumption of 1.25 mg per day. Mice were sacrificed at the end of 2 weeks' treatment. Plasma iFGF23 and cFGF23 levels were measured using ELISA (Quidel). Control mice received vehicle treatment. Data were analyzed using unpaired t-tests.

In vitro studies

MC3T3-E1 subclone four mouse preosteoblast cells were cultured and differentiated as previously described[26]. Osteogenic differentiation was initiated by culturing the cells in a medium containing 50 µg/ml ascorbic acid and 10 mM β-glycerophosphate for 6 days. Upon differentiation, cells were treated with corticosterone or hydrocortisone (0–200 nM) for 24 hours as noted in results section. *Fgf23* mRNA levels were quantified using qPCR. FGF23 protein levels in the supernatant were measured by ELISA after concentrating the cell supernatant as described earlier[26]. For GR inhibition studies, cells were pre-treated with 50 nM RU-486 (Biotechne) for 1 hour, followed by corticosterone treatment. All experiments were performed in triplicate.

RNA-seq analysis

Differentiated MC3T3 cells were treated with 100 nM corticosterone for 24 hours (n=3 biological replicates). RNA was extracted, and sequencing was performed commercially at Novogene using the Illumina platform[26]. Differential gene expression (DEG) analysis was conducted using DESeq2, with adjusted p-values ($p_{adj} < 0.05$) and $\log_2$ fold-change > 1 considered significant. Gene ontology (GO) analysis was performed to identify enriched biological pathways as previously described[26].

Statistical Analysis

Data were presented as mean ± SEM. Statistical significance was determined using t-tests for two-group comparisons. Correlation analyses were conducted using Pearson's correlation coefficient. All analyses were performed using GraphPad Prism or RStudio.

Data availability

The raw RNA-seq data used in this publication have been deposited in NCBI's Gene Expression Omnibus and are accessible through accession number GSE304562. All data are available from the authors upon request.

Acknowledgments

The authors sincerely thank Monique Carrel for her excellent technical assistance. We are grateful to Prof. Franz Weber (Centre of Dental Medicine, University of Zurich) for providing the MC3T3-E1 cells, and to Prof. Mariusz P. Kowalewski (Institute of Veterinary Anatomy, University of Zurich) for critically reading the manuscript.

Author contributions

G.P. conceptualized research; M.B.M., A.P.B., M.S.S., R.S.H., E.F., D.F. and G.P. designed research; M.B.M., K.K., A.P.B., M.S. and G.P. performed research/reanalyzed human cohort data; J.L. contributed new reagents/analytic tools; M.B.M., K.K., A.P.B., M.S.S., R.S.H., E.F., D.F., J.L. and G. P. analyzed the data; G.P. wrote the manuscript; M.B.M., K.K., A.P.B., M.S.S., R.S.H., E.F., D.F. and J.L. reviewed and edited the manuscript; G.P. and J.L. were responsible and funding acquisition; G.P. was responsible for supervision.

Funding and additional information

This work was supported by the Swiss National Science Foundation through the National Center of Competence in Research NCCR Kidney. C.H. [grant number N-403-03-55 (to G. P.)], funds from the Clinical Research Priority Program HYRENE of the University of Zurich (UZH) and intramural funding of the UZH (to J.L.). M.S.S. received funding from the Medical Research Council (United Kingdom) as part of a Clinical Research Training Fellowship (MR/T008172/1).

REFRENCES

1. Jobin, K., Müller, D. N., Jantsch, J. & Kurts, C. Sodium and its manifold impact on our immune system. *Trends Immunol* **42**, 469–479 (2021).

2. Kawarazaki, W., Mizuno, R., Nishimoto, M., Ayuzawa, N., Hirohama, D., Ueda, K., Kawakami-Mori, F., Oba, S., Marumo, T. & Fujita, T. Salt causes aging-associated hypertension via vascular Wnt5a under Klotho deficiency. *J Clin Invest* **130**, 4152–4166 (2020).

3. Hunter, R. W., Dhaun, N. & Bailey, M. A. The impact of excessive salt intake on human health. *Nature Reviews Nephrology 2022 18:5* **18**, 321–335 (2022).

4. Schweda, F. Salt feedback on the renin-angiotensin-aldosterone system. *Pflugers Arch* **467**, 565–576 (2015).

5. Lewicka, S., Nowicki, M. & Vecsei, P. Effect of sodium restriction on urinary excretion of cortisol and its metabolites in humans. *Steroids* **63**, 401–405 (1998).

6. Wambach, G., Bleienheuft, C. & Bönner, G. Sodium loading raises urinary cortisol in man. *J Endocrinol Invest* **9**, 257–259 (1986).

7. Kuro-o, M. & Moe, O. W. FGF23-αKlotho as a paradigm for a kidney-bone network. *Bone* **100**, 4–18 (2017).

8. Hu, M. C., Reneau, J. A., Shi, M., Takahashi, M., Chen, G., Mohammadi, M. & Moe, O. W. C-terminal fragment of fibroblast growth factor 23 improves heart function in murine models of high intact fibroblast growth factor 23. *Am J Physiol Renal Physiol* **326**, F584–F599 (2024).

9. Shimada, T., Kakitani, M., Yamazaki, Y., Hasegawa, H., Takeuchi, Y., Fujita, T., Fukumoto, S., Tomizuka, K. & Yamashita, T. Targeted ablation of Fgf23 demonstrates an essential physiological role of FGF23 in phosphate and vitamin D metabolism. *Journal of Clinical Investigation* **113**, 561–568 (2004).

10. Pathare, G., Anderegg, M., Albano, G., Lang, F. & Fuster, D. G. Elevated FGF23 Levels in Mice Lacking the Thiazide-Sensitive NaCl cotransporter (NCC). *Sci Rep* (2018) doi:10.1038/s41598-018-22041-1.

11. Pathare, G., Föller, M., Daryadel, A., Mutig, K., Bogatikov, E., Fajol, A., Almilaji, A., Michael, D., Stange, G., Voelkl, J., Wagner, C. A., Bachmann, S. & Lang, F. OSR1-sensitive renal tubular phosphate reabsorption. *Kidney Blood Press Res* **36**, (2012).

12. Pathare, G., Föller, M., Michael, D., Walker, B., Hierlmeier, M., Mannheim, J. G., Pichler, B. J. & Lang, F. Enhanced FGF23 serum concentrations and phosphaturia in gene targeted mice expressing WNK-resistant spak. *Kidney Blood Press Res* (2012) doi:10.1159/000343393.

13. Zhang, B., Umbach, A. T., Chen, H., Yan, J., Fakhri, H., Fajol, A., Salker, M. S., Spichtig, D., Daryadel, A., Wagner, C. A., Föller, M. & Lang, F. Up-regulation of FGF23 release by aldosterone. *Biochem Biophys Res Commun* (2016) doi:10.1016/j.bbrc.2016.01.034.

14. Humalda, J. K., Yeung, S. M. H., Geleijnse, J. M., Gijsbers, L., Riphagen, I. J., Hoorn, E. J., Rotmans, J. I., Vogt, L., Navis, G., Bakker, S. J. L. & de Borst, M. H. Effects of Potassium or Sodium Supplementation on Mineral Homeostasis: A Controlled Dietary Intervention Study. *J Clin Endocrinol Metab* **105**, e3246–e3256 (2020).

15. Dhayat, N. A., Schneider, L., Popp, A. W., Lüthi, D., Mattmann, C., Vogt, B. & Fuster, D. G.
Predictors of Bone Mineral Density in Kidney Stone Formers. *Kidney Int Rep* **7**, 558–567
(2021).

16. Pechère-Bertschi, A., Olivier, V., Burnier, M., Udwan, K., De Seigneux, S., Ponte, B.,
Maillard, M., Martin, P. Y. & Feraille, E. Dietary sodium intake does not alter renal
potassium handling and blood pressure in healthy young males. *Nephrol Dial Transplant*
37, 548–557 (2022).

17. Mifsud, K. R. & Reul, J. M. H. M. Acute stress enhances heterodimerization and binding of
corticosteroid receptors at glucocorticoid target genes in the hippocampus. *Proc Natl Acad*
Sci U S A **113**, 11336–11341 (2016).

18. Barnes, P. J. How corticosteroids control inflammation: Quintiles Prize Lecture 2005. *Br J*
Pharmacol **148**, 245–254 (2006).

19. Ishida, Y. & Heersche, J. N. M. Glucocorticoid-Induced Osteoporosis: Both In Vivo and In
Vitro Concentrations of Glucocorticoids Higher Than Physiological Levels Attenuate
Osteoblast Differentiation. *Journal of Bone and Mineral Research* **13**, 1822–1826 (1998).

20. Chen, A. X., Haas, A. V., Williams, G. H. & Vaidya, A. Dietary sodium intake and cortisol
measurements. *Clin Endocrinol (Oxf)* **93**, 539–545 (2020).

21. Jobin, K., Stumpf, N. E., Schwab, S., Eichler, M., Neubert, P., Rauh, M., Adamowski, M.,
Babyak, O., Hinze, D., Sivalingam, S., Weisheit, C., Hochheiser, K., Schmidt, S. V., …
Kurts, C. A high-salt diet compromises antibacterial neutrophil responses through
hormonal perturbation. *Sci Transl Med* **12**, (2020).

22. Costello, H. M., Krilis, G., Grenier, C., Severs, D., Czopek, A., Ivy, J. R., Nixon, M.,
Holmes, M. C., Livingstone, D. E. W., Hoorn, E. J., Dhaun, N. & Bailey, M. A. High salt
intake activates the hypothalamic-pituitary-adrenal axis, amplifies the stress response,
and alters tissue glucocorticoid exposure in mice. *Cardiovasc Res* **119**, 1740–1750
(2023).

23. Delucchi, Á., Toro, L., Alzamora, R., Barrientos, V., González, M., Andaur, R., León, P.,
Villanueva, F., Galindo, M., Las Heras, F., Montecino, M., Moena, D., Lazcano, A., …
Michea, L. Glucocorticoids Decrease Longitudinal Bone Growth in Pediatric Kidney
Transplant Recipients by Stimulating the FGF23/FGFR3 Signaling Pathway. *J Bone Miner*
Res **34**, 1851–1861 (2019).

24. Feger, M., Ewendt, F., Strotmann, J., Schäffler, H., Kempe-Teufel, D., Glosse, P., Stangl,
G. I. & Föller, M. Glucocorticoids dexamethasone and prednisolone suppress fibroblast
growth factor 23 (FGF23). *J Mol Med (Berl)* **99**, 699–711 (2021).

25. Edmonston, D., Grabner, A. & Wolf, M. FGF23 and klotho at the intersection of kidney and
cardiovascular disease. *Nature Reviews Cardiology 2023* 1–14 (2023)
doi:10.1038/s41569-023-00903-0.

26. Radvanyi, Z., Yoo, E. J., Kandasamy, P., Salas-Bastos, A., Monnerat, S., Refardt, J.,
Christ-Crain, M., Hayashi, H., Kondo, Y., Jantsch, J., Rubio-Aliaga, I., Sommer, L.,
Wagner, C. A., … Pathare, G. Extracellular sodium regulates fibroblast growth factor 23
(FGF23) formation. *J Biol Chem* 105480 (2023) doi:10.1016/J.JBC.2023.105480.

 FIGURE LEGENDS:

 Fig. 1: High salt intake in humans elevates cortisol signaling and suppresses FGF23
 formation.
 (A) Positive correlation between 24-hour urinary sodium excretion (a marker of dietary salt intake)
 and urinary cortisol levels in the BKSR cohort. (B) Inverse correlation between 24-hour urinary
 sodium excretion and plasma cFGF23 levels in the BKSR cohort. (C) Inverse correlation between
 plasma cFGF23 levels and 24-hour urinary cortisone levels in the BKSR cohort. (D) Inverse
 correlation between plasma cFGF23 levels and 24-hour urinary cortisol levels in the BKSR cohort.
 The p-value indicates the statistical significance of the observed correlation. n= 292. (G) Schematic
 representation of the dietary salt intervention study design. (H) Urinary sodium excretion; (I) urinary
 cortisol levels after dietary salt intervention (low-salt, normal-salt, and high-salt diets). (J) Plasma
 iFGF23 levels; (K) plasma cFGF23 levels; (L) plasma phosphate levels after dietary salt
 intervention. (M) Urinary phosphate excretion after dietary salt intervention. Data are presented as
 mean ± SEM. Statistical significance is denoted with p-value (paired t-test). n=12 per group.

 Fig. 2: Corticosterone suppresses FGF23 levels via GR activation and repressed
 osteoblast activity.
 (A) Plasma iFGF23 levels and (B) cFGF23 levels in C57Bl6 wild-type mice treated with oral
 corticosterone (n=12 per group). (C) *Fgf23* mRNA expression in cell lysates and (D) FGF23 protein
 levels in cell supernatant after 100 nM corticosterone treatment for 24 hours in MC3T3 osteoblasts
 (n=3 per group). (E) Dose-dependent suppression of *Fgf23* mRNA expression in MC3T3
 osteoblasts treated with corticosterone for 24 hours (n=3 per group). (F) Suppression of *Fgf23*
 mRNA expression by 100 nM hydrocortisone after 24-hour treatment (n=3 per group). (G) Rescue
 of corticosterone (100 nM)-induced suppression of *Fgf23* mRNA expression by pretreatment with
 RU-486, a glucocorticoid receptor antagonist (n=3 per group). (H) Volcano plot showing
 differentially expressed genes (DEGs) in corticosterone-treated MC3T3 osteoblasts, highlighting
 significantly upregulated and downregulated genes (padj < 0.05; |log2FoldChange| > 1). Key
 genes implicated in GR activation, anti-inflammatory responses, and osteoblast activity are
 indicated. (I) GO enrichment (biological processes) analysis of DEGs. Pathways related to bone
 mineralization, inflammatory response, and osteoblast differentiation are highlighted in red. Data
 are presented as mean ± SEM, with statistical significance denoted with p-value.

FIGURES

Fig. 1

Fig. 2