**Title: Human single-cell atlas analysis reveals heterogeneous endothelial signaling**

Zimo Zhu[1,2,*], Rongbin Zheng[1,2,*,#] , Yang Yu[1,2], Lili Zhang[1,2], Kaifu Chen[1,2,#]

[1] Department of Cardiology, Boston Children's Hospital.

[2] Department of Pediatrics, Harvard Medical School.

[*] Equal Contribution.

[#] Corresponding: Kaifu.Chen@childrens.harvard.edu; Rongbin.Zheng@childrens.harvard.edu

**Abstract**

Endothelial cells (ECs) play complex roles across tissues and vessel types. Yet, systematic

investigations of EC heterogeneity in the combined context of vessel type and tissue

microenvironment are still largely lacking. We integrated over three million single cells of

scRNA-seq datasets in 15 human tissues and found that ECs in some tissues (e.g., heart and

kidney) exhibited greater tissue specificity, while others displayed more substantial vessel

specificity. We developed a computational pipeline to analyze cell-cell communications (CCC)

mediated by metabolites or proteins to explore microenvironmental regulation. Interestingly, our

results showed that CCC events involving ECs varied vastly across tissues, highlighting tissue-

specific EC interactions. Using topic modeling, we identified CCC patterns, termed CCC topics,

representing metabolite- and protein-mediated interactions between ECs and other tissue-

resident cells. Most CCC topics exhibited high tissue specificity, potentially explaining

microenvironmental regulations for EC heterogeneity. The work systematically investigates EC

heterogeneity and provides insights into how EC heterogeneity was regulated across diverse

tissue microenvironments.

**Keywords:**

Cell-cell communication, tissue specificity, human endothelial, heterogeneity, single-cell

**Introduction**

Endothelial cells (EC) line the cardiovascular vessels throughout the human body, performing essential roles in various biological processes, including tissue homeostasis, immune responses, and vascular health[1–4]. Despite ECs' common presence, they exhibit remarkable diversity in structure and function that varies by vessel types, anatomic location, and developmental stage [5], demonstrating heterogeneity within the vascular system. Major vessel types include the arterial, venous, capillary, and lymphatic vessels. For example, arterial ECs express markers such as ephrin-B2, which marks arterial identity in the arteries [6]. Venous ECs express EphB4, establishing venous characteristics in the veins [7]. EC also shows differences between tissue types. These differences allow ECs to perform tissue-specific molecular mechanisms, such as cardiac vascular morphogenesis in the heart, gas exchange in the lung, or filtration in the kidney [8]. Studies using single-cell RNA sequencing (scRNA-seq) have shown that ECs exhibit cellular heterogeneity and metabolic plasticity across tissues in mice [9–11] and humans [12–14]. However, there is still a lack of systematic comparison of EC heterogeneity across combined contexts of different vessels and tissues.

It has been reported that the heterogeneity of ECs is regulated by various molecular factors, including transcription factors (TFs) and other signaling molecules [15]. Early in embryogenesis, transcriptional programs driven by TFs, such as GATA2 and ETV2 [16,17], establish the endothelial identity and guide early EC lineage commitment. The differentiated ECs are later refined based on the tissue context in embryonic development. Besides transcriptional regulation, the microenvironment regulates cellular heterogeneity and functions [18]. It is known that microenvironmental factors, including extracellular matrix composition, mechanical forces, and neighboring cell types, modulate gene expression regulation in ECs [19,20]. These regulations make them adapt to the functional needs of respective tissues [20,21]. For instance, ECs in the heart undergo unique shear stress and mechanical forces that upregulate KLF2, promoting an anti-inflammatory gene expression profile [22]. These studies together suggested that the tissue microenvironment is pivotal in establishing tissue-specific EC transcriptomes. Yet, the comprehensive investigation of microenvironmental mechanisms for human tissue-specific ECs remains elusive.

We hypothesized that tissue-specific microenvironments mediated by CCC may regulate human EC heterogeneity across tissue types. CCC is a fundamental process through which cells cross-talk via signaling molecules, such as cytokines and metabolites. The intact CCC network

63  coordinates cell functions, including differentiation, development, and metabolism [23]. For
64  instance, the blood-brain barrier in the brain explicitly influences EC response to neuronal
65  signaling and inflammatory cues [24]. A unique Car4-high EC population was identified in the
66  lung primed to communicate with other alveolar cells during alveolar regeneration after lung
67  injury [25].
68
This study integrated over 3 million single cells from human scRNA-seq datasets in 15 tissues.
First, we systematically assessed the EC heterogeneity across tissues and vessel types.
Intriguingly, a higher heterogeneity was observed across vessel types compared to tissue types
in most tissues, except the heart, kidney, and intestine. Further, we identified tissue-specific EC
topics for metabolite and protein-mediated CCC using a machine learning algorithm, Latent
Dirichlet Allocation (LDA). Diving into specific CCC topics, we found that most identified
metabolite-sensor and ligand-receptor pairs were tissue-specific and have been reported as
vital in carrying out tissue-specific functions. Our work showed a systematic EC heterogeneity
analysis across vessel and tissue types. Moreover, the identified tissue-specific CCC topics hold
the potential to explain the EC heterogeneity regulated by tissue microenvironmental factors
and further provide insights into therapeutic target discovery for endothelial-related diseases.
80
**Results**
**Heterogeneity of EC across tissue and vessel types**
We integrated scRNA-seq data from 15 human tissue types (**Supplementary Table 1**) and
visualized the EC population using UMAP to investigate the human EC heterogeneity. During
the integration, we observed that batch effects may confound the cell clustering as different
projects dominate the separation into different clusters (**Supplementary Fig.1**). Exemplified by
cells from the heart tissue, the cells in the same tissue type or vessel type from different projects
were separated into different clusters (**Fig. 1A, B**). However, the ECs of the same tissue type
from different projects were no longer clustered separately after correcting the batch effects
(**Supplementary Fig.2**). Instead, ECs became clustered by tissue or EC vessel types (**Fig. 1C,**
**D**). Thereafter, we performed all follow-up analyses based on batch-corrected data.
Interestingly, the principal component analysis indicated that all ECs in the heart and kidney,
regardless of vessel types, appeared to be very different from those in the other tissue types
(**Fig. 1E**). However, most ECs tended to be grouped by vessel types rather than by tissue types,
such as the ones grouped into arterial and lymphatic ECs (**Fig. 1E**). To further confirm this
observation, we next quantified the EC similarity among tissue and vessel types using

unsupervised hierarchical clustering method. Two primary clusters (cluster 1 and cluster 2) were

observed, as shown by the largest cophenetic distance based on the height of the dendrogram

(**Fig. 1F**). In cluster 1, the distance between ECs was significantly greater within tissue type

than within vessel type, indicating that ECs were more similar within vessel type than within

tissue type. However, the ECs in cluster 2 showed a larger distance within a vessel type than

within a tissue type, meaning that ECs within a tissue type are more similar. These results

suggested that the human EC posed significant heterogeneity across vessel and tissue types.

Furthermore, some ECs were more heterogeneous across vessel types than tissue types, while

some ECs were more heterogeneous across vessel types.

**A computational workflow to unravel tissue-specific microenvironmental signaling**

While the EC population in cluster 1 was predominantly heterogeneous across vessel types,

they still showed substantial heterogeneity across tissue types within a vessel type. This raised

the question of how tissue-specific signals were associated with EC tissue heterogeneity.

Previous reports show that CCC can regulate tissue-specific EC functions [26]. To study tissue-

specific CCC patterns, we developed a computational workflow by integratively analyzing two

types of CCC events, namely, metabolite-mediated and protein-mediated CCC (**Fig.2A**). First,

cutting-edge bioinformatics tools were implemented to detect CCC events in scRNA-seq data of

each tissue. Specifically, MEBOCOST [27] was used for metabolite-mediated CCC detection,

while LIANA+ [28,29] was used to detect protein-mediated CCC events. Next, we applied topic

modeling, an unsupervised machine learning algorithm, to detect CCC topics among all CCC

events across tissues. Topic modeling was extensively used in natural language processing

tasks, such as summarizing documents based on groups of words. Here, we defined

"document" as tissue and "words" as a collection of CCC events. The output "topics" represent

CCC patterns that could be either housekeeping or tissue-specific across tissue types (**Fig. 2B**).

The model also outputs two matrices reflecting the probability of a CCC belonging to a topic and

the probability of a tissue including a topic, respectively (**Fig. 2C**). We identify tissue-specific

CCC topics based on the CCC topic probability vector across tissues. Using probability vectors

across CCC events, we further determined if any groups of CCC were highly concentrated on a

tissue-specific topic. Finally, the workflow further evaluates gene expression specificities of

CCC-related genes across tissue types.

**Great heterogeneity of EC-related CCC across tissue types**

We sought to explore the number of EC-associated CCC events across tissue types.

Interestingly, the number of CCC for EC varied across different tissues (**Fig. 3A**). Some tissues,

such as the ovary and intestine, showed a much higher number of CCC events than other

tissues, such as kidney and adipose. The observation was highly consistent between

metabolite-mediated and protein-mediated CCC, suggesting tissue specificity of EC

communication with resident cells. Since the number of CCC might be affected by EC

population size and tissue cell type number, we computed the EC percentage and total cell type

number across different tissues (**Fig. 3B**). Surprisingly, the EC percentage in a tissue showed

little correlation (R=-0.05) with the total CCC number (**Fig. 3C**), while the number of cell types

was strongly correlated with the number of CCC events in the tissue (R=0.76, **Fig. 3D**). This

was consistent with the prior knowledge that the diversity of cell types and their spatial

organization within a tissue contribute to the richness of CCC networks [23]. However, the

observation of varying CCC numbers across tissue types remained even when it was

normalized by the number of cell types in the tissue (**Fig. 3E**). Taken together, these results

showed the diverse capability of EC to receive CCC signaling in different tissues, indicating

tissue-specific CCC patterns.

**Topic modeling revealed tissue-specific metabolite-mediated CCC patterns**

Functional metabolite-sensing machinery facilitates the crosstalk between the cellular biological

network and its tissue microenvironment cues [30]. Recent studies have highlighted dynamic

mechanisms with metabolites acting as signaling mediators to modulate processes like

inflammation and cancer progression [31]. To study EC tissue-specific heterogeneity under the

context of metabolite-mediated CCC, we applied topic modeling using communications detected

in each tissue and identified 15 metabolite CCC topics (**Supplementary Table 2**). Of 15 topics,

9 topics were highly tissue-specific. For instance, topic 11 was highly correlated with thymus.

Topic 5 was liver-dominant based on the topic-tissue probability vectors (**Fig. 4A**). Other less

tissue-specific topics were also identified, such as topic 7, topic 2, and topic 10, suggesting

some shared CCC networks for ECs across different tissues. Investigating each metabolite-

sensor pair in each topic, we observed that they were highly correlated with the specificity of

CCC topics (**Fig.4B**). To confirm this specificity further, we checked the number of

communications inferred by MEBOCOST for each tissue and found a matched pattern. The

metabolite-sensor pairs that appeared more frequently in a topic also tend to show a higher

number of CCC in the tissue related to that topic (**Fig.4C**). For example, we identified topic 13 to

be highly associated with the ovary (**Fig.4A**). Five metabolite-sensor pairs, including the

Cholesterol (metabolite)~RORA (sensor), L-Lysine~SLC7A1, Ornithine~SLC7A1,

Adenine~SLC35F5, and L-Cysteine~SLC43A2, exclusively showed high probability in topic 13

(**Fig.4B**), and showed more CCC events in the ovary (**Fig. 4C**). To inspect the results at the

expression level, we further assessed the communication score, which was inferred by the

metabolite enzyme and sensor expression. We found that all these pairs were highly expressed

in the ovary (**Fig.4D**). Interestingly, L-Lysine~SLC7A1 and Ornithine~SLC7A1 were the two

pairs that showed the highest specificity as they nearly solely expressed in the ovary but not in

other tissue. Consistently, many previous studies have demonstrated the importance of the

selected metabolite and sensor in carrying out tissue-specific functions. For example, a study

found lysine succinylation (Ksuc) as a newly discovered post-translational modification (PTM)

that plays an essential role in ovarian aging [32]. The paper pointed out that in the ovaries of

aged mice and those with premature ovarian insufficiency (POI), increased levels of Ksuc were

observed. Further, SLC7A1 was reported to be a key player in the ovarian tumor

microenvironment, influencing both cancer cell behavior and stromal fibroblast activation [33].

You et al. also found that in ovarian cancer (EOC), SLC7A1 inhibited the expression of the

CCL4 that was related to the immune infiltration microenvironment, suggesting a great

candidate for EOC therapy [34]. Together, we identified highly tissue-specific CCC topics

through topic modeling. The identified CCC events pose the potential to regulate EC

heterogeneity across tissue types.

**Topic modeling revealed tissue-specific protein-mediated CCC patterns**

Numerous studies have shown that protein ligand-receptor interactions are crucial in regulating

cell function and cell fate [35,36]. By applying topic modeling on protein-mediated CCC

landscapes inferred by LIANA+ [28,29], we identified 15 CCC topics involving ECs

(**Supplementary Table 2**). Interestingly, 13 of 15 protein-CCC topics were uniquely associated

with one tissue type (**Fig. 5A**). For example, topic 13 was strongly enriched with the thymus,

while topic 15 was strongly associated with the heart. Visualizing the ligand-receptor pairs that

participated in those communication events, we spotted that the ligand-receptor pairs were

highly topic-specific (**Fig. 5B**), further suggesting the tissue-specific CCC pattern. By checking

the number of communications associated with each pair (**Fig. 5C**), many ligand-receptor pairs

indeed showed a much higher number of communications in one tissue compared to other

tissues, consistent with the patterns observed above (**Fig. 5B**). These results suggested that the

identified ligand-receptor pairs in topics were tissue-specific, which indicated their unique,

important roles in regulating tissue-specific EC functions. Here, we have dived deeper into topic

15, which was a topic highly related to the heart (**Fig. 5A**). We focused on six ligand-receptor

pairs based on their significantly high probability score (topic-related) and high communication

number (tissue-related) (**Fig. 5B-C**), respectively. Those ligand-receptor pairs were

ADAM17~ITGB1, CALM1~RYR2, ADAM17~ITGA5, LAMA2~RPSA, ADAM17~NOTCH1, and

MAML2~NOTCH1. Interestingly, those pairs' ligand and receptor gene expression were also

highly expressed in the heart compared to other tissues (**Fig. 5D**). These identified ligands and

receptors have been reported to have important functions in the heart with in vivo and in vitro

experiments. For example, ADAM17 was essential to normal cardiovascular functions [37].

Elevated ADAM17 levels have been found in the lipid core of human atherosclerotic plaques

and mouse models with atherosclerotic lesions progression [38]. ADAM17 also plays a crucial

role in cardiac inflammation by cleaving and activating various proinflammatory cytokines and

their receptors [39]. It shows that ADAM17 is an important ligand that regulates normal heart

functions. Moreover, NOTCH1, which interacts with ADAM17 and MAML2 in the heart, was also

a crucial receptor in tissue formation and homeostasis. Ye et al. have revealed that the

NOTCH1 deficiency could cause defects in the proliferation of early human cardiomyocytes [40].

Roifman et al. found reduced NOTCH1 levels in both ventricular outflow tracts, suggesting that

NOTCH1 haploinsufficiency may cause bicuspid aortic valve and related cardiac anomalies [41].

A recent clinical study also showed an association between the NOTCH1 variants and cardiac

and extracardiac vascular anomalies [42]. This analysis provided insights into tissue-specific

communication patterns to EC for the ligand and receptor pairs and further poses the potential

to explain how EC heterogeneity was regulated at the tissue niche level.

**Discussion**

Tissue-specific EC functions have been widely studied within individual tissues and compared

between vessel types. It is unclear how human EC heterogeneity across different tissue types

was regulated systematically. An intriguing question is whether ECs should continue to be

grouped by vessel types or should they be further grouped by tissue types.

In this study, we investigated the microenvironments led by CCC as a potential factor to

regulate EC differences across vessel and tissue types. Analyzing human EC scRNA-seq data

specifically, we first identified different patterns of EC heterogeneity, which were shown by

tissue type and vessel type separately. Aside from the well-known knowledge that the EC

vessel types differed due to their specific locations and functions, ECs in some tissues, e.g., the

heart and kidney, were more different under the tissue context than vessel types. This finding

was comparable with other studies. Chen et al. showed that the embryonic EC heterogeneity
came from tissue origin and vascular-bed position [43]. They also found that embryonic ECs in
some tissues, such as the heart and liver, expressed tissue-selective markers.
235
In contrast, ECs in other tissues were more similar within each vessel type. Other groups also
identified EC heterogeneity, as shown by tissue-specific marker genes and enriched functions,
but with the mouse model [9,10]. In this study, we dived deeper into the microenvironment
regulation by CCC and found tissue-specific CCC patterns for EC in humans. We investigated
EC communications under various tissue contexts using an unsupervised machine-learning
algorithm and topic modeling. For metabolite-mediated and protein-mediated communications,
many CCC events inferred *in silico* were exclusively strong in one tissue compared to other
tissues. Based on the previous reports, we further verified that the identified ligands and
receptors were associated with essential functions in specific tissues. In summary, the identified
CCC events in each topic were promising candidates for further investigation to provide insights
into EC tissue-specific differences regulated by metabolite-mediated and protein-mediated
CCC.
248
**Method and materials**
**Data acquisition and process**
We downloaded over 3,000,000 single cells of 15 human tissues from the DISCO database[44],
including adipose, bladder, breast, gut, heart, intestine, kidney, liver, lung, ovary, skeletal
muscle, skin, stomach, testis, and thymus. The downloaded data in each tissue includes raw
gene expression counts and cell metadata annotation, such as cell type, project identification
number, disease status, etc. First, we filtered out disease samples to focus on data from normal
tissues, which resulted in over 2,000,000 single cells of 15 tissue types. The pre-processing,
including normalization, was done in Python by Scanpy (version 1.8.2) based on the raw count
matrix. The normalized data was used for the downstream analysis in this study, including EC
heterogeneity analysis and cell-cell communication analysis. In the EC heterogeneity analysis,
we retrieved single cells with cell types labeled as four major EC vessel types, namely, arterial,
capillary, lymphatic, and venous EC. We finally removed the batch effect using the Harmony
algorithm wrapped in the Scanpy package.
263
**Dimensionality reduction and clustering**

We did uniform manifold approximation and projection (UMAP), reducing the dimensionality to
show the clustering patterns across EC vessels and tissue types. We further investigated the
difference between EC vessel type versus tissue type by performing principal component (PC)
analysis using the Harmony-corrected PCs (50 in total). Results were visualized using PC1 and
PC2, which explained the most variance. Finally, hierarchical clustering based on tissue type
and vessel type was done with Scipy using 50PCs; the ward variance minimization algorithm
and Euclidean metric were used to build the model. Pairwise distance (Euclidean) was
calculated using the scikit-learn package for two major clusters for further quantifications.
273

**Cell-cell communication analysis**
To increase the computation efficiency, we down-sampled data from each tissue to contain only
10,000 cells and did normalization. Then, we utilized two bioinformatics tools to infer cell-cell
communications. MEBOCOST [27] predicted metabolite-mediated CCC events. LIANA+ [28,29]
was used to predict protein-mediated CCC. Each inference was performed on cell types within
each tissue type. Default parameters were used when running the framework. The input data
contained all cell types. We focused on communication events with a false discovery rate (FDR)
of less than 0.05. Then, we restricted the predicted communications with the type of
sender/source or receiver/target as ECs to further focus on EC-related CCC for the downstream
analysis.
284

**Topic modeling**
Focusing on how EC was regulated, we further filtered our data to only allow EC as the
receiver/target cells in a single communication event. Then, topic modeling was performed with
Latent Dirichlet Allocation (LDA) in R. The input matrix was tissue type by communication
events, filling with the number of communication events predicted in each tissue. The number of
topics in the LDA analysis was set at 15 based on the assumption that each tissue formed its
topic. The top 10 communication pairs (with the highest beta score from topic modeling) were
selected to visualize topic-specific patterns due to their strong probability in each topic. The
topic's word distribution, beta $\beta$, and communication number for each metabolite-sensor/protein-
receptor pair were normalized for each topic and tissue, respectively. To validate the tissue-
specific communication events, the communication score (product of the average metabolite
and sensor expression) from the MEBOCOST and the average expression of ligand-receptor
pairs from the CellPhoneDB were used, respectively.
298

**Quantification and statistical analysis**

Statistical analyses were performed using Python 3.10. The Euclidean distance matrix was used to calculate the correlation. The data were presented as mean in bar plots, and the box plots displayed quantiles and median.

**Code and data availability**

All the analyses were done using Python (version 3.8 for raw data processing and 3.10 for downstream analysis), except for the LDA analysis, which was performed using R (version 4.4.0). The script to reproduce all analyses in this study can be found at https://github.com/zhuzimoo/EC_project.

**Acknowledgments**

During the period of this project, K. C. is supported in part by the grants R01GM125632, R01GM138407, R01HL148338, R01HL133254, R01CA278832, and R01HL174928 to K. C., L. Z. is supported in part by the grants R01HL155632 to L. Z. and Additional Ventures (Single Ventricle Research Foundation) to L. Z, R. Z is supported in part by the grants K99HG013662 to R. Z. and NextGen Accelerator Award (Cell Discovery Network).

**Author Contributions**

K. C. and R. Z. conceived the project. Z. Z. performed all the analyses in this study and drafted the manuscript. All other co-authors read and revised the manuscript.

**References**

[1] Krüger-Genge A, Blocki A, Franke R-P, Jung F. Vascular endothelial cell biology: An
update. Int J Mol Sci 2019;20:4411.

[2] Pi X, Xie L, Patterson C. Emerging roles of vascular endothelium in metabolic homeostasis.
Circ Res 2018;123:477–94.

[3] Jalkanen S, Salmi M. Lymphatic endothelial cells of the lymph node. Nat Rev Immunol
2020;20:566–78.

[4] Iglesias MJ, Kruse LD, Sanchez-Rivera L, Enge L, Dusart P, Hong M-G, et al. Identification
of endothelial proteins in plasma associated with cardiovascular risk factors. Arterioscler
Thromb Vasc Biol 2021;41:2990–3004.

[5] Aird WC. Phenotypic heterogeneity of the endothelium: I. Structure, function, and
mechanisms. Circ Res 2007;100:158–73.

[6] Shin D, Garcia-Cardena G, Hayashi S, Gerety S, Asahara T, Stavrakis G, et al. Expression
of ephrinB2 identifies a stable genetic difference between arterial and venous vascular
smooth muscle as well as endothelial cells, and marks subsets of microvessels at sites of
adult neovascularization. Dev Biol 2001;230:139–50.

[7] Jadlowiec CC, Feigel A, Yang C, Feinstein AJ, Kim ST, Collins MJ, et al. Reduced adult
endothelial cell EphB4 function promotes venous remodeling. Am J Physiol Cell Physiol
2013;304:C627–35.

[8] Hellmut G A, Gou Young K. Organotypic vasculature: From descriptive heterogeneity to
functional pathophysiology. Science 2017;357. https://doi.org/10.1126/science.aal2379.

[9] Kalucka J, de Rooij LPMH, Goveia J, Rohlenova K, Dumas SJ, Meta E, et al. Single-cell
transcriptome atlas of Murine endothelial cells. Cell 2020;180:764–79.e20.

[10] Paik DT, Tian L, Williams IM, Rhee S, Zhang H, Liu C, et al. Single-cell RNA sequencing
unveils unique transcriptomic signatures of organ-specific endothelial cells. Circulation
2020;142:1848–62.

[11] Bondareva O, Rodríguez-Aguilera JR, Oliveira F, Liao L, Rose A, Gupta A, et al. Single-cell
profiling of vascular endothelial cells reveals progressive organ-specific vulnerabilities
during obesity. Nat Metab 2022;4:1591–610.

[12] Schupp JC, Adams TS, Cosme C Jr, Raredon MSB, Yuan Y, Omote N, et al. Integrated
single-cell atlas of endothelial cells of the human lung. Circulation 2021;144:286–302.

[13] Li Q, Zhu Z, Wang L, Lin Y, Fang H, Lei J, et al. Single-cell transcriptome profiling reveals
vascular endothelial cell heterogeneity in human skin. Theranostics 2021;11:6461–76.

354 [14] Geldhof V, de Rooij LPMH, Sokol L, Amersfoort J, De Schepper M, Rohlenova K, et al.

355   Single cell atlas identifies lipid-processing and immunomodulatory endothelial cells in

356   healthy and malignant breast. Nat Commun 2022;13:5511.

357 [15] De Val S, Black BL. Transcriptional control of endothelial cell development. Dev Cell

358   2009;16:180–95.

359 [16] Lugus JJ, Chung YS, Mills JC, Kim S-I, Grass J, Kyba M, et al. GATA2 functions at multiple

360   steps in hemangioblast development and differentiation. Development 2007;134:393–405.

361 [17] Shi X, Richard J, Zirbes KM, Gong W, Lin G, Kyba M, et al. Cooperative interaction of Etv2

362   and Gata2 regulates the development of endothelial and hematopoietic lineages. Dev Biol

363   2014;389:208–18.

364 [18] Potente M, Mäkinen T. Vascular heterogeneity and specialization in development and

365   disease. Nat Rev Mol Cell Biol 2017;18:477–94.

366 [19] Majewska A, Wilkus K, Brodaczewska K, Kieda C. Endothelial cells as tools to model tissue

367   microenvironment in hypoxia-dependent pathologies. Int J Mol Sci 2021;22:520.

368 [20] Leone P, Malerba E, Susca N, Favoino E, Perosa F, Brunori G, et al. Endothelial cells in

369   tumor microenvironment: insights and perspectives. Front Immunol 2024;15:1367875.

370 [21] Aird WC, Edelberg JM, Weiler-Guettler H, Simmons WW, Smith TW, Rosenberg RD.

371   Vascular bed-specific expression of an endothelial cell gene is programmed by the tissue

372   microenvironment. J Cell Biol 1997;138:1117–24.

373 [22] Dekker RJ, van Thienen JV, Rohlena J, de Jager SC, Elderkamp YW, Seppen J, et al.

374   Endothelial KLF2 links local arterial shear stress levels to the expression of vascular tone-

375   regulating genes. Am J Pathol 2005;167:609–18.

376 [23] Su J, Song Y, Zhu Z, Huang X, Fan J, Qiao J, et al. Cell-cell communication: new insights

377   and clinical implications. Signal Transduct Target Ther 2024;9:196.

378 [24] Armulik A, Genové G, Mäe M, Nisancioglu MH, Wallgard E, Niaudet C, et al. Pericytes

379   regulate the blood-brain barrier. Nature 2010;468:557–61.

380 [25] Niethamer TK, Stabler CT, Leach JP, Zepp JA, Morley MP, Babu A, et al. Defining the role

381   of pulmonary endothelial cell heterogeneity in the response to acute lung injury. Elife

382   2020;9. https://doi.org/10.7554/eLife.53072.

383 [26] Amersfoort J, Eelen G, Carmeliet P. Immunomodulation by endothelial cells - partnering up

384   with the immune system? Nat Rev Immunol 2022;22:576–88.

385 [27] Zheng R, Zhang Y, Tsuji T, Gao X, Wagner A, Yosef N, et al. MEBOCOST: Metabolite-

386   mediated Cell Communication Modeling by Single Cell Transcriptome. Bioinformatics 2022.

387 [28] Dimitrov D, Türei D, Garrido-Rodriguez M, Burmedi PL, Nagai JS, Boys C, et al.

Comparison of methods and resources for cell-cell communication inference from single-

cell RNA-Seq data. Nat Commun 2022;13:3224.

[29] Dimitrov D, Schäfer PSL, Farr E, Rodriguez-Mier P, Lobentanzer S, Badia-I-Mompel P, et

al. LIANA+ provides an all-in-one framework for cell-cell communication inference. Nat Cell

Biol 2024;26:1613–22.

[30] Wang Y-P, Lei Q-Y. Metabolite sensing and signaling in cell metabolism. Signal Transduct

Target Ther 2018;3:30.

[31] Wang Y-P, Li J-T, Qu J, Yin M, Lei Q-Y. Metabolite sensing and signaling in cancer. J Biol

Chem 2020;295:11938–46.

[32] Le M, Li J, Zhang D, Yuan Y, Zhou C, He J, et al. The emerging role of lysine succinylation

in ovarian aging. Reprod Biol Endocrinol 2023;21:38.

[33] You S, Han X, Xu Y, Sui L, Song K, Yao Q. High expression of SLC7A1 in high-grade

serous ovarian cancer promotes tumor progression and is involved in MAPK/ERK pathway

and EMT. Cancer Med 2024;13:e7217.

[34] You S, Zhu X, Yang Y, Du X, Song K, Zheng Q, et al. SLC7A1 Overexpression is involved

in energy metabolism reprogramming to induce tumor progression in epithelial ovarian

cancer and is associated with immune-infiltrating cells. J Oncol 2022;2022:5864826.

[35] An C, Wang X, Song F, Hu J, Li L. Insights into intercellular receptor-ligand binding kinetics

in cell communication. Front Bioeng Biotechnol 2022;10:953353.

[36] Funk MA, Leitner J, Gerner MC, Hammerler JM, Salzer B, Lehner M, et al. Interrogating

ligand-receptor interactions using highly sensitive cellular biosensors. Nat Commun

2023;14:7804.

[37] Xu J, Mukerjee S, Silva-Alves CRA, Carvalho-Galvão A, Cruz JC, Balarini CM, et al. A

disintegrin and metalloprotease 17 in the cardiovascular and central nervous systems.

Front Physiol 2016;7:469.

[38] Oksala N, Levula M, Airla N, Pelto-Huikko M, Ortiz RM, Järvinen O, et al. ADAM-9, ADAM-

15, and ADAM-17 are upregulated in macrophages in advanced human atherosclerotic

plaques in aorta and carotid and femoral arteries--Tampere vascular study. Ann Med

2009;41:279–90.

[39] Adu-Amankwaah J, Adzika GK, Adekunle AO, Ndzie Noah ML, Mprah R, Bushi A, et al.

ADAM17, A key player of cardiac inflammation and fibrosis in heart failure development

during chronic catecholamine stress. Front Cell Dev Biol 2021;9:732952.

[40] Ye S, Wang C, Xu Z, Lin H, Wan X, Yu Y, et al. Impaired human cardiac cell development

due to NOTCH1 deficiency. Circ Res 2023;132:187–204.

[41] Roifman M, Chung BHY, Reid DM, Teitelbaum R, Martin N, Nield LE, et al. Heterozygous
NOTCH1 deletion associated with variable congenital heart defects. Clin Genet
2021;99:836–41.

[42] Stanley KJ, Kalbfleisch KJ, Moran OM, Chaturvedi RR, Roifman M, Chen X, et al.
Expanding the phenotypic spectrum of NOTCH1 variants: clinical manifestations in families
with congenital heart disease. Eur J Hum Genet 2024;32:795–803.

[43] Chen J, Zhang X, DeLaughter DM, Trembley MA, Saifee S, Xiao F, et al. Molecular and
spatial signatures of mouse embryonic endothelial cells at single-cell resolution. Circ Res
2024;134:529–46.

[44] Li M, Zhang X, Ang KS, Ling J, Sethi R, Lee NYS, et al. DISCO: a database of Deeply
Integrated human Single-Cell Omics data. Nucleic Acids Res 2022;50:D596–602.

**Figure 1**

**Figure 1: EC demonstrated heterogeneity across tissue and vessel types.**

(**A-D**) UMAP clustering visualization of EC across tissue and vessel types before (**A**, **B**) and

after (**C, D**) batch effect removal, with color-coded for tissue type (**A, C**) and vessel types (**B, D**).

(**E**) PCA projection of EC across tissue type and vessel type. Color-coded for tissue type

(consistent with **1A** and **1B**) and shape-indicated EC vessel type.

(**F**) Hierarchical clustering showing tissue type and vessel type on the x-axis and cophenetic

distance on the y-axis. The dotted line separated the two major clusters. Colors and shapes

were used according to the PCA project in **E**.

(**G-H**) Pairwise distances were calculated for cluster 1 on the top (**G**) and cluster 2 on the

bottom (**H**). The box on the left shows the results grouping by tissue type and calculating the

vessel type distances, and the box on the right shows the results grouping by vessel types and

calculating the tissue distances. The median was shown with the bold line in the middle.

**Figure 2**

**Figure 2: Workflow for processing and analyzing multi-conditional scRNA-seq data**

(**A**) Scheme of analysis design.

(**B**) Explanation of topic modeling under the context of this study.

(**C**) Input and output matrix visualizations for the LDA algorithm.

**Figure 3**

**Figure 3: Analysis of metabolite-sensor and ligand-receptor CCC.**

(**A**) A bar chart shows the number of communications inferred *in silico*. The blue color indicates

the number of metabolite-mediated CCC (refers to the left y-axis scale). The orange color

indicates the number of ligand-receptor CCC (refers to the right y-axis scale).

(**B**) Bar chart showing the percentage of EC in the whole cell population and the number of cell

types included in a tissue. The red-dotted line indicates the average value under each category.

(**C**) The scatter plot shows the number of communications against EC percentage. Each dot

represents a tissue type. The regression line is shown in blue.

(**D**) The scatter plot shows the number of communications against the number of cell types.

Each dot represents a tissue type. The regression line is shown in blue.

(**E**) The bar chart shows the number of CCC events of every hundred cell types in each tissue to

normalize the effect of total cell type numbers across tissues.

**Figure 4**

**Figure 4: LDA inferred topics of metabolite-mediated CCC networks**

(**A**) Clustered heatmap showing the correlation of the LDA results using MEBOCOST inferred

CCC results. The X-axis indicated inferred topics and the y-axis labeled tissue type. The darker

the color, the stronger the correlation between the corresponding topic and tissue.

(**B**) Clustered heatmap of the normalized beta score of selected metabolite-sensor pairs (y-axis)

in each topic (x-axis). The x-axis label was aligned with that in **4A**. The warmer the color, the

higher the specificity, and vice versa.

(**C**) Clustered heatmap of the normalized number of communications for tissue-specific

metabolite-sensor pairs (x-axis). Tissue labels (y-axis) were aligned with that in **4A**. The warmer

the color, the higher the communication number, and vice versa.

(**D**) Violin plots of highly expressed metabolite-sensor pairs in topic 13. The color represented

the expression level based on the communication score. A missing violin indicated the

metabolite-sensor pair was not expressed in that tissue.

**Figure 5**

**Figure 5: LDA inferred topics of ligand-receptor CCC networks**

(**A**) Clustered heatmap showing the correlation of the LDA results using LIANA+ inferred CCC

results. The X-axis indicated inferred topics and the y-axis labeled tissue type. The darker the

color, the stronger the correlation between the corresponding topic and tissue.

(**B**) Clustered heatmap of the normalized beta score of selected ligand-receptor pairs (y-axis) in

each topic (x-axis). The x-axis label was aligned with that in **5A**. The warmer the color, the

higher the specificity, and vice versa.

(**C**) Clustered heatmap of the normalized number of communications for tissue-specific ligand-

receptor pairs (x-axis). Tissue labels (y-axis) were aligned with that in **5A**. The warmer the color,

the higher the communication number, and vice versa.

(**D**) Violin plots of highly expressed ligand-receptor pairs in topic 15. The color represented the

average of the pair expression level. A missing violin indicated the ligand-receptor pair was not

expressed in that tissue.

# Fig.1

# Fig.2

# Fig.3

# Fig.4

# Fig.5