African Pan Genome Contigs Expose Biologically Relevant Sequence Still Hidden from Human Reference Frameworks

Rachel Martini[1,2*], Abdulfatai Tijjani[3,4*], Kyriaki Founta[3,4,5*], Daniel Cha[6], Alexandria Awai[1,2,7], Sebastian Maurice[8], Jason A. White[1,2], Christopher E. Mason[9], Isidro Cortes-Ciriano[10,11], Nicolas Robine[7], Onyinye Balogun[9], Nyasha Chambwe[3,4,5+], Melissa B. Davis[1,2,7,12+]

[1]Institute of Translational Genomic Medicine, Morehouse School of Medicine, Atlanta, GA
[2]Department of Microbiology, Biochemistry & Immunology, Morehouse School of Medicine, Atlanta, GA
[3]Northwell, New Hyde Park, NY
[4]Feinstein Institutes for Medical Research, Northwell Health, Manhasset, NY
[5]Donald and Barbara Zucker School of Medicine at Hofstra/Northwell, Hempstead, NY
[6]Carnegie Mellon University, Pittsburgh, PA
[7]New York Genome Center, New York, NY
[8]City College of New York, New York, NY
[9]Weill Cornell Medicine, New York, NY
[10]European Molecular Biology Laboratory, European Bioinformatics Institute, Hinxton, Cambridge CB10 1SD, UK
[11]Somatic Genomics Programme, Wellcome Sanger Institute, Hinxton, UK
[12]Georgia Research Alliance, Atlanta, GA
*denotes equal contribution
+corresponding authors

Abstract

Human reference genomes underpin biomedical discovery but remain incomplete and biased toward European populations, constraining interpretation of genetic variation in underrepresented populations. Here we characterize African Pan Genome (APG) contigs totaling 296.5 Mb to define the sequence and functional landscape of genomic regions absent from current references. Most contigs align to the telomere-to-telomere (T2T-CHM13) genome and across 47 haplotype-resolved Human Pangenome Reference Consortium (HPRC) assemblies, with T2T-CHM13 placements enriched in centromeric and satellite repeats and overlapping 373 genes, including disease-associated loci. Mapping across HPRC assemblies revealed ancestry-associated contig enrichment, particularly in African genomes. Notably, 742 contigs remained unmapped under both stringent and relaxed criteria. These sequences are largely nonrepetitive and exhibit strong functional potential, including predicted protein-coding genes, CpG islands and transcriptional activity. Together, these results demonstrate that functionally relevant, ancestry-enriched genomic sequences remain absent from current references, with important implications for disease variant interpretation and precision medicine.

Word count: 149/150

Keywords: African Pan-genome, ancestry-associated genomic variation, reference bias, centromeric and satellite repeats, novel transcripts.

Introduction

The human reference genome determines which sequences align, what variants are called and annotated, and ultimately what biological data becomes visible to genomic research studies. Despite the availability of newer reference genome assemblies, GRCh37 and GRCh38 remain the primary references used for sequence alignment in human genomics research and clinical practice[1,2]. With unresolved gaps comprising approximately 7% of the GRCh38 genomes[3,4], it reflects a limited view of human diversity, primarily focusing on European ancestry[5]. This Eurocentric reference bias affects genomic medicine by essentially rendering data that is relevant to specific ancestry groups invisible[1,2,6,7]. A clear illustration of this bias comes from genome-wide association studies (GWAS) that use platforms designed from a Eurocentric reference, have historically been conducted in primarily European populations, and lead to limited applicable replication across any other ancestries[8–11]. This underrepresentation can hinder the mapping of genetic risk alleles, our understanding of regulatory gene networks that define disease biology and therefore limit the precision of genomic medicine.

Advances in long-read sequencing are transformative for genome science by overcoming the limitations of existing human reference genomes, resolving previously inaccessible regions of the genome. The Telomere-to-Telomere (T2T-CHM13) assembly provides a complete, gapless linear reference genome, including centromeres and segmental duplications[3]. However, both GRCh38 and T2T-CHM13 remain limited in ancestral diversity[5,12]. More recently, the Human Pangenome Reference Consortium (HPRC) has introduced a more diverse reference framework, initially generating 47 assemblies with wider representation across genetic ancestry groups. These assemblies also capture population structural variation allowing the building of a pangenome graph that offers better sensitivity in variant calling across diverse populations[4,13].

The African Pan Genome (APG) study by Sherman et al[14] provided a powerful test case for this challenge by identifying 296.5 million base pairs missing from GRCh38, prior to the release of the T2T-CHM13 and HPRC reference genomes. This landmark study performed whole-genome sequencing of 910 individuals of African descent, and subsequently assembled unmapped GRCh38 reads into 125,715 contigs[14]. While these contigs revealed previously uncharacterized genomic content specific to individuals of African ancestry, their biological importance and overlap with newer reference builds remained unclear.

In this study, we investigated whether unresolved APG contig sequences, absent in GRCh38, reflected low-complexity sequence of limited consequence or biologically meaningful genomic content still excluded from modern reference frameworks. We asked three questions: (1) how much ancestry-

enriched sequence absent from GRCh38 is recovered by T2T-CHM13 and HPRC assemblies; (2) whether recovered sequence overlaps functional genomic elements; and (3) whether contigs still missing from current references show evidence of coding, regulatory, or transcriptional activity. We determined that the majority of these contigs correspond to novel, functionally relevant regions of the genome. By evaluating the representation and functional roles of these sequences, our work emphasizes the risks of missing essential data in genomic research when we continue to rely on incomplete and Eurocentric reference genomes. Despite the options of more comprehensive reference genomes, GRCh38 is still the predominant genome reference used to date, which leads to persistent blind spots in our understanding of genome function, and perpetuates the omission of ancestry-specific biologically and clinically significant genomic content.

Results

To determine how much ancestry-enriched sequence remains outside of the current reference genome frameworks, we aligned 124,240 contigs assembled by Sherman et. al[14] against T2T-CHM13 and 47 HPRC linear assemblies and then layered functional annotation, repeat analysis, and RNA-seq evidence onto mapped and unmapped subsets **(see Methods, Figure 1, Figure S1)**. This design allowed us to distinguish sequences recovered by modern references from sequences that remain unresolved even after gapless reference genome advances.

APG contigs map to T2T-CHM13 regions absent from GRCh38

We first determined whether the 124,240 APG contig sequences missing from GRCh38 were recovered in the gapless T2T-CHM13 reference assembly. We found that 39.50% (49,070/124,240) of the APGs aligned to T2T-CHM13 at the Nearly Perfect (NP) threshold, with 94.45% of these T2T-CHM13 alignments mapping to regions unique from GRCh38 **(Figure 2A-B, Tables S1-3)**. Using the more relaxed Reasonably Good (RG) threshold increases the recovery of APG contigs mapping by 47.77% and likely reflects complexity in variation not represented as much in the assemblies available. The highest number of alignments to T2T-CHM13 were in Chromosomes 9, 21, and 14, and the least number of alignments to Chromosomes 18, 16 and Y **(Figure 2C)**. As anticipated, recovery was driven largely by previously unplaced contigs, where neither end of the contig sequence was previously resolved to a location on GRCh38[14] **(Figure 2D)**, underscoring how unmappable sequence to legacy references misses real sequence.

Re-evaluating APG contig placement from GRCh38 using the T2T-CHM13 reference

We further assessed APG contig alignment to the T2T-CHM13 assembly in the context of their prior placement status on the GRCh38 reference genome **(Table S4)**[14]. Of 124,240 contigs, 302 were previously classified as two-end (TE) placed, 1,246 as one-end (OE) placed, and 122,692 remained unplaced. Unplaced contigs showed the highest level of reasonably good (RG) alignments (48.08%), yet the lowest proportion of NP alignment (39.1%). A lower proportion of OE (26.00%) and TE (12.25%) had RG alignments, but had similar substantial NP alignment (51.32% and 55.06%, respectively) **(Table S4)**. These observations indicate that the contigs with greater sequence divergence were harder to align.

Chromosome concordance was nearly complete for TE contigs, with only one discordant NP alignment **(Figure 2E)**. Three TE placed contigs that had been placed on GRCh38 unknown chromosomes mapped to T2T-CHM13 Chromosomes 15 and 22 (n = 2, n = 1, respectively). In contrast, 53.21% of OE contigs with NP alignments mapped to different chromosomes in T2T-CHM13 than in GRCh38 **(Figure 2F)**, and 64.89% of these discordances involved acrocentric chromosomes. Similar patterns of discordant alignments were also observed among RG mapping contigs **(Figure S2)**. In line with our expectations, we find that the vast majority of APG contigs map to the T2T-CHM13 genome at locations that correspond to novel sequences not present in earlier versions of the reference genome.

APG contigs are recovered in gaps resolved by T2T-CHM13 that are repeat-rich and functionally important

As the overwhelming majority of the NP contigs (94.45%) mapped to regions unique to T2T-CMH13 **(Figure 2B)**, we sought to determine their potential overlap with known functional annotations. Most APG contigs recovered in T2T-CHM13 overlap centromeric and satellite-rich sequence (94.16%), consistent with longstanding limitations of short-read-based reference construction. The remaining contigs were annotated to genes (2.64%), CpG islands (3.61%) and composite elements (4.39%) **(Figure 3A, Table S5),** with similar enrichment patterns for the less stringent RG-aligned contigs **(Table S5, Figure S3)**.

T2T-CHM13-mapped APG contigs also overlap 373 annotated genes, with enrichment in immune, synaptic, and intracellular signaling pathways. Specifically, we found statistically significant enrichment in biological processes related to antigen presentation via the MHC class II protein complex. Cellular component and molecular function gene ontology processes related to neuronal recognition and the regulation or modulation of synaptic signaling were also enriched **(Figure 3B, Table S6)**. Interestingly, MHC class II enrichment was driven by the HLA locus on chromosome 6, where 84 contigs mapped to this 170Kb region on chromosome 6, with 44 contigs mapping to 4 HLA genes (*HLA-DRB3*, *HLA-DRB1*,

HLA-DQA1, *HLA-DQRB1*) **(Figure 3A)**; in line with the known challenges of read mapping to this highly

polymorphic region[15–17]. The main driver of contig overrepresentation in neuronal signaling processes

was liprin-alpha 1 (*PPFIA1)*, a scaffold protein involved in synaptic organization[18], with 33 contigs

mapping.

We also determined contig-gene overlaps with disease phenotypes in the Online Mendelian Inheritance

in Man (OMIM) database[19] and GWAS-linked traits. Specifically, 234 contigs mapped to 60 distinct genes

associated with OMIM disease phenotypes (**Table S7**), and 33 contigs overlapped with 113 previously

identified GWAS hits (**Figure S4A, Table S8**). Contig–GWAS hit overlaps were detected across multiple

chromosomes, with enrichment on chromosome 6, where 23 contigs overlapped GWAS hits linked to 23

distinct phenotypic traits (**Figure 3C**). Asthma was the top enriched GWAS trait with 13 Contig-GWAS

locus hits. This enrichment may reflect that the APG contigs were derived from analysis of an asthma

cohort (Consortium on Asthma among African-ancestry Populations (CAAPA))[14,20]. Asthma was the

second most enriched trait in RG mapped contigs, following autism spectrum disorder or schizophrenia

(91 Contig-GWAS hits) (**Figure S4B**). These results imply that functionally important traits or disease-

associated genetic variations in more diverse populations may be missed when using older versions of

the human reference genome due to mapping challenges, particularly in the context of short-read

datasets.

<u>HPRC assemblies recover APG contigs across diverse ancestral references</u>

Next, we aligned the APG contigs to the 47 haploid paternal assemblies from the Human Pangenome

Reference Consortium (HPRC) v1[4], which capture a broader range of population diversity than single

linear references, including complete genomes from African populations. Of the 47 HPRC assemblies,

24 are derived from individuals of African ancestry (AFR, **Table S9**).

Compared to T2T-CHM13 alignments, we observed a substantial improvement in APG contigs

placement. Overall 99.40% of the APG contigs achieved a RG alignment, and 82.91% met the NP

alignment threshold to at least one HPRC assembly (n = 123,498 and n = 103,003, respectively) **(Table**

S10). Notably, 80.44% of APG contigs have at least 1 NP alignment to an AFR ancestry HPRC assembly

(n = 99,944) **(Table S10)**.

Across all HPRC linear assemblies, mapping improvements were observed across previously defined

GRCh38 placement categories. At the NP threshold, 95.03% of TE contigs, 94.06% of OE contigs, and

82.76% of unplaced APG contigs aligned to at least one HPRC assembly (n = 287, n = 1172, n = 101,544,

respectively) **(Table S11-S13)**.

APG contig alignments to HPRC assemblies reveal population-associated enrichment

Comparing NP alignment results across reference assemblies, 49,020 contigs mapped to both T2T-CHM13 and the HPRC linear assemblies, whereas 53,983 contigs showed NP alignment exclusively to specific HPRC assemblies **(Figure 4A)**. Among these HPRC-only contigs, we observed a notable enrichment of alignments to assemblies from the AFR superpopulation (n = 7,969) ($p < 2.2$ e-16), whereas alignments to the other reference superpopulations were less frequent **(Figures 4B, Figure S5)**. Odds were 1.53x higher than to American (AMR) assemblies, and substantially greater versus East Asian (EAS, OR=2.74), European (EUR, OR=7.23), and South Asian (SAS, OR=8.55) assemblies **(Figure 4B)**.

Hierarchical clustering of the alignment patterns further revealed ancestry-associated grouping of individuals, with AFR assemblies clustering together and AMR assemblies forming a distinct secondary cluster of contig alignments **(Figure 4E)**. Unsupervised clustering of alignment patterns further highlighted contigs that mapped predominantly or exclusively to the AFR and AMR populations **(Figure S5A)**. The largest overlap between any two superpopulations consisted of 8,006 contigs that aligned to both the AFR and Admixed American (AMR) populations (**Figure S5B**). Although, AFR and AMR assemblies are more heavily represented in the HPRC v1 dataset (n = 24 and n = 16, respectively), this contig alignment overlap supports the population-associated nature of the APG contigs and suggests shared ancestry signals or structural variation between the AFR and AMR superpopulations. In contrast, shared mappings to the EAS, EUR, and SAS populations were considerably smaller, possibly indicating a limited representation of APG contigs within these genomes **(Figures 4B-C)**.

Expression quantification of the HPRC-mapped only contigs

We quantified strand-specific transcriptional coverage of the 53,983 contigs that mapped exclusively to HPRC assemblies using RNAseq data from three independent cohorts: 100 samples respectively from the 1000 Genomes Project (1KGP)[21] and the (TCGA) breast cancer (BRCA) project[22], and 44 samples from an African ancestry-enriched breast cancer cohort (Martini et al.)[23]. The 1KGP and TCGA cohorts included approximately balanced African (AFR) and European (EUR) ancestry samples, whereas the Martini et al. cohort was enriched for African ancestry samples (13 Ghanaian, 19 Ethiopian, 9 African American, and 3 European American patients, see **Methods**).

Overall, only a small fraction of contigs exhibited detectable transcription. At the lowest read depth threshold (T1, ≥1 read), approximately 5% of contigs showed non-zero strand-specific coverage in at

least one sample from 1KGP, with comparable rates in TCGA BRCA (**Figure S6A**). In the Martini et al.

cohort, Ethiopian samples exhibited the highest detection rate (~5.5%), followed by African American,

Ghanaian, and European American samples. Detection rates decreased with increasing read depth

stringency across all cohorts, with fewer than 1% of contigs retaining detectable coverage at T1000 or

above (**Figure S6A**). These patterns were consistent across both the forward and reverse strands.

We next assessed ancestry-associated differences in contig coverage using strand-specific Wilcoxon

rank-sum tests at the T100 threshold with Benjamini-Hochberg correction (**Figure S6B, Figure 4D**). In

1KGP, two contigs reached significance on the forward strand, including one enriched in AFR

(PDBU01024832.1) and one in EUR (PDBU01048398.1), with additional AFR-enriched signals observed

on the reverse strand (**Figure S7**). In TCGA BRCA samples, a larger number of differentially covered

contigs were identified on the reverse strand, predominantly enriched in AFR (**Figure S5B, Figure S8**).

Among 9 contigs showing differential coverage between the two populations (**Figure 4D**), all mapping to

AFR HPRC assemblies, whereas only 2 had previously been mapped to EUR assemblies. Notably, one

contig (PDBU01000224.1) mapped exclusively to AFR HPRC assemblies and not to any other reference

population assembly, consistent with its enriched expression in AFR samples, included Ethiopian and

Ghanaian samples in the Martini et al cohort (**Figure 4E**). Of these contigs, four were partially anchored

to GRCh38 reference, whereas six were unplaced.

<u>Sequence characteristics and functional landscape of below-threshold APG contigs</u>

21,187 APG contigs failed to meet the NP alignment criteria when mapped to the T2T-CHM13 and 47

HPRC linear reference assemblies (**Figure 4A**). We therefore re-assessed their mappability using the

more permissive RG alignment threshold. Under these criteria, 14,796 contigs (~25 Mbp) aligned to both

the T2T-CHM13 and HPRC assemblies, while 5,649 contigs (10.4 Mbp) aligned exclusively to HPRC

assemblies. The remaining 742 contigs, totaling ~1.5 Mbp, either failed to align or were aligned at below

RG threshold in all references. These sequences were designated as "below-threshold contigs" (BTCs)

(**Figure 5A, Table S14**).

To evaluate whether repeat content influenced this mappability pattern, we annotated repeat elements

across all contig subsets using RepeatMasker (**Table S15**). Contigs mapping to both T2T-CHM13 and

HPRC assemblies were highly enriched for repetitive sequences, with over 93% of their total length

composed of repeats (**Figure 5B**). These sequences were primarily comprised of satellite (~74%) and

simple (~17–19%) repeats, consistent with centromeric and subtelomeric genomic regions. In contrast,

the 742 BTCs exhibited substantially lower repeat content, averaging ~19%, with more than 80% lacking

detectable repetitive elements (**Figure 5B, Table S15**). LINEs (3.2%), SINEs (1.9%), and LTR elements

(0.89%) were the most prevalent repeats identified within BTCs.

Despite their limited mappability in linear reference space, BTCs showed substantial evidence of

functional potential. Of the 742 contigs analyzed, 473 (63.7%) contained at least one functional feature,

including predicted genes, CpG islands, or both **(Figure 5C, Table S16)**. Computational gene prediction

identified coding potential in 446 contigs (60.1%), while CpG island detection revealed regulatory features

in 324 contigs (43.7%). Notably, 297 contigs (40.0%) contained both predicted genes and CpG islands,

suggesting coordinated coding and regulatory architecture rather than low-complexity or artifactual

sequence. Only 269 contigs (36.3%) lacked detectable gene or CpG features. In total, these BTCs

encoded 544 predicted gene models and 573 CpG islands spanning 344.8 kb, indicating that sequences

poorly represented in linear references collectively harbor substantial predicted coding and regulatory

content.

Protein-level validation further supported the biological relevance of predicted genes encoded within

BTCs **(Figure 5D)**. Among predicted genes, 377 (70.8%) showed evidence of protein-coding potential

based on Pfam domain annotation and/or BLASTP homology. Strong support from both Pfam and

BLASTP was observed for 193 genes (36.2%), while 184 genes (34.6%) were supported by Pfam

domains alone and 11 genes (2.1%) by BLASTP alone. Only 156 predicted genes (29.2%) lacked

detectable protein support. The predominance of Pfam-supported predictions suggests that many

encoded proteins belong to conserved domain families, even when full-length homologs are not readily

identifiable by sequence similarity.

<u>RNA-seq mapping confirms that BTCs encode actively transcribed genes</u>

To evaluate whether BTCs contain transcriptionally active sequences, we aligned RNAseq reads that

failed to map to GRCh38 against a reference panel of BTC sequences using the predicted BTC genes

as annotations. BTC predicted gene expression was detected in 98/100 1KGP RNAseq samples, with 18

of the 544 predicted BTC genes expressed across multiple samples. Predicted gene g325, detected in

97 samples, showed the most widespread expression (**Figure 5E**). Several expressed BTC genes

exhibited BLAST and/or Pfam support for protein-coding potential, although *g325* lacked clear functional

annotation despite its ubiquitous expression (**Figure 5E**).

To assess functional potential in a disease context we examined TCGA BRCA RNAseq dataset. BTC

gene expression was detected in 99 of 100 samples across 36 predicted BTC genes, including 14 of the

18 genes observed in the 1000 Genomes samples. Expression patterns were concordant between
cohorts (**Figure 5E**), with predicted genes *g305* and *g131* showing both widespread expression and
protein-coding functional support. Analysis of the Martini et al. breast cancer cohort identified BTC
expression in 26/43 samples, with 18 samples showing gene-level expression across 10 BTC genes,
eight of which overlapped with the most commonly expressed genes in the 1000 Genomes and TCGA
cohorts. g325 remained the most frequently expressed BTC gene (**Figure 5D**).

Together, these results demonstrate that BTCs contain transcriptionally active loci absent from current
reference genomes, including the HPRC linear assemblies, highlighting persistent gaps in the human
genome representation, particularly for ancestrally diverse populations.

Discussion

Human reference genomes are foundational to genetic discovery, yet they are incomplete and
underrepresent global genomic diversity, particularly African diversity. A central implication of this study
is that completeness and representativeness are distinct. While T2T-CHM13 resolves sequence gaps
and HPRC improves recovery of ancestry-enriched APG contigs, neither fully captures the functional
breadth of human genomic diversity. Sequences present only in HPRC assemblies or remaining
unresolved are particularly consequential, as they are largely invisible to standard pipelines for read
alignment, annotation, and variant interpretation. Recent studies now make this point harder to dismiss
as a purely technical concern. Inclusive graph-based references reduce reference bias[24], ancestry-
matched references alter scRNA-seq gene assignment[25], diverse long-read transcriptomics reveals
systematic ancestry bias in gene annotation[21,26], and pangenome-based analyses improve sensitivity for
disease-relevant structural variants[27,28]. In this context, missing reference sequence represents a
mechanism by which biologically relevant variation is systemically obscured, particularly in populations
historically underrepresented in reference resource[29]. This distinction is critical as African genomes
harbor the greatest human genetic diversity, yet remain underrepresented in the reference frameworks
that shape discovery, annotation, and interpretation.

By systematically characterizing African Pan-Genome contigs derived from GRCh38 unmapped reads[14],
we show that even advanced linear assemblies, including T2T-CHM13 and HPRC assemblies,
incompletely capture ancestry-enriched genomic sequences. Alignment to T2T-CHM13 placed
approximately 40% of contigs, largely within genomic regions absent from GRCh38 and enriched for
centromeric satellites and repetitive elements, consistent with longstanding challenges in assembling

heterochromatic DNA using short-read technologies[30,31]. However, ancestry-specific sequences
remained incompletely represented, reinforcing the limitation of single linear references for structurally
complex and polymorphic loci. Mapping to HPRC assemblies substantially increased placement rates
and revealed ancestry-associated enrichment, particularly within African genomes, reflecting both
demographic history and the greater genetic diversity[32]. Importantly, many contigs overlapped annotated
genes, CpG islands, and pathways related to immune and cellular signaling, indicating that sequence
absent from current references can harbor biologically relevant elements. A subset of contigs remained
poorly mapped across all assemblies yet exhibited predicted coding features and transcriptional
evidence, demonstrating that functional sequence can remain undetectable in alignment-based analyses.
Such reference bias reduces sensitivity for variant detection and downstream analyses,
disproportionately affecting underrepresented populations[14,33,34].

Several limitations of our study should be considered. Functional annotation of BTCs relied on
computational predictions and may overestimate functional potential, particularly sequences lacking
homologs in existing databases[35]. Although RNA-seq data support the expression of a subset of
predicted genes, experimental validation across tissues and development contexts remains necessary.
Additionally, the APG contigs were derived from an admixed African American cohort, which may limit
generalizability to continental African populations. Uneven representation within the Phase I HPRC panel
may also influence observed ancestry-associated patterns.

Despite these limitations, our findings have important translational implications. Sequences absent from
widely used reference genomes can contain coding and regulatory elements, indicating that clinically
relevant variants within these regions are systematically overlooked[34]. Incorporating ancestry-enriched
sequence into population-aware and graph-based reference frameworks will improve variant discovery,
gene annotation, and transcriptomic profiling, ultimately enhancing the accuracy and equity of precision
medicine initiatives. In conclusion, characterization of APG contigs reveals that substantial functional,
ancestry-structured genomic sequence remains beyond current reference genomes. Addressing these
gaps will be critical to reducing reference bias, improving variant interpretation, and ensuring that
genomic discoveries benefit all global populations.

Methods

African Pan Genome Contigs Data Sourcing

We accessed the published African Pan Genome contigs sequences at GenBank with accession code PDBU01000000. All APG contig annotations were available via supplementary data from Sherman et al (2019)[14]. 125,715 APG contig sequences were published, however, 1475 were annotated as "dead" and were therefore excluded in all subsequent analyses (**Table S1**). In all analyses described, we proceeded with the remaining 124,240 APG contig sequences.

Mapping APG contigs to GRCh38, T2T-CHM13 and HPRC linear references

We sought to determine if any of the APG contig sequences are represented in the T2T-CHM13 and the Phase I HPRC linear reference assemblies (n = 47). We utilized a similar approach as outlined by Sherman et al[14], to map and assess the representation of APG contigs among unique reference genomes representative of African, South American, European, East and South Asian ancestries **(Figure S1)**. Specifically, we aligned the APG contigs to GRCh38.p10 as a benchmark to the Sherman et al analysis[14], the T2T-CHM13 v2.0 reference genome[3], and the Phase I HPRC haploid paternal linear assemblies v2[4]. Alignment to the various reference genomes and assemblies was performed using bwa-mem v0.7.17 with default parameters[36]. Alignments to individual locations within at least 300 bp were "chained" and analyzed as a single mapping output.

Assessment of APG contig mapping to GRCh38, T2T-CHM13 and the HPRC linear references

To assess the quality of APG contig alignment to the various reference genomes and assemblies, we determined the coverage and identity of each alignment using thresholds set by Sherman et al[14]. Coverage is defined as the fraction of aligned length over the entire contig length. Identity is the fraction of correctly aligned bases over the aligned length. For individual alignments within 300 bp, we took the aligned length to be the sum of each alignment length, and measured identity by taking the weighted average of identities across each alignment, using the following formulae:

$$Coverage \ = \ Aligned \ length \div \ Query \ Sequence \ Length$$

$$Identity \ = \ (Aligned \ length \ - \ Distance) \ \div \ Aligned \ Length$$

After calculating the coverage and identity, we applied two thresholds to determine if the mapping result was a "Reasonably Good" or "Nearly Perfect" alignment[14]. A Reasonably Good alignment was defined as having greater than or equal to 50% coverage, and greater than or equal to 80% identity.

$$Reasonably\ Good\ Alignment\ = \geq 50\%\ Coverage + \geq 80\%\ Identity$$

A Nearly Perfect alignment was defined as having greater than or equal to 80% coverage, and greater
than or equal to 90% identity.

$$Nearly\ Perfect\ Alignment\ = \geq 80\%\ Coverage + \geq 90\%\ Identity$$

Using these metrics and thresholds, we determined the number of contigs passing each threshold for
each reference assembly. A binary matrix was created to indicate the alignment status of each APG
contig relative to T2T-CHM13 and each HPRC assembly, with values of 1 indicating alignment and 0
indicating no alignment/threshold for alignment not met.

To further assess the representativeness of APG contigs across HPRC superpopulations, the HPRC
assemblies were categorized into superpopulation groups based on the designations from the 1000
Genomes Project: African (AFR), Admixed American (AMR), East Asian (EAS), European (EUR), and
South Asian (SAS)[37]. For HPRC samples that were not a part of the 1000 Genomes project (HG002 and
HG005), these were assigned to a corresponding superpopulation group (EUR and EAS, respectively)
(Table S9). A corresponding superpopulation-level and individual-level binary matrix was then generated,
summarizing whether each contig aligned with at least one assembly within each superpopulation **(Table
S12, S13)**. These matrices facilitated the evaluation of contig distribution and potential population-specific
enrichment across HPRC assemblies. Additionally, to statistically assess the population-specific
enrichment of APG contig alignments, a 2 × 5 contingency table was constructed, summarizing the counts
of aligned and unaligned contigs across the superpopulations. The alignment counts were normalized
according to the number of assemblies representing each superpopulation (AFR: 24, AMR: 16, EAS: 5,
EUR: 1, SAS: 1) to account for the differences in sample sizes. A chi-square test of independence was
conducted to determine the association between alignment status and population group. Odds ratios
(ORs) were calculated for each pairwise comparison between AFR and other populations (AMR, EAS,
EUR, SAS) to quantify the relative likelihood of contig alignment in AFR genomes.

<u>Functional enrichment of APG contigs across T2T-CHM13 genomic elements</u>
To explore the potential localization of contigs mapped to T2T-CHM13 regions within functional genomic
elements, we first obtained genomic coordinates for T2T-CHM13 unmasked CpG islands, NCBI RefSeq
genes, problematic regions, as well as T2T-CMH13 unique regions in comparison to GRCh38 from the
UCSC Genome Browser[38] Table Browser (https://genome.ucsc.edu/cgi-bin/hgTables) in *.bed* file format.
Additionally, we retrieved T2T-CHM13 RepeatMasker annotations generated by the T2T-CHM13

consortium (https://github.com/marbl/CHM13?tab=readme-ov-file). We then used BEDTools[39] v2.30.0 to

identify the presence of contigs across these various T2T-CHM13 genomic elements.

For co-localization of contigs with previously reported GWAS hits, we obtained GWAS associations from

the EMBL-EBI GWAS Catalog (https://www.ebi.ac.uk/gwas/docs/file-downloads). GRCh38 GWAS

variant positions were lifted over to obtain T2T-CHM13 coordinate system location by the T2T-CHM13

consortium (https://github.com/marbl/CHM13?tab=readme-ov-file). To identify genes associated with

phenotypes or diseases among those mapped by contigs, we downloaded OMIM's *Synopsis of the*

Human Gene Map file (https://www.omim.org/downloads/) following licensing and registration, and

extracted genes with reported phenotype or disease associations[19].

Gene ontology over-representation analysis

We extracted gene annotations using Ensembl gene models mapped to the T2T-CHM13 assembly,

identifying a total of 373 genes that overlap with the regions where the APG contigs aligned. We utilized

the R clusterProfiler package[40] v4.14.6 to compare our test gene set against a background gene set

composed of all annotated genes in the T2T-CHM13 assembly for enrichment. We evaluated the

overrepresentation of Gene Ontology (GO) terms across functional categories, including Biological

Process (BP), Cellular Components (CC), and Molecular Functions (MF). Statistical significance was

assessed using Benjamini-Hochberg-adjusted p-values with a false discovery rate (FDR) of less than

0.05. We generated bar plots to summarize the enriched GO terms related to the genes within the APG-

contig mapped regions.

Assessment of Repetitive Elements in APG Contigs

Repetitive DNA elements in the African pan-genome (APG) contigs were identified using RepeatMasker[41]

v4.1.6, a widely used tool for detecting and classifying interspersed repeats and low-complexity

sequences. The analysis was performed on three subsets of contig categories based on the Reasonably

Good alignment threshold. These are: (i) those aligning to both T2T-CHM13 and HPRC assemblies, (ii)

those aligning only to HPRC, and (iii) Below threshold contigs (those that failed to align or are aligned to

either reference below the reasonably good threshold. RepeatMasker was executed using the Dfam 3.7

and RepBase repeat libraries[42,43], utilizing the human-specific repeat database for precise classification.

The analysis was performed with the parameters-species set to human to confine the search to repeats

curated for the human genome.

Gene and CpG Island Predictions and Annotation

We deployed AUGUSTUS v3.5.0[44], an *ab initio* gene prediction software tool trained on human gene models to annotate potential protein-coding genes within BTCs. Contigs were individually processed, and gene structures including start and stop codons, exons, and coding sequences (CDS) were predicted. The resulting gene predictions were filtered to retain only transcripts with complete CDS features, representing putative protein-coding genes for downstream annotation (**Table S16**).

Additionally, to assess the epigenetic potential of APG contig sequences, we used CpGplot v6.6.0.0, a component of the European Molecular Biology Open Software Suite (EMBOSS, https://www.ebi.ac.uk/Tools/seqstats/emboss_cpgplot/). This tool detects CpG-rich sequences based on windowed GC content and the observed-to-expected CpG dinucleotide ratio. We used default thresholds commonly applied in genome-wide CpG prediction: a minimum island length of 200 bp, GC content ≥ 50%, an observed/expected CpG ratio of ≥ 0.6, and a 100 bp sliding window. The identified CpG islands were filtered to show only contigs with islands, where we quantified the number of islands predicted. The output was parsed and converted to BED format for comparative positional analysis and overlay with predicted genes (**Table S16**).

Translated protein sequences derived from AUGUSTUS gene models were queried against curated protein domain databases using profile hidden Markov models to identify conserved functional motifs and domain architectures, providing evidence for protein-coding plausibility beyond open reading frame structure alone[45]. In parallel, homology-based searches were performed against comprehensive protein sequence databases using sequence similarity alignment to detect significant matches to known proteins, allowing inference of evolutionary conservation and partial or divergent homology[46].

Quantification of Expressed Contig Sequences

To assess evidence of expressed sequences in APG contigs, we further analyzed RNA-seq data from three cohorts: (i) 1000 Genomes lymphoblastoid cell lines (n = 100; 52 African and 48 European)[37]; (ii) The Cancer Genome Atlas (TCGA) Breast Cancer (BRCA) primary tumors samples[22] (n = 100; 50 AFR and 50 EUR) using a genetic admixture level > 0.85)[47]; and (iii) primary breast tumor samples an African ancestry enriched cohort reported in Martini et al[23](n = 44; 9 African-American, 19 Ethiopian, 13 Ghanaian, 3 White). RNA-seq data was acquired in the following ways: 1000 Genomes *.fastq* files were compressed and aligned to GRCh38 genome version (*Homo_sapiens_assembly38.fasta* derived from the GATK resource bundle: https://console.cloud.google.com/storage/browser/genomics-public-data/resources/broad/hg38/v0;tab=objects?prefix=&forceOnObjectsSortingFiltering=false) using the *star_salmon* aligner and unmapped reads retrieved. For the TCGA BRCA samples, we obtained GRCh38

unmapped reads in a compressed *.fastq* format from the GRCh38 aligned *.bam* files we obtained from
GDC Commons, using *samtools* v1.21 to retrieve only the unmapped reads.

To assess whether HPRC aligned contigs (n = 53,983) harbored expressed sequences, we used STAR
v2.7.11b first to generate an index for the reference *.fasta* file containing the HPRC-mapped only contigs
sequences, and then to align the RNA-seq data from all three cohorts to the same contigs. Reads per
million mapped (RPM) normalized reads were obtained and stranded bedgraph output files were used
for downstream analyses. To evaluate expression of predicted BTC genes, GRCh38-unmapped reads
from the same cohorts were also aligned to a reference containing all BTCs. RNA-seq processing was
performed using the *Nextflow* v25.10.0 nf-core/rnaseq pipeline v3.21.0. The AUGUSTUS-derived gene
models provided in GTF format were used as gene annotations for the expression quantification step,
yielding read counts and transcripts per million (TPM) values per predicted gene and sample. Because
some samples were expected to lack detectable BTC expression, pipeline execution was configured to
continue processing samples for which STAR reported zero aligned reads to the BTC reference genome.

Acknowledgements

This work was made possible by the MacMillan Family Foundation as part of the MacMillan Center for
the Study of the Non-Coding Cancer Genome at the New York Genome Center. This work was performed
using the CPU and GPU compute nodes of the high-performance computing (HPC) cluster at Cold Spring
Harbor Laboratory (CSHL). This cluster is supported by US National Institutes of Health grant
S10OD028632-01, P01CA272295, R01CA266279.

We would like to acknowledge the Human Pangenome Reference Consortium (BioProject ID:
PRJNA730823) and its funder, the National Human Genome Research Institute (NHGRI). Some of the
results shown here are in part based upon data generated by the TCGA Research Network:
https://www.cancer.gov/tcga. We are grateful to our colleagues for their thoughtful discussions and
constructive feedback, which helped strengthen this study.

Author Contributions

464   **Conceptualization**: RM, AT, KF, MBD, NC; **Methodology**: RM, DC, AT, KF, AA, MBD, NC; **Software,**
465   **Formal analysis**: RM, AT, KF, DC, SM, AA; **Data Curation**: RM, AT, KF, DC, AA; **Visualization**: RM,
466   AT, KF, DC, AA; **Writing - Original Draft**: RM, AT, KF, NC; **Writing - Review & Editing**: RM, AT, KF,

467   DC, CEM, ICC, NR, SM, JW, OB, MBD, NC; **Supervision:** RM, NC, MBD; **Project administration:** RM,

468   NC; **Funding acquisition**: RM, NC, OB, MBD

Figures

Figure 1. Study Design

Figure 1. Study Design. APG contigs were aligned to the T2T-CHM13 reference genome and to 47 haploid paternal linear assemblies from the Human Pangenome Reference Consortium (HPRC; Year 1, Release 1). Based on predefined mapping quality thresholds, alignments were classified as "Reasonably Good" or "Nearly Perfect". Contigs that failed to align, or that produced only weak alignments to both T2T-CHM13 and HPRC references, were designated as "Below Threshold Contigs (BTC)". Contigs with "Nearly Perfect" alignments to T2T-CHM13 were further examined based on their functional annotations, whereas those aligning "Nearly Perfectly" to HPRC assemblies were used for continental ancestry analysis. For "Below Threshold Contigs (BTC)", we performed de novo gene prediction and CpG island identification, as well as quantification of expression of predicted genes. Repeat element analysis was conducted on both the "Below Threshold Contigs (BTC)" and the "Reasonably Good" alignments to any of the reference genomes.

Figure 2. Summary of APG Contig Alignment to T2T-CHM13

485   **Figure 2. Summary of APG Contig Alignment to T2T-CHM13.** (A) Pie chart showing distribution of
486   contigs with Nearly Perfect (NP, green), Reasonably Good (RG, blue) or below threshold (BTC, gray)
487   alignments to T2T-CHM13. (B) Stacked bar chart depicting T2T-CHM13 RG or NP alignments that map

to T2T-CHM13 unique regions compared to the GRCh38 reference (orange). Alignments to the non-
unique areas are shown in gray. (C) Butterfly plot of T2T-CHM13 alignment counts per chromosome
(right) and total aligned length (Mb) T2T-CHM13 unique content from GRCh38 (left, pink). All alignments
shown passed either the RG (blue) or NP (green) thresholds. (D) Bar chart of counts of unplaced APG
contigs mapping to T2T-CHM13 chromosomes. Heatmaps of GRCh38 chromosome placements and
T2T-CHM13 chromosome NP mapping results for (E) Two-End placed and (F) One-End placed APG
contigs.

Figure 3. Distribution of Functional Feature Overlap of T2T-CHM13 Mapped Contigs.

Figure 3. Distribution of Functional Feature Overlap of T2T-CHM13 Mapped Contigs. (A) Placement
of nearly perfect aligned contigs across T2T-CMH13 chromosomes overlaid across various annotation
categories. Contigs are highlighted in red. Enrichment was examined across T2T-CMH13 unique regions
in comparison to GRCh38 (yellow), CpG Islands (green), RefSeq annotated genes (blue), composite
elements (pink), and centromeres/satellite repeats (brown). A zoomed-in 170 Kb region (black dashed-
line box) at the chromosome 6 HLA locus with the highest number of mapped contigs across genes is
shown at the bottom of the panel. (B) Significantly overrepresented Gene Ontology terms (BP: Biological
Processes, CC: Cellular Component, MF: Molecular Function) among the 373 T2T-CMH13 genes with
at least one APG contig nearly perfectly aligned to them. Bars represent enrichment significance as -
log10(FDR), and color indicates fold enrichment. (C) Overlap of APG contigs with GWAS hits on
chromosome 6. Each dot represents a GWAS trait, with the y-axis indicating the number of contigs
overlapping with GWAS hits (polymorphisms) for that trait. Trait labels are shown, and all traits are
colored according to their parent trait category.

Figure 4. Summary of APG Contig Alignment and Expression Patterns across HPRC

Assemblies

Figure 4. Summary of APG Contig Alignment and Expression Patterns across HPRC Assemblies
(A) Venn diagram depicting the overlap of NP APG contig mapping to T2T-CHM13, HPRC linear assemblies. (B) Odds ratios (OR) comparing the likelihood of APG contigs alignment in African (AFR) genomes versus other superpopulations. (**C**) Heatmap showing alignment across individual linear assemblies from African (AFR) and Admixed American (AMR) HPRC samples. (D) Characterization of the 9 contigs with differential RNAseq read coverage between AFR and EUR samples across 1000 Genomes and TCGA BRCA. Each row represents a contig, while the left color bar indicates the cohort in which it was identified as significant (purple, 1000 Genomes; blue, TCGA BRCA). HPRC mapping information across AFR and EUR assemblies, as well as exclusive AFR assembly mapping status and placement type relative to GRCh38 are shown for each differentially enriched contig (dark grey indicates

a positive hit). (E) Violin plots showing the percentage of contig length covered at ≥100× RNA-seq coverage depth across all 3 studied cohorts for 2 of the 9 contigs identified as differentially covered between AFR and EUR individuals: PDBU01024832.1 (identified in 1000 Genomes cohort; top) and PDBU01000224.1 (identified in TCGA BRCA cohort; bottom). Each point represents an individual sample, and samples are stratified by population group. Crossbar shows the mean of the distribution. Asterisks denote statistically significant differences between groups (*p*-value less than 0.05 (*), 0.01 (**) or 0.001 (***)).

Figure 5. Functional characterization of APG contigs weakly or unmapped to gapless reference assemblies

Figure 5. Functional characterization of APG contigs weakly or unmapped to gapless reference assemblies. (A) Mapping categories of APG contigs to T2T and HPRC at Reasonably Good Threshold. (B) Repeat profiles of APG contigs across mapping categories. (C) Proportions of functional feature classes for below-threshold contigs (BTC), which lack confident alignment to any reference genomes utilized. Category classes are shown in the legend, including genes and/or CpG islands ("Both," "Gene only," "CpG Islands only," "Neither"). (D) Protein-level evidence of BTC predicted genes. Augustus gene

models were assessed using Pfam domain annotation and BLASTP homology and categorized as
supported by Pfam alone, BLASTP alone, by both, or unsupported. (E) Expression of BTC predicted
genes across 3 diverse cohorts: 1000 Genomes Project, TCGA breast cancer, and Martini et al. The x-
axis represents the number of samples per cohort expressing each of the displayed genes. Only the top
12 genes with the highest representation (TPM>0) across the 3 cohorts are shown. Functional prediction
per gene is also provided as a row annotation.

References

1. Li, H. *et al.* Exome variant discrepancies due to reference-genome differences. *Am. J. Hum. Genet.* **108**, 1239–1250 (2021).

2. Nie, J., Tellier, J., Tarasova, I., Nutt, S. L. & Smyth, G. K. T2T-CHM13 versus hg38: accurate identification of immunoglobulin isotypes from scRNA-seq requires a genome reference matched for ancestry. *NAR Genom. Bioinform.* **7**, lqaf074 (2025).

3. Nurk, S. *et al.* The complete sequence of a human genome. *Science* **376**, 44–53 (2022).

4. Liao, W.-W. *et al.* A draft human pangenome reference. *Nature* **617**, 312–324 (2023).

5. Miga, K. H. & Wang, T. The need for a human pangenome reference sequence. *Annu. Rev. Genomics Hum. Genet.* **22**, 81–102 (2021).

6. Clavell-Revelles, P. *et al.* Long-read transcriptomics of a diverse human cohort reveals widespread ancestry bias in gene annotation. *Genomics* (2025).

7. Rosenfeld, J. A., Mason, C. E. & Smith, T. M. Limitations of the human reference genome for personalized genomics. *PLoS One* **7**, e40294 (2012).

8. Fatumo, S. *et al.* A roadmap to increase diversity in genomic studies. *Nat. Med.* **28**, 243–250 (2022).

9. Popejoy, A. B. & Fullerton, S. M. Genomics is failing on diversity. *Nature* **538**, 161–164 (2016).

10. Martin, A. R. *et al.* Human demographic history impacts genetic risk prediction across diverse populations. *Am. J. Hum. Genet.* **100**, 635–649 (2017).

11. Carlson, C. S. *et al.* Generalization and dilution of association results from European GWAS in populations of non-European ancestry: the PAGE study. *PLoS Biol.* **11**, e1001661 (2013).

12. Aganezov, S. *et al.* A complete reference genome improves analysis of human genetic variation. *Science* **376**, eabl3533 (2022).

13. Aganezov, S. *et al.* A complete reference genome improves analysis of human genetic variation.

574   *Science* **376**, eabl3533 (2022).

14. Sherman, R. M. *et al.* Assembly of a pan-genome from deep sequencing of 910 humans of African

576   descent. *Nat Genet* **51**, 30–35 (2019).

15. Horton, R. *et al.* Variation analysis and gene annotation of eight MHC haplotypes: the MHC

578   Haplotype Project. *Immunogenetics* **60**, 1–18 (2008).

16. Meyer, D., C Aguiar, V. R., Bitarello, B. D., C Brandt, D. Y. & Nunes, K. A genomic perspective on

580   HLA evolution. *Immunogenetics* **70**, 5–27 (2018).

17. Robinson, J. *et al.* IPD-IMGT/HLA database. *Nucleic Acids Res.* **48**, D948–D955 (2020).

18. Ko, J., Na, M., Kim, S., Lee, J.-R. & Kim, E. Interaction of the ERC family of RIM-binding proteins

583   with the liprin-alpha family of multidomain proteins. *J. Biol. Chem.* **278**, 42377–42385 (2003).

19. McKusick-Nathans Institute of Genetic Medicine, John Hopkins University (Baltimore, MD). Online

585   Mendelian Inheritance in Man, OMIM. https://omim.org/.

20. Daya, M. *et al.* Association study in African-admixed populations across the Americas recapitulates

587   asthma risk loci in non-African populations. *Nat. Commun.* **10**, 880 (2019).

21. Taylor, D. J. *et al.* Sources of gene expression variation in a globally diverse human cohort. *Nature*

589   **632**, 122–130 (2024).

22. Cancer Genome Atlas Network. Comprehensive molecular portraits of human breast tumours.

591   *Nature* **490**, 61–70 (2012).

23. Martini, R. *et al.* African ancestry-associated gene expression profiles in triple-negative breast

593   cancer underlie altered tumor biology and clinical outcome in women of African descent. *Cancer*

594   *Discov.* **12**, 2530–2551 (2022).

24. Lin, M.-J., Iyer, S., Chen, N.-C. & Langmead, B. Measuring, visualizing, and diagnosing reference

596   bias with biastools. *Genome Biol* **25**, 101 (2024).

25. Yao, J. & Gazal, S. Evaluating genetic ancestry inference from single-cell transcriptomic datasets.

598   *HGG Adv.* **7**, 100564 (2026).

26. Clavell-Revelles, P. *et al.* Long-read transcriptomics of a diverse human cohort reveals ancestry

600  bias in gene annotation. *Nat Commun* **16**, 10194 (2025).

601 27. Groza, C. *et al.* Pangenome graphs improve the analysis of structural variants in rare genetic

602   diseases. *Nat Commun* **15**, 657 (2024).

603 28. Schloissnig, S. *et al.* Structural variation in 1,019 diverse humans based on long-read sequencing.

604   *Nature* **644**, 442–452 (2025).

605 29. Nassir, N. *et al.* A draft UAE-based Arab pangenome reference. *Nat. Commun.* **16**, (2025).

606 30. Miga, K. H. Centromere studies in the era of 'telomere-to-telomere' genomics. *Exp. Cell Res.* **394**,

607   112127 (2020).

608 31. Altemose, N. *et al.* Complete genomic and epigenetic maps of human centromeres. *Science* **376**,

609   eabl4178 (2022).

610 32. Campbell, M. C. & Tishkoff, S. A. African genetic diversity: implications for human demographic

611   history, modern human origins, and complex disease mapping. *Annu. Rev. Genomics Hum. Genet.*

612   **9**, 403–433 (2008).

613 33. Chen, N.-C., Solomon, B., Mun, T., Iyer, S. & Langmead, B. Reference flow: reducing reference

614   bias using multiple population genomes. *Genome Biol.* **22**, 8 (2021).

615 34. Sirugo, G., Williams, S. M. & Tishkoff, S. A. The missing diversity in human genetic studies. *Cell*

616   **177**, 26–31 (2019).

617 35. Yandell, M. & Ence, D. A beginner's guide to eukaryotic genome annotation. *Nat. Rev. Genet.* **13**,

618   329–342 (2012).

619 36. Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv [q-*

620   *bio.GN]* (2013).

621 37. 1000 Genomes Project Consortium *et al.* A global reference for human genetic variation. *Nature*

622   **526**, 68–74 (2015).

623 38. Perez, G. *et al.* The UCSC Genome Browser database: 2025 update. *Nucleic Acids Res* **53**,

624   D1243–D1249 (2025).

625 39. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features.

Bioinformatics **26**, 841–842 (2010).

40. Xu, S. *et al.* Using clusterProfiler to characterize multiomics data. *Nat. Protoc.* **19**, 3292–3320

(2024).

41. Smit, A. F. A., Hubley, R. & Green, P. *RepeatMasker Open-4.0.* (2013-2015).

42. Hubley, R. *et al.* The Dfam database of repetitive DNA families. *Nucleic Acids Res.* **44**, D81–9

(2016).

43. Jurka, J. *et al.* Repbase Update, a database of eukaryotic repetitive elements. *Cytogenet. Genome*

Res. **110**, 462–467 (2005).

44. Stanke, M., Steinkamp, R., Waack, S. & Morgenstern, B. AUGUSTUS: a web server for gene

finding in eukaryotes. *Nucleic Acids Res* **32**, W309–12 (2004).

45. Finn, R. D. *et al.* The Pfam protein families database. *Nucleic Acids Res.* **36**, D281–8 (2008).

46. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment search tool.

J. Mol. Biol. **215**, 403–410 (1990).

47. Carrot-Zhang, J. *et al.* Comprehensive analysis of genetic ancestry and its molecular correlates in

cancer. *Cancer Cell* **37**, 639–654.e6 (2020).

Supplemental Figures

Supplemental Figure 1. African Pan Genome Contig Alignment Computational
Workflow

Supplemental Figure 1. African Pan Genome Contig Alignment Computational Workflow.
Overview of the computational steps and resources utilized in the characterization of the
124,240 APG Contigs.

Supplemental Figure 2. GRCh38 chromosome placements and T2T-CHM13
chromosome mapping results for Reasonably Good mapping APG Contigs.

Supplemental Figure 2. GRCh38 chromosome placements and T2T-CHM13 chromosome
mapping results for Reasonably Good mapping APG Contigs. (A) Bar chart of RG mapping to
T2T-CHM13 chromosomes. Heatmaps of GRCh38 chromosome placements and T2T-CHM13
chromosome mapping results for (B) Two-End placed and (C) One-End placed APG contigs for
Reasonably Good alignments.

Supplemental Figure 3. Placement of nearly perfectly and reasonably good aligned
contigs across T2T-CMH13 chromosomes.

Supplemental Figure 3. Placement of nearly perfectly and reasonably good aligned contigs
across T2T-CMH13 chromosomes. Nearly perfectly aligned contigs are highlighted in red, while
reasonably good contigs are highlighted in turquoise. Enrichment was examined across T2T-
CMH13 unique regions in comparison to GRCh38 (yellow), CpG Islands (green), RefSeq
annotated genes (blue), composite elements (pink), and centromeres/satellite repeats (brown).

Supplemental Figure 4. Overlap of APG contigs with GWAS hits across

chromosomes.

Supplemental Figure 4. Overlap of APG contigs with GWAS hits across chromosomes. Each
dot represents a GWAS trait, with the y-axis indicating the number of contigs overlapping with
GWAS hits (polymorphisms) for that trait. Trait labels are shown, and all traits are colored
according to their parent trait category. Contig-GWAS hit overlap is visualized for (A) nearly
perfect (NP), and (B) reasonably good (RG) alignments to T2T-CHM13.

Supplemental Figure 5. Superpopulation-Level Summary of APG Contig Alignment
Patterns Across HPRC Assemblies

52
53　**Supplemental Figure 5.** Population-Level Summary of APG Contig Alignment Patterns Across
54　HPRC Assemblies. (A) Continental-level binary alignment patterns across HPRC linear
55　assemblies, where a contig is aligned to at least one member of each of the five human
56　superpopulations. (C) Upset plot depicting the number of intersections of contig alignments
57　across human superpopulations.

 Supplemental Figure 6. RNAseq alignment to HPRC-mapped only contigs.

Supplemental Figure 6. RNAseq Alignment to HPRC-mapped only contigs. (A) Percentage of

HPRC contigs (out of 53,983 total) exhibiting non-zero strand-specific coverage in at least one

sample, stratified by read depth threshold (T1–T50,000), RNA strand (forward/reverse), and

dataset. Three independent cohorts are shown: 1000 Genomes Project (AFR, n = 50; EUR, n =

50), TCGA breast cancer (AFR, n = 52; EUR, n = 48), and Martini et al. (African American, n =

9; Ethiopian, n = 19; Ghanaian, n = 13; White American, n = 3), (B) Volcano plots showing

differential strand-specific coverage between AFR and EUR populations at the T100 read depth

threshold in the 1000 Genomes and TCGA BRCA, assessed by row-wise Wilcoxon rank-sum

tests with Benjamini-Hochberg correction. The x-axis represents the difference in mean percent

coverage (AFR − EUR), and the y-axis represents statistical significance ($-\log_{10}$ adjusted p-

value). Dashed horizontal line indicates padj = 0.05. Orange points denote contigs enriched in

AFR; blue points denote contigs enriched in EUR; grey points are not significant (NS). Labeled

contigs are those reaching significance (padj < 0.05); contigs significant on both strands are

marked with asterisk.

Supplemental Figure 7. APG Contigs with Differential RNA-seq Coverage Across
Populations in the 1000 Genomes Cohort

79
Supplemental Figures 7. APG Contigs with Differential RNA-seq Coverage Across Populations
in the 1000 Genomes Cohort. Percentage of contig length covered at ≥100× RNA-seq coverage
depth across all 3 studied cohorts for the remaining contigs identified as significantly differentially
covered between AFR and EUR individuals not shown in Fig. 4E. Each point represents an
individual sample, and samples are stratified by population group. Crossbar shows the mean of
the distribution. Asterisks denote statistically significant differences between groups ($p < 0.05$ (*),
$p < 0.01$ (**), or $p < 0.001$ (***)).

 Supplemental Figure 8. APG Contigs with Differential RNA-seq Coverage Across

Populations in the TCGA BRCA Cohort

Supplemental Figures 8. APG Contigs with Differential RNA-seq Coverage Across Populations

in the TCGA BRCA Cohort. Percentage of contig length covered at ≥100× RNA-seq coverage

depth across all 3 studied cohorts for the remaining contigs identified as significantly differentially

covered between AFR and EUR individuals not shown in Fig. 4E. Each point represents an

individual sample, and samples are stratified by population group. Crossbar shows the mean of

the distribution. Asterisks denote statistically significant differences between groups ($p < 0.05$ (*),

$p < 0.01$ (**), or $p < 0.001$ (***)).

97