# Fine-Scale Structural Information Substantially Improves mRNA Therapeutic Stability Prediction

Soon Yi[1,2], Sara E. Ali[3], Yashrajsinh Jadeja[3], J. Wade Davis[3], Mihir Metkar[3,*]

[1]Center for RNA Science and Therapeutics, School of Medicine, Case Western Reserve University, Cleveland, OH, USA

[2]Department of Biochemistry, School of Medicine, Case Western Reserve University, Cleveland, OH, USA

[3]Moderna, Inc., Cambridge, MA, USA

[*]Correspondence should be addressed to M.M. (mihir.metkar@modernatx.com)

**ABSTRACT**

The success of COVID-19 mRNA vaccines has made the in-solution stability optimization

of mRNAs a key objective. However, we still lack a complete understanding of sequence

metrics that influence mRNA in-solution stability. RNA secondary structure plays a critical

role in protecting against hydrolysis, the primary degradation pathway under storage

conditions. Yet, the structural metrics that best guide stability-focused mRNA design

remain incompletely defined. Global metrics like minimum free energy and average

unpaired probability have improved mRNA stability but fail to capture local structural

variation relevant to RNA degradation. We demonstrate that base-pairing log odds

provide fine-scale, orthogonal insight that complements global metrics and improves

stability modeling. Further, by combining local and global features into a parsimonious

four-feature regression model, dubbed **St**ability **R**egression **A**nalysis using **N**ucleotide-

**D**erived features (STRAND), we achieve a greater than two-fold reduction in prediction

error compared to existing machine learning and deep learning approaches and

demonstrate robust generalization across diverse transcript contexts. This compact and

interpretable model provides an accurate and reliable framework for predicting mRNAs

in-solution stability.

## INTRODUCTION

Optimizing therapeutic mRNA requires not only effective protein expression but also in-vial stability that resists degradation during storage and delivery. Although the rapid mRNA-based vaccine development has highlighted the platform's potential[1], mRNA instability remains a critical barrier to broader deployment, particularly in settings lacking reliable cold-chain infrastructure such as developing countries[2]. Degradation primarily occurs through hydrolysis and transesterification[3,4] reactions in-solution during storage in-vial, typically initiating at structurally accessible nucleotides within unpaired loops or flexible regions[5,6]. Consequently, solution conditions such as temperature, pH, and ionic strength can drastically reduce mRNA half-life, thereby limiting therapeutic efficacy and significantly increasing manufacturing complexity and costs[7].

A key mRNA degradation determinant is its secondary structure, which arises from intramolecular base pairing. Base-paired regions can shield vulnerable sites from chemical or enzymatic attacks, extending RNA stability[6]. Accordingly, rational mRNA design has focused on engineering sequences that adopt favorable structural conformations. However, this design process requires reliable metrics that are predictive of RNA stability[8,9].

Traditional metrics such as free energy (FE), average unpaired probability (AUP), and GC content have been widely used to guide mRNA optimization[10,11]. Free energy minimization identifies the most thermodynamically stable structure an RNA can adopt[12,13]; AUP captures the average likelihood that nucleotides remain unpaired; and GC content correlates with duplex stability due to stronger G-C base pairing[8,14]. These metrics are computationally efficient and biologically relevant, and they have been

integrated into mRNA design tools[15,16] such as CDSFold[17], RiboTree[18], LinearDesign[19]

and DERNA[20]. However, despite their usefulness, these metrics do not fully explain

observed degradation differences between mRNA sequence variants as reflected by their

modest Spearman correlation coefficient (<0.6)[9], suggesting they fail to capture important

aspects of RNA structural ensembles[9].

Large-scale efforts have attempted to model mRNA degradation using both

statistical and deep learning approaches. For example, the dual crowd-sourcing study by

Wayment-Steele et al.[21] combined human- and machine-designed sequences to

benchmark predictive models of degradation, including those incorporating experimental

selective 2'-hydroxyl acylation analyzed by primer extension (SHAPE) data[6,22,23]. Despite

access to thousands of synthetic constructs and high-resolution structural probing data,

models achieved only moderate predictive accuracy when applied to full-length mRNAs,

with Spearman correlation coefficients (rho) ranging from 0.36 to 0.48[21]. These results

reflect the inherent difficulty of modeling mRNA degradation, which likely depends on a

combination of global and local structural features not fully captured by existing metrics.

To address this limitation, we propose a new metric: the base pairing log-odds

(LO), which quantifies the base-pairing likelihood at each position across the structural

ensemble by comparing the odds of being paired to unpaired. Unlike base-pairing

probability[24,25], which is bounded between 0 and 1 and compresses extreme values, the

log-odds transformation expands the dynamic range[26], particularly for high and low

probabilities, thereby amplifying meaningful structural differences that might otherwise be

obscured. LO reflects the local base-pairing probability distribution, offering finer

resolution than global metrics like FE or AUP. By analyzing mRNAs with similar global

scores but divergent in-solution half-lives, we show that LO captures functional

differences not apparent in traditional metrics. Finally, we demonstrate that incorporating

LO into a compact four-feature regression model, <u>S</u>tability <u>R</u>egression <u>A</u>nalysis using

<u>N</u>ucleotide-<u>D</u>erived features (STRAND), significantly improves mRNA stability prediction

over existing approaches and generalizes across diverse transcript contexts, providing

an interpretable framework for stability-focused mRNA design.

**RESULTS**

**Traditional RNA Structural Metrics Partially Explain In-Solution Stability**

A major goal in RNA therapeutic design is to extend mRNA half-life in-vial[27].

Existing studies have attempted to improve stability using sequence-derived structural

metrics, such as FE and AUP, which provide approximations of RNA secondary

structure[18]. However, whether these metrics alone are sufficient to predict mRNA

degradation rates remains an open question[9].

To assess the relationship between RNA structure and in-solution degradation, we

examined a dataset of Nano Luciferase (NLuc), enhanced green fluorescence protein

(eGFP), and multi-epitope vaccine (MEV) mRNAs with measured half-lives in-solution

(**Figure 1A**)[9]. NLuc sequences were selected for further analysis due to their broad half-

life range and the largest number of available sequence variants (N = 69), making them

an ideal test set for evaluating structural predictors.

Across the NLuc dataset, we observed moderate negative correlations between

half-life and FE (Pearson r = -0.52) as well as AUP (Pearson r = -0.47) (red and blue

points in **Figure 1B, 1C**). These correlations suggest that mRNAs with lower FE and AUP,

indicative of more stable secondary structures, tend to degrade more slowly. However,

while these metrics broadly capture structural trends, they fail to fully account for the

variability in observed degradation rates. For example, the NLuc sequence variants with

FE around -400 kcal/mol (N = 21, NLuc-21 set) showed a lower correlation than the global

comparison, with Pearson correlation values of -0.38 and 0.29, respectively (**Figure 1B,**

**1C**, highlighted in red). Moreover, linear regression models using FE and AUP as

predictors exhibited high residual errors, particularly at higher values of either metric

(**Figure 1D, 1E**). We also noted similar observations for eGFP CDS variants (N = 13;

**Figure S1**), in which eGFP sequences with FE near –500 kcal/mol show approximately

1.7-fold differences in their measured half-lives and sequences with AUP near 0.40 show

approximately 2-fold differences in their half-lives. These findings suggest that the

observed discrepancy between the traditional metrics and degradation rates is a general

phenomenon across different coding sequence sets. While global structure formation

plays a role in stability, additional sequence-dependent factors influence degradation in

ways that traditional metrics fail to capture.

To further investigate this variability, we applied Uniform Manifold Approximation

and Projection (UMAP) to reduce the three-dimensional structural feature space into a

lower-dimensional representation.[28] The resulting UMAP embedding, based on traditional

metrics, including minimum FE normalized by transcript length (FE/length), average

unpaired probability (AUP), and GC content, separated the NLuc sequences into distinct

clusters (**Figure 1F**). Normalizing FE by transcript length enabled comparison across

sequences of varying lengths, ensuring that the energy values reflect general structural

stability rather than absolute sequence length. The resulting UMAP clusters were

identified as high (H), medium (M), and low (L) based on the average in-solution half-life.

Despite the apparent separation in UMAP space, clusters H and M exhibited no

statistically significant difference in their half-life distributions (**Figure 1G**), underscoring

the inability of traditional metrics to resolve functionally distinct stability groups. This

suggests that additional sequence-derived features may be necessary to explain mRNA

degradation more accurately. To explore this further, we selected two NLuc sequences

with nearly identical FE and AUP values, but extreme differences in half-life, and aimed

to determine whether additional features could explain their stability discrepancies.

**Local Structural Features Differentiate Sequences with Similar Global Metrics**

The two sequences, X320004D1 and X930007D1, had FEs of -387.71 and -407.65

kcal/mol, a ~5% difference, while their AUP values differed by less than 3% (0.34 vs.

0.33). Notably, X930007D1 had a higher GC content (60.03%) than X320004D1 (52.04%),

though the direct effect of GC content on stability in this context remains uncertain. Their

measured half-lives varied by nearly three-fold (0.72 vs. 1.99 hours, respectively) (**Figure**

**2A**).

To investigate the structural basis for this difference, we predicted secondary

structures of both mRNAs using LinearPartition (**Figure 2A**)[29]. A heatmap visualization of

per-position base-pairing probabilities (BPPs) (**Figure 2B**) illustrated that, despite similar

global metrics, these sequences exhibited distinct base-pairing patterns along the

sequence and BPP compositions that may contribute to their degradation differences.

Specifically, X320004D1 formed a more branched secondary structure, with a higher

number of multiloops (41) and hairpin loops (24), which could introduce greater structural

flexibility and accessibility. In contrast, X930007D1 contained fewer multiloops (34) and

hairpin loops (17) but a higher number of inner loops (45 vs. 32 in X320004D1), forming

more extended helical regions (**Figure 2C**). To quantify these differences, we compared

the BPP distributions of the two sequences (**Figure 2D**). The BPP distribution of

X320004D1 showed a sharp bimodal pattern, with clear peaks near 0.0 (unpaired) and

1.0 (paired), suggesting more well-defined structural elements. In contrast, X930007D1

exhibited a broader, more continuous distribution, with a dominant peak near 1.0 but more

intermediate BPP values, indicating a less rigid structure.

These divergences between the global metrics and stability were not unique to this

sequence pair. Among sequences with higher stability (FE near -475 kcal/mol), we

observed similar disconnect between global metrics and measured half-lives. For

example, X310055T3 and X310005T1 showed highly distinct predicted structures

(**Figure S2A**) and measured half-lives (1.08 vs 2.03 h), despite having nearly identical

FE (–476.09 vs –499.53 kcal/mol), GC content (53.56 vs 52.16%), and AUP of 0.28. In

addition, the two sequences showed varying structural compositions (**Figure S2B**) and

BPP distributions (**Figure S2C, D**). Together, these examples demonstrate that global

metrics alone do not capture the local structural variation that may underlie differences in

RNA stability, motivating a more systematic analysis of BPPs across a broader sequence

set.

**Base-Pairing Probability Alone Does Not Fully Explain mRNA Stability**

To determine whether the structural patterns observed in X320004D1 and

X930007D1 extend to a broader dataset, we analyzed 21 NLuc variants (NLuc-21; red

points in **Figure 1B, C**) with similar FE and AUP values but differing in-solution half-lives.

Despite having nearly identical global structural metrics, these sequences displayed

distinct patterns of local base-pairing interactions (**Figure 3A, 3B**).

A heatmap of BPPs across these NLuc-21 sequences (**Figure 3A, 3B**) revealed

that, in general, each sequence possesses unique patterns of paired and unpaired

regions. Some sequences exhibited more extended single-stranded regions

(X320004D1), while others contained higher proportions of stable paired regions

(X310012T1). Interestingly, while some sequences with higher stability displayed

consistent base-pairing throughout the sequence (X930004D1), others with similar

stability had more localized variations in paired and unpaired regions (X930009D1 vs

X930007D1).

To quantify these differences, we examined whether the number of unpaired

positions (BPP < 0.2) or the presence of long unpaired patches (≥5 nucleotides)

correlated with half-life. Pearson correlation analysis showed that neither total unpaired

positions ($r = -0.47$, **Figure 3C**) nor long unpaired patches ($r = -0.32$, **Figure 3D**) provided

a significantly stronger correlation than FE or AUP. Moreover, our analysis showed no

significant improvement in Pearson correlation with different lengths of unpaired patches

(≥3 or ≥10 nucleotides; **Figure S3**).

These findings indicate that BPP alone does not account for mRNA stability

differences among sequences with similar global metrics. Although BPP provides a

continuous measure of pairing likelihood at each position, its bounded range (0 to 1)

compresses differences at the extremes, making it difficult to resolve functionally

meaningful distinctions, especially among positions with high or low pairing confidence.

Additionally, when summarized (e.g., by averaging in AUP), BPP can obscure the

contribution of unpaired positions, as zero-probability sites are effectively treated as

absence of signal rather than informative features. This suggests that a metric with

greater dynamic range may be needed to explain the stability variation that global metrics

and BPP miss.

**Log-Odds of Base-Pairing Captures RNA Stability Differences Beyond BPP**

To address this limitation, we employed a base-pairing log-odds (LO) metric, which

transforms BPP values into an unbounded scale ($-\infty$ to $+\infty$). Highly paired positions (BPP

> 0.5) yield positive LO values, while strongly unpaired positions (BPP < 0.5) yield

negative values. This log transformation expands the dynamic range at both extremes of

the BPP distribution, amplifying differences among positions with very high or very low

pairing confidence that are otherwise compressed on the 0-to-1 BPP scale.

To test whether LO captures RNA degradation patterns better than BPP, we first

examined per-position LO values for X320004D1 and X930007D1 (shown in **Figure 2**).

While both sequences had similar global structural metrics, LO values were distributed

differently, particularly in weakly structured regions (**Figure 4A)**. Notably, X320004D1

contained a greater number of strongly negative LO positions, indicating unpaired regions

that are likely more susceptible to degradation. In contrast, X930007D1 exhibited fewer

such sites, suggesting fewer vulnerable positions, consistent with its longer half-life.

A heatmap visualization of LO across NLuc-21 sequences (**Figure 4B, C**)

demonstrated that LO more effectively differentiated stable and unstable sequences

compared to BPP. Sequences with shorter half-lives exhibited more widespread strongly

negative LO values (red regions in **Figure 4C**), consistent with a greater number of

confidently unpaired positions. To quantify this, we focused on weakly paired regions (LO

< -1.363, derived from BPP curvature analysis) and found that the average LO (ALO) of

these positions showed a strong positive correlation with in-solution half-life (Pearson r =

0.78, **Figure 4D**), indicating that sequences with less extreme negative LO values in their

unpaired regions tend to be more stable.

However, when we computed the ALO across all positions, the correlation with

half-life was negative (Pearson r = -0.63, **Figure 4E**). Although this opposing direction

appeared counterintuitive, closer analysis of the heatmap in **Figure 4C** revealed that

sequences with shorter half-lives also contained more positions with extremely positive

LO values (blue regions; LO > 20). These sequences thus exhibit a more polarized LO

distribution, with greater representation at both extremes, while longer-lived sequences

show more moderate LO profiles overall. When averaged across all positions, the

extreme positive values in shorter-lived sequences inflate their ALO, producing the

observed negative correlation. Despite this reversal in sign, the magnitude of the ALO

correlation still exceeds that of FE or AUP, and the strong positive correlation in the

weakly paired regions (r = 0.78) provided a clear link between local structural features

and mRNA stability. Overall, these results demonstrate that the LO transformation reveals

structurally meaningful variation among sequences that appear similar to traditional

metrics.

**Multi-feature Regression with ALO Improves mRNA Stability Prediction**

Previous work by Wayment-Steele et al.[21] evaluated statistical and deep learning

models on these full-length transcripts (**Figure 1A**), achieving Spearman correlations of

0.36–0.48 on orthogonal degradation datasets[9,21]. Despite the advantage of large-scale,

high-resolution input, predictive performance on full-length sequences remained modest,

highlighting the challenge of generalizing across design contexts.

To test whether LO complements existing structural metrics, we incorporated ALO

alongside FE/length, AUP, and GC content in a unified UMAP embedding (**Figure 5A**).

This integrated model revealed improved separation between NLuc sequence clusters,

with all clusters achieving statistically significant differences in in-solution half-life (**Figure**

**5B**). Notably, the p-value between the high half-life (red) and medium half-life (green)

clusters improved from 0.62 without ALO (**Figure 1G**) to 0.023 with ALO (**Figure 5B**),

based on a Mann-Whitney test representing approximately a 27-fold increase in

significance. This demonstrates that LO contributes orthogonal information, enabling

better resolution of stability differences that traditional global metrics alone fail to capture.

Since no single metric fully explains RNA stability, we next sought to construct a

multi-feature regression model integrating key structural features. An ideal model should

balance explanatory power with generalizability while minimizing overfitting, particularly

given the small dataset size (N = 69). Given the strong correlations among features, we

selected a minimal, non-redundant feature set that captures complementary aspects of

RNA structure. FE (normalized by length) and AUP represent the global secondary

structure stability, which influences degradation. GC content captures sequence

composition effects on stability, as GC-rich regions tend to form stronger base-pairs. ALO

reflects local structural differences missed by traditional metrics, providing finer resolution

of base-pairing variability. This feature set ensures that we account for thermodynamic

stability, sequence composition biases, and fine-scale structural variability.

We next used Partial Least Squares Regression (PLSR)[30] to assess the linear

relationships between the feature sets and in-solution half-lives of NLuc sequences, as it

is well-suited for modeling datasets with few observations and collinear predictors. The

PLSR biplot revealed partially overlapping contributions from featured traditional

metrics—FE normalized by length, AUP, and GC content (**Figure 5C**). GC content

emerged as the dominant contributor, projecting strongly along both principal components

and aligning with samples exhibiting longer half-lives. In contrast, FE/length and AUP

were collinear and pointed in the opposite direction of GC, suggesting potential

redundancy in the information they provide. The ALO loading vector was clearly

separated from traditional metrics (both GC and FE/length), confirming its orthogonal

contribution.

To further interpret the contributions of each feature in the PLSR model, we

examined both the component weights and Variable Importance in Projection (VIP)

scores. Component 1 weights revealed that GC content and ALO contributed most

strongly to the positive direction of the latent axis associated with increasing half-life, while

FE/Length and AUP contributed in the opposite direction (**Figure 5D, top**). VIP scores,

which reflect the overall importance of each feature across all components, identified GC

content and FE/Length as the most influential predictors (VIP = 1.05 and 1.11,

respectively), with AUP and ALO contributing moderately (VIP = 0.96 and 0.86,

respectively) (**Figure 5D, bottom**).

The bivariate correlations between the variables further support the unique

informational content provided by ALO (**Figure S4**). While ALO exhibited strong negative

correlations with FE/Length (r = -0.87) and AUP (r = -0.90), it showed moderate positive

correlation with GC (r = 0.37) accompanied by a broad distribution of values. In contrast,

FE/Length and AUP both showed stronger, more consistent negative correlations with

GC (r = –0.66 and –0.47, respectively). This divergent relationship between ALO and GC

demonstrated a new pattern of covariance that was not captured by the initial set of

predictors (GC, AUP and FE/length). We refer to this sequence and structure feature-

based model as Stability Regression Analysis using Nucleotide-Derived features

(STRAND). STRAND integrates global (FE, AUP), local structural (ALO) and sequence

features (GC) into a compact, interpretable framework for predicting mRNA degradation.

We next aimed to evaluate the generalizability and predictive utility of STRAND.

Given the limited size of the NLuc dataset (N = 69), we did not perform a traditional train-

test split, as doing so would risk underpowering the model and obscuring true structure-

function relationships. Instead, we trained the models on the full NLuc dataset and tested

their ability to generalize to the eGFP variants (N = 13). To maximize the data utilized in

each training fold given our small dataset, we performed Leave-One-Out Cross-Validation

(LOOCV) for model tuning and evaluation (**Figure S5A**)**.** STRAND achieved strong

predictive performance across all metrics (Spearman *r* = 0.82, Pearson *r* = 0.72, RMSE

= 0.19, MAE = 0.17), highlighting the effectiveness of this compact feature set for mRNA

stability prediction (**Figure 5E**).

To compare STRAND's performance with state-of-the-art published approaches,

we evaluated half-life predictions using the benchmark DegScore-XGBoost model

described by Wayment-Steele et al.[21], along with the two top-performing deep learning

models from that study: Nullrecurrent, the overall first-place model, and Kazuki2, the

second-place model which achieved the best performance on the specific dataset we

used. For the same eGFP test set, all three models predicted degradation scores at the

nucleotide level, and the average per-position degradation scores (AUP) showed strong

correlation with experimental degradation rates (**Figure S5B**). Predicted half-lives for the

eGFP sequences for each model were then calculated using the relation half-life =

ln(2)/AUP and compared to the experimentally measured values.

The three models achieved comparable performances for the eGFP sequences,

with the DegScore-XGBoost model achieving a Spearman correlation of 0.56 (Pearson r

= 0.58) and RMSE of 0.46 (**Figure 5F**). NullRecurrent and Kazuki2 showed similar

performance, with a Spearman correlation of 0.58 and 0.60, respectively (Pearson r =

0.40 and 0.62), and RMSE of 0.42 and 0.48 (**Figure S5C**). By comparison, STRAND

achieved a Spearman correlation of 0.82, a Pearson r of 0.72, and RMSE of 0.19,

outperforming all three ML models with approximately a two-fold reduction in prediction

error relative to the best-performing comparison model. These results demonstrate that

integrating local and global structural features in a compact model like STRAND improves

both the accuracy and generalizability of mRNA degradation prediction.

To further assess STRAND's generalizability, we applied the model to sequences

derived from an independent study, VZV gE and SARS-CoV-2 Spike mRNA variants[19].

These differed substantially from the NLuc training set in both sequence length and

structural feature ranges (**Figure 6A**). Moreover, in-solution half-lives of VZV and Spike

sequences, measured under both 20 mM Mg$^{2+}$ (**Figure 6A)** and 10 mM Mg$^{2+}$ (**Figure**

**S6A**) buffer conditions, were substantially longer than those of the NLuc training set. The

LO distributions and per-position LO profiles of both VZV and Spike sequences showed

a clear trend: sequences with more positions at strongly positive LO values tended to

have longer half-lives (**Figures 6B, C**), consistent with the relationship between local

structural confidence and stability observed in the NLuc dataset. When STRAND

predictions were compared to experimentally measured half-lives, the model preserved

strong rank-order accuracy for both VZV (Spearman r = 0.86) and Spike (Spearman r =

0.95) sequences, although absolute prediction errors were large (RMSE = 20.36 and 5.25,

respectively) (**Figure 6D**). Notably, the FE values for both VZV and Spike sequences fall

well outside the range observed in the NLuc training set, and their GC content values lie

near the extremes of the NLuc distribution (**Figure 6A**). The large RMSE likely reflects

extrapolation beyond the training domain and the substantial difference in half-life scale

between these transcripts and the NLuc training data. This prediction trend was also

maintained for the half-lives measured in 10 mM $Mg^{2+}$ buffer condition (**Figure S6B**).

Overall, these results show that STRAND generalizes across proteins in predicting

stability rank order, although precise half-life prediction will require training data within the

relevant distribution.

**DISCUSSION**

In this study, we examined the relationship between RNA secondary structure and mRNA in-solution half-life, focusing on the predictive power of different structural metrics. Our findings highlight two key contributions. First, we show that ALO of base-pairing probability provides orthogonal information that complements traditional metrics such as FE, AUP, and GC content. This orthogonality enables ALO to differentiate sequences that have similar traditional structure feature values but exhibit markedly different stability. Second, we demonstrate that a multi-feature model integrating ALO with FE, AUP, and GC content outperforms previously published statistical and deep learning models in predicting mRNA stability across variants encoding multiple proteins, despite relying on a minimal and interpretable four-feature set. This compact feature combination balances complementary contributions while avoiding redundancy, resulting in a robust and generalizable framework for modeling RNA degradation.

While MFE has long been used as a proxy for RNA structural stability[12,31,32], it provides only a single, optimal conformation[13,33] and does not account for the diversity of structures an RNA sequence can adopt[10,11]. As illustrated in Figure 2, sequences with similar MFE values can exhibit vastly different structural topologies, such as the number and arrangement of stem-loops, bulges, or multibranch loops. These variations lead to distinct base-pairing patterns that can dramatically affect degradation rates[34,35], yet are not captured by MFE alone.

ALO and AUP are both derived from BPP, which ranges from 0 (unpaired) to 1 (paired). While BPP provides valuable insights, it is limited by its compressed range, making it difficult to detect subtle but functionally meaningful differences near the

extremes. Moreover, when BPP values are averaged across a sequence, positions with

BPP near zero, which represent confidently unpaired and degradation-prone sites, are

effectively treated as an absence of pairing signal rather than as informative structural

features. ALO addresses this by expanding BPP values into a logarithmic scale, ranging

from -∞ to +∞, thereby amplifying differences near 0 and 1 and improving interpretability

of pairing confidence[26]. For example, BPP values of 0.9, 0.99, and 0.999 are all close to

1 and difficult to distinguish on the original scale, but map to log-odds of 2.20, 4.60, and

6.91, respectively, offering clearer resolution of high pairing confidence. Similarly, low

BPP values (e.g., 0.01 vs. 0.1) become more distinguishable once transformed, allowing

better identification of confidently unpaired regions. This enhanced contrast allows ALO

to highlight structural differences between sequences that appear similar using traditional

metrics, as demonstrated by our binning analysis. These findings led us to explore

whether combining ALO with other established metrics could enhance predictive

modeling of mRNA stability.

Building on these insights, we developed STRAND, a multi-feature regression

model incorporating ALO, FE, AUP, and GC content. Despite its compact four-feature

design, this model performs competitively with previously published statistical and deep

learning approaches while maintaining greater interpretability. Importantly, it generalizes

well to independent data, including eGFP, VZV gE, and SARS-CoV-2 Spike constructs,

with strong rank-order performance even when applied to transcripts well outside the

training domain. The model's simplicity makes it especially suitable for early-stage

screening and optimization workflows, where interpretability and low feature complexity

are critical. Moreover, its performance without reliance on experimental probing data

highlights the value of well-chosen *in silico* features for robust prediction.

Beyond its predictive utility, STRAND and the LO metric can be readily

incorporated into existing mRNA design workflows (**Figure S7**). During the initial

sequence design stage, LO or ALO can be an additional objective or constraint within

optimization algorithms, complementing traditional metrics to guide the sequence

generation with favorable local structural profiles. Alternatively, for a given candidate

sequence pool, STRAND can be used to predict in-solution half-lives, or ALO can serve

as a filter to prioritize candidates with structural features associated with greater stability

prior to experimental validation. In either application, STRAND's computational simplicity

and its reliance on readily computed *in silico* features make it practical for integration into

high-throughput design pipelines.

While STRAND performs well overall, there are limitations that offer opportunities

for further improvement. As expected, prediction accuracy declines at the extremes of the

half-life range, likely due to limited representation of highly stable or unstable sequences

in the training set (**Figure 1A**). One of the most influential biological variables affecting

degradation is transcript length[7,36–38], yet STRAND was trained using variants of a single

open reading frame (ORF), as it was the only transcript in the dataset with a sufficient

number of sequence variants for training. While STRAND performed reasonably well

even for sequence sets of different length and feature values outside of the training

regime, expanding the training dataset to include transcripts encoding proteins of varying

lengths   and   sequence   features   would   improve   the   model's   robustness   and

generalizability. For example, STRAND predictions for MEV sequences were poor

(**Figure S8**). This is likely attributable to the short transcript length of MEV variants, for

which FE normalization by length becomes unreliable due to the non-linear relationship

between FE and sequence length at shorter lengths[39], combined with their narrow FE

distributions that fall outside the training regime. Increasing the diversity of the dataset

may uncover new features that contribute independently of both ALO and MFE, providing

further gains in predictive accuracy and biological insight. Moreover, STRAND currently

models CDS-level structural features and does not explicitly account for other important

dimensions of mRNA design, such as co-optimization of UTR features, or incorporating

structure predictions specific for relevant modified nucleotides. Investigating how these

factors interact with the structural features captured by STRAND represents an important

direction for future work.

For any given value of conventional metrics such as MFE, sequences often share

similar range of LO values but differ in their distributions, making it difficult for LO alone

to distinguish stability differences across broader structural contexts. LO effectively

resolves differences between sequences that share similar global metrics but differ in

degradation rates, capturing subtle structural features that traditional approaches often

overlook. However, it lacks the broad discriminatory power needed to classify sequences

across a wider spectrum of structural contexts. Therefore, the strength of LO lies in

complementing existing metrics, providing finer-scale insights particularly valuable in

cases where traditional metrics offer limited differentiation.

Additionally, a key limitation of our approach stems from inherent inaccuracies in

*in silico* RNA secondary structure predictions, which rely on imperfect thermodynamic

parameters and do not always fully capture biologically relevant structural ensembles[40–

[42]. As our stability model relies on these predicted structures, its accuracy is likely

influenced by the quality of the folding algorithms. Continued improvements in RNA

structure prediction methods could therefore further enhance the performance and

reliability of STRAND.

Across the NLuc, eGFP, VZV, and Spike datasets, we observed distinct patterns

of LO distributions that are associated with differences in degradation. Understanding

how the balance between highly structured and weakly structured regions contributes to

degradation across different sequence contexts remains an open question. Nevertheless,

STRAND demonstrates that well-chosen structural features can substantially reduce

prediction error for mRNA stability, and its compact, interpretable design makes it readily

adaptable as new training data and design contexts become available.

**METHODS**

**RNA Structure Prediction and Metric Calculation**

Data Acquisition

RNA sequence and corresponding half-life data were obtained from the study by Leppek et al.[9] accompanying the OpenVaccine: COVID-19 mRNA Vaccine Degradation Prediction dataset[9,21].

Feature Calculation using LinearPartition

Acquired RNA sequences were used as input for the LinearPartition[29] software to generate structural metrics, including the FE predictions, the BPP for each nucleotide position, the average base pairing probability across the sequence, and the AUP across the sequence. The BPP at each nucleotide position is calculated as the summed probability of that nucleotide forming a base pair with any other nucleotide across the entire predicted structural ensemble derived from the partition function.

Structure Visualization using RNAfold

For visual representation of predicted RNA secondary structures, sequences were submitted to the RNAfold[43] web server. Default options provided by the web server were used for both structure prediction and visualization.

Log Odds Calculation

Using the BPP calculated by LinearPartition[29], the Log Odds Per Position (LOPP) was determined for each nucleotide using the formula: $LOPP = \ln(p / (1-p))$, where p is the

BPP. To handle probabilities of exactly 0 or 1, a pseudocount of $1e^{-10}$ was added to p

when BPP = 0 and subtracted from p when BPP = 1 prior to the log odds calculation.

<u>BPP Curvature Analysis and Thresholding</u>

The relationship between BPP and LO was examined by calculating the mathematical

curvature of the BPP vs. LO curve. Curvature was determined by numerically calculating

the second derivative of LO with respect to BPP ($d^2(\ln(p/(1-p)))/dp^2$) within the boundary

conditions $0 \le p \le 1$. This analysis identified points of maximum curvature at BPP values

of approximately 0.20 and 0.80. Based on this, a BPP threshold of 0.2 was selected to

define positions considered likely to be unpaired for downstream analyses.

**Sequence Clustering Analysis**

Uniform Manifold Approximation and Projection (UMAP)[28] was employed to explore

potential clustering based on sequence features. Input features included FE normalized

by sequence length, Average Unpaired Probability (AUP), and GC content. Analyses

were performed both including and excluding the Average Log Odds (ALO) as an

additional feature. UMAP was performed using the UMAP package (Konopka, T. ; DOI:

<u>10.32614/CRAN.package.umap</u>) in R (v 4.4.0 ; https://www.r-project.org) utilizing default

parameters (including n_neighbors=15, min_dist=0.1, n_epochs=200). Due to the

stochastic nature of the algorithm, UMAP analysis was repeated 20 times for each feature

set; cluster consistency was assessed by visually confirming the presence of three distinct

clusters and the stability of sequence identities within each cluster across runs.

**Linear Regression Modeling**

497 Linear regression models using FE and AUP as predictors were generated using the stats

498 package in R (v 4.4.0).

**PLS Regression Modeling**

To model the relationship between sequence-derived features and in-solution half-life,

Partial Least Squares Regression (PLSR)[30,44] was implemented using the pls package in

R (v 4.4.0). The final set of predictor variables included FE normalized by sequence length,

Average Unpaired Probability (AUP), GC content, and Average Log Odds (ALO).

Predictor variables were centered and scaled prior to modeling. Given the limited training

dataset size (N = 69), Leave-One-Out Cross-Validation (LOOCV) was utilized on the

training set for model parameter tuning by minimizing the Root Mean Squared Error of

Prediction (RMSEP) (Fig. S5A). Variable Importance in Projection (VIP) scores were

calculated using the plsVarSel package[45] in R (v 4.4.0 ; https://www.r-project.org) to

assess the contribution of each predictor variable to the final PLSR model trained on the

full training set. Final model performance was evaluated on a separate, independent

EGFP dataset (N = 13) by calculating the Pearson and Spearman correlation coefficients,

and Root Mean Square Error (RMSE) between predicted and actual half-life values.

## DATA AVAILABILITY

The STRAND model described in this paper is publicly available to support reproducibility and broad use. The raw training data used in this study were obtained from the OpenVaccine: COVID-19 mRNA Vaccine Degradation Prediction dataset (Leppek et al., 2022). Any additional data required to verify and extend the research are available from the corresponding author upon request.

## AUTHOR CONTRIBUTIONS

**S.Y.:** Conceptualization (Lead), Formal analysis (Lead), Investigation (Lead), Validation (Lead), Visualization (Lead), Writing – original draft (Lead), Writing – review & editing (Lead). **S.E.A.:** Data curation (Supporting), Formal analysis (Supporting), Investigation (Supporting), Writing – original draft (Supporting), Writing – review & editing (Supporting). **Y.J.:** Formal analysis (Supporting), Investigation (Supporting), Validation (Supporting), Writing – review & editing (Supporting). **J.W.D.:** Conceptualization (Lead), Supervision (Supporting), Writing – review & editing (Supporting). **M.M.:** Conceptualization (Lead), Investigation (Lead), Project administration (Lead), Resources (Lead), Supervision (Lead), Writing – original draft (Lead), Writing – review & editing (Lead).

## KEYWORDS

RNA Secondary Structure, RNA Stability, Base Pairing Probability, Base Pairing Log Odds, Computational Prediction, RNA Therapeutics, mRNA Vaccines

**ACKNOWLEDGEMENTS**

The authors thank Chris Pepin for the preliminary discussions of this work, Alicia Bicknell

and Chiaowen Joyce Hsiao for critical reading of the manuscript. This work was supported

in part by the NIH under the award T32 GM152319 (S.Y.)

**DECLARATION OF INTERESTS**

M.M., S.E.A., J.W.D., Y.J. are currently employees of and shareholders in Moderna Inc.

**Declaration of generative AI and AI-assisted technologies in the manuscript preparation process**

During the preparation of this work the author(s) used Claude and ChatGPT to assist with drafting and editing portions of the manuscript. After using this tool/service, the author(s) reviewed and edited the content as needed and take(s) full responsibility for the content of the published article.

## REFERENCES

1. Baden, L. R. *et al.* Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine. *N. Engl. J. Med.* **384**, 403–416 (2021).

2. Crommelin, D. J. A., Anchordoquy, T. J., Volkin, D. B., Jiskoot, W. & Mastrobattista, E. Addressing the Cold Reality of mRNA Vaccine Stability. *J. Pharm. Sci.* **110**, 997–1001 (2021).

3. Li, Y. & Breaker, R. R. Kinetics of RNA Degradation by Specific Base Catalysis of Transesterification Involving the 2'-Hydroxyl Group. *J. Am. Chem. Soc.* **121**, 5364–5372 (1999).

4. Garneau, N. L., Wilusz, J. & Wilusz, C. J. The highways and byways of mRNA decay. *Nat. Rev. Mol. Cell Biol.* **8**, 113–126 (2007).

5. Pogocki, D. & Schöneich, C. Chemical stability of nucleic acid–derived drugs. *J. Pharm. Sci.* **89**, 443–456 (2000).

6. Mikkola, S., Kaukinen, U. & Lönnberg, H. The effect of secondary structure on cleavage of the phosphodiester bonds of RNA. *Cell Biochem. Biophys.* **34**, 95–119 (2001).

7. Chheda, U. *et al.* Factors Affecting Stability of RNA – Temperature, Length, Concentration, pH, and Buffering Species. *J. Pharm. Sci.* **113**, 377–385 (2024).

8. Mauger, D. M. *et al.* mRNA structure regulates protein expression through changes in functional half-life. *Proc. Natl. Acad. Sci.* **116**, 24075–24083 (2019).

9. Leppek, K. *et al.* Combinatorial optimization of mRNA structure, stability, and translation for RNA-based therapeutics. *Nat. Commun.* **13**, 1536 (2022).

10.     Ward, M., Courtney, E. & Rivas, E. Fitness functions for RNA structure design.

*Nucleic Acids Res.* **51**, e40 (2023).

11.     Dirks, R. M., Lin, M., Winfree, E. & Pierce, N. A. Paradigms for computational

nucleic acid design. *Nucleic Acids Res.* **32**, 1392–1403 (2004).

12.     Mathews, D. H., Sabina, J., Zuker, M. & Turner, D. H. Expanded sequence

dependence of thermodynamic parameters improves prediction of RNA secondary

structure1. *J. Mol. Biol.* **288**, 911–940 (1999).

13.     Mathews, D. H. *et al.* Incorporating chemical modification constraints into a

dynamic programming algorithm for prediction of RNA secondary structure. *Proc.*

*Natl. Acad. Sci.* **101**, 7287–7292 (2004).

14.     Bicknell, A. A. *et al.* Attenuating ribosome load improves protein output from

mRNA by limiting translation-dependent mRNA decay. *Cell Rep.* **43**, (2024).

15.     Cohen, B. & Skiena, S. Natural selection and algorithmic design of mRNA. *J.*

*Comput. Biol. J. Comput. Mol. Cell Biol.* **10**, 419–432 (2003).

16.     Ward, M., Richardson, M. & Metkar, M. mRNA Folding Algorithms for Structure

and Codon Optimization. Preprint at https://doi.org/10.48550/arXiv.2503.19273

(2025).

17.     Terai, G., Kamegai, S. & Asai, K. CDSfold: an algorithm for designing a protein-

coding sequence with the most stable secondary structure. *Bioinformatics* **32**, 828–

834 (2016).

18.     Wayment-Steele, H. K. *et al.* Theoretical basis for stabilizing messenger RNA

through secondary structure design. *Nucleic Acids Res.* **49**, 10604–10617 (2021).

19.    Zhang, H. *et al.* Algorithm for optimized mRNA design improves stability and

immunogenicity. *Nature* **621**, 396–403 (2023).

20.    Gu, X., Qi, Y. & El-Kebir, M. DERNA Enables Pareto Optimal RNA Design. *J.*

*Comput. Biol. J. Comput. Mol. Cell Biol.* **31**, 179–196 (2024).

21.    Wayment-Steele, H. K. *et al.* Deep learning models for predicting RNA

degradation via dual crowdsourcing. *Nat. Mach. Intell.* **4**, 1174–1184 (2022).

22.    Kaukinen, U., Lyytikäinen, S., Mikkola, S. & Lönnberg, H. The reactivity of

phosphodiester bonds within linear single-stranded oligoribonucleotides is strongly

dependent on the base sequence. *Nucleic Acids Res.* **30**, 468–474 (2002).

23.    Siegfried, N. A., Busan, S., Rice, G. M., Nelson, J. A. E. & Weeks, K. M. RNA

motif discovery by SHAPE and mutational profiling (SHAPE-MaP). *Nat. Methods* **11**,

959–965 (2014).

24.    Lu, Z. J., Gloor, J. W. & Mathews, D. H. Improved RNA secondary structure

prediction by maximizing expected pair accuracy. *RNA* **15**, 1805–1813 (2009).

25.    Hofacker, I. L., Bernhart, S. H. F. & Stadler, P. F. Alignment of RNA base pairing

probability matrices. *Bioinformatics* **20**, 2222–2227 (2004).

26.    Norton, E. C. & Dowd, B. E. Log Odds and the Interpretation of Logit Models.

*Health Serv. Res.* **53**, 859–878 (2018).

27.    Metkar, M., Pepin, C. S. & Moore, M. J. Tailor made: the art of therapeutic mRNA

design. *Nat. Rev. Drug Discov.* **23**, 67–83 (2024).

28.    McInnes, L., Healy, J. & Melville, J. UMAP: Uniform Manifold Approximation and

Projection for Dimension Reduction. Preprint at

https://doi.org/10.48550/arXiv.1802.03426 (2020).

29.    LinearPartition: linear-time approximation of RNA folding partition function and

base-pairing probabilities | Bioinformatics | Oxford Academic.

https://academic.oup.com/bioinformatics/article/36/Supplement_1/i258/5870487.

30.    Wold, S., Ruhe, A., Wold, H. & Dunn, III, W. J. The Collinearity Problem in Linear

Regression. The Partial Least Squares (PLS) Approach to Generalized Inverses.

*SIAM J. Sci. Stat. Comput.* **5**, 735–743 (1984).

31.    Flamm, C., Fontana, W., Hofacker, I. L. & Schuster, P. RNA folding at

elementary step resolution. *RNA* **6**, 325–338 (2000).

32.    Hofacker, I. L. *et al.* Fast folding and comparison of RNA secondary structures.

*Monatshefte Für Chem. Chem. Mon.* **125**, 167–188 (1994).

33.    Dowell, R. D. & Eddy, S. R. Evaluation of several lightweight stochastic context-

free grammars for RNA secondary structure prediction. *BMC Bioinformatics* **5**, 71

(2004).

34.    Zarringhalam, K., Meyer, M. M., Dotu, I., Chuang, J. H. & Clote, P. Integrating

Chemical Footprinting Data into RNA Secondary Structure Prediction. *PLOS ONE* **7**,

e45160 (2012).

35.    Cordero, P. & Das, R. Rich RNA Structure Landscapes Revealed by Mutate-and-

Map Analysis. *PLOS Comput. Biol.* **11**, e1004473 (2015).

36.    Agarwal, V. & Kelley, D. R. The genetic and biochemical determinants of mRNA

degradation rates in mammals. *Genome Biol.* **23**, 245 (2022).

37.    Cheng, F. *et al.* Research Advances on the Stability of mRNA Vaccines. *Viruses*

**15**, 668 (2023).

38.   Feng, L. & Niu, D.-K. Relationship Between mRNA Stability and Length: An Old

Question with a New Twist. *Biochem. Genet.* **45**, 131–137 (2007).

39.   Trotta, E. On the Normalization of the Minimum Free Energy of RNAs by

Sequence Length. *PLOS ONE* **9**, e113380 (2014).

40.   Chaturvedi, M., Rashid, M. A. & Paliwal, K. K. RNA structure prediction using

deep learning — A comprehensive review. *Comput. Biol. Med.* **188**, 109845 (2025).

41.   Akiyama, M. & Sato, K. RNA Secondary Structure Prediction Based on Energy

Models. in *RNA Structure Prediction* (eds Kawaguchi, R. K. & Iwakiri, J.) 89–105

(Springer US, New York, NY, 2023). doi:10.1007/978-1-0716-2768-6_6.

42.   Doshi, K. J., Cannone, J. J., Cobaugh, C. W. & Gutell, R. R. Evaluation of the

suitability of free-energy minimization using nearest-neighbor energy parameters for

RNA secondary structure prediction. *BMC Bioinformatics* **5**, 105 (2004).

43.   Gruber, A. R., Lorenz, R., Bernhart, S. H., Neuböck, R. & Hofacker, I. L. The

Vienna RNA Websuite. *Nucleic Acids Res.* **36**, W70–W74 (2008).

44.   Wold, S., Sjöström, M. & Eriksson, L. PLS-regression: a basic tool of

chemometrics. *Chemom. Intell. Lab. Syst.* **58**, 109–130 (2001).

45.   Mehmood, T., Liland, K. H., Snipen, L. & Sæbø, S. A review of variable selection

methods in Partial Least Squares Regression. *Chemom. Intell. Lab. Syst.* **118**, 62–69

(2012).

 **LIST OF FIGURE LEGENDS AND CAPTIONS**

**Figure 1. In-solution half-life distributions across protein mRNA variants and**
**structure–function analysis in the Nano Luciferase set. (A)** A boxplot distribution of
in-solution half-lives for mRNA variants of different proteins. The box represents the
interquartile range, the horizontal line indicates the median, and whiskers denote 1.5X
the interquartile range; individual sequences are plotted as points. The NanoLuc (NLuc;
bold) CDS variants were selected for further analysis due to their wide range of half-lives
and large number of sequences. **(B)** Correlation of FE with half-life in the NLuc dataset.
Red-highlighted sequences show similar FE but divergent half-lives. Pearson r for the
entire dataset (black) and for sequences with similar FE (red) are shown. **(C)** Same as
**(B)** but for AUP. **(D)** Residual plots for FE showing increased error for sequences with
higher half-lives. **(E)** Same as (D) but for AUP. **(F)** UMAP projection of NLuc sequences
based on global structural features (FE, AUP, and GC content), revealing three clusters.
Clusters are labeled as high (H), medium (M), and low (L) based on their in-solution half-
lives. **(G)** Boxplot distribution of half-lives for sequences in each UMAP-defined cluster in
**(F)**, with pairwise statistical comparisons shown. The Mann-Whitney test was used to
calculate p-values. The number of sequences within each cluster is noted in parentheses.

**Figure 2. Structure comparison between two RNA sequences with similar global**
**metrics but substantial differences in in-solution half-life. (A)** Predicted secondary
structures of two NLuc sequences, X320004D1 and X930007D1, which exhibit similar FE
and AUP but differ substantially in in-solution half-life. Global structure metrics and half-
life values are shown. **(B)** Heatmap of local BPP values by position for the two selected
sequences. **(C)** Barplot comparing structural components between the predicted
maximum expected accuracy structures, including counts of unpaired nucleotides, hairpin
loops, interior loops, and multiloops. **(D)** Heatmap comparison of BPPs binned at 0.05
resolution for the two selected sequences. Color scale indicates $\log_{10}$ frequency of
positions in each bin.

**Figure 3. Local RNA base-pairing patterns differ among similarly structured sequences and associate with in-solution half-life. (A)** Heatmap of local BPP values by position for sequences with similar global metrics (FE and AUP) but differing in-solution half-lives. Sequences are ordered by increasing half-life. **(B)** Heatmap comparisons of BPPs binned at 0.05 resolution across the same sequences as in **(A)**. Color scale indicates $\log_{10}$ frequency of positions in each bin. Numbers flanking each row indicate the count of positions in the lowest and highest BPP bins. **(C)** Correlation between the number of unpaired positions (defined as BPP < 0.2) and in-solution half-life. Pearson r for the NLuc-21 sequences (red) is shown. **(D)** Correlation between the number of unpaired patches (≥5 consecutive unpaired positions with BPP < 0.2) and in-solution half-life. Pearson r for the NLuc-21 sequences (red) is shown.

**Figure 4. Base-pairing probability Log-odds (LO) capture local structure features predictive of RNA stability. (A)** Histogram of LO values (binwidth = 0.5) for the two sequences (X320004D1 and X930007D1) with similar global structure metrics but divergent half-lives. **(B)** Heatmap comparisons of binned LO values across the subset of NLuc sequences (**Figure 3**), sorted by increasing in-solution half-life. Color scale indicates the frequency of positions in each bin. **(C)** Heatmap comparisons of LO values by position for the same sequences as in **(B)** with color scale indicating LO values and highlighting localized structural differences along the transcript. **(D)** Correlation of average LO for weakly paired regions (LO < -1.36, equivalent to BPP < 0.2) and in-solution half-life. Pearson r for the NLuc-21 sequences is shown. **(E)** Same as **(D)** but for average LO across all positions.

**Figure 5. Multi-feature regression model of RNA degradation shows improved prediction accuracy through orthogonal contribution from Average Log Odds (ALO). (A)** UMAP projection of NLuc sequences based on a multi-feature model including FE/length, AUP, GC content, and ALO, showing separation into three clusters. **(B)** Boxplot distributions of in-solution half-lives for each UMAP-derived cluster, with statistical comparisons indicating significant differences between groups. The Mann-Whitney test was used to calculate p-values. The number of sequences within each

cluster is noted in parentheses. **(C)** PLSR biplot of NLuc sequences using FE/length, AUP,

GC content, and ALO as features, plotted along Component 1 and Component 2. Arrows

indicate loading vectors for each feature. Points are colored by in-solution half-life. **(D)**

PLSR Component 1 weight of each feature (Top). Positive and negative weights indicate

the direction and magnitude of each feature's contribution to the first latent component.

Variable Importance in Projection (VIP) scores across all components (Bottom). The

dashed line indicates a canonically used threshold to define important variables. **(E)**

PLSR model predictions of in-solution half-life for eGFP sequences using models trained

on NLuc variants. A linear regression fit is shown with a 95% confidence interval

represented by the shaded region. **(F)** DegScore-XGBoost model predictions for the

same eGFP sequences as in **(E)**. A linear regression fit is shown with a 95% confidence

interval represented by the shaded region.

**Figure 6. STRAND generalizes to mRNAs encoding viral antigens with structural**

**features outside the training range. (A)** Distributions of features (AUP, FE, GC content)

and in-solution half-lives for Spike, VZV, and NLuc sequence sets, illustrating the

substantial differences in feature ranges between the training (NLuc) and test (VZV,

Spike) datasets. **(B)** Heatmaps of binned LO distributions (binwidth = 1) for VZV gE (top)

and Spike (bottom) sequences, sorted by increasing half-life. Color scale indicates

frequency of positions in each bin. **(C)** Per-position LO heatmaps for VZV gE (top) and

Spike (bottom) sequences. Color scale indicates LO values. **(D)** STRAND predicted half-

life versus experimentally measured half-life for VZV (left) and Spike (right) sequences.

Spearman r, Pearson r, and RMSE are shown.

**A** Predicted Secondary Structures Comparison

**X320004D1**
GC: 52.04%
FE: -387.71
AUP: 0.34
$t_{1/2}$: 0.72

**X930007D1**
GC: 60.03%
FE: -407.65
AUP: 0.33
$t_{1/2}$: 1.99

**B** BPP Comparison by Position

X320004D1
X930007D1

Nucleotide Position

BPP
1.0
0.8
0.6
0.4
0.2
0.0

**C** Structure Components Comparison

Counts

ss-Nucleotide  ds-Nucleotide

Multiloops  Hairpin Loops  Internal Loops

X320004D1
X930007D1

**D** BPP Comparison by Composition

X320004D1
X930007D1

0.0 0.1 0.2 0.3 0.4 0.5 0.6 0.7 0.8 0.9 1.0
BPP

$Log_{10}$(Freq.)

**A** Distributions of RNA vFeatures

**B** Distribution of LOs for VZV (top) and Spike (bottom) Sequences

**C** LO Comparisons by Position for VZV (top) and Spike (bottom) Sequences

**D** VZV Sequences STRAND Prediction — Spike Sequences STRAND Prediction