Recurrent structural variation and recent turnover at the 17q21.31 locus in humans and great apes

Samvardhini Sridharan[1,2], Runyang Nicolas Lou[3], Scott Ferguson[3], Joana L Rocha[3], Rishi De-Kayne[3], Matthew W Mitchell[4], Alison N Killilia[1], Victor Borda[5], Santiago G. Medina-Munoz[6], Simon Gravel[7], CAAPA2 PopGen Working Group, Brenna Henn[8], Peter H Sudmant[1,2,3]

1 Department of Molecular and Cell Biology, University of California, Berkeley, CA, USA
2 Center for Computational Biology, University of California, Berkeley, CA, USA
3 Department of Integrative Biology, University of California, Berkeley, CA, USA
4 Coriell Institute for Medical Research, Camden, NJ 08103
5 University of Maryland School of Medicine, Baltimore, MD, USA
6 CINVESTAV, Mexico City, Mexico
7 Department of Human Genetics, McGill University, Montreal, Canada
8 Department of Anthropology, University of California, Davis, CA, USA

Correspondence should be addressed to Peter H Sudmant: psudmant@berkeley.edu

ABSTRACT

The 17q21.31 locus in humans harbors several complex structural haplotypes including a ~970kb inversion. Different inversion haplotypes have been associated with susceptibility to microdeletions causing Koolen-de Vries syndrome and variation in fecundity and recombination rates. Here, using 210 haplotype-resolved human genome assemblies and pangenome graph-based approaches we characterize 11 distinct structural haplotypes, several of which have not been previously described. Extending our analyses to a set of haplotype-resolved great-ape genomes, we characterize the structure of an independent inversion in chimpanzees which extends an additional 650kb, encompasses 5 additional genes, and is ~2 million years younger than the human inversion. We further determine that gorillas exhibit an independent duplication of the *KANSL1* gene which may predispose them to Koolen-de Vries syndrome causing microdeletions. Using short read sequencing data we characterize 17q21.31 haplotype diversity worldwide in ~5174 individuals from 107 populations finding increased frequencies of *KANSL1* duplication-containing haplotypes in both European and South Asian populations as well as 8 double recombination events between inverted and non-inverted haplotypes ranging in size from 20-180kb. Finally, using 626 ancient Eurasian human genomes we show the frequency of haplotypes containing *KANSL1* duplications has increased ~6-fold over the past 12 thousand years in Europe. Together, our results highlight the dynamics, complexity, and recurrent, independent evolution of a medically relevant locus across humans and great apes.

INTRODUCTION

Complex structural variants (SVs) play critical roles in human disease[1], diversity[2,3], and evolution[4,5]. Inversions are a particularly intriguing and understudied class of SV which have historically been challenging to assay as they are often flanked by large, complex, segmental duplications[1,6]. Nevertheless, across taxa inversions are powerful drivers of genetic and phenotypic diversity influencing traits such as behavior[7], fitness[8,9], and morphology[8,10]. Inversions can also act as barriers to gene flow between populations, promoting speciation and adaptation to different ecological niches and environmental pressures[11–13].

One remarkable inversion in humans is found at the 17q21.31 locus spanning ~970kb and impacting patterns of linkage disequilibrium over more than a megabase[14]. Haplotypes at this locus group into two major clades based on their inversion status: H1 haplotypes which are the most common in humans, and H2 haplotypes, which are inverted with respect to the H1 haplotype. Since its discovery, this locus has been associated with signatures of positive selection[14], disease[15], and recurrent structural rearrangement ("toggling")[16] throughout the great ape lineage, highlighting its complex evolutionary history.

The 17q21.31 locus is of particular medical importance because it harbors several genes associated with human disease. These include *MAPT,* which encodes the tau protein found in aggregates in Alzheimer's dementias[17], and *KANSL1,* disruption of which results in Koolen-de Vries Syndrome (KdVS)(Koolen et al. 2006; Koolen et al. 2012). KdVS is characterized by developmental delay, intellectual disability, hypotonia, epilepsy, distinctive facial features, and congenital malformations across multiple organ systems[18]. While rare truncating mutations of *KANSL1* can cause KdVS[15] it is more commonly caused by 17q21.31 microdeletions which occur at a frequency of ~1/16,000 individuals. These microdeletions encompass 5 genes and are mediated by non-allelic homologous recombination (NAHR) between directly oriented segmental duplications which are present on the H2 (inverted) haplotype, but not the H1 haplotype. H2 carriers are thus at risk for KdVS. However, H2 haplotypes, which are at particularly high frequency in European populations, have also been associated with increased fecundity in some populations (Clark et al. 2022; Stefansson et al. 2005) and thus may be the target of positive selection[18].

In addition to the non-inverted (H1) and inverted (H2) haplotypes, recent studies have identified additional structural complexity at the 17q21.31 locus[19,20], including two independent partial duplications of the *KANSL1* gene on both the H1 and H2 haplotypes, as well as copy number variation of the *NSF* gene. Here, we set out to characterize the recent history and full extent of structural complexity at the 17q21.31 locus using both humans and non-human primate long-read haplotype-resolved genome assemblies, extensive human population diversity sequencing, and ancient genomes.

RESULTS

Resolving complex genome structures at the 17q21.31 locus with 210 haplotype-resolved genome assemblies

Recently generated long-read genomic resources provide an unprecedented opportunity to uncover structural diversity at complex loci such as the 17q21.31 locus. We collated 210 diverse haplotypes including the recently assembled T2T human genome to examine the structural complexity of this region. We first used PGRTK (Chin et al. 2023) (cite) to annotate the repeat architecture of these genomes (**Fig 1A, B, S1**). PGRTK identifies clusters of paralogous and homologous sequences within and between genomes enabling us to compare structures among haplotypes. Previous literature has established that alongside the major inversion haplotype groups H1 and H2, several additional regions of copy number variation exist on these haplotype backgrounds. These include distinct and independent partial duplications of the *KANSL1* gene on the H2 and H1 haplotypes, referred to as α and β respectively. Additionally partial duplications of the *NSF* gene, on either the H1 or H2 haplotype, are referred to as γ duplications.

Alongside our PGRTK-based classification of structures we used a pangenome-based approach (PGGB) (Garrison et al. 2024) to cluster haplotype architectures (see methods) identifying a total of 24 unique structures (**Fig S1**). Many of these structures were highly similar differing only subtly in the composition of the segmentally duplicated *LRRC37A* gene. We thus further clustered structures based on their pairwise jaccard distance into 11 unique summary structures (**Methods**). Structural differences between these haplotypes were confirmed by plotting a multiway structural alignment using SVbyEye (**Fig 1C**). These clusters exhibit finer resolution than previously developed nomenclature. For example, we identified three structures which can be classified as H1.β1.γ1, based on their single copies of the *KANSL1* and the *NSF* duplications, which we designate with additional numeric suffixes. We further identified two distinct H1.β1.γ2 haplotypes and three H1.β1.γ3 haplotypes. These distinct structures are mostly differentiated by copy number variation at the *LRRC37A2* gene, which flanks the inversion. Overall, the most commonly observed H1 structures were H1.β1.γ1.1 (37% of H1), H1.β1.γ2.2 (30% of H1), H1.β1.γ3.1 (19% of H1) and H1.β2.γ1 (8% of H1) - making up a total proportion of 94% of H1s (88% overall). Most of the variation in structural diversity we observed is located on H1.β1 haplotypes, which are the highest frequency. Only a single H1.β2 and H2.α2 structure were observed in our dataset and several previously described haplotype structures were not present in our long read assemblies (H1.β3.γ1, H2.α1.γ2, and H2.α2.γ1). However, these structures were observed at low frequency in our subsequent analyses of structural variation using short reads. Together, these data recapitulate previously described structures in higher detail and demonstrate additional complexity present in humans.

Recurrent but distinct inversions in humans and chimpanzees

The recent completion of several haplotype-resolved T2T ape genomes provides the opportunity to contextualize the evolutionary history of this locus in great apes. To explore the population-level diversity in these non-human species we further sequenced and assembled two high-quality haplotype resolved gorilla genomes (four haplotypes) and assessed a set of

recently assembled chimpanzee and bonobo genomes (Rocha et al. in prep). Previous work
has demonstrated that the H2 orientation is the ancestral architecture of the locus. Intriguingly,
fluorescent *in-situ* hybridization of a panel of several apes has suggested that the 17q21.31
inversion is segregating in orangutans, chimpanzees, and bonobos[16] emerging independently in
these species (i.e. inversion *toggling*).

We extracted the orthologous 17q21.31 sequence across five non-human great ape species
and constructed structural alignments using SVbyEye (**Fig. 2A,B**). These alignments
demonstrate the increasing complexity of segmental duplications in African great apes in
comparison to orangutans, as has been previously described[16]. To understand how these
structures relate to the human architecture we complemented these analyses by projecting
human-identified PGRTK sequence blocks onto the great ape genomes (**Fig. 2C**). These
projections lose resolution as a function of genetic distance particularly in rapidly evolving
sequence such segmental duplications but still highlight major architectural features like the
inversion and flanking regions.

Our analyses revealed several striking, lineage specific structural features at this locus. In
gorillas we identified a 500kb duplication overlapping the *KANSL1* gene. This is an independent
duplication of the gene distinct from that observed in human haplotypes. This duplication
corresponds to an annotated gene sequence in the recent T2T gorilla genome assembly,
though it is unclear if it is a functional duplication. Furthermore, these *KANSL1* duplications are
in direct orientation predisposing these haplotypes to NAHR-driven microdeletions/duplications.
Thus, gorillas may be at risk for Koolen-de-vries syndrome. We also identified two inverted
chimpanzee haplotypes. These inversions are in the same orientation as the human H1
haplotype. We refer to this haplotype as the H1p haplotype, and the ancestral primate haplotype
as H2p. The chimpanzee H1p/H2p inversion spans ~1.4Mbp (**Fig 2B, S2**) encompassing the
entirety of the human inversion and further extending an additional ~650kb overlapping several
genes including *WNT3, WNT9B, GOSR2,* and *RPML* (**Fig 2B**). The human and chimpanzee
inversions share one approximate breakpoint though the other is distinct (**Fig S2**). Thus, not
only is the chimpanzee inversion the result of an independent event, but it is a completely
distinct structural variant. This provides further evidence that the H2 orientation is the ancestral
structure in humans and chimpanzees, as while the H1 and H1p haplotypes differ structurally by
hundreds of kb the H2 and H2p haplotypes exhibit highly similar structures.

We next sought to compare the timing of the different inversion events by aligning ~400kbp of
unique (non-duplicated) sequence within the shared inversion block across humans and great
apes (**Fig. 2D, methods**). The human H1/H2 inversion exhibits a coalescence time of
approximately to 2.38 million years ago, confirming a relatively deep coalescence of the two
haplotypes[19]. The chimpanzee H1p/H2p inversion however emerged much more recently
~330,000 years ago. These analyses also revealed extensive incomplete lineage sorting across
humans, gorillas, chimpanzees, and bonobos. 54% of trees exhibited a topology with humans
as an outgroup to the *pan* and gorilla lineages with the canonical topology represented 46% of
the time. Together, these results demonstrate that the 17q21.31 locus is highly dynamic across

great apes with distinct but overlapping structural variants occurring independently in several
taxa over vastly different timescales.

Timing and recurrence of human 17q21.31 structural haplotypes

Long-read haplotype-resolved assemblies phase genome structures with linked SNPs which
can be used to trace the evolutionary history of complex genome architectures. Extended
suppression of recombination between the H1 and H2 haplotypes enables dating of the
coalescent time of the inversion to ~2.38 million years. However, recombination between H1
haplotypes will disrupt patterns of linkage disequilibrium (LD) making it challenging to assess
the emergence of specific H1 substructures. We sought to identify high LD blocks adjacent to
the distal region of H1 structural variation (containing the *KANSL1* and *NSF* duplications). Such
regions can be used to trace the history of these structures. We identified 100kb of sequence
adjacent to the structurally variable portion of H1 with a consistent tree topology indicating high
LD and reduced recombination (**Fig 3A, B**). Using this sequence we constructed a coalescent
tree across 210 human haplotypes with the T2T chimpanzee genome as the outgroup. This tree
reveals that H1 *KANSL1* duplications (i.e. H1.β2) are largely found in one cluster with a handful
of samples falling in an independent region of the tree, potentially resulting from an independent
origin, or a recombination event.
In contrast, *NSF* duplications (γ2, γ3) likely exhibit several independent origins throughout the
tree. One particularly large cluster of γ3 duplications was identified, within which all observed γ4
duplications were also found. We also find independent origins of the sub-haplotypes cataloged
in **Fig 1A** including multiple origins of the H1.β1.γ1.3 structure. Together, these results highlight
that the structure of H1 haplotypes at the 17q21.31 locus is highly mutable and predisposed to
recurrent structural variation. This includes large-scale gene duplications such as *KANSL1* and
NSF as well as smaller SVs including copy number changes of *LRRC37A.*

Population diversity of the 17q21.31 locus worldwide using short read data

Despite their immense power, existing long-read datasets remain limited in sample size and
population diversity. Nevertheless, large-scale short-read sequencing datasets from diverse
human populations have recently been generated. To characterize patterns of 17q21.31
variation across the world we analyzed short-read sequencing data from 5174 samples
encompassing 107 populations representative of human diversity worldwide (see **Methods,**
Figs S3-5). Using SNPs tagging the inversion as well as read-depth based copy number
quantification we estimated frequencies of four simplified haplotype architectures which could
be accurately determined from the short read data: H1.β1, representing H1 haplotypes without
KANSL1 duplications; H1.β2+, representing H1 haplotypes with *KANSL1* duplications; H2.α1
representing H2 haplotypes without *KANSL1* duplications; and H2.α2+ representing H2
haplotypes with *KANSL1* duplications (**Fig 4A-D**).
198
Worldwide, the most common haplotype is H1.β1,confirming previous work using more limited
genome sequencing datasets[19,20]. However, we observe extreme population stratification of
complex haplotypes at this locus. European populations exhibit the highest worldwide frequency
of H2.α2+ haplotypes (10%-30.58%) and intermediate haplotypes frequencies (10.9%-31.5%) of
H1.β2+. In contrast, South Asian populations, which have been historically underrepresented in

sequencing datasets, exhibit the highest worldwide frequency of H1.β2+ haplotypes (14.6%-
66.8%) and lower levels of H2.α2+ (1.4%-10.9%). East Asian populations are essentially fixed
for H1.β1 haplotypes. We also assessed whole genome sequencing data from 368 African
genomes from the CAAPA2 consortium demonstrating limited haplotype diversity in Africa, with
most populations exhibiting H1.β1. However, we observed H2.α1 haplotypes at low frequency in
Khomani San, Nama, Mbuti, Biaka, Bench, and Sheko populations. H2.α2+ haplotypes were
also present at low frequencies in some African populations. These results confirm previous
analyses highlighting the presence of rare H2.α1 haplotypes almost exclusively in African
populations[19] however also highlight that *KANSL1* duplication-containing haplotypes (including
the inverted H2.α2+ haplotypes) have increased dramatically in both Europe and South Asia.
214
While short-read sequencing data do not enable characterization of phased *NSF* duplications
(i.e. γ2+) we calculated diploid *NSF* copy number genotype frequencies worldwide confirming
the highest *NSF* copy numbers in East Asia (**Fig 4E**). We find that the overwhelming majority of
NSF duplications occur on the H1.β1 background. However, we do observe evidence of both
NSF duplications and deletions happening at low frequencies on H2.α2 background (**Fig 4F**).
Together, these results provide the most comprehensive catalog of 17q21.31 diversity in
humans to date and highlight the extreme stratification of haplotypes between continental
populations.

Double recombinants and complex gene conversion events between structural
haplotypes
Inversions can strongly suppress recombination impacting patterns of linkage disequilibrium
(LD), however genetic exchange between inversion alleles can occasionally occur through
double recombination or gene conversion events. To understand how the 17q21.31 inversion
has impacted patterns of recombination we used SNPRelate [(Xiuwen Zheng [aut, cre, cph],](#)
[Stephan...)](#), a windowed PCA-based analysis, to cluster 3135 short-read samples in 10kb
sliding windows across the locus (**Fig 5A, B**). Most individuals consistently clustered into one of
three groups corresponding to H1/H1, H1/H2, H2/H2 genotypes throughout the non-duplicated
part of the inversion and even ~80kb beyond the proximal inversion breakpoint, highlighting the
far-reaching effect of inversions in suppressing recombination. Despite the persistence of this
long 900kb LD block, we identified at least three instances of recombination at the proximal end
of the extended LD block (**Fig 5C**) occurring primarily in African individuals (32/56, 57%). We
also noted several individuals exhibiting 20-180 kb stretches showing switching between
genotype groups within the inversion region, indicating double crossover recombination events,
or possibly, long stretches of interlocus gene conversion (**Fig 5A-C**). Of the 8 identified switch
events, two exceeded 140 kb, occurring in a single European individual and a single South
Asian individual respectively, and represent cases of double crossover recombination. The
remaining events range from 20-70 kb and were found almost exclusively in African populations
(51/54, 94%). We also noted that this windowed PCA approach identified more complex
clustering patterns at the proximal and distal ends of the inversion locus that contain
duplications (**Fig 5B**). We found that these patterns corresponded strongly with the complex
structural haplotypes, thus suggesting that complex structural variation can be detected from
short read-based SNPsin some cases. Together these results demonstrate that despite

extensive inversion-driven recombination suppression, substantial exchange has happened
between vastly divergent haplotypes.

Ancient genomes track the rise in frequency of *KANSL1* duplications in Europe

To determine how 17q21.31 structural haplotype frequencies have changed in Europeans over
recent history we analyzed 626 ancient genomes spanning ~10 thousand years of human
history[21] **(Fig 6A)**. European Hunter Gatherer populations, dating from ~5000-10,000 years
before present (BP) exhibited extremely low haplotype diversity with H1.β1 making up >95% of
haplotypes **(Fig 6B)**. Higher frequencies of H2.α2+ haplotypes appear ~7 to ~5 kya BP in Early
and Neolithic Farmers (25%-31%). Steppe Pastoralists also exhibit H2.α2+ haplotypes (10%) as
well as high frequencies of the H1.β2+ haplotype (30%). To understand the temporal dynamics
of haplotype frequency changes over the last 10 kya we fit a multinomial logistic regression
model to these ancient genomic derived data **(Fig 6C)**. This model demonstrates that while
H1.β1 was the dominant haplotype ~11 thousand years ago with a frequency of >90%, it has
subsequently declined ~1.6-fold. Concomitantly, the H1.β2+ and H2.α2+ haplotypes have
increased from ~1% and ~7% to 29% and 21% respectively. Together, these data demonstrate
that 17q21.31 haplotype diversity in Europe today is the result of a massive increase in
frequencies of the *KANSL1* duplication in both H1 and H2 haplotypes over recent history.

DISCUSSION
Here we leverage long-read haplotype-resolved human genomes, newly assembled T2T and
population-scale long-read sequenced great ape genomes, diverse human short-read data, and
ancient DNA to assess the diversity and recent evolution of the 17q21.31 locus. Together, our
results provide the most comprehensive assessment to date of this remarkable region of the
genome providing new insights and clarifying previous works. For instance, while it has been
established that the H2 haplotype is the ancestral orientation of the 17q21.31 locus, the
presence of the inversions in chimpanzees, bonobos, and orangutans has been interpreted as
recurrent "toggling" of the inversion across great apes. Here, by using population-scale
haplotype resolved assemblies from chimpanzees we resolve the structure and sequence of the
chimpanzee 17q21.31 inversion. We find that the chimpanzee inversion is substantially larger
than the human inversion. While it overlaps the human inversion across ~750kb of sequence it
extends an additional ~650kb. Furthermore, we find that this inversion occurred far more
recently than the human inversion, ~330,000 years ago, in contrast to previous estimates[16].
Thus, "highly mutable and recurrent" might be a more apt description of the inversion locus than
"toggling" as independent inversion events with distinct breakpoints have occurred on different
lineages. However, additional population-scale long-read sequencing of primate genomes will
be required to determine the sequence and structures of the bonobo and orangutan inversions
and discern their evolutionary history.
283
The H2 haplotype has been associated with increased fecundity in Icelandic[14] and Polish[22]
populations and has thus been suggested to be the target of selection in Europeans. Indeed, H2
haplotype frequencies are highest in European populations. We used ancient genomes to
reconstruct allele frequency trajectories of 17q21.31 haplotypes finding that while H1.β1
haplotypes have declined in frequency markedly over the past 10,000 years, H1.β2+ and

H2.α2+ haplotypes have increased in frequency substantially: ~29-fold for H1.β2+ and 3-fold for
H2.α2+. Thus, the group of haplotypes containing duplications of the *KANSL1* gene has
increased in frequency ~6-fold. Prima facie such an increase in frequency would correspond to
a selection coefficient of approximately 0.015, similar to the average selection coefficients for
regions such as lactase persistence. Critically, however, this estimate is an extremely crude
oversimplification which neglects the small sample sizes in the ancient data and the importance
of modelling the complex demographic history of Europe. Nevertheless, these results provide an
intriguing alternate hypothesis: that duplications of the *KANSL1* gene may be the targets of
selection as opposed to the inversion itself.
298
The availability of increasingly diverse population-scale short-read sequencing of humans
enabled us to characterize 17q21.31 haplotype diversity worldwide. Our results confirm previous
work highlighting the prevalence of the H1.β1 haplotype in Africa with the ancestral H2.α1
haplotypes present at low frequencies in some populations such as the Khomani San. However,
our analyses also provide some of the first observations of 17q21.31 diversity in South Asian
populations, which have historically been undersampled. We find surprisingly high frequencies
of the H1.β2+ and H2.α2+ haplotypes across South Asia. These results mirror those of Europe
with the two *KANSL1* duplication-containing haplotypes showing increased frequencies
compared to African and East Asian populations, albeit with differing relative frequencies of the
H1.β2+ and H2.α2+. These results potentially suggest that a recent increase in duplicated
haplotypes may have also occurred in South Asia mirroring the increase in Europe, though
additional analyses are needed to confirm that hypothesis.
311
Finally, we discovered several double recombination events inside the 17q21.31 inversion
locus. Intriguingly, these events are overwhelmingly found in African populations, many of which
have very low H2 haplotype frequencies. Therefore, these double recombination events are
likely quite old, having occurred when H2 haplotypes were at higher frequencies in Africa. We
also identified two other much longer double recombination events present in two non-African
individuals. In addition, we identified a region intersecting the *CRHR1* gene where H1 and H2
haplotypes are completely homogenized and individuals cannot be assigned to one haplotype
or the other (**Fig 5A-C**). This finding has previously been reported and explained as the result of
an ancient double recombination event[19]. Crucially, unlike the other double recombinants that
we identified, this event has likely been fixed in H2 haplotype, and the original sequence
associated with the H2 haplotype has been lost in all human populations. Except for these rare
double recombination events, however, we found that recombination is completely suppressed
across the inversion. An additional consequence of segregating inversions is that by
suppressing recombination they can promote the accumulation of deleterious variation[9]. It
remains an unanswered question how the increased frequency of 17q21.31 inversion in specific
human populations may affect mutational load.
The importance of the 17q21.31 locus in human disease, human diversity, and evolution has
been the focus of many studies since its discovery. Nevertheless, long read sequencing, ancient
genomes, and additional human population datasets are continuing to increase our
understanding of this region. Our insights synthesize previous work with these new advances in

genetics and genomics highlighting both new insights and posing new avenues for future
research.

METHODS

Datasets

Short-read sequencing data were compiled from high-coverage resequencing of 5174
individuals from the 1,000 Genomes Project (1KG), the Human Genome Diversity Panel
(HGDP) and the Simons Genome Diversity Panel (SGDP). We also obtained unpublished short-
read sequencing data of 318 individuals from the CAAPA2 Data consortium. In total we were
able to infer the complex genotypes of 4585 individuals with high confidence from these
contemporary samples. Among these 697 are trios from 1KG and the remainder 3888 are
unrelated individual samples compiled from 1KG, HGDP, SGDP, and CAAPA2. The trios are
only used for method validation purposes. Genotypes from 626 historical samples from
between -12kya and 1000 ya were obtained from [23]. Historical samples were assigned to
populations based on their original publications [5,21]. 216 long-read haplotype assemblies were
compiled from the HPRC and HGSVC [3]. In addition we included the GRCh38 genome and
CHM13 reference genomes to our dataset[24]. Haplotype assemblies were only included if they
spanned the 17q21.31 locus. This excluded three assemblies `HG0015#hap1, HG00320#hap2,`
`and NA21093#hap1` from our dataset, although they were used in characterization of complex
structures in the locus. These three assemblies that were excluded appear to have been
misassembled across the locus as they were either discontiguous, or had a partial *KANSL1*
duplication that could not be cross-validated with short-read sequencing data. We obtained
great ape genome assemblies for orangutans, gorillas, bonobos, and chimpanzees from three
sources: GenomeArk (eight haplotypes, two per species), and 57 newly generated and
unpublished gorilla, chimpanzee, and bonobo genomes. For the purposes of this study, we
include the unpublished assemblies for the 17q21.31 region, with full genome releases planned
alongside separate publications currently in preparation.

Characterization of complex structures at the 17q21.31 locus in humans

We extracted the 17q21.31 region including flanking sequences from the GRCh38 reference
genome (chr17: 45275488-46921902). We then used minimap2[25] (-x asm5) to identify the
homologous sequences in all the samples in the HGSVC and HPRC genomes. We performed a
principal bundle decomposition using PanGenomic Research Tool Kit (PGR-TK) for 216 of the
HGSVC and HPRC genome assemblies with the parameters of `-w 80 -r 12 --min-span 32`
`--bundle-length-cutoff 10000 --min-branch-size 16 --min-cov 0`[26]. We examined
the results, and excluded five assemblies that were potentially misassembled. We further
excluded bundles that were unique to just one haplotype, or those whose length were in the
lowest quantile across all the remaining bundles (i.e. < 23kb). We grouped the haplotypes into
representative structures based on the order and orientation of the principal bundles. The
position of genes along haplotypes was determined by mapping coding sequences to
haplotypes using minimap2[25] (-c -x asm5). We chose one haplotype within each unique
structure and used PGGB to build a pangenome graph[27]. We then used odgi similarity to
generate a matrix of jaccard distance between each unique haplotype structure and made a

neighbor joining tree to cluster these haplotype structures. We ran minimap2[25] (`-x asm5 -c --`
`eqx -D -P --dual=no`) on pairs of assemblies to create sequence alignments, which were
then inputted to SVByEye[28] for visualization.

Characterization of complex structures at the 17q21.31 locus in great apes

We used minimap2[25] (-x asm20) to identify the orthologous sequences in the great ape
assemblies to the human 17q21.31 locus in GRCh38 reference genome (chr17:45275488-
46921902). Because we observed an inversion in chimpanzees that spans a longer region than
the inversion in humans, we then manually extended these sequences by 2Mb on either flank in
both great apes and all human haplotypes, so that these sequences span the entirety of the
chimpanzee inversion. We then reran PGRTK on these extended human sequences with the
same settings and projected the principal bundles to the great ape sequences. Within each
species, we categorized these haplotypes by their inversion, *KANSL1* and *NSF* duplication
status. We then selected the one representative haplotype for each unique structure in each
species for visualization. We ran minimap2[25] and SVByEye[28] on pairs of great ape and human
haplotypes with the same settings except -x asm20 was used as a flag in minimap2[25] instead of
-asm5. To estimate the divergence time of structural haplotypes across great apes, we first
identified the largest region of shared homology within the inverted segment (corresponding to
the blue bundle in Fig. 1A). We performed pairwise alignments using minimap2[25], with the
CHM13 human reference sequence as the query, against 17 great ape assemblies: all great
ape haplotypes from GenomeArk (including 4 orangutan, 2 gorilla, 2 bonobo, and 2 chimpanzee
haplotypes), 4 additional gorilla haplotypes, the inverted chimpanzee haplotype, and
representative human haplotypes H1.β1.γ1, H1.β2.γ1, and H2.α2.γ2. Multiple sequence
alignment was performed using kalign[29]. Phylogenetic trees were inferred using IQ-TREE with
the Jukes-Cantor substitution model and 1000 bootstrap replicates Divergence times were
estimated using the orangutan–human/chimpanzee/gorilla split `18.13` mya as a calibration
point[30]. Trees were visualized using the ggtree R package[31].

Evolutionary history of structural variation at the 17q21.31 locus in humans

To date the emergence of different H1 haplotype structures we first needed to find a unique
sequence at the 17q21.31 locus that is physically linked to the *KANSL1* and *NSF* duplications
and not broken down by recombination. The largest region of shared homology within the
inverted segment (corresponding to the blue bundle in Fig. 1A) in CHM13 is also the closest in
proximity to the *KANSL1* duplications. Therefore, we used similarities in phylogenetic tree
topology as a proxy for linkage disequilibrium across this region. This region was divided into
nine non-overlapping 50 kb windows using seqkit (`sliding -s 50000 -W 50000`)[32]. All 210
human haplotypes that were included in previous analyses were included to construct the trees,
as well as 2 GenomeArk chimpanzee haplotypes. For each of the nine windows, sequences
were aligned using kalign[29], and maximum likelihood trees were inferred with IQ-TREE[33] using
the Jukes–Cantor model and 1000 ultrafast bootstrap replicates (`-m JC -bb 1000 -nt AUTO`).
Trees were rooted using the two chimpanzee haplotypes as an outgroup. To quantify
topological similarity between trees, we computed pairwise Robinson–Foulds (RF) distances.
An RF distance matrix was then constructed for all pairwise tree comparisons, and the windows

with the lowest distance to the right-most window (i.e. the window that is closest to the *KANSL1*
duplication) were selected. We then merged the selected windows, and reran IQ-TREE using
the chimpanzee-human split time `7.74` mya as a calibration point for the emergence of
haplotype structures[33].

Using contemporary short read genomes to understand 17q21.31 population diversity in
modern human populations

Tag SNP based inversion genotyping

In order to determine the genotype status of individuals in our dataset, we used a total of 1271
tag SNPs to genotype the 17q21.31 locus based on the method used in[23] For each individual,
we constructed a genotype vector by coding observed genotypes at these SNPs as 0
(homozygous reference), 1 (heterozygous), or 2 (homozygous alternate). Then, Euclidean
distance was calculated between the observed genotype vector and expected genotype vectors
for H1/H1, H1/H2, and H2/H2. Each individual was assigned the inversion genotype with the
minimum Euclidean distance to their observed genotype.

Read-depth based copy number genotyping at NSF and KANSL1

Copy number genotypes for the 1000 Genomes, HGDP, and SGDP, and CAAPA2 individuals
were estimated using read depth as described in Bolognini et al[5]. In brief, read depths were
quantified from CRAM (CAAPA2) or BAM (1000G, HGDP, SGDP) files. CRAM files from
CAAPA2 were based on a different version of the GRCh38 reference genome, and to ensure
consistency across datasets, we remapped this data to the same version of the reference
genome as the other three datasets (GRCh38_full_analysis_set_plus_decoy_hla). We
generated d4 files from these BAM files. From the d4 files we calculated the average read
depths in 1,000 bp sliding windows in 100 bp steps across the 17q21.31 locus. These depths
were normalized to the average read depth within a control region chr17:42800000-46000000
within the 17q21.31 locus for each individual, in which no copy number variation was observed
in more than 3,000 individuals (i.e. individuals from the 1KGP and HGDP samples). To
determine the copy number of the α and β segments of the *KANSL1* duplication, we averaged
the copy number of windows that overlapped with the unique region of each duplication (α:
chr17:46143000-46238000 and β: chr17:46095000-46123000). In the case of *NSF* copy
number, since the GRCh38 reference genome has two copies of *NSF*, we summed the
normalized read depth across two target regions corresponding to the two *NSF* copies
(chr17:46,336,376–46,489,410 and chr17:46,564,311–46,707,123). CAAPA2 samples were
excluded from *NSF* analysis because of difficulties in aligning the sequencing reads to this
complex region. Lastly, we removed samples whose copy number estimates were too large to
be realistic (i.e. copy number > 20 for *KANSL1* and > 50 for *NSF*).

Determining haplotypes from 17q21.31 genotypes

We were able to associate the β and α duplications with the inversion status of their haplotype
background because the β duplication is always associated with the H1 haplotypes and that the
α duplication is always associated with the H2 haplotypes. Therefore, it was possible to
integrate the inversion genotypes (H1/H1, H1/H2, H2/H2) and copy numbers (based on read

depth in sector calls) to assign complex genotypes representing both inversion and *KANSL1*
duplication status as described in[23] **(Fig. S4)**. We then used R to plot the average copy number
across the β and α regions, and assigned genotypes based on their inversion status and copy
number. The samples clustered into duplication genotypes with the exception of a handful of
samples which did not match their inversion calls. In cases where multiple interpretations of
duplication status are possible, the more probable assignment is listed (e.g., H1D/H1D vs.
H1/H1DD we chose the former because H1.B3 is very rare). Since there is more diversity in
NSF copy numbers compared to *KANSL1* especially on the H1.β1 background, we were unable
to phase *NSF* duplications and only provide diploid copy number estimates. Complex genotypes
for the historical short-read genomes were obtained from Irving-Pease[23].

Sliding window PCAs reveal regions of double recombination

In order to determine recombination or gene conversion events within the inversion locus and
extended LD of 100kb, PC1 values were computed in non-overlapping 10kb windows for all
individuals using SNPRelate[34]. Tag SNP based inversion genotyping revealed three clusters:
H1 homozygous (bottom), H2 homozygous (top), and H1/H2 heterozygous (middle). We
computed the mean, median, and multiple quantile ranges of PC1 for individuals of each
inversion genotype in every window. To identify noisy or ambiguous positions, we evaluated
whether the variance for different genotypes overlapped at each position. Positions with
overlapping variance were flagged and excluded from downstream interpretation. The region of
interest was further refined to non-duplicated regions except for the unique *KANSL1*. To refine
outlier classification, we focused on runs of at least two adjacent outlier windows within the
same sample. For these regions, we reassigned the genotype based on directional deviation of
PC1: H1H2 genotypes with PC1 below the genotype's 5th percentile were reassigned to H1H1,
while those above the 95th percentile were reassigned to H2H2. Homozygous genotypes were
reassigned to H1H2 if their PC1 values deviated substantially in either the positive or negative
direction respectively. Further manual curation allowed us to determine single recombinant
events and double recombinant events (**Table S4**).

DECLARATION OF INTERESTS

The authors declare no competing interests.

ACKNOWLEDGEMENTS

We thank R Nielsen, BM Henn, P Moorjani, CT Miller, JM Vazquez and J Chin for helpful
discussion. This work was supported by NIH National Institute of General Medicine award
R35GM142916 to PHS and NIH National Human Genome Research Institute award
R01HG013017 to PHS and a Leakey Foundation award to SS. This manuscript is the result of
funding in whole or in part by the National Institutes of Health (NIH). It is subject to the NIH
Public Access Policy. Through acceptance of this federal funding, NIH has been given a right to
make this manuscript publicly available in PubMed Central upon the Official Date of Publication,
as defined by NIH. The content is solely the responsibility of the authors and does not
necessarily represent the official views of the National Institutes of Health.

AUTHOR CONTRIBUTIONS

PHS conceived of the study. SS, RNL, SF, JLR, and PHS performed all analyses and wrote the manuscript. MWM and ANK provided primate cell lines and tissue culture. VB, SGMM, SG, and BH provided data.

DATA AND CODE AVAILABILITY

All data used in this project are publicly available and described in the 'Datasets' section of the Methods and at the following github link https://github.com/sudmantlab/17q21.31_SV_and_evolution which is archived in Zenodo: https://doi.org/10.5281/zenodo.16883463 (Sridharan 2025). Copy numbers, genotypes, and structural haplotypes can be found in the Supplementary tables. The HPRC data can be obtained at: https://humanpangenome.org/data/. The HGSVC data can be obtained at: https://www.hgsvc.org/resources, The 1KG and HGDP data can be obtained at: https://www.internationalgenome.org/data/. The SGDP data can be obtained at: https://www.simonsfoundation.org/simons-genome-diversity-project/. The ancient data are available on the European Nucleotide Archive under the accession codes PRJEB64656 and PRJEB50857. The unpublished data for bonobos, chimpanzees and gorillas can be obtained at: https://github.com/sudmantlab/17q21.31_SV_and_evolution. The GenomeArk long read haplotype resolved assemblies for primates can be obtained at: https://www.genomeark.org/

FIGURE TITLES AND LEGENDS

Figure 1 – Distinct 17q21.31 structural haplotypes from human long-read genome assemblies: a) Schematic of haplotype structure at the 17q21.31 locus and flanking regions using PGRTK for the reference CHM13 genome. Large colored arrows represent principal bundles, which are individual units of repeats. Bundles of the same color represent homologous sequences. The directionality of the arrow represents the bundle orientation. Unique flanking regions are indicated in greyscale. Smaller labeled arrows denote genes and gene orientations.

b) Schematic of 11 unique structural haplotypes found within 210 long-read haplotype-resolved
assemblies. Each row represents a unique structural haplotype. Structural haplotypes are
arranged using hierarchical clustering based on jaccard distance, with tip sizes scaled by the
number of assemblies sharing each haplotype. **c)** Visualization of sequence alignment among
11 unique structural haplotypes. Each row represents a unique haplotype structure arranged in
the same order as in **b**. Alignments are colored by orientation: grey denotes alignments
between two structural haplotypes in the forward orientation and pink denotes alignment in the
reverse direction.
527

528
**Figure 2 – Characterizing the evolutionary context of the 17q21.31 inversion in the great
ape lineage: a)** Visualization of sequence alignments spanning the 17q21.31 locus and flanking

regions in great apes. The sequences are extended from **Fig 1** by 2 Mb on both ends to
characterize the shared homology of flanking regions across all species. Each row represents a
representative haplotype from each of the great apes. Alignments are colored by orientation
(gray = forward, pink = reverse). **b)** Alignment visualization comparing representative human
haplotypes H1.β1.γ3 and H2.α2.γ2 and chimpanzee haplotypes H1p and H2p (as seen in **a**),
reordered to highlight the distinct, and longer inversion found in chimpanzees. **c)** Unique
structural haplotypes in great apes with two representative human structural haplotypes shown
as reference. Human haplotypes are plotted along with gene labels and gene orientation.
Sequences are visualized with PGRTK trained on human structures with extended sequence as
seen in **a**. Unique flanking regions are shown in dark grey and lighter grey. Colors are broadly
consistent with **Fig 1b** although the longer genetic interval used here results in slight
differences. **d)** Dating the divergence of great ape and human 17q21.31 haplotypes. Trees were
inferred using a large non-duplicated sequence located within the human inversion (i.e. the blue
bundle in **c**, **see methods)**. Several important split times are annotated in the tree. The inset
shows two main topologies that are observed via bootstrapping and their frequencies.
546
547

Figure 3 – Timing and recurrence of human 17q21.31 structural haplotypes: a) Robinson-
Foulds (RF) distance clustering heatmap across phylogenetic trees constructed with non-

overapping 50 kb windows across all human long-read assemblies. Windows are extracted from
a large non-duplicated sequence located within the human inversion (i.e. the blue bundle in **Fig
1)**. Darker red indicates greater similarity (lower RF distance) between two trees. Dendogram
summarizes the topological similarity across all windows. Inset histogram displays the
distribution of RF distance values across all comparisons. **b)** Euclidean distance along PC1 and
PC2 axes between the phylogenetic tree constructed at window 9, the closest unique sequence
to *KANSL1*, and trees at all other windows. A loess-smoothed curve is shown in red. **c)** Dated
maximum-likelihood tree constructed using sequences at windows 8 and 9 and rooted with T2T
chimpanzee haplotypes as the outgroup. Human-chimpanzee split-time was used to date the
tree. Select internal nodes are labeled with divergence time. Tip symbols indicate copy-number
variation of *KANSL1* (blue squares) and *NSF* (green triangles).
562
563

564
Figure 4 – Population diversity of the 17q21.31 locus worldwide using short read data: a)
World map indicating the frequency of direct H1.β1, H1.β2+ and inverted H2.α1 and H2.α2+

17q21.31 haplotypes in 4585 individuals from 193 populations. Insets show the frequency
distribution in South Asia (**b**), Europe (**c**) and Africa (**d**). Pies are scaled by sample size in **b** and
c. Only populations with five or more samples are plotted in **a-c**, whereas **d** shows populations
with as few as a single sample. In **d**, only populations with H1.β1 frequencies less than 1 are
labeled. See **Supplementary Table 2** for exact sample sizes and haplotype frequencies of all
populations. **d)** World map indicating the average diploid *NSF* copy number in 3352 individuals
from 147 populations. Points are scaled based on sample size, with only populations with five or
more samples plotted. See **Supplementary Table 3** for the exact sample sizes and average
diploid copy numbers of all populations. **e)** The distribution of *NSF* copy numbers in individuals
with different inversion and *KANSL1* duplication genotypes.

577
578
Figure 5 – Population specific instances of recombination events at the 17q21.31 locus:
The x-axis represents genomic position across a subset of the 17q21.31 locus including the
inversion region and surrounding flanking sequences. A PGRTK-based haplotype structure of
the reference genome (hg38) is shown below the figure. **a)** 10kb sliding window PCA. Each
colored line represents the variation in the position of an individual on the first principal
component axis, and is colored by inversion genotype. **b)** 10kb sliding window PCA faceted by
inversion status and colored by complex genotype. **c)** All recombination events identified in the
region of interest. Gradient blocks represent single recombination events wherein the right edge
of the block represents the recombination breakpoint. Black boxes represent double

recombination events. Numbers next to each block indicate the number of individuals with that
recombination event, and pie charts indicate the population of these individuals.

590
Figure 6 – Ancient genomes track the rise in frequency of *KANSL1* duplication in Europe:
a) Geographic distribution of historical samples across Western Eurasia shown in 2.5 kya time
slices from 10 kya to the present. Samples are split based on their haplotype structure (H1.β1,
H1.β2+, H2.α1, and H2.α2+) and age in each subplot, with dots marking individual sampling

locations. **b)** Temporal distribution of 17q21.31 haplotypes across historical West Eurasian
populations. Stacked bar plots show the number of haplotypes present in each population over
a time bin. Time is shown on the x-axis in years before present (BP). Pie charts on the right
summarize the overall haplotype frequency for each population, with the. **c)** Modeled temporal
trajectories of 17q21.31 haplotype frequencies over the past 10 ky using a multinomial logistic
regression model. Each point represents the observed haplotype frequency in a time bin (all the
populations are pooled), with point size scaled by sample size.
602
Supplementary Figure 1 – 24 distinct 17q21.31 structural haplotypes from human long
read genome assemblies: Schematic of 24 unique structural haplotypes found within 210 long-
read haplotype-resolved assemblies. Each row represents a unique structural haplotype.
Structural haplotypes are arranged using hierarchical clustering based on jaccard distance, with
tip sizes scaled by the number of assemblies sharing each haplotype. The dashed red line
indicates where clades were collapsed into a single representative structure as shown in **Figure**
1B.
610
Supplementary Figure 2 - Comparative gene organization and segmental duplications at
the 17q21.31 locus in humans and chimpanzees: (Top) Gene names and orientation across
the human reference (hg38) 17q21.31 inversion locus and extended distal flanking sequence.
The black rectangle denotes the inversion breakpoints of the 17q21.31 locus; orange and yellow
rectangles denote the segmental duplications; protein coding genes are shown in blue; non-
coding transcripts are shown in green; and pseudogenes are shown in purple. (Bottom)
Orthologous inversion in chimpanzees shows a larger inversion region (by size) and additional
genes in the chimpanzee region that are not shown in the inversion region in humans.
619
620   **Supplementary Figure 3 – Inversion genotype assignments mapped atop copy numbers**
621   **in the β and α regions:** Each point represents an individual with the x-axis denoting the
622   normalized read depth across β and the y-axis denoting the normalized read depth across α.
623   Points are colored according to inversion genotype, assigned using 1271 tag SNPs. Jitter was
624   applied to both the x and y axes to improve visualization of overlapping points.
625
Supplementary Figure 4 – Table assigning complex genotype status based on α and β
regions: Cartesian coordinates (x,y) represent complex genotype assignments with numbers
indicating the average copy number as discrete values. The value before the comma
corresponds to region β, and the value after the comma corresponds to region α. Red cells
denote ambiguous cases which can only be distinguished based on 1271 tag SNPs that are
required to differentiate between inversion status. Blue cells represent the most conservative
assignment which was used in ambiguous cases within the direct or inverted haplotypes.
633
Supplementary Figure 5 – Complex genotype at the *KANSL1* region mapped atop copy
numbers in the β and α regions: Each point represents an individual with the x-axis denoting
the normalized read depth across β and the y-axis denoting the normalized read depth across
α. Points are colored according to complex genotype status, assigned based on the *KANSL1*

region using the criteria described in Supplementary Figures 2 and 3. Jitter was applied to both
the x and y axes to improve visualization of overlapping points.
640
Supplementary Figure 6 – Read depth across the 17q21.31 locus split by CNV assigned
by *KANSL1* calls: Each line represents an individual colored and grouped by complex
genotype. The x-axis denotes genome position, and the y-axis shows the average normalized
read depth calculated in sliding windows of 1000 bp. Grouping by complex genotype enables
visual comparison of CNV patterns across complex genotypes. Horizontal grey lines at the
bottom of each plot indicate the positions of the α and β regions, highlighting the differences in
read depth between complex genotypes. This visualization complements the discrete copy
number genotype assignments shown in Supplementary Figures 3 and 4, illustrating the
underlying read depth variation that defines α and β.

TABLES

Supplementary Table 1 - Inversion tagging SNPs
Supplementary Table 2 - Frequencies of H1.β1, H1.β2, H1.β3, H2.α1, and H2.α2 in 1000
Genomes, HGDP, SGDP and CAAPA2 samples
Supplementary Table 3 - Frequencies of diploid *NSF* copy numbers in 1000 Genomes, HGDP,
SGDP and CAAPA2 samples
Supplementary Table 4 - Recombinant individuals and recombinant breakpoints in 1000
Genomes, HGDP, and SGDP samples
Supplementary Table 5 - Frequencies of H1.β1, H1.β2, H1.β3, H2.α1, and H2.α2 in Ancient
European Genomes

REFERENCES

1. Porubsky, D. *et al.* Recurrent inversion polymorphisms in humans associate with genetic
instability and genomic disorders. *Cell* **185**, 1986–2005.e26 (2022).
2. Plender, E. G. *et al.* Structural and genetic diversity in the secreted mucins MUC5AC and
MUC5B. *Am. J. Hum. Genet.* **111**, 1700–1716 (2024).
3. Logsdon, G. A. *et al.* Complex genetic variation in nearly complete human genomes.
bioRxiv (2024) doi:10.1101/2024.09.24.614721.
4. Scheer, K. *et al.* Adaptive increase of amylase gene copy number in Peruvians driven by
potato-rich diets. *bioRxiv* (2025) doi:10.1101/2025.03.25.644684.
5. Bolognini, D. *et al.* Recurrent evolution and selection shape structural diversity at the
amylase locus. *Nature* **634**, 617–625 (2024).
6. Ebert, P. *et al.* Haplotype-resolved diverse human genomes and integrated analysis of
structural variation. *Science* **372**, (2021).
7. Thomas, J. W. *et al.* The chromosomal polymorphism linked to variation in social behavior
in the white-throated sparrow (Zonotrichia albicollis) is a complex rearrangement and
suppressor of recombination. *Genetics* **179**, 1455–1468 (2008).
8. Hager, E. R. *et al.* A chromosomal inversion contributes to divergence in multiple traits
between deer mouse ecotypes. *Science* **377**, 399–405 (2022).
9. Jay, P. *et al.* Mutation load at a mimicry supergene sheds new light on the evolution of
inversion polymorphisms. *Nat. Genet.* **53**, 288–293 (2021).

10.   Wang, H. *et al.* Chromosomal inversion polymorphisms shape human brain morphology.
Cell Rep. **42**, 112896 (2023).

11.   Le Moan, A. *et al.* Coupling of twelve putative chromosomal inversions maintains a strong
barrier to gene flow between snail ecotypes. *Evol. Lett.* **8**, 575–586 (2024).

12.   Harringmeyer, O. S. & Hoekstra, H. E. Chromosomal inversion polymorphisms shape the
genomic landscape of deer mice. *Nat. Ecol. Evol.* **6**, 1965–1979 (2022).

13.   Jamsandekar, M. *et al.* The origin and maintenance of supergenes contributing to
ecological adaptation in Atlantic herring. *Nat. Commun.* **15**, 9136 (2024).

14.   Stefansson, H. *et al.* A common inversion under selection in Europeans. *Nat. Genet.* **37**,
129–137 (2005).

15.   Koolen, D. A. *et al.* Mutations in the chromatin modifier gene KANSL1 cause the 17q21.31
microdeletion syndrome. *Nat Genet* **44**, 639–641 (2012).

16.   Zody, M. C. *et al.* Evolutionary toggling of the MAPT 17q21.31 inversion region. *Nat Genet*
40, 1076–1083 (2008).

17.   Pedicone, C., Weitzman, S. A., Renton, A. E. & Goate, A. M. Unraveling the complex role
of MAPT-containing H1 and H2 haplotypes in neurodegenerative diseases. *Mol.*
Neurodegener. **19**, 43 (2024).

18.   Karamik, G. *et al.* The clinical phenotype of Koolen-de Vries syndrome in Turkish patients
and literature review. *Am. J. Med. Genet. A* **191**, 1814–1825 (2023).

19.   Steinberg, K. M. *et al.* Structural diversity and African origin of the 17q21.31 inversion
polymorphism. *Nat Genet* **44**, 872–880 (2012).

20.   Boettger, L. M., Handsaker, R. E., Zody, M. C. & McCarroll, S. A. Structural haplotypes and
recent evolution of the human 17q21.31 region. *Nat Genet* **44**, 881–885 (2012).

21.   Allentoft, M. E. *et al.* Population genomics of post-glacial western Eurasia. *Nature* **625**,
301–311 (2024).

22.   Clark, J. S. C. *et al.* Association studies between chromosomal regions 1q21.3, 5q21.3,
14q21.2 and 17q21.31 and numbers of children in Poland. *Sci Rep* **12**, 18923 (2022).

23.   Irving-Pease, E. K. *et al.* The selection landscape and genetic legacy of ancient Eurasians.
Nature **625**, 312–320 (2024).

24.   Nurk, S. *et al.* The complete sequence of a human genome. *Science* **376**, 44–53 (2022).

25.   Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–
3100 (2018).

26.   Chin, C.-S. *et al.* Multiscale analysis of pangenomes enables improved representation of
genomic diversity for repetitive and clinically relevant genes. *Nat. Methods* **20**, 1213–1221
(2023).

27.   Garrison, E. *et al.* Building pangenome graphs. *Nat. Methods* **21**, 2008–2012 (2024).

28.   Porubsky, D. *et al.* SVbyEye: a visual tool to characterize structural variation among whole-
genome assemblies. *Bioinformatics* **41**, (2025).

29.   Lassmann, T. Kalign 3: multiple sequence alignment of large data sets. *Bioinformatics* **36**,
1928–1929 (2019).

30.   Yoo, D. *et al.* Complete sequencing of ape genomes. *Nature* **641**, 401–418 (2025).

31.   Guangchuang Yu [aut, cre, cph], Tommy Tsan-Yuk Lam [aut, ths], Justin Silverman [ctb].
Ggtree. (Bioconductor, 2017). doi:10.18129/B9.BIOC.GGTREE.

32.   Shen, W., Le, S., Li, Y. & Hu, F. SeqKit: A cross-platform and ultrafast toolkit for FASTA/Q
file manipulation. *PLoS One* **11**, e0163962 (2016).

33.   Minh, B. Q. *et al.* IQ-TREE 2: New models and efficient methods for phylogenetic inference
in the genomic era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

34.   Zheng, X. *et al.* A high-performance computing toolset for relatedness and principal
component analysis of SNP data. *Bioinformatics* **28**, 3326–3328 (2012).

730