Structure and dynamics of a muti-domain nitric oxide synthase regulated by a C2 domain

Dhruva Nair and Brian R. Crane[*]

AUTHOR ADDRESS:

Weill Institute for Cell and Molecular Biology

Department of Chemistry and Chemical Biology

Cornell University, Ithaca, NY 14853, USA.

*For correspondence: bc69@cornell.edu

Short Title: syNOS is a C2-domain-regulated multidomain NOS.

Abstract

Nitric oxide synthase (NOS) is a widely studied multidomain redox enzyme that produces the key signaling molecule and cytotoxic agent nitric oxide (NO) for functions that range from mammalian vasodilation to prokaryotic antibiotic resistance. NOS enzymes from metazoans and cyanobacteria rely on dynamic associations of their oxygenase and coupled di-flavin reductase domains that have largely evaded detailed structural characterization. CryoEM studies of a representative dimeric six-domain Synechococcus NOS reveal the architecture of the full-length enzyme, which contains an unusual regulatory C2 domain, and additional nitric oxide deoxygenase (NOD) and pseudo-globin modules. Five distinct structural states depict how pterin binding couples to tight and loose oxygenase conformations and how the Ca^{2+}-sensitive C2 domain moves over 85 Å to alternatively regulate either the NOS or NOD heme center. The extended C-terminal tail and its dynamic interactions highlight an added layer of regulation required by multidomain NOSs compared to other di-flavin reductases.

Teaser: syNOS is a highly dynamic multi-domain oxidoreductase that harnesses a Ca^{2+}-sensitive C2 domain to modulate activity.

Introduction

Nitric oxide (NO) is a potent free-radical signal and cytotoxic agent found throughout biology ([1–8]). Nitric Oxide Synthases (NOS) are homodimeric P450-like enzymes that produce NO through the 5-electron oxidation of L-arginine (L-arg) to L-citrulline ([9–12]). NOSs comprise a heme-containing oxygenase domain (NOS_{Oxy}) that is reductively activated by either an appended reductase domain (NOS_{Red}, characteristic of metazoans) or an auxiliary reductase partner (characteristic of prokaryotes) (Fig 1A) ([10–12]).

Mammalian NOSs (mNOSs) are multidomain enzymes composed of an N-terminal NOS_{Oxy} and a C-terminal NADPH-dependent NOS_{Red}, bridged by a calmodulin (CAM)-binding domain (NOS_{CBD}; Fig. 1A). NOS_{Red} belongs to the cytochrome P450 reductase (CYPOR) family and as such, is further divided into a NOS_{Oxy}-proximal FMN-binding flavodoxin domain (NOS_{Fld}) and an FAD- and NADPH-binding ferredoxin-$NADP^+$ reductase module (NOS_{FNR}) ([10–13]). NOS_{Oxy} utilizes tetrahydrobiopterin (BH_4) or tetrahydrofolate for rapid heme reduction during oxygen activation ([14]). To reduce heme for oxygen binding, electrons flow from NADPH -> FAD -> FMN(H)• -> Heme-Fe^{3+}. Ca^{2+}- CAM enhances heme reduction by facilitating the traverse of NOS_{Fld} ~70 Å from a shielded position in a conformationally locked state of NOS_{Red} to a productive binding interaction with NOS_{Oxy} ([15–18]).

Given their medicinal importance ([19]), NOSs have been extensively studied in mammals, which have three isoforms: 1) endothelial NOS (eNOS, regulation of the vasculature), 2) neuronal NOS (nNOS, regulation of neuronal function), and 3) inducible NOS (iNOS, immune responses to pathogens and cancer cells) ([11], [20–22]). Most prokaryotic NOSs have fewer domains than their metazoan counterparts and have diverse functions that include participation in stress responses, antibiotic resistance and biosynthesis of secondary metabolites ([23–27]).

To mitigate NO cytotoxicity, prokaryotes oxidize NO to NO_3^- with flavohemoprotein (Hmp), which combines a heme-reducing flavin reductase with an oxygen-activating globin domain. In

contrast, mammals oxidize NO with hemoglobin, cytoglobin and neuroglobin, which obtain

reduced equivalents from various environmental factors (*28–30*).

Cyanobacterial NOSs represent a bridge between metazoan and prokaryotic NOSs (*26, 31–34*).

They comprise an mNOS architecture fused to an N-terminal globin domain, as exemplified by

the 1468-residue NOS from *Synechococcus sp*. PCC 7335 (i.e. syNOS, Fig. 1A). Like mNOS, Ca^{2+}

activates syNOS, despite syNOS lacking a CBD and Synechococcus lacking an obvious CAM

homolog (*34*). The syNOS globin domain (NOS_{Glb}) oxidizes NO to NO_3^- as a nitric oxide

dioxygenase (NOD) and, when considered with $syNOS_{FNR}$, resembles Hmp (*34*).

Ca^{2+}-CAM regulates mNOS activity by controlling the motions and associations of NOS_{Oxy} and

NOS_{Red} that allow electron transfer (ET) between them. Owing to the dynamic nature of

turnover and inactivated states of full-length NOSs, they have evaded characterization by direct

structural methods. Single particle electron microscopy (EM) in conjunction with chemical

crosslinking and hydrogen-deuterium exchange mass spectrometry (HDX-MS) have yielded

informative, if incomplete, structural information (*35–39*). These studies, along with

biochemical and spectroscopic characterization (*12, 40–44*) have motivated a model of mNOS

conformational dynamics that involves three states: i) input, wherein NOS_{Fld} closely associates

with NOS_{FNR} so that the NOS_{Red} FAD can be reduced by NADPH; ii) intermediate, wherein NOS_{Fld}

transitions between its closed position next to NOS_{FNR} and an interaction with NOS_{Oxy}; and iii)

output, wherein Ca^{2+}-CAM promotes docking of NOS_{Fld} onto the NOS_{Oxy} of the adjacent subunit

(*44–47*). Although cryoEM has provided residue-level information on NOS_{Oxy}, NOS_{Red} positioning

has been challenging to discern owing to its conformational heterogeneity (*35, 46*).

Herein, we resolve domain configurations of the multi-domain syNOS by cryoEM in conjunction

with further characterizing ET properties associated with its NOS and NOD activities. We

capture resting and dynamic states of the enzyme and identify an atypical C2-like domain that

acts as a reversible lynchpin to mediate Ca^{2+} sensitivity. An elaborated C-terminal tail (CTT) (*22*)

of NOS_{Red} , key for regulation of mNOS, maintains a conformationally locked dimer and

represents an expansion of the regulatory mechanisms seen within the di-flavin reductase

family.

Results and Discussion

<u>SyNOS expression, purification and crosslinking mass spectrometry</u>

Full-length *Synechococcus PCC 7335* NOS (syNOS) was fused to an N-terminal-twin strep affinity
tag and co-expressed along with *E. coli* HSP90 and GAPDH, in BL21(DE3) and purified through a
series of affinity and size exclusion columns (SEC) (see Methods, Fig. S1) (*46*, *48*, *49*). As found
previously (*34*), syNOS purified as a mixture of monomers and dimers. SEC fractions were
tested for NO production with L-arg and NADPH and active fractions were applied to glow-
discharged Cu-Quantifoil grids, vitrified and imaged on either a Talos Arctica or Titan Krios. Prior
to freezing, samples were incubated with or without Ca^{2+}/L-arg/NADPH and excess BH_4 (Table
S1).

To complement cryoEM and gain further insight into domain dynamics we subjected purified
syNOS dimer fractions to disuccinimidyl dibutyric urea (DSBU) crosslinking mass spectrometry
(XL-MS), under inactive (Ca^{2+}-depleted) and turnover conditions (see methods for assay
conditions). Peptide coverage was relatively high for both (inactive state, 91%; and turnover
state, 93%) but only a limited number of intra/intersubunit Lys-to-Lys crosslinks were identified
(Table S2).

<u>Structure determination of inactivated full-length syNOS</u>

SyNOS data sets and the states within are summarized in Table S1 and described below. Two
datasets of syNOS samples lacking Ca^{2+}, L-arg and NADPH were collected on a Talos Arctica
equipped with a K3 detector. Dataset (DS) 1 (1215 micrographs) and DS2 (4055 micrographs)
were initially preprocessed using CryoSPARC (Fig. S2 and S3), then subjected to particle picking
using CRYOLO and finally refined in CryoSPARC (*50*, *51*). During 2D classification multiple classes
emerged resembling a symmetric full-length dimer (syNOS-Sym). DS1 produced a syNOS-Sym
overall resolution of 3.8 Å (47 K particles), and DS2 produced a NOS-Sym map of 4.7 Å
resolution (84 K particles).

In addition, an asymmetric class (syNOS-Asym) became apparent in both DS1 and DS2 that

lacked density for one $syNOS_{Red}$ and one $syNOS_{C2}$. DS1 produced a syNOS-Asym map with the

nominal resolution of 3.67 Å (67 K particles), whereas DS2 yielded a 4.00 Å resolution map (265

K particles). In syNOS-Asym, uninterpretable electron density surrounds the position of the

vacated $syNOS_{Red}$ indicative of conformational disorder for this subunit.

3D-refinement of syNOS-Sym gave clear electron density for secondary structure, cofactors and

in many cases, residue side chains. Portions of each subunit were less well defined yet still

allowed unambiguous placement of the constituent domains. Inspection of the electron density

indicated that the homodimer interface was structurally heterogeneous. To better reconstruct

the full-length dimer, we combined the syNOS-Sym particle stacks from DS1 (47 K particles) and

DS2 (84 K particles) and re-refined the density alignment for 3D-variability (VA) and 3D-Flex

analysis. We explored C1, C2 and C2 (relaxed) symmetries which yielded 3.99 Å, 3.95 Å and 3.95

Å nominal resolutions, respectively, with the C2(relaxed) refinement chosen for further

flexibility analysis (Fig. S4, discussed below). The C2 symmetry-imposed electron density was

sufficient for model building and refinement (Table S3).

<u>Structure determination of turnover syNOS with Ca^{2+} and L-arginine</u>

DS3 (Fig. S5) (5019 micrographs) represented syNOS incubated with Ca^{2+}, additional BH_4 and L-

arg and was collected on a Titan Krios equipped with a K3 detector. Micrographs were initially

preprocessed using CryoSPARC (Fig. S5), then subjected to particle picking with CRYOLO and

finally refined in CryoSPARC. During 2D classification, multiple classes emerged resembling NOS-

Asym but also a state that lacked density for $syNOS_{Red}$ on both subunits. The latter state

resembles previously determined nNOS cryoEM structures that resolved a NOS_{Oxy} homodimer

but lacked density for NOS_{Red} owing to conformational disorder (*36*). We refer to this class as

the a "loose" turnover state (NOS-TO$_L$), because the $syNOS_{Oxy}$ interface is expanded relative to a

typical "tight" $mNOS_{Oxy}$ dimer. 2D classes of NOS-TO$_L$ gave density plumes at the interface

characteristic of poor signal alignment, indicative of a conformationally heterogeneous particle,

hence a composite map from two local refinements was created(PDB:9Q15). syNOS-Asym

(PDB:9Q0Y) was refined to a nominal resolution of 3.19 Å (71 K particles), whereas syNOS-TO$_L$

(PDB:9Q0X) was resolved to 3.6 Å resolution (61 K particles, Table S3).

DS4 was collected from the same conditions on the 200 KV Talos Arctica equipped with a K3

detector. These grids, which had thicker cubic ice, produced an intermediate resolution

turnover state with a tight syNOS$_{Oxy}$ dimer interface (syNOS-TO$_T$) that more closely resembles

crystal structures of BH$_4$-bound mNOS$_{Oxy}$ (Fig. S6). SPA is possible from cubic ice conditions,

which is known to favor contraction (shrinkage) of protein structure, although the mechanisms

for this behavior are largely unknown (*52*).

<u>Structure determination of monomeric syNOS in turnover conditions</u>

DS5 (9036 micrographs) of syNOS incubated with additional BH$_4$, Ca^{2+}, L-arg and NADPH was

collected on a Titan Krios equipped with a Falcon 3 detector (Fig. S7). For this dataset, the

homodimers appeared primarily dissociated into monomers (syNOS-Mon). Unlike syNOS-Asym,

in syNOS-Mon, even syNOS$_{Oxy}$ appeared as a monomer. The micrographs were processed

through CryoSPARC with multiple rounds of 3D particle stack cleaning to remove damaged

particles (Fig. S7). The syNOS-Mon (PDB:9Q05) density was refined to 3.09 Å resolution (45 K

particles), with discernible sidechains and cofactors, although neither Ca^{2+}, nor L-arg could be

identified in the map (Table S3).

<u>Full-length syNOS (syNOS$_{FL}$) molecular architecture</u>

In the absence of Ca^{2+}, the syNOS-Sym(PDB:9Q06) state of the full-length enzyme (syNOS$_{FL}$)

forms a compact C2-symmetric homodimer composed of six domains per subunit (Fig. 1B,C,D);

the domain arrangements divide the protein into two redox modules: i) an Fld-FNR module with

an N-terminal C2 domain (residues 1-134) bound at the Oxy-FNR interface and ii) a Glb-Oxy

module, that also contains an 8-helical pseudo-globin domain that does not bind heme

(syNOS$_{pGlb}$, residues 179-339). Three extended linkers connect the most mobile domains, with

L1 (44 residues) connecting syNOS$_{C2}$ to syNOS$_{pGlb}$, L2 (10 residues) connecting syNOS$_{Oxy}$ to

syNOS$_{Fld}$ and L3 (13 residues) connecting syNOS$_{Fld}$ to syNOS$_{FNR}$ (Fig. S8).

In this symmetric "locked" state, syNOS dimerizes through interactions of syNOS$_{Oxy}$ (residues

474-852), and syNOS$_{pGlb}$, which bridges the gap between syNOS$_{Glb}$ (residues 340-473) and

syNOS$_{Oxy}$ (Fig. 1B, C). SyNOS does not contain a β-hairpin region and tetrahedral zinc-binding

site formed at the mNOS$_{Oxy}$ dimer interface; instead syNOS$_{pGlb}$ occupies this position, although

an insertion between αD/ αE (residues 245-263), that would superimpose on the mNOS zinc-

binding site is not well defined by the electron density (Fig. S9 A, B, C). The syNOS$_{Oxy}$ C-terminal

helical hairpin acts as an interaction hub at the center of the subunit by contacting syNOS$_{Fld}$,

syNOS$_{FNR}$, syNOS$_{Glb}$ and the syNOS$_{C2}$ domain.

3D variable analysis (3DVA) and 3D Flexible refinement (3DFlex) conducted on syNOS-Sym (

PDB:9Q06), revealed that syNOS subunits "breathe" between tight, ordered associations that

bind BH$_4$ within the helical lariats and more separated looser associations with lower density for

these regions (Movie S1), not unlike the "tight" and "loose" dimer states previously observed

for *B. subtilis* NOS (*53–55*). Most structures, except for NOS-TO$_T$, lacked density for the helical

lariats and BH$_4$, which compose the center of the syNOS$_{Oxy}$ dimer interface (Fig. S9). XL-MS

indicated that syNOS in turnover conditions produced three additional interfacial crosslinks

between syNOS$_{pGlb}$ and syNOS$_{Oxy}$ compared to the inactive protien (Fig S10 A – B). Thus, the

active state favors a tighter dimer interface, consistent with BH$_4$ binding. Samples collected in

thicker ice favored this tight association (syNOS-TO$_T$). In addition to an ordered NOS$_{Oxy}$ interface

and density for BH$_4$, NOS-TO$_T$ displayed new contacts between L1 and αA (syNOS$_{pGlb}$) on one

subunit and α9 (syNOS$_{Oxy}$) on the opposing subunit. Hence, L1 may influence syNOS$_{pGlb}$

positioning, which in turn propagates structural stability to the dimer interface (Fig. S9B).

The previously unannotated and uncharacterized N-terminal domain (residues 1-134) takes the

form of a tightly packed 8-stranded β-sandwich, capped by peripheral loops at each end (Fig.

1C, D). This structure strongly resembles a C2 domain (syNOS$_{C2}$), which is usually associated

with Ca^{2+}-dependent membrane targeting in eukaryotes and typically not found in prokaryotes.

(*56, 57*) SyNOS$_{C2}$ brackets the backside of the syNOS$_{Oxy}$ heme pocket and syNOS$_{FNR}$ (residues

1017-1468; Fig. 1C). In NOS-Sym, syNOS$_{FNR}$ latches onto syNOS$_{Glb}$ through the ordered CTT,

which also binds to syNOS$_{Fld}$ (residues 863-1003; Fig. 1B, C).

SyNOS$_{Fld}$ and syNOS$_{FNR}$ comprise a canonical NOS reductase unit in a closed conformation with

the FMN-FAD isoalloxazine cofactors aligned in a planar end-to-end configuration (Fig. 2). To

facilitate ET between subunits, mNOSs rearrange to position the NOS$_{Fld}$ domain < 15 Å from the

NOS$_{Oxy}$ heme on the opposing subunit (*58*). Residue conservation, cross-linking and HDX-MS

indicate that NOS$_{Fld}$ interacts dynamically on the backside of the heme pocket at a conserved

Trp residue (356 in human eNOS) (*36–38, 55*). SyNOS$_{C2}$ blocks this position and would have to

move to allow productive ET from syNOS$_{Fld}$, which also must displace from its shielded

interaction with syNOS$_{FNR}$ (*38*). Notably, syNOS lacks the mNOS CBD (~30 residues) that upon

CAM binding favors the ET competent 'output state' (Fig. 2A-C) (*17, 38, 59, 60*). However, the

syNOS interdomain linkers also likely mediate domain motions. These linkers generally have

extended conformations, with the longest, L1, traversing ~60 Å. The threading of L1 beneath L2

implies that any C2 movement will displace L2 and thereby cause syNOS$_{Fld}$ to move "up-and-

over" toward the adjacent syNOS$_{Oxy}$ subunit (Fig. S8). In comparison to the interdomain

interactions implied by the 25 Å resolution nNOS + CAM structure by negative-stain EM (*37*),

syNOS$_{Fld}$ likely undergoes a different trajectory to interact with syNOS$_{Oxy}$ in part because

syNOS$_{Glb}$ will block the CAM-directed conformational change proposed for nNOS.

SyNOS$_{Red}$ is distantly related, but structurally similar, to cytochrome P-450 reductase (CYPOR)

SyNOS$_{Red}$, mNOS$_{Red}$, sulfite reductase diflavin reductase (SIRFP), and methionine synthase

reductase (MTRR) belong to the cytochrome P450 reductase (CYPOR) family of di-flavin

reductases (Fig. S11) (*61–64*). Di-flavin reductases perform three-separate ET reactions i)

hydride transfer from NADPH/NADH to the FNR FAD cofactor ii) sequential ET from the FNR to

the Fld iii) ET from the Fld to an electron acceptor. Activities (i) and (ii) occur when the di-flavin

reductase is trapped in a "closed" conformation permitting first hydride and then electron

transfer along the sequence NADPH->FAD->FMN(H)• with activity (iii) requiring an "open"

conformation that frees the Fld to reduce the acceptor (e.g. NOS$_{Oxy}$ or cytochrome c).

Analysis by sequence clustering, phylogenetic estimation using maximum likelihood (PhyML)

and multiple structure alignments (MSTA) indicated that syNOS$_{Red}$ and mNOS$_{Red}$ are outgroups

relative to CYPORs (Fig. S11); yet, syNOS$_{Red}$ is more closely related in structure to CYPOR than to

nNOS$_{Red}$ or other di-flavin reductases (Fig. 2 and S11) (*22*). Nevertheless, key differences arise in

the structural and electrostatic features that stabilize the reductase closed conformation in

syNOS (NOS-Sym) compared to CYPOR (Fig. 2A-C). CYPORs and mNOS$_{Red}$ stabilize the closed

conformation by way of an interfacing-region 1 (IR1) between an acidic patch surrounding the

CYPOR$_{Fld}$ FMN cofactor and a complementary basic patch surrounding the CYPOR$_{FNR}$ FAD

cofactor (Fig. 2B,C). An additional IR2 associates the Fld internal helices to the FNR control

region. SyNOS$_{Red}$ lacks electrostatic complementarity within IR1 and residue complementary

within IR2. Instead, regions of the CTT (CTT-IR) and the C2 domain mediate interactions

between the Fld and FNR in the syNOS$_{Red}$ closed conformation (Fig. 3A).

The CTT, shared by syNOS$_{Red}$ and mNOS$_{Red}$, are important for redox flux in NOS but are not

found in canonical CYPORs (Fig. 2A-C) (*22*, *46*). These enzymes also share a β-finger (syNOS

residues 1100-1132) that resides closer to the Fld domain in syNOS$_{Red}$ (~7.5 Å) compared to

mNOS$_{Red}$ (~17 Å), thereby suggesting either a differing structural roles for this element or

revealing a conformational state that nNOS$_{Red}$ can also assume (Fig. 2C). syNOS$_{Red}$ also lacks the

mNOS$_{Red}$ Fld autoinhibitory regulatory loop (AI, nNOS residues 836-849) that destabilizes CAM

binding at low levels of Ca^{2+} (Fig 2B) (*61*, *65*). Although syNOS, nNOS$_{Red}$, and CYPOR stabilize

their closed conformations through different interactions, they all produce a coplanar

orientation and similar separation between the FAD and FMN cofactors (~13 Å from flavin N$_5$-to-

N$_5$).

<u>NOS CTTs are enzyme-specific conformational regulators of ET.</u>

CTTs, thought to be an mNOS addition within the conserved CYPOR family (*66*), are 20-30

residue helical extensions that auto-inhibit reduction of NOS$_{Fld}$ by NOS$_{FNR}$ (Fig. 3). Sequence

alignments show no discernable residue similarity between the syNOS CTT and CTTs of the

three mNOS isoforms or that of the most similar eukaryotic NOS sequence from the diatom *T.*

minus (Fig. S12). The lack of sequence similarity between syNOS$_{CTT}$ and mNOS$_{CTT}$, despite

substantial similarity in the preceding syNOS$_{FNR}$ domains, implies convergent evolution of these

CTTs, likely in concert with the addition of syNOS$_{Glb}$.

Unlike previous NOS$_{Red}$ structures that capture only a portion of the CTT, the syNOS structure

resolves the entire CTT and its interactions with syNOS$_{Fld}$ and syNOS$_{Glb}$ (Fig. 3, S13A). Compared

to other NOSs, the 27-residue syNOS CTT divides into two perpendicular helices (CTT$_1$ and CTT$_2$;

residues 1440-1468; Fig. 3A). CTT$_1$ locks the position of syNOS$_{Fld}$ in the canonical Fld-FNR closed

conformation. CTT$_1$ (residues 1446-1458) interacts with the FMN ligating loops of NOS$_{Fld}$

(residues 984-989 and 951-960). In this interface, key residue contacts likely influence ET-

competent conformational states of syNOS$_{Fld}$ (Fig. 3C-D). CTT$_1$ Lys987 likely modulates Fld

dynamics by ordering residues that coordinate the FMN pyrophosphate groups. Indeed, the

Lys987Ala variant, which disrupts the Lys987/Glu1454 salt-bridge, has a much more extended,

flexible structure by small angle X-Ray scattering (SAXS) than does the WT (Fig. S14). The second

CTT helix (CTT$_2$, residues 1460-1468) is absent in other NOSs and CYPORs, likely because it binds

to the atypical syNOS$_{Glb}$ (Fig. 3A). Indeed, an aliphatic patch on CTT$_2$ interacts with the

hydrophobic crevice between residues 358-392 on syNOS$_{Glb}$ (Fig 3A-B). Specific salt-bridges and

hydrogen bonds among CTT$_2$, syNOS$_{Glb}$ and syNOS$_{Fld}$, help to stabilize the Ca^{2+}-free closed

conformation of syNOS (Fig. 3D,E). In turnover conditions, XL-MS revealed a loss of crosslinks

between the CTT and FNR, which underscores the role of the CTT in mediating dynamics of

syNOS$_{Red}$ (Fig S10C , Table S2).

SyNOS$_{Red}$ has distinct reactivity toward the surrogate electron acceptor cytochrome c (Cc),

compared to mNOS$_{Red}$ and other CYPOR family members. The CTT lowers Cc reduction by NOS$_{Fld}$

in both syNOS and nNOS, but the Ca^{2+} dependence is opposite. In the absence of Ca^{2+}, removal

of the CTT does not affect Cc reduction by syNOS (Fig. S15). Ca^{2+} decreases Cc reduction by

almost 4-fold, but removal of the CTT reactivates Cc reduction (Fig. S15). In mNOS, Ca^{2+}-CAM

increases, rather than decreases Cc reduction (*67*, *68*) and CTT removal increases activity

further. Furthermore, replacing the mNOS$_{FNR}$ domain with CYPOR$_{FNR}$, which lacks a CTT, switches

the Ca^{2+}-CAM dependence of mNOS such that Cc reduction now decreases ~4-fold with Ca^{2+}-

CAM (*67*, *68*), much like for syNOS + Ca^{2+}. These Cc reduction behaviors among NOS proteins

likely reflect differences in the relative strength of interactions among NOS_{Fld}, NOS_{FNR}, the CTT,

NOS_{Oxy} and Cc itself. Interestingly, syNOS structures in the Ca^{2+}-depleted states (NOS-Sym, NOS-

Asym, NOS-Mon) revealed a stable, closed Fld, whereas the Ca^{2+} turnover states ($NOS-TO_{1,2}$) had

a mobile undiscerned Fld domain, seemingly *more* conducive to Cc reduction, not less.

However, in the Ca^{2+}-state, the Fld domain may partition more frequently toward NOS_{Oxy}, which

would compete with Cc and thereby reduce Cc reduction rates. In mNOS, if the Fld-to-Oxy

interaction is weaker than in syNOS. relieving a stronger CTT-to-Fld interaction with Ca^{2+}-CAM

may favor the Fld-to-Cc interaction.

<u>The CTT-stablizied closed conformation of syNOS$_{Red}$ limits flavin reduction.</u>

SyNOS$_{FL}$ and syNOS$_{Red}$ rates of flavin reduction by NADPH, measured through anaerobic

stopped-flow experiments, were biphasic and resembled the flavin reduction kinetics of CYPOR

and nNOS$_{Red}$ bound to Ca^{2+}-CAM (Fig. S16A) (*69*). However, NADPH reduces syNOS$_{FL}$ at a ~74-

fold slower rate than syNOS$_{Red}$ implying that the closed conformation of syNOS$_{FL}$ restricts

hydride transfer from NADPH to FAD (Fig. S16B-D).   The lower FAD reduction rate in syNOS$_{FL}$ is

surprising because syNOS$_{Red}$ lacks an NADP(H)-ligating Arg (1400 in rat nNOS) known to inhibit

electron flow in CAM-free nNOS/eNOS (*70*). In nNOS, the Arg1400Ser substitution increases

flavin reduction while decreasing NO synthesis (*70*). Because syNOS contains an Asn residue at

the 1400 position it would be predicted to have a faster flavin reduction rate than nNOS$_{Red}$

(*70*). However, the syNOS$_{Glb}$-to-CTT interaction, unique to syNOS, may especially stabilize

conserved CTT residue Trp1440 (*71*), which blocks the nicotinamide group of NADPH from

approaching the FAD isoalloxazine ring for hydride transfer (Fig. S16B). The CTT enforced rigidity

in syNOS (also reflected in the XL-MS data, Fig. S10C) may compensate for the modified $NADP^+$-

binding pocket that lacks an autoinhibitory Arg1400 equivalent. Notably, compared to nNOS and

CYPOR the nicotinamide moiety in syNOS-Mon adopts an atypical conformation that is

intermediate to the reduction incapable NADP$^+$ conformation seen in nNOS (PDB:1TLL) and the

NADP$^+$/FAD stacked conformation seen in the CYPOR W677X variants (PDB:1AJO, Fig. S16D).

<u>Ca^{2+} regulates syNOS activity through a C2-like domain</u>

The lack of a CAM-CBD interaction in syNOS renders its Ca^{2+}sensitivity enigmatic. However,

syNOS$_{C2}$ structurally resembles C2 domains that mediate membrane recruitment by binding

Ca^{2+} (Fig. 4) particularly; mouse perforin C2 (Fig. 4A) (*56*, *57*, *72*).

In the closed conformation, syNOS$_{C2}$ forms a tripartite interface with the syNOS$_{Oxy}$ domain and

the FNR control region (Fig. 4B). Two regions of salt-bridges mediate the C2-Oxy interface (Fig.

4B). In particular, the interaction between syNOS$_{C2}$ Arg107 and NOS$_{Oxy}$ Glu568, positions the C2

domain beside Trp715, the proposed site of mNOS$_{Fld}$ binding onto mNOS$_{Oxy}$ (*38*). A basic patch

on the C2-domain and a complementary acidic patch on α-helix 1162-1173 of the FNR-control

domain wedges the C2 between syNOS$_{Oxy}$ and syNOS$_{Fld}$ (Fig. 4C). Because syNOS$_{C2}$ blocks the

putative Fld-Oxy contact; displacement of syNOS$_{C2}$ would decrease conformational restraints on

syNOS$_{Red}$ (particularly the L2 linker) and relieve steric hindrance on the backside of the heme

pocket.

A variant devoid of the C2 domain (syNOS$_{\Delta C2}$, residues 179-1468) has no NOS activity but retains

Cc-reduction activity (Fig. S14). Strikingly, syNOS$_{\Delta C2}$ has a high WT-like Cc-reduction rate that is

uninhibited by Ca^{2+}. In keeping with an enzyme unconstrained by inhibitory interactions. SAXS

data indicates that the global flexibility of syNOS$_{\Delta C2}$ exceeds that of the WT and the more

flexible K987A variant (Fig. S14). Thus, the C2 domain serves to stabilize NOS$_{Oxy}$ and influences

redox domain mobility of syNOS in response to Ca^{2+}, akin to the functional role that Ca^{2+}-CAM

plays in the regulation of mNOSs. In other systems, Ca^{2+} binding to C2 domains neutralizes

acidic regions to allow association with negatively charged phospholipid head-groups (*73*).

Although, syNOS$_{C2}$ lacks the typically conserved Asp residues that coordinate Ca^{2+}, isothermal

calorimetry (Fig. S17) indicated that the syNOS$_{C2}$ binds Ca^{2+} with a 1.6 μM Kd, which compares

to the Ca^{2+} affinity of PLDβ C2 domains (0.8 μM) (*74*).

Various stimuli produce surges in cyanobacterial intracellular Ca concentrations from 100-200

nm to the μM range (*75, 76*). In vitro, syNOS activity responds to Ca^{2+} with a K_M of ~ 200 μM

(*34*). This larger value may reflect the conformational coupling of the C2 domain movement to

Ca^{2+} binding and may be affected by the cellular context. Interestingly, some types of

cyanobacteria uptake Ca^{2+} at very high concentrations to mineralize $CaCO_3$ (*77*).

<u>Ca^{2+} causes a large movement of the C2 domain</u>

In the presence of Ca^{2+} and L-arginine, syNOS adopts a more flexible state, wherein most of

$syNOS_{Red}$ is undiscerned by cryoEM (NOS-TO$_{L,T}$). In this turnover state, the C2 domain has moved

away from the Oxy/FNR interface and now binds at the entrance of the $syNOS_{Glb}$ heme pocket,

nearly 85 Å from its original position (Fig. 5A). The new C2 position blocks the $syNOS_{Glb}$ heme-

binding center and substantially shifts the αB-D helices and constituent residues that compose

the distal heme pocket where NO is oxidized (Fig. 5B). The highly conserved C2 basic patch

(Lys35, Lys37 and Arg81) that formally bound $syNOS_{FNR}$ in the locked dimer interacts with the

heme-propionic acid groups of $syNOS_{Glb}$ and the 81-89 loop, also conserved among $syNOS_{C2}$

domains, reaches over to block the distal heme pocket. Furthermore, the movement of $syNOS_{C2}$

displaces the L2 linker, which swivels by ~135° and now directs $syNOS_{Fld}$ toward the backside of

the heme pocket on the opposing subunit (Fig. 5C), consistent with the trans ET mechanism

characteristic of mNOSs (*58*). As $syNOS_{C2}$ now resides where the electron-donating FNR

modules of Hmps interact with their cognate globin domains, we tested whether removal of the

C2 domain ($syNOS_{ΔC2}$) affected $syNOS_{Glb}$ reduction in $syNOS_{FL}$. Surprisingly, we observed little

change in $syNOS_{Glb}$ reduction rates in $syNOS_{ΔC2,}$ nor did Ca^{2+} affect $syNOS_{Glb}$ reduction rates in

$syNOS_{FL}$ (Fig. S18).

<u>$syNOS_{Glb}$ reduction requires an altered configuration of $syNOS_{FNR}$</u>

Flavohemoglobins reductively activate dioxygen with an FNR domain in order to oxidize NO to

NO_3^-; however, in principle, either $syNOS_{Fld}$ or $syNOS_{FNR}$ could reduce $syNOS_{Glb}$. In the locked

state, both the syNOS$_{Fld}$ FMN and the syNOS$_{FNR}$ FAD are too far from the syNOS$_{Glb}$ Fe^{3+} to yield

reasonably fast ET rates (27 Å and 31 Å, respectively). When considering that syNOS$_{Glb}$ has a

lower potential (E$^{0'}$ = -250 mV) than syNOS$_{Fld}$ (E$_m$ = -170 mV), but not syNOS$_{FNR}$ (E$_m$ = -290 mV),

syNOS$_{FNR}$ is likely the better electron donor. However, the relatively low expected driving force

for reduction between syNOS$_{FNR}$ and syNOS$_{Glb}$ (Table S5) and a distance of separation of ~31 Å,

suggests that syNOS$_{FNR}$ displaces from its position in NOS-Sym to reduce syNOS$_{Glb}$ at the

observed rate (Fig. 6; (78)). Indeed, the isolated syNOS$_{FNR}$ domain reduces the syNOS$_{Glb}$ domain

with saturation behavior (K$_m$ = 11.5 μM) at nearly the same rate (k$_{cat}$ = 0.25 s^{-1}) as syNOS$_{Glb}$ is

reduced in full-length Ca^{2+}-free syNOS (k$_{obs}$ = 0.22 s^{-1}, Fig. S18). Furthermore, if syNOS$_{Fld}$ in the

locked confirmation can directly reduce syNOS$_{Glb}$, we would expect the conformationally

destabilized syNOS$_{\Delta C2}$ to show slower syNOS$_{Glb}$ reduction, but this is not the case (Fig. S16). A

modest decrease in syNOS NOD activity with Ca^{2+} suggests an inhibitory effect of syNOS$_{C2}$ (Fig.

S18), which would be expected from its relocation to the site where FNR domains typically bind

globin domains in Hmps.

<u>The asymmetric state may act as a conformational intermediate between NOS and NOD</u>

<u>activities</u>

The mNOS input, intermediate, and output states are not discrete, but sampled within a

continuum of motion (18, 35). In the output state, Fld-CAM moves 70 Å to dock onto an

opposing NOS$_{Oxy}$ domain at the backside of the heme pocket. Similarly, syNOS-Asym (

PDB:9Q0Y,9Q15) represents an intermediate structural state between the locked dimer and the

turnover state that is likely in dynamic equilibrium with NOS-Sym and NOS-TO.   An Alphafold3

prediction of the Oxy dimer unit with one Fld domain places syNOS$_{Fld}$ at the experimentally

validated position on syNOS$_{Oxy}$ found by HDX-MS and crosslinking studies of mNOS (38). Upon

superimposing this prediction onto either NOS-Asym or NOS-Sym, the Fld binding surface on

syNOS$_{Oxy}$ is blocked by the syNOS$_{C2}$ (Fig. 5C and S19). Furthermore, compared to Hmp,

syNOS$_{FNR}$ would have to rotate 90^o and move 28 Å to reduce syNOS$_{Glb}$ (Fig. 6). This configuration

is likely accessible in the dynamic asymmetric subunit but would be prevented in syNOS$_{TO}$

(PDB:9Q0X) by the rearranged syNOS$_{C2}$. Hence, it is likely that syNOS-Asym is capable of

syNOS$_{Glb}$ reduction but not syNOS$_{Oxy}$ reduction.

The intermediate conformation of syNOS-Asym (PDB:9Q0Y,9Q15) may be important for the

switch between NOS and NOD activities and explain the inhibitory effect of Ca^{2+} on Cc-

reduction. In the absence of Ca^{2+}, syNOS-Sym locks down and inhibits NOS activity, but NOS-

Asym allows syNOS$_{FNR}$ to reduce syNOS$_{Glb}$, and syNOS$_{Fld}$ to reduce Cc. In syNOS-Asym

(PDB:9Q0Y,9Q15), syNOS$_{Fld}$ cannot reduce syNOS$_{Oxy}$ because syNOS$_{C2}$ blocks syNOS$_{Fld}$ from the

syNOS$_{Oxy}$ interaction site on the opposing subunit. Ca^{2+} promotes the relocation of the C2

domain from syNOS$_{Oxy}$ to syNOS$_{Glb}$; however, some C2 displacement from NOS$_{Oxy}$ may occur in

the absence of Ca^{2+} because the C2 domain is not well defined in the more mobile subunit of

syNOS-Asym. In addition to preventing syNOS$_{FNR}$ from reducing the syNOS$_{Glb}$, C2 movement

disrupts L2, destabilizing the interaction of syNOS$_{Fld}$ with syNOS$_{FNR}$ and favors partitioning of

syNOS$_{Fld}$ toward the unblocked syNOS$_{Oxy}$ and away from Cc (Fig. 7).

Similar to mNOS, syNOS assumes a continuum of conformations that circumscribe defined

enzymatic states. Hence to generate changes in physiologically relevant activities, the relative

weighting of states within the conformational ensemble must be regulated and responsive to

cellular context. It is striking that the syNOS$_{C2}$ moves between a location that blocks syNOS$_{Oxy}$

reduction to one that blocks syNOS$_{Glb}$ reduction, thereby suggesting counter regulation of NO

synthesis and NO oxidation. Such a mechanism may allow the enzyme to tune the balance of

NO production versus NO detoxification, which would be of benefit for signal transduction,

nitrogen metabolism and cellular homeostasis. Overall, these syNOS structural studies

demonstrate how a multidomain reductase and its associated regulatory components can

undergo large scale rearrangements to modulate reactivity in response to environmental

conditions.

Methods

Molecular cloning

The native coding sequence for full-length syNOS (Uniprot ID : B4WU43**)** was PCR-amplified and

cloned into a modified pet28-a construct between restriction sites NdeI and XhoI using

NEBuilder HiFi DNA Assembly master mix (E2621L). The construct contained sequences for an

N-terminal Twin-Strep Tag with a TEV cleavage site. To produce the hptG and gapA construct,

htpG and gapA were amplified from K-12 *E. coli* and placed into the MCS-1 and the MCS2 of a

tagless pAYC-Duet vector using NEBuilder HiFi DNA Assembly master mix (E2621L). Individual

point mutants were constructed using Kinase-Ligase-DpnI (KLD) method using primers in Table

S4 (New England Biosciences). All plasmid molecular cloning and plasmid material generation

were carried out with NEB® 5-alpha competent *E. coli* (New England Biosciences). Successful

transformants were validated through Sanger Sequencing or Nanopore sequencing (Eurofins

Genomics).

Protein expression and purification

SyNOS and/or variants were expressed in BL21(DE3) as previously described (*34*) with the

modification of co-expression with hptG/gapA, which improved yield and reproducibility. Cell

pellets were stored at -80 °C prior to purification. The pellet was thawed and resuspended in

lysis buffer (50 mM HEPES (pH 7.5), 150 mM NaCl, 1 mM TCEP, 10% glycerol and 10 uM BH_4),

followed by high-speed centrifugation at 100 000 G and application on Strep-Tactin XT 4-Flow

resin (IBA Life Sciences). Elution of the affinity-bound protein fraction was carried out in batch

with 50 mM biotin supplemented lysis buffer. Post elution samples were concentrated using

Viva-Spin columns (100 kDa cut -off), before application onto 10/300 Superose 6 Increase

Column (Cytiva) using Gel Filtration Buffer 1 (GFB1, 25 mM HEPES (pH 7.5), 150 mM NaCl, 1 mM

TCEP, 10% glycerol and 10 uM BH_4).

Size exclusion chromotography - Small angle X-ray scattering (SEC-SAXS)

All SEC-SAXS was performed at ID7A1 beamline at Cornell High Energy Synchrotron Source

(CHESS). Photon flux was 2.3 x 10^{12} ph/s and scattering data was collected 0.005 Å^{-1} ≤ q ≤ 0.7 Å^{-1}

on an EIGER 4M detector (Dectris). Samples were injected into an inline 3.2/150 Superdex 200

column in gel filtration buffer. The entire SEC-SAXS trace was captured using a 2s per exposure

throughout the elution. The flow-cell format of the experiment should limit x-ray

photoreduction of the sample, but it is possible that the NOS cofactors undergo some

photoreduction during the experiment (*79*). All SEC-SAXS data processing was performed

through RAW (*80*).

Single turnover anaerobic globin reduction

Anerobic syNOS$_{Glb}$ heme reduction assays were carried out using an Agilent 8453 UV-Vis

spectrometer with an attached Hewlett Packard 89090A Peltier temperature controller for

maintaining constant temperature at 20 ^{o}C and constant stirring at 400 rpm (Agilent & Hewlett

Packard). Prior to reaction initiation with 1 mM NADPH, syNOS or variants (1 μM) were

incubated anaerobically in the presence of the Glucose Oxidase/Catalase (GOD/CAT) oxygen

scavenging system, as previously described (*81*). Samples were run in either in the presence of 1

mM Ca^{2+} or 1 mM EGTA, as noted. In the case of syNOS$_{FNR}$ (1027-1468)/syNOS$_{Glb}$ (337-469)

reduction trials, syNOS$_{Glb}$ was fixed at 2.2 μM and syNOS$_{FNR}$ was varied between 1 μM to 21 μM

with no EGTA/Ca^{2+} present. All data was processed using home-made scripts in R Studio (Posit

PBC) with rates of syNOS$_{Glb}$ reduction with to a single exponential to determine k$_{obs}$. For the

truncated domains k$_{cat}$ and K$_m$ were calculated by plotting k$_{obs}$ for each syNOS$_{FNR}$/syNOS$_{Glb}$ pair

against syNOS$_{FNR}$ concentration using Prisim (GraphPad).

Stopped-flow pre-steady state flavin reduction kinetics

Anerobic stopped-flow mixing of NADPH with syNOS$_{Red}$ or syNOS$_{Fl}$ in the absence of Ca^{2+} was

used to characterize flavin reduction rates. After purification, syNOS$_{Red}$ contains oxidized FAD

and FMN. Attempts to pre-reduce the enzyme to FMNH• with sodium dithionate, DTT and

ascorbate caused precipitation without an observable spectral shift to FMNH•. Thus, flavin

reduction rates reflect conversion from fully oxidized syNOS$_{Red}$. All stopped flow experiments

were carried out anaerobically using a Stopped Flow SX20 instrument (Applied Photophysics).

Samples were pre-incubated anaerobically in the presence of the GOD/CAT oxygen scavenging

system (see above). Changes in absorbance were detected using a single wavelength

Photomultiplier Tube detector. syNOS$_{FL}$ (10 μM) / syNOS$_{Red}$ (40 μM) were mixed with 4x excess

NADPH. All data was processed using home-made scripts in R Studio (Posit PBC).

CryoEM grid preparation and data collection - syNOS ligand free (Dataset 1 and 2)

A 4 µl volume of 1.0 mg ml^{-1} syNOS solution, in GFB1, was applied to glow-discharged grids (200

mesh Quantifoil Cu, R1.2/1.3; Electron Microscopy Sciences). After 10 s of incubation, the

excess solution was blotted for 3.5 s with filter paper by a Vitrobot Mark IV and plunged into

liquid ethane, for vitrification (ThermoFisher). CryoEM images of syNOS were collected on a

Talos Arctica (ThermoFisher) operated at 200 keV at a nominal magnification of 63,000× with a

Gatan GIF Quantum LS Imaging energy filter, using a Gatan K3 direct electron camera in super-

resolution counting mode, corresponding to a pixel size of 1.31 Å. A total of 1214 images stacks

were obtained with a defocus range of −0.8 to −2.0 µm by Serial EM (University of Colorado-

Boulder). Each stack video was recorded for a total dose of 50 electrons per Å^2.

CryoEM grid preparation and data collection - Ca^{2+} and L-arginine conditions (Dataset 3)

A 4 µl volume of 1.0 mg ml^{-1} syNOS in the presence of 1 mM Ca^{2+} and 1 mM L-arg solution was

applied to glow-discharged grids (300 mesh Quantifoil Au, R1.2/1.3). After 10s of incubation, the

excess solution was blotted for 2.5 s with filter paper by a Vitrobot Mark IV and plunged into

liquid ethane for vitrification (ThermoFisher). CryoEM images of syNOS were collected on a

Talos Krios (ThermoFisher) operated at 300 keV at a nominal magnification of 165,000× with a

Gatan GIF Quantum LS Imaging energy filter, using a Gatan K3 direct electron camera in super-

resolution counting mode, corresponding to a pixel size of 0.422 Å. A total of 5019 images stacks

were obtained with a defocus range of −0.6 to −2.0 µm by Leginon (*82*). Each stack video was

recorded for a total dose of 50.5 electrons per Å^2.

CryoEM grid preparation and data collection - turnover conditions (Dataset 5)

A 4 µl volume of 1.0 mg ml^{-1} syNOS-ligand free solution in GFB1 was applied to glow-discharged

grids (200 mesh Quantifoil Cu, R1.2/1.3). After 10s of incubation, the excess solution was

blotted for 3.5 s with filter paper by a Vitrobot Mark IV and plunged into liquid ethane for

vitrification (ThermoFisher). CryoEM images of syNOS were collected on a Talos Krios

(ThermoFisher) operated at 300 keV at a nominal magnification of × 165,000 with a Gatan GIF

Quantum LS Imaging energy filter, using a Gatan K3 direct electron camera in super-resolution

counting mode, corresponding to a pixel size of 0.730 Å. A total of 5001 images stacks were

obtained with a defocus rage of −0.6 to −2.0 μm by Leginon (*82*). Each stack video was recorded

for a total dose of 44.9 electrons per Å^2.

CryoEM Image processing and structural reconstruction

All cryoEM density refinement was performed using CryoSPARC (*50*). In brief, micrograpghs

were motion corrected using patch motion followed by CTF estimation using patch CTF. Particles

were picked using CRYOLO and run through 2D classification and initial density reconstructions.

Candidate densities were then selected for further refinement, including junk classes to

eliminate contaminating/damaged particles. For dataset dependant details please refer to Figs

.S2-S6. as illustrated by the procedures illustrated in Figs. S2-S6.

For syNOS-Asym (DS4, EMD-72110), non-uniform refinement obscured the smaller subunit. To

improve the density quality we locally refined each subunit. ChimeraX was used to create the

two masks, centered at each subunit, for the local refinement(Fig. S5). Gaussian noise with a

standard deviation of 3 was applied to EMD 72110, producing a new density that obscured

secondary structure features. The Segger tool (*83*) was used to segment the map into each

subunit. composite map (EMD-72116) produced from DS4, The electron density was resampled

into the appropriate box size and imported into CryoSPARC where masks were created using the

3D Volume tool and subsequently used for local refinement. The refined maps were aligned and

merged using the vop maximum command in ChimeraX to produce the composite map.

An Alphafold3 prediction of a syNOS homodimer (residues 1-1468), was used as the input

model and docked into the appropriate electron density using ChimeraX (*50*, *84*, *85*). Ligand and

poor fitting regions were built and real-space refined using with Coot (*85*). The resulting models

were passed through ISOLDE (*86*) to approach thermodynamically rational

rotamers/conformations. The outputs from these initial models were subjected to a series of

real-space refinement with PHENIX (*87*) and conformational outlier residues were then adjusted

manually using Coot (*85*).

Cytochrome c reduction assay

Cytochrome c (Cc) was used as an electron acceptor to measure the efficiency of electron flow

from NOS_{Red}. Cc accepts an electron from the fully reduced $FMNH_2$ and not from $FADH_2$/FADH⁻

•/FAD•/ FMNH•. Cc reduction assays were carried out as previously described (*88*) using an

Agilent 8453 UV-Vis Spectrometer with an attached Hewlett Packard 89090A Peltier

Temperature Controller for constant temperature at 20 °C and constant stirring at 400 rpm

(Agilent & Hewlett Packard). All reported reactions were carried out in CC buffer (100 mM Tris

(pH 7.5) and 200 mM NaCl). Cc was pre-oxidized with 1 mM potassium ferrocyanide and then

buffer exchanged into CC buffer. Reactions were conducted with 10 uM Cc and 3 nM of the

sample of interest. Reactions were initiated with 1 mM NADPH and were run either in the

presence of 1 mM Ca^{2+} or 1 mM EGTA as noted. All data was processed using home-made

scripts in R Studio (Posit PBC).

Nitric oxide synthase activity assay

All activity assays were carried out for 30 minutes in 100 mM Tris, (pH 7.5) 200 mM NaCl, 1 mM

DTT and 1 mM L-arginine and initiated with 1 mM NADPH (25 °C). Each sample was evaluated in

the presence of 1 mM Ca^{2+} or 1 mM EGTA. Post reaction, samples were boiled at 70 °C to halt

NOS activity. NADPH was then precipitated using the $ZnCl_2$/$NaCO_3$ method and subsequently

centrifuged at 15000 G at room temperature. The supernatant was then quantified for total

NO_2^- and NO_3^- content through the Griess assay as implemented in the Nitrate/Nitrite

Colorimetric Assay Kit (Cayman Chemicals).

Oxygen consumption assay to probe Nitric Oxide Dioxygenase activity

The Oxygraph+ system (Hansatech Instruments) was used to measure oxygen consumption of

syNOS in the presence of the NO donor, 3-(2-Hydroxy-1-methyl-2-nitrosohydrazino)-N-methyl-1-

propanamine (NOC-7). 1 μM of enzyme ($syNOS_{Fl}$ or $syNOS_{\Delta C2}$) was incubated in buffer

containing 50 mM Tris and 150 mM NaCl at pH 7.5, with 10 μM NOC-7 for 5 minutes. The

reaction was initiated with 1 mM NADPH. Each sample was evaluated in the presence of 1 mM

Ca^{2+} or 1 mM EGTA . Three separate controls were performed to rule out superoxide or non-

enzymatic NO oxidation: NOC-7 + NADPH, NOC-7 + $syNOS_{FL}$ and syNOS + NADPH.

Isothermal calorimetry

Thermodynamic binding properties between Ca^{2+} and $syNOS_{C2}$ (1-138) were measured on a

Nano ITC Low Volume isothermal titration calorimeter (TA instruments). $SyNOS_{C2}$ was prepared

in a 50 mM CHES and 150 mM NaCl buffer at pH 9.0 at 0.072 mM with 200 µL in the sample cell.

Lower pHs also gave indication of Ca^{2+} binding, but led to aggregation of the C2 domain that

prevented quantitative assessment. $CaCl_2$ was prepared in the same buffer at a concentration of

0.250 mM; 2.5uL was titrated every 250 s for a total of 30 injections. Thermal data were

analyzed using an independent binding model with the NanoAnalyze software.

Reduction potential determination

Measurements were performed using the isolated domains $syNOS_{Glb}$ (337-469), $syNOS_{FMN}$ (856-

1027) and $syNOS_{FNR}$ (1027-1468), which were expressed and purified separately. Reduction

potential measurements were performed using a dye calibration method (*89, 90*), wherein

reduction was driven by a deazaflavin/EDTA photochemical system in a 50 mM $NaPO_3$, 10%

Glycerol at pH 7.0, with the following modifications: i) 9 µm 5-deazariobflavin and 5 mM EDTA

was used as the chemical reductant ii) reactions were equilibrated anaerobically in a coy

chamber while supplemented with GOD/CAT oxygen scavenging system. All measurements

were carried out using an Agilent 8453 UV-Vis spectrometer with an attached Hewlett Packard

89090A Peltier Temperature Controller for maintaining constant temperature at 20 ^{o}C and

constant stirring at 400 rpm (Agilent & Hewlett Packard), with the reaction being photoinitiated

with a 15 mW 300 nm laser (ThorLabs) placed above the clear septum capping the cuvette.

Calibrating dyes per domain were selected based on literature values of the published redox

potentials (Table S5). Note that the quoted reduction potentials for $syNOS_{Fld}$ and $syNOS_{FNR}$ will

be the average of the single electron couples between semiquinone and oxidized, and

hydroquinone and semiquinone.

DSBU crosslinking, in-gel protein digestion and extraction

Purified syNOS that correspond to the dimer fraction from SEC was split into two conditions: 1)

inactive - GFB1, 2) turnover - GFB1 + 1mM Ca^{2+} + 1 mM NADPH + 100 µM BH_4. Both conditions

were then incubated with 1mM (100 molar excess) disuccinimidyl dibutyric urea (DSBU,

ThermoFisher) for 30 minutes at room temperature prior to quenching with 20 mM Tris pH 8.0.

Samples were then centrifuged and run on an SDS-PAGE gel using lameli sample buffer.

Bands corresponding to dimers were excised, digested and extracted as previously described

(*91*), apart from the proteases of choice being Lys-C/Trypsin, prior to LC- MS/MS analysis.

Crosslinks identification by nanoLC-MS/MS

The digested product was characterized by NanoLC-MS/MS analysis at the Cornell

Biotechnology Resource Center. The analysis was carried out using an Orbitrap FusionTM

TribridTM (Thermo-Fisher Scientific, San Jose, CA) mass spectrometer equipped with a

nanospray Flex Ion Source, and coupled with a Dionex UltiMate 3000 RSLCnano system

(Thermo, Sunnyvale, CA). Each sample was loaded onto a nano Viper PepMap C18 trapping

column (5 μm, 100 μm ´ 20 mm, 100 Å, Thermo Fisher Scientific) at 20 ⬚L/min flow rate for rapid

sample loading. After 3 minutes, the valve switched to allow peptides to be separated on an

Acclaim PepMap C18 nano column (2 μm, 75 μm x 25 cm, Thermo Fisher Scientific) at 35 °C in a

93 min gradient of 5% to 40% then 93-98 min of 40%-60% buffer B (95% ACN with 0.1% formic

acid) at 300 nL/min. The Orbitrap Fusion was operated in positive ion mode with nano spray

voltage set at 1.5 kV and source temperature at 275 °C. External calibrations for Fourier

transform, ion-trap and quadrupole mass analyzers were performed prior to the analysis.

Samples were analyzed using the HCD-MS2 workflow, in which MS scan range was set to 300-

1800 m/z and the resolution was set to 60,000. Precursor ions with charge states 2-8 were

selected for HCD MS2 acquisitions in Orbitrap analyzer with a resolution of 30,000 with collision

energy of 25% and 30%, normalized AGC of 200%. The precursor isolation width was 1.6 m/z

and the maximum injection time was 70 ms. All data were acquired under Xcalibur 4.4

operation software (Thermo-Fisher Scientific).

Crosslink identification and visualization

Raw spectra were searched using Proteome Discoverer 2.4 (Thermo-Fisher Scientific, San Jose,

CA) with XlinkX v2.0 algorithm for identification of crosslinked peptides. The search parameters

were as follow: four missed cleavages for either double digestion or triple digestion with fixed

carbamidomethyl modification of cysteine, variable modifications of methionine oxidation,

asparagine/glutamine deamidation, protein N-terminal acetylation, serine/threonine/tyrosine

phosphorylation and lysine di-glycine tag . Only lysine-to-lysine crosslinks were considered. The

peptide mass tolerance was 10 ppm, and MS2 fragment mass tolerance was 0.6 Da. A custom

database (246 sequences) containing syNOS, common *E.coli* chaperones and common

contaminates was used for the search, with > 1% FDR for report of crosslink results. Identified

crosslinked peptides were filtered for Max. XlinkX Score > 40, a confidence score that reflects

how well peptide spectrum matching, peptide fragmentation and crosslink doublet

identification. Spectra were manually analyzed to validate potential crosslinks. The crosslinked

peptides were analyzed using XMAS (*93*).

References

1.   R. M. Palmer, S. Moncada, A novel citrulline-forming enzyme implicated in the formation of
nitric oxide by vascular endothelial cells. *Biochem Biophys Res Commun* **158**, 348–352
(1989).

2.   B. R. Crane, Bacterial Nitric Oxide Synthases. *Annual Review Of Biochemistry* **79** (2010).

3.   J. A. McCleverty, Chemistry of Nitric Oxide Relevant to Biology. *Chem. Rev.* **104**, 403–418
(2004).

4.   M. Cossenza, R. Socodato, C. C. Portugal, I. C. L. Domith, L. F. H. Gladulich, T. G. Encarnaçao,
K. C. Calaza, H. R. Mendonça, P. Campello-Costa, R. Paes-de-Carvalho, "Nitric Oxide in the
Nervous System: Biochemical, Developmental, and Neurobiological Aspects" in *Nitric*
Oxide, G. Litwack, Ed. (2014)vol. 96 of *Vitamins and Hormones*, pp. 79–125.

5.   C. Farah, L. Y. M. Michel, J. L. Balligand, Nitric oxide signalling in cardiovascular health and
disease. *Nature Reviews Cardiology* **15**, 292–316 (2018).

6.   J. O. Lundberg, E. Weitzberg, Nitric oxide signaling in health and disease. *Cell* **185**, 2853–
2878 (2022).

7.   M. A. Marletta, Revisiting Nitric Oxide Signaling: Where Was It, and Where Is It Going?
Biochemistry **60**, 3491–3496 (2021).

8.   M. Shepherd, D. Giordano, C. Verde, R. K. Poole, The Evolution of Nitric Oxide Function:
From Reactivity in the Prebiotic Earth to Examples of Biological Roles and Therapeutic
Applications. *Antioxidants* **11** (2022).

9. K. A. White, M. A. Marletta, Nitric oxide synthase is a cytochrome P-450 type hemoprotein,
ACS Publications (2002). https://doi.org/10.1021/bi00144a001.

10. U. Förstermann, W. C. Sessa, Nitric oxide synthases: regulation and function. *Eur Heart J*
33, 829–837, 837a–837d (2012).

11. M. A. Cinelli, H. T. Do, G. P. Miley, R. B. Silverman, Inducible nitric oxide synthase:
Regulation, structure, and inhibition. *Med Res Rev* **40**, 158–189 (2020).

12. D. J. Stuehr, M. M. Haque, Nitric oxide synthase enzymology in the 20 years after the Nobel
Prize. *British Journal of Pharmacology* **176**, 177–188 (2019).

13. J. Reyes-Rivera, Y. Wu, B. G. H. Guthrie, M. A. Marletta, N. King, T. Brunet, Nitric oxide
signaling controls collective contractions in a colonial choanoflagellate. *Curr Biol* **32**, 2539-
2547.e5 (2022).

14. J. Tejero, D. Stuehr, Tetrahydrobiopterin in nitric oxide synthase. *IUBMB Life* **65**, 358–365
(2013).

15. D. J. Stuehr, J. Santolini, Z.-Q. Wang, C.-C. Wei, S. Adak, Update on mechanism and catalytic
regulation in the NO synthases. *J Biol Chem* **279**, 36167–36170 (2004).

16. H. M. Abu-Soud, D. J. Stuehr, Nitric oxide synthases reveal a role for calmodulin in
controlling electron transfer. *Proc Natl Acad Sci U S A* **90**, 10769–10772 (1993).

17. D. H. Craig, S. K. Chapman, S. Daff, Calmodulin Activates Electron Transfer through
Neuronal Nitric-oxide Synthase Reductase Domain by Releasing an NADPH-dependent
Conformational Lock *. *Journal of Biological Chemistry* **277**, 33987–33994 (2002).

18. N. G. H. Leferink, S. Hay, S. E. J. Rigby, N. S. Scrutton, Towards the free energy landscape for
catalysis in mammalian nitric oxide synthases. *The FEBS Journal* **282**, 3016–3029 (2015).

19. T. L. Poulos, H. Li, Nitric Oxide Synthase and Structure-Based Inhibitor Design. *Nitric Oxide*
63, 68–77 (2017).

20. L. Zhou, D.-Y. Zhu, Neuronal nitric oxide synthase: Structure, subcellular localization,
regulation, and clinical implications. *Nitric Oxide* **20**, 223–230 (2009).

21. V. Garcia, W. C. Sessa, Endothelial NOS: perspective and recent developments. *Br J*
Pharmacol **176**, 189–196 (2019).

22. N. A. Popova, S. K. Soodaeva, I. A. Klimanov, V. M. Misharin, A. A. Temnov, Autoregulation
and Autoinhibition of the Main NO Synthase Isoforms (Brief Review). *Sovremennye*
Tehnologii V Medicine **15**, 53–59 (2023).

23.   M. C. Surdel, B. F. Dutter, G. A. Sulikowski, E. P. Skaar, Bacterial Nitric Oxide Synthase Is
Required for the Staphylococcus aureus Response to Heme Stress. *ACS Infect. Dis.* **2**, 572–
578 (2016).

24.   I. Gusarov, K. Shatalin, M. Starodubtseva, E. Nudler, Endogenous Nitric Oxide Protects
Bacteria Against a Wide Spectrum of Antibiotics. *Science* **325**, 1380–1384 (2009).

25.   B. A. Patel, M. Moreau, J. Widom, H. Chen, L. Yin, Y. Hua, B. R. Crane, Endogenous nitric
oxide regulates the recovery of the radiation-resistant bacterium Deinococcus radiodurans
from exposure to UV light. *Proc Natl Acad Sci U S A* **106**, 18183–18188 (2009).

26.   N. Gupta, A. Srivastava, A. K. Mishra, Nitric Oxide Synthases in Cyanobacteria: Diversity,
Cellular Implications and Ecological Pertinence. *J Plant Growth Regul* **44**, 508–528 (2025).

27.   T. G. Popova, A. Teunis, H. Vaseghi, W. Zhou, V. Espina, L. A. Liotta, S. G. Popov, Nitric oxide
as a regulator of *B-anthracis* pathogenicity. *Frontiers in Microbiology* **6** (2015).

28.   P. R. Gardner, Nitric oxide dioxygenase function and mechanism of flavohemoglobin,
hemoglobin, myoglobin and their associated reductases. *J Inorg Biochem* **99**, 247–266
(2005).

29.   P. R. Gardner, A. M. Gardner, L. A. Martin, A. L. Salzman, Nitric oxide dioxygenase: An
enzymic function for flavohemoglobin. *Proceedings of the National Academy of Sciences*
95, 10378–10383 (1998).

30.   M. E. S. Lewis, H. A. Corker, B. Gollan, R. K. Poole, "A survey of methods for the purification
of microbial flavohemoglobins" in *Globins and Other Nitric Oxide-Reactive Proteins, Pt A*, R.
K. Poole, Ed. (2008)vol. 436 of *Methods in Enzymology*, pp. 169–186.

31.   F. Del Castello, N. Foresi, A. Nejamkin, C. Lindermayr, F. Buegger, L. Lamattina, N. Correa-
Aragunde, Cyanobacterial NOS expression improves nitrogen use efficiency, nitrogen-
deficiency tolerance and yield in Arabidopsis. *Plant Science* **307** (2021).

32.   N. Correa-Aragunde, N. Foresi, F. Del Castello, L. Lamattina, A singular nitric oxide synthase
with a globin domain found in Synechococcus PCC 7335 mobilizes N from arginine to
nitrate. *Sci Rep* **8**, 12505 (2018).

33.   M. M. Labarthe, G. A. Maroniche, L. Lamattina, C. M. Creus, Nitric oxide synthase
expression in Pseudomonas koreensis MME3 improves plant growth promotion traits.
Appl. Microbiol. Biotechnol. **108** (2024).

34.   A. L. Picciano, B. R. Crane, A nitric oxide synthase-like protein from Synechococcus
produces NO/NO3- from l-arginine and NADPH in a tetrahydrobiopterin- and Ca2+-
dependent manner. *J Biol Chem* **294**, 10708–10719 (`).

35. M. G. Campbell, B. C. Smith, C. S. Potter, B. Carragher, M. A. Marletta, Molecular
architecture of mammalian nitric oxide synthases. *Proc Natl Acad Sci U S A* **111**, E3614-
3623 (2014).

36. D. Felker, K. Lee, T. H. Pospiech, Y. Morishima, H. Zhang, M. Lau, D. R. Southworth, Y.
Osawa, Mapping interactions of calmodulin and neuronal NO synthase by crosslinking and
mass spectrometry. *Journal of Biological Chemistry* **300** (2024).

37. A. L. Yokom, Y. Morishima, M. Lau, M. Su, A. Glukhova, Y. Osawa, D. R. Southworth,
Architecture of the nitric-oxide synthase holoenzyme reveals large conformational changes
and a calmodulin-driven release of the FMN domain. *J Biol Chem* **289**, 16855–16865
(2014).

38. B. C. Smith, E. S. Underbakke, D. W. Kulp, W. R. Schief, M. A. Marletta, Nitric oxide synthase
domain interfaces regulate electron transfer and calmodulin activation. *Proc Natl Acad Sci*
U S A **110**, E3577-3586 (2013).

39. T. Jiang, M. C. Thielges, C. J. Feng, Emerging approaches to investigating functional protein
dynamics in modular redox enzymes: Nitric oxide synthase as a model system. *J. Biol.*
Chem. **301** (2025).

40. C. R. Pudney, D. J. Heyes, B. Khara, S. Hay, S. E. J. Rigby, N. S. Scrutton, Kinetic and
spectroscopic probes of motions and catalysis in the cytochrome P450 reductase family of
enzymes. *Febs Journal* **279**, 1534–1544 (2012).

41. T. M. Hedison, N. G. H. Leferink, S. Hay, N. S. Scrutton, Correlating Calmodulin Landscapes
with Chemical Catalysis in Neuronal Nitric Oxide Synthase using Time-Resolved FRET and a
5-Deazaflavin Thermodynamic Trap. *Acs Catalysis* **6**, 5170–5180 (2016).

42. T. M. Hedison, S. Hay, N. S. Scrutton, Trapping methods for probing functional
intermediates in nitric oxide synthases and related enzymes. *Frontiers in Bioscience-*
Landmark **23**, 1874–1888 (2018).

43. Y. He, M. M. Haque, D. J. Stuehr, H. P. Lu, Conformational states and fluctuations in
endothelial nitric oxide synthase under calmodulin regulation. *Biophysical Journal* **120**,
5196–5206 (2021).

44. J. Li, H. Zheng, C. Feng, Deciphering mechanism of conformationally controlled electron
transfer in nitric oxide synthases. *Frontiers in Bioscience-Landmark* **23**, 1803–1821 (2018).

45. C. J. Feng, Mechanism of nitric oxide synthase regulation: Electron transfer and
interdomain interactions. *Coordination Chemistry Reviews* **256**, 393–411 (2012).

46. D. J. Stuehr, J. Tejero, M. M. Haque, Structural and mechanistic aspects of flavoproteins:
electron transfer through the nitric oxide synthase flavoprotein domain. *The FEBS Journal*
276, 3959–3974 (2009).

47. M. M. Haque, C. Kenney, J. Tejero, D. J. Stuehr, A kinetic model linking protein
conformational motions, interflavin electron transfer and electron flux through a dual-
flavin enzyme-simulating the reductase activity of the endothelial and neuronal nitric oxide
synthase flavoprotein domains. *FEBS J* **278**, 4055–69 (Nov).

48. R. Chakravarti, K. S. Aulak, P. L. Fox, D. J. Stuehr, GAPDH regulates cellular heme insertion
into inducible nitric oxide synthase. *Proceedings of the National Academy of Sciences* **107**,
18004–18009 (2010).

49. Y. Morishima, M. Lau, W. B. Pratt, Y. Osawa, Dynamic cycling with a unique Hsp90/Hsp70-
dependent chaperone machinery and GAPDH is needed for heme insertion and activation
of neuronal NO synthase. *J Biol Chem* **299**, 102856 (2023).

50. A. Punjani, J. L. Rubinstein, D. J. Fleet, M. A. Brubaker, cryoSPARC: algorithms for rapid
unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290–296 (2017).

51. T. Wagner, F. Merino, M. Stabrin, T. Moriya, C. Antoni, A. Apelbaum, P. Hagel, O. Sitsel, T.
Raisch, D. Prumbaum, D. Quentin, D. Roderer, S. Tacke, B. Siebolds, E. Schubert, T. R.
Shaikh, P. Lill, C. Gatsogiannis, S. Raunser, SPHIRE-crYOLO is a fast and accurate fully
automated particle picker for cryo-EM. *Commun Biol* **2**, 218 (2019).

52. H. Shi, C. Wu, X. Zhang, Addressing compressive deformation of proteins embedded in
crystalline ice. *Structure* **31**, 213-220.e3 (2023).

53. B. R. Crane, The enzymology of nitric oxide in bacterial pathogenesis and resistance.
Biochemical Society Transactions **36**, 1149–1154 (2008).

54. K. Pant, A. M. Bilwes, S. Adak, D. J. Stuehr, B. R. Crane, Structure of a nitric oxide synthase
heme protein from Bacillus subtilis. *Biochemistry* **41**, 11071–9 (2002).

55. B. R. Crane, A. S. Arvai, D. K. Ghosh, C. Wu, E. D. Getzoff, D. J. Stuehr, J. A. Tainer, Structure
of nitric oxide synthase oxygenase dimer with pterin and substrate. *Science* **279**, 2121–6
(1998).

56. S. Corbalan-Garcia, J. C. Gómez-Fernández, Signaling through C2 domains: More than one
lipid target. *Biochimica et Biophysica Acta (BBA) - Biomembranes* **1838**, 1536–1547 (2014).

57. T. A. Leonard, "C2 Domain Proteins" in *Encyclopedia of Metalloproteins*, R. H. Kretsinger, V.
N. Uversky, E. A. Permyakov, Eds. (Springer, New York, NY, 2013;
https://doi.org/10.1007/978-1-4614-1533-6_39), pp. 309–318.

58. U. Siddhanta, A. Presta, B. Fan, D. Wolan, D. L. Rousseau, D. J. Stuehr, Domain Swapping in
Inducible Nitric-oxide Synthase: ELECTRON TRANSFER OCCURS BETWEEN FLAVIN AND
HEME GROUPS LOCATED ON ADJACENT SUBUNITS IN THE DIMER*. *Journal of Biological
Chemistry* **273**, 18950–18958 (1998).

59. J. Tejero, M. M. Haque, D. Durra, D. J. Stuehr, A Bridging Interaction Allows Calmodulin to
Activate NO Synthase through a Bi-modal Mechanism. *J Biol Chem* **285**, 25941–25949
(2010).

60. P.-F. Chen, K. K. Wu, Structural elements contribute to the calcium/calmodulin dependence
on enzyme activation in human endothelial nitric-oxide synthase. *J Biol Chem* **278**, 52392–
52400 (2003).

61. E. D. Garcin, C. M. Bruns, S. J. Lloyd, D. J. Hosfield, M. Tiso, R. Gachhui, D. J. Stuehr, J. A.
Tainer, E. D. Getzoff, Structural Basis for Isozyme-specific Regulation of Electron Transfer in
Nitric-oxide Synthase * [boxs]. *Journal of Biological Chemistry* **279**, 37918–37927 (2004).

62. B. Ghazi Esfahani, N. Walia, K. Neselu, Y. Garg, M. Aragon, I. Askenasy, H. A. Wei, J. H.
Mendez, M. E. Stroupe, Structure of dimerized assimilatory NADPH-dependent sulfite
reductase reveals the minimal interface for diflavin reductase binding. *Nat Commun* **16**,
2955 (2025).

63. H. Olteanu, R. Banerjee, Human Methionine Synthase Reductase, a Soluble P-450
Reductase-like Dual Flavoprotein, Is Sufficient for NADPH-dependent Methionine Synthase
Activation *. *Journal of Biological Chemistry* **276**, 35558–35563 (2001).

64. Z.-Q. Wang, R. J. Lawson, M. R. Buddha, C.-C. Wei, B. R. Crane, A. W. Munro, D. J. Stuehr,
Bacterial flavodoxins support nitric oxide production by Bacillus subtilis nitric-oxide
synthase. *J Biol Chem* **282**, 2196–2202 (2007).

65. S. Daff, I. Sagami, T. Shimizu, The 42-amino acid insert in the FMN domain of neuronal
nitric-oxide synthase exerts control over Ca(2+)/calmodulin-dependent electron transfer. *J
Biol Chem* **274**, 30589–30595 (1999).

66. M. Tiso, J. Tejero, K. Panda, K. S. Aulak, D. J. Stuehr, Versatile regulation of neuronal nitric
oxide synthase by specific regions of its C-terminal tail. *Biochemistry* **46**, 14418–28 (2007).

67. L. J. Roman, R. T. Miller, M. A. de La Garza, J. J. Kim, B. S. Siler Masters, The C terminus of
mouse macrophage inducible nitric-oxide synthase attenuates electron flow through the
flavin domain. *J Biol Chem* **275**, 21914–21919 (2000).

68. L. J. Roman, P. Martásek, R. T. Miller, D. E. Harris, M. A. de La Garza, T. M. Shea, J. J. Kim, B.
S. Masters, The C termini of constitutive nitric-oxide synthases control electron flow
through the flavin and heme domains and affect modulation by calmodulin. *J Biol Chem*
275, 29225–29232 (2000).

69. D. Hamdane, C. Xia, S.-C. Im, H. Zhang, J.-J. P. Kim, L. Waskell, Structure and Function of an
NADPH-Cytochrome P450 Oxidoreductase in an Open Conformation Capable of Reducing
Cytochrome P450*. *Journal of Biological Chemistry* **284**, 11374–11384 (2009).

70.   M. Tiso, D. W. Konas, K. Panda, E. D. Garcin, M. Sharma, E. D. Getzoff, D. J. Stuehr, C-
812   terminal Tail Residue Arg1400 Enables NADPH to Regulate Electron Transfer in Neuronal
813   Nitric-oxide Synthase *. *Journal of Biological Chemistry* **280**, 39208–39219 (2005).

814   71.   S. Adak, M. Sharma, A. L. Meade, D. J. Stuehr, A conserved flavin-shielding residue
815   regulates NO synthase electron transfer and nicotinamide coenzyme specificity.
816   *Proceedings of the National Academy of Sciences* **99**, 13516–13521 (2002).

817   72.   E. A. Nalefski, J. J. Falke, The C2 domain calcium-binding motif: structural and functional
818   diversity. *Protein Sci* **5**, 2375–2390 (1996).

819   73.   N. Verdaguer, S. Corbalan-Garcia, W. F. Ochoa, I. Fita, J. C. Gómez-Fernández, Ca2+ bridges
820   the C2 membrane-binding domain of protein kinase Cα directly to phosphatidylserine. *The
821   EMBO Journal* **18**, 6329–6338 (1999).

822   74.   L. Zheng, R. Krishnamoorthi, M. Zolkiewski, X. Wang, Distinct Ca2+ Binding Properties of
823   Novel C2 Domains of Plant Phospholipase Dα and β *. *Journal of Biological Chemistry* **275**,
824   19700–19706 (2000).

825   75.   M. González-Pleiter, M. Tamayo-Belda, G. Pulido-Reyes, G. Amariei, F. Leganés, R. Rosal, F.
826   Fernández-Piñas, Secondary nanoplastics released from a biodegradable microplastic
827   severely impact freshwater environments. *Environ. Sci.: Nano* **6**, 1382–1392 (2019).

828   76.   I. Torrecilla, F. Leganés, I. Bonilla, F. Fernández-Piñas, Light-to-dark transitions trigger a
829   transient increase in intracellular Ca2+ modulated by the redox state of the photosynthetic
830   electron transport chain in the cyanobacterium Anabaena sp. PCC7120. *Plant, Cell &
831   Environment* **27**, 810–819 (2004).

832   77.   A. De Wever, K. Benzerara, M. Coutaud, G. Caumes, M. Poinsot, F. Skouri-Panet, T. Laurent,
833   E. Duprat, M. Gugger, Evidence of high Ca uptake by cyanobacteria forming intracellular
834   CaCO3 and impact on their growth. *Geobiology* **17**, 676–690 (2019).

835   78.   C. C. Moser, J. M. Keske, K. Warncke, R. S. Farid, P. L. Dutton, Nature of biological electron
836   transfer. *Nature* **355**, 796–802 (1992).

837   79.   W.-C. Huang, J. Ellis, P. C. E. Moody, E. L. Raven, G. C. K. Roberts, Redox-Linked Domain
838   Movements in the Catalytic Cycle of Cytochrome P450 Reductase. *Structure* **21**, 1581–1589
839   (2013).

840   80.   J. B. Hopkins, R. E. Gillilan, S. Skou, BioXTAS RAW: improvements to a free open-source
841   program for small-angle X-ray scattering data reduction and analysis. *J Appl Cryst* **50**, 1545–
842   1553 (2017).

843   81.   C. E. Aitken, R. A. Marshall, J. D. Puglisi, An Oxygen Scavenging System for Improvement of
844   Dye Stability in Single-Molecule Fluorescence Experiments. *Biophys J* **94**, 1826–1835
845   (2008).

846   82.   A. Cheng, C. Negro, J. F. Bruhn, W. J. Rice, S. Dallakyan, E. T. Eng, D. G. Waterman, C. S.
847   Potter, B. Carragher, Leginon: New features and applications. *Protein Sci* **30**, 136–150
848   (2021).

849   83.   G. Pintilie, W. Chiu, Comparison of Segger and other methods for segmentation and rigid-
850   body docking of molecular components in cryo-EM density maps. *Biopolymers* **97**, 742–
851   760 (2012).

852   84.   J. Abramson, J. Adler, J. Dunger, R. Evans, T. Green, A. Pritzel, O. Ronneberger, L. Willmore,
853   A. J. Ballard, J. Bambrick, S. W. Bodenstein, D. A. Evans, C.-C. Hung, M. O'Neill, D. Reiman,
854   K. Tunyasuvunakool, Z. Wu, A. Žemgulytė, E. Arvaniti, C. Beattie, O. Bertolli, A. Bridgland, A.
855   Cherepanov, M. Congreve, A. I. Cowen-Rivers, A. Cowie, M. Figurnov, F. B. Fuchs, H.
856   Gladman, R. Jain, Y. A. Khan, C. M. R. Low, K. Perlin, A. Potapenko, P. Savy, S. Singh, A.
857   Stecula, A. Thillaisundaram, C. Tong, S. Yakneen, E. D. Zhong, M. Zielinski, A. Žídek, V.
858   Bapst, P. Kohli, M. Jaderberg, D. Hassabis, J. M. Jumper, Accurate structure prediction of
859   biomolecular interactions with AlphaFold 3. *Nature* **630**, 493–500 (2024).

860   85.   P. Emsley, K. Cowtan, Coot: model-building tools for molecular graphics. *Acta Crystallogr D
861   Biol Crystallogr* **60**, 2126–32 (2004).

862   86.   T. I. Croll, ISOLDE: a physically realistic environment for model building into low-resolution
863   electron-density maps. *Acta Cryst D* **74**, 519–530 (2018).

864   87.   (IUCr) Real-space refinement in PHENIX for cryo-EM and crystallography.
865   https://journals.iucr.org/d/issues/2018/06/00/ic5103/index.html.

866   88.   F. P. Guengerich, M. V. Martin, C. D. Sohl, Q. Cheng, Measurement of cytochrome P450 and
867   NADPH–cytochrome P450 reductase. *Nat Protoc* **4**, 10.1038/nprot.2009.121 (2009).

868   89.   Z. A. Maschmann, T. K. Chua, S. Chandrasekaran, H. Ibáñez, B. R. Crane, Redox properties
869   and PAS domain structure of the Escherichia coli energy sensor Aer indicate a multistate
870   sensing mechanism. *J Biol Chem* **298**, 102598 (2022).

871   90.   S. L. Christgen, S. M. Becker, D. F. Becker, Methods for determining the reduction potentials
872   of flavin enzymes. *Methods Enzymol* **620**, 1–25 (2019).

873   91.   C. Lin, S. Feng, C. C. DeOliveira, B. R. Crane, Cryptochrome-Timeless structure reveals
874   circadian clock timing mechanisms. *Nature* **617**, 194–199 (2023).

875   92.   D. Pan, A. Brockmeyer, F. Mueller, A. Musacchio, T. Bange, Simplified Protocol for Cross-
876   linking Mass Spectrometry Using the MS-Cleavable Cross-linker DSBU with Efficient Cross-
877   link Identification. *Anal. Chem.* **90**, 10990–10999 (2018).

878   93.   I. M. Lagerwaard, P. Albanese, A. Jankevics, R. A. Scheltema, Xlink Mapping and AnalySis
879   (XMAS) - Smooth Integrative Modeling in ChimeraX. bioRxiv [Preprint] (2022).
880   https://doi.org/10.1101/2022.04.21.489026.

Acknowledgements: We thank Angela Picciano for helpful discussions. We thank the National

Center for CryoEM Access and Training (NCCAT) and the Cornell Center for Material Science

(CCMR) for access to data collection facilities. We thank Yajie Xu and Dr Mariena Silvestry Ramos

for assistance in CryoEM training and data collection. We thank the Proteomics and

Metabolomics Facility of Cornell University for providing the mass spectrometry data and NIH

SIG grant 1S10 OD017992-01 support for the Orbitrap Fusion mass spectrometer. The Center for

High-Energy X-ray Sciences at CHESS (CHEXS) is supported by the NSF award DMR-2342336, and

the MacCHESS resource is supported by NIGMS award 1-P30-GM124166. All figures were

created or formated using Biorender.com.

Funding: This work was supported by grant MCB-2129728 from the National Science

Foundation to BRC. **Author contributions:** D.N. and B.R.C. designed research, conducted

research, analyzed data and wrote the manuscript. Competing interests: The authors declare

that they have no competing interests. Data and materials availability: The cryoEM data and

structures are available at the Protein Data Bank with accession codes: 9Q05, 9Q06, 9Q0Y,

9Q0X, and 9Q15. All other data needed to evaluate the conclusions in the paper are present in

the paper or the Supplementary Materials.

Figures

Figure 1. Architecture of Ca^{2+}-free locked syNOS (A) Gene architecture of Synechococcus NOS,

mammalian NOS and a typical bacterial NOS. Note that some non-cyanobacterial prokaryotic

NOSs have auxiliary domains. (B) Sideview of electron density (left, threshold = 0.0298, C2

symmetry applied) and corresponding ribbon representation (right) for the locked syNOS

homodimer (NOS-Sym). Subunit 1 (right) colored as designated by the gene architecture,

Subunit 2 (left) colored as follows: C2' - lime green, pGlb' - navy, Glb' - magenta, Oxy' - indian

red, Fld' – pale yellow and FNR' - bright orange . (C) Top view of (B). (D) Individual syNOS

domains with cofactors indicated.

Figure 2. Structural comparison of syNOS$_{Red}$, nNOS$_{Red}$, and CYPOR (A-C). The di-flavin closed

conformation is defined by the L3 loop (situated between the Fld and the FNR domain, blue in

Fig. 2A, B), the Fld domain, and what we define as Interfacing Region (IR) 1, IR2 and the CTT-IR.

Domain architectures (top), the corresponding ribbon diagrams (middle), and key interfaces

(lower) mediated by electrostatic interactions for syNOS$_{Red}$ (A), nNOS$_{Red}$ (PDB:1TLL, B) and

CYPOR (PDB:1AMO, C) Surfaces are colored by coulombic electrostatic potential (blue – positive;

white – neutral; red – negative)) with the Fld and FNR domains separated and rotated to view

the interface. Interfacing Regions (IR) 1 (basic-to-acidic patch interactions between the FAD and

FMN binding pockets), IR2 (Fld interaction-to-FNR-control region) and CTT-IR (CTT interaction

with Fld) are denoted by colored circles in the electrostatic surface representation) and semi-

transperant rectangles in ribbons.

Figure 3. CTT interactions that stabilize the NOS$_{Red}$-NOS$_{Glb}$ locked state. (A) Ribbon diagram of

syNOS$_{Red}$ and syNOS$_{Glb}$ from the boxed region of inactivated syNOS$_{FL}$ (upper left inset) showing

key stabilizing interfaces. (B) Aliphatic surface patch between syNOS$_{CTT}$ and syNOS$_{Glb}$ shown by

hydrophobicity (yellow) after separating and rotating each domain to view the interface. (C) Fld-

to-CTT lock interface mediated by Lys987(Fld)/Glu1454(CTT) salt bridge and

Asp984(Fld)/Tyr1458(FNR) and Thr953/Asn1457 hydrogen bonds. (D) Fld-to-Glb interface

mediated by Asn1457(CTT)/Thr953(Fld) hydrogen bond, Arg1460(CTT)/Arg382-carbonyl(Glb)

hydrogen bond and Asn1457-carbonyl(CTT)/arg382(Glb) hydrogen bond. (E) C-Terminal cap

interface mediated by Trp1419(CTT)/His1419(FNR) hydrogen bond and a Leu1468-

Carboxyl(CTT)/Lys1412 salt bridge.

Figure 4. syNOS$_{C2}$ interactions with NOS$_{FNR}$ and NOS$_{Oxy}$. (A) Superimposition of syNOS (C2,

green) with Perforin C2 (PDB: 4Y1S, yellow), C2A-CEP120 (PDB:6EWL, orange) and Munc13-C2B

(PDB:6NYT, magenta) as labeled, with root mean square deviation (RMSD) of Cα positions

(RMSD), and Template Modeling (TM) scores boxed. B) Ribbon diagram of a syNOS subunit,

colored by domain with key C2-Oxy interactions shown in insets b1 and b2. b1) Salt bridge

between Arg57(C2)/Glu523(Oxy) with Glu523 stabilized by Gln494. b2) Salt bridge between

Arg107(C2)/Glu568(Oxy) with the Arg107 amide stabilized by Asn575. (C) Ribbon diagram of a syNOS subunit showing the electrostatic lock interface between syNOS$_{C2}$ and syNOS$_{Oxy}$ in box inset c1. (c1) C2 rotated 90° (left) and FNR (right, orientation as in (C), colored by electrostatic surfaces (colored: blue-positive, white-intermediate and red –negative). C2 (green, ribbons) is also shown in its original position for reference.

Figure 5. Ca^{2+}-activated C2 mobility. (A) syNOS gene architecture colored by domain, with linkers labeled L1-3(above). Surface representation of three separate syNOS states with relative orientations of the C2 domain. (B) Superimposed surface representations of the turnover state (C2 – lime green) and inactivated states (C2 – forest green, Oxy – red and Glb – hot pink) reveal dramatically different C2 positions. (C) L2 in the turnover state (tan electron density) connects to the Fld (yellow ribbons) but then swivels 135° (green density) from its position in the inactivated state to direct the NOS_{Fld} to the adjacent NOS_{Oxy} subunit (purple ribbons of AlphaFold prediction).

Figure 6. syNOS$_{FNR}$ likely requires a large conformational change to reduce syNOS$_{Glb}$. (A) Cartoon

representation of E.*coli* Hmp (PDB: 1GVH) with the NOD domain (lime green) and FNR (metallic

blue) domains orientated to show the Heme-to-FAD separation of 17.8 Å. (B) syNOS$_{Glb}$ (purple)

and associated syNOS$_{FNR}$ domain (red, control region 1084-1224 not shown) superimposed on

the Hmp Glb (green) and associated FNR (blue). SyNOS$_{FNR}$ would need to rotate 90° and

translate 28 Å to adopt an equivalent Hmp conformation.

Figure 7. Model for NOS to NOD activity switching in syNOS. In the inactivated locked state syNOS$_{Oxy}$ equilibrates between a loose and tight dimer, the latter stabilized by BH$_4$. This state can transition into the asymmetric NOD active state which involves a release of a C2 domain from one subunit and increased mobility of NOS$_{Fld}$ and NOS$_{FNR}$, such that NOS$_{FNR}$ can reduce the syNOS$_{Glb}$ to activate NOD activity. SyNOS$_{Glb}$ reduction coincides with a loosening of the CTT to allow hydride transfer from NADPH to FAD and movement of syNOS$_{FNR}$. In the presence of Ca^{2+}/NADPH/L-arginine syNOS rearranges to facilitate NOS activity. The displacement of syNOS$_{C2}$ from NOS$_{Oxy}$ and its binding to NOS$_{Glb}$ relieves constraints on L2 and exposes the Trp715 site such that NOS$_{FNR}$ can relocate to NOSOxy on the opposing subunit and reduce the NOS$_{Oxy}$ heme for NO production.

1000

1001

Structure and dynamics of a muti-domain nitric oxide synthase regulated by a C2
domain

Dhruva Nair and Brian R. Crane

Supplementary Movies

Movie S1: 3D Flexible Refinement of syNOS dimer shows a 'breathing' motion

Supplementary Tables

Table S1: Description of datasets, conditions and refined classes.

Table S2: Identified and validated DSBU crosslinks

Table S3: Cryo-EM data collection, refinement, and validation statistics.

Table S4: DNA primers used in the study with the cloning approach.

Table S5: Reduction potentials ($E^{0'}$) for $syNOS_{Glb}$, $syNOS_{Fld}$ and $syNOS_{FNR}$ vs. the Normal Hydrogen
Electrode (NHE)

Supplementary Figures

Figure S1: syNOS purification

Figure S2: Dataset 1 processing scheme

Figure S3: Dataset 2 processing scheme

Figure S4: Merged data for NOS-Sym state, boxed class used for 3DVA and 3DFlex

Figure S5: Dataset 3 processing scheme and validation.

Figure S6: Dataset 4 processing scheme and validation

Figure S7: Dataset 5 processing scheme and validation

Figure S8: syNOS linker intertwining

Figure S9: syNOS tight and loose NOS_{Oxy} states

Figure S10: Identified cross-links plotted onto dimer interface Oxy+pGlb and the syNOS FNR+CTT

Figure S11: Phylogenetic sequence clustering in the context of $syNOS_{Red}$.

Figure S12: C-terminal tail (CTT) sequence alignment of syNOS, mNOSs and diatom NOS.

Figure S13: Density validation of key interactions

Figure S14: SEC-SAXS of syNOS and variants.=

Figure S15: Steady state cytochrome c reductase activities of WT syNOS and variants

Figure S16: Flavin reduction by NADPH

Figure S17: Isothermal calorimetric titration of Ca^{2+} with $syNOS_{C2}$

1034

1035   **Supplemental Movies**

1036

1037   **Movie S1.** 3D Variability analysis of one subunit of the syNOS asymmetric dimer showing the

1038   breathing motion of NOS_{Red} as a unit relative to motion about the NOS_{Oxy} interface.

1039

1040   **Supplementary Tables**

1041

Dataset	Conditions	Class 1	Class 2
1	GFB1	Symmetric Dimer (NOS-Sym)	Asymmetric Dimer (NOS-Asym)
2			
3	GFB1 + 1mM Ca^{2+}/ 1mM L-arg / BH_4*	Asymmetric Dimer (NOS-Asym)	Turnover (NOS-TO_L)
4	GFB1 + 1mM Ca^{2+}/ 1mM L-arg / BH_4*	Asymmetric Dimer (NOS-Asym)	Turnover (NOS-TO_T)
5	GFB1 + 1mM Ca^{2+}/1mM L-arg/ BH_4* / 1mM NADPH	Monomer (NOS-Mon)	none

1042

1043   GFB1: 25 mM HEPES (pH 7.5), 150 mM NaCl, 1 mM TCEP, 10% glycerol and 10 μM BH_4.

1044   *Additional 25 μM BH_4 was supplemented.

1045   **Table S1.** Description of datasets, conditions and refined classes.

1046

1047

1048

1049

1050

Inactive				
Residue 1	Residue 2	Intra Cα- Cα	Inter Cα- Cα	XlinkX Score
FNR - FNR				
1111	1017	**8**	143	124
1366	1447(CTT)	**20**	117	118
1053	1463(CTT)	**40**	121	75
1366	1053(CTT)	**21**	138	52
Interdomain				
191(pGlb)	538(Oxy)	**17**	**24**	78

Turnover				
Residue 1	Residue 2	Intra Cα- Cα	Inter Cα- Cα	XlinkX Score
FNR-FNR				
1019	1111	**6**	149	222
1111	1017	**8**	143	86
Interdomain				
196(pGlb)	538(Oxy)	**16**	**31**	97
191(pGlb)	538(Oxy)	**17**	**24**	78
470(Glb)	481(Oxy)	**17**	57	61
473(Glb)	481(Oxy)	**12**	58	50
196(pGlb)	481(Oxy)	**24**	44	48
413(Glb)	592(Oxy)	**19**	79	45
1017(FNR/L3)	994(Fld)	**23**	120	41
470(Glb)	538(Oxy)	40	55	41

1051 **Table S2.** Identified and validated DSBU crosslinks. Reasonable distances based on the cryo-EM

1052 structures are bolded and underlined. PDB:9Q15 was used to estimate distances.

Dataset			Ca2+/L-Arginine		Ca^{2+}/L-Arginine/NADPH
State	Symmetric Dimer	Asymmetric	Turnover	Asymmetric (Composite)	Monomer
Data Collection and Processing					
Microscope	Arctica	Krios			Krios
Voltage (keV)	200	300			300
Detector	K3	Gatan K3			Falcon4EC
Nominal magnification	63,000	105,000			105,000
Data Acquisition Software	Serial EM	Leginon			Leginon
Electron dose (e–/Å2)	50	50.5			44.95
Pixel Size (Å) (binned)	1.31	0.422(0.844)			0.7304
Defocus range (μm)	0.5-2	0.4-2.1			0.4-2.2
Number of movies (#)	1214 + 4055	5019			9036
Number of particles	131,000	70,944	61,553	70,944	45,964
Symmetry imposed	C2	C1	C1	C1	C1
Resolution (Å)	3.95	3.16	3.69	3.16	3.09
FSC threshold	0.143	0.143	0.143	0.143	0.143
Refinement					
Initial model used	PDB 9Q15	PDB 9Q15	PDB 9Q15	Alphafold	PDB 9Q15
Non-hydrogen atoms	22712	16319	11268	16319	11404
Protein residues	2796	2001	1358	2001	1398
Ligand(#)	Heme(4) FAD(2) FMN(2)	Heme(2) FAD(1) FMN(1)	Heme(4)	Heme(4) FAD(1) FMN(1)	Heme(2) FMN(1) FAD(1) NADPH(1)
RMS Deviation					
Bond lengths (Å)	0.02	0.005	0.003	0.003	0.003
Bond angles (°)	1.55	0.55	0.574	0.485	0.519
MolProbity score	1.54	1.58	1.5	1.74	1.54
Clashscore	2.18	3.49	4.65	4.66	4.93
Poor rotamers (%)	2.08	1.63	0.93	1.8	0.91
Ramachandran Favored (%)	95.1	96.0	96.1	95.5	96.0
Ramachandran Allowed (%)	4.6	3.9	3.7	4.3	3.8
Ramachandran Disallowed (%)	0.3	0.2	0.2	0.2	0.2
Fit to map (CCmask)	0.8	0.79	0.75	0.81	0.82
EMDB (maps)	EMD-72088	EMD-72110	EMD-72109	EMD-72104, EMD-72105,EMD-72116	EMD-72087
PDB (model)	9Q06	9Q0Y	9Q0X	9Q15	9Q05

1053

Table S3. Cryo-EM data collection, refinement, and validation statistics.

Primer Name	Primer	Cloning Approach
pet28_twin_strep_syNOS_fwd	tcgagcaccaccaccacc	Gibson/NEB HiFi
pet28_twin_strep_syNOS_rev	catatgtcctccggattggaagtac	Gibson/NEB HiFi
syNOS_fwd	tccaatccggaggacatatgatgcttgtcaacgactctc	Gibson/NEB HiFi
syNOS_rev	gtggtggtggtggtgctcgagctacaagttggctagcc	Gibson/NEB HiFi
Hptg_fwd_PAYC_mcs1	TAATGCTTAAGTCGAACAG	Gibson/NEB HiFi
Hptg_rev_PAYC_mcs1	GGTATATCTCCTTATTAAAGTTAAAC	Gibson/NEB HiFi
Hptg_fwd	CTTTAATAAGGAGATATACCATGAAAGGACAAGAAACTCGTG	Gibson/NEB HiFi
Hptg_rev	TCTGTTCGACTTAAGCATTATCAGGAAACCAGCAGCTG	Gibson/NEB HiFi
gapa_fwd_pAYC_mcs2	TAATTAACCTAGGCTGCTG	Gibson/NEB HiFi
gapa_rev_pAYC_mcs2	ATGTATATCTCCTTCTTATACTTAAC	Gibson/NEB HiFi
gapa_fwd	TATAAGAAGGAGATATACATATGACTATCAAAGTAGGTATCAAC	Gibson/NEB HiFi
gapa_rev	GCAGCAGCCTAGGTTAATTATTATTTGGAGATGTGAGC	Gibson/NEB HiFi
pet28_tws_delc2_fwd	ctcgagcaccaccaccac	Gibson/NEB HiFi
pet28_tws_delc2rev	catatgtcctccggattggaag	Gibson/NEB HiFi
syNOSdelc2_178_1468_fwd	tccaatccggaggacatatgacctactcagaagccgtc	Gibson/NEB HiFi
syNOSdelc2_178_1468_fwd	tggtggtggtggtgctcgagctacaagttggctagccatttttc	Gibson/NEB HiFi
pet28_tws_delctt_fwd	tagctcgagcaccaccac	Gibson/NEB HiFi
pet28_tws_delctt_rev	catatgtcctccggattggaag	Gibson/NEB HiFi
syNOS_delctt1_1_1440_fwd	tccaatccggaggacatatgatgcttgtcaacgactctc	Gibson/NEB HiFi
syNOS_delctt1_1_1440_fwd	tggtggtggtgctcgagctagtagttgtctttctccacttg	Gibson/NEB HiFi
syNOS_K1412E_fwd	GCAAATTGCTgaaACAGAAGGAG	KLD
syNOS_K1412E_rev	ATAAATACCTCAAACACATTG	KLD
syNOS_T953A_fwd	GATCGGCAGCgcgGTCTATGAAC	KLD
syNOS_T953A_rev	CCTAGTACGGAATAGTTGAG	KLD
syNOS_K987A_fwd	CGACGAAATCgcgGGACAAGCCG	KLD
syNOS_K987A_rev	CCTTTATGTAGGGGAAC	KLD

1055

1056 **Table S4**. Primers used in the study with the cloning approach (left).

Domain	Couple	Potential (mV)	n	Calibrating dye (mV)
Globin (337-467)	Fe^{3+}/Fe^{2+}	-250 ± 20	1	safranin T (-289, 12.5µM)
Fld (856-1027)	ox/hq	-173 ± 3	2	cresyl violet (-177.0, 60 µM)
FNR (1027-1468)	ox/hq	-290 ± 13	2	safranin T (-289, 12.5 µM)

1057

1058 **Table S5.** Reduction potentials ($E^{0'}$) for $syNOS_{Glb}$, $syNOS_{Fld}$ and $syNOS_{FNR}$ vs. the Normal
1059 Hydrogen Electrode (NHE). For $syNOS_{Glb}$ the single electron reduction (n_e=1) from Fe^{3+} to Fe^{2+}
1060 was measured whereas for the flavin-containing domains a midpoint potential for the 2-
1061 electron reduction (n_e=2) from oxidized (ox) to hydroquinone (hq) was measured as no spectral
1062 signature for the semiquinone was apparent in the titrations. (n = 3 for all measurements)

1063

1064

1065 **Supplementary Figures**

1066   **Figure S1.** syNOS Purification. Representative chromatogram from a Superdex 200 increase
1067   10/30 size exclusion chromatography column with an arrow denoting the syNOS dimer peak.
1068   The inset corresponds to a Coomassie blue stained 4-12% SDS-PAGE gel of fractions 5,6,7 and 8.

1069

1070

1071

Dataset 1

1073　**Figure S2.** Dataset 1 processing scheme. (A) CryoEM processing workflow for dataset 1: syNOS

1074　ligand free. NOS-Sym class used for 3DVA and 3DFlex boxed (dashed-red). (B) Representative

1075   2D Classes. CTF, contrast transfer function; NU, non-uniform refinement. (C) Overall Particle and

1076   Gold-standard Fourier Shell Correlation (GSFSC) curves for NOS-Asym and NOS-Sym.

1077

1078

Figure S3. Dataset 2 processing scheme. (A) CryoEM processing workflow for dataset 2: syNOS

ligand free. NOS-Sym class used for 3DVA and 3DFlex boxed (dashed-red). (B) Representative 2D

Classes. (C) GSFSC curves and positional distribution of particles.

Figure S4. Merged data for NOS-Sym state, boxed class used for 3DVA and 3DFlex.

Figure S5. Dataset 3 processing scheme and validation. (A) CryoEM processing workflow for

dataset 3: syNOS in the presence of Ca²⁺/L-arginine (B) Representative 2D Classes for NOS-Asym

state (C) Representative 2D classes for NOS turnover state (NOS-TO$_L$) (D) Asymmetric states

GSFSC curves, orientation validation, model density fit and map to model Fourier Shell

Correlation (FSC).

1088

1089

1090   **Figure S6.** Dataset 4 processing scheme and validation. (A) CryoEM processing workflow for

1091   dataset 4, syNOS with NADPH/Ca^{2+} including GSFSC curves. (B) Representative 2D classes for

1092   NOS-TO$_T$.

Dataset 5

A

Micrographs
9036

↓ Patch Motion
Patch CTF
Curate Micrographs

Micrographs
4989

↓ Blob Picker

Particles
909,465

↓ 2D Classification

↓ Ab Initio
Hetero Refinement
Rebalance Orientations
Topaz Training + Extraction
2D classification

2x

↓ Ab Initio
Hetero Refinement

Class 1   Class 2   Class 3

↓ Non-Uniform Refinement

y 180

3.09 Å
45K Particles
Monomer

B

150 Å

C

GSFSC Resolution: 3.15Å

No Mask (4.1Å)
Spherical (3.9Å)
Loose (3.4Å)
Tight (3.1Å)
Corrected (3.1Å)

DC   7.3Å   3.7Å   2.4Å   1.8Å

Elevation
π/2
π/4
0
−π/4
−π/2

−π −3π/4 −π/2 −π/4 0 π/4 π/2 3π/4 π
Azimuth

# of images

D

Map to Model

Unmasked
Masked

3.4 Å

FSC

1/Å

E

Globin Heme   CTT(1440-1468)   NADP+ Binding Site

FMN   FAD   Oxygenase Heme

1093

1094 **Figure S7**. Dataset 5 processing scheme and validation. (A) CryoEM processing workflow for

1095 dataset 5: syNOS with L-arginine, NADPH/Ca^{2+}/NADPH (B) Representative 2D classes for NOS-

1096 Mon. (C) GSFSC curves, orientation validation and (D) Map to Model FSC. (E) Density fit of

1097 model in map.

1098

1099

1100 **Figure S8.** SyNOS linker intertwining. Intermediate resolution homodimer cryoEM density with

1101 all three linkers colored the same as their preceding domains. Note how L1 threads beneath L2

1102 and on top of L3. Movement of the C2 domain (green) will release the Fld module for NOS$_{Oxy}$

1103 reduction.

Figure S9. SyNOS tight and loose NOS$_{Oxy}$ states. (A) syNOS cryoEM density (dataset 4) with the NOS$_{Oxy}$ dimer interface outlined (orange and green) to show each subunit. (B) Close up of the dimeric interface showing the close association of NOS$_{Oxy}$ α9 to an intersubunit pGlb C/D loop. Although at lower resolution, strong electron density for the core the NOS-TO$_T$ interface includes the helical lariats and BH$_4$. (C) Loose (green) and tight (blue) NOS$_{Oxy}$ dimers differ by a large 20 Å breathing motion normal to the dimer interface that also affects the positioning of the C2 domain (lower right peripheral electron density).

A

**Inactive
Dimer Interface**

Residue 1	Residue 2	Intra Cα- Cα	Inter Cα- Cα	XlinkX Score
191(pGlb)	538(Oxy)	**<u>17</u>**	**<u>24</u>**	78

B

**Turnover
Dimer Interface**

Residue 1	Residue 2	Intra Cα- Cα	Inter Cα- Cα	XlinkX Score
196(pGlb)	538(Oxy)	**<u>16</u>**	**<u>31</u>**	97
191(pGlb)	538(Oxy)	**<u>17</u>**	**<u>24</u>**	78
196(pGlb)	481(Oxy)	**<u>24</u>**	44	48

C

Inactive CTT

Residue 1	Residue 2	Intra Cα- Cα	Inter Cα- Cα	XlinkX Score
1366	1447(CTT)	**<u>20</u>**	117	118
1053	1463(CTT)	**<u>40</u>**	121	75
1366	1053(CTT)	**<u>21</u>**	138	52

Figure S10. DSBU crosslinks plotted onto the dimer interface of syNOS Oxy + pGlb and the syNOS FNR + CTT. (A) syNOS inactive dimer interface, subunit 1 Oxy (red) + pGlb (blue) and subunit 2 Oxy (bright red) + pGlb (navy), with magenta crosslinks corresponding to table (left). (B) syNOS inactive dimer interface, protomer 1 Oxy (red) + pGlb (blue) and subunit 2 Oxy (bright red) + pGlb (navy), with green crosslinks corresponding to table (left). (C) syNOS inactive CTT with magenta crosslinks corresponding to table (left). Crosslinks within range based on the structures are underlined and bolded.

1125

Cluster ID	Average Sequance Length	Standard Deviation of Sequence Length	Sequence Count
2	678	63	2521
39	702	49	1588
55	1117	103	278
3	1479	118	94
13	976	174	63
58	700	17	49
30	1126	283	47
7	638	35	45
18	601	74	44
33	652	36	44

1126   **Figure S11.** Phylogenetic sequence clustering in the context of syNOS$_{Red}$(residues 853-1468).

1127   Representative clusters were subjected to PhyML with clusters above 40 sequences represented

1128   in the tree. (B) Cluster composition by sequence length and count.

1129

1131   **Figure S12.** C-terminal tail (CTT) sequence alignment of syNOS, mNOSs and diatom NOS.

1132

1133

1134

Figure S13. Cryo-EM density validation of key interactions. (A) Density fit of syNOS-Asym

(9Q15), key interactions shown in Fig 3 C-E (B) Density fit of syNOS-Asym (9Q15), key

interactions shown in Fig 4. B1-B2.

1138

1139

1140

1141

1142   **Figure S14.** SEC-SAXS of syNOS and variants. (A) Dimensionless Kratky plots, binned by 4 qRg

1143   values, for samples in the presence and absence of Ca^{2+}. (B) Table of predicted and

1144   experientially characterized radius of gyrations (Rgs) and molecular weights calculated either by

1145   volumes of correlation (V_C) or Porod volumes (V_P). (Samples were measured once)

1146

1147

Figure S15. Steady state cytochrome c reductase activities of WT syNOS and variants expressed

as turnover numbers (Cc reduced (NOS subunit)$^{-1}$ min^{-1}) with standard errors shown as error

bars (see Methods). Statistical comparisons between groups were performed with a 2-way

ANOVA with post hoc analysis using Tukey's multiple comparisons test. Significant differences

are indicated as follows: ns (not significant), * ($p \leq 0.05$), ** ($p \leq 0.01$), *** ($p \leq 0.001$), **** ($p \leq$

0.0001)..Ca:WT vs. Ca:K987A (ns); Ca:WT vs. Ca:ΔC2 (ns); Ca:WT vs. Ca:K1412E (ns); Ca:WT vs.

Ca:ΔCTT (ns); Ca:WT vs. EGTA:WT (****) ; Ca:WT vs. EGTA:K987A (ns); Ca:WT vs. EGTA:ΔC2 (ns);

Ca:WT vs. EGTA:K1412E (****); Ca:WT vs. EGTA:ΔCTT (****) ;Ca:K987A vs. Ca:ΔC2 (ns);

Ca:K987A vs. Ca:K1412E (***); Ca:K987A vs. Ca:ΔCTT (****); Ca:K987A vs. EGTA:WT (**);

Ca:K987A vs. EGTA:K987A (**); Ca:K987A vs. EGTA:ΔC2 (ns); Ca:K987A vs. EGTA:K1412E (****);

Ca:K987A vs. EGTA:ΔCTT (****); Ca:ΔC2 vs. Ca:K1412E (ns); Ca:ΔC2 vs. Ca:ΔCTT (**); Ca:ΔC2 vs.

EGTA:WT (****); Ca:ΔC2 vs. EGTA:K987A (ns); Ca:ΔC2 vs. EGTA:ΔC2 (ns); Ca:ΔC2 vs.

EGTA:K1412E (****); Ca:ΔC2 vs. EGTA:ΔCTT (****);Ca:K1412E vs. Ca:ΔCTT (ns); Ca:K1412E vs.

EGTA:WT (****); Ca:K1412E vs. EGTA:K987A (ns); Ca:K1412E vs. EGTA:ΔC2 (ns); Ca:K1412E vs.

EGTA:K1412E (**); Ca:K1412E vs. EGTA:ΔCTT (****); Ca:ΔCTT vs. EGTA:WT (**); Ca:ΔCTT vs.

EGTA:K987A (*); Ca:ΔCTT vs. EGTA:ΔC2 (***); Ca:ΔCTT vs. EGTA:K1412E (ns); Ca:ΔCTT vs.

EGTA:ΔCTT (**); EGTA:WT vs. EGTA:K987A (****); EGTA:WT vs. EGTA:ΔC2 (****); EGTA:WT vs.

EGTA:K1412E (ns); EGTA:WT vs. EGTA:ΔCTT (ns); EGTA:K987A vs. EGTA:ΔC2 (ns); EGTA:K987A vs.

EGTA:K1412E (***); EGTA:K987A vs. EGTA:ΔCTT (***); EGTA:ΔC2 vs. EGTA:K1412E (****);

EGTA:ΔC2 vs. EGTA:ΔCTT (****); EGTA:K1412E vs. EGTA:ΔCTT (ns). (n=3, except for K1412E

+EGTA; n = 2).

1169

A

Reductase	λ_{obs}	A_1 (%)	k_{obs1} (s^{-1})	A_2 (%)	k_{obs2} (s^{-1})	A_3 (%)	K_{obs3} (s^{-1})	
syNOS$_{Red}$(-Ca^{2+})	450	89	178 (± 0.78)	11	4.12 (± 0.03)	-	-	This study
syNOS$_{FL}$(-Ca^{2+})	450	63	2.38 (± 0.03)	37	0.004 (± 0.001)	-	-	This study
nNOS$_{Red}$(-CAM)	458	28	57 (± 1)	52	5.2 (± 0.1)	20	570 (± 41)	Craig et al.[17]
nNOS$_{Red}$(+CAM)	458	44	608 (± 66)	56	86 (± 2)	-	-	Craig et al.[17]
CYPOR	450	90	42 (± 4)	10	3.8 (± 0.5)	-	-	Hamdane et al.[68]

1170

Figure S16. Flavin reduction by NADPH. (A) Table of flavin reduction rate constants for syNOS,

nNOS and CYPOR(n = 4). (B) Surface representation and model of NADP$^+$ (magenta) bound in

the pocket of syNOS-Mon with loop 1050-1062 (orange-red and circled in black) in comparison

to (C) nNOS bound to NADP$^+$ (grey) with corresponding loop 1002-1012 (aqua) circled Iin black.

(D) Overlay of NADP$^+$ bound conformations of syNOS, nNOS and CYPOR W677X (1-676) (PDB:

1JA0) (green) over a consensus position of FNR-bound FAD (yellow). syNOS CTT residue 1440

blocks access of the nicotinamide ring to the isoalloxazine ring.

1178

1179

1181

Figure S17. Isothermal calorimetric titration of Ca^{2+} with syNOS$_{C2}$ (residues 1-138). (A)

Endothermic, baseline corrected isotherm of Ca^{2+} titrated into syNOS$_{C2}$. (B) Binding model fit

with the accompanying thermodynamic parameters. (Samples were measured once)

Figure S18. SyNOS globin-based NOD activity and heme reduction rates. (A) NOD activity was

measured by O2 consumption in the presence of the NO-donor NOC-7 and NADPH for WT and

syNOS$_{\Delta C2}$ (n=3, except for syNOS$_{\Delta C2}$ + Ca^{2+} ; n = 2). (B) Heme reduction rate constants of

syNOS$_{Glb}$ in the presence or absence of Ca^{2+} for WT or syNOS$_{\Delta C2}$ and the truncated

syNOS$_{Glb}$/syNOS$_{FNR}$ couple in the absence of Ca^{2+}, with standard error represented. Statistical

comparisons between groups were performed with a 2-way ANOVA with post hoc analysis using

Tukey's multiple comparisons test, all comparisons were non-significant. (C) Michelis-Menten

plot of truncated syNOS$_{Glb}$ reduced by varied concentrations of truncated syNOS$_{FNR}$ until

saturation, with standard deviation represented. (Each data point reflects at least n = 3.)

1194

1195

Figure S19. Output state predicted by AlphaFold3 with FMN-to-Trp715-to-Heme distances

shown. Overlay of the C2 in the locked state (green) with the output Fld conformation (yellow)

illustrates that the Fld is sterically blocked by the C2 domain in the NOS-Asym state. The AF3

prediction was made with a minimal unit of two truncated syNOS chains: 1) residues 473-863

(Oxy) and 2) residues 473-1003 (Oxy + Fld). Including additional domains did not predict

interaction between syNOS$_{Fld}$ and syNOS$_{Oxy}$.

1202

1203

1204

1205

1206

1207

1208

1209

1210

1211

1212

1213

1214

1215