Structure and dynamics of a muti-domain nitric oxide synthase regulated by a C2 domain

Dhruva Nair and Brian R. Crane[*]

AUTHOR ADDRESS:
Weill Institute for Cell and Molecular Biology
Department of Chemistry and Chemical Biology
Cornell University, Ithaca, NY 14853, USA.

*For correspondence: bc69@cornell.edu

Short Title: syNOS is a C2-domain-regulated multi-domain NOS.

Abstract

Nitric oxide synthase (NOS) is a widely studied multi-domain redox enzyme that produces the key signaling molecule and cytotoxic agent nitric oxide (NO) for functions that range from mammalian vasodilation to prokaryotic antibiotic resistance. NOS enzymes from metazoans and cyanobacteria rely on dynamic associations of their oxygenase and coupled di-flavin reductase domains that have largely evaded detailed structural characterization. CryoEM studies of a representative dimeric six-domain Synechococcus NOS reveal the architecture of the full-length enzyme, which contains an unusual regulatory C2 domain, and additional nitric oxide deoxygenase (NOD) and pseudo-globin modules. Five distinct structural states depict how pterin binding couples to tight and loose oxygenase conformations and how the Ca^{2+}-sensitive C2 domain moves over 85 Å to alternatively regulate either the NOS or NOD heme center. The extended C-terminal tail and its stabilizing interactions highlight an added layer of regulation required by multi-domain NOSs compared to other di-flavin reductases.

Teaser: syNOS is a highly dynamic multi-domain oxidoreductase that harnesses a Ca^{2+}-sensitive C2 domain to modulate activity.

Introduction

Nitric oxide (NO) is a potent free-radical signal and cytotoxic agent found throughout biology ([1–8]). Nitric Oxide Synthases (NOS) are homodimeric P450-like enzymes that produce NO through the 5-electron oxidation of L-arginine (L-arg) to L-citrulline ([9–12]). NOSs comprise a heme-containing oxygenase domain (NOS_{Oxy}) that is reductively activated by either an appended reductase domain (NOS_{Red}, characteristic of metazoans) or an auxiliary reductase partner (characteristic of prokaryotes) (Fig 1A) ([10–12]).

Mammalian NOSs (mNOSs) are multi-domain enzymes composed of an N-terminal NOS_{Oxy} and a C-terminal NADPH-dependent NOS_{Red}, bridged by a calmodulin (CAM)-binding domain (NOS_{CBD}; Fig. 1A). NOS_{Red} belongs to the cytochrome P450 reductase (CYPOR) family and as such, is further divided into a NOS_{Oxy}-proximal FMN-binding flavodoxin domain (NOS_{Fld}) and an FAD- and NADPH-binding ferredoxin-NADP$^+$ reductase module (NOS_{FNR}) ([10–13]). NOS_{Oxy} utilizes tetrahydrobiopterin (BH_4) or tetrahydrofolate for rapid heme reduction during oxygen activation ([14]). To reduce heme for oxygen binding, electrons flow from NADPH -> FAD -> FMN(H)• -> Heme-Fe^{3+}. Ca^{2+}- CAM enhances heme reduction by facilitating the traverse of NOS_{Fld} ~70 Å from a shielded position in a conformationally locked state of NOS_{Red} to a productive binding interaction with NOS_{Oxy} ([15–18]).

Given their medicinal importance ([19]), NOSs have been extensively studied in mammals, which have three isoforms: 1) endothelial NOS (eNOS, regulation of the vasculature), 2) neuronal NOS (nNOS, regulation of neuronal function), and 3) inducible NOS (iNOS, immune responses to pathogens and cancer cells) ([11], [20–22]). Most prokaryotic NOSs have fewer domains than their metazoan counterparts and have diverse functions that include participation in stress responses, antibiotic resistance and biosynthesis of secondary metabolites ([23–27]).

To mitigate NO cytotoxicity, prokaryotes oxidize NO to NO_3^- with flavohemoprotein (Hmp), which combines a heme-reducing flavin reductase with an oxygen-activating globin domain. In contrast, mammals oxidize NO with hemoglobin, cytoglobin and neuroglobin, which obtain reduced equivalents from various environmental factors ([28–30]).

Cyanobacterial NOSs represent a bridge between metazoan and prokaryotic NOSs (*26, 31–34*).

They comprise an mNOS architecture fused to an N-terminal globin domain, as exemplified by

the 1468-residue NOS from *Synechococcus sp*. PCC 7335 (i.e. syNOS, Fig. 1A). Like mNOS, Ca^{2+}

activates syNOS, despite syNOS lacking a CBD and Synechococcus lacking an obvious CAM

homolog (*34*). The syNOS globin domain (NOS_{Glb}) oxidizes NO to NO_3^- as a nitric oxide

dioxygenase (NOD) and, when considered with $syNOS_{FNR}$, resembles Hmp (*34*).

Ca^{2+}-CAM regulates mNOS activity by controlling the motions and associations of NOS_{Oxy} and

NOS_{Red} that allow electron transfer (ET) between them. Owing to the dynamic nature of

turnover and inactivated states of full-length NOSs, they have evaded characterization by direct

structural methods. Single particle electron microscopy (EM) in conjunction with chemical

crosslinking and hydrogen-deuterium exchange mass spectrometry (HDX-MS) have yielded

informative, if incomplete, structural information (*35–39*). These studies, along with

biochemical and spectroscopic characterization (*12, 40–44*) have motivated a model of mNOS

conformational dynamics that involves three states: i) input, wherein NOS_{Fld} closely associates

with NOS_{FNR} so that the NOS_{Red} FAD can be reduced by NADPH; ii) intermediate, wherein NOS_{Fld}

transitions between its closed position next to NOS_{FNR} and an interaction with NOS_{Oxy}; and iii)

output, wherein Ca^{2+}-CAM promotes docking of NOS_{Fld} onto the NOS_{Oxy} of the adjacent subunit

(*44–47*). Although cryoEM has provided residue-level information on NOS_{Oxy}, NOS_{Red} positioning

has been challenging to discern owing to its conformational heterogeneity (*35, 46*).

Herein, we resolve domain configurations of the multi-domain syNOS by cryoEM in conjunction

with further characterizing ET properties associated with its NOS and NOD activities. We

capture resting and dynamic states of the enzyme and identify an atypical C2-like domain that

acts as a reversible lynchpin to mediate Ca^{2+} sensitivity. An elaborated C-terminal tail (CTT) (*22*)

of NOS_{Red} , key for regulation of mNOS, maintains a conformationally locked dimer and

represents an expansion of the regulatory mechanisms seen within the di-flavin reductase

family.

Results and Discussion

<u>SyNOS expression and purification</u>

Full-length *Synechococcus PCC 7335* NOS was fused to an N-terminal-twin strep affinity tag and co-expressed along with *E. coli* HSP90 and GAPDH, in BL21(DE3) and purified through a series of affinity and size exclusion columns (SEC) (see Methods, Fig. S1) (*46*, *48*, *49*). As found previously (*34*), syNOS purified as a mixture of monomers and dimers. SEC fractions were tested for NO production with L-arg and NADPH and active fractions were applied to glow-discharged Cu-Quantifoil grids, vitrified and imaged on either a Talos Arctica or Titan Krios. Prior to freezing, samples were incubated with or without Ca^{2+}/L-arg/NADPH (Table S1).

<u>Structure determination of inactivated full-length syNOS</u>

SyNOS data sets and the states within are summarized in Table S1 and described below. Two datasets of syNOS samples lacking Ca^{2+}, L-arg and NADPH were collected on a Talos Arctica equipped with a K3 detector. Dataset (DS) 1 (1215 micrographs) and DS2 (4055 micrographs) were initially preprocessed using CryoSPARC (Fig. S2 and S3), then subjected to particle picking using CRYOLO and finally refined in CryoSPARC (*50*, *51*). During 2D classification multiple classes emerged resembling a symmetric full-length dimer (NOS-Sym). DS1 produced a NOS-Sym overall resolution of 3.8 Å (47 K particles), and DS2 produced a NOS-Sym map of 4.7 Å resolution (84 K particles).

In addition, an asymmetric class (NOS-Asym) became apparent in both DS1 and DS2 that lacked density for one syNOS$_{Red}$ and one syNOS$_{C2}$. DS1 produced a NOS-Asym map with the nominal resolution of 3.67 Å (67 K particles), whereas DS2 yielded a 4.00 Å resolution map (265 K particles). In NOS-Asym, uninterpretable electron density surrounds the position of the vacated syNOS$_{Red}$ indicative of conformational disorder for this subunit.

3D-refinement of NOS-Sym gave clear electron density for secondary structure, cofactors and in many cases, residue side chains. Portions of each subunit were less well defined yet still allowed unambiguous placement of the constituent domains. Inspection of the electron density indicated that the homodimer interface was structurally heterogeneous. To better reconstruct

122 the full-length dimer, we combined the NOS-Sym particle stacks from DS1 (47 K particles) and

123 DS2 (84 K particles) and re-refined the density alignment for 3D-variability (VA) and 3D-Flex

124 analysis. We explored C1, C2 and C2 (relaxed) symmetries which yielded 3.99 Å, 3.95 Å and 3.95

125 Å nominal resolutions, respectively, with the C2(relaxed) refinement chosen for further

126 flexibility analysis (Fig. S4, discussed below). The C2 symmetry-imposed electron density was

127 sufficient for model building and refinement (Table S2).

129 <u>Structure determination of turnover syNOS</u>

130 DS3 (Fig. S5) (5019 micrographs) represented syNOS incubated with Ca^{2+}, BH_4, L-arg and NADPH

131 and was collected on a Titan Krios equipped with a K3 detector. Micrographs were initially

132 preprocessed using CryoSPARC (Fig. S5), then subjected to particle picking with CRYOLO and

133 finally refined in CryoSPARC. During 2D classification, multiple classes emerged resembling NOS-

134 Asym but also a state that lacked density for $syNOS_{Red}$ on both subunits. The latter state

135 resembles previously determined nNOS cryoEM structures that resolved a NOS_{Oxy} homodimer

136 but lacked density for $nNOS_{Red}$ owing to conformational disorder (*36*). We refer to this class as

137 the a "loose" turnover state (NOS-TO$_L$), because the NOS_{Oxy} interface is expanded relative to a

138 typical "tight" $mNOS_{Oxy}$ dimer. 2D classes of NOS-TO$_L$ gave density plumes at the interface

139 characteristic of poor signal alignment, indicative of a conformationally heterogeneous particle,

140 hence a composite map from two local refinements was created. NOS-Asym was refined to a

141 nominal resolution of 3.19 Å (71 K particles), whereas NOS-TO$_L$ was resolved to 3.6 Å resolution

142 (61 K particles, Table S2).

144 DS4 was collected from the same conditions on the 200 KV Talos Arctica equipped with a K3

145 detector. These grids, which had thicker crystalline ice, produced an intermediate resolution

146 turnover state with a tight NOS_{Oxy} dimer interface (NOS-TO$_T$) that more closely resembles crystal

147 structures of BH_4-bound $mNOS_{Oxy}$ (Fig. S6).

Structure determination of monomeric syNOS in turnover conditions

DS5 (9036 micrographs) of syNOS incubated with BH_4, Ca^{2+}, L-arg and NADPH was collected on a

Titan Krios equipped with a Falcon 3 detector (Fig. S7). For this dataset, the homodimers

appeared primarily dissociated into monomers (NOS-Mon). Unlike NOS-Asym, in NOS-Mon,

even NOS_{Oxy} appeared as a monomer. The micrographs were processed through CryoSPARC with

multiple rounds of 3D particle stack cleaning to remove damaged particles (Fig. S7). The syNOS

monomer density was refined to 3.09 Å resolution (45 K particles), with discernible sidechains

and cofactors, although neither Ca^{2+}, nor L-arg could be identified in the map (Table S2).

Full-length syNOS ($syNOS_{FL}$) molecular architecture

In the absence of Ca^{2+}, the NOS-Sym state of the full-length enzyme ($syNOS_{FL}$) forms a compact

C2-symmetric homodimer composed of six domains per subunit (Fig. 1B,C,D); the domain

arrangements divide the protein into two redox modules: i) an Fld-FNR module with an N-

terminal C2 domain (residues 1-134) bound at the Oxy-FNR interface and ii) a Glb-Oxy module,

that also contains an 8-helical pseudo-globin domain that does not bind heme ($syNOS_{pGlb}$,

residues 179-339). Three extended linkers connect the most mobile domains, with L1 (44

residues) connecting NOS_{C2} to NOS_{pGlb}, L2 (10 residues) connecting NOS_{Oxy} to NOS_{Fld} and L3 (13

residues) connecting NOS_{Fld} to NOS_{FNR} (Fig. S8).

In this symmetric "locked" state, syNOS dimerizes through interactions of $syNOS_{Oxy}$ (residues

474-852), and $syNOS_{pGlb}$, which bridges the gap between $syNOS_{Glb}$ (residues 340-473) and

$syNOS_{Oxy}$ (Fig. 1B, C). $SyNOS_{pGlb}$ lacks discernible density for an insertion between αD/ αE

(residues 245-263), which would superimpose on the position of the zinc-binding site found in

the $mNOS_{Oxy}$ dimer; the zinc site is not contained in syNOS (Fig. S9 A, B, C). The $syNOS_{Oxy}$ C-

terminal helical hairpin acts as an interaction hub at the center of the subunit by contacting

NOS_{Fld}, NOS_{FNR}, NOS_{Glb} and the C2 domain.

All structures, except for NOS-TO_T, lacked density for the helical lariats or BH_4, which compose

the center of the NOS_{Oxy} dimer interface (Fig. S9). However, 3D variable analysis (3DVA) and 3D

Flexible refinement (3DFlex) conducted on NOS-Sym, revealed that syNOS subunits "breathe"

between tight, ordered associations that bind BH_4 within the helical lariats and more separated

looser associations with weak density for these regions (Movie S1), not unlike the "tight" and

"loose" dimer states previously observed for *B. subtilis* NOS (*52–54*). Samples collected in

thicker ice favored the tight association (NOS-TO$_T$) for reasons unclear. In addition to an ordered

NOS$_{Oxy}$ interface and density for BH_4, NOS-TO$_T$ displayed new contacts between L1 and αA

(syNOS$_{pGlb}$) on one subunit and α9 (syNOS$_{Oxy}$) on the opposing subunit. Hence, L1 may influence

NOS$_{pGlb}$ positioning, which in turn propagates structural stability to the dimer interface (Fig.

S9B).

The previously unannotated and uncharacterized N-terminal domain (residues 1-134) takes the

form of a tightly packed 8-stranded β-sandwich, capped by peripheral loops at each end (Fig.

1C, D). This structure strongly resembles a C2 domain (syNOS$_{C2}$), which is usually associated

with Ca^{2+}-dependent membrane targeting in eukaryotes and typically not found in prokaryotes.

(*55, 56*) SyNOS$_{C2}$ brackets the backside of the syNOS$_{Oxy}$ heme pocket and syNOS$_{FNR}$ (residues

1017-1468; Fig. 1C). In NOS-Sym, syNOS$_{FNR}$ latches onto syNOS$_{Glb}$ through the ordered CTT,

which also binds to syNOS$_{Fld}$ (residues 863-1003; Fig. 1B, C).

SyNOS$_{Fld}$ and syNOS$_{FNR}$ comprise a canonical NOS reductase unit in a closed conformation with

the FMN-FAD isoalloxazine cofactors aligned in a planar end-to-end configuration (Fig. 2). To

facilitate ET between subunits, mNOSs rearrange to position NOS$_{Fld}$ domain < 15 Å from the

syNOS$_{Oxy}$ heme on the opposing subunit (*57*). Residue conservation, cross-linking and HDX-MS

indicate that NOS$_{Fld}$ interacts dynamically on the backside of the heme pocket at a conserved

Trp residue (356 in human eNOS) (*36–38, 54*). SyNOS$_{C2}$ blocks this position and would have to

move to allow productive ET from syNOS$_{Fld}$, which also must displace from its shielded

interaction with syNOS$_{FNR}$ (*38*). Notably, syNOS lacks the mNOS CBD (~30 residues) that upon

CAM binding favors the ET competent 'output state' (Fig. 2A-C) (*17, 38, 58, 59*). However, the

syNOS interdomain linkers also likely mediate domain motions. These linkers generally have

extended conformations, with the longest, L1, traversing ~60 Å. The threading of L1 beneath L2

implies that any C2 movement will displace L2 and thereby cause syNOS$_{Fld}$ to move "up-and-

over" toward the adjacent syNOS$_{Oxy}$ subunit (Fig. S8). In comparison to the interdomain

interactions implied by the 25 Å resolution nNOS + CAM structure by negative-stain EM (*37*),

syNOS$_{Fld}$ likely undergoes a different trajectory to interact with syNOS$_{Oxy}$ in part because

syNOS$_{Glb}$ will block the CAM-directed conformational change proposed for nNOS.

<u>SyNOS$_{Red}$ is distantly related, but structurally similar, to cytochrome P-450 reductase (CYPOR)</u>

SyNOS$_{Red}$, mNOS$_{Red}$, sulfite reductase diflavin reductase (SIRFP), and methionine synthase

reductase (MTRR) belong to the cytochrome P450 reductase (CYPOR) family of di-flavin

reductases (Fig. S10) (*60–63*). Di-flavin reductases perform three-separate ET reactions i)

hydride transfer from NADPH/NADH to the FNR FAD cofactor ii) sequential ET from the FNR to

the Fld iii) ET from the Fld to an enzymatically relevant electron sink. Activities (i) and (ii) occur

when the di-flavin reductase is trapped in a "closed" conformation permitting first hydride and

then electron transfer along the sequence NADPH->FAD->FMN(H)• with activity (iii) requiring an

"open" conformation that frees the Fld to reduce the electron sink (e.g. NOS$_{Oxy}$ or cytochrome

c).

We sought to place syNOS$_{Red}$ within the genetic and structural context of other di-flavin

reductases by subjecting the syNOS$_{Red}$ protein sequence to clustering, phylogenetic estimation

using maximum likelihood (PhyML) and multiple protein structure alignments (MSTA). Sequence

analyses indicated that syNOS$_{Red}$ and mNOS$_{Red}$ are outgroups relative to CYPORs (Fig. S11); yet,

syNOS$_{Red}$ is more closely related in structure to CYPOR than to nNOS$_{Red}$ or other di-flavin

reductases (Fig. 2 and S10) (*22*). Nevertheless, key differences arise in the structural and

electrostatic features that stabilize the reductase closed conformation in syNOS (NOS-Sym)

compared to CYPOR (Fig. 2A-C).

The di-flavin closed conformation is defined by the L3 loop (situated between the Fld and the

FNR domain, blue in Fig. 2A, B), the Fld domain, and what we define as Interfacing Region (IR) 1,

IR2 and the CTT-IR. IR1 of CYPOR is comprised of an acidic patch surrounding the CYPOR$_{Fld}$ FMN

cofactor and a basic patch surrounding the CYPOR$_{FNR}$ FAD cofactor (Fig. 2C). mNOS$_{Red}$ retains a

homologous acidic/basic IR1 (Fig. 2B). CYPOR and mNOS conserve a hydrogen binding network

between the FNR control region and the Fld (Fig. 2A-C). SyNOS$_{Red}$ lacks electrostatic

complementarity to mediate IR1 and residue complementary to maintain IR2. (In syNOS, IR2 is

composed of Fld residues Gly922/Leu920 binding against Trp1140/Glu1136 on the control

region; whereas in CYPOR, IR2 is composed of Fld Glu213 hydrogen bonding with

Arg382/Thr383). Hence, rather than IR1 and IR2, the CTT-IR and the C2 domain primarily

stabilize the $syNOS_{Red}$ closed conformation (Fig. 3A).

$syNOS_{Red}$ and $mNOS_{Red}$ share a β-finger (syNOS residues 1100-1132) and a CTT that are

important for redox flux in NOS but are not found in canonical CYPORs (Fig. 2A-C) (22, 46).

$syNOS_{Red}$ control region residues 1113-1132, which follow the β-finger, interact with the basic

Fld loop (residues 888-891) to substitute for a CYPOR-like IR2. In syNOS, the β-finger is closer to

the Fld domain (~7.5 Å) compared to in $mNOS_{Red}$ (~17 Å), thereby suggesting either a different

structural role for this element or revealing a conformational state that $nNOS_{Red}$ can also

assume (Fig. 2C). $syNOS_{Red}$ also lacks the $mNOS_{Red}$ Fld autoinhibitory regulatory loop (AI, nNOS

residues 836-849) that destabilizes CAM binding at low levels of Ca^{2+} (Fig 2B) (60, 64). Although

syNOS, $nNOS_{Red}$, and CYPOR stabilize their closed conformations through different interactions,

they all produce a coplanar orientation and similar separation between the FAD and FMN

cofactors (~13 Å from flavin N5-to-N5).

<u>NOS CTTs are enzyme-specific conformational regulators</u>

CTTs, thought to be an mNOS addition within the conserved CYPOR family (65), are 20-30

residue helical extensions that auto-inhibit reduction of NOS_{Fld} by NOS_{FNR} (Fig. 3). Sequence

alignments show no discernable residue similarity between the syNOS CTT and CTTs of the

three mNOS isoforms or that of the most similar eukaryotic NOS sequence from the diatom *T.*

minus NOS (Fig. S11). The lack of sequence similarity between $syNOS_{CTT}$ and $mNOS_{CTT}$, despite

substantial similarity in the preceding $syNOS_{FNR}$ domains, implies convergent evolution of these

CTTs, likely in concert with the addition of $syNOS_{Glb}$.

Unlike previous NOS_{Red} structures that capture only a portion of the CTT, the syNOS structure

resolves the entire CTT and its interactions with $syNOS_{FLd}$ and $syNOS_{Glb}$ (Fig. 3). Compared to

other NOSs, the 27-residue syNOS CTT can be divided into two perpendicular helices (CTT_1 and

CTT_2; residues 1440-1468; Fig. 3A). Several interactions provided by CTT_1 lock the position of

$syNOS_{Fld}$ in the canonical FNR-Fld closed conformation. CTT_1 (residues 1446-1458) interacts with

the FMN ligating loops of NOS_{Fld} (residues 984-989 and 951-960). In this interface, key residue

contacts likely influence ET-competent conformational states of $syNOS_{Fld}$ (e.g. Glu1454 salt-

bridges to Lys987, Tyr1458 hydrogen bonds with Asp984; and Asn1457 hydrogen bonds with

Thr953 (Fig. 3C-D)). Of these residues, CTT_1 Lys987 likely modulates Fld dynamics because it is

flanked by residues that coordinate the FMN pyrophosphate groups (Gln989 and

Glu985). Indeed, the Lys987Ala variant, which disrupts the Lys987/Glu1454 salt-bridge, has a

much more extended, flexible structure by small angle X-Ray scattering (SAXS) than does the WT

(Fig. S12).

The second CTT helix (CTT_2, residues 1460-1468) is absent in other NOSs and CYPORs, likely

because it binds to the atypical NOS_{Glb} (Fig. 3A). Indeed, an aliphatic patch on CTT_2 interacts

with the hydrophobic crevice between residues 358-392 on $syNOS_{Glb}$ (Fig 3A-B). Key interactions

between CTT_2 and NOS_{Glb} include Arg1460 hydrogen bonding to the mainchain carbonyl of

Arg382 and the mainchain carbonyl of Asn1457 hydrogen bonding with the side chain of

Arg382. FNR Lys1412 caps the carboxy-terminus of CTT_2 with a supporting interaction provided

by a hydrogen bond between Trp1464 and the FNR His1412 (Fig. 3E). Thus, CTT_1 and CTT_2

stabilize the associations of NOS_{Fld}, NOS_{FNR}, and NOS_{Glb} domains in the Ca^{2+}-free closed

conformation of syNOS.

$SyNOS_{Red}$ has distinct reactivity toward the surrogate electron acceptor cytochrome c (Cc),

compared to other CYPOR family members, including $mNOS_{Red}$. In the absence of Ca^{2+}, removal

of the CTT does not affect Cc reduction by syNOS (Fig. S13). Ca^{2+} decreases Cc reduction by

almost 4-fold, but, in contrast to the Ca^{2+}-depleted condition, removal of the CTT reactivates Cc

reduction (Fig. S13). In mNOS, CTT removal also activates Cc reduction, but Ca^{2+}-CAM further

increases, rather than decreases activity (*66, 67*). Furthermore, replacing the $mNOS_{FNR}$ domain

with $CYPOR_{FNR}$, which lacks a CTT, switches the Ca^{2+}-CAM dependence of mNOS such that Cc

reduction now decreases ~4-fold with Ca^{2+}-CAM (*66, 67*), much like for syNOS+Ca^{2+}. Thus, the

CTT reduces NOS_{Fld} interaction with Cc in both syNOS and nNOS, but the Ca^{2+} dependence is

opposite. These Cc reduction behaviors among NOS proteins likely reflect differences in the

relative strength of interactions among NOS_{Fld}, NOS_{FNR}, the CTT, NOS_{Oxy} and Cc

itself. Interestingly, syNOS structures in the Ca^{2+}-depleted states (NOS-Sym, NOS-Asym, NOS-

Mon) revealed a stable, closed Fld, whereas the Ca^{2+} turnover states ($NOS-TO_{1,2}$) had a mobile

undiscerned Fld domain, seemingly *more* conducive to Cc reduction, not less. However, in NOS-

TO, the Fld domain may partition more frequently toward NOS_{Oxy}, which would compete with Cc

and thereby reduce Cc reduction rates. In mNOS, relieving a stronger CTT-to-Fld interaction with

Ca^{2+}-CAM may favor the Fld-to-Cc interaction if the Fld-to-Oxy interaction is weaker than in

syNOS.

$SyNOS_{Red}$ has distinct flavin reduction behavior

$SyNOS_{FL}$ and $syNOS_{Red}$ rates of flavin reduction by NADPH, measured through anaerobic

stopped-flow experiments, were biphasic and resembled the flavin reduction kinetics of CYPOR

and $nNOS_{Red}$ bound to Ca^{2+}-CAM (Fig. S14A). The fast phase of $syNOS_{Red}$ reduction that accounts

for 90% of the amplitude change and corresponds to FAD reduction by NADPH has a pseudo

first order rate constant of k_1 ~178 s^{-1}, which is 4-fold greater than that of CYPOR (~42 s^{-1}) (*68*).

The slow phases for syNOS and CYPOR are nearly identical with k_2 ~4 s^{-1}. In contrast, the fast

phase of flavin reduction in $nNOS_{Red}$ (+ Ca^{2+}-CAM) is 3.5-fold greater than that of $syNOS_{Red}$.

Notably NADPH reduces $syNOS_{FL}$ at a ~74-fold slower rate than $syNOS_{Red}$ implying that the

closed conformation of $syNOS_{FL}$ restricts hydride transfer from NADPH to FAD (Fig. S14B-D).

The lower FAD reduction rate in $syNOS_{FL}$ is surprising because $syNOS_{Red}$ lacks an NADP(H)-

ligating Arg (1400 in rat nNOS) known to inhibit electron flow in CAM-free nNOS/eNOS (*69*). In

nNOS, the Arg1400Ser substitution increases flavin reduction while decreasing NO synthesis

(*69*). Because syNOS does not conserve the Arg1400 interaction (the corresponding residues is

an Asn) it would be predicted to have a slower flavin reduction rate than NOS_{Red} (*69*). However,

the NOS_{Glb}-to-CTT interaction, unique to syNOS, may especially stabilize conserved CTT residue

Trp1440 (*70*). As observed in the higher resolution structure of NOS-Mon, Trp1440 blocks the

nicotinamide group of NADPH from approaching the FAD isoalloxazine ring for hydride transfer

(Fig. S14B). The CTT enforced rigidity in syNOS may compensate for the modified $NADP^+$-binding

pocket that lacks an autoinhibitory Arg1400 equivalent. Notably, compared to nNOS and CYPOR

the nicotinamide moiety in syNOS adopts an atypical conformation that is intermediate to the

reduction incapable $NADP^+$ conformation seen in nNOS (PDB:1TLL) and the $NADP^+$/FAD stacked

conformation seen in the CYPOR W677X truncations (PDB:1AJO, Fig. S14D).

<u>Ca^{2+} regulates syNOS activity through a C2-like domain</u>

The lack of a CAM-CBD interaction in syNOS renders its Ca^{2+}sensitivity enigmatic. However,

$syNOS_{C2}$ structurally resembles C2 domains that mediate membrane recruitment by binding

Ca^{2+} (Fig. 4) particularly; mouse perforin C2 (Fig. 4A) (55, 56, 71).

In the closed conformation, $syNOS_{C2}$ forms a tripartite interface with the $syNOS_{Oxy}$ domain and

the FNR control region (Fig. 4B). Two regions of salt-bridges mediate the C2-Oxy interface. The

first (Fig. 4B), involving NOS_{C2} Arg107 and NOS_{Oxy} Glu568, reside beside Trp715, whose analog in

mNOS facilitates interactions important for ET from NOS_{Fld} (38). Asn575 further facilitates the

contact by stabilizing the Arg107 guanidium group for interaction with Glu568. The second

interface (Fig. 4B) involves the NOS_{Oxy} αA-to-αB loop and C2 the β4-β5 connection. C2 Arg57

salt-bridges with Glu523 in an interaction that is buttressed by an amide sidechain of Gln494. A

basic patch on the C2-domain and a complementary acidic patch on α-helix 1162-1173 of the

FNR-control domain wedges the C2 between NOS_{Oxy} and NOS_{Fld} (Fig. 4C). Because $syNOS_{C2}$

blocks the putative Fld-Oxy contact; displacement of $syNOS_{C2}$ would decrease conformational

restraints on $syNOS_{Red}$ (particularly the L2 linker) and relieve steric hindrance on the backside of

the heme pocket.

Removal of the C2 domain ($syNOS_{\Delta C2}$, residues 179-1468) abrogates NOS activity but retains Cc-

reduction activity (Fig. S13). Strikingly, $syNOS_{\Delta C2}$ has a high WT-like Cc-reduction rate that is

uninhibited by Ca^{2+}. In keeping with an enzyme unconstrained by inhibitory interactions, SAXS

data indicates that the global flexibility of $syNOS_{\Delta C2}$ exceeds that of the WT and the more

flexible K987A variant (Fig. S13). Thus, the C2 domain influences redox domain mobility of

syNOS in response to Ca^{2+}, akin to the functional role that Ca^{2+}-CAM plays in the regulation of

mNOSs. In other systems, Ca^{2+} binding to C2 domains neutralizes acidic regions to allow

association with negatively charged phospholipid head-groups (*72*). Although, syNOS_{C2} lacks

the typically conserved Asp residues that coordinate Ca^{2+}, isothermal calorimetry (Fig. S15)

indicated that the syNOS_{C2} binds Ca^{2+} with a 1.6 μM Kd, which compares to the Ca^{2+} affinity of

PLDβ C2 domains (0.8 μM) (*73*).

<u>Ca^{2+} causes a large movement of the C2 domain</u>

In the presence of Ca^{2+} and L-arginine, syNOS adopts a more flexible state, wherein most of

NOS_{Red} is undiscerned by cryoEM ($\text{NOS-TO}_{L,T}$). In this turnover state, the C2 domain has moved

away from the Oxy/FNR interface and now binds at the entrance of the syNOS_{Glb} heme pocket,

nearly 85 Å from its original position (Fig. 5A). The new C2 position blocks the syNOS_{Glb} heme-

binding center and substantially shifts the αB-D helices and constituent residues that compose

the distal heme pocket where NO is oxidized (Fig. 5B). The highly conserved C2 basic patch

(Lys35, Lys37 and Arg81) that formally bound NOS_{FNR} in the locked dimer interacts with the

heme-propionic acid groups of NOS_{Glb} and the 81-89 loop, also conserved among NOS_{C2}

domains, reaches over to block the distal heme pocket. Furthermore, the movement of NOS_{C2}

displaces the L2 linker, which swivels by ~135° and now directs syNOS_{Fld} toward the backside of

the heme pocket on the opposing subunit (Fig. 5C), consistent with the trans ET mechanism

characteristic of mNOSs (*57*). As syNOS_{C2} now resides where the electron-donating FNR

modules of Hmps interact with their cognate globin domains we tested whether removal of the

C2 domain ($\text{syNOS}_{\Delta C2}$) affected syNOS_{Glb} reduction in syNOS_{FL}. Surprisingly, we observed little

change in syNOS_{Glb} reduction rates in $\text{syNOS}_{\Delta C2,}$ nor did Ca^{2+} affect syNOS_{Glb} reduction rates in

syNOS_{FL} (Fig. S14).

syNOS_{FNR} and syNOS_{Fld} have 2-electron reduction potentials characteristic of flavodoxin

domains. Considering that syNOS_{Glb} has a lower potential ($E^{0'}$ = -250 mV) than syNOS_{Fld} ($E^{0'}$ = -

170 mV), but not syNOS_{FNR} ($E^{0'}$ = -290 mV), if a hydroquinone state reduces the globin, syNOS_{FNR}

is likely the better donor. Moreover, a difference of ~40 mV between syNOS_{FNR} and syNOS_{Glb}

(Table S4), a typical reorganization energy of ~ 1.0 eV and a distance of separation of ~31 Å,

suggests that syNOS_{FNR} displaces from its position in NOS-Sym to reduce syNOS_{Glb} at the

observed rates (Fig. 6).(*74*) A modest decrease in syNOS NOD activity with Ca^{2+} suggests an

inhibitory effect of $syNOS_{C2}$ (Fig. S16). $SyNOS_{\Delta C2}$ again showed little change relative to $syNOS_{FL}$,

but note that this variant is also inactive for NO synthesis.   A difference in $syNOS_{Glb}$ reactivity in

$syNOS_{\Delta C2}$ may not be pronounced because the cryoEM data reveals that the protein is never in a

pure conformational state; and the exchange rates among conformational states may exceed

the interdomain reduction rates. In vivo, other factors may stabilize the turnover vs the locked

states of syNOS and produce tighter regulation among them.

<u>The asymmetric state may act as a conformational intermediate between NOS and NOD</u>

<u>activities</u>

The mNOS input, intermediate, and output states are not discrete, but sampled within a

continuum of motion (*18, 35*). In the output state, Fld-CAM moves 70 Å to dock onto an

opposing NOS_{Oxy} domain at the backside of the heme pocket. Similarly, the syNOS flexible

asymmetric dimer class (NOS-Asym) represents an intermediate structural state between the

locked dimer and the turnover state that is likely in equilibrium with NOS-Sym and NOS-TO.   An

Alphafold3 prediction of the pGlb-Glb-Oxy dimer unit with one Fld domain places NOS_{Fld} at the

experimentally validated position on NOS_{Oxy} found by HDX-MS and crosslinking studies of mNOS

(*38*). Upon superimposing this prediction onto either NOS-Asym or NOS-Sym, the Fld binding

surface on NOS_{Oxy} is blocked by the NOS_{C2} (Fig. 5C and S17). Furthermore, compared to Hmp,

$syNOS_{FNR}$ would have to rotate 90º and move 28 Å to reduce $syNOS_{Glb}$ (Fig. 6). This configuration

is likely accessible in the dynamic asymmetric subunit but would not be possible in NOS-TO

because of the rearranged $syNOS_{C2}$. Hence, it is likely that NOS-Asym is capable of $syNOS_{Glb}$

reduction but not $syNOS_{Oxy}$ reduction.

The intermediate conformation of NOS-Asym may be important for the switch between NOS

and NOD activities and explain the inhibitory effect of Ca^{2+} on Cc-reduction. In the absence of

Ca^{2+}, NOS-Sym locks down and inhibits NOS activity, but NOS-Asym allows $syNOS_{FNR}$ to reduce

$syNOS_{Glb}$, and $syNOS_{Fld}$ to reduce Cc. In NOS-Asym, $syNOS_{Fld}$ cannot reduce $syNOS_{Oxy}$ because

$syNOS_{C2}$ blocks $syNOS_{Fld}$ from the $syNOS_{Oxy}$ interaction site on the opposing subunit. Ca^{2+}

promotes the relocation of the C2 domain from $syNOS_{Oxy}$ to $syNOS_{Glb}$; however, some C2

displacement from NOS_{Oxy} may occur in the absence of Ca^{2+} because the C2 domain is not well

defined in the more mobile subunit of NOS-Asym. In addition to preventing $syNOS_{FNR}$ from

reducing the $syNOS_{Glb}$, C2 movement disrupts L2, destabilizing the interaction of NOS_{Fld} with

NOS_{FNR} and favors partitioning of NOS_{Fld} toward the unblocked NOS_{Oxy} and away from Cc (Fig. 7).

Similar to mNOS, syNOS assumes a continuum of conformations that circumscribe defined

enzymatic states. Hence to generate changes in physiologically relevant activities, the relative

weighting of states within the conformational ensemble must be regulated and responsive to

cellular context. It is striking that the $syNOS_{C2}$ moves between a location that blocks $syNOS_{Oxy}$

reduction to one that blocks $syNOS_{Glb}$ reduction, thereby suggesting counter regulation of NO

synthesis and NO oxidation. Such a mechanism may allow the enzyme to tune the balance of

NO production versus NO detoxification, which would be of benefit for signal transduction,

nitrogen metabolism and cellular homeostasis. Overall, these syNOS structural studies

demonstrate how a multi-domain reductase and its associated regulatory components can

undergo large scale rearrangements to modulate reactivity in response to environmental

conditions.

Methods

Molecular cloning

The native coding sequence for full-length syNOS (Uniprot ID : B4WU43**)** was PCR-amplified and

cloned into a modified pet28-a construct between restriction sites NdeI and XhoI using

NEBuilder HiFi DNA Assembly master mix (E2621L). The construct contained sequences for an

N-terminal Twin-Strep Tag with a TEV cleavage site. To produce the hptG and gapA construct,

htpG and gapA were amplified from K-12 *E. coli* and placed into the MCS-1 and the MCS2 of a

tagless pAYC-Duet vector using NEBuilder HiFi DNA Assembly master mix (E2621L). Individual

point mutants were constructed using Kinase-Ligase-DpnI (KLD) method using primers in

supplementary Table 3 (New England Biosciences). All plasmid molecular cloning and plasmid

material generation were carried out with NEB® 5-alpha competent *E. coli* (New England

Biosciences). Successful transformants were validated through Sanger Sequencing or Nanopore

sequencing (Eurofins Genomics).

Protein expression and purification

SyNOS and/or variants were expressed in BL21(DE3) as previously described (*34*) with the

modification of co-expression with hptG/gapA, which improved yield and reproducibility. Cell

pellets were stored at -80 °C prior to purification. The pellet was thawed and resuspended in

lysis buffer (50 mM HEPES (pH 7.5), 150 mM NaCl, 1 mM TCEP, 10% glycerol and 10 uM BH_4),

followed by high-speed centrifugation at 100 000 G and application on Strep-Tactin XT 4-Flow

resin (IBA Life Sciences). Elution of the affinity-bound protein fraction was carried out in batch

with 50 mM biotin supplemented lysis buffer. Post elution samples were concentrated using

Viva-Spin columns (100 kDa cut -off), before application onto 10/300 Superose 6 Increase

Column (Cytiva) using Gel Filtration Buffer 1(GFB1, 25 mM HEPES (pH 7.5), 150 mM NaCl, 1 mM

TCEP, 10% glycerol and 10 uM BH_4).

Size exclusion chromotography - Small angle X-ray scattering (SEC-SAXS)

443   All SEC-SAXS was performed at ID7A1 beamline at Cornell High Energy Synchrotron Source

444   (CHESS). Photon flux was 2.3 x 10^{12} ph/s and scattering data was collected 0.005 Å^{-1} ≤ q ≤ 0.7 Å^{-1}

445   on an EIGER 4M detector (Dectris). Samples were injected into an inline 3.2/150 Superdex 200

446   column in gel filtration buffer. The entire SEC-SAXS trace was captured using a 2s exposure. All

447   SEC-SAXS data processing was performed through RAW (*75*).

Single turnover anaerobic globin reduction

Anerobic syNOS$_{Glb}$ heme reduction assays were carried out using an Agilent 8453 UV-Vis

spectrometer with an attached Hewlett Packard 89090A Peltier temperature controller for

maintaining constant temperature at 20 °C and constant stirring at 400 rpm (Agilent & Hewlett

Packard). Prior to reaction initiation with 1 mM NADPH, syNOS or variants (1 µM) were

incubated anaerobically in the presence of the Glucose Oxidase/Catalase (GOD/CAT) oxygen

scavenging system, as previously described (*76*). Samples were run in either in the presence of 1

mM Ca^{2+} or 1 mM EGTA, as noted. All data was processed using home-made scripts in R Studio

(Posit PBC).

Stopped-flow pre-steady state flavin reduction kinetics

Anerobic stopped-flow mixing of NADPH with $syNOS_{Red}$ or $syNOS_{Fl}$ in the absence of Ca^{2+} was

used to characterize flavin reduction rates. After purification, $syNOS_{Red}$ contains oxidized FAD

and FMN. Attempts to pre-reduce the enzyme to FMNH• with sodium dithionate, DTT and

ascorbate caused precipitation without an observable spectral shift to FMNH•. Thus, flavin

reduction rates reflect conversion from fully oxidized $syNOS_{Red}$. All stopped flow experiments

were carried out anaerobically using a Stopped Flow SX20 instrument (Applied Photophysics).

Samples were pre-incubated anaerobically in the presence of the GOD/CAT oxygen scavenging

system (see above). Changes in absorbance were detected using a single wavelength

Photomultiplier Tube detector. $syNOS_{FL}$ (10 μM) / $syNOS_{Red}$ (40 μM) were mixed with 4x excess

NADPH. All data was processed using home-made scripts in R Studio (Posit PBC).

CryoEM grid preparation and data collection - syNOS ligand free

A 4 μl volume of 1.0 mg ml^{-1} syNOS solution, in GFB1, was applied to glow-discharged grids (200

mesh Quantifoil Cu, R1.2/1.3; Electron Microscopy Sciences). After 10 s of incubation, the

excess solution was blotted for 3.5 s with filter paper by a Vitrobot Mark IV and plunged into

liquid ethane, for vitrification (ThermoFisher). CryoEM images of syNOS were collected on a

Talos Arctica (ThermoFisher) operated at 200 keV at a nominal magnification of 63,000× with a

Gatan GIF Quantum LS Imaging energy filter, using a Gatan K3 direct electron camera in super-

resolution counting mode, corresponding to a pixel size of 1.31 Å. A total of 1214 images stacks

were obtained with a defocus range of −0.8 to −2.0 μm by Serial EM (University of Colorado-

Boulder). Each stack video was recorded for a total dose of 50 electrons per Å^2.

CryoEM grid preparation and data collection - turnover conditions

A 4 μl volume of 1.0 mg ml^{-1} syNOS-ligand free solution in GFB1 was applied to glow-discharged

grids (200 mesh Quantifoil Cu, R1.2/1.3). After 10s of incubation, the excess solution was

blotted for 3.5 s with filter paper by a Vitrobot Mark IV and plunged into liquid ethane for

vitrification (ThermoFisher). CryoEM images of syNOS were collected on a Talos Krios

(ThermoFisher) operated at 300 keV at a nominal magnification of × 165,000 with a Gatan GIF

Quantum LS Imaging energy filter, using a Gatan K3 direct electron CAMera in super-resolution

counting mode, corresponding to a pixel size of 0.730 Å. A total of 5001 images stacks were

obtained with a defocus rage of −0.6 to −2.0 µm by Leginon (*77*). Each stack video was recorded

for a total dose of 44.9 electrons per $Å^2$.

CryoEM grid preparation and data collection - Ca^{2+} and L-arginine conditions

A 4 µl volume of 1.0 mg ml^{-1} syNOS in the presence of 1 mM Ca^{2+} and 1 mM L-arg solution was

applied to glow-discharged grids (300 mesh Quantifoil Au, R1.2/1.3). After 10s of incubation, the

excess solution was blotted for 2.5 s with filter paper by a Vitrobot Mark IV and plunged into

liquid ethane for vitrification (ThermoFisher). CryoEM images of syNOS were collected on a

Talos Krios (ThermoFisher) operated at 300 keV at a nominal magnification of 165,000× with a

Gatan GIF Quantum LS Imaging energy filter, using a Gatan K3 direct electron camera in super-

resolution counting mode, corresponding to a pixel size of 0.422 Å. A total of 5019 images stacks

were obtained with a defocus range of −0.6 to −2.0 µm by Leginon (*77*). Each stack video was

recorded for a total dose of 50.50 electrons per $Å^2$.

CryoEM Image processing and structural reconstruction

All cryoEM density refinement was performed using CryoSPARC (*50*) as illustrated in Figs. S2-S6.

An Alphafold3 prediction was used as the input model by being docked into the appropriate

densities using ChimeraX (*50, 78, 79*). Ligand and poor fitting regions were built and real space

refined using Coot (*79*). The resulting models were passed through ISOLDE (*80*) to approach

thermodynamically rational rotamers/conformations. The outputs from these initial models

were subjected to a series of real-space refinement using PHENIX (*81*) conformational outlier

residues were then adjusted manually using Coot (*79*)

Cytochrome c reduction assay

Cytochrome c (Cc) was used as an electron acceptor to measure the efficiency of electron flow

from NOS_{Red}. Cc accepts an electron from the fully reduced $FMNH_2$ and not from $FADH_2/FADH^-$

$^\bullet$/FAD$^\bullet$/ FMNH$^\bullet$. Cc reduction assays were carried out as previously described (82) using an

Agilent 8453 UV-Vis Spectrometer with an attached Hewlett Packard 89090A Peltier

Temperature Controller for constant temperature at 20 ^{o}C and constant stirring at 400 rpm

(Agilent & Hewlett Packard). All reported reactions were carried out in CC buffer (100 mM Tris

(pH 7.5) and 200 mM NaCl). Cc was pre-oxidized with 1 mM potassium ferrocyanide and then

buffer exchanged into CC buffer. Reactions were conducted with 10 uM Cc and 3 nM of the

sample of interest. Reactions were initiated with 1 mM NADPH and were run either in the

presence of 1 mM Ca^{2+} or 1 mM EGTA as noted. All data was processed using home-made

scripts in R Studio (Posit PBC).

Nitric oxide synthase activity assay

All activity assays were carried out for 30 minutes in 100 mM Tris, (pH 7.5) 200 mM NaCl, 1 mM

DTT and 1 mM L-arginine and initiated with 1 mM NADPH (25 °C). Each sample was evaluated in

the presence of 1 mM Ca^{2+} or 1 mM EGTA. Post reaction, samples were boiled at 70 ^{o}C to halt

NOS activity. NADPH was then precipitated using the $ZnCl_2$/$NaCO_3$ method and subsequently

centrifuged at 15000 G at room temperature. The supernatant was then quantified for total

NO_2^- and NO_3^- content through the Griess assay as implemented in the Nitrate/Nitrite

Colorimetric Assay Kit (Cayman Chemicals).

Oxygen consumption assay to probe Nitric Oxide Dioxygenase activity

The Oxygraph+ system (Hansatech Instruments) was used to measure oxygen consumption of

syNOS in the presence of the NO donor, 3-(2-Hydroxy-1-methyl-2-nitrosohydrazino)-N-methyl-1-

propanamine (NOC-7). 1 µM of enzyme (syNOS$_{FI}$ or syNOS$_{\Delta C2}$) was incubated in buffer

containing 50 mM Tris and 150 mM NaCl at pH 7.5, with 10 µM NOC-7 for 5 minutes. The

reaction was initiated with 1 mM NADPH. Three separate controls were performed to rule out

superoxide or non-enzymatic NO oxidation: NOC-7 + NADPH, NOC-7 + syNOS$_{FL}$ and syNOS +

NADPH.

Isothermal calorimetry

Thermodynamic binding properties between Ca^{2+} and $syNOS_{C2}$ (1-138) were measured on a

Nano ITC Low Volume isothermal titration calorimeter (TA instruments). $SyNOS_{C2}$ was prepared

in a 50 mM CHES and 150 mM NaCl buffer at pH 9.0 at 0.072 mM with 200 μL in the sample cell.

Lower pHs also gave indication of Ca^{2+} binding, but led to aggregation of the C2 domain that

prevented quantitative assessment. $CaCl_2$ was prepared in the same buffer at a concentration of

0.250 mM; 2.5uL was titrated every 250 s for a total of 30 injections. Thermal data were

analyzed using an independent binding model with the NanoAnalyze software.

Reduction potential determination

Reduction potential measurements were performed using a dye calibration method (*83*, *84*),

wherein reduction was driven by a deazaflavin/EDTA photochemical system in a 50 mM $NaPO_3$,

10% Glycerol at pH 7.0, with the following modifications: i) 9 μm 5-deazariobflavin and 5 mM

EDTA was used as the chemical reductant ii) reactions were equilibrated anaerobically in a coy

chamber while supplemented with GOD/CAT oxygen scavenging system. All measurements

were carried out using an Agilent 8453 UV-Vis spectrometer with an attached Hewlett Packard

89090A Peltier Temperature Controller for maintaining constant temperature at 20 ^{o}C and

constant stirring at 400 rpm (Agilent & Hewlett Packard), with the reaction being photoinitiated

with a 15 mW 300 nm laser (ThorLabs) placed above the clear septum capping the cuvette.

Calibrating dyes per domain were selected based on literature values of the published redox

potentials (Table S4). Note that the quoted reduction potentials for $syNOS_{Fld}$ and $syNOS_{FNR}$ will

be the average of the single electron couples between semiquinone and oxidized, and

hydroquinone and semiquinone.

References

References

1.   R. M. Palmer, S. Moncada, A novel citrulline-forming enzyme implicated in the formation of nitric
oxide by vascular endothelial cells. *Biochem Biophys Res Commun* **158**, 348–352 (1989).

2.   B. R. Crane, Bacterial Nitric Oxide Synthases. *Annual Review Of Biochemistry* **79** (2010).

3.   J. A. McCleverty, Chemistry of Nitric Oxide Relevant to Biology. *Chem. Rev.* **104**, 403–418 (2004).

4.   M. Cossenza, R. Socodato, C. C. Portugal, I. C. L. Domith, L. F. H. Gladulich, T. G. Encarnaçao, K. C.
Calaza, H. R. Mendonça, P. Campello-Costa, R. Paes-de-Carvalho, "Nitric Oxide in the Nervous
System: Biochemical, Developmental, and Neurobiological Aspects" in *Nitric Oxide*, G. Litwack, Ed.
(2014)vol. 96 of *Vitamins and Hormones*, pp. 79–125.

5.   C. Farah, L. Y. M. Michel, J. L. Balligand, Nitric oxide signalling in cardiovascular health and disease.
Nature Reviews Cardiology **15**, 292–316 (2018).

6.   J. O. Lundberg, E. Weitzberg, Nitric oxide signaling in health and disease. *Cell* **185**, 2853–2878
(2022).

7.   M. A. Marletta, Revisiting Nitric Oxide Signaling: Where Was It, and Where Is It Going?
Biochemistry **60**, 3491–3496 (2021).

8.   M. Shepherd, D. Giordano, C. Verde, R. K. Poole, The Evolution of Nitric Oxide Function: From
Reactivity in the Prebiotic Earth to Examples of Biological Roles and Therapeutic Applications.
Antioxidants **11** (2022).

9.   K. A. White, M. A. Marletta, Nitric oxide synthase is a cytochrome P-450 type hemoprotein, *ACS
Publications* (2002). https://doi.org/10.1021/bi00144a001.

10.   U. Förstermann, W. C. Sessa, Nitric oxide synthases: regulation and function. *Eur Heart J* **33**, 829–
837, 837a–837d (2012).

11.   M. A. Cinelli, H. T. Do, G. P. Miley, R. B. Silverman, Inducible nitric oxide synthase: Regulation,
structure, and inhibition. *Med Res Rev* **40**, 158–189 (2020).

12.   D. J. Stuehr, M. M. Haque, Nitric oxide synthase enzymology in the 20 years after the Nobel Prize.
British Journal of Pharmacology **176**, 177–188 (2019).

13.   J. Reyes-Rivera, Y. Wu, B. G. H. Guthrie, M. A. Marletta, N. King, T. Brunet, Nitric oxide signaling
controls collective contractions in a colonial choanoflagellate. *Curr Biol* **32**, 2539-2547.e5 (2022).

14.   J. Tejero, D. Stuehr, Tetrahydrobiopterin in nitric oxide synthase. *IUBMB Life* **65**, 358–365 (2013).

15.   D. J. Stuehr, J. Santolini, Z.-Q. Wang, C.-C. Wei, S. Adak, Update on mechanism and catalytic
regulation in the NO synthases. *J Biol Chem* **279**, 36167–36170 (2004).

16.   H. M. Abu-Soud, D. J. Stuehr, Nitric oxide synthases reveal a role for calmodulin in controlling
electron transfer. *Proc Natl Acad Sci U S A* **90**, 10769–10772 (1993).

17. D. H. Craig, S. K. Chapman, S. Daff, Calmodulin Activates Electron Transfer through Neuronal Nitric-
oxide Synthase Reductase Domain by Releasing an NADPH-dependent Conformational Lock *.
Journal of Biological Chemistry **277**, 33987–33994 (2002).

18. N. G. H. Leferink, S. Hay, S. E. J. Rigby, N. S. Scrutton, Towards the free energy landscape for
catalysis in mammalian nitric oxide synthases. *The FEBS Journal* **282**, 3016–3029 (2015).

19. T. L. Poulos, H. Li, Nitric Oxide Synthase and Structure-Based Inhibitor Design. *Nitric Oxide* **63**, 68–
77 (2017).

20. L. Zhou, D.-Y. Zhu, Neuronal nitric oxide synthase: Structure, subcellular localization, regulation, and
clinical implications. *Nitric Oxide* **20**, 223–230 (2009).

21. V. Garcia, W. C. Sessa, Endothelial NOS: perspective and recent developments. *Br J Pharmacol* **176**,
189–196 (2019).

22. N. A. Popova, S. K. Soodaeva, I. A. Klimanov, V. M. Misharin, A. A. Temnov, Autoregulation and
Autoinhibition of the Main NO Synthase Isoforms (Brief Review). *Sovremennye Tehnologii V*
Medicine **15**, 53–59 (2023).

23. M. C. Surdel, B. F. Dutter, G. A. Sulikowski, E. P. Skaar, Bacterial Nitric Oxide Synthase Is Required for
the Staphylococcus aureus Response to Heme Stress. *ACS Infect. Dis.* **2**, 572–578 (2016).

24. I. Gusarov, K. Shatalin, M. Starodubtseva, E. Nudler, Endogenous Nitric Oxide Protects Bacteria
Against a Wide Spectrum of Antibiotics. *Science* **325**, 1380–1384 (2009).

25. B. A. Patel, M. Moreau, J. Widom, H. Chen, L. Yin, Y. Hua, B. R. Crane, Endogenous nitric oxide
regulates the recovery of the radiation-resistant bacterium Deinococcus radiodurans from
exposure to UV light. *Proc Natl Acad Sci U S A* **106**, 18183–18188 (2009).

26. N. Gupta, A. Srivastava, A. K. Mishra, Nitric Oxide Synthases in Cyanobacteria: Diversity, Cellular
Implications and Ecological Pertinence. *J Plant Growth Regul* **44**, 508–528 (2025).

27. T. G. Popova, A. Teunis, H. Vaseghi, W. Zhou, V. Espina, L. A. Liotta, S. G. Popov, Nitric oxide as a
regulator of *B-anthracis* pathogenicity. *Frontiers in Microbiology* **6** (2015).

28. P. R. Gardner, Nitric oxide dioxygenase function and mechanism of flavohemoglobin, hemoglobin,
myoglobin and their associated reductases. *J Inorg Biochem* **99**, 247–266 (2005).

29. P. R. Gardner, A. M. Gardner, L. A. Martin, A. L. Salzman, Nitric oxide dioxygenase: An enzymic
function for flavohemoglobin. *Proceedings of the National Academy of Sciences* **95**, 10378–10383
(1998).

30. M. E. S. Lewis, H. A. Corker, B. Gollan, R. K. Poole, "A survey of methods for the purification of
microbial flavohemoglobins" in *Globins and Other Nitric Oxide-Reactive Proteins, Pt A*, R. K. Poole,
Ed. (2008)vol. 436 of *Methods in Enzymology*, pp. 169–186.

31. F. Del Castello, N. Foresi, A. Nejamkin, C. Lindermayr, F. Buegger, L. Lamattina, N. Correa-Aragunde,
Cyanobacterial NOS expression improves nitrogen use efficiency, nitrogen-deficiency tolerance and
yield in Arabidopsis. *Plant Science* **307** (2021).

32. N. Correa-Aragunde, N. Foresi, F. Del Castello, L. Lamattina, A singular nitric oxide synthase with a
globin domain found in Synechococcus PCC 7335 mobilizes N from arginine to nitrate. *Sci Rep* **8**,
12505 (2018).

33. M. M. Labarthe, G. A. Maroniche, L. Lamattina, C. M. Creus, Nitric oxide synthase expression in
Pseudomonas koreensis MME3 improves plant growth promotion traits. *Appl. Microbiol.*
Biotechnol. **108** (2024).

34. A. L. Picciano, B. R. Crane, A nitric oxide synthase-like protein from Synechococcus produces
NO/NO3- from l-arginine and NADPH in a tetrahydrobiopterin- and Ca2+-dependent manner. *J Biol*
Chem **294**, 10708–10719 (`).

35. M. G. Campbell, B. C. Smith, C. S. Potter, B. Carragher, M. A. Marletta, Molecular architecture of
mammalian nitric oxide synthases. *Proc Natl Acad Sci U S A* **111**, E3614-3623 (2014).

36. D. Felker, K. Lee, T. H. Pospiech, Y. Morishima, H. Zhang, M. Lau, D. R. Southworth, Y. Osawa,
Mapping interactions of calmodulin and neuronal NO synthase by crosslinking and mass
spectrometry. *Journal of Biological Chemistry* **300** (2024).

37. A. L. Yokom, Y. Morishima, M. Lau, M. Su, A. Glukhova, Y. Osawa, D. R. Southworth, Architecture of
the nitric-oxide synthase holoenzyme reveals large conformational changes and a calmodulin-
driven release of the FMN domain. *J Biol Chem* **289**, 16855–16865 (2014).

38. B. C. Smith, E. S. Underbakke, D. W. Kulp, W. R. Schief, M. A. Marletta, Nitric oxide synthase domain
interfaces regulate electron transfer and calmodulin activation. *Proc Natl Acad Sci U S A* **110**,
E3577-3586 (2013).

39. T. Jiang, M. C. Thielges, C. J. Feng, Emerging approaches to investigating functional protein
dynamics in modular redox enzymes: Nitric oxide synthase as a model system. *J. Biol. Chem.* **301**
(2025).

40. C. R. Pudney, D. J. Heyes, B. Khara, S. Hay, S. E. J. Rigby, N. S. Scrutton, Kinetic and spectroscopic
probes of motions and catalysis in the cytochrome P450 reductase family of enzymes. *Febs Journal*
279, 1534–1544 (2012).

41. T. M. Hedison, N. G. H. Leferink, S. Hay, N. S. Scrutton, Correlating Calmodulin Landscapes with
Chemical Catalysis in Neuronal Nitric Oxide Synthase using Time-Resolved FRET and a 5-Deazaflavin
Thermodynamic Trap. *Acs Catalysis* **6**, 5170–5180 (2016).

42. T. M. Hedison, S. Hay, N. S. Scrutton, Trapping methods for probing functional intermediates in
nitric oxide synthases and related enzymes. *Frontiers in Bioscience-Landmark* **23**, 1874–1888
(2018).

43. Y. He, M. M. Haque, D. J. Stuehr, H. P. Lu, Conformational states and fluctuations in endothelial
nitric oxide synthase under calmodulin regulation. *Biophysical Journal* **120**, 5196–5206 (2021).

44.   J. Li, H. Zheng, C. Feng, Deciphering mechanism of conformationally controlled electron transfer in
nitric oxide synthases. *Frontiers in Bioscience-Landmark* **23**, 1803–1821 (2018).

45.   C. J. Feng, Mechanism of nitric oxide synthase regulation: Electron transfer and interdomain
interactions. *Coordination Chemistry Reviews* **256**, 393–411 (2012).

46.   D. J. Stuehr, J. Tejero, M. M. Haque, Structural and mechanistic aspects of flavoproteins: electron
transfer through the nitric oxide synthase flavoprotein domain. *The FEBS Journal* **276**, 3959–3974
(2009).

47.   M. M. Haque, C. Kenney, J. Tejero, D. J. Stuehr, A kinetic model linking protein conformational
motions, interflavin electron transfer and electron flux through a dual-flavin enzyme-simulating the
reductase activity of the endothelial and neuronal nitric oxide synthase flavoprotein domains. *FEBS*
J **278**, 4055–69 (Nov).

48.   R. Chakravarti, K. S. Aulak, P. L. Fox, D. J. Stuehr, GAPDH regulates cellular heme insertion into
inducible nitric oxide synthase. *Proceedings of the National Academy of Sciences* **107**, 18004–
18009 (2010).

49.   Y. Morishima, M. Lau, W. B. Pratt, Y. Osawa, Dynamic cycling with a unique Hsp90/Hsp70-
dependent chaperone machinery and GAPDH is needed for heme insertion and activation of
neuronal NO synthase. *J Biol Chem* **299**, 102856 (2023).

50.   A. Punjani, J. L. Rubinstein, D. J. Fleet, M. A. Brubaker, cryoSPARC: algorithms for rapid
unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290–296 (2017).

51.   T. Wagner, F. Merino, M. Stabrin, T. Moriya, C. Antoni, A. Apelbaum, P. Hagel, O. Sitsel, T. Raisch, D.
Prumbaum, D. Quentin, D. Roderer, S. Tacke, B. Siebolds, E. Schubert, T. R. Shaikh, P. Lill, C.
Gatsogiannis, S. Raunser, SPHIRE-crYOLO is a fast and accurate fully automated particle picker for
cryo-EM. *Commun Biol* **2**, 218 (2019).

52.   B. R. Crane, The enzymology of nitric oxide in bacterial pathogenesis and resistance. *Biochemical*
Society Transactions **36**, 1149–1154 (2008).

53.   K. Pant, A. M. Bilwes, S. Adak, D. J. Stuehr, B. R. Crane, Structure of a nitric oxide synthase heme
protein from Bacillus subtilis. *Biochemistry* **41**, 11071–9 (2002).

54.   B. R. Crane, A. S. Arvai, D. K. Ghosh, C. Wu, E. D. Getzoff, D. J. Stuehr, J. A. Tainer, Structure of nitric
oxide synthase oxygenase dimer with pterin and substrate. *Science* **279**, 2121–6 (1998).

55.   S. Corbalan-Garcia, J. C. Gómez-Fernández, Signaling through C2 domains: More than one lipid
target. *Biochimica et Biophysica Acta (BBA) - Biomembranes* **1838**, 1536–1547 (2014).

56.   T. A. Leonard, "C2 Domain Proteins" in *Encyclopedia of Metalloproteins*, R. H. Kretsinger, V. N.
Uversky, E. A. Permyakov, Eds. (Springer, New York, NY, 2013; https://doi.org/10.1007/978-1-4614-
1533-6_39), pp. 309–318.

57.   U. Siddhanta, A. Presta, B. Fan, D. Wolan, D. L. Rousseau, D. J. Stuehr, Domain Swapping in
Inducible Nitric-oxide Synthase: ELECTRON TRANSFER OCCURS BETWEEN FLAVIN AND HEME

GROUPS LOCATED ON ADJACENT SUBUNITS IN THE DIMER*. *Journal of Biological Chemistry* **273**, 18950–18958 (1998).

58. J. Tejero, M. M. Haque, D. Durra, D. J. Stuehr, A Bridging Interaction Allows Calmodulin to Activate NO Synthase through a Bi-modal Mechanism. *J Biol Chem* **285**, 25941–25949 (2010).

59. P.-F. Chen, K. K. Wu, Structural elements contribute to the calcium/calmodulin dependence on enzyme activation in human endothelial nitric-oxide synthase. *J Biol Chem* **278**, 52392–52400 (2003).

60. E. D. Garcin, C. M. Bruns, S. J. Lloyd, D. J. Hosfield, M. Tiso, R. Gachhui, D. J. Stuehr, J. A. Tainer, E. D. Getzoff, Structural Basis for Isozyme-specific Regulation of Electron Transfer in Nitric-oxide Synthase * [boxs]. *Journal of Biological Chemistry* **279**, 37918–37927 (2004).

61. B. Ghazi Esfahani, N. Walia, K. Neselu, Y. Garg, M. Aragon, I. Askenasy, H. A. Wei, J. H. Mendez, M. E. Stroupe, Structure of dimerized assimilatory NADPH-dependent sulfite reductase reveals the minimal interface for diflavin reductase binding. *Nat Commun* **16**, 2955 (2025).

62. H. Olteanu, R. Banerjee, Human Methionine Synthase Reductase, a Soluble P-450 Reductase-like Dual Flavoprotein, Is Sufficient for NADPH-dependent Methionine Synthase Activation *. *Journal of Biological Chemistry* **276**, 35558–35563 (2001).

63. Z.-Q. Wang, R. J. Lawson, M. R. Buddha, C.-C. Wei, B. R. Crane, A. W. Munro, D. J. Stuehr, Bacterial flavodoxins support nitric oxide production by Bacillus subtilis nitric-oxide synthase. *J Biol Chem* **282**, 2196–2202 (2007).

64. S. Daff, I. Sagami, T. Shimizu, The 42-amino acid insert in the FMN domain of neuronal nitric-oxide synthase exerts control over Ca(2+)/calmodulin-dependent electron transfer. *J Biol Chem* **274**, 30589–30595 (1999).

65. M. Tiso, J. Tejero, K. Panda, K. S. Aulak, D. J. Stuehr, Versatile regulation of neuronal nitric oxide synthase by specific regions of its C-terminal tail. *Biochemistry* **46**, 14418–28 (2007).

66. L. J. Roman, R. T. Miller, M. A. de La Garza, J. J. Kim, B. S. Siler Masters, The C terminus of mouse macrophage inducible nitric-oxide synthase attenuates electron flow through the flavin domain. *J Biol Chem* **275**, 21914–21919 (2000).

67. L. J. Roman, P. Martásek, R. T. Miller, D. E. Harris, M. A. de La Garza, T. M. Shea, J. J. Kim, B. S. Masters, The C termini of constitutive nitric-oxide synthases control electron flow through the flavin and heme domains and affect modulation by calmodulin. *J Biol Chem* **275**, 29225–29232 (2000).

68. D. Hamdane, C. Xia, S.-C. Im, H. Zhang, J.-J. P. Kim, L. Waskell, Structure and Function of an NADPH-Cytochrome P450 Oxidoreductase in an Open Conformation Capable of Reducing Cytochrome P450*. *Journal of Biological Chemistry* **284**, 11374–11384 (2009).

69. M. Tiso, D. W. Konas, K. Panda, E. D. Garcin, M. Sharma, E. D. Getzoff, D. J. Stuehr, C-terminal Tail Residue Arg1400 Enables NADPH to Regulate Electron Transfer in Neuronal Nitric-oxide Synthase *. *Journal of Biological Chemistry* **280**, 39208–39219 (2005).

70. S. Adak, M. Sharma, A. L. Meade, D. J. Stuehr, A conserved flavin-shielding residue regulates NO
synthase electron transfer and nicotinamide coenzyme specificity. *Proceedings of the National*
Academy of Sciences **99**, 13516–13521 (2002).

71. E. A. Nalefski, J. J. Falke, The C2 domain calcium-binding motif: structural and functional diversity.
Protein Sci **5**, 2375–2390 (1996).

72. N. Verdaguer, S. Corbalan-Garcia, W. F. Ochoa, I. Fita, J. C. Gómez-Fernández, Ca2+ bridges the C2
membrane-binding domain of protein kinase Cα directly to phosphatidylserine. *The EMBO Journal*
18, 6329–6338 (1999).

73. L. Zheng, R. Krishnamoorthi, M. Zolkiewski, X. Wang, Distinct Ca2+ Binding Properties of Novel C2
Domains of Plant Phospholipase Dα and β *. *Journal of Biological Chemistry* **275**, 19700–19706
(2000).

74. C. C. Moser, J. M. Keske, K. Warncke, R. S. Farid, P. L. Dutton, Nature of biological electron transfer.
Nature **355**, 796–802 (1992).

75. J. B. Hopkins, R. E. Gillilan, S. Skou, BioXTAS RAW: improvements to a free open-source program for
small-angle X-ray scattering data reduction and analysis. *J Appl Cryst* **50**, 1545–1553 (2017).

76. C. E. Aitken, R. A. Marshall, J. D. Puglisi, An Oxygen Scavenging System for Improvement of Dye
Stability in Single-Molecule Fluorescence Experiments. *Biophys J* **94**, 1826–1835 (2008).

77. A. Cheng, C. Negro, J. F. Bruhn, W. J. Rice, S. Dallakyan, E. T. Eng, D. G. Waterman, C. S. Potter, B.
Carragher, Leginon: New features and applications. *Protein Sci* **30**, 136–150 (2021).

78. J. Abramson, J. Adler, J. Dunger, R. Evans, T. Green, A. Pritzel, O. Ronneberger, L. Willmore, A. J.
Ballard, J. Bambrick, S. W. Bodenstein, D. A. Evans, C.-C. Hung, M. O'Neill, D. Reiman, K.
Tunyasuvunakool, Z. Wu, A. Žemgulytė, E. Arvaniti, C. Beattie, O. Bertolli, A. Bridgland, A.
Cherepanov, M. Congreve, A. I. Cowen-Rivers, A. Cowie, M. Figurnov, F. B. Fuchs, H. Gladman, R.
Jain, Y. A. Khan, C. M. R. Low, K. Perlin, A. Potapenko, P. Savy, S. Singh, A. Stecula, A.
Thillaisundaram, C. Tong, S. Yakneen, E. D. Zhong, M. Zielinski, A. Žídek, V. Bapst, P. Kohli, M.
Jaderberg, D. Hassabis, J. M. Jumper, Accurate structure prediction of biomolecular interactions
with AlphaFold 3. *Nature* **630**, 493–500 (2024).

79. P. Emsley, K. Cowtan, Coot: model-building tools for molecular graphics. *Acta Crystallogr D Biol*
Crystallogr **60**, 2126–32 (2004).

80. T. I. Croll, ISOLDE: a physically realistic environment for model building into low-resolution electron-
density maps. *Acta Cryst D* **74**, 519–530 (2018).

81. (IUCr) Real-space refinement in PHENIX for cryo-EM and crystallography.
https://journals.iucr.org/d/issues/2018/06/00/ic5103/index.html.

82. F. P. Guengerich, M. V. Martin, C. D. Sohl, Q. Cheng, Measurement of cytochrome P450 and
NADPH–cytochrome P450 reductase. *Nat Protoc* **4**, 10.1038/nprot.2009.121 (2009).

83.   Z. A. Maschmann, T. K. Chua, S. Chandrasekaran, H. Ibáñez, B. R. Crane, Redox properties and PAS
775   domain structure of the Escherichia coli energy sensor Aer indicate a multistate sensing
776   mechanism. *J Biol Chem* **298**, 102598 (2022).

777   84.   S. L. Christgen, S. M. Becker, D. F. Becker, Methods for determining the reduction potentials of
778   flavin enzymes. *Methods Enzymol* **620**, 1–25 (2019).

Acknowledgements: We thank Angela Picciano for helpful discussions. We thank the National

Center for CryoEM Access and Training (NCCAT) and the Cornell Center for Material Science

(CCMR) for access to data collection facilities. We thank Yajie Xu and Dr Mariena Silvestry Ramos

for assistance in CryoEM training and data collection.

Funding: This work was supported by grant MCB-2129728 from the National Science

Foundation to BRC. **Author contributions:** D.N. and B.R.C. designed research, conducted

research, analyzed data and wrote the manuscript. Competing interests: The authors declare

that they have no competing interests. Data and materials availability: The cryoEM data and

structures are available at the Protein Data Bank with accession codes: 9Q05, 9Q06, 9Q0Y,

9Q0X, and 9Q15. All other data needed to evaluate the conclusions in the paper are present in

the paper or the Supplementary Materials.

Figures

Figure 1. Architecture of Ca^{2+}-free locked syNOS (A) Gene architecture of Synechococcus NOS,

mammalian NOS and a typical bacterial NOS. Note that some non-cyanobacterial prokaryotic

NOSs have auxiliary domains. (B) Sideview of electron density (left, threshold = 0.0298, C2

symmetry applied) and corresponding ribbon representation (right) for the locked syNOS

homodimer (NOS-Sym). Domains colored as designated with the two subunits distinguished by

the degree of shading. (C) Top view of (B). (D) Individual syNOS domains with cofactors

indicated.

Figure 2. Structural comparison of syNOS$_{Red}$, nNOS$_{Red}$, and CYPOR (A-C). Domain architectures

(top), the corresponding ribbon diagrams (middle), and key interfaces (lower) mediated by

electrostatic interactions for syNOS$_{Red}$ (A), nNOS$_{Red}$ (PDB:1TLL, B) and CYPOR (PDB:1AMO, C)

Surfaces are colored by coulombic electrostatic potential (blue – positive; white – neutral; red –

negative)) with the Fld and FNR domains separated and rotated to view the interface.

Interfacing Regions (IR) 1 (basic-to-acidic patch interactions between the FAD and FMN binding

pockets), IR2 (Fld interaction-to-FNR-control region) and CTT-IR (CTT interaction with Fld) are

denoted by colored circles.

Figure 3. CTT interactions that stabilize the NOS_{Red}-NOS_{Glb} locked state. (A) Ribbon diagram of $syNOS_{Red}$ and $syNOS_{Glb}$ from the boxed region of inactivated $syNOS_{FL}$ (upper left inset) showing key stabilizing interfaces. (B) Aliphatic surface patch between $syNOS_{CTT}$ and $syNOS_{Glb}$ shown by hydrophobicity (yellow) after separating and rotating each domain to view the interface. (C) Fld-to-CTT lock interface mediated by Lys987(Fld)/Glu1454(CTT) salt bridge and Asp984(Fld)/Tyr1458(FNR) hydrogen bond. (D) Fld-to-Glb interface mediated by Asn1457(CTT)/Thr953(Fld) hydrogen bond, Arg1460(CTT)/Arg382-carbonyl(Glb) hydrogen bond and Asn1457-carbonyl(CTT)/arg382(Glb) hydrogen bond. (E) C-Terminal cap interface mediated by Trp1419(CTT)/His1419(FNR) hydrogen bond and a Leu1468-Carboxyl(CTT)/Lys1412 salt bridge.

Figure 4. syNOS$_{C2}$ interactions with NOS$_{FNR}$ and NOS$_{Oxy}$. (A) Superimposition of syNOS (C2, green) with Perforin C2 (PDB: 4Y1S, yellow), C2A-CEP120 (PDB:6EWL, orange) and Munc13-C2B (PDB:6NYT, magenta) as labeled, with root mean square deviation (RMSD) of Cα positions (RMSD), and Template Modeling (TM) scores boxed. B) Ribbon diagram of a syNOS subunit, colored by domain with key C2-Oxy interactions shown in insets b1 and b2. b1) Salt bridge between Arg57(C2)/Glu523(Oxy) with Glu523 stabilized by Gln494. b2) Salt bridge between Arg107(C2)/Glu568(Oxy) with the Arg107 amide stabilized by Asn575. (C) Ribbon diagram of a syNOS subunit showing the electrostatic lock interface between syNOS$_{C2}$ and syNOS$_{Oxy}$ in box inset c1. (c1) C2 rotated 90° (left) and FNR (right, orientation as in (C), colored by electrostatic

849   surfaces (colored: blue-positive, white-intermediate and red –negative). C2 (green, ribbons) is

850   also shown in its original position for reference.

Figure 5. Ca^{2+}-activated C2 mobility. (A) syNOS gene architecture colored by domain, with linkers labeled L1-3(above). Surface representation of three separate syNOS states with relative orientations of the C2 domain. (B) Superimposed surface representations of the turnover state and inactivated states reveal dramatically different C2 positions. (C) L2 in the turnover state (tan electron density) connects to the Fld (yellow ribbons) but then swivels 135° (green density) from its position in the inactivated state to direct the NOS_{Fld} to the adjacent NOS_{Oxy} subunit (purple ribbons of AlphaFold prediction).

Figure 6. syNOS$_{FNR}$ requires a large conformational change to reduce syNOS$_{Glb}$. (A) Cartoon

representation of E.*coli* Hmp (PDB: 1GVH) with the NOD domain (lime green) and FNR (metallic

blue) domains orientated to show the Heme-to-FAD separation of 17.8 Å. (B) syNOS$_{Glb}$ (purple)

and associated syNOS$_{FNR}$ domain (red, control region 1084-1224 not shown) superimposed on

the Hmp Glb (green) and associated FNR (blue). SyNOS$_{FNR}$ would need to rotate 90° and

translate 28 Å to adopt an equivalent Hmp conformation.

Figure 7. Model for NOS to NOD activity switching in syNOS. In the inactivated locked state syNOS$_{Oxy}$ equilibrates between a loose and tight dimer, the latter stabilized by BH$_4$. This state can transition into the asymmetric NOD active state which involves a release of a C2 domain from one subunit and increased mobility of NOS$_{Fld}$ and NOS$_{FNR}$, such that NOS$_{FNR}$ can reduce the syNOS$_{Glb}$ to activate NOD activity. SyNOS$_{Glb}$ reduction coincides with a loosening of the CTT to allow hydride transfer from NADPH to FAD and movement of syNOS$_{FNR}$. In the presence of Ca^{2+}/NADPH/L-arginine syNOS rearranges to facilitate NOS activity. The displacement of NOS$_{C2}$ from NOS$_{Oxy}$ and its binding to NOS$_{Glb}$ relieves constraints on L2 and exposes the Trp715 site such that NOS$_{FNR}$ can relocate to NOSOxy on the opposing subunit and reduce the NOS$_{Oxy}$ heme for NO production.

Structure and dynamics of a muti-domain nitric oxide synthase regulated by a C2 domain

Dhruva Nair and Brian R. Crane

Supplementary Movies

Movie S1: 3D Flexible Refinement of syNOS dimer shows a 'breathing' motion

Supplementary Tables

Table S1: Description of datasets, conditions and refined classes.

Table S2: Cryo-EM data collection, refinement, and validation statistics.

Table S3: DNA primers used in the study with the cloning approach.

Table S4: Reduction potentials ($E^{0'}$) for $syNOS_{Glb}$, $syNOS_{Fld}$ and $syNOS_{FNR}$ vs. the Normal Hydrogen
Electrode (NHE)

Supplementary Figures

Figure S1: syNOS purification

Figure S2: Dataset 1 processing scheme

Figure S3: Dataset 2 processing scheme

Figure S4: Merged data for NOS-Sym state, boxed class used for 3DVA and 3DFlex

Figure S5: Dataset 3 processing scheme and validation.

Figure S6: Dataset 4 processing scheme and validation

Figure S7: Dataset 5 processing scheme and validation.

Figure S8: syNOS linker intertwining.

Figure S9: syNOS tight and loose NOS_{Oxy} states.

Figure S10: Phylogenetic sequence clustering in the context of syNOS (residues 853-1468).

Figure S11: C-terminal tail (CTT) sequence alignment of syNOS, mNOSs and diatom NOS.

Figure S12: SEC-SAXS of syNOS and variants.

Figure S13: Steady state cytochrome c reductase activities of WT syNOS and variants expressed as
turnover numbers (Cc reduced (NOS subunit)$^{-1}$).

Figure S14: Flavin reduction by NADPH.

Figure S15: Isothermal calorimetric titration of Ca^{2+} with $syNOS_{C2}$ (residues 1-138).

Figure S16: syNOS NOD activity and heme reduction rates

Figure S17: Output state predicted by AlphaFold3.

Supplemental Movies

Movie S1. 3D Variability analysis of one subunit of the syNOS asymmetric dimer

showing the breathing motion of NOS$_{Red}$ as a unit relative to motion about the

NOS$_{Oxy}$ interface.

Supplementary Tables

Dataset	Conditions	Class 1	Class 2
1	Heme/FAD/FMN	Symmetric Dimer (NOS-Sym)	Asymmetric Dimer (NOS-Asym)
2			
3	Ca^{2+}/L-arg/BH$_4$	Asymmetric Dimer (NOS-Asym)	Turnover (NOS-TO$_L$)
4	Ca^{2+}/L-arg/BH$_4$	Asymmetric Dimer (NOS-Asym)	Turnover (NOS-TO$_T$)
5	Ca^{2+}/L-arg/ BH$_4$/NADPH	Monomer (NOS-Mon)	none

931 **Table S1.** Description of datasets, conditions and refined classes.

Dataset		Ca2+/L-Arginine			Ca^{2+}/L-Arginine/NADPH
State	Symmetric Dimer	Asymmetric	Turnover	Asymmetric (Composite)	Monomer
Data Collection and Processing					
Microscope	Arctica	Krios			Krios
Voltage (keV)	200	300			300
Detector	K3	Gatan K3			Falcon4EC
Nominal magnification	63,000	105,000			105,000
Data Acquisition Software	Serial EM	Leginon			Leginon
Electron dose (e–/Å2)	50	50.5			44.95
Pixel Size (Å) (binned)	1.31	0.422(0.844)			0.7304
Defocus range (μm)	0.5-2	0.4-2.1			0.4-2.2
Number of movies (#)	1214 + 4055	5019			9036
Number of particles	131,000	70,944	61,553	70,944	45,964
Symmetry imposed	C2	C1	C1	C1	C1
Resolution (Å)	3.95	3.16	3.69	3.16	3.09
FSC threshold	0.143	0.143	0.143	0.143	0.143
Refinement					
Initial model used	PDB 9Q15	PDB 9Q15	PDB 9Q15	Alphafold	PDB 9Q15
Non-hydrogen atoms	22712	16319	11268	16319	11404
Protein residues	2796	2001	1358	2001	1398
Ligand(#)	Heme(4) FAD(2) FMN(2)	Heme(2) FAD(1) FMN(1)	Heme(4)	Heme(4) FAD(1) FMN(1)	Heme(2) FMN(1) / FAD(1) NADPH(1)
RMS Deviation					
Bond lengths (Å)	0.02	0.005	0.003	0.003	0.003
Bond angles (°)	1.55	0.55	0.574	0.485	0.519
MolProbity score	1.54	1.58	1.5	1.74	1.54
Clashscore	2.18	3.49	4.65	4.66	4.93
Poor rotamers (%)	2.08	1.63	0.93	1.8	0.91
Ramachandran Favored (%)	95.1	96.0	96.1	95.5	96.0
Ramachandran Allowed (%)	4.6	3.9	3.7	4.3	3.8
Ramachandran Disallowed (%)	0.3	0.2	0.2	0.2	0.2
Fit to map (CCmask)	0.8	0.79	0.75	0.81	0.82
EMDB (maps)	EMD-72088	EMD-72110	EMD-72109	EMD-72104, EMD-72105,EMD-72116	EMD-72087
PDB (model)	9Q06	9Q0Y	9Q0X	9Q15	9Q05

Table S2. Cryo-EM data collection, refinement, and validation statistics.

Primer Name	Primer	Cloning Approach
pet28_twin_strep_syNOS_fwd	tcgagcaccaccaccacc	Gibson/NEB HiFi
pet28_twin_strep_syNOS_rev	catatgtcctccggattggaagtac	Gibson/NEB HiFi
syNOS_fwd	tccaatccggaggacatatgatgcttgtcaacgactctc	Gibson/NEB HiFi
syNOS_rev	gtggtggtggtggtgctcgagctacaagttggctagcc	Gibson/NEB HiFi
Hptg_fwd_PAYC_mcs1	TAATGCTTAAGTCGAACAG	Gibson/NEB HiFi
Hptg_rev_PAYC_mcs1	GGTATATCTCCTTATTAAAGTTAAAC	Gibson/NEB HiFi
Hptg_fwd	CTTTAATAAGGAGATATACCATGAAAGGACAAGAAACTCGTG	Gibson/NEB HiFi
Hptg_rev	TCTGTTCGACTTAAGCATTATCAGGAAACCAGCAGCTG	Gibson/NEB HiFi
gapa_fwd_pAYC_mcs2	TAATTAACCTAGGCTGCTG	Gibson/NEB HiFi
gapa_rev_pAYC_mcs2	ATGTATATCTCCTTCTTATACTTAAC	Gibson/NEB HiFi
gapa_fwd	TATAAGAAGGAGATATACATATGACTATCAAAGTAGGTATCAAC	Gibson/NEB HiFi
gapa_rev	GCAGCAGCCTAGGTTAATTATTATTTGGAGATGTGAGC	Gibson/NEB HiFi
pet28_tws_delc2_fwd	ctcgagcaccaccaccac	Gibson/NEB HiFi
pet28_tws_delc2rev	catatgtcctccggattggaag	Gibson/NEB HiFi
syNOSdelc2_178_1468_fwd	tccaatccggaggacatatgacctactcagaagccgtc	Gibson/NEB HiFi
syNOSdelc2_178_1468_fwd	tggtggtggtggtgctcgagctacaagttggctagccatttttc	Gibson/NEB HiFi
pet28_tws_delctt_fwd	tagctcgagcaccaccac	Gibson/NEB HiFi
pet28_tws_delctt_rev	catatgtcctccggattggaag	Gibson/NEB HiFi
syNOS_delctt1_1_1440_fwd	tccaatccggaggacatatgatgcttgtcaacgactctc	Gibson/NEB HiFi
syNOS_delctt1_1_1440_fwd	tggtggtggtgctcgagctagtagttgtctttctccacttg	Gibson/NEB HiFi
syNOS_K1412E_fwd	GCAAATTGCTgaaACAGAAGGAG	KLD
syNOS_K1412E_rev	ATAAATACCTCAAACACATTG	KLD
syNOS_T953A_fwd	GATCGGCAGCgcgGTCTATGAAC	KLD
syNOS_T953A_rev	CCTAGTACGGAATAGTTGAG	KLD
syNOS_K987A_fwd	CGACGAAATCgcgGGACAAGCCG	KLD
syNOS_K987A_rev	CCTTTATGTAGGGGAAC	KLD

Table S3. Primers used in the study with the cloning approach (left).

Domain	Couple	Potential (mV)	n	Calibrating dye (mV)
Globin	Fe^{3+}/Fe^{2+}	-250 ± 20	1	safranin T (-289, 12.5μM)
Fld	ox/hq	-173 ± 3	2	cresyl violet (-177.0, 60 μM)
FNR	ox/hq	-290 ± 13	2	safranin T (-289, 12.5 μM)

Table S4. Reduction potentials ($E^{0'}$) for syNOS$_{Glb}$, syNOS$_{Fld}$ and syNOS$_{FNR}$ vs. the Normal Hydrogen Electrode (NHE). For syNOS$_{Glb}$ the single electron reduction (n_e=1) from Fe^{3+} to Fe^{2+} was measured whereas for the flavin-containing domains a midpoint potential for the 2-electron reduction (n_e=2) from oxidized (ox) to hydroquinone (hq) was measured as no spectral signature for the semiquinone was apparent in the titrations. (n = 3 for all measurements)

Supplementary Figures

945 **Figure S1.** syNOS Purification. Representative chromatogram from a Superdex 200 increase
946 10/30 size exclusion chromatography column with an arrow denoting the syNOS dimer peak.
947 The inset corresponds to a Coomassie blue stained 4-12% SDS-PAGE gel of fractions 5,6,7 and 8.

Dataset 1

Figure S2. Dataset 1 processing scheme. (A) CryoEM processing workflow for dataset 1: syNOS

ligand free. NOS-Sym class used for 3DVA and 3DFlex boxed (dashed-red). (B) Representative

2D Classes. CTF, contrast transfer function; NU, non-uniform refinement. (C) Overall Particle and

Gold-standard Fourier Shell Correlation (GSFSC) curves for NOS-Asym and NOS-Sym.

Figure S3. Dataset 2 processing scheme. (A) CryoEM processing workflow for dataset 2: syNOS

ligand free. NOS-Sym class used for 3DVA and 3DFlex boxed (dashed-red). (B) Representative 2D

Classes. (C) GSFSC curves and positional distribution of particles.

Figure S4. Merged data for NOS-Sym state, boxed class used for 3DVA and 3DFlex.

Figure S5. Dataset 3 processing scheme and validation. (A) CryoEM processing workflow for

dataset 3: syNOS in the presence of Ca^{2+}/L-arginine (B) Representative 2D Classes for NOS-Asym

state (C) Representative 2D classes for NOS turnover state (NOS-TO$_L$) (D) Asymmetric states

GSFSC curves, orientation validation, model density fit and map to model Fourier Shell

Correlation (FSC).

Figure S6. Dataset 4 processing scheme and validation. (A) CryoEM processing workflow for

dataset 4, syNOS with NADPH/Ca^{2+} including GSFSC curves. (B) Representative 2D classes for

NOS-TO$_T$.

Dataset 5

A

Micrographs
9036

Patch Motion
Patch CTF
Curate Micrographs

Micrographs
4989

Blob Picker

Particles
909,465

2D Classification

Ab Initio
Hetero Refinement
Rebalance Orientations
Topaz Training + Extraction
2D classification

2x

Ab Initio
Hetero Refinement

Class 1   Class 2   Class 3

Non-Uniform Refinement

y 180

3.09 Å
45K Particles
Monomer

B

150 Å

C

GSFSC Resolution: 3.15Å

No Mask (4.1Å)
Spherical (3.9Å)
Loose (3.4Å)
Tight (3.1Å)
Corrected (3.1Å)

DC   7.3Å   3.7Å   2.4Å   1.8Å

π/2
π/4
0
−π/4
−π/2

Elevation

# of images

−π   −3π/4 −π/2 −π/4   0   π/4   π/2 3π/4   π
Azimuth

D

Map to Model

Unmasked
Masked

FSC

3.4 Å

1/Å

E

Globin Heme   CTT(1440-1468)   NADP+ Binding Site

FMN   FAD   Oxygenase Heme

Figure S7. Dataset 5 processing scheme and validation. (A) CryoEM processing workflow for

dataset 5: syNOS with L-arginine, NADPH/Ca^{2+}/NADPH (B) Representative 2D classes for NOS-

Mon. (C) GSFSC curves, orientation validation and (D) Map to Model FSC. (E) Density fit of

model in map.

Figure S8. SyNOS linker intertwining. Intermediate resolution homodimer cryoEM density with

all three linkers colored the same as their preceding domains. Note how L1 threads beneath L2

and on top of L3. Movement of the C2 domain (green) will release the Fld module for NOS$_{Oxy}$

reduction.

Figure S9. SyNOS tight and loose NOS$_{Oxy}$ states. (A) syNOS cryoEM density (dataset 4) with the NOS$_{Oxy}$ dimer interface outlined (orange and green) to show each subunit. (B) Close up of the dimeric interface showing the close association of NOS$_{Oxy}$ α9 to an intersubunit pGlb C/D loop. Although at lower resolution, strong electron density for the core the NOS-TO$_T$ interface includes the helical lariats and BH$_4$. (C) Loose (green) and tight (blue) NOS$_{Oxy}$ dimers differ by a large 20 Å breathing motion normal to the dimer interface that also affects the positioning of the C2 domain (lower right peripheral electron density).

Cluster ID	Average Sequance Length	Standard Deviation of Sequence Length	Sequence Count
2	678	63	2521
39	702	49	1588
55	1117	103	278
3	1479	118	94
13	976	174	63
58	700	17	49
30	1126	283	47
7	638	35	45
18	601	74	44
33	652	36	44

995 **Figure S10.** Phylogenetic sequence clustering in the context of syNOS (residues 853-1468).

996 Representative clusters were subjected to PhyML with clusters above 40 sequences represented

997 in the tree. (B) Cluster composition by sequence length and count.

1000 **Figure S11.** C-terminal tail (CTT) sequence alignment of syNOS, mNOSs and diatom NOS.

1001

1002

1003 **Figure S12.** SEC-SAXS of syNOS and variants. (A) Dimensionless Kratky plots, binned by 4 qRg

1004 values, for samples in the presence and absence of Ca^{2+}. (B) Table of predicted and

1005 experientially characterized radius of gyrations (Rgs) and molecular weights calculated either by

1006 volumes of correlation (V_C) or Porod volumes (V_P). (Samples were measured once)

1007

1008

Figure S13. Steady state cytochrome c reductase activities of WT syNOS and variants expressed

as turnover numbers (Cc reduced (NOS subunit)$^{-1}$ min^{-1}) with standard errors shown as error

bars (see Methods). Statistical comparisons between groups were performed with a 2-way

ANOVA with post hoc analysis using Tukey's multiple comparisons test. Significant differences

are indicated as follows: ns (not significant), * ($p \leq 0.05$), ** ($p \leq 0.01$), *** ($p \leq 0.001$), **** ($p \leq$

0.0001)..Ca:WT vs. Ca:K987A (ns); Ca:WT vs. Ca:ΔC2 (ns); Ca:WT vs. Ca:K1412E (ns); Ca:WT vs.

Ca:ΔCTT (ns); Ca:WT vs. EGTA:WT (****) ; Ca:WT vs. EGTA:K987A (ns); Ca:WT vs. EGTA:ΔC2 (ns);

Ca:WT vs. EGTA:K1412E (****); Ca:WT vs. EGTA:ΔCTT (****) ;Ca:K987A vs. Ca:ΔC2 (ns);

Ca:K987A vs. Ca:K1412E (***); Ca:K987A vs. Ca:ΔCTT (****); Ca:K987A vs. EGTA:WT (**);

Ca:K987A vs. EGTA:K987A (**); Ca:K987A vs. EGTA:ΔC2 (ns); Ca:K987A vs. EGTA:K1412E (****);

Ca:K987A vs. EGTA:ΔCTT (****); Ca:ΔC2 vs. Ca:K1412E (ns); Ca:ΔC2 vs. Ca:ΔCTT (**); Ca:ΔC2 vs.

EGTA:WT (****); Ca:ΔC2 vs. EGTA:K987A (ns); Ca:ΔC2 vs. EGTA:ΔC2 (ns); Ca:ΔC2 vs.

EGTA:K1412E (****); Ca:ΔC2 vs. EGTA:ΔCTT (****);Ca:K1412E vs. Ca:ΔCTT (ns); Ca:K1412E vs.

EGTA:WT (****); Ca:K1412E vs. EGTA:K987A (ns); Ca:K1412E vs. EGTA:ΔC2 (ns); Ca:K1412E vs.

EGTA:K1412E (**); Ca:K1412E vs. EGTA:ΔCTT (****); Ca:ΔCTT vs. EGTA:WT (**); Ca:ΔCTT vs.

EGTA:K987A (*); Ca:ΔCTT vs. EGTA:ΔC2 (***); Ca:ΔCTT vs. EGTA:K1412E (ns); Ca:ΔCTT vs.

EGTA:ΔCTT (**); EGTA:WT vs. EGTA:K987A (****); EGTA:WT vs. EGTA:ΔC2 (****); EGTA:WT vs.

EGTA:K1412E (ns); EGTA:WT vs. EGTA:ΔCTT (ns); EGTA:K987A vs. EGTA:ΔC2 (ns); EGTA:K987A vs.

EGTA:K1412E (***); EGTA:K987A vs. EGTA:ΔCTT (***); EGTA:ΔC2 vs. EGTA:K1412E (****);

EGTA:ΔC2 vs. EGTA:ΔCTT (****); EGTA:K1412E vs. EGTA:ΔCTT (ns). (n=3, except for K1412E

+EGTA; n = 2).

1030

A

Reductase	λ_{obs}	A_1 (%)	k_{obs1} (s^{-1})	A_2 (%)	k_{obs2} (s^{-1})	A_3 (%)	K_{obs3} (s^{-1})	
syNOS$_{Red}$(-Ca^{2+})	450	89	178 (± 0.78)	11	4.12 (± 0.03)	-	-	This study
syNOS$_{FL}$(-Ca^{2+})	450	63	2.38 (± 0.03)	37	0.004 (± 0.001)	-	-	This study
nNOS$_{Red}$(-CAM)	458	28	57 (± 1)	52	5.2 (± 0.1)	20	570 (± 41)	Craig et al.[17]
nNOS$_{Red}$(+CAM)	458	44	608 (± 66)	56	86 (± 2)	-	-	Craig et al.[17]
CYPOR	450	90	42 (± 4)	10	3.8 (± 0.5)	-	-	Hamdane et al.[68]

Figure S14. Flavin reduction by NADPH. (A) Table of flavin reduction rate constants for syNOS,

nNOS and CYPOR(n = 4). (B) Surface representation and model of NADP$^+$ (magenta) bound in

the pocket of syNOS-Mon with loop 1050-1062 (orange-red and circled in black) in comparison

to (C) nNOS bound to NADP$^+$ (grey) with corresponding loop 1002-1012 (aqua) circled Iin black.

(D) Overlay of NADP$^+$ bound conformations of syNOS, nNOS and CYPOR W677X (1-676) (PDB:

1JA0) (green) over a consensus position of FNR-bound FAD (yellow). syNOS CTT residue 1440

blocks access of the nicotinamide ring to the isoalloxazine ring.

1039

1040

1041

Figure S15. Isothermal calorimetric titration of Ca^{2+} with $syNOS_{C2}$ (residues 1-138). (A)

Endothermic, baseline corrected isotherm of Ca^{2+} titrated into $syNOS_{C2}$. (B) Binding model fit

with the accompanying thermodynamic parameters. (Samples were measured once)

Figure S16. SyNOS globin-based NOD activity and heme reduction rates. (A) NOD activity was

measured by O2 consumption in the presence of the NO-donor NOC-7 and NADPH for WT and

$syNOS_{\Delta C2}$ (n=3, except for $syNOS_{\Delta C2}$ + Ca^{2+} ; n = 2). (B) Heme reduction rate constants of

$syNOS_{Glb}$ in the presence or absence of Ca^{2+} for WT or $syNOS_{\Delta C2}$, with standard error

represented. Statistical comparisons between groups were performed with a 2-way ANOVA with

post hoc analysis using Tukey's multiple comparisons test, all comparisons were non-significant.

1051

1052 **Figure S17.** Output state predicted by AlphaFold3 with FMN-to-Trp715-to-Heme distances shown.

1053 Overlay of the C2 in the locked state (green) with the output Fld conformation (yellow) illustrates that

1054 the Fld is sterically blocked by the C2 domain in the NOS-Asym state.

1055

1056

1057

1058

1059

1060

1061

1062

1063

1064

1065

1066

1067

1068