Gradual Warming Drives Life-History Shifts and Reveals Reproductive Ceilings in

Daphnia magna

Nikola Petkovic[1], Tuğçe Ünlü[1] and İsmail K. Sağlam[1]

[1] College of Sciences, Molecular Biology and Genetics Department, Koç University, Istanbul, Turkey

Abstract

Global warming often pushes species toward their physiological limits, yet the pace of change

may be as important as the absolute temperature reached. Most experiments rely on abrupt

shifts, leaving the demographic and evolutionary consequences of gradual warming poorly

understood. We conducted a 30-generation experiment with clonal populations of *Daphnia*

magna maintained either under constant temperature (26 °C) or gradual warming (+1 °C every

4–6 generations, up to 32 °C). We measured fecundity, growth rate, age at maturation, body

size, and lifespan across generations, and used a reciprocal transplant assay to disentangle

genetic and plastic responses.

Warming populations matured earlier, grew faster, and initially reproduced more than controls,

but suffered shorter lifespans and lost their reproductive advantage near 30 °C, a sub-lethal

ceiling that coincided with extinction events. This rate-dependent trajectory contrasts with one-

step warming studies that typically report uniformly reduced reproduction. Reciprocal transplants

revealed a persistent reduction in adult body size of warming-line offspring, regardless of rearing

environment, consistent with heritable or trans-generational change, while growth rate remained

largely plastic. Together, these results show that gradual warming can simultaneously elicit

plastic and heritable trait shifts, and that reproductive ceilings well below lethal limits act as

tipping points for population persistence.

Keywords: climate change; thermal tolerance; life-history evolution; phenotypic plasticity;

transgenerational effects; reproductive thresholds

1. Introduction

Adverse anthropogenic impacts on the Biosphere have produced environmental change at a rate unprecedented in the Cenozoic (Barnosky et al., 2012). One of the most alarming consequences of the widely recognized issue of global change (Falkowski, 1994; Chivian and Bernstein, 2008; Barnosky et al., 2012) is the shifting of environmental conditions outside of the physiological tolerance range of many species (Parmesan, 2006). Recent studies predict the extinction of 14-35 % of extant species in the following decades (Thomas et al., 2004; Weins and Zelinka, 2024). Hence, understanding the factors which affect adaptation to changing environments has become one of the central problems of contemporary science, with the main goal to help mitigate the detrimental effects of global change. One of the most threatening aspects of global change is the continuous rise of temperature from the pre-industrial era (Hawkins et al., 2017) onwards, a trend predicted to accelerate within the next 50 years (Walther et al., 2002). Consequently, persistence of many extant species may be threatened due to the altered interspecies interactions (for example, food availability) and the limited physiological tolerances to high temperatures (Cahill et al., 2013). The latter factor is particularly pronounced in the lower latitudes, as many tropical species are already experiencing conditions close to their thermal optima (Deutsch et al., 2008). Crucially, population collapse may occur even when lethal limits are not reached, if sub-lethal thresholds for key fitness traits, such as reproduction, are exceeded (Kingsolver and Huey, 2008; Seefeldt and Ebert, 2019).

Predicting the patterns of species' response to climate change requires tracking of the variation in fitness-related traits (i.e. life histories) during exposure to thermally changing environments, because these traits may strongly correlate with the probability of survival (Isaac, 2009). For example, slower life histories (long gestation and small litters) may enhance extinction risk due to the limited capacity of a population to recover from a decline

caused by the natural or anthropogenically induced factors (Purvis et al., 2000). Moreover,

these extinction-promoting traits are often negatively correlated with body size (McKinney

1997; Cardillo et al. 2005). Thus, in species with an intraspecific variation in body size, we

might expect a shift to a smaller body size due to a lower extinction risk (Isaac, 2009).

However, the opposite pattern can be predicted if an intraspecific variation in body size

positively correlates with competitiveness (Isaac, 2009), survival (Yoccoz & Mesnager 1998)

and fecundity (Du et al. 2005).

Given the complex interactions among life history traits, clarifying their relative contributions

to fitness and survival under rising temperatures requires experimental studies that

manipulate temperature and directly monitor trait changes. To date, most of these studies

have been carried out using insects as model organisms or various plankton species

cultivated in the outdoor mesocosms (Van Doorslaer et al., 2007; Van Doorslaer et al., 2010;

Musolin et al., 2010; Ciota et al., 2014; Geerts et al., 2015; Laughton et al., 2017;

Cambronero et al., 2018; Iltis et. al., 2019; de Leon et al., 2023; Albini et al., 2025).

Most experimental studies with insects revealed negative correlation of temperature with

lifespan, development time, and body size (e.g. Ciota et al., 2014; Laughton et al., 2017; Iltis

et. al., 2019). Importantly, these effects of an elevated temperature on the life history traits

may depend on the different phenological parameters such as season. For example, an

increase in temperature during the warmest part of the season was found to reduce

development and increase mortality in the heteropteran species *Nezara viridula*, while adults

showed a decline in body size as well as reduced life span and fecundity (Musolin et al.,

2010). The negative effects of elevated temperature on life history traits may also vary with

geographical distribution of species. In a study of four *Protaphorura* species (Collembola)

sampled across a latitudinal gradient, heat stress reduced fecundity in all species, but the

most severe effects were observed in the boreal species, which was the most heat-sensitive

(de Leon et al., 2023).

Another common experimental approach includes monitoring of freshwater populations

under semi-natural conditions (outdoor mesocosms). Mean body size of the plankton

species showed variation in response to the elevated temperatures, depending on their

cultivation environment. A consistent decline in mean body size has been observed under

temperature increases of up to 8°C compared to ambient conditions, including in both single-

species setups (Cambronero et al., 2018) and more complex multispecies zooplankton

communities (Albini et al., 2025). However, increases in body size at maturity have also

been reported under moderate warming (+4°C), likely due to interactions between abiotic

factors (e.g., temperature) and biotic factors such as interspecific competition (Van

Doorslaer et al., 2010).

In addition, evidence points to latitude-specific differences in how elevated temperature

affects fecundity, with contrasting patterns reported for boreal and temperate species.

Whereas boreal species typically show reduced fecundity at elevated temperatures

(Giovannini et al., 2023), the pattern in temperate-origin species is less consistent, having

been reported as positive (Van Doorslaer et al., 2007), negative (Cambronero et al., 2018)

or more complex (Wang et al., 2023). In contrast, the response of age at maturation to

elevated temperatures appears to be relatively uniform regardless of the environmental

conditions, with experimental studies repeatedly reporting earlier maturation at higher

temperatures (Van Doorslaer et al., 2007; Cambronero et al., 2018; Giovannini et al., 2023;

Wang et al., 2023).

Despite substantial experimental research, significant gaps remain in our understanding of

how life history traits respond to temperature change, particularly because most studies have

focused on abrupt shifts in temperature, typically involving a single-step increase. In reality,

many natural populations experience gradual warming, and the pace of change may be as

critical to outcomes as the absolute temperatures reached (Killeen et al., 2017). To better

reflect ecological reality, we need experimental designs that examine the effects of gradually

and continuously changing thermal environments over longer time scales. This is important

for two reasons. First, the rate and therefore the impact of global warming varies across

regions, showing positive correlation with both latitude (Root et al., 2003; Deutsch et al.,

2008) and altitude (Pepin et al., 2022). Second, experimental studies have shown that the

adaptive dynamics of populations to stressful environments are strongly influenced by the

rate of environmental change (Collins and de Meaux, 2009; Bell and Gonzalez, 2011;

Lindsey et al., 2013; Killeen et al., 2017; Petkovic and Colegrave, 2023). This means that

warming regimes of identical endpoints may produce qualitatively different demographic and

evolutionary trajectories depending on whether the change is abrupt or gradual (Collins and

de Meaux, 2009; Lindsey et al., 2013).

Experimental studies with microbial systems highlight the importance of warming rate for

adaptive dynamics. For example, populations exposed to the most gradual temperature

increases are more likely to persist and evolve higher thermal tolerance, consistent with

predictions of evolutionary rescue (Killeen et al., 2017). Similarly, slower environmental

change can promote growth at high temperatures but may come at a cost to long-term

survival due to correlated selection (Liukkonen et al., 2021). Such findings suggest that

gradual warming can initially enhance certain performance traits yet may eventually drive

collapse once sub-lethal thresholds for reproduction or other key functions are exceeded

(Huey and Kingsolver, 2019). However, such studies have primarily focused on growth rate

in unicellular organisms. We still lack a clear understanding of how more complex life history

traits, such as body size, fecundity, and age at maturation, respond to gradual thermal

change, particularly in multicellular species. Most experimental work on these traits has

been based on abrupt or one-step temperature increases, limiting ecological relevance.

In addition, the extent to which these trait changes result from phenotypic plasticity versus

genetic adaptation remains poorly understood. Organisms may respond to thermal shifts

through adaptive plasticity, adjusting trait expression without genetic change (Lalejini et al.,

2021). For example, *Daphnia magna* populations can modify hemoglobin expression in

response to temperature (Yampolsky et al., 2014). Yet, genetic responses to warming have

also been observed in *D. magna*, including short-term adaptation in laboratory populations

(Seefeldt and Ebert, 2019) and long-term responses in the wild (Geerts et al., 2015).

Importantly, plasticity itself can evolve (Schaum et al., 2022), making it difficult to

disentangle the roles of plasticity and genetic change in trait evolution. Properly

distinguishing between these mechanisms requires experiments that track life history traits

under common garden conditions. To date, studies that have attempted such distinctions

have almost exclusively focused on abrupt temperature shifts (for example, Van Doorslaer et

al., 2010), leaving the effects of gradual temperature deterioration largely unexplored.

To address these gaps, we conducted a controlled multigenerational evolutionary experiment

with three clonal lines of *D. magna* to test how multiple life-history traits respond to gradually

increasing temperatures, a scenario more representative of ongoing climate change than

abrupt thermal shifts. Populations were propagated through non-overlapping generations

under either constant conditions (control) or a warming regime that increased by 1 °C every

4–6 generations. We monitored fecundity, somatic growth rate, age at maturation, body size,

and extinction risk across generations, and used a reciprocal transplant experiment to

disentangle genetic adaptation from phenotypic plasticity. This design allowed us to evaluate

both the pace and nature of trait change under sustained warming. Our results provide

insight into how gradual thermal deterioration can alter life-history strategies in a key model

species, informing predictions of population responses to ongoing climate change. In doing

so, our study also identifies a potential tipping point in reproductive performance that

precedes mortality, providing a novel perspective on how gradual warming can restructure

life-history strategies well before lethal limits are reached.

2. Materials and Methods

2.1 Experimental procedures

Model Organism and Culture Conditions

Daphnia magna Straus (1820) is a planktonic freshwater crustacean (class Phyllopoda) that

reproduces via cyclic parthenogenesis, producing clutches of up to 100 eggs after each

adult molt (Ebert, 2005). Eggs develop directly into juveniles (neonates), bypassing a resting

stage. The generation time is temperature-dependent, with the first clutch typically produced

5–10 days after birth at the optimal temperature of 20°C (Ebert, 2005). Due to its clonal

reproduction and tractable life cycle, *D. magna* is a widely used model for studying

phenotypic plasticity and local adaptation (Yampolsky et al., 2014).

Three clonal lines were used: HU-HO-2 (hereafter referred to as Hu) and TR-EG-1

(hereafter referred to as Tr) (University of Basel), and An (Middle East Technical University,

Ankara). Cultures were maintained in 250 ml glass beakers with 150 ml of ADaM medium

(Klüttgen et al., 1994), replenished weekly. All populations were kept under a 16:8 h

light:dark cycle and fed with *Chlamydomonas reinhardtii* at 50,000 cells/ml three times per

week. Temperature treatments varied according to experimental design.

Preliminary Thermal Performance Assay

Before the main experiment, we conducted a baseline thermal tolerance assay following

Seefeldt and Ebert (2019) to evaluate the performance limits of the three *D. magna* clones

and to inform temperature treatment design.

Five replicate populations per clone (Hu, Tr and An) containing five adults were maintained

at 22°C in 100 ml beakers containing ADaM medium, replenished weekly. Populations were

acclimated to increasing thermal conditions by raising the temperature to 25°C in week one

and to 28°C in week two. Juvenile females (1–3 days old) were then sampled from each

replicate and individually exposed to one of four test temperatures (30°C, 31°C, 32°C, or

33°C) in 60 ml jars. Each temperature was replicated at least three times per clone. Over

two weeks, individuals were monitored daily for evidence of reproduction (presence of eggs

in brood chambers or neonates in the medium).

Experimental Design and Temperature Treatments

A total of 24 experimental populations were established, comprising 12 treatment and 12

control lines (3 clones × 4 replicates each). Each population consisted of 15 individuals per

generation. Non-overlapping generations were maintained by transferring the first 15

neonates to a fresh beaker of medium at each generation (see Figure 1 for more details

regarding the experimental design and transfer procedures).

In the treatment group (hereafter referred to as 'warming'), temperature increased gradually

over time, beginning at 26°C. Temperature was raised by 1°C every 4 generations until it

reached 29°C, and then by 1°C every 6 generations thereafter. This slowing of the rate

beyond 29°C was intended to minimize heat-induced extinction risk, based on prior findings

(Mitchell and Lampert, 2000) and the results of our baseline assay.

Control populations were maintained at a constant 26°C for the duration of the experiment.

Life History Trait Assays

To quantify evolutionary responses to gradually deteriorating (increasing) temperature, life history traits were measured at five timepoints: 26°C, 27°C, 28°C, 29°C, and 30°C, corresponding to generations 4, 8, 12, 18, and 24 in the treatment populations.

At each timepoint, the first three mature individuals (i.e., individuals with eggs visible in the brood chamber) were isolated from each population and placed individually in 60 ml jars containing ADaM at the population's current temperature. Each female's first and second clutches were counted to estimate fecundity. The first clutch was discarded after hatching to avoid size bias, as first-clutch offspring tend to be smaller (Ebert, 1991). After release of the second clutch, the mother was removed.

Six neonates from the second clutch were randomly selected and reared individually in the same jars. Pilot tests had shown that higher densities could trigger unwanted male production or diapause (ephippia) formation. Neonates were photographed at birth and again at maturation using brightfield microscopy (BRESSER Advance ICD Stereo Microscope with Full HD eyepiece camera at 10x magnification). Body size was measured via gut length (GL) of the relaxed animals in ImageJ and converted to dry mass (DM) using the equation: $DM = 0.00679 \times GL^{2.75}$ (Fossen et al., 2018).

Neonates were monitored daily for development and survival. Age at maturation was defined as the time when the first eggs were visible in their brood chambers. Lifespan was defined as the number of days from birth until death. Somatic growth rate was calculated as the difference in dry mass between birth and maturation, divided by the number of days between the two measurements. The clutch interval (time between first and second clutches) was also recorded.

Figure 1 – Transfer procedure and data collection regime of experimental populations;
(A) 15 neonates per beaker were cultivated until they reached maturation (B), defined as the
time when the first eggs appeared in their brood chambers (note that both neonates and adults
are downscaled relative to the beakers for an easier representation); C) after each round of
temperature increase, 3 adults were sampled for the main assays for quantification of life
history traits; D) hatching (the release of the newborn neonates from the brood chambers)
resulted in formation of pool of both adults and neonates (note that here, for simplicity, the
ratio of newborns vs adults is drawn as 1:1; typically, a larger number of neonates is present);
E) after each new generation of neonates appeared, 15 individuals were sampled and
transferred to a new beaker (i.e. the next generation of an experimental population has been
initiated.

Extinction Monitoring

Each population was visually inspected daily. Populations were recorded as extinct when no

live individuals remained in the beaker.

Reciprocal Transplant Assay

To disentangle the effects of phenotypic plasticity and genetic adaptation, a reciprocal transplant was conducted at generation 25. Three mature individuals were sampled from each control and treatment population. From each adult, up to six second-clutch neonates were reared in the environment opposite to their lineage's ancestral condition.

Control-derived neonates were raised at 30°C, representing novel heat exposure. Treatment-derived neonates were raised at 26°C, representing a return to ancestral temperature. These individuals were reared until maturation, and their adult body size and somatic growth rate were quantified using the same methods as described above.

2.2 Data Analysis

2.2.1 Preliminary Thermal Performance Assay

To assess the baseline thermal tolerance of ancestral clones, we fit a generalized linear mixed model (GLMM) with a binomial error distribution. The presence or absence of offspring (eggs or neonates) was treated as a binary response variable. Clone identity was included as a fixed categorical predictor, and temperature was included as a continuous covariate.

2.2.2 Extinction dynamics

The extinction dynamics of each treatment group were analysed by fitting a Weibull regression model with two fixed factors: 'treatment' and 'clone'. The interaction between the factors was included in the model. The best fit was obtained using the extreme hazard function.

2.2.3 Life History Trait Assays

The datasets containing life history trait values from the main assays were analyzed using linear mixed models (LMMs). The models included "treatment" and "clone" as categorical fixed effects, "generation" as a continuous fixed effect, and "population" and "individual" as categorical random effects to account for hierarchical structure. Each of the seven life history traits was modeled separately as a continuous response variable. Simplified models excluding interaction terms were favored over the more complex models that included the interactions if AIC values were lower (see Appendix section for details). Given the limited sample sizes within each dataset, model parameters were estimated using restricted maximum likelihood (REML). The potential outliers were identified using Grubb's test and were excluded from model comparison (see Appendix section for details).

The potential covariation among life history traits in the experimental populations was examined using Pearson's correlation test, based on pairwise comparisons between traits.

2.2.4 The Reciprocal Transplant Assay

The dataset from the reciprocal transplant assay was analyzed using linear mixed models (LMMs). Fixed effects included "environment" (ancestral or novel), "treatment" (control or thermal deterioration), and "clone" (all categorical). Random effects accounted for hierarchical structure by including "population" and "individual" as categorical variables. Each of the two monitored life history traits—age at maturation and growth rate—was modeled separately as a continuous response variable. Given the limited number of observations in each dataset, model parameters were estimated using restricted maximum likelihood (REML).

All the analyses were carried out using RStudio (version 2024.12.1; R Core Team, 2017). The packages: 'gplots' (v3.1.3; Warnes et al., 2022), 'ggplot2' (v3.4.2; Wickham, 2016),

'plothrix' (v3.8.2; Lemon, 2006), 'RColorBrewer' (v1.1-3; Neuwirth, 2022) were used for

visualization of the datasets: baseline thermal tolerance of ancestral clones; life history trait

assays; the reciprocal transplant assay. The package 'plyr' (v1.8.8; Wickham, 2011) was

used for summarizing the data prior to visualization. 'Survival' (v3.2-13; Therneau, 2021) and

'survminer' (v0.4.9; Kassambara et al., 2021) were used for the analysis and visualization,

respectively, of extinction dynamics. For fitting of the mixed models, the packages lme4

(Bates et al., 2014) and lmerTest (Kuznetzova et al., 2017) were used.

3. Results

3.1 Preliminary thermal performance assay

The reproductive ability of the ancestral clones varied with test temperature (Generalised

Linear Mixed Model; df = 1; $F_{1,247}$ = 11.76; P < 0.001). While the average successful

reproduction rate at the temperature range between 30°C and 32°C was 28%, only a single

individual (0.01%) successfully reproduced at 33°C (Figure 2A). Clone identity also

significantly affected reproduction (Generalised Linear Mixed Model; df = 2; $F_{2,246}$ = 6.7; P =

0.001) (Figure 2B), with the Tr clone showing 2.66 times higher percentage of successfully

reproduced individuals relative to the other two clones combined (z = 3.154; P = 0.002). This

assay established that the upper thermal limit for successful reproduction in the

experimental populations lies just below 33 °C, with sharply reduced success already

evident between 30-32 °C.

3.2 Main evolutionary experiment

Population extinction dynamics

Populations in the warming treatment tended to go extinct earlier than controls, although this

effect was marginal (Weibull regression; df = 1; χ^2 = 3.04; P = 0.08) (Figure 3A). Clone

identity had a similarly marginal effect on extinction rates (Weibull regression; df = 2; χ^2 =

5.02; P = 0.08) (Figure 3B).

Figure 2 – Thermal performance assay of the ancestral clones; A) Percentage of
successfully reproduced individuals per test temperature; **B)** Percentage of successfully
reproduced individuals per clone.

Figure 3 – Extinction dynamics of experimental populations: per treatment group (A);
per clone (B); extinctions are plotted against the generations upon each extinction event; 95%
confidence interval for each Kaplan-Meier survival curve is represented with the dotted lines
(upper and lower boundary).

Life-history trait responses

Reproductive traits

Fecundity declined significantly across generations in both treatments (LMM, df = 1; F = 41.33; P < 0.001) (Figure 4A) but remained 12% higher in warming populations than controls (LMM, df = 1; F = 8.3; P = 0.004) (Figure 4B). A significant clone-by-treatment interaction (LMM; df = 2; F = 4.06; P = 0.02) was driven by higher fecundity in Hu and Tr clones relative to An under warming (An vs Hu, t = 2.58, P = 0.01; An vs Tr, t = 2.35, P = 0.02; Figure 4D). A generation-by-treatment interaction (LMM; df = 1; F = 4.59; P = 0.03) revealed that this advantage was consistent except at 30 °C (24th generation), where warming-line fecundity was 34% lower than controls (Figure 4C).

We saw no overall change in span between clutches (LMM, df = 1; F = 1.33; P = 0.25), but a marginal clone-by-treatment interaction (LMM, df = 2; F = 2.96; P = 0.05) indicated shorter intervals for Tr under warming (t = 2.29, P =0.02) (Figure 5A).

Size traits

Neonate body size was unaffected by generation (LMM, df = 1; F = 1.27; P = 0.26), treatment (LMM, df = 1; F = 1.84; P = 0.18) or clone (LMM, df = 2; F = 1.22; P = 0.3). In contrast, adult body size declined significantly over generations (LMM, df = 1; F = 17.63; P < 0.001) (Figure 5B). Treatment had no main effect (LMM, df = 1; F = 1.48; P = 0.23) but clone identity was significant (LMM, df = 2; F = 3.37; P = 0.04) with mean body size of An declining more than Hu or Tr (Figure 5C). A clone-by-treatment interaction (LMM, df = 2; F = 3.12; P = 0.047) revealed that Tr maintained a size advantage in the warming treatment (Tr vs Hu, t = 2.29, P = 0.02; Tr vs An, t = 1.99, P = 0.048) (Figure 5D).

Development and Somatic growth rate

Development time increased with generation (LMM, df = 1; F = 12.56; P = 0.005) (Figure 6A)

but was on average 3 days shorter (23% faster maturation) in warming populations (LMM, df

= 1; F = 7.66; P = 0.006) (Figure 6B). Clone effects were marginal (LMM, df = 2; F = 2.93; P

= 0.05), with Tr maturing later than An (t = 2.41, P = 0.02) (Figure 6C).

Somatic growth rate declined overall (LMM, df = 1; F = 23.05; P < 0.001) (Figure 6D) but

was 14% faster in warming populations (LMM, df = 1; F = 6.23; P = 0.01) (Figure 6E). Clone

effects were marginal (LMM, df = 2; F = 2.43; P = 0.09), with An growing faster than Tr (t =

2.14, P =0.03) (Figure 6F).

Lifespan

Lifespan decreased significantly across generations (LMM, df = 1; F = 35.59; P < 0.001)

(Figure 7A) and was 21% shorter in warming populations than controls (LMM, df = 1; F =

24.92; P < 0.001) (Figure 7B). The largest treatment effect occurred at 26°C (4th generation)

(LMM, generation × treatment; df = 1; F = 10.21; P = 0.002; Figure 7C). Clone identity was

significant (LMM, df = 2; F = 7.43; P < 0.001), with An living longer than both Tr (t = - 3.83; P

< 0.001) and Hu (t = - 3.16; P = 0.002) (Figure 7D).

Life history trait covariation in warming populations

Somatic growth rate was strongly negatively correlated with development time (r = - 0.86; t

(60) = - 13.41; P < 0.001) (Figure 8A) and with both neonate body size (r = - 0.31; t (60) = -

2.52; P = 0.01), and adult body size (r = - 0.49; t (60) = - 4.38; P < 0.001) (Figure 8B).

Development time was positively correlated with adult body size (r = 0.64; t (60) = 6.42; P <

0.001) (Figure 8C), and adult body size correlated positively with fecundity (r = 0.43; t (67) =

3.92; P = 0.002) (Figure 8D). Other correlations were weak and non-significant (see

Appendix section for details).

Figure 4 – Mean fecundity (the number of neonates within the first two clutches) of the
populations: per generation (A); per treatment group (B); per treatment group during each assay
(C): X axis contains the number of generations that each assayed population underwent, with the
temperature degree that the warming populations experienced during the assay stated below; per
clone, per treatment group (D); the bars represent standard error of the mean.

Figure 5 – Mean span between the first two clutches (day) of the populations (A); Mean
adult body size (mg od DM) of the populations (B-D): per generation (B); per clone (C); per
clone per treatment group; the bars represent standard error of the mean.

Figure 6 – Mean development time (day) of the populations (A- C): per generation (A); per
treatment group (B); per clone (C); **Mean somatic growth rate (per day) of the population (D -**
F): per generation (D); per treatment group (E); per clone (F); the bars represent standard error
of the mean.

Figure 7 – Mean lifespan (day) of experimental populations: per generation (A); per treatment group (B); per treatment group during each assay (C): X axis contains the number of generations that each assayed population underwent, with the temperature degree that the warming populations experienced during the assay stated below; per clone (D); the bars represent standard error of the mean.

Figure 8 – Significant correlations between the pairs of the life history traits in warming populations; A) Somatic growth rate (per day) plotted against development time (day); **B)** Somatic growth rate (per day) plotted against adult body size (mg of DM); **C)** Adult body size (mg of DM) plotted against development time (day); **D)** Fecundity (number of neonates within the first two clutches) plotted against adult body size (mg of DM).

4.3 The reciprocal transplant experiment

Mean adult body size differed significantly between treatments (LMM, df = 1; F = 9.05; P = 0.004), with control-line offspring 17% larger than warming-line offspring, regardless of rearing environment (LMM, environment × treatment; df = 1; F = 0.32; P = 0.57) (Figure 9A).

Mean growth rate showed no significant treatment effect (LMM, df = 1; F = 0.05; P = 0.82) and no environment × treatment interaction (LMM; df = 1; F = 0.96; P = 0.33) (Figure 9B).

A

B

Figure 9 – The reciprocal transplant experiment; A) Mean adult body size (mg of DM) of control
and warming populations kept in their ancestral and novel environment; **B)** Mean somatic growth
rate (per day) of control and warming populations kept in their ancestral and novel environment;
the bars represent standard error of the mean.

5. Discussion

Our multigenerational experiment demonstrates that gradual thermal deterioration reshapes life-

history strategies in *Daphnia magna* in ways that differ from the patterns observed under abrupt

temperature shifts. The most striking pattern was a distinct, rate-dependent trajectory, with an

initial phase of elevated reproduction followed by the loss of the reproductive advantage at a

sub-lethal fecundity threshold well below lethal temperatures (typically 35 °C for *D. magna*;

Seefeldt & Ebert, 2019). This is consistent with other studies showing that demographic tipping

points can occur before lethal thermal limits are reached (Kingsolver & Huey, 2008; Seefeldt &

Ebert, 2019). Populations exposed to sustained warming matured earlier, grew faster, and

initially produced more offspring than controls, but also exhibited shorter lifespan, and the

eventual loss of reproductive advantage relative to control near 30 °C, well below their lethal

limit. These findings highlight that the rate of environmental change influences both the

trajectory and limits of adaptive responses, with implications for predicting persistence under

ongoing climate change.

Warming maintains fecundity until a threshold is reached

In contrast to many abrupt-shift experiments that report uniformly reduced fecundity of *D. magna*

under elevated temperatures (for example, Cambronero et al., 2018; Wang et al., 2023),

warming lines in our experiment maintained a fecundity advantage for most of the thermal

trajectory. This pattern is consistent with studies showing that moderate warming can boost

reproductive output via accelerated maturation and shorter generation times (for example,

Musolin et al., 2010). However, this advantage diminishes near 30 °C, a tipping point also

indicated by our thermal tolerance assay, where only 28% of individuals reproduced

successfully and reproduction nearly ceased at 33 °C. The timing of extinctions in the main

experiment closely paralleled this threshold, with 75% of warming populations lost between 30

and 32 °C. These results show that reproductive ceilings can be reached well before mortality

thresholds, reducing population viability during warm periods even if temperatures remain below

lethal limits.

Adult body size decline was unaffected by warming

We detected an overall reduction in mean adult body size, but without difference between

warming and control populations. A plausible explanation is that the temperature window that

our warming populations experienced (the 26 - 30°C range) lies near the upper end of the

thermal performance curve for *Daphnia* (Seefeldt and Ebert, 2019; Wang et al., 2023), where

body size plasticity may already be constrained. Furthermore, there is evidence that resource

limitations become stronger at higher temperatures and larger body size, causing growth to

decelerate at a smaller size at warmer temperatures (Hoefnagel et al., 2018).

The role of clutch interval

463 Some traits, such as clutch interval, showed limited average change but exhibited significant

464 genotype-by-treatment interactions: the Tr clone reduced its clutch interval under warming, a

465 shift not seen in the other clones. Because shorter intervals between clutches can substantially

466 increase lifetime reproductive output in *Daphnia* (Garbutt et al., 2014; Claska & Gilbert, 1998),

467 this clone-specific change may represent an important adaptive pathway under warming.

Warming reduces lifespan

Elevated temperature significantly reduced the lifespan of experimental populations relative to

control populations. Moreover, we observed a significant effect of clonal identity, with An clone

showing higher longevity relative to the other two clones. The negative correlation between

temperature and longevity has previously been reported (for example, Musolin et al., 2007;

Ciota et al, 2014; Giovaninni et al., 2023), including the studies with *D. magna* (for example,

MacArthur and Baillie, 1929; Hoefnagel et al., 2018). Our study refined these estimates by

providing detailed lifespan data across a narrower but ecologically critical thermal window (26 -

30°C), capturing a sharp decline relative to control populations during the first three assays (26 -

28°C), followed by the equalization of mean longevity between the two treatment groups at the

two highest temperatures.

Life-history traits correlations

Despite an overall reduction in mean adult body size in both treatment groups, there is evidence

that a larger body size was advantageous at elevated temperatures through its effect on

fecundity. Moreover, we detected a significant plastic response in both traits, manifested through

a simultaneous increase of both mean body size and fecundity of Tr clone when subjected to

elevated temperature. Warming populations showed both a significant decline of development

time and an acceleration of somatic growth rate relative to controls, consistent with findings in

other studies (Henning-Lucass et al., 2016; Hoefnagel et al., 2018). These two traits showed a

strong negative correlation, driven by divergent clone responses. We also saw a significant

positive correlation between development time and adult body size, and a significant negative

correlation between growth rate and body size. Together, these patterns suggest that delayed

maturation can be advantageous at high temperatures by allowing individuals to reach a larger

size coupled with a higher fecundity, whereas faster-growing clones (namely, An) show trade-off

between both size and fecundity with increased rate of development. By contrast, we detected

no covariation between fecundity and lifespan, nor between lifespan and development or

growth, suggesting that lifespan reductions were driven primarily by direct thermal stress rather

than by indirect trade-offs (Sun et al., 2023; Schwartz et al., 2016).

Clone-specific resilience

Marked differences among clones underline the role of standing genetic variation. The Tr clone

maintained larger body size and higher fecundity under warming. In contrast, the An clone was

characterized by both reduced mean body size and fecundity but had greater longevity. These

patterns echo mesocosm and laboratory findings showing that genotypes vary in their thermal

sensitivity and capacity for persistence under stress (Geerts et al., 2015; Henning-Lucass et al.,

2016). Such variation indicates the potential for evolutionary rescue (Gomulkiewicz and Holt,

1995) but also means that population-level means can mask the presence of more resilient

genotypes.

Plasticity versus genetic change

The reciprocal transplant experiment (RTE) revealed a trait-specific pattern that was not

apparent in the main assays. While mean adult body size declined similarly across treatments

during the main experiment, warming-line offspring were consistently smaller than controls in the

transplant assay, regardless of rearing environment. This suggests a heritable component or

persistent trans-generational effect in body size reduction, which was masked in the main

experiment by parallel size declines across both treatments. By contrast, growth rate showed no

significant treatment or environment interaction, indicating that it remained largely plastic within

the time frame of the study. Notably, control offspring exhibited an increase in growth rate when

reared in the novel (warming) environment, whereas warming-line offspring did not show a

corresponding change when returned to the cooler environment. This trait-specific pattern aligns

with other Daphnia studies showing that some traits respond primarily through evolution, others

through plasticity (Yampolsky et al., 2014; Seefeldt & Ebert, 2019).

More broadly, our main experiment and RTE combined highlight that gradual warming can

reveal distinct evolutionary trajectories across traits: some (like adult body size) showing

persistent lineage-level shifts, while others (like growth rate) remain flexible but not divergent.

The limited number of generations (25) prior to the transplant likely constrained the magnitude of

genetic change, but the divergence we detected suggests that even under relatively short time

spans, gradual thermal deterioration can produce heritable shifts in key life-history traits.

Rate of change as a key driver

The +1 °C per 4–6 generations regime produced trajectories distinct from those under single-

step warming: an initial period of reproductive advantage, which was lost in sublethal

temperature. This dynamic mirrors microbial experiments where gradual change extended

persistence without preventing eventual decline (for example, Killeen et al., 2017). Our results

reinforce that warming rate and not just absolute temperature should be treated as a primary

parameter in both experiments and models, as predicted by eco-evolutionary theory and

observed in microbial systems (Collins and de Meaux, 2009; Bell and Gonzalez, 2011; Killeen et

al., 2017).

Limitations

Our experiment used a limited number of genotypes and spanned only 25 generations before

the transplant assay. Longer-term experiments incorporating greater genetic diversity are

needed to fully capture the interplay between plasticity and evolution under gradual warming. In

addition, our design did not manipulate resources or community composition, which may

influence whether the fecundity advantage observed here emerges across a wider range of

ecological contexts.

Although some baseline differences were present (e.g., slightly higher fecundity in Tr under

warming and longer lifespan in controls at 26 °C), these did not alter the main trajectories:

warming populations maintained a fecundity advantage until the highest temperature and

consistently showed lifespan declines. This suggests that the observed patterns were robust to

initial conditions.

Broader implications and conclusions

From an applied perspective, our findings show that gradual warming can mask an impending

demographic decline by temporarily enhancing reproduction. Sub-lethal thresholds such as the

~30 °C fecundity ceiling may be crossed more often in the future as thermal safety margins

shrink, particularly in regions or seasons already near species' optima (Deutsch et al., 2008). In

D. magna, this tipping point occurred near 30 °C, well below the lethal limit, and was closely

aligned with extinction events. The fact that reproductive ceilings can be reached long before

mortality thresholds, and that their timing depends on the pace of warming, underscores the

need to treat warming rate as a fundamental driver in both experimental design and predictive

modelling, and to recognise that identical thermal endpoints reached at different rates can

produce qualitatively different demographic and evolutionary outcomes (Killeen et al., 2017). As

thermal safety margins shrink globally, such rate-dependent demographic ceiling may become a

common feature of population decline, making the detection of early-warning signals for

reproductive tipping points essential for anticipating and mitigating biodiversity loss under

climate change.

6. Appendix

6.1 Outlier detection results and model selection

Span between the clutches

Two extreme values were identified as potential outliers using Grubbs' test (p < 0.05) and were

excluded from model comparison.

Neonate body size

Four extreme values were identified as potential outliers using Grubbs' test (p < 0.05) and were

excluded from model. Model selection was based on a comparison of AIC values and a

likelihood ratio test. A simplified model excluding interaction terms (AIC = -1274.3) was favored

over a more complex model that included the interactions (AIC = -1263.1), as the later did not

significantly improve model fit according to a likelihood ratio test (χ^2, P = 0.9).

Adult body size

A single extreme value was identified as potential outliers using Grubbs' test (p < 0.05) and was

excluded from model.

The development time

Two extreme values were identified as potential outliers using Grubbs' test (p < 0.05) and were

excluded from model comparison. A simplified model excluding interaction terms (AIC = 1198.7)

was favored over the full interaction model (AIC = 1208.5; LRT P = 0.77).

Somatic growth rate

Model selection was based on a comparison of AIC values and a likelihood ratio test. A simplified model excluding interaction terms (AIC = - 107.35) was favored over a more complex model that included the interactions (AIC = -97.86), as the later did not significantly improve model fit according to a likelihood ratio test (χ^2, P = 0.72).

6.2 Non-significant correlations between life history traits

We found no significant correlation between lifespan and development time (r = - 0.04; t (79) = - 0.34; P = 0.73); lifespan and somatic growth rate (r = 0.02; t (33) = 0.09; P = 0.93); lifespan and fecundity (r = 0.06; t (69) = 0.52; P = 0.61); fecundity and span between clutches (r = - 0.28; t (272) = -0.29; P = 0.78); adult body size and span between clutches (r = - 0.03; t (67) = - 0.26; P = 0.79); somatic growth rate and fecundity (r = - 0.08; t (60) = - 0.6; P = 0.55).

Acknowledgements

This study has been funded by Tübitak (project code: TB.0064).

We are grateful to the former postgraduate student, Seray Altıntaş, for assistance in development of the protocol for algal quantification, and for her assistance with pilot experiments and maintenance of cultures. We thank the undergraduate students, including Helin Bahar Calı, Kerem Durdu, Gülnur Karabiber, and Ece Türksever, for their assistance with pilot experiments and maintenance of cultures. We are grateful to Dr Bebac Šmepčević for great illustrations within Figure and for assistance in development of the protocol for algal quantification.

Data Accessibility Statement

Data is available through the Dryad Digital Repository

(http://datadryad.org/share/LINK_NOT_FOR_PUBLICATION/ZkFUNcfZSiCaQRD6FLZIJFfQ5T

EIUAQLMD3KHbT3ars).

Competing Interests Statement

The authors declare no conflict of interest.

References

Albini, D., Ransome, E., Dumbrell, A.J. *et al.* 2025. Warming alters plankton body-size
distributions in a large field experiment. *Commun Biol* 8, 162.

Bates D, Mächler M, Bolker B, Walker S. 2015. Fitting Linear Mixed-Effects Models Using
lme4. *Journal of Statistical Software*, 67(1), 1–48

Barnosky, A. D., Hadly, E. A., Bascompte, J., Berlow, E. L., Brown, J. H., Fortelius, M., Getz, W.
M., Harte, J., Hastings, A., Marquet, P. A., Martinez, N. D., Mooers, A., Roopnarine, P., Vermeij,
G., Williams, J. W., Gillespie, R., Kitzes, J., Marshall, C., Matzke, N., Mindell, D. P., Revilla, E.,
and Smith, A. B. 2012. Approaching a state shift in Earth's biosphere. *Nature* 486 (7401):52-8.

Bell, G., & Gonzalez, A. 2011. Adaptation and evolutionary rescue in metapopulations
experiencing environmental deterioration. *Science*. 332(6035), 1327–1330.

Cahill A.E., Aiello-Lammens M.E., Fisher-Reid M.C., Hua X., Karanewsky C.J., Yeong Ryu H.,
Sbeglia G.C., Spagnolo F., Waldron J.B., Warsi O., Wiens J.J. 2013. How does climate change
cause extinction? *Proc R Soc B*. 280: 20121890.

Cuenca Cambronero, M., Marshall, H., De Meester, L. *et al.* 2018. Predictability of the impact of
multiple stressors on the keystone species *Daphnia*. *Sci Rep* **8**, 17572.

Cardillo M, Mace GM, Jones KE, Bielby J, Bininda-Emonds OR, Sechrest W, Orme CD, Purvis
A. 2005. Multiple causes of high extinction risk in large mammal species. *Science*. Aug
19;309(5738):1239-41.

Chivian, E. and Bernstein, A. 2008. *Sustaining Life: How Human Health Depends on Biodiversity*.
Oxford, UK: Oxford University Press.

Ciota AT, Matacchiero AC, Kilpatrick AM, Kramer LD. 2014. The effect of temperature on life
history traits of Culex mosquitoes. *J Med Entomol.* Jan;51(1):55-62.

Claska, M. E., & Gilbert, J. J. 1998. The effect of temperature on the response of *Daphnia* to toxic
cyanobacteria. *Freshwater Biology*. 39(2), 221–232.

Collins, S. and de Meaux, J. 2009. Adaptation to different rates of environmental change in
Chlamydomonas. *Evolution* 63 (11):2952-65.

Martínez-De León, Gerard & Marty, Arianne & Holmstrup, Martin & Thakur, Madhav. 2024.
Population resistance and recovery after an extreme heat event are explained by thermal effects
on life-history traits. *Oikos*. 10.1111/oik.10023.

Deutsch C.A., Tewksbury J.J., Huey R.B., Sheldon K.S., Ghalambor C.K., Haak D.C., Martin
P.R. 2008. Impacts of climate warming on terrestrial ectotherms across latitude. *Proc Natl Acad
Sci U S A*. 105(18):6668-72.

Du WG, Ji X, Zhang YP, Xu XF, Shine R. 2005. Identifying sources of variation in reproductive
and life-history traits among five populations of a Chinese lizard (Takydromus septentrionalis,
Lacertidae). *Biol J Linn Soc* 85:443–453

Ebert, D. 1991. The effect of size at birth, maturation threshold and genetic differences on the
life-history of Daphnia magna. *Oecologia*. Apr;86(2):243-250.

Ebert, D. 2005. *Ecology, Epidemiology, and Evolution of Parasitism in Daphnia*. Bethesda (MD):
National Center for Biotechnology Information (US).

Falkowski, P. G. 1994. The role of phytoplankton photosynthesis in global biogeochemical cycles.
Photosynthesis Research 39 (3):235-58.

Fossen, E.I.F., Pélabon, C., Einum, S. 2018. An empirical test for a zone of canalization in thermal
reaction norms. J. *Evol. Biol.*, 31: 936-943.

Geerts, A., Vanoverbeke, J., Vanschoenwinkel, B., Van Doorslaer, W., Feuchtmayr, H., Atkinson
D., Moss, B., Davidson, T. A., Sayer, C. D., De Meester, L. 2015. Rapid evolution of thermal
tolerance in the water flea *Daphnia*. *Nature Clim Change*. 5, 665–668.

Garbutt, J. S., Scholefield, J. A., Vale, P. F., & Little, T. J. 2014. Elevated maternal temperature
enhances offspring disease resistance in *Daphnia magna*. *Biology Open*. 3(8), 546–552.

Giovannini I, Manfrin C, Greco S, Vincenzi J, Altiero T, Guidetti R, Giulianini P and Rebecchi L.
2023. Increasing temperature-driven changes in life history traits and gene expression of an
Antarctic tardigrade species. *Front. Physiol*. 14:1258932

Gomulkiewicz, R. and Holt, R. D. 1995. When does evolution by natural selection prevent
extinction?. *Evolution* 49 (1):201-7.

Hawkins, E., Ortega, P., Suckling, E., Schurer, A., Hegerl, G., Jones, P., Joshi, M., Osborn, T. J.,
Masson-Delmotte, V., Mignot, J., Thorne, P., van Oldenborgh, G. J. 2017. Estimating Changes in
Global Temperature since the Preindustrial Period. *Bulletin of the American Meteorological*
Society. 98(9), 1841-1856.

Henning-Lucass, N., Cordellier, M., Streit, B. and Schwenk, K. 2016. Phenotypic plasticity in life-
history traits of *Daphnia galeata* in response to temperature – a comparison across clonal
lineages separated in time. *Ecol Evol*. 6: 881-891.

Hoefnagel KN, de Vries EHJ, Jongejans E, Verberk WCEP. 2018. The temperature-size rule in
Daphnia magna across different genetic lines and ontogenetic stages: Multiple patterns and
mechanisms. *Ecol Evol*.;8:3828–3841.

Huey, R. B., & Kingsolver, J. G. (2019). Climate warming, resource availability, and the
metabolic meltdown of ectotherms. American Naturalist, 194(6), E140–E150.

679
Iltis C., Louâpre P., Pecharová K., Thiéry D., Zito S., Bois B., Moreau J. 2019. Are life-history traits
equally affected by global warming? A case study combining a multi-trait approach with fine-grain
climate modeling. *J Insect Physiol*. 117:103916.

Isaac J.L. 2009. Effects of climate change on life history: implications for extinction risk in
mammals. *Endang Species Res* 7:115-123.

Kassambara, A., Kosinski, M., & Biecek, P. 2021. Survminer: Drawing survival curves using
"ggplot2". R Package Version, 0(4), 9

Killeen, J., Gougat-Barbera, C., Krenek, S., Kaltz, O. 2017. Evolutionary rescue and local
adaptation under different rates of temperature increase: a combined analysis of changes in
phenotype expression and genotype frequency in *Paramecium* microcosm. *Mol Ecol*, 26: 1734-
1746.

Kingsolver, J. G., & Huey, R. B. 2008. Size, temperature, and fitness: Three rules. *Evolutionary*
Ecology Research. 10(2), 251–268.

693
Klüttgen, B., Dülmer, U., Engels, M., Ratte, H.T. 1994. ADaM, an artificial freshwater for the cult
ure of zooplankton. *Water Res*. 28:743–746.

Kuznetsova A, Brockhoff PB, Christensen RHB. 2017. lmerTest Package: Tests in Linear Mixed
Effects Models. *Journal of Statistical Software*, 82(13), 1–26.

Lalejini A, Ferguson AJ, Grant NA and Ofria C. 2021. Adaptive Phenotypic Plasticity Stabilizes
Evolution in Fluctuating Environments. *Front. Ecol. Evol*. 9:715381

Laughton A.M., O'Connor C.O., Knell R.J. 2017. Responses to a warming world: Integrating life
history, immune investment, and pathogen resistance in a model insect species. *Ecol Evol*.
7(22):9699-9710.

Lemon, J. 2006. Plotrix: A package in the red light district of R. R-News., 6(4), 8–12.

Lindsey, H. A., Gallie, J., Taylor, S., and Kerr, B. 2013. Evolutionary rescue from extinction is
contingent on a lower rate of environmental change. *Nature*. 494 (7438):463-7.

Liukkonen M, Kronholm I, Ketola T. 2021. Evolutionary rescue at different rates of environmental
change is affected by trade-offs between short-term performance and long-term survival. *J Evol*
Biol.;34:1177–1184.

McKinney, M. L. 1997. Extinction vulnerability and selectivity: combining ecological and
paleontological views. *A. Rev. Ecol. System*. 28, 495^516.

Mitchell, S.E. and Lampert, W. 2000. Temperature adaptation in a geographically widespread
zooplankter, *Daphnia magna*. *Journal of Evolutionary Biology*. 13: 371-382.

Musolin, D.L., Tougou, D., Fujisaki, K. 2010. Too hot to handle? Phenological and life-history
responses to simulated climate change of the southern green stink bug *Nezara*
viridula (Heteroptera: Pentatomidae). *Global Change Biology*. 16: 73-87.

Neuwirth, E. 2022. RColorBrewer: ColorBrewer palettes. R package version 1.1-3.

Parmesan, C. 2006. Ecological and Evolutionary Responses to Recent Climate Change. *Annual*
Review of Ecology, Evolution and Systematics. 37 (1):637-669.

Pepin, N. C., Arnone, E., Gobiet, A., Haslinger, K., Kotlarski, S., Notarnicola, C., et al. 2022.
Climate changes and their elevational patterns in the mountains of the world. *Reviews of*
Geophysics, 60, e2020RG000730.

Petkovic, N. and Colegrave, N. 2023. The effects of sex on extinction dynamics of
Chlamydomonas reinhardtii depend on the rate of environmental change. *Journal of Evolutionary*
Biology, *00*, 1–13.

Purvis A, Gittleman JL, Cowlishaw G, Mace GM. 2000. Predicting extinction risk in declining
species. *Proc Biol Sci*. Oct 7;267(1456):1947-52.

R Core Team. 2017. R: A language and environment for statistical computing. R Foundation for
Statistical Computing http://www.R-proje ct.org/.

Root, T., Price, J., Hall, K. *et al.* 2003. Fingerprints of global warming on wild animals and plants.
Nature. 421, 57–60.

Schaum C-E, Buckling A, Smirnoff N, Yvon-Durocher G. 2022. Evolution of thermal tolerance and
phenotypic plasticity under rapid and slow temperature fluctuations. *Proc. R. Soc. B*. 289:
20220834.

Schwartz, T. S., Pearson, P., Dawson, J., Allison, D. B., & Gohlke, J. M. 2016. Effects of
fluctuating temperature and food availability on reproduction and lifespan. *Experimental*
Gerontology. 86, 62–72.

Seefeldt, L. and Ebert, D. 2019. Temperature- versus precipitation-limitation shape local
temperature tolerance in a Holarctic freshwater crustacean. *Proc. R. Soc. B.* 286:
20190929.20190929.

Sun, S.-J., Dziuba, M. K., Jaye, R. N., & Duffy, M. A. 2023. Transgenerational plasticity in a
zooplankton in response to elevated temperature and parasitism. *Ecology and Evolution*. 13,
e9767.

Therneau, T. 2021. A package for survival analysis in R. R package version 3.2-13.

Thomas C.D., Cameron A., Green R.E., Bakkenes M., Beaumont L.J., Collingham Y.C., Erasmus
B.F., De Siqueira M.F., Grainger A., Hannah L., Hughes L., Huntley B., Van Jaarsveld A.S.,
Midgley G.F., Miles L., Ortega-Huerta M.A., Peterson A.T., Phillips O.L., Williams S.E. 2004.
Extinction risk from climate change. *Nature*. 427(6970):145-8.

Van Doorslaer, W., Stoks, R., Jeppesen, E., De Meester, L. 2007. Adaptive microevolutionary
responses to simulated global warming in *Simocephalus vetulus*: a mesocosm study. *Global
Change Biology*, 13: 878-886.

Van Doorslaer, W., Stoks, R., Swillen, I., Feuchtmayr, H., Atkinson, D., Moss, B., Meester, L.
2010. Experimental thermal microevolution in community-embedded Daphnia
populations''. *Climate Research*. 43, 81-89.

Walther GR, Post E, Convey P, Menzel A, Parmesan C, Beebee TJ, Fromentin JM, Hoegh-
Guldberg O, Bairlein F. 2002. Ecological responses to recent climate change. *Nature*. Mar
28;416(6879):389-95.

Wang Y-J, Tüzün N, De Meester L, Feuchtmayr H, Sentis A, Stoks R. 2023. Rapid evolution of
unimodal but not of linear thermal performance curves in *Daphnia magna*. *Proc. R. Soc*. B 290:
20222289.

Warnes, G. R., Bolker, B., Bonebakker, L., Gentleman, R., Huber, W., Liaw, A., Lumley, T.,
Maechler, M., Magnusson, A., Moeller, S., Schwartz, M., & Venables, B. 2022. gplots: Various R
programming tools for plotting data. R package version 3.1.3.

Wickham, H. 2011. The Split-apply-combine strategy for data analysis. *Journal of Statistical
Software*. 40(1), 1–29.

Wickham, H. 2016. ggplot2: Elegant graphics for data analysis. Springer-Verlag.

Wiens, J.J. and Zelinka, J. 2024. How many species will Earth lose to climate change?. *Global
Change Biology*. 30, e17125

Yampolsky, L.Y., Schaer, T.M.M., Ebert, D. 2014. Adaptive phenotypic plasticity and local
adaptation for temperature tolerance in freshwater zooplankton. *Proc. R. Soc. B*. 281: 20132744

Yoccoz NG and Mesnager, S. 1998. Are alpine bank voles larger and more sexually dimorphic
because adults survive better?. *Oikos* 82:85–98

774