The centriculum, a membrane reticulum that surrounds *C. elegans* centrosomes,

may serve as a microtubule filter

Richa Maheshwari[1], Mohammad M Rahman[1], Abigail Ruddick[1,2], Seth Drey[1,3], Robert S

Mirabello[1], and Orna Cohen-Fix[1]*

[1]The Laboratory of Biochemistry and Genetics, National Institute of Diabetes and

Digestive and Kidney Disease, National Institutes of Health, Bethesda MD, 20892, USA

[2]Current address: University of Oregon, Department of Biology, Eugene, OR 97403,

USA

[3]Current address: University of Louisville School of Medicine, Louisville KY, 40202, USA

*Corresponding author

ornac@niddk.nih.gov,

(301) 549-2184

Running title: Centriculum as a microtubule filter

Key Words: centriculum, centrosome, *C. elegans*, microtubules, SPD-5;

Abstract

Centrosomes are cytoplasmic microtubule-nucleating structures. They are considered membraneless organelles, but in several cell types they are surrounded by ER-derived membrane. In *C. elegans* early embryos, this membrane forms a dense membrane reticulum, named the centriculum, that affects centrosome structure and microtubule nucleating capacity. The centriculum is adjacent to the centrosome's pericentriolar material (PCM) and to abundant, short peri-centrosomal microtubules. Here we show that when microtubule abundance is reduced, centriculum and PCM size decrease, while the density of the PCM protein SPD-5 increases, suggesting that the PCM can be compacted. We further show that centriculum size is determined by microtubules, and that the centriculum likely limits the length of peri-centrosomal microtubules. Finally, we find that the centriculum is more porous where spindle microtubules pass compared to where astral microtubules pass. These data are consistent with the centriculum serving as a microtubule filter, blocking the extension of most, but not all, centrosome-nucleated microtubules. Finally, if microtubule-centriculum collisions result in microtubule catastrophe, the filter function of the centriculum could also explain the high concentration of soluble tubulin at the *C. elegans* centrosome.

45 **Summary Statement**

47 The centriculum is a centrosome-associated membrane reticulum in *C. elegans*

48 embryos. This study shows that its size depends on microtubules and that it may act as

49 a filter, limiting microtubule growth.

Introduction

Centrosomes are cytoplasmic structures that nucleate microtubules, polymers made of α- and β-tubulin heterodimers that form linear, side-by-side, protofilaments, creating a hollow microtubule tube (reviewed in (Aljiboury & Hehnly, 2023)). Microtubules are polar; centrosome-nucleated microtubules have their minus (α-tubulin facing) ends embedded in the centrosome, and their plus (β-tubulin facing) ends extending away from the centrosome. Microtubules are dynamic structures that grow and shrink mainly at their plus end in a manner that is dependent on soluble tubulin concentrations and aided by microtubule associated proteins, such as motor proteins, microtubule polymerases and severing proteins, and more (reviewed in (Gudimchuk & McIntosh, 2021). In dividing cells, centrosomes nucleate microtubules that either form the mitotic spindle or extend to the cell cortex in the form of astral microtubules (reviewed in (Hoffmann, 2021)).

At the core of centrosomes are two centrioles; they are surrounded by a pericentriolar material (PCM), a proteinaceous structure whose properties are just beginning to emerge (reviewed in (Magescas *et al*, 2019; Pimenta-Marques & Bettencourt-Dias, 2020; Woodruff *et al*, 2014). The PCM is rich in coiled-coil proteins that form a lattice, or scaffold, onto which additional PCM proteins are recruited. One of the main components of the PCM in *C. elegans* is SPD-5 (Hamill *et al*, 2002), which is functionally homologous to Centrosomin/Cnn in *Drosophila* (Megraw *et al*, 2001; Megraw *et al*, 1999; Timothy *et al*, 1999) and CDK5RAP2 in vertebrates (Fong *et al*, 2008). The PCM also includes γ-tubulin (TBG-1 in *C. elegans*), which, along with associated proteins that form the γ-tubulin ring complex (γ -TuRC), serves to nucleate microtubules ((Moritz *et al*, 1995; Zheng *et al*, 1995) reviewed in (Gao *et al*, 2024)). At the onset of mitosis, the PCM increases in size in a process known as centrosome maturation ((Khodjakov & Rieder, 1999) reviewed in (Vasquez-Limeta & Loncarek, 2021)). The expansion of the *C. elegans* PCM, as measured by SPD-5 accumulation, involves the incorporation of soluble SPD-5 throughout the volume of the SPD-5 lattice (Laos *et al*, 2015; Woodruff *et al*, 2015). This process is facilitated by phosphorylation of SPD-5 by the polo-like kinase PLK-1 (Cabral *et al*, 2019; Nakajo *et al*, 2022; Rios *et al*, 2024; Woodruff *et al.*, 2015;

Wueseke *et al*, 2016). PLK-1 phosphorylation of SPD-5 is independently required to

promote γ-tubulin recruitment to the PCM (Ohta *et al*, 2021). The role of PLK-1 in

centrosome maturation is conserved throughout eukaryotes (Conduit *et al*, 2015; Lane

& Nigg, 1996; Lee & Rhee, 2011).

Centrosomes are considered membraneless organelles, although there are numerous

examples of endoplasmic reticulum (ER)-derived membrane accumulation adjacent to

the centrosome in *Drosophila*, sea urchin, the fish medaka, mammalian cultured cells,

and *C. elegans* (Araújo *et al*, 2023; Bergman *et al*, 2015; Bobinnec *et al*, 2003; Diaz *et*

al, 2019; Harris, 1975; Karabasheva & Smyth, 2019; Rollins & Blankenship, 2023;

Schlaitz, 2014; Terasaki, 2000; Waterman-Storer *et al*, 1993). This peri-centrosomal

membrane was analyzed in the *C. elegans* 1-cell embryo by volume electron

microscopy at 9 nm isotropic resolution and found to be made of a dense membrane

reticulum, leading to its name the centriculum (Maheshwari *et al*, 2023). While the

structure of peri-centrosomal membranes in other organisms has yet to be examined at

this resolution, the presence of the ER curvature-inducing proteins Rtn1 and ReepB at

peri-centrosomal membranes of *Drosophila* embryos (Diaz *et al.*, 2019) suggests a

similar structure. In *C. elegans*, down-regulation of atlastin, which forms ER-ER

junctions (Hu & Rapoport, 2016), led to an increase in centriculum size (Maheshwari *et*

al., 2023). This was accompanied by an increase in both centrosome size and the

amount of PCM material (as measured by the area and total amount of fluorescently

tagged PCM proteins), and increased microtubule-nucleating capacity (Maheshwari *et*

al., 2023). Likewise, in *Drosophila* embryos, alteration to ER structure, including

prominent changes to the membrane surrounding centrosomes, caused defects in

centrosome and spindle structure (Araújo *et al.*, 2023; Rollins & Blankenship, 2023).

Thus, at least in these systems, centrosome function is affected by the membrane

adjacent to it, indicating that the centrosome may not be as membraneless as

previously assumed.

It is well documented that during *C. elegans* early embryogenesis, the majority of

microtubules nucleated from the surface of the centrosome terminate in the vicinity of

the centrosome (for example, see (Albertson, 1984; Redemann *et al*, 2017) and Fig. 1A;

defined here as peri-centrosomal microtubules). A similar accumulation of short

microtubules around centrosomes is seen in embryos of medaka (Kiyomitsu *et al*,

2024), *Drosophila* (Rollins & Blankenship, 2023) and sea urchin (Xie *et al*, 2025), all of

which also have centrosome-associated ER membranes. The *C. elegans* centrosome is

also able to accumulate soluble tubulin at a concentration that is 10-fold higher than in

the cytoplasm (Baumgart *et al*, 2019). The precise mechanism that facilitates this

accumulation is not known. A condensate of SPD-5 with additional PCM proteins was

shown to concentrate soluble tubulin *in vitro* (Woodruff *et al*, 2017), although not to the

extent observed *in vivo* (Baumgart *et al.*, 2019). The centriculum provides an attractive

mechanism for these two phenomena: First, peri-centrosomal microtubule accumulation

could reflect collisions of a subset of microtubules with centriculum membranes that are

in their path and impede their elongation. In this way, the centriculum would act as a

microtubule filter, limiting the number of microtubules that can fully elongate. This could

reduce the competition for cytoplasmic tubulin pools and/or prevent the formation of a

poorly functioning spindle. Second, if these collided microtubules terminate at the

centriculum and undergo catastrophe (akin to the catastrophe that occurs when

microtubules encounter the plasma membrane (Bouvrais *et al*, 2021)), this could

provide a source for the exceedingly high soluble tubulin concentration at the

centrosome (Baumgart *et al.*, 2019). We therefore set out to determine the spatial

relationship between the centrosome, centriculum and peri-centrosomal microtubules,

and to examine whether the centriculum functions as a microtubule filter.

Results

The spatial relationship between the centrosome, centriculum and microtubules.

In our previous study ((Maheshwari *et al.*, 2023), we showed that the centriculum

surrounds the centrosome and the peri-centrosomal microtubules (Fig. 1A), but the

exact spatial relationship between the three was not quantified. To do so, we examined

these three structures at metaphase of the *C. elegans* 1-cell embryo, when parental

genomes are still encased in separate pronuclei (Fig. 1A). In 1-cell embryos, centricula

become visible as soon as centrosomes separate (while still associated with the male

pronucleus), and both centricula and centrosomes increase in size throughout the cell

cycle (Hamill *et al.*, 2002; Maheshwari *et al.*, 2023). Consequently, size measurements

must be done at the same cell cycle stage, which in our case is at metaphase. At this

stage, the two pronuclei are adjacent to each other, and the membrane interface

between them is still present (Fig. 1A). Microtubules that are nucleated from the outer

portion of the PCM (O'Toole *et al*, 2012) extend outwards, towards the cortex (as astral

microtubules), or chromosome (as spindle microtubules). However, only a fraction of

these microtubules extends well beyond the centrosome (Redemann *et al.*, 2017),

creating an accumulation of peri-centrosomal microtubules (Fig. 1A, top panel). To

determine the spatial relationship between the centrosome, the centriculum and the

peri-centrosomal microtubules, we used confocal imaging to capture the areas

encompassed by the structures' perimeter(s) (Fig. 1B), visualized at their largest cross

section; as these structures are nested within each other, the plane of their largest

cross-section is shared. For the centrosome, we define "PCM area" as the area

occupied by SPD-5, the outermost component of the PCM (Magescas *et al.*, 2019). For

the centriculum, visualized using fluorescently tagged SP12 (Poteryaev *et al*, 2005), we

defined two areas: the area encompassed by the inner perimeter of the centriculum,

referred to as the centriculum "void area", and the area encompassed by the outer

perimeter of the centriculum, referred to as "centriculum outer perimeter area", or

"COPA" (note that the area of the centriculum itself is the difference between COPA and

the centriculum void area). For peri-centrosomal microtubules, visualized using

fluorescently tagged α or β tubulin, we defined the area encompassed by the outer

perimeter of the fluorescent tubulin signal as "the peri-centrosomal microtubule area"

(referred to in figures as "pcMT area", for simplicity). Although peri-centrosomal

microtubules often appear as a ring due to the exclusion of microtubules from the center

of the centrosome (Strome *et al*, 2001), the difference in fluorescence between the ring

and its "hole" is only about 10-15% (likely due to the high concentration of soluble

tubulin), making "hole" size determination somewhat ambiguous. Thus, for our

discussion, the area occupied by peri-centrosomal microtubules includes the "hole". The

comparison of areas defined by the perimeters of the three structures was done using

pairwise combinations; imaging all three structures at once was not feasible due to a

weak signal when proteins were fused to blue/cyan fluorescent proteins. Parenthetically,

we observed that the absolute size of various structures sometimes differed between

strains, even when they were otherwise isogenic (see, for example, the sizes of the void

area and COPA in Figs. 1C and D). Thus, the relative sizes of two structures were

determined by comparisons within the same strain. In the next set of experiments,

worms were treated with control RNAi so that the measurements could be compared to

the various RNAi treatments, described below.

The results of the pairwise combinations are shown in Figs. 1C-E and Fig. S1B. As

expected, the area defined by the outer perimeter of the centriculum, COPA, was the

largest. The area defined by the inner perimeter of the centriculum, the void area, was

the smallest. The area defined by the outer perimeter of peri-centrosomal microtubules

was greater than the area of the PCM (using two different fluorescently tagged protein

combinations, see Fig. 1E and Fig S1B), and both were greater than the centriculum

void area, suggesting that they abut or even extend into the centriculum. For

microtubules, we know this to be the case based on EM tomography data (Maheshwari

et al., 2023). Based on these data, an approximation of the overall spatial relationship

between the centrosome, centriculum and peri-centrosomal microtubules in wild type

cells is shown in Fig. 1F.

Decrease in microtubule number or stability results in a smaller centriculum and

PCM compression.

The proximity between the centrosome, centriculum and peri-centrosomal microtubules

raised the question of size dependency: Which of the three structures determines the

dimensions of one or both other structures? We previously showed that reducing the

levels of KLP-7, a microtubule depolymerase whose down-regulation leads to enhanced

microtubule outgrowth from centrosomes (Srayko *et al*, 2005), resulted in a larger

centriculum and a larger SPD-5 area (Maheshwari *et al.*, 2023). This suggested that

microtubule stability or abundance affects centriculum and/or PCM size. To further

examine this possibility, we examined the consequences of reduced microtubule

elongation or nucleation. To this end, we down-regulated by RNAi either *zyg-9* or *tbg-1*.

ZYG-9 is a microtubule polymerase and a homologue of *Xenopus* XMAP215 and

human ch-TOG. It localizes to centrosomes and the mitotic spindle (Gard & Kirschner,

1987; Matthews *et al*, 1998). In *Xenopus*, XMAP215 has been shown to bind free

tubulin dimers and catalyze their addition to the growing microtubule plus-end (Brouhard

et al, 2008). TBG-1 is the *C. elegans* γ-tubulin, which localizes to centrosomes and

facilitates microtubule nucleation (note that in *C. elegans*, TBG-1/ γ-tubulin is not

essential for microtubule nucleation at centrosomes (Hamill *et al.*, 2002; Hannak *et al*,

2002; Strome *et al.*, 2001). Both conditions are expected to reduce the amount of peri-

centrosomal microtubules (as determined by the total fluorescence of tubulin around the

centrosome), as is indeed the case (Fig. S1C). The area occupied by peri-centrosomal

microtubules was also reduced, as was the size of the centriculum (Figs. 2A-C; *zyg-9*

and *tbg-1* RNAi data are compared to control RNAi data also shown in Fig. 1C). These

results further support our conclusion that microtubule abundance and/or stability

affects centriculum size.

The reduction in centriculum size following *zyg-9* or *tbg-1* RNAi led us to examine what

happens to SPD-5 under these conditions. Using a strain expressing the SP12 ER

marker fused to mCherry and GFP::SPD-5, we observed that down-regulation of *zyg-9*

or *tbg-1* was accompanied by a decrease in centriculum size, as before (Fig. S1D), and

in PCM size (Figs. 2D and E). Interestingly, while the area occupied by SPD-5

decreased, there was no loss of SPD-5 from the PCM (Fig. 2F), and SPD-5's density in

the PCM increased (Fig. 2G). Overall, the increase in SPD-5 density was inversely

proportional to size of the centriculum void area (Fig. 2H; R^2=0.2624 for all data points).

This result suggests that centrosome size is affected by either microtubule abundance

and/or the size of the centriculum. This further suggests that the PCM is compressible

(see Discussion).

The *spd-5* expansion mutant, in the presence of wild type *spd-5*, reduces overall

PCM size.

The possibility that the centriculum affects centrosome size was intriguing, because the

centrosome is considered a membraneless organelle. In this scenario, centriculum size

is determined by the abundance and/or stability of microtubules, and the PCM expands

to the boundaries of the centriculum (Fig. 3Ai). Alternatively, however, centriculum size

could be determined by PCM size, while PCM size is determined by microtubules

abundance and/or stability (Fig. 3Aii). This latter scenario seemed less likely, because

PCM maturation is thought to be microtubule-independent (Hannak *et al.*, 2002;

Khodjakov & Rieder, 1999). Nonetheless, we needed to establish whether, in our

system, centriculum size was determined by the PCM or microtubules. To do so, we

needed a condition that differentially alters the areas occupied by the PCM vs peri-

centrosomal microtubules. We could then determine with which of the two centriculum

size correlates. This condition was met by using a *spd-5* allele that is defective in

expanding the SPD-5 lattice.

SPD-5 has been shown to have two separable functions: (1) centrosome maturation

and (2) microtubule nucleation via γ-tubulin recruitment (Ohta *et al.*, 2021). Both

functions are regulated by the PLK-1 kinase, which phosphorylates different residues on

SPD-5. Ohta et al created an RNAi-resistant *spd-5* phospho-mutant allele, *spd-5^{exp}*,

carrying S653A and S658A substitutions, which they expressed as a transgene in the

presence of the endogenous wild type, RNAi sensitive, *spd-5*. As a control, wild type

spd-5 was expressed as an RNAi-resistant transgene. As expected, when SPD-5^{exp} was

expressed as the sole SPD-5 version (i.e. endogenous *spd-5* was down-regulated by

RNAi), the expansion of the PCM was greatly attenuated compared to a strain

expressing transgenic wild type SPD-5 as its sole source of SPD-5 ((Ohta *et al.*, 2021)

and Figs. 3B and C). However, when transgenic SPD-5^{exp} was expressed in the

presence of the endogenous SPD-5 (which is untagged), transgenic SPD-5^{exp} occupied

a greater area than without the endogenous protein (compare transgenic GFP::SPD-

5^{exp} in Fig. 3C (no endogenous SPD-5) with Fig. 3D (with endogenous SPD-5); see also

Fig. S2A and B for more examples). This suggests that SPD-5^{exp} integrates into the wild

type SPD-5 lattice despite lacking two PLK-1 phosphorylation sites that contribute to

PCM expansion. This is consistent with in vitro findings showing that additional SPD-5

incorporates throughout an existing SPD-5 lattice (Woodruff *et al.*, 2017) and with

findings by Wueseke et al, who made similar observation with a *spd-5* allele lacking four

PLK-1 phosphorylation sites, including the two in SPD-5^{exp} (Wueseke *et al.*, 2016). As in

Wueseke et al, we also observed a good overlap between the endogenous SPD-5

(fused to tagRFP) and transgenic SPD-5 or SPD-5^{exp} proteins (both fused to GFP) (Fig.

3E), suggesting that both transgenes are integrated throughout the SPD-5 lattice.

Interestingly, the expression of the SPD-5^{exp} transgene in an otherwise wild type

background led to a smaller PCM, as judged by the areas occupied by tagRFP-tagged

endogenous SPD-5 (Figs. 3E-G). Furthermore, the amounts of both endogenous and

transgene SPD-5 at the centrosome were reduced in the presence of the *spd-5^{exp}*

mutant (Figs. 3 H and I). Thus, SPD-5^{exp} may act as a dominant negative, impeding the

ability of the SPD-5 lattice to fully expand. Alternatively, or in addition, the presence of

SPD-5^{exp} could have led to reduced overall levels of wild type SPD-5, as proposed by

Wueseke et al (Wueseke *et al.*, 2016), resulting in a smaller PCM. Note, however, that

the fluorescently tagged endogenous SPD-5 may exhibit a genetic interaction with the

expansion allele (see below).

Centriculum size is determined by peri-centrosomal microtubules, not the PCM.

Having a condition that reduced the size of the PCM, we next asked whether and how

the presence of the *spd-5^{exp}* mutant affected the area occupied by peri-centrosomal

microtubules. This was done using strains expressing wild type or expansion mutant

spd-5 transgene fused to GFP in the presence of endogenous (untagged) SPD-5, and

α-tubulin fused to tagRFP (*tba-1::tagRFP*-T; Fig. 4A). Despite having a smaller PCM in

the presence of *spd-5^{exp},* as shown before (Fig. 4B), the amount of tubulin withing the

peri-centrosomal region was comparable in the presence of either transgene (Fig. 4C).

Moreover, the area occupied by peri-centrosomal microtubules did not decrease due

the presence of the expansion mutant transgene (Fig. 4D). As before (Fig. 1E and Fig

S1B), peri-centrosomal microtubules occupied a greater area than SPD-5. This area

differential increased in the presence of the *spd-5^{exp}* mutant (Fig. 4E) due to the smaller

area occupied by SPD-5 and an unchanged area occupied by peri-centrosomal

microtubules. Thus, peri-centrosomal microtubules were less affected by the presence

of the SPD-5^{exp} variant compared to the PCM.

This differential change in size of the PCM and peri-centrosomal microtubules due to

spd-5^{exp} allowed us to examine whether centriculum size follows the PCM or

microtubules. Specifically, by decreasing PCM size, the *spd-5^{exp}* allele increased the

distance (Δ) between the perimeter of the PCM and that of the peri-centrosomal

microtubules (Fig. 5Ai). We reasoned that measuring the effect of the *spd-5^{exp}* allele on

the centriculum void area (determined by the inner perimeter of the centriculum) would

allow us to determine if centriculum size tracks with the PCM or the peri-centrosomal

microtubules (Fig. 5Aii). To do so, we used the same pairwise combinations used to

determine the spatial relationship between the centriculum and the PCM or peri-

centrosomal microtubules (Fig. 1B-F), this time in the presence of the wild type or *spd-*

5^{exp} mutant transgene.

In strains expressing fluorescently tagged SP12 and TBA-1, the *spd-5^{exp}* mutant led to a

decrease in the area occupied by peri-centrosomal microtubules (Figs. 5B and C) but

not in the amount of peri-centrosomal tubulin fluorescence (Fig. S3A). This decrease in

peri-centrosomal microtubules area was unlike the situation in strains expressing

fluorescently tagged SPD-5 and TBA-1, where *spd-5^{exp}* did not affect peri-centrosomal

microtubule area (Figs. 4B and C). The reason for this difference is not known.

Nonetheless, *spd-5^{exp}* also led to a comparable decrease in centriculum size, as seen

by the decrease in void area (Fig. 5D) and the area encompassed by the centriculum

outer perimeter, COPA (Fig. S3B). Importantly, the reduction in microtubule and void

areas were the same (Fig. 5E, control RNAi). In contrast, when we examined the effect

of the *spd-5^{exp}* transgene on the centriculum in relation to the PCM using strains

expressing mCherry-tagged SP12 and GFP-tagged wild type or *spd-5^{exp}* transgenes,

the effect of *spd-5^{exp}* on the PCM was significantly greater than on the centriculum: in

these strains, *spd-5^{exp}* caused the area of the PCM, but not the centriculum, to shrink

(Figs. 5F-I, control RNAi). In fact, in several cases the size of the PCM was smaller than

that of the centriculum void (see negative values in Fig. 5I). Taken together, these data

suggest that centriculum size follows the behaviors of the peri-centrosomal microtubules

rather than the PCM.

To further examine the dependence of centriculum size on microtubules vs. the PCM,

we stabilized microtubules in strains expressing wild type or spd-5^{exp} transgene using

RNAi against klp-7. We reasoned that if centriculum size is determined by microtubules,

stabilizing microtubules in the presence of the spd-5^{exp} transgene might further increase

the distance between the PCM and the inner perimeter of the centriculum. As expected,

(Maheshwari et $al.$, 2023), klp-7 RNAi led to expansion of the centriculum, PCM and

peri-centrosomal microtubule area (for PCM and peri-centrosomal microtubules areas,

compare the values in Fig. 4F (klp-7 RNAi) with Fig. 4B and 4D (control RNAi). The

effect of klp-7 RNAi on the PCM in a different strain is shown in Fig. 5G; for the

centriculum void area and COPA, see Fig. 5H and Fig. S3B). Following klp-7 RNAi

treatment, the effect of spd-5^{exp} was again greater on the PCM than on the

pericentrosomal microtubules (Figs. 4F and G), while area differential between the

centriculum void area and the peri-centrosomal microtubules remained the same (Fig.

5E; pcMT and void areas in the presence of klp-7 RNAi are shown in Fig. S3C).

However, when strains expressing the transgenic wild type or spd-5^{exp} mutant and

mCherry::SP12 were treated with RNAi against klp-7, a visible "gap" appeared between

the PCM and the inner perimeter of the centriculum (Fig. 5F, see enlarged area). While

klp-7 RNAi led to an overall increase in PCM area, the presence of the spd-5^{exp} allele

led to reduced PCM area (Fig. 5G) without affecting the centriculum void area (Fig. 5H).

Consequently, stabilizing microtubules through klp-7 RNAi exacerbated the difference

between the void area and the PCM such that in all cases, the PCM was smaller than

the inner perimeter of the centriculum (Fig. 5I, note negative values in the presence of

the spd-5^{exp} allele following klp-7 RNAi). In fact, klp-7 RNAi alone was sufficient to

increase the area differential between the void area and the PCM (Fig. 5I). Taken

together, these data suggest that the centriculum behaves the same as microtubules,

not the PCM, and that centriculum size is determined by microtubules.

In the experiments described above, the size of the PCM was assessed by the

fluorescently tagged transgenes. While there was good agreement between the areas

occupied by the transgenic and the endogenous SPD-5s (Fig. 3F and G), we attempted

to examine the spatial relationship between the centriculum and the PCM using an

endogenously tagged SPD-5. Unexpectedly, the spd-5^{exp} transgene and the

endogenously tagged spd-5 exhibited a genetic interaction, resulting in abnormal

distribution of both proteins (Figs. 6A and Fig. S4), reminiscent of SPD-2 distribution

(see for example (Pelletier et al, 2004)). We assume that when the endogenous spd-5

was not tagged, its distribution was uniform, as judged by the uniform distribution of the

GFP::SPD-5^{exp} transgene, which on its own cannot expand (Fig. 3C and D and Fig.

S4A). However, when the endogenous spd-5 was fused to a fluorescent protein, its

distribution in the presence of the transgenic spd-5^{exp} mutant becomes abnormal, with a

high concentration in the center of the centrosome, and a much lower concentration in

the majority of the PCM (Fig. S4B and C). Under these conditions, the distribution of the

GFP::SPD-5^{exp} transgene was similarly abnormal (Fig. S4B), in further support of the

similar distributions of the endogenous and transgene proteins. Despite this abnormal

distribution, we were still able to determine the area occupied by endogenously

expressed SPD-5, including the fainter periphery, using the same thresholding method

applied to all other area measurements (Fig. S4D). As before, the presence of the spd-

5^{exp} mutant transgene decreased the amount of SPD-5 at the centrosome, and this

pattern was unaffected by down-regulating klp-7 (Fig. 6B). Also as before, down-

regulation of klp-7 increased PCM area, and the presence of the spd-5^{exp} mutant

decreased PCM areas relative to the wild type transgene in both control and klp-7 RNAi

conditions (Fig. 6C). Consistent with the observations above, the presence of the spd-

5^{exp} mutant did not reduce the size of the centriculum void volume, and down-regulation

of klp-7 even increased it (Fig. 6D). Consequently, the area differential between the

PCM and the centriculum increased in the presence of the spd-5^{exp} mutant, and this

difference was further exacerbated when klp-7 was down-regulated (Fig. 6E).

Taken together, these data suggest two things (Fig. 6F): First, that centriculum size is

likely determined by sum forces exerted on it by peri-centrosomal microtubules.

Second, because the area occupied by pericentrosomal microtubules change little or

none when the PCM "shrinks" due to the presence of the *spd-5^{exp}* mutant, and because

microtubules nucleate from the surface of the PCM, our data suggest that the length of

pericentrosomal microtubules is not an inherent property of these microtubules,

because their length, from the surface of the PCM to the inner perimeter of the

centriculum, must be different in wild type vs *spd-5^{exp}* expressing cells (Fig. 6F). Rather,

our data suggest that the majority of pericentrosomal microtubules extend from the

surface of the PCM until they encounter the centriculum, where most of them terminate

due to obstruction by the centriculum membrane; only a small fraction can extend past

the centriculum to form astral or spindle microtubules. As such, the centriculum may

serve as a microtubule filter, a possibility that is explored next.

Centriculum porosity correlates with the ability of microtubules to traverse the

centriculum.

As noted above, two types of centrosome-nucleated microtubules extend beyond the

centriculum: astral microtubules, which extend towards the cell cortex, and spindle

microtubules, which extend towards chromosomes. The density of spindle microtubules

is much greater than astral microtubules (Fig. 7A, metaphase). Spindle microtubules are

only evident in metaphase; prior to that, only astral microtubules can be observed

outside the centriculum, and no microtubules have entered the pronuclei (Fig. 7A,

prophase). If the centriculum serves as a microtubule filter, we would expect that at

metaphase, the centriculum would be more porous on the side facing the chromosomes

vs. the side facing the cortex. To examine this possibility, we measured the effective

pore size that is available for microtubules nucleated at centrosomes to pass through

the centriculum using volume electron microscopy data acquired via focused ion beam-

scanning electron microcopy (FIB-SEM; (Maheshwari *et al.*, 2023)).

We created a fiducial in the center of segmented metaphase centricula, and imaged a

portion of the centriculum, from that fiducial, against a black background (Fig. 7B and C;

images were taken from within the centrosome towards either the cortex or the

chromosomes). Qualitatively, it already appeared that the holes on the chromosomal

side are larger than those on the cortical side (compare the two central panels,

chromosome-facing 1 and 2, to the side panels, cortex-facing 1 and 2). To maximize

having the openings imaged *en face*, the angles between the center line and the top or

bottom of any image were kept to a minimum (for cortex facing areas: 23.8° ± 2.82°,

average and standard deviation, n=6; for chromosome facing areas: 22.8° ± 2.23°,

average and standard deviation, n=6). We then determined the area of each opening,

defined as "hole", that can be seen through the centriculum. Considering that the

diameter of a microtubule is ~25 nm, the area of the smallest hole through which a

microtubule can pass would be 490 nm^2 (π x 12.5^2) and probably larger since most

holes are not perfect circles. The fraction of cortex-facing centriculum area that was

open, namely the sum of all hole areas out of the total area imaged, was 7.85 ± 1.66%

(n=6, from 3 centricula), while on the chromosome side it was significantly greater:

20.66 ± 3.75%, (n=6 from 3 centricula; p=0.009 as determined by chi square test),

consistent with the chromosome-facing side of the centriculum having larger opening. In

the case of cortex-facing holes, the cross sections of 61.34% of the holes, accounting

for 7.5% of the total open area, were below 500 nm^2, likely too small to allow for a

microtubule to pass (Fig. 7D and E). The percent of chromosome facing holes that had

a cross section below 500 nm^2 was 41.88%, and this accounted for only 1.4% of the

total open area on the chromosomal side (Fig. 7D and E). On the other side of the hole-

size spectrum, 11.56% of the open area facing the cortex were from holes that were

larger than 10,000 nm^2 (0.8% of all holes), compared to 74.52% on the chromosome-

facing side (9.1% of all holes) (Fig. 7E). Thus, the centriculum side facing the

chromosomes, through which spindle microtubules pass, is significantly more porous

than the side facing the cortex, where astral microtubules pass. Whether the different

degrees of porosity are caused by centriculum geometry or by properties of spindle

microtubules remains to be determined (see Discussion).

Given that only a small fraction of the centriculum area facing the cortex is available for

the microtubules to go through, and the fact that the thickness of centriculum at

metaphase is around 1 μm (Maheshwari *et al.*, 2023), we predicted that the

microtubules that become fully elongated astral microtubules are those that take the

shortest path through the centriculum, and are thus less likely to encounter centriculum

membrane. To test this hypothesis, we used ER tomography datasets of 1-cell embryos

at metaphase (Maheshwari *et al.*, 2023; Redemann *et al.*, 2017) and measured the

angle of microtubules as they pass the inner perimeter of the centriculum relative to the

radial vector, the line that is perpendicular to the tangent line (Fig. 8A). The shortest

path through the centriculum would be that of the radial vector (angle= 0°); increasing

angles between a microtubule path and the radial vector would mean longer paths

through the centriculum. We then determined whether the microtubule passed through

the centriculum (denoted as "continuing") or whether the microtubule terminated within

the centriculum (denoted "stopped"). On the cortex-facing side, we measured angles for

113 microtubules, of which 21 continued (18.6%), and on the chromosome-facing side

we measured angles for 146 microtubules, of which 40 continued (27.4%). In both

cases, the distribution of angles of microtubules that passed all the way through the

centriculum was significantly narrower, and closer to the radial vector, than the range of

angles for microtubules that ended within the centriculum (Fig. 8B). Nonetheless, our

data suggest that the centriculum serves as a passive microtubule filter, blocking most

microtubules from extending past the centriculum.

To form a spindle, microtubules must enter the space around the chromosomes. Prior to

metaphase, an intact nuclear envelope prevents centrosome-nucleated microtubules

from entering the nucleoplasm. Just before metaphase, two events take place: the

nuclear envelope loses its integrity specifically next to centrosomes (Hachet *et al*,

2012), and the centriculum fuses with the nuclear membrane (Maheshwari *et al.*, 2023).

In this regard, the larger hole size of the centriculum on the chromosomal side is

unlikely to be the only reason why spindle microtubule density is greater than that of

astral microtubules. At prometaphase, the percent of open centriculum area (14.64 ±

4.14%, n=6 from 3 centricula), distribution of hole size, and percent area per hole size

(Fig. S5A and B) on the chromosomal side of the centriculum were similar to those of

metaphase centricula. Effectviely, however, the number of holes available for

microtubules to reach the chromosomes is significantly smaller, because at

prometaphase, remnants of the nuclear membrane are still present, creating an

additional barrier for microtubules to enter the nucleoplasm (Fig. 8C, note the presence

of nuclear membrane (in green) masking many of the openings). At prometaphase the

centriculum is adjacent to the nuclear membrane, with a few possible contact sites (see,

for example, Fig. 8D), but there are no membrane junctions that indicate actual fusion.

In contrast, at metaphase, the centriculum is completely fused with the remnants of the

nuclear membrane (Maheshwari *et al.*, 2023), possibly in an atlastin-dependent manner

((Araújo *et al.*, 2023) and see below). Therefore, at metaphase, there are no other

membranes between the centriculum and the chromosomes. Thus, the fusion of the

centriculum and mitotic remnants of the nuclear membrane may reduce membrane

barriers in the process of spindle formation.

A further indication that centriculum-nuclear membrane fusion is important for spindle

microtubule formation is the state of metaphase 1-cell embryos following atlastin down-

regulation. Atlastin is a GTPase needed for the fusion of ER tubules (Hu & Rapoport,

2016; Orso *et al*, 2009). We down-regulated atlastin using a degron construct, under

conditions that partially lower the amount of atlastin in the embryo (Maheshwari *et al.*,

2023). Under these conditions, for unknown reasons, pronuclei have difficulties aligning,

and they sometimes fuse with only one of the two centricula (Fig. 8E, see pronuclei

marked in blue asterisks in panels i and iii), leading to the formation of monopolar

spindles in those pronuclei. Centricula at metaphase are normally fused to both

pronuclei, resulting in two dense clusters of spindle microtubules on either side of the

membrane between the two pronuclei (for example, see Fig. 7A). However, when *atln-1*

is down-regulated, a centriculum may fuse with only one of the two pronuclei (Fig. 8E,

red arrows). Under these conditions, dense microtubules can be observed only where

the centriculum is fused to the pronuclei, while the rest of the centriculum is surrounded

by lower density, astral microtubules (Fig. 8E and enlarged areas from panels iii and iv).

This suggests that robust microtubule elongation past the centriculum to form the

spindle's dense microtubule array can only happen once the centriculum fuses with a

pronucleus. Taken together, we proposed that formation of high-density microtubules on

the chromosome-facing side of the centriculum is facilitated by two processes: the

presence of large openings that traverse the centriculum, and the fusion of the

centriculum with remnants of the nuclear membrane.

Discussion

Discussion

In this study, we tested the hypothesis that the centriculum acts as a partial barrier, or

filter, preventing the extension of some, but not all, centrosome nucleated microtubules.

Our measurements show that the PCM and peri-centrosomal microtubules either abut

or slightly penetrate the inner perimeter of the centriculum (Fig. 1). We further observed,

both here and in our previous study, that changing microtubule stability or number

affects both PCM size and centriculum size ((Maheshwari *et al.*, 2023) and Figs 2, 5G-I,

6 and Figs. S1 C and D, and S3)). This suggested that microtubules affect centriculum

size either directly, potentially by exerting force on the centriculum and thus determining

its size, or indirectly, by affecting PCM size, which in turn affects centriculum size. To

gain insight into the spatial dependencies of these three structures, we used a *spd-5*

allele, *spd-5exp*, that is defective in PCM expansion (Ohta *et al.*, 2021). Expressing both

spd-5exp, as a transgene, and the endogenous wild type *spd-5* resulted in a smaller

PCM. An inhibitory effect by SPD-5 mutant proteins on the SPD-5 lattice has been

observed previously: Nakajo et al. observed this phenotype with a 272 amino acid C-

terminal deletion, which did not encompass the residues mutated in SPD-5exp (Nakajo *et*

al., 2022), while Wueseke et al. observed it with a SPD-5 variant lacking four PLK-1

phosphorylation sites, SPD-54A, including the two in SPD-5exp (Wueseke *et al.*, 2016).

Rios et al found that PLK-1 phosphorylation of SPD-5 reduces intramolecular

interactions, allowing these domains to interact intermolecularly (Rios *et al.*, 2024). We

therefore propose that the SPD-5exp variant acts as a dominant negative by

incorporating into the SPD-5 lattice but not contributing fully to intermolecular

connections, thereby impeding lattice expansion. Alternatively, or in addition, the

presence of transgenic mutant *spd-5* may have led to reduced levels of endogenously

expressed *spd-5* (Wueseke *et al.*, 2016), thus resulting in a smaller PCM.

Importantly, expressing both transgenic *spd-5exp* together with the endogenous wild type

spd-5 resulted in a smaller PCM without reducing the area occupied by the peri-

centrosomal microtubules (Fig. 4A-D). The same was true when worms were treated

with RNAi against *klp-7*, which stabilizes microtubules: while the areas occupied by the

PCM and peri-centrosomal microtubules was overall greater due to *klp-7* down-

regulation (Maheshwari *et al.*, 2023), only PCM area, but not peri-centrosomal

microtubule area, was affected by the presence of SPD-5^{exp} (Fig 4F). In other words,

expression of the *spd-5^{exp}* allele led to a greater difference in the areas occupied by

peri-centrosomal microtubules compared to the PCM (Fig 5E and G). This experimental

setup allowed us to address two questions: the first was what determines centriculum

size. By examining centriculum size relative to the PCM vs per-centrosomal

microtubules in the presence of *spd-5^{exp}*, we showed that the centriculum correlates

with microtubules, not the PCM (Fig 5). In fact, under *klp-7* RNAi treatment, a gap was

visible between the inner perimeter of the centriculum and the PCM (Fig 5F). The

increase is distance between the centriculum and the PCM was true regardless of

whether PCM area was determined using the transgene or endogenous SPD-5 (Fig 6

E). Thus, our data are consistent with the possibility that centriculum size is determined

by microtubules, not by the PCM. This further suggests that PCM size could be affected

by the size of the centriculum, such that, for example, when the centriculum is unusually

small due to reduced microtubule number or stability, the centriculum could lead to PCM

compaction (Fig 2). This is an intriguing possibility, as it suggests that the PCM lattice is

malleable, able to expand or fold inwards in response to forces acting upon it, akin to a

Hoberman Sphere (https://www.hoberman.com/portfolio/hoberman-sphere-toy/). That

the PCM is malleable was also proposed by Laos et al, who noted that the PCM lattice

in vivo can grow by incorporating SPD-5 anywhere within the lattice (Laos *et al.*, 2015).

This property could contribute to the PCM's ability to deform in response to forces

exerted on it by astral microtubules (Enos *et al.*, 2018), where the PCM could stretch

rather than break. This property, known as ductility, was also proposed by Mittasch et al

based on the PCM's deformability during anaphase in response to external forces

(Mittasch *et al.*, 2020).

The second question this experimental set up allowed us to address is whether peri-

centrosomal microtubules are short due to an inherent property. Numerous in vitro

studies, using tubulin concentrations that were significantly lower than in the 1-cell *C.*

elegans embryo, have examined microtubule length either directly or indirectly, and in

none of them were microtubules as short as the *C. elegans* 1-cell peri-centrosomal

microtubules (Arnal *et al*, 2004; Belmont *et al*, 1990; Chrétien *et al*, 1995; Farrell *et al*,

1987; Fygenson *et al*, 1994; Gould & Borisy 1977; Grego *et al*, 2001; Kuriyama &

Borisy, 1981; Mitchison & Kirschner, 1984; Verde *et al*, 1990), suggesting that

something is preventing the extension of peri-centrosomal microtubules. Ideally, we

would have liked to remove the centriculum entirely and examine the effect on peri-

centrosomal microtubule length. Unfortunately, we have yet to identify a condition that

eliminates the centriculum. Nonetheless, the *spd-5^{exp}* allele created a condition where

limits to peri-centrosomal microtubule length could be examined. Microtubules are

nucleated from the outer portion of the PCM. In the presence of the *spd-5^{exp}* allele, the

PCM is smaller, but the area occupied by peri-centrosomal microtubules, as determined

by the outer periphery of this structure, remains unchanged (Fig 4B, D and F). This

suggests that in the presence of the *spd-5^{exp}* allele, peri-centrosomal microtubules are

longer than in the control (Fig 6F). If the length of peri-centrosomal microtubules were a

function of an inherent property of these microtubules, the area occupied by these

microtubules would have been smaller in the presence of SPD-5^{exp}. Thus, we propose

that the accumulation of peri-centrosomal microtubules is a consequence of the

presence of the centriculum, which blocks the extension of a subset of microtubules that

are nucleated by the centrosome. While elongating microtubules have the capacity to

distort a membrane sheet (for example, see (Zimmerman & Chang, 2005)), we propose

that the reticular nature of the centriculum, creating a membrane lattice, allows it to

resist forces exerted on it by microtubules. Consistent with this possibility, the

microtubules that manage to pass through the centriculum take, on average, a shorter

route than microtubules that stop within the centriculum (Fig 8A and B). If microtubules

were agnostic to the presence of the centriculum, there would be no difference in the

angle relative to the radial vector of microtubules that terminate within the centriculum

and those that extend past it. A similar accumulation of peri-centrosomal microtubules

was observed in other organisms that also have peri-centrosomal ER accumulation

(Kiyomitsu *et al.*, 2024; Rollins & Blankenship, 2023; Xie *et al.*, 2025).

While the centriculum may act as a barrier to the extension of peri-centrosomal

microtubules, it is still permissive to extension of astral and spindle microtubule

extension. The density of microtubules extending toward the chromosomes by far

exceeds the density of microtubules extending toward the cortex (Figs. 1A and 7A). This

difference in density correlates with the size of centriculum "holes" and the overall open

area, which is significantly greater on the side of the centriculum that faces the

chromosomes compared to the side facing the cortex (Fig 7). We imagine two non-

mutually exclusive reasons why centriculum holes are larger on the side facing the

chromosomes: first, prior to metaphase, this side of the centriculum is adjacent to the

pronuclear membrane and is not as wide as the parts facing the cytoplasm (Maheshwari

et al., 2023), perhaps due to a spatial restriction of ER-derived membrane

accumulation. If we assume that the centriculum is a passive barrier, the wider it is, the

more microtubules it will block from extending. Second, the centriculum is likely a

dynamic structure, much like the rest of the ER. At metaphase, microtubules that reach

the chromosomes are stabilized, likely restricting the ability of the centriculum to

remodel around these microtubules. Under these conditions, it is possible that additional

microtubules can pass alongside the stable microtubule, further reducing the ability of

the centriculum to remodel at that site and establishing a "hole" whose size may

continue to increase. Consistent with this possibility, we observe a high density of

microtubules extending past the centriculum only after the centriculum fuses with

remnants of the pronuclear membranes (Fig 8E). This explanation may also apply to

cortical microtubules, which although not stabilized, may still promote the passage of

additional microtubules through the centriculum alongside them. Indeed, the distribution

of astral microtubules is not uniform (Fig. 7A), perhaps due to the filter properties of the

centriculum.

Taken together, we propose that the centriculum acts as a partial barrier, or filter, of

centrosome-nucleated microtubule, blocking the extension of many centrosome-

nucleated microtubules, thereby limiting the number of astral and spindle microtubules.

This may limit competition for limited cytoplasmic pools of soluble tubulin, which might

only be sufficient for a limited number of astral microtubules to extend all the way to

cortex. Likewise, an over-abundance of microtubules might interfere with proper spindle

formation or function. Moreover, the termination of peri-centrosomal microtubules at the

centriculum would likely result in microtubule catastrophe, creating an environment of

high tubulin concentration, as observed by Baumgart et al (Baumgart *et al.*, 2019); the

free tubulin is then captured by the PCM, which is capable of concentrating tubulin *in*

vitro (Woodruff *et al.*, 2017). Thus, the barrier properties of the centriculum could explain

both the accumulation of short peri-centrosomal microtubules and the accumulation of

soluble tubulin at the centrosome.

To date, the only centrosome-adjacent membrane that has been examined at

nanometer resolution is the *C. elegans* centriculum. Similar membrane structures,

however, have been observed in other cell types (Araújo *et al.*, 2023; Bergman *et al.*,

2015; Bobinnec *et al.*, 2003; Diaz *et al.*, 2019; Harris, 1975; Karabasheva & Smyth,

2019; Terasaki, 2000; Waterman-Storer *et al.*, 1993). In many cases, centrosome-

associated membranes are most prominent in the embryo (e.g. *C. elegans*, *Drosophila*

and sea urchin). Early embryos typically contain stores of protein and mRNA that are

used in the first few embryonic divisions, before the onset of zygotic expression. The

centriculum and other centrosome-associated membranes may help limit the number of

astral microtubules under conditions where microtubule nucleating capacity exceeds the

needs of the embryo. Given the overall conservation in centrosome and ER-associated

proteins, it is possible that centrosome structure and function in additional cell types are

also affected by an adjacent ER-derived membrane reticulum. If this is the case, then

the repertoire of proteins and processes that affect centrosome function is broader than

previously appreciated, and defects in these proteins or processes might be a basis for

centrosome-related human disease.

Materials and Methods

C. elegans strains

The *C. elegans* strains used in this study were derived from the N2 strain (Bristol; (Brenner, 1974) and are listed in Table S1. Strains were maintained at 20°C on OP50 unless treated with RNAi (see below), using standard methods (Brenner, 1974).

Manipulations of gene and protein expression

Feeding RNAi

E. coli RNAi feeding clones against *smd-1* F47G4.1, *klp-7* K11D9.1, *zyg-9* F22B5.7 and *tbg-1* F58A4.8 were from a *C. elegans* RNAi feeding library (Open Biosystems, Huntsville, AL). For *klp-7*, *tbg-1*, *zyg-9* or *smd-1* (control (Golden *et al*, 2009)) feeding RNAi treatments, a 5 ml Luria Broth (LB) with 50 μg/ml ampicillin (Sigma Aldrich, Cat # A9518) were inoculated using 1:100 dilution of an overnight saturated culture (at 37°C) of *E. coli* expressing dsRNA of the gene of interest. Once the culture grew to OD_{600} of around 0.5 ($\sim$ 4 h at 37°C), 0.5 M IPTG (Sigma Aldrich, Cat # I6758, 1 mM final concentration) was added to induce the bidirectional transcription of the relevant gene for another 4 h. The culture was centrifuged at 5000 g for 5 min at room temperature, and the pellet was resuspended in 1 ml of fresh LB + ampicillin (50 μg/ml) media. 200 μl of this culture were spread on each RNAi plate (MYOB with 4 mM IPTG and ampicillin 50 μg/ml). For feeding RNAi treatment, 20-40 L4-stage larvae were transferred to RNAi plates at 20°C, and after 48 h (for *klp-7*), or 24 h (for *tbg-1* and *zyg-9*), the RNAi treated worms were dissected on a glass slide, mounted on a 2% agar pad, and imaged as described below. Control RNAi treatments (*smd-1*) were done for the same amount of time as the experimental ones.

dsRNA injection RNAi

RNAi against *spd-5* was done using injection of double stranded RNA (dsRNA). A region of *spd-5* from 500-949 bp, was PCR amplified from N2 genomic DNA using oligonucleotides containing T7 or T3 promoter sequence. The amplified region was gel purified and then reamplified with the same PCR primers. The PCR product was purified using the Qiagen MinElute Reaction Cleanup Kit (Cat # 28206). To prepare the RNA for

injection, in vitro RNA synthesis was carried out using MEGAscript™ T7 Transcription

Kit (Invitrogen Cat # AM1333; for forward strand) and MEGAscript™ T3 Transcription Kit

(Invitrogen Cat # AM1338; for the reverse strand) followed by purification using

Phenol:CHCl3:Isoamyl Alcohol (invitrogen Cat # 15593031; 25:24:1, v/v) and

precipitated using 100% ethanol. The RNA pellet was dissolved in water. To prepare

dsRNA, an equal amount (2 µg/µl) of both the ssRNAs were mixed and incubated at

85°C for 3 minutes in an aluminum heat block incubator followed by slow cooling to

room temperature for annealing. Injection of dsRNA was done according to Ohta

laboratory protocol (Ohta *et al.*, 2021). L4s (15-20 worms) were injected with ~1 µg/µl

dsRNA. These worms were maintained at 16°C for 48 h prior to live imaging of early

embryos by confocal microscopy.

Auxin-mediated degradation

Auxin mediated degradation of ATLN-1 was done as described previously (Maheshwari

et al., 2023). The *atln-1* gene was tagged with an auxin-inducible degron tag (atln-

1::degron) in cells expressing TIR1, an exogenous F-box protein. Worms were

transferred to bacteria seeded indole-3 acetic acid (IAA, Alfa Aesar Cat # A10556)

plates (MYOB plates with 4 mM IAA) for ~20-25 minutes, and embryos were imaged

immediately thereafter, as per Zhang et al 2015 (Zhang *et al*, 2015).

Confocal microscopy

Imaging

Images were taken using a Nikon confocal Ti2 with Yokagawa CSU-X1 spinning disk

and a photometrix Prime 95B camera using a Nikon water/oil 60X 1.2-NA Apo Plan

objective. Images were captured using Nikon Elements software version 5.20.00. For

imaging, embryos were mounted on 2% agarose pads prepared in standard M9 buffer.

Images were taken at z=1 µm intervals unless otherwise mentioned.

Image analysis

All images were analyzed using Fiji (Schindelin *et al.*, 2012), http://imagej.nih.gov/ij).

Measurements:

Measurement of areas and amount

To compare between structures, images were taken by confocal microscopy as described above, and the comparisons between relevant structures were done at their midplane, where these structures is at its largest cross section. Since the centriculum, peri-centrosomal microtubules and centrosome are symmetrically nested within each other, the midplane of any two structures was at the same image plane. To determine the area of a given structure, the boundaries of structure were roughly traced. In the case of the PCM this involved tracing the outer perimeter of SPD-5. For the centriculum and peri-centrosomal microtubule ring, this involved tracing both the inner and outer perimeters of the ring, excluding nuclear membranes and spindle microtubules, respectively, resulting in a toroidal shape (Fig. S1A). For each traced area, the minimum (min) and mean fluorescent values were determined. Empirically, the most consistently faithful representation of the different structures' areas was by using the formula (mean+min)/2 to set the lower threshold value, and Max (maximum threshold, found in same measurement results FIJI that also included mean and minimum intensity values) to set the upper threshold. These values were entered into the threshold function of FIJI, resulting in consistently accurate representation of the structure's area (in the case of the microtubule ring, since the difference in intensity between the ring and its center is only about 15%, this method did not result in a ring but rather a solid round area) (Fig. S1A). Of note, while in the present study the min/mean ratios were determined for each structure in each image, these values are highly consistent, and once determined can be applied to a given structure. For example, the min/mean ratio of the traced centriculum region for mCherry::SP12 were 0.79 ± 0.03 (control RNAi; n=16) and 0.76 ± 0.04 (*klp-7* RNAi; n=16); for GFP:: SP12 they were 0.75 ± 0.05 (control RNAi; n=16) and 0.74 ± 0.03 (*klp-7* RNAi; n=20).

For the PCM and peri-centrosomal microtubules, the area and total fluorescence (Raw Intensity) were measured in the thresholded region using the analyze > measure function in FIJI. For the centriculum, the inner perimeter of the thresholded centriculum was selected using wand tool in FIJI to get the void area. For the area defined by the

outer perimeter, we traced the thresholded outer edge of the centriculum thereby

determining the area bound by the centriculum's outer perimeter.

Microtubule angle measurement

In a zoomed-in screenshot of a 240-slice-thick section of the tomography data

containing the area of interest (Maheshwari *et al.*, 2023; Redemann *et al.*, 2017)

originally from Amira, two straight lines were drawn, marking the approximate interior

and exterior edges of the centriculum (Fig. 8A; while the edges of the centriculum are

curved, at this resolution the edge approaches a straight line that represents the tangent

line of the centriculum edge). Microtubules that crossed the interior edge were traced

using the straight line tool in FIJI. If a microtubule continued beyond the exterior

boundary line of the centriculum, it was categorized as a "continued MT." Microtubules

that stopped before reaching the exterior boundary were categorized as "stopped MT."

To ensure that microtubules defined as "stopped" did not continue outside the range of

the slice that was being analyzed, screenshots were taken of 300 slices above and

below the same area of the initial screenshot. These images were used to determine

whether the traced microtubules continued in another plane. Those that did not continue

on either the above or below slice set were considered stopped.

To measure microtubule angles, the angle between the inner centriculum edge line (the

tangent line) and the microtubule line was determined. This angle was then subtracted

from 90 degrees. The resulting value represents the angle between the microtubule and

the perpendicular to the tangent line, also referred to as the normal line (Fig. 8A).

Focused-Ion beam scanning electron microscope (FIB-SEM) imaging

The FIB-SEM data used in this study was previously published (Rahman *et al*, 2020).

3D segmentation of centriculum structure were done as previously published (Rahman

et al., 2020) using Amira 6 (version 3D 2022.2, FEI/Thermo Fisher Scientific) default

general scheme to segment selected ROI in a semi-autonomous method. Briefly, first,

we selected centriculum membranes using a threshold followed by at least three rounds

of manual inspection (slice by slice through X, Y, and Z separate planes) to remove

incorrect segments and add unsegmented membranes to the final membrane volume

(Maheshwari *et al.*, 2023; Rahman *et al.*, 2020).

To measure the size of centriculum openings (namely open space between reticular

membranes that extend all the way through the centriculum), we placed fiducials (on the

same plane) next membrane edges on the pronuclear and cortical size of the

centriculum. We took TIFF screenshots of the centriculum membrane from behind a

central fiducial (i.e., equal distance from the pronuclear and cortical centriculum

membrane fiducials) placed on the same plane as perinuclear/cortical fiducials. For

each TIFF image, we determined a scale bar separately by marking 11 consecutive

pixels as 100 nm. We measured the size of each hole using the threshold-based

selection and area measurement tool in Fiji (version 2.14.0/1.54f). To ensure that the

centriculum segments in these images are *en face*, we restricted the angle between the

center plane and the top and bottom of each image to less than 30°. The actual angle

was determined by drawing straight lines from the central fiducial to fiducials that were

placed on the top bottom edges of each image. Next, we took TIFF images of the

perpendicular view of the above-mentioned lines and utilized the angle measurement

tool in Fiji.

Statistical analysis

All analyses were done using GraphPad Prism (GraphPad Prism [Version 10.1.1

(270)]). For comparisons between two normally distributed samples, unpaired t-tests

were used; otherwise, a two-tailed Mann-Whitney test was used as a non-parametric

test. For comparison between multiple datasets, one-way ANOVA non-parametric

Kruskal-Wallis test with correction for multiple comparisons was used. The statistical

tests used, the number of samples analyzed, and the p-values are indicated in the

Figure legends. The criterion for statistical significance was set at $p < 0.05$. Results in all

graphs are represented as mean ± standard deviation unless indicated otherwise. The

binned frequency distribution graphs in Fig. 7, 8 and Supplemental Figure S3 were done

using Microsoft excel. The bin width was set at 500 nm^2 up to 1000 nm^2 then from 1001

to 10000 the data were binned at every 1000 nm^2 with overflow bin set at >10000 nm^2.

AI use

AI was not used in any part of this study or manuscript preparation.

Acknowledgments

We thank Karen Oegema (UC San Diego), Jessica Feldman (Stanford University), Jon

Audhya (University of Wisconsin-Madison) and the *Caenohrabditis* Genetics Center for

strains. We thank Kevin O'Connell (NIH, NIDDK), Antonina Roll-Mecak (NIH, NINDS),

Gunar Fabig (University of Dresden) and Alexey Khodjakov (Wadsworth Center, New

York State Department of Health) for helpful discussions. We also thank Kevin

O'Connell for comments on the manuscript. We thank Kedar Narayan for support in

obtaining FIB-SEM data. R.M., M.M.R., A.R., S.D., R.S.M. and O.C-F. were supported

by an NIDDK intramural grant to O.C-F., DK069012-18.

This research was supported by the Intramural Research Program of the National

Institute of Diabetes and Digestive and Kidney Diseases (NIDDK) within the National

Institutes of Health (NIH). The contributions of the NIH authors were made as part of

their official duties as NIH federal employees, are in compliance with agency policy

requirements, and are considered Works of the United States Government. However,

the findings and conclusions presented in this paper are those of the authors and do not

necessarily reflect the views of the NIH or the U.S. Department of Health and Human

Services

Author contribution: The premise of this study was conceptualized by R.M., M.M.R

and O.C-F. Experiments were designed, executed and analyzed by R.M., M.M.R, A.R.,

S.D. R.S.M, and O.C-F. Methodology was developed by R.M., M.M.R, A.R., and S.D.

O.C-F supervised the project and wrote the original draft. All authors reviewed and

commented on the original draft as well as all other manuscript iterations and accepted

the final manuscript.

Disclosure of competing interests

The authors declare that there are no competing interests.

References

Albertson DG (1984) Formation of the first cleavage spindle in nematode embryos. *Dev Biol* 101: 61-72

Aljiboury A, Hehnly H (2023) The centrosome - diverse functions in fertilization and development across species. *J Cell Sci* 136: jcs261387

Araújo M, Tavares A, Vieira DV, Telley IA, Oliveira RA (2023) Endoplasmic reticulum membranes are continuously required to maintain mitotic spindle size and forces. *Life Sci Alliance* 6: 202201540

Arnal I, Heichette C, Diamantopoulos GS, Chrétien D (2004) CLIP-170/Tubulin-Curved Oligomers Coassemble at Microtubule Ends and Promote Rescues. *Curr Biol* 14: 2086-2095

Baumgart J, Kirchner M, Redemann S, Bond A, Woodruff J, Verbavatz J-M, Jülicher F, Müller-Reichert T, Hyman AA, Brugués J (2019) Soluble tubulin is significantly enriched at mitotic centrosomes. *J Cell Biol* 218: 3977-3985

Belmont LD, Hyman AA, Sawin KE, Mitchison TJ (1990) Real-time visualization of cell cycle-dependent changes in microtubule dynamics in cytoplasmic extracts. *Cell* 62: 579-589

Bergman ZJ, Mclaurin JD, Eritano AS, Johnson BM, Sims AQ, Riggs B (2015) Spatial reorganization of the endoplasmic reticulum during mitosis relies on mitotic kinase cyclin A in the early *Drosophila* embryo. *PLoS One* 10: e0117859

Bobinnec Y, Marcaillou C, Morin X, Debec A (2003) Dynamics of the endoplasmic reticulum during early development of Drosophila melanogaster. *Cell Motil Cytoskeleton* 54: 217-225

Bouvrais H, Chesneau L, Le Cunff Y, Fairbrass D, Soler N, Pastezeur S, Pécot T, Kervrann C, Pécréaux J (2021) The coordination of spindle-positioning forces during the asymmetric division of the *Caenorhabditis elegans* zygote. *EMBO Reports* 22: e50770

Brenner S (1974) The genetics of *Caenorhabditis elegans*. *Genetics* 77: 71-94

Brouhard GJ, Stear JH, Noetzel TL, Al-Bassam J, Kinoshita K, Harrison SC, Howard J, Hyman AA (2008) XMAP215 is a processive microtubule polymerase. *Cell* 132: 79-88

Cabral G, Laos T, Dumont J, Dammermann A (2019) Differential requirements for centrioles in mitotic centrosome growth and maintenance. *Dev Cell* 50: 355-366. e356

Chrétien D, Fuller SD, Karsenti E (1995) Structure of growing microtubule ends: two-dimensional sheets close into tubes at variable rates. *J Cell Biol* 129: 1311-1328

903   Conduit PT, Wainman A, Raff JW (2015) Centrosome function and assembly in animal
904   cells. *Nat Rev Mol Cell Biol* 16: 611-624
905

906   Diaz U, Bergman ZJ, Johnson BM, Edington AR, de Cruz MA, Marshall WF, Riggs B
907   (2019) Microtubules are necessary for proper Reticulon localization during mitosis.
908   *PLoS One* 14: e0226327
909

910   Enos SJ, Dressler M, Gomes BF, Hyman AA, Woodruff JB (2018) Phosphatase PP2A
911   and microtubule-mediated pulling forces disassemble centrosomes during mitotic exit.
912   *Biology Open* 7: bio029777
913

914   Farrell K, Jordan M, Miller H, Wilson L (1987) Phase dynamics at microtubule ends: the
915   coexistence of microtubule length changes and treadmilling. *J Cell Biol* 104: 1035-1046
916

917   Fong K-W, Choi Y-K, Rattner JB, Qi RZ (2008) CDK5RAP2 Is a Pericentriolar Protein
918   That Functions in Centrosomal Attachment of the γ-Tubulin Ring Complex. *Mol Biol Cell*
919   19: 115-125
920

921   Fygenson DK, Braun E, Libchaber A (1994) Phase diagram of microtubules. *Physical*
922   *Review E* 50: 1579-1588
923

924   Gao Q, Vermeulen BJ, Würtz M, Shin H, Erdogdu D, Zheng A, Hofer FW, Neuner A,
925   Pfeffer S, Schiebel E (2024) The structure of the γ-TuRC at the microtubule minus end–
926   not just one solution. *Bioessays* 46: 2400117
927

928   Gard DL, Kirschner MW (1987) A microtubule-associated protein from Xenopus eggs
929   that specifically promotes assembly at the plus-end. *J Cell Biol* 105: 2203-2215
930

931   Golden A, Liu J, Cohen-Fix O (2009) Inactivation of the *C. elegans* lipin homolog leads
932   to ER disorganization and to defects in the breakdown and reassembly of the nuclear
933   envelope. *J Cell Sci* 122: 1970-1978
934

935   Gould R, Borisy G (1977) The pericentriolar material in Chinese hamster ovary cells
936   nucleates microtubule formation. *J Cell Biol* 73: 601-615
937

938   Grego S, Cantillana V, Salmon ED (2001) Microtubule Treadmilling in Vitro Investigated
939   by Fluorescence Speckle and Confocal Microscopy. *Biophys J* 81: 66-78
940

941   Gudimchuk NB, McIntosh JR (2021) Regulation of microtubule dynamics, mechanics
942   and function through the growing tip. *Nat Rev Mol Cell Biol* 22: 777-795
943

Hachet V, Busso C, Toya M, Sugimoto A, Askjaer P, Gönczy P (2012) The nucleoporin
Nup205/NPP-3 is lost near centrosomes at mitotic onset and can modulate the timing of
this process in *Caenorhabditis elegans* embryos. *Mol Biol Cell* 23: 3111

Hamill DR, Severson AF, Carter JC, Bowerman B (2002) Centrosome maturation and
mitotic spindle assembly in *C. elegans* require SPD-5, a protein with multiple coiled-coil
domains. *Dev Cell* 3: 673-684

Hannak E, Oegema K, Kirkham M, Gönczy P, Habermann B, Hyman AA (2002) The
kinetically dominant assembly pathway for centrosomal asters in *Caenorhabditis*
elegans is γ-tubulin dependent. *J Cell Biol* 157: 591-602

Harris P (1975) The role of membranes in the organization of the mitotic apparatus. *Exp*
Cell Res 94: 409-425

Hoffmann I (2021) Centrosomes in mitotic spindle assembly and orientation. *Curr Opin*
Struct Biol 66: 193-198

Hu J, Rapoport TA (2016) Fusion of the endoplasmic reticulum by membrane-bound
GTPases. *Semin Cell Dev Biol* 60: 105-111

Karabasheva D, Smyth JT (2019) A novel, dynein-independent mechanism focuses the
endoplasmic reticulum around spindle poles in dividing *Drosophila* spermatocytes. *Sci*
Rep 9: 1-13

Khodjakov A, Rieder CL (1999) The sudden recruitment of γ-tubulin to the centrosome
at the onset of mitosis and its dynamic exchange throughout the cell cycle, do not
require microtubules. *J Cell Biol* 146: 585-596

Kiyomitsu A, Nishimura T, Hwang SJ, Ansai S, Kanemaki MT, Tanaka M, Kiyomitsu T
(2024) Ran-GTP assembles a specialized spindle structure for accurate chromosome
segregation in medaka early embryos. *Nat Commun* 15: 981

Kuriyama R, Borisy GG (1981) Microtubule-nucleating activity of centrosomes in
Chinese hamster ovary cells is independent of the centriole cycle but coupled to the
mitotic cycle. *J Cell Biol* 91: 822-826

Lane HA, Nigg EA (1996) Antibody microinjection reveals an essential role for human
polo-like kinase 1 (Plk1) in the functional maturation of mitotic centrosomes. *J Cell Biol*
135: 1701-1713

Laos T, Cabral G, Dammermann A (2015) Isotropic incorporation of SPD-5 underlies
centrosome assembly in *C. elegans*. *Curr Biol* 25: R648-R649

Lee K, Rhee K (2011) PLK1 phosphorylation of pericentrin initiates centrosome
maturation at the onset of mitosis. *J Cell Biol* 195: 1093-1101

Magescas J, Zonka JC, Feldman JL (2019) A two-step mechanism for the inactivation of
microtubule organizing center function at the centrosome. *Elife* 8: e47867

Maheshwari R, Rahman MM, Drey S, Onyundo M, Fabig G, Martinez MA, Matus DQ,
Müller-Reichert T, Cohen-Fix O (2023) A membrane reticulum, the centriculum, affects
centrosome size and function in *Caenorhabditis elegans*. *Curr Biol* 33: 791-806. e797
995

996 Matthews LR, Carter P, Thierry-Mieg D, Kemphues K (1998) ZYG-9, a *Caenorhabditis*
997 *elegans* protein required for microtubule organization and function, is a component of
998 meiotic and mitotic spindle poles. *J Cell Biol* 141: 1159-1168
999

1000 Megraw TL, Kao L-R, Kaufman TC (2001) Zygotic development without functional
1001 mitotic centrosomes. *Curr Biol* 11: 116-120
1002

1003 Megraw TL, Li K, Kao LR, Kaufman TC (1999) The centrosomin protein is required for
1004 centrosome assembly and function during cleavage in *Drosophila*. *Development* 126:
1005 2829-2839
1006

1007 Mitchison T, Kirschner M (1984) Microtubule assembly nucleated by isolated
1008 centrosomes. *Nature* 312: 232-237
1009

1010 Mittasch M, Tran VM, Rios MU, Fritsch AW, Enos SJ, Ferreira Gomes B, Bond A,
1011 Kreysing M, Woodruff JB (2020) Regulated changes in material properties underlie
1012 centrosome disassembly during mitotic exit. *J Cell Biol* 219: e201912036
1013

1014 Moritz M, Braunfeld MB, Sedat JW, Alberts B, Agard DA (1995) Microtubule nucleation
1015 by γ-tubulin-containing rings in the centrosome. *Nature* 378: 638-640
1016

1017 Nakajo M, Kano H, Tsuyama K, Haruta N, Sugimoto A (2022) Centrosome maturation
1018 requires phosphorylation-mediated sequential domain interactions of SPD-5. *J Cell Sci*
1019 135: jcs259025
1020

1021 O'Toole E, Greenan G, Lange KI, Srayko M, Müller-Reichert T (2012) The role of γ-
1022 tubulin in centrosomal microtubule organization. *PLoS One* 7: e29795
1023

1024 Ohta M, Zhao Z, Wu D, Wang S, Harrison JL, Gómez-Cavazos JS, Desai A, Oegema
1025 KF (2021) Polo-like kinase 1 independently controls microtubule-nucleating capacity
1026 and size of the centrosome. *J Cell Biol* 220: e202009083
1027

1028 Orso G, Pendin D, Liu S, Tosetto J, Moss TJ, Faust JE, Micaroni M, Egorova A,
1029 Martinuzzi A, McNew JA *et al* (2009) Homotypic fusion of ER membranes requires the
1030 dynamin-like GTPase atlastin. *Nature* 460: 978-983
1031

1032 Pelletier L, Ozlu N, Hannak E, Cowan C, Habermann B, Ruer M, Muller-Reichert T,
1033 Hyman AA (2004) The Caenorhabditis elegans centrosomal protein SPD-2 is required
1034 for both pericentriolar material recruitment and centriole duplication. *Curr Biol* 14: 863-
1035 873
1036

Pimenta-Marques A, Bettencourt-Dias M (2020) Pericentriolar material. *Curr Biol* 30:
R687-R689

Poteryaev D, Squirrell JM, Campbell JM, White JG, Spang A (2005) Involvement of the
actin cytoskeleton and homotypic membrane fusion in ER dynamics in Caenorhabditis
elegans. *Mol Biol Cell* 16: 2139-2153

Rahman M, Chang IY, Harned A, Maheshwari R, Amoateng K, Narayan K, Cohen-Fix O
(2020) *C. elegans* pronuclei fuse after fertilization through a novel membrane structure.
J Cell Biol 219: e201909137

Redemann S, Baumgart J, Lindow N, Shelley M, Nazockdast E, Kratz A, Prohaska S,
Brugués J, Fürthauer S, Müller-Reichert T (2017) *C. elegans* chromosomes connect to
centrosomes by anchoring into the spindle network. *Nat Commun* 8: 1-13

Rios MU, Bagnucka MA, Ryder BD, Ferreira Gomes B, Familiari NE, Yaguchi K, Amato
M, Stachera WE, Joachimiak ŁA, Woodruff JB (2024) Multivalent coiled-coil interactions
enable full-scale centrosome assembly and strength. *J Cell Biol* 223: e202306142

Rollins KR, Blankenship JT (2023) Dysregulation of the endoplasmic reticulum blocks
recruitment of centrosome-associated proteins resulting in mitotic failure. *Development*
150: dev201917.

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
Rueden C, Saalfeld S, Schmid B (2012) Fiji: an open-source platform for biological-
image analysis. *Nat Methods* 9: 676-682

Schlaitz AL (2014) Microtubules as key coordinators of nuclear envelope and
endoplasmic reticulum dynamics during mitosis. *Bioessays* 36: 665-671

Srayko M, Kaya A, Stamford J, Hyman AA (2005) Identification and characterization of
factors required for microtubule growth and nucleation in the early *C. elegans* embryo.
Dev Cell 9: 223-236

Strome S, Powers J, Dunn M, Reese K, Malone CJ, White J, Seydoux G, Saxton W
(2001) Spindle dynamics and the role of γ-tubulin in early *Caenorhabditis elegans*
embryos. *Mol Biol Cell* 12: 1751-1764

Terasaki M (2000) Dynamics of the endoplasmic reticulum and Golgi apparatus during
early sea urchin development. *Mol Biol Cell* 11: 897-914

Timothy LM, Li K, Kao L-R, Kaufman TC (1999) The Centrosomin protein is required for
centrosome assembly and function during cleavage in *Drosophila*. *Development* 126:
2829-2839
1081

1082   Vasquez-Limeta A, Loncarek J (2021) Human centrosome organization and function in
1083   interphase and mitosis. *Semin Cell Dev Biol* 117: 30-41

1084
1085   Verde F, Labbé J-C, Dorée M, Karsenti E (1990) Regulation of microtubule dynamics by
1086   cdc2 protein kinase in cell-free extracts of Xenopus eggs. *Nature* 343: 233-238

1087
1088   Waterman-Storer CM, Sanger JW, Sanger JM (1993) Dynamics of organelles in the
1089   mitotic spindles of living cells: membrane and microtubule interactions. *Cell motil*
1090   *cytoskelet* 26: 19-39

1091
1092   Woodruff JB, Gomes BF, Widlund PO, Mahamid J, Honigmann A, Hyman AA (2017) The
1093   centrosome is a selective condensate that nucleates microtubules by concentrating
1094   tubulin. *Cell* 169: 1066-1077. e1010

1095
1096   Woodruff JB, Wueseke O, Hyman AA (2014)
1097   Pericentriolar material structure and dynamics. *Philos Trans R Soc B: Biol Sci* 369:
1098   20130459

1099
1100   Woodruff JB, Wueseke O, Viscardi V, Mahamid J, Ochoa SD, Bunkenborg J, Widlund
1101   PO, Pozniakovsky A, Zanin E, Bahmanyar S (2015) Regulated assembly of a
1102   supramolecular centrosome scaffold in vitro. *Science* 348: 808-812

1103
1104   Wueseke O, Zwicker D, Schwager A, Wong YL, Oegema K, Jülicher F, Hyman AA,
1105   Woodruff JB (2016) Polo-like kinase phosphorylation determines *Caenorhabditis*
1106   *elegans* centrosome size and density by biasing SPD-5 toward an assembly-competent
1107   conformation. *Biology open* 5: 1431-1440

1108
1109   Xie J, Najafi J, Nommick A, Lederer L, Salle J, Dmitrieff S, Lacroix B, Dumont J, Minc N
1110   (2025) Cell shape modulates mitotic spindle positioning forces via intracellular
1111   hydrodynamics. *Curr Biol*: S0960-9822(0924)01622-01621.

1112
1113   Zhang L, Ward JD, Cheng Z, Dernburg AF (2015) The auxin-inducible degradation (AID)
1114   system enables versatile conditional protein depletion in *C. elegans*. *Development* 142:
1115   4374-4384

1116
1117   Zheng Y, Wong ML, Alberts B, Mitchison T (1995) Nucleation of microtubule assembly
1118   by a γ-tubulin-containing ring complex. *Nature* 378: 578-583

1119
1120   Zimmerman S, Chang F (2005) Effects of γ-tubulin complex proteins on microtubule
1121   nucleation and catastrophe in fission yeast. *Mol Biol Cell* 16: 2719-2733

1122
1123

Figure Legends

Figure 1: The spatial relationship between the centrosome, centriculum and peri-centrosomal microtubules.

A) Top panel: a *C. elegans* 1-cell embryo (strain OCF193) at metaphase expressing GFP::SP12 and PH domain::GFP (ER and plasma membrane markers, respectively; left panel and in green in merged panel) and TBA-1::RFP (microtubules; middle panel and in magenta in merged panel). Three types of microtubules are indicated: astral microtubules, which are nucleated at the centrosome and extend toward the cortex, spindle microtubules, which extend toward the chromosomes (not shown, but are located in the gap between the two half spindles), and peri-centrosomal microtubules, which accumulate in the vicinity of the centrosome but do not extend much past it. The two centricula flanking the spindle are indicated by yellow arrows in the left and merged images. Lower panel: *C. elegans* embryo (strain OCF176) at metaphase expressing GFP::SP12 (ER marker; green in merged panel) and GFP::SPD-5 (PCM marker; magenta in merged panel). Scale bar=10 µm.

B) Schematic representation of the centrosome (PCM), with its area (light blue) defined by its outer perimeter (dark blue line); the centriculum, outlined by its inner and outer perimeters (dark orange lines; the area encompassed by the inner perimeter is defined as the centriculum "void", and the area enclosed by the outer perimeter of the centriculum is defined as the centriculum outer perimeter area, or COPA); and peri-centrosomal microtubule (pcMT) area (light green), defined by its outer perimeter (dark green).

C, **D and E)** Quantification of void area (light orange), peri-centrosomal microtubules (green), PCM (blue) and COPA (dark orange) in 1-cell embryos from control RNAi treated worms. The strains and their relevant markers were as follows: Panel C: OCF181 (mCherry::SP12; GFP::TBA-2; n=26); Panel D: OCF176 (mCherry::SP12; GFP::SPD-5; n=22 for void area, n=24 for PCM, and n=23 for COPA; Panel E: OCF247 (GFP::SPD-5; TBA-1:: tagRFP) n=20. Please refer to Fig. S1B for comparison with strain OCF240 (RFP::SPD-5; GFP::TBA-2). Statistical analyses: Panel C: p=0.0001 for void area vs pcMT area; p<0.0001 for void area vs COPA; p=0.0001 for pcMT area vs

COPA. Panel D: p=0.0004 for void area vs PCM area; p<0.0001 for void area vs COPA;

p=0.0002 for PCM area vs COPA. These analyses were done using Anova Kruskal-

Wallis test. Panel E: p=0.0043 using Anova Mann-Whitney test. Error bars indicate

mean and standard deviation.

F) Schematic representation of the spatial relationship between the perimeters of the

centriculum outer and inner perimeter (orange), peri-centrosomal microtubules (green)

and the PCM (blue).

1162

Figure 2: Decrease in microtubule abundance or stability leads to a smaller

centriculum and compression of the PCM.

A) Representative images of centricula from 1-cell embryos (OCF181) at metaphase

expressing the SP12 ER marker fused to mCherry (mCherry::SP12; green in merged

images) and α-tubulin fused to GFP (GFP::TBA-2; magenta in merged images), from

worms treated with RNAi against a control (top row), *zyg-9* (middle row) or *tbg-1* RNAi

(bottom row). Scale bar=5 μm.

B and C) Quantification of GFP::TBA-2 peri-centrosomal microtubule area, void area

and COPA, from embryos treated as shown in panel A. Panel B: n=26, 15, and 12 for

control, *zyg-9* and *tbg-1* RNAi treatments, respectively. p< 0.0001 for control vs *zyg-9*

and control vs *tbg-1* RNAi treatments using ordinary one-way ANOVA. Panel C: n=26,

16 and 12 for control, *zyg-9* and *tbg-1* RNAi treatments, respectively. For void area: p<

0.0001 for control vs *zyg-9* and control vs *tbg-1* RNAi treatments using Kruskal-Wallis

test. For COPA: p< 0.0001 for control vs *zyg-9* RNAi and p=0.0072 for control vs *tbg-1*

RNAi using ordinary one-way ANOVA. Error bars represent mean and standard

deviation. Note that the values for control RNAi are the same as shown in Fig. 1C.

D) Representative images of centricula and PCM from 1-cell embryos (OCF176) at

metaphase expressing mCherry::SP12 (green in merged images) and the SPD-5 PCM

protein fused to GFP (GFP::SPD-5; magenta in merged images), from worms treated

with RNAi against a control (top row), *zyg-9* (middle row) or *tbg-1* (bottom row). Scale

bar=5 μm.

E, F and G) Quantification of GFP::SPD-5 area, amount and density (determined by

SPD-5 total intensity divided (panel F) by SPD-5 area (panel E)) from embryos treated

as shown in panel D. n=24, 12, and 16 for control, *zyg-9* and *tbg-1* RNAi treatments,

respectively. For panel E: p<0.0001 for control vs *zyg-9* and for control vs *tbg-1* RNAi

treatments using ordinary one-way ANOVA. Note that the control RNAi values for SPD-5

area are the same as in Fig. 1D. Panel F: p=<0.0001 for control vs *zyg-9* and p=0.3290

for control vs *tbg-1* RNAi treatments using ordinary one-way ANOVA. Panel G:

p=<0.0001 for control vs *zyg-9* and for control vs *tbg-1* RNAi treatments using ordinary

one-way ANOVA. Error bars indicate mean and standard deviation.

H) GFP::SPD-5 density (as in panel G) as a function of void area (Fig. S1D) following

control (magenta), *zyg-9* (purple)*, or *tbg-1* (pink) RNAi treatments. Linear regression

was performed through all the datapoints. R^2=0.2624; slope was significantly different

from 0 (p=0.0002).

1197

Figure 3: *spd-5* expansion mutant inhibits PCM expansion in the presence of wild

type *spd-5*.

A) Diagram depicting possible relationships between microtubules, the centriculum, and

the PCM. See text for more detail.

B, C and D) Representative images of centricula and PCM from 1-cell embryos at

metaphase expressing mCherry::SP12 (green in merged images) and either (B) a

GFP::spd-5 transgene in the absence of endogenous protein (OCF187), (C) *GFP::spd-*

*5^exp^* in the absence of endogenous protein (OCF189), or (D) *GFP::spd-5^exp^* in the

absence of presence of endogenous protein (OCF189). The transgenic SPD-5 is shown

in magenta in the merged images. In panels B and C, the endogenous gene was down-

regulated using RNAi against *spd-5*; transgenes are codon modified such that they are

insensitive to the RNAi treatment (Ohta *et al.*, 2021). Additional examples of images of

embryos such as in panels C and D are shown in Fig S2A and B) Scale bar=5 μm.

E) Representative images of PCM from 1-cell embryos at metaphase expressing

endogenous RFP::SPD-5, and either transgenic GFP::SPD-5 wildtype (OCF201) or

transgenic GFP::SPD-5^exp^ (OCF200). Scale bar= 5 μm.

F and G) Quantification of areas occupied by endogenous RFP::SPD-5 and either

transgenic GFP::SPD-5 (panel F) or GFP::SPD-5^exp^ (panel G) from images such as

shown in panel E. n=11 and 13 for transgenic wild type and expansion mutant,

respectively. p<0.0001 using unpaired t test. Error bars indicate mean and standard

deviation.

H and I) Quantification of protein amount (as reflected by total intensity) occupied by

endogenous RFP::SPD-5 and either transgenic GFP::SPD-5 (panel H) or GFP::SPD-

5^{exp} (panel I) in the areas quantified in panels F and G. p=0.0004 (panel H) and 0.0034

(panel I), using Mann Whitney test.

1223

Figure 4: Microtubule nucleation remains unaffected in the presence of SPD-5

expansion mutant

A) Representative images of peri-centrosomal microtubules and PCM from 1-cell

embryos at metaphase expressing endogenous *tba-1::RFP* and either transgenic

GFP::spd-5 (OCF212) or transgenic *GFP::spd-5^{exp}* (OCF213), after control RNAi

treatments. Scale bar=5 μm.

B, C and D) Quantification of PCM area (panel B), amount of peri-centrosomal tubulin

(panel C) and the area occupied by peri-centrosomal microtubules (pcMT; panel D) from

control RNAi treated embryos such as shown in panel A. n=21 and n=29 for cells

expressing transgenic wild type (WT) or expansion mutant (Exp) *spd-5*, respectively.

p<0.0001 (panel B) and p=0.3192 (panel D) using unpaired t test, and p=0.6400

(panel C) using Mann Whitney test. Error bars indicate mean and standard deviation.

E) Quantification of area differential (pcMT area minus PCM area) derived from panels

B and D. p=0.0033 using Mann Whitney test.

F) Quantification of PCM area and pcMT area from the *klp-7* RNAi treated embryos

using the same strains as in panels A-E (OCF212 and OCF213). n=20 and n=19 for

strains expressing wild type or *spd-5^{exp}* transgenes, respectively. p<0.0001 for PCM

areas and p= 0.4056 for pcMT areas, using ordinary one-way ANOVA. Error bars

indicate mean and standard deviation.

G) Quantification of area differential (pcMT- PCM, as in panel E) derived from pcMT and

PCM areas (panel F) in *klp-7* RNAi treated embryos. p=0.0014 using the Mann Whitney

test.

1246

Figure 5: Centriculum size correlates with the position of peri-centrosomal microtubules, not the PCM.

A) Schematic representation of the spatial relationship between the perimeters of the centriculum (orange), PCM (blue) and the peri-centrosomal microtubules (green). (i) In the presence of $spd\text{-}5^{exp}$ mutant, the area of the PCM, but not the peri-centrosomal microtubules, decreases, and thus the distance between the edges of the two structures should increase. (ii) When considering how the centriculum size, and specifically the area bound but the centriculum's inner perimeter (the void area), is affected by the presence of $spd\text{-}5^{exp}$ mutant, two possibilities emerge: the edge of the void area follows the PCM (left) or the peri-centrosomal microtubules (right). See text for more detail.

B) Example of centricula and peri-centrosomal microtubules from 1-cell embryos expressing GFP::SP12 (green in merged image) and TBA-1::RFP (magenta in merged image) and either wild type (OCF214) or $spd\text{-}5^{exp}$ (OCF215) transgenes, following control or *klp-7* RNAi treatments. Scale bar=5 µm.

C and D) Quantification of pcMT areas (panel C) and void areas (panel D) from control RNAi treated embryos as in panel B. n=19 and n=15 for transgenic wild type and spd-5^{exp} expressing strain, respectively. $p<0.0001$ for pcMT area comparison using Mann Whitney test, and $p<0.0001$ for void area comparison using unpaired t test. Error bars indicate mean and standard deviation.

E) Quantification of area differential (pcMT area minus void area) derived from area values shown in panels C and D and Fig S3C. n=19, 15, 20, 23 for control RNAi, wild type and $spd\text{-}5^{exp}$ transgenes and *klp-7* RNAi, wild type and $spd\text{-}5^{exp}$ transgenes. p-values, from left to right, were 0.5653, 0.013, 0.2353 and 0.9945, using ordinary one-way ANOVA. Error bars indicate mean and standard deviation.

F) Representative images of centricula and PCM from 1-cell embryos at metaphase expressing *mCherry::SP12* (green in merged images) and either *GFP::spd-5* (OCF187) or *GFP::spd-5^{exp}* (OCF189) transgenes (magenta in merged images), following control or *klp-7* RNAi treatment. White arrow in enlarged image points to a gap between the centriculum and SPD-5. Scale bar=5 µm.

G and H) Quantification of PCM area (panel G) and void area (panel H) from 1-cell metaphase embryos as shown in panel F, following control or *klp-7* RNAi treatment.

n=12 all conditions except n=16 for *spd-5^exp* transgene following control RNAi. For PCM

area, $p<0.0001$ for all pairwise comparisons as determined by ordinary one-way

ANOVA. For void area, p-values, from left to right, are 0.6133, <0.0001, <0.0001 and

0.9999, as determined by ordinary one-way ANOVA. Error bars indicate mean and

standard deviation.

I) Quantification of the area differential (PCM area minus void area) derived from area

values shown in panels G and H. p-values, as determined by ordinary one-way

ANOVA, were <0.0001, except for WT vs Exp, control RNAi, where the p-value was

p=0.0003. Error bars indicate mean and standard deviation.

1287

Figure 6: Spatial relationship between the centriculum and the PCM using an

endogenously tagged *spd-5* in the presence of *spd-5^exp* mutant

A) Representative images of centricula and PCM from 1-cell embryos at metaphase

expressing RFP-tagged endogenous *spd-5* (RFP::SPD-5, in magenta) and GFP::SP12

(in green) in strains expressing either wild type (OCF234) or *spd-5^exp* (OCF233)

transgenes, following control or *klp-7* RNAi treatment. Scale bar=5 µm.

B, C and D) Quantification of SPD-5 amount, PCM areas and void areas from 1-cell

metaphase embryos treated as shown in panel A. n=16 and 11 for wild type and *spd-*

5^exp transgenes treated with control RNAi, and n=20 and n=16 for wild type and *spd-5^exp*

transgenes treated with *klp-7* RNAi. p-values, from left to right, were as follows: panel B:

<0.0001, 0.1795, 0.3578 and <0.0001, as determined by ordinary one-way ANOVA;

panel C: 0.0018, 0.0023, 0.0286 and <0.0001, as determined by Kruskal-Wallis test;

panel D: 0.0547, and the rest were <0.0001, as determined by ordinary one way

ANOVA. Error bars indicate mean and standard deviation.

E) Quantification of the area differential (PCM area minus void are) derived from data

shown in panels C and D. p-values, as determined by ordinary one-way ANOVA, were

<0.0001 except for p=0.0001 for wild type, control vs *klp-7* RNAi. Error bars indicate

mean and standard deviation.

F) Diagram illustrating the effect of the spd-5^exp transgene on the centriculum (orange),

peri-centrosomal microtubules (green) and PCM (blue). See text for more detail.

1308

Figure 7: Porousity of the centriculum facing the cortex.

A) Representative images of 1-cell *C. elegans* embryos expressing mCherry::SP12 to visualize the centriculum and GFP::TBA-2 (α-tubulin) to visualize microtubules (strain OCF181) at prophase (left side) and metaphase (right side). Note that at prophase, microtubules are excluded from the nucleoplasm, and that in metaphase, the density of spindle microtubules is greater than astral microtubules. Scale bar=10 µm.

B) Diagram depicting the centriculum (in orange) fused to the two pronuclei (brown lines) at a metaphase 1-cell embryo. Chromosomes (in red) and the cortex (black line) are also indicated. The areas analyzed for pore size are shown as blue (chromosome facing) and green (cortex facing) trapezoids. A central fiducial (in pink) that is equidistant (pink arrows) to the areas analyzed is also shown.

C) Cortical and pronuclear centriculum segments based on FIB-SEM data (Maheshwari *et al.*, 2023) from two centricula, shown side by side. The images were taken from the central fiducial. Scale bar=200 nm.

D) Binned frequency distribution of holes present on the cortical side (in green) and chromosome side (in blue) of centricula in 1-cell metaphase embryos. Bin size range, as hole area, is shown on the x-axis. n=308 and 507 for chromosomal and cortex side holes, respectively, each containing data pooled from 6 images taken from 3 centricula (2 images per centriculum).

E) Binned frequency distribution of the percentage of total open area per hole size range, for holes on the cortical side (green) and the pronuclear side (blue) using the same data as in panel D.

Figure 8:

A) Diagram depicting the angle of a microtubule (green line) passing through the centriculum (in orange) relative to the radial vector (dashed purple line). The actual measurement was done between the microtubule and the tangent line (dashed black line) and the value obtained was subtracted from 90 to obtain the angle to the radial vector, which is perpendicular to the tangent line.

B) Quantification of angles between of stopped (in orange) and continuing (in cyan) microtubule and radial vector for astral (cortex-facing) and spindle (chromosome-facing)

microtubules. Based on EM tomography data (Redemann *et al.*, 2017). p=0.0029 (for

cortex-facing) and 0.0018 (chromosome-facing) using one-way ANOVA with multiple

comparisons. Error bars indicate mean and standard deviation.

C) Left panel: Representative images of a segmented area based on FIB-SEM data

(Maheshwari *et al.*, 2023) from the pronuclear side of two prometaphase centricula (in

yellow), visualized from the center of the centrosome. Right panel: the same images

with segmentation of nuclear membrane remnants (in green) that are between the

centriculum and the prometaphase chromosomes (not shown). Scale bar=200 nm.

D) Areas of possible contact (indicated by red arrows) between a prometaphase

centriculum (in yellow) and remnants of the nuclear membrane (in green) based on FIB-

SEM data (Maheshwari *et al.*, 2023). In this image, the centrosome is to the upper right

side while the chromosomes are to the lower left side.

E) Two 1-cell embryos expressing mCherry::SP12 and GFP::TBA-2 from worms in

which atlastin was downregulated using an auxin-inducible degron (AID) system

(OCF183, see (Maheshwari *et al.*, 2023)). Red arrows point to sites where a centriculum

has fused to a single pronucleus. Green arrows point to sites where a centriculum has

fused to both pronuclei. The same arrows are shown in the bottom panels, where they

are adjacent to sites where spindle microtubules extend into the pronuclei. Blue

asterisks indicate pronuclei that fuse to only one centriculum. The enlarged areas show

sites of centriculum-pronuclear fusion. Note that unlike the wild type situation (Fig 7A),

under these conditions the two pronuclei fail to align and do not form extensive

pronuclear interfacial contacts. Scale bar=10 μm.

1362
1363
1364
1365

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A

RNAi condition

control   *klp-7*

SP12 SP-5

Wild type

Exp mut

Transgenic *spd-5* allele:

B

SPD-5 amount

Total intensity x 10^5, A.U.

ns

ns

**** ****

Transgene:	WT	Exp	WT	Exp
RNAi:	control		*klp-7*	

C

**PCM area
(endogenous SPD-5)**

Area, μm^2

*

**

** ****

Transgene:	WT	Exp	WT	Exp
RNAi:	control		*klp-7*	

D

Void area

Area, μm^2

****

****

ns ****

Transgene:	WT	Exp	WT	Exp
RNAi:	control		*klp-7*	

E

**Area differential
(PCM – void)**

Area, μm^2

****

***

**** ****

Transgene:	WT	Exp	WT	Exp
RNAi:	control		*klp-7*	

F

Centriculum
PCM
Peri-centromeric
microtubules

Wild type

spd-5^{exp} mutant

Figure 7

A

Prophase   Metaphase

ER

Microtubules

Spindle microtubules

Astral microtubules

B

centriculum   Pronuclear membranes

cortex

1
2
1
2

chromosomes

C

Cortex facing 1   Chromosome facing 1   Chromosome facing 2   Cortex facing 2

Centriculum 1

Centriculum 2

D

Distribution of hole number (%)

Cortex facing
Chromosome facing

Hole size (nm² X 1000)

E

% of area per hole size

Cortex facing
Chromosome facing

Hole size (nm² X 1000)

Figure 8

A Centriculum, Tangent line, Microtubule, Radial vector (perpendicular to tangent line), Measured angle

B Angles between microtubules and the radial vector

C

D

E *atln-1::*AID (+IAA)

SP12 i, iii, a, b

TBA-2 ii, iv, a, b