Hugin-AstA circuitry is a novel central energy sensor that directly regulates sweet sensation in *Drosophila* and mouse

Wusa Qin[1,6], Tingting Song[2,3,6], Zeliang Lai[4], Daihan Li[3], Liming Wang[1,2,3,*], and Rui Huang[3,5,*]

[1]Institute of Molecular Physiology, Shenzhen Bay Laboratory, Shenzhen, 518055, China.

[2]School of Public Health, Capital Medical University, Beijing, 100069, China.

[3]Chinese Institutes for Medical Research, Beijing, 100069, China.

[4] Center for Neurointelligence, School of Medicine, Chongqing University, Chongqing, 400030, China.

[5]Guangyang Bay Laboratory, Chongqing Institute for Brain and Intelligence, Chongqing, 400060, China.

[6]These authors contribute equally to this work.

[*]For correspondence: lmwang83@ccmu.edu.cn; huangrui0716@sina.com

Abstract

Taste sensation plays a crucial role in shaping feeding behavior and is intricately influenced by internal states like hunger or satiety. Despite the identification of numerous neural substrates regulating feeding behavior, the central neural substrate that linked energy-sensing and taste sensation remained elusive. Here, we identified a novel neural circuitry that could directly sense internal energy state and modulate sweet sensation in the Drosophila brain. Specifically, a subset of neuropeptidergic neurons expressing hugin directly detected elevated levels of circulating glucose via glucose transporter Glut1 and ATP-sensitive potassium channel. Upon activation, these neurons released hugin peptide and activated downstream Allatostatin A (AstA)$^+$ neurons via its cognate receptor PK2-R1. Subsequently, the activation of AstA$^+$ neurons then directly inhibited sweet sensation via AstA peptide and its cognate receptor AstA-R1 expressed in sweet-sensing Gr5a$^+$ neurons. We also showed that neuromedin U (NMU), the mammalian homolog of fly hugin, served as an energy sensor to suppress sweet sensation. Therefore, these data identify hugin$^+$ neuron as a central energy sensor responsible for regulating sweet sensation across species.

Keywords: satiety, sweet sensation, feeding, hugin, AstA

Introduction

Animal behaviors are tightly associated with the internal metabolic states (Munch et al., 2020).

When experiencing hunger, animals exhibit behaviors aimed at obtaining food, including

elevated sensitivity to appetitive cues and enhanced foraging and feeding behaviors

(Augustine et al., 2020; Marella et al., 2012; Yu et al., 2016). In addition, hunger tends to

increase anxiety levels and aggressive behaviors (de Rivaz et al., 2022), while inhibiting pain,

sleep, and mating behaviors (Alhadeff et al., 2018; Burnett et al., 2016; Sun et al., 2023).

These behavioral alterations upon hunger serve to sharpen the focus of starved animals on the

tasks related to food acquisition (Deem et al., 2022). Conversely, when animals are sated,

food-related behaviors diminish, redirecting their attention to other important behaviors such

as reproduction (Burnett et al., 2016; Karigo and Deutsch, 2022). Such adjustments in

behaviors help animals to fulfill various internal needs in a fast-changing environment, and

also suggests the presence of neural circuitry in the brain that detects and encodes hunger and

satiety states to guide individual behavioral choices and maintain metabolic homeostasis.

In mammals, agouti-related peptide (AgRP)- and pro-opiomelanocortin (POMC)-expressing

neurons in the hypothalamic arcuate nucleus (ARC) are key sensors of hunger and satiety (Fu

et al., 2019). When calories are scarce, AgRP neurons become active, promoting foraging and

feeding behaviors while suppressing competing behaviors such as thirst, innate fear, anxiety-

related behavior, and social interactions (Burnett et al., 2016; Dietrich et al., 2015).

Conversely, satiety signals gradually inhibit the activity of AgRP neurons in response to food

or food cues, resulting in reduced appetite (Bruning and Fenselau, 2023; Chen et al., 2015).

Antagonistic to AgRP neurons, POMC neurons play a critical role as satiety sensors,

suppressing feeding and increasing energy expenditure by releasing a α-melanocyte-

stimulating hormone (α-MSH) (Bruning and Fenselau, 2023; Koch et al., 2015).

AgRP/POMC neuron activity is regulated by various upstream signals, including hormones

like leptin, insulin, and ghrelin, as well as nutrients such as glucose, amino acids, and fatty

acids. Leptin, GLP-1, and insulin activate POMC neurons while inhibiting AgRP neurons,

reducing food intake (Belgardt et al., 2009; De Solis et al., 2024), whereas ghrelin, secreted

during fasting, activates AgRP neurons to promote feeding (Vivot et al., 2023). Furthermore,

AgRP and POMC neurons project to various brain regions inlcuding paraventricular

hypothalamus (PVH), dorsomedial hypothalamus (DMH), ventromedial hypothalamus

(VMH), and brainstem, exerting complex regulations on behavior and metabolism (Belgardt

et al., 2009; De Solis et al., 2024; Wei et al., 2018). These intertwined circuitry helps to

coordinate behaviors and physiological responses in a dynamic environment. Despite the

significant progress in the past decades, the direct link of these upstream internal energy

sensors, such as AgRP and POMC neurons, to various behavioral outputs, remains

incompletely understood. Current research highlights their central role in energy homeostasis

but also suggests a complex, multi-level circuitry that modulates these behaviors. Fully

elucidating the relative contributions of these neurons and circuits to behavioral and

metabolic regulations is an ongoing challenge in the field.

Fruit flies offer an excellent system for studying satiety/hunger sensing and mapping out the

complete circuitry towards behavioral consequences, owing to their simple nervous system

and comprehensive genetic and neurobiological tools (Lin et al., 2019). Hunger and satiety

states modulate a range of behaviors in *Drosophila*, including foraging, feeding, food

preference, sleep, and even memory formation (Chouhan et al., 2021; Lin et al., 2019).

Hungry flies tend to increase foraging, favor high-calorie food, and extend feeding durations,

while satiated flies reduce food intake and shift their food preferences (Lin et al., 2019).

Hunger also disrupts sleep, with starved flies becoming more alert to prioritize food-seeking

(Willyard, 2008). A constellation of neuromodulators has been shown to steer these

behavioral transitions. Dopamine navigates the feeding process by engaging reward pathways

upon food detection (Marella et al., 2012), whereas serotonin (5-HT) functions to quell

feeding behavior during satiety (Yao and Scott, 2022), simultaneously shaping food

preferences and foraging endeavors. Neuropeptide F (NPF), akin to mammalian neuropeptide

Y (NPY), is instrumental in instigating feeding bouts during periods of hunger (Malita et al.,

2022). Hugin, bearing a resemblance to mammalian neuropeptide U, influences satiety levels

and modulates food intake, regulated by gustatory neurons (Melcher and Pankratz, 2005;

Schoofs et al., 2014). Short neuropeptide F (sNPF) contributes to hunger-induced behaviors

by calibrating sensory reactions to food stimuli (Root et al., 2011). Allatostatin A (AstA)

emerges as a satiety signal that, likely through interactions with sweet-sensing neurons, and

suppresses feeding activities. (Chen et al., 2016; Hentze et al., 2015; Hergarden et al., 2012).

CCHamide, which is associated with growth modulation, potentially acts as a hunger signal,

whereas leucokinin adjusts meal size and manages the physiological aftermath of feeding (Lin

et al., 2019). Corazonin is perceived to be intertwined with the modulation of stress and

metabolic responses pertinent to hunger and satiety (Kubrak et al., 2016). SIFamide is

recognized for its role in translating hunger signals into both appetitive and consummatory

feeding actions (Martelli et al., 2017). The intricate interplay among these modulators

constitutes a sophisticated regulatory network, the intricacies of which are yet to be delineated

comprehensively, underscoring an exciting frontier for forthcoming research aimed at

unraveling the neural circuitry that underpins energy homeostasis.

Sugar serves as an important source of satiety signals. The body's precise responses to dietary

sugar involve both peripheral organs and the brain, which facilitates the regulation of various

behaviors, including sweet perception and food consumption (Oh et al., 2021; Qi et al., 2021;

Ugrankar et al., 2018; Zhao et al., 2022). When exposed to a high-sugar diet, the midgut

reacts by secreting Hedgehog (Hh) to suppress sweet sensation (Zhao et al., 2022).

Additionally, hugin$^+$ neurons in the ventral nerve cord (VNC) become sugar-sensitive in fed

states, leading to the suppression of food consumption via neuropeptide diuretic hormone 44

(DH44)$^+$ neurons(Oh et al., 2021). In the brain, insulin-producing cells (IPCs) and tackykinin

(TK)$^+$ neurons are also activated in the presence of sugar, contributing to appetite suppression

(Hergarden et al., 2012; Lin et al., 2019; Qi et al., 2021). In contrast to satiety sensing,

certain neurons, such as dopaminergic neurons, act as hunger sensors in fruit fly brains, being

activated during hunger states and involved in regulation of sleep, sugar memory expression

and sweet taste sensation (Marella et al., 2012; Senapati et al., 2019; Yurgel et al., 2019).

However, it remains unclear how these neurons interact to form a regulatory network on

energy homeostasis, especially how these neurons link the hunger/satiety state to direct

behavioral outputs.

We aimed to dissect out a complete neural pathway that directly sensed internal energy state

and modualted food-related behavioral output in the fly brain. More specifically, we focused

on the circuitry mediating the effect of satiety to inhibit sweet taste sensation. Among several

candidate neuropeptidergic neuron groups, we pinpointed hugin$^+$ neurons as a direct satiety

sensor in the fly brain. These hugin$^+$ neurons were activated by elevated levels of circulating

glucose after feeding, which in turn activated downstream AstA$^+$ neurons via the interaction

of hugin peptide and its cognate receptor PK2-R1. Subsequently, AstA$^+$ neurons exerted

direct inhibitory effect on sweet-sensing Gr5a$^+$ neurons via AstA peptide and its receptor

AstA-R1. As a result, sated flies exhibited a reduction in sweet sensitivity and hence food

consumption. In summary, we unveiled a novel satiety sensing mechanism in *Drosophila* that

detected circulating glucose and regulated sweet perception and feeding. We also showed that

NMU, the mammalian homolog of hugin, was also an energy sensor in mouse models

involved in regulation of sweet perception, highlighting the conserved regulatory mechanism

of central energy sensing and sweet sensation across species.

Results

hugin$^+$ neurons were a novel glucose sensor in the fly brain

Flies extend their proboscis in response to appetitive cues, a behavioral element known as the proboscis extension reflex (PER) (Marella et al., 2012; Song et al., 2023). As previously observed, hunger significantly enhances this reflex(Marella et al., 2012). In our study, we first replicated this phenomenon: In the state of hunger, flies exhibited elevated PER towards various concentrations of sugar compared to those under fed conditions (*Figure 1A-B*). These results confirmed the notion that sweet sensation was influenced by the internal energy state, elevated by starvation and suppressed by satiety (Inagaki et al., 2014; Marella et al., 2012). Sweet sensation in flies is mediated by sweet-sensing Gr5a$^+$ neurons located on the proboscis (Fujii et al., 2015). Consistent with the behavioral effect, we also confirmed that Gr5a$^+$ neurons exhibited elevated calcium responses to sugar under starvation conditions (*Figure supplementary 1A*).

Dopaminergic neurons in the subesophageal zone (SEZ) region of the fly brain acted as a hunger sensor, promoting sweet sensation via a specific dopamine receptor DopEcR expressed in Gr5a$^+$ neurons (Inagaki et al., 2014; Marella et al., 2012). However, although the gene knockout of tyrosine hydroxylase (TH) (Budnik and White, 1987), a key enzyme for dopamine biosynthesis, significantly suppressed sweet sensation in starved fruit flies (*Figure supplementary 1B*), these flies still showed a further reduction in sugar-induced PER under sated conditions (*Figure supplementary 1C-D*). These results suggest the existence of dopamine-independent mechanisms for sensing satiety and hunger in fruit flies that exerts

modulatory effect on sweet sensation.

In the terminating phase of feeding behavior, satiety signals play a crucial role to cease food

consumption and to ensure an appropriate amount of food ingestion. A number of

neuropeptidergic neurons were identified as putative satiety signals (Lin et al., 2019).

Notably, some of these neurons, including those secreting neuropeptides LK, TK, hugin,

AstA, CCHa1, and CCHa2, are also distributed in the SEZ region (Deng et al., 2019). It has

been reported that the sweet-sensing gustatory neurons primarily receive regulatory inputs

from neurons in the SEZ region (Marella et al., 2012; Song et al., 2023). Therefore, we

hypothesized that these neuropeptidergic neurons might function as satiety signals, imposing

inhibitory effect on sweet sensation after adequate food ingestion.

To test this hypothesis, we selectively expressed the temperature-sensitive ion channel TrpA1

in these neurons and activated them at 30 °C, triggering the release of corresponding

neuropeptides. Indeed, activation of LK-, TK-, hugin-, AstA-, CCHa1-, and CCHa2-

expressing neurons robustly suppressed sweet taste sensitivity in both fed and starved flies

(*Figure 1C–D*; *also see Figure supplementary 2*). Moreover, LK-, TK-, and hugin-

expressing neurons exerted particularly strong inhibition under satiated conditions (*Figure*

1C–D; also see Figure supplementary 2). These results reveal a multilayered, state-

dependent neuropeptidergic network that finely tunes gustatory processing and feeding

behavior. In contrast, activating dopaminergic neurons enhanced sweet perception in sated

flies as expected (*Figure 1C*). In starved flies, activating dopaminergic neurons elicited no

change in PER likely due to a ceiling effect (***Figure 1D***).

In mammals and flies, circulating sugar levels are directly linked to satiety/hunger state (Lin

et al., 2019; Qi et al., 2021; Yoon and Diano, 2021). We then asked whether changes in

circulating sugar levels exerted an effect on sweet sensation. Activating IPCs in the fly brain

released *Drosophila* Insulin-like Peptides (DILPs), the fly analog of mammalian insulin

(Brogiolo et al., 2001), into the hemolymph and reduced circulating sugar levels (***Figure***

supplementary 3A). As a result, such manipulation enhanced sweet sensitivity of fed flies

(***Figure supplementary 3B-C***). Therefore, it was plausible that changes in circulating sugar

might be the driver of changes in sweet sensation.

Indeed, allowing starved flies to re-feed on normal fly diet or sucrose alone for 15 minutes

both resulted in a significant decrease in sweet sensation (***Figure 1E***). To validate this effect

across sweet modalities, we assessed responses to trehalose and fructose after sucrose

refeeding and observed comparable reductions in sweet sensation (***Figure supplementary 4***).

Conversely, when starved flies were supplied with fly diet deprived of sucrose (but with

normal levels of polysaccharides, proteins, and lipids), there was no immediate change in

sweet sensation after re-feeding (***Figure 1E***). These results demonstrate that sugar intake

inhibits sweet sensation, probably via increasing circulating sugar levels.

Collectively, we thus hypothesized that the above neuropeptidergic neurons might directly

sense an increase of circulating sugar levels and suppress sweet sensation in response. If so,

these neurons should be activated by glucose, the main species of circulating sugars

associated with feeding behavior in flies (Ugrankar et al., 2018), at the concentrations

resembling sated state (>50 mM in fed flies (***Figure supplementary 5***)). We thus conducted

calcium imaging experiments in fly brains in an *ex vivo* preparation (Qi et al., 2021). Among

all the six groups of neuropeptidergic neurons we examined in the SEZ, only hugin$^+$ neurons

and AstA$^+$ neurons could be activated by 50 mM glucose (***Figure 1F-I, also see Figure***

supplementary 6). Remarkably, each of these populations subdivided into anatomically

distinct SEZ subclusters (***Figure supplementary 7A; Figure supplementary 9A***), yet glucose-

evoked Ca^{2+} responses occurred exclusively within a single subcluster (***Figure 1H,I; Figure***

supplementary 7B; Figure supplementary 9B). This selective activation highlighted

profound functional heterogeneity in nutrient sensing among these neuron groups. When

direct synaptic transmission was blocked with TTX, hugin$^+$ neurons, but not AstA$^+$ neurons,

still exhibited calcium responses towards 50 mM of glucose (***Figure 1G, green***), suggesting

that hugin$^+$ neurons are a direct glucose sensor. Furthermore, the level of circulating glucose

dropped to ~ 20 mM in starved flies (***Figure supplementary 5***). We found that 20 mM of

glucose did not activate hugin$^+$ and AstA$^+$neurons (***Figure 1G and I, orange***), suggesting that

these neurons can only be activated under sated conditions.

We also assessed the activity of hugin$^+$ neurons *in vivo* using calcium-modulated

photoactivatable ratiometric integrator (CaMPARI), a method employing a photoconvertible

fluorescent protein that allows the imaging of integrated calcium activity (Fosque et al.,

2015), under both sated and starved conditions. In the CaMPARI assay, when neurons are

activated, leading to an increase in free calcium ions, exposure to 405-nm light induces a

permanent green-to-red photoconversion (Fosque et al., 2015). We found an increased ratio of

red/green fluorescence in hugin[+] neurons in the sated state compared to that upon starvation

(*Figure 1J*). These findings strongly imply that hugin[+] neurons function as a direct satiety

sensor, sensing the levels of circulating glucose and modulating sweet sensation in response.

Glucose activated hugin[+] neurons via an ATP-sensitive potassium channel

We then investigated the cellular mechanism through which glucose could directly activate

hugin[+] neurons. In both mammals and flies, extracellular glucose is usually transported into

the cytosol to activate glucose-sensitive neurons (Dus et al., 2015; Oh et al., 2019; Parton et

al., 2007). When fly brains were pre-treated with phlorizin, a blocker of glucose transport

(Dus et al., 2015; Oh et al., 2019), hugin[+] neurons showed diminished calcium responses to

glucose, suggesting activation of hugin[+] neurons by glucose requires intracellular glucose

(*Figure 2A*).

The transport of glucose mediated by glucose transporter 1 (Glut1) can activate certain

neurons in the fruit fly brain (Dus et al., 2015; Oh et al., 2019). By specifically knocking

down Glut1 expression in hugin[+] neurons, we observed a significant reduction in calcium

responses to glucose (*Figure 2B*). Consistently, these flies showed a slight yet significant

increase in their PER responses to glucose (*Figure 2C*). Again, these results further confirm

that glucose needs to be transported into hugin[+] neurons to modulate neuronal activity and

behavioral output.

In pancreatic β-cells and certain fly neurons (Oh et al., 2019; Parton et al., 2007; Wang et al.,

2022), intracellular glucose modulates cell excitability primarily through its effect on

intracellular energy metabolism and ATP-sensitive potassium channels (K_{ATP}). We pre-treated

fly brains with a K_{ATP} inhibitor (glibenclamide) and observed a significant calcium response

in hugin[+] neurons (***Figure 2D***). Hexokinase, a key enzyme in glucose metabolism that

generates ATP, also plays a crucial role in this process. When hexokinase activity was

restricted by alloxan, glucose-induced calcium response in hugin[+] neurons was significantly

suppressed (***Figure 2E***). Similarly, when hexokinase was specifically knocked down in

hugin[+] neurons, a comparable reduction in glucose-induced calcium responses was observed

(***Figure 2F***). Pyruvate, the end product from glycolysis, can lead to ATP generation via the

tricarboxylic acid cycle. We also found that pyruvate stimulation led to a significant calcium

response in hugin[+] neurons (***Figure 2G***). As a specificity control, nonmetabolizable L-glucose

failed to activate hugin[+] neurons (***Figure 2H***).

Collectively, these findings suggest that extracellular glucose directly enters hugin[+] neurons

through specific glucose transporter and subsequently modulate the activity of K_{ATP} through

ATP production, ultimately leading to neuronal depolarization.

AstA[+] neurons were direct downstream target of hugin[+] neurons

Since hugin[+] neurons were directly involved in glucose sensing and regulated sweet sensation

(***Figure 1G and Figure supplementary 8A-B***), we sought to identify their downstream target.

We then asked whether the hugin peptide had a direct effect on sweet sensation. Upon

microinjection of synthetic hugin peptide, we observed a notable reduction in sweet sensation

under both sated and starved states (*Figure 3A*). In line with this observation, following the

injection of hugin peptide, we observed a significant reduction of sugar-induced calcium

responses in sweet-sensing Gr5a+ neurons (*Figure 3B*). Thus, the inhibitory function of

hugin+ neurons on sweet sensation was likely mediated by the secretion of hugin peptide.

There were two hugin receptors, PK2-R1 and PK2-R2, in fruit flies (Park et al., 2002). We

knocked down their expression by RNAi in the nervous system and then tested flies' PER to

sugar. Knocking down PK2-R1 but not PK2-R2 significantly enhanced sweet perception in

both sated and starved states (*Figure 3C-D*), suggesting a crucial role of PK2-R1 in receiving

the input of hugin+ neurons in satiety sensing. To validate this finding, we further tested PK2-

R1 null mutants and observed a similar enhancement in sweet sensitivity (*Figure*

supplementary 10). Interestingly, we did not observe any expression of PK2-R1 in the

proboscis where gustatory sensory neurons were distributed, indicating that sweet-sensing

Gr5a+ neurons could not directly receive input from hugin+ neurons via PK2-R1 (data not

shown).

AstA+ neurons served as a plausible relay between hugin+ neurons and sweet-sensing Gr5a+

neurons, especially based on a clear co-expression of hugin receptor PK2-R1 and AstA in the

SEZ region of the fly brain (*Figure 3E*). Notably, these double-labeled neurons corresponded

precisely to the AstA subcluster that exhibited glucose-specific Ca^{2+} responses (*Figure 1H*).

We also used optogenetic tools to further confirm that hugin$^+$ neurons acted upstream of

AstA$^+$ neurons. We expressed the light-sensitive neuronal activator CsChrimson in hugin$^+$

neurons and calcium indicator GCaMP6m in AstA$^+$ neurons, respectively (Klapoetke et al.,

2014). As shown in Figure 3F, opto-activation of hugin$^+$ neurons led to a robust calcium

transient in AstA$^+$ neurons.

We then investigated whether the hugin-AstA circuitry affected sweet sensation. As expected,

knocking-down PK2-R1 (but not PK2-R2) in AstA$^+$ neurons led to a significant increase in

sugar-induced PER (***Figure 3G-H***). Therefore, AstA$^+$ neurons are the direct downstream

target of hugin$^+$ neurons in satiety sensing. hugin$^+$ neurons can directly sense elevated

circulating glucose levels, promote the release of neuropeptide hugin, and then activate AstA$^+$

neurons via its cognate receptor PK2-R1.

hugin$^+$ neurons are distributed in both the brain and the ventral nerve cord (VNC). The hugin$^+$

neurons in the VNC sense sugar and inhibit feeding behavior by suppressing the activity of

DH44 neurons (Oh et al., 2021). To determine which hugin$^+$ neurons regulate the activity of

AstA neurons and sweet sensation, we performed the following experiments: 1, Physically

severing the neural connection between the brain and the VNC: After severing the neural

connection, activation of hugin$^+$ neurons were still able to downregulate the PER behavior,

suggesting that hugin neurons in the SEZ regulated sweet perception (***Figure supplementary***

11A-B). 2, Optogenetic experiments in isolated brains: We dissected the fly brains in Figure

3F, and upon light stimulation, we observed a significant calcium response in AstA neurons

(*Figure supplementary 11C*). This indicates that hugin[+] neurons located in the SEZ but not

VNC regulate the activity of AstA neurons. Therefore, we can infer that hugin neurons in the

SEZ are responsible for sugar sensing, energy signal transmission, and the output of PER

behavior.

AstA[+] neurons directly inhibited Gr5a[+] neurons.

Activation of AstA[+] neurons suppressed sweet sensation, resembling the effect of activating

hugin[+] neurons (Hergarden et al., 2012)(*Figure supplementary 8C-D*). We also evaluated the

behavioral effect of AstA peptide. Similar to hugin, the microinjection of AstA resulted in a

reduction in sweet sensitivity in fruit flies (*Figure 4A*) and the activity of Gr5a[+] neurons

(*Figure 4B*). We then speculated whether AstA[+] neurons and AstA peptide might function

directly on Gr5a[+] neurons. The cell bodies of sweet-sensing Gr5a[+] neurons are located in the

proboscis, whereas their axons extend to the SEZ region of the fly brain (Dahanukar et al.,

2007). We observed distributions of the projections of AstA[+] neurons and Gr5a[+] neurons both

in the SEZ region (*Figure 4C*), suggesting that Gr5a[+] neurons might directly receive AstA

signal in this region.

AstA has two cognate receptors in fruit flies, AstA-R1 and AstA-R2 (Larsen et al., 2001). Co-

labeling experiments in the proboscis confirmed the expression of AstA-R1 in Gr5a[+] neurons

(*Figure 4D, arrows*), while AstA-R2 was not expressed in the proboscis (data not shown).

Functionally, knockdown of AstA-R1 specifically in Gr5a[+] neurons, but not that of AstA-R2,

significantly enhanced sweet sensation of flies (*Figure 4E-F*). Similar results were observed

in AstA-R1 knockout flies (***Figure supplementary 12***).

Using optogenetic tools, we also confirmed the function of hugin$^+$ neurons and AstA$^+$ neurons

on Gr5a$^+$ neurons. Activating hugin$^+$ neurons and AstA$^+$ neurons both significantly reduced

sugar-induced responses of Gr5a$^+$ neurons (***Figure 4G-H***). Collectively, these results suggest

that satiety-sensing hugin-AstA circuitry can directly modulate sweet sensation of flies via

suppressing the activity of Gr5a$^+$ gustatory neurons.

hugin–AstA circuitry suppressed food consumption.

Collectively, the hugin-AstA circuitry senses satiety signals and inhibits sweet sensation.

Considering the crucial role of sweet sensation in regulating feeding behavior, we

hypothesized that hugin-AstA circuitry was also involved in the regulation of food

consumption.

We expressed a temperature-sensitive blocker of synaptic transmission (Shibire^{ts1}) in hugin$^+$

neurons or AstA$^+$ neurons (Kitamoto, 2001) and assayed the feeding behavior of flies under

both permissive temperature (***Figure supplementary 13A, blue***) and non-permissive

temperature (***Figure supplementary 13A, red***). We found that inhibiting the activity of either

hugin$^+$ neurons or AstA$^+$ neurons significantly increased food consumption (***Figure***

supplementary 13B-C), demonstrating the inhibitory role of the hugin-AstA circuitry on

feeding behavior.

We also examined the role of hugin and AstA receptors in feeding behavior by testing *PK2-R1*
and *AstA-R1* gene knockout flies. The results showed that knockout of both receptors
enhanced feeding behavior (***Figure supplementary 13D***), consistent with increased sweet
perception upon genetic manipulation of these two receptors (***Figure 3-4***). Moreover, these
receptor knockout flies exhibited increased lipid storage, which was likely a consequence of
their enhanced feeding behavior (***Figure supplementary 13E***). Additionally, we knocked
down the expression of AstA-R1 in sweet sensing Gr5a$^+$ neurons, and found that these flies
exhibited a significant increase in food consumption (***Figure supplementary 13F***), also likely
due to the enhanced sweet sensation (***Figure 4E***).

Mammalian homolog of hugin was also a putative satiety sensor

Our present work identified hugin-AstA circuitry as a novel satiety sensor and feeding
suppressor in fruit flies. Fruit flies and mammals share highly conserved regulatory elements
on feeding behavior and energy homeostasis (Song et al., 2023; Stocker, 2004; Yarmolinsky et
al., 2009). We thus asked whether this novel satiety-sensing mechanism was also conserved in
the mammalian system,

In rodent models, neuromedin U (NMU), a analogues of hugin, was also found to be an
appetite suppressor (Melcher et al., 2006; Schlegel et al., 2016). We found that synthetic
NMU could suppress sweet perception when injected into flies' thorax (***Figure
supplementary 14***), highlighting the analogy of fly hugin and mammalian NMU. In mouse
models, we observed a significant increase in NMU levels in the blood following glucose

feeding (***Figure 5A***). Using the sucrose preference test (SPT) to measure sweet preference,

we noted that NMU injection reduced sugar preference in mice, phenocopying the effect of

glucose feeding (***Figure 5B-C***). Conversely, NMU knockout enhanced sweet sensation in

mice (***Figure 5D***). These data demonstrated that both NMU gene and protein plays a crucial

role in regulation of sweet sensation.

Neuropeptide NMU is secreted by a group of neurons located in the VMH region of

hypothalamus (Melcher et al., 2006). To investigate whether NMU$^+$ neurons functioned as an

energy sensor in the mouse brain, we injected AAV-EF1α-DIO-GCaMP6m-WPRE into the

VMH of NMU-Cre mice, specifically expressing GCaMP6m in NMU$^+$ neurons. We then

monitored the dynamic activity of VMH^{NMU} neurons during glucose stimulation by using

fiber photometry (***Figure 5E***). As expected, these neurons showed significant calcium

responses following glucose administration in mice (***Figure 5F-G***). To confirm the specificity

of GCaMP6m expression in NMU$^+$ neurons, we co-expressed a Cre-dependent mCherry

reporter in the VMH of NMU-Cre mice and performed fluorescence in situ hybridization for

NMU mRNA. Colocalization analysis demonstrated that GCaMP6m fluorescence was

restricted exclusively to NMU$^+$ cells (***Figure supplementary 15A***).

Next, we conducted calcium imaging of brain slices *in vitro* to further assess these neurons'

ability to sense glucose (***Figure 5H***). Similar to hugin neurons in the fly brain, VMH^{NMU}

neurons in the mouse brain exhibited strong calcium responses to glucose stimulation, which

could not be inhibited by TTX (***Figure 5I***), indicating their direct glucose-sensing capability.

Also similar to flies, glucose activated these mouse neurons via entering the cells and

modulating K_{ATP} channels, as evidenced by the inhibition of calcium responses by alloxan and

phlorizin (*Figure 5J*).

Calbindin 2-positive neurons (Calb2) in the rostral nucleus of the solitary tract (rNST)

selectively respond to sweet tastants, facilitating sweet signal from the periphery into the

brain (Jin et al., 2021). Because attenuated sweet sensation should correspondingly diminish

central sweet-signal transmission, we asked whether NMU modulates this relay. To assess

potential anatomical connectivity between VMH NMU$^+$ neurons and rNST Calb2$^+$ neurons,

we employed a dual-virus tracing strategy. In NMU-Cre mice, we injected AAV9-EF1α-DIO-

EGFP into the VMH, which revealed dense GFP$^+$ projections in close apposition to Calb2$^+$

neurons in the rNST (*Figure supplementary 15B*). To further confirm this projection is

monosynaptic, we combined anterograde tracing by co-injecting AAV2/1-EF1α-DIO-EGFP

into the VMH and AAV2/1-hsyn-SV40-NLS-Cre into the rNST. This approach yielded GFP

expression in rNST Calb2$^+$ neurons, supporting a direct VMH→rNST NMU$^+$→Calb2$^+$

synaptic link (*Figure 5K*).

To evaluate the functional consequence of NMU on Calb2$^+$ neuronal activity during sweet

stimulation, we delivered AAVs encoding Cre-dependent GCaMP6m into the rNST of Calb2-

Cre mice, thereby enabling real-time calcium imaging in Calb2$^+$ neurons (*Figure*

supplementary 15C). Using fiber photometry, we recorded calcium responses to sugar

stimuli, both with and without NMU injection (*Figure 5L*). We found that NMU injection

significantly suppressed the calcium responses to sugar, aligning with observed behavioral

phenotypes (*Figure 5M*). These data indicated that the neuropeptide NMU plays a role in

regulating sweet taste perception during sugar feeding.

These findings indicate that NMU$^+$ neurons can indeed be a putative satiety sensor in the

mouse brain that suppresses sweet perception, shedding light on understanding the conserved

neural mechanism of feeding regulation in mammals including human.

Discussion

In summary, we identified hugin-AstA circuitry as a novel satiety sensor in the fly brain. It directly detects circulating sugar levels, inhibits sweet taste perception, and terminates feeding, helping to maintain the homeostasis of internal energy state (***Figure supplementary 16***). Previous research has shown that hugin[+] neurons project to the pharyngeal muscles and the ring gland, playing a role in regulating both feeding and sleep , and these neurons are also part of a neural circuit that modulates taste-mediated feeding behavior. (Melcher and Pankratz, 2005; Miroschnikow et al., 2020; Oh et al., 2021). AstA[+] neurons convey carbohydrate-satiety signals and shift dietary preference from sugar to protein (Hentze et al., 2015; Hergarden et al., 2012). . However, it was unclear whether they could sense the body's energy state, how they interacted with each other, or how they influenced sensory/motor neurons to regulate feeding. Our research has clarified the relationship between hugin and AstA: hugin functions upstream, sensing glucose, while AstA acts downstream, directly modulating sweet-sensing Gr5a[+] neurons to regulate sugar perception and feeding behavior. This discovery establishes a mechanistic link between glucose sensing and sweet sensation regulation, enhancing our understanding of how energy states influence nutrient sensing and feeding behavior.

Recent studies have highlighted the complex interplay of neural circuits and neuropeptides that regulate satiety and feeding behavior in Drosophila. For instance, dopaminergic and neuropeptide F (NPF) circuits play crucial roles in modulating the activity of Gr5a[+] sweet-sensing neurons during hunger and refeeding. These studies showed that dopamine signaling

adjusts the temporal integration of PER-driving inputs under fed conditions, while NPF

enhances Gr5a sensitivity during starvation (Inagaki et al., 2012; Inagaki et al., 2014; Marella

et al., 2012). Our findings complement these established mechanisms by identifying the

hugin–AstA pathway as a central satiety sensor that directly detects circulating glucose levels

and inhibits sweet taste perception to prevent overconsumption. Moreover, the hugin–AstA

circuitry fits within a broader regulatory framework of satiety signaling in Drosophila .

Peripheral satiety signals, such as enteroendocrine NPF$^+$ cells, detect luminal sugar and

suppress feeding via hormonal release (Malita et al., 2022). Central hunger circuits, including

dopaminergic neurons, adjust Gr5a responsiveness according to the energy state (Inagaki et

al., 2012). Our study reveals that the hugin–AstA loop acts as a central brake on Gr5a activity

following sugar intake, thus closing a feedback loop that protects against excessive sugar

ingestion. Together, these multilayered mechanisms—from gut to brain—ensure robust

regulation of sweet perception and energy balance.

Although multiple regulatory elements of energy homeostasis have been identified in both

flies and mammals, our present work reveals a novel mechanism that links energy sensing to

taste modulation. The energy sensor comprises three types of neurons: hugin$^+$ neurons as the

sensor, AstA$^+$ neurons as a relay, and sweet-sensing Gr5a$^+$ neurons as the behavioral

modulator. Given the highly conserved regulatory mechanism of energy homeostasis across

animal species (*Figure 5A-D*), our work paves a way to fully understand how internal state

sensing and behavioral alternations are tightly coupled in more complex organisms, and how

they are disrupted in metabolic disorders.

In addition to the novel mechanism we have uncovered in *Drosophila*, it is interesting to

consider how conserved these pathways might be in more complex organisms like mammals,

particularly concerning the function of the mammalian homologues, NMU (Melcher et al.,

2006). NMU is known to play critical roles in regulating energy homeostasis, feeding

behavior, insulin secretion, and immune responses in mammals (Teranishi and Hanada, 2021).

Our study has shown that NMU neurons may serve as an energy sensor in mammals as well,

which regulated nutrient sensing, much like the hugin neurons in *Drosophila*. Given the

functional parallels between the hugin-AstA circuitry in flies and NMU signaling in

mammals, it is plausible that NMU may interfere with taste signaling to the brain and

modulate the brain's reward circuits, affecting other behaviors beyond feeding.

Additionally, our findings open up the possibility of using NMU or its pathways as

therapeutic targets for appetite suppression and obesity treatment. Recent developments in

appetite-regulating peptides, such as glucagon-like peptide 1 (GLP-1) receptor agonists, have

shown efficacy in reducing food intake and promoting weight loss in humans (Aldawsari et

al., 2023). Given NMU's role in suppressing sweet taste perception and energy regulation, the

NMU system could be a complementary or alternative target for anti-obesity interventions.

Exploring how modulating NMU activity could synergize with GLP-1 therapies may provide

new strategies for controlling appetite and managing energy balance.

In conclusion, our discovery of the hugin-AstA circuitry in flies provides a foundation for

understanding the broader principles of energy sensing and behavioral regulation across

species. Further studies in mammals will be essential to explore the potential roles of NMU

neurons in these processes and their therapeutic relevance in metabolic disorders.

Materials and methods

Animals

Fly: Flies were maintained on a standard fly diet under conditions of 60% humidity and 25 °C with a 12-hour light and 12-hour dark cycle. Virgin female flies were chosen immediately after emerging and housed in standard fly food (ND) vials, with 20 flies per vial, for a period of 5-6 days before experiments. For temperature-sensitive manipulations (dTrpA1, Shibire^ts), flies were reared at 18 °C for 7–8 days, then transferred to 30 °C for 30 min to activate or inhibit targeted neurons; behavioral assays (PER or feeding) were conducted continuously at 30 °C. For starvation, the flies were kept on 2 % Agar for 12-hour.

The following UAS-RNAi lines: *UAS-AstA-R1 RNAi (#27280)*, *UAS-AstA-R2 RNAi (#25935)*, *UAS-PK2-R1 RNAi (#29624)*, *UAS-PK2-R2 RNAi (#28781)*, *UAS-CD8-GFP, LexAop-CD2-RFP (#67093)*, *UAS-GCaMP6m (#42748)*, *UAS-CaMPARI (#58764)*, Gr5a-GAL4 (#57592), *Gr5a-LexA* (#93014), *ilp2-GAL4 (#37516)*, *elav-GAL4 (#25750)* were obtained from the Bloomington Drosophila Stock Center at Indiana University. *TH-KO*, and all the other Gal4 and LexA lines were from Yi Rao (Deng et al., 2019) (Capital Medical University, Beijing). *UAS-dTrpA1* and *UAS-Shibire^{ts1}* were from David Anderson (Caltech). *UAS-GCaMP6m; LexAop-CsChrimson* was from Yufeng Pan (Southeast University, Nanjing). *LexAop-Chrimson* was from Wei Zhang (Tsinghua University, Beijing).

Mouse: Male C57BL/6J mice at the age of four weeks were procured from cyagen. The mice were kept in standard laboratory conditions at a temperature of 25 °C with a 12:12 light/dark

cycle. All experimental procedures were conducted in compliance with the guidelines set by

the Laboratory Animal Welfare and Ethics Committee of Chongqing University (CQU-

IACUC-RE-202401-004), adhering to both national and international standards.

Behavioral assays (Fly)

PER was evaluated using a method outlined in a previous study. Briefly, individual flies were

gently aspirated and positioned in a 200 μL pipette tip, allowing only the head and proboscis

to protrude while the dorsal thorax was lightly affixed to prevent escape. After a 3-minute

acclimation period, each fly was first presented with 0.5 μL water droplets on the labellum

twice. Flies that responded by extending their proboscis were allowed to drink and were

subsequently excluded from the experiment. Flies that did not extend their proboscis in

response to water were then stimulated sequentially with ascending sucrose concentrations

(6.25–800 mM). Each concentration was delivered as two brief (<1 s) touches of a 0.5 μL

droplet to the labellum, immediately withdrawn to prevent ingestion. A full proboscis

extension, defined as the complete uncoiling of the rostrum beyond the labellum plane, was

scored as positive. Partial or delayed (>1 s) extensions were not considered positive responses.

Responsiveness at each concentration was recorded if at least one full extension occurred in

the two trials.For Figure supplementary 11, the connection between the brain and VNC were

cut off by a dissecting scissors before the PER assay.

Feeding was assessed as previously described (Sun et al., 2017). Briefly, 5-day-old flies

underwent a 12-hour starvation period on 2% agar and were then moved to a new vial with

test food containing 0.5% Brilliant Blue (MACKLIN, China) for 10 minutes. After a rapid

freeze at -20 °C, flies were decapitated in PBS, homogenized, and centrifuged (13,000 g * 5

min). The resulting supernatants were diluted with PBS to a total volume of 1 ml, and

absorbance was measured at 620 nm.

Sucrose Preference Test

The two-bottle preference (TBP) behavioral test was conducted in a standard mouse cage.

Two bottles, one containing water and the other with a 10 mM sucrose solution, were placed

on top of the cage. Each tested mouse was individually caged and adopted to two bottles

containing distilled water or a 10 mM sucrose solution for one week. Throughout the training

period, the positions of the bottles were alternated every 24 hours to mitigate potential

position-related biases. Prior to the TBP tests for sucrose, mice underwent a 12-hour pre-

starvation period, followed by re-fed sucrose (20%) for 1 hour or intraperitoneal (IP)

administration of NMU (YFLFRPRN-NH$_2$, 4.5 μm/kg). The consumption of water and

sucrose solution over a 2-hour period was meticulously recorded. The sweetener preference

ratio was then calculated by dividing the weight of sweetener consumed by the weight of

water consumed.

NMU measurement

Blood was obtained from the lower jaw of the mice and clotted at room temperature for 1

hour. The samples were then centrifuged (1,500g, 4 °C) for 10 minutes and the supernatants

were collected for NMU measurement. All the manipulations were according to the

manufacturer's instructions (BMASSY,74030).

Hemolymph extraction and glucose measurement

Groups of 40 ice-anesthetized flies were decapitated and placed into a perforated 0.5 ml tube,

which was nested inside a 1.5 ml collection tube. Samples were centrifuged at $2,500 \times g$ for

10 min at 4 °C, yielding approximately 2 µl of hemolymph per batch. Hemolymph was

pooled to 4 µl per assay (n = 8 independent samples per group) and immediately analyzed for

glucose concentration using the Solarbio BC2500 kit (Solarbio, China) according to the

manufacturer's instructions.

Microinjection

Flies were delicately positioned and secured within a 200 ml pipette tip, ensuring their heads

were directed toward the tip's end. Subsequently, the tip was opened by cutting, exposing the

heads and a section of the thorax. About 20 nl of sterile saline, either with or without

synthesized peptides (hugin: SVPFKPRL-NH$_2$, 1 mg/ml (1.1 mM); AstA: LPVYNFGL-NH$_2$,

1mg/ml (1.1 mM); NMU: YFLFRPRN-NH$_2$, 1mg/ml (0.9 mM); Control peptide:

DYKDDDDKYPYDVPDYA, 1mg/ml (0.48 mM), were injected into the thorax of these flies

using a glass micropipette and a microinjector (3-000-207, Drummond Scientific Company

Instruments). The glass micropipette was created from thick-walled borosilicate capillaries (3-

000-203-G/X, Drummond Scientific Company Instruments).

Immunofluorescence staining

Fly brains were carefully dissected in PBS on ice and then fixed in 4% formaldehyde for 60

minutes. Following fixation, the brains underwent permeabilization and blocking using

Dilution/Blocking Buffer (PBS containing 10% Calf Serum and 0.5% Triton X-100) for 2

hours at room temperature. Subsequently, the samples were immersed in the appropriate

primary antibodies in Dilution/Blocking Buffer for 24 hours at 4 °C. Afterward, the samples

were subjected to a 60-minute wash with Washing Buffer (0.5% Triton X-100 in PBS) four

times at room temperature and were then incubated with secondary antibodies for 24 hours at

4 °C. The samples underwent three additional washes with Washing Buffer before being

mounted in Fluoroshield (Sigma-Aldrich).

Images were acquired using a scanning confocal microscope with Olympus objectives (20×

/0.7 and 40× /0.95w). Antibodies were employed at the following dilutions: mouse anti-nc82

(1:100, DSHB), rabbit anti-GFP (1:500, Abcam), Alexa Fluor 647 goat anti-mouse (1:500,

Invitrogen), and Alexa Fluor 488 goat anti-rabbit (1:500, Invitrogen).

In situ hybridization

Mice were anesthetized and perfused with 4% paraformaldehyde (PFA) dissolved in

phosphate-buffered saline (PBS, pH 7.4). Brains were carefully removed from the skull and

post-fixed in 4% PFA at 4 °C overnight. Subsequently, the brains were transferred to a 20%

sucrose solution (in 1 × PBS) at 4 °C until they were ready for sectioning. Twenty-

micrometer-thick sections were obtained using a cryostat. The detection of nmu was

performed using a fluorescent in situ hybridization technique (RNAscope, Pinpoease)

according to the manufacturer's instructions. Sections were mounted on SuperFrost Plus Gold

slides (ThermoFisher) and briefly rinsed in autoclaved Millipore water. They were then

subjected to gradient dehydration in 50%, 75%, and 100% ethanol for 5 minutes each. A

hydrophobic barrier was created around the sections using an ImmEdge hydrophobic barrier

pen (Cat No. 310018). All incubation steps were carried out at 40 °C using the Pinpoease

hybridization system (Cat No. SH-08). The subsequent hybridization, amplification, and

detection steps were performed strictly according to the manufacturer's instructions.

Stereotaxic brain surgeries and viral injection

Mice were anesthetized with 1-2% isoflurane during stereotaxic injections, performed using a

small animal stereotaxic instrument (RWD Life Science, #68030). Throughout the surgery,

core body temperature was maintained at 36 ± 1 °C using a feedback-controlled heating

system. Micro scissors were used for the incision, and a dental drill was employed to create

the cranial window. Viral injections were delivered using a glass microelectrode syringe

pump (RWD Life Science, #R-480). To express GCaMP6m and mCherry AAV2/9-EF1α-

DIO-GCaMP6m-WPRE (titer: 2.89×10^{12}, #H4955, Obio Biotechnology, Shanghai, China,

Co., Ltd.) and AAV2/9-EF1α-DIO-mCherry-WPRE (titer: 2.89×10^{12}, Obio Biotechnology,

Shanghai, China, Co., Ltd.) were bilaterally injected into the VMH and rNST, with

approximately 0.15 μL of virus delivered per site. The microelectrode was left in place for 10

minutes post-injection before being withdrawn. Injection coordinates were as follows: VMH

(AP, −1.46 mm; ML, ±0.4 mm; DV, −5.5 mm) and rNST (AP, −7.08 mm; ML, ±0.6 mm; DV,

−4.3 mm). After 3 weeks of AAV injections, optical fibers were chronically implanted in the

VMH or rNST and secured with dental cement (C&B Metabond®, Parkell, Japan). Injection

coordinates were determined based on the 4th edition of the mouse brain atlas by Franklin and

Paxinos (2008). Mice were allowed to recover for 5-7 days post fiber-implanted before any

behavioral or calcium imaging evaluations were conducted. After the experiments, mice were

perfused to verify virus expression and fiber placement. Data from mice with poor viral

expression or incorrect fiber placement (0-20% of cases) were excluded from the final

analysis.

For the anterograde tracing experiments, two viral vectors were used. AAV2/1-hsyn-DIO-

EGFP-WPRE (titer: 1×10^{13} viral particles/mL,Obio Biotechnology) was injected into the

ventromedial hypothalamus (VMH), and AAV2/1-hsyn-SV40 NLS-Cre (titer: 2.1×10^{12} viral

particles/mL, Brain Case) was injected into the rostral nucleus of the solitary tract (rNST).

Approximately 0.15 μL of virus was delivered per injection site. After 3 weeks, the brains

were perfused and fixed. Subsequently, the brain tissues were sliced and subjected to antibody

staining. The fluorescence signals were then recorded and analyzed.

Optical-fiber-based calcium recording in freely mice

Following injection of an AAV2/9-EF1α-DIO-GCaMP6m-WPRE viral vector, an optical

fiber (200 μm O.D., 0.37 numerical aperture, Inper Inc., China) was placed 150 μm above the

viral injection site. GCaMP6m$^+$ mice were implanted with the optic fiber 3 weeks post-AAV

injection and allowed to recover for 5–7 days prior to experimental testing.

For Figure 5K-L, mice were first trained to lick glucose (500 mM) from a petri dish. After

training, they were fasted for 12 hours and then received an injection of synthetic NMU

peptide, followed by a 30-minute waiting period. During the experimental session, the mice

were given 10 seconds to lick glucose from the petri dish while undergoing fiber photometry

recording. For Figure 5E-G, mice were fasted for 4 hours and then administered glucose via

gastric infusion, with fiber photometry recordings capturing the process. Fluorescence signals

were acquired using a dual-channel fiber photometry system (410 nm & 470 nm, RWD Life

Science). The laser power at the tip of the optical fiber was kept below 20 μW for both the

470 nm and 410 nm channels to minimize bleaching. $\Delta F / F$ was calculated as (470 nm signal

- fitted 410 nm signal) / fitted 410 nm signal. Data analysis was performed using software

from RWD Life Science.

Ex vivo calcium imaging

For Figure 1G, 1I, figure supplementary 6, and Figure 2freshly dissected fly brains were

placed in the sugar-free AHL buffer (108 mM NaCl, 8.2 mM $MgCl_2$, 4 mM $NaHCO_3$, 1 mM

NaH_2PO_4, 5 mM KCl, 2 mM $CaCl_2$, 5 mM HEPES, pH 7.3). Baseline recordings of the

samples in AHL buffer were taken over 1 minutes. Subsequently, the solutions were added

with 100 μM glibenclamide (*Figure 2D*), 50 mM pyruvate (*Figure 2G*) or switched to AHL

+ glucose (50 mM) with or without indicated chemicals (*Figure 2 A and E*, 1 mM phlorizin

or 10 um alloxan), and the pH was adjusted back to 7.3 through gentle perfusion for 3 minutes.

Solution flow in the perfusion chamber was controlled by a valve commander (Scientific

Instruments). Following stimulation, samples were washed out again with AHL. For the TTX

test, 2 μM TTX was added to the AHL solution. Prior to the assay, the samples were pre-

treated with 2 mM TTX for 15 minutes.

For Figure 5H-J, mice (6-9 weeks of age) were deeply anesthetized by avertin (1.25% avertin,

0.2 ml/10g) . Mouse brains were quickly extracted following decapitation and immediately

placed in ice-cold slicing buffer (110 mM Choline Cl, 2.5 mM KCl, 1 mM NaH_2PO_4, 25 mM

$NaHCO_3$, 5 mM glucose, 7 mM $MgCl_2$, 0.5 mM $CaCl_2$,1.3mM Na ascorbate,0.6mM Na

pyruvate bubbled with 95% oxygen and 5% CO_2 with an adjusted pH of 7.3). The brains were

then sectioned into 200 μm slices using a vibratome (Leica VT1200S). These slices were

incubated artificial cerebrospinal fluid (aCSF) (125 mM NaCl, 2.5 mM KCl, 1 mM NaH_2PO_4,

25 mM $NaHCO_3$, 5 mM glucose, 1.3 mM $MgCl_2$, 2 mM $CaCl_2$,1.3mM Na ascorbate, 0.6 mM

Na pyruvate bubbled with 95% oxygen and 5% CO_2 with an adjusted pH of 7.3) at 34 °C for

20 minutes, then transferred to a room temperature for 30 minutes before recording. For Ca^{2+}

imaging, the slices were placed in a recording chamber in the low sugar aCSF (125 mM NaCl,

2.5 mM KCl, 1 mM NaH_2PO_4, 25 mM $NaHCO_3$, 1 mM glucose, 1.3 mM $MgCl_2$, 2 mM $CaCl_2$,

1.3 mM Na ascorbate, 0.6mM Na pyruvate, pH 7.3) for 10 minutes (maintained the flow rate

for 1-3 ml/ minute). Baseline recordings, perfusion, and Ca^{2+} recording were performed as

detailed above.

All imaging was conducted on an Olympus confocal microscope (FVMPE-RS) with a water

immersion objective lens (25× /1.05w MP). Image analyses were performed using ImageJ to

calculate the mean intensity of the indicated neuron targeting ROIs, and then plotted in Excel

(Microsoft). Ratio changes were calculated using the formula: $\Delta F / F = [F - F_0] / F_0$, where F

represents the mean fluorescence of the cell body, and F_0 is the average baseline (before

stimulation).

Optogenetics and in vivo calcium imaging

Newly emerged virgin female flies were housed in a fresh vial with standard medium for 3

days and subsequently moved to a vial containing regular food supplemented with 400 μM

all-trans-retinal (Sigma R2500) for 2-4 days prior to experiments. These flies were then

immobilized on ice, affixed to transparent tape, and the dorsal cuticle of the fly head was

delicately removed with forceps to expose the brain, which was immersed in AHL.

For Figure 3F, an array of red LED (Thorlabs, M625L3) was positioned above the brain.

Calcium signals of Gr5a$^+$ neurons were captured for 60 frames (1 second per frame) in the

absence of light to establish a baseline. Subsequently, the next 20 frames were recorded

during opto-activation with the light (the light was switched on for 0.45 second and rested for

0.55 second per activation cycle, and the images were collected in rest time), followed by

another 60 frames recorded without light afterward.

For Figure 4G and 4H, to apply liquid food (5% sucrose) to the proboscis, a pipette was filled

with the taste solution and placed a few microns from the proboscis tip by using a

micromanipulator (MP225, Sutter Instruments) prior to recording. Then, we recorded calcium

signals for 60 seconds in the absence of light to establish a baseline (1 second per frame). To

activate neurons expressing CsChrimson, a red LED was positioned above the brain,

controlled by a custom computer program during opto-activation (0.45 second on and 0.55

second off from 60 second to 120 second). The pipette was placed on the proboscis from 80

second to 85 second. Calcium signals of Gr5a$^+$ neurons were recorded throughout these

processes (1 second per frame).

Image analyses about relative fluorescence changes $\Delta F / F$ were performed as ex vivo calcium

imaging.

Optogenetics and ex vivo calcium imaging

For Figure supplementary 11C, newly emerged virgin female flies were housed in a fresh vial

with standard medium for 3 days and subsequently moved to a vial containing regular food

supplemented with 400 µM all-trans-retinal (Sigma R2500) for 2-4 days prior to experiments.

The brains were dissected and placed under a red LED. Calcium signals of AstA$^+$ neurons

were recorded with or without LED light as above. Image analyses were performed as ex vivo

calcium imaging.

Calcium imaging with CaMPARI

The flies were then gently immobilized on ice, attached to transparent tape, and the dorsal

cuticle of the fly head was carefully removed with forceps to expose the brain, which was

immersed in AHL. Photoconversion (PC) was achieved for 30 seconds using a 405 nm LED

[Thorlabs, M405L2–UV (405 nm) Mounted LED, 1,000 mA, 410 mW], controlled by a LED

controller (Thorlabs, LEDDB1 driven with 1000 mA). Images were captured using an

Olympus confocal microscope. Image analyses were performed in ImageJ by manually

drawing ROIs covering individual neuronal cell bodies using the green channel. The

extent of CaMPARI photoconversion was determined as the Red: Green ratio.A '-405 nm'

control was included to demonstrate that scanning of the hugin neurons without exposure to

ultraviolet light does not convert green to red fluorescence.

Statistical analysis

Data are shown as means (± SEM). Data presented in this study were verified for normal

distribution by D'Agostino–Pearson omnibus test. Student's t test, one-way ANOVA and

two-way ANOVA (for comparisons among three or more groups and comparisons with more

than one variant) were used. The post hoc test with Bonferroni correction was performed for

multiple comparisons following ANOVA.

Acknowledgements

We express gratitude to all members of the Wang and Huang Lab for their valuable

discussions and technical support. Special thanks to Yi Rao and Bowen Deng for generously

providing *Drosophila* chemoconnectome lines, and to Yulong Li and Jianzhi Zeng for their

assistance with *in vivo* calcium imaging preparations. Appreciation is extended to the

members of the Neuroscience Pioneer Club for their insightful discussions throughout the

course of this study. This study was funded by the National Natural Science Foundation of

China (No. 32071006 for L.W. and No. 32271008 for R.H.), Guangdong Basic and Applied

Basic Research Foundation (2023A1515110454 for T.S.), and the startup funds from

Shenzhen Bay Laboratory and Chinese Institutes for Medical Research to L.W.

Author contributions

774 T.S., R.H., and L.W. designed the experiments and wrote the manuscript. W.Q., and T.S.

775 conducted and analyzed all the experiments. L.W. and R.H. supervised the project.

Competing interests

778 The authors declare no competing interest.

References

Aldawsari, M., Almadani, F.A., Almuhammadi, N., Algabsani, S., Alamro, Y., and Aldhwayan,

M. (2023). The Efficacy of GLP-1 Analogues on Appetite Parameters, Gastric Emptying, Food

Preference and Taste Among Adults with Obesity: Systematic Review of Randomized

Controlled Trials. Diabetes Metab Syndr Obes *16*, 575-595.

Alhadeff, A.L., Su, Z., Hernandez, E., Klima, M.L., Phillips, S.Z., Holland, R.A., Guo, C.,

Hantman, A.W., De Jonghe, B.C., and Betley, J.N. (2018). A Neural Circuit for the

Suppression of Pain by a Competing Need State. Cell *173*, 140-152 e115.

Augustine, V., Lee, S., and Oka, Y. (2020). Neural Control and Modulation of Thirst, Sodium

Appetite, and Hunger. Cell *180*, 25-32.

Belgardt, B.F., Okamura, T., and Bruning, J.C. (2009). Hormone and glucose signalling in

POMC and AgRP neurons. J Physiol *587*, 5305-5314.

Brogiolo, W., Stocker, H., Ikeya, T., Rintelen, F., Fernandez, R., and Hafen, E. (2001). An

evolutionarily conserved function of the Drosophila insulin receptor and insulin-like peptides in

growth control. Curr Biol *11*, 213-221.

Bruning, J.C., and Fenselau, H. (2023). Integrative neurocircuits that control metabolism and

food intake. Science *381*, eabl7398.

Budnik, V., and White, K. (1987). Genetic dissection of dopamine and serotonin synthesis in

the nervous system of Drosophila melanogaster. J Neurogenet *4*, 309-314.

Burnett, C.J., Li, C., Webber, E., Tsaousidou, E., Xue, S.Y., Bruning, J.C., and Krashes, M.J.

(2016). Hunger-Driven Motivational State Competition. Neuron *92*, 187-201.

Chen, J., Reiher, W., Hermann-Luibl, C., Sellami, A., Cognigni, P., Kondo, S., Helfrich-Forster,

C., Veenstra, J.A., and Wegener, C. (2016). Allatostatin A Signalling in Drosophila Regulates

Feeding and Sleep and Is Modulated by PDF. PLoS Genet *12*, e1006346.

Chen, Y., Lin, Y.C., Kuo, T.W., and Knight, Z.A. (2015). Sensory detection of food rapidly

modulates arcuate feeding circuits. Cell *160*, 829-841.

Chouhan, N.S., Griffith, L.C., Haynes, P., and Sehgal, A. (2021). Availability of food

determines the need for sleep in memory consolidation. Nature *589*, 582-585.

Dahanukar, A., Lei, Y.T., Kwon, J.Y., and Carlson, J.R. (2007). Two Gr genes underlie sugar

reception in Drosophila. Neuron *56*, 503-516.

de Rivaz, R., Swendsen, J., Berthoz, S., Husky, M., Merikangas, K., and Marques-Vidal, P.

(2022). Associations between Hunger and Psychological Outcomes: A Large-Scale

Ecological Momentary Assessment Study. Nutrients *14*.

De Solis, A.J., Del Rio-Martin, A., Radermacher, J., Chen, W., Steuernagel, L., Bauder, C.A.,

Eggersmann, F.R., Morgan, D.A., Cremer, A.L., Sue, M.*, et al.* (2024). Reciprocal activity of

AgRP and POMC neurons governs coordinated control of feeding and metabolism. Nat Metab

6, 473-493.

Deem, J.D., Faber, C.L., and Morton, G.J. (2022). AgRP neurons: Regulators of feeding,

energy expenditure, and behavior. FEBS J *289*, 2362-2381.

Deng, B., Li, Q., Liu, X., Cao, Y., Li, B., Qian, Y., Xu, R., Mao, R., Zhou, E., Zhang, W.*, et al.*

(2019). Chemoconnectomics: Mapping Chemical Transmission in Drosophila. Neuron *101*,

876-893 e874.

Dietrich, M.O., Zimmer, M.R., Bober, J., and Horvath, T.L. (2015). Hypothalamic Agrp

neurons drive stereotypic behaviors beyond feeding. Cell *160*, 1222-1232.

Dus, M., Lai, J.S., Gunapala, K.M., Min, S., Tayler, T.D., Hergarden, A.C., Geraud, E., Joseph,

C.M., and Suh, G.S. (2015). Nutrient Sensor in the Brain Directs the Action of the Brain-Gut

Axis in Drosophila. Neuron *87*, 139-151.

Fosque, B.F., Sun, Y., Dana, H., Yang, C.T., Ohyama, T., Tadross, M.R., Patel, R., Zlatic, M.,

Kim, D.S., Ahrens, M.B.*, et al.* (2015). Neural circuits. Labeling of active neural circuits in vivo

with designed calcium integrators. Science *347*, 755-760.

Fu, O., Iwai, Y., Narukawa, M., Ishikawa, A.W., Ishii, K.K., Murata, K., Yoshimura, Y.,

Touhara, K., Misaka, T., Minokoshi, Y.*, et al.* (2019). Hypothalamic neuronal circuits

regulating hunger-induced taste modification. Nat Commun *10*, 4560.

Fujii, S., Yavuz, A., Slone, J., Jagge, C., Song, X., and Amrein, H. (2015). Drosophila sugar

receptors in sweet taste perception, olfaction, and internal nutrient sensing. Curr Biol *25*, 621-

627.

Hentze, J.L., Carlsson, M.A., Kondo, S., Nassel, D.R., and Rewitz, K.F. (2015). The

Neuropeptide Allatostatin A Regulates Metabolism and Feeding Decisions in Drosophila. Sci

Rep *5*, 11680.

Hergarden, A.C., Tayler, T.D., and Anderson, D.J. (2012). Allatostatin-A neurons inhibit

feeding behavior in adult Drosophila. Proc Natl Acad Sci U S A *109*, 3967-3972.

Inagaki, H.K., Ben-Tabou de-Leon, S., Wong, A.M., Jagadish, S., Ishimoto, H., Barnea, G.,

Kitamoto, T., Axel, R., and Anderson, D.J. (2012). Visualizing neuromodulation in vivo:

TANGO-mapping of dopamine signaling reveals appetite control of sugar sensing. Cell *148*,

583-595.

Inagaki, H.K., Panse, K.M., and Anderson, D.J. (2014). Independent, reciprocal

neuromodulatory control of sweet and bitter taste sensitivity during starvation in Drosophila.

Neuron *84*, 806-820.

Jin, H., Fishman, Z.H., Ye, M., Wang, L., and Zuker, C.S. (2021). Top-Down Control of Sweet

and Bitter Taste in the Mammalian Brain. Cell *184*, 257-271 e216.

Karigo, T., and Deutsch, D. (2022). Flexibility of neural circuits regulating mating behaviors in

mice and flies. Front Neural Circuits *16*, 949781.

Kitamoto, T. (2001). Conditional modification of behavior in Drosophila by targeted expression

of a temperature-sensitive shibire allele in defined neurons. J Neurobiol *47*, 81-92.

Klapoetke, N.C., Murata, Y., Kim, S.S., Pulver, S.R., Birdsey-Benson, A., Cho, Y.K., Morimoto,

T.K., Chuong, A.S., Carpenter, E.J., Tian, Z.*, et al.* (2014). Independent optical excitation of

distinct neural populations. Nat Methods *11*, 338-346.

Koch, M., Varela, L., Kim, J.G., Kim, J.D., Hernandez-Nuno, F., Simonds, S.E., Castorena,

C.M., Vianna, C.R., Elmquist, J.K., Morozov, Y.M.*, et al.* (2015). Hypothalamic POMC

neurons promote cannabinoid-induced feeding. Nature *519*, 45-50.

Kubrak, O.I., Lushchak, O.V., Zandawala, M., and Nassel, D.R. (2016). Systemic corazonin

signalling modulates stress responses and metabolism in Drosophila. Open Biol *6*.

Larsen, M.J., Burton, K.J., Zantello, M.R., Smith, V.G., Lowery, D.L., and Kubiak, T.M. (2001).

Type A allatostatins from Drosophila melanogaster and Diplotera puncata activate two

Drosophila allatostatin receptors, DAR-1 and DAR-2, expressed in CHO cells. Biochem

Biophys Res Commun *286*, 895-901.

Lin, S., Senapati, B., and Tsao, C.H. (2019). Neural basis of hunger-driven behaviour in

Drosophila. Open Biol *9*, 180259.

Malita, A., Kubrak, O., Koyama, T., Ahrentlov, N., Texada, M.J., Nagy, S., Halberg, K.V., and

Rewitz, K. (2022). A gut-derived hormone suppresses sugar appetite and regulates food

choice in Drosophila. Nat Metab *4*, 1532-1550.

Marella, S., Mann, K., and Scott, K. (2012). Dopaminergic modulation of sucrose acceptance

behavior in Drosophila. Neuron *73*, 941-950.

Martelli, C., Pech, U., Kobbenbring, S., Pauls, D., Bahl, B., Sommer, M.V., Pooryasin, A.,

Barth, J., Arias, C.W.P., Vassiliou, C.*, et al.* (2017). SIFamide Translates Hunger Signals into

Appetitive and Feeding Behavior in Drosophila. Cell Rep *20*, 464-478.

Melcher, C., Bader, R., Walther, S., Simakov, O., and Pankratz, M.J. (2006). Neuromedin U

and its putative Drosophila homolog hugin. PLoS Biol *4*, e68.

Melcher, C., and Pankratz, M.J. (2005). Candidate gustatory interneurons modulating feeding

behavior in the Drosophila brain. PLoS Biol *3*, e305.

Miroschnikow, A., Schlegel, P., and Pankratz, M.J. (2020). Making Feeding Decisions in the

Drosophila Nervous System. Curr Biol *30*, R831-R840.

Munch, D., Ezra-Nevo, G., Francisco, A.P., Tastekin, I., and Ribeiro, C. (2020). Nutrient

homeostasis - translating internal states to behavior. Curr Opin Neurobiol *60*, 67-75.

Oh, Y., Lai, J.S., Mills, H.J., Erdjument-Bromage, H., Giammarinaro, B., Saadipour, K., Wang,

J.G., Abu, F., Neubert, T.A., and Suh, G.S.B. (2019). A glucose-sensing neuron pair regulates

insulin and glucagon in Drosophila. Nature *574*, 559-564.

Oh, Y., Lai, J.S., Min, S., Huang, H.W., Liberles, S.D., Ryoo, H.D., and Suh, G.S.B. (2021).

Periphery signals generated by Piezo-mediated stomach stretch and Neuromedin-mediated

glucose load regulate the Drosophila brain nutrient sensor. Neuron *109*, 1979-1995 e1976.

Park, Y., Kim, Y.J., and Adams, M.E. (2002). Identification of G protein-coupled receptors for

Drosophila PRXamide peptides, CCAP, corazonin, and AKH supports a theory of ligand-

receptor coevolution. Proc Natl Acad Sci U S A *99*, 11423-11428.

Parton, L.E., Ye, C.P., Coppari, R., Enriori, P.J., Choi, B., Zhang, C.Y., Xu, C., Vianna, C.R.,

Balthasar, N., Lee, C.E.*, et al.* (2007). Glucose sensing by POMC neurons regulates glucose

homeostasis and is impaired in obesity. Nature *449*, 228-232.

Qi, W., Wang, G., and Wang, L. (2021). A novel satiety sensor detects circulating glucose and

suppresses food consumption via insulin-producing cells in Drosophila. Cell Res *31*, 580-588.

Root, C.M., Ko, K.I., Jafari, A., and Wang, J.W. (2011). Presynaptic facilitation by

neuropeptide signaling mediates odor-driven food search. Cell *145*, 133-144.

Schlegel, P., Texada, M.J., Miroschnikow, A., Schoofs, A., Huckesfeld, S., Peters, M.,

Schneider-Mizell, C.M., Lacin, H., Li, F., Fetter, R.D.*, et al.* (2016). Synaptic transmission

parallels neuromodulation in a central food-intake circuit. Elife *5*.

Schoofs, A., Huckesfeld, S., Schlegel, P., Miroschnikow, A., Peters, M., Zeymer, M., Spiess,

R., Chiang, A.S., and Pankratz, M.J. (2014). Selection of motor programs for suppressing

food intake and inducing locomotion in the Drosophila brain. PLoS Biol *12*, e1001893.

Senapati, B., Tsao, C.H., Juan, Y.A., Chiu, T.H., Wu, C.L., Waddell, S., and Lin, S. (2019). A

neural mechanism for deprivation state-specific expression of relevant memories in

Drosophila. Nat Neurosci *22*, 2029-2039.

Shiraiwa, T., and Carlson, J.R. (2007). Proboscis extension response (PER) assay in

Drosophila. J Vis Exp, 193.

Song, T., Qin, W., Lai, Z., Li, H., Li, D., Wang, B., Deng, W., Wang, T., Wang, L., and Huang,

R. (2023). Dietary cysteine drives body fat loss via FMRFamide signaling in Drosophila and

mouse. Cell Res *33*, 434-447.

Stocker, R.F. (2004). Taste perception: Drosophila - a model of good taste. Curr Biol *14*,

R560-561.

Sun, J., Liu, C., Bai, X., Li, X., Li, J., Zhang, Z., Zhang, Y., Guo, J., and Li, Y. (2017).

Drosophila FIT is a protein-specific satiety hormone essential for feeding control. Nat

Commun *8*, 14161.

Sun, M., Ma, M., Deng, B., Li, N., Peng, Q., and Pan, Y. (2023). A neural pathway underlying

hunger modulation of sexual receptivity in Drosophila females. Cell Rep *42*, 113243.

Teranishi, H., and Hanada, R. (2021). Neuromedin U, a Key Molecule in Metabolic Disorders.

Int J Mol Sci *22*.

Ugrankar, R., Theodoropoulos, P., Akdemir, F., Henne, W.M., and Graff, J.M. (2018).

Circulating glucose levels inversely correlate with Drosophila larval feeding through insulin

signaling and SLC5A11. Commun Biol *1*, 110.

Vivot, K., Meszaros, G., Pangou, E., Zhang, Z., Qu, M., Erbs, E., Yeghiazaryan, G., Quinones,

M., Grandgirard, E., Schneider, A.*, et al.* (2023). CaMK1D signalling in AgRP neurons

promotes ghrelin-mediated food intake. Nat Metab *5*, 1045-1058.

Wang, M., Wu, J.X., Ding, D., and Chen, L. (2022). Structural insights into the mechanism of

pancreatic K(ATP) channel regulation by nucleotides. Nat Commun *13*, 2770.

Wei, Q., Krolewski, D.M., Moore, S., Kumar, V., Li, F., Martin, B., Tomer, R., Murphy, G.G.,

Deisseroth, K., Watson, S.J., Jr.*, et al.* (2018). Uneven balance of power between

hypothalamic peptidergic neurons in the control of feeding. Proc Natl Acad Sci U S A *115*,

E9489-E9498.

Willyard, C. (2008). Hungry for sleep. Nat Med *14*, 477-480.

Yao, Z., and Scott, K. (2022). Serotonergic neurons translate taste detection into internal

nutrient regulation. Neuron *110*, 1036-1050 e1037.

Yarmolinsky, D.A., Zuker, C.S., and Ryba, N.J. (2009). Common sense about taste: from

mammals to insects. Cell *139*, 234-244.

Yoon, N.A., and Diano, S. (2021). Hypothalamic glucose-sensing mechanisms. Diabetologia

64, 985-993.

Yu, Y., Huang, R., Ye, J., Zhang, V., Wu, C., Cheng, G., Jia, J., and Wang, L. (2016).

Regulation of starvation-induced hyperactivity by insulin and glucagon signaling in adult

Drosophila. Elife *5*.

Yurgel, M.E., Kakad, P., Zandawala, M., Nassel, D.R., Godenschwege, T.A., and Keene, A.C.

(2019). A single pair of leucokinin neurons are modulated by feeding state and regulate sleep-

metabolism interactions. PLoS Biol *17*, e2006409.

Zhao, Y., Khallaf, M.A., Johansson, E., Dzaki, N., Bhat, S., Alfredsson, J., Duan, J., Hansson,

B.S., Knaden, M., and Alenius, M. (2022). Hedgehog-mediated gut-taste neuron axis controls

sweet perception in Drosophila. Nat Commun *13*, 7810.

 Figure legends

Figure 1. hugin$^+$ neurons were a novel glucose sensor in the fly brain.

(**A**) Experimental procedure schematic. Newly eclosed flies were gathered and provided with

regular food for 5 days, followed by a 24-hour period of starvation. Proboscis Extension

Response (PER) assays were conducted. (**B**) Fraction of flies showing PER to different

concentrations of sucrose (n=4 groups, each of 10 flies). (**C-D**) Fraction of flies of the

indicated genotypes under 30 °C showing PER to 400 mM of sucrose (n=4-5 groups, each of

10 flies). (**E**) Fraction of indicated flies showing PER to different concentrations of sucrose

(ND represent standard fly food) (n=4-5 groups, each of 10 flies). (**F**) hugin expression in the

brain, illustrated by mCD8::GFP expression driven by *hugin^{GAL4}*. Scale bar represents 100

μm. (**G**) Representative traces and quantification of ex vivo calcium responses of hugin$^+$

neurons during the perfusion of glucose with or without TTX (n=6-7). Horizontal black bar

represents the duration indicated glucose solution stimulation. (**H**) AstA expression in the

brain, illustrated by mCD8::GFP expression driven by *AstA^{GAL4}*. Scale bar represents 100 μm.

(**I**) Representative traces and quantification of ex vivo calcium responses of AstA$^+$ neurons

during the perfusion of glucose with or without TTX (n=6). (**J**) Representative images of pre-

photoconversion (pre-PC) and post-photoconversion (post-PC) CaMPARI signal in hugin-

expressing neurons (upper). The Red:Green ratio represents intracellular Ca^{2+} concentrations

(lower). Scale bar represents 100 μm (n=18-23). ns, $P > 0.05$; *$P < 0.05$; **$P < 0.01$; ***$P <$

0.001; ****$P < 0.0001$. Student's *t-test*, one-way and two-way ANOVA followed by post hoc

test with Bonferroni correction were used for multiple comparisons when applicable.

Figure 2. Glucose activated hugin$^+$ neurons via Gltu1 and K$_{ATP}$ channel.

(**A**) Representative traces and quantification of ex vivo calcium responses of hugin$^+$ neurons

during the perfusion of glucose with or without phlorizin (1mM, n=6-7). Horizontal black bar

represents the duration indicated glucose solution stimulation. (**B**) Representative traces and

quantification of ex vivo calcium responses of hugin$^+$ neurons in indicated flies during the

perfusion of glucose. (**C**) Fraction of flies of the indicated genotypes showing PER to

different concentrations of sucrose (n=4 groups, each of 10 flies). (**D-E**) Representative traces

and quantification of ex vivo calcium responses of hugin$^+$ neurons during the perfusion of

glibenclamide (100 μM, D), glucose with alloxan (10 μM, E). (**F**) Representative traces and

quantification of ex vivo calcium responses of hugin$^+$ neurons in indicated flies during the

perfusion of glucose. (**G**) Representative traces and quantification of ex vivo calcium

responses of hugin$^+$ neurons during the perfusion of pyruvate (50 mM, n=6). (**H**)

Representative traces and quantification of ex vivo calcium responses of hugin$^+$ neurons

during the perfusion of D-glucose and L-glucose (50 mM, n=6) ns, $P > 0.05$; *$P < 0.05$; **P

< 0.01; ****$P < 0.0001$. Student's t-test and two-way ANOVA followed by post hoc test with

Bonferroni correction were used for multiple comparisons when applicable.

Figure 3. hugin⁺ neurons activated AstA⁺ neurons through PK2-R1.

(**A**) Fraction of flies showing PER to different concentrations of sucrose (n=4 groups, each of

10 flies). Flies were injected with saline or synthetic hugin for 30 minutes before the assay. (**B**)

Representative traces (upper) and quantification (lower) of peak calcium transients of Gr5a⁺

neurons in indicated flies upon 5% sucrose after injection of synthetic hugin (n=6). Horizontal

black bar represents the duration sucrose stimulation. (**C-D**) Fraction of flies of the indicated

genotypes showing PER to different concentrations of sucrose (n=4 groups, each of 10 flies).

(**E**) Co-localization (dashed box) of PK2-R1⁺ neurons (green) and AstA⁺ neurons (red) in the

SEZ region. Scale bar represents 100 μm. (**F**) Representative traces (upper) and quantification

(lower) of peak calcium transients of AstA⁺ neurons after the photo-activation of hugin⁺

neurons (n=6) from in vivo calcium imaging. Horizontal black bar represents the duration of

red-light stimulation. (**G-H**) Fraction of flies of the indicated genotypes showing PER to

different concentrations of sucrose (n=4 groups, each of 10 flies). ns, $P > 0.05$; *$P < 0.05$; **P

< 0.01; ***$P < 0.001$; ****$P < 0.0001$. Student's *t-test* and two-way ANOVA followed by

post hoc test with Bonferroni correction were used for multiple comparisons when applicable.

1008

Figure 4. AstA$^+$ neurons inhibited Gr5a$^+$ neurons through AstA-R1.

(**A**) Fraction of flies showing PER to different concentrations of sucrose (n=4 groups, each of

10 flies). Flies were injected with saline or synthetic AstA for 30 minutes before the assay. (**B**)

Representative traces (left) and quantification (right) of peak calcium transients of Gr5a$^+$

neurons in indicated flies upon 5% sucrose after injection of synthetic AstA (n=6). Horizontal

black bar represents the duration sucrose stimulation. (**C**) Localization of Gr5a$^+$ neurons

(green) and AstA$^+$ neurons (red) in the brain. Scale bar represents 100 μm. (**D**) Co-

localization of Gr5a$^+$ neurons (green) and AstA-R1$^+$ neurons (red). Arrows point to the

neurons co-expressing the two receptors. Scale bar represents 50 μm. (**E-F**) Fraction of flies

of the indicated genotypes showing PER to different concentrations of sucrose (n=4 groups,

each of 10 flies). (**G-H**) Representative traces (left) and quantification (right) of peak calcium

transients of Gr5a$^+$ neurons in indicated flies upon 5% sucrose during the process of photo-

activation of AstA$^+$ neurons (G, n=6) and hugin$^+$ neurons (H, n=7-8). Shadows represents the

duration of red-light stimulation. Horizontal black bar represents the duration of sucrose

stimulation. ns, $P > 0.05$; *$P < 0.05$; ***$P < 0.001$; ****$P < 0.0001$. Student's *t-test* and two-

way ANOVA followed by post hoc test with Bonferroni correction were used for multiple

comparisons when applicable.

1026

Figure 5. NMU$^+$ neurons were the central energy sensor in mouse.

(**A**) Blood NMU levels of starved mice re-fed with 20% sucrose (n=6). (**B**) Two-bottle preference tests for starved mice re-fed with sucrose (n=7). (**C**) Two-bottle preference tests for starved mice with or without intraperitoneal injection of NMU peptide (YFLFRPRN-NH$_2$, 4.5μm/kg) (n=7). (**D**) Two-bottle preference tests for indicated starved mice. (**E**) Fiber photometry to record the calcium dynamics of NMU$^+$ neurons in the VMH with GCaMP6m. (**F-G**) Representative trace (left) and heatmaps (right, n=5) showing calcium dynamics of NMU$^+$ neurons in the VMH in response to glucose perfusion. (**H**) Experimental approach to assess calcium signaling in NMU$^+$ neurons in the VMH in vitro. (**I-J**) Representative traces and quantification of ex vivo calcium responses of NMU$^+$ neurons during the perfusion of glucose with or without TTX (I), Alloxan (J) and Phlorizin (J) (n=6-7). Horizontal black bar represents the duration indicated glucose solution stimulation. (**K**) Anterograde tracing the downstream terget of NMU neurons. (**L**) Fiber photometry to record the calcium dynamics of Calb2$^+$ neurons in the rNST with GCaMP6m. (**M**) Representative traces and quantification of calcium responses of Calb2$^+$ neurons during glucose feeding with or without NMU injection. ns, $P > 0.05$; *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$; ****$P < 0.0001$. Student's *t-test* and one-way ANOVA followed by post hoc test with Bonferroni correction was used for multiple comparisons when applicable.

Figure supplementary 1. Satiety suppressed sweet sensitivity in a dopamine-independent

manner.

(**A**) Representative traces (left) and quantification (right) of peak calcium transients of Gr5a$^+$

neurons in indicated flies (n=6). Horizontal black bar represents the duration 5% sucrose

stimulation. (**B-D**) Fraction of flies of the indicated genotypes showing PER to different

concentrations of sucrose (B-C, n=4 groups, each of 10 flies). The Area Under the Curve

(AUC, D) represents sweet sensitivity in C. *P < 0.05; ****P < 0.0001. Student's *t-test* and

two-way ANOVA followed by post hoc test with Bonferroni correction were used for

multiple comparisons when applicable.

1055

Figure supplementary 2. The effect of satiety signals on taste modulation.

(**A**) Fraction of fed flies of the indicated genotypes and environmental temperatures showing

PER to different concentrations of sucrose (n=4 groups, each of 10 flies). (**B**) Fraction of

starved flies of the indicated genotypes and environmental temperatures showing PER to

different concentrations of sucrose (n=4-5 groups, each of 10 flies). ns, P > 0.05; *P < 0.05;

***P < 0.001; ****P < 0.0001. Two-way ANOVA followed by post hoc test with Bonferroni

correction was used for multiple comparisons when applicable.

1063

Figure supplementary 3. Activation of IPCs decreased the circulating sugar levels and

increased sweet sensitivity.

(**A**) Hemolymph glucose levels from the indicated genotypes and treatment temperatures

(n=8-9, 1 hour at 30 ºC). (**B-C**) Fraction of fed flies of the indicated genotypes and

environmental temperatures showing PER to different concentrations of sucrose (n=4 groups,

each of 10 flies). ns, P > 0.05; ****P < 0.0001. Student's *t-test* and two-way ANOVA

followed by post hoc test with Bonferroni correction were used for multiple comparisons

when applicable.

1072

Figure supplementary 4. Sugar intake suppressed sweet sensation.

(**A**) Fraction of indicated flies showing PER to different concentrations of fructose (n=4

groups, each of 10 flies). (**B**) Fraction of indicated flies showing PER to different

concentrations of trehalose (n=4 groups, each of 10 flies).****P < 0.0001. two-way ANOVA

followed by post hoc test with Bonferroni correction were used for multiple comparisons

when applicable.

1079

1080   **Figure supplementary 5. Starvation led to a decrease in the circulating sugar levels.**

1081   Hemolymph glucose levels from indicated flies (n=6-8). ***P < 0.001. Student's *t-test* was

1082   used for comparisons.

1083

Figure supplementary 6. The response of different neuropeptide-expressing neurons to

glucose.

(**A**) LK expression in the brain, illustrated by mCD8::GFP expression driven by LK^{GAL4}. Scale

bar represents 100 μm. (**B**) Representative traces and quantification of ex vivo calcium

responses of LK^+ neurons during the perfusion of glucose (n=6). (**C-D**) The distribution (C)

and calcium responses (D) of TK^+ neurons (n=6). (**E-F**) The distribution (E) and calcium

responses (F) of CCHa1[+] neurons (n=6). (**G-H**) The distribution (G) and calcium responses

(H) of CCHa2[+] neurons (n=6).

1092

1093 **Figure supplementary 7. The calcium response of hugin neurons upon glucose stimuli.**

1094 (**A**) Representative image of hugin neurons in SEZ. Scale bar represents 50 μm. (**B**) Calcium

1095 responses of different cluster of hugin neurons during the prefusion of glucose (n=6)..*P <

1096 0.05; ***P < 0.001. Student's *t-test* used for comparisons.

1097

1098 **Figure supplementary 8. hugin[+] or AstA[+] neurons suppressed sweet taste.**

1099 (**A-D**) Fraction of flies of the indicated genotypes and environmental temperatures showing

1100 PER to different concentrations of sucrose (n=4-5 groups, each of 10 flies). ns, P > 0.05; *P <

1101 0.05; ***P < 0.001; ****P < 0.0001. Two-way ANOVA followed by post hoc test with

1102 Bonferroni correction was used for multiple comparisons when applicable.

1103

1104 **Figure supplementary 9. The calcium response of AstA[+] neurons upon glucose treatment.**

1105 (**A**) Representative image of different part of AstA neurons .Scale bar represents 50 μm. (**B**)

1106 Calcium responses of different cluster of hugin neurons during the prefusion of glucose (n=6)

1107

1108 **Figure supplementary 10. Knockout of *PK2-R1* enhanced sweet sensation.**

1109 (**A-C**) Fraction of flies of the indicated genotypes showing PER to different concentrations of

1110 sucrose (n=4 groups, each of 10 flies). ***P < 0.001; ****P < 0.0001. Two-way ANOVA

1111 followed by post hoc test with Bonferroni correction were used for multiple comparisons

1112 when applicable.

1113

Figure supplementary 11. hugin$^+$ neurons in the brain activated AstA$^+$ neurons. (**A-B**)

Fraction of flies of the indicated genotypes and environmental temperatures showing PER to

different concentrations of sucrose (n=4 groups, each of 10 flies). Note connection between

the brain and VNC was cut off before the behavioral assay. (**C**) Representative traces (upper)

and quantification (lower) of peak calcium transients of AstA$^+$ neurons after the photo-

activation of hugin$^+$ neurons (n=6) from ex vivo calcium imaging. Horizontal black bar

represents the duration of red-light stimulation.

1121

1122 **Figure supplementary 12. Manipulating gene *AstA-R1* enhance PER.**

1123 (**A-C**) Fraction of flies of the indicated genotypes showing PER to different concentrations of

1124 sucrose (n=4 groups, each of 10 flies). ***$P < 0.001$; ****$P < 0.0001$. Two-way ANOVA

1125 followed by post hoc test with Bonferroni correction were used for multiple comparisons

1126 when applicable.

1127

Figure supplementary 13. hugin-AstA-Gr5a circuitry inhibited feeding behavior.

(**A**) The illustration of experimental design in Figure supplementary 13B-C. Note that all

flies were maintained at 20ºC before the feeding assay to prevent neuronal silencing by

Shibire^{ts1} during these procedures. (**B-C**) Food consumption of flies of the indicated

genotypes and environmental temperatures (n=8 groups, each of 4 flies). (**D**) Food

consumption of flies of the indicated genotypes (n=8 groups, each of 4 flies). (**E**) The TG

content of indicated genotypes (n=8 groups, each of 2 flies). (**F**) Food consumption of flies of

the indicated genotypes (n=8 groups, each of 4 flies). ns, P > 0.05; ***P < 0.001; ****P <

0.0001. Student's *t-test* and two-way ANOVA followed by post hoc test with Bonferroni

correction were used for multiple comparisons when applicable.

1138

1139 **Figure supplementary 14. NMU peptide suppressed sweet taste in fly.**

1140 Fraction of flies showing PER to different concentrations of sucrose (n=4 groups, each of 10

1141 flies). Flies were injected with saline or synthetic NMU for 30 mins before the assay.**P <

1142 0.01; ****P < 0.0001. Two-way ANOVA followed by post hoc test with Bonferroni

1143 correction were used for multiple comparisons when applicable.

1144

Figure supplementary 15. The expressing pattern of NMU neurons and Calb2

neurons(**A**) Co-labeling of NMU$^+$ neurons using a Cre-dependent mCherry reporter (red) and

in situ hybridization (green).Scale bar represents 200 μm. (**B**) NMU arbors was extended

near Calb2^{rNST} neurons. Scale bar represents 200 μm. (**C**) Co-labeling of Calb2$^+$ neurons with

a Cre-dependent mCherry reporter (red) and NMU antiboy staining (green). Scale bar

represents 200 μm.

1151

Figure supplementary 16. A working model

In the fly brain, a small group of neurons expressing hugin peptide is responsible for detecting

the circulating glucose levels. Following feeding, as the levels of glucose in the circulatory

system increased, these neurons are activated, leading to the release of hugin, which activated

AstA neurons via its receptor PK2-R1. Subsequently, the activation of AstA$^+$ neurons then

directly inhibited sweet perception via AstA peptide and its cognate receptor AstA-R1

expressed in sweet-sensing Gr5a$^+$ neurons. This neural circuitry is a novel energy sensor that

suppressed sweet perception and terminated feeding behavior, providing an efficient

mechanism for maintaining energy homeostasis.

1161

1162

1163

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure supplementary 1

Figure supplementary 2

Figure supplementary 3

A

B FED

C ST

Figure supplementary 4

Figure supplementary 5

Figure supplementary 6

Figure supplementary 7

A

B

Figure supplementary 8

Figure supplementary 9

A

B

Figure supplementary 10

A

B

C

Figure supplementary 11

A FED

B ST

- hugin > TrpA1
- + > TrpA1
- hugin > +

C

AstA-Gal4 > UAS-GCaMP6m
hugin-LexA > +

AstA-Gal4 > UAS-GCaMP6m
hugin-LexA > LexAop-CsChrimson

Figure supplementary 12

Figure supplementary 13

A

hugin/AstA-gal4
×
UAS-Shibiri^{ts1}

18 ℃ → ND 7 Days → 20 ℃ → Starved for 12 hr → 20 ℃

20 ℃ → 10-minute feeding (control)

30 ℃ → 10-minute feeding (experiment)

B
- hugin > Shibire^{ts1}
- + > Shibire^{ts1}
- hugin > +

Fold change — 20 ℃: ns, **; 30 ℃: ****, ****

C
- AstA > Shibire^{ts1}
- + > Shibire^{ts1}
- AstA > +

Fold change — 20 ℃: ****, ns; 30 ℃: ****, ****

D
- W1118
- PK2-R1 KO
- AstA-R1 KO

Fold change — ****, ***

E
- W1118
- PK2-R1 KO
- AstA-R1 KO

TG content (µg/fly) — ****, ****

F
- Gr5a > AstA-R1 i
- + > Gr5a
- AstA-R1 i > +

Fold change — ****, ****

A

B

Figure supplementary 15

A
AAV-DIO-mCherry injection — 3 weeks → Perfused Brain 4% PFA → FISH (NMU probe) → Antibody staining (mcherry antibody)

NMU-IRES-Cre mice

nmu
mCherry
VMH

merged | NMU | mCherry

B
AAV-EF1α-DIO-EGFP

VMH   rNST

NMU-IRES-Cre mice

merged | GFP | Calb2

C
AAV-DIO-mCherry injection — 3 weeks → Perfused Brain 4% PFA → Antibody staining (Calb2/mCherry antibody)

Calb2-IRES-Cre mice

Calb2
mCherry
rNST

merged | Calb2 | mCherry

Figure supplementary 16