Hugin-AstA circuitry is a novel central energy sensor that directly regulates sweet sensation in *Drosophila* and mouse

Wusa Qin[1,6], Tingting Song[2,3,6], Zeliang Lai[4], Daihan Li[3], Liming Wang[1,2,3,*], and Rui Huang[3,5,*]

[1]Institute of Molecular Physiology, Shenzhen Bay Laboratory, Shenzhen, 518055, China.

[2]School of Public Health, Capital Medical University, Beijing, 100069, China.

[3]Chinese Institutes for Medical Research, Beijing, 100069, China.

[4] Center for Neurointelligence, School of Medicine, Chongqing University, Chongqing, 400030, China.

[5]Guangyang Bay Laboratory, Chongqing Institute for Brain and Intelligence, Chongqing, 400060, China.

[6]These authors contribute equally to this work.

[*]For correspondence: lmwang83@ccmu.edu.cn; huangrui0716@sina.com

Abstract

Taste sensation plays a crucial role in shaping feeding behavior and is intricately influenced by internal states like hunger or satiety. Despite the identification of numerous neural substrates regulating feeding behavior, the central neural substrate that linked energy-sensing and taste sensation remained elusive. Here, we identified a novel neural circuitry that could directly sense internal energy state and modulate sweet sensation in the Drosophila brain. Specifically, a subset of neuropeptidergic neurons expressing hugin directly detected elevated levels of circulating glucose via glucose transporter Glut1 and ATP-sensitive potassium channel. Upon activation, these neurons released hugin peptide and activated downstream Allatostatin A (AstA)$^+$ neurons via its cognate receptor PK2-R1. Subsequently, the activation of AstA$^+$ neurons then directly inhibited sweet sensation via AstA peptide and its cognate receptor AstA-R1 expressed in sweet-sensing Gr5a$^+$ neurons. We also showed that neuromedin U (NMU), the mammalian homolog of fly hugin, served as an energy sensor to suppress sweet sensation. Therefore, these data identify hugin$^+$ neuron as a glucose-responsive central energy-sensing module that modulates sweet sensation across species.

Keywords: satiety, sweet sensation, feeding, hugin, AstA

Introduction

Animal behavior is tightly coupled to internal metabolic states[1]. Hunger profoundly alters feeding-related behaviors, enhancing sensitivity to appetitive cues and promoting food-seeking, whereas satiety actively suppresses these responses to shift behavioral priorities[2,3]. Such state-dependent modulation allows animals to dynamically allocate resources to maintain energy homeostasis[4]. While the neural mechanisms that drive feeding during hunger have been extensively characterized[5], the central circuits that sense satiety signals—specifically circulating glucose—to directly inhibit sensory processing remain less understood.

In both mammals and insects, "hunger signals" act as powerful accelerators for feeding. In the mammalian hypothalamus, AgRP neurons are activated by energy deficits to promote consumption[6,7], while in *Drosophila*, dopaminergic and Neuropeptide F (NPF) pathways increase the gain of sweet-sensing gustatory neurons to drive sugar intake[8-10]. These systems explain how animals ramp up feeding motivation when energy is low. However, to prevent overconsumption, the brain must also possess precise "braking" mechanisms that detect elevated energy states and rapidly dampen sensory sensitivity.

Dietary sugar is a primary satiety signal[11], yet the pathway linking central sugar sensing to behavioral inhibition remains elusive. In *Drosophila*, high-sugar diets are known to reduce sweet sensation via gut-derived signals or through central sensors like insulin-producing cells and DH44-expressing neurons that regulate general metabolic states[12-14]. Despite these advances, a critical gap remains: identifying a direct central sensor that detects acute rises in

circulating glucose and acts specifically to inhibit the peripheral sweet-sensing machinery.

Hugin-expressing neurons are promising candidates for this role, as they have been implicated

in satiety-related feeding regulation[15,16]. Furthermore, Allatostatin A (AstA) has been

proposed as a downstream signal capable of inhibiting sweet sensation[17]. However, whether

and how these components form a unified circuit to sense glucose and apply the "brake" on

feeding has not been established.

Here, we identify a complete neural circuitry that bridges this gap, directly linking internal

glucose levels to the modulation of sweet taste sensation. We show that a specific

subpopulation of hugin-expressing neurons functions as a central energy sensor, detecting

elevated circulating glucose via the glucose transporter Glut1 and ATP-sensitive potassium

channels. Upon activation, these neurons release hugin neuropeptide to activate downstream

AstA-expressing neurons via the PK2-R1 receptor. Subsequently, AstA neurons directly

inhibit sweet-sensing Gr5a$^+$ gustatory neurons through AstA–AstA-R1 signaling. This

pathway functions as a glucose-responsive inhibitory module whose endogenous activity is

elevated in fed flies, thereby contributing to the suppression of sweet sensitivity after feeding.

Finally, we demonstrate that Neuromedin U (NMU), the mammalian homolog of hugin[18],

similarly acts as a central energy sensor in mice to modulate sweet-responsive circuits,

revealing a conserved neural strategy for coupling metabolic state to sensory processing

across species.

Results

hugin[+] neurons were a novel glucose sensor in the fly brain

Flies extend their proboscis in response to appetitive cues, a behavioral element known as the proboscis extension reflex (PER) [20,51]. As previously observed, hunger significantly enhances this reflex[20]. In our study, we first replicated this phenomenon: In the state of hunger, flies exhibited elevated PER towards various concentrations of sugar compared to those under fed conditions (*Figure 1A-B*). These results confirmed the notion that sweet sensation was influenced by the internal energy state, elevated by starvation and suppressed by satiety [20,52]. Sweet sensation in flies is mediated by sweet-sensing Gr5a[+] neurons located on the proboscis [53]. Consistent with the behavioral effect, we also confirmed that Gr5a[+] neurons exhibited elevated calcium responses to sugar under starvation conditions (*Figure supplementary 1A*).

Dopaminergic neurons in the subesophageal zone (SEZ) region of the fly brain acted as a hunger sensor, promoting sweet sensation via a specific dopamine receptor DopEcR expressed in Gr5a[+] neurons [20,52]. However, although the gene knockout of tyrosine hydroxylase (TH) [54], a key enzyme for dopamine biosynthesis, significantly suppressed sweet sensation in starved fruit flies (*Figure supplementary 1B*), these flies still showed a further reduction in sugar-induced PER under sated conditions (*Figure supplementary 1C-D*). These results suggest the existence of dopamine-independent mechanisms for sensing satiety and hunger in fruit flies that exerts modulatory effect on sweet sensation.

In the terminating phase of feeding behavior, satiety signals play a crucial role to cease food

consumption and to ensure an appropriate amount of food ingestion. A number of

neuropeptidergic neurons were identified as putative satiety signals [36]. Notably, some of these

neurons, including those secreting neuropeptides LK, TK, hugin, AstA, CCHa1, and CCHa2,

are also distributed in the SEZ region [55]. It has been reported that the sweet-sensing gustatory

neurons primarily receive regulatory inputs from neurons in the SEZ region [20,51]. Therefore,

we hypothesized that these neuropeptidergic neurons might function as satiety signals,

imposing inhibititory effect on sweet sensation after adequate food ingestion.

To test this hypothesis, we selectively expressed the temperature-sensitive ion channel TrpA1

in these neurons and activated them at 30 °C, triggering the release of corresponding

neuropeptides. Indeed, activation of LK-, TK-, hugin-, AstA-, CCHa1-, and CCHa2-

expressing neurons robustly suppressed sweet taste sensitivity in both fed and starved flies

(*Figure 1C–D*; *also see Figure supplementary 2*). Moreover, LK-, TK-, and hugin-

expressing neurons exerted particularly strong inhibition under satiated conditions (*Figure*

1C–D; also see Figure supplementary 2). These results reveal a multilayered, state-

dependent neuropeptidergic network that finely tunes gustatory processing and feeding

behavior. In contrast, activating dopaminergic neurons enhanced sweet perception in sated

flies as expected (*Figure 1C*). In starved flies, activating dopaminergic neurons elicited no

change in PER likely due to a ceiling effect (*Figure 1D*).

In mammals and flies, circulating sugar levels are directly linked to satiety/hunger state [11,36,56].

We then asked whether changes in circulating sugar levels exerted an effect on sweet

sensation. Activating IPCs in the fly brain released *Drosophila* Insulin-like Peptides (DILPs),

the fly analog of mammalian insulin [57], into the hemolymph and reduced circulating sugar

levels (*Figure supplementary 3A*). As a result, such manipulation enhanced sweet sensitivity

of fed flies (*Figure supplementary 3B-C*). Therefore, it was plausible that changes in

circulating sugar might be the driver of changes in sweet sensation.

Indeed, allowing starved flies to re-feed on normal fly diet or sucrose alone for 15 minutes

both resulted in a significant decrease in sweet sensation (*Figure 1E*). Refeeding with

multiple caloric sugars similarly suppressed sweet sensation (*Figure supplementary 4A*),

indicating that energy intake contributes to the attenuation of sweet sensitivity. To examine

whether sweet taste experience itself also plays a role, we refed starved flies with arabinose, a

sweet but non-nutritive sugar. Arabinose feeding likewise led to a reduction in sweet

sensitivity (*Figure supplementary 4B*), suggesting that sweet taste experience can contribute

to the suppression of sweet perception, potentially through sensory adaptation. Consistent

with this interpretation, responses to other sweet compounds, including trehalose and fructose,

were also reduced following sucrose refeeding (*Figure supplementary 4C-D*). Conversely,

when starved flies were supplied with fly diet deprived of sucrose (but with normal levels of

polysaccharides, proteins, and lipids), there was no immediate change in sweet sensation after

re-feeding (*Figure 1E*). These results demonstrate that sugar intake inhibits sweet sensation,

probably via increasing circulating sugar levels.

Collectively, we thus hypothesized that the above neuropeptidergic neurons might directly

sense an increase of circulating sugar levels and suppress sweet sensation in response. If so,

these neurons should be activated by glucose, the main species of circulating sugars

associated with feeding behavior in flies [47], at the concentrations resembling sated state (>50

mM in fed flies (*Figure supplementary 5*)). We thus conducted calcium imaging experiments

in fly brains in an *ex vivo* preparation [11]. Among all the six groups of neuropeptidergic

neurons we examined in the SEZ, only hugin[+] neurons and AstA[+] neurons could be activated

by 50 mM glucose (*Figure 1F-I, also see Figure supplementary 6*). Remarkably, each of

these populations subdivided into anatomically distinct SEZ subclusters (*Figure

supplementary 7A; Figure supplementary 8A*), yet glucose□evoked Ca²□ responses

occurred exclusively within a single subcluster (*Figure 1H,I; Figure supplementary 7B;

Figure supplementary 8B*). This selective activation highlighted profound functional

heterogeneity in nutrient sensing among these neuron groups. When direct synaptic

transmission was blocked with TTX, hugin[+] neurons, but not AstA[+] neurons, still exhibited

calcium responses towards 50 mM of glucose (*Figure 1G, green*), suggesting that hugin[+]

neurons are a direct glucose sensor. Furthermore, the level of circulating glucose dropped to ~

20 mM in starved flies (*Figure supplementary 5*). We found that 20 mM of glucose did not

activate hugin[+] and AstA[+]neurons (*Figure 1G and I, orange*), suggesting that these neurons

can only be activated under sated conditions.

We also assessed the activity of hugin[+] neurons *in vivo* using calcium-modulated

photoactivatable ratiometric integrator (CaMPARI), a method employing a photoconvertible

fluorescent protein that allows the imaging of integrated calcium activity [58], under both sated

and starved conditions. In the CaMPARI assay, when neurons are activated, leading to an

increase in free calcium ions, exposure to 405-nm light induces a permanent green-to-red

photoconversion [58]. We found an increased ratio of red/green fluorescence in hugin+ neurons

in the sated state compared to that upon starvation (*Figure 1J*). These findings strongly imply

that hugin+ neurons function as a glucose-sensitive energy sensor whose activity tracks

internal metabolic state.

Glucose activated hugin+ neurons via an ATP-sensitive potassium channel

We then investigated the cellular mechanism through which glucose could directly activate

hugin+ neurons. In both mammals and flies, extracellular glucose is usually transported into

the cytosol to activate glucose-sensitive neurons [13,14,59]. When fly brains were pre-treated with

phlorizin, a blocker of glucose transport [13,14], hugin+ neurons showed diminished calcium

responses to glucose, suggesting activation of hugin+ neurons by glucose requires

intracellular glucose (*Figure 2A*).

The transport of glucose mediated by glucose transporter 1 (Glut1) can activate certain

neurons in the fruit fly brain [13,14]. By specifically knocking down Glut1 expression in hugin+

neurons, we observed a significant reduction in calcium responses to glucose (*Figure 2B*).

Consistently, these flies showed a slight yet significant increase in their PER responses to

glucose (*Figure 2C*). Again, these results further confirm that glucose needs to be transported

into hugin+ neurons to modulate neuronal activity and behavioral output.

In pancreatic β-cells and certain fly neurons [13,59,60], intracellular glucose modulates cell

excitability primarily through its effect on intracellular energy metabolism and ATP-sensitive

potassium channels (K_{ATP}). We pre-treated fly brains with a K_{ATP} inhibitor (glibenclamide)

and observed a significant calcium response in hugin$^+$ neurons (***Figure 2D***). Hexokinase, a

key enzyme in glucose metabolism that generates ATP, also plays a crucial role in this process.

When hexokinase activity was restricted by alloxan, glucose-induced calcium response in

hugin$^+$ neurons was significantly suppressed (***Figure 2E***). Similarly, when hexokinase was

specifically knocked down in hugin$^+$ neurons, a comparable reduction in glucose-induced

calcium responses was observed (***Figure 2F***). Pyruvate, the end product from glycolysis, can

lead to ATP generation via the tricarboxylic acid cycle. We also found that pyruvate

stimulation led to a significant calcium response in hugin$^+$ neurons (***Figure 2G***). As a

specificity control, nonmetabolizable L-glucose failed to activate hugin☐ neurons (***Figure***

2H).

Collectively, these findings suggest that extracellular glucose directly enters hugin$^+$ neurons

through specific glucose transporter and subsequently modulate the activity of K_{ATP} through

ATP production, ultimately leading to neuronal depolarization. We note that blocking glucose

transport or metabolism could, in principle, nonspecifically affect neuronal function, but the

rapid and selective nature of the observed responses argues against a generalized loss of

excitability.

AstA$^+$ neurons were direct downstream target of hugin$^+$ neurons

Since hugin[+] neurons were directly involved in glucose sensing and regulated sweet sensation

(*Figure 1G and Figure3A-B*), we sought to identify their downstream target. We then asked

whether the hugin peptide had a direct effect on sweet sensation. Upon microinjection of

synthetic hugin peptide, we observed a notable reduction in sweet sensation under both sated

and starved states (*Figure 3C*). In line with this observation, following the injection of hugin

peptide, we observed a significant reduction of sugar-induced calcium responses in sweet-

sensing Gr5a[+] neurons (*Figure 3D*). Thus, the inhibitory function of hugin[+] neurons on sweet

sensation was likely mediated by the secretion of hugin peptide.

There were two hugin receptors, PK2-R1 and PK2-R2, in fruit flies [61]. We knocked down

their expression by RNAi in the nervous system and then tested flies' PER to sugar. Knocking

down PK2-R1 but not PK2-R2 significantly enhanced sweet perception in both sated and

starved states (*Figure 3E-F*), suggesting a crucial role of PK2-R1 in receiving the input of

hugin[+] neurons in satiety sensing. To validate this finding, we further tested PK2-R1 null

mutants and observed a similar enhancement in sweet sensitivity (*Figure supplementary 9*).

Although PK2-R2 knockdown did not enhance PER, we observed a modest but reproducible

reduction in PER in starved flies (*Figure 3F*). This suggests potential functional

heterogeneity within hugin signaling pathways. Given the anatomical diversity of hugin

neurons, distinct subpopulations may engage different receptor-dependent mechanisms that

exert complementary or even opposing effects on feeding behavior. Further dissection of

PK2-R2-dependent signaling will be required to clarify this complexity.

236 Interestingly, we did not observe any expression of PK2-R1 in the proboscis where gustatory

237 sensory neurons were distributed, indicating that sweet-sensing Gr5a$^+$ neurons could not

238 directly receive input from hugin$^+$ neurons via PK2-R1 (*Figure supplementary 10*).AstA$^+$

239 neurons served as a plausible relay between hugin$^+$ neurons and sweet-sensing Gr5a$^+$ neurons

240 (*Figure 4A-B*), especially based on a clear co-expression of hugin receptor PK2-R1 and AstA

241 in the SEZ region of the fly brain (*Figure 4C*). Notably, these double-labeled neurons

242 corresponded precisely to the AstA subcluster that exhibited glucose□specific Ca²□

243 responses (*Figure 1H*). We also used optogenetic tools to further confirm that hugin$^+$ neurons

244 acted upstream of AstA$^+$ neurons. We expressed the light-sensitive neuronal activator

245 CsChrimson in hugin$^+$ neurons and calcium indicator GCaMP6m in AstA$^+$ neurons,

246 respectively [62]. As shown in Figure 4D, opto-activation of hugin$^+$ neurons led to a robust

247 calcium transient in AstA$^+$ neurons.

249 We then investigated whether the hugin-AstA circuitry affected sweet sensation. As expected,

250 knocking-down PK2-R1 (but not PK2-R2) in AstA$^+$ neurons led to a significant increase in

251 sugar-induced PER (*Figure 4E-F*). Therefore, AstA$^+$ neurons are the direct downstream

252 target of hugin$^+$ neurons in satiety sensing. hugin$^+$ neurons can directly sense elevated

253 circulating glucose levels, promote the release of neuropeptide hugin, and then activate AstA$^+$

254 neurons via its cognate receptor PK2-R1.

256 hugin$^+$ neurons are distributed in both the brain and the ventral nerve cord (VNC). The hugin$^+$

257 neurons in the VNC sense sugar and inhibit feeding behavior by suppressing the activity of

DH44 neurons [48]. To determine which hugin[+] neurons regulate the activity of AstA neurons

and sweet sensation, we performed the following experiments: First, we physically severed

the neural connection between the brain and the VNC. Under this condition, activation of

Hugin□ neurons still significantly reduced PER responses *(Figure supplementary 11A–B)*,

indicating that hugin□ neurons within the brain are sufficient to modulate sweet perception.

Second, in optogenetic experiments using isolated brain preparations, light stimulation of

hugin□ neurons elicited robust calcium responses in AstA□ neurons *(Figure supplementary*

11C), demonstrating a direct functional coupling between brain Hugin□ neurons and AstA□

neurons.

Notably, comparison of intact and severed preparations revealed that the magnitude of AstA□

neuronal activation was greater when VNC neurons were preserved, suggesting that VNC

hugin□ neurons may provide additional modulatory input that enhances AstA activation.

Together, these results indicate that Hugin□ neurons in the SEZ constitute a primary pathway

linking sugar sensing to AstA activation and PER regulation, while VNC hugin□ neurons

contribute to the overall strength of hugin signaling.

AstA[+] neurons directly inhibited Gr5a[+] neurons.

Activation of AstA[+] neurons suppressed sweet sensation, resembling the effect of activating

hugin[+] neurons [17]*(Figur 4A-B)*. We also evaluated the behavioral effect of AstA peptide.

Similar to hugin, the microinjection of AstA resulted in a reduction in sweet sensitivity in fruit

flies (*Figure 5A*) and the activity of Gr5a$^+$ neurons (*Figure 5B*). We then speculated whether

AstA$^+$ neurons and AstA peptide might function directly on Gr5a$^+$ neurons. The cell bodies of

sweet-sensing Gr5a$^+$ neurons are located in the proboscis, whereas their axons extend to the

SEZ region of the fly brain [63]. We observed distributions of the projections of AstA$^+$ neurons

and Gr5a$^+$ neurons both in the SEZ region (*Figure 5C*), suggesting that Gr5a$^+$ neurons might

directly receive AstA signal in this region.

AstA has two cognate receptors in fruit flies, AstA-R1 and AstA-R2 [64]. Co-labeling

experiments in the proboscis confirmed the expression of AstA-R1 in Gr5a$^+$ neurons (*Figure*

5D, arrows), while AstA-R2 was not expressed in the proboscis (*Figure supplementary 12*).

Functionally, knockdown of AstA-R1 specifically in Gr5a$^+$ neurons, but not that of AstA-R2,

significantly enhanced sweet sensation of flies (*Figure 5E-F*). Similar results were observed

in AstA-R1 knockout flies (*Figure supplementary 13*).

Using optogenetic tools, we also confirmed the function of hugin$^+$ neurons and AstA$^+$ neurons

on Gr5a$^+$ neurons. Activating hugin$^+$ neurons and AstA$^+$ neurons both significantly reduced

sugar-induced responses of Gr5a$^+$ neurons (*Figure 5G-H*). Collectively, these results suggest

that satiety-sensing hugin-AstA circuitry can directly modulate sweet sensation of flies via

suppressing the activity of Gr5a$^+$ gustatory neurons.

hugin–AstA circuitry suppressed food consumption.

Collectively, the hugin-AstA circuitry senses satiety signals and inhibits sweet sensation.

Considering the crucial role of sweet sensation in regulating feeding behavior, we

hypothesized that hugin-AstA circuitry was also involved in the regulation of food

consumption.

We expressed a temperature-sensitive blocker of synaptic transmission (Shibire[ts1]) in hugin[+]

neurons or AstA[+] neurons [65] and assayed the feeding behavior of flies under both permissive

temperature (*Figure6A, blue*) and non-permissive temperature (*Figur 6A, red*). In contrast to

the suppression of sweet sensitivity observed upon thermogenetic activation of these neurons,

silencing hugin□ neurons led to enhanced sweet-driven feeding behavior (*Figure 6B*).

Consistently, inhibiting synaptic transmission in either hugin□ or AstA□ neurons resulted in

a significant increase in food consumption at the non-permissive temperature compared to

controls (*Figure 6C–D*).

We also examined the role of hugin and AstA receptors in feeding behavior by testing *PK2-R1*

and *AstA-R1* gene knockout flies. The results showed that knockout of both receptors

enhanced feeding behavior (*Figure supplementary 14A*), consistent with increased sweet

perception upon genetic manipulation of these two receptors (*Figure 4-5*). Moreover, these

receptor knockout flies exhibited increased lipid storage, which was likely a consequence of

their enhanced feeding behavior (*Figure supplementary 14B*). Additionally, we knocked

down the expression of AstA-R1 in sweet sensing Gr5a[+] neurons, and found that these flies

exhibited a significant increase in food consumption (*Figure supplementary 14C*), also likely

due to the enhanced sweet sensation (*Figure 5E*).

Mammalian homolog of hugin was also a putative satiety sensor

Our present work identified hugin-AstA circuitry as a novel satiety sensor and feeding suppressor in fruit flies. Fruit flies and mammals share highly conserved regulatory elements on feeding behavior and energy homeostasis [51,66,67]. We thus asked whether this novel satiety-sensing mechanism was also conserved in the mammalian system,

In rodent models, neuromedin U (NMU), a analogues of hugin, was also found to be an appetite suppressor [15,18]. We found that synthetic NMU could suppress sweet perception when injected into flies' thorax (*Figure supplementary 15*), highlighting the analogy of fly hugin and mammalian NMU. In mouse models, we observed a significant increase in NMU levels in the blood following glucose feeding (*Figure 7A*). Using the sucrose preference test (SPT) to measure sweet preference, we noted that NMU injection reduced sugar preference in mice, phenocopying the effect of glucose feeding (*Figure 7B-C*). Conversely, NMU knockout enhanced sweet sensation in mice (*Figure 7D*). These data demonstrated that both NMU gene and protein plays a crucial role in regulation of sweet sensation.

Neuropeptide NMU is secreted by a group of neurons located in the VMH region of hypothalamus [18]. To investigate whether NMU^+ neurons functioned as an energy sensor in the mouse brain, we injected AAV-EF1α-DIO-GCaMP6m-WPRE into the VMH of NMU-Cre mice, specifically expressing GCaMP6m in NMU^+ neurons. We then monitored the dynamic activity of VMH^{NMU} neurons during glucose stimulation by using fiber photometry (*Figure 7E*). As expected, these neurons showed significant calcium responses following glucose

administration in mice (*Figure 7F-G*). To confirm the specificity of GCaMP6m expression in

NMU☐ neurons, we co-expressed a Cre-dependent mCherry reporter in the VMH of NMU-

Cre mice and performed fluorescence in situ hybridization for NMU mRNA. Colocalization

analysis demonstrated that GCaMP6m fluorescence was restricted exclusively to NMU☐

cells (*Figure supplementary 16A*).

Next, we conducted calcium imaging of brain slices *in vitro* to further assess these neurons'

ability to sense glucose (*Figure 7H*). Similar to hugin neurons in the fly brain, VMH^{NMU}

neurons in the mouse brain exhibited strong calcium responses to glucose stimulation, which

could not be inhibited by TTX (*Figure 7I*), indicating their direct glucose-sensing capability.

Also similar to flies, glucose activated these mouse neurons via entering the cells and

modulating K_{ATP} channels, as evidenced by the inhibition of calcium responses by alloxan and

phlorizin (*Figure 7J*). While supraphysiological glucose concentrations were used in slice

recordings to ensure reliable neuronal activation under ex vivo conditions, complementary in

vivo recordings demonstrate that VMH NMU☐ neurons are engaged by physiological sugar

ingestion.

Calbindin 2-positive neurons (Calb2) in the rostral nucleus of the solitary tract (rNST)

selectively respond to sweet tastants, facilitating sweet signal from the periphery into the

brain [68]. Because attenuated sweet sensation should correspondingly diminish central

sweet☐signal transmission, we asked whether NMU modulates this relay. To assess potential

anatomical connectivity between VMH NMU☐ neurons and rNST Calb2☐ neurons, we

employed a dual-virus tracing strategy. In NMU-Cre mice, we injected AAV9-EF1α-DIO-

EGFP into the VMH, which revealed dense GFP□ projections in close apposition to Calb2□

neurons in the rNST (*Figure supplementary 16B*). To further confirm this projection is

monosynaptic, we combined anterograde tracing by co-injecting AAV2/1-EF1α-DIO-EGFP

into the VMH and AAV2/1-hsyn-SV40-NLS-Cre into the rNST. This approach yielded GFP

expression in rNST Calb2□ neurons, supporting a direct VMH→rNST NMU□→Calb2□

synaptic link (*Figure 7K*).

To evaluate the functional consequence of NMU on Calb2□ neuronal activity during sweet

stimulation, we delivered AAVs encoding Cre-dependent GCaMP6m into the rNST of Calb2-

Cre mice, thereby enabling real-time calcium imaging in Calb2□ neurons (*Figure*

supplementary 16C).   Using fiber photometry, we recorded calcium responses to sugar

stimuli, both with and without NMU injection (*Figure 7L*). We found that NMU injection

elicited a significant, albeit partial, suppression of calcium responses to sugar *(Figure*

7M). Although the magnitude of this reduction was modest compared to the complete

silencing of activity, the effect was statistically robust and aligned with the behavioral

phenotypes, indicating that NMU acts to dampen sweet taste perception during feeding.

These findings indicate that NMU[+] neurons can indeed be a putative satiety sensor in the

mouse brain that suppresses sweet perception, shedding light on understanding the conserved

neural mechanism of feeding regulation in mammals including human.

Discussion

In this study, we identify a neural pathway that directly couples internal glucose levels to the

regulation of sweet taste perception and feeding behavior. In *Drosophila*, circulating glucose

activates a specific subset of hugin neurons, which signal to AstA neurons to suppress the

activity of peripheral sweet-sensing Gr5a neurons, thereby reducing sweet sensitivity and

terminating feeding. This pathway establishes a direct neural link between internal energy

state and early sensory processing (*Figure supplementary 17*).

A notable feature of this circuit is that its manipulation produces significant but relatively

modest changes in sweet sensitivity compared with the dramatic enhancement observed under

starvation[52,69]. This difference reflects the distinction between a specific satiety mechanism

and the global physiological state of hunger. Starvation is a systemic crisis that engages

multiple hunger-promoting pathways, including dopaminergic and NPF signaling, which

broadly elevate motivational drive and sensory gain[10,70]. In contrast, the hugin–AstA pathway

functions as a glucose-dependent inhibitory branch whose endogenous activity is elevated in

fed states but remains functionally capable of modulating sweet sensitivity across feeding

conditions.. Disrupting this inhibitory pathway prevents full suppression of sweet sensitivity

after feeding but does not recreate the potent hunger drive generated by dopaminergic

activation. Thus, energy homeostasis arises from the coordinated action between hunger

accelerators and satiety brakes.

This framework reconciles our findings with previous reports that silencing hugin neurons

does not alter proboscis extension in starved flies[20]. Although endogenous hugin neuronal

activity is reduced during starvation, our data show that further manipulation of this pathway

can still bidirectionally influence feeding behavior under starved conditions. This indicates

that the hugin–AstA circuitry is not exclusively restricted to fed states, but rather represents a

state-modulated inhibitory system whose baseline tone shifts with circulating glucose levels.

Under starvation, reduced endogenous activity may partially relieve this inhibitory influence;

however, the circuit retains functional capacity to modulate sweet sensitivity when

experimentally perturbed. In contrast, preventing the starvation-induced reduction of hugin

activity—such as by thermogenetic activation—significantly suppresses hunger-enhanced

sweet sensitivity. These results indicate that the dynamic downregulation of hugin neuron

activity is a critical component of the state-dependent modulation of taste. Therefore, rather

than functioning as a strictly "satiety-specific brake," the hugin–AstA axis is better

conceptualized as a glucose-responsive inhibitory module that biases sweet perception

according to metabolic state. Its activity increases following feeding to dampen sensory gain,

while decreased activity during starvation permits heightened sweet sensitivity. Such graded

modulation enables flexible integration of internal energy status with sensory processing.

Our data further reveal functional heterogeneity within peptidergic populations. Although

hugin and AstA neurons are broadly distributed in the central nervous system (including the

subesophageal zone and ventral nerve cord), only specific subpopulations respond to elevated

glucose[43,48]. While descending inputs from the VNC may potentiate signaling, our data

suggest that the glucose-responsive hugin neurons in the SEZ constitute a dedicated energy-

sensing module. This heterogeneity likely explains both the partial behavioral effects of

global manipulations and the ability of hugin neurons to regulate diverse behaviors beyond

feeding, such as locomotion.

Circulating sugars represent a complex internal signal. In *Drosophila*, trehalose serves as the

major circulating sugar supporting stress resistance, whereas glucose functions as the primary,

rapidly mobilized energy substrate[71]. Unlike receptor-based sensors (e.g., Gr43a for

fructose)[72], Hugin neurons rely on intracellular glucose metabolism and K^{ATP} channel activity

to modulate neuronal excitability. This mechanism ensures that the circuit activity is tightly

coupled to cellular energy status. Consistent with this, feeding with fructose, trehalose, or

non-nutritive sweeteners only weakly suppressed sweet sensitivity within the acute post-

feeding interval, whereas sucrose—which rapidly restores hemolymph glucose—robustly

activated this pathway. Thus, our work identifies glucose as the key metabolic signal that

engages this central brake.

The presence of multiple sugar-sensing systems in *Drosophila* reflects functional

specialization rather than redundancy. Peripheral and gut-derived signals regulate feeding

over longer timescales, while central hunger circuits adjust sensory gain during deprivation.

The hugin–AstA circuit adds a distinct layer by acting as a rapid central brake that prevents

overconsumption when energy stores are sufficient.

Strikingly, we find that this circuit logic is conserved in mammals. Although NMU has been

implicated in feeding suppression[73,74], whether NMU neurons sense internal nutrients directly

had remained unclear. Our data demonstrate that VMH NMU☐ neurons are activated by

elevations in internal glucose and form a functional circuit with sweet-responsive Calb2

neurons in the rostral nucleus of the solitary tract (rNST). Gastric glucose infusion

experiments established that NMU neurons respond to internal metabolic signals

independently of oral sensory input, while physiological sugar ingestion confirmed that this

pathway actively suppresses downstream sweet-driven activity. Together, these findings

define a previously unrecognized NMU-centered circuit linking internal glucose sensing to

the modulation of early taste processing. All mouse experiments in this study were performed

in male animals; whether sex-specific differences exist in NMU-mediated glucose sensing

and taste modulation remains an important question for future investigation.

The conservation of this "satiety brake" suggests that suppressing sensory gain via central

neuropeptidergic energy sensors is a fundamental strategy for maintaining energy balance. By

defining a direct link between internal glucose levels and sweet taste perception, our work

provides a mechanistic framework for understanding how metabolic state shapes sensory

experience and feeding behavior.

 Materials and methods

Animals

Fly: Flies were maintained on a standard fly diet under conditions of 60% humidity and 25 °C

with a 12-hour light and 12-hour dark cycle. Virgin female flies were chosen immediately

after emerging and housed in standard fly food (ND) vials, with 20 flies per vial, for a period

of 5-6 days before experiments. For temperature☐sensitive manipulations (dTrpA1, Shibire[ts]),

flies were reared at 18 °C for 7–8 days, then transferred to 30 °C for 30 min to activate or

inhibit targeted neurons; behavioral assays (PER or feeding) were conducted continuously at

30 °C. For starvation, the flies were kept on 2 % Agar for 12-hour. For refeeding, the fasted

flies were transfer to different food for 15 mins.

The following UAS-RNAi lines: *UAS-AstA-R1 RNAi (#27280)*, *UAS-AstA-R2 RNAi (#25935)*,

UAS-PK2-R1 RNAi (#29624), *UAS-PK2-R2 RNAi (#28781)*, *UAS-CD8-GFP, LexAop-CD2-*

RFP (#67093), *UAS-GCaMP6m (#42748)*, *UAS-CaMPARI (#58764)*, *Gr5a-GAL4 (#57592)*,

Gr5a-LexA (#93014), *ilp2-GAL4 (#37516)*, *elav-GAL4 (#25750)* were obtained from the

Bloomington Drosophila Stock Center at Indiana University. *TH-KO, PK2-R1-KO, AstA-R1-*

KO and all the other Gal4 and LexA lines (*AstA-Gal4, LK-Gal4, hugin-Gal4, TK-Gal4,*

CCHa1-Gal4, CCHa2-Gal4, TH-Gal4, PK2-R1-Gal4, AstA-LexA, AstA-R1-LexA) were from

Yi Rao (Deng et al., 2019) (Capital Medical University, Beijing). *UAS-dTrpA1* and *UAS-*

Shibire[ts1] were from David Anderson (Caltech). *UAS-GCaMP6m; LexAop-CsChrimson* was

from Yufeng Pan (Southeast University, Nanjing). *LexAop-Chrimson* was from Wei Zhang

(Tsinghua University, Beijing).

Mouse: Male C57BL/6J mice at the age of four weeks were procured from cyagen. The mice

were kept in standard laboratory conditions at a temperature of 25 □ with a 12:12 light/dark

cycle. All experimental procedures were conducted in compliance with the guidelines set by

the Laboratory Animal Welfare and Ethics Committee of Chongqing University (CQU-

IACUC-RE-202401-004), adhering to both national and international standards.

Behavioral assays (Fly)

PER was evaluated using a method outlined in a previous study{Marella, 2012 #22}. Briefly,

individual flies were gently aspirated and positioned in a 200 µL pipette tip, allowing only the

head and proboscis to protrude while the dorsal thorax was lightly affixed to prevent escape.

After a 3-minute acclimation period, each fly was first presented with 0.5 µL water droplets

on the labellum twice. Flies that responded by extending their proboscis were allowed to

drink and were subsequently excluded from the experiment. Flies that did not extend their

proboscis in response to water were then stimulated sequentially with ascending sucrose

concentrations (6.25−800 mM). Each concentration was delivered as two brief (<1 s) touches

of a 0.5 µL droplet to the labellum, immediately withdrawn to prevent ingestion. A full

proboscis extension, defined as the complete uncoiling of the rostrum beyond the labellum

plane, was scored as positive. Partial or delayed (>1 s) extensions were not considered

positive responses. Responsiveness at each concentration was recorded if at least one full

extension occurred in the two trials.For Figure supplementary 11, the connection between the

brain and VNC were cut off by a dissecting scissors before the PER assay.

Feeding was assessed as previously described [78]. Briefly, 5-day-old flies underwent a 12-hour

starvation period on 2% agar and were then moved to a new vial with test food containing 0.5%

Brilliant Blue (MACKLIN, China) for 10 minutes. After a rapid freeze at -20 °C, flies were

decapitated in PBS, homogenized, and centrifuged (13,000 g * 5 min). The resulting

supernatants were diluted with PBS to a total volume of 1 ml, and absorbance was measured

at 620 nm.

Sucrose Preference Test

The two-bottle preference (TBP) behavioral test was conducted in a standard mouse cage.

Two bottles, one containing water and the other with a 10 mM sucrose solution, were placed

on top of the cage. Each tested mouse was individually caged and adopted to two bottles

containing distilled water or a 10 mM sucrose solution for one week. Throughout the training

period, the positions of the bottles were alternated every 24 hours to mitigate potential

position-related biases. Prior to the TBP tests for sucrose, mice underwent a 12-hour pre-

starvation period, followed by re-fed sucrose (20%) for 1 hour or intraperitoneal (IP)

administration of NMU (YFLFRPRN-NH$_2$, 4.5 µm/kg). The consumption of water and

sucrose solution over a 2-hour period was meticulously recorded. The sweetener preference

ratio was then calculated by dividing the weight of sweetener consumed by the weight of

water consumed.

NMU measurement

Blood was obtained from the lower jaw of the mice and clotted at room temperature for 1

hour. The samples were then centrifuged (1,500g, 4 °C) for 10 minutes and the supernatants

were collected for NMU measurement. All the manipulations were according to the

manufacturer's instructions (BMASSY,74030).

Hemolymph extraction and glucose measurement

Groups of 40 ice☐anesthetized flies were decapitated and placed into a perforated 0.5 ml tube,

which was nested inside a 1.5 ml collection tube. Samples were centrifuged at $2,500 \times g$ for

10 min at 4 °C, yielding approximately 2 µl of hemolymph per batch. Hemolymph was

pooled to 4 µl per assay (n = 8 independent samples per group) and immediately analyzed for

glucose concentration using the Solarbio BC2500 kit (Solarbio, China) according to the

manufacturer's instructions.

Microinjection

Flies were delicately positioned and secured within a 200 ml pipette tip, ensuring their heads

were directed toward the tip's end. Subsequently, the tip was opened by cutting, exposing the

heads and a section of the thorax. About 20 nl of sterile saline, either with or without

synthesized peptides (hugin: SVPFKPRL-NH$_2$, 1 mg/ml (1.1 mM); AstA: LPVYNFGL-NH$_2$,

1mg/ml (1.1 mM); NMU: YFLFRPRN-NH$_2$, 1mg/ml (0.9 mM); Control peptide:

DYKDDDDKYPYDVPDYA, 1mg/ml (0.48 mM), were injected into the thorax of these flies

using a glass micropipette and a microinjector (3-000-207, Drummond Scientific Company

Instruments). The glass micropipette was created from thick-walled borosilicate capillaries (3-

000-203-G/X, Drummond Scientific Company Instruments).

Immunofluorescence staining

Fly brains were carefully dissected in PBS on ice and then fixed in 4% formaldehyde for 60

minutes. Following fixation, the brains underwent permeabilization and blocking using

Dilution/Blocking Buffer (PBS containing 10% Calf Serum and 0.5% Triton X-100) for 2

hours at room temperature. Subsequently, the samples were immersed in the appropriate

primary antibodies in Dilution/Blocking Buffer for 24 hours at 4 °C. Afterward, the samples

were subjected to a 60-minute wash with Washing Buffer (0.5% Triton X-100 in PBS) four

times at room temperature and were then incubated with secondary antibodies for 24 hours at

4 °C. The samples underwent three additional washes with Washing Buffer before being

mounted in Fluoroshield (Sigma-Aldrich).

Images were acquired using a scanning confocal microscope with Olympus objectives (20×

/0.7 and 40× /0.95w). Antibodies were employed at the following dilutions: mouse anti-nc82

(1:100, DSHB), rabbit anti-GFP (1:500, Abcam), Alexa Fluor 647 goat anti-mouse (1:500,

Invitrogen), and Alexa Fluor 488 goat anti-rabbit (1:500, Invitrogen).

In situ hybridization

Mice were anesthetized and perfused with 4% paraformaldehyde (PFA) dissolved in

phosphate-buffered saline (PBS, pH 7.4). Brains were carefully removed from the skull and

post-fixed in 4% PFA at 4 °C overnight. Subsequently, the brains were transferred to a 20%

sucrose solution (in $1 \times$ PBS) at 4 ˚C until they were ready for sectioning. Twenty-

micrometer-thick sections were obtained using a cryostat. The detection of nmu was

performed using a fluorescent in situ hybridization technique (RNAscope, Pinpoease)

according to the manufacturer᾽s instructions. Sections were mounted on SuperFrost Plus Gold

slides (ThermoFisher) and briefly rinsed in autoclaved Millipore water. They were then

subjected to gradient dehydration in 50%, 75%, and 100% ethanol for 5 minutes each. A

hydrophobic barrier was created around the sections using an ImmEdge hydrophobic barrier

pen (Cat No. 310018). All incubation steps were carried out at 40 ˚C using the Pinpoease

hybridization system (Cat No. SH-08). The subsequent hybridization, amplification, and

detection steps were performed strictly according to the manufacturer᾽s instructions.

Stereotaxic brain surgeries and viral injection

Mice were anesthetized with 1-2% isoflurane during stereotaxic injections, performed using a

small animal stereotaxic instrument (RWD Life Science, #68030). Throughout the surgery,

core body temperature was maintained at 36 ± 1 °C using a feedback-controlled heating

system. Micro scissors were used for the incision, and a dental drill was employed to create

the cranial window. Viral injections were delivered using a glass microelectrode syringe

pump (RWD Life Science, #R-480). To express GCaMP6m and mCherry AAV2/9-EF1α-

DIO-GCaMP6m-WPRE (titer: 2.89×10^{12}, #H4955, Obio Biotechnology, Shanghai, China,

Co., Ltd.) and AAV2/9-EF1α-DIO-mCherry-WPRE (titer: 2.89×10^{12}, Obio Biotechnology,

Shanghai, China, Co., Ltd.) were bilaterally injected into the VMH and rNST, with

approximately 0.15 µL of virus delivered per site. The microelectrode was left in place for 10

minutes post-injection before being withdrawn. Injection coordinates were as follows: VMH

(AP, −1.46 mm; ML, ±0.4 mm; DV, −5.5 mm) and rNST (AP, −7.08 mm; ML, ±0.6 mm; DV,

−4.3 mm). After 3 weeks of AAV injections, optical fibers were chronically implanted in the

VMH or rNST and secured with dental cement (C&B Metabond®, Parkell, Japan). Injection

coordinates were determined based on the 4th edition of the mouse brain atlas by Franklin and

Paxinos (2008). Mice were allowed to recover for 5-7 days post fiber-implanted before any

behavioral or calcium imaging evaluations were conducted. After the experiments, mice were

perfused to verify virus expression and fiber placement. Data from mice with poor viral

expression or incorrect fiber placement (0-20% of cases) were excluded from the final

analysis.

For the anterograde tracing experiments, two viral vectors were used. AAV2/1-hsyn-DIO-

EGFP-WPRE (titer: 1×10^{13} viral particles/mL,Obio Biotechnology) was injected into the

ventromedial hypothalamus (VMH), and AAV2/1-hsyn-SV40 NLS-Cre (titer: 2.1×10^{12}

viral particles/mL, Brain Case) was injected into the rostral nucleus of the solitary tract

(rNST). Approximately 0.15 µL of virus was delivered per injection site. After 3 weeks, the

brains were perfused and fixed. Subsequently, the brain tissues were sliced and subjected to

antibody staining. The fluorescence signals were then recorded and analyzed.

Optical-fiber-based calcium recording in freely mice

Following injection of an AAV2/9-EF1α-DIO-GCaMP6m-WPRE viral vector, an optical

fiber (200 µm O.D., 0.37 numerical aperture, Inper Inc., China) was placed 150 µm above

the viral injection site. GCaMP6m$^+$ mice were implanted with the optic fiber 3 weeks post-

AAV injection and allowed to recover for 5–7 days prior to experimental testing.

For Figure 5K-L, six mice were first trained to lick glucose (500 mM) from a petri dish. After

training, they were fasted for 12 hours and then received an intraperitoneal injection (ip) of

synthetic NMU peptide (4.5 µm/kg), followed by a 30-minute waiting period. During the

experimental session, the mice were given 10 seconds to lick glucose from the petri dish

while undergoing fiber photometry recording. For Figure 5E-G, mice were fasted for 4 hours

and then administered glucose via gastric infusion, with fiber photometry recordings

capturing the process. Fluorescence signals were acquired using a dual-channel fiber

photometry system (410 nm & 470 nm, RWD Life Science). The laser power at the tip of the

optical fiber was kept below 20 µW for both the 470 nm and 410 nm channels to minimize

bleaching. $\Delta F / F$ was calculated as (470 nm signal - fitted 410 nm signal) / fitted 410 nm

signal. Data analysis was performed using software from RWD Life Science.

Ex vivo calcium imaging

For Figure 1G, 1I, figure supplementary 6, and Figure 2freshly dissected fly brains were

placed in the sugar-free AHL buffer (108 mM NaCl, 8.2 mM $MgCl_2$, 4 mM $NaHCO_3$, 1 mM

NaH_2PO_4, 5 mM KCl, 2 mM $CaCl_2$, 5 mM HEPES, pH 7.3). Baseline recordings of the

samples in AHL buffer were taken over 1 minutes. Subsequently, the solutions were added

with 100 µM glibenclamide (*Figure 2D*), 50 mM pyruvate (*Figure 2G*) or switched to AHL

+ glucose (50 mM) with or without indicated chemicals (*Figure 2 A and E*, 1 mM phlorizin

or 10 um alloxan), and the pH was adjusted back to 7.3 through gentle perfusion for 3 minutes.

Solution flow in the perfusion chamber was controlled by a valve commander (Scientific

Instruments). Following stimulation, samples were washed out again with AHL. For the TTX

test, 2 μM TTX was added to the AHL solution. Prior to the assay, the samples were pre-

treated with 2 mM TTX for 15 minutes.

For Figure 7H-J, mice (6-9 weeks of age) were deeply anesthetized by avertin (1.25% avertin,

0.2 ml/10g) . Mouse brains were quickly extracted following decapitation and immediately

placed in ice-cold slicing buffer (110 mM Choline Cl, 2.5 mM KCl, 1 mM NaH_2PO_4, 25 mM

$NaHCO_3$, 5 mM glucose, 7 mM $MgCl_2$, 0.5 mM $CaCl_2$,1.3mM Na ascorbate,0.6mM Na

pyruvate bubbled with 95% oxygen and 5% CO_2 with an adjusted pH of 7.3). The brains were

then sectioned into 200 μm slices using a vibratome (Leica VT1200S). These slices were

incubated artificial cerebrospinal fluid (aCSF) (125 mM NaCl, 2.5 mM KCl, 1 mM NaH_2PO_4,

25 mM $NaHCO_3$, 5 mM glucose, 1.3 mM $MgCl_2$, 2 mM $CaCl_2$,1.3mM Na ascorbate, 0.6 mM

Na pyruvate bubbled with 95% oxygen and 5% CO_2 with an adjusted pH of 7.3) at 34 °C for

20 minutes, then transferred to a room temperature for 30 minutes before recording. For Ca^{2+}

imaging, the slices were placed in a recording chamber in the low sugar aCSF (125 mM NaCl,

2.5 mM KCl, 1 mM NaH_2PO_4, 25 mM $NaHCO_3$, 1 mM glucose, 1.3 mM $MgCl_2$, 2 mM $CaCl_2$,

1.3 mM Na ascorbate, 0.6mM Na pyruvate, pH 7.3) for 10 minutes (maintained the flow rate

for 1-3 ml/ minute). Baseline recordings, perfusion, and Ca^{2+} recording were performed as

detailed above.

All imaging was conducted on an Olympus confocal microscope (FVMPE-RS) with a water

immersion objective lens (25× /1.05w MP). Image analyses were performed using ImageJ to

calculate the mean intensity of the indicated neuron targeting ROIs, and then plotted in Excel

(Microsoft). Ratio changes were calculated using the formula: $\Delta F / F = [F - F_0] / F_0$, where F

represents the mean fluorescence of the cell body, and F_0 is the average baseline (before

stimulation).

Optogenetics and in vivo calcium imaging

Newly emerged virgin female flies were housed in a fresh vial with standard medium for 3

days and subsequently moved to a vial containing regular food supplemented with 400 µM

all-trans-retinal (Sigma R2500) for 2-4 days prior to experiments. These flies were then

immobilized on ice, affixed to transparent tape, and the dorsal cuticle of the fly head was

delicately removed with forceps to expose the brain, which was immersed in AHL.

For Figure 3F, an array of red LED (Thorlabs, M625L3) was positioned above the brain.

Calcium signals of Gr5a$^+$ neurons were captured for 60 frames (1 second per frame) in the

absence of light to establish a baseline. Subsequently, the next 20 frames were recorded

during opto-activation with the light (the light was switched on for 0.45 second and rested for

0.55 second per activation cycle, and the images were collected in rest time), followed by

another 60 frames recorded without light afterward.

For Figure 4G and 4H, to apply liquid food (5% sucrose) to the proboscis, a pipette was filled

with the taste solution and placed a few microns from the proboscis tip by using a

micromanipulator (MP225, Sutter Instruments) prior to recording. Then, we recorded calcium

signals for 60 seconds in the absence of light to establish a baseline (1 second per frame). To

activate neurons expressing CsChrimson, a red LED was positioned above the brain,

controlled by a custom computer program during opto-activation (0.45 second on and 0.55

second off from 60 second to 120 second). The pipette was placed on the proboscis from 80

second to 85 second. Calcium signals of Gr5a$^+$ neurons were recorded throughout these

processes (1 second per frame).

Image analyses about relative fluorescence changes $\Delta F / F$ were performed as ex vivo calcium

imaging.

Optogenetics and ex vivo calcium imaging

For Figure supplementary 11C, newly emerged virgin female flies were housed in a fresh vial

with standard medium for 3 days and subsequently moved to a vial containing regular food

supplemented with 400 µM all-trans-retinal (Sigma R2500) for 2-4 days prior to experiments.

The brains were dissected and placed under a red LED. Calcium signals of AstA$^+$ neurons

were recorded with or without LED light as above. Image analyses were performed as ex vivo

calcium imaging.

Calcium imaging with CaMPARI

The flies were then gently immobilized on ice, attached to transparent tape, and the dorsal

cuticle of the fly head was carefully removed with forceps to expose the brain, which was

immersed in AHL. Photoconversion (PC) was achieved for 30 seconds using a 405 nm LED

[Thorlabs, M405L2–UV (405 nm) Mounted LED, 1,000 mA, 410 mW], controlled by a LED

controller (Thorlabs, LEDDB1 driven with 1000 mA). Images were captured using an

Olympus confocal microscope. Image analyses were performed in ImageJ by manually

drawing ROIs covering individual neuronal cell bodies using the green channel. The

extent of CaMPARI photoconversion was determined as the Red: Green ratio.A '-405 nm'

control was included to demonstrate that scanning of the hugin neurons without exposure to

ultraviolet light does not convert green to red fluorescence.

Statistical analysis

Data are shown as means (± SEM). Data presented in this study were verified for normal

distribution by D'Agostino–Pearson omnibus test. Student's t test, one-way ANOVA and

two-way ANOVA (for comparisons among three or more groups and comparisons with more

than one variant) were used. The post hoc test with Bonferroni correction was performed for

multiple comparisons following ANOVA.

Acknowledgements

We express gratitude to all members of the Wang and Huang Lab for their valuable

discussions and technical support. Special thanks to Yi Rao and Bowen Deng for generously

providing *Drosophila* chemoconnectome lines, and to Yulong Li and Jianzhi Zeng for their

assistance with *in vivo* calcium imaging preparations. Appreciation is extended to the

members of the Neuroscience Pioneer Club for their insightful discussions throughout the

course of this study. This study was funded by the National Natural Science Foundation of

Author contributions

T.S., R.H., and L.W. designed the experiments and wrote the manuscript. W.Q., and T.S.

conducted and analyzed all the experiments. L.W. and R.H. supervised the project.

Competing interests

The authors declare no competing interest.

References

1.   Ritterhoff, J., and Tian, R. (2023). Metabolic mechanisms in physiological and

753   pathological cardiac hypertrophy: new paradigms and challenges. Nat Rev Cardiol *20*,

754   812–829. 10.1038/s41569-023-00887-x.

755   2.   Smith, N.K., Plotkin, J.M., and Grueter, B.A. (2022). Hunger dampens a nucleus

756   accumbens circuit to drive persistent food seeking. Curr Biol *32*, 1689–1702 e1684.

757   10.1016/j.cub.2022.02.034.

758   3.   Yu, Y., Huang, R., Ye, J., Zhang, V., Wu, C., Cheng, G., Jia, J., and Wang, L. (2016).

759   Regulation of starvation-induced hyperactivity by insulin and glucagon signaling in

760   adult Drosophila. Elife *5*. 10.7554/eLife.15693.

761   4.   Roh, E., and Kim, M.S. (2016). Brain Regulation of Energy Metabolism. Endocrinol

762   Metab (Seoul) *31*, 519–524. 10.3803/EnM.2016.31.4.519.

763   5.   Tan, H.L., Yin, L., Tan, Y., Ivanov, J., Plucinska, K., Ilanges, A., Herb, B.R., Wang, P.,

764   Kosse, C., Cohen, P., et al. (2024). Leptin-activated hypothalamic BNC2 neurons

765   acutely suppress food intake. Nature *636*, 198–205. 10.1038/s41586-024-08108-2.

766   6.   De Solis, A.J., Del Rio-Martin, A., Radermacher, J., Chen, W., Steuernagel, L.,

767   Bauder, C.A., Eggersmann, F.R., Morgan, D.A., Cremer, A.L., Sue, M., et al. (2024).

768   Reciprocal activity of AgRP and POMC neurons governs coordinated control of

769   feeding and metabolism. Nat Metab *6*, 473–493. 10.1038/s42255-024-00987-z.

770   7.   Sayar-Atasoy, N., Aklan, I., Yavuz, Y., Laule, C., Kim, H., Rysted, J., Alp, M.I., Davis,

771   D., Yilmaz, B., and Atasoy, D. (2024). AgRP neurons encode circadian feeding time.

772   Nat Neurosci *27*, 102–115. 10.1038/s41593-023-01482-6.

773   8.   Chung, B.Y., Ro, J., Hutter, S.A., Miller, K.M., Guduguntla, L.S., Kondo, S., and

774   Pletcher, S.D. (2017). Drosophila Neuropeptide F Signaling Independently Regulates

775   Feeding   and   Sleep-Wake   Behavior.   Cell   Rep   *19*,   2441   –   2450.

776   10.1016/j.celrep.2017.05.085.

777   9.   Shohat-Ophir, G., Kaun, K.R., Azanchi, R., Mohammed, H., and Heberlein, U. (2012).

778   Sexual deprivation increases ethanol intake in Drosophila. Science *335*, 1351–1355.

779   10.1126/science.1215932.

780   10.   Inagaki, H.K., Ben-Tabou de-Leon, S., Wong, A.M., Jagadish, S., Ishimoto, H.,

781   Barnea, G., Kitamoto, T., Axel, R., and Anderson, D.J. (2012). Visualizing

782   neuromodulation in vivo: TANGO-mapping of dopamine signaling reveals appetite

783   control of sugar sensing. Cell *148*, 583–595. 10.1016/j.cell.2011.12.022.

784   11.   Qi, W., Wang, G., and Wang, L. (2021). A novel satiety sensor detects circulating

785   glucose and suppresses food consumption via insulin-producing cells in Drosophila.

786   Cell Res *31*, 580–588. 10.1038/s41422-020-00449-7.

787   12.   Zhao, Y., Khallaf, M.A., Johansson, E., Dzaki, N., Bhat, S., Alfredsson, J., Duan, J.,

788   Hansson, B.S., Knaden, M., and Alenius, M. (2022). Hedgehog-mediated gut-taste

789   neuron axis controls sweet perception in Drosophila. Nat Commun *13*, 7810.

790   10.1038/s41467-022-35527-4.

791   13.   Oh, Y., Lai, J.S., Mills, H.J., Erdjument-Bromage, H., Giammarinaro, B., Saadipour,

792   K., Wang, J.G., Abu, F., Neubert, T.A., and Suh, G.S.B. (2019). A glucose-sensing

793   neuron pair regulates insulin and glucagon in Drosophila. Nature *574*, 559–564.

794   10.1038/s41586-019-1675-4.

14.   Dus, M., Lai, J.S., Gunapala, K.M., Min, S., Tayler, T.D., Hergarden, A.C., Geraud, E.,

Joseph, C.M., and Suh, G.S. (2015). Nutrient Sensor in the Brain Directs the Action

of the Brain-Gut Axis in Drosophila. Neuron *87*, 139 – 151.

10.1016/j.neuron.2015.05.032.

15.   Schlegel, P., Texada, M.J., Miroschnikow, A., Schoofs, A., Huckesfeld, S., Peters, M.,

Schneider-Mizell, C.M., Lacin, H., Li, F., Fetter, R.D., et al. (2016). Synaptic

transmission parallels neuromodulation in a central food-intake circuit. Elife *5*.

10.7554/eLife.16799.

16.   Schoofs, A., Huckesfeld, S., Schlegel, P., Miroschnikow, A., Peters, M., Zeymer, M.,

Spiess, R., Chiang, A.S., and Pankratz, M.J. (2014). Selection of motor programs for

suppressing food intake and inducing locomotion in the Drosophila brain. PLoS Biol

12, e1001893. 10.1371/journal.pbio.1001893.

17.   Hergarden, A.C., Tayler, T.D., and Anderson, D.J. (2012). Allatostatin-A neurons

inhibit feeding behavior in adult Drosophila. Proc Natl Acad Sci U S A *109*, 3967–

3972. 10.1073/pnas.1200778109.

18.   Melcher, C., Bader, R., Walther, S., Simakov, O., and Pankratz, M.J. (2006).

Neuromedin U and its putative Drosophila homolog hugin. PLoS Biol *4*, e68.

10.1371/journal.pbio.0040068.

19.   Munch, D., Ezra-Nevo, G., Francisco, A.P., Tastekin, I., and Ribeiro, C. (2020).

Nutrient homeostasis - translating internal states to behavior. Curr Opin Neurobiol *60*,

67–75. 10.1016/j.conb.2019.10.004.

20.   Marella, S., Mann, K., and Scott, K. (2012). Dopaminergic modulation of sucrose

acceptance   behavior   in   Drosophila.   Neuron   *73*,   941   –   950.

10.1016/j.neuron.2011.12.032.

21.   Augustine, V., Lee, S., and Oka, Y. (2020). Neural Control and Modulation of Thirst,

Sodium Appetite, and Hunger. Cell *180*, 25–32. 10.1016/j.cell.2019.11.040.

22.   de Rivaz, R., Swendsen, J., Berthoz, S., Husky, M., Merikangas, K., and Marques-

Vidal, P. (2022). Associations between Hunger and Psychological Outcomes: A

Large-Scale   Ecological   Momentary   Assessment   Study.   Nutrients   *14*.

10.3390/nu14235167.

23.   Alhadeff, A.L., Su, Z., Hernandez, E., Klima, M.L., Phillips, S.Z., Holland, R.A., Guo,

C., Hantman, A.W., De Jonghe, B.C., and Betley, J.N. (2018). A Neural Circuit for the

Suppression of Pain by a Competing Need State. Cell *173*, 140 – 152 e115.

10.1016/j.cell.2018.02.057.

24.   Sun, M., Ma, M., Deng, B., Li, N., Peng, Q., and Pan, Y. (2023). A neural pathway

underlying hunger modulation of sexual receptivity in Drosophila females. Cell Rep *42*,

113243. 10.1016/j.celrep.2023.113243.

25.   Burnett, C.J., Li, C., Webber, E., Tsaousidou, E., Xue, S.Y., Bruning, J.C., and

Krashes, M.J. (2016). Hunger-Driven Motivational State Competition. Neuron *92*, 187

–201. 10.1016/j.neuron.2016.08.032.

26.   Deem, J.D., Faber, C.L., and Morton, G.J. (2022). AgRP neurons: Regulators of

feeding,   energy   expenditure,   and   behavior.   FEBS   J   *289*,   2362 – 2381.

10.1111/febs.16176.

27.   Karigo, T., and Deutsch, D. (2022). Flexibility of neural circuits regulating mating

behaviors in mice and flies. Front Neural Circuits *16*, 949781.

10.3389/fncir.2022.949781.

28.   Fu, O., Iwai, Y., Narukawa, M., Ishikawa, A.W., Ishii, K.K., Murata, K., Yoshimura, Y.,

Touhara, K., Misaka, T., Minokoshi, Y., and Nakajima, K.I. (2019). Hypothalamic

neuronal circuits regulating hunger-induced taste modification. Nat Commun *10*, 4560.

10.1038/s41467-019-12478-x.

29.   Dietrich, M.O., Zimmer, M.R., Bober, J., and Horvath, T.L. (2015). Hypothalamic Agrp

neurons drive stereotypic behaviors beyond feeding. Cell *160*, 1222 – 1232.

10.1016/j.cell.2015.02.024.

30.   Bruning, J.C., and Fenselau, H. (2023). Integrative neurocircuits that control

metabolism and food intake. Science *381*, eabl7398. 10.1126/science.abl7398.

31.   Chen, Y., Lin, Y.C., Kuo, T.W., and Knight, Z.A. (2015). Sensory detection of food

rapidly modulates arcuate feeding circuits. Cell *160*, 829 – 841.

10.1016/j.cell.2015.01.033.

32.   Koch, M., Varela, L., Kim, J.G., Kim, J.D., Hernandez-Nuno, F., Simonds, S.E.,

Castorena, C.M., Vianna, C.R., Elmquist, J.K., Morozov, Y.M., et al. (2015).

Hypothalamic POMC neurons promote cannabinoid-induced feeding. Nature *519*, 45–

50. 10.1038/nature14260.

33.   Belgardt, B.F., Okamura, T., and Bruning, J.C. (2009). Hormone and glucose

signalling in POMC and AgRP neurons. J Physiol *587*, 5305 – 5314.

10.1113/jphysiol.2009.179192.

34.   Vivot, K., Meszaros, G., Pangou, E., Zhang, Z., Qu, M., Erbs, E., Yeghiazaryan, G.,

861   Quinones, M., Grandgirard, E., Schneider, A., et al. (2023). CaMK1D signalling in

862   AgRP neurons promotes ghrelin-mediated food intake. Nat Metab *5*, 1045–1058.

863   10.1038/s42255-023-00814-x.

864   35.   Wei, Q., Krolewski, D.M., Moore, S., Kumar, V., Li, F., Martin, B., Tomer, R., Murphy,

865   G.G., Deisseroth, K., Watson, S.J., Jr., and Akil, H. (2018). Uneven balance of power

866   between hypothalamic peptidergic neurons in the control of feeding. Proc Natl Acad

867   Sci U S A *115*, E9489–E9498. 10.1073/pnas.1802237115.

868   36.   Lin, S., Senapati, B., and Tsao, C.H. (2019). Neural basis of hunger-driven behaviour

869   in Drosophila. Open Biol *9*, 180259. 10.1098/rsob.180259.

870   37.   Chouhan, N.S., Griffith, L.C., Haynes, P., and Sehgal, A. (2021). Availability of food

871   determines the need for sleep in memory consolidation. Nature *589*, 582–585.

872   10.1038/s41586-020-2997-y.

873   38.   Willyard, C. (2008). Hungry for sleep. Nat Med *14*, 477–480. 10.1038/nm0508-477.

874   39.   Yao, Z., and Scott, K. (2022). Serotonergic neurons translate taste detection into

875   internal nutrient regulation. Neuron *110*, 1036 – 1050 e1037.

876   10.1016/j.neuron.2021.12.028.

877   40.   Malita, A., Kubrak, O., Koyama, T., Ahrentlov, N., Texada, M.J., Nagy, S., Halberg,

878   K.V., and Rewitz, K. (2022). A gut-derived hormone suppresses sugar appetite and

879   regulates food choice in Drosophila. Nat Metab *4*, 1532–1550. 10.1038/s42255-022-

880   00672-z.

881   41.   Melcher, C., and Pankratz, M.J. (2005). Candidate gustatory interneurons modulating

882   feeding behavior in the Drosophila brain. PLoS Biol *3*, e305.

10.1371/journal.pbio.0030305.

42.   Root, C.M., Ko, K.I., Jafari, A., and Wang, J.W. (2011). Presynaptic facilitation by

neuropeptide signaling mediates odor-driven food search. Cell *145*, 133 – 144.

10.1016/j.cell.2011.02.008.

43.   Chen, J., Reiher, W., Hermann-Luibl, C., Sellami, A., Cognigni, P., Kondo, S.,

Helfrich-Forster, C., Veenstra, J.A., and Wegener, C. (2016). Allatostatin A Signalling

in Drosophila Regulates Feeding and Sleep and Is Modulated by PDF. PLoS Genet

12, e1006346. 10.1371/journal.pgen.1006346.

44.   Hentze, J.L., Carlsson, M.A., Kondo, S., Nassel, D.R., and Rewitz, K.F. (2015). The

Neuropeptide Allatostatin A Regulates Metabolism and Feeding Decisions in

Drosophila. Sci Rep *5*, 11680. 10.1038/srep11680.

45.   Kubrak, O.I., Lushchak, O.V., Zandawala, M., and Nassel, D.R. (2016). Systemic

corazonin signalling modulates stress responses and metabolism in Drosophila.

Open Biol *6*. 10.1098/rsob.160152.

46.   Martelli, C., Pech, U., Kobbenbring, S., Pauls, D., Bahl, B., Sommer, M.V., Pooryasin,

A., Barth, J., Arias, C.W.P., Vassiliou, C., et al. (2017). SIFamide Translates Hunger

Signals into Appetitive and Feeding Behavior in Drosophila. Cell Rep *20*, 464–478.

10.1016/j.celrep.2017.06.043.

47.   Ugrankar, R., Theodoropoulos, P., Akdemir, F., Henne, W.M., and Graff, J.M. (2018).

Circulating glucose levels inversely correlate with Drosophila larval feeding through

insulin signaling and SLC5A11. Commun Biol *1*, 110. 10.1038/s42003-018-0109-4.

48.   Oh, Y., Lai, J.S., Min, S., Huang, H.W., Liberles, S.D., Ryoo, H.D., and Suh, G.S.B.

(2021). Periphery signals generated by Piezo-mediated stomach stretch and

Neuromedin-mediated glucose load regulate the Drosophila brain nutrient sensor.

Neuron *109*, 1979–1995 e1976. 10.1016/j.neuron.2021.04.028.

49.   Senapati, B., Tsao, C.H., Juan, Y.A., Chiu, T.H., Wu, C.L., Waddell, S., and Lin, S.

(2019). A neural mechanism for deprivation state-specific expression of relevant

memories in Drosophila. Nat Neurosci *22*, 2029–2039. 10.1038/s41593-019-0515-z.

50.   Yurgel, M.E., Kakad, P., Zandawala, M., Nassel, D.R., Godenschwege, T.A., and

Keene, A.C. (2019). A single pair of leucokinin neurons are modulated by feeding

state and regulate sleep-metabolism interactions. PLoS Biol *17*, e2006409.

10.1371/journal.pbio.2006409.

51.   Song, T., Qin, W., Lai, Z., Li, H., Li, D., Wang, B., Deng, W., Wang, T., Wang, L., and

Huang, R. (2023). Dietary cysteine drives body fat loss via FMRFamide signaling in

Drosophila and mouse. Cell Res *33*, 434–447. 10.1038/s41422-023-00800-8.

52.   Inagaki, H.K., Panse, K.M., and Anderson, D.J. (2014). Independent, reciprocal

neuromodulatory control of sweet and bitter taste sensitivity during starvation in

Drosophila. Neuron *84*, 806–820. 10.1016/j.neuron.2014.09.032.

53.   Fujii, S., Yavuz, A., Slone, J., Jagge, C., Song, X., and Amrein, H. (2015). Drosophila

sugar receptors in sweet taste perception, olfaction, and internal nutrient sensing.

Curr Biol *25*, 621–627. 10.1016/j.cub.2014.12.058.

54.   Budnik, V., and White, K. (1987). Genetic dissection of dopamine and serotonin

synthesis in the nervous system of Drosophila melanogaster. J Neurogenet *4*, 309–

314.

55. Deng, B., Li, Q., Liu, X., Cao, Y., Li, B., Qian, Y., Xu, R., Mao, R., Zhou, E., Zhang,

W., et al. (2019). Chemoconnectomics: Mapping Chemical Transmission in

Drosophila. Neuron *101*, 876–893 e874. 10.1016/j.neuron.2019.01.045.

56. Yoon, N.A., and Diano, S. (2021). Hypothalamic glucose-sensing mechanisms.

Diabetologia *64*, 985–993. 10.1007/s00125-021-05395-6.

57. Brogiolo, W., Stocker, H., Ikeya, T., Rintelen, F., Fernandez, R., and Hafen, E. (2001).

An evolutionarily conserved function of the Drosophila insulin receptor and insulin-like

peptides in growth control. Curr Biol *11*, 213–221. 10.1016/s0960-9822(01)00068-9.

58. Fosque, B.F., Sun, Y., Dana, H., Yang, C.T., Ohyama, T., Tadross, M.R., Patel, R.,

Zlatic, M., Kim, D.S., Ahrens, M.B., et al. (2015). Neural circuits. Labeling of active

neural circuits in vivo with designed calcium integrators. Science *347*, 755–760.

10.1126/science.1260922.

59. Parton, L.E., Ye, C.P., Coppari, R., Enriori, P.J., Choi, B., Zhang, C.Y., Xu, C., Vianna,

C.R., Balthasar, N., Lee, C.E., et al. (2007). Glucose sensing by POMC neurons

regulates glucose homeostasis and is impaired in obesity. Nature *449*, 228–232.

10.1038/nature06098.

60. Wang, M., Wu, J.X., Ding, D., and Chen, L. (2022). Structural insights into the

mechanism of pancreatic K(ATP) channel regulation by nucleotides. Nat Commun *13*,

2770. 10.1038/s41467-022-30430-4.

61. Park, Y., Kim, Y.J., and Adams, M.E. (2002). Identification of G protein-coupled

receptors for Drosophila PRXamide peptides, CCAP, corazonin, and AKH supports a

theory of ligand-receptor coevolution. Proc Natl Acad Sci U S A *99*, 11423–11428.

10.1073/pnas.162276199.

62.   Klapoetke, N.C., Murata, Y., Kim, S.S., Pulver, S.R., Birdsey-Benson, A., Cho, Y.K.,

Morimoto, T.K., Chuong, A.S., Carpenter, E.J., Tian, Z., et al. (2014). Independent

optical excitation of distinct neural populations. Nat Methods *11*, 338 – 346.

10.1038/nmeth.2836.

63.   Dahanukar, A., Lei, Y.T., Kwon, J.Y., and Carlson, J.R. (2007). Two Gr genes

underlie sugar reception in Drosophila. Neuron *56*, 503 – 516.

10.1016/j.neuron.2007.10.024.

64.   Larsen, M.J., Burton, K.J., Zantello, M.R., Smith, V.G., Lowery, D.L., and Kubiak, T.M.

(2001). Type A allatostatins from Drosophila melanogaster and Diplotera puncata

activate two Drosophila allatostatin receptors, DAR-1 and DAR-2, expressed in CHO

cells. Biochem Biophys Res Commun *286*, 895–901. 10.1006/bbrc.2001.5476.

65.   Kitamoto, T. (2001). Conditional modification of behavior in Drosophila by targeted

expression of a temperature-sensitive shibire allele in defined neurons. J Neurobiol

47, 81–92. 10.1002/neu.1018.

66.   Stocker, R.F. (2004). Taste perception: Drosophila - a model of good taste. Curr Biol

14, R560–561. 10.1016/j.cub.2004.07.011.

67.   Yarmolinsky, D.A., Zuker, C.S., and Ryba, N.J. (2009). Common sense about taste:

from mammals to insects. Cell *139*, 234–244. 10.1016/j.cell.2009.10.001.

68.   Jin, H., Fishman, Z.H., Ye, M., Wang, L., and Zuker, C.S. (2021). Top-Down Control

of Sweet and Bitter Taste in the Mammalian Brain. Cell *184*, 257–271 e216.

10.1016/j.cell.2020.12.014.

69.   Kain, P., and Dahanukar, A. (2015). Secondary taste neurons that convey sweet

taste and starvation in the Drosophila brain. Neuron *85*, 819 – 832.

10.1016/j.neuron.2015.01.005.

70.   Williams, M.J., Akram, M., Barkauskaite, D., Patil, S., Kotsidou, E., Kheder, S., Vitale,

G., Filaferro, M., Blemings, S.W., Maestri, G., et al. (2020). CCAP regulates feeding

behavior via the NPF pathway in Drosophila adults. Proc Natl Acad Sci U S A *117*,

7401–7408. 10.1073/pnas.1914037117.

71.   Matsushita, R., and Nishimura, T. (2020). Trehalose metabolism confers

developmental robustness and stability in Drosophila by regulating glucose

homeostasis. Commun Biol *3*, 170. 10.1038/s42003-020-0889-1.

72.   Ma, D., Hu, M., Yang, X., Liu, Q., Ye, F., Cai, W., Wang, Y., Xu, X., Chang, S., Wang,

R., et al. (2024). Structural basis for sugar perception by Drosophila gustatory

receptors. Science *383*, eadj2609. 10.1126/science.adj2609.

73.   Hanada, R., Teranishi, H., Pearson, J.T., Kurokawa, M., Hosoda, H., Fukushima, N.,

Fukue, Y., Serino, R., Fujihara, H., Ueta, Y., et al. (2004). Neuromedin U has a novel

anorexigenic effect independent of the leptin signaling pathway. Nat Med *10*, 1067–

1073. 10.1038/nm1106.

74.   Howard, A.D., Wang, R., Pong, S.S., Mellin, T.N., Strack, A., Guan, X.M., Zeng, Z.,

Williams, D.L., Jr., Feighner, S.D., Nunes, C.N., et al. (2000). Identification of

receptors for neuromedin U and its role in feeding. Nature *406*, 70 – 74.

10.1038/35017610.

75.   Miroschnikow, A., Schlegel, P., and Pankratz, M.J. (2020). Making Feeding Decisions

in the Drosophila Nervous System. Curr Biol *30*, R831 – R840.

10.1016/j.cub.2020.06.036.

76. Teranishi, H., and Hanada, R. (2021). Neuromedin U, a Key Molecule in Metabolic

Disorders. Int J Mol Sci *22*. 10.3390/ijms22084238.

77. Aldawsari, M., Almadani, F.A., Almuhammadi, N., Algabsani, S., Alamro, Y., and

Aldhwayan, M. (2023). The Efficacy of GLP-1 Analogues on Appetite Parameters,

Gastric Emptying, Food Preference and Taste Among Adults with Obesity:

Systematic Review of Randomized Controlled Trials. Diabetes Metab Syndr Obes *16*,

575–595. 10.2147/DMSO.S387116.

78. Sun, J., Liu, C., Bai, X., Li, X., Li, J., Zhang, Z., Zhang, Y., Guo, J., and Li, Y. (2017).

Drosophila FIT is a protein-specific satiety hormone essential for feeding control. Nat

Commun *8*, 14161. 10.1038/ncomms14161.

1005

1006

 Figure legends

 Figure 1. hugin^{+} neurons were a novel glucose sensor in the fly brain.

 (**A**) Experimental procedure schematic. Newly eclosed flies were gathered and provided with

 regular food for 5 days, followed by a 12-hour period of starvation. Proboscis Extension

 Response (PER) assays were conducted. (**B**) Fraction of flies showing PER to different

 concentrations of sucrose (two-way ANOVA ;*,p=0.0349;***,p=0.002;****,p<0.0001; n=4

 groups, each of 10 flies). (**C-D**) Fraction of flies of the indicated genotypes under 30 ºC

 showing PER to 400 mM of sucrose (one-way

 ANOVA ;*,p=0.0233;**,p=0.0054;***,p=0.001;****,p<0.0001; n=4-5 groups, each of 10

 flies). (**E**) Fraction of indicated flies showing PER to different concentrations of sucrose (ND

 represent standard fly food) (two-way ANOVA ; ****,p<0.0001; n=4-5 groups, each of 10

 flies). (**F**) hugin expression in the brain, illustrated by mCD8::GFP expression driven by

 hugin^{GAL4}. Scale bar represents 100 μm. (**G**) Representative traces and quantification of ex

 vivo calcium responses of hugin^{+} neurons during the perfusion of glucose with or without

 TTX (one-way ANOVA ;**,p=0.002; n=6-7). Horizontal black bar represents the duration

 indicated glucose solution stimulation. (**H**) AstA expression in the brain, illustrated by

 mCD8::GFP expression driven by *AstA^{GAL4}*. Scale bar represents 100 μm. (**I**) Representative

 traces and quantification of ex vivo calcium responses of AstA^{+} neurons during the perfusion

 of glucose with or without TTX (one-way ANOVA ;*,p=0.0187 or 0.022; n=6). (**J**)

 Representative images of pre-photoconversion (pre-PC) and post-photoconversion (post-PC)

 CaMPARI signal in hugin-expressing neurons (upper). The Red:Green ratio represents

 intracellular Ca^{2+} concentrations (lower). Scale bar represents 100 μm (one-way

1029   ANOVA ;****,p<0.0001; n=18-23). ns, $P > 0.05$; *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$;

1030   ****$P < 0.0001$. Student's *t-test*, one-way and two-way ANOVA followed by post hoc test

1031   with Bonferroni correction were used for multiple comparisons when applicable.

1032

Figure 2. Glucose activated hugin$^+$ neurons via Gltu1 and K$_{ATP}$ channel.

(**A**) Representative traces and quantification of ex vivo calcium responses of hugin$^+$ neurons

during the perfusion of glucose with or without phlorizin (1mM),(t-test;**,p=0.0052; n=6-7).

Horizontal black bar represents the duration indicated glucose solution stimulation. (**B**)

Representative traces and quantification of ex vivo calcium responses of hugin$^+$ neurons in

indicated flies during the perfusion of glucose (t-test;*,p=0.022; n=6). (**C**) Fraction of flies of

the indicated genotypes showing PER to different concentrations of sucrose (two-way

ANOVA ; ****,p<0.0001;n=4 groups, each of 10 flies). (**D-E**) Representative traces and

quantification of ex vivo calcium responses of hugin$^+$ neurons during the perfusion of

glibenclamide (100 μM, D), glucose with alloxan (10 μM, E) (t-test;**,p=0.0091; n=6). (**F**)

Representative traces and quantification of ex vivo calcium responses of hugin$^+$ neurons in

indicated flies during the perfusion of glucose (t-test;**,p=0.0011; n=6) . (**G**) Representative

traces and quantification of ex vivo calcium responses of hugin$^+$ neurons during the perfusion

of pyruvate (50 mM), (t-test;ns,p>0.05; n=6). (**H**) Representative traces and quantification of

ex vivo calcium responses of hugin$^+$ neurons during the perfusion of D-glucose and L-

glucose (50 mM), (t-test;***,p=0.0045; n=6) ns, P > 0.05; *P < 0.05; **P < 0.01; ****P <

0.0001. Student's t-test and two-way ANOVA followed by post hoc test with Bonferroni

correction were used for multiple comparisons when applicable.

1051

1052

1053

1054

Figure 3. hugin$^+$ neurons suppressed sweet taste through PK2-R1.

(**A-B**) Fraction of flies of the indicated genotypes and environmental temperatures showing PER to different concentrations of sucrose (two-way ANOVA ;*,p=0.0153 or 0.0237; ****,p<0.0001;n=4-5 groups, each of 10 flies). (**C**) Fraction of flies showing PER to different concentrations of sucrose (two-way ANOVA ;*,p=0.0486 or 0.0438 ;***,p=0.0007 or 0.0001; ****,p<0.0001; n=4 groups, each of 10 flies). Flies were injected with saline or synthetic hugin for 30 minutes before the assay. (**D**) Representative traces (upper) and quantification (lower) of peak calcium transients of Gr5a$^+$ neurons in indicated flies upon 5% sucrose after injection of synthetic hugin (t-test;***,p=0.0047; n=6). Horizontal black bar represents the duration sucrose stimulation. (**E-F**) Fraction of flies of the indicated genotypes showing PER to different concentrations of sucrose (two-way ANOVA ;**,p=0.0055 or 0.0016 or 0.003 ;***,p=0.0002 or 0.0005; ****,p<0.0001;n=4 groups, each of 10 flies)ns, P > 0.05; *P < 0.05; ***P < 0.001; ****P < 0.0001. Two-way ANOVA followed by post hoc test with Bonferroni correction was used for multiple comparisons when applicable.

Figure 4. hugin⁺ neurons activated AstA⁺ neurons through PK2-R1.

(**A-B**) Fraction of flies of the indicated genotypes and environmental temperatures showing

PER to different concentrations of sucrose (two-way ANOVA ;*,p=0.043 ;***,p=0.0007 or

0.0001; ****,p<0.0001;n=4-5 groups, each of 10 flies). (**C**) Co-localization (dashed box) of

PK2-R1⁺ neurons (green) and AstA⁺ neurons (red) in the SEZ region. Scale bar represents 100

μm. (**D**) Representative traces (upper) and quantification (lower) of peak calcium transients of

AstA⁺ neurons after the photo-activation of hugin⁺ neurons (t-test;*,p=0.0114; n=6) from in

vivo calcium imaging. Horizontal black bar represents the duration of red-light stimulation.

(**E-F**) Fraction of flies of the indicated genotypes showing PER to different concentrations of

sucrose (two-way ANOVA ;*,p=0.0486 or 0.033 or 0.0224;**,p=0.0043 or

0.0011 ;***,p=0.0001 or 0.0008; ****,p<0.0001; n=4 groups, each of 10 flies). ns, P > 0.05;

*P < 0.05; **P < 0.01; ***P < 0.001; ****P < 0.0001. Student's *t-test* and two-way ANOVA

followed by post hoc test with Bonferroni correction were used for multiple comparisons

when applicable.

1089

Figure 5. AstA$^+$ neurons inhibited Gr5a$^+$ neurons through AstA-R1.

(**A**) Fraction of flies showing PER to different concentrations of sucrose (two-way

ANOVA ;*,p=0.024;**,p=0.0072 or 0.002 ;***,p=0.0002; ****,p<0.0001; n=4 groups, each

of 10 flies). Flies were injected with saline or synthetic AstA for 30 minutes before the assay.

(**B**) Representative traces (left) and quantification (right) of peak calcium transients of Gr5a$^+$

neurons in indicated flies upon 5% sucrose after injection of synthetic AstA (t-test;*,p=0.0476;

n=6). Horizontal black bar represents the duration sucrose stimulation. (**C**) Localization of

Gr5a$^+$ neurons (green) and AstA$^+$ neurons (red) in the brain. Scale bar represents 100 μm. (**D**)

Co-localization of Gr5a$^+$ neurons (green) and AstA-R1$^+$ neurons (red). Arrows point to the

neurons co-expressing the two receptors. Scale bar represents 50 μm. (**E-F**) Fraction of flies

of the indicated genotypes showing PER to different concentrations of sucrose (two-way

ANOVA **,p=0.0088; ***,p=0.0004; ****,p<0.0001; n=4 groups, each of 10 flies). (**G-H**)

Representative traces (left) and quantification (right) of peak calcium transients of Gr5a$^+$

neurons in indicated flies upon 5% sucrose during the process of photo-activation of AstA$^+$

neurons (G, n=6; t-test;**,p=0.0058) and hugin$^+$ neurons (H, n=7-8;t-test;*,p=0.0351).

Shadows represents the duration of red-light stimulation. Horizontal black bar represents the

duration of sucrose stimulation. ns, P > 0.05; *P < 0.05; ***P < 0.001; ****P < 0.0001.

Student's *t-test* and two-way ANOVA followed by post hoc test with Bonferroni correction

were used for multiple comparisons when applicable.

1109

1110

1111

Figure 6. hugin-AstA circuitry inhibited feeding behavior.

(**A**) The illustration of experimental design in this Figure. Note that all flies were maintained

at 20ºC before the feeding assay to prevent neuronal silencing by Shibire[ts1] during these

procedures. (B) Fraction of flies of the indicated genotypes and environmental temperatures

showing PER to different concentrations of sucrose (two-way ANOVA ; ****,$p<0.0001$; n=4

groups, each of 10 flies). (**C-D**) Food consumption of flies of the indicated genotypes and

environmental temperatures (one-way ANOVA ;**,$p=0.004$; ****,$p<0.0001$; n=8 groups,

each of 4 flies). Food consumption is presented as fold change relative to the corresponding

genetic control.

1121

Figure 7. NMU⁺ neurons were the central energy sensor in mouse.

(**A**) Blood NMU levels of starved mice re-fed with 20% sucrose (t-test; *, p=0.0373; n=6).

(**B**) Two-bottle preference tests for starved mice re-fed with sucrose (t-test; ***, p=0.0002; n=7). (**C**) Two-bottle preference tests for starved mice with or without intraperitoneal injection of NMU peptide (YFLFRPRN-NH₂, 4.5μm/kg) (t-test; *, p=0.0293; n=7). (**D**) Two-bottle preference tests for indicated starved mice (t-test; **, p=0.0073; n=6). (**E**) Fiber photometry to record the calcium dynamics of NMU⁺ neurons in the VMH with GCaMP6m (left) and representative IHC image of expressing GCaMP6min VMH (right). (**F-G**) Representative trace (left) and heatmaps (right, n=5) showing calcium dynamics of NMU⁺ neurons in the VMH in response to gastric glucose infusion (20% sucrose, 200 μl), which elevates circulating glucose levels independently of oral sensory stimulation.. (**H**) Experimental approach to assess calcium signaling in NMU⁺ neurons in the VMH in vitro (left) and representative IHC image of expressing GCaMP6min VMH (right). (**I-J**) Representative traces and quantification of ex vivo calcium responses of NMU⁺ neurons during the perfusion of glucose with or without TTX (I), Alloxan (J) and Phlorizin (J) (one-way ANOVA ; **, p=0.0049;***, p=0.0001 or 0.0006; n=6-7). Horizontal black bar represents the duration indicated glucose solution stimulation. (**K**) Anterograde trans-synaptic tracing of downstream targets of NMU□ neurons. NMU□ neurons were labeled by GFP expression following injection of a Cre-dependent AAV2/1-DIO-GFP into NMU-Cre mice. This virus undergoes anterograde trans-synaptic transfer to postsynaptic neurons. To enable GFP expression specifically in downstream target regions, an AAV-Cre virus was locally injected into the rNST. As a result, postsynaptic neurons in the rNST receiving input from

NMU☐ neurons were labeled by GFP delivered anterogradely from upstream NMU☐

neurons. Representative images show that GFP-labeled downstream neurons in the rNST

colocalize with Calb2 immunoreactivity, indicating that NMU☐ neurons preferentially target

Calb2☐ rNST neurons. (**L**) Fiber photometry to record the calcium dynamics of

Calb2$^+$ neurons in the rNST with GCaMP6m (left) and representative IHC image of expressing

GCaMP6m in rNST (right). (**M**) Representative traces and quantification of calcium responses

of Calb2$^+$ neurons during glucose licking under physiological feeding conditions (500 mM

sucrose), with or without NMU administration, assessing downstream modulation of sweet-

responsive brainstem circuits(t-test; ****, $p<0.0001$; n=6). ns, $P > 0.05$; *$P < 0.05$; **$P <$

0.01; ***$P < 0.001$; ****$P < 0.0001$. Student's *t-test* and one-way ANOVA followed by post

hoc test with Bonferroni correction was used for multiple comparisons when applicable.

1155

Figure supplementary 1. Satiety suppressed sweet sensitivity in a dopamine-independent

manner.

(**A**) Representative traces (left) and quantification (right) of peak calcium transients of $Gr5a^+$

neurons in indicated flies (t-test; *, p=0.0373; n=6). Horizontal black bar represents the

duration 5% sucrose stimulation. (**B-D**) Fraction of flies of the indicated genotypes showing

PER to different concentrations of sucrose (two-way ANOVA ; *, p=0.0498;****, p<0.0001;

B-C, n=4 groups, each of 10 flies). The Area Under the Curve (AUC, D) represents sweet

sensitivity in C. *$P < 0.05$; ****$P < 0.0001$. Student's *t-test* and two-way ANOVA followed

by post hoc test with Bonferroni correction were used for multiple comparisons when

applicable.

1166

Figure supplementary 2. The effect of satiety signals on taste modulation.

(**A**) Fraction of fed flies of the indicated genotypes and environmental temperatures showing

PER to different concentrations of sucrose (two-way ANOVA ; *, p=0.012; ***,

p=0.0006;****, p<0.0001; n=4 groups, each of 10 flies). (**B**) Fraction of starved flies of the

indicated genotypes and environmental temperatures showing PER to different concentrations

of sucrose (two-way ANOVA; *, p=0.0154; **, p=0.0081 or 0.0011; ***,p=0.0005 or 0.0001;

****, p<0.0001; n=4-5 groups, each of 10 flies). ns, $P > 0.05$; *$P < 0.05$; ***$P < 0.001$;

****$P < 0.0001$. Two-way ANOVA followed by post hoc test with Bonferroni correction was

used for multiple comparisons when applicable.

1176

Figure supplementary 3. Activation of IPCs decreased the circulating sugar levels and

increased sweet sensitivity.

(**A**) Hemolymph glucose levels from the indicated genotypes and treatment temperatures (t-

test; ****, p<0.0001; n=8-9, 1 hour at 30 ℃). (**B-C**) Fraction of fed flies of the indicated

genotypes and environmental temperatures showing PER to different concentrations of

sucrose (two-way ANOVA;**, p=0.0048; ****, p<0.0001; n=4 groups, each of 10 flies). ns,

P > 0.05; ****P < 0.0001. Student's *t-test* and two-way ANOVA followed by post hoc test

with Bonferroni correction were used for multiple comparisons when applicable.

1185

Figure supplementary 4. Sugar intake suppressed sweet sensation.

(A) Fraction of indicated flies which refed with different sugar with energy showing PER to

different concentrations of sucrose (two-way ANOVA ; ****, p<0.0001; n=4 groups, each of

10 flies). (B) Fraction of indicated flies which refed with different sugar without energy

showing PER to different concentrations of sucrose (two-way ANOVA ; ****, p<0.0001; n=4

groups, each of 10 flies). (**C**)   Fraction of indicated flies showing PER to different

concentrations of fructose (two-way ANOVA ; *, p=0.0255; ***, p=0.0002; ****, p<0.0001;

n=4 groups, each of 10 flies). (**D**) Fraction of indicated flies showing PER to different

concentrations of trehalose (two-way ANOVA;**, p=0.0081 or 0.0013; ***, p=0.0002; ****,

p<0.0001; n=4 groups, each of 10 flies). ****P < 0.0001. two-way ANOVA followed by post

hoc test with Bonferroni correction were used for multiple comparisons when applicable.

1197

1198 **Figure supplementary 5. Starvation led to a decrease in the circulating sugar levels.**

1199 Hemolymph glucose levels from indicated flies (t-test; ***, p=0.00017; ****, p<0.0001; n=6-

1200   8). ***P < 0.001. Student's *t-test* was used for comparisons.

1201

Figure supplementary 6. The response of different neuropeptide-expressing neurons to

glucose.

(**A**) LK expression in the brain, illustrated by mCD8::GFP expression driven by LK^{GAL4}. Scale

bar represents 100 μm. (**B**) Representative traces and quantification of ex vivo calcium

responses of LK$^+$ neurons during the perfusion of glucose (n=6). (**C-D**) The distribution (C)

and calcium responses (D) of TK$^+$ neurons (n=6). (**E-F**) The distribution (E) and calcium

responses (F) of CCHa1$^+$ neurons (n=6). (**G-H**) The distribution (G) and calcium responses

(H) of CCHa2$^+$ neurons (n=6).

1210

1211   **Figure supplementary 7. The calcium response of hugin neurons upon glucose stimuli.**

1212   (**A**) Representative image of hugin neurons in SEZ. Scale bar represents 50 μm. (**B**) Calcium

1213   responses of different cluster of hugin neurons during the prefusion of glucose (t-test, *,

1214   p=0.0151; n=6).*P < 0.05; ***P < 0.001. Student's *t-test* used for comparisons.

1215

1216

1217   **Figure supplementary 8. The calcium response of AstA$^+$ neurons upon glucose treatment.**

1218   (**A**) Representative image of different part of AstA neurons .Scale bar represents 50 μm. (**B**)

1219   Calcium responses of different cluster of hugin neurons during the prefusion of glucose (n=6)

1220

Figure supplementary 9. Knockout of *PK2-R1* enhanced sweet sensation.

(**A-C**) Fraction of flies of the indicated genotypes showing PER to different concentrations of

sucrose (two-way ANOVA ; *, p=0.0464 or 0.0153; **, p=0.0015 or 0.0038 or 0.0035; ***,

p= 0.0002; ****, p<0.0001; n=4 groups, each of 10 flies). ***P < 0.001; ****P < 0.0001.

Two-way ANOVA followed by post hoc test with Bonferroni correction were used for

multiple comparisons when applicable.

1227

1228   **Figure supplementary 10. The expressing pattern of PK2-R1 in the proboscis**

1229   Representative image of PK2-R1 in the proboscis. Scale bar represents 50 μm.

1230

Figure supplementary 11. hugin[+] neurons in the brain activated AstA[+] neurons.

(**A-B**) Fraction of flies of the indicated genotypes and environmental temperatures showing

PER to different concentrations of sucrose (two-way ANOVA ; **, p=0.0082; ***, p= 0.0007

or 0.0001; ****, p<0.0001; n=4 groups, each of 10 flies). Prior to the behavioral assay, the

neural connection between the brain and the ventral nerve cord (VNC) was surgically severed,

while the flies remained physically intact and retained normal proboscis musculature and

motor output. PER was measured using the same protocol as in intact flies. (**C**)

Representative traces (upper) and quantification (lower) of peak calcium transients of AstA[+]

neurons after the photo-activation of hugin[+] neurons (t-test;*, p=0.0337; n=6) from ex vivo

calcium imaging. Horizontal black bar represents the duration of red-light stimulation.

1241

Figure supplementary 12. Manipulating gene *AstA-R1* enhance PER.

(**A-C**) Fraction of flies of the indicated genotypes showing PER to different concentrations of

sucrose (two-way ANOVA ;*, p=0.0404; **, p=0.009 or 0.0022 or 0.0019;***, p=0.0005;

****, $p < 0.0001$; n=4 groups, each of 10 flies). ***$P < 0.001$; ****$P < 0.0001$. Two-way

ANOVA followed by post hoc test with Bonferroni correction were used for multiple

comparisons when applicable.

1248

1249   **Figure supplementary 13. The expressing pattern of AstA-R2 in the proboscis**

1250   Representative image of AstA-R2 in the proboscis. Scale bar represents 50 μm.

1251

Figure supplementary 14. hugin-AstA-Gr5a circuitry inhibited feeding behavior.

(**A**) Food consumption of flies of the indicated genotypes (one-way ANOVA ; ***, p=0.0009;

****, $p < 0.0001$; n=8 groups, each of 4 flies). (**B**) The TG content of indicated genotypes

(one-way ANOVA ; ****,p<0.0001; n=8 groups, each of 2 flies). (**C**) Food consumption of

flies of the indicated genotypes (one-way ANOVA ; ****,p<0.0001; n=8 groups, each of 4

flies). ns, $P > 0.05$; ***$P < 0.001$; ****$P < 0.0001$. Student's *t-test* and two-way ANOVA

followed by post hoc test with Bonferroni correction were used for multiple comparisons

when applicable.

1260

Figure supplementary 15. NMU peptide suppressed sweet taste in fly.

Fraction of flies showing PER to different concentrations of sucrose (two-way ANOVA ;*,

p=0.106; **, p=0.0015 or 0.0029 or 0.0025; ****, $p < 0.0001$; n=4 groups, each of 10 flies).

Flies were injected with saline or synthetic NMU for 30 mins before the assay.**$P < 0.01$;

****$P < 0.0001$. Two-way ANOVA followed by post hoc test with Bonferroni correction

were used for multiple comparisons when applicable.

1267

Figure supplementary 16. The expressing pattern of NMU neurons and Calb2 neurons

(**A**) Co-labeling of NMU[+] neurons using a Cre-dependent mCherry reporter (red) and in situ

hybridization (green).Scale bar represents 200 μm. (**B**) NMU arbors was extended near

Calb2[rNST] neurons. Scale bar represents 200 μm. (**C**) Co-labeling of Calb2[+] neurons with a

Cre-dependent mCherry reporter (red) and NMU antiboy staining (green). Scale bar

represents 200 μm.

1274

Figure supplementary 17. A working model

Elevated circulating glucose following feeding directly activates hugin+ neurons via cell-

autonomous glucose sensing. Activated hugin neurons release Hugin neuropeptide, which

engages PK2-R1 receptors on AstA neurons, promoting AstA release. AstA then suppresses

the sensitivity of Gr5a-expressing gustatory sensory neurons through AstA-R1 signaling,

thereby reducing sweet-driven feeding behavior.

Under starvation, reduced circulating glucose decreases endogenous hugin neuronal activity,

partially relieving this inhibitory tone and permitting enhanced sweet sensitivity. However,

experimental manipulation of the circuit demonstrates that the hugin–AstA axis retains the

capacity to modulate feeding behavior across feeding states, indicating that it functions as a

glucose-responsive, state-modulated inhibitory system rather than a strictly satiety-exclusive

brake.

The modest phenotype observed upon PK2-R2 reduction suggests the potential existence of

parallel or complementary hugin-dependent pathways, possibly arising from functional

heterogeneity within hugin neuronal subpopulations. These additional branches remain to be

defined.

1291

1292

1293

1294

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure supplementary 1

Figure supplementary 2

Figure supplementary 3

A

B

FED

C

ST

Figure supplementary 4

Figure supplementary 5

Figure supplementary 6

Figure supplementary 7

A

B

Figure supplementary 8

A

B

Figure supplementary 9

A

B

C

Figure supplementary 10

Figure supplementary 11

A FED

Fraction of flies showing PER

- hugin > TrpA1
- + > TrpA1
- hugin > +

ns ns

20 ℃

30 ℃

****

Sucrose concentration (mM)
12.5 50 200 800

B ST

Fraction of flies showing PER

- hugin > TrpA1
- + > TrpA1
- hugin > +

ns ns

20 ℃

30 ℃

**** **** ***

Sucrose concentration (mM)
12.5 50 200 800

C

AstA-Gal4 > UAS-GCaMP6m
hugin-LexA > +

AstA-Gal4 > UAS-GCaMP6m
hugin-LexA > LexAop-CsChrimson

ΔF/F₀

Time (s)

Peak ΔF/F₀

*

Figure supplementary 12

Figure supplementary 13

Figure supplementary 14

Figure supplementary 15

A

B

Figure supplementary 16

A AAV-DIO-mCherry injection → 3 weeks → Perfused Brain 4% PFA → FISH (NMU probe) → Antibody staining (mcherry antibody)

NMU-IRES-Cre mice

nmu
mCherry

VMH

merged   NMU   mCherry

B AAV-EF1α-DIO-EGFP

VMH   rNST

NMU-IRES-Cre mice

merged   GFP   Calb2

C AAV-DIO-mCherry injection → 3 weeks → Perfused Brain 4% PFA → Antibody staining (Calb2/mCherry antibody)

Calb2-IRES-Cre mice

Calb2
mCherry

rNST

merged   Calb2   mCherry

Figure supplementary 17