Tuberculosis susceptibility and inbreeding depression hinder *ex-situ* conservation in a critically endangered rainforest bird.

Peri E. Bolton*[1,2,3], Dustin J. Foote*[1,4], Nancy E. Drilling[5], Susan B. McRae[1], Michael D. Sorenson[6], Christopher N. Balakrishnan[1,2,7]

[1] Department of Biology, East Carolina University, Greenville, North Carolina, USA
[2] Department of Vertebrate Zoology, National Museum of Natural History, Smithsonian Institution, DC
[3] Department of Evolution, Ecology & Organismal Biology, Ohio State University, Columbus, Ohio, USA
[4] Sylvan Heights Bird Park, Scotland Neck, North Carolina, USA
[5] Department of Fisheries, Wildlife and Conservation Biology, University of Minnesota, St. Paul, Minnesota, USA
[6],Department of Biology, Boston University, Boston, Massachusetts, USA
[7] Division of Environmental Biology, National Science Foundation, Alexandria, Virginia, USA

*these authors contributed equally to this work

correspondence to: christopher.balakrishnan@gmail.com

ABSTRACT

Captive breeding can be a key component of species conservation strategies, but also exposes these rare species to novel environments including the pathogen landscape. The critically endangered white-winged wood duck (WWWD) *Asarcornis scutulata* has experienced substantial population declines, local extirpations and fragmentation of its former range in South-East Asia, making it one of the rarest birds in the world. Like other rare species, WWWD declines have led to the initiation of captive breeding programs, but these have been hampered by the WWWDs' high susceptibility to *Mycobacterium avium*, avian tuberculosis (TB). In this study we describe genome-wide patterns of diversity to understand the WWWD's demographic and phylogeographic history, inbreeding in the wild and in captivity, and the causes of TB susceptibility. Captive birds, which originated from northeast India, are genetically differentiated from wild birds sampled in Sumatra, Indonesia, likely reflecting long-standing phylogeographic structure. Demographic analyses revealed that long-term (Pleistocene) population declines preceded anthropogenic declines, a pattern shared with other codistributed, forest-dependent species. All sampled WWWD populations had extremely low genetic diversity but wild-sampled birds retained higher Major Histocompatibility Complex (MHC) diversity, reflecting important functional diversity in the wild. Genetic diversity has eroded over time in captivity and importantly, birds with higher levels of inbreeding succumb earlier to TB infections, suggesting inbreeding depression. Finally, by comparing gene expression between susceptible WWWD and resistant redhead ducks *Aythya americana* we identify possible mechanisms of TB susceptibility in WWWD. Altogether our study provides genomically-guided objectives for future management and a cautionary tale for *ex-situ* conservation.

INTRODUCTION

Captive breeding is an important component of species conservation strategies. However, husbandry in captivity poses additional challenges as these rare species are exposed to novel environments, including pathogens (1). These novel exposures may interact with inbreeding to hinder captive breeding success. Conservation genomics has the potential to provide not only an understanding of the demographic history of rare and threatened species but also their ability to adapt and respond to novel threats in a changing environment (2). However, genomic efforts have mostly focussed on species in the global north (3), leaving many equally or more imperiled species in the tropics poorly understood.

Asarcornis scutulata, the white-winged wood duck (Figure 1A), is one of the rarest birds on earth, yet we know nothing about patterns of genetic variation in this species. Also known simply as the white-winged duck, the longer moniker is more commonly used within the species' range (4) so we use that nomenclature here, abbreviated as WWWD. The current WWWD range extends from Northeast India to Sumatra, though only small, fragmented populations persist across this range (Figure 1B) (4–7). Somewhat unusual for waterfowl, WWWDs are forest dwelling and cavity nesting. WWWDs are highly dependent on mature rainforest with slow moving water bodies,

habitats that have been particularly degraded over the last century. Deforestation and poaching remain the biggest threats to the survival of the species.

Recent estimates suggest only 150 to 450 mature adult WWWDs remain in the wild, with the largest populations in northern Myanmar and Arunachal Pradesh and Assam Provinces in India (8). The WWWD is now considered critically endangered by the International Union for Conservation of Nature (IUCN), but protection efforts in India began in 1937 and the species was placed on the Indian Special Protected List in 1952 (9). The World Wildlife Fund selected the WWWD for a focal species project in 1968. As a result, the International Waterfowl Research Bureau and the World Wildlife Fund recommended that immediate action was needed to ensure the survival of WWWD in Assam (6, 9, 10).

As part of a resulting conservation plan, captive populations were initiated in the United Kingdom, and then the United States. These captive populations were established using birds collected in Assam in 1969 and in 1970 (9). Both of these early wild collections took place on the same tea plantation in upper Assam (Figure 1B). Staff at two Wildfowl and Wetland Trust (WWT) centers, located at Slimbridge and Peakirk, UK, were able to rapidly grow their captive WWWD populations to a total of 86 individuals by 1976 (11). Soon after establishment of the captive breeding program, however, it became clear that *Mycobacterium avium* — avian tuberculosis (TB) — was the source of mortality for most individuals. Between 1976 and 1991, 102 out of 121 birds (84%) succumbed to TB (12). Originating from the same wild population in Assam, WWWDs were initially imported to various facilities in the US. Led by Sylvan Heights Bird Park (SHBP) in Scotland Neck, North Carolina, the US population expanded to just under 200 individuals (Figure 1C) but following this population growth (*circa* 1999), avian TB began to significantly affect the North American population, primarily due to an increase in mortality in birds aged 2-6 years (13). This rapid turn of fortune for the US captive population suggests the possibility that inbreeding depression may have interacted with TB exposure to reduce the viability of captive breeding. At present, the current captive population is descended from just two wild founders.

In this study, we set out to describe genetic variation in WWWDs, including wild birds sampled in Indonesia and captive birds in the US, originally sourced from India. We aimed to provide a first look at the extant genetic diversity in this species with an aim to guide future conservation efforts. We specifically explore genetic variation in immune system genes of the major histocompatibility complex (MHC) and transcriptome-wide expression patterns to provide mechanistic insights into TB susceptibility in WWWDs.

RESULTS

Whole genome sequencing

We generated whole genome sequencing data for samples from three populations: "Historical Captive" (n = 9) birds were hatched at multiple breeding facilities in the U.S. between 1985 and 1990, and were later housed together at SHBP. "Current Captive" (n

= 10) birds hatched between 2008 and 2014 at SHBP (Figure 1B). "Wild" Indonesian (n =
11) samples were collected between 1999 and 2001 in the Way Kambas National Park
in Lampung Province, Indonesia (southern tip of Sumatra, Figure 1C; Table S1). We
used two different reference genomes and read mapping strategies for different
analyses. For analyses of mitochondrial DNA (mtDNA) and pairwise sequentially
markovian coalescent (PSMC) analyses we used the draft quality WWWD genome
assembly, which includes a complete mtDNA genome (GCA_013398475.1 (14), see
methods). For other population genomic analyses, we used a chromosome-level tufted
duck (*Aythya fuligula*) genome assembly (GCF_009819795.1 (15)) to allow insights into
complex genomic regions like the major histocompatibility complex (MHC). Overall
mapping rates were 97.63% to the *A. fuligula* assembly, marginally higher than mapping
to the less contiguous WWWD assembly. Captive birds were sequenced to an average
of 10.5x depth and wild birds were sequenced to an average 16.5x coverage; we
downsampled to equal read-depths as needed for specific analyses (see methods). All
resequencing data have been deposited in the NCBI Short Read Archive under
PRJNA1258233. Three wild samples were removed from genome-wide analyses due to
poor data quality (mapping rates < 50%).

Indonesian and Indian birds are genetically distinctive.

Analyses of both mitochondrial DNA (mtDNA) and genome-wide variation indicate
clear, historical divergence of the Indian-derived, captive population and the wild
Indonesian population. Mitochondrial lineages are reciprocally monophyletic, with 46 to
48 differences (~0.28 % divergence; Figure 1C) between the captive and wild lineages.
A time-calibrated phylogeny (Figure S1) suggests that these lineages diverged during
the late Pleistocene at ~170,000 years ago (95% HPD range = 103-241 kya). We note
that this estimate is considerably more recent than the calibration point, such that it likely
overestimates the actual divergence time to some degree (16). Similarly, Principal
Components Analysis of genome-wide variants using PCAngsd clearly separated the
three sampled populations (Figure 1D). Captive (India-derived) and wild (Indonesian)
birds separated along PC1 (26.9% of variation), whereas the historical and current
samples of captive birds separated along PC2 (4.9% of the variation) (17). This
separation in PC space corresponds to moderately high autosomal F_{ST} values between
captive and wild birds (historical versus wild = 0.3, current versus wild = 0.4) and
relatively low genome-wide differentiation between current and historical captive birds
(0.04).

Genetic diversity is low and inbreeding is high

Demographic analyses using *PSMC* (18) suggest long-term population declines,
particularly for the sample of wild Indonesian birds, which show evidence of steady
population decline throughout the Pleistocene (Figure 2A). In contrast, the WWWD
population in northeast India may have maintained a larger population size during the
late Pleistocene. Interestingly, the point at which the demographic trajectories of these
two populations diverge, around 150,000 years ago, may correspond to the isolation of
these populations and their mtDNA lineage divergence. Effective population sizes for the

Indian population remained relatively large until about 100,000 years ago after which this
population also declined.
156
Consistent with both longer term and more recent, anthropogenically-driven declines,
nucleotide diversity (π) was very low in all three population samples, with current captive
birds (median π = 0.0008) harboring about 53% of the diversity of their captive ancestors
(median π = 0.0015). Surprisingly, genetic diversity in the wild-sampled birds was
similarly low (median π = 0.0010) (Figure S2). Autosomal heterozygosity estimates
showed similar trends, with the wild population having lower heterozygosity than the
historical captive population and the current captive population declining considerably in
heterozygosity (~50%) (wild het = 0.001, historical het = 0.002, current het = 0.001;
Figure S2). Consistent with patterns of nuclear diversity, a single mtDNA haplotype is
found in the captive population, whereas the wild sample included two closely related
haplotypes (Figure 1B). These patterns are also reflected in the lengths and abundance
of runs of homozygosity (ROHs), which reflect inbreeding (19). Recent inbreeding
produces longer segments, which may be broken into shorter segments by subsequent
recombination. Current captive birds had the largest number of long (>10Mb) segments
and highest inbreeding coefficients as measured by F_{ROH} ($F_{ROH} = \Sigma L_{ROH}/L_{autosomes} = 0.47$
median for the current captive population), whereas ROHs were observed in only two
historical captive individuals (median F_{ROH} = 0). Some of the wild WWWDs had many
long ROHs, indicating recent inbreeding, but with a lower level of inbreeding overall
(median F_{ROH} = 0.11). This is in broad agreement with individual inbreeding coefficients
from *ngsF* (Figure S3), which were very low in the historical captive sample (median F =
0.007), and much higher in contemporary captives (median F = 0.25). In contrast, wild
birds had the highest individual inbreeding coefficients (median F = 0.36), primarily due
to intermediate length ROHs, which may indicate ongoing inbreeding in a smaller and
declining population on the island of Sumatra.
181
For a sample of sixteen captive birds with known age at death (ten current and six
historical captives), we observed a strong and significant negative relationship between
the *ngsF* individual inbreeding coefficient and age at death (Figure 2C; r = - 0.8, β = -29,
p = 0.00014, see also F_{ROH}, Figure S5, Table S2). More highly inbred individuals died
earlier, usually from mycobacteriosis. This pattern is driven in large part by a difference
between the historical and recent captives, with the recent captives dying younger than
their historical counterparts. Notably, the two recent captives with the longest lifespans
were the least inbred.

MHC diversity is low in captivity

Given the effect of inbreeding on susceptibility to mycobacteriosis, we next examined
genetic diversity in the major histocompatibility complex, a part of the genome with a key
role in immunity broadly and TB specifically (20, 21). Due to MHC copy number variation
among bird species we estimated diversity with two approaches: 1) using reads aligned
to the heterospecific tufted duck reference genome, which may underestimate diversity
due to reads that fail to align, and 2) using sample-depth corrected MHC reads aligned to
a single MHC copy (see methods).

Using the latter dataset, we estimated MHC gene copy numbers for WWWDs.
Coverage of reads mapped to a single MHC copy was different among populations,
especially for MHC Class I (Table S3). Based on these differences in relative sequencing
depth, we estimate that wild Indonesian WWWDs have at least three MHC class I
copies, and 1-2 class IIβ copies. In contrast, captive ducks have approximately half as
many copies of class I (1-2), but a comparable number of class IIβ copies.

There was a significant loss of heterozygosity in the peptide binding region (exons 2-
3) of MHC class I genes in captive birds (Figure 3A; Figure S7), with a non-significant
loss in nucleotide diversity (Figure 3B; Figure S6). These patterns are similarly reflected
in patterns of amino acid diversity (Figure 3C-D), suggesting a loss of functional
diversity. MHC reads aligned to single gene copy suggest a reduction in MHC Class II
PBR (exon 2) heterozygosity and nucleotide diversity (Figure 3A), but this was not
evident when the data were aligned to the tufted duck genome (Figure S6), perhaps due
to variation in read depth across paralogs (Figure S8). Importantly, wild birds have
relatively high MHC diversity at both Class I and Class II loci.

Immune gene expression in white-winged wood ducks

Birds in captive environments, including those at Sylvan Heights Bird Park, are
frequently exposed to *M. avium* but many species are only rarely affected by infections
(22). To provide a window into the mechanism of susceptibility, we conducted RNA-seq
on whole blood from WWWDs and closely-related redhead (RH) ducks *Aythya*
americana, which rarely become sick from TB exposure. We examined both species-
level differences in expression in asymptomatic birds (WWWD = 6, RH = 7) and
differences in symptomatic (WWWD=2, RH=1) versus asymptomatic birds within species
(Table S4).

We found that 56% of the whole-blood transcriptome was differentially expressed
between asymptomatic redhead and asymptomatic WWWDs (5,797 genes, q<0.05,
Table S5). There were no GO Biological Processes or KEGG pathways that were
significantly enriched among these genes (Table S6; Table S7). There were 341
candidate immune genes (see methods) differentially expressed between species, which
was not a significant enrichment (p=0.9). These immune genes included 27 RIG-I-like
receptor signalling genes (ko04622), and 51 genes in the Tuberculosis KEGG pathway
(ko05152) (p>0.05, Table S7). Among the differentially expressed candidate immune
genes were four key MHC genes, MHC Class I⬚ and beta-2 microglobulin (B2M), which
dimerize to form the Class I receptor protein, and transporter associated with antigen
processing 1 and 2 (TAP1 and TAP2), which together form the antigen presentation
complex. All four of these genes are substantially upregulated in asymptomatic WWWDs
relative to redhead ducks (LFC=1.8-2.8, q<5x10^{-57}, Figure 3E). Within the TB pathway,
there was no evidence for differential expression in MHC IIβ (Figure 3F), but toll-like
receptor 2 (TLR2) was downregulated in WWWDs (Figure 3G).

Although limited in sample size, we also contrasted symptomatic and asymptomatic
WWWDs. In total, 186 genes were differentially expressed after excluding genes with
expression patterns affected by outliers (q*<0.05; Table S8). There was significant
enrichment for genes in the GO Biological Process "Regulation of Apoptotic Process"

(GO:0042981, q=0.04, 5 genes). There was also weaker, non-significant, enrichment of
genes involved in "Immune Response" (GO:0006955, q=0.1, 5 genes; Table S9),
Yersinia Infection (ko05135, 5 genes, p=0.03, q=0.8) and Chemokine Signalling Pathway
(ko04062, 5 genes, p=0.04, q=0.8, Table S10). We found fewer disease-related DEGs
in redhead ducks (34 genes q*<0.05) (Table S11), with significant enrichment only non-
immune KEGG "Replication and Repair" (ko03030, 3 genes, Table S12-S13). We found
seven immune genes had opposite expression direction in symptomatic individuals of the
different species (Figure S11). No genes with significant differences in disease
expression were shared between the two species.

DISCUSSION

With biodiversity disappearing faster than it can be described, we provide here the
first characterization of genetic diversity in a critically endangered bird species. The
white-winged wood duck represents a monotypic genus *Asarcornis (23, 24)* that
historically ranged widely from eastern India to Indonesia. Though subspecies are not
currently recognized (25), Indonesian WWWDs have been considered a separate
subspecies (*leucoptera*) by earlier authors (26, 27) based on an apparently greater
extent of white on their head and mantle plumage as compared to Indian ducks
(nominate) (28). We found that captive birds derived from northeast India were
substantially genetically differentiated from those sampled in Sumatra (Figure 1). The
level of genetic divergence between Indian and Indonesian birds, including reciprocal
monophyly of mitochondrial lineages, is consistent with the recognition of distinct
subspecies. Given our sparse geographic sampling, however, the phylogeographic
distribution of these lineages and whether and where there is a clear break between
them remains unresolved. Further work is needed to test for associations between
genetic and phenotypic variation.
266

We assessed genetic variation in WWWD populations, including temporal samples of
India-derived captive birds in the US and wild birds sampled in Indonesia. All three
WWWD populations had extremely low genetic diversity with levels similar to other
critically endangered species like the five kiwi lineages (*Apteryx* genus) (29), crested ibis
Nipponia nippon (30), and the channel island fox *Urocyon littoralis* (31). Unexpectedly, a
sample of wild Indonesian WWWD collected in 1999-2001 had similarly low genetic
diversity to the captive populations sourced from India, which have been affected by both
founder effects and many generations of inbreeding in captivity. Low genetic diversity in
the wild appears to be a result of long-term population decline driven by environmental
change during the Pleistocene (Figure 2), and perhaps compounded by founder effects
during colonization of Sumatra and/or subsequent isolation following sea-level rise and
insularisation (32). Therefore, the low observed genetic diversity in the wild is not solely
driven by recent anthropogenic effects, but signatures of recent inbreeding are likely
signals of recent fragmentation (33). Instead, environmental change during Pleistocene
glacial cycles and the associated reduction of forest habitats (34) appear to have initiated
the decline of WWWDs globally. This observed demographic trajectory of WWWDs is
similar to other tropical vertebrates (35, 36) , including the critically endangered
Sumatran rhino, which shares a similar geographic distribution and habitat requirements

(37). As a charismatic megafauna species, the Sumatran rhino is a "flagship" for
conservation efforts that may also promote the persistence of WWWDs.
287
The current captive US population of WWWDs is derived from a single pair of
founders from Assam, India. As a result, we observed that genetic diversity has eroded
substantially over time and that more highly inbred birds had shorter lifespans (Figure
2C). This pattern is largely explained by differences in lifespan between individuals in our
current and historical samples, and so could be explained in part by possible increases
in TB exposure over time. Among our "current" sample, however, the two birds with the
longest lifespans were the least inbred, suggesting that inbreeding depression, and not
just TB exposure, contributes to reduced lifespans. This finding suggests that genetic
improvements to the current captive stock will be an important part of future WWWD *ex-
situ* management (38). Most WWWDs breed at ~3 years of age, and early in the
breeding program, birds lived long enough to lay multiple clutches of eggs. This is no
longer the case (13).
300
Despite having low overall genetic diversity, wild Indonesian WWWDs had higher
MHC diversity than the captive population (Figure 3A-D), representing a potentially
important reservoir of functional genetic variation. Similar retention of MHC diversity has
been seen in other bird species that have suffered population bottlenecks during island
colonization (39, 40). The observed difference in MHC diversity between captive Indian
and wild Indonesian birds is due to an apparent difference in copy number, with further
reductions in per locus diversity through time. Due to our sampling, however, it remains
unclear whether the difference in copy number is an evolved difference between Indian
and Indonesian birds, or is due to the two US founders from the Indian population having
an atypically small number of loci. Sampling of wild, Indian birds would resolve this
uncertainty.
312
Given the low MHC diversity in captive birds, we sought further insights into the
mechanism driving differences in susceptibility between WWWDs and other birds
housed in the same conditions. By conducting RNA-seq on a small number of
asymptomatic WWWDs and closely-related redhead ducks, we found substantial
differences in the expression of immune genes, including many genes associated with
the RIG-I pathway, a critical component of innate immune response to TB (41), and the
human tuberculosis KEGG pathway. One TB pathway gene, TLR2, is down-regulated in
asymptomatic WWWDs relative to redhead ducks. TLR2 plays a critical role in the innate
immune response. In mice, exposure to *M. tuberculosis* downregulates TLR2 and, in
turn, interferes with the induction of the type I interferon response and cross processing
by MHC Class I (20, 42). *Mycobacterium tuberculosis* itself produces TLR2 agonists in
mice. We speculate that *M. avium* may have a similar effect in WWWD, whereas
redhead ducks may be resistant to this form of immune system evasion. Necropsies of
WWWDs have revealed broad dissemination of the disease across tissues, a high
concentration of mycobacteria and a lack of multinucleated giant cells within lesions, all
suggesting a failure of cell mediated immunity (43). MHC Class I, its heterodimer B2M,
and associated transport proteins TAP1 and TAP2 were all upregulated in asymptomatic
WWWDs relative to redheads, but no MHC genes were differentially expressed between

symptomatic and asymptomatic WWWDs. This suggests that WWWDs have persistently
engaged a core component (MHC Class I) of self–non-self recognition and adaptive
immune pathways. This may be due to ubiquitous TB exposure in captivity. In this
scenario, asymptomatic birds are actually pre-symptomatic; they are infected with TB
and are mounting an immune response that is not effective at staving off disease in the
long-term. Alternatively, these expression patterns could suggest underlying immune
dysregulation such as autoimmunity or immune overexpression (44, 45).
338
Our findings strongly suggest that the genetic health of the captive WWWD
population in the US is a key factor in their susceptibility to *M. avium*, but perhaps not the
only factor. Husbandry changes, such as breeding birds at lower density, may be
effective in reducing exposure to shed virus. Until now, vaccine development and
administration has been prohibitively expensive, and a previous vaccination effort in the
UK was not successful at preventing disease in adult WWWDs that had presumably
been exposed to TB prior to vaccination (46). With the advent of mRNA-based vaccines
for TB (47), cost-effective vaccines may be within reach and may be successful if aimed
at TB-naive hatchlings. We suggest that a combination of genetic improvement,
improved husbandry and vaccination should be pursued to improve captive breeding
success. These efforts should be coordinated internationally, with an emphasis on
working within the native range of the species, and maximising positive conservation
outcomes in the wild.
352

MATERIALS & METHODS

DNA samples

Our sampling design included captive, India-derived birds from SHBP, and wild birds
from Indonesia. Captive birds were sampled as part of routine health checks or during
necropsies performed by veterinary staff at SHBP. These samples represent two time
periods that we label "historical" and "current." The current captive sample comprised ten
individuals (six males and four females) hatched between 2008 and 2014. The historical
captive sample included nine birds that were blood-sampled at Sylvan Heights in 1994.
Among the birds in the historical sample, those with known studbook numbers hatched
between 1985 and 1990 at three locations in North America; SHBP, NC; Goodewood
Game Bird Farm, AL; and St. Louis Zoo, MO; all were subsequently moved to SHBP.
Three individuals in the historical sample could not be associated with their
corresponding studbook number and consequently have limited historical information.
366
Feather samples were collected from 11 wild WWWDs in Lampung Province, Indonesia,
primarily in Way Kambas National Park on the southern tip of the island of Sumatra,
between June 1999 and February 2001 by XXX. Samples were exported from Indonesia
with permission from the Directorate General of Forest Protection and Nature
Conservation in the Department of Forestry, and imported to the US under CITES permit
01US042484/9.

DNA extraction, sequencing and read mapping

The ten current captive samples included DNA extracted from blood, liver, lung, and spleen. Nine current WWWD samples were extracted in 2017, with a Qiagen DNeasy Blood & Tissue kit. Historical captive WWWD DNA samples were extracted from blood samples collected at SHBP; standard phenol-chloroform extractions were completed by XXX in 1995 and stored in a -80ºC freezer before being sent to XXXXXXXXXX in 2019. The 11 wild WWWD feather samples collected by XXX were sent to XXX in 2003; DNA extraction from feathers was completed using the Qiagen DNeasy Tissue and Blood kit with the addition of 30 µl of 100 mg/ml dithiothreitol to digest the feather keratin; these extracts were preserved in a -80ºC freezer until being sent to XXXXXXXXXX in 2018 for whole genome sequencing.

Due to project funding and sample availability, samples were sequenced in two batches. Fragment libraries for a first batch of seven current samples were prepared with an Illumina TruSeq PCR-free kit and sequenced at the University of Illinois Roy J. Carver Biotechnology Center on an Illumina HiSeq 4000 instrument. A second batch of 22 birds, including the remaining current, historical and wild samples, were also sequenced at the University of Illinois. These libraries were prepared with a Hyper Library construction kit from Kapa Biosystems and sequenced on a NovaSeq 6000.

For most of the population genomic analyses reported here, WWWD data were aligned to the highly contiguous tufted duck genome (*Aythya fuligula* GCF_009819795.1, contig N50=17.7Mb), but we also mapped the data to the short-read-based WWWD genome assembly (GCA_013398475.1, contig N50=91.2kb (14)). Reads were aligned to both reference genomes using *bwa mem* v0.7.17(default settings) (48). Alignment success was similarly high for both reference genomes (WWWD reference: historical=99%, current=99%, wild=94%; tufted reference: historical=99%, current=99%, wild=95%). We favoured the tufted duck-aligned dataset in most downstream analyses because of our interest in multi-copy immune gene diversity, the annotations for which are better in contiguous genomes with high Contig N50 (49, 50). Depending on the given analysis we used either called genotypes from *GATK* or genotype likelihoods inferred by *ANGSD* v0.911 (51) and using the *ANGSD-wrapper (52)*. We estimated genotype likelihoods in *ANGSD* with the following parameters: -doGLF 3 -GL 1 -doMaf 1 -SNP_pval 1e6 -doMajorMinor 1 -minMapQ 30 -minQ 20 (51). Called genotypes, based on depth-normalized aligned data, were obtained using *bcftools* (v 1.13) *mpileup* and *bcftools call -m -f GQ (53)*, and included invariant sites. VCF files were filtered using *vcftools* v0.1.15 to include only sites that were present in 70% of individuals, and excluded sites with minDP<5 and max-meanDP>50. Both approaches yielded comparable results in downstream population genetic analyses, such as estimates of inbreeding and population divergence (not shown).

Reconstruction of historical demography

We used *PSMC* (Li & Durbin 2011) to compare the demographic histories of the sampled WWWD populations. Demographic estimates from *PSMC* are impacted by variation in coverage, with lower coverage samples missing variants. Our captive birds

had lower depth than our wild birds, so for these analyses we used our two highest depth captive birds and downsampled all the wild birds to match. This procedure will tend to reduce estimates of *Ne*, so our *PSMC* findings provide relative estimates of effective population size and population trajectory. For this analysis, we aligned trimmed reads to the reference genome using *bowtie v*2.5.4 and *–very-sensitive* options, we used *Picard* v3.1.1 *MarkDuplicates* to remove PCR duplicate reads. We then called variants for *PSMC* using *samtools mpileup* (-Q 30 -q 30 -u -v -f) for the 15 longest scaffolds and converted the output into a fastq using *vcfutils vcf2fq* (-d 4 -D 80 -Q 30). Hilgers et al (54) note a common issue in which PSMC estimates unreasonably large effective population sizes in the most recent time interval under typical settings. Following their guidance we split the first atomic time interval (-N25 -t15 -r5 -p "2+2+25*2+4+6") which generated interpretable and reasonable demographic trajectories. PSMC results were scaled using a generation time of three years and a per-generation mutation rate of 2.5×10^{-9} (55).

mtDNA assembly & phylogenetic reconstruction

To assemble the mitochondrial genomes, we aligned short read data to an existing mtDNA assembly for WWWD (GenBank accession: MN356440) and imported the resulting reads into Geneious Prime v. 2023.2.1 (https://www.geneious.com). Given the presence of nuclear copies of mtDNA sequences in related duck species (Sorenson et al. 1996), we closely examined alignments for evidence of "numts" (Sorenson & Quinn, 1998). For the feather samples from the wild population (*n* = 11), there was little evidence of numts and substantial coverage of the mtDNA (>800x), allowing inference of complete, high confidence mtDNA sequences for all samples. Results were more variable for blood samples from the captive population (*n* = 19 including the sample used for the reference genome), with coverage ranging from 22x to over 500x. Divergent numt reads were evident in all samples, but represented a decreasing proportion of reads as total coverage increased. Thus, complete mtDNA sequences could be assembled for those blood samples that yielded relatively high coverage of the mtDNA; this included 1 historical sample and seven current samples with total coverage of 85x or more. To provide an estimate of the timing of divergence of mtDNA lineages, we augmented the WWWD data with available waterfowl mtDNA genomes (see Figure Sx) and completed a phylogenetic analysis in BEAST (v. 2.7.7(56)) using the 12 light-strand encoded mt protein-coding genes. We partitioned the data by codon position, employed model averaging and an optimized relaxed clock (Bouckaert & Drummond 2017; Bouckaert et al. 2019). Following Mitchell et al. (57) and using identical parameters values, we time-calibrated the phylogeny by setting a lognormal prior on the age of the dabbling duck lineage based on the middle-Miocene fossil *Anas soporata* (58).

Genetic diversity, inbreeding and differentiation

Using called genotypes, we estimated individual autosomal heterozygosity in each population using *vcftools* (--het) (59). We estimated nucleotide diversity (π) for each of the three populations and pairwise F_{ST} among populations using 100kb windows in *Pixy v2.0.0.beta6* (60). These analyses were based on VCF files with invariant sites included.

We used *ANGSD* and ANGSD-wrapper (51, 52) to visualize genetic differentiation
with PCA and estimate inbreeding from genotype likelihoods. We estimated individual
inbreeding coefficients using the expectation maximization algorithm in the ngsF
module(61). We used *PCangsd* to represent genetic data as principal components (17).

Finally, we estimated runs of homozygosity in the genome using *ROHan v1.0.1* (62)
directly from individual alignment files. We restricted the analysis to tufted duck
autosomes longer than 10Mb and used the following parameters: rohmu=1×10^{-3}, and
window size=500kb. The genomic inbreeding coefficient (F_{ROH}) was calculated as
$\sum L_{ROH}/L_{autosomes}$ where $\sum L_{ROH}$ is the total length of all autosomal ROHs in an individual
and $L_{autosomes}$ is the total length of the autosomal genome used in the analysis (63–65).

For captive birds only, we tested for associations between lifespan and observed
heterozygosity, individual inbreeding coefficients and F_{ROH} using a gaussian generalized
linear model in the R package *glmmTMB* and model checks in *DHARMa* (66, 67). We
compared two models, one which included a parameter for historical versus current
population and one with the genetic measure alone. Models were compared using AIC.
Models and all plotting were conducted in *R* (V 4.4.0)(68)

Major Histocompatibility Complex

We characterized diversity in functionally relevant peptide binding regions (PBR) of
MHC classes Iα and IIβ. Our ability to accurately assess heterozygosity and MHC copy
number may have been influenced by different factors depending on the reference
genome we used. The WWWD genome assembly is likely incomplete for MHC genes,
whereas the tufted duck genome may be divergent in copy number, such that some
copies may have insufficient sequencing depth for accurate SNP calling (Figure S7).
Therefore, we used two approaches to characterize MHC diversity in WWWDs. We first
re-annotated the tufted duck genome using previously published consensus sequences
of exons 2-4 derived from Anatidae MHC Class I and II PBRs (i.e., a total of six exons)
(50). We extracted putative matching sequences with e-values <1e10 using *tblastn v2.12*
(69), then extracted the hit with the lowest e-value when hits overlapped. We used
Geneious Prime 2022 to visualize these sequences in the genomic context and checked
for premature stop codons. We considered sequential copies of exons 2-4 as a complete
copy. Using this method, we estimated that the tufted duck has nine copies of Class I
and three copies of Class II, whereas the WWWD genome assembly has two and one
copies, respectively. Class I estimates differ from the tufted duck NCBI annotation, which
includes three genes, as many annotations contain multiple PBRs. Secondly, to ensure
we did not underestimate diversity, we extracted a representative gene (all exons and
introns) from the tufted duck genome and re-mapped all previous MHC-aligning depth-
normalized reads to the single copies representing Class I and Class II, respectively.
Using *samtools depth*, we assessed potential copy number by comparing the mean
aligned read depth at the single copy MHC against the average depth of reads mapped
to the tufted duck chromosome that contains MHC, chromosome 33. Copy number was
assessed by dividing the average depth of MHC reads aligned to each MHC class and
dividing by the average number of reads for chromosome 33.

Using this alignment, we called MHC SNPs for each class separately using *GATK*
HaplotypeCaller v4.2.0(70, 71). Because multiple MHC copies were aligned to a single
pseudoreference we doubled the copy number and used this number with the *--ploidy*
flag. Then we filtered the VCF with the following expression: "QD < 2.0 || FS > 60.0 || MQ
< 40.0 || MQRankSum < -12.5 || ReadPosRankSum < -8.0" (72). For this dataset, we
estimated both nucleotide and amino acid diversity in PBR exons using custom code in
R. For data aligned to the genome, we calculated nucleotide diversity using *pixy,* and
calculated individual heterozygosity directly from genotypes in PBR exons.

RNA samples

We sampled blood from eight WWWDs in 2022, and seven redhead ducks (*Aythya*
americana) to provide a comparison to a closely related species that is less susceptible
to avian TB. During the sampling period in 2022, two WWWDs were symptomatic for
mycobacteriosis, whereas none of the redhead ducks were symptomatic. We further
included two opportunistic samples from 2019: one redhead blood sample with
mycobacteriosis, and a paired sample without (total redhead sample size=9). All
instances of mycobacteriosis were fatal, and one WWWD was sampled approximately
two hours post-mortem. This individual's death was unexpected as it did not show
outward symptoms of mycobacteriosis, but necropsy revealed typical mycobacterial
lesions on its organs.

Birds were sampled at Sylvan Heights Bird Park using a 23 gauge needle and
syringe to draw blood from the brachial vein. Whole blood was immediately placed in
RNAlater®, and stored on ice until it was placed in a -80ºC freezer. All animal
procedures were reviewed and approved by XXXXXXXXXX's IACUC under Animal Use
Protocol D351.

RNA extraction, sequencing & read mapping

527 We extracted RNA from blood samples using TriZol and Qiagen RNeasy Kit, and all
528 samples had RIN>7. Libraries were prepared with TruSeq® Stranded mRNA Library
529 Prep, and sequenced on NextSeq 2000 P2 Reagents (200 cycles).
530
We generated 28.4-42.2 million paired-end reads per sample, retaining 89-97% of
reads after trimming. We mapped RNAseq reads to the tufted duck genome (*Aythya*
fuligula, GCF_009819795.1) using *STAR* v2.7.11b after trimming with *fastp* v0.23.2 (73,
74). Of these reads, 84-89% were uniquely mapped to the tufted duck genome (Figure
S9). Reads mapping to each gene were counted using the *geneCounts* function in
STAR.

Gene expression

We filtered counts by removing genes with counts of ≤10 in at least six individuals.

We identified three "potentially influential" individuals using Principal Component

Analysis (using *PCAtools* (75)); these individuals had the smallest library sizes (Figure

S9-S10). However, two of these three individuals represented two-thirds our disease
sample, and so we used a conservative approach to remove the outlier effect from our
list of differentially expressed genes (see below) We explored differences in normalized
gene expression using a negative binomial generalized linear model framework
implemented in *DESeq2* v1.44.0. First, we compared gene expression between
asymptomatic redhead and white-winged ducks (gene expression ~ species), using
seven redhead samples (excluding one outlier individual) and six WWWD samples.
Then, we compared diseased vs asymptomatic birds in each species separately (gene
expression ~ disease status), and compared the disease-related expression patterns
between species. To identify differentially expressed genes (DEGs), we applied the Wald
Test and considered genes to be significant with an FDR-corrected p-value (q) < 0.05.
Due to the small sample sizes of diseased individuals, we removed significant genes that
appeared to be affected by the outliers, rather than disease status, using custom code.
We refer to the FDR-corrected p-value of genes that passed this additional filter as q*.

Susceptibility in WWWD ducks is likely linked to immune gene regulation or function,
and thus we focused on immune DEGs. We extracted immune-related genes using a
previously published list of immune GO terms (76), using the tufted duck GO annotations
available on Genbank. We supplemented the immune gene list by including previously
annotated MHC genes and searching the tufted duck reference annotations for genes
relating to B-cells, T-cells, TLRs, immunoglobulin, interleukins, and interferons. We
tested whether our candidate gene list was enriched in DEGs using the hypergeometric
test function *phypher()* in R. To compare gene expression in the difficult multi-copy MHC,
we summed counts from MHC Class Iα and MHC Class IIβ paralogs into a single "gene"
to include in the analysis, but most reads mapped to a single copy in the tufted duck
genome so this did not change the signal.

Then, we explored enrichment of GO Biological Processes and KEGG pathways
using *clusterProfiler* V4.12.6 (77). We considered the full DEG list (q<0.05) as the
foreground and the whole blood transcriptome as the background. For KEGG
enrichment, we converted the NCBI gene-ids to functional ortholog ids ("ko") using the
KEGGREST v1.44.1 package (78), and used *enrichKEGG()* to identify enriched
pathways. We excluded "significant" enrichments that included ≤ two genes.

Data Availability Statement

573 Raw DNA sequence data are available under PRJNA1258233 (Table S1), raw gene
574 expression data is available under PRJNA1285963 (Table S4). All data and code are
575 provided in Dryad (https://doi.org/10.5061/dryad.crjdfn3ht).

References

1. J. D. Ballou, Assessing the Risks of Infectious Diseases in Captive Breeding and
Reintroduction Programs. *J. Zoo Wildl. Med.* **24**, 327–335 (1993).

2. T. van der Valk, L. Dalèn, From genomic threat assessment to conservation action.
Cell **187**, 1038–1041 (2024).

3. E. B. Linck, C. D. Cadena, A latitudinal gradient of reference genomes. *Mol. Ecol.*
e17551 (2024). https://doi.org/10.1111/mec.17551.

4. A. Choudhury, A Tryst With The White-Winged Wood Duck. *Sanctuary Asia* **43**
(2023).

5. A. Choudhury, The status of endangered species in northeast India. *J. Bombay Nat.*
Hist. Soc. **103**, 157–167 (2006).

6. A. Green, The status and conservation of the White-winged Wood Duck *Cairina*
scutulata. *International Waterfowl and Wetlands Research Bureau* 17 (1992).

7. P. K. Saikia, M. K. Saikia, Present distribution, status, and ecology of white-winged
wood duck and hornbills in nameri National Park, considering the tropical forest.
Zoos' Print **11** (2011).

8. BirdLife International, Species factsheet: White-winged Duck *Asarcornis scutulata*.
(2024). Available at: https://datazone.birdlife.org/species/factsheet/white-winged-
duck-asarcornis-scutulata [Accessed 11 April 2025].

9. M. J. S. Mackenzie, J. & Kear, The White-winged Wood Duck. *Wildfowl* **27**, 5–17
(1976).

10. A. Green, Progress in the White-winged Wood Duck *Cairina scutulata* Action Plan
Project: A call for information. *Wildfowl* **41**, 161–162 (1990).

11. A. Richardson, Aviculture at WWT over half a century. *Wildfowl* **47**, 264–272 (1996).

12. R. E. Cromie, M. J. Brown, J. L. Stanford, The epidemiology of avian tuberculosis
inWhite-winged wood ducks *Cairina scutulata* at The Wildfowl and Wetlands
Trust,Slimbridge Centre (1976-91). *Wildfowl* **43**, 211–214 (1992).

13. K. Cook, *Population Analysis and Breeding and Transfer Plan, White-winged*
WoodDuck AZA Species Survival Plan Yellow Program. AZA Population
Management Center (2016).

14. S. Feng, *et al.*, Dense sampling of bird diversity increases power of comparative
genomics. *Nature* **587**, 252–257 (2020).

15. R. C. Mueller, *et al.*, A high-quality genome and comparison of short- versus long-
read transcriptome of the palaearctic duck Aythya fuligula (tufted duck). *Gigascience*
10 (2021).

16. S. Y. W. Ho, M. J. Phillips, A. Cooper, A. J. Drummond, Time dependency of
molecular rate estimates and systematic overestimation of recent divergence times.
Mol. Biol. Evol. **22**, 1561–1568 (2005).

17. J. Meisner, A. Albrechtsen, Inferring population structure and admixture proportions

in low-depth NGS data. *Genetics* **210**, 719–731 (2018).

18. H. Li, R. Durbin, Inference of human population history from individual whole-
genome sequences. *Nature* **475**, 493–496 (2011).

19. D. Díez-Del-Molino, F. Sánchez-Barreiro, I. Barnes, M. T. P. Gilbert, L. Dalén,
Quantifying temporal genomic erosion in endangered species. *Trends Ecol. Evol.*
33, 176–185 (2018).

20. K. D. Witt, Role of MHC class I pathways in Mycobacterium tuberculosis antigen
presentation. *Front. Cell. Infect. Microbiol.* **13**, 1107884 (2023).

21. P. Bettencourt, *et al.*, Identification of antigens presented by MHC for vaccines
against tuberculosis. *NPJ Vaccines* **5**, 2 (2020).

22. C. L. Witte, L. L. Hungerford, R. Papendick, I. H. Stalis, B. A. Rideout, Investigation
of characteristics and factors associated with avian mycobacteriosis in zoo birds. *J.*
Vet. Diagn. Invest. **20**, 186–196 (2008).

23. K. P. Johnson, M. D. Sorenson, Phylogeny and biogeography of dabbling ducks
(genus: Anas): A comparison of molecular and morphological evidence. *Auk* **116**,
792–805 (1999).

24. J. Gonzalez, H. Düttmann, M. Wink, Phylogenetic relationships based on two
mitochondrial genes and hybridization patterns in Anatidae. *J. Zool. (1987)* **279**,
310–318 (2009).

25. F. E. Rheindt, *et al.*, AviList: a unified global bird checklist. *Biodivers. Conserv.*
(2025). https://doi.org/10.1007/s10531-025-03120-y.

26. W. Blyth, *Catalogue of the birds in the Museum Asiatic Society* (Baptist Mission
Press, 1849).

27. A. O. Hume, C. H. T. Marshall, *Game birds of India, Burma and Ceylon* (Hume and
Marshall, 1880).

28. A. J. Green, The biology of the White-winged Duck *Cairina scutulata*. *Forktail* **8**, 65–
82 (1993).

29. J. B. Bemmels, *et al.*, Demographic decline and lineage-specific adaptations
characterize New Zealand kiwi. *Proc. Biol. Sci.* **288**, 20212362 (2021).

30. S. Li, *et al.*, Genomic signatures of near-extinction and rebirth of the crested ibis and
other endangered bird species. *Genome Biol* **15**, 557 (2014).

31. J. A. Robinson, *et al.*, Genomic flatlining in the endangered island fox. *Curr. Biol.* **26**,
1183–1189 (2016).

32. L. Husson, F. C. Boucher, A.-C. Sarr, P. Sepulchre, S. Y. Cahyarini, Evidence of
Sundaland's subsidence requires revisiting its biogeography. *J. Biogeogr.* **47**, 843–
853 (2020).

33. A. Khan, *et al.*, Genomic evidence for inbreeding depression and purging of
deleterious genetic variation in Indian tigers. *Proc. Natl. Acad. Sci. U. S. A.* **118**,
e2023018118 (2021).

34. J. Louys, P. Roberts, Environmental drivers of megafauna and hominin extinction in
Southeast Asia. *Nature* **586**, 402–406 (2020).

35. R. Karjee, *et al.*, Species biology and demographic history determines species
vulnerability to climate change in tropical island endemic birds. (2025).

36. T.-T. Gu, *et al.*, Genomic analysis reveals a cryptic pangolin species. *Proc. Natl.*
Acad. Sci. U. S. A. **120**, e2304096120 (2023).

37. J. von Seth, *et al.*, Genomic insights into the conservation status of the world's last
remaining Sumatran rhinoceros populations. *Nat. Commun.* **12**, 2393 (2021).

38. E. Humble, *et al.*, Conservation management strategy impacts inbreeding and
mutation load in scimitar-horned oryx. *Proc. Natl. Acad. Sci. U. S. A.* **120**,
e2210756120 (2023).

39. D. J. Newhouse, C. N. Balakrishnan, High major histocompatibility complex class I
polymorphism despite bottlenecks in wild and domesticated populations of the zebra
finch (Taeniopygia guttata). *BMC Evol. Biol.* **15**, 265 (2015).

40. M. Stervander, E. G. Dierickx, J. Thorley, M. de L. Brooke, H. Westerdahl, High
MHC gene copy number maintains diversity despite homozygosity in a Critically
Endangered single-island endemic bird, but no evidence of MHC-based mate
choice. *Mol. Ecol.* **29**, 3578–3592 (2020).

41. Y. V. Skvortsova, O. S. Bychenko, T. L. Azhikina, Role of RIG-I-like receptors in the
activation of innate immunity in tuberculosis. *Russ. J. Bioorganic Chem.* **49**, 742–
750 (2023).

42. D. P. Simmons, *et al.*, Mycobacterium tuberculosis and TLR2 agonists inhibit
induction of type I IFN and class I MHC antigen cross processing by TLR9. *J.*
Immunol. **185**, 2405–2415 (2010).

43. M. D. Saggese, *et al.*, Gross and microscopic findings and investigation of the
aetiopathogenesis of mycobacteriosis in a captive population of white-winged ducks
(Cairina scutulata). *Avian Pathol.* **36**, 415–422 (2007).

44. B. Liu, Y. Shao, R. Fu, Current research status of HLA in immune-related diseases.
Immun. Inflamm. Dis. **9**, 340–350 (2021).

45. E. A. Eskew, D. Fraser, M. J. Vonhof, M. L. Pinsky, B. Maslo, Host gene expression
in wildlife disease: making sense of species-level responses. *Mol. Ecol.* **30**, 6517–
6530 (2021).

46. R. Cromie, "Development of an avian tuberculosis vaccine for wildfowl," University
College London. (1991).

47. B. J. B. Touray, M. Hanafy, Y. Phanse, R. Hildebrand, A. M. Talaat, Protective RNA
nanovaccines against Mycobacterium avium subspecies hominissuis. *Front.*
Immunol. **14**, 1188754 (2023).

48. H. Li, Aligning sequence reads, clone sequences and assembly contigs with BWA-
MEM. *arXiv [q-bio.GN]* (2013).

49. R. J. Driver, C. N. Balakrishnan, Highly contiguous genomes improve the
understanding of avian olfactory receptor repertoires. *Integr. Comp. Biol.* **61**, 1281–
1290 (2021).

50. K. He, P. Minias, P. O. Dunn, Long-read genome assemblies reveal extraordinary
variation in the number and structure of MHC loci in birds. *Genome Biol. Evol.* **13**
(2021).

51. T. S. Korneliussen, A. Albrechtsen, R. Nielsen, ANGSD: Analysis of next generation
sequencing data. *BMC Bioinformatics* **15**, 356 (2014).

52. A. Durvasula, *et al.*, Angsd-wrapper: Utilities for analysing next-generation
sequencing data. *Mol. Ecol. Resour.* **16**, 1449–1454 (2016).

53. H. Li, A statistical framework for SNP calling, mutation discovery, association
mapping and population genetical parameter estimation from sequencing data.
Bioinformatics **27**, 2987–2993 (2011).

54. L. Hilgers, *et al.*, Avoidable false PSMC population size peaks occur across
numerous studies. *Curr. Biol.* **35**, 927–930.e3 (2025).

55. S. Singhal, *et al.*, Stable recombination hotspots in birds. *Science* **350**, 928–932
(2015).

56. R. Bouckaert, *et al.*, BEAST 2.5: An advanced software platform for Bayesian
evolutionary analysis. *PLoS Comput. Biol.* **15**, e1006650 (2019).

57. K. J. Mitchell, J. R. Wood, R. P. Scofield, B. Llamas, A. Cooper, Ancient
mitochondrial genome reveals unsuspected taxonomic affinity of the extinct
Chatham duck (Pachyanas chathamica) and resolves divergence times for New
Zealand and sub-Antarctic brown teals. *Mol. Phylogenet. Evol.* **70**, 420–428 (2014).

58. N. V. Zelenkov, E. N. Kurochkin, Dabbling ducks (Aves: Anatidae) from the middle
Miocene of Mongolia. *Paleontol. J.* **46**, 421–429 (2012).

59. P. Danecek, *et al.*, The variant call format and VCFtools. *Bioinformatics* **27**, 2156–
2158 (2011).

60. K. L. Korunes, K. Samuk, pixy: Unbiased estimation of nucleotide diversity and
divergence in the presence of missing data. *Mol. Ecol. Resour.* **21**, 1359–1368
(2021).

61. F. G. Vieira, M. Fumagalli, A. Albrechtsen, R. Nielsen, Estimating inbreeding
coefficients from NGS data: Impact on genotype calling and allele frequency
estimation. *Genome Res.* **23**, 1852–1861 (2013).

62. G. Renaud, K. Hanghøj, T. S. Korneliussen, E. Willerslev, L. Orlando, Joint
estimates of heterozygosity and runs of homozygosity for modern and ancient
samples. *Genetics* **212**, 587–614 (2019).

63. F. C. Ceballos, P. K. Joshi, D. W. Clark, M. Ramsay, J. F. Wilson, Runs of
homozygosity: windows into population history and trait architecture. *Nat. Rev.*
Genet. **19**, 220–234 (2018).

64. C.-Z. Fu, X.-M. Guang, Q.-H. Wan, S.-G. Fang, Genome resequencing reveals
congenital causes of embryo and nestling death in crested ibis (Nipponia nippon).
Genome Biol. Evol. **11**, 2125–2135 (2019).

65. R. McQuillan, *et al.*, Runs of homozygosity in European populations. *Am. J. Hum.*
Genet. **83**, 359–372 (2008).

66. M. E. Brooks, *et al.*, glmmTMB Balances Speed and Flexibility Among Packages for
Zero-inflated Generalized Linear Mixed Modeling. *The R Journal* **9:2** (2017).

67. F. Hartig, *DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)*
Regression Models (2024).

68. R Core Team, R: A language and environment for statistical computing.
https://www.R-project.org/ (2025).

69. C. Camacho, *et al.*, BLAST+: architecture and applications. *BMC Bioinformatics* **10**,
421 (2009).

70. A. McKenna, *et al.*, The Genome Analysis Toolkit: a MapReduce framework for
analyzing next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303
(2010).

71. R. Poplin, *et al.*, Scaling accurate genetic variant discovery to tens of thousands of
samples. *bioRxiv* (2017).

72. D. Caetano-Anolles, Hard-filtering germline short variants. *GATK Technical*
Documentation (2024). Available at: https://gatk.broadinstitute.org/hc/en-
us/articles/360035890471-Hard-filtering-germline-short-variants.

73. A. Dobin, *et al.*, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21
(2013).

74. S. Chen, Y. Zhou, Y. Chen, J. Gu, fastp: an ultra-fast all-in-one FASTQ
preprocessor. *Bioinformatics* **34**, i884–i890 (2018).

75. K. Blighe, A. Lun, *PCAtools: PCAtools: Everything Principal Components Analysis*
(2024).

76. L. W. Silver, *et al.*, A targeted approach to investigating immune genes of an iconic
Australian marsupial. *Mol. Ecol.* **31**, 3286–3303 (2022).

77. G. Yu, L.-G. Wang, Y. Han, Q.-Y. He, clusterProfiler: an R package for comparing
biological themes among gene clusters. *OMICS* **16**, 284–287 (2012).

78. D. Tenenbaum, *KEGGREST* (Bioconductor, 2017).

79. G. O. User, Occurrence Download. The Global Biodiversity Information Facility.
https://doi.org/10.15468/DL.RJ8D95. Deposited 25 October 2024.

Figure 1: Endangered white-winged wood ducks have experienced population declines in the US captive breeding program, and are genetically differentiated from wild ducks.

A) White-winged wood ducks (*Asarcornis scutulata*) at Sylvan Heights Bird Park, North Carolina. Photo: Katie Lubbock. B) Time series showing US captive population size of male (black lines) and female (grey lines) white-winged wood ducks (left y-axis). Colored rectangles indicate the hatch years for captive birds, and sampling period for wild birds for the genetic samples used in this study (x-axis) and their height indicates the number of samples used in this study (right y-axis). Three additional wild birds were included in mtDNA phylogeny. C) Distribution of white-winged wood ducks in south-east Asia from Global Biodiversity Information Facility (GBIF) records (79). Black points indicate museum specimens, whereas grey indicate human observations such as from eBird. Triangles show observations prior to the year 2000 indicating likely extinction in Java and Malay Peninsula. Observations in central northern Thailand are from a reintroduction program [7]. Large colored circles indicate the approximate locations of the founders of the captive population in Assam (dark blue), and the feather samples taken from wild

790 ducks in Sumatra (light blue). Inset: Maximum parsimony mitochondrial DNA phylogeny
791 of wild and captive birds, where scale is the number of changes along each branch. D)
792 Principal components analysis of autosomal variants for individuals used in this study,
793 color coded by population.

Figure 2: Populations of white-winged wood ducks from India and Indonesia have declined during the Pleistocene, and have experienced ongoing inbreeding and inbreeding depression.

A) Estimates of effective population size through time from PSMC for wild (Indonesia) and historical (India) captive ducks. Grey shaded box shows the late Pleistocene period (11.7-115kya), with years scaled by generation time of three years and a per-generation mutation rate of 2.5×10^{-9} (55). B) Inbreeding coefficients calculated from runs of homozygosity (ROH) in genomic size-classes per individual. C) Inbred birds in captivity typically live shorter lives (see also Table S2 and Figure S4-S6).

Figure 3: White-winged wood ducks in captivity have lost nucleotide and amino acid diversity in Major Histocompatibility Complex genes, and show potential immune gene dysregulation.

A-D) Diversity of peptide binding exons in the Major Histocompatibility Complex (MHC). Estimates based on aligning individual data to single exon sequences for each class. A) Per-site heterozygosity, B) Nucleotide diversity, C) Number of variable amino acids, D) Pairwise amino acid diversity. Reductions in MHC Class IIB diversity are less dramatic when aligned to the tufted duck genome (Figure S7). E) MHC Class I and other key genes in the Class I antigen presentation pathway are significantly upregulated in WWWD compared to non-susceptible RH ducks. F) MHC Class II is not differentially expressed between these species. G) Among other differentially expressed immune genes, toll-like receptor 2 is downregulated in WWWD.