Octopamine and Tyramine Modulate Motor Program Competition in *Drosophila* larvae

William V. Smith[1] and Stefan R. Pulver[1][¶]

[1] School of Psychology and Neuroscience, University of St Andrews, St Mary's Quad, South Street, St Andrews KY16 9JP, UK

[¶] Corresponding author: sp96@st-andrews.ac.uk

Abstract

Dynamic Interactions amongst competing motor programs shape behavioural output. Here, we use a combination of opto- and electro-physiology to explore how the tdc^{2+} adrenergic-like system modulates competition amongst central pattern generating (CPG) networks controlling locomotion in *Drosophila* larvae. Bath application of octopamine (OA) promoted fictive forwards locomotion, suppressed backwards locomotion, and induced bouts of fictive head sweeps during wash period that were proportional to the promotion of forward waves. In contrast, tyramine (TA), the conceptualised inhibitory antagonist to OA, promoted collisions and overlap of motor programmes, activity-dependent inhibition, and bouts of silence during wash periods. Dual-colour calcium imaging of the tdc^{2+} system with motor neurons revealed that the tdc^{2+} neurons are recruited phasically prior to a majority of fictive locomotor behaviours. Optogenetic manipulation of the tdc^{2+} system recapitulated a subset of effects and further revealed that activity in tdc^{2+} system neurons necessary and sufficient for locomotor CPG activity in the system. Overall, our work generates testable predictions for computational and connectomics studies for *Drosophila* motor competition and provides insights into how adrenergic-like systems can modulate dynamics of motor competition among interacting CPGs within locomotor networks.

Introduction

Appropriate locomotion requires both stability and flexibility in neural networks. Locomotion is engendered by not just program initiation but the inhibition of competing motor programs [1]. To balance the need for stability and flexibility, motor systems need to enable appropriate initiation and persistence of locomotion without impairing the capacity to induce alternative motor programs. Given the essential nature of motor adaptability to organismal survival (e.g., escape responses), understanding the general principles of how motor systems manage modulation to motor competition may provide a universal paradigm to examine fundamental network output flexibility. While the role of circuit motifs in specific motor selection and coordination are gradually becoming better understood (e.g. [2]), how neuromodulation underpins competitive selection of motor programs is relatively poorly understood.

The adrenergic-like system in *Drosophila* provides a powerful testbed to evaluate motor competition by enabling controlled, acute modulation to motor bias. The *Drosophila* adrenergic-like system is composed of octopamine (OA) - an invertebrate analogous to adrenaline - and tyramine (TA) [3]. The biosynthetic pathways of OA/TA, their receptor pharmacology [4], [5], and the expression patterns of OA and TA neurons are well studied in multiple invertebrate species [4], [6], [7], [8], [9], [10]. The adrenergic-like system has been implicated in a variety of fundamental behaviours including learning

and memory[11], [12], social behaviours like aggression[13], [14], [15], motor initiation and

modulation [9], [16], [17], [18], [19], including fundamental processes underlying action like energy

metabolism [20], [21], synaptic plasticity [22], [23], and activity-mediated adaptation [24], [25]. The

adrenergic-like system is thus a crucial, multifaceted, and complex system consequential to

invertebrate movement and life, especially in its capacity to modulate selection of motor programs.

Traditionally, OA and TA are considered antagonistic neuromodulators [9], [26], [27]. Namely, OA has

a well-documented promotive and facilitatory role in locomotion [9], [16], [17], [18], [27], [28], [29],

[30], [31], [32] while TA appears to have an overall inhibitory effect on locomotion [6], [8], [27], [33],

[34] across invertebrates. However, the functional relationship between OA and TA's action is not

always exclusively mutually antagonistic nor definitely excitatory/inhibitory. For instance, the mutual-

antagonistic relationship of OA/TA appears to be conditional on their mutual relative concentrations

[26]. Further, OA and TA are not always excitatory or inhibitory, respectively. For example, octopamine

can have inhibitory effects on CPG activity through inhibition of CPG command interneurons[35], can

inhibit nerve-based locomotion stimulation [36], and can inhibit synaptic transmission at the

Drosophila larval neuromuscular junction [37]. Indeed, different OA receptors are linked to either

stimulatory (Octβ2R) or inhibitory (Octβ1R) cAMP/CREB pathways which putatively self-regulates

octopaminergic synaptogenesis of boutons at the larval neuromuscular junction [23]. On the other

hand, TA can have excitatory effects on neural networks underpinning behaviour. For example, TA can

evoke depolarisation through an indirect DA-mediated pathway in rat subthalamic neurons [38], at

low concentrations in hyperpolarized states increase the amplitude of locust oviduct excitatory

junction potentials stimulating rhythmic contractions [39], and facilitate sensitisation by delaying

cocaine-induced responsive hypoactivity [40]. Such research underscores how OA/TA

neuromodulation is not exclusively one extreme in a binary view of neuromodulation.

Nonetheless, a significant body of work points to a binary perspective on OA/TA within locomotory

systems. One cause of this perspective may be that a majority of prior studies on adrenergic-like

modulation have utilised powerful genetic tools (e.g., RNAi, genetic ablation) that often induces long-

term changes in the neuromodulatory environments and likely induce undetermined compensation

processes during development [41]. Given the highly adaptive, redundant, and complex nature of

neural systems, a diverse array of system configurations and features can be somewhat divorced from

their ecological contexts [42], [43], [44]. Consequently, chronic genetic manipulations may induce

outcomes with complicated interpretations. In contrast, acute modulation of adrenergic-like signalling

gives insights into effects that occur moment-to-moment and are not confounded by compensation

due to changes in gene expression. Given the high level of characterisation of the adrenergic-like

system in *Drosophila* [45] and the transferability of larval fictive activity to understanding intact

locomotion [46], [47], we opted to evaluate the effects of OA/TA on motor competition within CPG

networks in *Drosophila* 3rd instar larvae.

A crucial element of movement is action selection and the inhibition of competing motor programs.

Appropriate movement is a product of both appropriately sustained driving of CPGs as well as

precisely timed and coordinated inhibition of pre-motor networks which drive opposing CPG activity

patterns. Indeed, the failure of inhibition between CPG networks may be a direct contributor to

uncoordinated and dysrhythmic movement [1]. The role of the adrenergic-like system in modulating

action selection and motor pattern competition has been extensively studied in *C. elegans* through

tyraminergic RIM neurons [48], [49], [50] which has instantiated a fundamental adrenergic-like motif

(i.e., "master-and-slave") in *C. elegans* locomotion. However, while the role of OA/TA has been studied

in *Drosophila* with reference to specific program motor initiation and coordination [51], limited work

has been conducted on their coordinated role to generalisable motor competition. Through studying

the effects of acute modulation on action selection and motor competition, we may be able to gain a

more robust, nuanced, and applicable understanding of OA/TA neuromodulatory antagonism.

Here, we explore the role of the tdc^{2+} system modulating fictive motor competition in *Drosophila*

larvae to better understand the neural architecture behind CPG coordination. Firstly, we evaluate the

intrinsic and modulatory activity of tdc^{2+} system neurons on isolated central pattern generators in 3rd

instar *Drosophila* larvae. Specifically, we show that activity in tdc^{2+} neurons and motor neurons is

highly correlated during fictive locomotion. Secondly, we demonstrate how bath-applied exogenous

OA and TA induce acute changes in CPG acivity during after bath application. Specifically, OA induces

elevated fictive forward locomotion in isolated central pattern generators. In contrast, exogenous TA

induces increases in motor competition reflective of increased overlapping fictive patterns with

activity-dependent inhibitory effects on general fictive activity. Both neuromodulators also induce

long lasting effects following bath application that reflect activity changes in the network. Thirdly,

through optogenetic manipulation of tdc^{2+} neuronal activity, we demonstrate how intrinsic tdc^{2+}

neuronal activity appears sufficient and necessary for producing fictive rhythms. Finally we show how

regionally restricted depolarisation of tdc^{2+} neuronal activity in the posterior abdominal isolated

ganglion is sufficient to induce fictive forward activity. Overall, we demonstrate how the adrenergic-

like system is a valuable tool for understanding the secondary effects of modulating motor

competition

Methods

Animal Rearing & Dissection

Drosophila were raised at 18-25∘C in incubators on an approximate 12-hour light-dark cycle. All experiments were conducted with feeding 3rd instar *Drosophila* larva (2-3 days posthatching). The central nervous system (CNS) of *Drosophila* was isolated as reported in [46]. Activity of isolated CNSs was recorded with Baines External Saline (BES) submersion containing (in mM) 135 NaCl, 5 KCl, 2 CaCl$_2$, 4 MgCl$_2$, 5 TES, and 36 sucrose, pH 7.15 [52]. All CNS preparations were isolated from *Drosophila* larva using dorsal incision across the larva body using fine scissors with body walls pinned into a Sylgard-lined Petri dishes using fine tungsten wires (California Fine Wire, Grover Beach, CA). All projecting CNS nerves were cut, CNS removed, posterior nerves and peripheral gastric tissue attached were pinned using fine tungsten wires (see [46]).

Dual Colour Ca^{2+} Imaging of *Tdc*$^{2+}$ and Motor Neurons

The calcium activity patterns of motor neurons and *tdc*$^{2+}$ neurons [53] were assessed in isolated CNS feeding 3rd instar *Drosophila* larva using a genetic construct that drove RCaMP1a expression in glutamatergic neurons (VGlut-LexA combined with 13XLexAOP-RCaMP1a balanced over CyO, Tb-RFP), while also containing UAS-GCaMP6f (provided by K.L. Hibbard, janelia Reserch Campus) This construct was then crossed to w[*]; P{w[+mC]=Tdc2-GAL4.) Isolated CNS *Drosophila* were imaged using an OptoSplit Imaging system (Cairn Research Ltd, Kent, UK) with appropriate excitation and emission filters for RCAMP and GCAMP imaging. Preparations were imaged using an OLA light engine (Lumencor, OR, USA) and an ORCA-Fusion CMOS camera (Hamamatsu, Hamamatsu, Japan) with a XLUMPLFLN 20X dipping objective (Olympus, Tokyo, Japan). We imaged isolated dual-colour CNS *Drosophila* with BES using an X-light spinning disk confocal with a 70µm pinhole array (Crest Optics, Rome, Italy) for up to 30 minutes at 5fps with 150ms exposure. Images were collected using MicroManager (v2.0 gamma [54]) and post-processed using ImageJ [55], [56]. The channels displaying the calcium activity for motor neurons and *tdc*$^{2+}$ neurons were overlayed using each segment's nerve roots as a reference marker. All calcium signals were extracted using area-consistent regions of interest on nerve roots projecting from ventral nerve cord segments (T3-A6) (Figure 1Aii). All calcium signals were pre-processed (df/f) and smoothed using a custom Python script (see data repositor for details) Fictive events were determined through using Hill's Valley Analysis in DataView [1] and/or SciPy's peak detection algorithm based on criteria reported in [46].

Visualisation of dual-colour calcium data was performed in DataView and Python3 using scripts available in paper-associated repositories. The overlaid average calcium activity trace in *tdc*$^{2+}$ and

motor neurons (as shown in Figure 1C) was performed using DataView's event-scope view based on

segregated fictive events detected by Hill's Valley analysis across all preparations based on criteria

reported in [46]. The average waveform of each event was calculated by DataView and extracted into

Python for across preparation visualisation. All average waveforms were determined using a duration

threshold of 8300ms. The peak time lag calculation (as shown in Figure 1D) was the difference in peaks

from the average waveform of tdc^{2+} and motor neurons in each segment using SciPy's peak detection

algorithm (see repository code). Fictive events in calcium traces were determined using Hill's Valley

analysis and manually counted as reported in [46]. Peak and duration relation between tdc^{2+} and

motor neuron calcium signals (as shown in Figure 1G) was performed using Hill's Valley detection on

each segment across all preparations with duration calculated as 20% on-off from the centre of the

calcium signal. The correlation between tdc^{2+} and motor neuron calcium signals (as shown in Figure

1E) was conducted using Pearson Cross Correlation coefficient across all preparations through

Python's Scipy.Signal package (see repository code).

Ca^{2+} Imaging with Tyramine & Octopamine Perfusion

The effect of exogenous octopamine bath application on calcium signals in motor neurons was

assessed in isolated CNS feeding 3rd instar *Drosophila* larvae using a genetic line in which GCaMP6m is

driven in glutamatergic neurons by the OK371-GAL4 line. Isolated CNS preparations were pinned to

Sylgard-lined Petri dishes as described in "Dual Colour Ca^{2+} Imaging of tdc^{2+} and Motor Neurons".

Saline and OA or TA were applied during imaging using a gravity-based perfusion system with a

consistent drip inflow across preparations and removed continuously using a KCP3-B10W peristaltic

pump (Kamoer 60-120 RPM). Across all preparations, saline was bath applied onto the preparation for

at least 10 minutes followed by 10-minute OA or TA application then < 20 minutes saline reapplication

to wash away octopamine/tyramine from the Sylgard-lined Petri dish. All OA concentration ranges (5

- 400μM) were generated from a diluted 0.02M master mix (O0250-1G Sigma Aldrich). All

TAconcentration ranges (5 - 400μM) were generated from a diluted 0.01M master mix (T90344-5G

Sigma Aldrich). Images were collected using WinFluor (University of Strathclyde, UK) at 10fps and post-

processing using ImageJ [55], [56]. All isolated preparations were recorded using a UPlanFLN Olympus

10x, 0.3nA.

Tdc^{2+} Optogenetics

The effect of endogenous tdc^{2+} activity on fictive motor activity was studied using extracellular

electrophysiology on segmental nerve roots in isolated CNS from feeding 3rd instar *Drosophila* larvae.

Octopaminergic tdc^{2+} neurons were either depolarised using lines containing 20xUAS-CsChrimson-

mVenus inserted at the attP18 landing site or hyperpolarized using lines containing UAS-GtACR1

inserted in VK00005 driven by w[*]; P{w[+mC]=Tdc2-GAL4. Across all conditions, the activity of
thoracic nerve root (T3) and posterior abdominal nerve root (A6-A8) was measured by suction
electrode recording (AM System Differential AC Amplifier Model (#59721, 1700), 1-100kHz cut-off
frequency allowance). Borosilicate glass capillaries were pulled using an electrode puller (Narishege,
Japan), and tapered tips were broken to produce suction electrodes that fit tightly around single nerve
roots. Stimulation during the tdc^{2+}-CsChrimson conditions was performed with a downward-projecting
red LED light source (Cairn Research Ltd., 620 nm, 0.58mW/cm^2). The stimulation period was either
consistent for 10, 20, or 40s or discontinuous with 0.01s pulses generated at intervals of 40Hz for ~10s
total pulsed stimulation. Stimulation during the tdc^{2+}-GtACR1 conditions was performed with upward-
projecting white light (~2.5mW/cm^2). The stimulation period was consistent for ~15, 30, or 60s
induced by manual instantiation. All whole-prep stimulation used full-prep illumination using a
UPlanFLN Olympus 10x, 0.3nA lens.

Restricted illumination stimulation conditions involved restricting the downward-projecting red light
source using a light-path restrictor with ∞0.6cm aperture and 0.16mW/cm^2 power-to-area ratio.
Restricted stimulation involved a LUMPlanF Olympus 40x, (0.8 nA) dipping lens focused on posterior
abdominal, anterior abdominal, thoracic, and suboesophageal ganglion regions (see Figure 6). Aside
from direct restricted prep stimulation, all conditions were matched with control stimulation where
illumination of the left-flank of the preparation (< 1cm) was conducted during nerve root recording.

All electrophysiology recordings and light stimulations were made using LabChart 8.0 software (AD
Instruments, Colorado Springs, CO) and post-processed in Spike2 (Cambridge Electronic Design,
Cambridge, UK) and Python3. Post-processing involved rectification and smoothing using a consistent
0.2s time interval. All rectified and smoothed traces were imported into DataView where event peaks
and 20% on-off duration were determined by Hill's Valley analysis. Visualisation and further analysis
were conducted in Python3 (see repository code).

Analysis, Visualisation, & Statistical Evaluation

211 All calcium signal and electrophysiology nerve root data were initially analysed in DataView (version
212 12.2.2)[57] using Hill's Valley Analysis for peak detection and Event-Triggered Scope View for
213 averaging waveforms. All data were visualised in Python3 using custom scripts available in the
214 repository supplied by the authors. All statistical tests were performed using packages within Python
215 (see repository scripts and report tests in the results sections).

Results

*Tdc*²⁺ VNC Neuron Activity is Correlated to Motor Neuron Activity across all Fictive Motor Programs

The isolated *Drosophila* central nervous system (CNS) spontaneously generates fictive motor rhythms in the thoracic and abdominal segments that can be measured via calcium indicators expressed in neuronal populations[46]. Previous work has characterised the features and relationship between fictive activity patterns using different genetically-encoded calcium indicators [1], [46], [47], [58]. Succinctly, isolated CNS preparations display fictive forwards and backwards motor programmes characterised by anterior-to-posterior or posterior-to-anterior progression of activity in motor neuron segments, respectively. In addition, preparations also can display fictive bilaterally symmetric activity in the anterior thoracic and posterior abdominal segments indicative of head raising or tail grounding behaviours, respectively. Finally, preparations can display bilaterally asymmetric activity in the anterior thoracic indicative of headsweep-like behaviours in intact *Drosophila* larva. OA is manufactured and released from neuronal subpopulations in the suboesophageal ganglion and ventral nerve cord expressing the *tdc2+* gene [9] with staining suggesting the brain contains only TA, and no OA-immunoreactive cells [59]. Given both chronic and acute manipulations can have a profound effect on locomotion, we first evaluated if activity of OA neurons in the ventral nerve cord (VNC) - indicative of TA and OA release - was correlated with fictive motor rhythms. There exist subpopulations of distinct octopaminergic neurons that line the ventromedial (T3-A6) axis of the CNS[9]. Through expressing different calcium indicators in VNC motor neurons (VGlut, RCaMP1a) and ventromedial *tdc*²⁺ neurons (TDC²⁺, GCaMP6f) (Figure 1Aii), we explored the intrinsic relationship between VNC *tdc*²⁺ neurons and motor neurons during spontaneous fictive activity bouts. The calcium activity of motor neurons and *tdc*²⁺ neurons in each VNC segment showed high correlation across different fictive motor programs (Figure 1B).

By parsing all instances of fictive events across dual-colour preparations (N=7), we can see a robust similarity in *tdc*²⁺ and motor neuron temporal activity within each segment (Figure 1C). Specifically, if we take the average calcium activity trace for all segments for motor neurons and *tdc*²⁺ neurons for all fictive activity patterns, we see they have consistent similarity and phase locking for all fictive motor categories. The strong association between activity in motor neurons and *tdc*²⁺ neurons is more apparent when comparing activity peaks between each neuronal population in each segment per each unique fictive behaviour (Figure 1D). The temporal relationship between motor neuronal and *tdc*²⁺ neurons showed variations depending on the fictive motor program. At the initiating segment for fictive activity, mean±SEM motor neuron activity on average preceded *tdc*²⁺ neuron activity for

forward (462±704 ms at A6, N=5), posterior burst (449±614 ms at A6, N=5), and anterior burst

(235±352 ms at T3, N=2) fictive events. In contrast, tdc^{2+} neuron activity preceded motor neuron

activity in initiating segments for fictive backward (432±786 ms at T3, N=3), right asymmetric (9±38

ms at T3, N=2), and left asymmetric (99±57ms at T3, N=3) events. Importantly, any initiating segment

peak difference relationship between motor neurons and tdc^{2+} neurons is not consistent across

segments for fictive forward and backward activity events. For example, as fictive forward motor

activity progresses to the anterior regions (A2 - T3) of the CNS, average tdc^{2+} neuron peak activity now

precedes motor neuron activity (Figure 1B, D). In short, while there is some inter-prep variability

between the temporal organisation of motor neuron and tdc^{2+} neuron peak activity, there is a strong

coupling between the activity of these two neuron subpopulations in all fictive behaviours; in all fictive

patterns, there is a complete ipsilateral mirror-image of activity across all segments even if the lag

relationships show intersegmental differences.

Tdc^{2+} neurons show fictive-like activity patterns both commensurate with but also independent of

motor neuron activity. The majority of fictive motor events across preparations showed

commensurate tdc^{2+} neuron activity events. Across all preparations, 69% of fictive forward waves and

anterior asymmetric events, 76% of posterior burst events, and 100% fictive backward waves showed

commensurate tdc^{2+} activity in a close time window (< 5s) and same segment-to-segment relationship

that classically defines the fictive motor programs (Figure 1E). Noteably, there was a large variability

in recruitment between preparations that underestimates the overall level of recruitment displayed

by some preparations. For instance, in 4 out of 6 preparations there was >90% fictive forward

recruitment. In short, most fictive motor patterns have commensurate tdc^{2+} neuron patterns but there

are excess motor patterns not matched in tdc^{2+} patterns. Similarly, there are excess fictive-like tdc^{2+}

activity patterns that occur without commensurate fictive motor patterns in the network (Figure 1F).

Across preparations, there are excess tdc^{2+} activity patterns reminiscent of all types of fictive-like

activity events. For instance, across all preparations there were 7.8% more backwards-like activity

patterns in octopaminergic neurons than backward fictive motor activity patterns despite 100%

recruitment per preparation. Similarly, per preparation, there were 8.7%, 5.4%, 1.8%, 2.1% excess

tdc^{2+} recruitment for anterior burst, anterior asymmetric, fictive forward, and posterior burst activity,

respectively.

Independent of fictive events, the activity in motor neurons and tdc^{2+} neuron pools per segment are

highly correlated. Specifically, the quantity of motor neuron and tdc^{2+} neuron calcium activity peaks

(r_s = 0.82, p < .001) (Figure 1Gi) and duration (r_s = 0.81, p < .001) (Figure 1Gii) of activity within each

segment were highly correlated. Pearson correlation between motor neuron and tdc^{2+} neuron calcium

activity per segment demonstrated an average, moderately positive correlation (r = 0.4) but with

considerable intersegmental and lateral variability (Fisher's Z Score = 0.44 with 0.27 variance). Overall,

activity of motor neurons and tdc^{2+} neurons are highly correlated. When fictive motor events occur,

there is likely a commensurate, almost synchronous tdc^{2+} signal as the event propagates in the VNC

but there are occasions when both fictive motor events and tdc^{2+} fictive-like events appear to occur

independently.

Figure 1: High Activity Correlation between *tdc²⁺* VNC and Motor Neurons in the Same Segment and Fictive Pattern

(**Ai**) The isolated ventral nerve cord (VNC) demonstrates sequences of motor neuron activity resembling activity patterns underlying locomotion in intact animals (i.e., "FW" = fictive forward, "BW" = fictive backward, "AN" = anterior burst, "PB" = posterior burst, "ASR" = asymmetric right, "ASL" = asymmetry left). (**Aii**) Intrinsic calcium activity of motor neurons (MN) and octopaminergic neurons (TDC) is evaluated through dual calcium indicator expression at nerve root regions of interest (blue circles) T3 - A6. Dots above the traces indicate the initiation of the different fictive behaviours (light blue, fictive forward; orange, fictive backward; yellow, anterior symmetric burst; pink, posterior burst; blue, left anterior asymmetry; green, right anterior asymmetry). Lines indicate baseline-correct calcium fluorescence values (green/blue = right/left MN in each segment, magenta/red = right/left *tdc²⁺* VUM in each segment) (**B**) Exemplar calcium trace of MN (left = blue, green = right) and *tdc²⁺* neurons (left = magenta, right = red) demonstrating strong association between MNs and Tdc2+ neuron activity within each ventral nerve cord (VNC) segment. (**C**) Averaged calcium image traces of *VGlut RCaMP1a* (black)*, Tdc2⁺ GCaMP6f* (magenta) for fictive forward (FW), backward (BW), anterior burst (AB), left asymmetry (blue, ASL), right asymmetry (green, ASR), and posterior burst (PB) activity. Bold traces represent average across all preparations (N = 7) with each individual prep average calcium trace in non-bold. The calcium signal amplitudes are normalised within each fictive behavioural class. (**D**) The time difference between the peaks in the MN and TDC calcium trace within each preparation, across fictive patterns and ventral nerve cord segment (T3 - A6). Negative average peak time differences occur when the average prep MN trace proceeds the TDC signal in the segment, positive average peak time represents TDC proceeding MN signals in the segment. Green line represents the average peak time difference across all preparations in the most distal VNC segments. (**E**) The % event count where a motor neuron fictive pattern exists alongside a corresponding TDC activity pattern across different fictive activity categories (with standard error bars). (**F**) The count of fictive activity patterns demonstrated by MN and TDC VNC populations. Note in many fictive activity patterns there is an excess of TDC exhibitions compared to motor fictive patterns. (**G**) An agnostic measure of MN and TDC signals pooled across segments and preparations showing the Spearman's Rank for (**i**) peak count comparison and (**ii**) 20% on-off peak duration comparison. (**H**) Pearson correlation coefficient for the MN and TDC traces of each segment within all preparations for the left (blue) and right (green) axis of the VNC. Comparisons followed by post-hoc was conducted with p-values: < .5 (*), < .01 (**), <.001 (***) shown. Only statistically significant bars are shown.

Bath Application of OA induces Fictive Forward & Long-lasting Anterior Asymmetric Activity

Given the high correlation between intrinsic *tdc²⁺* neuron CNS and spontaneous motor activity, we next turned to evaluate the effects of OA on fictive motor bouts. Exogenous application of OA to intact invertebrates has a well-reported role as a promoter of locomotion aligned with fight-or-flight responses across species [4], [11]. Here, we explored how exogenous application of OA shapes locomotory networks devoid of sensory feedback to better understand the role of OA on coupled oscillatory circuits and motor competition.

The exogenous application of OA altered the fictive landscape of the isolated VNC preparation from fictive variability to nearly exclusive fictive forward motor patterns (Figure 2A). Application of OA concentrations ≥100μM significantly promoted fictive forward activity (100μM $5.4\pm2.0\text{m}^{-1}$ p = .003;

200µM 6.5$\pm$3.2m^{-1} p < .001; 400µM 14.9$\pm$6.4m^{-1} p = .004) compared to saline pre-application (100µM

0.7$\pm$0.4m^{-1}; 200µM 0.7$\pm$0.7m^{-1}; 400µM 0.8$\pm$0.7m^{-1}) (Figure 2Ci). At OA concentrations ≥200µM, the

induction of fictive forward bias came at the expense of fictive backward motor activity which was

significantly reduced (200µM 0.2$\pm$0.6m^{-1} p = .005; 400µM 0.6$\pm$0.7m^{-1} p = .004) compared to saline

pre-application (200µM 1.4$\pm$0.5m^{-1}; 400µM 2.3$\pm$0.9m^{-1}) (Figure 2Cii). Interestingly, anterior

asymmetric events were significantly promoted at higher OA concentrations >400µM (Figure 2Cv).

However, the significant elevation in anterior asymmetric activity was not induced during the 10-

minute OA application but only after the OA wash and thus consequent to the neuromodulator

induced fictive forward bias (Figure 2A, Cv). Importantly, while OA did not appear to significantly

potentiate anterior asymmetric fictive activity in a concentration-dependent manner, there was a

trend to increased anterior asymmetric activity with increasing OA concentration. Interestingly, OA

induced long-lasting elevated anterior asymmetric activity whose activity trend peaks 35.7$\pm$16.9

minutes post the start of the OA wash. Noteworthily, as anterior asymmetric activity is not promoted

at lower OA concentrations, we can be confident that the post-application effects are not an artefact

of diminishing OA during the gradual wash period. Taken together, we can conceptualise OA

neuromodulation as restraining motor competition via biasing the network to fictive forward

locomotion. By exploring fictive transitions (i.e., the probability of making a transition from one motor

program to another), we can see that OA's forward bias is reflective of a significant decrease in fictive

transitions programs that would transition the network away from, and thus reinforce programs

towards, fictive forward motor patterns (Supplementary Figure 1B). Overall, exogenous application of

OA appears to collapse fictive activity diversity, tuning the network into fictive forward bias mode with

consequent post-application motor bias to non-wave-like activity.

To understand the relationship between OA's induction of fictive forward bias and the post-

application anterior asymmetric fictive activity, we examined the relative changes in fictive frequency

between different motor programs across all concentrations (Figure 2E). Intriguingly, the more

exogenous OA promoted fictive forward bias, the greater the post-application anterior asymmetric

activity relative to pre-application (Figure 2Ei, r_s = 0.5, p < .001). This proportional relationship was not

exhibited between any other fictive event. Interestingly, the proportional increase in anterior

asymmetric activity with respect to fictive forward bias appears significantly conditional on the relative

promotion of anterior asymmetric activity during OA application (Figure 2Eii, r_s = -0.4, p = .002). Put

simply, the greater fictive forward bias promoted by OA, the more fictive anterior asymmetric activity

is delayed until the network is no longer forward biased (Figure 2Evi ,R^2 = 0.4 p < .001). In effect, it

appears that post-OA promotion of anterior asymmetric events is a product of OA evoking greater

fictive forward bias in the isolated preparation at the expense of anterior asymmetric events. During

OA application, the induced fictive forward bias is regularly interspersed with a few anterior asymmetric events (Figure 2A, Octopamine, "FW // AS Transitions"). Taken all together, this implies that fictive forward bias immediately potentiates anterior asymmetric activity, but the former dominates the fictive landscape over the latter until OA's removal removes the neuromodulatory potentiation of fictive forward activity.

Aside from changes in the frequency of fictive events at the CPG level, OA induces significant decrease in the amplitude of the calcium signal without a noticeable change in signal baseline as previously reported by [60] (Figure 2Aii, Supplementary Figure 1A-E). The reduction in calcium amplitude was reflected by a reduced duration in motor root bursting during OA application in both thoracic and abdominal motor pools demonstrating the amplitude change in the calcium signal is reflective of OA-induced shorter motor neuron bursting (Supplementary Figure 1F-H).

We next evaluated how the OA-induced faster rhythms exhibited in intact *Drosophila*[27] are represented in the isolated motor circuit. Exogenous bath application of OA qualitatively decreased intersegmental propagation delay for fictive forward and turn but not fictive backward activity (Figure 2D). Fictive forward wave (Figure 3Di) and turn wave (Figure 3Diii) duration is significantly shorter with exogenous OA application. Specifically, all OA concentrations ≥50μM significantly increased forward wave speed (50μM 4.4$\pm$2.4s p = .05; 100μM 3.6$\pm$1.3s p < .001; 200μM 3.7$\pm$1.4s p = .002; 400μM 1.7$\pm$0.7s p = .01) compared to pre-application saline perfusion (8.4$\pm$8.3s). Similarly, octopamine concentrations ≥200μM significantly increased turn wave speed (200μM 3.2$\pm$1.6s p = .01; 400μM 1.6$\pm$1.4s p = .03) compared to pre-application saline perfusion (7.0$\pm$1.9s). OA did not induce any significant changes in fictive backward wave speed. Thus, OA biases the isolated *Drosophila* CNS to fictive forward locomotion consequently restricting motor diversity, induces faster CPG bursting, and potentiates intersegmental propagation between segments that affects some system-wide activity progression. More subtly, given that lower OA concentrations could significantly affect selective intersegmental propagation but not overall fictive frequency, these data imply OA can modulate locomotory speed independent of modulating initiation bias.

Figure 2: Bath Application of OA Induces Fictive Forward Bias in the isolated larval CNS

(**A**) Exemplar calcium dynamics in left (blue) and right (green) motor neurons in the 3rd instar isolated *Drosophila* ventral nerve cord (VNC) over 45 minutes under saline, octopamine (OA) (200μM), wash bath applications. Difference trace for T3 (ΔT3) shown above all traces (left - right). Dots above the traces indicate the initiation of the different fictive behaviours (light blue, fictive forward; orange, fictive backward; yellow, anterior symmetric burst; pink, posterior burst; blue, left anterior asymmetry; green, right anterior asymmetry). Black lines indicate the shape of each fictive activity to aid user's recognition of fictive programmes. Note that octopamine application induces fictive forward bias with intermittent asymmetric activity with the post-application wash period exhibiting dominating anterior asymmetric activity. (**B**) Changes in fictive frequencies of different fictive behaviours under the application of OA concentrations (5 - 400μM) between saline, OA perfusion, and wash conditions for fictive: (**i**) forward , (**ii**) backward, (**iii**) anterior burst, (**iv**) posterior burst, (**v**) anterior asymmetric headsweeps, and (**vi**) turn. Note that the axis showing the rate of each fictive activity is different for each fictive activity. (**C**) Activity propagation speed from

posterior (A8) to anterior (T3) for (**i**) fictive forward and anterior-to-posterior for (**ii**) fictive backward and (**iii**) fictive

turning behaviours. (**Ei**) Ranked relationship between the relative effect of OA on promoting fictive forward activity

and the relative change in headsweep asymmetric. (**Eii**) Ranked relationship between the relative promotion of

anterior asymmetric activity by OA and the change in anterior asymmetric activity in wash relative to OA application

activity level. (**F**) Ranked relationship between relative promotion of fictive forward waves by octopamine during

application (X variable, FW (OA-S)), relative promotion of anterior asymmetric activity during octopamine

application compared to the activity in the pre-application time window (Y variable, HS (OA-S)), and relative change

in anterior asymmetric activity between the octopamine application and post-application time window (Z variable,

HS (W-OA)). Z = 0.45*X + -0.44*Y + 27.78. Note the fact that fictive forward promotion in octopamine comes at the

expense of asymmetric anterior headsweep activity.   Significant post-hoc comparisons are exclusively shown

denoted by □ saline vs wash, ✳ saline vs OA, Δ OA vs wash.

Tyramine Bath Application Induces Fictive Overlap & Motor Competition

Tdc²⁺-expressing neurons are capable of synthesising and releasing both OA and TA [7], [8], [10], [59].

Given the highly active coupling of intrinsic *tdc²⁺* activity and motor activity and OA's excitatory effect

on fictive activity, we next evaluated the role of TA on CPG activity and motor competition. TA is

considered antagonistic to OA modulation with evidence of TA's inhibitory role on intact locomotion

[27] and motor electrophysiology [34]. Thus, we turned to explore whether TA has a universal

inhibitory role on motor competition in isolated preparations to better understand how behavioural

inhibition manifests at a CPG and network level via *tdc²⁺* activity.

We explored the effect of TA on motor competition by exogenously bath applying TA onto isolated 3ʳᵈ

instar larva VNC as in our OA experiments. Unlike OA, TA did not induce a consistent motor bias across

applied concentrations compared to pre-application (Figure 3A, B). Across all fictive behavioural

patterns, the only significant fictive changes occurred post-TA application. Specifically, fictive forward

(200μM $0.6\pm0.4\text{m}^{-1}$ p < .001) and posterior burst (400μM $0.7\pm0.6\text{m}^{-1}$ p < .001) activity was

significantly reduced in the wash period relative to the TA application period (FW, 200μM $10.3\pm8.1\text{m}^{-1}$

$^{-1}$) or pre-application saline period (PB, 200μM $18.7\pm11.5\text{m}^{-1}$) respectively. Further, again in contrast

to OA, TA did not induce any significant changes to the intersegmental propagation of neither fictive

forward, backward, nor turn behaviours (Figure 3C). Reflective of this insignificant change in fictive

incidences, TA perfusion appears to not induce significant, widespread changes in the landscape of

fictive transitions (Supplementary Figure 2B). In fact, in some instances, TA perfusion significantly

increased motor transitions reflective of subtle increases in motor competition and thus overall fictive

program diversity.

While TA did not induce fictive bias in the isolated network, TA perfusion induced a significant increase

in the overlap of fictive motor programs (Figure 3A, Tyramine "Overlap") represented by increased

overlap in activity across at least one ventral nerve cord ganglion segment (Figure 3D). Perfusion of

200μM TA induced a significant increase in time the network demonstrates activity overlap across at least one segment ($6.2\pm0.4\%$ p = < .001) compared to the 0% overlap demonstrated during the pre-tyramine application period. Aside from overlap promotion, the application of ≥100μM TA induced a significant decrease in the activity in the post-application in a concentration-dependent manner (100μM $84.5\pm4.8\%$ p = .01; 200μM $49.8\pm31.7\%$ p = .002; 400μM $56.4\pm12.3\%$ p = .006) compared to pre-application period (100μM $95.4\pm3.8\%$; 200μM $93.2\pm9.8\%$; 400μM $97.7\pm2.2\%$) (Figure 3A, Wash "Silence"; Figure 3Ei). Similar to our prior analysis evaluating correlations between fictive changes by the OA and the post-application anterior asymmetric increase, we turned to correlation analysis between activity and silence to attempt to understand the long-lasting modulatory effects of TA. However, we did not find any significant relationship between activity changes by TA perfusion and the relative changes in activity in the post-application wash period (Figure 3Eii). Further, there was no significant relationship between the overlap and the post-application silence promoted by TA (Figure 3Eiii). Given prior work has drawn attention to the state-dependent effects of TA inhibition[26], we next evaluated if TA's effects on fictive activity were dependent upon the activity state prior to TA application. As shown in Figure Eiv, TA did exhibit proportional and increasing inhibitory effects as overall fictive activity levels rose beyond pre-application activity levels > $6.7m^{-1}$ (p = .04, r^2 = 0.2). Further more, when baseline activity is low, TA can actually have excitatory effects. Taken together, TA promotes overlaps between different fictive motor programs and demonstrates activity-dependent inhibition of overall fictive activity and

Given that previous work has indicated an inhibitory role for TA at an individual cellular level[26], [48] we evaluated the effect of TA perfusion on motor nerve root bursting activity. Unexpectedly, TA perfusion induced a trend, albeit marginally insignificant, towards an increase in motor root bursting (Supplementary Figure 3Ai, ii) with no significant effects on bursting duration (Supplementary Figure 3Aiii), overall duration of inactivity (Supplementary Figure 3Aiv), bursting amplitude (Supplementary Figure 3Av), or tonic baseline amplitude (Supplementary Figure 3Avi). 200μM TA application induced a significant reduction (p = .001) in the amplitude of calcium activity (Figure 3A, Supplementary Figure 3Bi, iii) without, similar to OA, any changes to the baseline calcium activity (Supplementary Figure 3Bii, iv).

Overall, TA application to the isolated network is inhibitory on rhythm generation at higher activity states but excitatory at lower activity states. I Further, TA appears to facilitate the overlapping of fictive motor programs with prolonged reduction in overall activity absent inhibition at the motor neuron population level.

Figure 3: Bath Application of TA Induces Fictive Motor Overlap and Competition

(**A**) Exemplar calcium dynamics in left (blue) and right (green) motor neurons in the 3rd instar isolated *Drosophila* ventral nerve cord (VNC) over 45 minutes under saline, tyramine (TA) (200µM), wash bath applications. Note the overlap between fictive forward and backward motor activity (red) during tyramine application and absence of fictive activity in the wash period. (**B**) Changes in fictive frequencies of different fictive behaviours under the application of TA concentrations (10 - 400µM) between saline, TA perfusion, and wash conditions for fictive: (**i**) forward , (**ii**) backward, (**iii**) anterior burst, (**iv**) posterior burst, (**v**) anterior asymmetric headsweeps, and (**vi**) turn. Note that the axis showing the rate of each fictive activity is different for each fictive activity. (**C**) Activity propagation speed from posterior (A8) to anterior (T3) for (**i**) fictive forward and anterior-to-posterior for (**ii**) fictive backward and (**iii**) fictive turning behaviours. (**D**) Percentage of time where overlap in activity between segments exists for pre-application, application, and post-application of TA across all concentration ranges. (**Ei**) Percentage of time where there is activity in at least one segment for pre-application, application, and post-application of TA across all concentration ranges. Note the concentration-dependent reduction in total activity within the wash period. (**Eii**) The effect of

relative promotion (green) or suppression (red) of activity during TA application against the percentage of time
where there is activity in at least one segment post TA application. Note the absence of a bifurcation or trend
between wash activity and TA's promotive or suppressive activity relative to pre-application. (**Eiii**) Non-significant
relationship between the % overlap induced by TA and post-application silence. (**Eiv**) The relationship between
overall pre-application fictive activity ("Saline Activity") and the change in overall fictive activity during TA application
("TA vs Saline"). Note the activity-dependent inhibitory effect of tyramine. Significant post-hoc comparisons are
exclusively shown denoted by □ saline vs wash, ✳ saline vs TA, Δ TA vs wash.

Tdc^{2+} Neurons are Necessary and Sufficient for Fictive Locomotor Activity

Intrinsic *tdc^{2+}* activity is highly correlated to segmental motor neuron activity with exogenous OA and

TA application modulating motor competition. We next evaluated if manipulating intrinsic *tdc^{2+}* system

activity can also induce fictive motor bias, anterior asymmetric circuits potentiation post-application,

and alter motor competition. We expressed either *CsChrimson* or *GtACR1* in *tdc^{2+}* neurons to

optogenetically manipulate the intrinsic adrenergic-like system and evaluate motor neuron activity

using nerve-root electrophysiology. Here, we show how intrinsic *tdc^{2+}* activity is necessary and

sufficient for fictive locomotor activity and exhibits most of the phenotypic effects displayed by

exogenous OA application.

Firstly, we optogenetically depolarised *tdc^{2+}* neurons by persistent red-light illumination (10, 20, 40s)

and recorded motor root activity from T3 and A6 VNC ganglia (Figure 4A). Optogenetic depolarisation

of *tdc^{2+}* neurons induced significantly increased motor root bursting for all: 10s (pre: 9.6 ± 7.5 m^{-1}, on:

55.4 ± 26.1 m^{-1}, p = .002), 20s (pre: 11.4 ± 4.6 m^{-1}, on: 46.3 ± 9.1 m^{-1}, p=.002), and 40s (pre: 11.2 ± 6.6 m^{-1}

^{1}, on: 41.0 ± 12.9 m^{-1}, p=.005) stimulation periods (Figure 4A, Ci). The increased bursting frequency in

A6 was accompanied by significant decrease in bursting duration for all: 10s (pre: 2.8 ± 1.3 m^{-1}, on:

1.0 ± 0.4 m^{-1}, p=.002) 20s (pre: 3.3 ± 1.2 m^{-1}, on: 0.9 ± 0.3 m^{-1}, p=.004), and 40s (pre: 3.2 ± 1.7 m^{-1}, on:

1.7 ± 1.9 m^{-1}, p=.005) stimulation periods (Figure 4Cii). Elevated A6 motor root bursting frequency was

also evocable by pulsed-based stimulation (Supplementary Figure 4). Across both continuous and

pulsed-based stimulation of *tdc^{2+}* neurons, intrinsic manipulation of the *tdc^{2+}* system induced fictive

forward bias akin to results demonstrated by exogenous OA application (Figure 4E). Specifically,

optogenetic depolarisation of *tdc^{2+}* neurons induced significantly increased fictive forward motor bias

for all: 10s (pre: 3.3 ± 2.2 m^{-1}, on: 64.2 ± 43.0 m^{-1}, p < .001), 20s (pre: 4.3 ± 4.0 m^{-1}, on: 36.3 ± 7.7 m^{-1}, p

< .001), and 40s (pre: 4.8 ± 5.1 m^{-1}, on: 36.8 ± 10.8 m^{-1}, p <.001) stimulation periods. In terms of fictive

bias, averaged across stimulation periods, optogenetic depolarisation of *tdc^{2+}* neurons induced fictive

forward waves to become $79\pm8\%$ of all fictive activity events during stimulation as opposed to

$36\pm13\%$ before or $36\pm14\%$ after red-light stimulation.

Counter to the linear frequency relationship found between promoted fictive forward and anterior asymmetric elevated activity post-application in exogenous octopamine application, anterior asymmetric fictive activity was not elevated in the post-stimulation relative to pre-stimulation period (Figure 4F). Further, while we witnessed a seemingly consistent elevation in silence (i.e., continuous lack of apparent rhythmic activity in any nerve root) post optogenetic stimulation (see Figure 4A, ~40s post-stimulation, 34.6±27.8 m) compared to control preparations (9.9±9.1 m), the duration of silence did not significantly differ (p = .58) nor covary with stimulation period (10s: p = .07; 20s: p = .27; 40s: p = .25) (Figure 4G).

Secondly, we optogenetically hyperpolarised tdc^{2+} neurons with pulses of light (15s, 30s, 60s) and recorded motor root activity from T3 and A6 VNC ganglia (Figure 4B). Optogenetic hyperpolarisation of tdc^{2+} neurons significantly reduced, and in many cases, completely collapsed, motor root bursting for all light-stimulation periods: 15s (pre: 1.6±2.1 m^{-1}, on: 0.5±1 m^{-1}, p = .04), 30s (pre: 1.8±1.8 m^{-1}, on: 0±0 m^{-1}, p=.009), and 60s (pre: 2.2±2.1 m^{-1}, on: 0.4±0.7 m^{-1}, p = .01) illumination periods with 37 of 49 (75.9±30.8%) trials across preparations (N=7) showing no fictive activity during the white-light stimulation period (Figure 4Bii, D). For longer period white-light stimulation (> 15s), there was weak recovery of rhythmic activity in the A6, indicative of post-inhibitory rebound. Interestingly, long-lasting (60s) white-light illumination in tdc^{2+}>GtACR1 no retinol control preparations induced significantly increased (pre: 1.8±2.0 m^{-1}, on: 3.7±1.5 m^{-1}, p = .01) A6 motor root activity potentially indicative of photoreceptor-based motor stimulation in the isolated VNC preparation.

Thus, tdc^{2+} depolarisation provokes elevated fictive forward activity akin to exogenous OA application. Further, tdc^{2+} hyperpolarisation collapsed all motor rhythms. Considering the strong correlation between intrinsic tdc^{2+} and motor neuron activity, such results demonstrate tdc^{2+} neurons are necessary and sufficient for fictive motor activity in isolated VNC 3rd *Drosophila* larval preparations.

540 **Figure 4: *tdc*²⁺ Neuronal Activity is Sufficient and Necessary for Fictive Locomotor Activity**

541 (**A**) Exemplar three nerve root (T3 (L), T3 (R), A6(R)) electrophysiology trace for continuous red-light depolarisation

542 of *tdc*²⁺ neurons using CsChrimson for preparations raised on retinol (experimental, left) and no retinol (control,

right). Before, during (red box), and post-stimulation periods are shown with rectified and smoothed traces shown. Black lines represent fictive activity motifs. (**B**) Exemplar two nerve root (T3 (L), A6 (R)) electrophysiology trace for continuous white-light hyperpolarisation (red box) of tdc^{2+} neurons using GtACR1 for preparations raised on retinol (experimental, left), and no retinol (control, right). (**C**) Quantification of abdominal (A6) motor root bursting for (**i**) burst rate and (**ii**) duration of bursts in tdc^{2+}>CsChrimson for 10s, 20s, and 40s depolarisation period. (**D**) Quantification of abdominal (A6) motor root bursting for burst rate in tdc^{2+}>GtACR1 for 15s, 30s, and 60s hyperpolarisation period. (**E**) Change in the fictive activity profile for tdc^{2+}>CsChrimson stimulation periods demonstrating inducing of fictive forward bias across stimulation period. Activity profile within stimulation periods and average across stimulation periods are shown. (**F**) Difference in activity rate of fictive forward activity between stimulation and pre-stimulation period (blue) as well as fictive anterior asymmetric activity between post-stimulation and pre-stimulation period (green). (**G**) Duration of no rhythmic activity quantified post stimulation periods in control (- retinol) and experimental (+ retinol, CsChrimson) preparations. Comparisons followed by post-hoc was conducted with p-values: < .5 (*), < .01 (**), <.001 (***) shown. Only statistically significant bars are shown.

Given tdc^{2+} VNC neurons are spatially segregated into each thoracic and abdominal ganglion, we next turned to see if spatially-restricted tdc^{2+} activity may differentially modulate fictive locomotion network behaviour. To evaluate any effect of spatially-restricted release of OA on fictive locomotion, we performed restricted red-light stimulation on tdc^{2+} neuronal populations in the ventromedial axis on the isolated ventral nerve cord. Qualitatively, direct red-light stimulation of tdc^{2+} neuronal populations induces elevated bursting rate in all restricted regions: SOG (Figure 5A), thoracic ganglion (Figure 5B), anterior abdominal ganglion (Figure 5C), and posterior abdominal ganglion (Figure 5D). Across all stimulation regions, both the thoracic and abdominal nerve root bursting rate is significantly increased during tdc^{2+} depolarisation (Figure 6Ai - Di). Specifically, direct SOG tdc^{2+} depolarisation induced a significant increase in T3 (pre: 8.3±4.8 m^{-1}, on: 30.0±6.9 m^{-1}, p = .001) and A6 (pre: 8.8±4.7 m^{-1}, on: 31.7±10.8 m^{-1}, p = .001) motor root burst rate (Figure 6Ai). Direct thoracic ganglion (T1-3) tdc^{2+} depolarisation induced a significant increase in T3 (pre: 6.5±3.4 m^{-1}, on: 30.0±6.0 m^{-1}, p < .001) and A6 (pre: 6.8±3.5 m^{-1}, on: 37.7±13.3 m^{-1}, p < .001) motor root burst rate (Figure 6Bi). Direct anterior abdominal ganglion (A1-3) tdc^{2+} depolarisation induced a significant increase in T3 (pre: 6.6±2.1 m^{-1}, on: 34.3±12.4 m^{-1}, p = .001) and A6 (pre: 6.8±3.5 m^{-1}, on: 32.6±17.3 m^{-1}, p = .022) motor root burst rate (Figure 6Ci). Finally, direct posterior abdominal ganglion (A6-8) tdc^{2+} depolarisation induced a significant increase in T3 (pre: 6.1±2.5 m^{-1}, on: 48.0±12.0 m^{-1}, p < .001) and A6 (pre: 6.5±2.1 m^{-1}, on: 45.4±9.7 m^{-1}, p < .001) motor root burst rate (Figure 6Di). Further, across all stimulation regions, both the thoracic and abdominal nerve root bursting duration was significantly reduced during tdc^{2+} depolarisation with the exception of the T3 motor root during thoracic and anterior abdominal ganglion (Figure 6Ai-Di). Specifically, direct SOG tdc^{2+} depolarisation induced a significant decrease in T3 (pre: 3.0±1.3s, on: 1.5±0.6s, p = .01) and A6 (pre: 3.0±1.3s, on: 1.3±0.4s, p = .03) motor root burst duration (Figure 6Ai). While direct thoracic ganglion (T1-3) tdc^{2+} depolarisation did not induced a significant decrease in T3 motor root burst rate (pre: 2.9±1.0s, on: 1.5±0.4s, p = n.s),

depolarisation of T1-T3 tdc^{2+} neurons did induce a significant decrease in the A6 motor root burst duration (pre: 3.1$\pm$1.9s, on: 1.4$\pm$0.6s, p = .02) (Figure 6Bi). While direct anterior abdominal ganglion (A1-3) tdc^{2+} depolarisation did not induced a significant decrease in T3 motor root burst rate (pre: 3.2$\pm$1.2s, on: 1.5$\pm$0.4 m^{-1}, p = n.s), depolarisation of A1-A3 tdc^{2+} neuron did induce a significant decrease in the A6 motor root burst duration (pre: 3.1$\pm$1.9s, on: 1.4$\pm$0.6s, p = .04) (Figure 6Ci). Finally, direct posterior abdominal ganglion (A6-8) tdc^{2+} depolarisation induced a significant decrease in T3 (pre: 3.5$\pm$1.3s, on: 1.1$\pm$0.2s, p = .001) and A6 (pre: 3.3$\pm$1.1s, on: 1.1$\pm$0.2s, p = .003) motor root burst duration (Figure 6Di). Thus, depolarisation of any ganglion tdc^{2+} neuron subpopulation is sufficient to induce elevated motor activity in the isolated ventral nerve curve.

We next examined if sub-region restricted tdc^{2+} neuron depolarisation induced fictive forward bias akin to our earlier results on exogenous OA application and whole-VNC tdc^{2+} depolarisation. Depolarisation of tdc^{2+} neurons in the SOG (Figure 6Aii), thoracic ganglion (Figure 6Bii) and anterior abdominal ganglion (Figure 6Cii) induced an, albeit insignificant, elevated mixture of fictive behavioural patterns. However, decomposing the population data down into individual iterations of restricted stimulation across preparations did demonstrate a significant variation in the fictive bias within each region of tdc^{2+} depolarisation (Supplementary Figure 5). Namely, restricted tdc^{2+} depolarisation in the SOG (Supplementary Figure 5A, $\chi^2(2)$ = 7.3, p = .007), thoracic ganglion (Supplementary Figure 5B, $\chi^2(2)$ = 7.3, p = .007), and anterior abdominal ganglion (Supplementary Figure 5C, $\chi^2(2)$ = 7.3, p = .007) all significant induced changes in fictive wave bias with some instances showing promotion of fictive forward whereas others fictive backward normalised bias. In other words, while averaging between preparations demonstrates a general trend of promoting both fictive forward and backward preparation in restricted tdc^{2+} depolarisation in the SOG, T1-3, and A1-3, in individual trials, there are many instances of either fictive forward or backward bias resultants. In contrast, posterior abdominal restricted tdc^{2+} depolarisation induced a significant increase in exclusively fictive forward motor activity (pre: 2.4$\pm$1.9m^{-1}, on: 38.6$\pm$13.8m^{-1}, p = .005) (Figure 6Dii). While posterior abdominal restricted tdc^{2+} depolarisation similarly induced increased fictive bias variation (Supplementary Figure 5D, $\chi^2(2)$ = 5.8, p = .02), a majority of stimulation instances were fictive forward biased. Thus, both abdominal posterior and whole tdc^{2+} depolarisation induced locomotor CPGs into a fictive forward bias.

Figure 5: Restricted Optogenetic Stimulation of *tdc²⁺* Neurons Induces Diversity of Activity

Restricted 10s optogenetic stimulation of *tdc2*+ neurons in the (**A**) SOG, (**B**) thoracic ganglion (T1-T3), (**C**) anterior abdominal ganglion (A1-A3), and (**D**) posterior abdominal ganglion (A6-A8). Representative electrophysiology

thoracic (T3) and posterior abdominal (A6) nerve root recording traces for direct and flanked stimulation conditions

shown with coloured-dots indicating occurrence of fictive motor pattern: fictive forward wave (blue dot), posterior

burst (purple dot), fictive backward wave (orange dot), anterior a/symmetric activity (green dot).

Figure 6: Optogenetic Stimulation of Posterior Abdominal *tdc^{2+}* Neurons Induces Fictive Forward
Locomotion
Quantification of (**i**) abdominal and thoracic burst rate, burst duration, and (**ii**) fictive activity ("FW" = fictive forward
wave, "HS/AB" = anterior symmetric or asymmetric, "BW" = fictive backward wave, "PB" = posterior bust) rate
during direct (colour) and flanked (gray) optogenetic stimulation of (**A**) SOG, (**B**) thoracic ganglion (T1-T3), (**C**)
anterior abdominal ganglion (A1-A3), and (**D**) posterior abdominal ganglion (A6-A8) *tdc^{2+}* neurons. Comparisons
followed by post-hoc was conducted with p-values: < .5 (*), < .01 (**), <.001 (***) shown. Only statistically significant
bars are shown.

Discussion

The adrenergic-like system is involved in a diversity of behaviours across invertebrates. Here, we show

that the adrenergic-like system is necessary and sufficient for generating rhythmic activity in

Drosophila larval CPG networks. Through acute manipulations to OA and TA signalling, we

demonstrate that the core adrenergic-like neuromodulators – OA and TA – may not always be

mutually-antagonistic for locomotion in *Drosophila* larvae. Finally, we discovered that a by-product

of extensive and sustained motor bias is an instantiation of antagonistic fictive activity after the end

of a prior bias episode. The crucial element of sustained movement is action selection and the

inhibition of competing motor programs. Here, we show how the isolated neural system appears to

have an innate disposition to rebalance motor diversity to antagonistic motor patterns in response to

sustained suppression of motor diversity through sustained motor bias.

The Fundamental Role of *tdc^{2+}* Activity in *Drosophila* Larvae Fictive Activity

The cellular architecture of adrenergic-like *tdc^{2+}* system in *Drosophila* is reported [45], [61] but the

broad functional role of the intact system, divorced from the inferences about the role of their

released neuromodulators, is less explored. Here, we showed how depolarisation of both the whole,

and subparts of, *tdc^{2+}* system does drive fictive locomotion and how system-wide hyperpolarisation

collapses all rhythm generation in the isolated network.

Despite the absence of a comprehensive list of specific presynaptic partners for the *tdc^{2+}* system, we

can infer that the *tdc^{2+}* system likely exists directly upstream of the VNC rhythm generating networks.

Based on current connectomics data, only the ascending and descending subparts of the *Drosophila*

1st instar larva *tdc^{2+}* system appear to have postsynaptic partners. The VNC *tdc^{2+}* system - the vast

majority of our activity manipulation - appears to exclusively project to the neuromuscular junction

[45], [61], [62]. Given the high correlation in *tdc^{2+}* system activity to system-wide segmental motor

activity alongside the fact that hyperpolarisation of the *tdc^{2+}* system evoked an immediate cessation

of fictive activity, our results imply the *tdc^{2+}* system may have a direct neuromodulatory roles that

might emerge after the 1st instar developmental stage. The complete connectome and architectural

makeup of the 3rd instar *Drosophila* larvae is undetermined; one testable prediction is that segmental

tdc^{2+} neuronal populations are connected, directly or indirectly, to reported GDL/A27h premotor

circuity, or intermediary accessory of descending command neuronal structures like MDN, and Pair1

interneurons [63], [64], [65]. Further, the *tdc*$^{2+}$ system may make direct synaptic connections to

pacemaker command neurons (i.e., AcN, A01j) underpinning locomotory speed [66]. Indeed, such

predictions are supported by the tyraminergic architecture in *C. elegans* wherein the activity of RIM

neurons - presynaptic to GABAergic premotor neurons that facilitate bending - occurs prior to changes

in motor neuron activity [33], [48], [49], [50], [51]. Further, the highly coordinated and rhythmic

behaviour of the *tdc*$^{2+}$ system itself implies the existence of a similar intersegmental regulatory

architecture which may include GDL-like feedforward inhibitory motifs [64]. Importantly, the *tdc*$^{2+}$-

GAL4 construct also incorporates large ventral unpaired median (VUM) motor neurons - a subset of

the *tdc*$^{2+}$ system and motor neurons system - that appear to exclusively innervate the larval muscles

[63]. While hyperpolarizing these neurons should undoubtedly reduce motor pool activity, the fact

that system-wide motor activity comprehensively collapses under mass hyperpolarisation suggests

still a fundamental role of the interneuronal or descending portions of the *tdc*$^{2+}$ system in fictive

rhythmic generation. Consequently, we might expect any realised or modelled architecture of the

tdc^{2+} system to incorporate a subpopulation motif for both internal and projecting outputs. Finally,

independent of architecture, the mechanism of modulation by the *tdc*$^{2+}$ system may be more complex

than simply adrenergic-like modulation by OA and TA exclusively. Indeed, we have only a partial OA/TA

receptor expression map restricted to the tissue, but not cellular level [4], [6], [67], [68]. Given that

different OA/TA receptor isoforms have fundamentally different downstream signalling effects [4] and

thus different effects on behaviour (e.g., [22]), any neuromodulatory mechanistic picture currently

lacks a firm foundation. Further, given a majority of adult *tdc*$^{2+}$ neurons co-express OA and glutamate

[69] and some *tdc*$^{2+}$ neurons modulate through gap junctions (e.g., APL neurons in the MB [70]), a

complex, potentially degenerate, neuromodulatory picture may underpin the mechanism behind *tdc*$^{2+}$

system fictive modulation. Overall, we show that the *tdc*$^{2+}$ system could be considered at least part of

a necessary and sufficient motif for rhythm generation in locmotor networks.

Our study uses the isolated VNC removing dynamic re-afferent and sensory feedback. Prior studies

have extensively shown how genetic manipulations to the adrenergic-like system in a variety of

invertebrates induces dysrhythmic movement [4], [9], [27], [28].Leaving aside potentially

developmental compensatory effects, the fact that genetic ablation of the *tdc*$^{2+}$ system does not arrest

all movement in intact animals[33], whereas our results on hyperpolarisation in isolated networks do

collapse rhythmic generation, implies re-afferent or sensory modulation may be one source of

compensation in intact animals with dysfunctional *tdc*$^{2+}$ system.s In future, the isolated

CNSpreparation and consequent analysis pipelines [1], [47] could be used to interrogate if there are

specific tdc^{2+} system deficits induced by genetic manipulations, further bridging our understanding

between intact and fictive locomotion.

The Adrenergic-like System in Modulating Motor Competition

Movement necessitates a balance between excitation-driven initiation of a rhythm and pattern

generating circuit alongside inhibition of competing motor program-related networks. Motor

competition - the dynamic interplay that underpins excitatory and inhibitory neural action - is one lens

through which the OA/TA modulatory dichotomy has yet to be examined. Here, we demonstrated that

the well-reported excitatory role of OA and the inhibitory role of TA can be considered as

neuromodulators underpinning motor network competition.

The locomotor effects of OA/TA neuromodulation are well-reported across a variety of invertebrate

species. Generally, the OA/TA are considered mutual-antagonistic in locomotor neuromodulation with

an excitatory/inhibitory binary [4], [26] .Complementing some studies that highlighted nuanced roles

for adrenergic-like neuromodulation [26], we showed that TA exhibited activity-dependent inhibitory

effects, was promotive of overlapping between fictive programs, thus not exclusively leading to

inhibitory effects. Thus, the slower and shorter locomotion exhibited when *Drosophila* larva are fed

TA [27] could be conceived as a failure to regulate motor competition rather than a direct, general

inhibitory effect. Those two competing explanations for TA effects are non-trivial differences. For

instance, while TA has a hyperpolarising effect on an isolated neuron [34] ,we found an overall trend

of TA to depolarise motor pool activity. Our indication that TA's inhibitory effects may be a product of

activity state can help marry the two bodies of thought. Namely, given that intrinsic tdc^{2+} activity is

highly coupled to fictive motor activity, tdc^{2+} activity is indicative of OA release [45], and that TA's

effects might be conditional on OA levels [26], one can see how TA/OA's antagonism is in response to

their neuromodulatory effects. At a functional level, we therefore could consider OA/TA as mutual-

antagonists altering motor competition rather than generally activity levels.

Our results and the role of the tdc^{2+} system in *Drosophila* larvae can be greatly contextualised by its

similarity to the *C. elegans* neuroarchitecture underpinning turning escape behaviour. Within *C.

elegans*, OA/TA play a significant and fundamental role in modulating escape behaviour and motor

competition through the AVA/AVB/RIM motif [26], [50]. Importantly, the proposed motif

demonstrates how TA/OA underpins motor competition between spontaneous forward and backward

locomotion through TA's disinhibition [51]. Taken together with the effect of increased overlap and

activity-dependent inhibition within our work, we could expect TA to have a similar disinhibitory role

in *Drosophila*. Namely, as reported in our recent work [1], core interneuronal inhibitory motifs might

be essential for all forms of motor selection and appropriate rhythmic output. Thus, we would expect

the tdc^{2+} system - itself releasing neuromodulators essential for escape responses - to be integral to

orchestrating and modulating the dynamics of motor selection at a fundamental level.

OA's excitatory effect on fictive motor behaviour is reflective of an underlying inhibition of motor

competition. Reminiscent of a large catalogue of prior studies [3], [8], [9], [26], [27], [32], we found

exogenously-applied OA was excitatory by being promotive of forward fictive locomotion with higher

intersegmental propagation of activity. Importantly, this promotion of locomotion, especially given

the network appeared perpetually active with little to no episodes of non-activity, was reflective of

significant inhibition of competing motor program occurrence. Indicative of a suppression of initiation,

we here contextualise the neuromodulator OA, in comparison to TA, as a suppressor of motor

competition. Reflective of a deeper role of the adrenergic-like system in motor competition, the

emergence of proportional post-application activity changes due to OA's neuromodulatory effects

indicates a potential foundational connection in CPG competition. Given that asymmetric anterior

activity can be considered a preliminary and important element of backward locomotion [1], the fact

OA's induction of an exclusive fictive forward bias precipitated an elevation of such asymmetric

anterior activity post-hoc suggests a potential underpinning dynamic in motor selection. Namely, a

long-lasting memory-like system may exist to compensate for extremes in fictive forward activity that

compensates if that extreme bias manifests. Indeed, the emergence of post-application elevation in

silence from TA perfusion may also be reflective of a long-lasting effect to motor competition

promotion. Unsurprisingly, inhibition is a vital element in necessitating variability as for a system to

be able to devolve into different output modes it must first have a configuration that allows it to sit at

that potential junction (i.e., conception of criticality) [1], [71]. Notably, there was no apparent post-

asymmetric elevation in our electrophysiology-based stimulation as exhibited in our exogenous

perfusion data. Thus, we would predict "long-lasting bias" could be denoted within a motor system as

dominance of one fictive motif beyond a 60s episode.

Taken together, here we provide a case of how the tdc^{2+} system in *Drosophila* may be a necessary and

sufficient component in rhythm generation underlying locomotion. Further, we nuance the role of

adrenergic-like neuromodulation in motor competition by demonstrating how OA/TA operate to

either restrain or promote diversity of fictive motor programmes. Our results provide constraints on

modelling studies and a framework for further connectomics exploration into how the tdc^{2+} activity is

integrated into the pre-motor system. Finally, we highlight a novel potential relationship within motor

competition between induced locomotion bias and long-lasting fictive antagonism that may
demonstrate a fundamental relationship of CPG networks not currently considered.

752 Conflicts of Interest

753 No declared conflict of interest.

754 Data Avaliability

755 The research data supporting this publication can be accessed at https://doi.org/10.17630/84df17ac-
756 77d3-4a61-9bf0-fa6e78a802e1.

757 Author's Contribution

758 S.R.P. and W.V.S. conceived the study. W.V.S conducted all experiments, analysis, and visualisation.
759 W.V.S. and S.R.P finalised the figures. S.R.P. supervised the project.

760 Acknowledgements

761 This project was made possible by an Industrial CASE PhD studentship (UKRI Biotechnology and
762 Biological Sciences Research Council (BBSRC) grant number BB/T00875X/1) to W.V.S.

 Supplementary Figures

Supplementary Figure 1: Octopamine Decreases Calcium Signal Amplitude and Motor Root Burst Duration Despite No Baseline Change

(**A**) Representative calcium signal trace from A3 left motor neuron before (blue), during 200μM octopamine application (magenta), and after (blue). (**B**) Exemplar activity in the left A3 calcium signal before, during 200μM octopamine application and after. Traces in B are subsections of the total trace denoted by *. (**C**) The average fluorescence value from calcium signal trace before, during, and after 200μM octopamine application. (**D**) The averaged peak amplitude from all A3 motor neuron calcium activity from calcium signal trace before, during, and after 200μM octopamine application across full ventral nerve cord (VNC) perfusion experiments (N=12). (**E**) All traces from full VNC perfusion experiments (grey, dashed line) before, during (magenta box), and after 200μM octopamine application. Note the absence of a visual change in baseline during octopamine application (magenta

box). (**F**) Nerve root recordings for full VNC before (blue) and during (magenta) 200μM octopamine bath application. Yellow lines indicate fictive patterns. Note the elevated frequency of fictive forward wave activity during octopamine application (yellow lines, magenta traces). (**G**) The average amplitude of peaks in A6 motor neurons before and after 200μM octopamine application (N=3). (**H**) The average burst duration of peaks in A7 motor neurons before and after 200μM octopamine application (N=3). Note, the decreased burst duration in motor activity represents calcium activity as a lower signal amplitude. Post-hoc comparison were conducted with p-values: < .5 (*), < .01 (**), <.001 (***) shown. Only statistically significant bars are shown.

Supplementary Figure 2: Octopamine and Tyramine Alters Motor Competition by Modulation Fictive Transition Probabilities

(**A**) The probability of any fictive behavioural transition (column) to a subsequent fictive program (row) within OK371>GCaMP6m control data (N=28) within the 10 to 20 minute time period ("On") representative of the time period of the neuromodulator perfusion period and > 20 minute time period ("Post") representative of the time period of the neuromodulator wash period. "On" statistical comparisons made between the average transition probabilities

 in the 10-20 minute period across all control isolated 3rd instar ventral nerve cords compared to the average

transition probabilities in the 0-10 minute period. "Post" statistical comparisons are made between the average

transition probabilities in the 20-30 minute period across all control data compared to the average transition

probabilities in the 0-10 minute period. (**B**) The average fictive transition probabilities across all the octopamine

(OA) concentrations within the application ("On") and post-application ("Wash") period. (**C**) The average fictive

transition probabilities across all the tyramine (TA) concentrations within the application ("On") and post-application

("Wash") period. Post-hoc comparison were conducted with p-values: < .5 (*), < .01 (**), <.001 (***) shown. Only

statistically significant bars are shown.

798
Supplementary Figure 3: Tyramine Shows No Inhibition of Motor Root Activity nor Calcium Baseline

(**A**) Effects of 50µM tyramine bath application on posterior abdominal motor nerve root activity demonstrating (**Ai**)

subtle changes to firing patterns (N=3). The effect of 50µM tyramine bath application on posterior abdominal motor

root (**Aii**) bursting frequency, (**Aiii**) average burst duration, (**Aiv**), proportion of time (%) without activity > 5%

change from baseline, (**Av**) average burst amplitude, and (**Avi**) baseline average for the pre-application ("Saline"),

application ("TA"), and post-application ("Wash") periods. (**B**) Representative effects of 200µM tyramine bath

application (dark red) (N=11) on the (**Bi**) calcium activity on the motor neuron pool (abdominal segment 3, left: A3

(L)). (**Bii**) The baseline calcium fluorescence for all isolated ventral nerve cord preparations under 200µM tyramine

bath application (dark red). The effect of 200µM tyramine bath application on (**Biii**) average calcium amplitude
across fictive programs and (**Biv**) baseline calcium signal. Post-hoc comparison were conducted with p-values: <
.5 (*), < .01 (**), <.001 (***) shown. Only statistically significant bars are shown.

Supplementary Figure 4: Pulsed Stimulation of *tdc*²⁺ Neurons Induces Fictive Forward Bias

(**A**) Representative T3(L) and A6(R) motor root electrophysiology trace demonstrating promotion of fictive forward

activity during pulsed stimulation of CsChrimson>*tdc*²⁺ (0.01s, 40Hz, 400 repeats = 10s total stimulation) for

experimental (+ retinol) compared to control (- retinol) 3rd instar *Drosophila* larva. (**B**) Quantification of motor root

bursting at A6 demonstrating (**i**) peak rate and (**ii**) burst duration changes upon pulsed stimulation (On10) relative

to before (Pre10) and after (Post10) stimulation period. (**C**) Quantification of fictive bias before, during, and after

pulsed stimulation in preparations where fictive activity could be accurately inferred (N=6). (**D**) The relative fictive

activity difference for fictive forward activity during (on) vs before (pre) stimulation (blue) and for anterior asymmetric

fictive activity after (post) vs before (pre) stimulation (green). Comparisons followed by post-hoc was conducted

with p-values: $< .5$ (*), $< .01$ (**), $<.001$ (***) shown. Only statistically significant bars are shown.

Supplementary Figure 5: Relative Fictive Wave Bias Induced by Restricted Stimulation of *tdc*²⁺ Neurons

Restricted *tdc*²⁺ depolarisation in the (A) SOG, (B), thoracic ganglion (T1-3), (C) anterior abdominal ganglion (A1-

3), and (D) posterior abdominal ganglion (A6-8) induces different fictive wave bias in different stimulation periods

(N > 2) across all preparations (N = 7). Plots show normalised bias of fictive forward activity rate (m^{-1}) / fictive

backward activity rate (m^{-1}) with gray dashed line indicative of relative fictive wave bias before restricted stimulation.

Note, across all stimulation periods there are a diverse array of bias changes (i.e., below the level of normalised

bias before stimulation = more fictive backward wave bias, above the level of normalised bias before stimulation =

more fictive forward wave bias). Comparisons followed by post-hoc was conducted with p-values: < .5 (*), < .01

(**), <.001 (***) shown. Only statistically significant bars are shown.

834 **References**

[1] J. Francis *et al.*, 'Inhibitory circuit motifs in Drosophila larvae generate motor program
diversity and variability', *PLOS Biol.*, vol. 23, no. 4, p. e3003094, Apr. 2025, doi:
10.1371/journal.pbio.3003094.
[2] T. Jovanic *et al.*, 'Competitive Disinhibition Mediates Behavioral Choice and Sequences
in Drosophila.', *Cell*, vol. 167, no. 3, pp. 858-870.e19, Oct. 2016, doi:
10.1016/j.cell.2016.09.009.
[3] T. Roeder, 'The control of metabolic traits by octopamine and tyramine in
invertebrates', *J. Exp. Biol.*, vol. 223, no. 7, p. jeb194282, Apr. 2020, doi:
10.1242/jeb.194282.
[4] T. Farooqui, 'Review of octopamine in insect nervous systems', *Open Access Insect
Physiol.*, p. 1, Jun. 2012, doi: 10.2147/OAIP.S20911.
[5] Y. Pei, A. Asif-Malik, and J. J. Canales, 'Trace Amines and the Trace Amine-
Associated Receptor 1: Pharmacology, Neurochemistry, and Clinical Implications',
Front. Neurosci., vol. 10, Apr. 2016, doi: 10.3389/fnins.2016.00148.
[6] M. Kutsukake, A. Komatsu, D. Yamamoto, and S. Ishiwa-Chigusa, 'A tyramine receptor
gene mutation causes a defective olfactory behavior in Drosophila melanogaster',
Gene, vol. 245, no. 1, pp. 31–42, Mar. 2000, doi: 10.1016/S0378-1119(99)00569-7.
[7] M. Monastirioti, M. Gorczyca, J. Rapus, M. Eckert, K. White, and V. Budnik,
'Octopamine immunoreactivity in the fruit fly *Drosophila melanogaster*', *J. Comp.
Neurol.*, vol. 356, no. 2, pp. 275–287, May 1995, doi: 10.1002/cne.903560210.
[8] Y. Nagaya, M. Kutsukake, S. I. Chigusa, and A. Komatsu, 'A trace amine, tyramine,
functions as a neuromodulator in Drosophila melanogaster', *Neurosci. Lett.*, vol. 329,
no. 3, pp. 324–328, Sep. 2002, doi: 10.1016/S0304-3940(02)00596-7.
[9] M. Selcho, D. Pauls, B. El Jundi, R. F. Stocker, and A. S. Thum, 'The Role of
octopamine and tyramine in *Drosophila* larval locomotion', *J. Comp. Neurol.*, vol. 520,
no. 16, pp. 3764–3785, Nov. 2012, doi: 10.1002/cne.23152.
[10] M. Vömel and C. Wegener, 'Neuroarchitecture of Aminergic Systems in the Larval
Ventral Ganglion of Drosophila melanogaster', *PLoS ONE*, vol. 3, no. 3, p. e1848, Mar.
2008, doi: 10.1371/journal.pone.0001848.
[11] T. Farooqui, 'Octopamine-Mediated Neuromodulation of Insect Senses', *Neurochem.
Res.*, vol. 32, no. 9, pp. 1511–1529, Jul. 2007, doi: 10.1007/s11064-007-9344-7.
[12] M. Selcho, 'Octopamine in the mushroom body circuitry for learning and memory',
Learn. Mem., vol. 31, no. 5, p. a053839, May 2024, doi: 10.1101/lm.053839.123.
[13] S. C. Hoyer *et al.*, 'Octopamine in Male Aggression of Drosophila', *Curr. Biol.*, vol. 18,
no. 3, pp. 159–167, Feb. 2008, doi: 10.1016/j.cub.2007.12.052.
[14] I. K. Yakovlev, 'Effects of Octopamine on Aggressive Behavior in Red Wood Ants',
Neurosci. Behav. Physiol., vol. 48, no. 3, pp. 279–288, Mar. 2018, doi:
10.1007/s11055-018-0561-0.
[15] P. A. Stevenson, V. Dyakonova, J. Rillich, and K. Schildberger, 'Octopamine and
Experience-Dependent Modulation of Aggression in Crickets', *J. Neurosci.*, vol. 25, no.
6, pp. 1431–1441, Feb. 2005, doi: 10.1523/JNEUROSCI.4258-04.2005.
[16] K. Mizutani, H. Ogawa, J. Saito, and K. Oka, 'Fictive locomotion induced by octopamine
in the earthworm', *J. Exp. Biol.*, vol. 205, no. 2, pp. 265–271, Jan. 2002, doi:
10.1242/jeb.205.2.265.
[17] I. Orchard, J. M. Ramirez, and A. B. Lange, 'A Multifunctional Role for Octopamine in
Locust Flight', *Annu. Rev. Entomol.*, vol. 38, no. 1, pp. 227–249, Jan. 1993, doi:
10.1146/annurev.en.38.010193.001303.

[18]   J. C. Ormshaw and C. J. H. Elliott, 'Octopamine boosts snail locomotion: behavioural
883   and cellular analysis', *Invert. Neurosci.*, vol. 6, no. 4, pp. 215–220, Nov. 2006, doi:
884   10.1007/s10158-006-0031-1.
885   [19]   T. Roeder, 'TYRAMINE AND OCTOPAMINE: Ruling Behavior and Metabolism', *Annu.*
886   *Rev. Entomol.*, vol. 50, no. 1, pp. 447–477, Jan. 2005, doi:
887   10.1146/annurev.ento.50.071803.130404.
888   [20]   D. J. Candy, 'The regulation of locust flight muscle metabolism by octopamine and
889   other compounds', *Insect Biochem.*, vol. 8, no. 3, pp. 177–181, Jan. 1978, doi:
890   10.1016/0020-1790(78)90070-7.
891   [21]   Y. Li *et al.*, 'Octopamine controls starvation resistance, life span and metabolic traits in
892   Drosophila', *Sci. Rep.*, vol. 6, no. 1, p. 35359, Oct. 2016, doi: 10.1038/srep35359.
893   [22]   A. C. Koon and V. Budnik, 'Inhibitory Control of Synaptic and Behavioral Plasticity by
894   Octopaminergic Signaling', *J. Neurosci.*, vol. 32, no. 18, pp. 6312–6322, May 2012, doi:
895   10.1523/JNEUROSCI.6517-11.2012.
896   [23]   A. C. Koon *et al.*, 'Autoregulatory and paracrine control of synaptic and behavioral
897   plasticity by octopaminergic signaling', *Nat. Neurosci.*, vol. 14, no. 2, pp. 190–199, Feb.
898   2011, doi: 10.1038/nn.2716.
899   [24]   Z. Yang, Y. Yu, V. Zhang, Y. Tian, W. Qi, and L. Wang, 'Octopamine mediates
900   starvation-induced hyperactivity in adult *Drosophila*', *Proc. Natl. Acad. Sci.*, vol. 112,
901   no. 16, pp. 5219–5224, Apr. 2015, doi: 10.1073/pnas.1417838112.
902   [25]   A. Sujkowski, D. Ramesh, A. Brockmann, and R. Wessells, 'Octopamine Drives
903   Endurance Exercise Adaptations in Drosophila', *Cell Rep.*, vol. 21, no. 7, pp. 1809–
904   1823, Nov. 2017, doi: 10.1016/j.celrep.2017.10.065.
905   [26]   B. Brembs, F. Christiansen, H. J. Pflüger, and C. Duch, 'Flight Initiation and
906   Maintenance Deficits in Flies with Genetically Altered Biogenic Amine Levels', *J.*
907   *Neurosci.*, vol. 27, no. 41, pp. 11122–11131, Oct. 2007, doi:
908   10.1523/JNEUROSCI.2704-07.2007.
909   [27]   S. Saraswati, L. E. Fox, D. R. Soll, and C. Wu, 'Tyramine and octopamine have
910   opposite effects on the locomotion of *Drosophila* larvae', *J. Neurobiol.*, vol. 58, no. 4,
911   pp. 425–441, Mar. 2004, doi: 10.1002/neu.10298.
912   [28]   L. E. Fox, D. R. Soll, and C.-F. Wu, 'Coordination and Modulation of Locomotion
913   Pattern Generators in *Drosophila* Larvae: Effects of Altered Biogenic Amine Levels by
914   the Tyramine β Hydroxlyase Mutation', *J. Neurosci.*, vol. 26, no. 5, pp. 1486–1498, Feb.
915   2006, doi: 10.1523/JNEUROSCI.4749-05.2006.
916   [29]   H. Hashemzadeh-Gargari and W. Otto Friesen, 'Modulation of swimming activity in the
917   medicinal leech by serotonin and octopamine', *Comp. Biochem. Physiol. Part C Comp.*
918   *Pharmacol.*, vol. 94, no. 1, pp. 295–302, Jan. 1989, doi: 10.1016/0742-8413(89)90182-
919   5.
920   [30]   B. Johnson and R. Harris-Warrick, 'Aminergic modulation of graded synaptic
921   transmission in the lobster stomatogastric ganglion', *J. Neurosci.*, vol. 10, no. 7, pp.
922   2066–2076, Jul. 1990, doi: 10.1523/JNEUROSCI.10-07-02066.1990.
923   [31]   S. C. Kinnamon, L. W. Klaassen, A. E. Kammer, and D. Claassen, 'Octopamine and
924   chlordimeform enhance sensory responsiveness and production of the flight motor
925   pattern in developing and adult moths', *J. Neurobiol.*, vol. 15, no. 4, pp. 283–293, Jul.
926   1984, doi: 10.1002/neu.480150405.
927   [32]   J. M. Ramirez and K. G. Pearson, 'Octopaminergic modulation of interneurons in the
928   flight system of the locust', *J. Neurophysiol.*, vol. 66, no. 5, pp. 1522–1537, Nov. 1991,
929   doi: 10.1152/jn.1991.66.5.1522.
930   [33]   M. J. Alkema, M. Hunter-Ensor, N. Ringstad, and H. R. Horvitz, 'Tyramine Functions
931   Independently of Octopamine in the Caenorhabditis elegans Nervous System', *Neuron*,
932   vol. 46, no. 2, pp. 247–260, Apr. 2005, doi: 10.1016/j.neuron.2005.02.024.
933   [34]   N. Schützler *et al.*, 'Tyramine action on motoneuron excitability and adaptable
934   tyramine/octopamine ratios adjust *Drosophila* locomotion to nutritional state', *Proc. Natl.*
935   *Acad. Sci.*, vol. 116, no. 9, pp. 3805–3810, Feb. 2019, doi: 10.1073/pnas.1813554116.

[35] B. Mulloney, L. D. Acevedo, and A. G. Bradbury, 'Modulation of the crayfish swimmeret rhythm by octopamine and the neuropeptide proctolin', *J. Neurophysiol.*, vol. 58, no. 3, pp. 584–597, Sep. 1987, doi: 10.1152/jn.1987.58.3.584.

[36] A. D. McClellan, G. D. Brown, and P. A. Getting, 'Modulation of swimming in Tritonia: excitatory and inhibitory effects of serotonin', *J. Comp. Physiol. A*, vol. 174, no. 2, Feb. 1994, doi: 10.1007/BF00193792.

[37] K. Nishikawa and Y. Kidokoro, 'Octopamine inhibits synaptic transmission at the larval neuromuscular junction in Drosophila melanogaster', *Brain Res.*, vol. 837, no. 1–2, pp. 67–74, Aug. 1999, doi: 10.1016/S0006-8993(99)01676-5.

[38] Z.-T. Zhu, A. C. Munhall, and S. W. Johnson, 'Tyramine excites rat subthalamic neurons in vitro by a dopamine-dependent mechanism', *Neuropharmacology*, vol. 52, no. 4, pp. 1169–1178, Mar. 2007, doi: 10.1016/j.neuropharm.2006.12.005.

[39] A. Donini and A. B. Lange, 'Evidence for a possible neurotransmitter/neuromodulator role of tyramine on the locust oviducts', *J. Insect Physiol.*, vol. 50, no. 4, pp. 351–361, Apr. 2004, doi: 10.1016/j.jinsphys.2004.02.005.

[40] C. McClung and J. Hirsh, 'The trace amine tyramine is essential for sensitization to cocaine in Drosophila', *Curr. Biol.*, vol. 9, no. 16, pp. 853–860, Aug. 1999, doi: 10.1016/S0960-9822(99)80389-3.

[41] M. A. El-Brolosy and D. Y. R. Stainier, 'Genetic compensation: A phenomenon in search of mechanisms', *PLOS Genet.*, vol. 13, no. 7, p. e1006780, Jul. 2017, doi: 10.1371/journal.pgen.1006780.

[42] A. A. Prinz, D. Bucher, and E. Marder, 'Similar network activity from disparate circuit parameters', *Nat. Neurosci.*, vol. 7, no. 12, pp. 1345–1352, Dec. 2004, doi: 10.1038/nn1352.

[43] E. Marder, 'Neuromodulation of Neuronal Circuits: Back to the Future', *Neuron*, vol. 76, no. 1, pp. 1–11, Oct. 2012, doi: 10.1016/j.neuron.2012.09.010.

[44] E. Marder, 'Variability, compensation, and modulation in neurons and circuits', *Proc. Natl. Acad. Sci.*, vol. 108, no. supplement_3, pp. 15542–15548, Sep. 2011, doi: 10.1073/pnas.1010674108.

[45] M. Selcho, D. Pauls, A. Huser, R. F. Stocker, and A. S. Thum, 'Characterization of the octopaminergic and tyraminergic neurons in the central brain of *Drosophila* larvae', *J. Comp. Neurol.*, vol. 522, no. 15, pp. 3485–3500, Oct. 2014, doi: 10.1002/cne.23616.

[46] S. R. Pulver, T. G. Bayley, A. L. Taylor, J. Berni, M. Bate, and B. Hedwig, 'Imaging fictive locomotor patterns in larval Drosophila.', *J. Neurophysiol.*, vol. 114, no. 5, pp. 2564–2577, Nov. 2015, doi: 10.1152/jn.00731.2015.

[47] W. C. Lemon *et al.*, 'Whole-central nervous system functional imaging in larval Drosophila.', *Nat. Commun.*, vol. 6, p. 7924, Aug. 2015, doi: 10.1038/ncomms8924.

[48] A. Sordillo and C. I. Bargmann, 'Behavioral control by depolarized and hyperpolarized states of an integrating neuron', *eLife*, vol. 10, p. e67723, Nov. 2021, doi: 10.7554/eLife.67723.

[49] J. K. Pirri, A. D. McPherson, J. L. Donnelly, M. M. Francis, and M. J. Alkema, 'A Tyramine-Gated Chloride Channel Coordinates Distinct Motor Programs of a Caenorhabditis elegans Escape Response', *Neuron*, vol. 62, no. 4, pp. 526–538, May 2009, doi: 10.1016/j.neuron.2009.04.013.

[50] Y. Kagawa-Nagamura, K. Gengyo-Ando, M. Ohkura, and J. Nakai, 'Role of tyramine in calcium dynamics of GABAergic neurons and escape behavior in Caenorhabditis elegans', *Zool. Lett.*, vol. 4, no. 1, p. 19, Dec. 2018, doi: 10.1186/s40851-018-0103-1.

[51] J. Meng *et al.*, 'A tonically active master neuron modulates mutually exclusive motor states at two timescales', *Sci. Adv.*, vol. 10, no. 15, p. eadk0002, Apr. 2024, doi: 10.1126/sciadv.adk0002.

[52] R. Marley and R. A. Baines, 'Dissection of Third-Instar *Drosophila* Larvae for Electrophysiological Recording from Neurons', *Cold Spring Harb. Protoc.*, vol. 2011, no. 9, p. pdb.prot065656, Sep. 2011, doi: 10.1101/pdb.prot065656.

[53] S. H. Cole, G. E. Carney, C. A. McClung, S. S. Willard, B. J. Taylor, and J. Hirsh, 'Two
Functional but Noncomplementing Drosophila Tyrosine Decarboxylase Genes', *J. Biol.*
Chem., vol. 280, no. 15, pp. 14948–14955, Apr. 2005, doi: 10.1074/jbc.M414197200.
[54] A. Edelstein, N. Amodaj, K. Hoover, R. Vale, and N. Stuurman, 'Computer Control of
Microscopes Using µManager', *Curr. Protoc. Mol. Biol.*, vol. 92, no. 1, Oct. 2010, doi:
10.1002/0471142727.mb1420s92.
[55] C. A. Schneider, W. S. Rasband, and K. W. Eliceiri, 'NIH Image to ImageJ: 25 years of
image analysis', *Nat. Methods*, vol. 9, no. 7, pp. 671–675, Jul. 2012, doi:
10.1038/nmeth.2089.
[56] J. Schindelin *et al.*, 'Fiji: an open-source platform for biological-image analysis', *Nat.*
Methods, vol. 9, no. 7, pp. 676–682, Jul. 2012, doi: 10.1038/nmeth.2019.
[57] W. J. Heitler, 'Practical tools for analysing rhythmic neural activity', *J. Neurosci.*
Methods, vol. 185, no. 1, pp. 151–164, Dec. 2009, doi:
10.1016/j.jneumeth.2009.09.009.
[58] J. Berni, 'Genetic dissection of a regionally differentiated network for exploratory
behavior in Drosophila larvae.', *Curr. Biol. CB*, vol. 25, no. 10, pp. 1319–1326, May
2015, doi: 10.1016/j.cub.2015.03.023.
[59] M. Monastirioti, C. E. Linn, Jr., and K. White, 'Characterization of **Drosophila**
Tyramine β-Hydroxylase Gene and Isolation of Mutant Flies Lacking Octopamine', *J.*
Neurosci., vol. 16, no. 12, pp. 3900–3911, Jun. 1996, doi: 10.1523/JNEUROSCI.16-12-
03900.1996.
[60] A. Debnath, P. D. E. Williams, and B. A. Bamber, 'Reduced Ca2+ transient amplitudes
may signify increased or decreased depolarization depending on the neuromodulatory
signaling pathway', *Front. Neurosci.*, vol. 16, p. 931328, Jul. 2022, doi:
10.3389/fnins.2022.931328.
[61] M. Winding *et al.*, 'The connectome of an insect brain', *Science*, vol. 379, no. 6636, p.
eadd9330, Mar. 2023, doi: 10.1126/science.add9330.
[62] T. Stürner *et al.*, 'Comparative connectomics of the descending and ascending neurons
of the *Drosophila* nervous system: stereotypy and sexual dimorphism', Jun. 06, 2024.
doi: 10.1101/2024.06.04.596633.
[63] A. A. Zarin, B. Mark, A. Cardona, A. Litwin-Kumar, and C. Q. Doe, 'A multilayer circuit
architecture for the generation of distinct locomotor behaviors in *Drosophila*', *eLife*, vol.
8, p. e51781, Dec. 2019, doi: 10.7554/eLife.51781.
[64] A. Fushiki, M. F. Zwart, H. Kohsaka, R. D. Fetter, A. Cardona, and A. Nose, 'A circuit
mechanism for the propagation of waves of muscle contraction in Drosophila', *eLife*,
vol. 5, p. e13253, Feb. 2016, doi: 10.7554/eLife.13253.
[65] A. Carreira-Rosario *et al.*, 'MDN brain descending neurons coordinately activate
backward and inhibit forward locomotion', *eLife*, vol. 7, p. e38554, Aug. 2018, doi:
10.7554/eLife.38554.
[66] A. Zhao *et al.*, 'High-ambition climate action in all sectors can achieve a 65%
greenhouse gas emissions reduction in the United States by 2035', *Npj Clim. Action*,
vol. 3, no. 1, p. 63, Jul. 2024, doi: 10.1038/s44168-024-00145-x.
[67] U. Černe *et al.*, 'Ca^{2+} excitability of glia to neuromodulator octopamine in *Drosophila*
living brain is greater than that of neurons', *Acta Physiol.*, vol. 241, no. 2, p. e14270,
Feb. 2025, doi: 10.1111/apha.14270.
[68] S. El-Kholy, F. Stephano, Y. Li, A. Bhandari, C. Fink, and T. Roeder, 'Expression
analysis of octopamine and tyramine receptors in Drosophila', *Cell Tissue Res.*, vol.
361, no. 3, pp. 669–684, Sep. 2015, doi: 10.1007/s00441-015-2137-4.
[69] L. M. Sherer *et al.*, 'Octopamine neuron dependent aggression requires dVGLUT from
dual-transmitting neurons', *PLOS Genet.*, vol. 16, no. 2, p. e1008609, Feb. 2020, doi:
10.1371/journal.pgen.1008609.
[70] C.-L. Wu, M.-F. M. Shih, P.-T. Lee, and A.-S. Chiang, 'An Octopamine-Mushroom Body
Circuit Modulates the Formation of Anesthesia-Resistant Memory in Drosophila', *Curr.*
Biol., vol. 23, no. 23, pp. 2346–2354, Dec. 2013, doi: 10.1016/j.cub.2013.09.056.

[71]   J. Li and W. L. Shew, 'Tuning network dynamics from criticality to an asynchronous
1044   state', *PLOS Comput. Biol.*, vol. 16, no. 9, p. e1008268, Sep. 2020, doi:
1045   10.1371/journal.pcbi.1008268.
1046