Multimodal learning decodes the global binding landscape of chromatin-associated proteins

Jimin Tan[1,2,3#*], Xi Fu[4,5*], Xinyu Ling[1*], Shentong Mo[6], Jiangshan Bai[1], Raúl Rabadán[4,5], David Fenyö[2,7], Jef D. Boeke[2,7,8], Aristotelis Tsirigos[3,9#], Bo Xia[1,2,10#]

Gene Regulation Observatory, Broad Institute of MIT and Harvard, Cambridge, MA, USA
Institute for Systems Genetics, NYU Grossman School of Medicine, New York, NY, USA
Division of Precision Medicine, NYU Grossman School of Medicine, New York, NY, USA
Program of Mathematical Genomics, Department of System Biology, Columbia University, New York, NY, USA
Department of Biomedical Informatics, Columbia University, New York, NY, USA
School of Computer Science, Carnegie Mellon University, Pittsburgh, PA, USA
Department of Biochemistry and Molecular Pharmacology, NYU Grossman School of Medicine, New York, NY, USA
Department of Biomedical Engineering, NYU Tandon School of Engineering, Brooklyn, NY, USA
Department of Pathology, NYU Grossman School of Medicine, New York, NY, USA
Society of Fellows, Harvard University, Cambridge, MA, USA
*Co-first Authors
#Corresponding Authors: Jimin.Tan@nyulangone.org; Aristotelis.Tsirigos@nyulangone.org; xiabo@broadinstitute.org

Abstract

Chromatin-associated proteins (CAPs), including over 1,600 transcription factors, bind directly or indirectly to the genomic DNA to regulate gene expression and determine a myriad of cell types. Mapping their genome-wide binding and co-binding landscape is essential towards a mechanistic understanding of their functions in gene regulation and resulting cellular phenotypes. However, due to the lack of techniques that effectively scale across proteins and biological samples, their genome-wide binding profiles remain challenging to obtain, particularly in primary cells. Here we present Chromnitron, a multimodal foundation model that accurately predicts CAP binding landscapes across hundreds of proteins in unseen cell types. Via *in silico* perturbation experiments, we show that the model learned principles of CAP binding from multimodal features including DNA sequence motifs, chromatin accessibility levels, and protein functional domains. Applying Chromnitron to study cell fate transitions, we discovered novel CAPs regulating the T cell exhaustion process. Furthermore, Chromnitron can predict the dynamic CAP binding landscapes during development, revealing the global orchestration of protein and regulatory element activities in neurogenesis. We expect Chromnitron to accelerate discovery and engineering in regulatory genomics, particularly in human primary cells, and empower future therapeutic opportunities.

INTRODUCTION

How does the genome encode gene activities to determine the full spectrum of cell types and their functions in a multicellular organism? While DNA sequence carries the shared basal genetic information across all cell types, chromatin-associated proteins (CAPs) establish cell-type-specific gene regulatory programs in both normal and disease states[1–5]. In the human genome, there are an estimated 2,500 CAPs, including approximately 1,600 transcription factors (TF) and around 900 other transcriptional co-factors and chromatin regulators[6–8]. These proteins interact with over one million putative *cis*-regulatory elements (CREs), mediated by chromatin states, to confer gene regulatory programs[9,10]. Such CAP-DNA interactions can be achieved both directly and indirectly. Through DNA binding domains (DBDs), TFs directly recognize and bind to short consensus DNA sequence motifs within appropriate chromatin states[11–16]. Transcriptional co-factors and chromatin regulators typically interact with DNA indirectly via histones, other CAPs, or chromatin-associated RNAs[17–20]. Together, the specificity of CAP binding across the genome and cell types is collectively determined by DNA sequence, cell-type-specific chromatin states, and protein structural features encoded in the amino acid sequence.

Mapping the genome-wide binding landscape of all CAPs across cell types is crucial for decoding cell-type-specific gene regulatory programs and predicting cell fate transitions[21–23]. However, traditional experimental approaches for mapping CAP binding profiles, such as ChIP-seq, have limited scalability across proteins and biological samples, especially primary cells due to their heterogeneity and rarity[24,25]. As a result, only around 1,200 human CAPs have been mapped, primarily in cultured cell lines[26–29]. Compared to experimental approaches, *in silico* predictions are highly scalable and hold the promise to generalize to primary cells that are otherwise not measurable. Recent advances in machine learning have shown early promises in predicting CAP binding profiles from DNA sequence or chromatin features[30–33]. However, existing models are mostly unimodal and limited in their ability to generalize across proteins and cell types. We hypothesize that a multimodal foundation model is required to learn a comprehensive set of rules governing CAP-chromatin interactions and enable transferable prediction across unseen cell types and chromatin states.

Here we present Chromnitron (Chromatin omni-modal transformer), a biologically informed
multimodal deep neural network that integrates information from DNA sequence, chromatin
accessibility signals, and CAP residues to predict cell type-specific CAP binding landscapes (Fig.
1). Chromnitron was pre-trained on 1,105 CAP ChIP-seq datasets across five cell types to learn
the general principles of CAP-chromatin interaction and individually fine-tuned for each CAP to
further improve prediction performance. The trained model can predict the global CAP binding
landscape across 767 distinct CAPs in cell types that were unseen during training. We show that
Chromnitron can be used to discover critical regulators at key regulatory elements and categorize
co-regulation structures across CAPs and CREs. Applying Chromnitron to study cell fate
transitions, we discovered ZNF865 and ZNF766 as novel regulators of the T cell exhaustion
process. Furthermore, Chromnitron can predict global CAP binding landscape and reveal the
orchestration of CAPs and CRE activities during continuous cell fate transitions in developmental
systems. Together, these results demonstrate the effectiveness of a multimodal foundation model
in understanding the principles of CAP-chromatin interactions in gene regulatory programs and
accelerating discoveries in cell fate determination.

85
Fig. 1: Chromnitron predicts the binding landscape of chromatin-associated proteins.

a. Schematic of varied CAP types and their interaction modes with chromatin. **b.** Chromnitron
model architecture. Chromnitron uses a cascaded architecture across multimodal encoding,
modality fusion, transformer, and decoding modules to predict CAP binding profiles. **c.** Two-phase
training strategy showing cross-CAP pre-training using shared weights (grey) followed by
individual fine-tuning for each CAP with LoRA, visualized as weight deltas (ΔW). **d.** Predicted
CAP-binding landscape, presented as two-dimensional heatmap of binding intensity across
genomic regions for 767 CAPs (CAP axis). CAPs are colored by clusters based on binding
similarity. **e.** Downstream applications of Chromnitron to accelerate discoveries. Top: discovering
CRE-specific regulators at single locus. Middle: genome-wide discovery of regulatory structures
across CAPs and CREs. Bottom: mapping global CAP-binding dynamics across continuous cell
fate transition trajectories.

RESULTS

Multimodal learning of CAP binding principles

To create a model that learns the principles of CAP binding, we designed a multimodal architecture

that integrates information from DNA sequence features, cell-type-specific chromatin states, and

CAP structural features encoded in the amino acid sequences (Fig.1b and Supplementary Fig. 1).

This design allows Chromnitron to learn features from all modalities, enabling accurate predictions

of CAP binding that generalize across new genomic sequences and cell types. To efficiently learn

and integrate input information across different modalities, Chromnitron employs a cascaded

architecture including encoding, modality fusion, transformer, and decoding modules (Fig. 1b and

Supplementary Fig. 1). Each input modality was first processed by its own encoder module. DNA

sequence and chromatin accessibility signals were separately encoded by two convolutional

modules, while CAP residues were encoded with a pre-trained protein language model[34] (Methods).

These encoders yield latent embeddings that were subsequently merged in two stages. First, DNA

sequence and chromatin accessibility embeddings were concatenated to form an integrated

chromatin embedding aligned on the genomic axis. Second, we fused the protein embedding to the

integrated chromatin embedding on the length dimension (Methods). This fusion enables the

following transformer to attend every protein residue to every location in the chromatin embedding

and allow implicit learning of both intra- and cross-modality attentions. In the final decoder

module, the chromatin embedding was processed by a transposed convolutional module to

generate predicted CAP binding intensities at single nucleotide resolution (Fig. 1b and

Supplementary Fig. 1).

Training a high-performance foundation model requires large-scale and standardized data[34–37]. To

minimize experimental noise across proteins and data sources, we first curated a collection of

1,105 qualified ChIP-seq from five different cell types, covering 767 unique CAPs, as well as the

corresponding cell-type-specific ATAC-seq datasets (Supplementary Fig. 2a-d, Methods). Next,

we developed a robust bioinformatic infrastructure, Chrom2Vec, to convert raw sequencing data

(i.e. FASTQ files) to training-ready data vectors (Supplementary Fig. 2e, Methods). Chrom2Vec

generates CAP binding profiles that are comparable across proteins and cell types, effectively

minimizes noise-induced errors to ensure model performance and robustness[35,38,39].

Next, we trained Chromnitron in two-phases: cross-CAP pre-training and individual fine-tuning

for each CAP. Compared with individualized training across each CAP from scratch, this two-

phase training strategy substantially reduced the amount of total training time and allows the model

to learn integrated binding principles across CAPs (Methods). However, the pre-training step can

be computationally heavy due to the compounded sampling space across genomic region, cell type,

and CAP dimensions (Supplementary Fig. 3a-c). To improve pre-training efficiency, we developed

an algorithm, OuterMix, to reuse embeddings generated by each encoding step (Supplementary

Fig. 3d-f , Methods). This approach dramatically reduced sampling complexities and accelerated

data loading and encoding by five folds during pre-training. We then used low-rank adaptation

(LoRA) to fine-tune individual sub-models for each CAP[40]. During pre-training we observed

distinct learning dynamics across CAP groups, possibly driven by their differential binding

mechanisms. For instance, performance of cohesin-related CAPs quickly improved in early stages,

whereas MAF-related CAPs did not increase in performance until the middle stage of pre-training

(Supplementary Fig. 4a). After pre-training, individual CAP fine-turning substantially increased

prediction performance overall (Supplementary Fig. 4a-b).

Fig. 2: High-quality prediction across chromatin contexts.

a. Paired Chromnitron prediction (dark) and ChIP-seq (light) tracks for 16 CAPs in leave-out chromosome region. **b.** Dimensionality reduction (t-SNE) visualization of predicted (circle) and experimental (squares) binding profiles from a subset of CAPs. Colors define CAPs clustered with similar binding profiles. **c-d.** CAP-binding prediction performance distribution (**c**), measured by Pearson's *r*, and their correlation with training data quality (**d**, Methods). Green, blue, and grey colors mark high- (302), medium- (233) and low- (232) performance CAP categories, respectively (Methods, and Supplementary Fig. 5-6). **e.** Distribution of correlation coefficients (Pearson's *r*) between experimental ChIP-seq signal and chromatin accessibility in HepG2 cells. **f.** Scatter plots of CAP binding versus chromatin accessibility signals. Each CAP presents both experimental ChIP-seq signal (grey) and predicted binding signal (blue). **g.** Predicted and experimental CAP binding signals in both high- and low-accessibility regions. Presented heatmaps include only 2kb genomic regions in leave-out chromosome (chr10). Acc: accessibility. **h.** Example CAP binding tracks comparing Chromnitron, motif accessibility, maxATAC and Enformer predictions, all aligned together with the experimental ChIP-seq data in leave-out chromosome 10. **i.** Performance comparison (Pearson's *r*) between Chromnitron and other methods across CAPs.

High-quality and cross-modality CAP binding predictions

Accurate prediction is the foundation for downstream biological discovery. Chromnitron was trained to predict high-quality CAP binding profiles across 767 proteins at single-nucleotide resolution and can generalize to unseen chromatin and cell types. In leave-out genomic regions

(chromosome 10), Chromnitron accurately predicts CAP-binding profiles comparable to
experimental measurements (Fig. 2a). Projecting both Chromnitron predictions and experimental
ChIP-seq profiles into t-distributed stochastic neighbor embedding (t-SNE) space, the predicted
data form clusters that align closely with their experimental counterparts, demonstrating high-
fidelity prediction performance (Fig. 2b). We quantified prediction performance by calculating
Pearson's correlation (r) between Chromnitron-predicted and experimental signals in the leave-
out chromosome. Across the 767-CAP catalogue, more than half of the CAPs have Pearson's
correlation coefficients larger than 0.5 (Fig. 2c, Methods). We found that Chromnitron's
predictions match or outperform the quality of experimental replicates (Supplementary Fig. 4c-e).
Chromnitron also achieves high-quality prediction when generalizing to new cell types and CAPs
via ablation studies (Supplementary Fig. 4f-g). We notice that Pearson's correlation is highly
sensitive to noise at background regions of the genome. Correlating with data quality, we found
that prediction performance of individual CAP improves consistently as training-data quality
increases (i.e. reduced background noise level) (Fig. 2d and Methods). Despite the modest
correlation coefficients in some CAPs ($r < 0.5$), Chromnitron-predicted CAP-binding peaks have
high correspondence to ChIP-seq peaks (Supplementary Fig. 5-6).

Most existing tools for inferring candidate CAP binding from ATAC-seq signals are more sensitive
to highly accessible region and less so to lowly accessible regions[41–43]. We found that
experimentally measured CAP binding profiles includes substantial binding events at low-
accessibility or nominally inaccessible regions (Fig. 2e-g). Comparing experimental CAP binding
profiles with chromatin accessibility, we observed a wide range of relationships, including positive
correlation (e.g. MAX and MYC), negative correlation (e.g. CEBPB and EZH2), and non-linear
patterns (e.g. CTCF and FOSL2, Fig. 2f). Since Chromnitron is not restricted by accessibility
signals but rather by learning the binding principles across different chromatin states, we observed
that Chromnitron faithfully captures different CAP binding modes as observed in experimental
data (Fig. 2f-g). Together, these results demonstrate that Chromnitron, with its multimodal design,
successfully learned the intricate relationships between CAP binding and local chromatin
accessibility states.

Chromnitron is a multimodal foundation model for predicting cell-type-specific CAP-binding

profiles in unseen cell types during training. Although no other model employs the same

multimodal framework, we nevertheless benchmarked Chromnitron against several approaches

that aimed to infer or predict CAP binding profiles. These approaches include "motif accessibility"

(MA) score, which multiplies TF motif strength by chromatin accessibility as a first-order

approximation of CAP binding. Comparing with ground truth ChIP-seq data, the MA score tends

to be noisy and shows weak correlation due to its frequent false positive and false negative signals

(Fig. 2h, Supplementary Fig. 7). Chromnitron prediction outperforms MA score by a large margin

(Fig. 2i). Next, we compared Chromnitron with two machine learning approaches, maxATAC and

Enformer[31,44]. MaxATAC was trained on sparse CAP-binding peak files and predicts the

probability of CAP binding, while Enformer is a DNA sequence-based model trained with multi-

task learning and unable to generalize to new cell types or CAPs[31,44]. Using experimental ChIP-

seq data as the reference, we found that Chromnitron substantially outperforms both methods

across various metrics (Fig. 2h-i, and Supplementary Fig. 7-9).

Fig. 3: *In silico* **perturbation reveals principles governing CAP binding.**

a. Quantification of CAP-specific sequence impact scores through base-resolution *in silico* mutations and Chromnitron prediction (Methods). **b.** Representative impact motifs for CEBPB, MYC, JUNB, and MAFF. **c.** Impact motifs generated from the same locus for two different CAPs (CTCF and FOS). **d.** Hierarchical clustering using impact motif cross-correlations of CAPs. **e.** Chromatin accessibility perturbation using *in silico* scaling multipliers (Methods). **f-g.** Accessibility scaling results at CTCF (**f**) and ARID3A (**g**) binding sites. Left: predicted CAP binding changes responding to perturbation radius and accessibility multiplier (Methods). Middle: heatmap of predicted binding intensity upon perturbing accessibility at fixed 500bp window, centered at binding peaks. Right: CAP binding changes (top) and the rate of change (bottom) as a function of accessibility multiplier. **h.** Schematic of *in silico* mutation for CAP protein sequences. Upon mutation, predicted CAP binding intensities of mutant CAPs were compared to reference CAP to calculate protein residue impact scores (Methods). **i.** Violin plot distribution of impact scores across CAP residues, stratified by domain annotations. **j.** Protein–protein network inferred from predicted CAP co-localization patterns, colored by groups. **k.** Per-residue binding impact

231 scores for SOX13 and CREB1, with AlphaFold-predicted structures highlighting high-impact
232 residues (HMG-box, KID, b-ZIP domains) which bind proximately to DNA.
233

Multimodal *in silico* perturbations reveal principles of CAP binding

The multimodal design allows Chromnitron to learn CAP binding principles from different

modalities and improve prediction performance. To investigate whether the learned principles

conform to known biological mechanisms, we performed *in silico* perturbations to interrogate the

model across all three input modalities, starting with *in silico* DNA sequence mutagenesis. For a

locus with CAP binding, we sequentially mutated every nucleotide to each of the three other

nucleotides, followed by quantification of the mutation impact by computing the resulting changes

in predicted binding (Fig. 3a). We found that these single-locus, per-nucleotide impact scores

recapitulate TF consensus motifs that are typically derived from hundreds to thousands of

experimental binding sequences[45,46] (Fig. 3b). Applying *in silico* mutagenesis across all CAPs at a

given locus, Chromnitron can identify multiple, spatially distinct impact motifs within the locus,

reflecting the precise sequence-dependent CAP binding syntax (Fig. 3c).

Chromnitron's sensitivity to individual nucleotide alternations enables systematic discovery of

uncharacterized CAP-DNA interactions. By collecting impact motifs across multiple loci for each

CAP and clustering them by similarity, we found several modules corresponding to distinct CAP

families, such as the C/EBP, bHLH, and AP-1 families, and a large set of unique impact motifs,

many uncharacterized (Fig. 3d, and Supplementary Fig. 10, Methods). Moreover, *in silico*

mutagenesis with Chromnitron identified impact motifs for proteins that lack canonical DNA

binding domains, such as cohesin subunits resembling CTCF motif (Fig. 3d and Supplementary

Fig. 10). We found these motifs often resemble the sequence features of their frequent co-binding

partners, suggesting a promising strategy for discovering co-binding partners by comparing impact

motif matrices (Fig. 3d and Supplementary Fig. 10). Last, Chromnitron is sensitive to non-coding

variants and captures their effect on CAP binding in both gain-of-binding and loss-of-function

mutation events (Supplementary Fig. 10).

Chromatin accessibility differences represent different epigenetic states across genomic regions

and cell types. Therefore, we evaluated the effect of perturbing chromatin accessibility levels (Fig.

3e). By scaling accessibility intensity *in silico* across different window sizes, we found that binding

prediction is indeed sensitive to accessibility changes, and in a CAP-dependent manner. For

example, the relationship between CTCF binding and accessibility level follows a logarithmic

curve, saturating at relatively low accessibility levels (Fig. 3f), whereas ARID3A binding

intensifies linearly with increasing accessibility (Fig. 3g). These results confirmed that

Chromnitron learned both the non-linearity and specificity in CAP binding, echoing our previous

results correlating ChIP-seq to chromatin accessibility (Fig. 2e-g).

Last, we carried out *in silico* mutagenesis on CAP amino acid sequences to assess whether

Chromnitron learned informative features from protein residues. We randomly mutated amino

acids in segments with a sliding window approach and quantified the impact of mutant residues

by calculating the resulting changes in predicted CAP binding (Fig. 3h, Methods). Scanning

through the entire protein sequence, we observed strong enrichment in evolutionary conserved

DNA-binding domains such as bZIP and E2F (Fig. 3i-j, Supplementary Fig. 11). For examples,

we observed high impact scores within the direct DNA binding regions for both SOX13 and

CREB1, which were confirmed by both domain annotation and AlphaFold-predicted protein-DNA

binding structures (Fig. 3k). In contrast, functional domains that do not contribute to sequence

specificity of CAP binding, such as bromodomain and PHD finger domain, tends to have much

lower impact scores (Fig. 3k, Supplementary Fig. 11). Collectively, these *in silico* perturbation

results demonstrate that Chromnitron effectively integrates multimodal information to uncover

both canonical and hidden principles of CAP binding.

Fig. 4: Predicting novel regulatory CAPs at individual regulatory element.

a. Predicting cell-type-specific CAP binding using ATAC-seq datasets across conditions. **b.** Dimensionality reduction (UMAP) showing T cell fate trajectory from naïve (T0) to exhausted (T3) states (top), together with chromatin accessibility changes at a *TOX* promoter-proximal regulatory element. **c.** Predicted CAP binding signals at this *TOX* promoter-proximal CRE, highlighting ZNF865 binding. **d.** ZNF865 binding stood out among other CAPs at the *TOX* CRE during late pseudo-time stage. **e.** Scatter plot of ZNF865 experimental ChIP-seq binding (grey) and predicted binding (blue) versus chromatin accessibility, indicating that ZNF865 preferentially binds at low-accessibility regions. **f.** Chromatin accessibility signals at *TOX* promoter and the ZNF865-bound CRE across pseudo-time. **g.** Chromatin accessibility at the ZNF865-bound CRE (top) and the corresponding ZNF865 impact motif (bottom). **h.** Schematic of *in vitro* model of T cell exhaustion through persistent anti-CD3/CD28 stimulation, coupled with CRISPR/Cas9-based gene knockout in primary T cells. **i-j.** Flow cytometry quantification of T cell exhaustion markers (**i**, PD1, TIGIT, LAG3 and TIM3) and cytokines (**j,** IFN-γ, IL-2, and TNF-α) across negative control, non-targeting control (NTC), *TOX* knockout, or *ZNF865* knockout conditions. Bar plots show mean ± s.d. for n = 3 independent replicates. Statistical comparisons performed using two-sided unpaired Student's t-tests with significance denoted as *, P < 0.05; **, P < 0.01; ***, P < 0.001.

Discovering CRE-specific regulators at single locus

The collection of CAPs binding at a cis-regulatory element (CRE) determines its regulatory

function. Given the chromatin accessibility signals at a genomic locus, Chromnitron predicts high-

quality CAP-binding profiles that enables the discovery of key regulators (Fig. 4a). We applied

Chromnitron to study T cell fate transition during chronic stimulation. Persistent antigen exposure

pushes tumor-infiltrated T cells into a dysfunctional or exhausted state characterized by enhanced

expression of inhibitory cell surface receptors, reduced cytokine production, and activation of

TOX[47–54]. Single-cell ATAC-seq (scATAC-seq) of tumor-infiltrated T cells has captured the

transition from naïve T cell to exhausted T cells[55]. To pinpoint the locus-specific regulators that

control *TOX* expression, we applied Chromnitron to track CAP binding dynamics at the *TOX* locus

across the T cell exhaustion trajectory derived from scATAC-seq data (Fig. 4b, Methods).

Surprisingly, ZNF865, an uncharacterized C2H2 zinc-finger protein, stands out as a unique

regulator at a *TOX* promoter-proximal regulatory element with a steady increase of binding along

the exhaustion pseudo-time (Fig. 4c-d). We noticed that ZNF865 preferentially binds at relatively

inaccessible regions, including this *TOX* promoter-proximal element, whose accessibility is

substantially lower than that of the promoter region (Fig. 4e-f). Further *in silico* mutagenesis

revealed that ZNF865 recognizes a previously uncharacterized CA-repeat binding pattern, which

explains in part of its unique binding to the *TOX* locus (Fig. 4g). These results underscore

Chromnitron's ability to leverage fine-grained DNA sequence features and nuanced changes in

accessibility for discovering novel regulators of gene expression.

To validate this finding, we employed an established *in vitro* T cell exhaustion model using

persistent anti-CD3 antibody or anti-CD3/CD28 cocktail stimulation[56–58] (Methods). After

persistent stimulation, human T cells were analyzed by flow cytometry for cell surface exhaustion

markers and cytokine production (Methods). As expected, exhaustion markers including PD1,

TIGIT, LAG3, and TIM3 were significantly upregulated in the persistently stimulated group, while

IFN-γ, IL-2, and TNF-α expression level were significantly reduced, confirming successful

induction of T cell exhaustion phenotype *in vitro* (Supplementary Fig. 12). Consistently, both

ZNF865 and *TOX* expression increase along the progression of persistent T cells stimulation

(Supplementary Fig. 13a-b). To evaluate the function of ZNF865, we electroporated Cas9

ribonucleoprotein with sgRNAs targeting *ZNF865*, *TOX* (positive control), and a non-targeting

control (NTC) before persistent anti-CD3/CD28 stimulation of the T cells (Methods). Consistent

with the hypothesis, we found that knockout *ZNF865* reduced *TOX* expression level (Supplementary Fig. 13c-d). Notably, *ZNF865* knockout led to a significant reduction in surface exhaustion markers compared to NTC, closely resembling the effects observed in the *TOX* knockout group (Fig. 4i). Moreover, ZNF865-deficient T cells produced elevated cytokine production, suggesting restoration of effector function (Fig. 4j).

Fig. 5: Genome-wide characterization and discovery of novel regulatory CAPs driving T cell exhaustion.

a. Workflow of genome-wide characterization of co-regulation structures across CAPs and CREs between paired cell conditions. **b.** The distribution of effector T cells (Teff, blue) and exhausted T cells (Tex, red) in UMAP (top) and the chromatin accessibility differences between the two cell population (bottom, Methods). **c.** Predicted CAP binding difference matrix between Teff and Tex cells, clustered by both co-regulating CAPs (rows) and co-regulated gene (promoter) groups (columns). A zoom-in plot highlights a subset of CAP modules (C0, C2, C3) and gene groups (G0, G4, G9) enriching reported Tex marker genes (Methods). **d.** Elbow plot of enriched CAPs ranked

by log2 fold change of Tex over Teff. **e.** Distribution of reported Tex marker genes (CTLA4,
PDCD1, TIGIT, HAVCR2) in co-regulated CRE groups (top) and their corresponding co-
regulating CAP clusters (bottom). **f.** CRISPR dropout screens in human CD8[+] T cells using *in vitro*
chronic stimulation model (Methods). **g.** Scatter plot of targeting-gene sgRNA enrichment p-
values in the knockout (x-axis) and knockdown (y-axis) CRISPR screens, highlighting *ZNF766* as
the top hit in both conditions. **h-i.** Flow cytometry quantification of T cell exhaustion markers (**i**,
PD1, TIGIT, LAG3 and TIM3) and cytokine secretion levels (**j**, IFN-γ, IL-2, and TNF-α) across
negative control, non-targeting control (NTC), or *ZNF766* knockout conditions. Bar plots show
mean ± s.d. for n = 3 independent replicates. Statistical comparisons performed using two-sided
unpaired Student's t-tests with significance denoted as *, P < 0.05; **, P < 0.01; ***, P < 0.001.
362
363

Genome-wide discovery of CAP-dependent regulatory structures

Having demonstrated Chromnitron's ability to discover candidate regulators at a single regulatory

element of interest, we next asked whether it could generalize these insights to uncover genome-

wide regulatory structures. Cell fate transitions are regulated by groups of CAPs interacting with

CREs across the genome. Given chromatin accessibility datasets of differential cell states,

Chromnitron can predict their global CAP binding changes to identify co-regulating CAPs and co-

regulated CREs (Fig. 5a). We applied this *in silico* framework to dissect the genome-wide

regulatory structures driving T cell exhaustion, focusing on the group of co-regulated genes

corresponding to differentially accessible gene promoters between effector T cells and exhausted

T cells (Fig. 5b). Using Chromnitron, we predicted CAP binding profiles across these promoters

in both groups of T cell population (Methods). Comparing the CAP binding signals between the

two populations, we calculated the differential CAP binding matrix (CRE-by-CAP) and clustered

on both axes to uncover modules of co-regulating CAPs and co-regulated CREs (Fig. 5c). The top

co-regulating CAP module (C0) includes both previously identified CAPs such as BHLHE40 and

KDM6A[58,59], and uncharacterized CAPs such as ZNF766 (Fig. 5d). Mapping known exhaustion

surface markers onto our co-regulated CRE modules revealed strong enrichment in group G4, G9

and G0. Corresponding to these CRE groups, we found that co-regulating CAP clusters C0, C2,

and C3 exhibit the strongest differential binding, suggesting potential regulatory relationship (Fig.

5e).

We hypothesized that these co-regulating CAPs in the above modules include uncharacterized

upstream regulators of T cell exhaustion progress. To assess the functions of these candidate CAPs,

we conducted a small-scale CRISPR-based pooled screen using the *in vitro* T cell exhaustion

model. We designed two separates sgRNA libraries comprising 370 sgRNAs targeting 65 genes,

respectively optimized for gene knock-out and transcriptional knock-down via lentiviral

transduction (Methods). Following sgRNA delivery, human primary CD8$^+$ T cells were subjected

to persistent anti-CD3 treatment, with T cell proliferation serving as the phenotypic readout (Fig.

5f, Methods). Across both knock-out and knock-down screens, ZNF766 consistently stood out as

the most prominent regulator, exhibiting the strongest impact on T cell persistence during chronic

stimulation (Fig. 5g). To independently validate this result, we performed targeted gene knockout

of ZNF766 via RNP electroporation (Supplementary Fig. 14, Methods). Subsequent flow

cytometry analyses revealed significant decreases in surface exhaustion markers, along with

increases in effector cytokine production (Fig. 5h-i). RNA-seq following *ZNF766* knockout

showed gene expression changes that reflect a shift toward reduced exhaustion (Supplementary

Fig. 14). Together, these findings suggest that ZNF766 may play an important role in regulating T

cell exhaustion.

Fig. 6: Global dynamics of CAP binding and CRE activities during cell differentiation.

a. Single cell ATAC-seq is used to compute pseudo-time and identify differentially accessible CREs across the genome for Chromnitron inference. **b.** Three-dimensional tensor representing predicted CAP binding landscapes across developmental time. **c.** Heatmap of predicted CAP binding signals across CREs grouped by developmental stages. Rows and columns represent CAPs and CREs activities, respectively. **d.** T-SNE projection of CREs colored by activity clusters across pseudo-time. Clusters were defined by CAP binding profiles and sorted from inactive (blue) to active (red) based on the average binding activity. **e-h**. Vector fields of directional CRE activity transition to their following developmental stage, projected on the t-SNE map in **d**.

Global orchestration of CAP binding and CRE activities

Cell fate transition typically goes through drastic state changes to achieve functional maturity in developmental process. To characterize the shifting gene regulatory programs, we applied Chromnitron to predict genome-wide CAP-binding landscapes along cell fate transition trajectory using single cell chromatin accessibility data (Fig. 6a-b). The predicted three-dimensional CAP-

binding tensor across genomic position, CAP, and time offers unprecedented analytical scales to

characterize how CAP binding and CRE activities are globally orchestrated (Fig. 6b). We applied

this analytical framework to Purkinje neuron maturation during first trimester neurodevelopment[60].

After generating the CAP binding landscapes across Purkinje neuron maturation, we first

transformed the three-dimensional data tensor into a CAP-by-CRE matrix with sub-grouping by

time (Fig. 6c, Methods). We observed a clear transition of CRE activities, measured by global

CAP occupancy, from neural progenitor cells to mature Purkinje cells, with a sharp transition

between neuroblast to early Purkinje state (Fig. 6c).

To dissect stage-specific transitions in CRE activities along the developmental time trajectory, we

embedded all CREs in a t-SNE map using their CAP-binding features. Subsequent clustering of

all CREs revealed the global distribution of CRE identities ranging from inactive to active states

across genomic space and temporal stages (Fig. 6d, Supplementary Fig. 15a-d, Methods). Next,

we tracked each CRE from its current developmental stage to the following stage, computing a

vector field that visualizes its trajectory (Methods). The resulting vector field uncovers the

migration snapshots of CRE identities (Fig. 6e-h). From neural progenitor (Prog) to neuroblast

state 1 (NBL1), stem-cell-specific active CREs start to drift toward the inactive region (Fig. 6e).

From neuroblast state 2 (NBL2) to early Purkinje cells (Purk1), most inactive CREs start to shift

towards activation, representing the strongest wave of CAP-driven CRE activity changes, while

those already active CREs continue to migrate to the inactive state (Fig. 6f). Purk1 to Purk2 interval

marks the final major wave of CRE activity shift, where the last set of inactive CREs complete

their transition (Fig. 6g). Starting from Purk2, all CREs converge to a relatively stable

configuration to form the terminal regulatory landscape (Fig. 6h).

The three-dimensional tensor can be interrogated along any axis to offer new insights about CAP

binding and CRE activities. Using developmental time as the organizing axis, we can track the

binding trend for each CAP on every CRE. This analysis allows us to uncover cohorts of early-

and late-acting regulators during Purkinje cell maturation trajectory, most of which follow

monotonic gain- or loss-of-binding trends (Supplementary Fig. 15e). In parallel, the tensor can be

transformed using the CRE dimension as the feature, followed by re-clustering CAPs. We found

distinct CAP binding signatures across developmental time. For example, unlike transcription

activators, cohesin-related protein, chromatin remodelers, and repressors assume relatively stable,
factor-specific binding patterns (Supplementary Fig. 15f-g). These CAP bindings tend to have
minor changes within their own group and rarely shift between groups, consistent with the
regulatory constraints imposed by their associated functions on chromatin (Supplementary Fig.
15g). Together, these multidimensional analyses illustrate how the CAP-binding tensor offers an
unprecedented lens on the dynamic orchestration of CAP binding and CRE activities, setting the
stage for systematic discovery of gene regulation programs that controls cell fate transition, and
ultimately, guiding precision rewiring of cell fate.

DISCUSSION
A major challenge in understanding the molecular mechanisms of human health and disease is to
systematically characterize how chromatin-associated proteins configure gene regulatory
programs. Here we show that Chromnitron, a biologically informed multimodal foundation model,
can learn the core principles of CAP-chromatin interactions and achieve accurate prediction of
global CAP-binding landscapes across hundreds of proteins. The multimodal and cell-type-
specific design allows the model to generalize to virtually any cell type, enabling systematic
discovery of core transcriptional regulators and their downstream impacts on cellular phenotypes.
Chromnitron acts as a prism to transform single chromatin accessibility signal into a spectrum of
functional genomics data with high-fidelity. By harnessing the power of such high-throughput
prediction, we discovered previously uncharacterized regulators of T-cell exhaustion and mapped
the global dynamics of CAP binding and CRE activities during neurodevelopment with
unprecedented scale. Chromnitron exemplifies how biologically grounded genomics foundation
models can accelerate hypothesis generation and enable targeted experimental validation,
revealing new biological insights.

Beyond its practical applications, Chromnitron also uncovers both known CAP-chromatin binding
rules and non-canonical interactions at low-accessibility or absence-of-accessibility regions with
previously unrecognized sequence motifs. Compared with sequence-based models that excel at
pinpointing nucleotide-level attribution of regulatory activity, Chromnitron learns the fundamental
rules of CAP binding through a multimodal architecture, enabling generalizable high-performance
prediction across cell types and chromatin states. Moreover, Chromnitron's multimodal design

allows the integration of pre-trained DNA language models as alternative or additional encoders

to further increase training efficiency and performance[32–34,36,43,44,61].

Despite its generalizability and high prediction performance, Chromnitron nonetheless has

limitations. As a machine learning model, Chromnitron's prediction accuracy depends on the

training data quality. Current training datasets are derived primarily from conventional ChIP-seq

experiments, which are subject to variable antibody quality, lack of dosage considerations, and

batch effects. Although substantial computational efforts were made to denoise and standardize

the data, not all processed ChIP-seq reach high quality. These challenges underscore the need for

generating standardized, high-quality CAP-binding datasets. Importantly, the pre-trained

foundation model can be readily updated by fine-tuning with improved CAP-binding datasets.

Generating standardized CAP-binding datasets covering the remaining uncharacterized human

CAPs will further improve Chromnitron's prediction performance and scalability across diverse

biological systems.

Biologically informed multimodal foundation models like Chromnitron offer a new paradigm in

human biology research. By accurately predicting large-scale functional data from information-

rich measurement, these models can enable high-throughput *in silico* screening to aid hypotheses

generation and validation, therefore accelerating biological discovery and new therapeutic

development.

Methods:

Model architecture

We designed Chromnitron (Chromatin Omni-modal Transformer) to integrate three information

streams that jointly determine CAP-chromatin binding: DNA sequence, local chromatin

accessibility, and the amino-acid sequence of the assayed CAP. Chromnitron comprises three

successive components: modality-specific encoders, a multimodal self-attention module and a

decoder (Supplementary Fig. 1).

The model has three modality-specific encoders. The encoders for DNA sequence and chromatin

accessibility consists of similar convolutional blocks (ConvBlocks). Each block begins with a

strided Conv1D layer that covers large local windows while halving the sequence length. The

convolution is followed by group normalization (groups = 1) and a Gaussian Error Linear Unit

(GELU) activation function[62]. Feature maps then traverse 3 residual ConvBlocks (kernel = 9;

stride = 1) that maintain length while expanding the hidden dimensionality to 384. Outputs from

the DNA and accessibility encoders were projected to the same embedding of 512×384. The two

embeddings were fused on the feature dimension to form a joint embedding of size 512×768. The

joint embedding was scaled down to a 512×384 chromatin embedding with a header for

subsequent fusion with the protein embedding.

For the protein encoder, we leveraged a pre-trained protein language model ESM2 as a protein

feature extractor[34]. Running ESM2 inference during training can add to the computational

overhead. To increase model training efficiency, we extracted and cached the protein features

before the training step. A non-redundant list of human transcription factors was assembled by

merging all proteins for which we had ChIP-seq data and the available proteins in UniProt database.

To encode the embeddings, we used the publicly available ESM2 transformer with 36 layers and

3 billion parameters (esm2_t36_3B_UR50D, hidden size = 2560). For each protein, we requested

the hidden representations of the final, 36th transformer block. Since ESM2 accepts up to 1,024

tokens, we split any long proteins into non-overlapping 1,024-residue segments, each embedded

independently. The resulting matrices were concatenated along the sequence dimension to

reconstitute the full-length protein representation. The number of residues was capped at 4,096 to

accommodate most of the full-length CAP sequences while keeping a low compute cost. The per-

residue embeddings were converted to NumPy arrays and stored in a compressed npz file. These

cached embeddings are loaded at run-time, down-sampled by the Chromnitron protein header, and

fused with genomic features in the multimodal attention module.

We implemented the attention mechanism in transformer to fuse contextual information both

within and across modalities[63]. Chromatin feature embeddings and protein embeddings were

concatenated along the length dimension, separated by a single zero vector that marks the modality

boundary. Positional information was added via fixed sinusoidal encodings. The resulting fused

feature embedding has a length of 769 (512 from chromatin embedding, 256 from protein

embedding, and a separator with a length of 1). The fused embedding was processed by 16 layers

of pre-layer-normalized transformer. Each transformer layer has 8 attention heads, 384 hidden
state dimension, 2,048 feed-forward layer dimension, and a dropout rate of 0.1. Self-attention
allows every genomic position to attend not only to distant nucleotides but also cross-modal to any
residue of the corresponding CAP.
546
547 After transformer module, we only kept features corresponding to genomic positions and
548 forwarded them to the decoder. The decoder has four ConvTranspose blocks successively up-
549 sample the latent representation to the original 8,192-bp resolution while preserving contextual
550 information through residual connections. A final convolution header produces a one-dimensional
551 per-base binding profile for each target CAP.
552
553 Pre-trained Chromnitron model has 72 million parameters. Each LoRA fine-tuned sub-model
554 weight 9 million parameters. The fully fine-tuned Chromnitron model with pre-trained and LoRA
555 weights has over 6.9 billion parameters in total.
556
Genomic data collection
To obtain training data with more standardized quality, we primarily focused on consortium-
generated datasets for consistency. We developed a systematic approach to collect and curate high-
quality CAPs ChIP-seq datasets from the ENCODE database[9]. Our data collection strategy focused
on obtaining non-revoked and non-archived datasets with sufficient sequencing depth to ensure
reliable modeling. We required a minimum of 40 million reads to achieve adequate genome
coverage for CAPs ChIP-seq and ATAC-seq, while immunoprecipitation control experiments
required 100,000 reads. Chromatin accessibility information is essential for understanding the
regulatory context of transcription factor binding. To ensure experimental coherence and enable
integrative analysis, we retained only those CAPs ChIP-seq datasets that had good-quality ATAC-
seq data in the corresponding cell type, resulting to 1,105 qualified ChIP-seq from five different
cell types, covering 767 unique CAPs.
569
Genomic data preprocessing with Chrom2Vec
To prepare precise and highly consistent training data, we designed Chrom2Vec, a modular
bioinformatic platform that streamlines training data processing for Chromnitron. Chrom2Vec
platform starts from raw FASTQ files and delivers training-ready data tensors. Chrom2Vec is
based on Apache Airflow and orchestrates next-generation-sequencing processing for both ChIP-
seq and ATAC-seq. It implements configurable pipelines that handle distributed workers, tracking,
and reproducible containers, performs robust assay-specific normalization, and signal
compression to optimize data storage. Pipelines are described by lightweight Directed Acyclic
Graphs (DAGs) that call a library of preprocessing operators, which are Airflow wrappers around
containerized shell or Python scripts to ensure maximum reproducibility. While tasks are being
scheduled and executed by Airflow, the metadata are streamed to Neo4j for full data-lineage
tracking. To reduce storage and I/O overheads without sacrificing data quality on key genomic
regions, we implemented an adaptive dynamic-bin compression algorithm. We observed as high
as 50% reduction in storage for some of the high-quality tracks. The final compressed training

dataset is around 952 GB in size. This architecture was optimized for machine-learning workflow
as it guarantees reproducibility, supports scaling, and writes outputs in chunked Zarr format which
is directly ingestible by the model and supports fast random-access during training.

To enable quantitative comparisons across cell types and experimental conditions, we performed
peak-based quantile normalization on bulk ATAC-seq, ChIP-seq and pseudo-bulk single cell
ATAC-seq data. We followed a two-step approach that first removed noise and then rescaled data.
The noise level can be represented by variations in the background, non-active genomic regions.
To first define active regions, we used cPeaks from Meng et al., a union of accessible peaks across
human cell types[64]. The background region is defined as its complement regions on the genome,
excluding ENCODE blacklist[65].

For ChIP-seq, we performed a two-stage correction that explicitly models both biochemical
background and sequencing bias. First, chromatin signal was extracted from immunoprecipitated
(IP) and matched input libraries. Using the pre-defined background regions, we calculated the
mean IP and input signal in the background bins. The input was rescaled to reach similar
background signal intensity as IP. The rescaled input was then subtracted from the IP track,
effectively removing experimental bias across the genome. To harmonize dynamic range across
ChIP-seq experiments, the corrected IP signal was divided by the average of the 99 to 99.9
percentile values with 1 percent step sizes, computed over the pre-defined active peak regions from
cPeaks. This normalization puts the strongest peaks near a value of one while retaining linearity at
regions with lower signal intensities. The result is a bias-corrected, unit-less collection of ChIP-
seq datasets that are comparable across CAPs and experiments, dramatically facilitating training
stability. After scaling the training datasets, we further measured a background-noise level for each
dataset by calculating standard deviation in the background chromatin regions defined above. The
training data quality metric was calculated by (1 – noise) for downstream model performance
evaluation.

For ATAC-seq, where a control library was not available, we removed noise from background
directly followed by percentile scaling. The algorithm first estimates systematic noise by averaging
the coverage signal over the same background regions. This background mean was subtracted
from every genomic position, eliminating basal accessibility level. We collected mean cPeak
signals, and computed the scaling constant based on top 99 to 99.9 percentiles similar to ChIP-seq.
Dividing the background-corrected track by this constant compresses the dynamic range so that
the signals of typical accessible loci fall between 0 and 1 (though exceptionally strong regulatory
regions can exceed 1), producing normalized ATAC-seq datasets that are consistent and
comparable across batches and cell types.

For single cell ATAC-seq data, we started with fragment files and processed them through a
pipeline with shared preprocessing steps as Chrom2Vec and generates quantitative chromatin
accessibility profiles. Fragment coordinates were standardized to the hg38 reference genome. Cells
with 1,000-50,000 fragments per cell were retained to exclude low-quality cells with insufficient

chromatin accessibility signal and potential doublets with artificially inflated fragment counts.
Individual cells were subsequently aggregated into pseudo-bulk profiles based on cell cluster
annotations (i.e. cell types or states) in original publication, ensuring each pseudo-bulk has over
10 million unique sequencing reads (or 20 million Tn5 transposase insertions). These accessibility
reads from all cells within each cluster were summed to generate high-resolution chromatin
accessibility tracks with improved signal-to-noise ratios compared to individual cells. Each
pseudo-bulk data represents a unique cell type or state, and was normalized by the same pipeline
described in the data normalization section.
634

Model pre-training and fine-tuning
Training data were assembled from the three input feature modalities and the target CAP binding
profiles. Each training sample unit was set to an 8,192-bp genomic region. For the two chromatin
input features, we extracted a five-channel (A, C, G, T, N) one-hot representation of the hg38 DNA
sequence and a single-channel chromatin-accessibility track (ATAC-seq Tn5-transposase insertion
frequency signal). To stabilize training, the ATAC-seq signal was transformed with a monotonic
log1p function. CAP binding profiles were retrieved from the normalized ChIP-seq coverage for
each CAP in every cell type. In a separate preprocessing step, we embedded the primary amino-
acid sequence of every CAP with the 36-layer ESM2 protein language model
(esm2_t36_3B_UR50D). This embedding was cached and loaded on to the protein header during
training.
646

Random samples from different modalities are recombined to form the final training batch.
Batched tensors were shaped as (batch, length, channels) for DNA and ATAC, and (batch, residues,
channels) for ESM2 embeddings. The model takes in these inputs and predicts the target CAP
binding profile of shape (batch, length). Validation loss was calculated as the mean-squared error
(MSE) between predicted and observed log1p-signal. We used Adam[66] as the optimizer with a
learning rate of 0.00002 and weight decay of 0. The scheduler used ReduceLROnPlateau strategy
with a patience of 5 epochs. After validation loss stabilizes at 390 epochs, we fine-tuned the model
for another 10 epochs for each CAP with low-rank adaptation (LoRA), updating the CAP-specific
sub-model parameters[67]. LoRA layers were directly applied to feedforward layers in the attention.
For convolution and convolution transpose layers, we custom-designed the LoRA layers, with
details described in the following sections.
658

The model was pre-trained on a node with 4 NVIDIA A100 80GB for 300 Hours with FP32 full
precision. The pre-training stage took around 1,200 GPU hours and has a total compute of 8.7e19
FLOPs (floating-point operations). The fine-tuned model was trained in parallel on a cluster with
32 NVIDIA A100 GPUs using TF32 precision, which has much higher throughput. Each fine-
tuning was set to run 8 hours for 10 epochs. The fine-tuning step was performed on all 767 CAPS,
leading to a total compute of 3.4e21 FLOPs, and took around 6,000 GPU hours using the lower
precision TF32 tensor format. The total amount of compute for training and fine-tuning is around
3.5e21 FLOPs.
667

Compared with previous methods that trained separate models for individual CAPs, Chromnitron
training employed a cross-CAP pretraining with per-CAP fine-tuning strategy which dramatically
increased both training efficiency and model performance. To measure the training efficiency
difference between individualized training approach versus LoRA finetuning, we used CTCF as
an example to compare the two approaches. We found that training the model from scratch took
more than 30 epochs to reach good performance, whereas fine-tuning from a pre-trained model
only took 1 epoch to reach comparable performance. As the pre-training compute is only a small
fraction of the fine-tuning compute ($< 3\%$ by estimation), assuming all CAPs requires a similar
level of compute training from scratch, we estimated that the pre-training with fine-tuning strategy
is at least one magnitude more efficient compared with individually training models for each CAP
from scratch. The stored LoRA weight matrix is also 8 times smaller than full weight matrix.
Meanwhile, pre-training substantially increases overall model performance through its generalized
learning of CAP-binding principles across different CAPs.
681
Efficient multimodal sampling with OuterMix during pre-training
To improve model training efficiency at the cross-CAP pre-training phase, we developed
OuterMix, an algorithm to generate Cartesian product of encoded features from randomly sampled
data points on each dimension as the training units. OuterMix maximizes throughput by re-using
modality-specific embeddings generated by each encoding step, thus dramatically reduces training
complexities, and increases data loading and encoding efficiency exponentially.
688
First of all, a naïve sampling approach for multimodal training through randomly selecting a fixed
number of examples for each batch would not be practical due to exponentially increased
complexities. Each sample consists of random examples from different modalities. Given a fixed
U number of samples from each modality with V modalities in total, the compute complexity
would be $O(U^V)$. Formally, if we are sampling $N \cdot M \cdot K$ samples in a batch representing the total
amount of genomic loci, chromatin feature, and CAPs during pre-training, the three encoders need
to encode all the samples. Using l, t, c to represent locus, cell type and CAP, respectively, the total
random sample space for a batch $\mathcal{B}_{rand}$ can be defined as:

$$\mathcal{B}_{rand} = \bigcup_{i=1}^{NMK}\{(l_i, t_i, c_i)\}, \ (l_i, t_i, c_i) \sim \mathcal{U}(\mathcal{L} \times \mathcal{T} \times \mathcal{C})$$

698 where each encoder needs to run across all samples.
699
700 OuterMix algorithm was designed to reuse embeddings generated by each encoder and reduce an
701 exponential complexity $O(U^V)$ down to a linear $O(UV)$ complexity, increasing data loading and
702 encoding efficiency exponentially, especially with large encoders and a high number of modalities.
703 With OuterMix, we started by randomly selecting fixed number of samples from each dimension
704 and produce a Cartesian product of loci (l_i), cell types (t_j) and proteins (c_k):

$$\mathcal{B}_{OuterMix} = \bigcup_{j=1}^{N}\bigcup_{i=1}^{M}\bigcup_{k=1}^{K}\{(l_i, t_j, c_k)\}$$

Here, each encoder only runs through samples from their predefined subset (Supplementary Fig.
3). In our implementation, the genome was first divided into non-overlapping 100kb chunks. In
each iteration, the data loader selects one genomic chunk per cell type, followed by sampling
8,192bp windows and applying lightweight augmentations. For each window, we randomly
selected up to 4 CAPs with their corresponding DNA sequence, ChIP-seq, and ESM2 data.
Modality-specific encoders then generate feature-specific embeddings, which are tiled and
concatenated to form the Cartesian product samples ready to be fed into the Transformer. This
design allows the same sequence-ATAC tensors to be paired with multiple proteins, increasing
Transformer trunk throughput while avoiding additional disk I/O and encoder overheads.
715
LoRA implementation during fine-tuning
Since Low-Rank Adaption (LoRA) was originally developed for fine-tuning natural language
applications such as Large Language Models (LLMs), its implementation for one-dimensional
convolution and convolution-transpose layers used in Chromnitron is not trivial[67]. Here we
implemented LoRA for both Conv1D and ConvTranspose1D. On a high level, LoRA for
convolution layers were implemented by freezing the original kernel and learning a low-rank
update that can be added onto the kernel at evaluation time. The 1-dimensional convolutional layer
weights are 3D tensors representing input dimension, kernel size and output dimension. In this
implementation, we reshaped input and kernel size into one dimension and used it to construct two
low-rank matrices.
726
For Conv1D, the convolutional kernel weights $W \in \mathbb{R}^{C_{in} \times C_k \times C_{out}}$ is kept frozen during fine-tuning,
with C_{in}, C_k, and C_{out} representing input size, number of kernels and output size, respectively. The
weight is separated into two low-rank matrices A and B with rank r, defined as:
$$A \in \mathbb{R}^{(r \cdot k) \times (C_{in} \cdot k)}, B \in \mathbb{R}^{C_{out} \times (r \cdot k)}$$
and is updated through $\Delta W = B A$. In the forward pass, the weight is added to the frozen weight
to apply fine-tuning changes:
$$y = \text{conv1d}\left(x, W + \frac{\alpha}{r} \cdot \Delta W\right)$$
where y is the convolution activation and α defines the strength for LoRA weights in which the
default is set to 1.
736
For ConvTranspose1D, the same decomposition is applied with a custom conv_transpose1d
forward pass applying the ΔW, similar to conv1d:
$$y = \text{conv1d\_transpose}\left(x, W + \frac{\alpha}{r} \cdot \Delta W\right)$$
We tested a range of ranks from 4 to 16 and did not observe significant performance difference.
To reduce weight size and fine-tuning efficiency, we used r = 4 for the final large-scale fine-tuning.
The custom LoRA implementation is available on https://github.com/tanjimin/LoRA.
743
Predicting CAP-binding profiles
During prediction, the LoRA checkpoint corresponding to each CAP was loaded together with the
backbone weights. Since Chromnitron prediction operates on individual 8,192bp windows, to

predict CAP-binding signals across the entire chromosome, we used a sliding window approach
with 5,120bp step size to create overlap between adjacent predictions. Chromnitron's predictions
are robust to window shifts, but the predicted signal at the window edge can have lower quality
due to the absence of flanking chromatin context. To mitigate this effect, we retained only the
central 6,144bp predicted profiles out of the 8,192bp as valid predictions. With a 5,120bp step size
and a 6,144 effective window size, adjacent predictions will have a 1,024bp effective overlap. To
further ensure a high-quality transition between the overlapped regions between adjacent
predictions, we implemented a weighted average in the overlapping regions to give more weights
to bases that are closer to the center of their prediction windows. The weighted signal $\bar{S}$ is an
average of left signal S_l and right signal S_r, weighted by their relative base location i in the region:
$\bar{S} = \frac{i}{1024} \cdot S_l + \frac{1024-i}{1024} S_r.$
758

Prediction performance evaluation
All prediction performance evaluations were conducted on the leave-out chromosome 10. To
compare with experimentally measured ChIP-seq profiles, we first predicted CAP-binding profiles
across all five cell types on chromosome 10. For each CAP, we extracted profiles from both
prediction and ChIP-seq profiles on the chromosome 10p arm between 6 Mb and 30 Mb. We
evaluated model performance by calculating a set of metrics (detailed in the following section)
between prediction and experimental ChIP-seq profiles in the leave-out chromosome across all
cell types. For cell-type-specific prediction performance evaluation, we first calculated the
experimental data difference and the predicted CAP-binding difference between HepG2 and K562
cells, the two cell types with the highest number of overlapping CAP profiles. We then applied the
same set of evaluation metrics to compare the difference signal between the two groups.
770

For a comprehensive comparison, we computed in total eight statistics, including Pearson's
correlation (r), Spearman's correlation (ρ), the coefficient of determination (R squared), F1 score,
area under the receiver-operating characteristic curve (AUROC), and area under the precision-
recall curve (AUPR) with both target-centric and prediction-centric labelling. We repeated the
statistics over a grid of different bin sizes and top-percentile subset of the bins. Predicted or ChIP-
seq signals were first converted to numerical vectors, with missing values set to zero.
Subsequently, the vectors were mean-reduced into non-overlapping bins of 50, 250, 500 or
1000bp. This multi-scale representation allows assessment of both fine-grained and coarse
regulatory patterns. The top-percentile bins measure how well the model can differentiate between
peaks and their relative enrichment level. For this, we set an exponential increment of percentile
using a list of 100% (1), 50% (1/2), 25% (1/4), 12.5% (1/8), …, 0.05% (1/2048). For AUROC and
AUPR calculation, we used the median value of the selected bins as threshold to separate positive
and negative labels. We also tested an alternative strategy to select most characteristic bins by
selecting both top and bottom percentile bins (Supplementary Fig. 6).
785

To categorize CAP-binding prediction performance, we focused on the Pearson's correlation (r)
as a simple metric to help interpreting the prediction results. We categorized CAP-binding
prediction performance with both validation performance and cell-type-specific performance on

the leave-out chromosome (chr10) for all CAPs in cell types where training data are available. The
validation performance was defined as the highest performance across all cell types, while the cell-
type-specific performance was calculated as described above. To establish a relationship between
validation performance and cell-type-specific performance, we correlated the validation
performance against cell-type-specific performance and found a general positive correlation. The
cell-type-specific performance is a relatively more challenging metric due to additional
experimental noises across cell types. We selected 187 CAPs that are shared between HepG2 and
K562 and fitted a line between validation performance and cell-type-specific performance. After
custom evaluation of prediction performance, we assigned CAPs to different performance groups
based on both validation and cell-type-specific prediction performance metrics. The specific
performance cutoff was chosen based on the fitted curve between the two metrics (Supplementary
Fig. 4f). We classified a CAP into high-performance group if its prediction performance has either
a validation performance > 0.6 or the corresponding cell-type-specific performance. CAPs with
either a validation performance > 0.4 or the corresponding cell-type-specific performance were
assigned in medium-performance group. The rest of CAPs were categorized into low-performing
group. Notably, although some CAPs were classified into low-performance group by Pearson's
correlation, their predicted binding profiles recapitulate ChIP-seq peaks (Supplementary Fig. 5-6).
Thus, the proposed CAP-performance categorization should be regarded as a practical metric for
interpreting prediction results, rather than a strict measure of biological validity.
808
Global comparison of prediction to experimental data at CAP-binding regions
To evaluate prediction performance across different chromatin backgrounds, we compared CAP-
binding signals by aligning Chromnitron predictions with matched ChIP-seq and ATAC-seq
profiles centered in peak regions. For each CAP of a particular cell type, three one-dimensional
signal tracks were collected: (i) Chromnitron-predicted CAP-binding signal, (ii) experimental
ChIP-seq data (reference), and (iii) ATAC-seq data to represent chromatin accessibility.
Chromnitron-predicted CAP-binding data on chromosome 10 were generated and documented, as
described in the previous performance evaluation section. Peaks were independently identified in
prediction, ChIP-seq and ATAC-seq by thresholding the signal at 0.5 (half of very high peaks),
merging contiguous bins above this level, and discarding events narrower than 100 bp. The mean
signal within each retained interval was recorded to generate a peak set annotated by genomic
coordinates and intensity.
821
To visualize and directly compare prediction to the reference CAP-binding data, all peaks detected
in Chromnitron prediction were stratified into two groups according to chromatin accessibility
signal. Each predicted peak was labelled according to its overlap with ATAC-seq peaks, generating
an accessible set and an inaccessible set. For every peak in each group, we extracted a 2kb genomic
window centered at the highest score in each peak. Finally, we visualized the predicted CAP-
binding signal, ChIP-seq signal, and ATAC-seq signals as heatmaps to evaluate global prediction
performance across chromatin background.
829
Ablation study

We performed ablation studies to measure how the model performs under different constraints.
This analysis includes performance evaluation across models trained without fine-tuning, and
models trained with missing cell types or CAPs to test their generalizability across modalities in
unseen contexts. We performed ablation studies on a random subset of chromosome 10 to reduce
computation and storage consumption. Our preliminary calculation found that for one set of
experiments, inferring 767 CAPs on chromosome 10 for five cell types could take two weeks and
6.5 TB of storage. Instead, we used Pearson's correlation coefficient calculated during the
validation step embedded in the evaluation process to save computation and storage. This
evaluation was performed on 500 random loci on chromosome 10 across all cell types and works
as a proxy of the genome-wide prediction performance statistics. The performance statistics used
for both the full Chromnitron model and ablated models were all calculated from validation step
using the same metrics.
843
844 To evaluate the performance improvements of fine-tuning, we extracted performance metrics from
845 the pre-trained model at its last epoch (390) and metrics from fine-tuned models at their last fine-
846 tuning epoch (≤ 10). The last fine-tuned epoch can change due to our early stopping strategy. Then,
847 we compared the performance individually across cell types and CAPs.
848
For cell-type-specific prediction evaluation through ablation study, we kept data from HepG2 and
K562 cells as the training set, and left out data from IMR-90, GM12878 and HCT116 cells as
testing. Chromosome 10 region was again used as leave-out chromosome. This model was then
pre-trained (i.e. without fine-tuning) and compared with the full Chromnitron pre-training model
in leave-out cell types. For CAP-specific evaluation, we randomly left out half of the CAPs (n=382)
and kept the other half (n=385) in the training set. Then we extracted the performance metrics from
the ablated model and compared with the full Chromnitron pre-training model on leave-out CAPs
(again in chromosome 10 regions).
857
858
Benchmarking prediction performance
Although there was no directly comparable model that used the same multimodality framework,
we benchmarked performance of Chromnitron against previous machine learning approaches that
aimed for similar goals, including Enformer and maxATAC, and the motif accessibility score as a
baseline approximation[31,44]. Detailed inference of CAP binding signal for each approach was
described below. For each benchmark, we compared the inferred CAP-binding tracks against the
experimental data reference (Chrom2Vec processed ENCODE ChIP-seq profiles). We calculated
eight different metrics across bins and top binding regions, similar to the procedure described in
the model performance evaluation section.
868
Enformer: We used GRCh38 reference genome sequence as input to Enformer. Chromosome 10
was divided into non-overlapping 100 kb windows; for each window, 196,608 bp centered on the
window mid-point were retrieved (zero-padded at chromosome ends) and one-hot encoded. Each
sequence was forwarded through the PyTorch Enformer implementation (EleutherAI/enformer-

official-rough) which has similar performance as the TensorFlow implementation. The model
returned an 896-bin output tensor per window which was cached on disk. CHIP channels relevant
to HepG2 or K562 cells were identified from human ChIP-seq targets. For every requested TF
channel, the central 50,176 bp (±50,176 bp) of each window were extracted to avoid model
padding artefacts. The selections were written into an array representing chromosome 10 at 1bp
resolution. Overlaps occurred only at window boundaries and were overwritten by later windows.
The final arrays were exported to BigWig files for visualization and comparison with other
approaches.
881
MaxATAC: Raw ATAC-seq data were obtained from ENCODE as BAM files (HepG2:
ENCFF939JOT, K562: ENCFF804MXV). These files were down-sampled to 40M reads and
deduplicated to generate the BAM files required for maxATAC pipeline. MaxATAC pre-
processing ("maxatac prepare") was performed with "-c chr10" to build the ATAC coverage tracks
necessary its prediction. For every CAP that is included in both Chromnitron and maxATAC
model, we used the corresponding sub-model to predict the CAP binding profile as a BigWig file
using "maxatac predict".
889
Motif-Accessibility: Non-redundant motif from Jeff Viestra
(https://resources.altius.org/~jvierstra/projects/motif-clustering/releases/v1.0/) were filtered to
only retain human CAPs. Motif scanning results on chr10 were converted into a dense 1-bp signal
for each motif with the corresponding matching scores. A 20-bp moving average was applied to
the dense track. The final results were stored as BigWig files. To weight motif by chromatin
accessibility, we loaded ATAC-seq for all five cell types (HepG2, K562, GM12878, IMR-90,
HCT116). For each cell type, the smoothed motif track was multiplied elementwise by the
corresponding ATAC-seq signal and stored as BigWig files.
898
In silico DNA sequence mutagenesis
To identify the impact of DNA sequence on the binding of a CAP, we focused on regions where
Chromnitron predicted intense CAP-binding signals. For each CAP-binding peak, we extended the
locus by ±3 kb and records its midpoint base location. We defined a 300-bp "mutation window"
(±150 bp) flanking the midpoint and systematically substituted every base within the window.
Subsequently, we performed _in silico_ saturated DNA base mutagenesis of the 300bp window, one
base at a time. For every position, we generated four sequences, each carrying a single A, C, G or
T substitution, and predicted the corresponding CAP-binding signal for each sequence. We
recorded the original sequence and its predicted CAP-binding signal as wild-type (WT) reference.
We then calculated the effect of every mutation as the difference between mutant and WT
predictions. The resulting inferred CAP-binding data at this window was stored as a $300 \times 4 \times$
8,192 tensor, with the three dimensions representing the number of mutation positions, number of
substitution nucleotides, and the window size of the CAP-binding prediction, respectively. The
signals represent the predicted CAP-binding intensities. To represent only the CAP-binding region,
the tensor was cropped from 8,192bp to the central 1kb and averaged, generating a 300×4 matrix
whose values represent CAP-binding intensities across the 300bp mutated positions and 4

nucleotides. This matrix served as the nucleotide-resolution perturbation signals for downstream
motif analyses and interpretation. For visualization purposes, the impact motif was normalized by
subtracting the median CAP-binding intensities across the 4 nucleotides (i.e. the mean binding
intensities of the middle two nucleotides) at each position. In essence, the impact motif is computed
from a single locus, differing from the conventional motif calling which computes consensus
sequence from a large number of loci.
921
Cross-CAP motif discovery and clustering
To analyze motifs across all CAPs, we generated weight matrices (represented by median-
subtracted CAP-binding signals as described above) for all CAPs at their top binding loci. These
weight matrices were used as input for TF-MoDISco to generate motifs for each individual CAP[68].
For every CAP, the tool scans up to 20,000 high-scoring 200-bp windows ("seqlets", or high-
importance windows of the sequences, as referred in TF-MoDISco), clustering them by k-mer
similarity and reports positive and negative patterns (i.e. sequels). After identifying relevant seqlets
for a CAP, we mapped their corresponding seqlets back to the genomic coordinates and applies a
strict quality control procedure: a pattern must contain at least one seqlet whose high (90th-
percentile) absolute weight difference is nontrivial (>0.1) after median subtraction. This procedure
excluded noisy results from the analyses. We then stored the remaining patterns for every CAP.
To pick the most representative seqlets for each CAP, we ranked them by (1) the number of
retained seqlets, (2) the overall motif intensity represented by the median difference value and (3)
the highest motif intensity represented by median 90th percentile difference. Patterns whose
seqlets overlapped more than ten distinct loci and repeated in different CAPs were discarded to
avoid trivial repeats.
938
Then, we built normalized weight matrices for cross-CAP comparison from the top ranked seqlets.
For each seqlet, we subtracted its per-position median and scaling by its mean absolute difference.
Each motif was flattened into a single vector (length by 4). Pairwise distances were computed as
the maximal cross-correlation between motif A and motif B or B's reverse complement. We
performed hierarchical clustering with Ward linkage based on the distance metrics, generating the
clustering heatmap for different motifs.
945
In silico **chromatin accessibility perturbations**
We measured the changes of predicted CAP binding upon *in silico* scaling of chromatin
accessibility signal to examine Chromnitron's sensitivity to local chromatin states. We employed
a scaling approach through multiplying the ATAC-seq profiles by a defined factor (multiplier) to
alter accessibility intensities while preserving peak shape. Meanwhile, the size of the perturbation
window (i.e. number of bases) also matters in this *in silico* perturbation. Thus, we created an *in*
silico perturbation grid by combining multiplicative factor between 0 to 2, and with a perturbation
radius that ranged between 50 and 1,000bps. For a given radius, the slice of the ATAC vector
centered on the locus was multiplied by the selected factor; the remainder of the signal was kept
the same. For each combination of multiplier and perturbation radius size values, we predicted the
corresponding CAP-binding signal for perturbation and calculated the binding differential by

subtracting the unperturbed condition, generating a data tensor of $40 \times 20 \times 8{,}192$ (multipliers ×
window-sizes × predicted profile). Similar to *in silico* DNA sequence mutagenesis, we averaged
the center 400bp region of difference between perturbed prediction and wild-type prediction to
generate a 40×20 matrix that measures the accessibility perturbation effects.
961
***In silico* protein residue mutagenesis**
To investigate how the model leverages protein embedding features in prediction, we performed
systematic mutagenesis directly on the input CAP residues without referring to particular protein
domain or functional data. Each perturbation operates on a 30-amino acid sliding window with
single-residue steps, replacing the window with random amino acids to generate the corresponding
mutant protein sequence. Mutated CAP sequence was feed into the same protein encoder (ESM2)
to generate the corresponding mutant embeddings, which were used in downstream prediction of
mutant binding profile and comparison with wild-type protein.
970
To ensure more robust comparison of protein mutation effects between the mutant and wild-type
CAP, we selected the top 200 Chromnitron-predicted binding peaks for each CAP in HepG2 cells,
filtered for ATAC-seq overlap, and equally extended to the flanking sequences up to 8,192bp to
facilitate prediction. Chromnitron predictions of the corresponding CAP were generated for wild-
type and mutant embeddings across all selected loci. We then extracted maximum scores from the
center 400bp region of each window (recorded as "wildtype_max" and "mutant_max",
respectively). CAP mutation impact scores were calculated as ($\frac{|\text{mutant\_max} - wildtype\_max|}{wildtype\_max}$). For
protein domain interpretation, we retrieved InterPro domain annotations and computed precision-
recall curves using domain boundaries as labels and mutation scores as predictions[69].
Domain enrichment was quantified using AUPR compared to random baselines.
981
Inferring CAP interactions through colocalization analysis
To investigate whether model prediction serves as a potential source to study protein-protein
interaction between CAPs, we followed our previous approach and applied linear non-Gaussian
acyclic model (LiNGAM) causal-discovery algorithm to a selection of high-quality CAPs
prediction across HepG2 ATAC-seq peaks[37,70]. We found that, as expected, the strongest
colocalizations were often between transcription factors that belong to the same family as they
usually share similar conserved DNA-binding domains. To systematically evaluate whether
colocalization calculated from experimental ChIP-seq or predicted CAP-binding data reconciles
with known physical protein-protein interactions, we benchmarked our casual discovery-based
interaction graph with Pearson's correlation-based interaction graph. This analysis was performed
on both experimental and predicted ChIP-seq data using confident interaction annotation from
STRING database (using scores over 700 as positive, and 400 to 700 as negative)[71]. The casual
discovery on predicted data yields the highest AUPR (0.426), followed by causal discovery on
observed data (0.372), while the correlation-based approaches achieved AUPR scores of 0.351 and
0.321 for predicted and experimental ChIP-seq data, respectively (Supplementary Fig. 11). The
lower performance when using ChIP-seq data could be a result from heterogeneous noise

distribution in the experimental data compared to the predicted results, leading to prediction of
only strongest interactions.
1000

Single cell differential ATAC-seq peak calling

To identify differentially accessible chromatin regions between pairs of cell types from single-cell
ATAC-seq data, we employed a computational workflow utilizing SCANPY[72]. For each pairwise
comparison, we first extracted the cells from the two specified populations the from complete
peak-by-cell matrix. Peaks with low signal, defined as those with fewer than 500 total fragment
counts across the selected cells, were removed from consideration. To account for technical
variations in sequencing depth, we normalized the counts for each cell to a total of 100,000,
followed by a log1p transformation to stabilize the variance. Subsequently, we identified highly
variable peaks between the two cell populations by using a dispersion-based method implemented
in SCANPY (scanpy.pp.highly_variable_genes), retaining peaks that exhibited a normalized
dispersion of at least 0.1. The genomic coordinates of these candidate regions were used as
candidate differentially accessible regions and exported for downstream analysis.
1013

Predicting CAP-binding at single locus across T cell states

We used a public single-cell ATAC-seq data of T cell exhaustion collected from human patient
PBMC samples[55]. Four representative T cell stages (from effector to exhausted states) were
prepared to relate CAP binding changes to T-cell states. Pseudo-bulk ATAC-seq data for each
stage was generated by sampling 10M reads from cells in the vicinity of selected center. The
pseudo-bulk ATAC-seq was then processed and normalized as described in the single cell data
processing method section. Differential accessible peaks of the corresponding cell populations
were called as described in the above section. These peak regions served as putative *cis*-regulatory
elements that might play important functions in T cell states. Next, for a given regulatory element
of interest, we predicted CAP-binding profiles at an element-centered 8,192bp window across the
cell states. As single locus is highly susceptible to noise, we selected a set of inference CAPs using
a more stringent cutoff of 0.5 (Pearson's *r*) validation performance for this analysis. Predictions
for every T cell state were cropped to a 300bp window around the CRE center. The binding signal
was normalized by median on the genome dimension to remove global scaling differences.
Trajectories were plotted as ridge plots that overlay the complete track profile per CAP. The mean
CAP-binding signals across trajectory were smoothed with Gaussian kernel and visualized with
line charts tracing binding dynamics across cell state transition trajectory.
1031

Genome-wide identification of regulatory CAPs in T cell exhaustion

The same T cell exhaustion scATAC-seq and annotation in single-locus-analysis were used for
this analysis. We first isolated reference pseudo-bulks for both effector T cells and exhausted T
cells according to the original cell type annotations[55]. To obtain a cleaner causal relationship
between CAPs and target genes, we focused on promoter regions as the putative regulatory
elements. We used GENCODE v45 transcript annotation (GFF3) and the hg38.chrom.sizes file
hosted by IGV to define promoter coordinates and chromosome boundaries[73]. Promoter regions
(±500 bp around the transcription-start site, TSS) were extracted from the GENCODE annotation.

Genes on sex chromosomes or mitochondrial DNA were discarded. We also removed transcripts
not matching a whitelist of supported biotypes (protein-coding, lincRNA, miRNA, snRNA,
snoRNA, rRNA). The maximal and mean ATAC-seq accessibility signal were computed in both
T-cell states for each promoter. Promoters with a cross-condition maximum accessibility ≤ 1 were
removed to exclude inactive loci. We then calculated log2 fold change (log2FC) of accessibility
for all the loci, and partitioned them into three categories: (1) up-regulated promoters, defined as
top 2,000 positive log2FC values, (2) down-regulated promoters defined as top 2,000 negative
log2FC values, and (3) stable promoters, defined as 100 promoters with the smallest absolute
log2FC. We also considered alternative TSS as separate promoters and implemented additional
constraints to merge overlapping TSS loci.

1050

Due to the high compute requirement for this task, Chromnitron prediction was sharded by cell
type and all CAPs, and executed in parallel on a GPU cluster. For this task, we used the full set of
CAPs since genome-wide analysis can better tolerate noisy predictions. After all shards completed,
we merged the predictions by iterating over chromosomes, stacking the individual CAP arrays to
generate genome_length × CAPs matrices. This matrix was rechunked to optimize downstream
biological analyses for characterizing putative regulator CAPs.

1057

Promoter-level average CAP-binding signals were extracted by recording the maximum predicted
signal in the 1kb window centered at TSS, generating a three-dimensional tensor with dimension
(cell types × promoters × CAPs). Downstream analyses used the log-transformed values. A
promoter-specific differential CAP-binding score was calculated by subtracting the predicted
CAP-binding signals of the exhaustion group by that of the effector T cell group. To account for
basal CAP-specific variability, the difference matrix was z-standardized by subtracting the mean
and dividing by the standard deviation of the difference values measured at stable promoters. To
increase signal-to-noise ratio, mid-range values between the 10th and 90th percentiles were set to
zeros separately across loci and CAPs, thereby highlighting the most differential – potentially
functional – signals for informing downstream analysis. Next, we performed k-means clustering
on both the loci and CAP dimensions. Silhouette analysis over k-means clustering using k ranging
from 3 to 40 was used to derive optimal number of clusters. To identify important CAPs and loci
within each cluster, we ranked loci and CAPs by revaluating the average binding against the
opposing cluster assignments iteratively on both dimension until convergence.

1072

Global CAP and CRE dynamics characterization during Purkinje cell differentiation
Single-cell ATAC-seq datasets for Purkinje cell development were obtained from Mannens et al[60].
We adopted the original assignment of Purkinje cell differentiation trajectory encompassing seven
stages: progenitor cells (Prog), two neuroblast stages (NBL1, NBL2), and four Purkinje cell
maturation stages (Purk1 to Purk4). All datasets were processed using the scATAC-seq
preprocessing steps described earlier. Only autosomes were included for this analysis.

1079

To identity genomic regions of interest, we used multiple filtering approaches based on chromatin
accessibility dynamics across developmental stages. Initial candidate regions were generated from

cPeaks representing the repertoire of accessible regions across human cell types[64]. We extended
each peak center by 500 bp flanking region on each side to create 1 kb inference windows to
include the complete peak region. Then, we filtered these loci by classifying them into two
categories based on temporal accessibility patterns: differential and stable regions. Differential loci
were defined as regions exhibiting significant accessibility changes across developmental stages
(log2 fold-change larger than 0.1 compared with average accessibility), representing dynamic
regulatory elements. Stable loci have relatively minimal accessibility change, defined as having
absolute log2FC smaller than 0.1. Similar to the genome-wide regulator screening in T cell
exhaustion, these stable loci served as normalization controls for downstream analysis. We
selected the top 1,000 differential loci and top 100 most stable loci. To ensure non-redundant
genomic coverage, overlapping regions were merged. Regions with insufficient chromatin
accessibility across all conditions were excluded to prevent analysis of inactive or noisy genomic
elements. The final dataset comprised 1,073 high-confidence genomic loci distributed across
autosomes.
1096
This task was computationally intensive and was thus parallelized on a GPU cluster. To reduce
compute, we only keep a set of high-performance CAPs similar to the single locus T cell
exhaustion inference experiment. Individual CAP predictions were concatenated to construct a 3D
data tensor of developmental time × genomic loci × CAPs. For each genomic locus, we calculated
the maximum peak heights within 1 kb windows centered at peak summits. The resulting 3D tensor
was filtered for missing values, infinite numbers, and outliers. Data normalization was performed
using stable loci as internal controls to account for technical variation and global accessibility
changes across developmental stages. We performed z-score normalization using the mean and
standard deviation of differential patterns calculated from stable loci. Outlier detection and
removal were implemented using percentile-based clipping (1st and 99th percentiles) to prevent
extreme values from dominating downstream clustering analyses.
1108
Regulatory pattern discovery was performed using multiple complementary approaches to capture
different aspects of the temporal regulatory landscape. The 3D tensor was reshaped into matrices
emphasizing different analytical perspectives. There are three ways to transform the 3D tensor to
2D matrix for specific analyses, each enabling discovery of distinct regulatory relationships. First,
we can use CAP as features, reshaping the tensor to a matrix with dimension (time × loci) by CAPs
to identify loci whose activities change across time. Second, we can use loci as features to generate
the (time × CAPs) by loci matrix. This approach can represent how CAP binding profiles change
across time. Finally, using time as the feature, the reshaped matrix can tell how a binding event
specific to a CAP and a locus varies across time.
1118
We focused on the first analysis using CAP as features. To observe CRE activity shifts across time,
we performed hierarchical clustering of CREs on the first pseudo-time point. This approach
grouped CREs with similar regulatory dynamics. To further elucidate the relationship between
CRE and time, we applied k-means clustering using a k between 5 and 20 to identify regulatory
locus modules across all time points and CREs. The optimal cluster number was determined using

silhouette score. Clustering was performed on unit-normalized data using Euclidean distance
metrics, with multiple random initializations to ensure stable cluster assignments. For visualization
and interpretation, t-distributed Stochastic Neighbor Embedding (t-SNE) was used to generate 2D
embeddings of high-dimensional regulatory patterns. Each dot on the t-SNE embedding represents
a CRE at a specific time point and was colored by clustering results representing its identity.
1129
1130 To visualize transition of CREs between adjacent pseudo-time points in Purkinje cell development,
1131 we calculated a vector field where the start and end were defined by the t-SNE location of each
1132 locus at first and second time points. This velocity field represents how CREs change their
1133 regulatory identity between consecutive developmental stages. We applied Gaussian kernel
1134 smoothing to the vector field to reduce noise while preserving temporal trends.
1135

Cell culture and primary cell isolation
HEK293T was obtained from the American Type Culture Collection (ATCC). HEK293T cells
were maintained in Dulbecco's Modified Eagle Medium (DMEM; Gibco, 11965092)
supplemented with 10% fetal bovine serum (FBS; Gibco, A5670701) and 1% penicillin–
streptomycin (Pen/Strep; Gibco, 15140122). Human Peripheral blood mononuclear cells (PBMCs)
were isolated from healthy donor leukopaks (Charles River Laboratories Cell Solutions, 103051-
034) using Ficoll-Paque PLUS density gradient centrifugation (Cytiva, 17144002) according to
the manufacturer's protocol. Isolated PBMCs were cryopreserved in 90% human AB serum (MP
Biomedicals, 092938249) plus 10% dimethyl sulfoxide (Sigma-Aldrich, D4540) at a density of
30M cells per ml. For in vitro T cell assays, thawed PBMCs or purified CD8$^+$ T cells (isolated
using EasySep™ Human CD8+ T Cell Isolation Kit, STEMCELL Technologies, 17953) were
cultured in X-VIVO 20 medium (Lonza, BW04-448Q) supplemented with 5% heat-inactivated
human AB serum (MP Biomedicals, 092938249) and 10 ng/ml recombinant human IL-2
(PeproTech, 200-02-50UG). T cell cultures were maintained at 37 °C in a humidified incubator
with 5% CO_2.
1151

sgRNA library preparation
For each selected gene, five sgRNAs were designed to either knock out or repress gene expression
using the CRISPick online tool (Broad Institute). The complete list of sgRNA sequences is
provided in Supplementary File 1. A total of 370 DNA oligonucleotides corresponding to sgRNAs
were synthesized as pooled libraries (500 ng per library) by Twist Bioscience. For gene knockout
experiments, sgRNAs were cloned into the lentiviral vector pXPR_023 vector (Broad Institute,
also available through Addgene #52961), which encodes Cas9, a puromycin resistance gene, and
the sgRNA scaffold. For gene repression (CRISPRi), sgRNAs were cloned into pXPR_066 vector
(obtained via Broad Institute Genetic Perturbation Platform), which expresses KRAB-dCas9 under
the EFS promoter along with a puromycin resistance cassette and sgRNA scaffold.
Oligonucleotides were cloned into the respective backbone vectors using Golden Gate assembly
with BsmBI restriction enzyme (New England Biolabs, R0739S), following the protocol described
by the Feng Zhang lab (Broad Institute). Ligation products were transformed into NEB Stable
Competent *E. coli* (New England Biolabs, C3040H) and plated on LB-agar plates supplemented

with carbenicillin (100 μg/ml). Colonies were scraped from the plates and pooled, and plasmid
DNA was extracted using the ZymoPURE II Midiprep Kit (Zymo Research, D4201) and
quantified by NanoDrop for downstream lentivirus production.
1169
Lentivirus production
Lentivirus was produced in HEK293T cells (ATCC; passage number <20) seeded at 4×10^7 cells
per 150-mm tissue culture dish. Twenty-four hours after seeding, the culture medium was replaced
with fresh pre-warmed DMEM containing 10% FBS. Transfection was performed using
polyethyleneimine (PEI 40K; Polysciences, 24765) as follows: 13.3 μg of the sgRNA vector, 10 μg
of packaging plasmid psPAX2, and 6.7 μg of envelope plasmid pMD2.G were mixed in 400 μl of
Opti-MEM (Thermo Fisher Scientific, 31985070) (Tube A). Separately, 90 μl of PEI (1 mg/ml)
was diluted in 310 μl of Opti-MEM (Tube B). The two solutions were incubated separately for
5 min at room temperature, combined, and incubated for an additional 15 min before being added
dropwise to the HEK293T cells. Seventy-two hours post-transfection, the culture supernatant was
harvested and centrifuged at 3,000g for 15 min to remove cellular debris. Lentiviral particles were
precipitated by adding polyethylene glycol 8000 (PEG8000; Sigma-Aldrich, P2139-500G) to a
final concentration of 8% (w/v) and incubating the mixture overnight at 4 °C. The precipitated
lentiviral particles were pelleted by centrifugation at 1,500g for 30 min and resuspended in 500 μl
of X-VIVO 20 medium (Lonza, BW04-448Q). Aliquots were stored at −80 °C until use.
1185
T cell stimulation assays
Cryopreserved human CD8+ T cells were thawed and resuspended in T cell culture medium (X-
VIVO 20 supplemented with 5% human AB serum and 10 ng ml^{-1} recombinant human IL-2) at a
density of 1×10^6 cells per ml in 12-well plates. Cells were stimulated by the addition of Human
CD3/CD28 T Cell Activation Beads (BioLegend, 422603) at a 1:1 bead-to-cell ratio and incubated
for 48 h at 37 °C with 5% CO_2. Following activation, beads were removed using a magnetic
separation rack according to the manufacturer's instructions. Cells were then washed twice with
fresh T cell medium and split into two different stimulation conditions. For the acute stimulation
group, cells were maintained in T cell medium alone. For the chronic stimulation group with
antibody cocktail, cells were cultured in T cell medium supplemented with ImmunoCult Human
CD3/CD28 T Cell Activator (Stem Cell Technologies, 10971), following the manufacturer's
protocol. For the chronic stimulation group with anti-CD3 antibody (eBioscience, 16-0037-81),
the antibody was coated at 12 well plate with 5 ug/ml concentration. Cells in the chronic condition
were passaged every 48 h, with medium replacement and re-activate for three consecutive
stimulation rounds. At the end point, cells were harvested and divided into two groups for either
surface staining of exhaustion markers or intracellular cytokine analysis by flow cytometry.
1202
Lentiviral transduction and titration in CD8$^+$ T cells
To determine optimal lentiviral transduction conditions for primary human CD8$^+$ T cells, cells
were activated for 48 h using CD3/CD28 beads as described above. After bead removal, cells were
plated in 96-well plates at 2×10^5 cells per well in 200 μl of T cell medium (X-VIVO 20
supplemented with 5% human AB serum and 10 ng/ml IL-2). Serial dilutions of lentivirus

encoding puromycin resistance gene were added to the wells in the presence of 8 μg/ml polybrene
(Millipore, TR-1003-G). Plates were centrifuged at 1,500g for 90 min at 37 °C, the medium was
changed to T cell medium after additional 4h incubation, the cell was continued culture at 37 °C,
5% CO_2. At 48 h post-transduction, cells were diluted to 1×10^5 cells per ml and cultured in the
presence of 1 μg/ml puromycin for 48 h. cell viability was assessed by counting viable cells using
trypan blue. Viral titer and multiplicity of infection (MOI) were calculated based on the fraction
of puromycin-resistant cells relative to total input.
1215
CRISPR screening in primary CD8+ T cells
Human CD8+ T cells were isolated from PBMCs by negative selection and activated with
CD3/CD28 beads for 24 h. After bead removal, cells were transduced with pooled lentiviral
sgRNA libraries at an MOI of 0.3 in the presence of 8 μg/ml polybrene. Cells were spinoculated
at 1000g and 32 °C for 90 min. 6h after spin infection, the cell were resuspended to T cell medium
for overnight culture. 24 hours after spin infection, the cell were treated with puromycin (1.5 μg/ml)
to select the transduced cells. Following selection, cells were washed and resuspended into two
treatment conditions corresponding to acute and chronic stimulation, as described previously. Cells
were maintained under these conditions for 10 days, with medium refreshed and IL-2 re-
supplemented every 2 days. On the final day of the screen, genomic DNA was harvested from each
condition using the QIAamp DNA Blood Maxi Kit (QIAGEN, 51192) following the
manufacturer's instructions. Genomic DNA was quantified using Qubit dsDNA BR Assay Kit
(Thermo Fisher Scientific, Q32850), and sgRNA sequences were PCR-amplified from extracted
DNA using a one-step amplification protocol compatible with next-generation sequencing, as
previously described by Broad Institute Genetic Perturbation Platform ("sgRNA/shRNA/ORF
PCR for Illumina Sequencing" linked through
https://portals.broadinstitute.org/gpp/public/resources/protocols), PCRprimers were provided in
Supplementary File 2. PCR products were purified using AMPure XP beads (Beckman Coulter,
A63880). Libraries were quantified, normalized, and sequenced on an Illumina NovaSeq X
platform using a 25B flow cell, generating paired-end 150 bp reads. Raw sequencing data were
demultiplexed using bcl2fastq (Illumina) and further processed as described below.
1237
CRISPR screening analysis
Sequencing reads from CRISPR screens were first demultiplexed and trimmed to extract sgRNA
sequences. Reads were aligned to the sgRNA reference library using the MAGeCK count function
in the MAGeCK software package with default parameters[74]. Only sgRNAs with perfect matches
to the reference library were retained for downstream analysis. Read count matrices were
generated for all samples and normalized using median normalization. Differential sgRNA
abundance between experimental conditions (Chronic vs. Acute stimulation) was analyzed using
the MAGeCK test module, which applies a negative binomial model to assess gene-level
enrichment or depletion. For each gene, MAGeCK aggregates sgRNA-level statistics to compute
a robust gene-level score and associated p value using a robust ranking aggregation (RRA)
algorithm. Genes were ranked by their adjusted p values (Benjamini–Hochberg FDR correction),

and significance thresholds were defined as indicated in the figure legends. All analyses were
performed using MAGeCK with default settings unless otherwise noted.
1251

RNP electroporation in primary human CD8$^+$ T cells
To perform Cas9 RNP-mediated gene knockout, two synthetic sgRNAs targeting the same gene
were used per condition (sequence available in Supplementary File 2). Each sgRNA was
resuspended to a final concentration of 100 µM in RNase-free water and stored at −80 °C until use.
Cas9 protein (EnGen® SpyCas9 NLS, 20 µM stock, New England Biolabs, M0646T) was used
for complex formation. For each reaction, 60 pmol of Cas9 protein was complexed with 180 pmol
total sgRNA by mixing 3 µl of Cas9 protein, 0.9 µl of each sgRNA, and 0.4 µl of 10× NEBuffer
3.1 in a total volume of 5.2 µl. The mixture was incubated at room temperature for 15 min to allow
RNP formation. Primary CD8$^+$ T cells were activated for 48 h with CD3/CD28 beads as described
above, then washed once with 1× PBS and resuspended in Lonza P3 electroporation buffer at a
density of 1×10^6 cells per 20 µl. The entire 5.2 µl RNP mixture was added to 20 µl of cell
suspension, mixed gently, and transferred into a well of a 16-well Nucleocuvette™ strip (Lonza).
Air bubbles were removed by tapping the strip on the benchtop to prevent disruption of the electric
pulse. Electroporation was performed using the Lonza 4D-Nucleofector system with the program
EH115, optimized for primary human T cells. Immediately after electroporation, 80 µl of pre-
warmed T cell medium (X-VIVO 20 supplemented with 5% human AB serum and 10 ng ml^{-1} IL-
2) was added to each well. The cells were incubated in the cuvette strip at 37 °C for 10 min to
enhance recovery before being gently transferred to pre-warmed 24-well plates.
1270

Flow cytometry staining and analysis
For intracellular cytokine staining, T cells were stimulated for 3 h at 37 °C with Cell Activation
Cocktail containing Brefeldin A (BioLegend, 423303). Cells were washed with Cell Staining
Buffer (BioLegend, 420201), fixed and permeabilized with Fixation/Permeabilization solution
(BD Biosciences, AB_2869008) for 20 min at 4 °C, and washed twice with 1× Perm/Wash buffer.
After Fc receptor blocking (BioLegend, 422302), cells were stained for 2 h at 4 °C in the dark with
fluorophore-conjugated antibodies in staining buffer (BioLegend, 420201) along with NIR
Live/Dead dye (1:10000 dilution, ThermoFisher, L34975). Antibodies included APC anti-human
CD8 Antibody (BioLegend, 344722), Brilliant Violet 785™ anti-human IFN-γ Antibody
(BioLegend, 502541), Brilliant Violet 605™ anti-human TNF-α Antibody (BioLegend, 502935),
and Brilliant Violet 421™ anti-human IL-2 Antibody (BioLegend, 500327).
1282

For surface marker staining, cells were blocked with FC receptor blocking reagent (BioLegend,
422302), then stained with antibodies and NIR Live/Dead dye (1:10000 dilution, ThermoFisher,
L34975) at 4 °C for 1 h. Antibodies included CD8 Antibody (BioLegend, 344722), Brilliant Violet
785™ anti-human TIGIT (VSTM3) Antibody (BioLegend, 372735), Brilliant Violet 605™ anti-
human CD366 (Tim-3) Antibody (BioLegend, 345017), Brilliant Violet 510™ anti-human CD223
(LAG-3) Antibody (BioLegend, 369317), and Brilliant Violet 421™ anti-human CD279 (PD-1)
Antibody (BioLegend, 329920). Flow cytometry was performed on a CytoFLEX LE (Beckman
Coulter, six-laser configuration). Gating strategy included sequential gating on FSC-A/SSC-A

(lymphocytes), FSC-A/FSC-H (single cells), and exclusion of dead cells using viability dye.
Fluorescence compensation was calculated using single-stained compensation beads (BioLegend).
Data were initially analyzed with FlowJo v10.8, with gates being defined using isotype controls,
followed by custom reproducing of the analysis for visualization purpose only.
1295

Surveyor assay for genome editing efficiency
To evaluate genome editing efficiency following RNP electroporation, genomic DNA was
extracted from T cells 72 h post-nucleofection using the Monarch® Genomic DNA Purification
Kit (New England Biolabs, T3010S), following the manufacturer's instructions. Genomic regions
flanking the Cas9 cleavage site were PCR-amplified using 2× Phanta Max Master Mix (Vazyme,
P525-02) in a 50 μl reaction volume (PCR primers provided in Supplementary File 2). PCR
products were verified by agarose gel electrophoresis and purified using the Zymo DNA Clean &
Concentrator Kit (Zymo Research, D4014), with final elution in 30 μl of nuclease-free water. To
generate heteroduplex DNA for mismatch detection, 200 ng of purified PCR product was
combined with 2 μl of 10× NEBuffer 2 (NEB) and nuclease-free water to a total volume of 19 μl.
The mixture was denatured and reannealed in a thermocycler using the following program: 95 °C
for 5 min; ramp at −2 °C/s to 85 °C; then ramp at −0.1 °C/s to 25 °C; and held at 4 °C. After
annealing, 1 μl of T7 Endonuclease I (NEB, M0302) was added to each reaction and incubated at
37 °C for 60 min. Reactions were terminated by adding 1.5 μl of 0.25 M EDTA and analyzed by
electrophoresis on a 1% agarose gel to visualize cleavage products.
1311

mRNA knockdown analysis by quantitative PCR
To assess transcript-level gene knockdown following RNP electroporation, edited T cells were
collected at the end point for RNA extraction. Total RNA was isolated using a RNeasy Mini Kit
(Qiagen, 74104), and RNA concentration was quantified by nanodrop. Reverse transcription was
performed using the HiScript® III 1st Strand cDNA Synthesis Kit (+gDNA wiper) (Vazyme,
R312-01) according to the manufacturer's instructions. Gene-specific primers targeting the
transcript of interest were used during amplification (Supplementary File 2). Quantitative PCR
was performed using PowerUp™ SYBR® Green Master Mix (Thermo Fisher Scientific, A25742)
on a CFX96 Real-Time PCR Detection System (Bio-Rad). ACTB (β-actin) was used as the internal
control gene for normalization. Each sample was run in triplicates, and specificity of amplification
was confirmed by melt-curve analysis. Relative gene expression was quantified using the ΔΔCt
method, and results were reported as fold change compared to the control condition.
1324

RNA-seq analyses
Total RNA was extracted on the final day from ZNF766 knockout and NTC control groups using
the RNeasy Mini Kit (Qiagen, 74004). Two biological replicates were collected per condition.
RNA-seq libraries were generated with the NEBNext Ultra II Directional RNA Library Prep Kit
(NEB, E7765L) following the manufacturer's poly(A) mRNA workflow, including enrichment
with the NEBNext Poly(A) mRNA Magnetic Isolation Module (NEB, E7490L). Raw RNA-seq
reads were mapped to the human reference genome (GRCh38/hg38) using STAR (v2.7.2a) with
default settings. Strand-specific read counts were aggregated into a gene-level count matrix for

subsequent analyses. Transcriptomic variation across samples was evaluated by principal
component analysis (PCA) based on the 500 most variable genes. Differential expression analysis
was conducted using DESeq2 (v1.40.2), and genes with an absolute log2 fold change greater than
0.5 and an adjusted P-value below 0.05 were defined as significantly differentially expressed.
Functional enrichment analyses were performed using clusterProfiler (v4.10.0), together with
org.Hs.eg.db (v3.18.0), enrichplot (v1.22.0), and DOSE (v3.28.2). Significant genes were
converted to Entrez Gene identifiers and assessed for enrichment in Gene Ontology biological
processes and KEGG pathways. Enrichment results were visualized as dot plots (GO) and bar plots
(KEGG).
1342

Data Availability

All ATAC-seq, ChIP-seq used in this study are public data collected from the ENCODE data portal (https://www.encodeproject.org). T cell scATAC-seq data is obtained from Gene Expression Omnibus (GEO) under accession GSE129785. Purkinje neuron developmental scATAC-seq data is obtained from https://github.com/linnarsson-lab/fetal_brain_multiomics. CRISPR screen data is available on GEO under accession number GSE305611. RNA-seq data on ZNF766 KO is available on GEO under accession number GSE305610.

Code Availability

The code for Chromnitron is available at https://github.com/tanjimin/Chromnitron.

Competing Interests

J.T., X.F., X.L., B.X., and A.T. are inventors on provisional patents covering the models and tools reported herein. A.T. is a scientific advisor to Intelligencia AI. J.T. is a consultant for Flagship Pioneering. J.D.B is a Founder and Director of CDI Labs, Inc., a Founder of and consultant to Opentrons LabWorks/Neochromosome, Inc, and serves or served on the Scientific Advisory Board of CZ Biohub New York, LLC, Logomix, Inc., Rome Therapeutics, Inc., SeaHub, Seattle, WA, Tessera Therapeutics, Inc., and the Wyss Institute. All these services are unrelated to this work.

Acknowledgement

This work is supported by NIH grants DP5OD033430 and R03DE033371 (to B.X.), R01CA252239 and P01CA229086 (to A.T.). X.F. and R.R. are supported by R35CA253126 and P30CA013696. J.T. was supported by the Judith & Stewart Colton Center for Autoimmunity at NYU Langone and Arthritis National Research Foundation Grant Award (No. 1471427). X.L. is a Fellow of The Jane Coffin Childs Fund for Medical Research. B.X. is a Junior Fellow of the Harvard Society of Fellows. We thank the High Performance Computing Core Facility at NYU School of Medicine and the Broad Institute for the support of computation. We thank the Broad Genetic Perturbation Platform and Broad Clinical Labs for the support of genetic and genomic experiments. We also acknowledge Milk Bar, where Xi and Jimin enjoyed coffee and met weekly for over two years to work on Chromnitron. We thank Huiyuan Zhang, Chenzhen Zhang, Hanqing Liu, Lloyd Bod, Jason D. Buenrostro, Romanos S. Pistofidis, Liz Gaskell, Wenke Liu, Jose U. Scher, and the members of the Xia lab and Tsirigos lab for their valuable suggestions and discussion. Lastly, we are grateful to the ENCODE consortium for generating, organizing, and sharing standardized genomics data that enabled this study.

Author Contribution

B.X. and J.T. conceived the project. J.T., B.X. and A.T. designed the computational and experimental components and interpreted the results with inputs from X.F. and X.L. J.T., F.X. and S.M. designed, implemented and optimized the neural network. J.T. built the preprocessing platform with input from X.F. X.F. performed single cell data preprocessing and protein feature analysis. J.T. trained the model, performed inference and downstream computational analysis with input from X.F, B.X. and A.T. J.B. performed RNA-seq analysis. X.L. performed the experimental

References

References

1. Arendt, D. *et al.* The origin and evolution of cell types. *Nat. Rev. Genet.* **17**, 744–757 (2016).

2. Xia, B. & Yanai, I. A periodic table of cell types. *Dev. Camb. Engl.* **146**, dev169854 (2019).

3. Gurdon, J. B. Cell Fate Determination by Transcription Factors. *Curr. Top. Dev. Biol.* **116**,

445–454 (2016).

4. Chen, X. *et al.* Integration of external signaling pathways with the core transcriptional

network in embryonic stem cells. *Cell* **133**, 1106–1117 (2008).

5. Boyer, L. A. *et al.* Core transcriptional regulatory circuitry in human embryonic stem cells.

Cell **122**, 947–956 (2005).

6. Lander, E. S. *et al.* Initial sequencing and analysis of the human genome. *Nature* **409**, 860–

921 (2001).

7. Lambert, S. A. *et al.* The Human Transcription Factors. *Cell* **172**, 650–665 (2018).

8. Vaquerizas, J. M., Kummerfeld, S. K., Teichmann, S. A. & Luscombe, N. M. A census of

human transcription factors: function, expression and evolution. *Nat. Rev. Genet.* **10**, 252–263

(2009).

9. Gerstein, M. B. *et al.* Architecture of the human regulatory network derived from ENCODE

data. *Nature* **489**, 91–100 (2012).

10.   Li, Y. E. *et al.* A comparative atlas of single-cell chromatin accessibility in the human

brain. *Science* **382**, eadf7044 (2023).

11.   Weirauch, M. T. *et al.* Determination and inference of eukaryotic transcription factor

sequence specificity. *Cell* **158**, 1431–1443 (2014).

12.   Jolma, A. *et al.* DNA-binding specificities of human transcription factors. *Cell* **152**, 327–

339 (2013).

13.   Vierstra, J. *et al.* Global reference mapping of human transcription factor footprints.

Nature **583**, 729–736 (2020).

14.   Murre, C. Helix-loop-helix proteins and the advent of cellular diversity: 30 years of

discovery. *Genes Dev.* **33**, 6–25 (2019).

15.   Doughty, B. R. *et al.* Single-molecule states link transcription factor binding to gene

expression. *Nature* **636**, 745–754 (2024).

16.   Durdu, S. *et al.* Chromatin-dependent motif syntax defines differentiation trajectories.

Mol. Cell **85**, 2900-2918.e16 (2025).

17.   Kim, S. & Wysocka, J. Deciphering the multi-scale, quantitative cis-regulatory code.

Mol. Cell **83**, 373–392 (2023).

18.   Kim, S. *et al.* DNA-guided transcription factor cooperativity shapes face and limb

mesenchyme. *Cell* **187**, 692-711.e26 (2024).

19.   Statello, L., Guo, C.-J., Chen, L.-L. & Huarte, M. Gene regulation by long non-coding

RNAs and its biological functions. *Nat. Rev. Mol. Cell Biol.* **22**, 96–118 (2021).

20.   Smith, E. & Shilatifard, A. The chromatin signaling pathway: diverse mechanisms of

recruitment of histone-modifying enzymes and varied biological outcomes. *Mol. Cell* **40**, 689–

701 (2010).

21.   Yosef, N. *et al.* Dynamic regulatory network controlling TH17 cell differentiation.

Nature **496**, 461–468 (2013).

22.   Johnson, D. S., Mortazavi, A., Myers, R. M. & Wold, B. Genome-wide mapping of in

vivo protein-DNA interactions. *Science* **316**, 1497–1502 (2007).

23.   Tsankov, A. M. *et al.* Transcription factor binding dynamics during human ES cell

differentiation. *Nature* **518**, 344–349 (2015).

24.   Park, P. J. ChIP-seq: advantages and challenges of a maturing technology. *Nat. Rev.*

Genet. **10**, 669–680 (2009).

25.   Skene, P. J. & Henikoff, S. An efficient targeted nuclease strategy for high-resolution

mapping of DNA binding sites. *eLife* **6**, e21856 (2017).

26.   Partridge, E. C. *et al.* Occupancy maps of 208 chromatin-associated proteins in one

human cell type. *Nature* **583**, 720–728 (2020).

27.   Hammal, F., de Langen, P., Bergon, A., Lopez, F. & Ballester, B. ReMap 2022: a

database of Human, Mouse, Drosophila and Arabidopsis regulatory regions from an

integrative analysis of DNA-binding sequencing experiments. *Nucleic Acids Res.* **50**, D316–

D325 (2022).

28.   Moyers, B. A. *et al.* Characterization of human transcription factor function and patterns

of gene regulation in HepG2 cells. *Genome Res.* **33**, 1879–1892 (2023).

29.   Wang, J. *et al.* Sequence features and chromatin structure around the genomic regions

bound by 119 human transcription factors. *Genome Res.* **22**, 1798–1812 (2012).

30.   Avsec, Ž. *et al.* Base-resolution models of transcription-factor binding reveal soft motif

syntax. *Nat. Genet.* **53**, 354–366 (2021).

31.   Cazares, T. A. *et al.* maxATAC: Genome-scale transcription-factor binding prediction

from ATAC-seq with deep neural networks. *PLOS Comput. Biol.* **19**, e1010863 (2023).

32.   Linder, J., Srivastava, D., Yuan, H., Agarwal, V. & Kelley, D. R. Predicting RNA-seq

coverage from DNA sequence as a unifying model of gene regulation. *Nat. Genet.* **57**, 949–

961 (2025).

33.   Avsec, Ž. *et al.* AlphaGenome: advancing regulatory variant effect prediction with a

unified DNA sequence model.

34.   Lin, Z. *et al.* Evolutionary-scale prediction of atomic-level protein structure with a

language model.

35.   Abramson, J. *et al.* Accurate structure prediction of biomolecular interactions with

AlphaFold 3. *Nature* **630**, 493–500 (2024).

36.   Nguyen, E. *et al.* Sequence modeling and design from molecular to genome scale with

Evo. *Science* **386**, eado9336 (2024).

37.   Fu, X. *et al.* A foundation model of transcription across human cell types. *Nature* (2025)

doi:10.1038/s41586-024-08391-z.

38.   Hayes, T. *et al.* Simulating 500 million years of evolution with a language model.

Preprint at https://doi.org/10.1101/2024.07.01.600583 (2024).

39.   Theodoris, C. V. *et al.* Transfer learning enables predictions in network biology. *Nature*

618, 616–624 (2023).

40.   Hu, E. J. *et al.* LoRA: Low-Rank Adaptation of Large Language Models. Preprint at

https://doi.org/10.48550/arXiv.2106.09685 (2021).

41.   Bravo González-Blas, C. *et al.* SCENIC+: single-cell multiomic inference of enhancers

and gene regulatory networks. *Nat. Methods* **20**, 1355–1367 (2023).

42.   Hu, Y. *et al.* Multiscale footprints reveal the organization of cis-regulatory elements.

Nature **638**, 779–786 (2025).

43.   Pampari, A. *et al.* ChromBPNet: bias factorized, base-resolution deep learning models of

chromatin accessibility reveal cis-regulatory sequence syntax, transcription factor footprints

and regulatory variants. Preprint at https://doi.org/10.1101/2024.12.25.630221 (2024).

44.   Avsec, Ž. *et al.* Effective gene expression prediction from sequence by integrating long-

range interactions. *Nat. Methods* **18**, 1196–1203 (2021).

45.   Stoltenburg, R., Reinemann, C. & Strehlitz, B. SELEX—A (r)evolutionary method to

generate high-affinity nucleic acid ligands. *Biomol. Eng.* **24**, 381–403 (2007).

46.   Ren, B. *et al.* Genome-Wide Location and Function of DNA Binding Proteins. *Science*

290, 2306–2309 (2000).

47.   Khan, O. *et al.* TOX transcriptionally and epigenetically programs CD8+ T cell

exhaustion. *Nature* **571**, 211–218 (2019).

48.   Seo, H. *et al.* TOX and TOX2 transcription factors cooperate with NR4A transcription

factors to impose CD8$^+$ T cell exhaustion. *Proc. Natl. Acad. Sci.* **116**, 12410–

12415 (2019).

49.   Yao, C. *et al.* Single-cell RNA-seq reveals TOX as a key regulator of CD8+ T cell

persistence in chronic infection. *Nat. Immunol.* **20**, 890–901 (2019).

50.   Mann, T. H. & Kaech, S. M. Tick-TOX, it's time for T cell exhaustion. *Nat. Immunol.*

20, 1092–1094 (2019).

51.   Scott, A. C. *et al.* TOX is a critical regulator of tumour-specific T cell differentiation.

Nature **571**, 270–274 (2019).

52.   Alfei, F. *et al.* TOX reinforces the phenotype and longevity of exhausted T cells in

chronic viral infection. *Nature* **571**, 265–269 (2019).

53.   Hashimoto, M. *et al.* CD8 T Cell Exhaustion in Chronic Infection and Cancer:

Opportunities for Interventions. *Annu. Rev. Med.* **69**, 301–318 (2018).

54.   Baessler, A. & Vignali, D. A. A. T Cell Exhaustion. *Annu. Rev. Immunol.* **42**, 179–206

(2024).

55.   Satpathy, A. T. *et al.* Massively parallel single-cell chromatin landscapes of human

immune cell development and intratumoral T cell exhaustion. *Nat. Biotechnol.* **37**, 925–936

(2019).

56.   Jenkins, E., Whitehead, T., Fellermeyer, M., Davis, S. J. & Sharma, S. The current state

and future of T-cell exhaustion research. *Oxf. Open Immunol.* **4**, iqad006 (2023).

57.   Tay, T. *et al.* Degradation of IKZF1 prevents epigenetic progression of T cell exhaustion

in an antigen-specific assay. *Cell Rep. Med.* **5**, 101804 (2024).

58.   Belk, J. A. *et al.* Genome-wide CRISPR screens of T cell exhaustion identify chromatin

remodeling factors that limit T cell persistence. *Cancer Cell* **40**, 768-786.e7 (2022).

59.   Wu, J. E. *et al.* In vitro modeling of CD8+ T cell exhaustion enables CRISPR screening

to reveal a role for BHLHE40. *Sci. Immunol.* **8**, eade3369 (2023).

60.   Mannens, C. C. A. *et al.* Chromatin accessibility during human first-trimester

neurodevelopment. *Nature* (2024) doi:10.1038/s41586-024-07234-1.

61.   Chen, K. M., Wong, A. K., Troyanskaya, O. G. & Zhou, J. A sequence-based global map

of regulatory activity for deciphering human genetics. *Nat. Genet.* **54**, 940–949 (2022).

62.   Hendrycks, D. & Gimpel, K. Gaussian Error Linear Units (GELUs). Preprint at

https://doi.org/10.48550/arXiv.1606.08415 (2023).

63.   Vaswani, A. *et al.* Attention Is All You Need. *ArXiv170603762 Cs* (2017).

64.   Meng, Q. *et al.* A generic reference defined by consensus peaks for scATAC-seq data

analysis. Preprint at https://doi.org/10.1101/2023.05.30.542889 (2023).

65.   Amemiya, H. M., Kundaje, A. & Boyle, A. P. The ENCODE Blacklist: Identification of

Problematic Regions of the Genome. *Sci. Rep.* **9**, 9354 (2019).

66.   Kingma, D. P. & Ba, J. Adam: A Method for Stochastic Optimization. Preprint at

https://doi.org/10.48550/arXiv.1412.6980 (2017).

67.   Hu, E. J. *et al.* LoRA: Low-Rank Adaptation of Large Language Models. Preprint at

https://doi.org/10.48550/arXiv.2106.09685 (2021).

68.   Shrikumar, A. *et al.* Technical Note on Transcription Factor Motif Discovery from

Importance Scores (TF-MoDISco) version 0.5.6.5. Preprint at

https://doi.org/10.48550/arXiv.1811.00416 (2020).

69.   Blum, M. *et al.* InterPro: the protein sequence classification resource in 2025. *Nucleic*

Acids Res. **53**, D444–D456 (2025).

70.   Ikeuchi, T., Ide, M., Zeng, Y., Maeda, T. N. & Shimizu, S. Python package for causal

discovery based on LiNGAM.

71.   Szklarczyk, D. *et al.* The STRING database in 2023: protein–protein association

networks and functional enrichment analyses for any sequenced genome of interest. *Nucleic*

Acids Res. **51**, D638–D646 (2023).

72.   Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression

data analysis. *Genome Biol.* **19**, 15 (2018).

73.   Robinson, J. T. *et al.* Integrative genomics viewer. *Nat. Biotechnol.* **29**, 24–26 (2011).

74.   Li, W. *et al.* MAGeCK enables robust identification of essential genes from genome-

scale CRISPR/Cas9 knockout screens. *Genome Biol.* **15**, 554 (2014).

1542

Supplementary Figures

 Supplementary Fig. 1: Chromnitron model architecture.

Left: overall model structure, convolution and transformer modules, and layer configurations. The feature size of embeddings (including batch as the first dimension) are denoted in brackets.

1549

1550

Supplementary Fig. 2: Training data statistics and the Chrom2Vec platform.

a. CAP-binding ChIP-seq datasets and the corresponding percentages across five human cell lines.

b. Heatmap of pairwise overlaps of CAP-binding datasets across cell lines. Diagonal entries are removed.

c. CAP categorization by their Chromnitron-prediction performance.

d. Distribution of CAPs across prediction-performance categories in each cell type.

e. Workflow of Chrom2Vec platform for processing training datasets. Detailed workflow was also described in Methods sections. Slurm workload controller was used to initialize schedulers

1559 tracking tools including Airflow and Neo4j webservers. Databases used for information relay

1560 (Redis in-memory DB, PostgreSQL DB, Neo4j graph DB) were initialized separately. Celery

1561 workers perform the containerized (Singularity, DockerHub) bioinformatic pipeline, including

1562 Download FastQ, FastQC quality control, Trimming, Alignment, Merge BAM, Subsample,

1563 Coverage, Normalization, and Compression steps. Each action includes a completion callback that

1564 logs through Airflow to the Neo4j database.

1565

Supplementary Fig. 3: OuterMix algorithm increases multimodal training efficiency via factorized batch sampling.

a-c. Sampling and computation with random sampling strategy. A randomly sampled batch has $N{\times}M{\times}K$ combinations of genome sequence (N), chromatin feature (M), and CAP (K), selected from the full three-dimensional data manifold (**a**). This random process generates large data matrices for encoding (**b**). Thus, each encoder for the corresponding modality would process every combination independently, yielding a computational complexity of O($N{\times}M{\times}K$) per encoder (**c**). **d-f.** Sampling and computation with OuterMix strategy. OuterMix first samples *N, M, K* training samples from each dimension, effectively sampling a dense data cube from the full three-dimensional data manifold (**d**). Each sampling was encoded, with the encoded embeddings being recombined via a Cartesian product to form $N{\times}M{\times}K$ training samples (**e**). These encoded features are reused in the final Cartesian product and avoids redundant computation, reducing encoder computation complexity down to O(N) + O(M) + O(K).

1579

1580

1581
1582 **Supplementary Fig. 4: Chromnitron model performance evaluation and ablation studies.**
1583 **a.** Model performance across pre-training and fine-tuning epochs, grouped by CAP categories.
1584 Line and the shade represent scores of mean ± s.d. for each CAP category.
1585 **b.** Scatter plot of model performance (Pearson's correlation between prediction and experimental
1586 data of individual CAPs) between fine-tuned model and pre-trained model.

c-e. Evaluation of Chromnitron predictions against experimental ChIP-seq replicates for CTCF (top) and MAX (bottom). Each evaluation was performed by comparing prediction performance with that of replicating experimental measurements. These evaluations include: (1) calculating Pearson's correlation to the reference experimental data across different percentiles of genomic bins (**c**), (2) scatter plot visualization of predicted or experimental replicate binding signals (x-axis) to the reference experimental data (y-axis, **d**), and (3) direct visualization of CAP-binding tracks of prediction alongside two other experimental replicates (Rep2, Rep3), all aligning to the reference experimental data (Ref, **e**).

f. Establishment of CAP performance groups. Two cutoffs (0.4 and 0.6) based on validation performance were selected on the regression line between validation performance and cell-type-specific performance. Regression coefficient and offset is shown. Areas split by the horizontal and vertical line segments represent regions for high, medium, and low performance CAP groups (annotated on the right of y axis). The grouping rules were later generalized to all CAPs.

g. Ablation study to test *de novo* prediction of CAP-binding in unseen cell types. Ablation-study model was trained with ChIP-seq data from only HepG2 and K562 cells. Top: scatter plot of prediction performance (Pearson's r) comparing ablation-study model trained using only two cell types (HepG2 and K562) versus the full Chromnitron model trained on all five cell types (HepG2, K562, GM12878, IMR-90, HCT116). Bottom: performance of ablation-study model versus training data quality. Blue and grey points represent CAPs in *de novo* cell types and in training cell types, respectively.

h. Ablation study to test *de novo* CAP prediction. Ablation-study model was trained with ChIP-seq corresponding to 385 CAPs. Top: scatter plot of ablation-study model performance for model trained with 385 CAPs versus the full Chromnitron model trained on all 767 CAPs. Bottom: performance of ablation-study model versus data quality. Blue and grey points represent *de novo* CAPs and training CAPs, respectively.

i. Training curves comparisons across three model-training strategies: fine-tuning a fully pre-trained model (dark blue), fine-tuning after 19 epochs pre-training (light blue), and training from scratch (green). The comparison includes overall validation loss (across all cell types) and Pearson's r relating to reference data in each cell type across training epochs.

a

Pred
ChIP
CAP Name: Pearson's r

SAP130: 0.91
RAD21: 0.88
HOXA3: 0.85
ZNF407: 0.82
SOX13: 0.81
TFAP4: 0.81
ZNF48: 0.78
ELF3: 0.78
SMAD4: 0.77
FOXP4: 0.76
HNRNPLL: 0.73
JUND: 0.69

b

PHF5A: 0.68
ZNF547: 0.67
ZNF83: 0.67
RCOR2: 0.66
ZNF784: 0.64
DNMT1: 0.62
SSRP1: 0.61
ZNF124: 0.55
TMF1: 0.52
ZNF12: 0.50
NACC2: 0.48
KLF12: 0.41

c

HNRNPK: 0.57
ZC3H13: 0.46
CHCHD3: 0.45
PCBP2: 0.44
HNRNPH1: 0.44
WIZ: 0.40
FUBP1: 0.38
SNAI1: 0.28
PBX2: 0.24
ZNF30: 0.19
CENPX: 0.16
ZNF850: 0.14

chr10: 89,266,500-89,366,500

1617

1625

Supplementary Fig. 6: Peak detection-based performance evaluation for low-correlation CAPs.

Performance metrics for four randomly selected low-correlation (Pearson's $r < 0.5$) CAPs, PCBP2, FUBP1, PBX2, ZNF850. Pearson's correlation was calculated by comparing to reference experimental ChIP-seq data. For each CAP, four statistics were calculated, including area under the receiver operating characteristic curve (AUROC) and area under the precision-recall curve (AUPR) using both target-centric and prediction-centric labelling. Metrics were calculated using bin sizes across 50, 250, 500, and 1,000 bp (color intensity denotes bin size). Red curves indicate selection of bins from the top percentile only, while blue curves indicate selection of bins from both the top and bottom percentiles of bins.

1638

Supplementary Fig. 7: Performance comparison of CAP binding predictions.
Relating to Fig. 2h, 10 more randomly selected CAPs were presented at the same 500kb genomic window in the leave-out chromosome 10. Each CAP includes predicted binding tracks from Chromnitron, motif accessibility, maxATAC and Enformer models, all referring to the experimental ChIP-seq data in leave-out chromosome 10. Min-max scaling of the binding signal was applied to each track separately.

1645

Supplementary Fig. 8: Performance metrics comparison between Chromnitron and three other models, using MAX as an example.

All performance metrics were performed against reference experimental ChIP-seq data of MAX. In total, eight statistics were calculated, including Pearson's correlation (r), Spearman's correlation (ρ), the coefficient of determination (R squared), F1 score, AUROC and AUPR

curves using both target-centric and prediction-centric labelling. Each of these methods was
calculated using bins sizes of 50, 250, 500, and 1,000 bp (color intensity denotes bin size). Each
plot shows a performance metric (y-axis) dynamic as the different percentages of genomic bins
included in the analysis (x-axis, log-scaled). Red lines indicate selection of bins from the top
percentiles only, while blue lines indicate selection of bins from both the top and bottom
percentiles of bins.
1657

1658
Supplementary Fig. 9: Global statistics of Chromnitron benchmark.

a. Benchmarking prediction performance in the leave-out genomic region (chromosome 10) of
Chromnitron. The benchmarking was performed in HepG2 cells. Each dot represents a CAP
prediction performance by Chromnitron (y-axis) versus the comparison model (x-axis). Area under
the precision–recall curve (AUPR) and area under the receiver operating characteristic curve
(AUROC) were calculated by using ChIP-seq (top) or prediction (bottom) as the reference label
of binarized binding status.
b. Performance benchmarking by predicting cell type-specific CAP-binding. The benchmarking
was performed by comparing model-predicted binding difference between HepG2 and K562 cells
(HepG2 – K562) to the experimental ChIP-seq difference (HepG2 – K562). Pearson's r and
Spearman's ρ were included as evaluation metrics in addition to the AUPR and AUROC as in (**a**).
1670

1671
Supplementary Fig. 10: *In silico* **DNA sequence mutagenesis identifies CAP impact motifs**
and enables variant effect prediction.

a-c. Chromnitron prediction-based *in silico* DNA sequence mutagenesis identifies CAP-specific
impactful DNA bases (i.e. impact motifs) that resembles both known (**a**) and novel (**b**) motifs, as
well as indirect binding motifs that resemble their interaction partner CAPs which binds DNA(**c**).
Impact scores were presented in the form of sequence logos which were normalized per base to
median value. Each impact motif was predicted at a specific locus as denoted in panel title.
d. Chromnitron predicts gain-of-binding effect at *TERT* promoter G250A variant (hg38
chr5:1295135 G>A). Left: CAPs ranked by mutation impact score with vertical dashed colored
lines marking ETS-family TFs. Right: top ETS-family candidate ETV4 binding profiles on wild
type (WT) and mutant (Mut) sequences and their differential binding (Diff: Mut – WT).
e. Chromnitron predicts loss-of-binding effect at a CTCF-binding site (hg38 chr10:25081160 C>T).
Left: CAPs ranked by mutation impact score with red dashed line indicating CTCF. Right: CTCF
binding profiles across WT and Mut sequences and their differential binding.

1691
1692
Supplementary Fig. 11: Chromnitron-based *in silico* CAP residue mutagenesis captures
protein functional domains and protein–protein interactions.
a. Visualization of LiNGAM causal-discovery algorithm using predicted CAP binding profiles
(Methods). Edges represent high interaction potential between adjacent CAPs. Nodes were colored
by CAP community clusters.
b. *In silico* CAP residue mutagenesis by Chromnitron captures protein residues from annotated
functional domains. Each dot in the scatter plot represents a CAP's AUPR scores against protein
domain annotations using *in silico* mutagenesis-identified residues (y-axis) and compared with
that using randomly selected residues (x-axis).
c. Performance comparison between different methods for discovering protein-protein interactions.
We calculated both causal-inference-based interaction graph and Pearson's correlation-based
interaction graph for both Chromnitron prediction and ChIP-seq and compared them against
experimental STRING database-annotated interactions to calculate AUPR (Methods).
1706

1707
Supplementary Fig. 12: *In vitro* chronic stimulation of human primary CD8[+] T cells to model
T cell exhaustion trajectory.
a. Schematic of the *in vitro* treatment of human CD8[+] T cells across acute stimulation (top), chronic
stimulation by anti-CD3 antibodies (middle), and chronic stimulation by anti-CD3/CD28
antibodies (bottom).
b. Flow cytometry quantification of cell surface markers of T cell exhaustion (PD1, LAG3, TIM3,
and TIGIT) and intracellular cytokine production (IFN-γ, TNF-α, IL-2) in different stimulation
groups at day 12. Bar plots show mean ± s.d. for n = 3 independent replicates. Statistical
comparisons performed using two-sided unpaired Student's t-tests with significance denoted as *,
P < 0.05; **, P < 0.01; ***, P < 0.001.
1718

1719

Supplementary Fig. 13: Validation of CRISPR-mediated perturbation and transcript-level
perturbation of *TOX* and *ZNF865*.
a. RT–qPCR quantification of *TOX* and *ZNF865* mRNA expression level changes across chronic
stimulation of human CD8$^+$ T cells from day 4 to day 12, indicating increased expression along
the trajectory. Relative expression levels were normalized to day 2 using the ΔΔCt method. Each
dot represents an independent biological replicate (n = 4); error bars indicate mean ± SEM.
b. Surveyor assay confirmed efficient CRISPR knockout gRNA targeting *TOX* and *ZNF865*. Each
gene includes two gRNAs for knock-out perturbation. Genomic DNA was amplified around the
gRNA target sites and the fragments were digested with T7 Endonuclease I.
c. RT–qPCR quantification of *ZNF865* and *TOX* mRNA level changes across chronic stimulation
days with individual gene knockouts (*ZNF865* KO and *TOX* KO). Relative expression levels were
normalized to day 2. Each dot represents an independent biological replicate (n = 4); error bars
indicate mean ± SEM.
1733

1734
Supplementary Fig. 14: Validation of CRISPR-mediated *ZNF766* knock-out and
transcriptomic profiling of perturbation effects on chronic-stimulated T cells.

a. RT–qPCR quantification of *ZNF766* mRNA level changes across chronic stimulation of CD8[+]
T cells from day 4 to day 12, indicating increased expression along the trajectory. Relative
expression levels were normalized to day 2 using the ΔΔCt method. Each dot represents an
independent biological replicate (n = 4); error bars indicate mean ± SEM.

b. Surveyor assay confirmed efficient CRISPR knockout gRNA targeting *ZNF766*.

c. RT–qPCR quantification of *ZNF766* mRNA expression levels after gene knockout. Primers
were designed flanking the CRISPR cutting site to confirm transcript disruption. Increased ΔCt
number after perturbation indicates efficient knock down of unperturbed *ZNF766* mRNA.

d. Heatmap of differentially expressed genes between *ZNF766* knockout and non-targeting control
(NTC) T cells.

e. Volcano plot of differentially expressed genes between *ZNF766* knockout and NTC groups. Red
and blue dots indicate significantly upregulated and downregulated genes, respectively, in the
knock-out condition ($|\log_2 FC| > 1$, p.adj < 0.05). Representative genes involved in T cell activation
(e.g., *GZMB*, *MMP25*, *SLC7A5*) and exhaustion-associated regulators (e.g., *PDCD4*, *NT5E*,
TXNIP) were individually labeled.
f. Gene Ontology (Biological Process) enrichment analysis of significantly upregulated genes in
ZNF766 knockout group. Dot size corresponds to gene count per term; color indicates adjusted p-
value. Enriched pathways highlight enhanced DNA replication, chromatin organization, and
mitochondrial respiration, suggesting a functional restoration of exhausted T cells.

1756

Supplementary Fig. 15: Dynamics of CAP binding profile and CRE activities during Purkinje cell differentiation.

a. CAP binding profiles across CRE groups. Representative CAPs were labeled. Color scale indicates normalized binding.

b. T-SNE embedding of CREs colored by activity clusters (same as Fig. 6d).

c. Heatmap of CRE identity transitions across Purkinje cell differentiation stages.

d. Average binding dynamics for four different CRE sets across pseudo-time. Each CRE set includes three CAP classes (Repressor: e.g. TCF7, RCOR1; Early CAP: e.g. BCL11A, MEF2A, ATF2; Late CAP: e.g. FOXA1, FOXA2). Shading denotes 1 standard deviation.

e. T-SNE embedding of individualized CAP binding across loci using time trajectory as feature, colored by trajectory trend cluster. Each dot represents a CAP binding at a single locus across all pseudo-time stages. Average binding trajectories for each cluster across time were presented at bottom.

f. T-SNE embedding of CAP across time using CRE (binding profile) as feature, colored by CAP cluster identities. Each dot represents a CAP's binding profile across all loci at a single time point.

g. Alluvial diagram showing CAP cluster identity changes across time. Representative CAPs in each cluster were annotated on the right.