Decoding the gene regulatory landscape through multimodal learning of protein-DNA interactions

Jimin Tan[1,2,3#*], Xi Fu[4,5*], Xinyu Ling[1,6*], Shentong Mo[7], Jiangshan Bai[1,6], Raúl Rabadán[4,5], David Fenyö[2,8], Jef D. Boeke[2,8,9], Aristotelis Tsirigos[3,10#], Bo Xia[1,2,6,11#]

Gene Regulation Observatory, Broad Institute of MIT and Harvard, Cambridge, MA, USA
Institute for Systems Genetics, NYU Grossman School of Medicine, New York, NY, USA
Division of Precision Medicine, NYU Grossman School of Medicine, New York, NY, USA
Program of Mathematical Genomics, Department of System Biology, Columbia University, New York, NY, USA
Department of Biomedical Informatics, Columbia University, New York, NY, USA
Department of Pathology and Krantz Family Center for Cancer Research, Massachusetts General Hospital and Harvard Medical School, Boston, MA, USA
School of Computer Science, Carnegie Mellon University, Pittsburgh, PA, USA
Department of Biochemistry and Molecular Pharmacology, NYU Grossman School of Medicine, New York, NY, USA
Department of Biomedical Engineering, NYU Tandon School of Engineering, Brooklyn, NY, USA
Department of Pathology, NYU Grossman School of Medicine, New York, NY, USA
Society of Fellows, Harvard University, Cambridge, MA, USA
*Co-first Authors
#Corresponding Authors: Jimin.Tan@nyulangone.org; Aristotelis.Tsirigos@nyulangone.org; xiabo@broadinstitute.org

Abstract

The identity of a cell is governed by regulatory proteins binding to the genome to control gene expression. Mapping these genome-wide binding events across thousands of proteins and cell types is essential for understanding development and disease at scale, yet has remained a major experimental and computational barrier. Here we present Chromnitron, a multimodal foundation model that learns the rules of protein-DNA binding from protein sequence, DNA sequence, and context-specific chromatin states. Unlike prior single-task and multi-task learning approaches, Chromnitron implements a multimodal learning framework that accurately predicts the binding landscape for proteins and cell types not seen during training. Using Chromnitron, we discovered and experimentally validated new protein regulators of T cell exhaustion. Chromnitron also uncovered previously uncharacterized dynamic shifts in the binding landscape of regulatory proteins during neurogenesis. This marks a critical step toward a predictive model of interpretable gene regulatory programs across cell types, enabling rapid discovery of regulatory circuits and identification of new therapeutic targets.

INTRODUCTION

A central challenge in biology is to understand the regulatory logic that gives rise to diverse cell types and states. While the DNA sequence itself provides the basal genetic blueprint shared across all cells, this logic is executed by thousands of chromatin-associated proteins (CAPs) that interact with the genomic DNA to control cell type-specific gene expression in multicellular organisms[1–5]. These proteins – including transcription factors (TFs) that bind directly to DNA and indirectly recruited co-factors and chromatin regulators – engage thousands to millions of *cis*-regulatory elements (CREs) to orchestrate cell-type-specific gene expression programs (Fig. 1A)[6–21]. Elucidating the genome-wide landscape of protein-DNA interactions is therefore crucial for understanding the rules of gene regulation in disease and cell fate transitions[22–24].

Despite its importance, creating such a protein-DNA interaction landscape in a particular cell type has been prohibitive. Experimentally, methods such as ChIP-seq are resource intensive, measuring only one protein at a time and does not scale to primary samples due to cellular heterogeneity and scarcity in the tissue[25,26]. As a result, our knowledge of CAPs remains far from complete: genome-wide binding maps exists for fewer than half of the around 2,500 predicted human CAPs, and this data is mainly generated from cultured cell lines rather than primary cells[27–30]. Computational approaches, including recent machine learning tools that predict CAP binding from DNA sequence, have made progress but do not generalize across different cell types[31–34]. These limitations underscore the need for a predictive tool that can chart the CAP binding landscapes for any protein in any cell.

Here, we present Chromnitron (Chromatin omni-modal transformer), a biologically grounded multimodal foundation model that learns the principles of protein-DNA interaction by integrating three core modalities: DNA sequence, cell-type-specific chromatin accessibility, and protein amino acid sequences (Fig. 1B). Chromnitron was pretrained on an atlas of over 1,100 curated ChIP-seq CAP binding maps, covering 767 distinct proteins, and then fine-tuned for each CAP to capture protein-specific nuances. This cross-protein pretraining on multimodal data is key for Chromnitron to learn generalizable principles of protein-DNA interaction. Consequently, the model achieved remarkable performance in predicting CAP binding landscapes for proteins and cell types not seen during training (Fig. 1C). Chromnitron establishes a generalizable framework

70   for decoding the global gene regulatory landscape across cell types. We demonstrate that the

71   accurate and cell-type-specific predictions can be used to reconstruct causal regulatory programs

72   linking CAPs to CREs, discover key regulatory factors during immune cell fate transition, and

73   track dynamic CAP binding events along developmental trajectories (Fig. 1D).

Fig. 1: A multimodal framework for decoding the protein-DNA binding landscape.
A. Chromatin-associated proteins (CAPs) can bind DNA directly (transcription factors) and
indirectly (co-factors and chromatin regulators). **B.** A multimodal deep learning architecture that
integrates DNA sequence, cell-type-specific chromatin accessibility, and protein sequence to
predict CAP binding profiles. **C.** Chromnitron predicts genome-wide binding landscape for
hundreds of proteins. CAPs are colored by clusters based on binding similarity. **D.** Chromnitron-
predicted CAP binding landscapes enable downstream discovery applications, such as linking
CAPs to CREs to identify causal regulatory factors and modules, and tracking dynamic CAP
binding events across temporal trajectories.

RESULTS

Chromnitron learns multimodal principles of protein-DNA interactions

The specificity of CAP binding across the genome arises from protein functional domains,

genomic sequence, and local chromatin context. To learn these binding principles in a single model,

we designed a multimodal architecture that incorporates all three modalities as inputs (Fig. 1B and

Supplementary Fig. 1). In this architecture, DNA sequence and chromatin accessibility signals

were encoded by two independent convolutional modules, while CAP residues were encoded with

a pretrained protein language model (ESM)[35]. The resulting embeddings are subsequently merged

and processed by attention layers to allow implicit learning of both intra- and cross-modality

dependencies. The final decoder module processed the chromatin embedding with a transposed

convolutional module to produce CAP binding at single nucleotide resolution (Fig. 1B and

Supplementary Fig. 1).

Training a high-performance foundation model requires large-scale and high-quality data[35–38]. To

reduce technical variation across proteins and data sources, we first curated an atlas of 1,105 ChIP-

seq datasets covering 767 unique CAPs across five different cell types (Supplementary Fig. 2,

Methods). To improve consistency in data quality across CAPs and batches, we developed a fully

automated and scalable preprocessing infrastructure, Chrom2Vec, which prepares raw sequencing

data for training with custom denoising and signal-scale standardization across CAPs and cell

types (Supplementary Fig. 2)[36,39,40].

Next, we trained Chromnitron in two phases: cross-CAP pretraining and per-CAP fine-tuning. Our

training procedure differs from traditional paradigms such as single-task learning where separate

models were trained for each CAP, and multi-task learning where all CAPs were predicted

simultaneously from different output heads[32,33,41,42]. The multimodal learning implemented by

Chromnitron at the cross-CAP pretraining step allows the model to learn shared CAP-binding

principles across proteins, establishing a single foundation model[41]. Because multimodal

pretraining is computationally intensive due to its combinatorial nature, we developed an algorithm,

OuterMix, that reduces the exponential sampling complexity down to linear, yielding roughly a

five-fold increase in training efficiency (Supplementary Fig. 3).

Accurate and cell-type-specific prediction is essential for downstream biological discoveries. To

assess whether multimodal architecture enables generalizable prediction in unseen cell-type

contexts, we first trained a prototype using data from only two cell types and evaluated its *de novo*

prediction performance in three leave-out cell types on leave-out chromosome 10. This prototype

reliably   outperformed   leading   unimodal   frameworks,   such   as   Enformer,   Borzoi,   and

AlphaGenome, when predicting cell-type-specific CAP binding signals (Supplementary Fig. 4)
[32,33,42].

These results highlight the value of modeling CAP binding with multimodal learning incorporating
cell-type-specific chromatin features. We therefore scaled training to all 1,105 quality-controlled
ChIP-seq datasets across five cell types. During cross-CAP pretraining, we observed distinct
learning dynamics across CAP groups, potentially reflecting their differential binding mechanisms
(Supplementary Fig. 5). After pretraining, we performed low-rank adaptation (LoRA) fine-tuning
to produce individual sub-models for each CAP, which further increased overall performance and
training efficiency (Supplementary Fig. 6)[43].

Fig. 2: Chromnitron performance and interpretability.
A. Paired Chromnitron prediction (Pred) and experimental (Exp) ChIP-seq tracks in leave-out
chromosome. **B.** Benchmarking of prediction performance in cell-type-specific CAP-binding

regions, measured with Pearson's correlation (left) and average precision (AUPR, right). Acc. Motif, accessibility-weighted motif score. Bar plots in this figure show mean ± s.e. **C.** Scatter plots illustrating the relationship between CAP binding and chromatin accessibility for representative CAPs, and comparing experimental ChIP-seq signal (Exp, grey) with Chromnitron predictions (Pred, blue). **D.** Heatmaps comparing predicted and experimental CAP-binding signals across three classes of genomic regions: accessible with CAP-binding (Joint), inaccessible with binding (Binding-only), and accessible without binding (Accessible-only). Each row in the heatmaps represents a 2 kb genomic window centered on a peak region in leave-out chromosome. Corresponding chromatin accessibility (ATAC-seq) signals are shown in parallel. **E.** DNA sequence impact motifs learned by Chromnitron, identified through *in silico* single-nucleotide mutagenesis. Top: representative impact motifs. Bottom: heatmap and hierarchical clustering of cross-correlations among impact motifs across all pairs of CAPs. **F.** Identification of critical CAP amino acid residues via *in silico* protein residue perturbation. Left: per-residue impact scores along the protein sequences of SOX13 and CREB1. Right: corresponding impact scores mapped onto AlphaFold-predicted protein-DNA complex structures, using the same color map of binding impact scores. **G.** Distribution of per-residue impact scores across the top ten annotated protein domains, ranked by average impact score. Points in the violin plots denote individual CAPs, with the mean shown for each violin.

Chromnitron accurately predicts cell-type-specific CAP binding

We systematically evaluated Chromnitron on multiple prediction tasks using chromosome 10 as a leave-out test chromosome. We found that Chromnitron accurately predicts CAP-binding profiles on par with experimental replicates (Supplementary Fig. 7). The predictions recapitulate experimental measurements across large genomic regions with varying CAP binding intensities and chromatin states (Fig. 2A, Supplementary Fig. 8-10). To assess cell-type-specific prediction performance, we benchmarked against other models on regions differentially active between liver carcinoma line HepG2 and leukemia line K562. In this context, the fully trained Chromnitron markedly outperformed both sequence-based multi-task models and other single-task models that use cell-type-specific features (Fig. 2B, Supplementary Fig. 11–13). This performance gap is particularly notable given that our evaluation was performed on a leave-out chromosome that was part of the training sets of the other models, suggesting a larger true performance improvement.

Notably, Chromnitron learns the non-linear, CAP-specific relationship between binding and chromatin accessibility, which is essential for generalizing across cellular contexts. Comparing experimental CAP binding signals (e.g. ChIP-seq) to chromatin accessibility, we found that many CAPs showed weak correspondence between these signals. While CAP binding is often enriched

at highly accessible chromatin regions, some CAPs, such as pioneer factors, show strong binding

in regions with low accessibility (Fig. 2A and 2C, Supplementary Fig. 14). Existing tools that infer

CAP binding by combining motifs with accessibility peak signals are dependent on peak strength

and therefore prone to false negative predictions in low-accessibility regions[44,45]. In contrast,

Chromnitron accurately predicts binding signal in both high- and low-accessibility regions, owing

to its ability to leverage a richer and non-linear representation of local chromatin contexts (Fig.

2D). These results highlight Chromnitron's utility for characterizing key regulators that

preferentially bind at lowly accessible regions, such as pioneer factors that drive cell fate

transitions[46,47].

***In silico* perturbations reveal principles of protein-DNA interaction**

To further interrogate the CAP binding principles learned by Chromnitron, we performed

systematic *in silico* perturbation across all three input modalities. We first applied single-

nucleotide mutagenesis along genomic loci and quantified the effect of each mutation on predicted

CAP binding (Supplementary Fig. 15). These per-base impact scores recapitulate canonical TF

consensus motifs derived from high-throughput experimental measurements[48,49], which suggests

that Chromnitron learns sequence features that determine CAP binding (Fig. 2E). Extending such

in silico DNA mutagenesis across all CAPs revealed motif modules corresponding to distinct CAP

families, along with unique ones that are previously uncharacterized (Fig. 2E and Supplementary

Fig. 15). Notably, we identified coherent impact motifs for proteins lacking recognizable DNA

binding domains, such as cohesin subunits, whose impact motifs resemble that of their known

partner CTCF (Supplementary Fig. 15). Using the same sequence perturbation method,

Chromnitron can predict cell-type-specific CAP binding changes resulting from non-coding

variants in both gain-of-binding and loss-of-binding mutation events (Supplementary Fig. 16).

Perturbing chromatin accessibility revealed a non-linear relationship between CAP binding and

accessibility levels (Supplementary Fig. 17), with binding intensity of different CAPs saturating

at varied levels. These results echoed our observation from the experimental data that many CAPs

bind efficiently to inaccessible or lowly accessible chromatin regions, and support that

Chromnitron learned accessibility-related features and incorporated them for prediction

(Supplementary Fig. 17, Fig. 2C).

Lastly, to examine important features encoded in the CAP amino acid sequences, we applied

saturated *in silico* mutagenesis to CAP amino acid sequences (Supplementary Fig. 18). This *in*

silico experiment revealed that high-impact residues, which correspond to high prediction changes,

align precisely with known DNA binding domains, as illustrated by SOX13 and CREB1 (Fig. 2F).

Generalizing mutagenesis across CAPs revealed strong enrichments in evolutionarily conserved

DNA-binding domains such as bZIP and E2F, and depletion in domains that do not contribute to

DNA sequence specificity (Fig. 2G, Supplementary Fig. 18). These results suggest that

Chromnitron learns protein domain features relevant for their binding specificity directly from

CAP amino acid sequence. Consistent with this analysis, we found Chromnitron enables *de novo*

prediction of CAP binding profiles for previously unseen CAPs. When trained on a subset of the

CAPs and evaluated performance on the leave-out ones, the model can perform *de novo* predictions

on unseen CAPs and outperforms the best baseline inference using motif score multiplied by

accessibility (Supplementary Fig. 19). Together, these multimodal perturbation analyses show that

Chromnitron captures core principles of CAP binding: (i) DNA sequence-encoded motif features

and co-binding syntax; (ii) CAP-specific and nonlinear dependencies on chromatin accessibility;

and (iii) protein-structural determinants of DNA recognition. These learned rules enable the model

to effectively predict functional genomic data at scale and to generalize across unseen genomic

loci, cell types, and CAPs.

Fig. 3: Chromnitron discovers novel regulators of T cell exhaustion.

A. Dimensionality reduction (UMAP) of a T cell fate trajectory from naïve (T0) to exhausted (T3)
states (top), with corresponding chromatin accessibility profiles highlighting a promoter-proximal

regulatory element at the *TOX* locus (bottom). **B.** Chromnitron-predicted CAP-binding dynamics
at the *TOX* promoter-proximal CRE across exhaustion pseudo-time, highlighting a unique and
progressive increase in binding of an uncharacterized protein, ZNF865 (blue). **C.** Flow cytometry
quantification of cytokines (IL-2, IFN-γ, and TNF-α) and T cell exhaustion markers (PD1, LAG3,
TIGIT, and TIM3) across negative control (Neg Ctrl), non-targeting control (NTC), and *ZNF865*
knockout (*ZNF865* KO) conditions. **D.** Schematic of the Chromnitron-based workflow for
genome-wide characterization of co-regulation modules linking CAPs to genes by comparing
effector and exhausted T cell conditions. **E.** Differential CAP-binding matrix between effector
(Teff) and exhausted (Tex) T cells, clustered by both co-regulating CAPs (rows) and co-regulated
gene groups (columns). A zoom-in plot highlights a subset of CAP modules (C0, C2, C3) and gene
groups (G0, G4) enriching reported T cell exhaustion marker genes. **F.** Ranking of CAPs by
differential binding intensity (log2 fold change between Tex over Teff), highlighting both known
(grey) and unknown (blank) top candidate regulators. **G.** Scatter plot of targeting-gene sgRNA
enrichment p-values from CRISPR knockout (x-axis) and knockdown (y-axis) screens,
highlighting *ZNF766* as the top hit in both assays. **H.** Flow cytometry quantification of cytokines
(IL-2, IFN-γ, and TNF-α) and T cell exhaustion markers (PD1, LAG3, TIGIT, and TIM3) across
negative control (Neg Ctrl), non-targeting control (NTC), and *ZNF766* knockout (*ZNF766* KO)
conditions. Bar plots in (**C**) and (**H**) show mean ± s.d. for *n* = 3 independent replicates. Statistical
comparisons were performed using two-sided unpaired Student's *t*-tests with significance denoted
as *, P < 0.05; **, P < 0.01; ***, P < 0.001.
247

Chromnitron identifies ZNF865 as a *TOX* regulator during T cell exhaustion

The cell-type-specificity of Chromnitron allows the model to predict CAP binding in unseen

cellular contexts. To demonstrate its utility for dissecting disease relevant gene regulation in

human, we applied Chromnitron to study T cell exhaustion, an immune suppression mechanism

common in cancer. T cell exhaustion is a progressive cell state transition driven by chronic antigen

stimulation, leading to a dysfunctional state characterized by enhanced expression of inhibitory

cell surface receptors, reduced cytokine production, and activation of *TOX*[50–57]. Single-cell ATAC-

seq (scATAC-seq) of tumor-infiltrating T cells has captured the chromatin state transition from

naïve to exhausted T cell but lacks mechanistic insight into the underlying regulation[58].

To pinpoint candidate locus-specific regulators of *TOX* expression, we applied Chromnitron to

track predicted CAP binding dynamics at the *TOX* locus across the T cell exhaustion trajectory

derived from scATAC-seq data (Fig. 3A, Supplementary Fig. 20). Surprisingly, ZNF865, an

uncharacterized C2H2 zinc-finger protein, emerged as a unique regulator at a *TOX* promoter-

proximal regulatory element with a steady increase in binding along the exhaustion pseudo-time

(Fig. 3B). Notably, ZNF865 show preferential binding at low-accessibility regions, including this

TOX promoter-proximal element, whose chromatin is markedly less accessible than the promoter

itself (Supplementary Fig. 20). Further *in silico* mutagenesis revealed that ZNF865 recognizes a

previously uncharacterized CA-repeat sequence pattern, which explains in part its unique binding

to the *TOX* promoter-proximal element (Supplementary Fig. 20).

The timing and specificity of ZNF865 binding suggested that it may act upstream of *TOX* during

T cell exhaustion. To test this, we employed an established *in vitro* T cell exhaustion model using

persistent anti-CD3 antibody or anti-CD3/CD28 cocktail stimulation together with CRISPR-

mediated knockout of *ZNF865*[59–61] (Supplementary Fig. 21). Consistent with the hypothesis,

knockout of *ZNF865* reduced *TOX* expression levels and consequently led to reduction in surface

exhaustion markers and elevated effector cytokine production, suggesting partial restoration of the

effector function (Fig. 3C, Supplementary Fig. 22).

A genome-wide screen with Chromnitron uncovers ZNF766 as a regulator of T cell

exhaustion

Having shown that Chromnitron can resolve a CAP-mediated regulatory circuit at one locus, we

next asked whether it could dissect the genome-wide regulatory program driving T cell exhaustion.

We predicted CAP binding across promoters whose accessibility changes between effector and

exhausted T cells. Comparing the two states generated a differential gene-by-CAP matrix in which

each entry represents differential binding of a CAP at a gene promoter (Fig. 3D). Clustering this

matrix revealed distinct modules of co-regulated genes and CAP groups predicted to act on them

(Fig. 3E). Gene modules enriched for known exhaustion marker genes aligned with CAP modules

showing the strongest differential binding (C0, C2, C3; Fig. 3E, Supplementary Fig. 23). These

CAP modules contained both established regulators of T cell exhaustion and several

uncharacterized ones, including the KRAB zinc finger protein ZNF766 (Fig. 3F). This analysis

suggested that these CAP groups may cooperate as upstream regulators of the exhaustion program.

To assess the functions of these candidate CAPs from the top regulator groups, we conducted a

focused CRISPR screen targeting 65 regulators genes nominated by Chromnitron using the *in vitro*

T cell exhaustion model. We designed two separate sgRNA libraries, optimized for lentivirus-

based gene knockout and transcriptional knockdown, respectively. Following library transduction,

human primary CD8[+] T cells were subjected to persistent anti-CD3 treatment, with T cell

proliferation serving as the phenotypic readout. Across both knockout and knockdown screens,
ZNF766 consistently showed the strongest functional effect, reducing T cell persistence under
chronic stimulation (Fig. 3G). Independent validation via targeted gene knockout of ZNF766
confirmed the screening results, with ZNF766-deficient T cells express significantly lower surface
exhaustion markers, along with increases in effector cytokine production (Fig. 3H, Supplementary
Fig. 23). RNA-seq following *ZNF766* knockout revealed broad transcriptional remodeling under
chronic stimulation, consistent with the functional rescue (Supplementary Fig. 24). Together,
these findings demonstrate that Chromnitron can identify genome-wide regulatory modules and
uncover previously unrecognized CAPs involved in T cell exhaustion.

Fig. 4: Chromnitron maps the dynamic regulatory landscape of neurogenesis.
A. Single-cell ATAC-seq revealed differential CRE activities during Purkinje neuron development.
B. Chromnitron predicts a three-dimensional tensor of CAP-binding landscapes spanning the
genomic position, CAP identity, and developmental time. **C.** Heatmap of predicted CAP-binding
signals at key CREs, with CAPs organized by binding similarity (rows) and CREs grouped by

developmental stage (columns). **D.** Dimension reduction (t-SNE) projection of CREs across
pseudo-time based on predicted CAP-binding profiles, revealing a continuum of regulatory
element identities ranging from inactive (blue) to active (red). **E**. Vector field representation of the
directional CRE state transitions between consecutive developmental stages. The vector fields
were projected onto the t-SNE map in (**D**).
316

Chromnitron maps the dynamic regulatory landscape during neurogenesis

To demonstrate its power in developmental contexts, we applied Chromnitron to predict the

genome-wide dynamics of CAP-mediated regulation during neurogenesis using single-cell

chromatin accessibility dataset (Fig. 4A–B). This yields a three-dimensional CAP-binding tensor

spanning genomic position, CAP identity, and developmental time, providing a scalable

framework to characterize how CAP binding and CRE activities are globally orchestrated (Fig.

4B).

We applied this analytical framework to a trajectory describing Purkinje neuron maturation during

first-trimester neurodevelopment[62]. When projected as CAP-by-CRE matrices grouped by

developmental stage, the data revealed a clear transition in CRE activity, with a sharp shift

occurring between neuroblast and early Purkinje states (Fig. 4C). To dissect stage-specific

transitions in CRE activities along the developmental time trajectory, we performed

dimensionality reduction on CAP features and clustered CREs. This mapping revealed a spectrum

of CRE identities ranging from inactive to active states, revealing a genome-wide redistribution of

regulatory element identity during maturation (Fig. 4D, Supplementary Fig. 25). Tracking

individual CREs across consecutive stages generated a vector field that visualizes their "migration"

through regulatory space, showing that stem-cell-specific and Purkinje-specific elements shift

dynamically during early stages before stabilizing later in development (Fig. 4E–H).

Interrogating the binding tensor along its remaining axes provided additional insights into

regulatory dynamics. Using developmental time as the organizing axis, we identified cohorts of

early-acting and late-acting CAPs along the Purkinje cell maturation trajectory, many following

monotonic gain- or loss-of-binding trends (Supplementary Fig. 26). Using the CREs as the feature

dimension, we found distinct CAP binding signatures across developmental time, with cohesin-

related proteins, chromatin remodelers, and repressors exhibiting more stable binding patterns

across time than transcriptional activators (Supplementary Fig. 26). With its unprecedented scale

of functional genomic prediction, Chromnitron enables systematic and multidimensional analyses

required to elucidate CAP-mediated regulatory dynamics underlying cell-fate transitions during

development.

DISCUSSION

Understanding how CAPs configure cell-type-specific gene regulatory programs remains a major

challenge in human biology. By integrating DNA sequence, chromatin state, and protein features,

Chromnitron learns the multimodal determinants of protein-DNA interactions and enables highly

accurate prediction of CAP-binding landscapes in unseen cell types. The model's high predictive

performance enables large-scale *in silico* experimentation, substantially narrowing the candidate

search space while generating interpretable predictions that can directly guide laboratory studies.

Leveraging high-throughput *de novo* prediction, we discovered previously uncharacterized

regulators of T cell exhaustion and mapped global dynamics of CAP binding and CRE activities

during neurodevelopment. Chromnitron demonstrates how foundation models grounded in

biological principles can accelerate hypothesis generation, enable targeted experimental validation,

and reveal new biological insights.

Beyond its practical applications, Chromnitron also uncovers both known CAP-chromatin binding

rules and non-canonical interactions at low-accessibility or inaccessible regions with previously

unrecognized sequence motifs. While sequence-based models excel at pinpointing nucleotide-

level determinants of regulatory activity, Chromnitron addresses a complementary challenge:

learning the fundamental rules of CAP binding through a multimodal architecture, enabling

generalizable and high-performance prediction across unseen cell types and chromatin states.

Chromnitron's modular design allows the integration of pretrained DNA language models as

alternative or additional encoders to further increase training efficiency and

performance[32,33,35,37,42,63,64].

In addition, this multimodal framework provides a natural foundation for incorporating CAP

binding data generated from perturbation experiments. Whereas conventional single- or multi-task

architectures typically require specialized model modifications to accommodate perturbed CAP

profiles, Chromnitron can natively integrate protein-residue perturbation datasets and learn

relationships between amino acid sequence variation and DNA binding at scale. As more diverse

and perturbed CAPs are incorporated in future iterations, Chromnitron's ability to perform *de novo*

predictions for previously unseen factors can further improve.

Despite its high prediction accuracy, Chromnitron's performance is dependent on training data

quality. The current training datasets are generated primarily from conventional ChIP-seq

experiments, which are subject to variable antibody quality, lack of dosage measurement, and

batch effects. Although substantial computational efforts were made to denoise and standardize

the data, not all processed ChIP-seq datasets reach high quality. These challenges underscore the

need for generating standardized, high-quality CAP-binding datasets. Importantly, the pretrained

foundation model can be readily updated by fine-tuning with improved CAP-binding datasets.

Generating standardized CAP-binding datasets covering the remaining uncharacterized human

CAPs will further expand the scope and power of Chromnitron's prediction catalogue.

Predicting how a cell will behave under a given perturbation remains one of the most important

and challenging problems in regulatory genomics. Like many other biological processes, cellular

responses are intrinsically multifaceted, emerging from molecular interactions that span across

multiple hierarchies and typically result in regulatory changes in gene expression. CAPs sit at the

core of these regulatory mechanisms, directly controlling the induced changes in gene expression

and cell state transitions that underlie development and disease. Chromnitron represents a

substantive step toward a predictive, interpretable, and mechanistic framework for understanding

gene regulation, highlighting the potential of multimodal foundation models to transform

functional genomics. By accurately modeling the CAP binding landscape, Chromnitron provides

a powerful tool for dissecting the mechanisms that drive cellular identity changes, building

mechanistic virtual cell models, and enabling rational cell-fate engineering.

Materials and Methods:

Model architecture

We designed Chromnitron (Chromatin Omni-modal Transformer) to integrate three information
streams that jointly determine CAP-chromatin binding: DNA sequence, local chromatin
accessibility, and the amino-acid sequence of the assayed CAP. Chromnitron comprises three
successive components: modality-specific encoders, a multimodal self-attention module, and a
decoder (Supplementary Fig. 1).

The model has three modality-specific encoders. The encoders for DNA sequence and chromatin
accessibility consist of similar convolutional blocks (ConvBlocks). Each block begins with a
strided Conv1D layer that covers large local windows while halving the sequence length. The
convolution is followed by group normalization (groups = 1) and a Gaussian Error Linear Unit
(GELU) activation function[65]. Feature maps then traverse 3 residual ConvBlocks (kernel = 9; stride
= 1) that maintain length while expanding the hidden dimensionality to 384. Outputs from the
DNA and accessibility encoders were projected to the same embedding of 512 × 384. The two
embeddings were fused on the feature dimension to form a joint embedding of size 512 × 768. The
joint embedding was scaled down to a 512 × 384 chromatin embedding with a header for
subsequent fusion with the protein embedding.

For the protein encoder, we leveraged a pretrained protein language model ESM2 as a protein
feature extractor[35]. As running ESM2 inference during training can add to the computational
overhead, we extracted and cached the protein features before the training step to increase model
training efficiency. A non-redundant list of human transcription factors was assembled by merging
all proteins for which we had ChIP-seq data and the available proteins in UniProt database. To
encode the embeddings, we used the publicly available ESM2 transformer with 36 layers and 3
billion parameters (esm2_t36_3B_UR50D, hidden size = 2560). For each protein, we requested
the hidden representations of the final, 36th transformer block. Since ESM2 accepts up to 1,024
tokens, we split any long proteins into non-overlapping 1,024-residue segments, each embedded
independently. The resulting matrices were concatenated along the sequence dimension to
reconstitute the full-length protein representation. The number of residues was capped at 4,096 to
accommodate most of the full-length CAP sequences while keeping a low compute cost. The per-
residue embeddings were converted to arrays and stored in a compressed npz file. During training
these cached embeddings are directly loaded, transformed by a header, and fused with genomic
features in the multimodal attention module.

We implemented the attention mechanism in transformer to fuse contextual information both
within and across modalities[66]. Chromatin feature embeddings and protein embeddings were
concatenated along the length dimension, separated by a single zero vector that marks the modality
boundary. Positional information was added via fixed sinusoidal encodings. The resulting fused
feature embedding has a length of 769 (512 from chromatin embedding, 256 from protein
embedding, and a separator with a length of 1). The fused embedding was processed by 16 layers

of pre-layer-normalized transformer. Each transformer layer has 8 attention heads, 384 hidden
state dimension, 2,048 feed-forward layer dimension, and a dropout rate of 0.1. Self-attention
allows every genomic position to attend not only to distant nucleotides but also cross-modal to any
residue of the corresponding CAP.

After transformer module, we only kept features corresponding to genomic positions and
forwarded them to the decoder. The decoder has four ConvTranspose blocks successively up-
sample the latent representation to the original 8,192-bp resolution while preserving contextual
information through residual connections. A final convolution header produces a one-dimensional
per-base binding profile for each target CAP.

Pretrained Chromnitron model has 72 million parameters. Each LoRA fine-tuned sub-model
weighs 9 million parameters. The fully fine-tuned Chromnitron model with pretrained and LoRA
weights has over 6.9 billion parameters in total.

Genomic data collection
To obtain training data with more standardized quality, we primarily focused on consortium-
generated datasets for consistency. We developed a systematic approach to collect and curate high-
quality CAPs ChIP-seq datasets from the ENCODE database[6]. Our data collection strategy focused
on obtaining non-revoked and non-archived datasets with sufficient sequencing depth to ensure
reliable modeling. We required a minimum of 40 million reads to achieve adequate genome
coverage for CAPs ChIP-seq and ATAC-seq, while immunoprecipitation control experiments
required 100,000 reads. Chromatin accessibility information is essential for understanding the
regulatory context of transcription factor binding. To ensure experimental coherence and enable
integrative analysis, we retained only those CAPs ChIP-seq datasets that had good-quality ATAC-
seq data in the corresponding cell type, resulting in 1,105 qualified ChIP-seq from five different
cell types, covering 767 unique CAPs.

Genomic data preprocessing with Chrom2Vec
To prepare precise and highly consistent training data, we designed Chrom2Vec, a modular
bioinformatic platform that streamlines training data processing for Chromnitron. The Chrom2Vec
platform starts from raw FASTQ files and delivers training-ready data tensors. Chrom2Vec is
based on Apache Airflow and orchestrates next-generation-sequencing processing for both ChIP-
seq and ATAC-seq. It implements configurable pipelines that handle distributed workers, tracking,
and reproducible containers, performs robust assay-specific normalization, and signal
compression to optimize data storage. Pipelines are described by lightweight Directed Acyclic
Graphs (DAGs) that call a library of preprocessing operators, which are Airflow wrappers around
containerized shell or Python scripts to ensure maximum reproducibility. While tasks are being
scheduled and executed by Airflow, the metadata are streamed to Neo4j for full data-lineage
tracking. To reduce storage and I/O overheads without sacrificing data quality on key genomic
regions, we implemented an adaptive dynamic-bin compression algorithm. We observed as high
as 50% reduction in storage for some of the high-quality tracks. The final compressed training

dataset is around 952 GB in size. This architecture was optimized for machine-learning workflow
as it guarantees reproducibility, supports scaling, and writes outputs in chunked Zarr format which
is directly ingestible by the model and supports fast random-access during training.
491
To enable quantitative comparisons across cell types and experimental conditions, we performed
peak-based quantile normalization on bulk ATAC-seq, ChIP-seq and pseudo-bulk single cell
ATAC-seq data. We followed a two-step approach that first removed noise and then rescaled data.
The noise level can be represented by signal variations in the background genomic regions. To
define these regions, we used cPeaks from Meng et al., a union of accessible peaks across human
cell types[67]. The background region is defined as its complementary regions on the genome,
excluding ENCODE blacklist[68].
499
For ChIP-seq, we performed a two-stage correction that explicitly models both biochemical
background and sequencing bias. First, chromatin signal was extracted from immunoprecipitated
(IP) and matched input libraries. Using the pre-defined background regions, we calculated the
mean IP and input signal in the background bins. The input was rescaled to reach similar
background signal intensity as IP. The rescaled input was then subtracted from the IP track,
effectively removing experimental bias across the genome. To harmonize dynamic range across
ChIP-seq experiments, the corrected IP signal was divided by the average of the 99 to 99.9
percentile values with 1 percent step sizes, computed over the pre-defined active peak regions from
cPeaks. This normalization puts the strongest peaks near a value of one while retaining linearity at
regions with lower signal intensities. The result is a bias-corrected, unit-less collection of ChIP-
seq datasets that are comparable across CAPs and experiments, dramatically facilitating training
stability. After scaling the training datasets, we further measured a background-noise level for each
dataset by calculating standard deviation in the background chromatin regions defined above. The
training data quality metric was calculated by (1 – noise) for downstream model performance
evaluation.
515
For ATAC-seq, where a control library was not available, we removed noise from background
directly followed by percentile scaling. The algorithm first estimates systematic noise by averaging
the coverage signal over the same background regions. This background mean was subtracted
from every genomic position, eliminating basal accessibility level. We collected mean cPeak
signals, and computed the scaling constant based on top 99 to 99.9 percentiles similar to ChIP-seq.
Dividing the background-corrected track by this constant compresses the dynamic range so that
the signals of typical accessible loci fall between 0 and 1 (though exceptionally strong regulatory
regions can exceed 1), producing normalized ATAC-seq datasets that are consistent and
comparable across batches and cell types.
525
For single cell ATAC-seq data, we started with fragment files and processed them through a
pipeline with shared preprocessing steps as Chrom2Vec and generates quantitative chromatin
accessibility profiles. Fragment coordinates were standardized to the hg38 reference genome. Cells
with 1,000-50,000 fragments per cell were retained to exclude low-quality cells with insufficient

chromatin accessibility signal and potential doublets with artificially inflated fragment counts.
Individual cells were subsequently aggregated into pseudo-bulk profiles based on cell cluster
annotations (i.e. cell types or states) in original publication, ensuring each pseudo-bulk has over
10 million unique sequencing reads (or 20 million Tn5 transposase insertions). These accessibility
reads from all cells within each cluster were summed to generate high-resolution chromatin
accessibility tracks with improved signal-to-noise ratios compared to individual cells. Each
pseudo-bulk data represents a unique cell type or state, and was normalized by the same pipeline
described in the data normalization section.
538
Model pretraining and fine-tuning
Training data were assembled from the three input feature modalities and the target CAP binding
profiles. Each training sample unit was set to an 8,192-bp genomic region. For the two chromatin
input features, we extracted a five-channel (A, C, G, T, N) one-hot representation of the hg38 DNA
sequence and a single-channel chromatin-accessibility track (ATAC-seq Tn5-transposase insertion
frequency signal). To stabilize training, the ATAC-seq signal was transformed with a monotonic
log1p function. CAP binding profiles were retrieved from the normalized ChIP-seq coverage for
each CAP in every cell type. In a separate preprocessing step, we embedded the primary amino-
acid sequence of every CAP with the 36-layer ESM2 protein language model
(esm2_t36_3B_UR50D). This embedding was cached and loaded on to the protein header during
training.
550
Random samples from different modalities are recombined to form the final training batch.
Batched tensors were shaped as (batch, length, channels) for DNA and ATAC, and (batch, residues,
channels) for ESM2 embeddings. The model takes in these inputs and predicts the target CAP
binding profile of shape (batch, length). Validation loss was calculated as the mean-squared error
(MSE) between predicted and observed log1p-signal. We used Adam[69] as the optimizer with a
learning rate of 0.00002 and weight decay of 0. The scheduler used ReduceLROnPlateau strategy
with a patience of 5 epochs. After validation loss stabilizes at 390 epochs, we fine-tuned the model
for another 10 epochs for each CAP with low-rank adaptation (LoRA), updating the CAP-specific
sub-model parameters[70]. LoRA layers were directly applied to feedforward layers in the attention.
For convolution and convolution transpose layers, we custom-designed the LoRA layers, with
details described in the following sections.
562
The model was pretrained on a node with 4 NVIDIA A100 80GB for 300 Hours with FP32 full
precision. The pretraining stage took around 1,200 GPU hours and has a total compute of 8.7e19
FLOPs (floating-point operations). The fine-tuned model was trained in parallel on a cluster with
32 NVIDIA A100 GPUs using TF32 precision, which has much higher throughput. Each fine-
tuning was set to run 8 hours for 10 epochs. The fine-tuning step was performed on all 767 CAPS,
leading to a total compute of 3.4e21 FLOPs, and took around 6,000 GPU hours using the lower
precision TF32 tensor format. The total amount of compute for training and fine-tuning is around
3.5e21 FLOPs.
571

Compared with previous methods that trained separate models for individual CAPs, Chromnitron
training employed a cross-CAP pretraining with per-CAP fine-tuning strategy which dramatically
increased both training efficiency and model performance. To measure the training efficiency
difference between individualized training approach versus LoRA finetuning, we used CTCF as
an example to compare the two approaches. We found that training the model from scratch took
more than 30 epochs to reach good performance, whereas fine-tuning from a pretrained model
only took 1 epoch to reach comparable performance. As the pretraining compute is only a small
fraction of the fine-tuning compute ($< 3\%$ by estimation), assuming all CAPs requires a similar
level of compute training from scratch, we estimated that the pretraining with fine-tuning strategy
is at least one order of magnitude more efficient compared with individually training models for
each CAP from scratch. The stored LoRA weight matrix is also 8 times smaller than full weight
matrix. Meanwhile, pretraining substantially increases overall model performance through its
generalized learning of CAP-binding principles across different CAPs.
585
Efficient multimodal sampling with OuterMix
To improve model training efficiency at the cross-CAP pretraining phase, we developed OuterMix,
an algorithm to generate Cartesian product of encoded features from randomly sampled data points
on each dimension as the training units. OuterMix maximizes throughput by re-using modality-
specific embeddings generated by each encoding step, thus dramatically reduces training
complexities, and increases data loading and encoding efficiency exponentially.
592
First, a naïve sampling approach for multimodal training through randomly selecting a fixed
number of examples for each batch would not be practical due to exponentially increased
complexities. Each sample consists of random examples from different modalities. Given a fixed
U number of samples from each modality with V modalities in total, the compute complexity
would be $O(U^V)$. Formally, if we are sampling $N \cdot M \cdot K$ samples in a batch representing the total
amount of genomic loci, chromatin feature, and CAPs during pretraining, the three encoders need
to encode all the samples. Using l, t, c to represent locus, cell type and CAP, respectively, the total
random sample space for a batch $\mathcal{B}_{rand}$ can be defined as:

601
$$\mathcal{B}_{rand} = \bigcup_{i=1}^{NMK}\{(l_i, t_i, c_i)\}, \ (l_i, t_i, c_i) \sim \mathcal{U}(\mathcal{L} \times \mathcal{T} \times \mathcal{C})$$

602 where each encoder needs to run across all samples.
603
604 OuterMix algorithm was designed to reuse embeddings generated by each encoder and reduce an
605 exponential complexity $O(U^V)$ down to a linear $O(UV)$ complexity, increasing data loading and
606 encoding efficiency exponentially, especially with large encoders and a high number of modalities.
607 With OuterMix, we started by randomly selecting fixed number of samples from each dimension
608 and produce a Cartesian product of loci (l_i), cell types (t_j) and proteins (c_k):

609
$$\mathcal{B}_{OuterMix} = \bigcup_{j=1}^{N}\bigcup_{i=1}^{M}\bigcup_{k=1}^{K}\{(l_i, t_j, c_k)\}$$

Here, each encoder only runs through samples from their predefined subset (Supplementary Fig.
3). In our implementation, the genome was first divided into non-overlapping 100kb chunks. In
each iteration, the data loader selects one genomic chunk per cell type, followed by sampling 8,192
bp windows and applying lightweight augmentations. For each window, we randomly selected up
to 4 CAPs with their corresponding DNA sequence, ChIP-seq, and ESM2 data. Modality-specific
encoders then generate feature-specific embeddings, which are tiled and concatenated to form the
Cartesian product samples ready to be fed into the Transformer. This design allows the same
sequence-ATAC tensors to be paired with multiple proteins, increasing Transformer trunk
throughput while avoiding additional disk I/O and encoder overheads.
619
LoRA implementation during fine-tuning
Since Low-Rank Adaptation (LoRA) was originally developed for fine-tuning natural language
applications such as Large Language Models (LLMs), its implementation for one-dimensional
convolution and convolution-transpose layers used in Chromnitron is not trivial[70]. Here we
implemented LoRA for both Conv1D and ConvTranspose1D. On a high level, LoRA for
convolution layers was implemented by freezing the original kernel and learning a low-rank update
that can be added onto the kernel at evaluation time. The 1-dimensional convolutional layer
weights are 3D tensors representing input dimension, kernel size, and output dimension. In this
implementation, we reshaped input and kernel size into one dimension and used it to construct two
low-rank matrices.
630
For Conv1D, the convolutional kernel weights $W \in \mathbb{R}^{C_{\text{in}} \times C_k \times C_{\text{out}}}$ are kept frozen during fine-
tuning, with C_{in}, C_k, and C_{out} representing input size, number of kernels, and output size,
respectively. The weight is separated into two low-rank matrices A and B with rank r, defined as:
$$A \in \mathbb{R}^{(r \cdot k) \times (C_{\text{in}} \cdot k)}, B \in \mathbb{R}^{C_{\text{out}} \times (r \cdot k)}$$
and is updated through $\Delta W = B A$. In the forward pass, the weight is added to the frozen weight
to apply fine-tuning changes:
$$y = \text{conv1d}\left(x, W + \frac{\alpha}{r} \cdot \Delta W\right)$$
where y is the convolution activation and α defines the strength for LoRA weights in which the
default is set to 1.
640
For ConvTranspose1D, the same decomposition is applied with a custom conv_transpose1d
forward pass, applying the ΔW, similar to conv1d:
$$y = \text{conv1d\_transpose}\left(x, W + \frac{\alpha}{r} \cdot \Delta W\right)$$
We tested a range of ranks from 4 to 16 and did not observe significant performance difference.
To reduce weight size and fine-tuning efficiency, we used r = 4 for the final large-scale fine-tuning.
The custom LoRA implementation is available on https://github.com/tanjimin/LoRA.
647
Predicting CAP-binding profiles
During prediction, the LoRA checkpoint corresponding to each CAP was loaded together with the
backbone weights. Since Chromnitron prediction operates on individual 8,192 bp windows, to

predict CAP-binding signals across the entire chromosome, we used a sliding window approach
with 5,120 bp step size to create overlap between adjacent predictions. Chromnitron's predictions
are robust to window shifts, but the predicted signal at the window edge can have lower quality
due to the absence of flanking chromatin context. To mitigate this effect, we retained only the
central 6,144 bp predicted profiles out of the 8,192 bp as valid predictions. With a 5,120 bp step
size and a 6,144 effective window size, adjacent predictions will have a 1,024 bp effective overlap.
To further ensure a high-quality transition between the overlapped regions between adjacent
predictions, we implemented a weighted average in the overlapping regions to give more weights
to bases that are closer to the center of their prediction windows. The weighted signal $\bar{S}$ is an
average of left signal S_l and right signal S_r, weighted by their relative base location i in the region:
$\bar{S} = \frac{i}{1024} \cdot S_l + \frac{1024-i}{1024} S_r$.
662
Prediction performance evaluation
All prediction performance evaluations were conducted on the leave-out chromosome 10. To
compare with experimentally measured ChIP-seq profiles, we first predicted CAP-binding profiles
across all five cell types on chromosome 10. For each CAP, we extracted profiles from both
prediction and ChIP-seq profiles on the chromosome 10p arm between 6 Mb and 30 Mb. We
evaluated model performance by calculating a set of metrics (detailed in the following section)
between prediction and experimental ChIP-seq profiles in the leave-out chromosome across all
cell types. For cell-type-specific prediction performance evaluation, we first calculated the
experimental data difference and the predicted CAP-binding difference between HepG2 and K562
cells, the two cell types with the highest number of overlapping CAP profiles. We then applied the
same set of evaluation metrics to compare the difference signal between the two groups.
674
For a comprehensive comparison, we computed in total eight statistics, including Pearson's
correlation (r), Spearman's correlation (ρ), the coefficient of determination (R squared), F1 score,
area under the receiver-operating characteristic curve (AUROC), and area under the precision-
recall curve (AUPR) with both target-centric and prediction-centric labelling. We repeated the
statistics over a grid of different bin sizes and top-percentile subset of the bins. Predicted or ChIP-
seq signals were first converted to numerical vectors, with missing values set to zero.
Subsequently, the vectors were mean-reduced into non-overlapping bins of 50, 250, 500 or 1000
bp. This multi-scale representation allows assessment of both fine-grained and coarse
regulatory patterns. The top-percentile bins measure how well the model can differentiate between
peaks and their relative enrichment level. For this, we set an exponential increment of percentile
using a list of 100% (1), 50% (1/2), 25% (1/4), 12.5% (1/8), …, 0.05% (1/2048). For AUROC and
AUPR calculation, we used the median value of the selected bins as threshold to separate positive
and negative labels. We also tested an alternative strategy to select most characteristic bins by
selecting both top and bottom percentile bins (Supplementary Fig. 6). The performance metrics
across methods and all bin sizes were in included in Supplementary File 2.
690
To categorize CAP-binding prediction performance, we focused on the Pearson's correlation (r)
as a simple metric to help interpret the prediction results. We categorized CAP-binding prediction

performance with both validation performance and cell-type-specific performance on the leave-
out chromosome (chr10) for all CAPs in cell types where training data are available. The validation
performance was defined as the highest performance across all cell types, while the cell-type-
specific performance was calculated as described above. To establish a relationship between
validation performance and cell-type-specific performance, we correlated the validation
performance against cell-type-specific performance and found a general positive correlation. The
cell-type-specific performance is a relatively more challenging metric due to additional
experimental noises across cell types. We selected 187 CAPs that are shared between HepG2 and
K562 and fitted a line between validation performance and cell-type-specific performance. After
custom evaluation of prediction performance, we assigned CAPs to different performance groups
based on both validation and cell-type-specific prediction performance metrics. The specific
performance cutoff was chosen based on the fitted curve between the two metrics (Supplementary
Fig. 4f, and summarized Supplementary File 1). We classified a CAP into high-performance group
if its prediction performance has either a validation performance > 0.6 or the corresponding cell-
type-specific performance. CAPs with either a validation performance > 0.4 or the corresponding
cell-type-specific performance were assigned in medium-performance group. The rest of CAPs
were categorized into low-performing group. Notably, although some CAPs were classified into
low-performance group by Pearson's correlation, their predicted binding profiles recapitulate
ChIP-seq peaks (Supplementary Fig. 5–6). Thus, the proposed CAP-performance categorization
should be regarded as a practical metric for interpreting prediction results, rather than a strict
measure of biological validity.
714
Global comparison of prediction to experimental data at CAP-binding regions
To evaluate prediction performance across different chromatin backgrounds, we compared CAP-
binding signals by aligning Chromnitron predictions with matched ChIP-seq and ATAC-seq
profiles centered in peak regions. For each CAP of a particular cell type, three one-dimensional
signal tracks were collected: (i) Chromnitron-predicted CAP-binding signal, (ii) experimental
ChIP-seq data (reference), and (iii) ATAC-seq data to represent chromatin accessibility.
Chromnitron-predicted CAP-binding data on chromosome 10 were generated and documented, as
described in the previous performance evaluation section. Peaks were independently identified in
prediction, ChIP-seq and ATAC-seq by thresholding the signal at 0.5 (half of very high peaks),
merging contiguous bins above this level, and discarding events narrower than 100 bp. The mean
signal within each retained interval was recorded to generate a peak set annotated by genomic
coordinates and intensity.
727
To visualize and directly compare prediction to the reference CAP-binding data, all peaks detected
in Chromnitron prediction were stratified into two groups according to chromatin accessibility
signal. Each predicted peak was labelled according to its overlap with ATAC-seq peaks, generating
an accessible set and an inaccessible set. For every peak in each group, we extracted a 2kb genomic
window centered at the highest score in each peak. Finally, we visualized the predicted CAP-
binding signal, ChIP-seq signal, and ATAC-seq signals as heatmaps to evaluate global prediction
performance across chromatin background.

Ablation study

We performed ablation studies to measure how the model performs under different constraints. This analysis includes performance evaluation across models trained without fine-tuning, and models trained with missing cell types or CAPs to test their generalizability across modalities in unseen contexts. We performed ablation studies on a random subset of chromosome 10 to reduce computation and storage consumption. Our preliminary calculation found that for one set of experiments, inferring 767 CAPs on chromosome 10 for five cell types could take two weeks and 6.5 TB of storage. Instead, we used Pearson's correlation coefficient calculated during the validation step embedded in the evaluation process to save computation and storage. This evaluation was performed on 500 random loci on chromosome 10 across all cell types and works as a proxy of the genome-wide prediction performance statistics. The performance statistics used for both the full Chromnitron model and ablated models were all calculated from validation step using the same metrics.

To evaluate the performance improvements of fine-tuning, we extracted performance metrics from the pretrained model at its last epoch (390) and metrics from fine-tuned models at their last fine-tuning epoch (≤ 10). The last fine-tuned epoch can change due to our early stopping strategy. Then, we compared the performance individually across cell types and CAPs.

For cell-type-specific prediction evaluation through ablation study, we kept data from HepG2 and K562 cells as the training set, and left out data from IMR-90, GM12878 and HCT116 cells as testing. Chromosome 10 region was again used as leave-out chromosome. This model was then pretrained (i.e. without fine-tuning) and compared with the full Chromnitron pretraining model in leave-out cell types. For CAP-specific evaluation, we randomly left out half of the CAPs (n=382) and kept the other half (n=385) in the training set. Then we extracted the performance metrics from the ablated model and compared with the full Chromnitron pretraining model on leave-out CAPs (again in chromosome 10 regions).

Benchmarking prediction performance

Although there was no directly comparable model that used the same multimodality framework, we benchmarked performance of Chromnitron against previous machine learning approaches that aimed for similar goals, including Enformer, Borzoi, AlphaGenome, maxATAC, and accessibility-weighted motif score as a baseline approximation[41,42]. We compared inference of CAP binding tracks from different models against the experimental data reference (Chrom2Vec processed ENCODE ChIP-seq profiles). We describe first the overall setup for different benchmarks and then the procedure for obtaining CAP-binding predictions from each model.

The *de novo* cell type performance benchmark was performed using a prototype model separately trained on HepG2 and K562 and evaluated on leave-out HCT116, IMR-90 and GM12878 cell types and at leave-out chromosome 10. The prototype model is a pretrained model and not fine-

tuned for each CAP. The ATAC-seq corresponding to each cell type is used as input for prototype
Chromnitron. Since sequence-based models do not accept cell-type-specific information as input
and do not generalize to new cell types, the most robust estimation is computing average prediction
profiles across cell types. For each CAP, we averaged prediction profiles across all cell lines except
for the cell type that is under evaluation. Since most of the region on the genome have rather
conserved binding across cell types, we selected cell-type-specific regions by performing pair-
wise differential analysis for 3 pairs between HCT116, IMR-90, and GM12878. The genome is
binned at 1 kb. The top 1% most differential bins based on signal log2 fold change was used as
regions for evaluation. We note that all selected regions are on Chromnitron leave-out
chromosome 10 for stringent evaluation, but many of which are in the training set of other
sequence-based models due to different data splitting strategies. We compared performance of
prototype Chromnitron against aggregated prediction from sequence-based models and
accessibility-weighted motif baseline on these cell-type-specific regions. Since each model has
different numbers of prediction target, we selected the joint CAP subset shared between
Chromnitron, AlphaGenome, Borzoi and Enformer for benchmarking purpose.

The overall evaluation with the full Chromnitron model is performed with a similarly setup on
HepG2 and K562. The model used in this evaluation is pretrained on all cell types and fine-tuned
for each CAP as described in the model training method section. The same prediction aggregation
is performed for all sequence-based models. The same region selection strategy is used to obtain
differential regions between HepG2 and K562 on Chromnitron leave-out chromosome 10. The
chromosome-wide evaluation has the same model prediction setup but is performed on the entire
chromosome 10.

Enformer: We used GRCh38 reference genome sequence as input to Enformer. Chromosome 10
was divided into non-overlapping 100 kb windows; for each window, 196,608 bp centered on the
window mid-point were retrieved (zero-padded at chromosome ends) and one-hot encoded. Each
sequence was forwarded through the PyTorch Enformer implementation (EleutherAI/enformer-
official-rough) which has similar performance as the TensorFlow implementation. The model
returned an 896-bin output tensor per window which was cached on disk. CHIP channels relevant
to HepG2 or K562 cells were identified from human ChIP-seq targets. For every requested TF
channel, the central 50,176 bp (±50,176 bp) of each window were extracted to avoid model
padding artefacts. The selections were written into an array representing chromosome 10 at 1 bp
resolution. Overlaps occurred only at window boundaries and were overwritten by later windows.
The final arrays were exported to BigWig files for visualization and comparison with other
approaches.

Borzoi: We used the same reference sequence and processed chromosome 10 in non-overlapping
523,264 bp windows. For each window, a 524,288 bp sequence centered on the window midpoint
was retrieved (zero-padded at chromosome ends if needed) and one-hot encoded via the
Baskerville DNA utilities. The sequence was fed to a single-replicate Borzoi model (SeqNN)
initialized from the published parameters/targets (params_pred.json, targets_human.txt) and

restored fold0 weights (model0_best.h5). The model returned per-track predictions for the full
window; raw output tensors were cached for downstream analysis. Processing advanced across
chromosome 10 in 523,264 bp steps; trailing partial windows were skipped to avoid padding
artefacts. Optional post-processing (not run in the main loop) converts selected tracks to dense 1
bp resolution arrays and exports BigWig files for visualization and downstream comparison.
824
AlphaGenome: We tiled chromosome 10 in non-overlapping 1,048,576 bp windows and skipped
trailing partial windows. For each window, we called the AlphaGenome DNA API
(dna_client.create) with organism HOMO_SAPIENS, OutputType.CHIP_TF only, and no
ontology terms. We requested predictions via API over the exact interval without padding. The
prediction server returned a CHIP-TF tensor (output.chip_tf.values) plus accompanying track
metadata. We cached the raw prediction arrays and metadata CSVs. The resulting prediction arrays
are merged and converted to individual bigwig files for further analysis.
832
MaxATAC: Raw ATAC-seq data were obtained from ENCODE as BAM files (HepG2:
ENCFF939JOT, K562: ENCFF804MXV). These files were down-sampled to 40M reads and
deduplicated to generate the BAM files required for maxATAC pipeline. MaxATAC pre-
processing ("maxatac prepare") was performed with "-c chr10" to build the ATAC coverage tracks
necessary its prediction. For every CAP that is included in both Chromnitron and maxATAC
model, we used the corresponding sub-model to predict the CAP binding profile as a BigWig file
using "maxatac predict".
840
Accessibility-weighted Motif (Acc. Motif) score: Non-redundant motif from Jeff Vierstra
(https://resources.altius.org/~jvierstra/projects/motif-clustering/releases/v1.0/) were filtered to
only retain human CAPs. Motif scanning results on chr10 were converted into a dense 1-bp signal
for each motif with the corresponding matching scores. A 20-bp moving average was applied to
the dense track. The final results were stored as BigWig files. To weight motif by chromatin
accessibility, we loaded ATAC-seq for all five cell types (HepG2, K562, GM12878, IMR-90,
HCT116). For each cell type, the smoothed motif track was multiplied elementwise by the
corresponding ATAC-seq signal and stored as BigWig files.
849
***In silico* DNA sequence mutagenesis**
To identify the impact of DNA sequence on the binding of a CAP, we focused on regions where
Chromnitron predicted intense CAP-binding signals. For each CAP-binding peak, we extended the
locus by ±3 kb and records its midpoint base location. We defined a 300-bp "mutation window"
(±150 bp) flanking the midpoint and systematically substituted every base within the window.
Subsequently, we performed *in silico* saturated DNA base mutagenesis of the 300 bp window, one
base at a time. For every position, we generated four sequences, each carrying a single A, C, G or
T substitution, and predicted the corresponding CAP-binding signal for each sequence. We
recorded the original sequence and its predicted CAP-binding signal as wild-type (WT) reference.
We then calculated the effect of every mutation as the difference between mutant and WT
predictions. The resulting inferred CAP-binding data at this window was stored as a 300 × 4 ×

8,192 tensor, with the three dimensions representing the number of mutation positions, number of
substitution nucleotides, and the window size of the CAP-binding prediction, respectively. The
signals represent the predicted CAP-binding intensities. To represent only the CAP-binding region,
the tensor was cropped from 8,192 bp to the central 1 kb and averaged, generating a 300×4 matrix
whose values represent CAP-binding intensities across the 300 bp mutated positions and 4
nucleotides. This matrix served as the nucleotide-resolution perturbation signals for downstream
motif analyses and interpretation. For visualization purposes, the impact motif was normalized by
subtracting the median CAP-binding intensities across the 4 nucleotides (i.e. the mean binding
intensities of the middle two nucleotides) at each position. In essence, the impact motif is computed
from a single locus, differing from the conventional motif calling which computes consensus
sequence from a large number of loci.
872
Cross-CAP motif discovery and clustering
To analyze motifs across all CAPs, we generated weight matrices (represented by median-
subtracted CAP-binding signals as described above) for all CAPs at their top binding loci. These
weight matrices were used as input for TF-MoDISco to generate motifs for each individual CAP[71].
For every CAP, the tool scans up to 20,000 high-scoring 200-bp windows ("seqlets", or high-
importance windows of the sequences, as referred in TF-MoDISco), clustering them by k-mer
similarity and reports positive and negative patterns (i.e. seqlets). After identifying relevant seqlets
for a CAP, we mapped their corresponding seqlets back to the genomic coordinates and applies a
strict quality control procedure: a pattern must contain at least one seqlet whose high (90th-
percentile) absolute weight difference is nontrivial (>0.1) after median subtraction. This procedure
excluded noisy results from the analyses. We then stored the remaining patterns for every CAP.
To pick the most representative seqlets for each CAP, we ranked them by (1) the number of
retained seqlets, (2) the overall motif intensity represented by the median difference value and (3)
the highest motif intensity represented by median 90th percentile difference. Patterns whose
seqlets overlapped more than ten distinct loci and repeated in different CAPs were discarded to
avoid trivial repeats.
889
Then, we built normalized weight matrices for cross-CAP comparison from the top ranked seqlets.
For each seqlet, we subtracted its per-position median and scaling by its mean absolute difference.
Each motif was flattened into a single vector (length by 4). Pairwise distances were computed as
the maximal cross-correlation between motif A and motif B or B's reverse complement. We
performed hierarchical clustering with Ward linkage based on the distance metrics, generating the
clustering heatmap for different motifs.
896
***In silico* chromatin accessibility perturbations**
We measured the changes of predicted CAP binding upon *in silico* scaling of chromatin
accessibility signal to examine Chromnitron's sensitivity to local chromatin states. We employed
a scaling approach through multiplying the ATAC-seq profiles by a defined factor (multiplier) to
alter accessibility intensities while preserving peak shape. Meanwhile, the size of the perturbation
window (i.e. number of bases) also matters in this *in silico* perturbation. Thus, we created an *in*

silico perturbation grid by combining multiplicative factor between 0 to 2, and with a perturbation
radius that ranged between 50 and 1,000 bp. For a given radius, the slice of the ATAC vector
centered on the locus was multiplied by the selected factor; the remainder of the signal was kept
the same. For each combination of multiplier and perturbation radius size values, we predicted the
corresponding CAP-binding signal for perturbation and calculated the binding differential by
subtracting the unperturbed condition, generating a data tensor of $40 \times 20 \times 8,192$ (multipliers $\times$
window-sizes $\times$ predicted profile). Similar to *in silico* DNA sequence mutagenesis, we averaged
the center 400 bp region of difference between perturbed prediction and wild-type prediction to
generate a 40×20 matrix that measures the accessibility perturbation effects.
912
In silico protein residue mutagenesis
To investigate how the model leverages protein embedding features in prediction, we performed
systematic mutagenesis directly on the input CAP residues without referring to particular protein
domain or functional data. Each perturbation operates on a 30-amino acid sliding window with
single-residue steps, replacing the window with random amino acids to generate the corresponding
mutant protein sequence. Mutated CAP sequence was feed into the same protein encoder (ESM2)
to generate the corresponding mutant embeddings, which were used in downstream prediction of
mutant binding profile and comparison with wild-type protein.
921
To ensure more robust comparison of protein mutation effects between the mutant and wild-type
CAP, we selected the top 200 Chromnitron-predicted binding peaks for each CAP in HepG2 cells,
filtered for ATAC-seq overlap, and equally extended to the flanking sequences up to 8,192 bp to
facilitate prediction. Chromnitron predictions of the corresponding CAP were generated for wild-
type and mutant embeddings across all selected loci. We then extracted maximum scores from the
center 400 bp region of each window (recorded as "wildtype_max" and "mutant_max",
respectively). CAP mutation impact scores were calculated as ($\frac{|\text{mutant\_max} - wildtype\_max|}{wildtype\_max}$). For
protein domain interpretation, we retrieved InterPro domain annotations and computed precision-
recall curves using domain boundaries as labels and mutation scores as predictions[72].
Domain enrichment was quantified using AUPR compared to random baselines.
932
Inferring CAP interactions through colocalization analysis
To investigate whether model prediction serves as a potential source to study protein-protein
interaction between CAPs, we followed our previous approach and applied linear non-Gaussian
acyclic model (LiNGAM) causal-discovery algorithm to a selection of high-quality CAPs
prediction across HepG2 ATAC-seq peaks[38,73]. We found that, as expected, the strongest
colocalizations were often between transcription factors that belong to the same family as they
usually share similar conserved DNA-binding domains. To systematically evaluate whether
colocalization calculated from experimental ChIP-seq or predicted CAP-binding data reconciles
with known physical protein-protein interactions, we benchmarked our causal discovery-based
interaction graph with Pearson's correlation-based interaction graph. This analysis was performed
on both experimental and predicted ChIP-seq data using confident interaction annotation from
STRING database (using scores over 700 as positive, and 400 to 700 as negative)[74]. The causal

discovery on predicted data yields the highest AUPR (0.426), followed by causal discovery on
observed data (0.372), while the correlation-based approaches achieved AUPR scores of 0.351 and
0.321 for predicted and experimental ChIP-seq data, respectively (Supplementary Fig. 11). The
lower performance when using ChIP-seq data could be a result from heterogeneous noise
distribution in the experimental data compared to the predicted results, leading to prediction of
only strongest interactions.
951

952   **Dosage correction of predicted CAP binding**
953   Here we describe the post-inference dosage-correction module to adjust predicted CAP binding by
954   the inferred availability of the corresponding CAP, using either CAP promoter accessibility or
955   CAP RNA abundance as a proxy for dosage.
956

For each CAP and cell type we obtain (i) promoter/TSS chromatin accessibility (log-transformed
read depth or signal at the TF's annotated TSS window) or (ii) bulk or pseudo-bulk RNA-seq
expression (log-transformed TPM/CPM or normalized counts). These scalars were matched to the
CAP predictions for the same cell type. These scalars can be used to either scale the prediction or
gate it. When used as a dosage multiplier, we multiply the log-transformed scalar to the
corresponding predicted CAP binding profile. When used as gating, we define a threshold
representing zero dosage of the CAP. Thresholds can be chosen empirically from the distribution
of accessibility or expression across TFs to balance false suppression of true binders against
removal of non-expressed CAPs. Predicted binding profile of CAPs that fall below the threshold
are set to zero.
967

968   Dosage correction is applied post inference and does not affect prediction. It can be toggled
969   between the scaling and gating modes depending on whether a graded or binary adjustment to TF
970   availability is desired. This module was not used in the cell-type-specific analyses to avoid
971   potential bias and turned off by default.
972

Single-cell differential ATAC-seq peak calling
To identify differentially accessible chromatin regions between pairs of cell types from single-cell
ATAC-seq data, we employed a computational workflow utilizing SCANPY[75]. For each pairwise
comparison, we first extracted the cells from the two specified populations the from complete
peak-by-cell matrix. Peaks with low signal, defined as those with fewer than 500 total fragment
counts across the selected cells, were removed from consideration. To account for technical
variations in sequencing depth, we normalized the counts for each cell to a total of 100,000,
followed by a log1p transformation to stabilize the variance. Subsequently, we identified highly
variable peaks between the two cell populations by using a dispersion-based method implemented
in SCANPY (scanpy.pp.highly_variable_genes), retaining peaks that exhibited a normalized
dispersion of at least 0.1. The genomic coordinates of these candidate regions were used as
candidate differentially accessible regions and exported for downstream analysis.
985

Predicting CAP-binding at single locus across T cell states

We used a public single-cell ATAC-seq data of T cell exhaustion collected from human patient
PBMC samples[58]. Four representative T cell stages (from effector to exhausted states) were
prepared to relate CAP binding changes to T cell states. Pseudo-bulk ATAC-seq data for each
stage was generated by sampling 10M reads from cells in the vicinity of selected center. The
pseudo-bulk ATAC-seq was then processed and normalized as described in the single cell data
processing method section. Differential accessible peaks of the corresponding cell populations
were called as described in the above section. These peak regions served as putative *cis*-regulatory
elements that might play important functions in T cell states. Next, for a given regulatory element
of interest, we predicted CAP-binding profiles at an element-centered 8,192 bp window across the
cell states. As single locus is highly susceptible to noise, we selected a set of inference CAPs using
a more stringent cutoff of 0.5 (Pearson's *r*) validation performance for this analysis. Predictions
for every T cell state were cropped to a 300 bp window around the CRE center. The binding signal
was normalized by median on the genome dimension to remove global scaling differences.
Trajectories were plotted as ridge plots that overlay the complete track profile per CAP. The mean
CAP-binding signals across trajectory were smoothed with Gaussian kernel and visualized with
line charts tracing binding dynamics across cell state transition trajectory.
1003
Genome-wide identification of regulatory CAPs in T cell exhaustion
The same T cell exhaustion scATAC-seq and annotation in single-locus-analysis were used for
this analysis. We first isolated reference pseudo-bulks for both effector T cells and exhausted T
cells according to the original cell type annotations[58]. To obtain a cleaner causal relationship
between CAPs and target genes, we focused on promoter regions as the putative regulatory
elements. We used GENCODE v45 transcript annotation (GFF3) and the hg38.chrom.sizes file
hosted by IGV to define promoter coordinates and chromosome boundaries[76]. Promoter regions
(± 500 bp around the transcription-start site, TSS) were extracted from the GENCODE annotation.
Genes on sex chromosomes or mitochondrial DNA were discarded. We also removed transcripts
not matching a whitelist of supported biotypes (protein-coding, lincRNA, miRNA, snRNA,
snoRNA, rRNA). The maximal and mean ATAC-seq accessibility signal were computed in both
T cell states for each promoter. Promoters with a cross-condition maximum accessibility ≤ 1 were
removed to exclude inactive loci. We then calculated log2 fold change (log2FC) of accessibility
for all the loci, and partitioned them into three categories: (1) up-regulated promoters, defined as
top 2,000 positive log2FC values, (2) down-regulated promoters defined as top 2,000 negative
log2FC values, and (3) stable promoters, defined as 100 promoters with the smallest absolute
log2FC. We also considered alternative TSS as separate promoters and implemented additional
constraints to merge overlapping TSS loci.
1022
Due to the high compute requirement for this task, Chromnitron prediction was sharded by cell
type and all CAPs, and executed in parallel on a GPU cluster. For this task, we used the full set of
CAPs since genome-wide analysis can better tolerate noisy predictions. After all shards completed,
we merged the predictions by iterating over chromosomes, stacking the individual CAP arrays to
generate genome_length × CAPs matrices. This matrix was rechunked to optimize downstream
biological analyses for characterizing putative regulator CAPs.

1029

Promoter-level average CAP-binding signals were extracted by recording the maximum predicted
signal in the 1 kb window centered at TSS, generating a three-dimensional tensor with dimension
(cell types × promoters × CAPs). Downstream analyses used the log-transformed values. A
promoter-specific differential CAP-binding score was calculated by subtracting the predicted
CAP-binding signals of the exhaustion group by that of the effector T cell group. To account for
basal CAP-specific variability, the difference matrix was z-standardized by subtracting the mean
and dividing by the standard deviation of the difference values measured at stable promoters. To
increase signal-to-noise ratio, mid-range values between the 10th and 90th percentiles were set to
zeros separately across loci and CAPs, thereby highlighting the most differential – potentially
functional – signals for informing downstream analysis. Next, we performed k-means clustering
on both the loci and CAP dimensions. Silhouette analysis over k-means clustering using k ranging
from 3 to 40 was used to derive optimal number of clusters. To identify important CAPs and loci
within each cluster, we ranked loci and CAPs by revaluating the average binding against the
opposing cluster assignments iteratively on both dimension until convergence.

1044

Global CAP and CRE dynamics characterization during Purkinje cell differentiation
Single-cell ATAC-seq datasets for Purkinje cell development were obtained from Mannens et al[62].
We adopted the original assignment of Purkinje cell differentiation trajectory encompassing seven
stages: progenitor cells (Prog), two neuroblast stages (NBL1, NBL2), and four Purkinje cell
maturation stages (Purk1 to Purk4). All datasets were processed using the scATAC-seq
preprocessing steps described earlier. Only autosomes were included for this analysis.

1051

To identity genomic regions of interest, we used multiple filtering approaches based on chromatin
accessibility dynamics across developmental stages. Initial candidate regions were generated from
cPeaks representing the repertoire of accessible regions across human cell types[67]. We extended
each peak center by 500 bp flanking region on each side to create 1 kb inference windows to
include the complete peak region. Then, we filtered these loci by classifying them into two
categories based on temporal accessibility patterns: differential and stable regions. Differential loci
were defined as regions exhibiting significant accessibility changes across developmental stages
(log2 fold-change larger than 0.1 compared with average accessibility), representing dynamic
regulatory elements. Stable loci have relatively minimal accessibility change, defined as having
absolute log2FC smaller than 0.1. Similar to the genome-wide regulator screening in T cell
exhaustion, these stable loci served as normalization controls for downstream analysis. We
selected the top 1,000 differential loci and top 100 most stable loci. To ensure non-redundant
genomic coverage, overlapping regions were merged. Regions with insufficient chromatin
accessibility across all conditions were excluded to prevent analysis of inactive or noisy genomic
elements. The final dataset comprised 1,073 high-confidence genomic loci distributed across
autosomes.

1068

This task was computationally intensive and was thus parallelized on a GPU cluster. To reduce
compute, we only keep a set of high-performance CAPs similar to the single locus T cell

exhaustion inference experiment. Individual CAP predictions were concatenated to construct a 3D
data tensor of developmental time × genomic loci × CAPs. For each genomic locus, we calculated
the maximum peak heights within 1 kb windows centered at peak summits. The resulting 3D tensor
was filtered for missing values, infinite numbers, and outliers. Data normalization was performed
using stable loci as internal controls to account for technical variation and global accessibility
changes across developmental stages. We performed z-score normalization using the mean and
standard deviation of differential patterns calculated from stable loci. Outlier detection and
removal were implemented using percentile-based clipping (1st and 99th percentiles) to prevent
extreme values from dominating downstream clustering analyses.
1080

Regulatory pattern discovery was performed using multiple complementary approaches to capture
different aspects of the temporal regulatory landscape. The 3D tensor was reshaped into matrices
emphasizing different analytical perspectives. There are three ways to transform the 3D tensor to
2D matrix for specific analyses, each enabling discovery of distinct regulatory relationships. First,
we can use CAP as features, reshaping the tensor to a matrix with dimension (time × loci) by CAPs
to identify loci whose activities change across time. Second, we can use loci as features to generate
the (time × CAPs) by loci matrix. This approach can represent how CAP binding profiles change
across time. Finally, using time as the feature, the reshaped matrix can tell how a binding event
specific to a CAP and a locus varies across time.
1090

We focused on the first analysis using CAP as features. To observe CRE activity shifts across time,
we performed hierarchical clustering of CREs on the first pseudo-time point. This approach
grouped CREs with similar regulatory dynamics. To further elucidate the relationship between
CRE and time, we applied k-means clustering using a k between 5 and 20 to identify regulatory
locus modules across all time points and CREs. The optimal cluster number was determined using
silhouette score. Clustering was performed on unit-normalized data using Euclidean distance
metrics, with multiple random initializations to ensure stable cluster assignments. For visualization
and interpretation, t-distributed Stochastic Neighbor Embedding (t-SNE) was used to generate 2D
embeddings of high-dimensional regulatory patterns. Each dot on the t-SNE embedding represents
a CRE at a specific time point and was colored by clustering results representing its identity.
1101

1102   To visualize transition of CREs between adjacent pseudo-time points in Purkinje cell development,
1103   we calculated a vector field where the start and end were defined by the t-SNE location of each
1104   locus at first and second time points. This velocity field represents how CREs change their
1105   regulatory identity between consecutive developmental stages. We applied Gaussian kernel
1106   smoothing to the vector field to reduce noise while preserving temporal trends.
1107

Cell culture and primary cell isolation
HEK293T cells were obtained from the American Type Culture Collection (ATCC). HEK293T
cells were maintained in Dulbecco's Modified Eagle Medium (DMEM; Gibco, 11965092)
supplemented with 10% fetal bovine serum (FBS; Gibco, A5670701) and 1% penicillin–
streptomycin (Pen/Strep; Gibco, 15140122). Human Peripheral blood mononuclear cells (PBMCs)

were isolated from healthy donor leukopaks (Charles River Laboratories Cell Solutions, 103051-
034) using Ficoll-Paque PLUS density gradient centrifugation (Cytiva, 17144002) according to
the manufacturer's protocol. Isolated PBMCs were cryopreserved in 90% human AB serum (MP
Biomedicals, 092938249) plus 10% dimethyl sulfoxide (Sigma-Aldrich, D4540) at a density of
30M cells per ml. For in vitro T cell assays, thawed PBMCs or purified CD8$^+$ T cells (isolated
using EasySep™ Human CD8+ T Cell Isolation Kit, STEMCELL Technologies, 17953) were
cultured in X-VIVO 20 medium (Lonza, BW04-448Q) supplemented with 5% heat-inactivated
human AB serum (MP Biomedicals, 092938249) and 10 ng/ml recombinant human IL-2
(PeproTech, 200-02-50UG). T cell cultures were maintained at 37 °C in a humidified incubator
with 5% CO_2.
1123
sgRNA library preparation
For each selected gene, five sgRNAs were designed to either knock out or repress gene expression
using the CRISPick online tool (Broad Institute). The complete list of sgRNA sequences is
provided in Supplementary File 3. A total of 370 DNA oligonucleotides corresponding to sgRNAs
were synthesized as pooled libraries (500 ng per library) by Twist Bioscience. For gene knockout
experiments, sgRNAs were cloned into the lentiviral vector pXPR_023 vector (Broad Institute,
also available through Addgene #52961), which encodes Cas9, a puromycin resistance gene, and
the sgRNA scaffold. For gene repression (CRISPRi), sgRNAs were cloned into pXPR_066 vector
(obtained via Broad Institute Genetic Perturbation Platform), which expresses KRAB-dCas9 under
the EFS promoter along with a puromycin resistance cassette and sgRNA scaffold.
Oligonucleotides were cloned into the respective backbone vectors using Golden Gate assembly
with BsmBI restriction enzyme (New England Biolabs, R0739S), following the protocol described
by the Feng Zhang lab (Broad Institute). Ligation products were transformed into NEB Stable
Competent *E. coli* (New England Biolabs, C3040H) and plated on LB-agar plates supplemented
with carbenicillin (100 μg/ml). Colonies were scraped from the plates and pooled, and plasmid
DNA was extracted using the ZymoPURE II Midiprep Kit (Zymo Research, D4201) and
quantified by NanoDrop for downstream lentivirus production.
1141
Lentivirus production
Lentivirus was produced in HEK293T cells (ATCC; passage number <20) seeded at 4×10^7 cells
per 150-mm tissue culture dish. Twenty-four hours after seeding, the culture medium was replaced
with fresh pre-warmed DMEM containing 10% FBS. Transfection was performed using
polyethyleneimine (PEI 40K; Polysciences, 24765) as follows: 13.3 μg of the sgRNA vector, 10 μg
of packaging plasmid psPAX2, and 6.7 μg of envelope plasmid pMD2.G were mixed in 400 μl of
Opti-MEM (Thermo Fisher Scientific, 31985070) (Tube A). Separately, 90 μl of PEI (1 mg/ml)
was diluted in 310 μl of Opti-MEM (Tube B). The two solutions were incubated separately for
5 min at room temperature, combined, and incubated for an additional 15 min before being added
dropwise to the HEK293T cells. Seventy-two hours post-transfection, the culture supernatant was
harvested and centrifuged at 3,000g for 15 min to remove cellular debris. Lentiviral particles were
precipitated by adding polyethylene glycol 8000 (PEG8000; Sigma-Aldrich, P2139-500G) to a
final concentration of 8% (w/v) and incubating the mixture overnight at 4 °C. The precipitated

lentiviral particles were pelleted by centrifugation at 1,500g for 30 min and resuspended in 500 μl
of X-VIVO 20 medium (Lonza, BW04-448Q). Aliquots were stored at −80 °C until use.
1157
T cell stimulation assays
Cryopreserved human CD8+ T cells were thawed and resuspended in T cell culture medium (X-
VIVO 20 supplemented with 5% human AB serum and 10 ng ml^{-1} recombinant human IL-2) at a
density of 1×10^6 cells per ml in 12-well plates. Cells were stimulated by the addition of Human
CD3/CD28 T cell Activation Beads (BioLegend, 422603) at a 1:1 bead-to-cell ratio and incubated
for 48 h at 37 °C with 5% CO_2. Following activation, beads were removed using a magnetic
separation rack according to the manufacturer's instructions. Cells were then washed twice with
fresh T cell medium and split into two different stimulation conditions. For the acute stimulation
group, cells were maintained in T cell medium alone. For the chronic stimulation group with
antibody cocktail, cells were cultured in T cell medium supplemented with ImmunoCult Human
CD3/CD28 T Cell Activator (Stem Cell Technologies, 10971), following the manufacturer's
protocol. For the chronic stimulation group with anti-CD3 antibody (eBioscience, 16-0037-81),
the antibody was coated at 12-well plate with 5 μg/ml concentration. Cells in the chronic condition
were passaged every 48 h, with medium replacement and re-activate for three consecutive
stimulation rounds. At the end point, cells were harvested and divided into two groups for either
surface staining of exhaustion markers or intracellular cytokine analysis by flow cytometry.
1174
Lentiviral transduction and titration in CD8$^+$ T cells
To determine optimal lentiviral transduction conditions for primary human CD8$^+$ T cells, cells
were activated for 48 h using CD3/CD28 beads as described above. After bead removal, cells were
plated in 96-well plates at 2×10^5 cells per well in 200 μl of T cell medium (X-VIVO 20
supplemented with 5% human AB serum and 10 ng/ml IL-2). Serial dilutions of lentivirus
encoding puromycin resistance gene were added to the wells in the presence of 8 μg/ml polybrene
(Millipore, TR-1003-G). Plates were centrifuged at 1,500g for 90 min at 37 °C, the medium was
changed to T cell medium after additional 4 h incubation, the cell was continued culture at 37 °C,
5% CO2. At 48 h post-transduction, cells were diluted to 1×10^5 cells per ml and cultured in the
presence of 1 μg/ml puromycin for 48 h cell viability was assessed by counting viable cells using
trypan blue. Viral titer and multiplicity of infection (MOI) were calculated based on the fraction
of puromycin-resistant cells relative to total input.
1187
CRISPR screening in primary CD8$^+$ T cells
Human CD8$^+$ T cells were isolated from PBMCs by negative selection and activated with
CD3/CD28 beads for 24 h After bead removal, cells were transduced with pooled lentiviral sgRNA
libraries at an MOI of 0.3 in the presence of 8 μg/ml polybrene. Cells were spinoculated at 1000g
and 32 °C for 90 min. 6 h after spin infection, the cell were resuspended to T cell medium for
overnight culture. 24 hours after spin infection, the cell were treated with puromycin (1.5 μg/ml)
to select the transduced cells. Following selection, cells were washed and resuspended into two
treatment conditions corresponding to acute and chronic stimulation, as described previously. Cells
were maintained under these conditions for 10 days, with medium refreshed and IL-2 re-

supplemented every 2 days. On the final day of the screen, genomic DNA was harvested from each
condition using the QIAamp DNA Blood Maxi Kit (QIAGEN, 51192) following the
manufacturer's instructions. Genomic DNA was quantified using Qubit dsDNA BR Assay Kit
(Thermo Fisher Scientific, Q32850), and sgRNA sequences were PCR-amplified from extracted
DNA using a one-step amplification protocol compatible with next-generation sequencing, as
previously described by Broad Institute Genetic Perturbation Platform ("sgRNA/shRNA/ORF
PCR for Illumina Sequencing" linked through
https://portals.broadinstitute.org/gpp/public/resources/protocols), PCR primers were provided in
Supplementary File 4. PCR products were purified using AMPure XP beads (Beckman Coulter,
A63880). Libraries were quantified, normalized, and sequenced on an Illumina NovaSeq X
platform using a 25B flow cell, generating paired-end 150 bp reads. Raw sequencing data were
demultiplexed using bcl2fastq (Illumina) and further processed as described below.
1209
CRISPR screening analysis
Sequencing reads from CRISPR screens were first demultiplexed and trimmed to extract sgRNA
sequences. Reads were aligned to the sgRNA reference library using the MAGeCK count function
in the MAGeCK software package with default parameters[77]. Only sgRNAs with perfect matches
to the reference library were retained for downstream analysis. Read count matrices were
generated for all samples and normalized using median normalization. Differential sgRNA
abundance between experimental conditions (Chronic vs. Acute stimulation) was analyzed using
the MAGeCK test module, which applies a negative binomial model to assess gene-level
enrichment or depletion. For each gene, MAGeCK aggregates sgRNA-level statistics to compute
a robust gene-level score and associated p value using a robust ranking aggregation (RRA)
algorithm. Genes were ranked by their adjusted p values (Benjamini–Hochberg FDR correction),
and significance thresholds were defined as indicated in the figure legends. All analyses were
performed using MAGeCK with default settings unless otherwise noted.
1223
RNP electroporation in primary human CD8+ T cells
To perform Cas9 RNP-mediated gene knockout, two synthetic sgRNAs targeting the same gene
were used per condition (sequence available in Supplementary File 4). Each sgRNA was
resuspended to a final concentration of 100 μM in RNase-free water and stored at −80 °C until use.
Cas9 protein (EnGen® SpyCas9 NLS, 20 μM stock, New England Biolabs, M0646T) was used
for complex formation. For each reaction, 60 pmol of Cas9 protein was complexed with 180 pmol
total sgRNA by mixing 3 μl of Cas9 protein, 0.9 μl of each sgRNA, and 0.4 μl of 10× NEBuffer
3.1 in a total volume of 5.2 μl. The mixture was incubated at room temperature for 15 min to allow
RNP formation. Primary CD8+ T cells were activated for 48 h with CD3/CD28 beads as described
above, then washed once with 1× PBS and resuspended in Lonza P3 electroporation buffer at a
density of 1×10^6 cells per 20 μl. The entire 5.2 μl RNP mixture was added to 20 μl of cell
suspension, mixed gently, and transferred into a well of a 16-well Nucleocuvette™ strip (Lonza).
Air bubbles were removed by tapping the strip on the benchtop to prevent disruption of the electric
pulse. Electroporation was performed using the Lonza 4D-Nucleofector system with the program
EH115, optimized for primary human T cells. Immediately after electroporation, 80 μl of pre-

warmed T cell medium (X-VIVO 20 supplemented with 5% human AB serum and 10 ng ml^{-1} IL-
2) was added to each well. The cells were incubated in the cuvette strip at 37 °C for 10 min to
enhance recovery before being gently transferred to pre-warmed 24-well plates.
1242
Flow cytometry staining and analysis
For intracellular cytokine staining, T cells were stimulated for 3 h at 37 °C with Cell Activation
Cocktail containing Brefeldin A (BioLegend, 423303). Cells were washed with Cell Staining
Buffer (BioLegend, 420201), fixed and permeabilized with Fixation/Permeabilization solution
(BD Biosciences, AB_2869008) for 20 min at 4 °C, and washed twice with 1× Perm/Wash buffer.
After Fc receptor blocking (BioLegend, 422302), cells were stained for 2 h at 4 °C in the dark with
fluorophore-conjugated antibodies in staining buffer (BioLegend, 420201) along with NIR
Live/Dead dye (1:10000 dilution, ThermoFisher, L34975). Antibodies included APC anti-human
CD8 Antibody (BioLegend, 344722), Brilliant Violet 785™ anti-human IFN-γ Antibody
(BioLegend, 502541), Brilliant Violet 605™ anti-human TNF-α Antibody (BioLegend, 502935),
and Brilliant Violet 421™ anti-human IL-2 Antibody (BioLegend, 500327).
1254
For surface marker staining, cells were blocked with FC receptor blocking reagent (BioLegend,
422302), then stained with antibodies and NIR Live/Dead dye (1:10000 dilution, ThermoFisher,
L34975) at 4 °C for 1 h. Antibodies included CD8 Antibody (BioLegend, 344722), Brilliant Violet
785™ anti-human TIGIT (VSTM3) Antibody (BioLegend, 372735), Brilliant Violet 605™ anti-
human CD366 (Tim-3) Antibody (BioLegend, 345017), Brilliant Violet 510™ anti-human CD223
(LAG-3) Antibody (BioLegend, 369317), and Brilliant Violet 421™ anti-human CD279 (PD-1)
Antibody (BioLegend, 329920). Flow cytometry was performed on a CytoFLEX LE (Beckman
Coulter, six-laser configuration). Gating strategy included sequential gating on FSC-A/SSC-A
(lymphocytes), FSC-A/FSC-H (single cells), and exclusion of dead cells using viability dye.
Fluorescence compensation was calculated using single-stained compensation beads (BioLegend).
Data were initially analyzed with FlowJo v10.8, with gates being defined using isotype controls,
followed by custom reproduction of the analysis for visualization purpose only.
1267
Surveyor assay for genome editing efficiency
To evaluate genome editing efficiency following RNP electroporation, genomic DNA was
extracted from T cells 72 h post-nucleofection using the Monarch® Genomic DNA Purification
Kit (New England Biolabs, T3010S), following the manufacturer's instructions. Genomic regions
flanking the Cas9 cleavage site were PCR-amplified using 2× Phanta Max Master Mix (Vazyme,
P525-02) in a 50 μl reaction volume (PCR primers provided in Supplementary File 4). PCR
products were verified by agarose gel electrophoresis and purified using the Zymo DNA Clean &
Concentrator Kit (Zymo Research, D4014), with final elution in 30 μl of nuclease-free water. To
generate heteroduplex DNA for mismatch detection, 200 ng of purified PCR product was
combined with 2 μl of 10× NEBuffer 2 (NEB) and nuclease-free water to a total volume of 19 μl.
The mixture was denatured and reannealed in a thermocycler using the following program: 95 °C
for 5 min; ramp at −2 °C/s to 85 °C; then ramp at −0.1 °C/s to 25 °C; and held at 4 °C. After
annealing, 1 μl of T7 Endonuclease I (NEB, M0302) was added to each reaction and incubated at

37 °C for 60 min. Reactions were terminated by adding 1.5 μl of 0.25 M EDTA and analyzed by
electrophoresis on a 1% agarose gel to visualize cleavage products.
1283

mRNA knockdown analysis by quantitative PCR
To assess transcript-level gene knockdown following RNP electroporation, edited T cells were
collected at the end point for RNA extraction. Total RNA was isolated using a RNeasy Mini Kit
(Qiagen, 74104), and RNA concentration was quantified by nanodrop. Reverse transcription was
performed using the HiScript® III 1st Strand cDNA Synthesis Kit (+gDNA wiper) (Vazyme,
R312-01) according to the manufacturer's instructions. Gene-specific primers targeting the
transcript of interest were used during amplification (Supplementary File 4). Quantitative PCR
was performed using PowerUp™ SYBR® Green Master Mix (Thermo Fisher Scientific, A25742)
on a CFX96 Real-Time PCR Detection System (Bio-Rad). ACTB (β-actin) was used as the internal
control gene for normalization. Each sample was run in triplicates, and specificity of amplification
was confirmed by melt-curve analysis. Relative gene expression was quantified using the ΔΔCt
method, and results were reported as fold change compared to the control condition.
1296

RNA-seq analyses
Total RNA was extracted on the final day from ZNF766 knockout and NTC control groups using
the RNeasy Mini Kit (Qiagen, 74004). Two biological replicates were collected per condition.
RNA-seq libraries were generated with the NEBNext Ultra II Directional RNA Library Prep Kit
(NEB, E7765L) following the manufacturer's poly(A) mRNA workflow, including enrichment
with the NEBNext Poly(A) mRNA Magnetic Isolation Module (NEB, E7490L). Raw RNA-seq
reads were mapped to the human reference genome (GRCh38/hg38) using STAR (v2.7.2a) with
default settings. Strand-specific read counts were aggregated into a gene-level count matrix for
subsequent analyses. Transcriptomic variation across samples was evaluated by principal
component analysis (PCA) based on the 500 most variable genes. Differential expression analysis
was conducted using DESeq2 (v1.40.2), and genes with an absolute log2 fold change greater than
0.5 and an adjusted P-value below 0.05 were defined as significantly differentially expressed.
Functional enrichment analyses were performed using clusterProfiler (v4.10.0), together with
org.Hs.eg.db (v3.18.0), enrichplot (v1.22.0), and DOSE (v3.28.2). Significant genes were
converted to Entrez Gene identifiers and assessed for enrichment in Gene Ontology biological
processes and KEGG pathways. Enrichment results were visualized as dot plots (GO) and bar plots
(KEGG).
1314

Data Availability

All ATAC-seq, ChIP-seq used in this study are public data collected from the ENCODE data portal (https://www.encodeproject.org). T cell scATAC-seq data is obtained from Gene Expression Omnibus (GEO) under accession GSE129785. Purkinje neuron developmental scATAC-seq data is obtained from https://github.com/linnarsson-lab/fetal_brain_multiomics. CRISPR screen data is available on GEO under accession number GSE305611. RNA-seq data on ZNF766 KO is available on GEO under accession number GSE305610.

Code Availability

The code for Chromnitron is available at https://github.com/tanjimin/Chromnitron.

Competing Interests

J.T., X.F., X.L., A.T., and B.X. are inventors on provisional patents covering the models and tools reported herein. A.T. is a scientific advisor to Intelligencia AI. J.T. is a consultant for Flagship Pioneering. J.D.B is a Founder and Director of CDI Labs, Inc., a Founder of and consultant to Opentrons LabWorks/Neochromosome, Inc, and serves or served on the Scientific Advisory Board of CZ Biohub New York, LLC, Logomix, Inc., Rome Therapeutics, Inc., SeaHub, Seattle, WA, Tessera Therapeutics, Inc., and the Wyss Institute. All these services are unrelated to this work.

Acknowledgement

This work is supported by NIH grants DP5OD033430 and R03DE033371 (to B.X.), R01CA252239 and P01CA229086 (to A.T.). X.F. and R.R. are supported by R35CA253126 and P30CA013696. J.T. was supported by the Judith & Stewart Colton Center for Autoimmunity at NYU Langone and Arthritis National Research Foundation Grant Award (No. 1471427). X.L. is a Fellow of The Jane Coffin Childs Fund for Medical Research. B.X. is a Junior Fellow of the Harvard Society of Fellows. We thank the High Performance Computing Core Facility at NYU Grossman School of Medicine and the Broad Institute for the support of computation. We thank the Broad Genetic Perturbation Platform and Broad Clinical Labs for the support of genetic and genomic experiments. We thank Sonya Craig from scientific editing team at NYU Langone for editing the manuscript. We also acknowledge Milk Bar, where X.F. and J.T. enjoyed coffee and met weekly for over two years to work on Chromnitron. We thank Huiyuan Zhang, Chenzhen Zhang, Hanqing Liu, Lloyd Bod, Jason D. Buenrostro, Eric S. Lander, Anshul Kundaje, Romanos S. Pistofidis, Liz Gaskell, Wenke Liu, Jose U. Scher, and the members of the Xia lab and Tsirigos lab for their valuable suggestions and discussion. Lastly, we are grateful to the ENCODE consortium for generating, organizing, and sharing standardized genomics data that enabled this study.

Author Contribution

B.X. and J.T. conceived the project. J.T., B.X. and A.T. designed the computational and experimental components and interpreted the results, with inputs from X.F. and X.L.. J.T., F.X. and S.M. designed, implemented, and optimized the neural network. J.T. and X.F. built the preprocessing platform. X.F. performed single cell data preprocessing and protein feature analysis.

References

1. Arendt, D. *et al.* The origin and evolution of cell types. *Nat. Rev. Genet.* **17**, 744–757 (2016).

2. Xia, B. & Yanai, I. A periodic table of cell types. *Dev. Camb. Engl.* **146**, dev169854 (2019).

3. Gurdon, J. B. Cell Fate Determination by Transcription Factors. *Curr. Top. Dev. Biol.* **116**, 445–454 (2016).

4. Chen, X. *et al.* Integration of external signaling pathways with the core transcriptional network in embryonic stem cells. *Cell* **133**, 1106–1117 (2008).

5. Boyer, L. A. *et al.* Core transcriptional regulatory circuitry in human embryonic stem cells. *Cell* **122**, 947–956 (2005).

6. Gerstein, M. B. *et al.* Architecture of the human regulatory network derived from ENCODE data. *Nature* **489**, 91–100 (2012).

7. Li, Y. E. *et al.* A comparative atlas of single-cell chromatin accessibility in the human brain. *Science* **382**, eadf7044 (2023).

8. Lander, E. S. *et al.* Initial sequencing and analysis of the human genome. *Nature* **409**, 860–921 (2001).

9. Lambert, S. A. *et al.* The Human Transcription Factors. *Cell* **172**, 650–665 (2018).

10. Vaquerizas, J. M., Kummerfeld, S. K., Teichmann, S. A. & Luscombe, N. M. A census of human transcription factors: function, expression and evolution. *Nat. Rev. Genet.* **10**, 252–263 (2009).

11. Weirauch, M. T. *et al.* Determination and inference of eukaryotic transcription factor sequence specificity. *Cell* **158**, 1431–1443 (2014).

12. Jolma, A. *et al.* DNA-binding specificities of human transcription factors. *Cell* **152**, 327–339 (2013).

13.   Vierstra, J. *et al.* Global reference mapping of human transcription factor footprints.

Nature **583**, 729–736 (2020).

14.   Murre, C. Helix-loop-helix proteins and the advent of cellular diversity: 30 years of

discovery. *Genes Dev.* **33**, 6–25 (2019).

15.   Doughty, B. R. *et al.* Single-molecule states link transcription factor binding to gene

expression. *Nature* **636**, 745–754 (2024).

16.   Durdu, S. *et al.* Chromatin-dependent motif syntax defines differentiation trajectories.

Mol. Cell **85**, 2900-2918.e16 (2025).

17.   Kim, S. & Wysocka, J. Deciphering the multi-scale, quantitative cis-regulatory code.

Mol. Cell **83**, 373–392 (2023).

18.   Kim, S. *et al.* DNA-guided transcription factor cooperativity shapes face and limb

mesenchyme. *Cell* **187**, 692-711.e26 (2024).

19.   Statello, L., Guo, C.-J., Chen, L.-L. & Huarte, M. Gene regulation by long non-coding

RNAs and its biological functions. *Nat. Rev. Mol. Cell Biol.* **22**, 96–118 (2021).

20.   Smith, E. & Shilatifard, A. The chromatin signaling pathway: diverse mechanisms of

recruitment of histone-modifying enzymes and varied biological outcomes. *Mol. Cell* **40**, 689–

701 (2010).

21.   Moore, J. E. *et al.* An expanded registry of candidate cis-regulatory elements. *Nature*

https://doi.org/10.1038/s41586-025-09909-9 (2026) doi:10.1038/s41586-025-09909-9.

22.   Yosef, N. *et al.* Dynamic regulatory network controlling TH17 cell differentiation.

Nature **496**, 461–468 (2013).

23.   Johnson, D. S., Mortazavi, A., Myers, R. M. & Wold, B. Genome-wide mapping of in

vivo protein-DNA interactions. *Science* **316**, 1497–1502 (2007).

24.   Tsankov, A. M. *et al.* Transcription factor binding dynamics during human ES cell

differentiation. *Nature* **518**, 344–349 (2015).

25.   Park, P. J. ChIP-seq: advantages and challenges of a maturing technology. *Nat. Rev.*

Genet. **10**, 669–680 (2009).

26.   Skene, P. J. & Henikoff, S. An efficient targeted nuclease strategy for high-resolution

mapping of DNA binding sites. *eLife* **6**, e21856 (2017).

27.   Partridge, E. C. *et al.* Occupancy maps of 208 chromatin-associated proteins in one

human cell type. *Nature* **583**, 720–728 (2020).

28.   Hammal, F., de Langen, P., Bergon, A., Lopez, F. & Ballester, B. ReMap 2022: a

database of Human, Mouse, Drosophila and Arabidopsis regulatory regions from an

integrative analysis of DNA-binding sequencing experiments. *Nucleic Acids Res.* **50**, D316–

D325 (2022).

29.   Moyers, B. A. *et al.* Characterization of human transcription factor function and patterns

of gene regulation in HepG2 cells. *Genome Res.* **33**, 1879–1892 (2023).

30.   Wang, J. *et al.* Sequence features and chromatin structure around the genomic regions

bound by 119 human transcription factors. *Genome Res.* **22**, 1798–1812 (2012).

31.   Avsec, Ž. *et al.* Base-resolution models of transcription-factor binding reveal soft motif

syntax. *Nat. Genet.* **53**, 354–366 (2021).

32.   Linder, J., Srivastava, D., Yuan, H., Agarwal, V. & Kelley, D. R. Predicting RNA-seq

coverage from DNA sequence as a unifying model of gene regulation. *Nat. Genet.* **57**, 949–

961 (2025).

33.   Avsec, Ž. *et al.* AlphaGenome: advancing regulatory variant effect prediction with a

unified DNA sequence model.

34.   Avsec, Ž. *et al.* Base-resolution models of transcription-factor binding reveal soft motif

syntax. *Nat. Genet.* **53**, 354–366 (2021).

35.   Lin, Z. *et al.* Evolutionary-scale prediction of atomic-level protein structure with a

language model.

36.   Abramson, J. *et al.* Accurate structure prediction of biomolecular interactions with

AlphaFold 3. *Nature* **630**, 493–500 (2024).

37.   Nguyen, E. *et al.* Sequence modeling and design from molecular to genome scale with

Evo. *Science* **386**, eado9336 (2024).

38.   Fu, X. *et al.* A foundation model of transcription across human cell types. *Nature*

https://doi.org/10.1038/s41586-024-08391-z (2025) doi:10.1038/s41586-024-08391-z.

39.   Hayes, T. *et al.* Simulating 500 million years of evolution with a language model.

Preprint at https://doi.org/10.1101/2024.07.01.600583 (2024).

40.   Theodoris, C. V. *et al.* Transfer learning enables predictions in network biology. *Nature*

618, 616–624 (2023).

41.   Cazares, T. A. *et al.* maxATAC: Genome-scale transcription-factor binding prediction

from ATAC-seq with deep neural networks. *PLOS Comput. Biol.* **19**, e1010863 (2023).

42.   Avsec, Ž. *et al.* Effective gene expression prediction from sequence by integrating long-

range interactions. *Nat. Methods* **18**, 1196–1203 (2021).

43.   Hu, E. J. *et al.* LoRA: Low-Rank Adaptation of Large Language Models. Preprint at

https://doi.org/10.48550/arXiv.2106.09685 (2021).

44.   Bravo González-Blas, C. *et al.* SCENIC+: single-cell multiomic inference of enhancers

and gene regulatory networks. *Nat. Methods* **20**, 1355–1367 (2023).

45.   Hu, Y. *et al.* Multiscale footprints reveal the organization of cis-regulatory elements.

Nature **638**, 779–786 (2025).

46.   Barral, A. & Zaret, K. S. Pioneer factors: roles and their regulation in development.

Trends Genet. **40**, 134–148 (2024).

47.   Balsalobre, A. & Drouin, J. Pioneer factors as master regulators of the epigenome and

cell fate. *Nat. Rev. Mol. Cell Biol.* **23**, 449–464 (2022).

48.   Stoltenburg, R., Reinemann, C. & Strehlitz, B. SELEX—A (r)evolutionary method to

generate high-affinity nucleic acid ligands. *Biomol. Eng.* **24**, 381–403 (2007).

49.   Ren, B. *et al.* Genome-Wide Location and Function of DNA Binding Proteins. *Science*

290, 2306–2309 (2000).

50.   Khan, O. *et al.* TOX transcriptionally and epigenetically programs CD8+ T cell

exhaustion. *Nature* **571**, 211–218 (2019).

51.   Seo, H. *et al.* TOX and TOX2 transcription factors cooperate with NR4A transcription

factors to impose CD8^{+} T cell exhaustion. *Proc. Natl. Acad. Sci.* **116**, 12410–

12415 (2019).

52.   Yao, C. *et al.* Single-cell RNA-seq reveals TOX as a key regulator of CD8+ T cell

persistence in chronic infection. *Nat. Immunol.* **20**, 890–901 (2019).

53.   Mann, T. H. & Kaech, S. M. Tick-TOX, it's time for T cell exhaustion. *Nat. Immunol.*

20, 1092–1094 (2019).

54.   Scott, A. C. *et al.* TOX is a critical regulator of tumour-specific T cell differentiation.

Nature **571**, 270–274 (2019).

55.   Alfei, F. *et al.* TOX reinforces the phenotype and longevity of exhausted T cells in

chronic viral infection. *Nature* **571**, 265–269 (2019).

56.   Hashimoto, M. *et al.* CD8 T Cell Exhaustion in Chronic Infection and Cancer:

Opportunities for Interventions. *Annu. Rev. Med.* **69**, 301–318 (2018).

57.   Baessler, A. & Vignali, D. A. A. T Cell Exhaustion. *Annu. Rev. Immunol.* **42**, 179–206

(2024).

58.   Satpathy, A. T. *et al.* Massively parallel single-cell chromatin landscapes of human

immune cell development and intratumoral T cell exhaustion. *Nat. Biotechnol.* **37**, 925–936

(2019).

59.   Jenkins, E., Whitehead, T., Fellermeyer, M., Davis, S. J. & Sharma, S. The current state

and future of T cell exhaustion research. *Oxf. Open Immunol.* **4**, iqad006 (2023).

60.   Tay, T. *et al.* Degradation of IKZF1 prevents epigenetic progression of T cell exhaustion

in an antigen-specific assay. *Cell Rep. Med.* **5**, 101804 (2024).

61.   Belk, J. A. *et al.* Genome-wide CRISPR screens of T cell exhaustion identify chromatin

remodeling factors that limit T cell persistence. *Cancer Cell* **40**, 768-786.e7 (2022).

62.   Mannens, C. C. A. *et al.* Chromatin accessibility during human first-trimester

neurodevelopment. *Nature* https://doi.org/10.1038/s41586-024-07234-1 (2024)

doi:10.1038/s41586-024-07234-1.

63.   Pampari, A. *et al.* ChromBPNet: bias factorized, base-resolution deep learning models of

chromatin accessibility reveal cis-regulatory sequence syntax, transcription factor footprints

and regulatory variants. Preprint at https://doi.org/10.1101/2024.12.25.630221 (2024).

64.   Chen, K. M., Wong, A. K., Troyanskaya, O. G. & Zhou, J. A sequence-based global map

of regulatory activity for deciphering human genetics. *Nat. Genet.* **54**, 940–949 (2022).

65.   Hendrycks, D. & Gimpel, K. Gaussian Error Linear Units (GELUs). Preprint at

https://doi.org/10.48550/arXiv.1606.08415 (2023).

66.   Vaswani, A. *et al.* Attention Is All You Need. *ArXiv170603762 Cs*

http://arxiv.org/abs/1706.03762 (2017).

67.   Meng, Q. *et al.* A generic reference defined by consensus peaks for scATAC-seq data

analysis. Preprint at https://doi.org/10.1101/2023.05.30.542889 (2023).

68.   Amemiya, H. M., Kundaje, A. & Boyle, A. P. The ENCODE Blacklist: Identification of

Problematic Regions of the Genome. *Sci. Rep.* **9**, 9354 (2019).

69.   Kingma, D. P. & Ba, J. Adam: A Method for Stochastic Optimization. Preprint at

https://doi.org/10.48550/arXiv.1412.6980 (2017).

70.   Hu, E. J. *et al.* LoRA: Low-Rank Adaptation of Large Language Models. Preprint at

https://doi.org/10.48550/arXiv.2106.09685 (2021).

71.   Shrikumar, A. *et al.* Technical Note on Transcription Factor Motif Discovery from

Importance Scores (TF-MoDISco) version 0.5.6.5. Preprint at

https://doi.org/10.48550/arXiv.1811.00416 (2020).

72.   Blum, M. *et al.* InterPro: the protein sequence classification resource in 2025. *Nucleic*

Acids Res. **53**, D444–D456 (2025).

73.   Ikeuchi, T., Ide, M., Zeng, Y., Maeda, T. N. & Shimizu, S. Python package for causal

discovery based on LiNGAM.

74.   Szklarczyk, D. *et al.* The STRING database in 2023: protein–protein association

networks and functional enrichment analyses for any sequenced genome of interest. *Nucleic*

Acids Res. **51**, D638–D646 (2023).

75.   Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression

data analysis. *Genome Biol.* **19**, 15 (2018).

76.   Robinson, J. T. *et al.* Integrative genomics viewer. *Nat. Biotechnol.* **29**, 24–26 (2011).

77.   Li, W. *et al.* MAGeCK enables robust identification of essential genes from genome-

scale CRISPR/Cas9 knockout screens. *Genome Biol.* **15**, 554 (2014).

1523

1524 **Supplementary Figures**

1525

1526

Supplementary Fig. 1: Chromnitron model architecture.

Left, Overview of model structure, showing modality-specific encoders for DNA sequence,

chromatin accessibility, and protein sequence, followed by multimodal fusion, transformer trunk,

and a CAP-specific decoder that outputs base-resolution binding profiles.

Right, Detailed layer configurations for the convolution encoders, transformer attention blocks,

and decoder modules. Convolutional blocks include Group Normalization, Gaussian Error Linear

Unit (GELU) activation, and residual connections with strided down-scaling or transposed up-
scaling layers as indicated. Transformer layers use a pre-layer-normalized (Pre-LN) architecture
with multi-head self-attention and feedforward sublayers.
Dimensions of inputs, intermediate representations, and outputs are indicated in brackets as (batch
size, length, channels).
1538

1539
Supplementary Fig. 2: Training data composition and the Chrom2Vec preprocessing
platform.
a. Distribution of curated CAP-binding ChIP-seq datasets across five human cell lines.
b. Pairwise overlap of CAP-binding datasets between cell lines, displayed as a heatmap indicating
the number of shared CAPs. Diagonal entries are removed.

c. Workflow of Chrom2Vec platform for processing training datasets. The workflow integrates
job scheduling, data storage, execution tracking, containerized bioinformatic pipelines, and
automated normalization and compression steps. Full details about the platform were described in
Methods.

1549

Supplementary Fig. 3: OuterMix algorithm improves multimodal training efficiency via factorized batch sampling.

a-c. Sampling and computation with random sampling strategy. A randomly sampled batch has $N \times M \times K$ combinations of genome sequence (N), chromatin feature (M), and CAP (K), selected from the full three-dimensional data manifold (**a**). This random processing approach generates large data matrices for encoding (**b**). Thus, each encoder for the corresponding modality would process every combination independently, yielding a computational complexity of O($N \times M \times K$) per encoder (**c**).

d-f. Sampling and computation with OuterMix strategy. OuterMix first samples *N, M, K* training samples from each dimension, effectively sampling a dense data cube from the full three-dimensional data manifold (**d**). Each sampling was encoded, with the encoded embeddings being recombined via a Cartesian product to form $N \times M \times K$ training samples (**e**). These encoded features are reused in the final Cartesian product and avoids redundant computation, reducing encoder computation complexity down to O(N) + O(M) + O(K).

1564

1565

1566

Supplementary Fig. 4: Cell-type-specific CAP-binding prediction performance of Prototype Chromnitron.

a. Performance comparison between Prototype Chromnitron, trained on datasets from two cell types (HepG2 and K562), and other models in pair-wise cell-type-specific regions among GM12878, HCT116 and IMR-90. Cell-type-specific ATAC-seq was used as input for prediction and accessibility-weighted motif score (Acc. Motif) baseline. Average prediction across cell lines excluding the target cell type was used for sequence-based models (AlphaGenome, Borzoi, and Enformer).

b. The same benchmark as in (**a**), except that sequence-based model predictions were averaged across all available cell types, including the validation cell line.

1577

1578

1579

Supplementary Fig. 5: Full-scale Chromnitron pretraining and finetuning.

a. Two-phase training strategy showing cross-CAP pre-training using shared weights (grey)
followed by individual fine-tuning for each CAP with low-rank adaptation (LoRA), visualized as
weight deltas (ΔW).
b. Model performance across pretraining and fine-tuning epochs, grouped by CAP categories. Line
and the shade represent scores of mean ± s.d. for each CAP category.
c. Scatter plot of model performance (Pearson's correlation between prediction and experimental
data of individual CAPs) between fine-tuned model and pretrained model.
1588

1589

1590
Supplementary Fig. 6: Model-training efficiency comparison between finetuning and training from scratch.
Training curves comparisons across three model-training strategies: fine-tuning a fully pretrained model (dark blue), fine-tuning a checkpoint after 19 epochs of pretraining (light blue), and training from scratch (green). The comparison includes overall validation loss (across all cell types) and Pearson's *r* relating to reference data in each cell type across training epochs.

1597

1598

Supplementary Fig. 7: Chromnitron predictions match experimental ChIP-seq replicates.
a-c. Evaluation of Chromnitron predictions against experimental ChIP-seq replicates for CTCF (top) and MAX (bottom). Each evaluation was performed by comparing prediction performance with that of replicating experimental measurements. These evaluations include Pearson's correlation to the reference experimental data across different percentiles of genomic bins ranked by mean signal strength (**a**), scatter plot of predicted or experimental replicate binding signals (x-axis) versus the reference experimental data (y-axis, **b**), and direct visualization of CAP-binding tracks of prediction alongside two other experimental replicates (Rep2, Rep3), all aligning to the reference experimental data (Ref, **c**).
d. Dimensionality reduction (t-SNE) visualization of predicted (circle) and experimental (squares) binding profiles from a subset of CAPs. Colors define CAPs clustered with similar binding profiles.

1610

1611

Supplementary Fig. 8: Paired visualization of Chromnitron-predicted and ChIP-seq-measured CAP binding profiles.

Paired CAP-binding tracks of Chromnitron prediction (top) and experimental data (bottom) for 36 randomly selected CAPs across the leave-out genomic region (chr 10). CAP-binding tracks were colored by their performance (Pearson's *r*).

1617
Supplementary Fig. 9: Performance metrics comparison between Chromnitron and three other models, using MAX as an example.
All performance metrics were performed against reference experimental ChIP-seq data of MAX. A total of eight statistics were calculated, including Pearson's correlation (r), Spearman's correlation (ρ), the coefficient of determination (R squared), F1 score, and area under the receiver-

operating characteristic (AUROC) and precision–recall (AUPR) curves, computed using both target-centric and prediction-centric labelling. Each of these methods was calculated using bins sizes of 50, 250, 500, and 1,000 bp (color intensity denotes bin size). Each plot shows a performance metric (y-axis) dynamic as the different percentages of genomic bins included in the analysis (x-axis, log-scaled). Red lines indicate selection of bins from the top percentiles only, while blue lines indicate selection of bins from both the top and bottom percentiles of bins.

1630

Supplementary Fig. 10: Robust peak-level performance for CAPs with low global correlation.
Performance metrics for four randomly selected low-correlation (Pearson's *r* < 0.5) CAPs,
including PCBP2, FUBP1, PBX2, and ZNF850. Despite the modest correlation coefficients in
these CAPs , Chromnitron-predicted CAP-binding peaks show high correspondence with
reference ChIP-seq peaks, measured by top and bottom percentile AUPR and AUROC (blue line).
Pearson's correlation was calculated by comparing to reference experimental ChIP-seq data. For
each CAP, four statistics were calculated, including AUROC and AUPR scores using both target-
centric and prediction-centric labelling. Metrics were calculated using bin sizes across 50, 250,
500, and 1,000 bp (color intensity denotes bin size). Red curves indicate selection of bins from the
top percentile only, while blue curves indicate selection of bins from both the top and bottom
percentiles of bins.

1645

Supplementary Fig. 11: Chromnitron performance benchmark.

a. Performance comparison between Chromnitron and other models in cell-type-specific regions between HepG2 and K562. Cell-type-specific ATAC-seq was used as input for Chromnitron, maxATAC and accessibility-weighted motif score (Acc. Motif) baseline. Average prediction across all cell lines was used for sequence-based models.

b. Performance comparison between Chromnitron and other models across the entire Chromosome 10 in HepG2 and K562 cells, including both cell-type-specific and non-specific binding regions.

c. Representative CAP binding tracks comparing Chromnitron, other models and experimental ChIP-seq measurement on chromosome 10.

80.05Mb   80.30Mb   80.55Mb

chr10:80,050,000-80,550,001

— ChIP-seq
— Chromnitron
— Motif Accessibility
— maxATAC
— Enformer

1656
Supplementary Fig. 12: Comparison of CAP-binding predictions across models.

Predicted CAP-binding profiles for 10 randomly selected CAPs were presented at the same 500kb
genomic window in the leave-out chromosome 10. Each CAP includes predicted binding tracks
from Chromnitron, accessibility-weighted motif score (Acc. Motif), maxATAC and Enformer
models, all referring to the experimental ChIP-seq data in leave-out chromosome 10. Min-max
scaling of the binding signal was applied to each track separately.
1663

1664

Supplementary Fig. 13: Performance statistics across cell types.

a. Benchmarking prediction performance on Chromnitron's leave-out genomic region

(chromosome 10) in HepG2 cells. Each dot represents one CAP, plotted as Chromnitron prediction

performance (y-axis) versus the comparison model (x-axis). Performance was quantified using

AUPR and AUROC, calculated by using ChIP-seq (top) or prediction (bottom) as the reference

label of binarized binding status.

b. Performance benchmarking by predicting cell type-specific CAP-binding. The benchmarking

was performed by comparing model-predicted binding difference between HepG2 and K562 cells

(HepG2 – K562) to the experimental ChIP-seq difference (HepG2 – K562). Pearson's r and

Spearman's ρ were included as evaluation metrics in addition to the AUPR and AUROC as in (**a**).

1675
Supplementary Fig. 14: Comparing chromatin accessibility, CAP binding, and Chromnitron
prediction.
a. Distribution of correlation coefficients (Pearson's *r*) between experimental ChIP-seq signal and
chromatin accessibility in HepG2 cells.
b. Scatter plots of CAP binding versus chromatin accessibility signals for experimental ChIP-seq
signal (Exp, grey) and Chromnitron predictions (Pred, blue).
1682

a

CAP-binding region

WT ...CCACTAGGTGGCGC...

...CCACTAGATGGCGC...
...CCACTAGCTGGCGC...
...CCACTAGTTGGCGC...

Per-Base mutation

Prediction

Base impact

...CCACTAGGTGGCGC...

Mutations in a region

Impact motif

b   Known motifs

NFE2
chr10:34,978,905-34,982,180

NFYB
chr10:102,241,971-102,245,990

USF1
chr10:97,580,922-97,584,580

ZNF384
chr10:28,377,279-28,380,782

ZNF146
chr10:116,930,298-116,933,679

ZNF770
chr10:103,659,674-103,663,182

c   Potential new motifs

MIER2
chr10:86,303,364-86,307,298

TRIM24
chr10:7,697,578-7,700,962

ZNF512
chr10:90,926,919-90,930,262

ZNF629
chr10:79,070,058-79,073,358

d   Indirect-binding motifs

RAD21
chr10:79,371,567-79,374,910

SMC3
chr10:95,427,368-95,430,796

EP300
chr10:22,680,088-22,683,758

MED1
chr10:116,053,898-116,057,330

Position

Position

1683

Supplementary Fig. 15: *In silico* **DNA sequence mutagenesis identifies CAP impact motifs**
and enables variant effect prediction.
a. Quantification of CAP-specific sequence impact scores through base-resolution *in silico*
mutations and Chromnitron prediction.
b-d. Chromnitron prediction-based *in silico* DNA sequence mutagenesis identifies CAP-specific
impactful DNA bases (i.e. impact motifs) that resembles both known (**b**) and novel (**c**) motifs, as
well as indirect binding motifs that resemble their interaction partner CAPs which binds DNA(**d**).
Impact scores were presented in the form of sequence logos which were normalized per base to
median value. Each impact motif was predicted at a specific locus as denoted in panel title.
1693

1694
Supplementary Fig. 16: Chromnitron learns DNA sequence features at single-nucleotide
resolution to enable cell-type-specific variant effect prediction.
a. Chromnitron predicts gain-of-binding effect at *TERT* promoter G250A variant (hg38
chr5:1295135 G>A). Left: CAPs ranked by mutation impact score with a vertical dashed colored
line marking ETV4, an ETS-family transcription factor. Right: binding profiles of ETV4 on wild
type (WT), mutant (Mut) sequences, and their corresponding differential binding (Diff: Mut – WT).
b. Chromnitron predicts loss-of-binding effect at a CTCF-binding site (hg38 chr10:25081160
C>T). Left: CAPs ranked by mutation impact score with red dashed line indicating CTCF. Right:
CTCF binding profiles across WT and Mut sequences and their differential binding.
1704

1705

Supplementary Fig. 17: *In silico* **chromatin accessibility perturbation revealed non-linear**
relationship between accessibility and CAP binding.
a. Chromatin accessibility perturbation using *in silico* scaling multipliers.
b-c. *In silico* chromatin accessibility perturbation results at CTCF (**b**) and ARID3A (**c**) binding
sites. Left: predicted CAP binding changes responding to perturbation radius and accessibility
multiplier. Middle: heatmap of predicted binding intensity upon perturbing accessibility at fixed
500 bp window, centered at binding peaks. Right: CAP binding changes (top) and the rate of
change (bottom) as a function of accessibility multiplier.
1714

1715

Supplementary Fig. 18: Chromnitron-based *in silico* CAP residue mutagenesis captures protein functional domains.

a. Schematic of *in silico* mutation for CAP protein sequences. Upon mutation, predicted CAP binding intensities of mutant CAPs were compared to reference CAP to calculate protein residue impact scores (Methods).

b. Violin plot distribution of impact scores across CAP residues, stratified by domain annotations. Top and bottom 10 domains are shown.

c. *In silico* CAP residue mutagenesis by Chromnitron captures protein residues from annotated functional domains. Each dot in the scatter plot represents a CAP's AUPR scores against protein domain annotations using *in silico* mutagenesis-identified residues (y-axis) and compared with that using randomly selected residues (x-axis).

1727

1728
Supplementary Fig. 19: Chromnitron enables inference of protein–protein interactions and
de novo **binding prediction across proteins.**
a. Visualization of LiNGAM causal-discovery algorithm using predicted CAP binding profiles.
Edges represent high interaction potential between adjacent CAPs. Nodes were colored by CAP
community clusters.
b. Performance comparison between different methods for discovering protein-protein interactions.
Causal-inference-based and Pearson's correlation-based interaction graphs derived from
Chromnitron predictions and from ChIP-seq data were compared against curated interactions from
the STRING database using AUPR (Methods).
c. Evaluation of *de novo* CAP-binding prediction for unseen proteins. Performance evaluation of
prototype Chromnitron pretrained on 385 CAPs (in HepG2 and K562 cells) and evaluated in leave-
out CAPs and leave-out genomic region (chromosome 10). Corresponding CAP protein sequence
was used for Chromnitron to generalize to new CAPs. Performance was compared with
accessibility-weighted motif score (Acc. Motif) baseline, demonstrating the viability of CAP-
sequence-encoded structural features that enables generalizable CAP-binding prediction.
1744

1745
Supplementary Fig. 20: Profiling ZNF865 binding at *TOX* promoter-proximal locus.

a. Predicting cell-type-specific CAP binding using ATAC-seq datasets across conditions.
b. Characterization of ZNF865 binding profile at the *TOX* promoter-proximal regulatory element.
Left: Scatter plot of ZNF865 experimental ChIP-seq binding (grey) and predicted binding (blue)
versus chromatin accessibility (x-axis), indicating that ZNF865 preferentially binds at low-
accessibility regions. Right: Chromatin accessibility signals at *TOX* promoter and the ZNF865-
bound CRE across T cell exhaustion pseudo-time.
c. Chromatin accessibility at the ZNF865-bound CRE (top) and the corresponding ZNF865 impact
motif revealed by *in silico* mutagenesis (bottom).
1755

1756
Supplementary Fig. 21: *In vitro* **chronic stimulation of human primary CD8⁺ T cells to model**
T cell exhaustion trajectory.
a. Schematic of the *in vitro* treatment of human CD8⁺ T cells across acute stimulation (top), chronic
stimulation by anti-CD3 antibodies (middle), and chronic stimulation by anti-CD3/CD28
antibodies (bottom).
b. Flow cytometry quantification of cell surface markers of T cell exhaustion (PD1, LAG3, TIGIT,
and TIM3) and intracellular cytokine production (IFN-γ, IL-2, and TNF-α) in different stimulation
groups at day 12. Bar plots show mean ± s.d. for n = 3 independent replicates. Statistical
comparisons performed using two-sided unpaired Student's *t*-tests with significance denoted as *,
P < 0.05; **, P < 0.01; ***, P < 0.001.
1767

1768

Supplementary Fig. 22: Validating ZNF865 as an upstream regulator of *TOX* for T cell exhaustion.

a. Schematic of *in vitro* model of T cell exhaustion through persistent anti-CD3/CD28 stimulation, coupled with CRISPR/Cas9-based gene knockout in primary T cells.

b. RT-qPCR quantification of *TOX* and *ZNF865* mRNA expression level changes across chronic stimulation of human CD8[+] T cells from day 4 to day 12, indicating increased expression along the trajectory. Relative expression levels were normalized to day 2 using the ΔΔCt method. Each dot represents an independent biological replicate (n = 4); error bars indicate mean ± s.e.m.

c. Surveyor assay confirmed efficient CRISPR knockout gRNA targeting *TOX* and *ZNF865*. Each gene includes two gRNAs for knock-out perturbation. Genomic DNA was amplified around the gRNA target sites and the fragments were digested with T7 Endonuclease I.

d-e. Flow cytometry quantification of T cell exhaustion markers (**e**, PD1, TIGIT, LAG3 and TIM3)
and cytokines (**f,** IFN-γ, IL-2, and TNF-α) across negative control, non-targeting control (NTC),
TOX knockout, or *ZNF865* knockout conditions. Bar plots show mean ± s.d. for n = 3 independent
replicates. Statistical comparisons performed using two-sided unpaired Student's *t*-tests with
significance denoted as *, P < 0.05; **, P < 0.01; ***, P < 0.001.
f. RT–qPCR quantification of *ZNF865* and *TOX* mRNA level changes across chronic stimulation
days with individual gene knockouts (*ZNF865* KO and *TOX* KO). Relative expression levels were
normalized to day 2. Each dot represents an independent biological replicate (n = 4); error bars
indicate mean ± s.e.m.
1789
1790

1791

Supplementary Fig. 23: Validation of CRISPR-mediated *ZNF766* knock-out effects on
chronic-stimulated T cells.

a. Distribution of reported exhausted T cell (Tex) marker genes (CTLA4, PDCD1, TIGIT,
HAVCR2) in co-regulated CRE groups (top) and their corresponding co-regulating CAP clusters
(bottom).

b. Schematic of CRISPR dropout screens in human CD8[+] T cells using *in vitro* chronic stimulation
model.

c-d. Flow cytometry quantification of T cell exhaustion markers (**c**, PD1, TIGIT, LAG3 and TIM3)
and cytokine secretion levels (**d,** IFN-γ, IL-2, and TNF-α) across negative control, non-targeting
control (NTC), or *ZNF766* knockout conditions. Bar plots show mean ± s.d. for n = 3 independent
replicates. Statistical comparisons performed using two-sided unpaired Student's *t*-tests with
significance denoted as *, P < 0.05; **, P < 0.01; ***, P < 0.001.

1804

1805
Supplementary Fig. 24: Transcriptomic profiling of *ZNF766* knock-out effects in chronic-
stimulated T cells.
a. RT-qPCR quantification of *ZNF766* mRNA level changes across chronic stimulation of CD8+
T cells from day 4 to day 12, indicating increased expression along the trajectory. Relative
expression levels were normalized to day 2 using the ΔΔCt method. Each dot represents an
independent biological replicate (n = 4); error bars indicate mean ± s.e.m.
b. Surveyor assay confirmed efficient CRISPR knockout gRNA targeting *ZNF766*.
c. RT-qPCR quantification of *ZNF766* mRNA expression levels after gene knockout. Primers were
designed flanking the CRISPR cutting site to confirm transcript disruption. Increased ΔCt number
after perturbation indicates efficient knock down of unperturbed *ZNF766* mRNA. Error bars
indicate mean ± s.e.m. Statistical significance was determined using a two-tailed paired Student's
t-test (P < 0.001).
1818

d. Heatmap of differentially expressed genes between *ZNF766* knockout and non-targeting control
(NTC) T cells.
e. Volcano plot of differentially expressed genes between *ZNF766* knockout and NTC groups. Red
and blue dots indicate significantly upregulated and downregulated genes, respectively, in the
knock-out condition ($|\log_2FC| > 1$, p.adj < 0.05). Representative genes involved in T cell activation
(e.g., *GZMB*, *MMP25*, *SLC7A5*) and exhaustion-associated regulators (e.g., *PDCD4*, *NT5E*,
TXNIP) were individually labeled.
f. Gene Ontology (Biological Process) enrichment analysis of significantly upregulated genes in
ZNF766 knockout group. Dot size corresponds to gene count per term; color indicates adjusted p-
value. Enriched pathways highlight enhanced DNA replication, chromatin organization, and
mitochondrial respiration, suggesting a functional restoration of exhausted T cells.
1830

1831

Supplementary Fig. 25: Dynamics of CAP binding and CRE activities during Purkinje cell differentiation.

a. CAP binding profiles across CRE groups. Representative CAPs were labeled. Color scale indicates normalized binding.

b. T-SNE embedding of CREs colored by activity clusters (same as Fig. 4D).

c. Heatmap of CRE identity transitions across Purkinje cell differentiation stages.

d. Average binding dynamics for four different CRE sets across pseudo-time. Each CRE set includes three CAP classes (Repressor: e.g. TCF7, RCOR1; Early CAP: e.g. BCL11A, MEF2A, ATF2; Late CAP: e.g. FOXA1, FOXA2). Shading denotes 1 standard deviation.

1841

1842
Supplementary Fig. 26: Characterizing Purkinje cell differentiation dynamics using pseudo-
time and CRE as organizing features.
a. T-SNE embedding of individualized CAP binding across loci using time trajectory as feature,
colored by trajectory trend cluster. Each dot represents a CAP binding at a single locus across all
pseudo-time stages.
b. Average CAP-binding trajectories for each cluster across time, illustrating both early- and late-
acting regulatory programs.
c. T-SNE embedding of CAPs across time using CRE-level binding profiles as features, colored
by CAP cluster identities. Each dot represents a CAP's binding profile across all loci at a given
time point.
d. Alluvial diagram showing transitions in CAP cluster identity across developmental stages.
Representative CAPs in each cluster were annotated on the right, highlighting coordinated shifts
in regulatory factor usage during Purkinje cell maturation.