Determinants of mutation load in birds

7   Fidel Botero-Castro [1], Jochen B. W. Wolf [1*]

10   [1] Department of Evolutionary Biology, Faculty of Biology, LMU Munich, Germany

18   * Address for correspondence: Jochen Wolf, Department of Evolutionary Biology, Faculty of
19   Biology, Ludwig Maximilian University of Munich, Großhaderner Straße 2, 82152 Planegg-
20   Martinsried, Germany, Phone: 0049-89-2180-74102, E-mail j.wolf@bio.lmu.de; Fidel
21   Botero-Castro, fidel.boteroc@gmail.com

<u>Running title:</u> Mutation load and recombination

<u>Keywords:</u> Selection, effective population size, recombination, distribution of fitness effects,
purging.

<u>Summary.</u> Harmful genetic mutations can accumulate in populations, affecting individual-
and population-level health. The authors analyzed genomic data of 24 populations of 19 bird
species to understand the evolutionary determinants of such mutation load. They confirm that
small populations tend to experience a heavier mutation burden. Moreover, deleterious
mutations accumulate more readily in regions of low recombination and are further elevated
by GC-biased gene conversion (*gBGC*), which is offset by purging efficacy in large, but not
in small populations. This study suggests that genetic risk factors in small populations are
fueled by *gBGC* and cluster in regions of low recombination.

Abstract

Many mutations have detrimental effects. The mutation load in a population depends on the efficacy of purifying selection in removing deleterious genetic variation. Here, we estimated the proportion of deleterious mutations segregating in 24 population samples of 19 bird species. Exploiting the conserved avian karyotype with high variation in recombination rate and GC content, we quantified the joint effects of effective population size (N_e), recombination (r) and GC-biased gene-conversion ($gBGC$). In agreement with the nearly-neutral theory of molecular evolution, mutation load was substantially higher in populations with small N_e. Purging efficacy increased with recombination rate resulting in more than a two-fold difference of genetic load between large and small chromosomes. GC-biased mutations contributed about one third to the pool of deleterious mutations. Their expected accumulation in regions of high recombination was offset by purging efficacy in large, but not small populations. This study provides insight into how the interaction of evolutionary processes shapes mutation load. It suggests that genetic risk factors in small populations are fueled by $gBGC$ and cluster in regions of low recombination.

Introduction

Novel mutations enter a population each generation. Many are neutral, but of those affecting the fitness of its carrier most are deleterious, some are advantageous (Keightley and Lynch 2003). The overall frequency of selective effects is summarized in the genome-wide distribution of fitness effects (DFE, **Figure 1**) reflecting properties of the organism and the distance to its global fitness optimum (Eyre-Walker and Keightley 2007; Lanfear et al. 2014; Chen et al. 2022). Lethal or strongly deleterious mutations are quickly removed from a population by purifying selection, and beneficial mutations are promoted to fixation. Neutral

and weakly deleterious mutations, however, can persist for many generations and constitute

the bulk of segregating genetic variation. The time they spend in a population depends on the

interaction of their fitness effects (s) with the effective size of the population (N_e), the

population selection coefficient $N_e s$ (Kimura 1983; Ohta 1992; Charlesworth 2009; Akashi et

al. 2012). As a consequence, deleterious mutations can rise to high frequencies in populations

with small N_e (**Figure 1**).

These considerations of the nearly-neutral theory of molecular evolution are of real-world

importance (Dussex et al. 2023). The accumulation of deleterious mutations maintained by

mutation-selection balance, the mutation load (Muller 1950), can eventually elevate the risk

of population extinction (Lande 1994; Frankham 1998; Soulé and Mills 1998; Agrawal and

Whitlock 2012). There is growing empirical evidence that mutation load indeed scales

negatively with proxies for N_e as it has been predicted by theory (Kimura et al. 1963; Lynch

et al. 1995). One of such proxies is the size of the geographical range occupied by a given

population, which is expected to be directly correlated to the census size of said population,

and, ultimately, to N_e (Peart et al. 2020). Indeed, across a broad taxonomic range, the

efficacy of purifying selection tends to be enhanced in species and populations characterized

by a large number of effectively breeding individuals (Slotte et al. 2010; Deinum et al. 2015;

Chen et al. 2017; Castellano et al. 2018; Galtier and Rousselle 2020). As a consequence,

mutation load tends to be increased in small, isolated island populations as compared to

populations with larger distribution areas (Kutschera et al. 2020), even when comparing

island and continental populations of the same species (Leroy et al. 2021).

While the effect of population size on mutation load has been well studied, other factors have

received less attention. Recombination, for instance, is known to modulate the efficacy of

selection at the intragenomic scale. In regions of low recombination, Hill-Robertson

interference acts as to reduce the local effective population size, thus increasing the

probability of deleterious alleles rising to fixation (Hill and Robertson 1966; Felsenstein

1974; Comeron et al. 2008). Similarly, the purging effect of synergistic epistasis between

deleterious mutations, though contentious, is influenced by recombination (Agrawal and

Whitlock 2012). As a consequence, mutation load is expected to differ within the genome as

a function of recombination intensity (Comeron et al. 2008). Empirical evidence comes from

exploiting variation in recombination rates within and across chromosomes (Gossmann et al.

2011; Webster and Hurst 2012; Corcoran et al. 2017; Chase et al. 2024). Due to obligate

crossover, small chromosomes have high recombination rates per physical unit (cM/Mb)

(Hillier et al, 2004; Kawakami et al. 2014, Backström et al 2010, Bascón-Cardozo et al 2023)

and are expected to carry the smallest load, whereas large, low-recombining chromosomes

are expected to accumulate more deleterious mutations. Non-recombining, uniparentally

inherited sex-chromosomes or organellar genomes are predicted hotspots of mutation load

and have accordingly received most attention (Charlesworth and Charlesworth 2000; Marais

2007; Warmuth et al. 2022).

Importantly, recombination is also known to be associated with GC-biased gene conversion

(*gBGC*), referring to the preferential reparation of double-strand breaks towards G and C

alleles, which eventually results in an increase of GC content in highly recombining regions

(Galtier et al. 2001; Duret and Galtier 2009). Most of these mutations are prone to deleterious

effects but are nevertheless constantly replenished through the segregational bias in regions

undergoing high recombination, mimicking the effect of positive selection (Berglund et al.

2009). *gBGC* is thus predicted to counteract efficient purging of weakly-deleterious

mutations in regions of high recombination (Bolívar et al. 2018). Yet, it is unclear at which

rate these opposing forces operate, in particular, as the strength of *gBGC* also seems to

interact with the overall effective population size (Bolívar et al. 2019) (**Figure 1**).

Avian genomes are well suited to study these interacting effects, as they provide the

necessary variation in N_e across a stable, yet highly variable, recombination landscape within

and between chromosomes (**Figure 2**) (Ellegren 2010). To assess the interaction of 2/4/2026

5:57:00 AMrecombination, effective population size and GC-biased gene conversion, and

their influence on mutation load, we collated genome-wide resequencing data across 24

populations from 19 avian species varying in effective population size. We estimated the

DFE for individual chromosomes with substantial variation in recombination rate to obtain

estimates of the proportion of segregating, mildly-deleterious mutations ($N_e s \in$ (-1;0]),

serving as proxy for mutation load. According to theory, we expect the following: mutation

load should be: 1) higher in populations with low N_e, 2) reduced in regions of high

recombination and 3) increased for the mutational spectrum subject to *gBGC* (weak-to-strong

mutations: [A,T]->[C,G]). 4) Relative rates of the two recombination-mediated, opposing

processes of purging and *gBGC* may differ and be additionally modified by N_e (**Figure 1**).

Materials and Methods

Sequence Data

We collated publicly available resequencing data for 443 individuals from 47 populations of

30 passerine (sub)species and utilized *.vcf* files generated by comparable pipelines described

in (Leroy et al. 2021) and (Kutschera et al. 2020) (see **Supplementary Table 1** for accession

numbers of NCBI's Sequencing Read Archive). In summary, reads were first mapped to the

closest available reference (see **Supplementary Table 1**). After the usual steps of adding

read group information, merging bam files per individual, marking duplicates and

realignment of indels, both studies used GATK (v3.4.0 in Kutschera et al (2020), v3.7.0 in

Leroy et al. (2021)) to call variants using joint genotyping with *GenotypeGVCF*. This

resulted in *vcf* files including both variant and invariant sites.

While Leroy et al. (2021) directly proceeded to filter the output of *GATK* on the basis of

quality by depth (QD < 2.0), Fisher Strand bias (FS > 60), mapping quality (MQ < 40),

MQranksum < 2 or ReadPosRankSum < 2 or Raw Mapping Quality (Raw_MQ < 45,000),

Kutschera et al (2020), performed an additional step of base-quality score recalibration

(BQSR) before jointly genotyping, using a set of "true" variants resulting from the highest

quality variants identified in the intersection of the outputs of *samtools* v1.3 (Li et al. 2009),

FreeBayes v1.0.2 (Garrison and Marth 2012), and *GATK*'s HaplotypeCaller. Then, after joint

genotyping, variants were again filtered using variant quality score recalibration (VQSR)

implemented in *GATK* and only those passing the filter were retained. Given that both studies

include samples from both island and mainland areas, and that the number of resulting total

sites and variants is comparable across species and populations, the observed differences

between island and mainland species are unlikely to result from biased sampling across

studies or differences in variant filtering.

We filtered all *vcf* files such that only sites with a minimum Qval of 20 and belonging to

autosomal scaffolds/chromosomes were retained. Next, sites were marked as 0-fold, 2-fold,

3-fold and 4-fold degenerate on the basis of the genome annotation of the corresponding

reference genome (**Supplementary Table 1**) using the script *NewAnnotateRef.py* by

Williamson   et   al.   (2014)   available   at

https://github.com/fabbyrob/science/tree/master/pileup_analyzers. The output file was then

converted into bed files for each type of site (0-, 2-, 3-, 4-fold degenerated) using *bedtools,*

and a *vcf* file containing these sites was generated. The four resulting *vcf* files were merged

and split by chromosome. An additional dataset was produced by excluding CpG-prone sites

and weak-to-strong mutations (A-to-G, A-to-C, T-to-G, T-to-C substitutions) following the

approach of Kutschera et al. (2020). This dataset was used to assess the effect of *gBGC* by

comparison to the dataset including all sites.

Relatedness and Population Structure

The presence of closely-related individuals, as well as population structure and gene flow

between populations, can distort inference of the DFE by mimicking changes in population

size, although the effect can be partially absorbed by explicitly modelling a shift in

population size, as implemented in Keightley and Eyre-Walker's (2007) two-epoch model

(2e) (see also (Kutschera et al. 2020)). To reduce any effects arising from relatedness or

population structure, we first removed related individuals within each species. As a first step,

we filtered from the full autosomal dataset sites with linkage-disequilibrium above 60% with

bcftools (bcftools +prune -l 0.6 -w 50kb -n 5 --AF-tag), and then used plink (v2.0) on the

filtered *vcf* file to produce a KING-formatted table of kinship coefficients (option –make-

king-table). All individuals with kinship coefficients higher than 0.0442, corresponding to 3[rd]

degree relatives were excluded (Manichaikul et al. 2010).

In order to control for population structure, we additionally performed a PCA for each

species using the full set of coding autosomal SNPs and all individuals in *plink* (v1.90b6.21).

Many of the individuals appearing as outliers in the first two axes of variation corresponded

to those identified as being closely related, providing further arguments for their exclusion

from the final dataset (**Supplementary Figure 1**). Moreover, the Italian population of

collared flycatchers (*Ficedula albicollis*) and the Spanish population of pied flycatchers

(*Ficedula hypoleuca*) were analyzed on their own since they clustered separately from the

remaining populations, as has also been previously documented (Burri et al. 2015). Two

species of white eyes (*Zosterops borbonicus* and *Z. virens*) were excluded due to significant

structure and gene flow between the subpopulations (Bourgeois et al. 2020). Furthermore, as

the inference of the site frequency spectrum (SFS) requires sites to be genotyped for all

individuals in the sample (i.e. no missing data) we removed individuals with lower quality of

sequencing data at the cost of sample size (**Supplementary Table 1**): given that small

chromosomes display an increased GC-content and Illumina sequencing biases against them,

they are more likely to have missing data (i.e. sites not genotyped for some individuals). Yet,

these chromosomes are important for the intended analyses, as they 'provide' the high

recombination rates, and cannot be discarded. We therefore maximized the number of sites

sampled by setting a very conservative threshold for missing data: any individual displaying a

fraction of missing data of more than 5% of sites missing in any chromosome was discarded.

Moreover, SFS required a minimum of eight chromosomal copies to be sampled, i.e. four

individuals with sufficient sequencing coverage (cf. Kutschera et al. 2020). As only

exception, the Spanish population of *Ficedula hypoleuca* was included with six

chromosomes.

After applying these individual and population filters, the final dataset consisted of 24

populations from 19 species, from which eight were mapped to a reference of the same

species, and sixteen to that of the closest relative available in the genus or family. Divergence

times between sample and query were within 6.5 mya (**Supplementary Figure 2**) (McTavish

et al. 2025). Each population is represented by a median average of 1,604,1053 total and

46,342 segregating sites (see **Supplementary Table 2** for sample sizes of individuals and

segregating sites).

Inference of the DFE

We inferred the folded SFS for each population and each of 28 autosomes separately using

the scripts *vcfSummarizer.py* and *bootstrap.py* available at

(https://github.com/fabbyrob/science/tree/master/pileup_analyzers). By running these two

scripts, all sites with missing data, indels, or multi-nucleotide polymorphisms are discarded.

The inferred folded SFS was used to model the DFE of each species/population both per-

chromosome and using the full autosomal dataset using *DFEalpha v2.16* (Keightley and

Eyre-Walker 2007). This approach requires a set of sites under selection, in which we

included the 0-fold, 2-fold and 3-fold degenerated sites, and a set of sites evolving under or

close to neutrality, for which we used the set of 4-fold degenerated sites. Two demographic

models, one with constant size (1-epoch model) and one with change in population size (2-

epoch model) - either an expansion or a contraction - were run for comparison and the best

fitting model was selected using a likelihood ratio test. Note, however, that single estimates

of long-term N_e are likely not capturing more complex demographic histories including

repeated changes in population size with contributions of gene flow. These are likely only

controlled in part by the model estimating the DFE from the SFS.

Inference of the best-fitting demographic model differed by chromosome (**Supplementary**

Figure 3). This may be due to differences in power but may also suggest that different

genomic regions experience demographic variations in different ways, likely due to varying

linked selection (see e.g. (Boitard et al. 2022)). Estimates of the DFE can in consequence also

be expected to change across chromosomes. Inference of the DFE using the full autosomal

dataset was largely dominated by the largest chromosomes, masking different inferences for

the smaller chromosomes. Given the variability among chromosomes and our interest in

comparing the DFE to recombination rate estimates on a per-chromosome basis, we chose to

select the statistically preferred model per chromosome and population rather than imposing a

single model to all genomic regions. The fit of the inferred DFE was assessed through the

plots of the observed SFS and the ones expected under the best inferred demographic model.

These plots are provided in the *zenodo* repository.

While there are alternatives for DFE estimation, like *polyDFE* (Tataru et al. 2017) or *grapes*

(Galtier 2016), they turned out to be less suited to the analyses we intended to perform.

PolyDFE requires an unfolded SFS which is prone to errors (Tataru et al. 2017) and requires

a significant amount of data (around 20Mb), impossible to reach for the smaller

chromosomes. Grapes is mainly aimed to estimating the fraction of adaptive substitutions

(alpha), and although the DFE is also estimated, the slices it provides (proportions of weakly-

and strongly-deleterious mutations) are not directly comparable with the ones provided by

DFE-alpha.

Estimation of the Effective Population Size

Effective populations sizes were obtained from Kutschera et al. (2020) and Leroy et al.

(2021), inferred as the harmonic mean of N_e estimates derived from the sequential Markovian

coalescent (McKenna et al. 2010; Schiffels and Durbin 2014) weighted by the length of each

time period. For details see Supplementary Text S1 in Kutschera et al. (2020) and Method

detail in Leroy et al. (2021). According to the neutral theory, long-term coalescent N_e should

be similarly reflected by $\theta_\pi = 4N_e\mu$. In agreement with this prediction, estimates from the

sequential Markovian coalescent covaried strongly with estimates of synonymous nucleotide

diversity (π_s) as in Leroy et al (2021). The estimates were obtained from pairs of genomic

sequences reconstructed for each individual, applying filters for depth of coverage, and

removing sites with missing data (**Supplementary Figure 4**). Note that both estimates of N_e

comprise the entire population history likely including a mainland phase prior to island

colonization in several cases. Since the DFE was equally shaped during the entire history of

the population, including the mainland phase, these long-term estimates of N_e, however, seem

an appropriate choice.

Besides these coalescent-based estimates, we used geographical range (Island vs. Mainland)

as a proxy to N_e (see also discussion). This assumption is based on the expectation that

species occupying mainland territories have larger population sizes than those living on the

smaller territories located in islands (**Supplementary Figure 5**). Moreover, this link is

further supported by the documented observation, in agreement with the predictions of

neutral theory, that populations with large N_e should be more genetically diverse than those

with reduced effective population sizes (Kutschera et al. 2020; Leroy et al. 2021)

(**Supplementary Figure 4**). Life history traits influence N_e and evolutionary rates, and might

also affect the DFE (Romiguier et al. 2014). Since all species under considerations here are

passerines with comparable life histories, we abstained from including this in the statistical

models to avoid overparameterization.

Recombination Rate Estimation

The average recombination rate (*r*) per chromosome was approximated in centimorgans per

mega base pair [cM/Mb] as follows: we extracted estimates of the per-base population

recombination rate ρ[1/bp] obtained by Vijay et al. (2016) in bins of 50 kb windows for 3

populations of hooded and carrion crows (*Corvus (corone) ssp.*). We then calculated, for each

population, the median across the 50 kb windows to obtain per-chromosome estimates. Last,

we averaged the estimates across the three populations. We then estimated genetic variation

along each chromosome and converted the population recombination rate estimates ρ[1/bp] to

recombination estimates in centimorgan using the relationship $\rho \, / \, \pi \, = \, 4N_er \, / \, 4Ne\mu \, = \, r \, / \, \mu$.

Multiplication with the mutation rate and scaling to cM (10^2) and Mb (10^6) then yields

r[cM/Mb]. We chose to use a single mutation rate of 4.6×10^{-9} mutations/site/generation as

estimated by Smeds et al. (2016) for the collared flycatcher. The resulting per-chromosome

estimates range between 2.46 and 11.5 cM/Mb (mean weighted by chromosome length: 3.57)

adding up to a total genetic map length of 3,500 cM which is comparable to previous,

pedigree-based estimates of passerine birds (Backström et al. 2010; Kawakami et al. 2014).

The choice of a single mutation rate for all species seems justified, as the estimate for

flycatcher is highly comparable to that of zebra finch (5.0×10-9 mutations/site/generation)

(Prentout et al. 2025). Flycatchers (Muscicapidae) and zebra finches (Estrildidae) diverged

~21 mya ago, which approximates the range of 28 mya of divergence for our study organisms

(**Supplementary Figure 2**) (McTavish et al. 2025).

Central to our study is the assumption that the karyotype of passerine species considered here

is stable enough to reflect comparable recombination rates across syntenic chromosomes

(Dutoit et al. 2017; Vijay et al. 2017). To test the assumption that a single set of

recombination rates is representative for the species involved in the analyses, we compared

chromosome-level recombination rates of hooded crow (Vijay et al. 2016) with those of zebra

finch (*Taeniopygia guttata*) (two estimates from Singhal et al. (2015) and Stapley et al.

(2010)), collared flycatcher (*Ficedula albicollis*) (Chase et al. 2024), and Eurasian blackcap

(*Sylvia atricapilla*) (Bascón-Cardozo et al. 2024). Analogous to the mutation rate estimates

for flycatcher and zebra finch described above, these species represent approximately 23

million years of avian evolution (divergence time Sylvidae – Estrildidae) and thus encompass

most of the divergence among the species included in this study. As the estimates of zebra

finch in Singhal et al (2015) and the ones of the Eurasian blackcap were provided as

estimates of the population recombination rate (ρ / bp), we used the values of genetic

diversity provided in the respective studies (genomic estimate of 0.082 for zebra finch; per

chromosome estimates in Eurasian blackcap as in Table S3 of (Bascón-Cardozo et al. 2024)),

and the avian mutation rate as in Smeds et al. (2016)(4.6×10^{-9} mutations/site/generation), to

convert them into cM/Mb using the same approach as for hooded crow outlined above.

Chromosome-level recombination rates were significantly correlated across all comparisons

(median R^2=0.668, median p-value = 1.838787^{-07}, **Supplementary Figure 6**) supporting that,

at chromosome-level, recombination rates are comparable across passerine birds. A single

recombination estimate, here from crows, appears thus suitable for comparative analysis,

although it will introduce noise which may conceal the effect of recombination rate on the

DFE. Moreover, chromosome length strongly correlates with recombination rate and is

highly conserved across avian species (Ellegren 2010, this study; Bascón-Cardozo et al.

2024). We therefore also used chromosome length as a proxy for recombination rate.

In order to ensure that the chromosomes correspond to the names assigned in the different

references used here (**Supplementary Table 1**), we verified synteny based on the

corresponding annotations using custom scripts. First, all annotated genes for each

chromosome were text-mined from annotation files for hooded crow, zebra finch, collared

flycatcher, great tit and chaffinch. The gene lists were then compared for each chromosome

between each possible pair of references. The annotations were highly consistent and synteny

could be verified, with few exceptions, for all references. Most mismatches occurred

between the reference for hooded crow and the other references, mainly due to crow scaffolds

belonging to a) chromosomes that were not present in the crow assembly version

GCF_000738735.1, to which the *vcf* files utilized here correspond, being assigned to other

chromosomes (e.g. 25, LGE22) or to b) small chromosomes (e.g. 23, 27). The identified

mismatches were corrected and the scaffolds included in the correct chromosomes, if they

were confirmed to be syntenic across all the other analyzed references; or excluded, if the

chromosomes were not present in the hooded crow assembly. A few mismatches from the

used zebra finch reference (GCF_000151805.1) have been already corrected and verified as

syntenic in a more recent annotation (GCF_003957565.2).

Statistical Analyses

We used a mixed linear model framework to test the influence of recombination (*r*), effective

population size (N_e) and GC-biased gene conversion (*gBGC*) on the proportion of mildly

deleterious alleles ($N_e s \in$ (-1;0]). Recombination and effective population size were

approximated by two alternative measures (*r*: recombination rate [cM/Mb] or chromosome

length (see above); N_e: species restricted to island yes/no or log(N_e)). The main results were

largely unaffected by the choice of measure. In the main text, we present the results for

recombination rate [cM] and island yes/no. Results for the alternative measures can be found

in **Supplementary Table 3.** The effect of *gBGC* was assessed as categorical variable by

comparing the estimate of DFE using all polymorphisms (with *gBGC*) or the subset including

only weak-to-weak (A-T), and strong-to-strong mutations (C-G) (without *gBGC*).

We fitted a mixed effect model including all three parameters and a choice of meaningful

interactions as fixed effects using the lmer function of the R package lme4 (Bates et al. 2015).

We were interested whether the effect of recombination on the DFE depended on N_e or *gBGC*

(pairwise interactions: $r*N_e$, $r*gBGC$), as well as whether the effect of *gBGC* differed by N_e

(pairwise interaction $N_e *gBGC$). In addition, to understand whether the relative rates of the

two recombination-mediated, opposing processes of purging and *gBGC* differed by N_e, we

also investigated the interaction of all three factors (three-way interaction: $r*N_e*gBGC$). We

further incorporated chromosome (chr) and population identity (pop) as random effects

including variation in slope for the latter. More than a single population was available for a

subset of five species, which led us to also include species (spec) as a nested random effect

(population nested in species). Model choice was then performed largely following (Zuur et

al. 2009): we started with a model containing the full set of random factors and all relevant

fixed factors and their interactions: *DFE ~ r + Ne + gBGC + r*Ne + r*gBGC + Ne*gBGC*

+ r*Ne*gBGC + (1|chr) + (1+r|pop/spec). We then selected the best combination of random

parameters using restricted maximum likelihood estimation (REML) minimizing the Akaike

Information Criterion (AIC) which eliminated the nested population/species component and

reduced the random effects to *(1+r|pop)+(1|chr)*. Next, we formulated a set of 10 models

exploring the influence of *r*, N_e, *gBGC* and their interactions with inclusion of the null model

(see **Supplementary Table 3**) and selected the model minimizing AIC using full maximum

likelihood which provides information about the goodness of fit of the respective model to

the observed data and penalizes overly complex models (Johnson and Omland 2004). Models

differing by less than two units of AIC were considered to be supported. For parameter

estimation, we then re-ran the best model using REML estimation and conducted an analysis

of variance as implemented in the R package car (Fox and Weisberg 2019). Finally, we

assessed normality of residuals by visual inspection and model performance by plotting

observed to predicted values (**Supplementary Figure 7**).

In order to assess the impact of phylogenetic dependency, we built a mitochondrial

phylogeny for all 19 species using part of the sequences included in Leroy et al (2021), to

which we added missing sequences for the two crow species (NC062298.1, NC051471.1).

We inferred a maximum likelihood (ML) tree using *IQtree* (v2.1.3) (Minh et al. 2020) under

a GTR+GAMMA+I substitution model to account for base content and rate heterogeneity

across sites (**Supplementary Figure 8**). We then fitted Bayesian mixed models using the

MCMCglmm package in R (Hadfield 2010). We modeled the DFE as a function of

recombination rate (parametrized as r [cM/Mb]) and N_e (parameterized as island yes/no),

gBGC (yes/no) and their interactions, while including a phylogenetic random effect whose

covariance structure was specified by the inverse of the phylogeny-derived relatedness

matrix. Following the mixed model above, chromosome was added as additional random

factor. We focused model comparison on the null model and the two best models as inferred

by the mixed model described above (three-way interaction, all two-way interactions).

Models, including only one population per species (see **Supplementary Table 2**) were run

under Gaussian error structure using weakly informative priors (G: $V = 1$, $v = 0.002$; R: $V = 1$,

$v = 0.002$) with 150,000 iterations, a 50,000-iteration burn-in, and thinning interval of 50 for

which no evidence of autocorrelation remained. Model choice was based on the Deviance

Information Criterion (DIC) and visual inspection of posterior distributions. In this Bayesian

framework, effects with posterior distributions that exclude zero correspond to high posterior

probability of an effect, analogous to low p-values in frequentist analyses, while broadly

overlapping distributions indicate weak or no support.

Results

We calculated the DFE for each of the 24 populations from 19 different species based on

segregating sites in coding sequence. The DFE was calculated separately for each of 28

autosomes by contrasting the site frequency spectrum (SFS) of neutral mutations at 4-fold

degenerate sites with the frequency of non-synonymous mutations subject to selection

controlling for demographic perturbation where necessary (see methods). Recombination

rate, approximated in cM/Mb, covaried negatively with chromosome length and (*gBGC-*

impacted) GC content (log-log correlations: R^2=0.77 and 0.81, respectively; both p<0.001)

(**Figure 2**).

In order to establish a baseline without the influence of *gBGC*, we first performed the

analyses on the SFS after excluding CpG-prone sites and weak-to-strong mutations [A,T]-

>[C,G]. The median, genome-wide proportion of mildly deleterious mutations differed

substantially between populations (min/median/max: 0.09/0.18/0.25), as well as between

species (0.09/0.19/0.25). This variation in mutation load appears to be well explained by

population size, regardless of whether it was approximated by coalescence-based estimates of

N_e or by species distribution (island vs. mainland) (**Figure 3A, Supplementary Figure 5**).

Species restricted to islands had a median average N_e of approx. 90,000, while for mainland

species this value was approx. 160,000. This 2-fold difference in N_e was reflected by a

median increase in the mutation load of approximately 1.6-fold. Mutation load was not only

dependent on N_e of the species, but also varied along the genomes of nearly all studied

populations as a function of recombination rate: low-recombining chromosomes carried

substantially more load than high-recombining genomes for both island and mainland

populations (**Figure 3A**).

gBGC increased the mutation load by approximately 1.5-fold on median average (**Figure**

3B). At first glance, the patterns of mutation load in relation to recombination rate and

species-level effective population size (N_e) appear similar whether or not biased gene

conversion is considered (**Figure 3A** vs. **Figure 3B**). To formally evaluate the effects of

population size, recombination, *gBGC*, and their interactions on mutation load, we employed

a linear mixed-effects model (**Supplementary Table 3**). Chromosome and population

identity were included as random effects, as they accounted for a significant proportion of

additional variance. The best-fitting model (**Table 1, Supplementary Figure 7**) indicates a

strong influence of all three factors, summarized as follows: (i) mutation load was

substantially higher in small populations, (ii) was further increased for mutations affected by

gBGC, and (iii) was markedly reduced in genomic regions with high recombination rates.

The model also predicts that the contribution of *gBGC* to deleterious variation is greater in

small populations (i.e., a significant N_e x *gBGC* interaction), with island populations showing

a 9.6% higher proportion of deleterious mutations compared to a 6.4% increase in mainland

populations. Nonetheless, the relative increase due to *gBGC* remained consistent across

population sizes, with island and mainland populations exhibiting a similar fold-increase in

deleterious mutations (1.44-fold vs. 1.46-fold, respectively).

The data further allowed us to assess the relative rates by which the two opposing,

recombination-dependent processes of purging and *gBGC* affect mutation load. If both

processes took place with the same intensity, then a doubling in the increase of deleterious

mutations by *gBGC* should be fully offset by a proportional increase in purging efficacy. The

relative contribution of both processes may further depend on the species' N_e (**Figure 1**). We

found that the purging efficacy of recombination on mutation load was reduced in small

populations for the mutational spectrum affected by *gBGC*. In other words, small populations

suffer from a disproportionate introduction of deleterious [A,T]->[C,G] mutations that is not

fully offset by purging efficacy in regions of high recombination (**Figure 3C**). This effect

(three-way interaction *r* x N_e x *gBGC*) was weak but statistically supported when using a

binary measure of population size (island/mainland) (**Table 1**, ΔAIC to 2[nd] best model

=2.229), but not when parametrizing over a continuum of coalescent based estimates of N_e

(**Supplementary Table 3**). The three-way interaction using the island/mainland contrast still

had support when controlling for phylogenetic dependence using a dataset limited to the 19

species in a Bayesian approach (ΔAIC to 2^{nd} best model =1.492, p-value analogue=0.059, see

Supplementary Table 3).

Discussion

In this study, we estimated the distribution of fitness effects for 24 populations from 19 avian

species including representatives from across the major songbird clades (Corvoidea,

Muscicapoidea, Passeroidea, Sylvoidea) (Jønsson and Fjeldså 2006). Oscine songbirds are

the most speciose avian clade arising broadly around the Cretaceous-Palaeogene boundary

some 66 million years ago (Sibley and Ahlquist 1990; Barker et al. 2004; Jarvis et al. 2014).

The study thus claims some generality at the higher end of evolutionary timescales compared

to previous comparative work in the area (within species: (Eyre-Walker et al. 2006; Slotte et

al. 2010; Deinum et al. 2015; Castellano et al. 2018), genera: (Kutschera et al. 2020),

families: (Castellano et al. 2019), and beyond (Chen et al. 2017; Leroy et al. 2021; Chen et al.

2022)). With the exception of few closely related species, the chosen species in this study are

overall phylogenetically distant enough as not to share ancestral polymorphism (isolation > 9

Ne generations Hudson et al. 2002; Mugal et al. 2020). As some evolutionary dependency

may remain, we additionally fitted a model taking phylogenetic contrasts into account.

Focusing on the mildly deleterious proportion of the DFE, we tested the effects of N_e,

recombination rate, *gBGC*, and their interaction, on mutation load. The effective population

size, as parameterized by species distribution (island/mainland) or coalescent-based

estimates, was a strong predictor. Coalescent effective population sizes of island populations

were approximately half of the mainland populations. In accordance with the central tenet of

the nearly-neutral theory and previous studies, mutation load was significantly higher in

small effective populations (Ohta 1992; Charlesworth 2009; Agrawal and Whitlock 2012;

Kutschera et al. 2020; Leroy et al. 2021). Our model predicts this difference to translate into

an increase in the genome-wide mutation load of approximately 1.6-fold on islands (cf.

(Castellano et al. 2018) for a similar estimate considering local genomic N_e variation within

the *Drosophila* genome). While the difference in mutation load on islands is mediated by

small population size, a rather discrete change between mainland and island populations may

also suggest additional factors – such as bottlenecks or different positions in the fitness

landscape (see below) – that are not fully accounted for by the coalescent-based N_e estimates

(**Supplementary Figure 4**). Moreover, single estimates of long-term N_e are likely not

capturing more complex demographic histories including repeated changes in population size

with contributions of gene flow. These are likely only controlled in part by the model

estimating the DFE from the SFS.

Despite these limitations, these results add to the evidence that species living on islands

accumulate mildly deleterious mutations more readily than their mainland counterparts

(Johnson and Seger 2001; Woolfit and Bromham 2005; Kutschera et al. 2020), making them

more vulnerable to extinction in the long term (Frankham 1998). Our results further suggest

that mutation load is unevenly distributed across the genome. In accordance with relatively

few previous studies touching upon the subject (Gossmann et al. 2011; Leroy et al. 2021;

Chase et al. 2024), we find a clear reduction of mutation load with an increase in

recombination. In our model, the consistent ~ 5-fold range of recombination rate across avian

chromosomes (Backström et al. 2010; Kawakami et al. 2014; Bascón-Cardozo et al. 2024,

this study) translates to a ~1.7-fold change in mutation load. The order of this change is

comparable to the effect of N_e-change between island and mainland. Importantly, however,

the effect of recombination was independent of N_e (no statistical interaction N_e x r)

suggesting the same relative purging efficiency between chromosomes regardless of

population size. This result is in conflict with conclusions from Leroy et al. (2021). While

they also find reduced mutation load in genomic regions of high recombination, the beneficial

effect of recombination was stronger in island populations. This disconnect may be explained

by a combination of factors like using GC content at the third position of the codon (GC3) as

proxy for recombination, a different set of species and presence of population structure in

several of their datasets that can influence the DFE (Kutschera et al. 2020) and which we

removed for this study (**Supplementary Figure 1**).

In addition to N_e and intragenomic heterogeneity of recombination rate, the quality of

mutations also affected mutation load. When including mutations subject to *gBGC* the

genome-wide proportion of mildly deleterious variation increased by ~1.5 fold, both in island

and mainland populations. An increase of mutation load in this fraction of mutations is

expected, as they are replenished through the segregational bias mimicking the effect of

positive selection (Berglund et al. 2009). Yet its interaction with recombination and overall

population size remains largely unclear. Since *gBGC* relies on double-strand breaks

associated with increased levels of recombination, *gBGC* mediated mutation load is expected

to be more pronounced in regions of high recombination (Duret and Galtier 2009; Kostka et

al. 2012; Pouyet et al. 2018). We thus expected to see relatively more deleterious mutations

in regions of high recombination with *gBGC* than in those with lower recombination rates

(flatter slope for DFE with *gBGC*, **Figure 1**). We observed the effect only in island

populations, while the effect may be too slight to pick up in the mainland with an overall

increased efficacy of purifying selection. Alternatively, the relative increase in *gBGC*-

contributed mutation load in small island populations, may reflect the reduced overall

intensity of *gBGC*. Genes experiencing high levels of *gBGC* in regions of high recombination

tend to incur a fitness disadvantage (Joseph 2024), but can still accumulate deleterious AT-

GC mutations in large populations where the *gBGC* pressure is high. In small populations,

reduced *gBGC* pressure might facilitate segregation of GC→AT back mutations in regions of

high recombination. The outcome depends on the relative effect of recombination on *gBGC*

and selection efficacy.

Returning to the hypotheses of the introduction, we conclude that mutation load 1) is elevated

in (island) populations with low N_e, 2) is reduced in highly-recombining regions, and 3) is

further increased for the mutational spectrum subject to *gBGC*. All three effects are of

comparable magnitude across the relevant biological range of variation in our sample.

Relative rates of the two recombination-mediated, opposing processes of purging and *gBGC*

differ in populations with low N_e. Only few studies have hitherto explored these interactions

(Jiang et al. 2025). In our view, they deserve more empirical and theoretical attention, as the

fundamentally affect the overall magnitude and genomic distribution of mutation load

segregating in natural populations.

In addition, non-negligible variance for the species random effect is consistent with

additional components affecting a population's' DFE, such as its current position in the

fitness landscape (Lanfear et al. 2014). For species at or close to the fitness optimum, most

mutations can be expected to have deleterious effects, whereas in species further removed

from their fitness optimum, a subset of mutations may have beneficial effects. This may also

impact our proxy for N_e. While geographic restriction has been suggested as a valuable proxy

for the effective population sizes (Kutschera et al. 2020; Leroy et al. 2021), island or

mainland populations may qualitatively differ in their position in the fitness landscape and

thereby in the mutation load they experience.

Overall, this study highlights that assuming a homogeneous DFE across the genome is a

substantial oversimplification of the complex interplay among evolutionary forces shaping

genomic diversity. As a consequence, genetic risk factors are non-homogeneously distributed

across the genome (Kardos et al. 2021). Emerging medical, population and conservation

genomic studies screening for deleterious mutations and their fitness effects (Bertorelle et al.

2022; Hoffman et al. 2024; Chen et al. 2025) may benefit from taking the interaction between

recombination, *gBGC* and N_e into account.

Data Availability

575 Whole genome resequencing data are available through the National Center of Biotechnology

576 Information (NCBI). Accession numbers are provided in **Supplementary Table 1**. Analysis

577 scripts, processed *vcf* files and plots of the SFS for individual chromosomes for each

578 population are available at the zenodo repository (doi: 10.5281/zenodo.17822666).

Acknowledgements

581 We thank Benoit Nabholz and Verena Kutschera who provided access to raw data and *vcf*

582 files. We also acknowledge valuable feed-back from Dirk Metzler on the Bayesian statistics.

Study Funding

585 Funding was provided to J.B.W.W. by the European Research Council (ERCStG-336536

586 FuncSpecGen), the Swedish Research Council Vetenskapsrådet (621-2013-4510), the Knut

587 and Alice Wallenberg Foundation and LMU Munich. HPC computing was performed on the

588 BioHPC hosted at Leibniz Rechenzentrum Munich funded by the German Research

589 Foundation (grant INST 86/2050-1 FUGG).

Conflict of Interest

592 We declare no conflict of interest.

Author Contributions

595 JBWW and FB conceived of the study. FB conducted all bioinformatic analyses, JBWW

596 conducted the statistical analyses. JBWW and FB wrote the manuscript.

References

Agrawal AF, Whitlock MC. 2012. Mutation Load: The Fitness of Individuals in Populations
Where Deleterious Alleles Are Abundant. Annu Rev Ecol Evol Syst. 43(1):115–135.
doi:10.1146/annurev-ecolsys-110411-160257.

Akashi H, Osada N, Ohta T. 2012. Weak selection and protein evolution. Genetics.
192(1):15–31. doi:10.1534/genetics.112.140178.

Backström N, Forstmeier W, Schielzeth H, Mellenius H, Nam K, Bolund E, Webster MT,
Öst T, Schneider M, Kempenaers B, et al. 2010. The recombination landscape of the zebra
finch Taeniopygia guttata genome. Genome Res. 20(4):485–495. doi:10.1101/gr.101410.109.

Barker FK, Cibois A, Schikler P, Feinstein J, Cracraft J. 2004. Phylogeny and diversification
of the largest avian radiation. Proc Natl Acad Sci U S A. 101(30):11040–11045.
doi:10.1073/pnas.0401892101.

Bascón-Cardozo K, Bours A, Manthey G, Durieux G, Dutheil JY, Pruisscher P, Odenthal-
Hesse L, Liedvogel M. 2024. Fine-Scale Map Reveals Highly Variable Recombination Rates
Associated with Genomic Features in the Eurasian Blackcap. Genome Biol Evol.
16(1):evad233. doi:10.1093/gbe/evad233.

Bates D, Mächler M, Bolker B, Walker S. 2015. Fitting Linear Mixed-Effects Models Using
lme4. J Stat Softw. 67(1). doi:10.18637/jss.v067.i01. [accessed 2016 Aug 22].
http://www.jstatsoft.org/v67/i01/.

Berglund J, Pollard KS, Webster MT. 2009. Hotspots of biased nucleotide substitutions in
human genes. Plos Biol. 7:e1000026(1).

Bertorelle G, Raffini F, Bosse M, Bortoluzzi C, Iannucci A, Trucchi E, Morales HE, van
Oosterhout C. 2022. Genetic load: genomic estimates and applications in non-model animals.
Nat Rev Genet. 23(8):492–503. doi:10.1038/s41576-022-00448-x.

Boitard S, Arredondo A, Chikhi L, Mazet O. 2022. Heterogeneity in effective size across the
genome: effects on the inverse instantaneous coalescence rate (IICR) and implications for

demographic inference under linked selection. Genetics. 220(3):iyac008.
doi:10.1093/genetics/iyac008.

Bolívar P, Guéguen L, Duret L, Ellegren H, Mugal CF. 2019. GC-biased gene conversion
conceals the prediction of the nearly neutral theory in avian genomes. Genome Biol. 20(1):1–
13. doi:10.1186/s13059-018-1613-z.

Bolívar P, Mugal CF, Rossi M, Nater A, Wang M, Dutoit L, Ellegren H. 2018. Biased
Inference of Selection Due to GC-Biased Gene Conversion and the Rate of Protein Evolution
in Flycatchers When Accounting for It. Mol Biol Evol. 35(10):2475–2486.
doi:10.1093/molbev/msy149.

Bourgeois YXC, Bertrand JAM, Delahaie B, Holota H, Thébaud C, Milá B. 2020.
Differential divergence in autosomes and sex chromosomes is associated with intra-island
diversification at a very small spatial scale in a songbird lineage. Mol Ecol. 29(6):1137–1153.
doi:10.1111/mec.15396.

Burri R, Nater A, Kawakami T, Mugal CF, Olason PI, Smeds L, Suh A, Dutoit L, Bureš S,
Garamszegi LZ, et al. 2015. Linked selection and recombination rate variation drive the
evolution of the genomic landscape of differentiation across the speciation continuum of
Ficedula flycatchers. Genome Res. 25:1656–1665.

Castellano D, James J, Eyre-Walker A. 2018. Nearly Neutral Evolution across the Drosophila
melanogaster Genome. Mol Biol Evol. 35(11):2685–2694. doi:10.1093/molbev/msy164.

Castellano D, Macià MC, Tataru P, Bataillon T, Munch K. 2019. Comparison of the Full
Distribution of Fitness Effects of New Amino Acid Mutations Across Great Apes. Genetics.
213(3):953–966. doi:10.1534/genetics.119.302494.

Charlesworth B. 2009. Fundamental concepts in genetics: Effective population size and
patterns of molecular evolution and variation. Nat Rev Genet. 10(3):195–205.

Charlesworth B, Charlesworth D. 2000. The degeneration of Y chromosomes. Philos Trans R
Soc Lond Ser B-Biol Sci. 355(1403):1563–1572.

Chase MA, Vilcot M, Mugal CF. 2024. The role of recombination dynamics in shaping
signatures of direct and indirect selection across the   Ficedula   flycatcher
genome. Proc R Soc B Biol Sci. 291(2015). doi:10.1098/rspb.2023.2382.

Chen J, Bataillon T, Glémin S, Lascoux M. 2022. What does the distribution of fitness effects
of new mutations reflect? Insights from plants. New Phytol. 233(4):1613–1619.
doi:10.1111/nph.17826.

Chen J, Glémin S, Lascoux M. 2017. Genetic Diversity and the Efficacy of Purifying
Selection across Plant and Animal Species. Mol Biol Evol. 34(6):1417–1428.
doi:10.1093/molbev/msx088.

Chen RS, Soulsbury CD, Hench K, van Oers K, Hoffman JI. 2025. Predicted deleterious
mutations reveal the genetic architecture of male reproductive success in a lekking bird. Nat
Ecol Evol. 9(10):1924–1937. doi:10.1038/s41559-025-02802-8.

Comeron JM, Williford A, Kliman RM. 2008. The Hill-Robertson effect: evolutionary
consequences of weak selection and linkage in finite populations. Heredity. 100(1):19–31.
doi:10.1038/sj.hdy.6801059.

Corcoran P, Gossmann TI, Barton HJ, Great Tit HapMap Consortium, Slate J, Zeng K. 2017.
Determinants of the Efficacy of Natural Selection on Coding and Noncoding Variability in
Two Passerine Species. Genome Biol Evol. 9(11):2987–3007. doi:10.1093/gbe/evx213.

Deinum EE, Halligan DL, Ness RW, Zhang Y-H, Cong L, Zhang J-X, Keightley PD. 2015.
Recent Evolution in Rattus norvegicus Is Shaped by Declining Effective Population Size.
Mol Biol Evol. 32(10):2547–2558. doi:10.1093/molbev/msv126.

Duret L, Galtier N. 2009. Biased Gene Conversion and the Evolution of Mammalian
Genomic Landscapes. Annu Rev Genomics Hum Genet. 10(1):285–311.
doi:10.1146/annurev-genom-082908-150001.

Dussex N, Morales HE, Grossen C, Dalén L, Van Oosterhout C. 2023. Purging and
accumulation of genetic load in conservation. Trends Ecol Evol. 38(10):961–969.
doi:10.1016/j.tree.2023.05.008.

Dutoit L, Vijay N, Mugal CF, Bossu CM, Burri R, Wolf JBW, Ellegren H. 2017. Covariation
in levels of nucleotide diversity in homologous regions of the avian genome long after
completion of lineage sorting. Proc R Soc B. 284(1849):20162756.
doi:10.1098/rspb.2016.2756.

Ellegren H. 2010. Evolutionary stasis: the stable chromosomes of birds. Trends Ecol Evol.
25(5):283–291. doi:10.1016/j.tree.2009.12.004.

Eyre-Walker A, Keightley PD. 2007. The distribution of fitness effects of new mutations. Nat
Rev Genet. 8(8):610–618. doi:10.1038/nrg2146.

Eyre-Walker A, Woolfit M, Phelps T. 2006. The Distribution of Fitness Effects of New
Deleterious Amino Acid Mutations in Humans. Genetics. 173(2):891–900.
doi:10.1534/genetics.106.057570.

Felsenstein J. 1974. The evolutionary advantage of recombination. Genetics. 78(2):737–756.

Fox J, Weisberg S. 2019. An R Companion to Applied Regression. 3rd ed. Thousand Oaks
CA: Sage. https://socialsciences.mcmaster.ca/jfox/Books/Companion/.

Frankham R. 1998. Inbreeding and extinction: island populations. Conserv Biol. 12(3):665–
675. doi:10.1111/j.1523-1739.1998.96456.x.

Galtier N. 2016. Adaptive Protein Evolution in Animals and the Effective Population Size
Hypothesis. PLoS Genet. 12(1):e1005774. doi:10.1371/journal.pgen.1005774.

Galtier N, Piganeau G, Mouchiroud D, Duret L. 2001. GC-Content Evolution in Mammalian
Genomes: The Biased Gene Conversion Hypothesis. Genetics. 159(2):907–911.
doi:10.1093/genetics/159.2.907.

Galtier N, Rousselle M. 2020. How Much Does Ne Vary Among Species? Genetics.
216(2):559–572. doi:10.1534/genetics.120.303622.

Garrison E, Marth G. 2012. Haplotype-based variant detection from short-read sequencing.
Report No.: 1207.3907. [accessed 2013 June 11]. http://arxiv.org/abs/1207.3907.

Gossmann TI, Woolfit M, Eyre-Walker A. 2011. Quantifying the variation in the effective
population size within a genome. Genetics. 189(4):1389–1402.
doi:10.1534/genetics.111.132654.

Hadfield JD. 2010. MCMC Methods for Multi-Response Generalized Linear Mixed Models:
The MCMCglmm R Package. J Stat Softw. 33:1–22. doi:10.18637/jss.v033.i02.

Hill WG, Robertson A. 1966. The effect of linkage on limits to artificial selection. Genet Res.
8(3):269–294.

Hoffman JI, Vendrami DLJ, Hench K, Chen RS, Stoffel MA, Kardos M, Amos W,
Kalinowski J, Rickert D, Köhrer K, et al. 2024. Genomic and fitness consequences of a near-
extinction event in the northern elephant seal. Nat Ecol Evol. 8(12):2309–2324.
doi:10.1038/s41559-024-02533-2.

Hudson RR, Coyne JA, Huelsenbeck J. 2002. Mathematical consequences of the genealogical
species concept. Evolution. 56(8):1557–1565. doi:10.1554/0014-
3820(2002)056%5B1557:MCOTGS%5D2.0.CO;2.

Jarvis ED, Mirarab S, Aberer AJ, Li B, Houde P, Li C, Ho SYW, Faircloth BC, Nabholz B,
Howard JT, et al. 2014. Whole-genome analyses resolve early branches in the tree of life of
modern birds. Science. 346(6215):1320–1331. doi:10.1126/science.1253451.

Jiang J, Chen J-F, Li X-T, Wang L, Mao J-F, Wang B-S, Guo Y-L. 2025. Incorporating
genetic load contributes to predicting Arabidopsis thaliana's response to climate change. Nat
Commun. 16(1):2752. doi:10.1038/s41467-025-58021-z.

Johnson JB, Omland KS. 2004. Model selection in ecology and evolution. Trends Ecol Evol.
19(2):101–108. doi:10.1016/j.tree.2003.10.013.

Johnson KP, Seger J. 2001. Elevated Rates of Nonsynonymous Substitution in Island Birds.
Mol Biol Evol. 18(5):874–881. doi:10.1093/oxfordjournals.molbev.a003869.

Jønsson KA, Fjeldså J. 2006. A phylogenetic supertree of oscine passerine birds (Aves:
Passeri). Zool Scr. 35(2):149–186. doi:10.1111/j.1463-6409.2006.00221.x.

Joseph J. 2024. Increased Positive Selection in Highly Recombining Genes Does not
Necessarily Reflect an Evolutionary Advantage of Recombination. Mol Biol Evol.
41(6):msae107. doi:10.1093/molbev/msae107.

Kardos M, Armstrong EE, Fitzpatrick SW, Hauser S, Hedrick PW, Miller JM, Tallmon DA,
Funk WC. 2021. The crucial role of genome-wide genetic variation in conservation. Proc
Natl Acad Sci. 118(48):e2104642118. doi:10.1073/pnas.2104642118.

Kawakami T, Smeds L, Backström N, Husby A, Qvarnström A, Mugal CF, Olason P,
Ellegren H. 2014. A high-density linkage map enables a second-generation collared
flycatcher genome assembly and reveals the patterns of avian recombination rate variation
and chromosomal evolution. Mol Ecol. 23(16):4035–4058. doi:10.1111/mec.12810.

Keightley PD, Eyre-Walker A. 2007. Joint Inference of the Distribution of Fitness Effects of
Deleterious Mutations and Population Demography Based on Nucleotide Polymorphism
Frequencies. Genetics. 177(4):2251–2261. doi:10.1534/genetics.107.080663.

Keightley PD, Lynch M. 2003. Toward a Realistic Model of Mutations Affecting Fitness.
Evolution. 57(3):683–685. doi:10.1111/j.0014-3820.2003.tb01561.x.

Kimura M. 1983. The neutral theory of molecular evolution. Cambridge (UK): Cambridge
University Press.

Kimura M, Maruyama T, Crow JF. 1963. The mutation load in small populations. Genetics.
48(10):1303–1312. doi:10.1093/genetics/48.10.1303.

Kostka D, Hubisz MJ, Siepel A, Pollard KS. 2012. The Role of GC-Biased Gene Conversion
in Shaping the Fastest Evolving Regions of the Human Genome. Mol Biol Evol. 29(3):1047–
1057. doi:10.1093/molbev/msr279.

Kutschera VE, Poelstra JW, Botero-Castro F, Dussex N, Gemmell NJ, Hunt GR, Ritchie MG,
Rutz C, Wiberg RAW, Wolf JBW. 2020. Purifying selection in corvids is less efficient on
islands. Mol Biol Evol. 37(2):469–474. doi:10.1093/molbev/msz233.

Lande R. 1994. Risk of population extinction from fixation of new deleterious mutations.
Evolution. 48(5):1460–1469. doi:10.2307/2410240.

Lanfear R, Kokko H, Eyre-Walker A. 2014. Population size and the rate of evolution. Trends
Ecol Evol. 29(1):33–41. doi:10.1016/j.tree.2013.09.009.

Leroy T, Rousselle M, Tilak M-K, Caizergues AE, Scornavacca C, Recuerda M, Fuchs J,
Illera JC, De Swardt DH, Blanco G, et al. 2021. Island songbirds as windows into evolution
in small populations. Curr Biol. 31(6):1303-1310.e4. doi:10.1016/j.cub.2020.12.040.

Lynch M, Conery J, Burger R. 1995. Mutation Accumulation and the Extinction of Small
Populations. Am Nat. 146(4):489–518. doi:10.1086/285812.

Manichaikul A, Mychaleckyj JC, Rich SS, Daly K, Sale M, Chen W-M. 2010. Robust
relationship inference in genome-wide association studies. Bioinformatics. 26(22):2867–
2873. doi:10.1093/bioinformatics/btq559.

Marais GAB. 2007. The Hill-Robertson effects extend from nucleus to mitochondria.
Heredity. 99:357–358.

McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K,
Altshuler D, Gabriel S, Daly M, et al. 2010. The Genome Analysis Toolkit: A MapReduce
framework for analyzing next-generation DNA sequencing data. Genome Res. 20(9):1297–
1303. doi:10.1101/gr.107524.110.

McTavish EJ, Gerbracht JA, Holder MT, Iliff MJ, Lepage D, Rasmussen PC, Redelings BD,
Sánchez Reyes LL, Miller ET. 2025. A complete and dynamic tree of birds. Proc Natl Acad
Sci. 122(18):e2409658122. doi:10.1073/pnas.2409658122.

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A,
Lanfear R. 2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference
in the Genomic Era. Mol Biol Evol. 37(5):1530–1534. doi:10.1093/molbev/msaa015.

Mugal CF, Kutschera VE, Botero-Castro F, Wolf JBW, Kaj I. 2020. Polymorphism data
assist estimation of the nonsynonymous over synonymous fixation rate ratio ω for closely
related species. Mol Biol Evol. 37(1):260–279. doi:10.1093/molbev/msz203.

Muller HJ. 1950. Our load of mutations. Am J Hum Genet. 2(2):111–176.

Ohta T. 1992. The Nearly Neutral Theory of Molecular Evolution. Annu Rev Ecol Evol Syst.
23(1):263–286. doi:10.1146/annurev.es.23.110192.001403.

Peart CR, Tusso S, Pophaly SD, Botero-Castro F, Wu C-C, Aurioles-Gamboa D, Baird AB,
Bickham JW, Forcada J, Galimberti F, et al. 2020. Determinants of genetic variation across
eco-evolutionary scales in pinnipeds. Nat Ecol Evol. 4(8):1095–1104. doi:10.1038/s41559-
020-1215-5.

Pouyet F, Aeschbacher S, Thiéry A, Excoffier L. 2018. Background selection and biased
gene conversion affect more than 95% of the human genome and bias demographic
inferences. Veeramah K, Wittkopp PJ, Gronau I, editors. eLife. 7:e36317.
doi:10.7554/eLife.36317.

Prentout D, Bykova D, Hoge C, Hooper DM, McDiarmid CS, Wu F, Griffith SC, Manuel M
de, Przeworski M. 2025. Germline mutation rates and fine-scale recombination parameters in
zebra finch. PLOS Genet. 21(4):e1011661. doi:10.1371/journal.pgen.1011661.

Romiguier J, Gayral P, Ballenghien M, Bernard A, Cahais V, Chenuil A, Chiari Y, Dernat R,
Duret L, Faivre N, et al. 2014. Comparative population genomics in animals uncovers the
determinants of genetic diversity. Nature. 515(7526):261–263. doi:10.1038/nature13685.

Schiffels S, Durbin R. 2014. Inferring human population size and separation history from
multiple genome sequences. Nat Genet. 46(8):919–925. doi:10.1038/ng.3015.

Sibley CG, Ahlquist JE. 1990. Phylogeny and Classification of the Birds: A Study in
Molecular Evolution. New Haven (CT): Yale University Press.

Singhal S, Leffler EM, Sannareddy K, Turner I, Venn O, Hooper DM, Strand AI, Li Q,
Raney B, Balakrishnan CN, et al. 2015. Stable recombination hotspots in birds. Science.
350(6263):928–932. doi:10.1126/science.aad0843.

Slotte T, Foxe JP, Hazzouri KM, Wright SI. 2010. Genome-Wide Evidence for Efficient
Positive and Purifying Selection in Capsella grandiflora, a Plant Species with a Large
Effective Population Size. Mol Biol Evol. 27(8):1813–1821. doi:10.1093/molbev/msq062.

Smeds L, Qvarnström A, Ellegren H. 2016. Direct estimate of the rate of germline mutation
in a bird. Genome Res. 26(9):1211–1218. doi:10.1101/gr.204669.116.

Soulé ME, Mills LS. 1998. No need to isolate genetics. Science. 282(5394):1658–1659.
doi:10.1126/science.282.5394.1658.

Stapley J, Birkhead TR, Burke T, Slate J. 2010. Pronounced inter- and intrachromosomal
variation in linkage disequilibrium across the zebra finch genome. Genome Res. 20(4):496–
502. doi:10.1101/gr.102095.109.

Tataru P, Mollion M, Glémin S, Bataillon T. 2017. Inference of Distribution of Fitness
Effects and Proportion of Adaptive Substitutions from Polymorphism Data. Genetics.
207(3):1103–1119. doi:10.1534/genetics.117.300323.

Vijay N, Bossu CM, Poelstra JW, Weissensteiner MH, Suh A, Kryukov AP, Wolf JBW.
2016. Evolution of heterogeneous genome differentiation across multiple contact zones in a
crow species complex. Nat Commun. 7:13195. doi:10.1038/ncomms13195.

Vijay N, Weissensteiner M, Burri R, Kawakami T, Ellegren H, Wolf JBW. 2017.
Genomewide patterns of variation in genetic diversity are shared among populations, species
and higher-order taxa. Mol Ecol. 26(16):4284–4295. doi:10.1111/mec.14195.

Warmuth VM, Weissensteiner MH, Wolf JBW. 2022. Accumulation and ineffective silencing
of transposable elements on an avian W Chromosome. Genome Res. 32(4):671–681.
doi:10.1101/gr.275465.121.

Webster MT, Hurst LD. 2012. Direct and indirect consequences of meiotic recombination:
implications for genome evolution. Trends Genet. 28(3):101–109.
doi:10.1016/j.tig.2011.11.002.

Williamson RJ, Josephs EB, Platts AE, Hazzouri KM, Haudry A, Blanchette M, Wright SI.
2014. Evidence for Widespread Positive and Negative Selection in Coding and Conserved
Noncoding Regions of Capsella grandiflora. PLOS Genet. 10(9):e1004622.
doi:10.1371/journal.pgen.1004622.

Woolfit M, Bromham L. 2005. Population size and molecular evolution on islands. Proc R
Soc B Biol Sci. 272(1578):2277–2282. doi:10.1098/rspb.2005.3217.

Zuur AF, Ieno EN, Walker N, Saveliev AA, Smith GM. 2009. Mixed effects models and
extensions in ecology with R. New York, NY: Springer (Statistics for Biology and Health).

Figure captions

Figure 1 | Theoretical considerations on mutation load. The curves outlined in black

illustrate a mock example of a genome-wide distribution of fitness effects (DFE) at a given

point in the fitness landscape. We here only consider the fraction of mutations with selection

coefficient s<0 reducing fitness (left of the vertical dashed line marking s=0). The efficacy of

selection in removing those crucially depends on the effective population size N_e (left

column) and the recombination rate r (bottom row) changing the fraction of mutations

behaving as strongly deleterious (red: $N_e s < -10$), deleterious (orange: $N_e s \in (-10;-1]$), or

mildly deleterious (grey: $N_e s \in (-1;0]$). The latter are the focus of this study. *Central figure,*

solid lines: The y-axis shows the fraction of all mutations falling into the mildly deleterious

category $N_e s \in (-1;0]$. This corresponds to the mutation load which is expected to be highest

in populations with small N_e. Mutation load is further expected to be elevated in genomic

regions experiencing low recombination rates (x-axis). Line colours correspond to N_e

categories small (light blue), intermediate (blue) and large (dark blue). *gBGC* is a

recombination-associated process expected to increase mutation load overall, and particularly

so in regions of high recombination *(dashed lines)*. Difference in slope between solid

(without *gBGC*) and dashed lines (with *gBGC*) is expected if the relative rate by which

recombination introduces mildly deleterious mutations through *gBGC* differs from the rate by

which recombination increases the efficacy of selection removing these. Such difference in

relative rates of *gBGC* and purging efficacy may further interact with the effective population

size (length of arrows) mediating the overall and/or recombination-dependent contribution of

gBGC to mutation load.

Figure 2 | Relationship between recombination rate [cM/Mb], chromosome length [Mb]

and GC content [%] in Eurasian crows, *Corvus (corone) ssp*. A near 5-fold difference in

recombination rate is closely associated with chromosome length which strongly covaries

with GC content (R^2=0.90, p<0.001). In small, highly recombining chromosomes GC content

is increased by up to 15% reflecting an evolutionary history of *gBGC*. The heterogenous

recombination landscape is conserved across the avian species investigated in this study.

872
873
874
Figure 3. Relationship between recombination rate and mutation load estimated as the

fraction of mildly deleterious mutations with $N_e s \in [-1;0)$. Correlation between

recombination rate and mutation load A) excluding and B) including mutations subject to

gBGC (right). Each regression line represents a population sample of a species differing in

effective population sizes (N_e) as is indicated by the blue colour gradient on a logarithmic

scale. Dashed and solid lines illustrate island and mainland populations, respectively. C) as

above, but the y-axis displays the difference between estimates of mutation load with and

without mutations subject to *gBGC*. Note that the slopes differ between island and mainland

populations indicating that relative rates of the two recombination-mediated processes, *gBGC*

and purging efficacy, may interact with the effective population size of a species.

Table 1: Summary of the four best statistical models and the null model exploring the

relationship between the mutation load quantified as the proportion of mildly deleterious

alleles (DFE) with recombination rate (r), effective population size (Ne) and GC-biased gene

conversion (*gBGC*). *r* was here approximated by chromosome length, N_e was here

approximated by whether a population was restricted to an island (yes/no), *gBGC* was

estimated for all sites (with *gBGC*) and for a subset of sites excluding weak-to-strong

mutations (without gBCG). For other proxies of r and N_e and the full set of models with

parameter estimates see **Supplementary Table 3**. Model selection is based on Akaike's

Information criterion (AIC) with models by $\Delta AIC_c < 2$ are considered supported (best model in

bold). In addition, thresholds of Type 1 error probabilities of $p < 0.001$ (***), $p < 0.01$ (**),

$p < 0.05$ (*) and $p > 0.05$ (n.s) are indicated for all fixed effects and their interactions included

in a specific model.

Model	r [cM/Mb]	Ne [yes/no]	*gBGC* [with/without]	Ne*gBGC	r*Ne	r*gBGC	r*Ne*gBGC	k	AIC	ΔAIC
10	***	***	***	***	n.s.	n.s.	*	**13**	**-4479.5754**	**0**
9	***	***	***	***	n.s.	n.s.	-	12	-4477.2763	2.299
8	***	***	***	***	-	-	-	10	-4476.8571	2.718
7	***	***	***	-	-	-	-	11	-4440.2616	39.31
NULL	-	-	-	-	-	-	-	6	-3555.9041	923.671