Single cell long read genotyping of transcripts reveals discrete mechanisms of
clonal evolution in post-MPN AML.

Julian Grabek[1,2*], Jasmin Straube[1,2*], Leanne Cooper[1], Rohit Haldar[1], Ranran Zhang[1,2],
Inken Dulige[1,3], Matthew Barker[1,2], Will Gatehouse[1], Helen Christensen[1], Gerlinda
Amor[1], Victoria Y. Ling[1,2,4], Caroline McNamara[5], David M. Ross[6,7], Andrew Perkins[8,9],
Megan J. Bywater[1,2φ] & Steven W. Lane[1,2,5φ]

Affiliations:

[1]Cancer Research Program, QIMR Berghofer Medical Research Institute, Brisbane,
QLD, Australia.
[2] The University of Queensland, St Lucia, Brisbane, QLD, Australia.
[3]University of Lübeck, Lübeck, Schleswig-Holstein, Germany.
[4]Department of Haematology, Princess Alexandra Hospital, Brisbane, Australia.
[5]Cancer Care Services, Royal Brisbane and Women's Hospital, Brisbane, QLD,
Australia.
[6]Department of Haematology, Royal Adelaide Hospital, Adelaide, SA, Australia.
[7]Acute Leukaemia Laboratory, Centre for Cancer Biology, SA Pathology and University
of South Australia, Adelaide, SA, Australia.
[8]Alfred Hospital, Melbourne, Victoria, Australia.
[9]Australian Centre for Blood Diseases, Monash University, Melbourne, Victoria,
Australia.
21

*these authors contributed equally to the study

φco-corresponding authors:

megan.bywater@qimrb.edu.au

steven.lane@qimrb.edu.au

Words 3967/4000

Abstract (194/250)
Myeloproliferative neoplasms (MPNs) are caused by acquired mutations in hematopoietic
stem and progenitor cells (HSPCs). The acquisition of additional mutations like *TP53* and
the overall mutational burden influence a patient's risk of disease progression toward
lethal post-MPN acute myeloid leukemia (AML).

Recent technological advancements in linking single-cell gene expression with genotype
have improved our understanding of tumor heterogeneity. However, current

methodologies have limitations in simultaneously genotyping low-expression genes (such

as *JAK2*) alongside other pathogenic loci.

To address this, we developed a novel <u>lo</u>ng read genotyping pipeline of cDNA <u>tr</u>anscripts

called LOTR-Seq, which can genotype the full length of expressed transcripts of 30 genes

at once. Using LOTR-Seq, we genotyped HSPCs at the *JAK2*V617 locus in 9,075 single

cells from eight patients with chronic phase MPN (CP-MPN) and in 5,016 cells from four

patients with post-MPN AML. We then linked the mutations to the single cell transcriptome

of 29,712 *JAK2*V617F-driven CP-MPN cells and 16,895 post-MPN AML cells.

In our analysis of post-MPN AMLs, we identified nine mutated loci across six genes

(*JAK2, IDH1/2, TP53, SRSF2, U2AF1*) and linked these mutations to specific

transcriptional phenotypes. Overall, LOTR-Seq provides novel insights into the evolution

of post-MPN AML.

Introduction

Myeloproliferative Neoplasms (MPN) are clonal blood cancers characterized by the

overproduction of mature, functional myeloid elements. MPNs are caused by driver

mutations within the hematopoietic stem and progenitor (HSPC) compartment[1], with the

most common being *JAK2*V617F[2]. *JAK2*V617F gives rise to a clonal stem cell population[3]

and results in pathology mediated by mature myeloid cells. Both healthy and MPN HSPCs

exhibit both transcriptional evidence of myeloid and lymphoid lineage priming[4] and self-

renewal capacity[5]. Exactly how this clonal outgrowth affects the transcriptional diversity

of the HSPC compartment is unclear.

In contrast to MPN, acute myeloid leukemias (AML) are characterized by the

overproduction of immature myeloid cells. Genetic complexity is a common feature of

*JAK2*V617F post-MPN AML, with patients having multiple additional mutations identified

at AML transformation[6]. Disease progression from chronic phase MPN (CP-MPN) to AML

is variable and presumed to be dictated by the acquisition of mutations in HSPCs in

addition to the MPN driver and subsequent clonal expansion[7,8].

Single cell technologies have provided unique insights behind this evolution, however

current methods are constrained by the cell number, and the number of genetic loci that

64 can be concurrently analyzed (Supplemental Figure 1)[7-10]. We have developed an

65 innovative pipeline called LOTR-Seq, which allows for concurrent single-cell transcription

66 plus genotyping of a comprehensive panel of full-length transcripts from 30 genes that

67 are recurrently mutated in post-MPN AML. This full-length approach facilitates the

68 detection of rare variants, the phasing of mutations, and the identification of different

69 isoforms.

70 Here we used LOTR-Seq to identify clonal evolution during transformation from MPN to

71 AML and to determine the impact of these additional mutations on *JAK2*-mutant HSPCs

72 in driving disease progression.

Materials and methods

Ethics approval and Primary Human Sample processing

Patient bone marrow (BM) samples were obtained from the Royal Brisbane Women's

Hospital and from the South Australian Cancer Research Biobank (SACRB), and

analyzed with approvals from QIMR Berghofer HREC (p1382). Patients provided written

informed consent for the collection, storage, and use of their samples in research.

Detailed patient information can be found in Supplemental Table 1. Samples were stained

with CD3-PE/Cy7, CD34-PE, CD14-FITC and CD15-APC (Supplemental Table 2).

CD34+ HSPCs were FACS sorted, and single cell 3' RNA sequencing was performed

with the 10x Chromium v3.1 on the Illumina NextSeq 500/550 Platform. CD34+ BM bulk

genomic DNA capture panel sequencing and mutation calling were performed as

described in Magor et al, 2016[11] on 7 of the 12 samples.

Long read sequencing of transcripts: LOTR-Seq

We first designed a capture panel on the Roche HyperDesign platform targeting all exonic

regions of 30 recurrently mutated MPN/AML genes: *ASXL1*, *BCOR*, *CALR*, *CBL*, *CSF3R*,

DNMT3A, *EZH2*, *FLT3*, *GATA2*, *IDH1*, *IDH2*, *JAK2*, *KIT*, *KRAS*, *MPL*, *MYC*, *NPM1*,

NRAS, *PPM1D*, *PTPN11*, *RUNX1*, *SF3B1*, *SH2B3*, *SRSF2*, *STAG2*, *TET2*, *TP53*, *TERT*,

U2AF1, and *ZRSR2*. 10x barcoded cDNA underwent pre-capture PCR and read

extension using custom primers (Supplemental Table 3)[12]. Amplified and extended cDNA

was used to capture genes of interest using the default KAPA HyperCap Workflow v3,

followed by post-capture PCR amplification. Further detailed information can be found in

the Supplemental methods. Oxford Nanopore Technology (ONT) ligation kits, LSK110 or

LSK114, were used in accordance with the manufacturer's protocols, with modifications

to the incubation time from 10 minutes to 20 minutes[13] to improve adapter ligation and

ONT Flow Cell loading amount (50-100 fmol; Supplemental Table 4).

Single cell short read sequencing processing

MPN and post-MPN AML sequenced reads were adapter trimmed and processed through

cellranger (v6.0.1, or MPN10 v3.0.2) with GRCh38 genome build to obtain a counts per

single cell and gene matrix. 10x single cell RNA-seq data from healthy human CD34+ BM

HSPCs from Ainciburu et al. 2023[14] were downloaded from GEO (Supplemental Table

5). Healthy, CP-MPN, and post-MPN AML samples were preprocessed in R (v3.6.3)

following the Seurat v4.3 pipline[15]. Clusters were annotated according to key lineage

marker gene expression. For analyzing lineage skewing, clusters were broadly grouped

by adding up clusters into HSC/LMPP (HSC+LMPP), Mega/Ery-Primed (MEP+MkP+e-

Ery+l-Ery+Ery), Gran/Mono-Primed (GMP+Mono+Baso+DC), and Lymphoid-Primed

(CLP+B-primed). Pseudo time prediction and cell fate scoring were performed using

Palantir (v1.3.0)[16] within Python (v3.11.4). Differential gene expression and gene set

enrichment analysis are detailed in Supplemental methods.

ONT sequencing, read processing and mutational profiling analysis

Base calling and chimeric read separation are detailed in Supplemental methods. Reads

were assigned a 10x barcode using FLAMES[17], allowing for two mismatches. Fastqs were

split into individual barcode fastqs using zgrep to identify FLAMES-assigned barcodes in

the fastq header line. Deconvoluted reads were mapped against the GRCh38 genome

build using minimap2 (v2.27)[18] with option `-L -ax splice`. Reads filtering and mutation

calling are detailed in the Supplemental methods. Expressed variant allele frequency

(eVAF) was calculated in R as the ratio of the number of reads with bases matching the

mutant allele to the total number of reads at the locus. Finally, eVAFs were assigned to a

single cell transcriptome by matching barcodes recovered by long read and short read

sequencing.

Post-MPN AML differentiation stage assessment

ScRNASeq raw counts from 11 post-MPN AML *JAK2*V617F/*TP53* bi-allelic loss[8] and our

post-MPN AML samples were projected on the Bone marrow map as described in

https://github.com/andygxzeng/BoneMarrowMap[21] (accessed 27/04/2025). Raw counts

of bulk RNA-Seq data from 45 CP MPN, 34 post-MPN AML, and 11 healthy controls were

downloaded from GEO with GSE283710[20]. Single cell RNA-Seq (scRNASeq) data of 11

*JAK2*V617F/*TP53 bi-allelic loss* post-MPN AML and 5 healthy controls (GSE226340)[8]

were extracted and raw counts pseudo-bulked. Data were normalized (edgeR v4.2.2) and

log2 transformed before differentiation stage scoring was performed as described[21]. AML

differentiation scores from bulked and pseudo-bulked RNASeq data were centered, and

patient-specific scores Euclidean distance underwent hierarchical clustering (hclust; stats

R base package) with default parameters. Data were visualized with heatmap3.R

https://github.com/obigriffith/biostar-tutorials/blob/master/Heatmaps/heatmap.3.R.

Results

HSPC heterogeneity is preserved in chronic phase MPN but with lineage skewing

To determine whether the expression of *JAK2*V617F alters the transcriptional
heterogeneity of the CD34+ HSPC compartment, we compared the single cell
transcriptional landscape of CD34+ HSPC-enriched populations (Figure 1A) from normal
human bone marrow samples[14] (Supplemental Figure 2A; n=6) to that of CD34+ HSPCs
from eight patients with *JAK2*V617F-driven CP-MPN (Figure 1B). Transcriptional
heterogeneity was largely preserved in CP-MPN CD34+ cell populations, comprising
HSC/LMPPs, megakaryocyte-erythroid-, lymphoid- or granulocyte-monocyte-primed
populations, and consistent with that seen in the healthy CD34+ populations (Figure 1B-
C; Supplemental Figure 2B). However, CP-MPN CD34+ populations demonstrated a
higher percentage of cells exhibiting megakaryocyte/erythroid-primed gene expression
profiles (Figure 1D-E: Mega/Ery-Primed mean±SD: 22.5±6.9% healthy vs 34.1±10.3%
CP-MPN, p=0.028; Supplemental Figure 2C).

These findings demonstrate that, consistent with healthy CD34+ HSPCs, transcriptional
lineage-priming is present in primitive hematopoietic cell populations in CP-MPN;
however, CP-MPN exhibits an HSPC compartment with a higher abundance of cells with
a megakaryocyte/erythroid transcriptional phenotype, consistent with the pathology of
CP-MPN, characterized by the overproduction of mature megakaryocytes and
erythrocytes.

MEP lineage priming in HSPCs is linked to the expansion of *JAK2*V617F mutant cells

We were interested to determine how clonal heterogeneity within CP-MPN altered this
lineage bias within the CD34+ HSPC population. To identify *JAK2*V617F-mutated cells
within CP-MPN CD34+ HSPCs, we developed a technique for long read genotyping of
transcripts (LOTR-Seq). LOTR-Seq uses the single cell barcoded cDNA derived by the

10x Chromium platform to capture the cDNA of target genes, like *JAK2*, followed by ONT

long read sequencing (Figure 2A) to correlate expressed mutational information to the

transcriptome.

When LOTR-Seq was performed on the CP-MPN CD34+ HSPCs, we were able to

demonstrate that *JAK2*V617F pseudo-bulked eVAF strongly correlated with both genomic

VAF (r=0.9; p=0.005; Supplemental Figure 3A) and short read-derived eVAF (r=0.96;

p<000.1; Supplemental Figure 3B) of the CD34+ compartment, validating the

reproducibility of mutation calling from long read sequencing. LOTR-Seq enabled us to

genotype up to 51% of cells at the *JAK2*V617 locus, improving upon previous techniques

(Supplemental Figure 1)[10,22]. Notably, the percentage of genotyped cells at the *JAK2*V617

locus correlated with sequencing depth (r=0.8, p=0.0006; Supplemental Figure 3C).

We then assessed the contribution of *JAK2*V617F-mutated cells to the CD34+ HSPC

compartment using the density distribution of *JAK2*V617F-mutated and *JAK2*-wild type

cells within the clusters identified by uniform manifold approximation and projection

(UMAP) (Figure 2B). The percentage of *JAK2*V617F mutated HSPCs within the

Mega/Ery-primed CD34+ compartment correlated with the relative size of the Mega/Ery-

primed compartment (R=0.75; p=0.031; Figure 2C) and inversely correlated with that of

the Gran/Mono-primed compartment (R=-0.79; p=0.02). PV patients displayed higher

*JAK2*V617F clone percentages in the Mega/Ery-Primed compartments[10] (Supplemental

Figure 3D), conversely, *JAK2*V617F lineage quantification was stably distributed in

patients with ET or MF.

These findings suggest that *JAK2*V617F either skews lineage bias towards the Mega/Ery

fate or confers a proliferative advantage exclusively to Mega/Ery-primed HSPCs. To

resolve this, we used Palantir analysis of differentiation in pseudotime to examine the

effect of *JAK2*V617F on HSPC cell fate potential[16]. Here, HSPCs of PV patients

containing the *JAK2*V617F mutation show a higher probability of converging towards an

erythroid cell fate, evidenced by a higher average erythroid cell fate score compared to

both healthy CD34+ and *JAK2*-wild type cells (Figure 2D-E, Supplemental Figure 3E),

while sMF patients exhibit a higher probability of converging towards a megakaryocytic

progenitor fate (Supplemental Figure 3F-G). This supports the finding that *JAK2*V617F

favors erythroid/megakaryocyte differentiation. Furthermore, cell cycle analysis revealed
that *JAK2*V617F mutant cells contained a greater proportion of cells in G2M phase
(Figure 2F, Supplemental Figure 4A).

We next sought to determine the impact of the *JAK2*V617F mutation on canonical
signaling pathway activation. Compared to healthy CD34+, CP-MPN CD34+
demonstrated increased JAK-STAT signaling across all compartments (Figure 2G,
Supplemental Figure 4B) in both *JAK2*V617F, *JAK2*-wild type cells, suggesting that the
activation of JAK-STAT signaling within HSPC is also regulated by cell-extrinsic factors,
such as increased activation of cytokine pathways[23]. The impact of this *JAK2*V617F
mutation appears most pronounced in the Mega/Ery-Primed compartment with Myc
targets and the heme metabolism pathway activated in CP-MPN. Interestingly,
inflammatory pathways are suppressed in PV but activated in MF patients, particularly at
the HSC/LMPP level (Supplemental Figure 4C).

These data validate the LOTR-Seq pipeline to identify *JAK2*V617F mutant vs *JAK2*-wild
type HSPCs within individual CP-MPN samples. *JAK2*V617F mutated cells demonstrate
increased proliferation and preferential expansion towards the erythroid and
megakaryocyte lineages and context-specific transcriptional effects of *JAK2*V617F by
lineage-primed compartment and MPN subtype, reflecting the heterogeneity observed in
the clinical presentation of MPNs.

Transformation to post-MPN AML is characterized by the loss of stem cell
transcriptional heterogeneity and is driven by genetic complexity

We next sought to determine whether the LOTR-Seq pipeline could be used to investigate
the relationship between genetic and transcriptional heterogeneity within the HSPC
compartment during progression from CP-MPN to post-MPN AML. In dramatic contrast
to the transcriptional heterogeneity observed in CP-MPN, the CD34+ HSPC compartment
in post-MPN AML is largely dominated by a single transcriptional profile (Figure 3A-B).
For example, MPN10 AML exhibits a dominant HSC/LMPP-like transcriptional signature,

representing 79.4% of all cells (Figure 3A), while MPN20 AML shows a dominant

Mega/Ery primed-like transcriptional signature (84.8%; Figure 3B).

A limitation of existing protocols integrating single cell RNA-Seq and genotyping is the

inability to simultaneously genotype across numerous loci with sufficient coverage and

representation to draw inferences on cells with multiple mutations within the CD34+HSPC

compartment (Supplemental Figure 1)[22]. The LOTR-Seq pipeline overcomes this

limitation by using enrichment for a panel of 30 genes, enabling the genotyping of

expressed genes across a range from 0% to 100% (Supplemental Figure 5A). We

therefore sought to apply this technology to determine the relationship between the

genomic architecture of post-MPN AML and the transcriptional consequences linked with

disease progression.

MPN10 contained a dominant, HSC/LMPP-like expanded population in post-MPN AML.

LOTR-Seq identified the MPN-driver *JAK2*V617F mutation together with an *IDH2*R140W

mutation and *SRSF2*P95_R103 in-frame deletion (Figure 3A, C-E, Supplemental Figure

5B) in 56.7%, 92.4 %, and 83.4% of genotyped cells, respectively, and expressed at high

VAF in the HSC/LMPP-like CD34+ cells. Additionally, in *JAK2*V617F, *IDH2*R140W and

*SRSF2*P95R co-mutated cells we observed the predominant inclusion of a poison exon

cassette[24] , in comparison to cells from a post-MPN AML containing *JAK2*V617F and

*IDH1*R132S without a mutation in *SRSF2* (Supplemental Figure 5C).

In a second post-MPN AML, MPN20, we identified a dominant Mega/Ery primed-like

HSPC population containing *JAK2*V617F (89.6% of genotyped cells) with biallelic

*TP53*C238F mutation (93.9% of genotyped cells) (Figure 3B, F-G, Supplemental Figure

5D). This identification of a Mega/Ery-signature in *JAK2*V617F multi-hit-*TP53* post-MPN

AML is consistent with other published human data[8] and aligns with data from murine

post-MPN AML with this same combination of mutations[25,26].

Altogether, these data validate the use of LOTR-Seq in clonal analysis of human post-

MPN AML and demonstrate that post-MPN AMLs exhibit loss of HSPC heterogeneity with

the expansion of a dominant transcriptional cluster that appears to be linked to the identity

of co-existing secondary mutations.

Post-MPN AML demonstrates lineage restricted, recurrent differentiation priming

Recent findings suggest that molecular subtypes of de novo AML are defined by unique

transcriptional profiles that reflect lineage bias[21,27,28]. Our initial analysis of the HSPC

compartment in post-MPN AML suggests lineage priming analogous to that seen in de

novo AML[21,28]. To determine if these findings can be applied more generally to post-MPN

AML, we analyzed bulk RNA-Seq of 34 CD34+ post-MPN AML samples[20] together with

single cell RNA-Seq CD34+ HSPCs of a further 11 post-MPN AML cases[8]. Here, we

adopted methods developed by Zeng et al 2025[21] to assess differentiation stages in post-

MPN AML. Both methods of projecting AML on a detailed, healthy bone marrow

hierarchical hematopoiesis map[21] (Figure 4A-B) and AML differentiation stage scoring

(Figure 4C) suggest that even genetically homogenous subgroups, such as *JAK2V617F*

with multi-hit *TP53*, can exhibit expansion in different lineage-primed compartments.

Compared to adult de novo AML, as reported by Zeng et al 2025[21] (n=28 of 139), post-

MPN AML exhibited a significantly higher overrepresentation of HSC/LMPP states (n=23

of 45, Chi-Square p=0.0001). In the 45 post-MPN AML samples, 51% showed an

immature differentiation stage (HSC/LMPP/MEP), with HSC/LMPP/early lymphoid stage

or MEP/MKP/erythroid stage observed in 31% and 18%, respectively. Strikingly, there

was an absence of a promonocytic/monocytic biased subgroup in *JAK2*V617F mutated

post-MPN AML, a dominant subgroup in de novo AML[21] (Figure 4C). These data

demonstrate that post-MPN AML is not only genetically distinct from de novo AML, but

also phenotypically and transcriptionally distinct with a more restricted differentiation

repertoire.

High-risk mutations in post-MPN AML are already present in chronic phase

disease.

Post-MPN AML is intrinsically chemotherapy resistant, and novel strategies are urgently

required to prevent the progression of high-risk CP-MPN to AML. We therefore sought to

determine whether LOTR-Seq could identify pre-leukemic high-risk populations within

CD34+ cells from patients with CP-MPN. To do this, we analyzed a patient with matched

CP-MPN and post-MPN AML samples (Figure 5). The *JAK2*V617F mutation was stable

between CP-MPN and post-MPN AML (Figure 5B, E), however deep sequencing analysis

of the CP-MPN CD34+ cells identified a rare subpopulation of co-mutated *JAK2*V617F

and *IDH1*R132S (8.2%) in CP-MPN, which expanded to 64.5% in post-MPN AML (Figure

5C, F). In contrast to what was observed in the analysis of the previous post-MPN AML

samples (Figure 4), the expansion of the double-mutated population resulted in the

relative expansion of the HSC/LMPP clusters with partially maintained representation of

the more committed lineage-primed cells.

The co-occurrence of *JAK2*V617F and *IDH1*R132S clones in chronic phase, with

expansion of the double-mutated clone during AML progression, suggests that this

leukemia was driven by the linear clonal evolution of the double-mutant cells (Figure 5G).

Thus, the LOTR-Seq protocol can identify genetic complexity at the single cell level,

enabling the identification of pre-leukemic clones that later give rise to post-MPN AML.

Additional mutations in *JAK2*V617F cells exhibit mutation and context specific
gene expression signatures
297

To determine how the presence of additional mutations can impact MPN stem cells and

if this changes with leukemic transformation, we first compared *IDH1*R132-mutant cells

(n=122) with *JAK2*V617F/*IDH1*WT cells (n=1562), in the matched chronic phase sample,

for differential gene expression and gene set enrichment across the major transcriptional

compartments in CD34⁺ bone marrow (Figure 6A-C, Supplemental Table 9-10).

*IDH1*R132S cells within the Mega/Ery-Primed compartment displayed heightened

activation of inflammatory pathways, notably TNFα and IL2–STAT5A, alongside an

increase in *STAT5A* gene expression, known to have potential roles in MPN

pathophysiology[20,29]. Concurrently, we observed an upregulation of hypoxia-related

genes (e.g., *BCL2*) and a downregulation of oxidative phosphorylation (OXPHOS)-related

genes, suggesting a metabolic shift in the *IDH1*-mutant cellular environment.

Interestingly, even within chronic phase disease, the gene expression changes in *IDH1*-

mutant cells showed similarities to gene expression changes in *IDH*-mutant de novo

AML[30], also revealing characteristics such as leukemia stem cell priming and

quiescence[27,31]. This occurred alongside the suppression of genes involved in heme

metabolism, E2F-targets, DNA metabolic processes, and replication pathways. However,

when comparing matched *JAK2*V617F and *IDH1*R132S co-mutated cells between

chronic phase disease and post-MPN AML (Figure 6A), we identified more pronounced

alterations across all HSPC compartments (Figure 6D-E), including that TNFα signaling

and de novo *IDH*-mutant core pathways were upregulated in the leukemic phase, while

E2F-targets and DNA replication processes were suppressed. Of note, we observed

significant upregulation of TP53 pathway in the Mega/Ery-Primed compartment and other

malignancy associated gene expression[20,29,32,33].

Through linking mutations to transcriptomic profiles via LOTR-Seq, we provide novel

insights into the pathways that contribute to *IDH1*-mediated progression to post-MPN

AML, highlighting the striking similarities with de novo *IDH*-mutant AML. These findings

argue strongly for further investigations into the therapeutic potential of IDH inhibitors

during the chronic phase and post-MPN AML settings[34,35].

Hyperactivation of JAK-STAT signaling is maintained in *JAK2*V617F-negative post-

MPN AML

It has been observed that in 16-50% of post-MPN AML cases, the MPN driver is absent

at the time of leukemia transformation, even when the original disease contained a

mutation in a canonical MPN driver gene [36,37,6]. The origin of this MPN driver-negative

post-MPN AML is incompletely understood. We reasoned that this category of post-MPN

AML must have a different mechanism of transformation compared to the classical linear

clonal evolution model.

To examine this further, we used LOTR-Seq to analyze paired, sequential samples from

a patient who presented with *JAK2*V617F CP-MPN and *JAK2*V617F-negative post-MPN

AML (Figure 7A-H, Supplemental Figure 6A-D). Here, the CP-MPN contained a dominant

clone with *TP53*Y236C mutations in addition to a *JAK2*L393V mutation, previously

described to result in gain of function[38] (Figure 6C, Supplemental Figure 7B,D). Within

the *TP53*Y236C, *JAK2*L393V population, there was further heterogeneity with two

identifiable sub-clonal populations containing either *TP53*R282W (Figure 7D) or

*U2AF1*S34Y and *JAK2*V617F (Supplemental Figure 6C; Figure 7B). With evolution to

AML, the *TP53*Y236C, *JAK2*L393V *TP53*R282W containing cells demonstrated selective

clonal advantage and are dominant in the post-MPN AML stage (Figure 7H). Both *TP53*

mutations identified are located within the DNA-binding domain (Supplemental Figure

7A), and thus are predicted to confer a loss-of-function phenotype. Importantly, we were

also able to demonstrate that the *TP53* mutations occurred on independent alleles within

the same cell and exhibited random monoallelic expression (i.e., somatically acquired in

trans; Supplemental Figure 7B-C). These findings identify leukemic transformation that

arose from a genetically complex, high-risk ancestral clone and demonstrate parallel

clonal evolution resulting in *JAK2*V617F-negative AML (Figure 5I; Supplemental Figure

5E). Here, the hyperactivation of JAK-STAT signaling evident in the chronic phase sample

was maintained post leukemic transformation, despite the loss of the *JAK2*V617F clone

(Supplemental Figure 7E-F). Of note, maintained hyperactivation of JAK-STAT signaling

post leukemic transformation was also observed in the case of linear evolution, patient

SA2 (Supplemental Figure 7E-F).

In summary, using LOTR-Seq to combine single cell transcriptional analysis and panel-

enriched genotyping of transcripts from loci commonly associated with myeloid

malignancies, we have been able to demonstrate that the transformation of CP-MPN to

post-MPN AML is associated with the acquisition of recurrent secondary mutations that

drive loss of transcriptional heterogeneity and the immortalization of a genotypically

defined, transformed progenitor cell population.

Discussion

Mutational complexity in MPN is common, with more than 40% of CP-MPN patients

harboring a mutation relevant to myeloid malignancies in addition to an MPN driver.

Importantly, the acquisition of additional mutations in CP-MPN is associated with reduced

event-free survival. Consistent with this, AML arising from antecedent MPN is associated

with 2 or more genetic mutations in the vast majority of patients[6], strongly supporting the

hypothesis that the acquisition of these additional mutations drives leukemic

transformation of CP-MPN. However, our capacity to determine the functional

consequences of mutational complexity in MPN stem cells has been limited by our ability

to multiplex single cell phenotyping technology with mutational status. Specifically, single

cell RNA-sequencing approaches have provided a high-resolution map of the substantial

amount of transcriptional heterogeneity within the HSPC compartment in the BM of

patients with CP-MPN but have had limited capacity to associate these differences with

an individual cell's genotype across the spectrum of loci recurrently mutated in post-MPN

AML[9,10,22].

Here, we describe a novel pipeline, LOTR-Seq, combining 10x single cell RNA-

sequencing of CP-MPN and post-MPN AML, together with broad, panel-based

enrichment and long read genotyping of the vast majority of genetic loci frequently

mutated in MPN. We demonstrate that cells expressing the MPN-driver *JAK2*V617F can

exhibit transcriptional lineage priming diversity similar to that of their healthy counterparts,

but are heavily biased towards a megakaryocyte/erythroid program. Strikingly, this

transcriptional diversity is lost upon leukemic transformation, with the HSPC compartment

of post-MPN AML exhibiting a homogenous transcriptional profile that appears to be

dictated by the identity of the co-occurring mutations.

In the case of the *JAK2*V617F-positive, *TP53*-mutant AML, compound mutant cells were

found almost exclusively within the megakaryocyte/erythroid-primed clusters,

corroborating recent findings in additional human samples and mouse models with this

same genetic combination. Notably, however, the *JAK2*V617F and *IDH*-mutated AMLs

exhibited transcriptional features of more primitive HSCs, suggesting that post-MPN is

transcriptionally diverse. Testing this hypothesis using a bulk RNA sequencing dataset

with a larger number of patients, we demonstrated that post-MPN AML comprises a group

of leukemias with three transcriptionally distinct profiles. Importantly, the identity of these

transcriptional profiles both diverges and overlaps with what is observed in de novo AML.

Patients with post-MPN AML are excluded from AML clinical trials for targeted agents,

attributed to the comparatively dismal outcomes in post-MPN AML. The finding that *IDH-*

mutated post-MPN AMLs exhibit transcriptional similarity with the corresponding de novo

AMLs should warrant reconsideration of this approach. This is particularly relevant for

IDH inhibitors, as the transcriptional consequences of mutant *IDH1* appear to be amplified

as either a cause or consequence of leukemic transformation to post-MPN AML.

The progression of CP-MPN to post-MPN AML is rare, difficult to predict, and can occur

over decades. Consequently, clinical studies evaluating new MPN therapies are often

underpowered to use leukemic transformation as an endpoint. Given the association

between mutational complexity and reduced event-free survival, monitoring the

emergence of compound mutational clones could prove valuable and should be

investigated as surrogate endpoint for CP-MPN disease progression. LOTR-Seq

improves upon conventional amplicon-based genomic mutational profiling here by

demonstrating conclusively whether TP53 mutations are monoallelic or biallelic, which is

highly predictive of prognosis in CP-MPN. LOTR-Seq has also enabled the identification

of additional mutations in the genes of interest that are not routinely assessed for in MPN.

The identification of *JAK2*L393V has obvious implications for our understanding of, and

the evolution of, MPN driver-negative post-MPN AML and CP-MPN.

Furthermore, the additional dimension of transcriptional profiling to the clonal analysis of

MPN disease progression allows for both the monitoring of the emergence of high-risk

mutations and the assessment of their functional consequence. This information could be

used to both inform treatment decisions and allow updated prognostication. It also

enables the identification of unique features of high-risk clones that can hopefully be

leveraged into new treatments that can prevent the development of post-MPN AML.

In aggregate, the development of LOTR-Seq has both advanced our understanding of the

functional consequences of mutational complexity associated with CP-MPN disease

progression and provides a new tool to be exploited in CP-MPN clinical trial design to

progress the field towards surrogate endpoints of disease progression and ultimately

improve long-term outcomes for these patients.

Acknowledgement

We are grateful for the assistance of the QIMR Berghofer facilities, including flow cytometry and sequencing, and for the helpful comments from members of the Lane Lab, particularly Stacey Anderson. We thank Rachel Thjissen for the fruitful discussion on ONT barcode recovery, primer sequences, and ONT read extension protocols. We gratefully acknowledge the support of Neil Herron and the Herron Family Trust, and the Gordon and Jessie Gilmour Family Trust as dedicated supporters of leukemia research in Queensland. We gratefully acknowledge the contribution of the MPN Alliance Australia for the valuable patient perspective they provided for this study. SWL was funded by an NHMRC Investigator Grant (1195987 2021-2025) and by a philanthropic donation from the MPN Alliance of Australia. ID was supported by a PROMOS, DAAD stipend. JS was funded by a Cancer Council Queensland fellowship (2025829). JG was supported by an HSANZ/Leukaemia Foundation PhD scholarship. This research was supported by the Australian Cancer Research Foundation Centre for Optimised Cancer Therapy.

Authorship

SWL, MJB, JS conceptualized, designed, and supervised wet lab experiments and bioinformatics analysis. JG, LC, RH, RZ performed wet lab experiments. JS, JG, ID, MB performed bioinformatic analysis. CM, DR, SWL coordinated primary patient sample collection. GA, HC, WG coordinated the collection, consent, and storage of patient sample information. VYL helped with gene panel design. AP performed genomic DNA NGS analysis of primary patient samples. JG, JS, MJB, SWL wrote the manuscript. All authors contributed to and edited the manuscript.

Conflict of Interest Disclosure

SWL, MJB, and JS have received research funding from Bristol Myers Squibb for unrelated projects. We acknowledge receipt of reagents from PharmaEssentia for unrelated projects. MJB has received research funding from Cylene Pharmaceuticals for unrelated projects. DMR has consulted for or received honoraria from GSK, Jubilant,

Keros, Merck, Menarini, Novartis, Prelude, and Takeda, for unrelated projects. SWL has

consulted for AbbVie, Novartis, Astellas, and GSK, also for unrelated projects.

Data Availability Statement

The 10x sequencing raw and processed data generated in this study is deposited in

ArrayExpress with accession E-MTAB-15981. Processed data will be made available via

figshare/zendo. Code is available via GitHub https://github.com/JStrau/LOTRSeq. ONT,

raw data, due to its size, will be made available upon request.

References

1.   Jamieson CH, Gotlib J, Durocher JA, et al. The JAK2 V617F mutation occurs in hematopoietic
stem cells in polycythemia vera and predisposes toward erythroid differentiation. *Proceedings of the*
National Academy of Sciences of the United States of America. Apr 18 2006;103(16):6224-9.
doi:10.1073/pnas.0601462103
2.   Prins D, González Arias C, Klampfl T, Grinfeld J, Green AR. Mutant Calreticulin in the
Myeloproliferative Neoplasms. *HemaSphere*. Feb 2020;4(1):e333. doi:10.1097/hs9.0000000000000333
3.   Beer PA, Delhommeau F, LeCouédic JP, et al. Two routes to leukemic transformation after a
JAK2 mutation-positive myeloproliferative neoplasm. *Blood*. Apr 8 2010;115(14):2891-900.
doi:10.1182/blood-2009-08-236596
4.   Challen GA, Boles NC, Chambers SM, Goodell MA. Distinct hematopoietic stem cell subtypes are
differentially regulated by TGF-beta1. *Cell stem cell*. Mar 5 2010;6(3):265-78.
doi:10.1016/j.stem.2010.02.002
5.   Yamamoto R, Morita Y, Ooehara J, et al. Clonal analysis unveils self-renewing lineage-restricted
progenitors generated directly from hematopoietic stem cells. *Cell*. Aug 29 2013;154(5):1112-1126.
doi:10.1016/j.cell.2013.08.007
6.   McNamara CJ, Panzarella T, Kennedy JA, et al. The mutational landscape of accelerated- and
blast-phase myeloproliferative neoplasms impacts patient outcomes. *Blood Adv*. Oct 23
2018;2(20):2658-2671. doi:10.1182/bloodadvances.2018021469
7.   Calabresi L, Carretta C, Romagnoli S, et al. Clonal dynamics and copy number variants by single-
cell analysis in leukemic evolution of myeloproliferative neoplasms. *American journal of hematology*.
Oct 2023;98(10):1520-1531. doi:10.1002/ajh.27013
8.   Rodriguez-Meira A, Norfo R, Wen S, et al. Single-cell multi-omics identifies chronic inflammation
as a driver of TP53-mutant leukemic evolution. *Nature genetics*. Sep 2023;55(9):1531-1541.
doi:10.1038/s41588-023-01480-1
9.   Rodriguez-Meira A, Buck G, Clark SA, et al. Unravelling Intratumoral Heterogeneity through
High-Sensitivity Single-Cell Mutational Analysis and Parallel RNA Sequencing. *Mol Cell*. Mar 21
2019;73(6):1292-1305 e8. doi:10.1016/j.molcel.2019.01.009

10.   Van Egeren D, Escabi J, Nguyen M, et al. Reconstructing the Lineage Histories and Differentiation
Trajectories of Individual Cancer Cells in Myeloproliferative Neoplasms. *Cell stem cell*. Mar 4
2021;28(3):514-523.e9. doi:10.1016/j.stem.2021.02.001
11.   Magor GW, Tallack MR, Klose NM, et al. Rapid Molecular Profiling of Myeloproliferative
Neoplasms Using Targeted Exon Resequencing of 86 Genes Involved in JAK-STAT Signaling and
Epigenetic Regulation. *J Mol Diagn*. Sep 2016;18(5):707-718. doi:10.1016/j.jmoldx.2016.05.006
12.   Thijssen R, Tian L, Anderson MA, et al. Single-cell multiomics reveal the scale of multilayered
adaptations enabling CLL relapse during venetoclax therapy. *Blood*. Nov 17 2022;140(20):2127-2141.
doi:10.1182/blood.2022016040
13.   Wei S, Williams Z. Rapid Short-Read Sequencing and Aneuploidy Detection Using MinION
Nanopore Technology. *Genetics*. Jan 2016;202(1):37-44. doi:10.1534/genetics.115.182311
14.   Ainciburu M, Ezponda T, Berastegui N, et al. Uncovering perturbations in human hematopoiesis
associated with healthy aging and myeloid malignancies at single-cell resolution. *Elife*. Jan 11
2023;12doi:10.7554/eLife.79363
15.   Hao Y, Hao S, Andersen-Nissen E, et al. Integrated analysis of multimodal single-cell data. *Cell*.
2021/06/24/ 2021;184(13):3573-3587.e29. doi:https://doi.org/10.1016/j.cell.2021.04.048
16.   Setty M, Kiseliovas V, Levine J, Gayoso A, Mazutis L, Pe'er D. Characterization of cell fate
probabilities in single-cell data with Palantir. *Nature biotechnology*. Apr 2019;37(4):451-460.
doi:10.1038/s41587-019-0068-4
17.   Tian L, Jabbari JS, Thijssen R, et al. Comprehensive characterization of single-cell full-length
isoforms in human and mouse with long-read sequencing. *Genome biology*. Nov 11 2021;22(1):310.
doi:10.1186/s13059-021-02525-6
18.   Li H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics*. 2018;34(18):3094-
3100. doi:10.1093/bioinformatics/bty191
19.   Koboldt DC, Zhang Q, Larson DE, et al. VarScan 2: somatic mutation and copy number alteration
discovery in cancer by exome sequencing. *Genome Res*. Mar 2012;22(3):568-76.
doi:10.1101/gr.129684.111
20.   Kong T, Laranjeira ABA, Letson CT, et al. RSK1 is an exploitable dependency in myeloproliferative
neoplasms and secondary acute myeloid leukemia. *Nat Commun*. Jan 16 2025;16(1):492.
doi:10.1038/s41467-024-55643-7
21.   Zeng AGX, Iacobucci I, Shah S, et al. Single-cell Transcriptional Atlas of Human Hematopoiesis
Reveals Genetic and Hierarchy-Based Determinants of Aberrant AML Differentiation. *Blood Cancer
Discov*. Jul 1 2025;6(4):307-324. doi:10.1158/2643-3230.BCD-24-0342
22.   Nam AS, Kim KT, Chaligne R, et al. Somatic mutations and cell identity linked by Genotyping of
Transcriptomes. *Nature*. Jul 2019;571(7765):355-360. doi:10.1038/s41586-019-1367-0
23.   Austin RJ, Straube J, Bruedigam C, et al. Distinct effects of ruxolitinib and interferon-alpha on
murine JAK2V617F myeloproliferative neoplasm hematopoietic stem cell populations. *Leukemia*. Apr
2020;34(4):1075-1089. doi:10.1038/s41375-019-0638-y
24.   Kim E, Ilagan JO, Liang Y, et al. SRSF2 Mutations Contribute to Myelodysplasia by Mutant-
Specific Effects on Exon Recognition. *Cancer Cell*. May 11 2015;27(5):617-30.
doi:10.1016/j.ccell.2015.04.006
25.   Li B, An W, Wang H, et al. BMP2/SMAD pathway activation in JAK2/p53-mutant
megakaryocyte/erythroid progenitors promotes leukemic transformation. *Blood*. Jun 23
2022;139(25):3630-3646. doi:10.1182/blood.2021014465
26.   Rampal R, Ahn J, Abdel-Wahab O, et al. Genomic and functional analysis of leukemic
transformation of myeloproliferative neoplasms. *Proceedings of the National Academy of Sciences of the
United States of America*. Dec 16 2014;111(50):E5401-10. doi:10.1073/pnas.1407792111

27.   Zeng AGX, Bansal S, Jin L, et al. A cellular hierachy framework for understanding heterogeneity and predicting drug response in acute myeloid leukemia. *Nat Med*. Jun 2022;28(6):1212-1223. doi:10.1038/s41591-022-01819-x

28.   van Galen P, Hovestadt V, Wadsworth Ii MH, et al. Single-Cell RNA-Seq Reveals AML Hierarchies Relevant to Disease Progression and Immunity. *Cell*. Mar 7 2019;176(6):1265-1281 e24. doi:10.1016/j.cell.2019.01.031

29.   Tefferi A, Vaidya R, Caramazza D, Finke C, Lasho T, Pardanani A. Circulating interleukin (IL)-8, IL-2R, IL-12, and IL-15 levels are independently prognostic in primary myelofibrosis: a comprehensive cytokine profiling study. *J Clin Oncol*. Apr 1 2011;29(10):1356-63. doi:10.1200/JCO.2010.32.9490

30.   Sirenko M, Lee S, Sun Z, et al. Deconvoluting clonal and cellular architecture in IDH-mutant acute myeloid leukemia. *Cell stem cell*. Jul 3 2025;32(7):1102-1121 e5. doi:10.1016/j.stem.2025.04.012

31.   Ng SW, Mitchell A, Kennedy JA, et al. A 17-gene stemness score for rapid determination of risk in acute leukaemia. *Nature*. Dec 15 2016;540(7633):433-437. doi:10.1038/nature20598

32.   Li R, Colombo M, Wang G, et al. A proinflammatory stem cell niche drives myelofibrosis through a targetable galectin-1 axis. *Sci Transl Med*. Oct 9 2024;16(768):eadj7552. doi:10.1126/scitranslmed.adj7552

33.   Dunbar AJ, Kim D, Lu M, et al. CXCL8/CXCR2 signaling mediates bone marrow fibrosis and is a therapeutic target in myelofibrosis. *Blood*. May 18 2023;141(20):2508-2519. doi:10.1182/blood.2022015418

34.   Goldberg LA, Yoon JJ, Johnston H, et al. Response and outcomes of patients with IDH1/2-mutated accelerated/blast-phase myeloproliferative neoplasms. *Blood Neoplasia*. May 2025;2(2):100089. doi:10.1016/j.bneo.2025.100089

35.   Chifotides HT, Masarova L, Alfayez M, et al. Outcome of patients with IDH1/2-mutated post-myeloproliferative neoplasm AML in the era of IDH inhibitors. *Blood Adv*. Nov 10 2020;4(21):5336-5342. doi:10.1182/bloodadvances.2020001528

36.   Venton G, Courtier F, Charbonnier A, et al. Impact of gene mutations on treatment response and prognosis of acute myeloid leukemia secondary to myeloproliferative neoplasms. *American journal of hematology*. Mar 2018;93(3):330-338. doi:10.1002/ajh.24973

37.   Theocharides A, Boissinot M, Girodon F, et al. Leukemic blasts in transformed JAK2-V617F-positive myeloproliferative disorders are frequently negative for the JAK2-V617F mutation. *Blood*. Jul 1 2007;110(1):375-9. doi:10.1182/blood-2006-12-062125

38.   Lanikova L, Babosova O, Swierczek S, et al. Coexistence of gain-of-function JAK2 germ line mutations with JAK2V617F in polycythemia vera. *Blood*. Nov 3 2016;128(18):2266-2270. doi:10.1182/blood-2016-04-711283

Figure legends

Figure 1. Single cell RNA sequencing identifies expanded megakaryocyte-erythroid primed cells in chronic phase MPN.

(A) Schematic of MPN bone marrow hematopoietic stem and progenitor cell (HSPC) enrichment followed by 10x Chromium single cell separation and 3' cDNA barcoding.

(B) Chronic phase MPN UMAPs (n=4, polycythemia vera, PV; n=1, essential

thrombocythemia, ET; n=1 primary myelofibrosis, PMF; n=2 secondary myelofibrosis,

sMF) of short read sequencing derived whole transcriptomics of HSPCs. Clusters

annotated by gene expression associated with HSC, hematopoietic stem cells; LMPP,

lympho-myeloid multipotent progenitor; CLP, common lymphoid progenitors; GMP,

granulocyte-monocyte progenitors; DC; Dendritic Cell progenitors; Baso,

eosinophil/basophil/mast cells progenitors; MEP, megakaryocyte-erythroid progenitor;

MkP, megakaryocyte progenitor, e-Ery; early erythroid and l-Ery, late erythroid priming.

(C) Pooled healthy sample UMAPs (n=6) showing the expression of key lineage-

defining transcriptional markers including HSC-CRHBP, LMPP-HOPX, B-primed-

VPREB1, myeloid-MPO, monocyte-LYZ, DC-SPIB, basophil-LMO4, erythroid-KLF1 and

megakaryocyte-ITGA2B.

(D) Schematic of summarized clusters analysed in (E).

(E) Bargraph with mean (black dot) and standard deviation (black line) comparing the

percentage (%) of clustered cells in healthy (n=6) vs chronic phase (CP) MPN (n=8).

Each grey or blue dot represents data from an individual sample. P-value derived from

pairwise, two-sided Welch t-test. (A, D) Created in BioRender. Lane, S. (2025) (A)

https://BioRender.com/vygkh71, (D) https://BioRender.com/9t72apn).

Figure 2. Target enrichment and long read sequencing demonstrates the impact

of mutational status on the gene transcription in chronic phase MPN.

(A) Schematic showing the long read genotyping of transcripts (LOTR-Seq) pipeline that

enriches single cell cDNA of 30 MPN/AML associated genes followed by long read

sequencing and genotyping of transcripts, with an example of one MPN patient's UMAP

colored by *JAK2*V617F expressed variant allele frequency (eVAF). (A) Created in

BioRender. Lane, S. (2025) https://BioRender.com/vygkh71.

(B) UMAPs of the CD34+ HSPC single cells (light grey) of n=8 MPNs with the density

distribution of *JAK2*V617F mutation (red), *JAK2*-wild type (blue) and non-genotyped

cells (NA; grey).

(C) Scatterplots of HSC/LMPP and key lineage-primed compartment percentage (%; x-

axis) and the % of *JAK2*V617F genotyped cells (y-axis) within the compartment (n=8

chronic phase MPN). Pearson's correlation coefficient R and test of association p-value.

The blue line represents the linear regression line with a 95%-confidence interval (grey

band).

(D) Smoothed regression lines of the Palantir predicted erythroid cell fate score (y-axis)

in dependency of the pseudotime (x-axis) for n=5 healthy, n=4 Polycythemia vera

patients (PV) *JAK2*V617F and *JAK2* wild type cells.

(E) Bargraph of the mean erythroid cell fate for each sample comparing n=5 healthy to

n=4 PV *JAK2*V617F and *JAK2*-wild type (*JAK2*WT*)* summarized in (D). Black dot and

line represent the group mean and standard deviation, respectively. P-values derived

from pairwise, two-sided Welch t-tests.

(F) Bargraph of the percentage (%) of cell cycle phase (G1, S, G2M) for the Mega/Ery-

Primed compartment comparing n=6 healthy to n=8 CP-MPN *JAK2*V617F and *JAK2*

wild type (*JAK2*WT*)*. Black dot and line represent the group mean and standard

deviation, respectively. P-values derived from pairwise, two-sided Welch t-tests.

(G) Bargraph of the KEGG JAK-STAT signaling pathway average single cell gene set

enrichment score comparing n=6 healthy to n=8 MPN *JAK2*V617F and *JAK2* wild type

(*JAK2*WT). Black dot and line represent the group mean and standard deviation,

respectively. P-values derived from pairwise, two-sided Welch t-tests.

Figure 3. Loss of heterogeneity occurs in leukemic transformation and is driven

by additional mutations in the *JAK2*V617F mutant clone.

(A-B) UMAPs of post-MPN AML samples (A) MPN10 and (B) MPN20 showing

transcriptionally defined cells clusters and a bargraph depicting the relative percentage

of cells per cluster,

(C-E) UMAPs of post-MPN AML sample MPN10 showing expressed variant allele

frequencies (eVAF) of mutations (C) *JAK2*V617F (D) *IDH2*R140W and (E)

*SRSF2*P95_R103del.

(F-G) UMAPs of post-MPN AML sample MPN20 showing expressed variant allele

frequencies (eVAF) of mutations (F) *JAK2*V617F and (G) *TP53*C238F.

Grey colored dots indicate non-genotyped cells.

Figure 4. Post-MPN AML exhibit lineage-biased hematopoiesis

(A) Bone marrow hematopoietic stem cell atlas UMAP with cell type annotations as

generated by Zeng et al. 2025[21].

(B) UMAPs highlighting key lineage gene expression markers HSC-*CRHBP*, LMPP-

HOPX, erythroid-*KLF1*, megakaryocyte-*ITGA2B*, Baso-*LMO4*, and B-primed-*VPREB1*.

(C) Projection of our generated n=3 *JAK2*V617F post-MPN AML single cell samples

(SA2LT, MPN20, MPN10)and n=11 *JAK2*V617F TP53-multi-hit post-MPN AMLs from

Rodriguez-Meira et al. 2023[8] on to the healthy bone marrow atlas UMAP.

(D) Hierarchical clustering of lineage-defining gene expression scores (y-axis) of n= 11

*JAK2*V617F post-MPN AML and n=6 healthy samples from Rodriguez-Meira et al.

2023[8], n=45 chronic phase MPN, n=34 post-MPN AML and n=11 healthy from Kong et

al 2025[20] (x-axis).

Figure 5. Leukemic transformation from chronic phase MPN showing a linear

pattern of clonal expansion of a high risk *JAK2*V617F and *IDH1*R132S mutant

clone.

UMAPs of the same patient at diagnosis of (A-C) chronic phase MPN (SA2CP) and (D-

F) post-MPN AML (SA2LT), showing (A,D) transcriptionally defined clusters with

bargraphs showing the percentage of cells in the transcriptionally defined clusters and

(B-C,E-F) expressed variant allele frequencies (eVAF) of mutations (B,E) *JAK2*V617F

and (C,F) *IDH1*R132S. Grey colored dots indicate non-genotyped cells.

(G) Fish plot illustrating the linear clonal evolution of the *JAK2*V617F and *IDH1*R132S

mutant clone.

Figure 6. IDH1 mutations in *JAK2*V617F cells exhibit mutation and context

specific gene expression signatures

(A) Schema of analysis performed in B-E, Created in BioRender. Lane, S. (2026)

https://BioRender.com/1kuev8s

(B) Volcanoplot displaying the $\log_2$ fold change (x-axis) and $\log_{10}$ P-value (y-axis) from

comparing gene expression of *JAK2*V617F *IDH1*R132S (higher expression marked in

darkblue) vs *JAK2*V617F *IDH1* wildtype (WT) (higher expression marked in darkred)

cells of a patient with chronic phase MPN (SA2CP). Colored are significant different

genes (p<0.05) associated with msigdb hallmark signatures and signatures of interest:

TNFA signaling via NFKB (TNF; green), Hypoxia (purple), epithelial to mesenchymal

transition (EMT; blue), heme metabolism (Heme; red), E2F targets (yellow) and

Leukemia stem cell and progenitor-primed (LSPC-primed; black). Genes up regulated in

IDH-mutant de novo AML are indicated as *.

(C) Heatmap of gene set normalized enrichment scores (NES) from the comparison of

the average fold changes *JAK2*V617F *IDH1*R132S vs *JAK2*V617F *IDH1*WT cells of a

patient with chronic phase MPN (SA2CP) in key transcriptional compartments. Darkblue

indicates higher NES in *JAK2*V617F *IDH1*R132S while dark red indicates higher NES in

*JAK2*V617F WT. Columns indicate key transcriptional compartments in CD34+ namely

hematopoietic stem cells and lympho-myeloid multipotent progenitor (HSC/LMPP),

megakaryocyte/erythroid-primed (Mega/Ery-Primed) and granulocytic/monocytic-primed

(Gran/Mono-Primed). False discovery rate (FDR) significance levels are indicated as

follows *<0.05; **<0.01; ***<0.001; ****<0.0001.

(D) Volcanoplot displaying the $\log_2$ fold change (x-axis) and $\log_{10}$ FDR (y-axis) from

comparing gene expression of a patients *IDH1*R132S mutant cells in chronic phase

MPN (SA2CP, higher expression marked in darkgreen) vs leukemia transformation

(SA2LT; higher expression marked in yellow). Colored and marked are top 70

significant different genes (FDR<0.05) as described in (B).

(E) Heatmap of gene set NES from the comparison of the average fold changes of

*IDH1*R132S cells in a patient at chronic phase MPN (SA2CP) and after leukemia

transformation (SA2LT) in key transcriptional compartments. Darkgreen indicates higher

NES in SA2CP while yellow indicates higher NES in SA2LT. Columns indicate key

transcriptional compartments in CD34+ namely HSC/LMPP, Mega/Ery-Primed and

Gran/Mono-Primed. FDR-significance levels are indicated as follows *<0.05; **<0.01;

<0.001; *<0.0001.

Figure 7. Leukemic transformation from chronic phase MPN showing parallel

clonal evolution of a *JAK2*V617F*-negative TP53-multi-hit* clone.

UMAPs of the same patient at diagnosis of (A-D) chronic phase MPN (SA1CP) and (E-

H) post-MPN AML (SA1LT), showing (A,E) transcriptionally defined clusters and

bargraphs with the percentage of cells in the clusters and (B-D,F-H) expressed variant

allele frequencies (eVAF) of mutations (B,F) *JAK2*V617F, (C,G) *TP53*Y236C and (D,H)

*TP53*R282W. Grey colored dots indicate non-genotyped cells.

(I) Fish plot illustrating the parallel clonal evolution of the *JAK2*V617F and *TP53*R282W

mutant clones.

Figure 1

Supplemental Figure 1

Study	Geno-typing molecule	#Total *JAK*2V617F MPN samples: # MPN subtypes	# *JAK*2V617F post-MPN AML samples	%*JAK*2 genotyped cells (Mean # JAK2 genotyped cells)	Phasing/ Isoform detection	Detection of rare variants	Typical # genes genotyped (coverage)
LOTR-Seq Grabek, Straube 2026	RNA	8: 1 ET 4 PV 1 PMF 2 MF	3	13-55% (1174)	+/+	+	30 (full length)
GoT Nam 2019	RNA	1 (ET)	0	7.3% (290)	(+)/(+) (limited to loci and amplicon size)	-	3-4 (selected loci)
Van Egeren 2021	RNA	7: 4 ET 3 PV	0	5-15% (733)	(+)/(+) (limited to loci and amplicon size)	-	3 (selected loci)
TargetSeq Rodriguez–Meira 2019	RNA+ DNA	12: 1 ET 1 PV 5 PMF 4 sMF 1 MDS/MPN MF	0	90-100% (168)	(+)/(+) (limited to loci and amplicon size)	-	4 (selected loci)
TargetSeq Rodriguez–Meira 2023	RNA+ DNA	2: 2 sMF (TP53 mutant from 2019 study)	11 (only TP53 mutant)	100% (748)	(+)/(+) (limited to loci and amplicon size)	-	2 (selected loci)
Tapestri Calabresi 2023	DNA (no transcript-omics)	6: 2 ET, 2 PMF 2 sMF	4	NA	(+)/(+) (limited to loci and amplicon size)	-	29-45 (selected loci)

Abbreviations: MPN: Myeloproliferative Neoplasms; PV: Polycythemia Vera; ET: Essential Thrombocythemia; PMF: Primary Myelofibrosis; sMF: secondary Myelofibrosis; MDS: Myelodysplastic Syndrome

Supplemental Figure 2

A

HealthyCD34+Young1 HealthyCD34+Young2 HealthyCD34+Young3 HealthyCD34+Elderly HealthyCD34+Elderly1 HealthyCD34+Elderly2

UMAP2

UMAP1

● HSC ● LMPP ● CLP ● B-primed ● GMP ● Mono
● DC ● Baso ● MEP ● MkP ● e-Ery ● l-Ery ● Ery

B

Split Identity

HSC
LMPP
CLP
B-primed
GMP
Mono
DC
Baso
MEP
MkP
e-Ery
l-Ery
Ery

CRHBP HOPX SPINK2 IGHM VPREB1 CD79B CSF3R ELANE MPO AZU1 CSTA LGALS1 IRF8 SPIB LMO4 CLC PBX1 ITGA2B KLF1 GATA1 HBD HBB AHSP CA1

Features

%
expressed

· 0
● 25
● 50
● 75
● 100

C

■ HSC/LMPP ■ Lymphoid-Primed ■ Gran/Mono-Primed ■ Mega/Ery-Primed

Celltype %

Healthy Young 1 Healthy Young 2 Healthy Young 3 Healthy Elderly Healthy Elderly1 Healthy Elderly2 MPN25 MPN15 MPN32 MPN36 SA2CP SA1CP MPN27 MPN18

Figure 2

Supplemental Figure 3

A CD34+ BM *JAK2V617F*
r=0.9
p=0.005
NGS gVAF%
Long Read eVAF%
CP-MPN
post-MPN AML

B CD34+ BM *JAK2V617F*
r=0.96
p<0.0001
Short Read eVAF%
Long Read eVAF%
CP-MPN
post-MPN AML

C *JAK2V617* locus
r=0.8
p=0.0006
% of cells genotyped
Gigabases PASS
PromethION 24
MinION Mk1b

D
PV ET PMF
sMF post-MPN AML
JAK2V617F % of genotyped cells
HSC/LMPP, Gran/Mono-Primed, Mega/Ery-Primed, Lymphoid-Primed

MPN25, MPN36, MPN27, MPN10, MPN15, SA2CP, MPN18, SA2LT, MPN32, SA1CP, MPN20, SA1LT

E PV
Healthy, JAK2V617F, JAK2WT
MkP cell fate score
Palantir pseudotime
Average MkP cell fate score
Healthy, JAK2V617F, JAK2WT
0.4, 0.4, 0.99

F sMF
Healthy, JAK2V617F, JAK2WT
MkP cell fate score
Palantir pseudotime
Average MkP cell fate score
Healthy, JAK2V617F, JAK2WT
0.00037, 0.036, 0.87

G sMF
Healthy, JAK2V617F, JAK2WT
Erythroid cell fate score
Palantir pseudotime
Average Erythroid cell fate score
Healthy, JAK2V617F, JAK2WT
0.12, 0.033, 0.34

Supplemental Figure 4

Figure 3

Supplemental Figure 5

A

	Median MinION	Median PromethION	
Gigabases PASS	5.25	45.15	
Assessed Gene Loci	Median MinION % genotyped cells	Median PromethION % genotyped cells	P-value Welch t-test, two-sided: MinION vs PromethION
*NPM1*W288	93.2	97.9	**0.0108**
*U2AF1*S34	90.9	94.1	0.0672
*CALR*K385	83.0	89.5	**0.0287**
*PTPN11*A72	58.5	85.3	**<0.0001**
*IDH2*R140	77.3	84.5	0.0626
*KRAS*G12	60.3	83.7	**0.0002**
*SRSF2*P95	71.0	82.0	**0.0080**
*SF3B1*K666	37.4	67.8	**<0.0001**
*ZRSR2*R19	32.8	61.2	**0.0219**
*MYC*P74	46.8	53.4	0.5358
*DNMT3A*R882	32.1	52.7	**0.0209**
*GATA2*A318	22.9	49.6	0.1743
*RUNX1*R201	28.8	48.5	**0.0194**
*NRAS*G12	25.9	48.4	**0.0042**
*EZH2*Y646	22.6	38.2	**0.0040**
*IDH1*R132	19.2	37.7	**0.0411**
*TET2*R1440	17.0	35.3	**0.0263**
*JAK2*V617	16.8	33.2	**0.0066**
*FLT3*D835	25.4	31.8	0.2886
*SH2B3*E208	22.0	28.3	0.0631
*PPM1D*Q150	11.8	21.9	**0.0028**
*KIT*D816	9.0	19.4	**0.0070**
*CSF3R*T618	8.1	18.9	0.0564
*STAG2*R614	7.4	16.4	**0.0150**
*TP53*C238	10.4	15.9	0.0512
*MPL*W515	1.5	5.5	0.1840
*BCOR*S336	1.6	5.0	**0.0082**
*CBL*C404	1.8	3.0	0.3707
*ASXL1*G642	1.2	2.8	**0.0134**

Figure 4

A — Bone Marrow Map - Zeng et al. 2025

B — CRHBP, HOPX, MPO, LYZ, KLF1, ITGA2B, LMO4, VPREB1

C — *JAK2*V617F post-MPN AML

HSC/LMPP/MEP/MKP/CLP — SA1LT *IDH1*R132S — Grabek,Straube 2026 — IF0392, IF0131, GR005 — TP53-multi-hit — Rodriguez-Meira 2021

MEP/Early/Late Erythroid — MPN20 *TP53*-multi-hit — Grabek,Straube 2026 — GR002, GR003, IF0393, GR007, GR006 — TP53-multi-hit — Rodriguez-Meira 2021

HSC/LMPP/CLP — MPN10 *IDH2*R140W *SRSF2*P95R — Grabek,Straube 2026 — GR004, GH001, IF0391 — TP53-multi-hit — Rodriguez-Meira 2021

D — Value -4 0 4

Study, Treatment, MPN, TP53mut, MPN Driver, Stage

scAML_HSCMPP, scAML_LMPP, scAML_MEPMkP, scAML_EarlyEry, scAML_LateEry, scAML_EoBasoMast, scAML_GMP, scAML_ProMono, scAML_Monocyte, scAML_cDC, scAML_EarlyLymphoid, scAML_ProBPreB, scAML_pDC

Study: Kong et al. 2025, Rodriguez-Meira et al. 2023
MPN treatment: Ruxo, HU, HU+Pipo, HU+Asp
Preceding MPN: PMF, SMF, PV, ET
TP53 mutation: Yes, No
MPN Driver: CALR52DEL, CALR5INS, JAK2V617F, MPL
Stage: Healthy, ET, PV, MF, post-MPN AML

Figure 5

Figure 6

Figure 7

Legend:
HSC, LMPP, CLP, B-primed, GMP, Mono
DC, Baso, MEP, MkP, e-Ery, l-Ery, Ery

A SA1CP
#cells 2072
% cells in cluster
100
0
SA1CP

B SA1CP
*JAK*2V617F eVAF
1.0
0.5
0.1
0.0
12.5% genotyped
44.3% *JAK*2V617F

C SA1CP
*TP53*Y236C eVAF
1.0
0.5
0.1
0.0
21.1% genotyped
54% *TP53*Y236C

D SA1CP
*TP53*R282W eVAF
1.0
0.5
0.1
0.0
23.2% genotyped
16% *TP53*R282W

E SA1LT
#cells 4061
% cells in cluster
100
0
SA1LT

F SA1LT
*JAK*2V617F eVAF
1.0
0.5
0.1
0.0
37% genotyped
0.8% *JAK*2V617F

G SA1LT
*TP53*Y236C eVAF
1.0
0.5
0.1
0.0
20.3% genotyped
52.1% *TP53*Y236C

H SA1LT
*TP53*R282W eVAF
1.0
0.5
0.1
0.0
22.3% genotyped
62% *TP53*R282W

I
TP53R282W
JAK2L393V
TP53Y236C
U2AF1S34Y
JAK2V617F
ET
Post-MPN AML

Supplemental Figure 6

Supplemental Figure 7