Curcumin-mediated photodynamic inactivation of *Escherichia coli*,
Pseudomonas fluorescens*, and *Candida auris

Jaime A. Garcia-Diosa,[a,1] Bhanu K. Pothineni,[a,2] and Adrian Keller[a,*]

[a] Paderborn University, Technical and Macromolecular Chemistry, Warburger Str. 100,
33098 Paderborn, Germany

* Correspondence: adrian.keller@uni-paderborn.de

[1] present address: Martin-Luther-Universität Halle-Wittenberg, Institut für Pharmazeutische
Technologie und Biopharmazie, Weinbergweg 22, D-06120 Halle (Saale), Germany

[2] present address: Programmable Biomaterials Laboratory, Institute of Materials, School of
Engineering, École Polytechnique Fédérale de Lausanne, Lausanne 1015, Switzerland

Abstract

Photodynamic therapy (PDT) is an emerging antimicrobial strategy that uses light to activate
photosensitizers, generating reactive oxygen species (ROS) that induce microbial cell death.
This study investigates the antimicrobial efficacy of curcumin-mediated PDT against the
microbial pathogens *Escherichia coli*, *Pseudomonas fluorescens*, and *Candida auris*. The
minimum inhibitory concentration (MIC) for each microorganism is determined at different
light fluence rates within a clinically relevant range. Our results demonstrate that curcumin-
mediated PDT effectively inactivates *E. coli* and *C. auris*, while exhibiting only partial
inhibition against *P. fluorescens* at curcumin concentrations up to 60 µM. Notably, within the
tested fluence rate range, the total light dose appears to be a more critical determinant of
antimicrobial efficacy than the specific fluence rate itself. The observed variations in
microbial susceptibility highlight the importance of species-specific factors, such as cell
wall/membrane structure. These findings provide guidelines for the application of curcumin-
mediated PDT in combatting antimicrobial resistance.

Keywords: curcumin; photosensitizer; antimicrobial photodynamic therapy; bacteria; fungi

1. Introduction

The escalating challenge of antimicrobial resistance (AMR) calls for novel therapeutic strategies to combat infections caused by bacteria, fungi, viruses, and parasites (Laxminarayan et al. 2016). Traditional antibiotic treatments are becoming increasingly ineffective, posing a significant threat to global health. In this context, photodynamic therapy (PDT) has emerged as a promising therapeutic modality that utilizes light to activate photosensitizing agents, generating reactive oxygen species (ROS) that damage essential cellular components and ultimately cause the death of the microorganism (Dougherty et al. 1998; Ghorbani et al. 2018). PDT has demonstrated efficacy against a broad spectrum of microorganisms, including multidrug-resistant bacterial and fungal strains, positioning it as a valuable tool in combating AMR (Youf et al. 2021). Unlike conventional antibiotics, the multi-target mechanism of action of PDT reduces the likelihood of resistance development (Wainwright 1998; Youf et al. 2021). Resistance to antimicrobial PDT (aPDT) is rarely reported, suggesting that while the possibility of microbial adaptation and escape from this treatment may exists, it is currently limited (Cieplik et al. 2018). This is a crucial advantage in the face of rapidly evolving resistance mechanisms in pathogenic bacteria and fungi.

Despite its advantages, aPDT faces a number of challenges that impede its widespread clinical adoption. Common photosensitizers (PSs) often suffer from issues such as limited tissue penetration, lack of selectivity, aggregation and quenching, and dark toxicity, resulting in intense research efforts dedicated to the development and discovery of new PSs (Klausen et al. 2020; Polat and Kang 2021; Huis In 't Veld et al. 2023). Curcumin, a naturally occurring polyphenol derived from turmeric (*Curcuma longa*), has emerged as a promising alternative due to several key advantages. These include high biocompatibility, low systemic toxicity, and synergistic therapeutic effects, such as anti-proliferative, anti-inflammatory, and anti-angiogenic activities (Dias et al. 2020). Consequently, it is investigated in various implementations of PDT, most importantly in cancer therapy (Xie et al. 2022) but also in the treatment of oral infections (Santezi et al. 2018). Furthermore, since curcumin is a common food ingredient and food additive, it is frequently used in the photodynamic inactivation of foodborne pathogens in various food products (Yu et al. 2022; Lan et al. 2023).

The PDT effect depends on the interaction of three essential components: molecular oxygen, the PS, and light. The PDT dose can be controlled by adjusting the latter two components. It is generally assumed that the biological effect of light in PDT is determined by the total light dose, regardless of the exposure time (Gao et al. 2020). Consequently, PDT studies often present data in terms of total light dose to compare the photoinactivation of microorganisms (Dias et al. 2020; Jao et al. 2023; Law et al. 2023). However, the fluence rate, *i.e.*, the power density of light, may influence PDT efficacy by affecting metabolic oxygen consumption and, consequently, the efficiency of ROS production (Nichols and Foster 1994; Henderson et al.

2006; Ghorbani et al. 2018). Indeed, some studies on the photodynamic treatment of cancer
cells and tumors have shown that lower fluence rates can enhance the cytotoxic effect of the
treatment by optimizing the depletion of the cell's antioxidant defenses (Angell-Petersen et
al. 2006; Yamamoto et al. 2006; Wang et al. 2007; Woodhams et al. 2007; Grossman et al.
2016; Dos Santos et al. 2020). However, there are only few reports on this phenomenon in
aPDT. Qin *et al.* conducted a study on the effect of different parameters of toluidine blue-
mediated PDT effectiveness, including fluence rate, on the inactivation of periodontal
pathogens. They observed that a high fluence rate (159 mW cm^{-2}) increases the cytotoxicity
of the treatment (Qin et al. 2008). A similar conclusion was reported by Prates *et al.*, who
studied the effects of fluence rate and light dose on the treatment of three different strains
of *Candida sp.* using methylene blue as the PS. In these experiments, a higher fluence rate
(300 mW cm^{-2}) lead to a higher cytotoxicity of the treatment, although the production of ROS
is inversely proportional to the fluence rate (Prates et al. 2009). Due to the limited number of
studies, the role of the fluence rate in aPDT is not fully understood. Additionally, these studies
used fluence rates that are much higher than those recommended to avoid possible damage
to host tissue in clinical applications (Sitnik and Henderson 1998; Jori et al. 2006).

In this study, we evaluate the efficacy of curcumin-mediated photodynamic inactivation of
three relevant pathogens by determining their minimum inhibitory concentrations (MICs) at
different fluence rates within a clinically relevant range. It has been observed before that
different bacterial species show markedly different susceptibilities to curcumin-mediated
photodynamic inactivation (Penha et al. 2017), which have to be considered in the
development of effective treatment protocols. For the current investigation, we have
selected *Escherichia coli*, *Pseudomonas fluorescens*, and *Candida auris* as the target
pathogens. *E. coli* is a Gram-negative bacterium that causes a wide range of infections
(Nataro and Kaper 1998; Rangel et al. 2005) and thus is a frequent target of aPDT. *P.*
fluorescens is a common food contaminant capable of growing at a wide range of
temperatures, making it a significant issue in both refrigerated and unrefrigerated food items
(Kumar et al. 2019). Curcumin-mediated inactivation of *P. fluorescens*, therefore, has been
investigated almost exclusively in food items (Saraiva et al. 2021; Saraiva et al. 2024; Zheng
et al. 2025). However, *P. fluorescens* can also cause severe infections in
immunocompromised individuals, most importantly bacteremia due to contaminated
medical equipment (Gershman et al. 2008), but also skin and soft tissue infections, as well
as respiratory tract infections (Liu et al. 2021), which warrants investigations also in the
context of aPDT settings. *C. auris* is a newly emerged fungal pathogen that has become a
global threat due to its rapid spread and multidrug resistance (Du et al. 2020). Photodynamic
inactivation of *C. auris* has already been demonstrated using several PSs (Bapat and Nobile
2021; Grizante Barião et al. 2022; Silva et al. 2023), but not with curcumin. By investigating
the combined effects of curcumin concentration and fluence rate on these target pathogens,

this study aims to contribute to a better understanding of the factors influencing the efficacy
of curcumin-mediated aPDT and its potential as a novel antimicrobial strategy to combat the
growing challenge of antimicrobial resistance.

2. Results and discussion

Figure 1 evaluates *E. coli* growth after 420 nm irradiation with a fluence rate of 17.8 mW cm^{-2}
and 76.3 mW cm^{-2}, respectively. The control samples incubated in the dark exhibit a relatively
stable ΔOD_{600} value of approximately 0.4, indicative of normal bacterial growth and no dark
cytotoxicity (see Figure 1A). In contrast, the samples irradiated at 420 nm with both fluence
rates (17.8 mW cm^{-2} and 76.3 mW cm^{-2}) demonstrate a sharp decline from ΔOD_{600} values of
0.4 to almost zero at curcumin concentrations of 7.5 µM and above (see Figure 1A). Notably,
no significant difference in bacterial growth inhibition is observed between the two fluence
rates. This suggests that the maximum aPDT effect is reached once the curcumin
concentration is 7.5 µM or above, regardless of the fluence rate within the therapeutic
window. The images of *E. coli* cultures on agar plates after 24 h of aPDT provide a visual
confirmation of the quantitative ΔOD_{600} data (see Figure 1B and Figure S1). The control plates
(black outline) show confluent bacteria at the maximum curcumin concentration
investigated, consistent with the ΔOD_{600} values. After aPDT, plates corresponding to
curcumin concentrations of 1.88 µM and 7.5 µM (blue and red outlines) show a clear
reduction in colony forming units compared to the control, consistent with the ΔOD_{600}
measurements.

Figure 1. *E. coli* growth after curcumin-mediated aPDT using 420 nm light at a dose of 20 J cm^{-2} with fluence rate of 17.8 mW cm^{-2} and 76.3 mW cm^{-2}, respectively, at different curcumin concentrations. A) Optical density change (ΔOD_{600}) after 24 h of incubation at 37 °C. Error bars indicate the standard deviation (n = 3). Controls were incubated in the dark. B) *E. coli* cultures (24 h) on agar after aPDT at different curcumin concentrations and fluence rates. Control (incubated in the dark, black outline), low fluence rate (blue outline), and high fluence rate (red outline).

Figure 2 presents the results of curcumin-mediated aPDT on *P. fluorescens* with both fluence rates at different curcumin concentrations. Again, the control samples exhibit a stable

ΔOD_{600} value, indicating consistent bacterial growth and no dark cytotoxicity. In contrast,

samples treated with aPDT demonstrate a concentration-dependent decrease of ΔOD_{600}

values, indicative of a reduction of bacterial viability (see Figure 2A). At the lower fluence rate

(17.8 mW cm^{-2}), a remarkable linear decrease of ΔOD_{600} is observed at curcumin

concentration higher than 15 µM. Similarly, at the highest fluence rate (76.3 mW cm^{-2}), ΔOD_{600}

values decrease with increasing curcumin concentration. However, this decrease starts

already at the lowest curcumin concentration investigated and appears to saturate for the

highest concentrations. Bacterial growth is detected even at 60 µM curcumin. This is

consistent with the *P. fluorescens* cultures on agar plates shown in Figure 2B and Figure S2.

Samples treated with PDT at 15 µM and 60 µM curcumin at both fluence rates (blue and red

outlines), show a clear reduction in colony forming units compared to the control, but no

complete bacterial inhibition is observed.

Figure 2. *P. fluorescens* growth after curcumin-mediated aPDT using 420 nm light at a dose of 20 J cm^{-2} with fluence rate of 17.8 mW cm^{-2} and 76.3 mW cm^{-2}, respectively, at different curcumin concentrations. A) Optical density change (ΔOD_{600}) after 24 h of incubation at 30 °C. Error bars indicate the standard deviation (n = 3). Controls were incubated in the dark. B) *P. fluorescens* cultures (48 h) on agar after aPDT at different curcumin concentrations and fluence rates. Control (incubated in the dark, black outline), low fluence rate (blue outline), and high fluence rate (red outline).

The results of curcumin-mediated aPDT on *C. auris* at low and high fluence rates are shown in Figure 3. The control samples exhibit a relatively stable ΔOD_{600} value around 0.65 – 0.70

across varying curcumin concentrations, showing consistent growth of the yeast and no dark

cytotoxicity (see Figure 3A). On the other hand, irradiated samples show a markedly

concentration-dependent decrease in ΔOD_{600} values. At both fluence rates (17.8 mW cm^{-2}

and 76.3 mW cm^{-2}), a strong reduction of the fungal growth is observed even at the lowest

curcumin concentration used (1.87 µM). When the curcumin concentration reaches 15 µM

and more, the ΔOD_{600} values approaches zero, indicating a near-complete inhibition of fungal

growth (see Figure 3A). Notably, there is again no significant difference between the two

fluence rates at any of the tested curcumin concentrations. This suggests that within the

tested fluence range, the curcumin concentration is the primary parameter of the

photodynamic effect. *C. auris* cultures on agar plates after aPDT visually confirm the trends

observed in the ΔOD_{600} measurements (see Figure 3B and Figure S3). The control group

shows dense *C. auris* growth, consistent with the high ΔOD_{600} values. In the case of the

samples irradiated at both fluence rates (blue and red outlines), a strong reduction in the

number of colonies is observed at 3.75 µM, whereas at 15 µM, visible colonies are almost

complete absent (see Figure 3B and Figure S3), corroborating the trend of the ΔOD_{600} values.

Figure 3. *C. auris* growth after curcumin-mediated aPDT using 420 nm light at a dose of 20 J cm^{-2} using with fluence rate of 17.8 mW cm^{-2} and 76.3 mW cm^{-2}, respectively, at different curcumin concentrations. A) Optical density change (ΔOD_{600}) after 24 h of incubation at 37 °C. Error bars indicate the standard deviation (n = 3). Controls were incubated in the dark. B) *C. auris* cultures (24 h) on agar after aPDT at different curcumin concentrations and fluence rates. Control (incubated in the dark, black outline), low fluence rate (blue outline), and high fluence rate (red outline).

Table 1 summarizes the MIC of curcumin during aPDT for *E. coli*, *P. fluorescens*, and *C. auris* as determined from the OD_{600} results. The data reveal a remarkable difference in the

susceptibility of the three microorganisms to curcumin-mediated aPDT. *E. coli* exhibits the
highest sensitivity of the treatment, with a MIC of 7.5 µM, implying that a relatively low
concentration of curcumin is sufficient to inhibit *E. coli* growth. *C. auris* demonstrates
intermediate susceptibility with a MIC of 15 µM, requiring a higher curcumin concentration
to effectively inhibit the growth of the yeast. On the other hand, *P. fluorescens* shows the
lowest susceptibility to curcumin-mediated aPDT with a MIC greater than 60 µM, indicating
that *P. fluorescens* is significantly more resistant to the photodynamic effects of the PS
curcumin under the studied conditions.

Table 1. Minimum inhibitory concentrations (MICs) of curcumin during aPDT using a
fluence rate range of 17.8 – 76.3 mW cm^{-2}, and a total light dose of 20 J cm^{-2}.

Microorganism	MIC (µM)
E. coli	7.5
P. fluorescens	> 60.0
C. auris	15.0

Our results demonstrate that curcumin-mediated aPDT is effective in inactivating *E. coli* and
C. auris, and partially inactivating *P. fluorescens* at curcumin concentrations up to 60 µM.
These results are consistent with previous studies, which have reported varying degrees of
susceptibility to curcumin-mediated aPDT among different microorganisms (Dovigo,
Pavarina and Ribeiro et al. 2011; Andrade et al. 2013; Penha et al. 2017; Saraiva et al. 2021;
Saraiva et al. 2024). For instance, *P. fluorescens* has been shown to be more resistant to
curcumin-mediated aPDT compared to *E. coli* and *C. albicans* (Saraiva et al. 2021; Saraiva et
al. 2024). This difference in susceptibility may be attributed to several factors, including
variations in cell wall/membrane structure and composition, and variations in the uptake and
intracellular accumulation of curcumin among the different microorganisms. In general,
curcumin can interact with microorganisms through several mechanisms, including charge
interactions, which are pH-dependent due to keto-enol tautomerism (Priyadarsini 2009),
binding to various enzymes such as sortase A (Hu et al. 2013; Praditya et al. 2019) and FtsZ
(Kaur et al. 2010), and membrane adsorption and subsequent uptake into the cell. Species-
specific differences may play a role in all these mechanisms but are particularly important
for cellular uptake (Annunzio et al. 2018).

This study highlights the importance of species-specific variations in susceptibility to
curcumin-mediated aPDT. Despite sharing a common Gram-negative cell wall structure with

an outer membrane containing lipopolysaccharides (LPSs), *E. coli* and *P. fluorescens* display
markedly different responses to aPDT with curcumin (see Table 1). While the precise
mechanisms governing the interaction of curcumin with these species remain to be fully
elucidated, the high efficacy of curcumin-mediated aPDT against *E. coli* suggests that
curcumin can effectively penetrate the outer membrane, likely via charge-mediated
interactions, leading to subsequent membrane disruption by the generated ROS (Oliveira et
al. 2018; Gao et al. 2019). In contrast, the significantly elevated MIC for *P. fluorescens*
suggests a potential limitation in curcumin accumulation within the cell, probably due to
reduced uptake or increased efflux mechanisms.

In the case of *C. auris*, a cell wall composed of chitin, glucans, and mannans is present,
which differs significantly from bacterial cell walls (Free 2013). However, the intermediate
susceptibility to curcumin-mediated aPDT compared to *E. coli* and *P. fluorescens* (see Table
1), suggests that curcumin can effectively penetrate the *C. auris* cell wall and accumulate
within the cell at a higher rate than for *P. fluorescens*. This is in line with previous observations
of effective curcumin-mediated photodynamic inactivation of various *Candida* species
(Dovigo, Pavarina and Carmello et al. 2011; Dovigo, Pavarina and Ribeiro et al. 2011; Andrade
et al. 2013). The observed variations in susceptibility among the different microorganisms
underscore the importance of considering species-specific factors, such as cell
wall/membrane structure, and curcumin uptake mechanisms, when evaluating the
effectiveness of aPDT.

Regarding the fluence rate effect on the PDT efficiency, several studies in cancer treatment
suggest that lower fluence rates at the same total light dose with a longer treatment time can
be more efficient in PDT compared to higher fluence rates (Henderson et al. 2006; Dos
Santos et al. 2020; Gao et al. 2020). The reasons behind this effect of the fluence rate are still
under investigation, but some studies suggest that oxygen availability plays an important
role. Lower fluence rates would allow for more efficient oxygen diffusion into the target tissue,
whereas high fluence rates lead to the depletion of available oxygen, thereby limiting ROS
generation and ultimately reducing PDT efficacy (Dougherty et al. 1998; Alvarez and Sevilla
2024). Conversely, some reports suggest an inverse relationship between fluence rate and
aPDT efficiency (Qin et al. 2008; Prates et al. 2009). While higher fluence rates may lead to
increased PS bleaching, they have also been associated with greater microbial inhibition
(Prates et al. 2009). This apparent contradiction may be explained by a potential initial surge
in ROS formation at higher fluence rates, followed by photobleaching. However, increasing
the fluence rate infinitely to enhance aPDT efficacy is not feasible. Photon absorption by the
PS is limited, restricting ROS production, and the photobleaching effect becomes
increasingly significant (Qin et al. 2008). Furthermore, to minimize host tissue damage, the

therapeutic window for aPDT is generally considered to be below approximately 50 mW cm^{-2} (Jori et al. 2006).

In this study, the tested fluence rates (17.8 – 76.3 mW cm^{-2}) did not significantly influence the antimicrobial activity within the tested concentration range. This finding suggests that the moderate changes of the fluence rate possible within the therapeutic window are insufficient to induce significant variations in ROS generation. Under these conditions, the total light dose thus appears to be the primary determinant of aPDT efficacy. Nevertheless, future studies should investigate a wider range of fluence rates and total light doses to further elucidate the impact of these parameters on aPDT efficacy. Furthermore, investigating the underlying mechanisms of curcumin-mediated aPDT in these and other microorganisms, including the role of specific cellular targets and the generation of ROS, will provide valuable insights for optimizing this approach and developing novel antimicrobial strategies.

3. Experimental

3.1. Bacteria culture

E. coli W1485 (DSM 5695) and *P. fluorescens* 1175 (DMS 1976) were purchased from Leibniz Institute DSMZ. For bacteria culture preparation, two or three bacterial colonies were suspended in 30 mL LB medium (MP Biomedicals) and incubated overnight at 37 °C for *E. coli* or 24 h at 30 °C for *P. fluorescens*. In the case of *E. coli*, 250 μL overnight suspension was used to inoculate fresh LB medium and incubated for 4-6 h at 37 °C. For *P. fluorescens*, the culture incubated for 24 h was used directly. Lastly, the bacteria suspensions were centrifuged, washed, and resuspended in 5 % dimethyl sulfoxide (DMSO, Carl Roth), adjusting the optical density at 600 nm (OD$_{600}$) to 0.2. DMSO was used as a solvent to ensure the solubility of curcumin (Saraiva et al. 2021). At a concentration of 5 %, DMSO has been reported to exert minimal cytotoxic effects on bacteria (Ansel et al. 1969; Tunçer Çağlayan and Gurbanov 2024).

3.2. Fungi culture

C. auris (DSM 105990) was purchased from Leibniz Institute DSMZ. The fungi culture was prepared using two or three colonies, which were resuspended in 30 mL yeast extract peptone dextrose medium (YPD medium, SERVA) and incubated at 180 rpm and 37 °C overnight. Finally, the culture was centrifuged, washed, and resuspended in 5 % dimethyl sulfoxide (DMSO), adjusting the optical density (OD$_{600}$) to 0.2.

3.3. Antimicrobial photodynamic inactivation

Photodynamic inactivation was performed using a Lumidox II 96-position LED array 420 nm
with cooling base (Analytical Sales and Services) with Curcumin (Sigma Aldrich) as the PS.
The experiments were performed in triplicate using the microdilution method in sterile 96-
wells microplates. 50 μL aliquots of microbial suspension in 5 % DMSO (OD_{600} = 0.2) were
added to 50 μL of curcumin in 5 % DMSO. The microplates were incubated for 30 min at 25 °C
before photodynamic inactivation. Control samples without curcumin and light irradiation
were also prepared to analyze the curcumin toxicity and light-induced inhibition.

The fluence rate effect in the photodynamic inactivation was evaluated at a constant light
dose of 20 J cm^{-2} using the lowest and highest fluence rate levels available in the Lumidox II
equipment. Since the 96-well microplate was held at 17 mm from the light source, the
samples were irradiated for 1123 s at the lowest fluence rate (17.8 mW cm^{-2}), and for 262 s at
highest fluence rate (76.3 mW cm^{-2}), obtaining a light dose of 20 J cm^{-2} at both fluence rates.
At these short exposure times, the potential impact of prolonged DMSO exposure on
bacterial growth is minimized. For subsequent minimum inhibitory concentration (MIC)
assays, the DMSO concentration was reduced to 1%, a level generally considered to have
negligible or no effect on bacterial viability (Ansel et al. 1969; Liu et al. 2023). Furthermore,
all photodynamic inactivation experiments were conducted in the absence of bacterial
growth media to prevent light absorption by medium components, and to minimize
interactions with ROS, which could ultimately reduce the effectiveness of PDT.

3.4. Optical density measurements

Photodynamic inactivation at the different fluence rates was studied by measuring the
change in the OD_{600} (ΔOD_{600}) of the treated samples after 24 h of incubation in their respective
media. After treatment, a 20 μL aliquot of each sample was added to 80 μL (1 % DMSO final
concentration) of LB (*E. coli and P. fluorescens*) or YPD medium (*C. auris*), in sterile 96-well
microplates. Then, the microplates were incubated at 37 °C (*E. coli* and *C. auris*) or at 30 °C
(*P. fluorescens*). The OD_{600} values were measured at the beginning and after 24 h of
incubation using a microplate reader TrisStar^2 S LB 942 (Berthold technologies). The ΔOD_{600}
value is an indicator of the growth or inhibition of microbes.

3.5. Agar plate culture

For visual inspection of photodynamic inactivation, 80 μL aliquots of treated samples were
spread on agar plates and incubated. *E. coli* was incubated on LB agar plates at 37 °C for 24
h. *P. fluorescens* was incubated on LB agar plates at 30 °C for 48 h. *C. auris* was incubated
on YPD agar plates at 37 °C for 24 h.

3.6. Statistical analysis

The change in optical density at 600 nm (ΔOD_{600}) was measured in triplicate for each
experimental condition (n = 3). Data are presented as mean ± standard deviation (SD) of the
three values.

4. Conclusions

This study investigated the antimicrobial efficacy of curcumin-mediated aPDT against *E. coli*,
P. fluorescens, and *C. auris* using a range of curcumin concentrations and two distinct
fluence rates. Our findings demonstrate that curcumin-mediated aPDT is able to effectively
inactivate *E. coli* and *C. auris*, while leading only to partial inhibition of *P. fluorescens* at
curcumin concentrations up to 60 µM. Notably, within the tested fluence rate range (17.8 –
76.3 mW cm^{-2}), no significant difference in antimicrobial activity was observed, suggesting
that within the therapeutic window, the total light dose is a more critical determinant of aPDT
efficacy than the specific fluence rate itself.

The observed variations in microbial susceptibility to curcumin-mediated aPDT highlight the
importance of species-specific factors, such as cell wall/membrane structure, and
curcumin uptake and accumulation. While the mechanisms responsible for the observed
differences remain to be fully elucidated, our results suggest that curcumin effectively
penetrates the bacterial cell wall of *E. coli* and the fungal cell wall of *C. auris*, leading to
significant growth inhibition. In contrast, the limited efficacy against *P. fluorescens* may be
attributed to reduced curcumin uptake or increased efflux mechanisms.

These findings provide valuable insights into the potential of curcumin-mediated aPDT as an
antimicrobial strategy. However, further research is warranted in order to optimize treatment
parameters, including a comprehensive evaluation of the effects of different total light doses
and a wider range of fluence rates. Additionally, investigating the underlying mechanisms of
curcumin-mediated aPDT in these microorganisms, including the specific cellular targets
and the generation of ROS, will be crucial for developing more effective and targeted
antimicrobial therapies**.**

Disclosure statement: The authors report there are no competing interests to declare.

Funding details: This work was supported by the Alexander von Humboldt Foundation
(grant number COL-1224696-GF-P, to J.A.G.-D.).

References

Alvarez N, Sevilla A. 2024. Current Advances in Photodynamic Therapy (PDT) and the Future
Potential of PDT-Combinatorial Cancer Therapies. Int. J. Mol. Sci. 25:1023. Epub 2024
Jan 13. eng. doi:10.3390/ijms25021023.

Andrade MC, Ribeiro APD, Dovigo LN, Brunetti IL, Giampaolo ET, Bagnato VS, Pavarina AC.
2013. Effect of different pre-irradiation times on curcumin-mediated photodynamic
therapy against planktonic cultures and biofilms of Candida spp. Arch. Oral Biol.
58:200–210. Epub 2012 Nov 13. eng. doi:10.1016/j.archoralbio.2012.10.011.

Angell-Petersen E, Spetalen S, Madsen SJ, Sun C-H, Peng Q, Carper SW, Sioud M,
Hirschberg H. 2006. Influence of light fluence rate on the effects of photodynamic
therapy in an orthotopic rat glioma model. J. Neurosurg. 104:109–117. eng.
doi:10.3171/jns.2006.104.1.109.

Annunzio SR de, Freitas LM de, Blanco AL, da Costa MM, Carmona-Vargas CC, Oliveira KT
de, Fontana CR. 2018. Susceptibility of Enterococcus faecalis and Propionibacterium
acnes to antimicrobial photodynamic therapy. J. Photochem. Photobiol., B. 178:545–
550. Epub 2017 Dec 5. eng. doi:10.1016/j.jphotobiol.2017.11.035.

Ansel HC, Norred WP, Roth IL. 1969. Antimicrobial activity of dimethyl sulfoxide against
Escherichia coli, Pseudomonas aeruginosa, and Bacillus megaterium. J. Pharm. Sci.
58:836–839. eng. doi:10.1002/jps.2600580708.

Bapat PS, Nobile CJ. 2021. Photodynamic Therapy Is Effective Against Candida auris
Biofilms. Front. Cell Infect. Microbiol. 11:713092. Epub 2021 Sep 3. eng.
doi:10.3389/fcimb.2021.713092.

Cieplik F, Deng D, Crielaard W, Buchalla W, Hellwig E, Al-Ahmad A, Maisch T. 2018.
Antimicrobial photodynamic therapy - what we know and what we don't. Crit. Rev.
Microbiol. 44:571–589. Epub 2018 May 11. eng. doi:10.1080/1040841X.2018.1467876.

Dias LD, Blanco KC, Mfouo-Tynga IS, Inada NM, Bagnato VS. 2020. Curcumin as a
photosensitizer: From molecular structure to recent advances in antimicrobial
photodynamic therapy. J. Photochem. Photobiol., C. 45:100384.
doi:10.1016/j.jphotochemrev.2020.100384.

Dos Santos AF, Almeida DRQ de, Terra LF, Wailemann RAM, Gomes VM, Arini GS, Ravagnani
FG, Baptista MS, Labriola L. 2020. Fluence Rate Determines PDT Efficiency in Breast
Cancer Cells Displaying Different GSH Levels. Photochem Photobiol. 96:658–667. Epub
2019 Dec 17. eng. doi:10.1111/php.13182.

Dougherty TJ, Gomer CJ, Henderson BW, Jori G, Kessel D, Korbelik M, Moan J, Peng Q. 1998.
Photodynamic therapy. J. Natl. Cancer Inst. 90:889–905. eng.
doi:10.1093/jnci/90.12.889.

Dovigo LN, Pavarina AC, Carmello JC, Machado AL, Brunetti IL, Bagnato VS. 2011.
Susceptibility of clinical isolates of Candida to photodynamic effects of curcumin.
Lasers Surg. Med. 43:927–934. eng. doi:10.1002/lsm.21110.
Dovigo LN, Pavarina AC, Ribeiro APD, Brunetti IL, Costa CAdS, Jacomassi DP, Bagnato VS,
Kurachi C. 2011. Investigation of the photodynamic effects of curcumin against Candida
albicans. Photochem Photobiol. 87:895–903. Epub 2011 Jun 13. eng.
doi:10.1111/j.1751-1097.2011.00937.x.
Du H, Bing J, Hu T, Ennis CL, Nobile CJ, Huang G. 2020. Candida auris: Epidemiology,
biology, antifungal resistance, and virulence. PLoS Pathog. 16:e1008921. Epub 2020 Oct
22. eng. doi:10.1371/journal.ppat.1008921.
Free SJ. 2013. Fungal cell wall organization and biosynthesis. Adv. Genet. 81:33–82. eng.
doi:10.1016/B978-0-12-407677-8.00002-6.
Gao R, Xu H, Liu L, Zhang Y, Yin T, Zhou H, Sun M, Chen N, Ren Y, Chen T, et al. 2020.
Photoacoustic visualization of the fluence rate dependence of photodynamic therapy.
Biomed. Opt. Express. 11:4203–4223. Epub 2020 Jul 10. eng. doi:10.1364/BOE.395562.
Gao Y, Wu J, Li Z, Zhang X, Lu N, Xue C, Leung AW, Xu C, Tang Q. 2019. Curcumin-mediated
photodynamic inactivation (PDI) against DH5α contaminated in oysters and cellular
toxicological evaluation of PDI-treated oysters. Photodiagnosis Photodyn. Ther. 26:244–
251. Epub 2019 Apr 3. eng. doi:10.1016/j.pdpdt.2019.04.002.
Gershman MD, Kennedy DJ, Noble-Wang J, Kim C, Gullion J, Kacica M, Jensen B, Pascoe N,
Saiman L, McHale J, et al. 2008. Multistate outbreak of Pseudomonas fluorescens
bloodstream infection after exposure to contaminated heparinized saline flush prepared
by a compounding pharmacy. Clin. Infect. Dis. 47:1372–1379. eng. doi:10.1086/592968.
Ghorbani J, Rahban D, Aghamiri S, Teymouri A, Bahador A. 2018. Photosensitizers in
antibacterial photodynamic therapy: an overview. Laser Ther. 27:293–302. eng.
doi:10.5978/islsm.27_18-RA-01.
Grizante Barião PH, Tonani L, Brancini GTP, Nascimento E, Braga GÚL, Wainwright M, Zeska
Kress MR von. 2022. In vitro and in vivo photodynamic efficacies of novel and
conventional phenothiazinium photosensitizers against multidrug-resistant Candida
auris. Photochem. Photobiol. Sci. 21:1807–1818. Epub 2022 Jul 11. eng.
doi:10.1007/s43630-022-00258-4.
Grossman CE, Carter SL, Czupryna J, Le Wang, Putt ME, Busch TM. 2016. Fluence Rate
Differences in Photodynamic Therapy Efficacy and Activation of Epidermal Growth
Factor Receptor after Treatment of the Tumor-Involved Murine Thoracic Cavity. Int. J.
Mol. Sci. 17:101. Epub 2016 Jan 14. eng. doi:10.3390/ijms17010101.
Henderson BW, Busch TM, Snyder JW. 2006. Fluence rate as a modulator of PDT
mechanisms. Lasers Surg. Med. 38:489–493. eng. doi:10.1002/lsm.20327.

Hu P, Huang P, Chen MW. 2013. Curcumin reduces Streptococcus mutans biofilm
formation by inhibiting sortase A activity. Arch. Oral Biol. 58:1343–1348. Epub 2013 Jun
16. eng. doi:10.1016/j.archoralbio.2013.05.004.

Huis In 't Veld RV, Heuts J, Ma S, Cruz LJ, Ossendorp FA, Jager MJ. 2023. Current Challenges
and Opportunities of Photodynamic Therapy against Cancer. Pharmaceutics. 15:330.
Epub 2023 Jan 18. eng. doi:10.3390/pharmaceutics15020330.

Jao Y, Ding S-J, Chen C-C. 2023. Antimicrobial photodynamic therapy for the treatment of
oral infections: A systematic review. J. Dent. Sci. 18:1453–1466. Epub 2023 Jul 11. eng.
doi:10.1016/j.jds.2023.07.002.

Jori G, Fabris C, Soncin M, Ferro S, Coppellotti O, Dei D, Fantetti L, Chiti G, Roncucci G.
2006. Photodynamic therapy in the treatment of microbial infections: basic principles
and perspective applications. Lasers Surg. Med. 38:468–481. eng.
doi:10.1002/lsm.20361.

Kaur S, Modi NH, Panda D, Roy N. 2010. Probing the binding site of curcumin in Escherichia
coli and Bacillus subtilis FtsZ--a structural insight to unveil antibacterial activity of
curcumin. Eur. J. Med. Chem. 45:4209–4214. Epub 2010 Jun 22. eng.
doi:10.1016/j.ejmech.2010.06.015.

Klausen M, Ucuncu M, Bradley M. 2020. Design of Photosensitizing Agents for Targeted
Antimicrobial Photodynamic Therapy. Molecules. 25:5239. Epub 2020 Nov 10. eng.
doi:10.3390/molecules25225239.

Kumar H, Franzetti L, Kaushal A, Kumar D. 2019. Pseudomonas fluorescens: a potential
food spoiler and challenges and advances in its detection. Ann. Microbiol. 69:873–883.
doi:10.1007/s13213-019-01501-7.

Lan X, Liu Y, Wang L, Wang H, Hu Z, Dong H, Yu Z, Yuan Y. 2023. A review of curcumin in food
preservation: Delivery system and photosensitization. Food Chem. 424:136464. Epub
2023 May 26. eng. doi:10.1016/j.foodchem.2023.136464.

Law SK, Leung AWN, Xu C. 2023. Photodynamic Action of Curcumin and Methylene Blue
against Bacteria and SARS-CoV-2-A Review. Pharmaceuticals. 17:13. Epub 2023 Dec 25.
eng. doi:10.3390/ph17010034.

Laxminarayan R, Matsoso P, Pant S, Brower C, Røttingen J-A, Klugman K, Davies S. 2016.
Access to effective antimicrobials: a worldwide challenge. Lancet. 387:168–175. Epub
2015 Nov 18. eng. doi:10.1016/S0140-6736(15)00474-2.

Liu J, Zhang H, Zhang L, Li T, Liu N, Liu Q. 2023. Effect of various concentrations of common
organic solvents on the growth and proliferation ability of Candida glabrata and their
permissible limits for addition in drug susceptibility testing. PeerJ. 11:e16444. Epub
2023 Nov 21. eng. doi:10.7717/peerj.16444.

Liu X, Xiang L, Yin Y, Li H, Ma D, Qu Y. 2021. Pneumonia caused by Pseudomonas
fluorescens: a case report. BMC Pulm. Med. 21:212. Epub 2021 Jul 5. eng.
doi:10.1186/s12890-021-01573-9.

Nataro JP, Kaper JB. 1998. Diarrheagenic Escherichia coli. Clin. Microbiol. Rev. 11:142–201.
eng. doi:10.1128/CMR.11.1.142.

Nichols MG, Foster TH. 1994. Oxygen diffusion and reaction kinetics in the photodynamic
therapy of multicell tumour spheroids. Phys. Med. Biol. 39:2161–2181. eng.
doi:10.1088/0031-9155/39/12/003.

Oliveira EF de, Tosati JV, Tikekar RV, Monteiro AR, Nitin N. 2018. Antimicrobial activity of
curcumin in combination with light against Escherichia coli O157:H7 and Listeria
innocua: Applications for fresh produce sanitation. Postharvest Biol. Biotechnol.
137:86–94. doi:10.1016/j.postharvbio.2017.11.014.

Penha CB, Bonin E, da Silva AF, Hioka N, Zanqueta ÉB, Nakamura TU, Abreu Filho BA de,
Campanerut-Sá PAZ, Mikcha JMG. 2017. Photodynamic inactivation of foodborne and
food spoilage bacteria by curcumin. LWT. 76:198–202. doi:10.1016/j.lwt.2016.07.037.

Polat E, Kang K. 2021. Natural Photosensitizers in Antimicrobial Photodynamic Therapy.
Biomedicines. 9:584. Epub 2021 May 21. eng. doi:10.3390/biomedicines9060584.

Praditya D, Kirchhoff L, Brüning J, Rachmawati H, Steinmann J, Steinmann E. 2019. Anti-
infective Properties of the Golden Spice Curcumin. Front. Microbiol. 10:912. Epub 2019
May 3. eng. doi:10.3389/fmicb.2019.00912.

Prates RA, Silva EG, Yamada AM, Suzuki LC, Paula CR, Ribeiro MS. 2009. Light parameters
influence cell viability in antifungal photodynamic therapy in a fluence and rate fluence-
dependent manner. Laser Phys. 19:1038–1044. doi:10.1134/S1054660X09050284.

Priyadarsini KI. 2009. Photophysics, photochemistry and photobiology of curcumin: Studies
from organic solutions, bio-mimetics and living cells. J. Photochem. Photobiol., C.
10:81–95. doi:10.1016/j.jphotochemrev.2009.05.001.

Qin Y, Luan X, Bi L, He G, Bai X, Zhou C, Zhang Z. 2008. Toluidine blue-mediated
photoinactivation of periodontal pathogens from supragingival plaques. Lasers Med.
Sci. 23:49–54. Epub 2007 Mar 15. eng. doi:10.1007/s10103-007-0454-x.

Rangel JM, Sparling PH, Crowe C, Griffin PM, Swerdlow DL. 2005. Epidemiology of
Escherichia coli O157:H7 outbreaks, United States, 1982-2002. Emerg. Infect. Dis.
11:603–609. eng. doi:10.3201/eid1104.040739.

Santezi C, Reina BD, Dovigo LN. 2018. Curcumin-mediated Photodynamic Therapy for the
treatment of oral infections-A review. Photodiagnosis Photodyn. Ther. 21:409–415. Epub
2018 Feb 22. eng. doi:10.1016/j.pdpdt.2018.01.016.

Saraiva BB, Campanholi KdSS, Machado RRB, Nakamura CV, Silva AA, Caetano W, Pozza
MSdS. 2024. Reducing Pseudomonas fluorescens in milk through photodynamic

inactivation using riboflavin and curcumin with 450 nm blue light-emitting diode. Int.
Dairy J. 148:105787. doi:10.1016/j.idairyj.2023.105787.

Saraiva BB, Rodrigues BM, da Silva Junior RC, Scapim MRdS, Lancheros CAC, Nakamura
CV, Caetano W, Pereira PCdS, Santana EHW de, Pozza MSdS. 2021. Photodynamic
inactivation of Pseudomonas fluorescens in Minas Frescal cheese using curcumin as a
photosensitizer. LWT. 151:112143. doi:10.1016/j.lwt.2021.112143.

Silva AR, Cabral FV, Silva CR, Silva DFT, Freitas AZ, Fontes A, Ribeiro MS. 2023. New Insights
in Phenothiazinium-Mediated Photodynamic Inactivation of Candida Auris. J. Fungi.
9:717. Epub 2023 Jun 30. eng. doi:10.3390/jof9070717.

Sitnik TM, Henderson BW. 1998. The Effect of Fluence Rate on Tumor and Normal Tissue
Responses to Photodynamic Therapy. Photochem Photobiol. 67:462–466.
doi:10.1111/j.1751-1097.1998.tb05228.x.

Tunçer Çağlayan S, Gurbanov R. 2024. Modulation of bacterial membranes and cellular
macromolecules by dimethyl sulfoxide: A dose-dependent study providing novel
insights. Int. J. Biol. Macromol. 267:131581. Epub 2024 Apr 12. eng.
doi:10.1016/j.ijbiomac.2024.131581.

Wainwright M. 1998. Photodynamic antimicrobial chemotherapy (PACT). J. Antimicrob.
Chemother. 42:13–28. eng. doi:10.1093/jac/42.1.13.

Wang H-W, Rickter E, Yuan M, Wileyto EP, Glatstein E, Yodh A, Busch TM. 2007. Effect of
photosensitizer dose on fluence rate responses to photodynamic therapy. Photochem
Photobiol. 83:1040–1048. eng. doi:10.1111/j.1751-1097.2007.00139.x.

Woodhams JH, Macrobert AJ, Bown SG. 2007. The role of oxygen monitoring during
photodynamic therapy and its potential for treatment dosimetry. Photochem. Photobiol.
Sci. 6:1246–1256. Epub 2007 Sep 21. eng. doi:10.1039/b709644e.

Xie L, Ji X, Zhang Q, Wei Y. 2022. Curcumin combined with photodynamic therapy,
promising therapies for the treatment of cancer. Biomed. Pharmacother. 146:112567.
Epub 2021 Dec 22. eng. doi:10.1016/j.biopha.2021.112567.

Yamamoto J, Yamamoto S, Hirano T, Li S, Koide M, Kohno E, Okada M, Inenaga C, Tokuyama
T, Yokota N, et al. 2006. Monitoring of singlet oxygen is useful for predicting the
photodynamic effects in the treatment for experimental glioma. Clin. Cancer Res.
12:7132–7139. eng. doi:10.1158/1078-0432.CCR-06-0786.

Youf R, Müller M, Balasini A, Thétiot F, Müller M, Hascoët A, Jonas U, Schönherr H,
Lemercier G, Montier T, et al. 2021. Antimicrobial Photodynamic Therapy: Latest
Developments with a Focus on Combinatory Strategies. Pharmaceutics. 13:1995. Epub
2021 Nov 24. eng. doi:10.3390/pharmaceutics13121995.

Yu X, Zou Y, Zhang Z, Wei T, Ye Z, Yuk H-G, Zheng Q. 2022. Recent advances in antimicrobial
applications of curcumin-mediated photodynamic inactivation in foods. Food Control.
138:108986. doi:10.1016/j.foodcont.2022.108986.

Zheng S, Yan H, Deng P, Zhu J, Lu H. 2025. Enhancement of curcumin-mediated
photodynamic inactivation against Pseudomonas fluorescens on fish fillets combined
with ε-polylysine hydrochloride. LWT:117344. doi:10.1016/j.lwt.2025.117344.
542