Sleep Renormalizes Negative Emotional Generalization

Ella Bar[1,2,3], Max Bringmann[1], Gennadiy Belonosov[2.3], Kristoffer Aberg[1], Reut Weissgross[1], Yuval Nir[2, 4,5,6, *], Rony Paz[1, *]

[1] Department of Brain Sciences, Weizmann Institute of Science, Rehovot, Israel.
[2] Department of Physiology and Pharmacology, Gray Faculty of Medical and Health Sciences, Tel Aviv University, Israel, Tel Aviv, Israel.
[3] Sagol School of Neuroscience, Tel Aviv University, Tel Aviv, Israel. [4] Sagol Brain Institute, Tel Aviv Sourasky Medical Center, Tel Aviv, Israel.
[5] Department of Biomedical Engineering, Faculty of Engineering, Tel Aviv University, Tel Aviv, Israel.
[6] The Sieratzki-Sagol Center for Sleep Medicine, Tel Aviv Sourasky Medical Center, Tel Aviv, Israel.
* Equal senior contribution

Abstract

Overgeneralization of negative experiences, in which aversive responses spread to otherwise safe stimuli, often co-occurs with sleep disruption, and both are central features of anxiety and posttraumatic stress disorder (PTSD). Here, we show that sleep shifts emotional generalization away from the negative and toward the positive. Across three experiments combining behavior, fMRI, and sleep electrophysiology, participants learned to associate three faces with positive, negative, or neutral outcomes. Participants were tested using morphed faces that blended the original stimuli in varying proportions. Immediate generalization was assessed post-learning, and delayed generalization was assessed after overnight sleep or daytime wakefulness. Behaviorally, we found that sleep selectively promotes positive generalization, whereas prolonged daytime wakefulness favors generalization of the negative face. In the fMRI scanner, amygdala and limbic activity during outcome processing predicted stronger immediate generalization of the negative face, and subsequent shift from negative to positive face generalization that occurred only after sleep. Overnight sleep spindle activity, extracted from high-density sleep EEG, positively correlated both with positive face generalization and with the shift from negative to positive generalization following sleep. These findings reveal a potential neural mechanism by which sleep attenuates negative bias and enhances positive representations, suggesting a potential method to buffer against maladaptive generalization in anxiety and PTSD.

Introduction

Imagine walking down the street to your favorite café when a dog suddenly jumps out from around a corner, knocking you to the ground. Upon cautiously returning to the same street, you see a large dog; even if only vaguely resembling the original, you may believe it is the same dog and instinctively retreat. This response reflects emotional generalization*:* an adaptive process that allows individuals to extend past experiences to similar, novel situations[1,2]. Generalization can occur consciously or unconsciously, often influencing how we interpret current stimuli based on past outcomes[2,3]. It is particularly sensitive to emotionally salient events, with a stronger bias toward negative associations. For example, tones paired with negative reinforcement produce broader generalization gradients than those linked to positive reinforcement, which in turn produce broader generalization than neutral stimuli [3,4].

While generalization is considered adaptive in many contexts, overgeneralization, especially of negative experiences, can become maladaptive. For example, a single dog attack may lead to fear of all dogs or even avoidance of outdoor settings altogether. Such behavioral patterns are implicated in psychiatric disorders, including major depression, phobias, generalized anxiety disorder, and posttraumatic stress disorder (PTSD)[1,4,5].

Sleep disturbances are also a hallmark of psychiatric disorders**,** reflecting the bidirectional relationship between sleep and mental health[6–9]. Sleep disruption can impair emotional regulation and increase anxiety, while anxiety itself can interfere with sleep quality [10]. This raises a critical question: *What role does sleep play in shaping emotional generalization?*

Research increasingly shows that sleep facilitates the consolidation of both rewarding[11–15] and aversive[16,17] emotionally salient memories, and supports the extraction of patterns from experience[18–20]. Thus, sleep may help to regulate emotional generalization by limiting the spread of negative associations. Accordingly, we hypothesized that sleep serves a corrective function, reducing negative overgeneralization while enhancing the generalization of positive events.

Participants learned associations between neutral faces and positive, negative, or neutral outcomes. Generalization was assessed using a face discrimination task involving morphs between the original stimuli, to capture how participants extended their learned associations to novel yet similar inputs. Participants completed this task before learning, immediately after, and following a consolidation period of either overnight sleep or daytime wakefulness. We found that

63 sleep reduced the generalization of negative stimuli and enhanced the generalization of positive
64 stimuli. Subsequent fMRI and sleep electrophysiology experiments revealed potential neural
65 mechanisms subserving these core behavioral findings. Overall, our work aims to clarify how
66 sleep shapes emotional generalization, with implications for understanding and treating anxiety
67 and PTSD [21].

Material and Methods

Experimental tasks

Face stimuli used in the tasks: Twenty-six participants (14 women) rated faces with neutral
expression sourced from an online database (the siblings DB)[22], arranging them along a
continuum from most malicious to most kind. Three males with similar average ratings as well
as similar skin tones, hair, and eye colors were selected as the original faces (Figure 1A). These
faces were positioned as vertices of a triangle space comprising 66 faces that were morphed
composites of the three original (vertex) faces (FantaMorph software). The proximity of each
face to each vertex corresponded to the proportion of the original face in the morph. Thus, the
face positioned in the center of the triangle was an average representation derived from all three
original faces (Figure 1A). The assignment of original faces to negative, positive, or neutral
valence vertices was counterbalanced across participants. Twenty-nine (18 women) participants
completed an online discrimination experiment to ensure no bias in perceiving morphed faces
as similar to the original faces (see also 'Discrimination Task' below).

Learning task. We used a probabilistic learning task (two-armed bandit[23]) to associate valence
with the original face stimuli. After a short training to ensure understanding, participants were
instructed to choose one face in each trial and maximize their points according to feedback.
During each trial, participants were presented with two out of three original faces simultaneously
on the screen, with the left and right positions randomized for each trial. All possible pairs of
faces were presented throughout the task. Each original face was probabilistically assigned
(p=0.9) to either positive, negative, or neutral feedback (gain +10, loss -10, or zero points), and
participants learned these associations as the task progressed. After their selection, the chosen
face was highlighted by a yellow frame on the screen. Feedback regarding the participant's
response (gain +10, loss -10, or zero points) was then provided based on the selected face (see
Figure 1B and Supplementary Figure 1A-C). Because face–feedback associations were highly
predictable (p=0.9), participants quickly learned the value of each face and consistently chose
the more rewarding option. This might lead to disproportionate exposure to positive feedback

and limited exposure to negative feedback. To overcome this potential bias, we included
'helpless' trials in which both presented stimuli were identical faces associated with negative or
neutral feedback, ensuring balanced exposure across feedback types. Trial composition
included a total of 132 trials comprising 30 positive-negative, 30 positive-neutral, 30 negative-
neutral, 18 neutral-neutral, and 24 negative-negative face pairs trial.

In the fMRI experiment, fixation crosses appeared both between face selection and feedback,
and during intervals between trials. The duration of the fixation cross was jittered uniformly
between 1 and 7 seconds, with a mean of 4 seconds (Supplementary Figure 1B). The task
lasted approximately one hour for the fMRI experiment, and 15 minutes for both the behavioral
and sleep EEG experiments. In the sleep EEG experiment, participants first viewed two faces;
after 1500 ms, arrows appeared beneath the faces, cueing participants to make a choice
(Supplementary Figure 1C).

Discrimination task. The discrimination task operationalized generalization as the perceived
similarity between images within the face space and each one of the original vertex faces. This
task was performed in three sessions at different time points throughout the experiment: (i)
before the first session, participants received instructions and practiced the task in a short
training session. In the behavioral and fMRI experiments, each trial in the discrimination task
begins with a presentation of one morphed face image from within the face space. Then,
participants are asked to choose one of the three original faces that resemble the presented
morphed face image the most (Figure 1C, Supplemental Figure 1D). Participants indicated their
choice by pressing one of three arrows. As a result, the same screen was shown with a yellow
rectangle surrounding the chosen face. The same trial was repeated if participants didn't
indicate a choice within three seconds. The correct choice granted the participants points, while
the wrong choice caused a loss of points. Notably, feedback on a given trial was not presented
regarding the correctness of the choice. Instead, the final score of points collected throughout
the task was presented at the end of each session. The accumulated points from all sessions of
the discrimination task and the gained points from the learning task were translated to a bonus
monetary reward at the end of the experiment.

In the EEG version of the experiment, participants initially had the opportunity to observe three
original faces, and memorize their appearance before the trials began. Subsequently, in each
trial, the three original faces are presented briefly, followed by the appearance of the morphed
face (random order). After 1000 ms (while the morph is still visible), arrows appear on the
screen, allowing participants to make their selections (refer to Supplemental Figure 1E). In all

experiments, each morph was presented six times (a total of 396 trials). The task lasted
approximately 25 minutes in all experiments. (See Figure 1C and Supplementary Figure 1D, E).

Psychomotor Vigilance Task (PVT). Participants were instructed to press a mouse
buttonwhen numbers, representing a running RT timer, appeared on a screen. The inter-
stimulus interval (ISI) varied randomly between 2–10 seconds, and the task lasted five
minutes[24]. Sustained attention performance was assessed using mean and SD of RT, 1/RT, as
well as lapse counts (RT > 500 ms) and errors (button presses without a stimulus). These
metrics are highly sensitive to sleepiness, making the PVT a standard tool for vigilance-related
assessments [25–27].

Experiments:

Behavioral Experiment

Participants: Sixty healthy participants took part in the behavioral experiment, randomly
assigned to a **day group** (n=30) or a **night group** (n=30). Written informed consent was
obtained from each participant, and the study was approved by the IRB of the Weizmann
Institute of Science. No participants were excluded. The final sample included 30 participants in
the night group (19 females, mean age=25.57, SD=3.11) and 30 in the day group (14 females,
mean age=28.13, SD=6.4).

Procedure: Participants completed two experimental sessions ~8–12 hours apart. Day group
participants were tested in the morning and again in the evening; night group participants were
tested in the evening and returned the next morning after sleeping at home. In the first session
(~1 hr 15 min), participants completed sleepiness (SSS) and vigilance (PVT) assessments (~10
min), followed by two discrimination task sessions (~25 min each) separated by a learning task
(~15 min). In the second session (~1 hr 15 min), they completed another SSS and PVT (~10
min) and a final discrimination task (~25 min), before receiving monetary compensation.
Session started at 8:00–9:30 a.m. or 8:00–9:30 p.m., depending on group assignment. Sleep in
the night group and wake activity in the day group were monitored via actigraphy between the
sessions. (See Figure 1D).

fMRI Experiment

Participants: Ninety healthy, right-handed participants were recruited for the fMRI experiment.
They were assigned to a **day group** (n=45) or a **night group** (n=45). Written informed consent
was obtained, and all procedures were approved by the IRB of the Weizmann Institute of

Science and the Medical IRB of the Tel-Aviv Sourasky Medical Center. Ten participants were
excluded due to technical issues with the scanner or task scripts (n=5), falling asleep during
scanning (n=1), voluntary withdrawal (n=2), or excessive motion inside the scanner (n=1), and
their data were neither preprocessed nor analyzed. The final analysis included 40 participants in
the night group (21 females, mean age=26.97, SD=5.66) and 40 in the day group (25 females,
mean age=26.1, SD=3.71).

Procedure: Each participant completed two sessions spaced ~8–10 hours apart. The day
group was tested in the morning (7:00–9:00 a.m.) and returned that evening; the night group
was tested in the evening (7:00–9:00 p.m.) and returned the following morning. The first session
(~2.5 hrs) included sleepiness and vigilance assessments (~10 min), the first discrimination task
(~25 min), and the learning task conducted inside the MRI scanner (~1 hour + preparation).
After scanning, participants completed a second discrimination task (~25 min) and left the lab.
The second session matched the behavioral format, including SSS, PVT, and the final
discrimination task. Sleep behavior in the night group and wake behavior in the day group were
tracked using an actigraph watch between the sessions. (See Figure 1D).

Sleep EEG Experiment

Participants: Forty-five healthy participants were recruited for the sleep EEG experiment.
Written informed consent was obtained, and procedures were approved by the Medical IRB of
the Tel Aviv Sourasky Medical Center. Three participants were excluded due to noisy or
corrupted EEG data, resulting in a final sample of 42 participants (26 females, mean age=25.71,
SD=3.17).

Procedure: All participants were assigned to a **night group** and slept in the lab under full
polysomnographic and high-density EEG monitoring. They arrived ~3 hours before their regular
bedtime. Setup included EEG net application (~1 hr), followed by SSS and PVT assessments
(~10 min), and two discrimination task sessions (~25 min each) separated by a learning task
(~15 min), totaling ~2.5 hours. After overnight sleep and a short recovery period, participants
completed a second sleepiness and vigilance assessment (~10 min), and a final discrimination
task (~25 min). The EEG net was removed, and participants were compensated. (See Figure
1D).

Actigraphy

In the fMRI and behavioral experiments, we provided the participants with an actigraph watch (Ambulatory Monitoring, Inc. motionlogger Micro Watch). This watch measured acceleration, light, and temperature. These measures were used to validate sufficient sleep in the night groups and the absence of daytime naps in the day groups.

fMRI

fMRI acquisition: Functional imaging data were collected using a 3T Siemens MAGNETOM Tim-Trio scanner with a 32-channel head coil. T2*-weighted images were acquired with a gradient-echo EPI sequence (TR=1450 ms, TE=30 ms, flip angle=90°), with 60 descending slices (3 mm thick, 10% gap) at a 30° coronal angle from the ACPC plane. Voxel size was 2.3 mm³ (FOV=220), and phase encoding was anterior-to-posterior. Functional scanning occurred in two sessions with up to a 2-minute interval. High-resolution anatomical T1-weighted images were also acquired (TR=2300 ms, TE=2.29 ms, flip angle=8°, voxel size=0.9 mm³, FOV=240). Functional scans covered nearly the entire brain, excluding a small dorsal parietal region. The anatomical scan covered the whole brain. **fMRI preprocessing:** Preprocessing was performed using *fMRIPrep* 21.0.0 [35,36], based on Nipype 1.6.1 framework [37,38]. Fieldmaps for B0 distortion correction were estimated using topup on EPI reference images [39]. T1-weighted anatomical images were corrected for intensity nonuniformity (N4BiasFieldCorrection [40] implemented in ANTs version 2.3.3.[41]), skull-stripped (Nipype implementation of ANTs-based brain extraction using the OASIS30ANTs template), and segmented into gray matter (GM), white matter (WM), and cerebrospinal fluid (CSF) using FSL's FAST [42]. Nonlinear spatial normalization to a standard space (MNI152NLin2009cAsym [43]) was completed using ANTs (antsRegistration). For each BOLD run, a reference volume was generated for motion correction (FSL mcflirt [44]). Slice timing correction was applied (AFNI's 3dTshift [45]). Co-registration to the T1-weighted image was done using FreeSurfer's *bbregister* with boundary-based registration [46]. Functional images were resampled to MNI space using antsApplyTransforms (with Lanczos interpolation to minimize smoothing). Functional images were resampled to standard space using *antsApplyTransforms* with Lanczos interpolation [50], and surface-based resampling was done with FreeSurfer's *mri_vol2surf* [47].Confound regressors included framewise displacement (Power [48] and Jenkinson [44]), DVARS, and global signals (CSF, WM, whole brain). Temporal and anatomical CompCor were computed [49] temporal (tCompCor) components were derived from the top 2% most variable voxels, and anatomical (aCompCor) components were extracted from masks of CSF, WM, and combined compartments, with GM faction minimized by excluding voxels with >5%

GM partial volume [49]. For each CompCor method, components explaining 50% of nuisance
variance were retained. All regressors, including motion estimates and their
derivatives/quadratics, were included in the confound file. Volumes exceeding FD > 0.9 mm or
DVARS > 1.5 SD were flagged as motion outliers.

Whole-brain GLM analysis: Voxel-wise statistical analyses were conducted across the whole
brain. At the first level, individual events were modeled using a canonical hemodynamic
response function (HRF), and framewise displacement estimates from preprocessing were
included as nuisance regressors to control for motion. Each participant's two imaging sessions
were modeled separately using a first-level general linear model (GLM), then combined in a
second-level group analysis. The event-related design included six regressors, corresponding to
the appearance of the face's stimulus and the feedback, each with positive, negative, or neutral
valence. For the face stimuli, valence was determined by the expected feedback associated
with the participant's choice. Regressors were time-locked to event onset and modeled over the
full event duration: three seconds for the stimulus and two seconds for the feedback. All
participants were included in the group-level model regardless of sleep condition (day/night),
enabling unbiased voxel selection for downstream analysis. Group-level statistical maps were
corrected for multiple comparisons using a family-wise error rate (FWER) threshold of $p < 0.001$,
with a minimum cluster size of 10 voxels.

ROI selection: A predefined list of 28 regions of interest (ROIs) was compiled based on the
Harvard-Oxford Atlas, guided by prior research implicating these regions in reward/punishment
processing, decision-making, emotion regulation, and face perception. ROIs were included in
the final analysis only if they (1) passed the corrected GLM threshold for both contrasts
mentioned below and (2) were part of the predefined list. Fourteen regions met these criteria.
See Table 1.

ROI name	Included?	ROI name	Included?
Caudate	V	Cuneus	X
Putamen	V	Thalamus	X
Pallidum	V	Posterior Superior temporal gyrus	X
Hippocampus	V	Occipital Fusiform Gyrus	X
Amygdala	V	Fusiform Face Area	X
Accumbens	V	Occipital Face Area	X
Precuneus	V	Posterior Temporal Fusiform Cortex	X

Orbitofrontal Cortex	V	Anterior Temporal Fusiform Cortex	X
Medial Frontal Cortex	V	Temporo-occipital fusiform cortex	X
Subcallosal Cortex	V	Superior Temporal Sulcus	X
Anterior Cingulate Cortex	V	Anterior Middle Temporal Gyrus	X
Posterior Cingulate Gyrus	V	Posterior Middle temporal Gyrus	X
Anterior Superior Temporal Gyrus	V	Postcentral Gyrus and the Brainstem	X
Insula	V	Brainstem	X

Table 1: List of 28 a priori regions of interest (ROIs) based on the Harvard-Oxford Atlas. Fourteen ROIs that passed statistical thresholding (FWE-corrected $p < 0.001$, cluster size ≥ 10 voxels) were included in the final analysis (left column).

For each of the 14 ROIs, a 3 mm-radius sphere was defined around the peak activation voxel. Activity was extracted from both the peak voxel and its surrounding sphere using each participant's unthresholded statistical maps. If a sphere extended beyond anatomical ROI boundaries, only the portion within the ROI was retained. To assess differences in neural responses, we computed two contrasts: Gain (positive > neutral feedback) and Loss (negative > neutral feedback). Contrast values were averaged across both scanning sessions for each participant and used in subsequent correlational analyses with behavioral generalization measures.

Sleep EEG

Data Acquisition: Continuous high-density EEG and full polysomnography were recorded throughout the entire experimental session, which included both cognitive tasks and sleep periods. A 256-channel hydrogel Geodesic Sensor Net (Electrical Geodesics, Inc., EGI) was used for data acquisition. Each electrode consisted of a silver chloride carbon fiber pellet, a lead wire, and a gold-plated pin, and was filled with conductive gel (Electro-Cap International). Signals were amplified using an AC-coupled, high-input-impedance amplifier (NetAmps 300, EGI) with an integrated antialiasing analog filter. Recordings were referenced to Cz and digitized at a sampling rate of 1000 Hz. Prior to recording, electrode impedance was verified to be below 50 kΩ for all sensors.

Sleep Scoring: Sleep stages were manually scored offline according to the American Academy of Sleep Medicine (AASM) guidelines. Each recording was scored by two trained raters to

ensure reliability. Standard sleep parameters, including sleep latency, total sleep time, and time
spent in each sleep stage, were calculated for subsequent analysis.

Preprocessing: Analyses were conducted in Python using the sleepEEGpy package[50], an
open-source "wrapper" pipeline that integrates MNE-Python[51], PyPREP[52], YASA[53],
SpecParam[54], and other tools. Each EEG recording was down-sampled from 1000 Hz to 250 Hz
and band-pass filtered between 0.75–40 Hz using a zero-phase FIR filter with a Hamming
window. Manual inspection of raw signals was performed using the "butterfly" view,
complemented by topographical maps to identify and exclude abnormal channels.
Subsequently, Independent Component Analysis (ICA) was applied using the extended infomax
algorithm[55] with 40 Principal Component Analysis (PCA) components. Components associated
with ECG artifacts were identified through visual inspection and removed.

Power Spectral Analysis: Power Spectral Density (PSD) was computed separately for each
sleep stage using the Welch method as implemented in SleepEEGpy [50]. Parameters included a
Fast Fourier Transform (FFT) length of 2048 samples, a Hamming window, and a 50% segment
overlap (1024 samples). For each participant, the maximum sigma band power (12.5–15 Hz)
across all channels was identified. The sigma power from the peak channel was used in
subsequent correlation analyses.

Sleep Spindle Detection: Analyses were conducted in Python using sleepEEGpy[50] employing
the YASA algorithm [53]. Spindles were detected during Non-rapid eye movement (NREM) sleep
from each participant's peak sigma channel. Fast spindles were defined within the 13–15 Hz
frequency range, with durations constrained between 0.5–2.0 seconds and a minimum inter-
spindle interval of 500ms. Detection criteria included: Correlation ≥ 0.65 (moving correlation
between the raw and sigma-filtered signal), Relative power ≥ 0.2 (ratio of power in the spindle
band to broadband power), root-mean-square (RMS) amplitude ≥ 1.5 standard deviations above
the mean of the moving RMS of the sigma-filtered signal.

Generalization analysis

Generalization assessment: We analyzed each participant's response patterns separately for
each session. We calculated the proportion of responses classifying each morphed face as
most similar to one of the original faces, organizing them into three triangular spaces based on
valence: positive, negative, or neutral. As face–valence pairings were counterbalanced,
triangles were labeled by valence rather than face identity (Figure 1E, top left). We interpolated

each triangle to generate continuous response gradients (Figure 1E, top right). A 50% threshold
defined perceptual boundaries, areas where faces were most likely judged similar to the
original. The size of these areas served as a measure of generalization (Figure 1E, bottom left;
for visualization, all three are shown on one triangle). To assess changes over time, we
computed two metrics: (1) immediate generalization: the area change from Session 1 to
Session 2 (after learning), and (2) delayed generalization: the change from Session 2 to
Session 3 (after sleep or wake; Figure 1E, bottom right).
Note: While a 66% threshold was used in the figures for clearer visualization, all analyses were
conducted using the 50% threshold.

Generalization score: In the fMRI experiment, we computed a single score to capture the
balance between generalization of positive and negative faces. This generalization score was
calculated by subtracting negative face generalization from positive face generalization. Scores
above zero indicate a bias toward generalizing the positive face, while scores below zero
indicate greater generalization of the negative face. This measure reflects the idea that
increased generalization of one emotional valence occurs at the expense of the other.

Renormalization index: To quantify changes in generalization over time, we computed a
renormalization index for each participant by subtracting the immediate generalization score
from the delayed generalization score. In the fMRI experiment, we used the generalization bias
score (positive minus negative) for this calculation. In the EEG experiment, we focused on
generalization of the positive face only.

Overgeneralization Calculation: Overgeneralization was defined as instances where
participants incorrectly classified morphed faces as resembling one of the original faces, beyond
the boundaries of the correct generalization zone. In the generalization task, such incorrect
responses resulted in point loss and were considered maladaptive. Regions of the triangular
face space that extended beyond the adaptive generalization boundaries were designated as
overgeneralization areas. See Supplementary Figure 7A, B for illustration.

Statistics: Group-level effects in generalization were assessed using mixed-model ANOVAs.
Significant interactions were followed by paired or independent-samples two-tailed t-tests,
depending on the comparison. To examine whether brain ROI activity predicted generalization,
we applied both linear regression and LASSO (least absolute shrinkage and selection operator)
regression. All predictors were z-scored prior to LASSO to ensure comparability. The

regularization parameter was optimized using 10-fold cross-validation and averaged across
folds. Linear regression models were then constructed using the subset of predictors selected
by LASSO.

Pearson correlations were used to assess associations between behavioral generalization
scores and neural measures. Specifically, we examined correlations between amygdala
activation (from ROI-based fMRI contrasts) and generalization indices, as well as between
EEG-derived metrics (e.g., spindle count and sigma power) and changes in generalization
across sessions. Statistical significance was set at $p < .05$.

Learning task analysis

To confirm that participants learned the associations between faces and outcomes, we
calculated the percentage of correct choices for each participant across all trials. In all
experiments, all participants performed above chance level, with a minimum accuracy of 60%.
The average accuracy across all three experiments was 93.1% (SD=5.9%). The learning curve
shown in Figure 1B reflects data from all participants across the three experiments (N=182),
smoothed using a moving average over a window of five trials.

Results

To quantitatively study the role of sleep in the generalization of sensory stimuli with different
emotional valence in healthy participants, we first employed a series of repeated behavioral
discrimination tasks (N=60). Participants were presented with face images generated through
morphing, depicting intermediate levels along a continuum between three original faces (Figure
1A). For each image, participants judged which original face it most closely resembled (Figure
1C). In a second task, using a probabilistic reinforcement-learning procedure (two-arm bandit),
participants learned to associate the original faces with positive, negative, or neutral feedback
(Figure 1B). Participants completed the discrimination task three times (Figure 1D): before
(session 1) and immediately after (session 2) learning the associations between faces and
feedback, as well as following an 8-10 hour interval of nighttime sleep or daytime wakefulness
(session 3).

Behavioral responses from each session were organized into a triangular face space
representing perceived similarity between each morph and the original faces. Areas around
each vertex where similarity exceeded a 50% threshold were compared between two sessions

to quantify generalization (Methods, Figure 1E). Put simply, generalization was marked by a
wider area of morphed faces in the triangle that was perceived as identical to the original face in
the vertex. Throughout the paper, we refer to differences in generalization while learning
associations as 'immediate generalization' (session 2 - session 1) and refer to differences
across the long daytime wake or nighttime sleep intervals as 'delayed generalization' (session 3
- session 2).

Figure 1: Experimental design: A. Face Space: Top left: the three original faces used in the experiment. Right: The
triangular "face space" generated from the three original faces. Faces at the vertices are the original faces, while
intermediate positions represent morphed faces created by blending the three originals. Each morph's proximity to a
vertex indicates the proportion of that original face in the morph. The face at the center is the average of all three
original faces. **B.** Learning task trial timeline. Left: During each trial of the learning task, two faces (selected from the
three original faces) were displayed. Participants chose one face and received feedback (+10, -10, or 0 points) based
on their selection. Each original face was probabilistically associated with positive, negative, or neutral feedback
(90% probability per face for each category). Feedback-face associations were randomized across participants.
Detailed timing for different experiments is provided in Supplemental Figure 1. Top-right: Feedback association
illustration for the original faces at the vertices of the face space (randomized for each participant). Bottom-right:
learning curve from participants across all three experiments (behavioral, fMRI, EEG). **C.** Discrimination task trial.
Left: Each trial began with a figure of a morphed face from within the face space. Then, participants were asked to
choose from the three original faces the face that most resembled the presented morph. Selections were indicated by

Sleep selectively promotes generalization of positive associations

We observed distinct generalization patterns between day and night groups, finding that
wakefulness and sleep had different effects on generalization. Specifically, participants in the
night group were more likely to generalize the positive associated face, whereas participants in
the day group more often generalized the negative associated face (see Figure 2A for
representative examples from each group). To quantify and assess the effect of the sleep/wake
interval on generalization, we conducted a two-way, mixed-model ANOVA on delayed
generalization, with Group (Day/Night, between subjects) and Face (Positive/Negative/Neutral,
within-subjects) as factors. The analysis revealed a significant Face × Group interaction ($F_{(2,116)}=3.61$, p=0.03, Figure 2B, 2C). Post-hoc comparisons indicated that the negative face
was generalized more during daytime wakefulness than during overnight sleep (t(58)=-2.3,
p=0.025). By contrast, the positive face was more generalized during sleep than wake
(t(58)=2.09, p=0.049).

To further understand this effect, we conducted separate two-way, mixed-model ANOVAs for
the negative and positive faces, considering the total classified area in each session (rather than
only the session-to-session difference; Methods, Figure 2D). For negative face generalization, a
mixed-model ANOVA with Group (Day/Night) and Session (Session 2/Session 3) as factors
revealed a significant Group × Session interaction ($F(1, 59)=2.34$, p=0.02; Figure 2D, left). Post-
hoc comparisons indicated a significant increase in the area classified as negative in Session 3
compared to Session 2, but only in the daytime wake group (t(29)=–2.44, p=0.02). A similar
analysis for positive face generalization also revealed a significant Group × Session interaction
($F(1, 59)=2.04$, p=0.04; Figure 2D, right). Here, post-hoc comparisons showed a significant

increase in the area classified as positive in Session 3 relative to Session 2, but only in the
sleep group (t(29)=–2.14, p=0.04). No significant group differences were observed for neutral
face generalization (Figure 2E, 2F), nor were there differences in immediate generalization prior
to the sleep/wake interval (Figure 2G, 2H). Together, these results suggest a dissociation in
generalization patterns: overnight sleep preferentially enhances generalization of positive
stimuli, whereas daytime wakefulness enhances generalization of negative stimuli.

To control for potential confounds, we compared subjective sleepiness (SSS) and vigilance
(PVT) between groups. No significant group differences were found in either measure during the
first meeting. In the second meeting, participants in the day group reported higher sleepiness
than the night group (Day: 3.87 ± 1.63 vs. Night: 3.03 ± 1.34; t(58)=2.15, p=0.036); however, no
group differences were observed in vigilance as measured in the PVT (a highly sensitive
measure of sleepiness[26,56]). Additionally, neither SSS nor PVT scores were significantly
correlated with generalization measures in either group (see Discussion).

A. Examples for delayed generalization ⏱️

(i) Example from night group 🌙

Negative face

Session 2   Session 3

Delayed generalization

Positive face

Session 2   Session 3

Delayed generalization

(ii) Example from day group ☀️

Negative face

Session 2   Session 3

Delayed generalization

Positive face

Session 2   Session 3

Delayed generalization

Less than threshold
Negative face
Positive face
Neutral face
No difference
After>before
After<before

B. Averages for day and night groups (all faces) ⏱️
delayed generalization

Night group 🌙

Session 2   Session 3

Delayed generalization

Day group ☀️

Session 2   Session 3

Delayed generalization

C. Delayed generalization: negative and positive faces ⏱️

Generalization

Negative face   Positive face

D. Sessions 2 and 3: negative and positive faces ⏱️

Negative face 🌙 ☀️   Positive face 🌙 ☀️

Classified Face Area

Session 2   Session 3   Session 2   Session 3

E. Delayed generalization: Neutral face ⏱️

Generalization

F. Sessions 2 and 3: Neutral face ⏱️

🌙 ☀️

Classified Face Area

Session 2   Session 3   Session 2   Session 3

G. Averages for day and night groups (all faces) ⏱️
Immediate generalization

Night group 🌙

Session 1   Session 2

Immediate generalization

Day group ☀️

Session 1   Session 2

Immediate generalization

H. Immediate generalization: all faces ⏱️

Generalization

Negative   Positive   Neutral

Sleep reverses amygdala-associated generalization of negative stimuli

Next, we set out to investigate the neural correlates associated with generalization of positive and negative faces and their dynamics using fMRI, employing a modified version of the learning task performed in the scanner (Methods). We could not replicate the group-level behavioral differences between generalization after extended wakefulness versus sleep (Supplementary Figure 2, see also Discussion), observing considerable inter-individual variability in generalization patterns. During immediate testing, some participants exhibited stronger generalization of the positive face (Figure 3A), whereas others showed a bias toward generalization of the negative face (Figure 3B). Similar individual variability was observed in delayed generalization after daytime wakefulness or after overnight sleep. We focused subsequent analysis on testing whether these individual variabilities are associated with specific brain activities.

First, we tested whether brain activity during learning, specifically when receiving feedback linked to point gains or losses, could predict immediate and delayed generalization behavior. To quantify generalization, we computed a composite score calculated as the difference between positive- and negative-face generalization scores (Methods), with values above zero indicating greater generalization of the positive face and below zero indicating greater generalization of the negative face. This score was used in multiple linear regression models to predict either (i) immediate generalization (for both groups combined, as it is prior to the sleep/wake

consolidation period) or (ii) delayed generalization following daytime wake or overnight sleep
(separated between the groups). fMRI blood oxygenation level dependent (BOLD) activity
underwent whole-brain general linear model (GLM) analysis to identify task-related activity, and
14 Regions of Interest (ROIs) were selected as predictors (Methods). To identify the most
predictive ROIs, we applied Lasso regression [57] (elastic net regularization) with 10-fold cross-
validation for model optimization.

We found that greater activation for losses (versus neutral feedback) in limbic regions,
specifically the Amygdala, Caudate, Putamen, and Subcallosal Cortex, was correlated with
increased immediate generalization of the negative face, but decreased immediate
generalization of the positive face, This corresponds to a negative correlation with the
generalization composite score (regression model: N=80, R^2=0.18, $F_{(4,75)}$=4.17, p=0.004;
Figure 3C, D). A similar pattern emerged during gains (versus neutral feedback): activity in the
Amygdala, Orbitofrontal cortex, and Insula was positively correlated with generalization of the
negative face and negatively with the positive face (positive correlation to the generalization
score), whereas activity in the Precuneus (default mode network, DMN), showed the opposite
trend: (regression model: N=80, R^2=0.18, $F_{(4,75)}$=4.14, p=0.004, Figure 3E, F). Thus,
regardless of feedback valence (gaining or losing points associated with monetary reward),
limbic activity biases immediate generalization toward negatively associated stimuli, whereas
engagement of the DMN is associated with positive stimuli immediate generalization. These
effects were also evident when modeling positive and negative generalization separately, rather
than using the generalization composite score. Specifically, limbic activity during learning mostly
predicted increased immediate generalization of the negative face (see Supplementary Figure
3).

Next, we analyzed brain activity associated with delayed generalization after overnight sleep.
Greater limbic activation during both loss- and gain-related feedback in the learning session was
associated with a renormalization from negative to positive face generalization following
overnight sleep. Specifically, activity in the Amygdala, Nucleus Accumbens, and Anterior
Superior Temporal Gyrus during loss-related feedback predicted increased generalization of the
positive face after sleep (regression model: N=40, R^2=0.27, $F_{(3,36)}$=4.45, p=0.009; Figure 3G,
H). A similar trend was observed for gain-related feedback, where higher activation in the
Amygdala and Anterior Cingulate Cortex during gains was associated with a greater shift in
generalization from the negative to the positive face following sleep (regression model: N= 40,
R^2=0.18, $F_{(2,37)}$=4.14, p=0.02; Figure 3, I-J). Similarly, separate models for delayed positive

and negative generalization, instead of the generalization composite score, showed that limbic
activation during learning predicted a sleep-dependent shift toward greater generalization of the
positive face (see Supplementary Figure 4).

Together, these findings suggest that heightened limbic engagement during emotionally salient
learning—regardless of outcome valence—may promote immediate negative generalization,
followed by a sleep-dependent renormalization that shifts toward positive generalization. In
contrast, applying the same Lasso regression approach to predict delayed generalization after
daytime wakefulness did not select any ROI predictors for either gain or loss, suggesting that
this renormalization process may be specific to sleep-related consolidation (see Discussion).

Finally, no significant group differences in subjective sleepiness (SSS) or vigilance (PVT) were
observed at either time point. Additionally, neither SSS nor PVT scores were significantly
correlated with generalization measures in either group.

Figure 3. Immediate and late generalization are correlated with Amygdala and limbic activity.
A–B. Individual variability in immediate generalization. **A.** A participant who generalized the positive face immediately after learning. Left: areas classified as the negative face during Sessions 1 and 2. Right: areas classified as the positive face. Larger triangles represent changes in classification between sessions, reflecting immediate generalization. **B.** regions classified as the negative face during Sessions 1 and 2. Right: regions classified as the positive face.
C-D. Regression results for predicting immediate generalization from neural activity during loss-related feedback. **C.** Coefficient estimates for ROIs selected via Lasso regression: Subcallosal Cortex, Amygdala, Putamen, and Caudate. All coefficients are negative, indicating that greater activity in these limbic regions was associated with reduced generalization of the positive face and increased generalization of the negative face. Inset (top right): model fit (x-axis=predicted generalization from ROI activity; y-axis=observed generalization). Red line=regression fit; gray shading=95% confidence intervals. **D**. Mean ROI activity during loss feedback for regions in (C). Black circle marks a 3-voxel-radius sphere centered on the peak voxel within each anatomical ROI. **E-F**. Immediate generalization: gain-

Amygdala activity consistently emerged as an important predictor for immediate and delayed generalization when activated for both positive and negative emotional valence stimuli, being the only ROI repeatedly selected by LASSO in all prediction models. Specifically, amygdala activation during losses was significantly correlated with immediate generalization of the negative face (negative correlation to the composite score) across both experimental groups (N=80, r_p=-0.26, p=0.02, Figure 4A). However, for delayed generalization following overnight sleep, this activity predicted the positive face generalization (N=40, r=0.36, *p*=0.02; Figure 4B), a phenomenon that did not occur after daytime wakefulness (Figure 4B, bottom-right, r_p=0.04, p=0.8).

To quantify this effect, we computed a renormalization index for each participant by subtracting the immediate generalization score from the delayed generalization score for each participant (Methods). Positive renormalization indices indicate that, after sleep, the generalization is shifted towards more positive values than those at baseline. In the night group, greater amygdala activation during loss-related feedback predicted stronger renormalization following sleep. Participants with higher amygdala activity not only exhibited greater immediate generalization of the negative face but also demonstrated a clear shift toward positive face generalization after sleep, as reflected by a significant correlation between the renormalization index and amygdala activity during loss (N=40, r=0.43, *p*=0.005; Figure 4C). No such correlation was observed in the daytime group (Figure 4C, bottom-right, r=0.12, p=0.45).

A similar pattern emerged when examining amygdala activity during reward-related feedback. Higher amygdala activation while gaining points was associated with greater immediate generalization of the negative face across both groups (N=80, r=-0.25, *p*=0.02; Figure 4F). However, the amygdala activity showed trend toward a significant correlation with positive face

generalization following sleep (N=40, r=0.28, p=0.08; Figure 4G). Although the correlation was

not significant, it was significantly different than the day group that showed no correlation

between the amygdala activity and the generalization (Figures 4G bottom-right panel, r_p=-0.16,

p=0.31, fisher test to compare night and day correlations: Z=1.9, p=0.03).

Furthermore, the renormalization index showed a strong correlation with Amygdala activity

during gains (N=40, r=0.4, p=0.009; Figure 4H), indicating that individuals with higher initial

amygdala engagement for gain-related feedback also underwent a greater shift from negative to

positive generalization after sleep. As was found for loss-related activity, renormalization effects

were specific to overnight sleep, no correlation to the renormalization index in the daytime wake

group (4H, bottom, right panel, r_p=-0.01, p=0.95).

Sleep-dependent renormalization was consistently observed for Amygdala activity, but not

unique to this brain region. Direct correlations with other limbic ROIs revealed a consistent

pattern whereby higher activation during learning (either reward or loss) was associated with

increased immediate generalization of the negative face, followed by a renormalization toward

the positive face after sleep (Supplementary Figures 5 and 6).

In addition, we examined overgeneralization, defined as generalization that is no longer

adaptive (i.e., leads to point loss). We found significant correlations between amygdala activity

during both loss- and gain-related feedback and two behavioral outcomes: immediate

overgeneralization of the negative face and delayed overgeneralization of the positive face

following sleep. Notably, overgeneralization scores were highly correlated with overall

generalization scores, and the observed associations with amygdala activity did not remain

significant after controlling for this shared variance (Supplementary Figure 7).

Figure 4. Direct correlations between amygdala activity during learning and generalization behavior. A. Immediate generalization**:** Amygdala activity during loss feedback in the learning phase is negatively correlated with immediate generalization across both groups. Higher amygdala activity is associated with increased generalization of the negative face. The grey line represents the fitted regression line; the gray shaded area indicates 95% confidence intervals. **B.** Delayed generalization**.** Left: In the night group, higher amygdala activity during loss is associated with positive face generalization after sleep. *Bottom right*: No such correlation is observed in the day group, suggesting the effect is sleep-specific. **C.** Correlation with the renormalization index**.** Left: In the night group, higher amygdala activity during loss is significantly correlated with a greater renormalization index, indicating a stronger shift from negative to positive generalization after sleep. Bottom right: No such relationship is observed in the day group. **D.** Activation map showing the amygdala response to loss feedback. The black circle marks a sphere (radius=3 voxels) centered on the peak activation voxel. Activity used for analysis is the average across voxels within this sphere that fall within the anatomical ROI **E.** Same as D for gain feedback. **F–H**. Same analyses as in panels A–C, but for amygdala activity during gain feedback. **F.** Higher amygdala activity during gains is associated with increased immediate generalization of the negative face. **G.** Left: Amygdala activity during gains predicts a renormalization toward positive-face generalization after sleep in the night group. Bottom right: No such effect is observed in the day

Sleep spindle activity correlates with renormalization of negative-to-positive emotional generalization

To further investigate the role of sleep in promoting positive generalization and reversing
negative emotional bias, we conducted an overnight sleep study with polysomnography and
high-density (256-channel) EEG in a separate cohort of healthy volunteers (n=42, Methods,
Figure 5A, B). Participants completed the same behavioral paradigm used in prior experiments,
both before and after sleep. Sleep scoring was performed to assess whether sleep architecture
was related to generalization outcomes. On average, participants exhibited a typical profile for
overnight sleep in healthy young adults, with a mean (SD) total sleep time of 400.2 (69.7) min,
sleep efficiency (sleep time/time in bed) of 92.7% (5.2) and a mean sleep latency of 18.3 (15.5)
min. Non-rapid eye movement (NREM) sleep, REM sleep, and wake after sleep onset (WASO)
constituted 302.2 (32.3), 96.8 (32.3), and 19.7 (17.7) minutes, respectively. We tested for
associations between time spent in specific sleep stages and generalization, but no significant
relationships emerged (Supplementary Table 1).

Next, we examined whether specific electrophysiological activities during sleep may be
correlated with generalization. We focused on sleep spindle activities over centro-occipital
regions occurring during NREM sleep in the EEG sigma frequency band (12-15Hz, Methods;
Figure 5C)[58] as such activities are implicated in memory consolidation and cortical plasticity[59–63]
We found that greater sigma power during NREM sleep was significantly correlated with an
increase in delayed generalization of the *positive* face following sleep (r_p=0.38, p=0.01; Figure
5E). We also observed a trend toward positive correlation between *immediate* generalization of
the negative face and subsequent sigma power (r_p=0.25, p=0.09), as well as a trend toward
negative correlation between immediate generalization of the positive face and sigma power
(r_p=–0.26, p=0.09; Figure 5D). These trends suggest that individuals who initially generalized
the negative face more strongly than the positive face exhibited increased spindle activity during
subsequent sleep. Could this pattern reflect a compensatory mechanism, wherein enhanced
spindle activity supports a renormalization of negative emotional generalization? Consistent with
this notion, the *renormalization* in positive face generalization (i.e., the increase from immediate
to delayed generalization) was also significantly correlated with sigma power during sleep
(r_p=0.40, p=0.009; Figure 5F).

To extend this analysis, we went beyond power spectral analysis to detect and quantify individual spindle events using SleepEEGpy[50], incorporating YASA[53] detection (Methods, Figure 5G). We found that the number of fast (13–15 Hz) spindles during NREM sleep was positively correlated with post-sleep generalization of the positive face (r=0.33, p=0.03; Figure 5I) and with the renormalization in positive generalization (r=0.031, p=0.04; Figure 5J). Notably, we also found that a higher number of fast spindles was associated with reduced generalization of the neutral face after sleep (r=–0.38, p=0.013; figure 5H). Since neutral faces were not associated with reward or punishment, this finding suggests that sleep spindles may help suppress the generalization of emotionally irrelevant or neutral stimuli, highlighting a possible role of spindles in valence-selective memory processing (see Discussion).

Figure 5: Sleep spindle activity facilitates positive generalization A. Time–frequency spectrogram (log power) from a representative participant recorded at electrode Pz, illustrating spectral dynamics across the night. The overlaid hypnogram depicts transitions between sleep stages. **B.** Power Spectral Density (PSD) plots of preprocessed data from electrode Pz from a representative participant, separated by sleep stage. Colored lines

denote individual stages. The gray-shaded region indicates frequencies removed by the high-pass filter. **C.** Topographical map of sigma power (12–15 Hz) during N2 sleep from a representative participant, demonstrating typical centroparietal distribution of sleep spindles. **D.** Correlation between sigma power during NREM sleep (from each participant's peak-sigma channel) and immediate generalization of the positive face. Participants with lower immediate positive generalization exhibited greater subsequent sigma power. **E.** Higher sigma power during NREM sleep is positively correlated with delayed generalization of the positive face, measured after sleep. **F.** Sigma power during NREM sleep is significantly correlated with the renormalization of the positive face generalization from the pre-sleep (immediate) to post-sleep (delayed) session. **G.** A representative example of a fast spindle (13–15 Hz) recorded during NREM sleep, shown in the time domain. Scale bars indicate 1 second (horizontal) and 20μV (vertical). **H.** Greater spindle count during NREM sleep (from the peak-sigma channel) is associated with reduced delayed generalization of the neutral faces, suggesting selective suppression of non-valenced stimuli generalization. **I.** The number of fast spindles during NREM sleep is positively correlated with delayed generalization of the positive face. **J.** Fast spindle count is also positively correlated with the renormalization in positive generalization from immediate to delayed sessions.

Discussion

In this study, we tested the hypothesis that sleep reduces emotional generalization of negative stimuli while boosting the generalization of positive stimuli. Using a face discrimination task with morphed faces, we found that overnight sleep selectively promoted generalization of positive associations, whereas daytime wakefulness was linked to generalization of negative associations (Figure 2). fMRI analysis revealed that greater limbic responses to emotional feedback during learning (gains or losses) were associated with broader immediate generalization of the negative face. This activity was associated with a reversal of this pattern after sleep: participants exhibited greater generalization of positive faces following overnight sleep, but not after a day awake (Figure 3). Amygdala activity consistently emerged as a main predictor of both immediate and delayed generalization, for both positive and negative emotional valence (Figure 4). Participants with high amygdala activation during learning exhibited the strongest immediate generalization of the negative face and showed the greatest sleep-associated renormalization. Finally, we found that sleep spindle activity was significantly correlated with a shift from negative to positive generalization (Figure 5).

Data from the behavioral experiment revealed a clear divergence in delayed generalization profiles following overnight sleep or daytime wakefulness. Participants who remained awake after learning tended to generalize the negatively associated face, reinforcing a negative bias, while those who slept were more likely to generalize the positively associated face. These findings align with previous research showing that negative stimuli often elicit broader generalization curves than neutral or positive stimuli [1,3,4], a pattern thought to reflect an adaptive threat-detection mechanism. Our results extend this literature by showing that sleep does not

merely preserve these biases but actively reshapes the generalization landscape, attenuating
negative overgeneralization and enhancing positive generalization

fMRI data showed that greater amygdala activity during processing of either positive or negative
outcomes was linked to broader generalization of negative stimuli immediately after learning.
After sleep, this pattern reversed: higher amygdala activity was now associated with greater
generalization of positive faces, suggesting that sleep supports a renormalization of amygdala-
related emotional memory. This effect was absent in the wake group, reinforcing its sleep
dependency. Notably, participants with high amygdala activation showed, after sleep, even
greater positive generalization than at baseline, extending beyond simple renormalization.
These findings align with previous research linking amygdala activation to loss encoding and the
broadening of generalization curves for negatively associated stimuli[1,4]. Moreover, sleep
deprivation has been shown to amplify amygdala reactivity to emotional stimuli and impair
discrimination between threat and safety cues [9,64], potentially contributing to overgeneralization
in anxiety disorders. Our results suggest that sleep may help restore the negative bias by
shifting amygdala-linked generalization away from negative and toward positive stimuli.

To investigate the role of sleep-specific neurophysiology in this reorganization, we recorded
high-density EEG during overnight sleep. We found that both sigma power and the number of
fast spindle events during NREM sleep were positively correlated with the generalization of
positively associated faces. Additionally, we observed a trend suggesting that individuals who
showed greater negative generalization immediately after learning exhibited increased spindle
activity during subsequent sleep, pointing to a possible compensatory or recovery-related
renormalization mechanism. Both measures of spindle activity were significantly correlated with
the renormalization in positive face generalization, reflecting a shift from negative to positive
bias after sleep. These findings extend our understanding of the association between sleep
spindles and memory consolidation to show that spindles play a role in emotionally adaptive
reorganization of generalized learned associations.

In addition to modulating emotional memory, our results suggest that sleep spindles may
contribute to the selective suppression of emotionally neutral information. Specifically, greater
fast spindle activity was associated with reduced generalization of neutral faces, which were not
linked to reward or punishment. This pattern suggests that spindles may serve a dual function,
supporting the generalization of emotionally salient memories while promoting forgetting of

behaviorally irrelevant content. This interpretation extends prior work proposing that sleep
supports the prioritization of salient or goal-relevant information during consolidation[12,16,65].

Some limitations of this study should be explicitly acknowledged. First, circadian factors may
influence the observed group differences. Because testing sessions occurred at opposite times
of day (morning vs. evening), it is difficult to fully disentangle the effects of sleep from those of
time of day. In the behavioral experiment, participants in the day group reported higher
subjective sleepiness in the evening session than the night group did in the morning, although
objective vigilance (PVT) scores did not differ. Previous research has linked fatigue to more
negative interpretation biases[9,64], which may in principle partly explain the negative
generalization in the day group. However, neither SSS nor PVT scores were significantly
correlated with generalization measures, arguing against a major role for circadian factors.
Further arguing against circadian confounds, robust correlations between individual sleep
spindle activity in the sleep EEG experiment and shifts in emotional generalization support the
interpretation that sleep plays an active, corrective role in modulating emotional
overgeneralization. Such renormalization may be mediated in part via spindle-related
mechanisms operating during NREM sleep. Second, group-level generalization differences in
the behavioral experiment were not replicated in the fMRI experiment, where no group
differences in subjective sleepiness were found. While we can only speculate about the source
of differences between the two experiments, the learning in the fMRI experiment was much
longer with extended inter-stimulus and inter-trial intervals (a total of over 1 hour compared with
15 minutes in the behavioral task), and this could potentially have influenced the delayed
generalization effect[66].

These findings carry important translational implications. While both negative
overgeneralization[1,4,67–69] and disrupted sleep[6,8,9,70–72] have been linked to anxiety and PTSD,
few studies have examined how they interact to shape emotional memory and vulnerability to
psychopathology [17]. Our results suggest that sleep, particularly NREM spindle activity, may
serve as a protective mechanism. Sleep spindles may not only reduce negative amygdala-
related overgeneralization but also promote adaptive generalization of positive content.
Previous studies have shown that sleep spindle density increases following stress[73] and is
elevated in individuals with PTSD[74–76]. While trauma may trigger these increases, our data
suggests that spindles may also play a role in emotional recovery. Indeed, sleep spindles in
individuals with PTSD have been linked to healthier sleep**,** reduced anxiety, and improved
emotional recovery [73,77–79]. In contrast, lower spindle density has been associated with greater

vulnerability to sleep disturbances following stress[67]. Future work could build on these findings,

testing whether enhancing sleep, and particularly sleep spindles, can reduce emotional

overgeneralization in clinical populations. Promising interventions could include targeted

memory reactivation (TMR), which reintroduces learning cues during sleep to reactivate specific

memories [63,80–82], brain stimulation techniques designed to boost sleep spindle activity[83,84], and

pharmacological treatments to elevate sleep spindle activity[85,86] Applying these methods to

emotional generalization paradigms could help advance sleep-based interventions for anxiety

and PTSD.

References

1.   Dunsmoor, J. E. & Paz, R. Fear Generalization and Anxiety: Behavioral and Neural
Mechanisms. *Biol Psychiatry* **78**, 336–343 (2015).

2.   Guttman, N. & Kalish, H. I. Discriminability and stimulus generalization. *J Exp Psychol* **51**,
79–88 (1956).

3.   Schechtman, E., Laufer, O. & Paz, R. Negative valence widens generalization of learning.
Journal of Neuroscience **30**, 10460–10464 (2010).

4.   Laufer, O., Israeli, D. & Paz, R. Behavioral and neural mechanisms of overgeneralization
in anxiety. *Current Biology* **26**, 713–722 (2016).

5.   Barlow, D. H., Allen, L. B. & Choate, M. L. Toward a Unified Treatment for Emotional
Disorders – Republished Article. *Behav Ther* **47**, 838–853 (2016).

6.   Cox, R. C. & Olatunji, B. O. A systematic review of sleep disturbance in anxiety and
related disorders. *J Anxiety Disord* **37**, 104–129 (2016).

7.   Breslau, N., Roth, T., Rosenthal, L. & Andreski, P. Sleep disturbance and psychiatric
disorders: A longitudinal epidemiological study of young Adults. *Biol Psychiatry* **39**, 411–
418 (1996).

8.   Shanahan, L., Copeland, W. E., Angold, A., Bondy, C. L. & Costello, E. J. Sleep problems
predict and are predicted by generalized anxiety/depression and oppositional defiant
disorder. *J Am Acad Child Adolesc Psychiatry* **53**, 550–558 (2014).

9.   Ben Simon, E., Vallat, R., Barnes, C. M. & Walker, M. P. Sleep Loss and the Socio-
Emotional Brain. *Trends Cogn Sci* **24**, 435–450 (2020).

10.   Staner, L. Sleep and anxiety disorders. *Dialogues Clin Neurosci* **5**, 249–258 (2003).

11.   Oudiette, D., Antony, J. W., Creery, J. D. & Paller, K. a. The Role of Memory Reactivation
during Wakefulness and Sleep in Determining Which Memories Endure. *Journal of*
Neuroscience **33**, 6672–6678 (2013).

12.   Igloi, K., Gaggioni, G., Sterpenich, V. & Schwartz, S. A nap to recap or how reward
789   regulates hippocampal-prefrontal memory networks during daytime sleep in humans.
790   *Elife* **4**, 1–16 (2015).

791   13.   Gruber, M. J., Ritchey, M., Wang, S. F., Doss, M. K. & Ranganath, C. Post-learning
792   Hippocampal Dynamics Promote Preferential Retention of Rewarding Events. *Neuron* **89**,
793   1110–1120 (2016).

794   14.   Fischer, S. & Born, J. Anticipated Reward Enhances Offline Learning During Sleep. *J Exp*
795   *Psychol Learn Mem Cogn* **35**, 1586–1593 (2009).

796   15.   Feld, G. B., Besedovsky, L., Kaida, K., Münte, T. F. & Born, J. Dopamine D2-like
797   Receptor Activation Wipes Out Preferential Consolidation of High over Low Reward
798   Memories during Human Sleep Gordon. *J Cogn Neurosci* **26**, 2310–2320 (2014).

799   16.   Payne, J. D., Stickgold, R., Swanberg, K. & Kensinger, E. A. Sleep preferentially
800   enhances memory for emotional components of scenes. *Psychol Sci* **19**, 781–788 (2008).

801   17.   Davidson, P., Carlsson, I., Jönsson, P. & Johansson, M. Sleep and the generalization of
802   fear learning. *J Sleep Res* **25**, 88–95 (2016).

803   18.   Witkowski, S., Schechtman, E. & Paller, K. A. Examining sleep's role in memory
804   generalization and specificity through the lens of targeted memory reactivation. *Curr Opin*
805   *Behav Sci* **33**, 86–91 (2020).

806   19.   Ellenbogen, J. M., Hu, P. T., Payne, J. D., Titone, D. & Walker, M. P. *Human Relational*
807   *Memory Requires Time and Sleep.* www.pnas.org/cgi/content/full/ (2007).

808   20.   Davidson, P., Carlsson, I., Jönsson, P. & Johansson, M. A more generalized fear
809   response after a daytime nap. *Neurobiol Learn Mem* **151**, 18–27 (2018).

810   21.   Talamini, L. M. & Juan, E. Sleep as a window to treat affective disorders. *Current Opinion*
811   *in Behavioral Sciences* vol. 33 99–108 Preprint at
812   https://doi.org/10.1016/j.cobeha.2020.02.002 (2020).

813   22.   Vieira, T. F., Bottino, A., Laurentini, A. & De Simone, M. Detecting siblings in image pairs.
814   *Visual Computer* **30**, 1333–1345 (2014).

815   23.   Daw, N. D., O'Doherty, J. P., Dayan, P., Seymour, B. & Dolan, R. J. Cortical substrates
816   for exploratory decisions in humans. *Nature* **441**, 876–879 (2006).

817   24.   Roach, G. D., Dawson, D. & Lamond, N. Can a shorter psychomotor vigilance task be
818   used as a reasonable substitute for the ten-minute psychomotor vigilance task?
819   *Chronobiol Int* **23**, 1379–1387 (2006).

820   25.   Drummond, S. P. A. *et al.* The neural basis of the psychomotor vigilance task. *Sleep* **28**,
821   1059–1068 (2005).

822   26.   Basner, M. & Dinges, D. F. Maximizing Sensitivity of the Psychomotor Vigilance Test
823   (PVT) to Sleep Loss. *Sleep* **34**, 581–591 (2011).

824   27.   Dinges, D. F. & Powell, J. W. Microcomputer analyses of performance on a portable,
825   simple visual RT task during sustained operations. *Behavior Research Methods,*
826   *Instruments, & Computers* **17**, 652–655 (1985).

827   28.   Hoddes, E., Zarcone, V. & Dement, W. Stanford Sleepiness Scale (SSS). *Enzyklopädie*
828   *der Schlafmedizin* **1184**, (1972).

829   29.   Beck, A. T., Steer, R. A. & Brown, G. Beck Depression Inventory–II. *APA PsycTests*
830   (1996).

831   30.   Marteau, T. M. & Bekker, H. The development of a six-item short-form of the state scale
832   of the Spielberger State—Trait Anxiety Inventory (STAI). *British Journal of Clinical*
833   *Psychology* **31**, 301–306 (1992).

834   31.   Torrubia, R., Ávila, C., Moltó, J. & Caseras, X. The Sensitivity to Punishment and
835   Sensitivity to Reward Questionnaire (SPSRQ) as a measure of Gray's anxiety and
836   impulsivity dimensions. *Pers Individ Dif* **31**, 837–862 (2001).

837   32.   CLONINGER, C. R. THE TRIDIMENSIONAL PERSONALITY QUESTIONNAIRE: U.S.
838   NORMATIVE DATA. *Psychol Rep* **69**, 1047 (1991).

839   33.   Carver, C. S. & White, T. L. Behavioral inhibition, behavioral activation, and affective
840   responses to impending reward and punishment: The BIS/BAS Scales. *J Pers Soc*
841   *Psychol* **67**, 319–333 (1994).

842   34.   Baron-Cohen, S., Wheelwright, S., Skinner, R., Martin, J. & Clubley, E. The autism-
843   spectrum quotient (AQ): evidence from Asperger syndrome/high-functioning autism,
844   males and females, scientists and mathematicians. *J Autism Dev Disord* **31**, 5–17 (2001).

845   35.   Esteban, O. *et al.* fMRIPrep: a robust preprocessing pipeline for functional MRI. *Nat*
846   *Methods* **16**, 111–116 (2019).

847   36.   Esteban, O. *et al.* fMRIPrep. *Software* (2018).

848   37.   Krzysztof J Gorgolewski *et al.* Nipype. *Software* (2018).

849   38.   Gorgolewski, K. *et al.* Nipype: A flexible, lightweight and extensible neuroimaging data
850   processing framework in Python. *Front Neuroinform* **5**, (2011).

851   39.   Andersson, J. L. R., Skare, S. & Ashburner, J. How to correct susceptibility distortions in
852   spin-echo echo-planar images: Application to diffusion tensor imaging. *Neuroimage* **20**,
853   870–888 (2003).

854   40.   Tustison, N. J. *et al.* N4ITK: Improved N3 bias correction. *IEEE Trans Med Imaging* **29**,
855   1310–1320 (2010).

856   41.   Avants, B. B., Epstein, C. L., Grossman, M. & Gee, J. C. Symmetric diffeomorphic image
857   registration with cross-correlation: Evaluating automated labeling of elderly and
858   neurodegenerative brain. *Med Image Anal* **12**, 26–41 (2008).

859   42.   Zhang, Y., Brady, M. & Smith, S. Segmentation of brain MR images through a hidden
860   Markov random field model and the expectation-maximization algorithm. *IEEE Trans Med*
861   *Imaging* **20**, 45–57 (2001).

43. Fonov, V. S., Evans, A. C., McKinstry, R. C., Almli, C. R. & Collins, D. Unbiased nonlinear
average age-appropriate brain templates from birth to adulthood. *Neuroimage* (2009).

44. Jenkinson, M., Bannister, P., Brady, M. & Smith, S. Improved Optimization for the Robust
and Accurate Linear Registration and Motion Correction of Brain Images. *Neuroimage* **17**,
825–841 (2002).

45. Cox, R. W. & Hyde, J. S. Software tools for analysis and visualization of fMRI data. *NMR*
Biomed **10**, 171–178 (1997).

46. Greve, D. N. & Fischl, B. Accurate and robust brain image alignment using boundary-
based registration. *Neuroimage* **48**, 63–72 (2009).

47. Lanczos, C. *EVALUATION OF NOISY DATA*. ANAL. Ser. B*
https://epubs.siam.org/terms-privacy (1964).

48. Power, J. D. *et al.* Methods to detect, characterize, and remove motion artifact in resting
state fMRI. *Neuroimage* **84**, 320–341 (2014).

49. Behzadi, Y., Restom, K., Liau, J. & Liu, T. T. A component based noise correction
method (CompCor) for BOLD and perfusion based fMRI. *Neuroimage* **37**, 90–101 (2007).

50. Falach, R. *et al.* SleepEEGpy: a Python-based software integration package to organize
preprocessing, analysis, and visualization of sleep EEG data. *Comput Biol Med* **192**,
110232 (2025).

51. Gramfort, A. *et al.* MEG and EEG data analysis with MNE-Python. *Front Neurosci* (2013)
doi:10.3389/fnins.2013.00267.

52. Bigdely-Shamlo, N., Mullen, T., Kothe, C., Su, K. M. & Robbins, K. A. The PREP pipeline:
Standardized preprocessing for large-scale EEG analysis. *Front Neuroinform* **9**, 1–19
(2015).

53. Vallat, R. & Walker, M. P. An open-source, high-performance tool for automated sleep
staging. *Elife* **10**, 70092 (2021).

54. Donoghue, T. *et al.* Parameterizing neural power spectra into periodic and aperiodic
components. *Nat Neurosci* **23**, 1655–1665 (2020).

55. Lee, T. W., Girolami, M. & Sejnowski, T. J. Independent component analysis using an
extended infomax algorithm for mixed subgaussian and supergaussian sources. *Neural*
Comput **11**, 417–441 (1999).

56. Rupp, T. L. *Psychomotor Vigilance Performance. Encyclopedia of Sleep* (Published by
Elsevier Inc., 2013). doi:10.1016/B978-0-12-378610-4.00148-0.

57. Tibshirani, R. *Regression Shrinkage and Selection via the Lasso. Source: Journal of the*
Royal Statistical Society. Series B (Methodological) vol. 58 (1996).

58. Andrillon, T. *et al.* Sleep Spindles in Humans: Insights from Intracranial EEG and Unit
Recordings. *Journal of Neuroscience* **31**, 17821–17834 (2011).

59. Fernandez, L. M. J. & Lüthi, A. Sleep spindles: Mechanisms and functions. *Physiol Rev*
100, 805–868 (2020).

60.   Geva-Sagiv, M. *et al.* Augmenting hippocampal–prefrontal neuronal synchrony during
sleep enhances memory consolidation in humans. *Nat Neurosci* **26**, 1100–1110 (2023).

61.   Seibt, J. *et al.* Cortical dendritic activity correlates with spindle-rich oscillations during
sleep in rodents. *Nat Commun* doi:10.1038/s41467-017-00735-w.

62.   Schmidig, F. J. *et al.* A visual paired associate learning (vPAL) paradigm to study
memory consolidation during sleep. *J Sleep Res* **33**, (2024).

63.   Bar, E. *et al.* Local Targeted Memory Reactivation in Human Sleep. *Current Biology* **30**,
(2020).

64.   Motomura, Y. *et al.* Sleepiness induced by sleep-debt enhanced amygdala activity for
subliminal signals of fear. *BMC Neurosci* **15**, (2014).

65.   Stickgold, R. & Walker, M. P. Sleep-dependent memory triage: Evolving generalization
through selective processing. *Nat Neurosci* **16**, 139–145 (2013).

66.   Vassiliadis, P., Lete, A., Duque, J. & Derosiere, G. Reward timing matters in motor
learning. *iScience* **25**, (2022).

67.   Lissek, S. *et al.* Overgeneralization of Conditioned Fear as a Pathogenic Marker of Panic
Disorder. *American Journal of Psychiatry* **167**, 47–55 (2010).

68.   Dymond, S., Dunsmoor, J. E., Vervliet, B., Roche, B. & Hermans, D. Fear Generalization
in Humans: Systematic Review and Implications for Anxiety Disorder Research. *Behav*
Ther **46**, 561–582 (2015).

69.   Lissek, S. *et al.* Generalized Anxiety Disorder Is Associated With Overgeneralization of
Classically Conditioned Fear. *Biol Psychiatry* **75**, 909–915 (2014).

70.   Murkar, A. L. A. & De Koninck, J. Consolidative mechanisms of emotional processing in
REM sleep and PTSD. *Sleep Med Rev* (2018) doi:10.1016/j.smrv.2018.03.001.

71.   Babson, K. A. & Feldner, M. T. Temporal relations between sleep problems and both
traumatic event exposure and PTSD: A critical review of the empirical literature. *J Anxiety*
Disord **24**, 1–15 (2010).

72.   Vanderheyden, W. M. *et al.* Sleep alterations following exposure to stress predict fear-
associated memory impairments in a rodent model of PTSD. *Exp Brain Res* **233**, 2335–
2346 (2015).

73.   Natraj, N. *et al.* Sleep Spindles Favor Emotion Regulation Over Memory Consolidation of
Stressors in Posttraumatic Stress Disorder. *Biol Psychiatry Cogn Neurosci Neuroimaging*
8, 899–908 (2023).

74.   Van Der Heijden, A. C. *et al.* Sleep spindle dynamics suggest over-consolidation in
posttraumatic stress disorder. *Sleep* **45**, (2022).

75.   Wang, C. *et al.* Increased oscillatory frequency of sleep spindles in combat-exposed
veteran men with posttraumatic stress disorder. *Sleep* **43**, (2020).

76.   Denis, D. *et al.* Sleep Power Spectral Density and Spindles in PTSD and Their
Relationship to Symptom Severity. *Front Psychiatry* **12**, (2021).

77. Azza, Y. *et al.* Sleep's role in updating aversive autobiographical memories. *Transl*
Psychiatry **12**, (2022).

78. Dang-Vu, T. T. *et al.* Sleep spindles predict stress-related increases in sleep
disturbances. *Front Hum Neurosci* **9**, (2015).

79. Natraj, N. & Richards, A. Sleep spindles, stress and PTSD: The state of the science and
future directions. *Neurobiol Stress* **23**, (2023).

80. Cellini, N. & Capuozzo, A. Shaping memory consolidation via targeted memory
reactivation during sleep. *Ann N Y Acad Sci* 1–20 (2018) doi:10.1111/nyas.13855.

81. Oudiette, D. & Paller, K. a. Upgrading the sleeping brain with targeted memory
reactivation. *Trends Cogn Sci* **17**, 142–9 (2013).

82. Rasch, B., Büchel, C., Gais, S. & Born, J. Odor cues during slow-wave sleep prompt
declarative memory consolidation. *Science* **315**, 1426–9 (2007).

83. Lustenberger, C. *et al.* Feedback-Controlled Transcranial Alternating Current Stimulation
Reveals a Functional Role of Sleep Spindles in Motor Memory Consolidation. *Current*
Biology **26**, 2127–2136 (2016).

84. Choi, J., Kwon, M. & Jun, S. C. A Systematic Review of Closed-Loop Feedback
Techniques in Sleep Studies—Related Issues and Future Directions. *Sensors* **20**, 2770
(2020).

85. Carbone, J. *et al.* The effect of zolpidem on targeted memory reactivation during sleep.
Learning & Memory **28**, 307–318 (2021).

86. Zhang, J., Yetton, B., Whitehurst, L. N., Naji, M. & Mednick, S. C. The effect of zolpidem
on memory consolidation over a night of sleep. *Sleep* **43**, (2020).

960