The cervicovaginal microbiome of pregnant people living with HIV on antiretroviral therapy in the Democratic Republic of Congo: A Pilot Study and Global Meta-analysis.

Kimberley S. Ndlovu[1,2,4]*, Ricardo R. Pavan[1,2,4]*, Jacqueline Corry[1,3,4], Samia Mahamed[1], Natalia Zotova[5], Martine Tabala[6], Pelagie Babakazo[6], Nicholas T. Funderburg[7], Marcel Yotebieng[5], Nichole R. Klatt[8], Jesse J. Kwiek[1,3,4#], & Matthew B. Sullivan[1,2,4,9#]

AFFILIATIONS

[1]Department of Microbiology, The Ohio State University, Columbus, Ohio, USA.

[2]Centre of Microbiome Science, The Ohio State University, Columbus, Ohio, USA.

[3]Center for Retrovirus Research, The Ohio State University, Columbus, Ohio, USA.

[4]The Infectious Disease Institute, The Ohio State University, Columbus, Ohio, USA.

[5]Albert Einstein College of Medicine, Bronx, New York, USA.

[6]Kinshasa School of Public Health, Kinshasa, DRC

[7]Division of Medical Laboratory Science, School of Health and Rehabilitation Sciences, The Ohio State University, Columbus Ohio, USA

[8]Department of Surgery, Division of Surgical Outcomes and Precision Medicine

Research, University of Minnesota, Minneapolis, Minnesota, USA

[9]Department of Civil, Environmental and Geodetic Engineering, The Ohio State

University, Columbus, Ohio, USA.

*These authors contributed equally to this work.

#corresponding authors **Matthew B. Sullivan** sullivan.948@osu.edu, **Jesse J. Kwiek**

kwiek.2@osu.edu

Abstract

Recent studies are revealing that a suboptimal cervicovaginal microbiome (CVMB),

including enrichment of anaerobic bacteria associated with multiple female genital

disorders, and adverse pregnancy and birth outcomes in pregnant people. Problematically,

however, the majority of the available data to date are biased towards highly developed,

Global North countries, leaving underrepresented populations like the Democratic Republic

of Congo (DRC) poorly characterised. Here, we investigate the CVMB from a cohort of 82

pregnant people living with HIV (PLWH) on antiretroviral therapy (ART) from the DRC.

Specifically, we explore the associations between the CVMB via 16S rRNA gene sequencing

and maternal peripheral immune factors. Additionally, we compare the CVMB of PLWH-ART

from DRC to publicly available CVMB data (5 studies, 1861 samples) in a meta-analysis to

elucidate the impact of HIV on the CVMB. Combined, these analyses revealed differences

in community structure and predicted function of the microbiota between PLWH-ART and

pregnant people without HIV (PWoH). Taxonomically, the CVMB of DRC PLWH-ART were enriched for *Lactobacillus iners*-dominated CVMBs (53%) or a diverse, polymicrobial CVMB, i.e., bacterial vaginosis (BV) (43%). Functional predictions made from these taxa suggested that protein-coupled receptors, amino sugar and nucleotide sugar metabolism, fatty acid metabolism, and polycyclic aromatic hydrocarbon degradation pathways were differentially abundant between communities. Correlation with host plasma immune factors revealed putative links between some CVMB metrics (e.g., alpha diversity and species abundance) that have been linked to adverse pregnancy and birth outcomes.

Importance: HIV remains prevalent in sub-Saharan Africa, where it has been linked to adverse birth outcomes. . Suboptimal CVMBs have shown similar links. This pilot study fills critical gaps in understanding how HIV interacts with the pregnant CVMB in populations underrepresented in microbiome research, like the Democratic Republic of Congo. We identified maternal systemic immune factors associated with suboptimal CVMBs that have been linked to poor birth outcomes. In a global meta-analysis, we found significant taxonomic and functional difference in the CVMBs between pregnant people living with and without HIV, revealing potential biomarkers that for increased risks for adverse birth outcomes. These findings provide crucial insights into CVMB features that may influence pregnancy health in pregnant people living with HIV, guiding future research and tailored interventions to support safer pregnancies in the DRC and similar populations.

Introduction

Human immunodeficiency virus (HIV) is transmitted through sexual contact, blood, needles, or from mother to infant and it attacks immune cells and leads to chronic systemic infection. In July 2024, UNAIDS estimated that women in sub-Saharan Africa account for approximately 62% of HIV infections (1). Several studies have established that pregnant PLWH on ART (PLWH-ART) have a higher risk of adverse pregnancy and birth outcomes, such as low birth weight (2–4) and preterm birth (5–7). This association with adverse birth outcomes also extends to the cervicovaginal microbiome (CVMB), a community of microbes residing in the vagina, which when perturbed towards dysbiosis and bacterial vaginosis (BV) has been associated with preterm birth (8,9) and factors that can lead to preterm birth like premature rupture of membranes (10,11), chorioamnionitis (12,13), and preeclampsia (14).

Currently, the CVMB is characterized using a community state type (CST) framework which can be utilized to assess the compounding impacts of the CVMB and/or HIV infection on pregnancy and birth outcomes. Under this framework, high diversity, non-optimal CVMB, clinically referred to as BV, are classified as CST IV, and *Lactobacillus*-dominated, optimal CVMBs are classified as CSTs I, II, III, and V (15). CST III is the transitional state type that oscillates between being optimal and non-optimal (16). In sub-Saharan Africa, a high percentage of women, particularly those of African ancestry, have CST IV and CST III CVMBs, which are associated with a higher rate of HIV acquisition (17,18) and adverse pregnancy and birth outcomes (19). Therefore, there is a need to elucidate the role of how HIV-associated changes to the CVMB may modulate maternal and infant health outcomes.

The impact of HIV on the CVMB has yielded conflicting results. Some studies report

changes in composition, and therefore CST, in the CVMB of PLWH-ART (20,21) while others

show no significant differences when compared to people without HIV (PWoH) (22,23). In

contrast to these conflicting results, there are some more consistent findings, including that

CST IV and HIV-associated CVMBs are linked to increased genital inflammation marked by

increased levels of proinflammatory cytokines and chemokines (24,25). In pregnant PLWH-

ART such phenotypes are associated with adverse birth and pregnancy outcomes like

preterm birth (26–28). Similarly, systemic inflammation PLWH-ART has been well

documented (29,30) and is also linked to adverse health outcomes like cardiovascular

disease (31,32).

To date, no study has investigated the potential relationship between the CVMB and the

soluble systemic immune factors in pregnant PLWH-ART. Here, we present an exploratory

study of the CVMB of pregnant PLWH-ART from Kinshasa, Democratic Republic of Congo

and to identify CVMB - associated circulating immune factors. Furthermore, we

contextualize the results from this with other CVMB datasets that include pregnant people

in a meta-analysis, providing a more global perspective on the interactions between the

CVMB and PLWH-ART. Finally, we discuss how these findings relate to pregnancy and birth

outcomes in PLWH-ART.

Results and discussion

CQI-PMTCT study population, sampling strategy, data generation and meta-analysis

In this pilot study, we sought to generate 16S rRNA gene sequencing data to investigate the CVMB of PLWH-ART in the CQI-PMTCT cohort and its association to the systemic immune system. To that end, we collected cervicovaginal swabs and peripheral blood from a subset of 82 pregnant PLWH-ART living in Kinshasa, DRC to generate 16S rRNA gene data and plasma immune factor levels, respectively (see **Methods, Fig. 1A**). Participant characteristics, stratified by CST, including participant age, gestational age, ART regimen, gravidity, socioeconomic status, and adverse birth outcomes are summarized in **Table 1.** The CQI-PMTCT cohort included only PLWH-ART participants, and since pregnant PWoH groups were lacking for comparison, the conclusions we can make from these analyses are limited. Hence, we also conducted a meta-analysis where we integrated CVMB data from the CQI-PMTCT cohort with other published and publicly available pregnant CVMB datasets to investigate the differences in the CVMB between PLWH-ART and PWoH (**Fig. 1B**).

Fig 1. Study schematic and CVMB composition of the CQI-PMTCT cohort A) Study schematic of the CQI-PMTCT study. Peripheral blood and cervicovaginal swabs were collected from PLWH-ART in the Democratic Republic of Congo and processed in Columbus, OH. **B)** CVMB data from the CQI-PMTCT cohort were combined with other publicly available pregnant CVMB data in a meta-analysis **C)** Heatmap of RCLR abundances of top 30 most abundant microbial taxa from 81 women. CSTs were determined using VALENCIA. HIV viremia, ART regimen, and per sample Shannon Index values are shown.

Table 1: Participant demographics at enrolment stratified by community state type[a]

Characteristic	No. of participant (%)[b] with characteristic		
	All	**CST III**	**CST IV[d]**
	(n = 81)	**(n = 43)**	**(n= 35)**
Median age in years (IQR[c]**)**	32 (28-35)	33 (30-34)	30 (26-35)
HIV-1 Viral load			
Detected (> 40 copies/mL)	8 (10)	6 (14)	2 (6)
Not Detected (≤ 40 copies/mL)	42 (52)	20 (47)	19 (54)
Missing	31 (38)	17 (40)	12 (34)
Gestational age (weeks)			
Second Trimester (13-26)	28 (35)	14 (33)	13 (37)
Third Trimester (27+)	32 (40)	19 (44)	11 (31)
Missing	21 (26)	10 (23)	11 (31)

ART regimen[e]

TDF + 3TC	1 (1)	1 (2)	0 (0)
TDF + 3TC + EFV	19 (23)	9 (21)	10 (29)
TDF + 3TC + TLD	59 (73)	32 (74)	24 (69)
Missing	2 (2)	1 (2)	1 (3)

Gravidity

Primigravida	13 (16)	5 (12)	8 (23)
Multigravida	68 (84)	38 (88)	27 (77)

Adverse birth outcome

(preterm birth, stillbirth, low birth weight)

Yes	22 (27)	12 (28)	8 (23)
No	56 (67)	26 (60)	26 (74)
Missing	3 (6)	5 (12)	1 (3)

SES quartile[f]

1(lowest)	1 (1)	0 (0)	1 (3)
2	15 (19)	9 (21)	6 (17)
3	37 (46)	20 (47)	14 (40)
4 (highest)	28 (35)	14 (33)	14 (40)

[a]CST, community state type

[b]Percentages rounded off to the nearest whole number.

[c] IQR, interquartile range.

^dCST IV combined subtypes A (n=4), B (n=6) and C (n =25). Statistics of three participants in

CST II (n=2) and CST V (n=1) are not reported.

^eART, antiretroviral therapy; 3TC, lamivudine; TDF, tenofovir disoproxil fumarate; EFV,

efavirenz; TLD, tenofovir/lamivudine/dolutegravir

^fSES, socioeconomic status

In the CQI-PMTCT cohort, CST III and CST IV types are predominant in the CVMB of pregnant PLWH-ART in the DRC.

To date, no study has investigated the CVMB of people in the DRC. We generated 16S

rRNA gene sequencing data and survey the CVMB of pregnant PLWH-ART. Focusing on the

top 30 most abundant species for CVMB analyses revealed 53% of the participants (n = 43)

were dominated by *Lactobacillus iners,* while another 43% of the participants (n=35) were

dominated by a consortia of polymicrobial anaerobic bacteria (i.e., *Bifidobacterium*

leopoldii , Bifidobacterium vaginae, Fannyhesea vaginae, Prevotella spp. and *UBA629*

sp0054587) (**Fig. 1C**). These findings classify the CVMBs of PLWH-ART to CST III (transitional

CST) or CST IV (the non-optimal CST), respectively.

Evaluating diversity across these datasets revealed the following. First, within

sample diversity showed significant differences between identified CSTs (p = 0.0023,

Kruskal-Wallis) (**Fig. 2A**). This agrees with the paradigm that CST IV CVMB are more diverse

compared to *Lactobacillus*-dominated CSTs (15). Second, alpha diversity demonstrated

that samples cluster by CST (ANOSIM, R = 0.206, p = 0.001, **Fig. 2B**), with *L. iners* driving the

highest variation in the CST III cluster and *B. vaginale, B. leopoldii and F vaginae,* are driving

the variation in the CST IV cluster.

Fig 2. Ecological statistics of the CVMB in the CQI-PMTCT cohort. A). Shannon entropy

diversity stratified by CSTs (Kruskal-Wallis p =0.00234). **B)** PCA biplot of Robust Aitchison

distances coloured by CSTs (ANOSIM, R = 0.206,p = 0.001). The loadings of the top 5 most

abundant species across all samples are labelled to primary highlight drivers of variation

amongst CST clusters with black arrows to indicate the magnitude and directionality. **C)**

Shannon indices aggregated by ART regimen (Kruskal-Wallis p=0.695). **D)** PCA of Robust

Aitchison distances colored by ART regimen for 79 participants (ANOSIM R =-0.077, p =

0.825). **E)** Aggregated Shannon entropy values of HIV viremia for 50 participants (Wilcox test

p = 0.2853). **F)** PCA of robust Aitchison distances colored by detected viral load (ANOSIM R=

-0.106, p = 0.783).

Within each boxplot, the horizontal lines denote median values and the boxes extend from

the 25th to the 75th percentile. The vertical lines in each box plot show the values with 1.5

IQR. **p<0.01, ns>0.05

These results are consistent with other studies that show sub-Saharan African

women are more likely to harbor a high risk, non-optimal CST IV CVMB (17,33,34).

Additionally, other studies with similar cohorts in geography and size have shown that

pregnant PWH have CVMBs largely dominated by *Gardnerella spp.* and *L. iners* (21,28).

Incidentally, *L. iners* is the common *Lactobacilli* in CVMBs of sub-Saharan African people

and people across the globe (15,33), however it is not the most optimal *Lactobacillus*

compared to *L. crispatus* or *L. jensenii*.

Despite its high prevalence in the global CVMB, the role of *L. iners* in vaginal health

and in pregnancy and birth outcomes remains ambiguous. For example, in pregnant PLWH-

ART, *L. iners*-dominated CVMBs were associated with risk of preterm birth (21,28), however,

in pregnant PWoH, *L. iners* was associated with lower risk of preterm birth (35,36). These

studies underscore the complexity of *L. iners* role in vaginal health and pregnancy

outcomes. *L. iners* can exist with other 'non-optimal' CST IV bacteria, is less stable in the

vaginal microbiome and so can easily allow the transition to non-*Lactobacillus* dominated,

CST IV communities (i.e. BV) (37). Additionally, *L. iners* has been shown to have pathogenic

potential; for example, it produces the less protective L-lactic acid isomer (38–40), and has

highly expressed homologs of cholesterol-dependent lysins, pore forming exotoxins, which

have been found in other pathogenic bacteria and not in other *Lactobacilli* (41–43). However,

its presence even in healthy, asymptomatic people, along with its existence as a variety of

strains, demonstrates that it is also a vaginal symbiont and is able to persist in the CVMB

despite other more beneficial or protective vaginal *Lactobacilli* (44,45). Further studies are

needed to clarify the enigmatic roles of *L. iners* in vaginal health and pregnancy outcomes

using strain resolved metagenomics.

The other two most abundant bacteria in the CQI-PMTCT cohort, *B. leopoldii* and *B.*

vaginale (both formerly *G. vaginalis* until a recent amendment (46)), have been shown to be

in high abundance in sub-Saharan populations (21,28), even in healthy, asymptomatic

CVMBs (47–49). These CST IV bacteria have been associated with adverse pregnancy and

birth outcomes like preterm birth (8,50) and fetal death (51–53). Overall, our results are in

accordance with other studies showing the CVMBs of sub-Saharan African PLWH-ART have

non-*Lactobacillus* dominated (CSTs IV) or *L.iners*-dominated (CST III) CVMBs, and may

contribute to higher rates of HIV and sexually transmitted infections (54,55) as well as

preterm birth (20,21).

CVMB structure is not associated with HIV viremia and ART regimen in the CQI-PMTCT cohort.

Next, we sought to investigate clinical characteristics that might influence the CVMB

in the CQI-PMTCT. We focused particularly on type of ART regimen and HIV viremia.

Currently, there are inconclusive results on whether ART impacts the CVMB (56–58). The

same is true for HIV viremia, particularly in chronic HIV infection. (21,22,59). Therefore, we

asked whether the type of ART regimen or HIV viremia in PLWH-ART had influence in the

diversity and structure of the CVMB. We hypothesized that type of ART will impact the CVMB

and that participants with detectable HIV viremia (n = 8, **Table 1**) will have more diverse

CVMB, with communities that were distinct from those with un-detectable HIV viremia (n

=42, **Table 1**).

Contrary to our hypotheses, we detected no significant differences in alpha diversity

at the species level between ART regimen (Kruskal Wallis test p = 0.695, **Fig. 2C)** and no

differences in CVMB structure between types of ART regimen (ANOSIM R= -0.077, p = 0.825,

Fig. 2D). We also did not see differences in alpha diversity between detected and

undetected HIV viremia (Wilcoxon Test, p =0.285, **Fig. 2E**) and in beta diversity (ANOSIM R=

-0.106, p = 0.783, **Fig. 2F**).

Studies with similar cohorts in size and geographic location have shown that ART use

did not impact the CVMB of Ugandan PLWH-ART (56). However, CVMBs have been

associated with varying concentrations of ART in the female genital tract and plasma

meaning there is a possibility that long term ART use could have some influence in the CVMB

(60). Currently, it is known that topical ARTs, like PreP and tenofovir, are metabolized by CST

IV cervicovaginal bacteria thereby reducing their efficacy (61–63) but these ARTs do not

seem to alter the CVMB composition and structure (56–58). While further research might be

necessary to investigate the link between ART use and CVMB composition, our results

contribute to the growing knowledge of the impact, or lack thereof, between ART use and the

CVMB (64).

Studies have shown before that there are no differences in vaginal communities due

to HIV viremia or status (22,59). Conversely, other 16S rRNA gene studies have shown that

HIV-associated CVMBs have a lower abundance of *Lactobacillus* species but there was no

conclusion on how CVMB structure (i.e. beta diversity) is different (55,65,66). Curiously,

there is inconclusive evidence on how HIV influences CVMB composition. On one hand,

cervicovaginal bacteria have been shown to interact with the HIV virus or host cells to

increase risk of contracting HIV. For example, CST IV bacteria like *G. vaginalis* induce a local

inflammation response, reduce cervicovaginal epithelial cell barrier and integrity hence

allowing bacterial and viral pathogens, like HIV, to infect the host (17,67,68). A recent study

also showed that bacterial lectins specific to O-glycans, bind to the HIV-1 virion and

envelope glycoproteins to increase HIV infectivity and resistance to antibodies (69).

However, the mechanisms by which HIV directly impact bacterial physiology and

metabolism remain undefined and unexplored. It is possible that the HIV viral load in the

vagina is sufficiently suppressed for there to be a detectable interaction between the virus

and vaginal bacteria (70,71), however studies on this subject are highly limited. The

ambiguity of the impact of chronic and systemic viral sexually transmitted infections like HIV

on the CVMB remains to be fully explored however, based on our results for the CQI-PMTCT

cohort, we can tentatively conclude and recapitulate the findings of studies that state HIV

viremia does not influence the CVMB (22,23,59). The high percentage of people with

undetectable HIV viremia is a testament to the effectiveness of ART use in this cohort.

Meta-analysis reveals that CVMB are distinct between PLWH-

ART and PWoH globally.

The CQI-PMTCT study discussed above has limitations including lack of a pregnant

PWoH group for comparison. Hence, to better understand the role of HIV on the CVMB, we

integrated this study with five other publicly available pregnant CVMB datasets. To reduce

batch effects, data from studies were selected to be as similar to the CQI-PMTCT CVMB data

as possible (see **Methods** and **Fig. S1**).

After adjusting for study specific batch effects using PLSDA (see **Methods**), we found

that within sample (alpha) species diversity was significantly higher in samples from PLWH-

ART compared to PWoH (**Fig. 3A**) Additionally, CVMB community structure was significantly

different between pregnant PLWH-ART and pregnant PWoH CVMBs (ANOSIM R= 0.518, p

=0.001, **Fig. 3B**). When stratified by CST, within samples species diversity was highest in CST

IV samples, which is suboptimal in the CVMB (**Fig. 3C)** and CVMB community structure was

different by CSTs (ANOSIM R= 0.399, p =0.001, **Fig. 3D**). All this indicates that there are

different bacterial communities between the PLWH-ART and PWoH groups.

Fig 3. Ecological statistics of the integrated datasets. A) Shannon Entropy values between

CVMBs of PWoH and PLWH-ART separated by study. **B)** PCA of Aitchison Distances showing

different bacterial communities by health status (ANOSIM R= 0.518, p= 0.001). **C)** Shannon

Entropy values stratified by CST (Kruskal-Wallis p = 1.023e-35). **D)** PCA of Aitchison

Distances showing different bacterial communities by CST (ANOSIM R=0.399, p= 0.001).

*p<0.05, ****p<10^{-4}

To further investigate this claim, we used differential abundance analyses (see

Methods) to identify bacterial species that were differentially enriched between the two

groups. We found that 21 cervicovaginal bacterial species from 14 families were

differentially abundant between the PLWH-ART and PWoH groups (**Fig. 4**). *Lactobacilli spp.,*

namely beneficial bacteria such as *L. crispatus, L. gasseri, L. hominis* and *L. jensenii* and

Limosilactobacilus reuteri were more abundant in the CVMB of PWoH. In contrast, the

PLWH-ART group had high abundance of CST IV, non-optimal, BV-associated bacteria

compared to the PWoH group. For example, species in the *Bacteroidaceae* family,

particularly *P. amnni* and *P.timonensis spp.* were more abundant in the PLWH-ART group.

Other CST IV associated bacteria that were more abundant in the PLWH-ART group were *B.*

leopoldii, M. lornae, KA00274 sp001552885 (AKA *Amygdalobacter nucleatus*), *F.magna, F.*

vaginae, M. hominis and *C. tuberculostearicum* (**Fig. 4**).

Fig 4. Boxplots showing differentially expressed species in pregnant PLWH-ART and

pregnant PWoH as determined using three differential abundance analysis methods (see

Methods). Within each box, the vertical lines denote median values, and the boxes extend

from the 25th to the 75th percentile. The horizontal lines in each box plot show the values with

1.5 IQR.

Other studies have found *Lactobacillus spp.* to be more dominant in PLWH not on ART

compared to those on ART, regardless of pregnancy state (55,65,66). This is corroborated by

the CST IV, non-optimal bacteria being more differentially abundant in PLWH-ART group

compared to the PWoH. It is possible that this differential abundance in (non)-

Lactobacillus-dominated CVMBs between PLWH-ART and PWoH existed prior to and during

HIV acquisition, as such differences are associated with risk of HIV acquisition (72). For that

reason, these communities may have remained stable in their respective states since.

Indeed, one study determined that in 79.7% of CVMBs from two longitudinal cohorts (16,73),

CSTs were predominantly mono-stable and rarely transitioned to completely different CSTs,

which could explain the persistence of these bacteria in the CVMB even after HIV acquisition

(37). It is also possible that the influence of maternal age, racioethnicity and geography (**Fig.**

S2) distorts the impact of HIV status on the CVMB in this meta-analysis as seen before

(47,48,74,75). Therefore, we tentatively conclude that HIV infection is associated with

changes in the CVMB of pregnant people as seen in a recent study with similar populations

(76). However, future research will have to explore the causative associations of HIV on the

pregnant CVMB and how this alters pregnancy outcomes.

Predicted metabolic pathways were differentially abundant among CSTs in the CQI-PMTCT cohort

310   The CST framework classifies the CVMB based on taxonomic composition (15). However, even when microbial communities differ in their taxonomic composition, they might have functional redundancy (77) or conversely, diverse functional capabilities in the same species (78). Hence, functional prediction of the CVMB may yield more specific insights into potential phenotypic differences between CST, and thus host responses. Here, we predicted the function of the CVMB, stratified by CST, using PICRUSt2 based on the KEGG (Kyoto Encyclopedia of Genes and Genomes) database (79,80). We predicted 6466 KEGG Orthologs (KOs), which were assigned to 162 KEGG pathways. Of note, these are predicted functions and not direct readouts by the bacterial species.

319   Using multiple differential abundance analyses (see **Methods**), we identified four (4) KEGG pathways that were differentially abundant between CSTs (**Fig. 5**), namely G protein-coupled receptors (GPCRs, KO04030), amino sugar and nucleotide sugar metabolism (KO00520), fatty acid metabolism (KO01212) and polycyclic aromatic hydrocarbon degradation (KO00624). Fatty acid metabolism and polycyclic aromatic hydrocarbon degradation had higher relative abundance in CST IV compared to CST III, with the highest abundance of both in CST II. G protein-coupled receptors and amino sugar and nucleotide sugar metabolism were more abundant in CST III, followed by CST IV and lastly CST II (**Fig. 5**).

Fig 5. Predicted KEGG pathways differentially abundant between CSTs in the CQI-PMTCT

331 cohort as determined using three differential abundance analysis methods (see **Methods).**

Because there were only two samples that were assigned CST II, we will focus our discussion on CST III and IV only. A combination of predicted CVMB genotypes suggests a plausible mechanism for disruption of host homeostasis. Short-chain fatty acid (SFCA) metabolism was more abundant in CST IV compared to CST III CVMBs, indicating greater presence of bacteria capable of producing SFCAs. CST IV bacteria play a large role in SCFA production, and the increased presence of SFCAs like propionate, acetate and butyrate have been associated with cervicovaginal dysbiosis (81). Furthermore, the production of SCFA leads to decreased *Lactobacillus* abundance and increased pH in the vagina (81–83). Based on this, we can posit that there were high concentrations of cervicovaginal SFCAs in this cohort. GPCRs were slightly, yet still significantly, more abundant in CST III compared to CST IV CVMSs. GPCRs mediate the immunoregulatory effects of SFCAs (82) and specifically, GPCRs 41, 43 and 109A, can inhibit histone deacetylases (HDACs) which subsequently induce a cellular and humoral immune response (82,84). Simultaneously increased SFCAs and decreased GPCRs may induce a local pro-inflammatory response in this cohort. Given that inflammation in the cervicovaginal tract is associated with adverse birth outcomes, like preterm birth in pregnant PLWH-ART (21), which may be linked to increases in SFCA production and decreases in the immunoregulators in the CVMB.

Amino sugar and nucleotide sugar metabolism was another pathway that might be implicated in birth outcomes in pregnant PLWH-ART. For bacteria, these are core intermediates involved in peptidoglycan and lipopolysaccharide, among other macromolecule, biosynthesis, energy metabolism, and other processes. In this cohort, this

pathway was more abundant in CST III compared to CST IV CVMBs. One study found that

amino sugar and nucleotide metabolism was negatively associated with mammalian target

of rapamycin (mTOR), a kinase that regulates cell growth and proliferation (85). In the same

study, mTOR was negatively correlated with *L. iners* and positively correlated with

polymicrobial, CST IV bacteria (85). While some of this association may be due to major

structural features of the CVMB, i.e., the presence of a lipopolysaccharide layer of some

members but not others, these metabolic products from the CVMB of pregnant PLWH-ART

may interact with the mTOR pathway, activating it (by CST III bacteria) or inhibiting it (by CST

IV bacteria). This interaction could maintain or disrupt cervicovaginal epithelial integrity,

potentially determining whether birth outcomes are adverse or normal through a

proinflammatory or anti-inflammatory response and either increasing or decreasing the risk

of STIs.

Lastly, polycyclic aromatic hydrocarbon (PAH) degradation was also differentially

more abundant in CST IV than in CST III CVMBs. PAHs, which are ubiquitous in the

environment can impact the CVMB (9) which can lead to adverse birth outcomes like

intrauterine growth restrictions and preterm birth (86,87). The polymicrobial state type, CST

IV, tends to have varying abundances of putative PAH-degrading bacterial genera including

BVAB2 and *Megasphaera, Sphingomonas*, *Acinetobacter*, *Micrococcus*, *Pseudomonas*,

and *Ralstonia,* and two of the genera, *Megasphaera* and *Pseudomonas* are present in the

CQI-PMTCT cohort (88–90). Therefore, somewhat unexpectedly, a more diverse,

polymicrobial CST IV CVMB that may be better able to degrade PAHs might offer some

protection against their harmful effects. Perhaps, because the Sub-Saharan African region

is exposed to a substantial amount of PAHs (91,92), a CST IV CVMB might be advantageous

and help explain its prevalence in Sub Saharan African populations (17,48,93). However, a

multidisciplinary research study, across life and social sciences, would need to be

conducted to investigate this claim (94,95), which is an important next step. Overall, these

findings provide the direction for further studies to clarify these differences with more

granularity and the role of bacterial metabolic pathways on reproductive and birth outcomes

in pregnant PLWH-ART in vulnerable and underrepresented populations. Additionally, future

multi-omics studies should be conducted to progress the CST framework from taxonomy-

only based classification to microbial function and taxonomy-based classification.

Meta-analysis reveals predicted metabolic pathways were

differentially abundant between PLWH-ART and PWoH.

We used PICRUSt2 to predict 4,534 KOs which were assigned to 130 KEGG pathways.

Of those, 62 KEGG pathways were differentially abundant between pregnant PWoH and

PLWH-ART CVMBs (**Fig. 6**).

Fig 6. KEGG pathways differentially abundant between pregnant PLWH-ART and pregnant

PWoH as determined using three differential abundance analysis methods (see **Methods)**.

Most of these pathways were primarily involved in metabolism. This could indicate

that HIV-associated changes to the pregnant CVMB may alter microbial function, which

could influence pregnancy and birth outcomes and susceptibility to other cervicovaginal

infections in PLWH-ART. For this study, we will focus on only a few pathways that might have

relevance with HIV, and pregnancy and birth outcomes. Lysine degradation (KO00310) was

differentially more abundant in the PLWH-ART group compared to the PWoH group.

Products of lysine degradation — cadaverine and pipecolate — were increased in

cervicovaginal fluids of women with BV (96). BV, indicative of CST IV, has been associated

with adverse birth outcomes in PLWH-ART (20,21,28) and increased risk of HIV acquisition

(17). Cysteine and methionine metabolism (KO00270) was differentially more abundant in

the PLWH-ART group compared to the PWoH group, and the opposite is true for folate

biosynthesis (KO00790). Levels of cysteine in maternal plasma or cervicovaginal fluid have

been associated with adverse pregnancy and birth outcomes (97–99). These findings

provide direction for further research to investigate how and if HIV-associated changes to

the CVMB function impact pregnancy and birth outcomes. One pathway, mucin O-glycan

synthesis (KO00512) was differentially more abundant in PLWH-ART and this has been

shown to increase the infectivity of sexually transmitted viral pathogens like HIV (69). Future

research should investigate how this pathway is involved in pregnancy and birth outcomes

in PLWH-ART.

CVMB characteristics are correlated with some plasma immune

factors in the CQI-PMTCT cohort

418   Next, we asked if there was an association between plasma immune factors and the

CVMB in the CQI-PMTCT cohort. The correlation and causative associations between CVMB

and cervicovaginal cytokines and chemokines have been well studied (17,24,67,100). Few

studies have identified the link between the systemic immune system and the CVMB in

PLWH-ART. To this end, we used linear models to identify the CVMB metrics (Shannon

diversity, CST and bacterial species abundance) that were associated with the systemic

immune system (see **Methods**). We hypothesized that CST IV bacteria or highly diverse

CVMBs would be associated with plasma immune factors that are linked to adverse birth

outcomes.

Shannon diversity was positively correlated with plasma MMP-2 ($r = 0.214$, $p = 0.051$)

and CXCL13 ($r = 0.353$, $p = 0.005$) (**Fig. 7A-B**). When looking at individual taxa, *L. iners* was

negatively correlated with sCD163 ($r = -0.32$, $p = 0.013$ (**Fig. 7C**). In contrast, sTNFRSF1A

concentrations were significantly higher in individuals with the CST III CVMBs vs those with

CST IV CVMBs (**Fig. 7D**). However, in most cases, CST IV associated bacteria were positively

correlated with immune markers of inflammation (**Fig. S3**). *L. crispatus,* a bacterium

associated with low diversity in the CVMB, was negatively correlated with MMP-2 and

CXCL13 and other biomarkers of inflammation (**Fig. S3**). In contrast, *L. gasseri,* also in low

diverse CVMBs, was positively correlated with pro- and anti- inflammatory immune factors

(**Fig. S3**).

Fig 7. Correlations of CVMB metrics to plasma immune mediators A-B) Spearman

correlation of MMP2 and CXCL13 to Shannon diversity **C)** sCD163 to *L. iners* abundance and

D) Levels of sTNFRSF1A between CST III and IV (Wilcoxon Test p = 0.0188). Only statistically

significant correlations (not corrected) are shown and only the top 3 abundant taxa are

shown. Linear regression lines of A-C are shown in blue, and the shaded regions represent

95% confidence intervals. Full abbreviations of immume factors are in **Table S3**

Genital inflammation, linked to cervicovaginal dysbiosis, has been associated with

adverse pregnancy and birth outcomes (21,101,102). Here, we investigate whether systemic

inflammation is also linked to the CVMB. MMP-2 and CXCL13 were positively correlated with

Shannon diversity. Abnormal concentrations and varying expression of MMP-2 during

pregnancy have been implicated in adverse pregnancy outcomes like preeclampsia

(103,104), recurrent spontaneous abortion (105), preterm birth (106) and gestational

trophoblastic disease (107) and pregnancy outcomes may lead to adverse infant outcomes

that impact infant mortality and development (108). The same is true for CXCL13 as it has

been associated with placental inflammation (109) and preterm birth (110). Similarly,

adverse pregnancy and birth outcomes, have been linked with highly diverse CVMBs,

particularly those with CST IV communities (8,14,21,28). Altogether, this suggests that there

is a complex interplay between the CVMB and systemic inflammation in PLWH-ART that

impacts pregnancy and birth outcomes.

L. iners was negatively correlated with sCD163, an immune factor that has also been

linked to preterm birth (111) and is a predictor of mortality in PLWH-ART (112,113) However,

one study showed that sCD163 concentrations varied during pregnancy in PLWH-ART,

declining in the first trimester and then remaining stable (114). All of this suggests that the

role cervicovaginal bacteria have in modulating the systemic immune system in pregnant

PLWH-ART — and therefore pregnancy and birth outcomes — is complex and may depend

on the host's immunological state. Indeed, the impact gut bacteria have on the systemic

immune system might greatly confound the effect of cervicovaginal bacteria on the systemic

immune system (115–117). Given that the local cervicovaginal immune profile is distinct

from the systemic immune profile, cervicovaginal bacteria may have very little effect on the

systemic immune profile (118). However, these associations between the plasma immune

factors and maternal CVMB may be important for understanding the biological mechanisms

underlying HIV-associated pregnancy and associated infant complications.

Limitations, future opportunities and conclusions

The power of our study is limited by the lack of a pregnant PWoH group for

comparison. This study was nested within a larger longitudinal study from the CQI-PMTCT

study that was evaluating the long-term effects of ART usage in pregnant and chestfeeding

people (119). As such, samples were only collected from pregnant PLWH-ART. Subsequent

studies will have samples from pregnant PLWH-ART and PWoH and will also include larger

sample sizes for better statistical inference. We did not present the impact of CVMB on birth

outcomes in the CQI-PMTCT cohort as there were not enough samples in each adverse birth

outcome to reach statistical significance.

Even acknowledging these limitations, our study revealed some important details about

the CVMB of PLWH-ART in DRC and globally. We showed that the CVMB of pregnant PLWH-

ART is largely CST III and IV both of which differ in their predicted functions. We found

associations between the systemic immune system and the CVMB, providing direction for more studies to investigate this association as it might have implications on maternal and infant health. In a meta-analysis, we found that the HIV-associated CVMB and predicted function in pregnant PLWH-ART is distinct from PWoH, contrary to previous studies. Subsequent studies with larger cohorts in the same population in Kinshasa, DRC, will utilize shotgun metagenomic sequencing for genome-resolved analyses and better functional inference of bacterial and viral populations and their association with adverse birth outcomes in PLWH-ART. This will allow us to infer any strain level differences in the CVMB that may cause different host phenotypes, therefore ecosystem roles that are not detected with 16S rRNA gene sequencing. The CST framework would benefit to move on from taxonomy only based classification to microbial function and taxonomy-based classification that will give a better insight on the role of the CVMB on health and disease.

Materials and methods

Ethics approval and consent to participate

The Ohio State Institutional Review Board (number 2015H0440), Albert Einstein College of Medicine Institutional Review Board (number 2020-12018), and the University of Kinshasa School of Public Health Ethical Committee, and the University of Kinshasa School of Public Health Institutional Ethical Committee (ESP/CE/046/2018) granted approval for conducting this study and for protocol related specimen testing. All research was conducted in accordance with relevant guideline and regulations. Participants in the CQI-PMTCT study provided signed consent before undergoing study procedures

Study cohort and sample collection

Participants in this analysis were enrolled in Kinshasa, Democratic Republic of Congo. This was a pilot study nested within the larger in the Continuous Quality Intervention: Preventing Mother to child transmission (CQI-PMTCT) study, a longitudinal study evaluation the long term outcomes of ART on pregnant people (NCT03048669) (120). From October 2020 to May 2021, we collected cervicovaginal swabs and peripheral blood from 82 participants at enrolment in their second or third trimesters who could access medical facilities during the SARS-CoV-2 pandemic. The 82 participants represent 40% of total participants who consented to provide cervicovaginal swabs. Clinicians collected cervicovaginal swabs and stored them in Puritan DNA/RNA shield (ZymoResearch, R1108) and transported to the central laboratory at Protestant University of Congo to be stored at -80C until shipped to Columbus, OH on dry ice for processing. Plasma collected from the peripheral blood was processed immediately and stored at -80C until shipment to Columbus, OH for processing (**Fig. 1A**).

Demographic data including participant age, gestational age, ART regimen, gravidity, socioeconomic status, depression scores were collected at enrolment. The first principal component analysis (PCA) of the factors indicative of the participants socioeconomic status (SES) was used to categorize the SES status to quartiles with 0 as the lowest and 3 as the highest (121). Adverse birth outcomes including preterm birth, stillbirth and low birth weight were collected at delivery.

Study selection, search strategy and criteria for meta-analysis

16S rRNA gene CVMB studies were selected by conducting a thorough and manually

curated search of the Web of Science Core Collection of Thomson Reuters (WOS) and the

European Nucleotide Archive (ENA) for studies published until December 2022. The search

terms, hits and justification for exclusion or inclusion are shown in **Fig. S1** and information

about the selected studies are in **Table S1**. The inclusion criteria were as follows: (i)

pregnant healthy or living with HIV (iii) availability of raw data and (iii) Illumina paired-end

short read amplicon sequencing of the V3-V4 hypervariable region. Only studies with

sufficient information for comparison were included (i.e., metadata and minimally

reproducible 16S rRNA gene sequence processing pipeline). Longitudinal studies were

excluded from this analysis. Metadata from each study was obtained from NCBI, and/or via

correspondence with the authors of the selected studies. Among these studies, three were

from Europe –one from Italy (122) and two from Finland (123,124), one from the United

States (125), and one from Uganda (126). Only the Ugandan study included samples from

PLWH-ART (n = 5). In total, including this study, 1861 samples were analyzed. More details

on the studies can be found in **Table S1.**

DNA extraction, 16S rRNA gene sequencing and processing

Bacterial DNA from cervicovaginal swabs in the CQI-PMTCT cohort were extracted

using the Qiagen AllPrep DNA/RNA Mini kit (Qiagen, 80204) following the manufacturer's

manual. These samples were low microbial biomass, hence, to reduce contamination,

samples were extracted in a clean room and all materials were treated with 70% ethanol,

RNAse and UV light for ~15minutes before DNA extraction. Additionally, reagent-only

controls were extracted in parallel with each DNA extraction batch. The PCR products of the

reagent-only controls were examined on 1.2% gels, and no bands were detected.

Samples were sent to SeqCenter (https://www.seqcenter.com/) for amplification of

the V3-V4 hypervariable region of the 16S rRNA gene using the 341F (5'-

CCTACGGGDGGCWGCAG-3') and 806R (5'-GACTACNVGGGTMTCTAATCC-3') primers

using the Zymo Research's Quick-16S kit. The V3-V4 region was chosen as it has been shown

to identify the most taxa in cervicovaginal samples (127). The PCR products were sequenced

on an Illumina NextSeq2000 to generate 2x301bp paired-end reads.

16S rRNA gene reads were processed following the DADA2 SOP (128). Sequence

quality was assessed using FASTQC v0.11.5 and any adapters or primer sequences were

trimmed using Trimmomatic v0.36 (129,130). Generally, sequences were trimmed at the

first base with a quality score of Q<25. Forward and reverse reads were merged using the

mergePairs function with default parameters (minOverlap = 12, maxMismatch = 0), and

chimeras removed using removeBimeraDenovo with default parameters. Taxonomy was

assigned using the Greengenes2 database (131). Sequences were filtered to remove

sequences of mitochondria and chloroplast, samples with less that 100 reads and taxa with

less than 10 reads, less than 0.5% abundance across all samples and not present more than

2 times in 10% of samples were removed. The resultant ASV table was agglomerated to

species-level and robust centered log-ratio (RCLR) transformation was performed on the

table using the decostand function on the vegan package v2.6-6. This transformation was to

mitigate the inherent negative correlation bias in compositional data (132,133) The

transformed table was used in all subsequent analyses.

For the five (5) selected studies for the meta-analysis, sequences were processed on

a per-study basis as described above. Reads were filtered based on read quality for each

study. Per study processing parameters are shown in **Table S2.** Generated ASV tables per

study were combined and taxonomy was assigned on the combined table (34,871 ASVs and

1861 samples). The aggregated table was further filtered by removing ASVs not assigned to

any taxonomic level. To reduce data complexity (134), bacterial ASVs were filtered if

observed at frequencies of 0.0005% study-wide and observed in only 2 samples. Samples

with fewer than 2000 reads or with excess zeroes were removed using the PreFL function

from the PLSDAbatch package v1.0 (135). After filtering, ASVs were agglomerated at the

species level resulting in an abundance table with 74 species-level taxa and 664 samples. A

centered log-ratio (CLR) transformation was performed on the abundance table. Before the

CLR transformation, zeros were replaced with a constant value smaller than the detection

limit (e.g., 65% of the detection limit) (136,137).

PICRUSt2 version v2.4.2 (79) was used to predict functional composition from the

16S rRNA gene sequences. Data were normalized based on the abundance of 16S rRNA

gene copy numbers. Metagenome predictions were used to estimate KEGG pathways and

any unassigned KEGG IDs were mapped using KEGGREST v3.19 (138).

Quantification of plasma cytokine, chemokine and soluble factors (immune factors)

591
592

Levels of cytokines, chemokines and soluble factors were measured from maternal

plasma were quantified using the LEGENDplex COVID-19 Cytokine Storm Panel 1 & 2

(BioLegend, 741095), LEGENDplex Human Proinflammatory Chemokine Panel 1

(BioLegend, 740984) and LEGENDplex Vascular Inflammation panel 1 (BioLegend, 740551).

Additional details related to measurement of immune factors are described in

Supplemental Methods.

Statistical analyses

For the CQI-PMTCT study, statistical analyses were as follows. Community state

types (CST) were assigned to each sample using VALENCIA, which assigns CSTs to each

samples based on their similarity to a reference centroid (139). Alpha diversity indices

(Chao1, Simpson, and Shannon) were estimated using an asymptotic statistical approach

(140,141). Differences in alpha diversity were tested using the Kruskal-Wallis test, followed

by Wilcoxon pairwise comparison tests if significant.

Beta diversity analyses were computed using principal component analysis (PCA) on

Robust Aitchison distances. Significance of PCAs used ANOSIM from the vegan package

v2.6-10. To identify associations between CVMB metrics (alpha diversity and bacterial

abundance) and the log transformed plasma cytokine, chemokine and soluble factors

concentrations, linear regression models were computed with the with Spearman's rank

correlation. Adjustments for multiple hypothesis testing were not performed due to this

being an exploratory study unless as a sensitivity analysis for our main findings (142)

For the meta-analysis, data were analyzed as following. CSTs were assigned as

described above using VALENCIA. Batch effects were accounted for using Partial Least

Square Discriminant Analysis (PLSDA-batch) after evaluation of several batch removal

methods (see Supplemental Methods for more details). Alpha diversity indices and beta

diversity on Aitchison distances were computed as above.

To identify differentially abundant species and predicted functions, three statistical

methods were used: LinDA from the MicrobiomeStat package v.1.1 (143,144) , ANCOM-BC

v.2.2.1(145) and ALDEx2 v.1.32.0 (146) with the formula "abundance ~ health_status +

study". A species was considered differentially abundant if it met the following criteria: p-

adjusted value < 0.05 in LinDA; q-value < 0.05 in ANCOM-BC; or effect size > 0.5 in ALDEx2.

Additional details on differential abundance analysis are described in Supplemental

Methods. The STORMS checklist (147) (**Table S5**) is included for standardization and

reproducibility of data.

Data availability

627 The datasets supporting the conclusions of this article are available in the Sequence

628 Read Archive (SRA) under BioProject PRJNA1234224 and scripts and data used to analyses

629 all the data are available on https://github.com/SulliVagOSU/16S_meta_analysis.

Acknowledgments

This work was funded by the National Institute of Child Health and Human Development (NICHD) under the award numbers R01HD087993 and R01HD105526. The funders had no role in study design, data collection and analysis, interpretation and preparation of this manuscript. The authors declare no competing interests.

We thank all the people who participated and contributed to this study. We also thank the contribution of the CQI-PMTCT study team, the staff of the participating clinics and the provincial and national health authorities. We would also like to thank members of the Sullivan and Kwiek labs for their support and critical feedback. Bioinformatics analysis was supported in part by the Ohio Supercomputer Center. Figure 1A and 1B were created in BioRender.

References

1. Global HIV & AIDS Statistics-2024 Fact Sheet | UNAIDS [Internet]. [cited 2024 Oct 4]. Available from: https://www.unaids.org/en/resources/fact-sheet

2. Fentie EA, Yeshita HY, Bokie MM. Low birth weight and associated factors among HIV positive and negative mothers delivered in northwest Amhara region referral hospitals, Ethiopia,2020 a comparative crosssectional study. PloS One. 2022;17(2):e0263812.

3. Sofeu CL, Warszawski J, Ndongo FA, Penda IC, Nadiang ST, Guemkam G, et al. Low Birth Weight in Perinatally HIV-Exposed Uninfected Infants: Observations in Urban Settings in Cameroon. PLOS ONE. 2014 Apr 3;9(4):e93554.

4. Xiao PL, Zhou YB, Chen Y, Yang MX, Song XX, Shi Y, et al. Association between maternal HIV infection and low birth weight and prematurity: a meta-analysis of cohort studies. BMC Pregnancy Childbirth. 2015 Oct 8;15(1):246.

656   5.   Duffy CR, Herlihy JM, Zulu E, Mwananyanda L, Forman L, Heeren T, et al. Preterm birth
657   among women with HIV: impact of preconception cART initiation. AIDS. 2024 Oct
658   1;38(12):1749.

659   6.   Shehu AY, Ogboghodo EO, Enaruna N, Jamda MA. Outcome of Delivery among HIV-
660   Positive Women that Attended the Antenatal Clinic at the University of Benin Teaching
661   Hospital, Benin City, Nigeria. Niger J Med. 2022 Apr;31(2):212.

662   7.   Twabi HS, Manda SO, Small DS. Assessing the effects of maternal HIV infection on
663   pregnancy outcomes using cross-sectional data in Malawi. BMC Public Health. 2020
664   Jun 22;20(1):974.

665   8.   Fettweis JM, Serrano MG, Brooks JP, Edwards DJ, Girerd PH, Parikh HI, et al. The
666   vaginal microbiome and preterm birth. Nat Med. 2019;25(6):1012–21.

667   9.   Kindschuh WF, Baldini F, Liu MC, Liao J, Meydan Y, Lee HH, et al. Preterm birth is
668   associated with xenobiotics and predicted by the vaginal metabolome. Nat Microbiol
669   2023. 2023 Jan 12;1–14.

670   10.   dos Anjos Borges LG, Pastuschek J, Heimann Y, Dawczynski K, Bergner M, Haase R, et
671   al. Vaginal and neonatal microbiota in pregnant women with preterm premature
672   rupture of membranes and consecutive early onset neonatal sepsis. BMC Med. 2023
673   Mar 13;21(1):92.

674   11.   Yan C, Hong F, Xin G, Duan S, Deng X, Xu Y. Alterations in the vaginal microbiota of
675   patients with preterm premature rupture of membranes. Front Cell Infect Microbiol.
676   2022;12:858732.

677   12.   Prince AL, Ma J, Kannan PS, Alvarez M, Gisslen T, Harris RA, et al. The placental
678   microbiome is altered among subjects with spontaneous preterm birth with and
679   without chorioamnionitis. Am J Obstet Gynecol. 2016 May;214(5):627.e1-627.e16.

680   13.   Urushiyama D, Ohnishi E, Suda W, Kurakazu M, Kiyoshima C, Hirakawa T, et al.
681   Vaginal microbiome as a tool for prediction of chorioamnionitis in preterm labor: a
682   pilot study. Sci Rep. 2021 Sep 23;11(1):1–10.

683   14.   Lin CY, Lin CY, Yeh YM, Yang LY, Lee YS, Chao A, et al. Severe preeclampsia is
684   associated with a higher relative abundance of Prevotella bivia in the vaginal
685   microbiota. Sci Rep. 2020 Oct 26;10(1):18249.

686   15.   Ravel J, Gajer P, Abdo Z, Schneider GM, Koenig SSK, McCulle SL, et al. Vaginal
687   microbiome of reproductive-age women. Proc Natl Acad Sci U S A. 2011 Mar
688   15;108(SUPPL. 1):4680–7.

16.   Gajer P, Brotman RM, Bai G, Sakamoto J, Schütte UME, Zhong X, et al. Temporal
Dynamics of the Human Vaginal Microbiota. Sci Transl Med. 2012 May
2;4(132):132ra52.

17.   Gosmann C, Anahtar MN, Handley SA, Farcasanu M, Abu-Ali G, Bowman BA, et al.
Lactobacillus-Deficient Cervicovaginal Bacterial Communities Are Associated with
Increased HIV Acquisition in Young South African Women. Immunity. 2017 Jan
17;46(1):29–37.

18.   Murewanhema G, Musuka G, Moyo P, Moyo E, Dzinamarira T. HIV and adolescent girls
and young women in sub-Saharan Africa: A call for expedited action to reduce new
infections. IJID Reg. 2022 Aug 28;5:30–2.

19.   Juliana NCA, Peters RPH, Al-Nasiry S, Budding AE, Morré SA, Ambrosino E.
Composition of the vaginal microbiota during pregnancy in women living in sub-
Saharan Africa: a PRISMA-compliant review. BMC Pregnancy Childbirth. 2021 Sep
3;21(1):596.

20.   Price JT, Vwalika B, Hobbs M, Nelson JAE, Stringer EM, Zou F, et al. Highly diverse
anaerobe-predominant vaginal microbiota among HIV-infected pregnant women in
Zambia. PLOS ONE. 2019 Oct 1;14(10):e0223128.

21.   Price JT, Vwalika B, France M, Ravel J, Ma B, Mwape H, et al. HIV-associated vaginal
microbiome and inflammation predict spontaneous preterm birth in Zambia. Sci Rep.
2022;12(1).

22.   Mehta SD, Donovan B, Weber KM, Cohen M, Ravel J, Gajer P, et al. The Vaginal
Microbiota over an 8- to 10-Year Period in a Cohort of HIV-Infected and HIV-
Uninfected Women. PLOS ONE. 2015 Feb 12;10(2):e0116894.

23.   Reimers LL, Mehta SD, Massad LS, Burk RD, Xie X, Ravel J, et al. The Cervicovaginal
Microbiota and Its Associations With Human Papillomavirus Detection in HIV-Infected
and HIV-Uninfected Women. J Infect Dis. 2016 Nov 1;214(9):1361–9.

24.   Anahtar MN, Byrne EH, Doherty KE, Bowman BA, Yamamoto HS, Soumillon M, et al.
Cervicovaginal bacteria are a major modulator of host inflammatory responses in the
female genital tract. Immunity. 2015 May 5;42(5):965.

25.   Jimenez N, Norton T, Diadala G, Bell E, Valenti M, Farland LV, et al. Vaginal and rectal
microbiome contribute to genital inflammation in chronic pelvic pain. BMC Med. 2024
Jul 8;22(1):283.

26.   Foessleitner P, Petricevic L, Boerger I, Steiner I, Kiss H, Rieger A, et al. HIV infection as
a risk factor for vaginal dysbiosis, bacterial vaginosis, and candidosis in pregnancy: A
matched case-control study. Birth. 2021 Mar 1;48(1):139–46.

27. Gudza-Mugabe M, Havyarimana E, Jaumdally S, Garson KL, Lennard K, Tarupiwa A, et
al. Human Immunodeficiency Virus Infection Is Associated With Preterm Delivery
Independent of Vaginal Microbiota in Pregnant African Women. J Infect Dis. 2020 Apr
1;221(7):1194–203.

28. Short CES, Brown RG, Quinlan R, Lee YS, Smith A, Marchesi JR, et al. Lactobacillus-
Depleted Vaginal Microbiota in Pregnant Women Living With HIV-1 Infection Are
Associated With Increased Local Inflammation and Preterm Birth. Front Cell Infect
Microbiol [Internet]. 2021 Feb 11 [cited 2023 Apr 9];10. Available from:
https://pubmed.ncbi.nlm.nih.gov/33643930/

29. Babu H, Ambikan AT, Gabriel EE, Svensson Akusjärvi S, Palaniappan AN, Sundaraj V,
et al. Systemic Inflammation and the Increased Risk of Inflamm-Aging and Age-
Associated Diseases in People Living With HIV on Long Term Suppressive
Antiretroviral Therapy. Front Immunol [Internet]. 2019 Aug 27 [cited 2024 Oct 8];10.
Available from:
https://www.frontiersin.org/journals/immunology/articles/10.3389/fimmu.2019.0196
5/full

30. Guerville F, Vialemaringe M, Cognet C, Duffau P, Lazaro E, Cazanave C, et al.
Mechanisms of systemic low-grade inflammation in HIV patients on long-term
suppressive antiretroviral therapy: the inflammasome hypothesis. AIDS Lond Engl.
2023 Jun 1;37(7):1035–46.

31. Durand M, Sheehy O, Baril JG, Lelorier J, Tremblay CL. Association between HIV
infection, antiretroviral therapy, and risk of acute myocardial infarction: a cohort and
nested case-control study using Québec's public health insurance database. J Acquir
Immune Defic Syndr 1999. 2011 Jul 1;57(3):245–53.

32. Kuller LH, Tracy R, Belloso W, De Wit S, Drummond F, Lane HC, et al. Inflammatory
and coagulation biomarkers and mortality in patients with HIV infection. PLoS Med.
2008 Oct 21;5(10):e203.

33. Mehta SD, Nannini DR, Otieno F, Green SJ, Agingu W, Landay A, et al. Host Genetic
Factors Associated with Vaginal Microbiome Composition in Kenyan Women.
mSystems. 2020 Jul 28;5(4):e00502-20.

34. Bisanz JE, Enos MK, PrayGod G, Seney S, Macklaim JM, Chilton S, et al. Microbiota at
Multiple Body Sites during Pregnancy in a Rural Tanzanian Population and Effects of
Moringa-Supplemented Probiotic Yogurt. Appl Environ Microbiol. 2015
Aug;81(15):4965–75.

35. Baud A, Hillion KH, Plainvert C, Tessier V, Tazi A, Mandelbrot L, et al. Microbial
diversity in the vaginal microbiota and its link to pregnancy outcomes. Sci Rep. 2023
Jun 4;13(1):9061.

36. Goltsman DSA, Sun CL, Proctor DM, DiGiulio DB, Robaczewska A, Thomas BC, et al.
Metagenomic analysis with strain-level resolution reveals fine-scale variation in the
human pregnancy microbiome. Genome Res. 2018 Oct;28(10):1467–80.

37. Lee CY, Diegel J, France MT, Ravel J, Arnold KB. Evaluation of vaginal microbiome
equilibrium states identifies microbial parameters linked to resilience after menses
and antibiotic therapy. PLOS Comput Biol. 2023 Aug 1;19(8):e1011295.

38. France MT, Mendes-Soares H, Forney LJ. Genomic Comparisons of Lactobacillus
crispatus and Lactobacillus iners Reveal Potential Ecological Drivers of Community
Composition in the Vagina. Appl Environ Microbiol. 2016 Dec 12;82(24):7063.

39. Tachedjian G, Aldunate M, Bradshaw CS, Cone RA. The role of lactic acid production
by probiotic *Lactobacillus* species in vaginal health. Res Microbiol. 2017 Nov
1;168(9):782–92.

40. Witkin SS, Mendes-Soares H, Linhares IM, Jayaram A, Ledger WJ, Forney LJ. Influence
of Vaginal Bacteria and d- and l-Lactic Acid Isomers on Vaginal Extracellular Matrix
Metalloproteinase Inducer: Implications for Protection against Upper Genital Tract
Infections. mBio. 2013 Aug 6;4(4):e00460-13.

41. Macklaim JM, Gloor GB, Anukam KC, Cribby S, Reid G. At the crossroads of vaginal
health and disease, the genome sequence of Lactobacillus iners AB-1. Proc Natl Acad
Sci. 2011 Mar 15;108(supplement_1):4688–95.

42. Macklaim JM, Fernandes AD, Di Bella JM, Hammond JA, Reid G, Gloor GB.
Comparative meta-RNA-seq of the vaginal microbiota and differential expression by
Lactobacillus iners in health and dysbiosis. Microbiome. 2013 Apr 12;1(1):12.

43. Rampersaud R, Planet PJ, Randis TM, Kulkarni R, Aguilar JL, Lehrer RI, et al. Inerolysin,
a cholesterol-dependent cytolysin produced by Lactobacillus iners∇. J Bacteriol.
2011;193(5):1034–41.

44. Carter KA, Fischer MD, Petrova MI, Balkus JE. Epidemiologic Evidence on the Role of
Lactobacillus iners in Sexually Transmitted Infections and Bacterial Vaginosis: A
Series of Systematic Reviews and Meta-Analyses. Sex Transm Dis. 2023
Apr;50(4):224.

45. Li L, Zhang J, Zhou L, Shi H, Mai H, Su J, et al. The First Lanthipeptide from
Lactobacillus iners, Inecin L, Exerts High Antimicrobial Activity against Human Vaginal
Pathogens. Appl Environ Microbiol. 2023 Feb 27;89(3):e02123-22.

46. Vaneechoutte M, Guschin A, Van Simaey L, Gansemans Y, Van Nieuwerburgh F, Cools
P. Emended description of Gardnerella vaginalis and description of *Gardnerella*
leopoldii sp. nov., Gardnerella piotii sp. nov. and Gardnerella swidsinskii sp. nov., with

delineation of 13 genomic species within the genus Gardnerella. Int J Syst Evol
Microbiol. 2019 Mar 1;69(3):679–87.

47. Borgdorff H, Van Der Veer C, Van Houdt R, Alberts CJ, De Vries HJ, Bruisten SM, et al.
The association between ethnicity and vaginal microbiota composition in Amsterdam,
the Netherlands. PLoS ONE. 2017 Jul 1;12(7).

48. Serrano MG, Parikh HI, Brooks JP, Edwards DJ, Arodz TJ, Edupuganti L, et al.
Racioethnic diversity in the dynamics of the vaginal microbiome during pregnancy.
Nat Med 2019 256. 2019 May 29;25(6):1001–11.

49. Wei X, Tsai MS, Liang L, Jiang L, Hung CJ, Jelliffe-Pawlowski L, et al. Population genetic
analyses of longitudinal vaginal microbiome reveal racioethnic evolutionary dynamics
and prevailing positive selection of Lactobacillus adhesins [Internet]. bioRxiv; 2023
[cited 2024 Jul 30]. p. 2023.10.11.561685. Available from:
https://www.biorxiv.org/content/10.1101/2023.10.11.561685v1

50. Liao J, Shenhav L, Urban JA, Serrano M, Zhu B, Buck GA, et al. Microdiversity of the
vaginal microbiome is associated with preterm birth. Nat Commun 2023 141. 2023
Aug 17;14(1):1–14.

51. Grewal K, Lee YS, Smith A, Brosens JJ, Bourne T, Al-Memar M, et al. Chromosomally
normal miscarriage is associated with vaginal dysbiosis and local inflammation. BMC
Med. 2022 Jan 28;20(1):38.

52. Nelson DB, Hanlon AL, Wu G, Liu C, Fredricks DN. First Trimester Levels of BV-
Associated Bacteria and Risk of Miscarriage Among Women Early in Pregnancy.
Matern Child Health J. 2015 Dec 1;19(12):2682–7.

53. Seale AC, Blencowe H, Bianchi-Jassir F, Embleton N, Bassat Q, Ordi J, et al. Stillbirth
With Group B Streptococcus Disease Worldwide: Systematic Review and Meta-
analyses. Clin Infect Dis Off Publ Infect Dis Soc Am. 2017 Nov 6;65(suppl_2):S125–32.

54. Cherpes TL, Meyn LA, Krohn MA, Lurie JG, Hillier SL. Association between Acquisition
of Herpes Simplex Virus Type 2 in Women and Bacterial Vaginosis. Clin Infect Dis.
2003 Aug 1;37(3):319–25.

55. Borgdorff H, Tsivtsivadze E, Verhelst R, Marzorati M, Jurriaans S, Ndayisaba GF, et al.
Lactobacillus-dominated cervicovaginal microbiota associated with reduced HIV/STI
prevalence and genital HIV viral load in African women. ISME J 2014 89. 2014 Mar
6;8(9):1781–93.

56. Liu CM, Packman ZR, Abraham AG, Serwadda DM, Nalugoda F, Aziz M, et al. The
Effect of Antiretroviral Therapy Initiation on the Vaginal Microbiome in HIV-Infected
Women. Open Forum Infect Dis. 2019 Sep 1;6(9):ofz328.

57.   Mazibuko-Motau N, Sobia P, Xu J, Elsherbini JA, San JE, Lewis L, et al. Vaginal
832   microbial shifts are unaffected by oral pre-exposure prophylaxis in South African
833   women. Sci Rep. 2022 Sep 28;12(1):16187.

834   58.   Thurman AR, Ravel J, Gajer P, Marzinke MA, Ouattara LA, Jacot T, et al. Vaginal
835   Microbiota and Mucosal Pharmacokinetics of Tenofovir in Healthy Women Using a 90-
836   Day Tenofovir/Levonorgestrel Vaginal Ring. Front Cell Infect Microbiol [Internet]. 2022
837   Mar 8 [cited 2024 Jun 5];12. Available from:
838   https://www.frontiersin.org/articles/10.3389/fcimb.2022.799501

839   59.   Chávez-Torres M, Gómez-Palacio-Schjetnan M, Reyes-Terán G, Briceño O, Ávila-Ríos
840   S, Romero-Mora KA, et al. The vaginal microbiota of women living with HIV on
841   suppressive antiretroviral therapy and its relation to high-risk human papillomavirus
842   infection. BMC Microbiol. 2023 Jan 19;23(1):21.

843   60.   Donahue Carlson R, Sheth AN, Read TD, Frisch MB, Mehta CC, Martin A, et al. The
844   Female Genital Tract Microbiome Is Associated With Vaginal Antiretroviral Drug
845   Concentrations in Human Immunodeficiency Virus–Infected Women on Antiretroviral
846   Therapy. J Infect Dis. 2017 Nov 15;216(8):990–9.

847   61.   Cheu RK, Gustin AT, Lee C, Schifanella L, Miller CJ, Ha A, et al. Impact of vaginal
848   microbiome communities on HIV antiretroviral-based pre-exposure prophylaxis
849   (PrEP) drug metabolism. PLOS Pathog. 2020 Dec 3;16(12):e1009024.

850   62.   Klatt NR, Cheu R, Birse K, Zevin AS, Perner M, Noël-Romas L, et al. Vaginal bacteria
851   modify HIV tenofovir microbicide efficacy in African women. Science. 2017 Jun
852   2;356(6341):938–45.

853   63.   Taneva E, Sinclair S, Mesquita PMM, Weinrick B, Cameron SA, Cheshenko N, et al.
854   Vaginal microbiome modulates topical antiretroviral drug pharmacokinetics. JCI
855   Insight [Internet]. 2018 Jul 12 [cited 2024 Jun 5];3(13). Available from:
856   https://insight.jci.org/articles/view/99545

857   64.   Michel KG, Ma B, Weber K, McClellan L, Sheth A, Gange S, et al. 4117 UNIQUE
858   VAGINAL MICROBIOME POPULATIONS AND MICROBIAL GENE CONTENT AMONG
859   WOMEN WHO NATURALLY CONTROL HIV PROGRESSION. J Clin Transl Sci. 2020
860   Jun;4(s1):20–1.

861   65.   Dareng EO, Ma B, Adebamowo SN, Famooto A, Ravel J, Pharoah PP, et al. Vaginal
862   microbiota diversity and paucity of Lactobacillus species are associated with
863   persistent hrHPV infection in HIV negative but not in HIV positive women. Sci Rep.
864   2020 Nov 5;10(1):19095.

865   66.   Kiama TN, Vaneechoutte M, Mbugua PM, Verstraelen H, Estambale B, Temmerman M,
866   et al. Dynamics of the vaginal microbiome during the menstrual cycle of HIV positive

and negative women in a sub-urban population of Kenya. Afr J Biotechnol. 2023 Nov
30;22(11):291–304.

67.　Anton L, Ferguson B, Friedman ES, Gerson KD, Brown AG, Elovitz MA. Gardnerella
vaginalis alters cervicovaginal epithelial cell function through microbe-specific
immune responses. Microbiome. 2022 Dec 1;10(1):1–20.

68.　Joseph A, Anton L, Guan Y, Ferguson B, Mirro I, Meng N, et al. Extracellular vesicles
from vaginal Gardnerella vaginalis and Mobiluncus mulieris contain distinct
proteomic cargo and induce inflammatory pathways. Npj Biofilms Microbiomes. 2024
Mar 21;10(1):1–12.

69.　Heindel DW, Acosta DMF, Goff M, Yengo CK, Jan M, Liu X, et al. HIV-1 interaction with
an O-glycan-specific bacterial lectin enhances virus infectivity and resistance to
neutralizing antibodies. iScience [Internet]. 2024 Aug 16 [cited 2024 Aug 30];27(8).
Available from: https://www.cell.com/iscience/abstract/S2589-0042(24)01615-8

70.　Frenkel LM, Morrison RL, Fuller TL, Gouvêa MI, Benamor Teixeira M de L, Coombs RW,
et al. Brief Report: Vaginal Viral Shedding With Undetectable Plasma HIV Viral Load in
Pregnant Women Receiving 2 Different Antiretroviral Regimens: A Randomized
Clinical Trial. JAIDS J Acquir Immune Defic Syndr. 2021 Dec 1;88(4):361.

71.　Olesen R, Swanson MD, Kovarova M, Nochi T, Chateau M, Honeycutt JB, et al. ART
influences HIV persistence in the female reproductive tract and cervicovaginal
secretions. J Clin Invest. 2016 Mar 1;126(3):892–904.

72.　McClelland RS, Lingappa JR, Srinivasan S, Kinuthia J, John-Stewart GC, Jaoko W, et al.
Key Vaginal Bacteria Associated with Increased Risk of HIV Acquisition in African
Women: A Nested Case-Control Study. Lancet Infect Dis. 2018 May 1;18(5):554.

73.　Ravel J, Brotman RM, Gajer P, Ma B, Nandy M, Fadrosh DW, et al. Daily temporal
dynamics of vaginal microbiota before, during and after episodes of bacterial
vaginosis. Microbiome. 2013 Dec 2;1(1):29.

74.　Hotkani ZG, Ghaedmohammadi S, Mozdoori N. Meta-analysis of race and age
influence on the vaginal microbiome in pregnant and nonpregnant healthy women.
Httpsdoiorg102217fmb-2021-0209. 2022 Aug 11;17(14):1147–59.

75.　Odogwu NM, Olayemi OO, Omigbodun AO. Biogeographical effect on the diversity of
vaginal microbiome in preterm birth: A systematic review and meta-analysis. Med
Microecol. 2023 Jun 1;16:100080.

76.　Li KT, Li F, Jaspan H, Nyemba D, Myer L, Aldrovandi G, et al. Changes in the Vaginal
Microbiome During Pregnancy and the Postpartum Period in South African Women: a
Longitudinal Study. Reprod Sci. 2024 Jan 1;31(1):275–87.

77. Tian L, Wang XW, Wu AK, Fan Y, Friedman J, Dahlin A, et al. Deciphering functional
redundancy in the human microbiome. Nat Commun. 2020 Dec 4;11(1):6217.

78. Holm JB, France MT, Gajer P, Ma B, Brotman RM, Shardell M, et al. Integrating
compositional and functional content to describe vaginal microbiomes in health and
disease. Microbiome. 2023 Nov 30;11(1):259.

79. Douglas GM, Maffei VJ, Zaneveld JR, Yurgel SN, Brown JR, Taylor CM, et al. PICRUSt2
for prediction of metagenome functions. Nat Biotechnol. 2020 Jun;38(6):685–8.

80. KEGG PATHWAY Database [Internet]. [cited 2022 Oct 12]. Available from:
https://www.genome.jp/kegg/pathway.html

81. Ceccarani C, Foschi C, Parolin C, D'Antuono A, Gaspari V, Consolandi C, et al.
Diversity of vaginal microbiome and metabolome during genital infections. Sci Rep.
2019;9(1):1–12.

82. Amabebe E, Anumba DOC. Female Gut and Genital Tract Microbiota-Induced
Crosstalk and Differential Effects of Short-Chain Fatty Acids on Immune Sequelae.
Front Immunol. 2020 Sep 10;11:2184.

83. Vitali B, Cruciani F, Picone G, Parolin C, Donders G, Laghi L. Vaginal microbiome and
metabolome highlight specific signatures of bacterial vaginosis. Eur J Clin Microbiol
Infect Dis. 2015 Dec 1;34(12):2367–76.

84. Liu G, Barczak W, Lee LN, Shrestha A, Provine NM, Albayrak G, et al. The HDAC
inhibitor zabadinostat is a systemic regulator of adaptive immunity. Commun Biol.
2023 Jan 26;6(1):1–14.

85. Berard AR, Brubaker DK, Birse K, Lamont A, Mackelprang RD, Noël-Romas L, et al.
Vaginal epithelial dysfunction is mediated by the microbiome, metabolome, and
mTOR signaling. Cell Rep. 2023 May 30;42(5):112474.

86. Choi H, Rauh V, Garfinkel R, Tu Y, Perera FP. Prenatal Exposure to Airborne Polycyclic
Aromatic Hydrocarbons and Risk of Intrauterine Growth Restriction. Environ Health
Perspect. 2008 May;116(5):658–65.

87. Freije SL, Enquobahrie DA, Day DB, Loftus C, Szpiro AA, Karr CJ, et al. Prenatal
exposure to polycyclic aromatic hydrocarbons and gestational age at birth. Environ
Int. 2022 Jun;164:107246.

88. Bukavina L, Isali I, Ginwala R, Sindhani M, Calaway A, Magee D, et al. Global Meta-
analysis of Urine Microbiome: Colonization of Polycyclic Aromatic Hydrocarbon–
degrading Bacteria Among Bladder Cancer Patients. Eur Urol Oncol. 2023 Apr
1;6(2):190–203.

89. Huang L, Guo R, Li S, Wu X, Zhang Y, Guo S, et al. A multi-kingdom collection of
33,804 reference genomes for the human vaginal microbiome. Nat Microbiol. 2024
Jun 21;1–16.

90. Vivekanandan V, Khan ZH, Venugopal G, Musunuru B, Mishra P, Srivastava S, et al.
VagiBIOM Lactobacillus suppository improves vaginal health index in perimenopausal
women with bacterial vaginosis: a randomized control trial. Sci Rep. 2024 Feb
9;14(1):3317.

91. Ephraim-Emmanuel BC, Ordinioha B. Exposure and Public Health Effects of
Polycyclic Aromatic Hydrocarbon Compounds in Sub-Saharan Africa: A Systematic
Review. Int J Toxicol. 2021 May 1;40(3):250–69.

92. Munyeza C, Rohwer E, Forbes P. A review of monitoring of airborne polycyclic
aromatic hydrocarbons: An African perspective. Trends Environ Anal Chem. 2019 Jul
1;24:e00070.

93. Roachford OSE, Alleyne AT, Nelson KE. Insights into the vaginal microbiome in a
diverse group of women of African, Asian and European ancestries. PeerJ [Internet].
2022 Nov 29 [cited 2023 Apr 25];10. Available from: /pmc/articles/PMC9744153/

94. De Wolfe TJ, Arefin MR, Benezra A, Rebolleda Gómez M. Chasing Ghosts: Race,
Racism, and the Future of Microbiome Research. mSystems [Internet]. 2021 Oct 26
[cited 2023 Apr 13];6(5). Available from:
https://journals.asm.org/doi/10.1128/mSystems.00604-21

95. Greenhough B, Read CJ, Lorimer J, Lezaun J, McLeod C, Benezra A, et al. Setting the
agenda for social science research on the human microbiome. Palgrave Commun.
2020 Jan 31;6(1):1–11.

96. Srinivasan S, Morgan MT, Fiedler TL, Djukovic D, Hoffman NG, Raftery D, et al.
Metabolic Signatures of Bacterial Vaginosis. mBio. 2015 Apr 14;6(2):e00204-15.

97. Bergen NE, Jaddoe VWV, Timmermans S, Hofman A, Lindemans J, Russcher H, et al.
Homocysteine and folate concentrations in early pregnancy and the risk of adverse
pregnancy outcomes: the Generation R Study. BJOG Int J Obstet Gynaecol. 2012 May
1;119(6):739–51.

98. Feehily C, Crosby D, Walsh CJ, Lawton EM, Higgins S, McAuliffe FM, et al. Shotgun
sequencing of the vaginal microbiome reveals both a species and functional potential
signature of preterm birth. Npj Biofilms Microbiomes. 2020;6(1):1–9.

99. Ghartey J, Bastek JA, Brown AG, Anglim L, Elovitz MA. Women with preterm birth have
a distinct cervicovaginal metabolome. Am J Obstet Gynecol. 2015 Jun;212(6):776.e1-
776.e12.

100. Florova V, Romero R, Tarca AL, Galaz J, Motomura K, Ahmad MM, et al. Vaginal host
immune-microbiome interactions in a cohort of primarily African-American women
who ultimately underwent spontaneous preterm birth or delivered at term. Cytokine.
2021 Jan 1;137:155316.

101. Amabebe E, Chapman DR, Stern VL, Stafford G, Anumba DOC. Mid-gestational
changes in cervicovaginal fluid cytokine levels in asymptomatic pregnant women are
predictive markers of inflammation-associated spontaneous preterm birth. J Reprod
Immunol. 2018 Apr 1;126:1–10.

102. Kindinger LM, MacIntyre DA, Lee YS, Marchesi JR, Smith A, McDonald JAK, et al.
Relationship between vaginal microbial dysbiosis, inflammation, and pregnancy
outcomes in cervical cerclage. Sci Transl Med. 2016 Aug 3;8(350):350ra102-350ra102.

103. Eleuterio NM, Palei ACT, Rangel Machado JS, Tanus-Santos JE, Cavalli RC, Sandrim
VC. Positive correlations between circulating adiponectin and MMP2 in preeclampsia
pregnant. Pregnancy Hypertens. 2015 Apr;5(2):205–8.

104. Montagnana M, Lippi G, Albiero A, Scevarolli S, Salvagno GL, Franchi M, et al.
Evaluation of metalloproteinases 2 and 9 and their inhibitors in physiologic and pre-
eclamptic pregnancy. J Clin Lab Anal. 2009 Mar 13;23(2):88–92.

105. Yan Y, Fang L, Li Y, Yu Y, Li Y, Cheng JC, et al. Association of *MMP2* and *MMP9* gene
polymorphisms with the recurrent spontaneous abortion: A meta-analysis. Gene.
2021 Jan 30;767:145173.

106. Ulrich CC, Arinze V, Wandscheer CB, Copley Salem C, Nabati C, Etezadi-Amoli N, et
al. Matrix metalloproteinases 2 and 9 are elevated in human preterm laboring uterine
myometrium and exacerbate uterine contractility. Biol Reprod. 2019 Jun;100(6):1597–
604.

107. Petignat P, Laurini R, Goffin F, Bruchim I, Bischof P. Expression of matrix
metalloproteinase-2 and mutant p53 is increased in hydatidiform mole as compared
with normal placenta. Int J Gynecol Cancer Off J Int Gynecol Cancer Soc.
2006;16(4):1679–84.

108. Masho SW, Archer PW. Does Maternal Birth Outcome Differentially Influence the
Occurrence of Infant Death among African Americans and European Americans?
Matern Child Health J. 2011 Nov;15(8):1249–56.

109. Kim MJ, Romero R, Kim CJ, Tarca AL, Chhauy S, LaJeunesse C, et al. Villitis of
Unknown Etiology is Associated with a Distinct Pattern of Chemokine Up-regulation in
the Feto-maternal and Placental Compartments: Implications for Conjoint Maternal
Allograft Rejection and Maternal Anti-fetal Graft-versus-Host Disease. J Immunol
Baltim Md 1950. 2009 Mar 15;182(6):3919–27.

110. Nhan-Chang CL, Romero R, Kusanovic JP, Gotsch F, Edwin SS, Erez O, et al. A role for CXCL13 (BCA-1) in pregnancy and intra-amniotic infection/inflammation. J Matern-Fetal Neonatal Med Off J Eur Assoc Perinat Med Fed Asia Ocean Perinat Soc Int Soc Perinat Obstet. 2008 Nov;21(11):763–75.

111. Shivakoti R, Gupte N, Kumar NP, Kulkarni V, Balasubramanian U, Bhosale R, et al. Intestinal Barrier Dysfunction and Microbial Translocation in Human Immunodeficiency Virus-Infected Pregnant Women Are Associated With Preterm Birth. Clin Infect Dis Off Publ Infect Dis Soc Am. 2018 Sep 14;67(7):1103–9.

112. Hunt PW. Soluble CD163 and Clinical Outcomes in Treated HIV Infection: Insights Into Mechanisms. J Infect Dis. 2016 Oct 15;214(8):1132–3.

113. Knudsen TB, Ertner G, Petersen J, Møller HJ, Moestrup SK, Eugen-Olsen J, et al. Plasma Soluble CD163 Level Independently Predicts All-Cause Mortality in HIV-1–Infected Individuals. J Infect Dis. 2016 Oct 15;214(8):1198–204.

114. Schnittman SR, Byakwaga H, Boum Y, Kabakyenga J, Matthews LT, Burdo TH, et al. Changes in Immune Activation During Pregnancy and the Postpartum Period in Treated HIV Infection. Open Forum Infect Dis. 2021 Jun 1;8(6):ofab245.

115. Lin CH, Chen CC, Chiang HL, Liou JM, Chang CM, Lu TP, et al. Altered gut microbiota and inflammatory cytokine responses in patients with Parkinson's disease. J Neuroinflammation. 2019 Jun 27;16(1):129.

116. Schirmer M, Smeekens SP, Vlamakis H, Jaeger M, Oosting M, Franzosa EA, et al. Linking the Human Gut Microbiome to Inflammatory Cytokine Production Capacity. Cell. 2016 Nov 3;167(4):1125-1136.e8.

117. Wang LJ, Li SC, Li SW, Kuo HC, Lee SY, Huang LH, et al. Gut microbiota and plasma cytokine levels in patients with attention-deficit/hyperactivity disorder. Transl Psychiatry. 2022 Feb 23;12(1):1–9.

118. Adapen C, Réot L, Nunez N, Cannou C, Marlin R, Lemaître J, et al. Local Innate Markers and Vaginal Microbiota Composition Are Influenced by Hormonal Cycle Phases. Front Immunol. 2022 Mar 25;13:841723.

119. Yotebieng M, Mpody C, Ravelomanana NL, Tabala M, Malongo F, Kawende B, et al. HIV viral suppression among pregnant and breastfeeding women in routine care in the Kinshasa province: a baseline evaluation of participants in CQI-PMTCT study. J Int AIDS Soc. 2019 Sep;22(9):e25376.

120. Yotebieng M, Behets F, Kawende B, Ravelomanana NLR, Tabala M, Okitolonda EW. Continuous quality improvement interventions to improve long-term outcomes of antiretroviral therapy in women who initiated therapy during pregnancy or

breastfeeding in the Democratic Republic of Congo: design of an open-label, parallel,
group randomized trial. BMC Health Serv Res. 2017 Apr 26;17:306.

121. Zotova N, Familiar I, Kawende B, Kasindi FL, Ravelomanana N, Parcesepe AM, et al.
HIV disclosure and depressive symptoms among pregnant women living with HIV: a
cross-sectional study in the Democratic Republic of Congo. J Int AIDS Soc.
2022;25(2):e25865.

122. Severgnini M, Morselli S, Camboni T, Ceccarani C, Laghi L, Zagonari S, et al. A Deep
Look at the Vaginal Environment During Pregnancy and Puerperium. Front Cell Infect
Microbiol [Internet]. 2022 May 17 [cited 2024 Dec 18];12. Available from:
https://www.frontiersin.org/journals/cellular-and-infection-
microbiology/articles/10.3389/fcimb.2022.838405/full

123. Kervinen K, Holster T, Saqib S, Virtanen S, Stefanovic V, Rahkonen L, et al. Parity and
gestational age are associated with vaginal microbiota composition in term and late
term pregnancies. eBioMedicine. 2022 Jun 24;81:104107.

124. Livson S, Virtanen S, Lokki AI, Holster T, Rahkonen L, Kalliala I, et al. Cervicovaginal
Complement Activation and Microbiota During Pregnancy and in Parturition. Front
Immunol. 2022 Jul 25;13:925630.

125. Ho M, Moon D, Pires-Alves M, Thornton PD, McFarlin BL, Wilson BA. Recovery of
microbial community profile information hidden in chimeric sequence reads. Comput
Struct Biotechnol J. 2021 Sep 3;19:5126–39.

126. Movassagh M, Bebell LM, Burgoine K, Hehnly C, Zhang L, Moran K, et al. Vaginal
microbiome topic modeling of laboring Ugandan women with and without fever. Npj
Biofilms Microbiomes. 2021 Sep 10;7(1):1–10.

127. Graspeuntner S, Loeper N, Künzel S, Baines JF, Rupp J. Selection of validated
hypervariable regions is crucial in 16S-based microbiota studies of the female genital
tract. Sci Rep 2018 81. 2018 Jun 26;8(1):1–7.

128. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-
resolution sample inference from Illumina amplicon data. Nat Methods. 2016;13(7).

129. Andrews S. FastQC: A quality control analysis tool for high throughput sequencing
data [Internet]. 2010 [cited 2022 Oct 12]. Available from:
https://www.bioinformatics.babraham.ac.uk/projects/fastqc/

130. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence
data. Bioinformatics. 2014 Aug 1;30(15):2114–20.

131. McDonald D, Jiang Y, Balaban M, Cantrell K, Zhu Q, Gonzalez A, et al. Greengenes2
unifies microbial data in a single reference tree. Nat Biotechnol 2023. 2023 Jul
27;14:1–4.

132. Aitchison J, Egozcue JJ. Compositional Data Analysis: Where Are We and Where
Should We Be Heading? Math Geol. 2005 Oct 1;37.

133. Gloor GB, Macklaim JM, Pawlowsky-Glahn V, Egozcue JJ. Microbiome datasets are
compositional: And this is not optional. Front Microbiol. 2017 Nov 15;8(NOV):294209.

134. Cao Q, Sun X, Rajesh K, Chalasani N, Gelow K, Katz B, et al. Effects of Rare
Microbiome Taxa Filtering on Statistical Analysis. Front Microbiol. 2021 Jan
12;11:607325.

135. Wang Y, Lê Cao KA. PLSDA-batch: a multivariate framework to correct for batch
effects in microbiome data. Brief Bioinform. 2023 Mar 1;24(2):bbac622.

136. Bastiaanssen TFS, Quinn TP, Loughman A. Bugs as Features (Part II): A Perspective on
Enriching Microbiome-Gut-Brain Axis Analyses. Nat Ment Health. 2023 Dec
5;1(12):930–8.

137. Lubbe S, Filzmoser P, Templ M. Comparison of zero replacement strategies for
compositional data with large numbers of zeros. Chemom Intell Lab Syst. 2021 Mar
15;210:104248.

138. Tenenbaum D, Maintainer. Bioconductor. 2024 [cited 2024 Oct 12]. KEGGREST:
Client-side REST access to the Kyoto Encyclopedia of Genes and Genomes (KEGG). R
package version 1.44.1. Available from:
http://bioconductor.org/packages/KEGGREST/

139. France MT, Ma B, Gajer P, Brown S, Humphrys MS, Holm JB, et al. VALENCIA: a
nearest centroid classification method for vaginal microbial communities based on
composition. Microbiome. 2020 Dec 1;8(1):1–15.

140. Ch C, A C. Estimating and comparing microbial diversity in the presence of
sequencing errors. PeerJ [Internet]. 2016 Feb 1 [cited 2024 Sep 27];4. Available from:
https://pubmed.ncbi.nlm.nih.gov/26855872/

141. Chao A, Henderson PA, Chiu CH, Moyes F, Hu KH, Dornelas M, et al. Measuring
temporal change in alpha diversity: A framework integrating taxonomic, phylogenetic
and functional diversity and the iNEXT.3D standardization. Methods Ecol Evol.
2021;12(10):1926–40.

142. Bender R, Lange S. Adjusting for multiple testing--when and how? J Clin Epidemiol.
2001 Apr;54(4):343–9.

143. Lu Y, Zhou G, Ewald J, Pang Z, Shiri T, Xia J. MicrobiomeAnalyst 2.0: comprehensive
statistical, functional and integrative analysis of microbiome data. Nucleic Acids Res.
2023 Jul 5;51(W1):W310–8.

144. Zhou H, He K, Chen J, Zhang X. LinDA: linear models for differential abundance
analysis of microbiome compositional data. Genome Biol. 2022 Apr 14;23(1):95.

145. Lin H, Peddada SD. Analysis of compositions of microbiomes with bias correction.
Nat Commun. 2020 Jul 14;11(1):3514.

146. Fernandes AD, Macklaim JM, Linn TG, Reid G, Gloor GB. ANOVA-Like Differential
Expression (ALDEx) Analysis for Mixed Population RNA-Seq. PLOS ONE. 2013 Jul
2;8(7):e67019.

147. Mirzayi C, Renson A, Zohra F, Elsafoury S, Geistlinger L, Kasselman LJ, et al. Reporting
guidelines for human microbiome research: the STORMS checklist. Nat Med. 2021
Nov;27(11):1885–92.

1122