Cell cycle dynamics regulate H3K27 and H3K9 histone modifications

Liyne Nogay [1, 2], Ananthakrishnan Vijayakumar Maya § [1, 2], Lara Heckmann § [1, 2], Francesco Cardamone [1, 2, 3], Isabelle Grass [1, 4], Aakriti Singh [1, 2], Anna Frey [1, 5], Laurin Ernst [1, 2], Nicola Iovino [3, 4], Anne-Kathrin Classen [1, 4], *

§ shared co-authors
* corresponding author: anne.classen@biologie.uni-freiburg.de

1 Faculty of Biology, University of Freiburg, Freiburg, Germany
2 International Max Planck Research School for Epigenetics, Biophysics and Metabolism, Freiburg, Germany
3 Max Planck Institute of Immunobiology and Epigenetics, Freiburg, Germany
4 CIBSS Centre for Integrative Biological Signalling Studies, University of Freiburg, Freiburg, Germany
5 Spemann Graduate School of Biology and Medicine (SGBM), University of Freiburg Germany

ABSTRACT

Cell cycle progression presents a fundamental challenge to genome integrity, particularly due to the need to reestablish post-translational histone modifications (PTMs) following DNA replication. Although proliferative and differentiating tissues exhibit markedly different cell cycle dynamics, how these differences shape the histone modification landscape in vivo remains largely unexplored. Here, we show that levels of H3K27ac, H3K27me3, and H3K9me3 are tightly linked to cell cycle dynamics in the *Drosophila* wing imaginal disc. We demonstrate that both physiological and pathological elongation of the cell cycle led to an accumulation of H3K9me3 and H3K27me3, whereas cell cycle acceleration reduces their levels. In contrast, H3K27ac exhibits the opposite pattern: levels decrease in arrested cells and increase with faster cycling. Genome-wide CUT&Tag analysis reveals that these changes predominantly affect genomic loci already modified in normally proliferating tissue. Importantly, the regulation of methylation levels at H3K9 and H3K27 is not solely mediated by the cell cycle machinery but reflects a metabolically guided process in which the rate of methylation is coupled to the rate of cell proliferation through metabolic activity, including signaling via the Insulin/PI3K/Akt pathway. Our study thus reveals key principles for understanding histone methylation in proliferating, senescent, and differentiating cells. In contrast, H3K27 acetylation is regulated through a distinct, cell cycle-coupled mechanism. We find that CBP/Nejire-mediated acetylation of H3K27 peaks during S-phase and is reversed by HDAC1, as cells exit replication. Disruption of this acetylation cycle leads to replication stress and a G2 cell cycle arrest via a DNA damage checkpoint. Notably, this genome-protective function of CBP/Nejire depends specifically on acetylation of the H3K27 residue itself, revealing a novel role for H3K27ac beyond its well-established function in transcriptional activation. Together, our findings establish a robust link between cell cycle progression and histone modification dynamics, highlighting the necessity of maintaining balanced PTM levels under varying proliferative states. These insights have broad implications for our understanding of development, aging, and tumor growth.

  INTRODUCTION

  The cell cycle and the regulation of chromatin need to be tightly coordinated preserve epigenetic

  integrity. Specifically, progression through S-phase poses several challenges: chromatin must become

  transiently accessible for DNA replication, pre-existing post-translationally modified histones must be

  evenly distributed between daughter genomes and newly incorporated histones must acquire

  appropriate modifications to compensate for the semi-conservative dilution of modifications. These

  demands are complicated by the fact that cell cycle progression and dynamics differ drastically across

  biological contexts, ranging from rapid divisions in embryonic and regenerating tissues to complete

  arrest in terminally differentiated cells. How chromatin states are maintained across diverse proliferative

  tissue environments *in vivo* is still insufficiently understood.

  In actively proliferating tissues, entry into the cell cycle at the G1/S transition, and progression through

  subsequent cycle phases, are tightly regulated. This regulation is mediated by Cyclins and Cyclin-

  dependent kinases (CDKs), as well as retinoblastoma (Rb) proteins and E2F transcription factors.

  Throughout the cell cycle, specific checkpoints respond to growth factors or cell size to modulate the

  rate of proliferation, as well as to DNA damage or chromosome misalignment to maintain the integrity

  of the genome (Matthews et al., 2022, Tapon et al., 2001, Lee and Orr-Weaver, 2003). As cells enter

  stages of differentiation, quiescence or senescence, they typically withdraw from active cycling by either

  prolonging their G1 phase or entering a stable cell cycle arrest known as G0. This transition away from

  proliferation often involves cyclin-dependent kinase (CDK) inhibitors, such as p21 and p27, which are

  intricately linked to gene expression programs that guide cell fate determination and differentiation

  (Buttitta and Edgar, 2007, Huang et al., 2022, de Morree and Rando, 2023, Ma et al., 2015). Progression

  through the cell cycle presents significant challenges for chromatin organization and epigenetic

  regulation - especially during S-phase but also during mitosis. Specifically, during S-phase, chromatin

  must become highly accessible to facilitate replication fork progression, nucleosome redistribution and

  de novo nucleosome incorporation in the replicated genomes (Stewart-Morgan et al., 2020, Alabert et

  al., 2015, Reveron-Gomez et al., 2018, Ma et al., 2015, Escobar et al., 2021, Ramos-Alonso and

  Chymkowitch, 2024). In contrast, in mitosis, chromatin must be tightly compacted to ensure accurate

  chromosome segregation (Ramos-Alonso and Chymkowitch, 2024, Zhou and Heald, 2020). Central to

  the regulation of chromatin organization are post-translational histone modifications (PTMs), which are

  also epigenetic regulators of transcriptional silencing and gene activation (Millan-Zambrano et al., 2022,

  Bhatnagar et al., 2023, Lopez-Hernandez et al., 2025). It is well-established that PTM dysregulation

  can lead to cell cycle defects by disrupting chromatin organization and entire gene regulatory networks

  (Brookes and Shi, 2014, Cavalli and Heard, 2019, Feinberg et al., 2016, Peterson and Laniel, 2004).

  While much attention has been dedicated to understanding how PTM dysregulation can drive

  pathogenesis by altering the expression of genes central to cell cycle regulation, less is known about

  how cell cycle dynamics connect to the maintenance and re-establishment of histone modifications.

  Yet, studies of the cell cycle-dependent regulation of the histone methyltransferase PR-Set7 and

histone demethylase PHF8 underscore the capacity of the cell cycle machinery to control specific
histone modifications and thereby maintain chromatin function (Oda et al., 2010, Liu et al., 2010).

Some of the most abundant PTMs that regulate chromatin accessibility and transcriptional activity are
acetylation and methylation of lysine residues on histone tails. While several lysine residues in histones
can be acetylated or methylated, research has most strongly focused on dynamic acetylation and
methylation of lysines 9 and 27 on histone H3 (H3K9 and H3K27), which represent some of the
functionally most important histone modifications. In most species, trimethylation of H3K27 is mediated
by Polycomb Repressive Complexes and relies on histone methyltransferases of the Enhancer of zeste
E(z) family. H3K27me3 maintains the silencing of many genes required for cell fate specification during
development, most famously Hox genes (Blackledge and Klose, 2021, Sparmann and van Lohuizen,
2006, Muller et al., 1995, Beuchle et al., 2001, Cao et al., 2002). In contrast, trimethylation of H3K9
(H3K9me3) is mediated by the enzymatic activity of Su(var)3-9 and is central to the formation and
maintenance of constitutive heterochromatin at telomeres, centromeres and repetitive repeats
(Padeken et al., 2022, Aagaard et al., 1999, Ebert et al., 2004). In *Drosophila*, H3K27 acetylation is
mediated by the histone acetyltransferase (HAT) Nejire (Nej), a homolog of mammalian p300/CBP, and
is reversed by the histone deacetylase HDAC1/Rpd3 (Willnow and Teleman, 2024, Petruk et al., 2001,
Tie et al., 2009, Ciabrelli et al., 2023). H3K27ac is strongly associated with active promoters and
enhancers, where it is thought to facilitate chromatin opening and transcriptional activation (Holmqvist
and Mannervik, 2013, Philip et al., 2015, Boija et al., 2017, McConnell et al., 2012, Chen et al., 2022).
As a consequence of these important functions, dysregulation of H3K9me3, H3K27me3 and H3K27ac
is a common feature in cancer or other diseases characterized by alterations and defects in cell
proliferation. Yet, importantly, it often remains unclear if defects in epigenetic modifications cause
altered proliferation defects or if vice versa proliferation defects cause defects in epigenetic
modifications (Brookes and Shi, 2014, Cavalli and Heard, 2019, Feinberg et al., 2016, Classen et al.,
2009). Disentangling this relationship is key to understanding how epigenetic and proliferative states
influence one another in development and disease.

Importantly, the semi-conservative nature of DNA replication during S-phase has a profound impact on
the landscape of these histone modifications. As the DNA is duplicated, pre-existing histones bearing
PTMs are equally distributed onto the newly synthesized strands. Naïve, newly synthesized histones
are integrated to ensure proper packaging of the duplicated genomes but also necessitating the
reestablishment of the now diluted PTM code (Escobar et al., 2021, Alabert and Groth, 2012, Escobar
et al., 2019, Wenger et al., 2023). To reestablish the epigenetic histone modification landscape after
incorporation of newly synthesized, naive histones, histone-modifying enzymes act post-replication by
using the modifications on old recycled histones as a template to modify the newly incorporated histones
(Stewart-Morgan et al., 2020, Annunziato, 2015, Flury and Groth, 2024). Importantly, in cultured cells,
histone acetylation marks are typically restored immediately after replication often via a transcription-
dependent process, while histone methylation is generally re-established during subsequent gap
phases (Alabert et al., 2015, Zee et al., 2012, Reveron-Gomez et al., 2018, Petruk et al., 2013).

Of note, strong acetylation of newly synthesized histones plays a specific cell-cycle-dependent role

during S-phase. Hyperacetylation, for example at H4K5, H4K12, H3K14, H3K23 or H3K56, is observed

during S-phase and facilitates recognition of newly synthesized histones by histone chaperone

complexes, promoting their correct incorporation into newly synthesized chromatin. Hyperacetylation of

histones presumably occurs in the cytoplasm and may be mediated by B-type acetyltransferases like

Hat1 (Li et al., 2008, Masumoto et al., 2005, Tessarz and Kouzarides, 2014, Benson et al., 2006, Falbo

and Shen, 2009, Jackson et al., 1976, Lande-Diner et al., 2009, Ruiz-Carrillo et al., 1975, Annunziato

and Hansen, 2000, Sobel et al., 1995, Verreault et al., 1998).

Despite these insights, questions remain about how cell cycle progression influences histone

modification dynamics, especially in *in vivo* settings. For instance, to what extent are histone

modifications regulated in a cell cycle-dependent manner in developing tissues? How does the

maintenance of these modifications differ between rapidly proliferating cells and cells in quiescent,

senescent, or post-mitotic states? Moreover, how is the activity of histone-modifying enzymes

coordinated with cell cycle checkpoints? To begin to address these questions we used the developing

Drosophila wing imaginal disc to characterize cell cycle dependent histone modifications and reveal

pronounced dynamics of H3K9 and H3K27 modifications in a developing tissue.

RESULTS

(1) Post-translational modifications of H3K27 and H3K9 are linked to cell cycle dynamics

To understand how cell cycle dynamics may affect chromatin dynamics, we aimed to identify post-

translational histone modifications that are sensitive to changes in cell cycle progression. Using the *rn-*

GAL4 driver **(Fig S1.1A)**, under the control of a temperature-sensitive GAL80ts, we genetically

manipulated cell cycle progression in the pouch of developing wing imaginal discs for 24 hours by

accelerating or decelerating cell cycle dynamics. Within this timeframe of 24 h, we expect to disturb at

least one cell cycle, which lasts between 12 and 18 h in third instar wing imaginal discs (Neufeld et al.,

1998, Milan et al., 1996, Prober and Edgar, 2000, Crucianelli et al., 2022). First, we induced a cell cycle

arrest in late G2 by expressing *Cdk1-RNAi (Cosolo et al., 2019)*. Second, ectopic expression of the

ERK-responsive ETS transcription factor Pointed-P1 (PntP1) causes cells to arrest in either G1 or in

G2 (Ito and Igaki, 2021). We confirmed the arrest in our genotypes by observing the cell cycle reporter

FUCCI and the absence of EdU incorporation, a nucleotide analogue visualizing DNA replication, in

nuclei of the *rn-GAL4* expression domain **(Fig S1.1B-L)** (Zielke et al., 2014). Conversely, we

accelerated the cell cycle by co-expressing of dE2F1 and dDP (Buttitta et al., 2007, Neufeld et al.,

1998). This did not alter the proportion of cells in G1 or G2 but resulted in elevated EdU incorporation,

reflecting the accelerated S-phase dynamics - both in frequency and speed (Crucianelli et al., 2022) -

of this genotype **(Fig S1.1M-Q)**.

As post-translational histone modifications are numerous, we focused on prominent modifications
mediated by well-known histone-modifying complexes. We analyzed modifications associated with
gene regulation, such as H3K27ac (mediated by the p300/CBP nejire), H3K9ac (mediated by SAGA-
HAT complex), H3K4me3 (mediated by the Trithorax (Trx)) and H3K27me3 (mediated by the Polycomb
protein E(z)), or modifications associated with genome maintenance, such as H3K9me3 (mediated by
Su(var)3-9), as well as examples of less well characterized modifications, such as H3K18ac and
H4K8ac (mediated by p300/CBP, Gcn5, SAGA-HAT), H3K18 crotonylation (mediated by the p300/CBP
nejire), and pan-H3K9 methylation or total lysine acetylation. We performed immunofluorescence
staining for these modifications in wing imaginal discs with the induced cell cycle alterations described
above.
177
Many of the histone modifications analyzed did not change upon manipulating cell cycle dynamics. For
example, H3K18ac and H3K4me3 levels were not sensitive to cell cycle modulation **(Fig 1A-D, I-L, U-**
W, Fig S1.3A), and neither were H4K8ac, H3K9ac, H3K18 crotonylation, total H3K9 methylation nor
levels of total lysine acetylation **(Fig S1.2, Fig S1.3B)**. Yet, strikingly, we found that H3K27ac levels
were strongly reduced in cell cycle arrested genotypes, whereas levels of this modification were
elevated when the cell cycle was accelerated **(Fig 1E-H, U-W)**. In contrast, H3K9me3 and H3K27me3
levels were elevated in arrested cells, whereas levels of both modifications were reduced when the cell
cycle was accelerated **(Fig 1M-W)**. Importantly, alterations observed in cell cycle arrested cells were
independent of which cell cycle phase the arrest occurred in. Specifically, H3K27ac reduction and
H3K9me3 and H3K27me3 elevation was observed in both G1 and G2 arrested cell populations upon
expression of *pntP1* **(Fig S1.1I, Fig 1G,O,S)**, suggesting that the elongation of either gap phase can
produce the observed effects.
190
Taken together, we conclude that cell cycle dynamics have a surprising impact on histone modification
in a proliferating tissue. First, only three tested modifications responded significantly to cell cycle
dynamics in our assays, suggesting that only a subset of modifications is strongly linked to the cell cycle
machinery. Second, these changes were not dependent to cells arresting in G1 or in G2, suggesting
that overall cell cycle length is a defining factor rather than the gap phase position within a cell cycle.
Third, if levels of histone modifications were primarily determined by the kinetics of restoring histone
modifications after semi-conservative replication and the resulting histone PTM dilution, one may
predict that modifications simply accumulate as the cell cycle lengthens. However, the reduction of
H3K27ac in arrested cells defies a scenario in which continuous HAT activity restores H3K27ac levels
after DNA replication. We thus wanted to better understand the relationship between histone
methylation, histone acetylation and cell cycle dynamics.
202
203　**(2) Cell cycle dependent changes in H3K27ac, H3K27me3 or H3K9me3 levels occur at**
204　**preexisting target loci**
205

To first identify the genomic regions affected by the cell-cycle-dependent changes in bulk levels of
H3K27ac, H3K27me3, and H3K9me3, we performed CUT&Tag analysis on wing imaginal discs
expressing *Cdk1-RNAi* under the control of *rn-GAL4* for 24 hours. Although *Cdk1-RNAi* is expressed in
only a subset of cells within the wing disc (see **Fig S1.1A**) - limiting the magnitude of detectable
differences - we consistently observed a decrease in H3K27ac levels, and an increase in levels of both
H3K27me3 and H3K9me3 modifications. These changes occurred at genomic regions already carrying
these modifications in wild type conditions, as evident from examples of genomic snapshots **(Fig 2 A,B)**
and changes in average levels of modification on and around peaks identified in both wild type and
Cdk1-RNAi-expressing discs **(Fig 2 C-F, Fig S2.1A, E)**. These findings are consistent with prior
observations in mESCs, where H3K27me3 levels increase at pre-existing sites during prolonged G1
phases (Trouth et al., 2025).
217
Of note, many H3K27ac peaks identified only in wild type samples exhibited a significant reduction in
H3K27ac levels in *Cdk1-RNAi*-expressing discs, suggesting that H3K27ac was indeed lost from pre-
existing acetylated regions **(Fig S2.1B,E)**. In agreement with this conclusion, genomic sequences not
identified as peaks by our algorithm in any sample showed little or no change in H3K27ac levels **(Fig
S2.1D, E)**. In contrast, although increases in H3K27me3 and H3K9me3 levels in *Cdk1-RNAi* discs
occurred at many pre-existing methylated sites **(Fig S2.1A, E)**, a substantial number of H3K27me3 and
H3K9me3 peaks identified only in *Cdk1-RNAi* discs, as well as sequences in other genomic regions,
also showed significantly elevated methylation compared to wild-type controls **(Fig S2.1C,D,E)**. This
indicates that the accumulation of H3K27me3 and H3K9me3 in *Cdk1-RNAi* discs may extend beyond
canonical methylation target sites identified in wild type discs. We conclude that cell-cycle-dependent
changes to bulk levels of H3K27ac, H3K27me3 or H3K9me3 predominantly affect existing target sites;
however, the increase in methylation may also affect other genomic regions. Importantly, the magnitude
of changes compared to levels of methylation in control discs is similar for all genomic regions,
suggesting that changes to H3K27me3 or H3K9me3 levels occur at the same rate across the genome.
232
Our conclusions are further supported by our observations that the histone-modifying enzymes
responsible for H3K27 acetylation and H3K27methylation do not compete for the same histone residues
in vivo. Specifically, when we performed a knock-down of the H3K27 methyltransferase E(z), we
observed an expected loss of H3K27me3 **(Fig S2.2 A,B)**, but importantly, no corresponding increase
in H3K27ac **(Fig S2.2 C,D)**. Conversely, when we performed a knock-down of the H3K27
acetyltransferase CBP/Nej (Nejire), we observed an expected loss of H3K27ac **(Fig S2.2 E,F)**, but no
corresponding increase in H3K27me3 **(Fig S2.2 G,H)**. We conclude that, even though studies propose
local antagonism of H3K27me3 and H3K27ac at the same residues (Tie et al., 2016, Philip et al., 2015,
Tie et al., 2009), the cell-cycle-length-dependent decrease of H3K27ac and increase of H3K27me3 bulk
levels upon expression of *Cdk1-RNAi* are not driven by competition for the same H3K27 residues but
instead reflect their natural occurrence of these distinct PTMs in distinct genomic contexts.
244
(3) H3K27 or H3K9 modifications are linked to cell cycle dynamics during development

If bulk histone modifications in wing discs are affected by cell cycle progression, can we provide
evidence that this also occurs in a physiologically relevant state? During larval development, cells in
the wing imaginal disc actively proliferate. However, the future wing margin forms a zone of non-
proliferating cells (ZNC), where cells arrest in either G1 or G2 **(Fig S3.1 A-C)** (Johnston and Edgar,
1998). In addition to the ZNC, two clusters of G2-arrested cells with senescent markers can be found
surrounding the wing pouch **(Fig S3.1 D)** (Zang et al., 2025). Strikingly, arrested cells in the ZNC **(Fig**
3 A-C) and senescent clusters **(Fig 3 D-F)** show a decrease in H3K27ac levels and an increase in
H3K9me3 and H3K27me3, consistent with our previous conclusions upon cell cycle elongation by
expression of *Cdk1-RNAi* or *UAS-pntP1.* Importantly, none of the other histone modifications
recapitulated this pattern, again reinforcing the specific association between H3K9me3, H3K27me3 and
H3K27ac levels and the dynamics of cell cycle progression **(Fig S3.1 E-H)**.

To begin to understand if there are consequences for cell-cycle-dependent PTM modulation during
development, we analyzed, as an example of three PTMs, H3K27me3-dependent regulation of cell
fates. H3K27me3 mediates Polycomb-dependent silencing of genes typically required for cell fate
specification (Blackledge and Klose, 2021, Sparmann and van Lohuizen, 2006, Beuchle et al., 2001).
In imaginal disc with either arrested (*Cdk1-RNAi*) or accelerated (*Dp,E2F1*) cell cycle dynamics, the
expression of Polycomb targeted genes required for wing disc differentiation, such as Antp, Ubx/AbdA,
En/inv, Nub, Wg, Ptc and Cut was not altered **(Fig S3.2)**. These observations demonstrate that reduced
or elevated H3K27me3 levels observed upon changing cell cycle dynamics still ensure robust Polycomb
target gene outputs, suggesting that a H3K27me3 levels buffer gene silencing against a range of cell
cycle dynamics.

Surprisingly, our subsequent experiments revealed that the cell cycle-dependent regulation of
H3K9me3, H3K27me3, and H3K27ac arises through mechanistically distinct pathways. Specifically,
methylation and acetylation marks exhibit fundamentally different dynamics. Given these divergent
mechanisms, we present our findings in two consecutive sections: first, we detail how H3K9me3 and
H3K27me3 levels are coupled to the cell cycle via growth signaling pathways **(Fig. 4)**, followed by an
independent analysis of the function of H3K27ac in preventing replication stress during S-phase **(Figs.**
5 and 6).

(4) H3K27me3 or H3K9me3 are coupled to cell cycle dynamics via metabolic signaling

Previous studies suggest that H3K27me3 and H3K9me3 modifications after DNA-replication are
reestablished after DNA replication through slow and continuous methyltransferase activity in
subsequent gap phases (Alabert et al., 2015, Zee et al., 2012, Reveron-Gomez et al., 2018). Our
findings are consistent with these observations: in all our imaginal disc models, levels of H3K9me3 and
H3K27me3 increase in conditions with elongated cell cycles (and thus infrequent S-phase events,
reducing replication-associated dilution of these marks), and decrease in conditions with shortened cell

cycles (increasing the frequency of S-phase and thereby enhancing dilution). These results suggest

that H3K9 and H3K27 methylation is driven by a relatively constant rate of histone methyltransferase

activity, which appears not to be affected by genetic alterations of the cell cycle machinery. If the rate

of histone methylation can indeed be genetically separated from the frequency of cell cycling, what

mechanisms then govern the restoration of histone methylation to the appropriate levels, particularly for

H3K9me3, which is essential for heterochromatin architecture, and H3K27me3, which is critical for gene

silencing? To address this question, we decided to manipulate cell cycle progression upstream of direct

regulators of the cell cycle by activating a proliferative pathway with important metabolic functions.

Specifically, we expressed a constitutively active Insulin receptor (InR^{DA}) under the control of *rn-GAL4*

for 24 h, which promotes PI3K/Akt signaling. The resulting metabolic changes can be visualized by

higher rates of protein synthesis reflecting higher biosynthetic capacity, accelerated cell proliferation

visualized by high EdU incorporation and ultimately, larger wing pouch size **(Fig 4A-C, Fig S4 A-E)**. Of

note, expression of InR^{DA} did not alter the FUCCI cell cycle profile, demonstrating that the higher

metabolic activity shortens the overall cell cycle length, rather than affecting specific cell cycle phases

(Fig S4F,G).

Importantly, we found that expression of InR^{DA} alone did not give rise to changes in H3K27 or H3K9

methylation levels, despite the accelerated cell divisions **(Fig 4 D,E,H, I)**. This demonstrates that the

metabolic potential can tune the rate of H3K27 and H3K9 methylation, to keep levels of H3K27 or H3K9

methylation comparable to wild type levels, despite accelerated cell cycle progression. To now

functionally separate metabolic potential from cell cycle progression, we also arrested the cell cycle by

co-expressing the constitutively active Insulin receptor with a *Cdk1-RNAi*. As expected from our

previous experiments, levels of H3K27me3 and H3K9me3 increased, but most strikingly, levels of

H3K27me3 and H3K9me3 were even higher than in discs expressing *Cdk-1-RNAi* alone **(Fig 4F,G,J,K)**.

These results demonstrate that rates of H3K27 or H3K9 methylation are intimately linked to the

metabolic potential of a cell - with the purpose to match the rate of H3K27 or H3K9me3 methylation

with the speed of cell cycle progression and thus the frequency of S-phase linked histone dilution **(Fig**

4 P,Q). Of note, levels of H3K4me3 were not affected by these manipulations **(Fig 4 L-O)**, indicating

that a modification maintaining steady levels of bulk transcriptional activity is not - and even should not

be - linked to metabolic rates or the rate of cell cycle progression.

We wanted to better understand the functional implications of our findings. The link between levels of

H3K9me3 and cell cycle progression becomes relevant in physiological and pathological setting of

senescence, where cytotoxic stress or oncogenic mutations provide metabolic activation that is

juxtaposed to a senescent cell cycle arrest program. Accordingly, in senescent G2-arrested cells at the

center of inflammatory tissue damage in wing imaginal discs induced by ectopic expression of the TNFα

homologue eiger (egr) **(Fig S4.2 A-F)** (Cosolo et al., 2019, Floc'hlay et al., 2023, Jaiswal et al., 2023,

Smith-Bolton et al., 2009), we observed high levels of H3K9 methylation **(Fig S4.2G-I, purple)**.

Importantly, levels in senescent cells with inflammatory metabolism were higher than in quiescent cells

at the disc periphery, which exhibit low insulin-signaling activity **(Fig S4.2G-I, yellow)** (Maya et al.,

2024). These observations support our conclusions that high and low metabolic activity coupled to cell

cycle arrest produce different levels of H3K9me3. We thus propose that the widely reported

senescence-associated heterochromatinization associated with elevated H3K9me3 during oncogene-

induced senescence may not be a feature of the senescence program *per se* but rather a consequence

of continuous H3K9 trimethylation driven by metabolic activation from oncogenic mutations in the

absence of cell division (Hernandez-Segura et al., 2018, Huang et al., 2022, Narita et al., 2003). Finally,

in agreement with our previous conclusions, we found that cells undergoing fast regenerative

proliferation around the site of inflammatory damage have lowest levels of H3K9me3 **(Fig S4.2G-I,**

cyan).

(5) CBP/nejire-mediated H3K27 acetylation is high during S-phase

Having revealed how H3K9me3 and H3K27me3 dynamics are coupled via metabolic cues to the cell

cycle, we now wanted to characterize the observed cell cycle dynamics of H3K27 acetylation. The

downregulation of H3K27ac by cell cycle elongation and the upregulation of H3K27ac by cell cycle

acceleration contradicts the simple explanation that an enzymatic activity acetylates H3K27 at a

constant rate. To better understand H3K27ac dynamics, we performed a detailed analysis of H3K27ac

levels throughout the cell cycle by correlating H3K27ac with FUCCI reporters in wing imaginal discs.

We specifically excluded mitotic cells from this analysis, as the highly condensed nature of their

chromosomes and the cells' extreme apical position prevented a direct comparison to other cell cycle

stages.

Using this approach, we found that H3K27ac levels were significantly increased in nuclei of early and

late S-phase cells if compared to G1 and G2 phases **(Fig 5A-C)**. This increase, and especially the

much higher levels in late S-phase in comparison to G2, could only partially be recapitulated by changes

in total H3 or total acetyl-Lysine levels **(Fig S5.1 A-F)**. Of note, fluctuation of H3 and total lysine

acetylation in our microscopy assays can be mostly explained by S-phase dependent histone synthesis

and histone acetylation, as well as fluctuating morphology of interphase nuclei (Stein et al., 1975,

Mendiratta et al., 2019, Verreault et al., 1998). Our finding that H3K27ac levels are higher throughout

S-phase are consistent with our earlier observations: cells which cycle fast repeatedly enter S-phase

and cells which cycle slowly are not often found in S-phase, hence bulk H3K27ac levels in a proliferating

and arrested tissue will appear high and low, respectively.

These results suggest that the enzymes responsible to regulate H3K27 acetylation may exhibit cell-

cycle-dependent regulation. Knock-down of the H3K27ac HAT CBP/nej via RNAi (*neijire-RNAi*)

expressed for 24 h under the control *rn-GAL4*, completely abolished H3K27ac staining in the wing

pouch, confirming that CBP/nej is the main source of H3K27 acetylation in the wing disc **(Fig 5 D-E)**

(Willnow and Teleman, 2024). To provide specific proof that CBP/nej mediates S-phase-dependent

H3K27 acetylation, we overexpressed CBP/nej for 24 h under the control of *rn-GAL4* and correlated

H3K27ac levels with cell-cycle phases using the FUCCI reporter. While global H3K27ac increased upon

CBP/nej overexpression **(Fig 5F,G,J; Fig S5.1 G,H)**, we observed a disproportionate increase of

H3K27ac in late S-phase cells **(Fig 5J')**. This demonstrates that CBP/nej-driven acetylation of H3K27

is especially pronounced during S-phase and indicates that H3K27ac accumulates ectopically in late

S-phase cells when CBP/nej levels are too high. In support of a model where CBP/nej is specifically

active during S-phase, we found that levels of two other CBP/nej-mediated modification (H3K18ac and

H4K8ac) showed a less pronounced but still detectable increase in S-phase cells of normally

proliferating wild type tissue **(Fig S5.2 A-F)**, which is consistent with the reduced CBP/nej substrate

specificity for these two residues, as well as for H3K18 crotonylation, observed in CBP/nej-RNAi

experiments **(Fig S5.2 G-N)**.

As these targeted overexpression experiments suggested that CBP/nej levels may influence S-phase-

dependent activity, we examined a CBP/nej-GFP fusion protein expressed from the endogenous locus

to assess if CBP levels fluctuate during the cell cycle (Marsh et al., 2025). We found that, in proliferating

wing discs, nuclear GFP levels were mostly uniform in all cells **(Fig S5.2 O)**, indicating that regulation

of CBP/nej enzymatic activity, rather than cell cycle dependent changes in protein levels, underlies the

S-phase-specific activity. Indeed, in cultured cells, CBP/p300 activity may be regulated by cyclin-

dependent kinases at the G1/S transition (Ait-Si-Ali et al., 1998).

The pronounced reduction of H3K27ac in late-S-phase, and subsequently in G2-phase or G1-phase

cells, which we observed in development, in arrested cell populations and upon CBP/nej overexpression

or knock-down, demonstrates that the removal of H3K27ac in late S-phase must be a highly controlled

process. Removal of acetylation from the H3K27 residue would likely be mediated by HDAC activity.

We thus depleted the primary H3K27 deacetylase HDAC1 (Rpd3) via expression of an RNAi construct

under the control of *rn-GAL4* for 24 h (Willnow and Teleman, 2024). HDAC1 knockdown led to a mild

but detectable overall increase in H3K27ac, consistent with its role in H3K27ac removal **(Fig 5 H, I, K;**

Fig S5.1 I,J). Yet, importantly, we observed a disproportionate increase of H3K27ac in late S-phase

cells, supporting a model where HDAC1 is required to specifically remove H3K27ac in late S-phase

(Fig 5K').

(6) CBP/nej activity and H3K27acetylation is required to prevent replication stress during S-

phase

We finally wanted to understand what function elevated CBP/nej activity during S-phase may serve. To

analyze if elevated CBP/nej activity may reflect a specific requirement during DNA replication, we

conducted a knock-down of *CBP/nej* in the wing pouch under the control of *rn-GAL4* for 24 h. We first

analyzed DNA replication by EdU incorporation and found a wide-spread decrease in the number of

cells entering S-phase and in DNA replication activity, suggesting that both S-phase entry and

progression was severely slowed within 24 h of *CBP/nej* knock-down **(Fig 6A-C)**. To determine if DNA

replication during *CBP/nej* knock-down had proceeded successfully, we investigated if *CBP/nej-RNAi-*

expressing cell experienced replication stress. We thus analyzed levels of phosphorylated γH2A, a

marker of DNA double strand breaks which can arise at stalled replication forks (Madigan et al., 2002,
Petermann et al., 2010). In all wild type tissues, we observed a similar number of high intensity nuclei
positive for γH2A, likely representing physiological cell death or (unspecific) binding of the antibody in
mitotically active cells (Khan et al., 2017). However, the majority of nuclei are negative for γH2A **(Fig
6D,F)**. Strikingly, in all nuclei of *CBP/nej-RNAi* -expressing cells, we observed a pronounced increase
in nuclear γH2A levels - which were represented by medium fluorescence intensities. The γH2A signal
covered the entire nuclei, suggesting wide-spread DNA damage **(Fig 6E,G,H)**. We conclude that
CBP/nej is specifically required to prevent genome-wide replication stress during S-phase. Accordingly,
CBP/nej-RNAi-expressing cells cannot progress through the cell cycle and completely arrest in G2 **(Fig
6I,J)**. In tissue damage models of wing discs, cells arrest in G2 as a consequence of stress-dependent
activation of JNK-signaling, which downregulates *cdc2/stg* (Cosolo et al., 2019). Yet, co-expressing a
dominant-negative JNKK (bskDN) in *CBP/nej-RNAi*-expressing caused pronounced downregulation of
JNK activation **(Fig S6A-C)**, but still resulted in G2 arrested cells **(Fig S6D)**. This strongly indicates that
a very specific function of CBP/nej in DNA replication caused a G2 cell cycle arrest by activation of a
DNA damage checkpoint, rather than the non-specific activation of cellular stress signaling through
JNK. These observations are thus consistent with a model were high levels of CBP/nej-mediated
H3K27ac are specifically required to ensure faithful DNA replication and to prevent DNA damage and
consequently, a DNA damage induced cell cycle arrest in G2. Of note, analysis of *CBP/nej-*
overexpressing tissues revealed normal patterns of EdU incorporation and distribution, suggesting that
cells with high - but still cell-cycle-dependent - levels of H3K27ac proliferate normally **(Fig S6E-G)**.
426
CBP/nej targets a wide range of substrates, including several non-histone proteins (Stauffer et al., 2007,
Cazzalini et al., 2014, Dutto et al., 2018, Tillhon et al., 2012, Ramadan et al., 2025, Ogiwara et al.,
2011). We thus wanted to determine, if the function of CBP/nej in preventing replication stress is
mediated by its substrate H3K27. Especially, since H3K27 was the histone substrate most sensitive to
the loss of CBP/nej activity in our assays **(**refer to **Fig S5.2 G-N)**. To test this, we generated mosaic
clones homozygous for a deletion of the entire histone gene cluster (ΔHisC), whose viability was
rescued by 12 copies of a histone gene unit in which wild type H3 was replaced with a non-modifiable
H3K27R amino acid substitution (Pengelly et al., 2013). As previously reported, these clones lose
Polycomb gene silencing function normally mediated by H3K27me3, and exhibit derepression of Hox
genes such as *AbdB* **(Fig S6H,I)** (Pengelly et al., 2013). When we assessed if these cells also exhibit
replication stress, we observed broad and elevated levels of γH2A in the H3K27R-substituted clones
(Fig 6K). These findings suggest that the function of CBP/nej in mitigating replication stress during DNA
replication may be, at least in part, mediated through its modification of H3K27 **(Fig 6I)**.
440
441 **DISCUSSION**
442
Our work dissects the link between H3K27ac, H3K27me3, and H3K9me3 modification dynamics and
dynamics of the cell cycle in proliferating tissues *in vivo*. We demonstrate that both physiological and
genetically induced elongation of the cell cycle leads to elevated bulk levels of H3K27me3 and

H3K9me3. Conversely, accelerated cell cycle progression specifically decreases these histone
marks. Genome-wide analyses using CUT&tag assays revealed that these changes occur
predominantly at their distinct pre-existing genomic target loci, implicating canonical histone readers
and modifiers in mediating these cell cycle-dependent changes. Importantly, despite cell cycle-
dependent changes in bulk H3K27me3 levels, expression patterns of Polycomb target genes remained
unaffected, implying that H3K27me3-dependent gene silencing at target loci is buffered even under
altered cell cycle conditions. Mechanistically, we found a tight coupling between histone methylation
rates and cell division frequency, mediated by cellular metabolic activity. Specifically, H3K27me3 and
H3K9me3 levels remained normal upon expression of a constitutively active Insulin receptor,
suggesting that cells channeled the higher metabolic activity into accelerated cell divisions and
maintained H3K27me3 and H3K9me3 levels proportional to cell division rates. Remarkably, active
Insulin receptor-expressing cells forced to arrest their cell cycle accumulated levels of H3K27me3 and
H3K9me3 to much higher levels than arrested wild type cells, demonstrating that metabolic rate tunes
the rate of histone methylation to match the frequency of S-phase occurrence, i.e. the frequency of cell
divisions. These observations have important implications for chromatin dysregulation in pathological
conditions, such as cancer, where the tight coordination between metabolic and cell cycle rates
becomes compromised. An important implication of our findings relates to the emergence senescence-
associated heterochromatin, a phenomenon well-documented during oncogene-induced senescence
(Hernandez-Segura et al., 2018, Huang et al., 2022, Narita et al., 2003). Our data suggest that
H3K9me3 accumulation in senescence-associated heterochromatin may reflect persistent
methyltransferase activity driven by aberrant metabolic stimuli in the absence of replication-dependent
dilution. Thus, the observed chromatin state in senescent cells may be a consequence of the mismatch
between metabolic and cycling rate, rather than an intrinsic feature of the senescence phenotype itself.
An important implication of our findings is that post-mitotic and differentiating cells, such as neurons,
must tightly regulate histone-modifying activities to prevent the aberrant accumulation of repressive
marks, such as H3K9me3 and H3K27me3.
472
Furthermore, we observed that H3K27ac exhibits a unique, cell cycle–dependent profile, peaking
significantly during S-phase. While hyperacetylation of newly synthesized histones at various residues
during S-phase is well-established, an S-phase dependent increase in nuclear acetylation of H3K27
has not been reported (Li et al., 2008, Masumoto et al., 2005, Tessarz and Kouzarides, 2014, Benson
et al., 2006, Falbo and Shen, 2009, Jackson et al., 1976, Lande-Diner et al., 2009, Ruiz-Carrillo et al.,
1975, Annunziato and Hansen, 2000, Sobel et al., 1995). We propose this peak in acetylation during
S-phase is mediated by CBP/nej, potentially regulated by cyclin-dependent kinases at the G1/S
transition (Ait-Si-Ali et al., 1998). Correspondingly, our data is consistent with the idea that H3K27
becomes deacetylated during late S-phase by HDAC1. Generally, HDAC activity is reported to remove
hyperacetylation on newly synthesized histone at the end of S-phase to reestablish higher order
chromatin organization (Annunziato and Seale, 1983). However, whether the H3K27 acetylation cycle
mediated by CBP and HDAC 1 occurs on old histones, newly synthesized histones, or on newly

incorporated histones cannot be resolved with our methods. Yet, the observed H3K27 acetylation cycle
reveals novel cell cycle dependent targets of CBP/nej and HDAC1.

Functionally, we demonstrate that CBP/nej activity is essential during S-phase to prevent DNA damage
from replication stress and consequently delays in cell cycle progression mediated by a replication
checkpoint. In fact, previous studies find that CBP/nej and ATR mei-41 are required for the replication
checkpoint itself to function properly (Smolik and Jones, 2007, Stauffer et al., 2007). This may be
explained by acetylation non-histone substrates by CBP/p300, which stabilizes key DNA damage repair
factors, including PCNA, ATR and ATM, thereby facilitating DNA repair following damage (Stauffer et
al., 2007, Cazzalini et al., 2014, Dutto et al., 2018, Tillhon et al., 2012, Ramadan et al., 2025, Ogiwara
et al., 2011). Consistent with this, mutations in EP300 result in persistent replication stress, replication
fork collapse, genomic instability, and embryonic proliferation defects (Barreto-Galvez et al., 2023; Yao
et al., 1998). (Barreto-Galvez et al., 2023, Yao et al., 1998). Collectively, these findings underscore the
essential role of CBP/p300-mediated acetylation in regulating replication integrity and genomic stability
beyond traditional cell cycle regulation. Importantly, our findings extend this previous work by providing
mechanistic insight, indicating that *in vivo* this function is mediated by acetylation of H3K27, a target
commonly regulated by CBP/nej and HDAC1. Notably, cells carrying the H3K27R mutation exhibit
derepression of Hox genes, closely resembling the phenotype associated with loss-of-function
mutations in PRC1 and PRC2 complexes(Pengelly et al., 2013, Classen et al., 2009, Beuchle et al.,
2001). However, H3K27R cells also display a pronounced, albeit not complete, reduction in overall
H3K27ac levels – in contrast to the nearly complete loss of H3K27ac observed after 24 h of CBP/nej
RNAi-expression. This suggests that CBP/nej-mediated acetylation of K27 may occur on both H3 and
H3.3 variants, with the anti-H3K27ac antibody likely recognizing both forms and the broad substrate
specificity of CBP/nej likely targeting both histone variants. Importantly, acetylation of H3 and H3.3 at
K27 may carry distinct functional implications. H3.3 variants appear to be particularly critical for
enhancer-associated H3K27ac thereby serving roles in interphase cells (Trovato et al., 2020), whereas
acetylation of canonical H3 may regulate chromatin function and DNA replication in S-phase. Indeed,
associations of H3K27ac with Origin Recognition Complex-binding sites and its requirement for gene
amplification during endoreplication (Aggarwal and Calvi, 2004, Eaton et al., 2011, Hua and Orr-
Weaver, 2017, McConnell et al., 2012), suggests that elevated levels of H3K27ac may be essential to
establish an open chromatin environment necessary for efficient DNA replication. Collectively, these
findings position H3K27ac as a central regulator integrating chromatin structure, DNA replication and
genome integrity.

In conclusion, our findings establish a robust mechanistic link between cell cycle dynamics, metabolic
signaling and histone modification of H3K27 and H3K9, and specifically implicate H3K27ac as critical
mediator of replication fidelity and chromatin organization. These insights highlight the necessity of
maintaining balanced histone modification levels to preserve genomic stability, particularly under
varying proliferative and metabolic states, with broad implications for developmental biology, aging, and
cancer biology.

 EXPERIMENTAL PROCEDURES

Drosophila **genetics**

All experiments were performed on *Drosophila melanogaster*. Fly strains (see Table S1) were

maintained on standard fly food (10L water, 74,5g agar, 243g dry yeast, 580g corn flour, 552ml

molasses, 20.7g Nipagin, 35ml propionic acid) at 18°C – 22°C. Larvae from experimental crosses were

allowed to be fed on Bloomington formulation (175.7g Nutry-Fly,1100ml water 20g dry yeast, 1.45g

Nipagin in 15ml Ethanol, 4.8ml Propionic acid) and raised at 18°C or 30°C to control GAL80ts-

dependent induction of GAL4/UAS. To drive expression of UAS-constructs, such as *UAS-Cdk1-RNAi*,

under the control of *rn-GAL4* in the wing pouch, experiments were conducted following protocols from

Smith-Bolton et al. (2009), La Fortezza et al. (2016), and Cosolo et al. (2019), with slight modifications.

Briefly, larvae with the genotype *rn-GAL4, tub-GAL80^{ts}* and appropriate UAS-transgenes were staged

by a 6 hours egg collection and raised at 18°C with a density of 50 larvae per vial. Transgene expression

was activated by shifting the temperature to 30°C for 24 hours on day 7 after egg deposition (AED).

Unless otherwise noted, dissections were performed immediately after this induction period (recovery

time point R0). Control genotypes were derived from either rn^{ts} control crosses or sibling animals

(+/TM6c). All experiments were performed with at least ≥ 2 independent experimental replicates. Our

experimental design did not consider differences between sexes unless for genetic crossing schemes.

FLP/FRT-Mediated Clone Induction

Mitotic recombination to generate mosaic clones of the *Drosophila* histone gene cluster (ΔHisC) was

induced using the FLP/FRT system. Adult flies were allowed to lay eggs for 72 hours at 25 °C. FLP

recombinase expression was triggered by a 1-hour heat shock at 37 °C. Larvae were dissected either

48 or 72 hours after heat shock for subsequent analysis. Transgene cassettes providing a total of 12

copies of the Histone gene unit with a H3K27R point mutation (12×His-GUH3-K27R) rescues cell

viability of (ΔHisC) mutant clones (see (Pengelly et al., 2013)).

Immunohistochemistry of wing imaginal discs

To visualize histone modifications, wing discs from third instar larvae were dissected and fixed for 10

minutes at room temperature in 4% paraformaldehyde in PBS. Washes were performed in PBS

containing 0.1% TritonX-100 (PBT). The discs were then incubated with primary antibodies (**Table S1**)

in 0.1% PBT, gently mixing 2-overnights at 4°C. During incubation with cross-absorbed secondary

antibodies coupled to Alexa Fluorophores (Table S1) at 4°C for 1-overnight, tissues were

counterstained with DAPI (0.25 ng/μL, Sigma, D9542). Tissues were mounted using SlowFade Gold

Antifade (Invitrogen, S36936). To visualize DNA damage using the anti-H2AvD phosphoS137 antibody,

wing discs were fixed 5 minutes at room temperature in 4% paraformaldehyde in PBS. Washes were

performed in PBS containing 0.5% TritonX-100. The primary antibody was incubated in 0.5% PBT

overnight at 4°C and the secondary antibody in 0.5% PBT for 2 hours at room temperature. For all other

staining protocols, discs were fixed for 15 minutes at room temperature in 4% paraformaldehyde in

PBS. All washes and incubations were performed in PBS containing 0.1% TritonX-100 as described

above.To maintain consistency in staining across genotypes, control and experimental wing discs were
processed together in the same vial throughout the protocol and mounted on the same slide whenever
possible. Images were acquired using the Leica TCS SP8 Microscope, using the same confocal settings
for linked samples and processed using tools in Fiji. Figure panels were assembled in Affinity Designer
2.
571
EdU labelling
EdU incorporation was performed using the Click-iT Plus EdU Alexa Fluor 647 Imaging Kit (Table S1).
Briefly, larval cuticles were inverted in Schneider's medium and incubated with EdU (10µM final
concentration) at RT for 15 minutes. Cuticles were then fixed in 4% PFA/PBS for 15 minutes, washed
for 30 minutes in PBT 0.5%. EdU-Click-iT labeling was performed according to manufacturer's
guidelines. Tissues were washed in PBT 0.1%, after which immunohistochemistry, sample processing
and imaging were carried out as described above.
579
SA-β-Gal staining
CellEvent senescence detection kit from Invitrogen (Table S1) was used to check senescence-
associated β-galactosidase activity, following the manufacturer instructions. Briefly, larvae were
dissected in PBS, fixed with 4% PFA, washed with 1% BSA (in PBS), and then incubated in working
solution for 2h at 37°C. Washing steps were performed in PBS and PBS containing 0.1% TritonX-100
(PBT). Tissues were counterstained with DAPI (0.25 ng/µl). Further immunohistochemistry analysis and
sample mounting was performed as described above.
587
Protein synthesis assay using OPP- Click-iT staining
OPP Assays were performed using Click-iT® Plus OPP Protein Synthesis Assay Kits (Invitrogen
Molecular Probe) according to manufacturer's instructions. Briefly, larvae were dissected and inverted
cuticles were incubated with a 1:1000 dilution of Component A in Schneider's medium for 15 minutes
on a nutator. Larval cuticles were fixed with 4% paraformaldehyde for 15 minutes, rinsed twice in 0.1%
PBT, and permeabilized with 0.5% PBT for 15 minutes. The cuticles were then stained with the Click-
iT® cocktail for 30 minutes at room temperature, protected from light. Further immunohistochemistry
analysis and sample mounting was performed as described above.
596
CUT&Tag sample preparation
Briefly, for each genotype around 80-100 wing discs were dissected in ice cold Schneider's medium,
collected, snap frozen and stored at -80°C until further use. Nuclei were isolated with the help of a
mechanical pestle and processed for CUT&Tag analysis as previously described (Zenk et al., 2023,
Cardamone et al., 2025). For normalizing the levels of histone H3 modifications and as a proxy of the
starting material in each condition, we also generated samples for total histone H3. Libraries were
prepared by supplementing the NEBNext HighFidelity 2× PCR Master Mix with 1 pg of Tn5-tagmented
lambda DNA (New England Biolabs, #N3011S) as a spike-in normalizer. The library indexing was
performed using Illumina i5 and i7 (Buenrostro et al., 2015) through 15 cycles (1 x 5 min at 72 °C, 1 x

30 sec at 98°C, 13 x 10 s at 98 °C, 30 sec at 63 °C, 1 x 1 min at 72 °C, hold at 4 °C). The libraries were

purified using Nucleomag NGS Clean-up and Size Select beads (Macherey-Nagel, #744970.50).

Quality checks were done with Qbit DNA HS Assay (ThermoScientific, #Q32854) and Bioanalyzer

(Agilent). Finally, two independent library replicates were sent for next generation sequencing.

QUANTIFICATIONS, BIOINFORMATIC ANALYSIS AND STATISTICS

General comments

For all quantifications of imaging data, control and experimental samples were processed together and

imaged in parallel, using the same confocal settings. Images were processed, analyzed and quantified

using tools in Fiji (ImageJ 2.9.0) (Schindelin et al., 2012). Extreme care was taken to apply consistent

methods (i.e., number of projected sections, thresholding methods, processing) for image analysis.

Figure panels were assembled using Affinity Designer 2.6.2. Statistical analyses were performed in

Graphpad Prism. Illustrations were created in Biorender.

Quantification of histone modifications

For quantification of levels of histone modifications, an xy-representation containing the maximum

number of pouch and hinge cells was generated using a sum projection of the relevant channel. Three

square regions of interest (ROIs) measuring 15 × 15 µm were placed in distinct areas of either the

pouch or the hinge, carefully excluding regions with developmentally arrested cells. The mean intensity

of the modification signal was measured within each ROI. For each disc, the average pouch signal was

normalized to the average hinge signal and reported as a pouch-to-hinge intensity ratio.

Quantification of EdU incorporation

Figure S1.1: To first create a nuclear mask, an xy- representation containing the maximum number of

pouch nuclei was generated using a sum projection of the DAPI channel. The background signal was

reduced using the "Subtract Background" function with a rolling ball radius of 100 pixels. Local contrast

was enhanced with the "Enhance Local Contrast (CLAHE)" function, using a block size of 127, 256

histogram bins, and a maximum slope of 3. Automatic thresholding was performed using the "Moments"

method with the "dark" option selected, and the image was then converted into a binary mask of DAPI-

positive nuclear areas. A nuclear ROI was generated from this mask using the selection tool. The pouch

region was manually selected based on wing fold morphology. To isolate the nuclei within the pouch, a

binary "AND" operation was used to combine nuclear mask and pouch selection. This final pouch-

specific DAPI mask was used to measure EdU signal intensity.

Figure 6: To create an EdU mask, an xy- representation containing the maximum number of pouch

nuclei was generated using a sum projection of the EdU channel. A "Gaussian blur" (σ = 2) was applied

to reduce noise and smooth the signal. Automatic thresholding was then performed using the

"Moments" method with the "dark" option selected. The image was converted into a binary EdU mask

with a black background. A ROI was generated from this mask using the selection tool to define EdU-

positive pixels. Separately, the pouch region was manually selected based on wing fold morphology.

To isolate EdU-positive pixels specifically within the pouch, a binary "AND" operation was applied to
combine the EdU selection with the pouch selection. This final pouch-specific EdU mask was used to
measure EdU signal intensity.
To assess changes in the number of proliferating cells, pouch-specific DAPI and EdU masks were
generated as described above. A binary "AND" operation was used to extract EdU-positive nuclei
located within DAPI-positive regions in the pouch. The area of EdU-positive nuclei was then divided by
the total area of EdU and DAPI-positive regions within the pouch to calculate the fraction of proliferating
cells. This EdU-to-DAPI area ratio was calculated per disc and used for comparative analysis.
654

655   **Quantification of OPP levels in wing imaginal discs**
656   An xy-representation containing the maximum number of pouch cells was generated using a sum
657   projection of the relevant channels. In both control and constitutively active insulin receptor-expressing
658   discs, the pouch region was manually selected based on wing fold morphology. Mean OPP intensity
659   was then measured within the defined pouch region for each disc.
660

661   **Quantification of pouch size**
662   For both control and constitutively active insulin receptor-expressing discs, a slice containing the
663   maximum number of pouch cells was selected. In each case, the pouch region was manually defined
664   based on wing fold morphology. The area size of the selected pouch region was then measured for
665   each disc.
666

Cell cycle phase classification using FUCCI and quantifications of H3, Total Acetyl Lysine,
H3K27ac, H3K18ac and H4K8ac level
To classify nuclei in the wing pouch according to cell cycle phase and to quantitatively correlate levels
of H3 and acetylation levels with each phase, a custom image analysis workflow was implemented in
ImageJ/Fiji. A single z-slice containing the highest number of pouch cells was selected from each
control disc. The image stack was split into individual channels (e.g., DAPI, GFP-E2f1[1-230], mRFP-NLS-
CycB[1-266], H3K27ac). To generate a nuclear mask, the DAPI, GFP, RFP, and other nuclear markers
were averaged sequentially using the Image Calculator. The resulting image was thresholded using the
Otsu method (upper limit set to 255) and converted into a binary mask defining nuclear pixels. A nuclear
pixel selection was generated from the binary image and added to the ROI Manager. The pouch region
was manually defined based on wing fold morphology. A binary "AND" operation was used to restrict
the nuclear pixel mask to the pouch, producing a final pouch-specific nuclear mask. This mask was
overlaid on the GFP, RFP, and other nuclear marker channels, and the X/Y coordinates of each ROI
were used to extract signal intensities for each nuclear pixel. Threshold values for GFP and RFP
determined based on average intensity of 5 nuclei in early S-phase. Each nuclear pixel was classified
into a cell cycle phase using these thresholds, following FUCCI-based criteria: G2 phase: GFP >
threshold, RFP > threshold; G1 phase: GFP > threshold, RFP ≤ threshold; Early/mid S phase: GFP ≤
threshold, RFP ≤ threshold; Late S phase: GFP ≤ threshold, RFP > threshold. For each nuclear pixel,
the intensity of the corresponding nuclear markers (e.g., H3, acetyl-lysine, H3K27ac) was recorded. To

account for inter-sample variability, intensity values within each cell cycle phase were normalized by

dividing them by the total nuclear marker signal summed across all phases (G1, early S, late S, and

G2) within the same disc. This normalization provided the relative contribution of each cell cycle phase

to the total signal, allowing for direct comparisons across samples.

γH2Av intensity quantifications

To create a nuclear mask, an xy- representation containing the maximum number of pouch nuclei was

generated using a sum projection of the DAPI channel. The background signal was reduced using the

"Subtract Background" function with a rolling ball radius of 100 pixels. Local contrast was enhanced

with the "Enhance Local Contrast (CLAHE)" function, using a block size of 127, 256 histogram bins,

and a maximum slope of 3. Automatic thresholding was performed using the "Moments" method with

the "dark" option selected, and the image was then converted into a binary mask. A nuclear ROI was

generated from this mask using the selection tool. The pouch region was manually selected based on

wing fold morphology. To isolate the nuclear pixels within the pouch, a binary "AND" operation was

used to combine nuclear mask and pouch selection. This final mask was used to measure γH2Av and

DAPI intensities for each disc. γH2Av intensities were normalized to the corresponding DAPI intensity.

CUT&Tag Bioinformatic Analysis

The CUT&Tag data processing was performed as described before with some modifications

(Atinbayeva et al., 2024, Ibarra-Morales et al., 2021, Cardamone et al., 2025). As our quantitative

CUT&Tag sequencing libraries contained DNA from *D. melanogaster* and spike-in of *lambda* phage

genome (Genbank: J02459.1), we mapped the raw paired-end reads to a constructed hybrid dm6 and

Lambda phage genome using *snakePipes-v3.0.0 DNA-mapping* (Bhardwaj et al., 2019) with updated

Bowtie2 mapping parameters, as well as adapter trimming and MAPQ ≥ 1 filtering (*DNAmapping --mapq*

1 --trim --fastqc --properPairs --dedup --cutntag --DAG --alignerOpts='--local --very-sensitive-local --no-

discordant --no-mixed -I 10 -X 700'). To assess reproducibility of both replicates, we utilised the

multiBamSummary and *plotCorrelation* from *deepTools-v.2.5.7* to compute the Spearman correlation

between the two replicates. Aligned replicates were merged before normalization to both H3 and spike-

in signals. The code used for the H3 and *Lambda* normalization analysis was based on work done by

Yinxiu Zhan (https://github.com/zhanyinx/atinbayeva_paper_2023) for (Atinbayeva et al., 2024).

CUT&Tag peaks were called using *snakePipes-v3.0.0 ChIPseq* with *MACS2-v 2.2.9.1* (*ChIPseq -d*

Output_spikein hybrid.yaml H3_spikein_config.yaml --peakCaller MACS2 --cutntag --

peakCallerOptions "--broad --qvalue 0.5 -f BAMPE" --DAG). Normalized bigWig and bam files were

generated with *deepTools-v3.5.6* and used to compute matrices of signal enrichment +/− 5kb around

transcription start sites (TSS) or peak centers of shared or unique peaks for *wt* and *Cdk1KD*. Coverage

heatmaps and profiles were created using *plotHeatmap* or *plotProfile* from *deepTools-v.2.5.7*.

For the quantification, the number of PE reads were counted in each shared or unique peak for *wt* and

Cdk1KD (*multiBamSummary BED-file --bamfiles <input files> --extendReads --outRawCounts <file>*)

as well as in 500 bp bins across the dm6 genome (*multiBamSummary bins --bamfiles <input files> --*

binSize 500 --extendReads --outRawCounts <file>). The *DESeq2-v1.38.3* analysis was performed

using the previously computed H3 and spike-in scaling factors from merged data for size factor. Bins and peaks with basemean (average read count across samples) of less than or equal to 25 read counts were discarded. The design matrix was setup to compare the samples by condition and correct for replicate effects (*design = ~replicate + condition*). Finally, we executed the *DESeq2* shrinkage of log2 fold changes (*type = 'normal'*).

CODE AND DATA AVAILABILITY

All data, workflows and FIJI based algorithms necessary to interpret the imaging data are included within the manuscript. Original imaging data and any additional information required to reanalyze the data reported in this paper will be shared by AKC upon request. The code used for the mapping, normalization, peak calling and signal quantification is available on Github: https://github.com/LaraH9/nogay_et_al_2025. The CUT&Tag sequencing data generated in this study is provided on NCBI SRA: https://www.ncbi.nlm.nih.gov/sra/PRJNA1300380.

AUTHOR CONTRIBUTIONS

Conceptualization LN, AKC; Data curation LN, LH, AKC; Formal analysis LN, LH, AKC; Funding acquisition AKC; Investigation LN, AVM; Methodology LN, AVM, FC, IG, AS, AF, LE, AKC; Project administration IG, AKC; Supervision NI, AKC; Validation LN, AVM, FC, IG; Visualization LN, LH, AKC; Writing – original draft LN, AKC; Writing – review & editing LN, AVM, AKC

ACKNOWLEDGEMENTS

We thank the staff of the Life Imaging Center (LIC) in the Hilde Mangold House (HMH) of the Albert-Ludwigs-University of Freiburg for help with their confocal microscopy resources, and the excellent support in image recording. We specifically thank the DFG for supporting our imaging work through project number 414136422. We thank the Bioinformatics and Sequencing facilities at the MPI-IE for supporting this work. We thank Galina Erikson for supporting our bioinformatic analysis. We thank Laura Bauttita, Melissa Harrison, Iswar Hariharan, Dirk Bohmann and Jürg Müller for sharing reagents and the Bloomington *Drosophila* Stock Center (BDSC), the Vienna *Drosophila* Stock Collection (VDRC), the University of Zurich ORFeome Project (FlyORF) and the Developmental Studies Hybridoma Bank (DSHB) for providing fly stocks and antibodies. We thank the IMPRS-EBM and SGBM graduate schools for supporting our doctoral researchers.

FUNDING

Funding for this work was provided by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany´s Excellence Strategy (CIBSS – EXC-2189), the DFG Heisenberg Program to AKC (668189), as well as DFG grants to AKC (667603) and the Boehringer Ingelheim

766 Foundation (BIF Plus3 & Rise Up) to AKC. The funders had no role in study design, data collection and
767 analysis, decision to publish, or preparation of the manuscript.
768
769 **DECLARATION OF INTEREST**

770
771 The authors declare no competing interests.
772

REFERENCES

AAGAARD, L., LAIBLE, G., SELENKO, P., SCHMID, M., DORN, R., SCHOTTA, G., KUHFITTIG, S., WOLF, A.,
LEBERSORGER, A., SINGH, P. B., REUTER, G. & JENUWEIN, T. 1999. Functional mammalian
homologues of the Drosophila PEV-modifier Su(var)3-9 encode centromere-associated proteins which
complex with the heterochromatin component M31. *EMBO J,* 18**,** 1923-38.

AGGARWAL, B. D. & CALVI, B. R. 2004. Chromatin regulates origin activity in Drosophila follicle cells. *Nature,*
430**,** 372-6.

AIT-SI-ALI, S., RAMIREZ, S., BARRE, F. X., DKHISSI, F., MAGNAGHI-JAULIN, L., GIRAULT, J. A., ROBIN, P.,
KNIBIEHLER, M., PRITCHARD, L. L., DUCOMMUN, B., TROUCHE, D. & HAREL-BELLAN, A. 1998.
Histone acetyltransferase activity of CBP is controlled by cycle-dependent kinases and oncoprotein E1A.
Nature, 396**,** 184-6.

ALABERT, C., BARTH, T. K., REVERON-GOMEZ, N., SIDOLI, S., SCHMIDT, A., JENSEN, O. N., IMHOF, A. &
GROTH, A. 2015. Two distinct modes for propagation of histone PTMs across the cell cycle. *Genes Dev,*
29**,** 585-90.

ALABERT, C. & GROTH, A. 2012. Chromatin replication and epigenome maintenance. *Nat Rev Mol Cell Biol,* 13**,**
153-67.

ANNUNZIATO, A. T. 2015. The Fork in the Road: Histone Partitioning During DNA Replication. *Genes (Basel),* 6**,**
353-71.

ANNUNZIATO, A. T. & HANSEN, J. C. 2000. Role of histone acetylation in the assembly and modulation of
chromatin structures. *Gene Expr,* 9**,** 37-61.

ANNUNZIATO, A. T. & SEALE, R. L. 1983. Histone deacetylation is required for the maturation of newly replicated
chromatin. *J Biol Chem,* 258**,** 12675-84.

ATINBAYEVA, N., VALENT, I., ZENK, F., LOESER, E., RAUER, M., HERUR, S., QUARATO, P., PYROWOLAKIS,
G., GOMEZ-AULI, A., MITTLER, G., CECERE, G., ERHARDT, S., TIANA, G., ZHAN, Y. & IOVINO, N.
2024. Inheritance of H3K9 methylation regulates genome architecture in Drosophila early embryos. *EMBO*
J, 43**,** 2685-2714.

BARRETO-GALVEZ, A., NILJIKAR, M., GAGLIARDI, J., ZHANG, R., KUMAR, V., JURUWALA, A., PRADEEP, A.,
SHAIKH, A., TIWARI, P., SHARMA, K., GERHARDT, J., CAO, J., KATAOKA, K., DURBIN, A., QI, J., YE,
B. H. & MADIREDDY, A. 2023. Acetyl transferase EP300 deficiency leads to chronic replication stress
mediated by defective fork protection at stalled replication forks. *bioRxiv*.

BENSON, L. J., GU, Y., YAKOVLEVA, T., TONG, K., BARROWS, C., STRACK, C. L., COOK, R. G., MIZZEN, C.
A. & ANNUNZIATO, A. T. 2006. Modifications of H3 and H4 during chromatin replication, nucleosome
assembly, and histone exchange. *J Biol Chem,* 281**,** 9287-96.

BEUCHLE, D., STRUHL, G. & MULLER, J. 2001. Polycomb group proteins and heritable silencing of Drosophila
Hox genes. *Development,* 128**,** 993-1004.

BHARDWAJ, V., HEYNE, S., SIKORA, K., RABBANI, L., RAUER, M., KILPERT, F., RICHTER, A. S., RYAN, D. P.
& MANKE, T. 2019. snakePipes: facilitating flexible, scalable and integrative epigenomic analysis.
Bioinformatics, 35**,** 4757-4759.

BHATNAGAR, A., KARNAY, A. M. & ELEFANT, F. 2023. Drosophila Epigenetics. *Handbook of Epigenetics.*

BLACKLEDGE, N. P. & KLOSE, R. J. 2021. The molecular principles of gene regulation by Polycomb repressive
complexes. *Nat Rev Mol Cell Biol,* 22**,** 815-833.

BOIJA, A., MAHAT, D. B., ZARE, A., HOLMQVIST, P. H., PHILIP, P., MEYERS, D. J., COLE, P. A., LIS, J. T.,
STENBERG, P. & MANNERVIK, M. 2017. CBP Regulates Recruitment and Release of Promoter-
Proximal RNA Polymerase II. *Mol Cell,* 68**,** 491-503 e5.

BROOKES, E. & SHI, Y. 2014. Diverse epigenetic mechanisms of human disease. *Annu Rev Genet,* 48**,** 237-68.

BUENROSTRO, J. D., WU, B., LITZENBURGER, U. M., RUFF, D., GONZALES, M. L., SNYDER, M. P., CHANG,
H. Y. & GREENLEAF, W. J. 2015. Single-cell chromatin accessibility reveals principles of regulatory
variation. *Nature,* 523**,** 486-90.

BUTTITTA, L. A. & EDGAR, B. A. 2007. Mechanisms controlling cell cycle exit upon terminal differentiation. *Curr
Opin Cell Biol,* 19**,** 697-704.

BUTTITTA, L. A., KATZAROFF, A. J., PEREZ, C. L., DE LA CRUZ, A. & EDGAR, B. A. 2007. A double-assurance
mechanism controls cell cycle exit upon terminal differentiation in Drosophila. *Dev Cell,* 12**,** 631-43.

CAO, R., WANG, L., WANG, H., XIA, L., ERDJUMENT-BROMAGE, H., TEMPST, P., JONES, R. S. & ZHANG, Y.
2002. Role of histone H3 lysine 27 methylation in Polycomb-group silencing. *Science,* 298**,** 1039-43.

CARDAMONE, F., PIVA, A., LOSER, E., EICHENBERGER, B., ROMERO-MULERO, M. C., ZENK, F., SHIELDS,
E. J., CABEZAS-WALLSCHEID, N., BONASIO, R., TIANA, G., ZHAN, Y. & IOVINO, N. 2025. Chromatin
landscape at cis-regulatory elements orchestrates cell fate decisions in early embryogenesis. *Nat
Commun,* 16**,** 3007.

CAVALLI, G. & HEARD, E. 2019. Advances in epigenetics link genetics to the environment and disease. *Nature,*
571**,** 489-499.

CAZZALINI, O., SOMMATIS, S., TILLHON, M., DUTTO, I., BACHI, A., RAPP, A., NARDO, T., SCOVASSI, A. I.,
NECCHI, D., CARDOSO, M. C., STIVALA, L. A. & PROSPERI, E. 2014. CBP and p300 acetylate PCNA
to link its degradation with nucleotide excision repair synthesis. *Nucleic Acids Res,* 42**,** 8433-48.

CHEN, Y. C., KOUTELOU, E. & DENT, S. Y. R. 2022. Now open: Evolving insights to the roles of lysine acetylation
in chromatin organization and function. *Mol Cell,* 82**,** 716-727.

CIABRELLI, F., RABBANI, L., CARDAMONE, F., ZENK, F., LOSER, E., SCHACHTLE, M. A., MAZINA, M.,
LOUBIERE, V. & IOVINO, N. 2023. CBP and Gcn5 drive zygotic genome activation independently of their
catalytic activity. *Sci Adv,* 9**,** eadf2687.

CLASSEN, A. K., BUNKER, B. D., HARVEY, K. F., VACCARI, T. & BILDER, D. 2009. A tumor suppressor activity
of Drosophila Polycomb genes mediated by JAK-STAT signaling. *Nat Genet,* 41**,** 1150-5.

COSOLO, A., JAISWAL, J., CSORDAS, G., GRASS, I., UHLIROVA, M. & CLASSEN, A. K. 2019. JNK-dependent
cell cycle stalling in G2 promotes survival and senescence-like phenotypes in tissue stress. *Elife,* 8.

CRUCIANELLI, C., JAISWAL, J., VIJAYAKUMAR MAYA, A., NOGAY, L., COSOLO, A., GRASS, I. & CLASSEN,
A. K. 2022. Distinct signaling signatures drive compensatory proliferation via S-phase acceleration. *PLoS
Genet,* 18**,** e1010516.

DE MORREE, A. & RANDO, T. A. 2023. Regulation of adult stem cell quiescence and its functions in the
maintenance of tissue integrity. *Nat Rev Mol Cell Biol,* 24**,** 334-354.

DUTTO, I., SCALERA, C. & PROSPERI, E. 2018. CREBBP and p300 lysine acetyl transferases in the DNA
damage response. *Cell Mol Life Sci,* 75**,** 1325-1338.

EATON, M. L., PRINZ, J. A., MACALPINE, H. K., TRETYAKOV, G., KHARCHENKO, P. V. & MACALPINE, D. M.
2011. Chromatin signatures of the Drosophila replication program. *Genome Res,* 21**,** 164-74.

EBERT, A., SCHOTTA, G., LEIN, S., KUBICEK, S., KRAUSS, V., JENUWEIN, T. & REUTER, G. 2004. Su(var)
genes regulate the balance between euchromatin and heterochromatin in Drosophila. *Genes Dev,* 18**,**
2973-83.

ESCOBAR, T. M., LOYOLA, A. & REINBERG, D. 2021. Parental nucleosome segregation and the inheritance of
cellular identity. *Nat Rev Genet,* 22**,** 379-392.

ESCOBAR, T. M., OKSUZ, O., SALDANA-MEYER, R., DESCOSTES, N., BONASIO, R. & REINBERG, D. 2019.
Active and Repressed Chromatin Domains Exhibit Distinct Nucleosome Segregation during DNA
Replication. *Cell,* 179**,** 953-963 e11.

FALBO, K. B. & SHEN, X. 2009. Histone modifications during DNA replication. *Mol Cells,* 28**,** 149-54.

FEINBERG, A. P., KOLDOBSKIY, M. A. & GONDOR, A. 2016. Epigenetic modulators, modifiers and mediators in
cancer aetiology and progression. *Nat Rev Genet,* 17**,** 284-99.

FLOC'HLAY, S., BALAJI, R., STANKOVIC, D., CHRISTIAENS, V. M., BRAVO GONZALEZ-BLAS, C., DE
WINTER, S., HULSELMANS, G. J., DE WAEGENEER, M., QUAN, X., KOLDERE, D., ATKINS, M.,
HALDER, G., UHLIROVA, M., CLASSEN, A. K. & AERTS, S. 2023. Shared enhancer gene regulatory
networks between wound and oncogenic programs. *Elife,* 12.

FLURY, V. & GROTH, A. 2024. Safeguarding the epigenome through the cell cycle: a multitasking game. *Curr*
Opin Genet Dev, 85**,** 102161.

HERNANDEZ-SEGURA, A., NEHME, J. & DEMARIA, M. 2018. Hallmarks of Cellular Senescence. *Trends Cell*
Biol, 28**,** 436-453.

HOLMQVIST, P. H. & MANNERVIK, M. 2013. Genomic occupancy of the transcriptional co-activators p300 and
CBP. *Transcription,* 4**,** 18-23.

HUA, B. L. & ORR-WEAVER, T. L. 2017. DNA Replication Control During Drosophila Development: Insights into
the Onset of S Phase, Replication Initiation, and Fork Progression. *Genetics,* 207**,** 29-47.

HUANG, W., HICKSON, L. J., EIRIN, A., KIRKLAND, J. L. & LERMAN, L. O. 2022. Cellular senescence: the good,
the bad and the unknown. *Nat Rev Nephrol,* 18**,** 611-627.

IBARRA-MORALES, D., RAUER, M., QUARATO, P., RABBANI, L., ZENK, F., SCHULTE-SASSE, M.,
CARDAMONE, F., GOMEZ-AULI, A., CECERE, G. & IOVINO, N. 2021. Histone variant H2A.Z regulates
zygotic genome activation. *Nat Commun,* 12**,** 7002.

ITO, T. & IGAKI, T. 2021. Yorkie drives Ras-induced tumor progression by microRNA-mediated inhibition of cellular
senescence. *Sci Signal,* 14.

JACKSON, V., SHIRES, A., TANPHAICHITR, N. & CHALKLEY, R. 1976. Modifications to histones immediately
after synthesis. *J Mol Biol,* 104**,** 471-83.

JAISWAL, J., EGERT, J., ENGESSER, R., PEYROTON, A. A., NOGAY, L., WEICHSELBERGER, V.,
CRUCIANELLI, C., GRASS, I., KREUTZ, C., TIMMER, J. & CLASSEN, A. K. 2023. Mutual repression

between JNK/AP-1 and JAK/STAT stratifies senescent and proliferative cell behaviors during tissue
regeneration. *PLoS Biol,* 21**,** e3001665.

JOHNSTON, L. A. & EDGAR, B. A. 1998. Wingless and Notch regulate cell-cycle arrest in the developing
Drosophila wing. *Nature,* 394**,** 82-4.

KHAN, C., MULIYIL, S., AYYUB, C. & RAO, B. J. 2017. The initiator caspase Dronc plays a non-apoptotic role in
promoting DNA damage signalling in D. melanogaster. *J Cell Sci,* 130**,** 2984-2995.

LANDE-DINER, L., ZHANG, J. & CEDAR, H. 2009. Shifts in replication timing actively affect histone acetylation
during nucleosome reassembly. *Mol Cell,* 34**,** 767-74.

LEE, L. A. & ORR-WEAVER, T. L. 2003. Regulation of cell cycles in Drosophila development: intrinsic and extrinsic
cues. *Annu Rev Genet,* 37**,** 545-78.

LI, Q., ZHOU, H., WURTELE, H., DAVIES, B., HORAZDOVSKY, B., VERREAULT, A. & ZHANG, Z. 2008.
Acetylation of histone H3 lysine 56 regulates replication-coupled nucleosome assembly. *Cell,* 134**,** 244-
55.

LIU, W., TANASA, B., TYURINA, O. V., ZHOU, T. Y., GASSMANN, R., LIU, W. T., OHGI, K. A., BENNER, C.,
GARCIA-BASSETS, I., AGGARWAL, A. K., DESAI, A., DORRESTEIN, P. C., GLASS, C. K. &
ROSENFELD, M. G. 2010. PHF8 mediates histone H4 lysine 20 demethylation events involved in cell
cycle progression. *Nature,* 466**,** 508-12.

LOPEZ-HERNANDEZ, L., TOOLAN-KERR, P., BANNISTER, A. J. & MILLAN-ZAMBRANO, G. 2025. Dynamic
histone modification patterns coordinating DNA processes. *Mol Cell,* 85**,** 225-237.

MA, Y., KANAKOUSAKI, K. & BUTTITTA, L. 2015. How the cell cycle impacts chromatin architecture and
influences cell fate. *Front Genet,* 6**,** 19.

MADIGAN, J. P., CHOTKOWSKI, H. L. & GLASER, R. L. 2002. DNA double-strand break-induced phosphorylation
of Drosophila histone variant H2Av helps prevent radiation-induced apoptosis. *Nucleic Acids Res,* 30**,**
3698-705.

MARSH, A. J., PIROGOV, S., KAUR, Y., RUFFRIDGE, A. J., SAJWAN, S., GIBSON, T. J., HUNT, G., HARRISON,
M. M. & MANNERVIK, M. 2025. Catalytic-dependent and independent functions of the histone
acetyltransferase CBP promote pioneer-factor-mediated zygotic genome activation. *Mol Cell*.

MASUMOTO, H., HAWKE, D., KOBAYASHI, R. & VERREAULT, A. 2005. A role for cell-cycle-regulated histone
H3 lysine 56 acetylation in the DNA damage response. *Nature,* 436**,** 294-8.

MATTHEWS, H. K., BERTOLI, C. & DE BRUIN, R. A. M. 2022. Cell cycle control in cancer. *Nat Rev Mol Cell Biol,*
23**,** 74-88.

MAYA, A. V., NOGAY, L., HECKMANN, L., GRASS, I., KIERDORF, K., BüSCHER, J. & CLASSEN, A.-K. 2024.
PDK-1/S6K and mTORC1 bypass systemic growth restrictions to promote regeneration. *bioRxiv***,**
2024.08.05.606658.

MCCONNELL, K. H., DIXON, M. & CALVI, B. R. 2012. The histone acetyltransferases CBP and Chameau integrate
developmental and DNA replication programs in Drosophila ovarian follicle cells. *Development,* 139**,** 3880-
90.

MENDIRATTA, S., GATTO, A. & ALMOUZNI, G. 2019. Histone supply: Multitiered regulation ensures chromatin
dynamics throughout the cell cycle. *J Cell Biol,* 218**,** 39-54.

MILAN, M., CAMPUZANO, S. & GARCIA-BELLIDO, A. 1996. Cell cycling and patterned cell proliferation in the
wing primordium of Drosophila. *Proc Natl Acad Sci U S A,* 93**,** 640-5.

MILLAN-ZAMBRANO, G., BURTON, A., BANNISTER, A. J. & SCHNEIDER, R. 2022. Histone post-translational
modifications - cause and consequence of genome function. *Nat Rev Genet,* 23**,** 563-580.

MULLER, J., GAUNT, S. & LAWRENCE, P. A. 1995. Function of the Polycomb protein is conserved in mice and
flies. *Development,* 121**,** 2847-52.

NARITA, M., NUNEZ, S., HEARD, E., NARITA, M., LIN, A. W., HEARN, S. A., SPECTOR, D. L., HANNON, G. J.
& LOWE, S. W. 2003. Rb-mediated heterochromatin formation and silencing of E2F target genes during
cellular senescence. *Cell,* 113**,** 703-16.

NEUFELD, T. P., DE LA CRUZ, A. F., JOHNSTON, L. A. & EDGAR, B. A. 1998. Coordination of growth and cell
division in the Drosophila wing. *Cell,* 93**,** 1183-93.

ODA, H., HUBNER, M. R., BECK, D. B., VERMEULEN, M., HURWITZ, J., SPECTOR, D. L. & REINBERG, D.
2010. Regulation of the histone H4 monomethylase PR-Set7 by CRL4(Cdt2)-mediated PCNA-dependent
degradation during DNA damage. *Mol Cell,* 40**,** 364-76.

OGIWARA, H., UI, A., OTSUKA, A., SATOH, H., YOKOMI, I., NAKAJIMA, S., YASUI, A., YOKOTA, J. & KOHNO,
T. 2011. Histone acetylation by CBP and p300 at double-strand break sites facilitates SWI/SNF chromatin
remodeling and the recruitment of non-homologous end joining factors. *Oncogene,* 30**,** 2135-46.

PADEKEN, J., METHOT, S. P. & GASSER, S. M. 2022. Establishment of H3K9-methylated heterochromatin and
its functions in tissue differentiation and maintenance. *Nat Rev Mol Cell Biol,* 23**,** 623-640.

PENGELLY, A. R., COPUR, O., JACKLE, H., HERZIG, A. & MULLER, J. 2013. A histone mutant reproduces the
phenotype caused by loss of histone-modifying factor Polycomb. *Science,* 339**,** 698-9.

PETERMANN, E., ORTA, M. L., ISSAEVA, N., SCHULTZ, N. & HELLEDAY, T. 2010. Hydroxyurea-stalled
replication forks become progressively inactivated and require two different RAD51-mediated pathways
for restart and repair. *Mol Cell,* 37**,** 492-502.

PETERSON, C. L. & LANIEL, M. A. 2004. Histones and histone modifications. *Curr Biol,* 14**,** R546-51.

PETRUK, S., BLACK, K. L., KOVERMANN, S. K., BROCK, H. W. & MAZO, A. 2013. Stepwise histone modifications
are mediated by multiple enzymes that rapidly associate with nascent DNA during replication. *Nat
Commun,* 4**,** 2841.

PETRUK, S., SEDKOV, Y., SMITH, S., TILLIB, S., KRAEVSKI, V., NAKAMURA, T., CANAANI, E., CROCE, C. M.
& MAZO, A. 2001. Trithorax and dCBP acting in a complex to maintain expression of a homeotic gene.
Science, 294**,** 1331-4.

PHILIP, P., BOIJA, A., VAID, R., CHURCHER, A. M., MEYERS, D. J., COLE, P. A., MANNERVIK, M. &
STENBERG, P. 2015. CBP binding outside of promoters and enhancers in Drosophila melanogaster.
Epigenetics Chromatin, 8**,** 48.

PROBER, D. A. & EDGAR, B. A. 2000. Ras1 promotes cellular growth in the Drosophila wing. *Cell,* 100**,** 435-46.

RAMADAN, W. S., AHMED, S. B. M., TALAAT, I. M., LOZON, L., MOUFFAK, S., GEMOLL, T., MANSOUR, W. Y.
& EL-AWADY, R. 2025. The histone acetyltransferase CBP participates in regulating the DNA damage
response through ATM after double-strand breaks. *Genome Biol,* 26**,** 89.

RAMOS-ALONSO, L. & CHYMKOWITCH, P. 2024. Maintaining transcriptional homeostasis during cell cycle.
Transcription, 15**,** 1-21.

REVERON-GOMEZ, N., GONZALEZ-AGUILERA, C., STEWART-MORGAN, K. R., PETRYK, N., FLURY, V.,
GRAZIANO, S., JOHANSEN, J. V., JAKOBSEN, J. S., ALABERT, C. & GROTH, A. 2018. Accurate
Recycling of Parental Histones Reproduces the Histone Modification Landscape during DNA Replication.
Mol Cell, 72**,** 239-249 e5.

RUIZ-CARRILLO, A., WANGH, L. J. & ALLFREY, V. G. 1975. Processing of newly synthesized histone molecules.
Science, 190**,** 117-28.

SCHINDELIN, J., ARGANDA-CARRERAS, I., FRISE, E., KAYNIG, V., LONGAIR, M., PIETZSCH, T., PREIBISCH,
S., RUEDEN, C., SAALFELD, S., SCHMID, B., TINEVEZ, J. Y., WHITE, D. J., HARTENSTEIN, V.,
ELICEIRI, K., TOMANCAK, P. & CARDONA, A. 2012. Fiji: an open-source platform for biological-image
analysis. *Nat Methods,* 9**,** 676-82.

SMITH-BOLTON, R. K., WORLEY, M. I., KANDA, H. & HARIHARAN, I. K. 2009. Regenerative growth in Drosophila
imaginal discs is regulated by Wingless and Myc. *Dev Cell,* 16**,** 797-809.

SMOLIK, S. & JONES, K. 2007. Drosophila dCBP is involved in establishing the DNA replication checkpoint. *Mol*
Cell Biol, 27**,** 135-46.

SOBEL, R. E., COOK, R. G., PERRY, C. A., ANNUNZIATO, A. T. & ALLIS, C. D. 1995. Conservation of deposition-
related acetylation sites in newly synthesized histones H3 and H4. *Proc Natl Acad Sci U S A,* 92**,** 1237-
41.

SPARMANN, A. & VAN LOHUIZEN, M. 2006. Polycomb silencers control cell fate, development and cancer. *Nat*
Rev Cancer, 6**,** 846-56.

STAUFFER, D., CHANG, B., HUANG, J., DUNN, A. & THAYER, M. 2007. p300/CREB-binding protein interacts
with ATR and is required for the DNA replication checkpoint. *J Biol Chem,* 282**,** 9678-9687.

STEIN, G., PARK, W., THRALL, C., MANS, R. & STEIN, J. 1975. Regulation of cell cycle stage-specific
transcription of histone genes from chromatin by non-histone chromosomal proteins. *Nature,* 257**,** 764-7.

STEWART-MORGAN, K. R., PETRYK, N. & GROTH, A. 2020. Chromatin replication and epigenetic cell memory.
Nat Cell Biol, 22**,** 361-371.

TAPON, N., MOBERG, K. H. & HARIHARAN, I. K. 2001. The coupling of cell growth to the cell cycle. *Curr Opin*
Cell Biol, 13**,** 731-7.

TESSARZ, P. & KOUZARIDES, T. 2014. Histone core modifications regulating nucleosome structure and
dynamics. *Nat Rev Mol Cell Biol,* 15**,** 703-8.

TIE, F., BANERJEE, R., FU, C., STRATTON, C. A., FANG, M. & HARTE, P. J. 2016. Polycomb inhibits histone
acetylation by CBP by binding directly to its catalytic domain. *Proc Natl Acad Sci U S A,* 113**,** E744-53.

TIE, F., BANERJEE, R., STRATTON, C. A., PRASAD-SINHA, J., STEPANIK, V., ZLOBIN, A., DIAZ, M. O.,
SCACHERI, P. C. & HARTE, P. J. 2009. CBP-mediated acetylation of histone H3 lysine 27 antagonizes
Drosophila Polycomb silencing. *Development,* 136**,** 3131-41.

TILLHON, M., CAZZALINI, O., NARDO, T., NECCHI, D., SOMMATIS, S., STIVALA, L. A., SCOVASSI, A. I. &
PROSPERI, E. 2012. p300/CBP acetyl transferases interact with and acetylate the nucleotide excision
repair factor XPG. *DNA Repair (Amst),* 11**,** 844-52.

TROUTH, A., RAVICHANDRAN, K., GAFKEN, P. R., MARTIRE, S., BOYLE, G. E., VERONEZI, G. M. B., LA, V.,
NAMCIU, S. J., BANASZYNSKI, L. A., SARTHY, J. F. & RAMACHANDRAN, S. 2025. The length of the
G1 phase is an essential determinant of H3K27me3 landscapes across diverse cell types. *PLoS Biol,* 23**,**
e3003119.

TROVATO, M., PATIL, V., GEHRE, M. & NOH, K. M. 2020. Histone Variant H3.3 Mutations in Defining the
Chromatin Function in Mammals. *Cells,* 9.

VERREAULT, A., KAUFMAN, P. D., KOBAYASHI, R. & STILLMAN, B. 1998. Nucleosomal DNA regulates the
core-histone-binding subunit of the human Hat1 acetyltransferase. *Curr Biol,* 8**,** 96-108.

WENGER, A., BIRAN, A., ALCARAZ, N., REDO-RIVEIRO, A., SELL, A. C., KRAUTZ, R., FLURY, V., REVERON-
GOMEZ, N., SOLIS-MEZARINO, V., VOLKER-ALBERT, M., IMHOF, A., ANDERSSON, R., BRICKMAN,
J. M. & GROTH, A. 2023. Symmetric inheritance of parental histones governs epigenome maintenance
and embryonic stem cell identity. *Nat Genet,* 55**,** 1567-1578.

WILLNOW, P. & TELEMAN, A. A. 2024. Nuclear position and local acetyl-CoA production regulate chromatin state.
Nature, 630**,** 466-474.

YAO, T. P., OH, S. P., FUCHS, M., ZHOU, N. D., CH'NG, L. E., NEWSOME, D., BRONSON, R. T., LI, E.,
LIVINGSTON, D. M. & ECKNER, R. 1998. Gene dosage-dependent embryonic development and
proliferation defects in mice lacking the transcriptional integrator p300. *Cell,* 93**,** 361-72.

ZANG, Y., YOSHIMOTO, M. & IGAKI, T. 2025. Programmed cell senescence is required for sensory organ
development in Drosophila. *iScience,* 28**,** 112048.

ZEE, B. M., BRITTON, L. M., WOLLE, D., HABERMAN, D. M. & GARCIA, B. A. 2012. Origins and formation of
histone methylation across the human cell cycle. *Mol Cell Biol,* 32**,** 2503-14.

ZENK, F., CARDAMONE, F., IBARRA MORALES, D. A., ATINBAYEVA, N., ZHAN, Y. & IOVINO, N. 2023.
Analyzing the Genome-Wide Distribution of Histone Marks by CUT&Tag in Drosophila Embryos. *Methods*
Mol Biol, 2655**,** 1-17.

ZHOU, C. Y. & HEALD, R. 2020. Emergent properties of mitotic chromosomes. *Curr Opin Cell Biol,* 64**,** 43-49.

ZIELKE, N., KORZELIUS, J., VAN STRAATEN, M., BENDER, K., SCHUHKNECHT, G. F. P., DUTTA, D., XIANG,
J. & EDGAR, B. A. 2014. Fly-FUCCI: A versatile tool for studying cell proliferation in complex tissues. *Cell*
Rep, 7**,** 588-598.
1033

1034

1035   **Table S1 Experimental strains, reagents and software**

1036

Tool	Source	ID
w^{118}	David Bilder, University of California, Berkeley	FBal0018157
UAS-GFP^{S56T}	Bloomington *Drosophila* Stock Center	BDSC: 1521
rn$^{GAL4-DeltaS}$, tubGAL80^{ts}	Bloomington *Drosophila* Stock Center	BDSC: 8142; 7018 recombinant
rn^{GAL4-5}, UAS-egr, tubP-GAL80^{ts}	Iswar Hariharan, University of California, Berkeley	BDSC: 8142; 7018 recombinant
'E2F1-FUCCI, CycB-FUCCI'; Ubi-GFP.E2f1$^{1-230}$, Ubi-mRFP1.NLS.CycB$^{1-266}$	Bloomington *Drosophila* Stock Center	BDSC: 55123
UAS-Cdk1 RNAi [TRiP.HMS01531]	Bloomington *Drosophila* Stock Center	BDSC: 36117
If/CyO-GFP; UAS-E2F, UAS-DP/TM6b	Laura Buttitta, University of Michigan	PMID: 9657151
UAS-pnt.P1	Bloomington *Drosophila* Stock Center	BDSC: 869
UAS-E(z)-RNAi (TRiP.HMS00066)	Bloomington *Drosophila* Stock Center	BDSC: 33659
UAS-nejire-RNAi(TRiP.HMS01507)	Bloomington *Drosophila* Stock Center	BDSC: 37489
UAS-InR-DA (UAS-InR.A1325D)	Bloomington *Drosophila* Stock Center	BDSC: 8263
'TRE-RFP': TRE-DsRed.T4	Dirk Bohmann, University of Rochester Medical Center	PMID: 22509270
UAS-nejire	Bloomington *Drosophila* Stock Center	BDSC: 32573
UAS-HDAC1-RNAi (TRiP.HMS00607)	Bloomington *Drosophila* Stock Center	BDSC: 33725
CBP-GFP	Melissa Harrison, University of Wisconsin School of Medicine and Public Health	PMID: 40441155
w; Df(2L)HisC FRT40A/CyO, twi-GAL4, UAS-GFP; 6xHisGU^{H3K27R}/6xHisGU^{H3K27R}	Jürg Müller, Max Planck Institute of Biochemistry	PMID: 23393264

yw hs-flp122; hs-nGFP FRT40A; 6xHisGU$^{H3-K27R}$/SM5^TM6B	Jürg Müller, Max Planck Institute of Biochemistry	PMID: 23393264
hs-flp122; FRT40A ubi-GFP/CyO (GFP); Dr/TM6c		
UAS-bsk[DN]	Bloomington *Drosophila* Stock Center	BDSC: 6409
Chicken anti-GFP	Abcam	Cat. #: ab13970
Rabbit monoclonal anti-GFP	Invitrogen	Cat. #: G10362
Rat monoclonal anti-RFP	ChromoTek	Cat. #: 5F8 RRID: AB_2336064
Rabbit polyclonal anti-H3K9ac	Active motif	Cat. #: 39017 RRID: AB_2616593
Rabbit polyclonal anti-H4K8ac	Active motif	Cat. #: 61104 RRID: AB_2793506
Rabbit polyclonal anti-acetyl Lysine	Abcam	Cat. #: ab80178
Rabbit monoclonal anti-crotonyl Histone H3	PTMab	Cat. #: PTM-517RM
Rabbit polyclonal anti-H3K9me1me2me3	Active motif	Cat. #: 39242 RRID: AB_2793200
Rabbit polyclonal anti-H3K18ac	Active motif	Cat. #: 39756 RRID: AB_2714186
Mouse monoclonal anti-H3K27ac	Active motif	Cat. #: 39085 RRID: AB_2793305
Rabbit polyclonal anti-H3K4me3	Abcam	Cat. #: ab8580
Mouse monoclonal anti-H3K27me3	Abcam	Cat. #: ab6002
Rabbit polyclonal anti-H3K9me3	Abcam	Cat. #: ab8898
Rabbit polyclonal anti-H3	Abcam	Cat. #: ab1791
Rabbit polyclonal anti-Histone H2AvD phosphoS137	Rockland	Cat. #: 600-401-914
Mouse monoclonal anti-Cut	DSHB	Cat. #: 2B10 RRID: AB_528186
Mouse monoclonal anti-Engrailed/Invected	DSHB	Cat. #: 4D9 RRID: AB_528224
Mouse monoclonal anti-Nubbin	DSHB	Cat. #: 2D4 RRID: AB_2722119
Mouse monoclonal anti-Antp	DSHB	Cat. #: 8C11 RRID: AB_528083
Mouse monoclonal anti-Ptc	DSHB	Cat. #: Apa1

		RRID: AB_528441
Mouse monoclonal anti Abd-B	DSHB	Cat. #: 1A2E9 RRID: AB_528061
Mouse monoclonal anti-Wg	DSHB	Cat. #: 4D4 RRID: AB_528512
Mouse monoclonal anti-Histone H3 (for CUT&Tag)	Active motif	Cat. #: 39763 RRID: AB_2650522
Rabbit polyclonal anti-H3K27ac (for CUT&Tag)	Diagenode	Cat. #: C15410196
Rabbit polyclonal anti-H3K27me3 (for CUT&Tag)	Diagendoe	Cat. #: C15410195
Rabbit monoclonal anti-H3K9me3 (for CUT&Tag)	Abcam	Cat. #: ab176916
DAPI	Sigma Aldrich	Cat #: D9564
Goat anti-mouse IgG Alexa Flour 488	Invitrogen	Cat. #: A-11001
Goat anti-chicken IgY Alexa Flour 488	Invitrogen	Cat. #: A-11039
Goat anti-rabbit IgG Alexa Flour 488	Invitrogen	Cat. #: A-11008
Goat anti-mouse IgG Alexa Flour 555	Invitrogen	Cat. #: A-21422
Goat anti-rat IgG Alexa Flour 555	Invitrogen	Cat. #: A-21434
Goat anti-mouse Alexa Flour 647	Invitrogen	Cat. #: A-21235
Donkey anti-mouse Alexa Flour 647 preabsorbed	Abcam	Cat. #: ab150111
Goat anti-rat Alexa Flour 647	Invitrogen	Cat. #: A-21247
Goat anti-rabbit Alexa Flour 647	Invitrogen	Cat. #: A-21244
Click-iT Plus EdU Alexa Fluor 647 Imaging Kit	Invitrogen	Cat. #: C10640
Click-iT Plus OPP Alexa Fluor 647 Imaging Kit	Invitrogen	Cat. #: C10458
CellEvent Senescence Green Detection Kit	Invitrogen	Cat. #: C10850
FIJI (ImageJ 1.54f)	(Schindelin et al., 2012)	https://fiji.sc/
GraphPad Prism	Graph Pad	version 9.5.0
Affinity Designer	Affinity	Up to version 2.6.3
Biorender	Science Suite Inc. DBA BioRender	Biorender

1037

1038

1039

1040

	MAIN FIGURE LEGENDS

	Figure 1. H3K27ac, H3K27me3, and H3K9me3 levels respond to changes in cell cycle dynamics

	Icons represent the experimentally verified FUCCI cell cycle status in each condition: G2-phase arrest
	in *Cdk1-RNAi*-expressing discs, either G1 or G2-phase arrest in *pntP1*-expressing discs, and cell cycle
	acceleration in *Dp, E2F1*-overexpressing discs. For better guidance through subsequent figures,
	schemes already introduce FUCCI reporter read-outs for phase G1 (green), early S-phase (grey), late
	S-phase (red) and G2 (yellow).

	A-D. Immunostaining for H3K18ac in control **(A)**, *Cdk1-RNAi*-expressing **(B)**, *pntP1*-expressing **(C)** and
	Dp, E2F1-expressing **(D)** discs. Expression was induced for 24 h in the wing pouch using *rn-GAL4*. The
	rn-GAL4 expressing pouch domain is represented by the tissue above the yellow line (rn^{ts}>). The tissue
	below the yellow line represents wild type tissues of the hinge.

	E-T. Immunostaining for H3K27ac **(E-H)**, H3K4me3 **(I-L)**, H3K27me3 **(M-P)**, H3K9me3 **(Q-T)** in control
	(E,I,M,Q), *Cdk1-RNAi*-expressing **(F,J,N,R)**, *pntP1*-expressing **(G,K,O,S)** and *Dp, E2F1*-expressing
	(H,L,P,T) discs.

	U. Quantification of relative signal intensities for H3K18ac, H3K27ac, H3K4me3, H3K27me3, and
	H3K9me3, presented as pouch-to-hinge ratios (i.e., rn^{ts}-to-wild-type-cell ratio) in *Cdk1-RNAi*-expressing
	discs. Mean and 95% CI is shown. Statistical significance was tested using two-tailed Mann-Whitney
	test, p value= 0.9591 for H3K18ac (control discs: n=8, *Cdk1-RNAi*-expressing discs: n=8); two-tailed
	Unpaired t test, p value=0.0012 for H3K27ac (control discs: n=6, *Cdk1-RNAi*-expressing discs: n=6);
	two-tailed Unpaired t test, p value=0.7334 for H3K4me3 (control discs: n=12, *Cdk1-RNAi*-expressing
	discs: n=11); two-tailed Unpaired t test, p value<0.0001 for H3K27me3 (control discs: n=9, *Cdk1-RNAi*-
	expressing discs: n=9); two-tailed Mann-Whitney test, p value=0.0473 for H3K9me3 (control discs:
	n=11, *Cdk1-RNAi*-expressing discs: n=11).

	V. Quantification of relative signal intensities for H3K18ac, H3K27ac, H3K4me3, H3K27me3, and
	H3K9me3, presented as pouch-to-hinge ratios (i.e., rn^{ts}-to-wild-type-cell ratio) in *pntP1*-expressing
	discs. Mean and 95% CI is shown. Statistical significance was tested using two-tailed Mann-Whitney
	test, p value= 0.5671 for H3K18ac (control discs: n=10, *pntP1*-expressing discs: n=15); two-tailed
	Welch's t test, p value=0.0031 for H3K27ac (control discs: n=6, *pntP1*-expressing discs: n=6); two-
	tailed Unpaired t test, p value=0.1614 for H3K4me3 (control discs: n=11, *pntP1*-expressing discs:
	n=14); two-tailed Unpaired t test, p value<0.0001 for H3K27me3 (control discs: n=9, *pntP1*-expressing
	discs: n=9); two-tailed Unpaired t test, p value=0.0002 for H3K9me3 (control discs: n=8, *pntP1*-
	expressing discs: n=11).

W. Quantification of relative signal intensities for H3K18ac, H3K27ac, H3K4me3, H3K27me3, and
H3K9me3, presented as pouch-to-hinge ratios (i.e., rn^{ts}-to-wild-type-cell ratio) in *Dp, E2F1-*
coexpressing discs. Mean and 95% CI is shown. Statistical significance was tested using two-tailed
Unpaired t test, p value= 0.0609 for H3K18ac (control discs: n=9, *Dp, E2F1*-coexpressing discs: n=9);
two-tailed Unpaired t test, p value=0.0019 for H3K27ac (control discs: n=7, *Dp, E2F1*-coexpressing
discs: n=6); two-tailed Unpaired t test, p value=0.4837 for H3K4me3 (control discs: n=12, *Dp, E2F1-*
coexpressing discs: n=11); two-tailed Unpaired t test, p value=0.0003 for H3K27me3 (control discs:
n=9, *Dp, E2F1*-coexpressing discs: n=9); two-tailed Mann-Whitney test, p value<0.0001 for H3K9me3
(control discs: n=10, *Dp, E2F1*-coexpressing discs: n=10).

1090

1091 Fluorescence intensities are reported as arbitrary units.
1092 Sum projections of multiple confocal sections are shown in **C**, **K**, **M-N**, **O** and **Q-R.**
1093 Maximum projections of multiple confocal sections are shown in **A-B**, **E-F**, **G**, **H**, **I-J**, **P**, **S** and **T**.
1094 Scale bars: 100 μm.
1095 Please refer to **(Fig S1.3A)** for corresponding DAPI images of all discs shown.

1096

1097

1098 **Figure 2. Cell cycle-dependent changes in H3K27ac, H3K27me3, and H3K9me3 occur at**
1099 **preexisting genomic sites**

1100

A-B. Two examples of genomic snapshots of CUT&Tag profiles comparing levels of H3K27ac,
H3K27me3 and H3K9me3 in wild type wing discs and in wing discs having expressed *Cdk1*-RNAi for
24 h in the pouch region under the control of rn-GAL4. Tracks for H3K27ac (orange), H3K27me3
(green), and H3K9me3 (red) were normalized against H3 as described in the Materials and Methods
and were then visualized using IGV. Black tracks represent the change between control and knockdown
conditions computed by subtracting number of sequence reads in *Cdk1-RNAi*-expressing discs from
wild type discs for each histone modification. Tracks in **(B)** cover the BX-C Hox gene cluster encoding
Ubx, Abd-A and Abd-B targeted by Polycomb silencing.

1109

1110 **C-F.** Profile plots and heatmaps showing normalized CUT&Tag signals at peaks shared between wild
1111 type control and *Cdk1-RNAi*-expressing discs for H3 **(C)**, H3K27ac **(D)**, H3K27me3 **(E)**, and H3K9me3
1112 **(F)** (see Materials and Methods). Averaged and individual signal intensities are centered ±5 kb around
1113 the shared peak.

1114

1115

Figure 3. Developmentally arrested cell populations under physiological conditions exhibit changes in H3K27ac, H3K27me3, and H3K9me3 levels

A-C. Immunostaining for H3K27ac **(A)**, H3K27me3 **(B)**, and H3K9me3 **(C)** in the zone of non-proliferating cells (ZNC) of developing wing discs. Cells in the ZNC (white brackets) are arrested in either G1 or G2 phase, and can be identified by the Fly-FUCCI reporter. Among the histone modifications examined, only H3K27ac, H3K27me3, and H3K9me3 exhibit cell cycle arrest associated changes in the ZNC.

D-F. Immunostaining for H3K27ac **(D)**, H3K27me3 **(E)** and H3K9me3 **(F)** in a hinge cell population undergoing programmed senescence in developing wing discs.

Discs were stained with DAPI to visualize nuclei.
Scale bars: 50 µm

Figure 4. Metabolic state sets the rate of H3K27 and H3K9 methylation

A-B. Control wing disc expressing GFP (magenta) for 24 h in the wing pouch domain under the control of *rn-GAL4* **(A)**. Wing disc expressing GFP and a constitutively active Insulin receptor (*InR-DA*) under the same conditions **(B)**. GFP expression visualizes the enlarged wing pouch domain in *InR-DA* expressing discs.

C. Quantification of pouch size area in control and *InR-DA*-expressing discs. Mean and 95% CI is shown. Statistical significance was tested using two-tailed Unpaired t test, p value=0.0432. (control discs: n=6, *InR-DA*-expressing discs: n=6).

D-G. Immunostaining for H3K27me3 in control **(D)**, *InR-DA*-expressing **(E)**, *Cdk1-RNAi*-expressing **(F)** and *InR-DA, Cdk1-RNAi*-coexpressing **(G)** discs.

H-K. Immunostaining for H3K9me3 in control **(H)**, *InR-DA*-expressing **(I)**, *Cdk1-RNAi*-expressing **(J)** and *InR-DA, Cdk1-RNAi*-coexpressing **(K)** discs.

L-O. Immunostaining for H3K4me3 in control **(L)**, *InR-DA*-expressing **(M)**, *Cdk1-RNAi*-expressing **(N)** and *InR-DA, Cdk1-RNAi*-coexpressing **(O)** discs. Expression was induced for 24 h in the wing pouch using *rn-GAL4.* Yellow dashed lines indicate the boundary between the wing pouch and hinge regions. The area above the line corresponds to the *rn-GAL4* expression domain, while the area below represents wild-type cells.

P. Illustration of the dual role of metabolism in regulating histone methylation levels. Metabolism contributes to the dilution of histone PTMs by accelerating the cell cycle, thereby increasing the frequency of DNA replication events. Simultaneously, it supports the re-establishment of these PTMs by supplying essential precursors and cofactors required for methyltransferase activity.

Q. Hypothetical model depicting the relationship between histone methylation levels, metabolic rate, and cell division rate. When metabolic activity and cell division are proportionally coupled (black diagonal), histone methylation levels are maintained at stable levels. However, when metabolism outpaces cell division, methylation accumulates.

Discs were stained with DAPI to visualize nuclei.
Sum projections of multiple confocal sections are shown in **A-B.**
Maximum projections of multiple confocal sections are shown in **D-K.**
Scale bars: 100 µm.

Figure 5. CBP/nejire-mediated H3K27 acetylation peaks during S phase

A-B. Immunostaining for H3K27ac in a control disc expressing the FUCCI reporters *GFP-E2F1*$^{1\text{-}230}$ (green) and *mRFP-NLS-CycB*$^{1\text{-}266}$ (red) to visualize cell cycle phases **(A).** Magnified region in **(B)** is demarcated by a white dashed square in **(A).**

C. Quantification of normalized H3K27ac intensity level in different cell cycle phases in the pouch region of control discs. Normalization was performed against the average fluorescence intensity across all cell cycle phases. Mean and 95% CI is shown. Statistical significance was tested using Repeated Measures One-Way ANOVA followed by Tukey's post-hoc test for multiple comparison (n=11 discs). Adjusted $p<0.0001$ for G1 vs. Early S; $p=0.0838$ for G1 vs. Late S; $p<0.0001$ for G1 vs. G2; $p<0.0001$ for Early S vs. Late S; $p<0.0001$ for Early S vs G2 and $p<0.0001$ for Late S vs G2.

D-E. Immunostaining for H3K27ac in control **(D)** and *nejire-RNAi*-expressing **(E)** discs. Expression was induced for 24 h in the wing pouch using *rn-GAL4.* Yellow dashed lines indicate the boundary between the wing pouch and hinge regions.

F-G. Immunostaining for H3K27ac in control **(F)** and *nejire*-expressing **(G)** discs also expressing the FUCCI reporters *GFP-E2F1*$^{1\text{-}230}$ (green) and *mRFP-NLS-CycB*$^{1\text{-}266}$ (red) to visualize cell cycle phases. Expression of *nejire* was induced for 24 h in the wing pouch using *rn-GAL4.*

H-I. Immunostaining for H3K27ac in control **(H)** and *HDAC1-RNAi*-expressing **(I)** discs also expressing the FUCCI reporters *GFP-E2F1*$^{1\text{-}230}$ (green) and *mRFP-NLS-CycB*$^{1\text{-}266}$ (red) to visualize cell cycle phases. Expression of *HDAC1-RNAi* was induced for 24 h in the wing pouch using *rn-GAL4.*

J. Quantification of mean H3K27ac intensity in the pouch region of control or *nejire*-expressing discs. Expression was induced for 24 h in the wing pouch using *rn-GAL4.* Mean and 95% CI is shown. Statistical significance was tested using two-tailed Welch's t test, p value<0.0001. (control discs: n=9, *nejire*-expressing discs: n=9).

J'. Quantification of normalized H3K27ac in G1 phase and Late S phase in the pouch region of control or *nejire*-expressing discs. Expression was induced for 24 h in the wing pouch using *rn-GAL4.* Mean and 95% CI is shown. Statistical significance was tested using Repeated Measures Two-Way ANOVA followed by with Šidák's post hoc test for multiple comparisons (control discs: n=9, *nejire*-expressing discs: n=9). Adjusted p= 0.9482 for Control G1 vs nejire-OE G1; p=0.0064 for Control Late S vs nejire-OE Late S.

K. Quantification of mean H3K27ac intensity in the pouch region of control or *HDAC1-RNAi*-expressing discs. Expression was induced for 24 h in the wing pouch using *rn-GAL4.* Mean and 95% CI is shown. Statistical significance was tested using two-tailed Mann-Whitney test, p value<0.0001. (control discs: n=10, *HDAC1-RNAi*-expressing discs: n=9).

K'. Quantification of normalized H3K27ac intensity level in G1 and Late S phases in the pouch region of control or *HDAC1-RNAi*-expressing discs. Expression was induced for 24 h in the wing pouch using *rn-GAL4.* Mean and 95% CI is shown. Statistical significance was tested using Repeated Measures Two-Way ANOVA followed by with Šidák's post hoc test for multiple comparisons (control discs: n=10, *HDAC1-RNAi*-expressing discs: n=9). Adjusted p= 0.1907 for Control G1 vs HDAC1-RNAi G1; p<0.0001 for Control Late S vs HDAC1-RNAi Late S.

Fluorescence intensities are reported as arbitrary units.

Sum projections of multiple confocal sections are shown in **D-E.**

Scale bars: 100 µm in **A** and **D-E**; 50 µm in **F-G** and **H-I**; 25 µm in **B**.

Figure 6. CBP/Nejire activity and H3K27 acetylation prevent replication stress during S phase

A-B. EdU incorporation to visualize DNA replication in the pouch region of control **(A)** and *nejire-RNAi*-expressing **(B)** discs. Dashed yellow squares demarcate magnified regions shown in **(A', B').** Expression was induced for 24 h in the wing pouch using *rn-GAL4.*

C. Quantification of mean EdU intensity per EdU area in the pouch region of control and *nejire-RNAi*-expressing discs, serving as a proxy speed of DNA replication. Mean and 95% CI are shown. Statistical significance was tested using two-tailed Mann Whitney test, p value=0.0001 (control discs: n=11, *nejire-RNAi*-expressing discs: n=11).

1235

1236 **C'.** Quantification of EdU area per DAPI area in the pouch region of control and *nejire-RNAi*-expressing

1237 discs, serving as a proxy number of cells undergoing DNA replication. Mean and 95% CI are shown.

1238 Statistical significance was tested using two-tailed Mann Whitney test, p value<0.0001 (control discs:

1239 n=11, *nejire-RNAi*-expressing discs: n=11).

1240

1241 **D-G.** Immunostaining for γH2Av in control **(F)** and *nejire-RNAi*-expressing **(G)** discs. Expression was

1242 induced for 24 h in the wing pouch using *rn-GAL4.* γH2Av serves as a marker of DNA double strand

1243 breaks. DAPI staining was used to visualize the nuclei in control **(D)** and *nejire-RNAi*-expressing **(E)**

1244 discs. The *rn-GAL4* expression domain is represented by the tissue above the yellow line (rn^{ts}>). The

1245 tissue below the yellow line represents wild type tissues of the hinge.

1246

1247 **H.** Quantification of mean γH2Av intensity in the pouch region of control or *nejire-RNAi*-expressing

1248 discs. Mean and 95% CI are shown. Statistical significance was tested using two-tailed Unpaired t test,

1249 p value=0.0126 (control discs: n=10, *nejire-RNAi*-expressing discs: n=10).

1250

1251 **K.** Immunostaining for H3K27ac (red) and γH2Av (cyan) in wing imaginal discs with clones of Δ*HisC*

1252 mutant clones deleting the histone cluster and rescued by a *12×His-GU$^{H3-K27R}$* transgene.

1253

1254 **L.** Illustration of the role of CBP/nej and HDAC1 in modulating levels of H3K27ac during S-phase to

1255 prevent replication stress

1256

1257 Discs were stained with DAPI to visualize nuclei.

1258 Fluorescence intensities are reported as arbitrary units.

1259 Sum projections of multiple confocal sections are shown in **A-B** and **D-G.**

1260 Maximum projections of multiple confocal sections are shown in **K.**

1261 Scale bars: 100 μm in **A-B**, **D-G**, **I-J** and **K.**

1262

     SUPPLEMENTARY FIGURE LEGENDS

     Figure S1.1. Cell cycle manipulation models for assessing histone modification dynamics

     A. A control wing disc after 24 h of *UAS-GFP*-expression in the pouch, under the control of the *rn-GAL4*

     (*rotund-GAL4*) driver. GFP-expression visualizes the tissue domain subject to manipulation throughout

     this study (magenta).

     B. Schematic representation of the Fly-FUCCI cell cycle reporter system, utilizing the degradable *GFP-*

     E2F1$^{1-230}$ (green) and *mRFP-NLS-CycB*$^{1-266}$ (red) as sensors to visualize cell cycle phases. Individual

     fluorophore expression indicates cells in G1 or late S-phase, respectively. Combined expression of both

     GFP and RFP labels cells is observed in early and late G2. Cells in late mitosis and early S-phase lack

     expression of either GFP or RFP. Cells in early S-phase are specifically detected by EdU incorporation

     (Crucianelli et al., 2022; Zielke et al., 2014). Cells in mitosis are positioned in very apical positions of

     the tissue and are not represented in our study.

     C-G. Cell cycle dynamics in control discs **(C, E)** and discs expressing *Cdk1-RNAi* for 24 h under the

     control of *rn-GAL4* **(D, F)**. FUCCI reporters *GFP-E2F1*$^{1-230}$ (green) and *mRFP-NLS-CycB*$^{1-266}$ (red) were

     used to visualize cell cycle phases **(C, D)** and EdU incorporation was used to detect DNA replication in

     S-phase cells **(E, F).** Cdk1 knockdown in the wing pouch domain leads to a cell cycle phase shift toward

     G2 phase **(D)**, and a corresponding loss of EdU incorporation **(F)** combined demonstrating a

     pronounced arrest in G2. **(G)** Quantification of mean EdU intensity per DAPI area in the pouch region

     of control and *Cdk1-RNAi*-expressing discs, serving as a proxy for relative DNA replication activity.

     Mean and 95% CI are shown. Statistical significance was tested using two-tailed Welch's test, p value

     <0.0001 (control discs: n=11, *Cdk1-RNAi*-expressing discs: n=12).

     H-L. Cell cycle dynamics in control discs **(H, J)** and discs expressing *pntP1* **(I, K)** for 24 h under the

     control of *rn-GAL4.* FUCCI reporters *GFP-E2F1*$^{1-230}$ (green) and *mRFP-NLS-CycB*$^{1-266}$ (red) were used

     to visualize cell cycle phases **(H, I)** and EdU incorporation was used to detect DNA replication in S-

     phase cells **(J, K)**. Pointed-P1 overexpression in the wing pouch domain leads to a cell cycle phase

     shift towards either the G1 or the G2 phase **(I)**, correlating with a loss of EdU incorporation **(K)**.

     Combined, this demonstrates a pronounced arrest in either G1 or G2. Mechanistically, the choice for

     either cell cycle phase is not known. **(L)** Quantification of mean EdU intensity per DAPI area in the

     pouch region of control and *pntP1*-expressing discs, serving as a proxy for relative DNA replication

     activity. Mean and 95% CI is shown. Statistical significance was tested using two-tailed Mann-Whitney

     test, p value=0.0043 (control discs: n=6, *pntP1*-expressing discs: n=6).

     M-Q. Cell cycle dynamics in control discs **(M, O)** and discs expressing *Dp, E2F1* **(N, P)** for 24 h under

     the control of *rn-GAL4.* FUCCI reporters *GFP-E2F1*$^{1-230}$ (green) and *mRFP-NLS-CycB*$^{1-266}$ (red) were

     used to visualize cell cycle phases **(M, N)** and EdU incorporation was used to detect DNA replication in

S-phase cells **(O,P).** dDp and dE2F1 co-expression in the wing pouch domain reduces the number of

cells in either G1 or G2 **(N)**, and an increase in S-phase cells as well as an elevation of EdU

incorporation **(P)** in the wing pouch region, confirming frequent entry and rapid progression through the

cell cycle. **(Q)** Quantification of mean EdU intensity per DAPI area in the pouch region of control and

Dp, E2F1-coexpressing discs, serving as a proxy for relative DNA replication activity. Mean and 95%

CI is shown. Statistical significance was tested using two-tailed Welch's t test, p value=0.0069 (control

discs: n=6, *Dp, E2F1*-expressing discs: n=5).

1310

Discs were stained with DAPI to visualize nuclei.

Fluorescence intensities are reported as arbitrary units.

Yellow dashed lines represent the pouch region of wing discs.

Sum projections of multiple confocal sections are shown in **A, E-F, J-K, M-N** and **O-P.**

Maximum projections of multiple confocal sections are shown in **C-D** and **H-I.**

Scale bars: 100 μm.

1317

1318

1319   **Figure S1.2. Histone modifications are largely insensitive to cell cycle manipulations**

1320

A-D. Immunostaining for H3K9ac in control **(A)**, *Cdk1-RNAi*-expressing **(B)**, *pntP1*-expressing **(C)** and

Dp, E2F1-expressing **(D)** discs. Expression was induced for 24 h in the wing pouch using *rn-GAL4*. The

rn-GAL4 expression pouch domain is represented by the tissue above the yellow line (rn^{ts}>). The tissue

below the yellow line represents wild type tissues of the hinge. Icons represent the experimentally

verified FUCCI cell cycle status (see **Fig S1.1**) in each condition: G2-phase arrest in *Cdk1-RNAi*-

expressing discs **(B)**, combined G1 and G2-phase arrest in *pntP1*-expressing discs **(C)**, and cell cycle

acceleration in *Dp, E2F1*-coexpressing discs **(D)**.

1328

1329   **E-H**. Immunostaining for H4K8ac in control **(E)**, *Cdk1-RNAi*-expressing **(F)**, *pntP1*-expressing **(G)** and

1330   *Dp, E2F1*-coexpressing **(H)** discs.

1331

1332   **I-L.** Immunostaining for total acetylated lysines in control **(I)**, *Cdk1-RNAi*-expressing **(J)**, *pntP1*-

1333   expressing **(K)** and *Dp, E2F1*-coexpressing **(L)** discs.

1334

1335   **M-P.** Immunostaining for H3K18crotonylation in control **(M)**, *Cdk1-RNAi*-expressing **(N)**, *pntP1*-

1336   expressing **(O)** and *Dp, E2F1*-coexpressing **(P)** discs.

1337

1338   **Q-T.** Immunostaining for total H3K9me1/2/3 methylation in control **(I)**, *Cdk1-RNAi*-expressing **(J)**,

1339   *pntP1*-expressing **(K)** and *Dp, E2F1*-coexpressing **(L)** discs.

1340

U. Quantification of relative signal intensities for H3K9ac, H4K8ac, total acetylated lysine,

H3K18crotonylation, and H3K9me1/2/3, presented as pouch-to-hinge ratios (rn^{ts}-to-wild-type-cell ratio)

in *Cdk1-RNAi*-expressing discs. Mean and 95% CI is shown. Statistical significance was tested using

two-tailed Unpaired t test, p value= 0.4268 for H3K9ac (control discs: n=5, *Cdk1-RNAi*-expressing discs:

n=5); two-tailed Unpaired t test, p value=0.8936 for H4K8ac (control discs: n=5, *Cdk1-RNAi*-expressing

discs: n=5); two-tailed Unpaired t test, p value=0.5553 for total acetylated lysines (control discs: n=6,

Cdk1-RNAi-expressing discs: n=6); two-tailed Mann-Whitney test, p value=0.4779 for H3K18crot

(control discs: n=11, *Cdk1-RNAi*-expressing discs: n=11); two-tailed Unpaired t test, p value=0.9619 for

H3K9me1/2/3 (control discs: n=9, *Cdk1-RNAi*-expressing discs: n=10).

1350

V. Quantification of relative signal intensities for H3K9ac, H4K8ac, total acetylated lysine,

H3K18crotonylation, and H3K9me1/2/3, presented as pouch-to-hinge ratios (rn^{ts}-to-wild-type-cell ratio)

in *pntP1*-expressing discs. Mean and 95% CI is shown. Statistical significance was tested using two-

tailed Unpaired t test, p value= 0.1975 for H3K9ac (control discs: n=8, *pntP1*-expressing discs: n=10);

two-tailed Unpaired t test, p value=0.2752 for H4K8ac (control discs: n=12, *pntP1*-expressing discs:

n=12); two-tailed Unpaired t test, p value=0.0012 for total acetylated lysines (control discs: n=6, *pntP1*-

expressing discs: n=6); two-tailed Unpaired t test, p value=0.6621 for H3K18crot (control discs: n=7,

pntP1-expressing discs: n=7); two-tailed Mann-Whitney test, p value=0.2403 for H3K9me1/2/3 (control

discs: n=6, *pntP1*-expressing discs: n=6).

1360

W. Quantification of relative signal intensities for H3K9ac, H4K8ac, total acetylated lysine,

H3K18crotonylation, and H3K9me1/2/3, presented as pouch-to-hinge ratios (rn^{ts}-to-wild-type-cell ratio)

in *Dp, E2F1*-coexpressing discs. Mean and 95% CI is shown. Statistical significance was tested using

two-tailed Unpaired t test, p value= 0.0282 for H3K9ac (control discs: n=8, *Dp, E2F1*-coexpressing

discs: n=8); two-tailed Mann-Whitney test, p value=0.0759 for H4K8ac (control discs: n=11, *Dp, E2F1*-

coexpressing discs: n=11); two-tailed Mann-Whitney test, p value=0.2403 for total acetylated lysines

(control discs: n=6, *Dp, E2F1*-coexpressing discs: n=6); two-tailed Mann-Whitney test, p value=0.0728

for H3K18crot (control discs: n=7, *Dp, E2F1*-coexpressing discs: n=7); two-tailed Unpaired t test, p

value=0.0154 for H3K9me1/2/3 (control discs: n=10, *Dp, E2F1*-coexpressing discs: n=10).

1370

1371   Fluorescence intensities are reported as arbitrary units.

1372   Sum projections of multiple confocal sections are shown in **C-T.**

1373   Scale bars: 100 µm.

1374   Please refer to **(Fig S1.3B)** for corresponding DAPI images of all discs shown.

1375

1376   **Figure S1.3 Corresponding DAPI images for Fig 1 and Fig S1.2**

1377

1378   Representation of DAPI images matched to the images of histone modifications in control, *Cdk1-RNAi*-

1379   expressing, *pntP1*-expressing and *Dp, E2F1*-coexpressing wing discs shown in Fig 1 and Fig S1.2.

1380

1381   Discs were stained with DAPI to visualize nuclei.

1382   Sum and max projections of multiple confocal sections are shown.

1383   Scale bars: 100 μm.

1384

1385

1386   **Figure S2.1. Comparative analysis of Histone H3 and H3K27ac, H3K27me3, and H3K9me3**
1387   **histone modifications by CUT&Tag in control and Cdk1 RNAi- expressing discs**

1388

A-D. MA plots displaying differential enrichment of histone modifications analysed by by CUT&Tag for
H3, H3K27ac, H3K27me3, and H3K9me3 (in rows). Each point represents a shared or unique genomic
peak identified by our analysis in control and Cdk1-RNAi expressing discs (shared peaks **(A)**, unique
peaks in control **(B)**, unique peaks in cdk1-RNAi **(C)**), or a 500 bp genomic fragment not called as any
peak in our analysis (other genomic regions **(D)**). The x-axis displays the mean normalized read count
on a log10 scale, representing an average signal intensity across samples or conditions. The y-axis
displays the log2 fold change (log2 ratio of read counts) between the Cdk1-RNAi and control conditions,
reflecting the magnitude and direction of differential enrichment. Peaks are color-coded by modification
type: blue for H3, orange for H3K27ac, green for H3K27me3, and red for H3K9me3. Regions with an
adjusted p-value<0.1 are colored accordingly, highlighting statistically significant differential peaks; non-
significant peaks (ns, adjusted p-value ≥0.1) are shown in gray.

1400

E. Each subplot depicts the distribution of log2 fold change values (Cdk1-RNAi compared to control)
for genomic regions categorized as Shared (peaks detected in both conditions), Control-Unique (peaks
specific to control), Cdk1-RNAi-Unique (peaks specific to Cdk1-RNAi expressing discs), and Other
(genomic fragments of 500bp not included in any called peaks). The x-axis indicates these categorical
groups, and the y-axis shows the log2 fold change of normalized counts between Cdk1-RNAi and
control samples. Each point represents an individual genomic region, colored by histone mark: blue
(H3), orange (H3K27ac), green (H3K27me3), and red (H3K9me3). Black horizontal bars indicate the
mean log2 fold change within each group. Peaks are color-coded by modification type: blue for H3,
orange for H3K27ac, green for H3K27me3, and red for H3K9me3. Regions with an adjusted p-
value<0.1 are colored accordingly, highlighting statistically significant differential peaks; non-significant
peaks (ns, adjusted p-value ≥0.1) are shown in gray.

1412

1413

1414   **Figure S2.2. Cell cycle–dependent fluctuations of H3K27ac and H3K27me3 are not due to**
1415   **competitive residue occupancy**

1416

1417   **A-B.** H3K27me3 staining in control **(A)** or *E(z)-RNAi*-expressing **(B)** discs. Knockdown of the
1418   H3K27me3 writer E(z) causes a reduction in H3K27me3 level in the pouch of the wing disc.

1419

1420   **C-D.** H3K27ac staining in control **(C)** or *E(z)-RNAi*-expressing **(D)** discs. Knockdown of the H3K27me
1421   writer E(z) causes no corresponding increase in H3K27ac level.

1422

E-F. H3K27ac staining in control **(E)** and *nejire-RNAi*-expressing **(F)** discs. Knockdown of the H3K27ac writer CBP/nej causes a reduction of H3K27ac level in the pouch of the wing disc.

G-H. H3K27me3 staining in control **(G)** and *nejire-RNAi*-expressing **(H)** discs. Knockdown of the H3K27ac writer CBP/nej causes no corresponding increase in H3K27me3 level.

Discs were stained with DAPI to visualize nuclei.

Maximum projections of multiple confocal sections are shown in **A-B.**

Sum projections of multiple confocal sections are shown in **C-D, E-F** and **G-H.**

Scale bars: 100 μm.

Figure S3.1. Visualization of cell cycle-arrested cell populations and histone modifications in developing wing imaginal disc

---

A. Schematic representation of the zone of non-proliferation cells (ZNC) (adapted from (Johnston and Edgar, 1998)).

B-C. The Fly-FUCCI system and EdU incorporation assay were used to visualize the spatial pattern of cell cycle arrest within the ZNC. In the anterior ZNC, central margin cells arrest in G1 phase, while adjacent cells arrest in G2. Cells in the posterior ZNC uniformly arrest in G1. The entire ZNC region also lacks EdU incorporation, indicating absence of S-phase entry. White arrowheads indicate the ZNC.

D. Elevated SA-β-gal activity marks two clusters of cell cycle–arrested cells undergoing programmed senescence (indicated by yellow arrowheads) in developing wing discs.

E,F. Immunostaining for H3K18ac **(E)** and H3K4me3 **(F)** in the zone of non-proliferation cells (ZNC) of developing wing discs. Neither histone modification shows detectable changes in this region.

G,H. Immunostaining for H3K18ac **(G)** and H3K4me3 **(H)** in senescent cell populations of the hinge in developing wing discs. Neither histone modification shows detectable changes in this region.

Discs were stained with DAPI to visualize nuclei.

Maximum projection of multiple confocal sections is shown in **C.**

Sum projections of multiple confocal sections are shown in **A-B and D.**

Scale bars: 100 μm in **B-D**; 50 μm in **E-H.**

Figure S3.2. Polycomb target gene expression remains stable despite altered cell cycle dynamics

---

1463

1464   **A-L.** Immunostaining for proteins expressed by Polycomb target genes: Cut **(A, B),** Engrailed/invected

1465   **(C, D),** Nubbin **(E, F),** Antennapedia **(G, H),** Patched **(I, J)** and Wingless **(K, L)** in control **(A, C, E, G, I,**

1466   **K)** and *Cdk1-RNAi*-expressing **(B, D, F, H, J, L)** wing discs.

1467

1468   **M-R.** Immunostaining for proteins expressed by Polycomb target genes: Antennapedia **(M, N),**

1469   Engrailed/**invected (O, P)** and Patched **(Q, R)** in control **(M, O, Q)** and *Dp, E2F1*-coexpressing **(N, P,**

1470   **R)** wing discs.

1471

1472   **S.** A control wing disc after 24 h of *UAS-GFP*-expression in the pouch (magenta), under the control of

1473   the *rn-GAL4* (*rotund-GAL4*) driver, displayed to reference the manipulated pouch domain in Figures A-

1474   R. Please note that this is the same disc shown in **Fig S1.1 A**.

1475

1476   Discs were stained with DAPI to visualize nuclei.

1477   Sum projections of multiple confocal sections are shown in **A-S.**

1478   Scale bars: 100 µm.

1479

1480

1481   **Figure S4.1. Increased insulin signaling promotes faster proliferation and growth independent**

1482   **of changes to the distribution of cell cycle phases**

1483

1484   **A-B.** Protein synthesis visualized by OPP incorporation in control **(A)** and *InR-DA*-expressing **(B)** discs.

1485   Yellow dashed lines indicate the boundary between the wing pouch and hinge regions. The area above

1486   the line corresponds to the *rn-GAL4* expression domain, while the area below represents wild-type cells.

1487   The *InR-DA*-expressing wing pouch exhibits higher rates of protein synthesis, reflecting an increased

1488   biosynthetic capacity promoted by higher InR/PI3K/Akt activity.

1489

1490   **C.** Quantification of mean OPP intensity in the pouch region of control and *InR-DA*-expressing discs.

1491   Mean and 95% CI is shown. Statistical significance was tested using two-tailed Unpaired t test, p

1492   value<0.0001. (control discs: n=5, *InR-DA*-expressing discs: n=6).

1493

1494   **D-E**. EdU incorporation visualizes DNA replication in the pouch region of control **(D)** and *InR-DA*-

1495   expressing **(E)** discs. Dashed yellow squares highlight the magnified regions shown in **(D',E')**. *InR-DA*-

1496   expressing wing pouch shows higher levels of EdU intensities, reflecting accelerated speed of DNA

1497   replication. Rates of EdU incorporation in *InR-DA*-expressing discs were previously reported and

1498   quantified (Crucianelli et al., 2022).

1499

1500   **F-G.** Cell cycle dynamics in control **(F)** and *InR-DA*-expressing **(G)** discs. FUCCI reporters *GFP-E2F1*$^{1-}$

1501 ^{230} (green) and *mRFP-NLS-CycB*$^{1-266}$ (red) were used to visualize cell cycle phases. Note that

1502   expression of a constitutively active Insulin receptor does not alter the cell cycle phase profile of the

1503   wing pouch compared to control disc.

1504

1505   Discs were stained with DAPI to visualize nuclei.

1506   Fluorescence intensities are reported as arbitrary units.

1507   Maximum projections of multiple confocal sections are shown in **F-G.**

1508   Sum projections of multiple confocal sections are shown in **A-B** and **D-E.**

1509   Scale bars: 100 µm in **A,B**, **D,E**, **F,G;** 50 µm in zoomed-in **D,E.**

1510

1511

1512   **Figure S4.2. Persistence methyltransferase activity drives H3K9me3 accumulation through**

1513   **metabolic changes coupled to a senescent cell cycle arrest**

1514

1515   **A-B.** Senescence-associated β-galactosidase (SA-β-gal) activity (cyan or gray) in control **(A)** and *egr-*

1516   expressing **(B)** discs. TRE-RFP reporter (magenta) visualizes JNK-pathway activity. SA-β-gal activity

1517   in *egr*-expressing discs was previously reported and quantified (Jaiswal et al., 2023).

1518

1519   **C-D. EdU incorporation (gray or cyan) visualizes DNA replication in control (B) and *egr-***

1520   **expressing (D) discs. TRE-RFP reporter (magenta) visualizes JNK-pathway activity.** EdU

1521   incorporation was previously reported and quantified (Jaiswal et al., 2023, Floc'hlay et al., 2023) .

1522

1523   **E-F.** Cell cycle dynamics in control **(E)** and *egr*-expressing **(F)** discs*.* FUCCI reporters *GFP-E2F1*[1-230]

1524   *(mRFP-NLS-CycB*[1-266] (red) were used to visualize cell cycle phases.

1525

G. A model illustrating inflammatory tissue damage in *Drosophila* wing imaginal discs. Eiger expression

activates the JNK pathway, an early stress response, which induces a senescence-like cell-cycle arrest

in the G2 phase at center of the tissue damage and triggers a senescence program (Cosolo et al.,

2019). These arrested cells secrete Unpaired cytokines, which activate the JAK/STAT pathway in

surrounding cells, promoting compensatory proliferation during tissue regeneration (Crucianelli et al.,

2022; Floc'hlay et al., 2023; Herrera & Bach, 2019; Jaiswal et al., 2023; La Fortezza et al., 2016;

Santabarbara-Ruiz et al., 2015; Worley et al., 2022). At the periphery of the damage site, cells remain

in a quiescent state (Maya et al., 2024).

1534

1535   **H-I.** Immunostaining for H3K9me3 in *egr*-expressing disc **(H).** The magenta frame highlights G2-

1536   arrested senescent cells, which show elevated H3K9me3 levels. The cyan frame marks cells

1537   undergoing compensatory proliferation, where H3K9me3 levels are reduced. The yellow frame

1538   indicates quiescent cells, which have higher H3K9me3 levels than proliferating cells **(I).**

1539

1540   Discs were stained with DAPI to visualize nuclei.

1541   Maximum projections of multiple confocal sections are shown in **H.**

1542   Sum projections of multiple confocal sections are shown in **A-D** and **E-F.**

1543   Scale bars: 100 µm in **A-B**, **C-D**, **E-F** and **H**; 10 µm in **I**.

1544

1545

Figure S5.1. Total H3 and acetyl-lysine dynamics do not recapitulate the S phase-associated

increase in H3K27ac

____

A-B. Immunostaining for H3 in a developing control disc **(A)**, with a magnified region (demarcated by a

white dashed square) shown in panel B, also expressing the FUCCI reporter (*GFP-E2F1*$^{1-230}$ (green)

and *mRFP-NLS-CycB*$^{1-266}$ (red)) to visualize cell cycle phases.

1552

C. Quantification of normalized H3 intensity level with respect to different cell cycle phases in the pouch

region of control discs. Please note that normalization was performed against the average fluorescence

intensity across all cell cycle phases. Mean and 95% CI is shown. Statistical significance was tested

using Repeated Measures One-Way ANOVA followed by Tukey's post-hoc test for multiple comparison

(n=11 discs). Adjusted p= 0.0192 for G1 vs. Early S; p<0.0001 for G1 vs. Late S; p<0.0001 for G1 vs.

G2; p<0.0001 for Early S vs. Late S; p<0.0001 for Early S vs G2 and p=0.4147 for Late S vs G2.

1559

1560   **D-E.** Immunostaining for total acetylated lysine level in a developing control disc **(D),** with a magnified

1561   region (demarcated by a white dashed square) in panel E, also expressing the FUCCI reporters (*GFP-*

1562   *E2F1*$^{1-230}$ (green) and *mRFP-NLS-CycB*$^{1-266}$ (red)) to visualize cell cycle phases.

1563

F. Quantification of normalized total acetylated lysine level in different cell cycle phases in the pouch

region of control discs. Please note that normalization was performed against the average fluorescence

intensity across all cell cycle phases. Mean and 95% CI is shown. Statistical significance was tested

using Repeated Measures One-Way ANOVA followed by Tukey's post-hoc test for multiple comparison

(n=13 discs). Adjusted p<0.0001 for G1 vs. Early S; p<0.0001 for G1 vs. Late S; p<0.0001 for G1 vs.

G2; p<0.0001 for Early S vs. Late S; p<0.0001 for Early S vs G2 and p<0.0001 for Late S vs G2.

1570

1571   **G-H.** Immunostaining for H3K27ac in control **(G)** and *nejire*-expressing **(H)** discs, also expressing the

1572   FUCCI reporters (*GFP-E2F1*$^{1-230}$ (green) and *mRFP-NLS-CycB*$^{1-266}$ (red)) to visualize cell cycle phases.

1573   The location of magnified regions shown in Main Figure 5F and G is indicated by white dashed squares.

1574

1575   **I-J.** Immunostaining for H3K27ac in control **(I)** and *HDAC1-RNAi*-expressing **(J)** discs also expressing

1576   the FUCCI reporters (*GFP-E2F1*$^{1-230}$ (green) and *mRFP-NLS-CycB*$^{1-266}$ (red)) to visualize cell cycle

1577   phases. The location of magnified regions shown in Main Figure 5H and I is indicated by white dashed

1578   squares.

1579

1580   Fluorescence intensities are reported as arbitrary units.

1581   Maximum projections of multiple confocal sections are shown in **D-E.**

1582   Scale bars: 100 μm.

1583

1584

Figure S5.2. Total H3K18ac and H4K8ac dynamics only partially recapitulate the S phase-
associated increase in H3K27ac

____

A-B. Immunostaining for H3K18ac in a developing control disc **(A),** with a magnified region (demarcated
by a white dashed square) in panel B, in the background of FUCCI reporters (*GFP-E2F1*[1-230] (green)
and *mRFP-NLS-CycB*[1-266] (red)) to visualize cell cycle phases.

1591

C. Quantification of normalized H3K18ac intensity level in different cell cycle phases in the pouch region
of control discs. Please note that normalization was performed against the average fluorescence
intensity across all cell cycle phases. Mean and 95% CI is shown. Statistical significance was tested
using Repeated Measures One-Way ANOVA followed by Tukey's post-hoc test for multiple comparison
(n=13 discs). Adjusted $p=0.0003$ for G1 vs. Early S; $p<0.0001$ for G1 vs. Late S; $p<0.0001$ for G1 vs.
G2; $p<0.0001$ for Early S vs. Late S; $p<0.0001$ for Early S vs G2 and $p=0.0042$ for Late S vs G2.

1598

1599   **D-E.** Immunostaining for H4K8ac in a normally developing control disc **(D),** with a magnified region
1600   (demarcated by a white dashed square) in panel E, also expressing the FUCCI reporters (*GFP-E2F1*[1-
1601   230] (green) and *mRFP-NLS-CycB*[1-266] (red)) to visualize cell cycle phases.

1602

F. Quantification of normalized H4K8ac intensity level in different cell cycle phases in the pouch region
of control discs. Please note that normalization was performed against the average fluorescence
intensity across all cell cycle phases. Mean and 95% CI is shown. Statistical significance was tested
using Repeated Measures One-Way ANOVA followed by Tukey's post-hoc test for multiple comparison
(n=12 discs). Adjusted $p=0.0018$ for G1 vs. Early S; $p<0.0001$ for G1 vs. Late S; $p<0.0001$ for G1 vs.
G2; $p<0.0001$ for Early S vs. Late S; $p<0.0001$ for Early S vs G2 and $p=0.1076$ for Late S vs G2.

1609

1610   **G-H.** Immunostaining for H3K27ac in control **(G)** and *nejire-RNAi*-expressing **(H)** discs. Same disc is
1611   also shown in Figure 5D and E but repeated here to allow for direct comparison with **(I-N)**.

1612

1613   **I-J.** Immunostaining for H3K18ac in control **(I)** and *nejire-RNAi*-expressing **(J)** discs.

1614

1615   **K-L.** Immunostaining for H4K8ac in control **(K)** and *nejire-RNAi*-expressing **(L)** discs.

1616

1617   **M-N.** Immunostaining for H3K18crot in control **(M)** and *nejire-RNAi*-expressing **(N)** discs. Yellow dashed
1618   lines indicate the boundary between the wing pouch and hinge regions.

1619

1620   **O.** Expression of a Crisper-tagged CBP/nej-GFP (Marsh et al., 2025) in a developing disc.

1621

1622 Discs were stained with DAPI to visualize nuclei.

1623 Fluorescence intensities are reported as arbitrary units.

1624 Maximum projections of multiple confocal sections are shown in **A-B** and **D-E.**

1625 Sum projections of multiple confocal sections are shown in **G-H, I-J, K-L** and **M-N.**

1626 Scale bars: 100 µm

1627

1628

1629 **Figure S6. CBP/nej and H3K27ac are required to prevent replication stress**

1630

1631 **A-C.** Expression of the JNK pathway reporter TRE-RFP in control (A), *nejire-RNAi*-expressing (B) and

1632 *bsk^{DN}, nejire-RNAi*-coexpressing (C) discs.

1633

1634 **D.** Cell cycle dynamics in *bsk^{DN}, nejire-RNAi*-coexpressing (D) disc*.* FUCCI reporters *GFP-E2F1*$^{1-230}$

1635 (green) and *mRFP-NLS-CycB*$^{1-266}$ (red) were used to visualize cell cycle phases. Even when JNK

1636 pathway activity is suppressed, one can still observe a shift of cells to the G2 phase of the cell cycle

1637 *nejire-RNAi*-expressing discs.

1638

1639 **E-F.** EdU incorporation to visualize DNA replication in the pouch region of control (E) and *nejire*-

1640 expressing (F) discs.

1641

1642 **G.** Quantification of mean EdU intensity per EdU area in the pouch region of control and *nejire*-

1643 expressing discs, serving as a proxy for speed of DNA replication. Mean and 95% CI are shown.

1644 Statistical significance was tested using two-tailed Mann Whitney test, p value=0.5737 (control discs:

1645 n=8, *nejire*-expressing discs: n=8).

1646

1647 **G'.** Quantification of EdU area per DAPI area in the pouch region of control and *nejire*-expressing discs,

1648 serving as a proxy number of cells undergoing DNA replication. Mean and 95% CI are shown. Statistical

1649 significance was tested using a two-tailed Mann Whitney test, p value=0.2345 (control discs: n=8, *nejire*-

1650 expressing discs: n=8).

1651

1652 **H-I.** Immunostaining for Abd-B in control discs **(H)** and in discs expressing *12×His-GU$^{H3-K27R}$* transgenes

1653 and containing clones homozygous mutant for a deletion of the histone gene cluster Δ*HisC* **(I)**.

1654

1655 Discs were stained with DAPI to visualize nuclei.

1656 Fluorescence intensities are reported as arbitrary units.

1657 Sum projections of multiple confocal sections are shown in **A-C**, **E-F** and **H-I**.

1658 Scale bars: 100 µm.

1659

1660

1661

Main Figure 1

Main Figure 2

2

DAPI	E2F1-FUCCI CycB-FUCCI	Histone modification
A		H3K27ac
B		H3K27me3
C		H3K9me3
D		H3K27ac
E		H3K27me3
F		H3K9me3

Main Figure 3

Main Figure 4

5

Main Figure 5

Main Figure 6