Data-driven strategies for drug repurposing

Susanna Savander[1], Nurettin Nusret Curabaz[1], Asifullah Khan[2], Khalid Saeed[1,3], Ziaurrehman Tanoli[1,4,*]

[1] Institute for Molecular Medicine Finland (FIMM), HiLIFE, University of Helsinki, Finland
[2] Pakistan Institute of Engineering & Applied Sciences (PIEAS), Islamabad, Pakistan
[3] King Abdullah International Medical Research Center (KAIMRC), King Saud bin Abdulaziz University for Health Sciences
[4] BioICAWtech, Helsinki, Finland
*Correspondence: ziaurrehman.tanoli@helsinki.fi, phone: +358503185639

ABSTRACT

Drug discovery is a complex, time-intensive, and costly process, often requiring more than a decade and substantial financial investment to bring a single therapeutic to market. Drug repurposing, the systematic identification of new indications for existing approved drugs, offers a cost-effective and expedited alternative to traditional pipelines, with the potential to address unmet clinical needs. In this study, we present a comparative analysis of drug–target interaction data from three extensively curated resources: ChEMBL, BindingDB, and GtoPdb, evaluating their release histories, curation methodologies, and coverage of approved drugs and targets. To facilitate therapeutic interpretation, we manually classified ChEMBL targets into 12 high-level biological families and mapped 817 clinically approved drug indications into 28 broader therapeutic groups. This structured framework enabled systematic profiling of the physicochemical properties of approved drugs across these groups. Leveraging this dataset, we established threshold values for 15 key physicochemical descriptors, including partition coefficient (alogP) and hydrogen bond donor/acceptor counts for each therapeutic group, providing practical benchmarks for indication-specific compound prioritization and filtering. We also analyzed cross-indication approvals to identify therapeutic areas with high repurposing potential. Furthermore, we introduce a pathway-based computational pipeline to predict repositioning opportunities for FDA-approved drugs across 10 major cancer types, which can be readily extended to other therapeutic domains. Collectively, our work integrates curated drug–target data, therapeutic-specific physicochemical profiling, and computational repurposing into a scalable, data-driven framework for accelerating drug discovery applications.

KEYWORDS:
Drug repurposing, drug discovery, drug-target interactions, drug repurposing pipeline, drug development

1. INTRODUCTION

The development of a new drug is an extraordinarily complex, resource-intensive, and time-consuming process. It typically requires $1–3 billion in investment and spans 9 to 15 years from target identification to market approval [1]. This multi-stage process begins with target identification, followed by hit identification, lead optimization, clinical evaluation across three trial phases, and post-marketing surveillance (commonly referred to as phase 4) [2]. To accelerate the hit identification stage, DNA-encoded libraries (DELs) have emerged as a transformative technology, enabling ultra-high-throughput screening of millions of compounds against selected molecular targets [3]. While DELs are highly efficient at identifying initial binders, they primarily produce binary outcomes (active/inactive) and do not provide detailed quantitative affinity measurements. In contrast, experimental techniques such as Proteome Integral Solubility Alteration (PISA) can offer proteome-wide insights into ligand-induced thermal stability shifts, indirectly informing on binding affinity and selectivity across multiple targets [4]. However, PISA assays are typically low throughput and more complex to implement in early screening pipelines.

These limitations of both DELs and PISA highlight a critical gap that can be addressed by computational methods capable of quantifying drug–target interactions (DTI) with scalability and precision. As a result, computational models have emerged as indispensable tools for refining high throughput screening output and guiding downstream prioritization. For example, MMAtt-DTA [5], an advanced attention-based architecture, predicts binding affinities for over 452,000 compounds and 1,251 human protein targets with high accuracy. Similarly, TransDTI [6] evaluated a diverse array of protein embeddings, demonstrating the utility of Evolutionary Scale Modeling (ESM) [7] and AlphaFold-derived features [8] as effective alternatives to conventional BERT-style embeddings. Drug-target affinity prediction methods should undergo rigorous validation, either through experimental assays or computational benchmarking. This principle is comprehensively discussed in the review by Tanoli et al. [9], which outlines best practices and current standards for validating DTI.

Most deep learning approaches in DTI prediction depend heavily on high-quality training data sourced from manually curated databases, including ChEMBL [10], BindingDB [11], IUPHAR/BPS Guide to Pharmacology (GtoPdb) [12], DrugTargetCommons [13], among others. As summarized in previous reviews [14, 15], dozens of publicly available DTI resources now exist, though their quality, coverage, and data curation standards vary significantly. In a recent landmark review published in Nature Reviews Drug Discovery [16], Tanoli et al. systematically categorized 181 in silico tools using a standardized

ontology, classifying them into 15 pharmacologically relevant categories. Accompanying this work, a web-based catalogue, iDRC-R4A (https://www.idrc-r4a.com/index.php) was developed to provide up-to-date information on in silico tools. This catalogue includes expert-driven rankings (via 27 field specialists), in which ChEMBL, BindingDB, and GtoPdb were ranked among the high-quality DTI databases (with ranking scores of 10, 9, and 8, respectively). Collectively, these three resources are among the most mature and robust, with more than 20 public releases (**Supplementary Figure 1**) each over the past two decades. These databases provide comprehensive, expertly curated datasets that encompass both primary and off-target interactions. Given their depth, consistency, and expert curation, the present study focuses exclusively on ChEMBL, BindingDB, and GtoPdb, utilizing their datasets to enable a thorough exploration of DTI.

While efficacy is a critical parameter, lead optimization also requires a thorough evaluation of Absorption, Distribution, Metabolism, Excretion, and Toxicity (ADMET) properties [17]. In addition, physicochemical descriptors such as partition coefficient (alogP), hydrogen bond donors (HBD) and acceptors (HBA), as well as topological polar surface area, play a significant role in determining a compound's viability in subsequent stages of drug development. Although this study does not explore ADMET characteristics, we analyzed physicochemical properties for indication-specific profiling of small molecules. This study also evaluates the drug repurposing potential of approved drugs across 28 high-level therapeutic indication groups. Drug repurposing has emerged as a cost-effective strategy to accelerate development timelines and address unmet medical needs, particularly in rare or neglected diseases [16]. Notable examples include minoxidil, originally an anti-hypertensive agent, now widely repurposed for male-pattern hair loss [18], and imatinib, first approved for chronic myeloid leukemia (CML) and later found effective against gastrointestinal stromal tumors (GISTs) through inhibition of the KIT (CD117) tyrosine kinase [19].

Expanding on this foundation, we integrate comparative DTI analyses from ChEMBL, BindingDB, and GtoPdb with high-level classifications of targets and therapeutic indications, and indication-specific physicochemical profiling of approved drugs. Leveraging threshold-based property analysis, cross-indication usage mapping, and a pathway-driven pipeline, we identify FDA-approved drugs with repurposing potential across major cancer types, delivering a unified framework for compound prioritization, profiling, and repurposing strategies applicable to diverse therapeutic domains.

Key points:

1. A timeline and curation strategies used by ChEMBL, BindingDB, and GtoPdb, highlighting their evolution and dataset characteristics.

2. A comparative analysis of drug–target interaction profiles across three databases.

3. A classification of all targets in ChEMBL into 12 high-level biological categories, examining binding profiles of both approved and investigational compounds.

4. Classification of approved drug indications in ChEMBL into 28 high-level disease groups, including cancer, cardiovascular, infectious, and neurological disorders.

5. A physicochemical profiling of approved drugs, leading to threshold values across 15 key properties (e.g., alogP, HBD/HBA) for each of the 28 disease groups. These thresholds may serve as decision-support tools for medicinal chemists by enabling the prioritization of compounds aligned with indication-specific profiles.

6. An analysis of cross-indication drug usage, revealing shared therapeutics across disease groups and identifying areas with high repurposing potential.

7. The implementation of a pathway-based computational pipeline to identify FDA-approved drugs with repurposing potential across 10 major cancer types, demonstrating a strategy that can be extended to other therapeutic areas.

2. MATERIALS AND METHODS

2.1. Curation strategies and general overview of ChEMBL, BindingDB and GtoPdb

ChEMBL, BindingDB, and the GtoPdb are three of the most comprehensive and widely utilized public repositories for DTI data with more than 25 releases each (**Supplementary Figure 1**). Collectively, they provide an essential foundation for computational pharmacology, drug repurposing, virtual screening, and the development of machine learning models for tasks such as binding affinity estimation, and compound prioritization.

ChEMBL, maintained by EMBL–EBI and first publicly released in 2009, has since progressed through 35 major version updates. It now contains over 21 million bioactivity measurements involving >2.4 million small molecules and biological ligands across >16,000 targets [10]. **Supplementary Figure 2** presents the number of first approvals per decade for small molecules and biologics (including oligosaccharides, proteins, and other modalities) recorded in the ChEMBL database. The highest number of small molecule approvals occurred in the 1980s (n = 314), while biologics peaked in the 2000s

(n = 13). **Supplementary File 1** lists the drug names and identifiers, together with their approval year and drug type.

BindingDB developed in the early 2000s at the University of North Carolina, focuses specifically on experimentally determined binding affinities (e.g., K_i, Kd, IC_{50}). The current dataset contains over 2.4 million measurements covering ~1.3 million unique ligands and nearly 9 thousand targets [11]. GtoPdb, the successor to IUPHAR-DB, became publicly accessible in 2014 and has since followed a quarterly release schedule. Its data are based on expert curation of pharmacological literature, emphasizing clinically relevant target classes such as GPCRs, ion channels, and nuclear receptors. As of the latest release, GtoPdb contains curated data on 3,039 targets and 12,163 ligands [12].

All three databases incorporate manual curation and accept direct data submissions from published literature. Chemical structures are represented using standardized formats such as SMILES, InChI, and InChIKey, with cross-references to external resources including PubChem [20], UniChem [21], and DrugBank [22]. Discrepancies in structure assignment, assay annotation, or target linkage are resolved through expert review by in-house teams. Version-controlled releases ensure data traceability, reproducibility, and transparency. **Supplementary Figure 1** shows the release timelines for ChEMBL, BindingDB, and GtoPdb. While release histories for ChEMBL and GtoPdb are directly obtained from their respective websites, BindingDB does not maintain publicly accessible version logs. To address this, we contacted the BindingDB team, who provided annual data growth statistics, enabling us to approximate its release trajectory. It is important to note that the analyses conducted in this study was initiated in 2024 and may not reflect the most recent updates to these databases.

To assess the journal coverage of drug–target activity data curated by ChEMBL, BindingDB, and GtoPdb, we extracted article-level metadata (PubMed IDs) from each database. In total, 80,286 articles are curated by ChEMBL (V33), 43,038 by BindingDB, and 8,183 by GtoPdb. Journal names corresponding to each PubMed ID are retrieved using the Entrez programming utilities (E-utilities) via the Biopython package. Among these, 2,360 articles are common to all three databases, while 38,912 are shared between ChEMBL and BindingDB, 104 between ChEMBL and GtoPdb, and 374 between GtoPdb and BindingDB (**Figure 1D**). ChEMBL curates articles from 215 unique journals, BindingDB from 197, and GtoPdb from 657. A comparative analysis of the top 10 most frequently cited journals in each database revealed considerable overlap; for example, Journal of Medicinal Chemistry, Bioorganic & Medicinal Chemistry Letters, Bioorganic & Medicinal Chemistry, and ACS Medicinal Chemistry Letters appeared in the top 10 lists of all three resources (**Figures 1A-C**).

Figure 1: Top 10 journals contributing bioactivity data curation across three databases: **A)** ChEMBL, **B)** BindingDB, and **C)** GtoPdb. **D)** Total number of overlapping articles curated by all three databases.

2.2. Preprocessing drug-target data in ChEMBL, BindingDB and GtoPdb

We downloaded the complete database dump for ChEMBL (V33) and standalone files for BindingDB and GtoPdb to perform a comparative analysis. In ChEMBL, ligands are annotated according to their maximum clinical development stage (e.g., approved, phase 3, phase 2, phase 1, and preclinical). For this study, we focused on approved drugs and investigational compounds (phase 1–3), excluding preclinical entries to maintain consistency with the scope of clinically relevant molecules. BindingDB and GtoPdb do not specify clinical phases but categorize compounds more broadly as either approved or investigational (in clinics). To enable cross-database comparisons, we standardized compound representation using InChIKeys and target identifiers using UniProt IDs.

Binding activity data are harmonized across the databases using pChEMBL values, which represent the negative logarithm of molar activity measurements such as IC_{50}, K_i, Kd, and EC_{50}. For instance, a pChEMBL value of 6 corresponds to an activity of 1,000 nM, while a value of 7 represents 100 nM. Both ChEMBL and GtoPdb provide pChEMBL values directly. For BindingDB, where activity values are reported in nanomolar concentrations, we manually converted them to pChEMBL. Entries that could not be converted, such as those reporting percentage inhibition or activity are excluded from analysis to ensure data

comparability. **Supplementary File 2** shows preprocessed datasets, including both approved and investigational ligands, for ChEMBL, BindingDB and GtoPdb.

2.3. Classification of target into 12 family groups for ChEMBL data

As BindingDB and GtoPdb do not offer target classification (in stand-alone files), our family-level analysis was conducted exclusively using ChEMBL data. ChEMBL classifies protein targets into 897 subfamilies present in the 'Protein_classification' table in the ChEMBL database (see [ChEMBL schema documentation](#)).

To facilitate consistent and biologically meaningful comparisons, we manually classified ChEMBL targets into 12 high-level functional target families: enzymes (excluding kinases), kinases, G-protein-coupled receptors (GPCRs), nuclear receptors, ion channels, epigenetic regulators, adhesion receptors, transporters, cytochromes, phosphodiesterases, proteases, and an 'other' category for unclassified or relatively smaller families. Smaller subfamilies are grouped together under broader categories; for example, tyrosine kinases and MAP kinases are classified under 'kinase', while 7TM and rhodopsin-like receptors are grouped under 'GPCR'. Conversely, in some cases, subgroups are separated to enhance resolution and interpretability, for instance, kinases are treated as distinct from the broader enzyme class, and adhesion receptors, typically considered a GPCR subclass, are assigned their own category.

Figure 2 illustrates the distribution of ChEMBL targets, drugs, and interactions for approved and investigational compounds across these 12 target families, without applying any potency cutoff on pChEMBL values. A more stringent version of this analysis, using a potency threshold of pChEMBL ≥ 6, is shown in **Supplementary Figure 3**. Among the 12 target families, enzymes, kinases, and GPCRs exhibited the broadest binding coverage for both approved and investigational compounds. The complete mapping of 3083 ChEMBL targets to their assigned high-level families is provided in **Supplementary File 3**.

Figure 2: Frequencies of drugs, targets, and their interactions are shown across 12 target families associated with approved drugs (blue) and investigational compounds (red) in the ChEMBL database. Target families with limited representation or undefined classification are grouped under the 'Other' category. The percentage values above each bar indicate the proportion of each target class within the set of approved drugs and investigational compounds. An activity threshold of pChEMBL ≥ 0 was applied for inclusion in this analysis. A more stringent version of this analysis, using a pChEMBL threshold of ≥ 6, is presented in **Supplementary Figure 3**.

2.4. Classification of indications into families

ChEMBL (V 33) links 2,330 approved drugs to 817 clinical indications, manually curated from reputable sources including ClinicalTrials.gov, DailyMed, the FDA, EMA, and others. This drug–indication mapping is accessible via the 'Drug_indication' table in ChEMBL's relational schema. To support higher-level pharmacological interpretation and cross-disease analyses, we manually grouped these 817 indications into 28 broad disease categories. These include infections, cardiovascular, mental, neoplasms, respiratory, cancer, nervous system, musculoskeletal, digestive system, eye, nutritional and metabolic, skin and connective tissue, hematologic, immunological, urogenital diseases, pathological conditions, dermatologic, renal, reproductive system, parasitic, ear, genetic

and congenital, dental, endocrine system, head and neck, oral diseases, biliary tract and sleep disorders indication groups. **Supplementary File 4** presents the comprehensive mapping of 817 indications into 28 indication groups, along with their associated 2,330 approved drugs and corresponding physicochemical properties**.**

Figure 3 illustrates the distribution of approved drugs across the indication groups, with six smaller categories combined under "Other." The infection indication group accounts for the largest number of approved drugs (n=751), followed by cardiovascular-related indications (n=666). Since some drugs are approved for multiple indications, overlaps exist across indication groups, which is further visualized via a chord diagram in **Figure 6**. Together, this structured classification offers a practical resource for future research in drug repurposing, indication-based compound prioritization, and pharmacological landscape modeling.

Figure 3: Number of drugs approved across different indication groups (smaller groups are combined under 'Other' category)

2.5. Comparative analysis of drug target data in ChEMBL, BindingDB and GtoPdb

To enable a comparative analysis across the three drug–target databases, ChEMBL, BindingDB, and GtoPdb, we first examined target coverage for both approved and investigational compounds. A word cloud plot (**Supplementary Figure 4**) visualizes the top 50 approved drugs represented in the three databases, with the font size reflecting the number of targets each drug is associated with. Sunitinib was the most broadly

profiled drug in both ChEMBL (290 targets) and BindingDB (178 targets), while flufenamic acid ranked highest in GtoPdb with 33 targets.

Without applying any threshold on binding affinity (i.e., pChEMBL ≥ 0), the average number of targets per approved drug is highest in ChEMBL (mean = 10), followed by BindingDB (mean = 7), and GtoPdb (mean = 3), as shown in **Figure 4**. For investigational compounds, a similar trend is observed with mean target counts of 9 (ChEMBL), 5 (BindingDB), and 2 (GtoPdb), respectively. These results indicate that ChEMBL provides broader proteome-level target coverage than the other two databases, a pattern that persisted under a stricter activity threshold (pChEMBL ≥ 6; **Supplementary Figure 5)**, indicating robust target profiling across confidence levels.

Figure 4: Mean number of targets across three databases at a pChEMBL threshold ≥ 0. **A)** approved drugs, **B)** investigational compounds. 'n' indicates the total number of drugs or compounds included in each analysis.

We further evaluated the overlap in drug, target, and interaction data across the three databases for both approved and investigational compounds (**Figure 5;** pChEMBL ≥ 0). Among approved drug–target pairs, 371 drugs, 283 targets, and 535 interactions are found to be common across all three databases. For investigational compounds, 255 compounds, 521 targets, and 316 interactions overlapped. Overall, ChEMBL demonstrated the most comprehensive proteome-scale annotation, consistently surpassing BindingDB and GtoPdb in both breadth and depth.

Figure 5: Overlap of drugs, targets, and interactions across **A)** approved drugs and **B)** investigational compounds in the three binding affinity databases. Target overlap is determined using UniProt IDs, and compound overlap is based on standard InChIKey identifiers.

2.6. Physiochemical properties

To better understand the chemistry associated with approved drugs, we extracted 15 physicochemical properties from the ChEMBL database. These properties include: alogP, cx_logD, cx_logP, cx_most_apKa, cx_most_bpKa, full_mwt, hba, hba_lipinski, hbd, hbd_lipinski, mw_freebase, np_likeness_score, psa, qed_weighted, and rtb. Detailed definitions and computational methodologies for these descriptors are available in ChEMBL's Compound_Properties table (ChEMBL Schema Documentation). Supplementary material further provides a section on analysis of drug's likeliness properties for approved drugs and investigational drugs present in ChEMBL database.

3. RESULTS AND DISCUSSION
3.1. Indications groups with shared drug repurposing potential

Section 2.4 describes the categorization of 817 indications from the ChEMBL database into 28 broad disease groups and highlights the number of approved drugs associated with each group (**Figure 3**). In this section, we investigate the extent of drug sharing across these indication groups, an analysis that may offer insights into therapeutic overlaps and repurposing potential. **Figure 6** provides a visual summary of drug overlap between indication groups, where the width of each connecting chord represents the number of shared approved drugs. While this figure conveys the relative scale of overlap,

the exact numbers are detailed in **Supplementary File 5**. For example, the cancer group is associated with 386 approved drugs, 103 of which are also approved for neoplasms, 30 for hematologic diseases, and 15 for nutritional and metabolic disorders. These overlaps reflect well-established biological relationships. For example, hematologic malignancies and solid tumors may share key molecular targets, most notably tyrosine kinases such as BCR-ABL and KIT, which can be targeted by the same drugs across multiple cancer types. A prominent example is imatinib, originally developed for chronic myeloid leukemia (CML) via BCR-ABL inhibition, and later approved for gastrointestinal stromal tumors (GISTs) through its inhibition of KIT and PDGFR [23, 24]. Furthermore, metabolic reprogramming is a recognized hallmark of cancer and is observed across both hematologic and solid malignancies [25, 26] This commonality supports the observed therapeutic overlap across cancer, hematologic, and metabolic disease indications. The largest indication group, infections (n = 751 approved drugs), also shows substantial overlap, most notably with respiratory (n = 67 shared drugs) and eye diseases (n = 55). This overlap is biologically plausible: many respiratory pathogens such as adenoviruses, haemophilus influenzae, streptococcus pneumoniae, and moraxella catarrhalis can infect both the lungs and the ocular surface, causing conjunctivitis, keratitis, and pneumonia [27]. These conditions caused by bacterial infections are often managed using shared classes of antimicrobials and anti-inflammatory agents [28].

This overlap analysis reveals disease clusters sharing pharmacologically active compounds, such as broad-spectrum agents targeting conserved mechanisms across pathogens [29], providing a rational basis for drug repurposing. These insights can guide candidate prioritization and cross-indication screening, accelerating therapy development for related conditions.

Figure 6: The number of drugs shared among 28 indication groups is represented, with the width of each connecting patch proportional to the number of shared approved drugs. **Supplementary File 5** presents this information in a tabular format.

3.2. Thresholds on physiochemical properties based on indication groups

Previously, we categorized approved ChEMBL drugs into 28 therapeutic indication groups based on their annotated clinical uses. As a logical extension, we aimed to explore whether distinct physicochemical trends exist within each indication group, with the goal of identifying optimal property ranges for drug design tailored to specific disease areas. We analyzed the distribution of the 15 physicochemical properties across all 28 indication groups. **Figure 7** presents the property distributions for the seven indication groups with the highest numbers of associated drug approvals, while **Supplementary Figures 6–8** provide similar data for the remaining 21 groups. Based on this analysis, we established lower and upper threshold values for each property within every indication group, defined as ±5% around the mean value observed for drugs in each group (see **Supplementary File 6**).

Figure 7: Distribution of 15 physicochemical properties (columns) for approved drugs across 7 indication groups (rows): infections, cardiovascular, mental, neoplasms, respiratory, cancer, and nervous system diseases. Orange lines represent the median values for each indication group across the 15 properties.

This property-based profiling enables the derivation of indication-specific physicochemical boundaries, which can be particularly useful in guiding early-stage drug design and optimization. For example, recognizing that drugs approved for nervous system diseases typically exhibit lower polar surface area (PSA) and fewer hydrogen bond donors (HBD) can guide the prioritization of compound libraries [23], This aligns with our findings, which show PSA values ranging from 54.65 to 60.41, the second lowest among the 28 groups and HBD values between 0.95 and 1.05, the lowest observed (**Supplementary File 6**). In conclusion, this analysis provides a systematic framework linking physicochemical profiles to therapeutic areas, with thresholds that can guide the design of compounds more likely to succeed in specific indications.

3.3. Top 10 drugs approved for most indications

We conducted a systematic analysis of the top repurposed drugs across diverse disease indications. The ten most frequently repurposed drugs identified were: dexamethasone, prednisone, prednisolone, cortisone acetate, menthol, naproxen, lidocaine, methotrexate, doxorubicin hydrochloride, and ibuprofen (**Table 1**). These drugs demonstrate substantial versatility in their therapeutic applications. Notably, dexamethasone is a synthetic glucocorticoid that primarily exerts its effects through the glucocorticoid receptor (UniProt ID: P04150), a ligand-activated transcription factor. It received FDA approval in 1958 and is subsequently introduced into clinical use for hematologic malignancies, including acute lymphoblastic leukemia (ALL) [31, 32]. Since its approval, dexamethasone has been repurposed for over 70 distinct indications spanning a broad range of indication groups, including oncology, respiratory diseases, endocrinology, dermatology, and autoimmune disorders [33] (see **Supplementary File 7**). Mechanistically, dexamethasone binds to cytosolic glucocorticoid receptors and translocates to the nucleus, modulating gene transcription involved in inflammatory, immune, metabolic, and apoptotic pathways, serving as a potent ligand-activated transcription factor [34]. ChEMBL data indicate interactions with 11 distinct protein targets, underscoring its pharmacological flexibility (**Table 1**).

Interestingly, 7 of the 10 most frequently repurposed drugs in our analysis interact with more than 10 distinct protein targets, highlighting polypharmacology as a common trait among highly repurposable therapeutics. This aligns with broader evidence that multi-target activity is a hallmark of efficacious agents used across multiple indications [35]. **Table 1** summarizes the top 10 drugs along with associated indications, targets, and target superfamilies. A more detailed version of this table is presented in **Supplementary File 7**.

Table 1: Top 10 repurposed drugs showing number of associated indications and targets.

Drug	Indications	Targets	Target families
Dexamethasone	74	11	Nuclear receptor, GPCR, Cytochrome, Other
Prednisone	62	7	Enzyme, Nuclear receptor, Other, Phosphodiesterase
Prednisolone	58	13	Enzyme, GPCR, Nuclear receptor, Other, Protease, Transporter
Cortisone acetate	44	3	Enzyme, GPCR, Nuclear receptor
Menthol	30	4	Ion channel
Lidocaine	25	11	Cytochrome, Enzyme, GPCR, Ion channel, Kinase, Other

Naproxen	24	14	Cytochrome, Enzyme, GPCR, Transporter
Methotrexate	22	50	Enzyme, Epigenetic regulator, Other, Phosphodiesterase, Protease, Transporter
Ibuprofen	22	24	Cytochrome, Enzyme, Other, Transporter
Doxorubicin hydrochloride	22	35	Cytochrome, Enzyme, Epigenetic regulator, GPCR, Kinase, Nuclear receptor, Other, Phosphodiesterase, Protease, Transporter

3.5. A novel pathway driven strategy to repurpose drugs for 10 cancer types

We developed a novel pathway-driven in silico strategy to identify repurposed drugs for ten cancer types: prostate, endometrial, breast, non-small cell lung (NSCLC), colorectal, pancreatic, hepatocellular cancer (HCC), gastric, thyroid, and acute myeloid leukemia (AML). These cancer types are chosen due to the availability of well-curated and disease-specific pathway information in the Kyoto Encyclopedia of Genes and Genomes (KEGG) database [36], enabling a biologically grounded approach to target identification. For each cancer type, we retrieved the KEGG disease ID, MeSH ID, and the associated KEGG disease pathways. From these pathways, we extracted gene symbols functionally implicated in the disease mechanisms. These gene symbols are then mapped to UniProt identifiers using the UniProt API (reviewed entries), providing a standardized and biologically relevant protein target set. A total of 377 unique genes are associated with the ten cancer types, of which 375 are successfully mapped to UniProt IDs. The number of pathway-associated protein targets identified for each cancer type is thyroid (37), NSCLC (72), AML (67), endometrial (58), pancreatic (76), prostate (105), colorectal (86), breast (147), gastric (148), and HCC (168). **Supplementary Table 3** shows names of gene targets involved in disease pathways in the KEGG database.

To link these targets to potential therapeutics, we retrieved drug–target bioactivity data from ChEMBL (v33), filtering for approved drugs with known activity against single human protein targets and available pChEMBL values. In cases where multiple bioactivity measurements existed for a given drug-target pair across different publications, we computed the median pChEMBL value to ensure data robustness. This resulted in a final dataset comprising 1,914 unique approved drugs and 16,256 drug-target interaction pairs. To identify viable drug repurposing candidates, we applied a potency threshold of pChEMBL ≥ 6, indicative of moderate to strong binding affinity. A drug is considered a repurposing candidate for a cancer type if it targets at least 10% of the protein targets

involved in that cancer's KEGG-defined pathways [37]. Using this threshold-driven approach, we identified three repurposing candidates each for thyroid cancer and NSCLC, two for AML, and one candidate each for endometrial and pancreatic cancers (**Figure 8**). **Supplementary Figure 9** is the extension of Figure 6 using a more stringent threshold of pChEMBL ≥5. **Supplementary Table 1 & 2** list the repurposed drugs alongside their corresponding target proteins involved in cancer-specific pathways for each cancer type. These results illustrate the potential of leveraging pathway-level biological context to systematically uncover repurposing therapeutic opportunities. We excluded from consideration any drug already approved for the same cancer type. This is done by cross-referencing each candidate against known approved indications as listed in **Supplementary File 4**. **Supplementary Figure 10** presents the number of approved drugs for each cancer type, while **Supplementary Table 4** lists their corresponding names. Of the ten cancers analyzed, breast cancer is associated with the largest number of approved drugs, while endometrial, gastric, and hepatocellular carcinoma (HCC) have no approved therapies recorded in the ChEMBL database. This case study introduces a scalable, pathway-informed drug repurposing pipeline that integrates curated biological knowledge and quantitative drug-target interaction data. Although demonstrated here across ten cancer types, the framework is applicable to any indication for which pathway information is available. By targeting biologically validated networks rather than isolated proteins, this strategy increases the likelihood of identifying repurposing candidates with mechanistic relevance and therapeutic efficacy [38].

Figure 8: Number of repurposed drugs identified across ten cancer types (using threshold on pChEMBL ≥6). The labels above each bar indicate the total number of pathway-associated targets for each cancer type, as well as the subset of those targets that are known to bind to the repurposed drugs.

4. CONCLUSION

This study presents a systematic, data-driven framework for drug repurposing that integrates high-quality drug–target data, indication-specific physicochemical profiling, and

a novel pathway-driven computational pipeline. By leveraging the most mature DTI databases, ChEMBL, BindingDB, and GtoPdb, we provide an in-depth comparative analysis of their curation practices, coverage, and binding data (see **Figures 1, 2, 4,** and **5**). Our indication-specific profiling of physicochemical properties offers practical thresholds to guide compound prioritization during early-stage discovery (**Figure 7**, **Supplementary Figures 6–8 and Supplementary File 6**). This approach enhances the efficiency of medicinal chemistry efforts by aligning compound properties with those historically successful in each therapeutic area. However, such profiling may oversimplify pharmacological complexity and potentially overlook novel chemotypes or emerging drug modalities that deviate from established norms [39].

Furthermore, the pathway-driven use case (**section 3.5**) applied to ten cancer types demonstrates how curated biological networks can be systematically leveraged to uncover promising repurposing candidates based on mechanistic relevance and target coverage (see **Figure 8 and Supplementary Figure 9**). The added value of this work lies in integrating drug–target data, physicochemical profiles, and indication-specific context into a unified framework. By linking these curated datasets to pathway-level and mechanistic insights, our approach supports more precise identification and prioritization of drug repurposing opportunities with strong biological relevance.

This study has several limitations. Although several rare diseases are included among the 817 indications (**Supplementary File 4**), we did not classify them separately as a distinct indication group. For example, Tourette syndrome, a rare neurological disorder, was grouped under *nervous system diseases*, while gaucher disease and wilson disease, both rare conditions, are assigned to the *nutritional and metabolic diseases* category. Another limitation arises from the use case on KEGG pathways, which provide a structured representation of disease biology but may not fully reflect tissue- or patient-specific molecular variation. In addition, our analysis concentrated primarily on pharmacodynamics, without incorporating pharmacokinetics, target expression patterns, or toxicity factors that are critical for translational relevance [38]. Data availability also constrained our focus largely to small-molecule drugs, with only a limited number of examples from other modalities such as peptides and proteins. This limitation could be addressed in future work as more comprehensive datasets on biological ligands become available. Future extensions could integrate transcriptomic, proteomic, and clinical outcome data to enable context-aware precision medicine [40], as well as incorporate adverse event profiles, drug-induced gene expression signatures [41], and network-based proximity metrics to refine prioritization as done in [42]. Applying this pathway-driven repurposing framework to non-oncological indications, particularly rare, neglected, or complex diseases, could further expand its clinical utility [43].

CONFLICTS OF INTEREST

The authors declare no conflicts of interest.

ACKNOWLEDGMENTS
The work was funded by the Academy of Finland (No. 351507 to ZT).

Susanna Savander is a master's student at the Institute for Molecular Medicine Finland (FIMM), HiLIFE, University of Helsinki, Finland.

Nurettin Nusret Curabaz is an erasmus student at FIMM, HiLIFE, University of Helsinki, Finland.

Asifullah Khan is full professor at Pakistan Institute of Engineering & Applied Sciences (PIEAS), Islamabad, Pakistan

Khalid Saeed is a principal investigator and associate scientist at King Abdullah International Medical Research Center (KAIMRC), King Saud bin Abdulaziz University for Health Sciences.

Ziaurrehman Tanoli is a principal Investigator at FIMM, HiLIFE, University of Helsinki, Finland.

REFERENCES

1.   Wouters OJ, McKee M, Luyten J. Estimated Research and Development Investment Needed to Bring a New Medicine to Market, 2009-2018, JAMA 2020;323:844–853.

2.   Hughes JP, Rees S, Kalindjian SB et al. Principles of early drug discovery, Br J Pharmacol 2011;162:1239–1249.

3.   Satz AL, Brunschweiger A, Flanagan ME et al. DNA-encoded chemical libraries, Nature Reviews Methods Primers 2022;2.

4.   Gaetani M, Zubarev RA. Proteome Integral Solubility Alteration (PISA) for High-Throughput Ligand Target Deconvolution with Increased Statistical Significance and Reduced Sample Amount, Methods Mol Biol 2023;2554:91–106.

5.   Schulman A, Rousu J, Aittokallio T et al. Attention-based approach to predict drug–target interactions across seven target superfamilies, Bioinformatics 2024;40.

6.   Kalakoti Y, Yadav S, Sundar D. TransDTI: Transformer-Based Language Models for Estimating DTIs and Building a Drug Recommendation Workflow, ACS Omega 2022;7:2706–2717.

7.   Lin Z, Akin H, Rao R et al. Evolutionary-scale prediction of atomic-level protein structure with a language model, Science 2023;379:1123–1130.

8.   Jumper J, Evans R, Pritzel A et al. Highly accurate protein structure prediction with AlphaFold, Nature 2021;596:583–589.

9.   Tanoli Z, Schulman A, Aittokallio T. Validation guidelines for drug-target prediction methods, Expert Opin Drug Discov 2025;20:31–45.

10.   Zdrazil B, Felix E, Hunter F et al. The ChEMBL Database in 2023: a drug discovery platform spanning multiple bioactivity data types and time periods, Nucleic Acids Res 2024;52:D1180–D1192.

11.   Liu T, Hwang L, Burley SK et al. BindingDB in 2024: a FAIR knowledgebase of protein-small molecule binding data, Nucleic Acids Res 2025;53:D1633–D1644.

12.   Harding SD, Armstrong JF, Faccenda E et al. The IUPHAR/BPS Guide to PHARMACOLOGY in 2024, Nucleic Acids Res 2024;52:D1438–d1449.

13.   Tanoli Z, Alam Z, Vaha-Koskela M et al. Drug Target Commons 2.0: a community platform for systematic analysis of drug-target interaction profiles, Database (Oxford) 2018;2018:1–13.

14.   Tanoli Z, Vaha-Koskela M, Aittokallio T. Artificial intelligence, machine learning, and drug repurposing in cancer, Expert Opin Drug Discov 2021;16:977–989.

15.   Tanoli Z, Seemab U, Scherer A et al. Exploration of databases and methods supporting drug repurposing: a comprehensive survey, Brief Bioinform 2021;22:1656–1678.

16.   Tanoli Z, Fernandez-Torras A, Ozcan UO et al. Computational drug repurposing: approaches, evaluation of in silico resources and case studies, Nat Rev Drug Discov 2025;24:521–542.

17.   Hodgson J. ADMET--turning chemicals into drugs, Nat Biotechnol 2001;19:722–726.

18.   Zappacosta AR. Reversal of baldness in patient receiving minoxidil for hypertension, N Engl J Med 1980;303:1480–1481.

19.   Joensuu H. Treatment of inoperable gastrointestinal stromal tumor (GIST) with Imatinib (Glivec, Gleevec), Med Klin (Munich) 2002;97 Suppl 1:28–30.

20.   Kim S, Chen J, Cheng T et al. PubChem 2025 update, Nucleic Acids Res 2025;53:D1516–D1525.

21.   Chambers J, Davies M, Gaulton A et al. UniChem: a unified chemical structure cross-referencing and identifier tracking system, J Cheminform 2013;5:3.

22.   Knox C, Wilson M, Klinger CM et al. DrugBank 6.0: the DrugBank Knowledgebase for 2024, Nucleic Acids Res 2024;52:D1265–D1275.

23.   van Oosterom AT, Judson I, Verweij J et al. Safety and efficacy of imatinib (STI571) in metastatic gastrointestinal stromal tumours: a phase I study, Lancet 2001;358:1421–1423.

24.   Capdeville R, Silberman S. Imatinib: a targeted clinical drug development, Semin Hematol 2003;40:15–20.

25.   Hindes MT, McElligott AM, Best OG et al. Metabolic reprogramming, malignant transformation and metastasis: Lessons from chronic lymphocytic leukaemia and prostate cancer, Cancer Lett 2025;611:217441.

26.   Phan LM, Yeung SC, Lee MH. Cancer metabolic reprogramming: importance, main features, and potentials for precise targeted anti-cancer therapies, Cancer Biol Med 2014;11:1–19.

27.   Lalbiaktluangi C, Yadav MK, Singh PK et al. A cooperativity between virus and bacteria during respiratory infections, Front Microbiol 2023;14:1279159.

28.   Inchingolo R, Pierandrei C, Montemurro G et al. Antimicrobial Resistance in Common Respiratory Pathogens of Chronic Bronchiectasis Patients: A Literature Review, Antibiotics (Basel) 2021;10.

29.   He W, Huan X, Li Y et al. A broad-spectrum antibiotic targets multiple-drug-resistant bacteria with dual binding targets and no detectable resistance, Nat Commun 2025;16:7048.

30.   Pajouhesh H, Lenz GR. Medicinal chemical properties of successful central nervous system drugs, NeuroRx 2005;2:541–553.

31.   Freireich EJ, Gehan E, Frei E et al. The Effect of 6-Mercaptopurine on the Duration of Steroid-induced Remissions in Acute Leukemia: A Model for Evaluation of Other Potentially Useful Therapy, Blood 1963;21:699–716.

32.   Relling MV, Dervieux T. Pharmacogenetics and cancer therapy, Nat Rev Cancer 2001;1:99–108.

33.   Vohra M, Sharma AR, Satyamoorthy K et al. Pharmacogenomic considerations for repurposing of dexamethasone as a potential drug against SARS-CoV-2 infection, Per Med 2021;18:389–398.

34.   Cain DW, Cidlowski JA. After 62 years of regulating immunity, dexamethasone meets COVID-19, Nat Rev Immunol 2020;20:587–588.

35.   Ryszkiewicz P, Malinowska B, Schlicker E. Polypharmacology: promises and new drugs in 2022, Pharmacol Rep 2023;75:755–770.

36.   Kanehisa M, Furumichi M, Sato Y et al. KEGG: biological systems database as a model of the real world, Nucleic Acids Res 2025;53:D672–D677.

37.   Kanehisa M, Furumichi M, Tanabe M et al. KEGG: new perspectives on genomes, pathways, diseases and drugs, Nucleic Acids Res 2017;45:D353–D361.

38.   Barabasi AL, Gulbahce N, Loscalzo J. Network medicine: a network-based approach to human disease, Nat Rev Genet 2011;12:56–68.

39.   Miethke M, Pieroni M, Weber T et al. Towards the sustainable discovery and development of new antibiotics, Nat Rev Chem 2021;5:726–749.

40.   Subramanian A, Narayan R, Corsello SM et al. A Next Generation Connectivity Map: L1000 Platform and the First 1,000,000 Profiles, Cell 2017;171:1437–1452 e1417.

41.   Paci P, Fiscon G, Conte F et al. Comprehensive network medicine-based drug repositioning via integration of therapeutic efficacy and side effects, NPJ Syst Biol Appl 2022;8:12.

42.   Guney E, Menche J, Vidal M et al. Network-based in silico drug efficacy screening, Nat Commun 2016;7:10331.

43.   Jonker AH, O'Connor D, Cavaller-Bellaubi M et al. Drug repurposing for rare: progress and opportunities for the rare disease community, Front Med (Lausanne) 2024;11:1352803.