SynaptoTagMe: A Toolkit for *In Vivo* Mapping and Modulating Neurotransmission
at Single-Cell Resolution

Authors: Andrea Cuentas-Condori[1], Patricia Chanabá-López[1], Matthew Thomas[1], Likui Feng[2],
Aaron Wolfe[1], Peter Agoba[1], Matthew L. Schwartz[3], Maximillian Brown[2], Margaret Ebert[2,4], Erik
Jorgensen[3], Cornelia I. Bargmann[2], Daniel Colón-Ramos[1, 5, 6 *]

Affiliations:
[1] Department of Neuroscience and Department of Cell Biology, Yale University School of
Medicine; New Haven, CT 06536, USA.
[2] Lulu and Anthony Wang Laboratory of Neural Circuits and Behavior, The Rockefeller
University. New York, NY 10065, USA.
[3] Howard Hughes Medical Institute and School of Biological Sciences, University of Utah, Salt
Lake City, UT 84112, USA.
[4] Current address: Omenn-Darling Bioengineering Institute, Princeton University, Princeton, NJ
08540, USA.
[5] Wu Tsai Institute, Yale University; New Haven, CT 06510, USA.
[6] Instituto de Neurobiología, Recinto de Ciencias Médicas, Universidad de Puerto Rico, San
Juan 00901, Puerto Rico.

*Corresponding author and Lead Contact. Email: daniel.colon-ramos@yale.edu

ABSTRACT

Understanding the organization and regulation of neurotransmission at the level of individual neurons and synapses requires tools that can track and manipulate transmitter-specific vesicles *in vivo*. Here, we present SynaptoTagMe, a suite of genetic tools in *Caenorhabditis elegans* to fluorescently label and conditionally ablate the vesicular transporters for glutamate, GABA, acetylcholine, and monoamines. Using a structure-guided approach informed by protein topology and evolutionary conservation, we engineered endogenously tagged versions for each transporter that maintain their physiological function while allowing for cell-specific, bright, and stable visualization. We also developed conditional knockout strains that enable targeted disruption of neurotransmitter synthesis or packaging in single neurons. We applied this toolkit to map co-expression of vesicular transporters across the *C. elegans* nervous system, revealing that over 10% of neurons exhibit co-transmission. Using the ADF sensory neuron as a case study, we demonstrate that serotonin and acetylcholine are trafficked in partially distinct vesicle pools. Our approach provides a powerful platform for mapping, monitoring, and manipulating neurotransmitter identity and use *in vivo*. The molecular strategies described here are likely applicable across species, offering a generalizable approach to dissect synaptic communication *in vivo*.

INTRODUCTION

Understanding how the nervous system generates behavior requires tools that can resolve the molecular identity, spatial localization, and functional contribution of neurotransmitters *in vivo*. Neurotransmitters are the primary means by which neurons communicate, and their synthesis, packaging, and release are governed by evolutionarily conserved molecular pathways shared from *Caenorhabditis elegans* to vertebrates (Südhof, 2021). These transmitters shape the strength, kinetics (tonic vs. phasic), and polarity (excitatory vs inhibitory) of synaptic transmission, thereby influencing how information is processed and how behavior is regulated (Crawford & Kavalali, 2015; Gouwens et al., 2020; Kamalova & Nakagawa, 2021; Liu et al., 2021). Because neurotransmitters are central to defining the functional properties of synapses, understanding their identity and dynamics is essential for interpreting circuit function. Even in organisms with complete connectomes—such as *C. elegans* (White et al., 1986) and *Drosophila melanogaster* (Scheffer et al., 2020; Seggewisse & Winding, 2024; Yi et al., 2024) anatomical connectivity alone cannot explain how neural circuits generate behaviors. To build accurate, testable models of

circuit function, it is necessary to also determine which neurotransmitters are used at specific
synapses and how their release is spatiotemporally organized and regulated *in vivo*. Yet, despite
the centrality of neurotransmitters to circuit logic, the field still lacks broadly applicable tools to
visualize and manipulate transmitter-specific vesicle pools with the precision needed to study their
roles in intact, living animals.

Traditional approaches—such as in situ hybridization, immunohistochemistry, and
transcriptomics—have been instrumental in mapping neurotransmitter identity. These methods,
however, often lack cell-specific control, temporal resolution, or the ability to monitor transmitter
usage dynamically within intact circuits. Moreover, neurotransmitter identity can change in
response to environmental or physiological cues. For example, neurons may co-release multiple
transmitters or modulate transmitter usage depending on stress, activity, or developmental stage,
and these changes have consequences in animal behavior and circuit function (Chen et al., 2023;
Li et al., 2024; Maddaloni et al., 2024; Sitko et al., 2025; Wu et al., 2020). Tracking and
manipulating these physiological changes require new tools that allow endogenous, live imaging
and functional interrogation of neurotransmitters in single neurons.

Vesicular transporters offer a strategic entry point for such investigations. These multi-pass
membrane proteins package specific neurotransmitters—such as glutamate, GABA,
acetylcholine, and monoamines—into synaptic vesicles and are necessary and sometimes
sufficient for defining a neuron's transmitter phenotype (Edwards, 2007). Because they are
genetically encoded and highly conserved (Alfonso et al., 1993; Bellocchio et al., 1998; Chaudhry
et al., 1998; Lee et al., 1999; McIntire, Jorgensen, & Horvitz, 1993; McIntire et al., 1997; Roghani
et al., 1994), vesicular transporters provide a powerful molecular handle for developing
generalizable tools that probe synaptic identity and function across species. Tagging these
transporters can offer direct, real-time readouts of presynaptic signaling and enable manipulations
that dissect the functional contribution of specific neurotransmitters *in vivo* (Li et al., 2020). Yet
for these tools to be broadly useful, it is essential that tagging does not disrupt the localization or
function of the transporter. If appropriate insertion sites can be identified and validated
functionally, the evolutionary conservation of vesicular transporters suggests that such designs
could serve as generalizable platforms across systems and species.

Here, we present SynaptoTagMe, a comprehensive toolkit for tracking and manipulating
transmitter-specific vesicles in *C. elegans*. Using a structure-guided approach informed by
predicted protein topology and sequence conservation, we engineered endogenously tagged

versions of the vesicular transporters for glutamate (EAT-4/VGLUT), GABA (UNC-47/VGAT),
acetylcholine (UNC-17/VAChT), and monoamines (CAT-1/VMAT). We validated *in vivo* that the
tagged transporters retain functionality and enables bright, cell-specific imaging. In parallel, we
developed conditional knockout strains that enabled spatiotemporal access to the ablation of the
packaging or synthesis of specific neurotransmitters in defined neurons, allowing causal tests of
neurotransmitter function at the single-cell level within behaving animals.

We applied this toolkit to identify neurons that co-express multiple vesicular transporters,
revealing that 10% of *C. elegans* neurons contain the machinery for co-transmission. Focusing
then on the ADF sensory neuron, we validate that ADF expresses the machinery for co-
transmission of serotonin and acetylcholine. We demonstrate that serotonin and acetylcholine are
packaged in partially distinct vesicle populations. Together, our observations suggest that co-
transmission can be spatially organized, offering a refined view of how individual neurons diversify
their signaling output *in vivo*. Our discoveries also highlight co-transmission as a widespread and
previously underappreciated feature of nervous system organization, rather than a rare or
specialized exception. Co-transmission is not unique to *C. elegans*; in *Drosophila* the VAChT
protein can be modulated in GABAergic and glutamatergic neurons by microRNAs (Chen et al.,
2023); in mammals, serotonergic neurons in the dorsal raphe co-release glutamate or GABA
depending on context (Li et al., 2024), while starburst amacrine cells in the retina release both
acetylcholine and GABA with distinct calcium sensitivities (Lee et al., 2010; Morrie & Feller, 2015).
These examples, along with our findings, underscore the evolutionary conservation of co-
transmission as a mechanism for expanding the functional repertoire of single neurons.

By enabling simultaneous visualization of different transmitter-specific vesicle pools within the
same neuron, our tools uncover molecular heterogeneity at individual synapses and reveal new
layers of synaptic plasticity. More broadly, our findings establish a functional framework for
probing neurotransmitter dynamics, synaptic architecture, and co-transmission *in vivo*. The
strategies developed here are generalizable to other model systems and open new avenues for
dissecting neural circuit logic with molecular and cellular precision.

RESULTS

A systematic strategy for tagging and manipulating transmitter-specific vesicles *in vivo*

All synaptic vesicle transporters are multi-pass transmembrane proteins with structural loops facing either the cytosolic or luminal space. To visualize transmitter-specific vesicle pools *in vivo*, we developed a suite of fluorescently tagged, functional versions of the vesicular transporters for glutamate (EAT-4/VGLUT), GABA (UNC-47/VGAT), acetylcholine (UNC-17/VAChT), and monoamines (CAT-1/VMAT) in *C. elegans*. We chose these four neurotransmitter classes because they are used by more than 90% of the neurons in *C. elegans* (Wang et al., 2024). We used a systematic design pipeline that integrated (1) protein topology predictions, (2) evolutionary conservation, and (3) structure-guided fluorophore placement to identify regions of each transporter suitable for tagging without disrupting function. These approaches were used to generate endogenous knock-in alleles with bright, cell-specific labeling through Flippase recombinase systems (Schwartz & Jorgensen, 2016) or self-assembling split-GFP tags (He et al., 2019) (Figure 1). When possible, tools were developed for both green and red-based fluorophores to allow for multi-color imaging. For each transporter, we also created matched conditional knockout strains by inserting FRT-flanked cassettes to disrupt neurotransmitter packaging or synthesis in defined cells, adding to the existing cell-specific knockout tools available in the field (Huang et al., 2023; Lopez-Cruz et al., 2019) (Figure 1). To drive expression of Flippase in GABAergic and Cholinergic neurons we inserted Flippase into the *unc-47* and *unc-17* locus after a self-cleaving 2A peptide sequence (Ahier & Jarriault, 2014), and used available Flippase drivers for Serotonergic and Dopaminergic neurons (Muñoz-Jiménez et al., 2017). Together, these new tools allow precise labeling and *loss-of-function* analysis of transmitter-specific vesicles in intact circuits and behaving animals (summarized in Table 1). All strains created in this study are accessible upon request from the Caenorhabditis Genetics Center (CGC) and the respective sequences are available at https://www.intralab.app/research-papers/cuentas-condori_etal-2026.

Table 1. Cellular, genetic and molecular tools to probe neurotransmission in single cells

Genotype	Purpose
Toolkit for examining Glutamatergic transmission	
eat-4::gfp FLP-on(syb8568)	Cell-specific GFP knock-in (C-terminus end)
eat-4::mRuby3 FLP-on(syb9139)	Cell-specific mRuby3 knock-in (C-terminus end)
eat-4 (kySi76 kySi77)	Cell-specific knockout
Toolkit for examining GABAergic transmission	
unc-47::gfp (syb6990)	Endogenous full body GFP tagging
unc-47::mKate2 (syb7358)	Endogenous full body mKate2 tagging

unc-47::gfp11 (syb7313)	Endogenous and cell-specific GFP labeling
unc-47::gfp11x3 (syb7849)	Endogenous and cell-specific GFPx3 labeling
unc-25 (syb5949 syb6275)	Cell-specific knockout
Toolkit for examining Cholinergic transmission	
unc-17::mKate2 (ot907)	Endogenous full body mKate2 tagging
unc-17::GFP FLP-on (ola503)	Cell-specific GFP knock-in (C-terminus end)
unc-17::mRuby3 FLP-on (syb7882)	Cell-specific mRuby3 knock-in (C-terminus end)
GFP FLP-on::unc-17 (syb7251)	Cell-specific GFP knock-in (N-terminus end)
unc-17 (syb5779 syb5987)	Cell-specific knockout
Toolkit for examining Monoaminergic transmission	
cat-1::gfp11x3 (syb7239)	Endogenous and cell-specific GFPx3 labeling
cat-1 (ky1101 ky1118)	Cell-specific knockout
Flippase drivers	
P*eft-3*::Flippase (*sybIs9614*)	Flippase expression in all cells
unc-47::T2A::Flippase(*syb8125*)	Flippase expression in GABAergic neurons
unc-17::T2A::Flippase (*syb8059*)	Flippase expression in Cholinergic neurons
P*tph-1*::Flippase (*bqSi488*)	Flippase expression in Serotonergic neurons
P*dat-1*::Flippase (*bqSi614*)	Flippase expression in Dopaminergic neurons

Functional labeling of glutamatergic vesicles via EAT-4/VGLUT

Glutamate functions as a key excitatory neurotransmitter in the nervous system, and its packaging into synaptic vesicles requires the conserved Vesicular Glutamate Transporter (VGLUT) (Bellocchio et al., 1998; Lee et al., 1999), which is sufficient to confer glutamatergic identity to a neuron. In *C. elegans*, the VGLUT homolog EAT-4 is expressed in 43 of the 118 neuronal classes catalogued (Wang et al., 2024). EAT-4/VGLUT is predicted to have 12 transmembrane domains (Figure 2A), and prior tools have allowed for cell-specific knockout of its full coding sequence (Lopez-Cruz et al., 2019). A previously reported transgene with EAT-4/VGLUT fused to GFP demonstrated localization to synapses (Yu & Chang, 2022); here we extend this approach to an endogenously tagged allele.

To generate a bright, functional reporter that reflects endogenous EAT-4/VGLUT localization *in*

vivo, we inserted a GFP tag into the protein C-terminal cytoplasmic domain using a FLP-on

cassette (Schwartz & Jorgensen, 2016). The C-terminus was chosen based on conservation

analysis and AlphaFold structural predictions (Jumper et al., 2021; Pei & Grishin, 2001), which

identified it as a cytosolic and weakly conserved region, minimizing the risk of disrupting

conserved protein functions (Figure S1A-B). GFP was inserted just before the STOP codon

(Figure 2B). To examine if introduction of GFP into the endogenous EAT-4/VGLUT gene affected

function, we assessed NaCl chemotaxis (Figure S1C) —an EAT-4–dependent learning behavior

mediated by the ASE neurons (Figure S1C-D) (Bargmann & Horvitz, 1991; Sato et al., 2021;

Uchida et al., 2003). EAT-4::GFP FLP-on strains, with or without Flippase expression in all

neurons or selectively in ASE neurons, displayed normal chemotaxis behavior (Figure 2C and

S1D), in contrast to *eat-4(ky5)* loss-of-function mutants or mutant animals with defects in

chemosensory neurons, including ASE (*che-1* mutants; Figure 2C, S1C-D). Our findings suggest

that the tagged protein remains functional and capable of sustaining known glutamate-dependent

behaviors in the organism.

The insertion of the FLP-on cassette enables expression of EAT-4/VGLUT::GFP upon cell-

specific expression on the FLP recombinase. To validate its use, we expressed pan-cellular

flippase via the *eft-3* promoter (Seydoux & Fire, 1994) in the worms engineered with the EAT-

4::GFP FLP-on cassette. We observed bright EAT-4::GFP signal throughout the nervous system,

especially in the nerve ring and sensory neurons (Figure 2D), consistent with earlier

transcriptional reporters of the *eat-4* gene (Lee et al., 1999; Serrano-Saiz et al., 2013). Moreover,

when Flippase was driven specifically in ASE neurons (using the ASE-specific promoter, P*flp-6*),

we observed punctate labeling along ASE axons, matching the distribution of presynaptic sites

identified by serial electron microscopy (Figure 2E), and cataloged in NeuroSC (Koonce et al.,

2024; White et al., 1986). Consistent with this, we observed that endogenous GFP::RAB-3 in ASE

similarly localizes in a punctate pattern along the axons in a pattern that is reminiscent to that

seen for endogenous EAT-4::GFP (Figure S2 and Figure 2E).

To expand the utility of this tool for multicolor imaging, we also generated a red-shifted FLP-on

reporter for the *eat-4/VGLUT* gene by inserting mRuby3 (Bajar et al., 2016) with a *C. elegans*-

optimized sequence at the same C-terminal site (Figure 1B). These spectrally distinct reporters,

when combined with the previously developed *eat-4* conditional knockout (Lopez-Cruz et al.,

2019) provide a comprehensive toolkit for dissecting glutamatergic transmission in a cell-specific
manner *in vivo*.

Cell-specific imaging and silencing of GABAergic neurotransmission

GABAergic neurons package GABA into synaptic vesicles via the conserved vesicular GABA
transporter VGAT(Chaudhry et al., 1998; McIntire et al., 1997). In *C. elegans*, the VGAT homolog
UNC-47 is expressed in 11 of the 118 neuronal classes (Gendrel et al., 2016; Wang et al., 2024).
Based on *in vivo* data, the N-terminus of VGAT is cytoplasmic while the C-terminus is luminal
(Martens et al., 2008). The N-terminus contains dileucine motifs critical for proper trafficking
(Santos et al., 2013), and to preserve transporter function we focused on tagging long cytoplasmic
loops. Structural predictions from AlphaFold indicate that UNC-47 has 11 transmembrane
domains (Figure 3A), and we identified the cytosolic loop between transmembrane domains 2
and 3— a long (13 amino acids) region (Figure S3A-B)—as an optimal tagging site.
201
Within this loop, AlphaFold predicts two beta-sheet regions with high confidence. We inserted
GFP between amino acids E145 and N146, immediately following the first predicted beta sheet,
to avoid disrupting secondary structures (Figure 3A, S3A-B). To assess functionality of the newly
engineered UNC-47/VGAT::GFP strain, we performed thrashing assays on *unc-47 (e307)*
mutants, which show impaired locomotion due to loss of GABA signaling at neuromuscular
junctions (McIntire, Jorgensen, Kaplan, et al., 1993). Expression of UNC-47::GFP from an
extrachromosomal array rescued the thrashing defect to wild-type levels (Figure S3C and D), as
expected. We next generated endogenous knock-ins of GFP and mKate2 at the same site but in
the endogenous *unc-47* locus, and observed wild-type locomotion for these strains, consistent
with the insertion of the fluorophores not affecting endogenous function of the transporter (Figure
S3D). These strains showed bright, punctate fluorescence in the nerve ring and along the dorsal
and ventral nerve cords (Figure S3E-F), consistent with previously reported *unc-47* expression
patterns (McIntire et al., 1997). Together, these results demonstrate that inserting a fluorescent
protein at position E145 results in a functionally tagged UNC-47/VGAT reporter that enables
endogenous visualization of the protein.
217
To enable *in vivo* visualization of GABAergic vesicles in single-cells, we next generated two UNC-
47::splitGFP alleles by inserting either one or three tandem copies of GFP11 at the E145 position
(Figure 3B). We used the splitGFP approach to avoid disruptions of the protein structures due to
the introduction of the FRT-cassettes at an internal sequence site. By leveraging the self-

assembling property of the GFP beta barrel, knock-in of the eleventh beta strand (GFP11) results
in labeling of a protein that is only visible when the complementary GFP1-10 is expressed in the
same cell. This property results in a combinatorial labeling strategy, in which cell-specific labeling
is achieved only in those cells that express both the GFP11 and the GFP 1-10 (He et al., 2019).
To validate these tools, we first achieved pan-cellular expression of GFP1-10 (*eft-3* promoter)
(Seydoux & Fire, 1994) in animals carrying the UNC-47::GFP11x3 *(syb7849)* allele. We observed
GABAergic synapses throughout the nerve ring and nerve cords (Figure 3C), similar to full-body
knock-in strains (Figure S3E-F). To then visualize GABAergic vesicles in subsets of cells, we
expressed GFP1-10 in the GABAergic DD motor neurons using the *flp-13* promoter and in animals
carrying the GFP11 (*syb7313*) or UNC-47::GFP11x3 *(syb7849)* alleles. We observed punctate
reconstituted signal in the dorsal nerve cord of both GFP11 and GFP11x3 strains, consistent with
the known distribution of DD synapses (Figure S4E). The triple GFP11 version produced
significantly brighter signal (Figure S4E-G), in line with reports of enhanced fluorescence from
multimerized tags (He et al., 2019).
236
To then visualize GABAergic synapses in single cells, we expressed GFP1-10 under an RIB-
interneuron, cell-specific promoter. We selected RIB because it is a GABAergic interneuron
embedded in the nerve ring and proximal to three other GABAergic interneurons with overlapping
neurites that impede visualization of RIB-specific synapses *in vivo* when using traditional
approaches (Figure S4A-C). Reconstituted fluorescence using our tools enabled visualization of
RIB-specific synapses, and the observed synaptic pattern was consistent with the pattern
expected from electron microscopy reconstructions (Figure 3D) (Koonce et al., 2024; White et al.,
1986) and from expression of endogenous mScarlet::RAB-3 in RIB neurons (Figure S4D). These
findings underscore the value of the tool in labeling GABAergic synapses in individual cells *in*
vivo.
247
To functionally dissect GABAergic transmission, we developed a conditional knockout of *unc-25*,
the gene encoding Glutamic Acid Decarboxylase (GAD), which catalyzes GABA synthesis. We
decided to target *unc-25*/GAD because it results in the elimination of GABA from neurons
(Gendrel et al., 2016). We flanked the *unc-25* coding sequence with FRT sites and inserted a
nuclear mCherry reporter downstream to indicate successful recombination (Figure 3B). We then
validated the tool by using thrashing assays. Consistent with previous findings, *unc-25* null
mutants show severely reduced locomotion in the thrashing assays (McIntire, Jorgensen, Kaplan,
et al., 1993) (Figure 3E). We observed that animals carrying the conditional knockout allele

behaved normally in the absence of Flippase, but that pan-cellular Flippase expression in all
GABAergic neurons (*unc-47* promoter), which is expected to result in loss of GABAergic
neurotransmission, phenocopied the thrashing defect of *unc-25* mutants (Figure 3E). These
results confirm that this tool effectively eliminates *unc-25/*GAD activity, with the capacity to be
activated in a cell-specific manner and allows investigation of how GABAergic transmission
contributes to neural function and behavior.

Functional labeling of cholinergic vesicles via UNC-17/VAChT

The cholinergic identity of neurons is defined by the expression of a conserved gene locus, shared
from nematodes to vertebrates, that includes both the acetylcholine synthesis enzyme Choline
Acetyltransferase (ChAT) and the Vesicular Acetylcholine Transporter (VAChT) (Eiden, 1998). In
C. elegans, the VAChT homolog UNC-17 is expressed in 57 of the 118 neuronal classes (Wang
et al., 2024). Based on prior studies, both the N- and C-termini of VAChT face the cytosol and
contain regulatory motifs important for trafficking (Fei et al., 2008). AlphaFold predicts UNC-17
has 12 transmembrane domains, but most cytosolic loops are very short (<5 amino acids), except
for the third cytoplasmic loop, which contains 28 amino acids and exhibits relatively high sequence
conservation (Figure S5A-B).

To determine suitable tagging sites for imaging UNC-17 without disrupting function, we tested
three locations: the conserved third cytosolic loop (site 1), the N-terminus (site 2), and the C-
terminus (site 3) (Figure S5B-C and S5E). We used a thrashing assay to assess function, as *unc-*
17(e245) null mutants fail to thrash in liquid and are rescued by the re-expression of untagged
UNC-17 under its own promoter (Figure S5F). We generated transgenic strains with a rescue
array containing the *unc-17* gene with GFP inserted into these three sites. We observed that GFP
insertion into site 1 failed to rescue the phenotype, suggesting disruption of UNC-17 function. In
contrast, tagging either the N-terminus or C-terminus restored normal thrashing (Figure S5E–F),
indicating that these positions tolerate modification. Informed by these rescue experiments, we
then generated two FLP-on conditional knockout alleles (Schwartz & Jorgensen, 2016) with GFP
inserted at either the N-terminus (*syb7251*) or C-terminus (*ola503*) (Figure 4A-B). To test whether
the FLP-on cassettes affected protein function, we examined behavior before Flippase expression
and found that both alleles behaved like wild-type animals (Figure S5F). Because splicing
regulatory elements are located near the 5' end of the *unc-17* gene and are required for successful
splicing of the cholinergic locus (both *unc-17* and *cha-1* transcripts) (Mathews et al., 2015), we

proceeded with the C-terminally tagged *ola503* allele, which leaves the 5' region intact. We tested
whether pan-cellular expression of Flippase in this strain impairs behavior. Animals with global
GFP-tagged UNC-17 showed wild-type thrashing in liquid (Figure 4C), confirming that the tagged
transporter is functional and validating this approach for cell-specific labeling of cholinergic vesicle
pools.

At the cellular level, GFP labeling of UNC-17/VAChT using the *ola503* allele and pan-cellular
Flippase expression (driven by the *eft-3* promoter) (Seydoux & Fire, 1994) resulted in
fluorescence in the nerve ring, dorsal cord, ventral cord, and sublateral cords (Figure 4D). This
expression pattern matched that of a full-body mKate2 knock-in of endogenous UNC-17/VAChT
(Figure S5D) (Pereira et al., 2019) and was consistent with prior transcriptional reporters (Pereira
et al., 2015) and anti-UNC-17 antibody staining (Duerr et al., 2008). The *ola503* allele also enables
cell-specific labeling. When Flippase was expressed specifically in ADF neurons using the P*srh-*
142 promoter (Maicas et al., 2021), we observed UNC-17::GFP puncta along the ADF axon
(Figure 4E). This punctate pattern aligned with known ADF presynaptic sites from electron
microscopy reconstructions of L4 animals (Koonce et al., 2024; White et al., 1986). Together,
these results demonstrate that the UNC-17::GFP FLP-on allele provides a reliable tool to label
cholinergic vesicles in individual neurons *in vivo*, and that this labeling does not affect function in
our assays. To enable multicolor imaging, we generated a red, fluorescent version of the UNC-
17/VAChT tool by replacing GFP with *C. elegans*-optimized mRuby3 (Figure 4B). This mRuby-
tagged allele performs as wild type animals in a thrashing assay and labels synaptic structures
along the nerve ring and nerve cords (Figure S6). Combined with the previously described cell-
specific *unc-17* knockout strain (Huang et al., 2023) (Figure 4B), these tools allow precise tracking
and manipulation of cholinergic neurotransmission *in vivo*.

Probing Monoaminergic Transmission

Monoamines such as serotonin, dopamine, norepinephrine, epinephrin, octopamine, and
tyramine are transported into vesicles by the conserved Vesicular Monoamine transporter (Duerr
et al., 1999; Erickson et al., 1992). In *C. elegans*, the VMAT homolog CAT-1 is expressed in 16
of the 118 neuronal classes (Wang et al., 2024). Recently, CAT-1 was tagged at its C-terminus
with a split GFP11x3 reporter (Figure 5A–B) (Huang et al., 2023). When GFP is reconstituted
pan-cellularly, this fusion produces a punctate signal enriched in the nerve ring—including the
characteristic and well-known serotonergic neuron NSM—and in serotonergic neurons that are
part of the reproductive organs (Figure 5C).

To determine whether the CAT-1::GFP11x3 fusion maintains protein function, we used a
behavioral assay based on the role of serotonin to prevent animal exploration on a lawn of bacteria
(Flavell et al., 2013). Since serotonin is packaged into vesicles by CAT-1/VMAT, similar to mutants
of serotonin production (Flavell et al., 2013), *cat-1* mutants display increased exploration behavior
compared to wild-type animals (Figure 5D). We found that animals with reconstituted CAT-
1::GFP11x3 explore bacterial lawns at levels comparable to wild-type (Figure 5D), indicating that
the GFP11x3 tag nor its reconstitution impair CAT-1 function. To complement this tool, we also
developed a cell-specific *cat-1* knockout allele in which the full coding sequence is excised upon
Flippase expression (Figure 5B). Removal of serotonin production (*tph-1*) specifically from ADF
neurons result in increased aversion from wild type *E. coli* bacteria lawns (Feng et al., 2025).
Consistent with this, cell-specific KO of *cat-1* in ADF neurons (via expression of ADF::Flippase)
results in increased *E. coli* aversion (Figure 5E). Together, these tools now enable the cell-specific
tracking and silencing of monoaminergic synapses in living animals.

***In-Vivo* Identification of Co-Transmitter Neurons**

Co-transmission is a conserved feature of neurons across the animal kingdom (Granger et al.,
2017; Trudeau & El Mestikawy., 2018; Vaaga et al., 2014), including *C. elegans* (Duerr et al.,
2008; Gendrel et al., 2016; Pereira et al., 2015; Pocock & Hobert, 2010; Serrano-Saiz et al., 2017;
Serrano-Saiz et al., 2013; Taylor et al., 2021; Wang et al., 2024), but the prevalence of co-
transmission *in vivo* for any given organism is not well understood. To validate the utility of our
tools and to map the architecture of co-transmission in *C. elegans*, we developed an intersectional
genetic strategy using Flippase recombinase and FRT-flanked fluorescent reporters (Figure 6A).
We focused on identifying neurons that co-transmit glutamate or acetylcholine—the two most
abundant excitatory neurotransmitters in *C. elegans*—in combination with other transmitters
(Figure 6B).

We reasoned that if two neurotransmitters were co-expressed in the same neuron, driving
Flippase under the promoter of one transmitter would activate the conditional reporter—resulting
in fluorescence—only in cells also expressing a second neurotransmitter identity (Figure 6A-B).
To achieve this, we used the engineered alleles for each vesicular transporter that we developed
(Figures 2B, 3B, 4B, and 5B) (Table 1) and developed additional Flippase driver by modifying the
locus of genes involved in the packaging of acetylcholine (*unc-17*) and GABA (*unc-47*) (Table 1).
Additionally, we used available Flippase driver lines for serotonin and dopamine (Figures S7A–B,
Table 1) (Muñoz-Jiménez et al., 2017).

We first used a conditional *eat-4/VGLUT* reporter strain in which cytosolic mCherry is expressed
upon Flippase-mediated recombination (Lopez-Cruz et al., 2019). When Flippase was driven by
the *unc-17/VAChT* promoter (cholinergic), we observed five mCherry-positive neurons in the head
and tail, consistent with co-expression of *unc-17/VAChT* and *eat-4/VGLUT*. Based on cell
position, neurite morphology, transcriptomic data (Taylor et al., 2021), and anatomical maps
(Wang et al., 2024), we identified these neurons as AFDL, AFDR, M5, DVA, and PVN (Figure
6C–D).

Flippase expression from the GABAergic *unc-47* promoter activated *eat-4*-driven mCherry
expression in a single pharyngeal neuron, identified as I2L (Figure 6E, S8A). Driving Flippase
from the dopaminergic *dat-1* promoter labeled the PDE neuron (Figure 6F, S8B), while
serotonergic *tph-1*-driven Flippase did not produce any mCherry-positive neurons. These results
are summarized in Figure S7C.

We applied a similar strategy by using our Flippase-dependent *unc-17*::GFP reporter to identify
candidate neurons that co-release acetylcholine with other neurotransmitters. In this context,
when Flippase was driven from the GABAergic *unc-47* locus, we observed GFP expression in the
M4, SDQR, and SMD neurons (Figure 6G, S8D). Flippase expression from the serotonergic *tph-
1* promoter revealed previously described acetylcholine/serotonin co-transmitting neurons,
including ADF, HSN, and VC4/VC5 (Figure 6H, S8C), consistent with prior findings (Pereira et al.,
2015). All neurons identified through this dual-reporter approach are summarized in Figure S7C.

We compared our findings with previous reports and have compiled a list of potential co-
transmitter neurons that are consistently identified across independent studies (Figure S9),
suggesting these are likely *bona fide* co-transmitter neurons. Together, we observe that *C.
elegans* has 35 neurons exhibiting molecular signatures of co-transmission (Table 2)—
representing ~ 10% of the *C. elegans* nervous system (Figure S9 and Table 3). Strikingly, the
pharyngeal nervous system—analogous to the vertebrate enteric nervous system (Albertson &
Thomson, 1976)—had the highest density of co-transmitter neurons: 30% (6 of 20 neurons)
displayed co-expression of multiple vesicular transporters. Across the entire nervous system, co-
transmission was prevalent among sensory neurons, with 12% (10 of 83), compared to 11% of
interneurons (9 of 81) and 7% of motor neurons (8 of 116) (Table 2). We also compiled a list of
neurons that were suggested to be co-transmitters (RIB, AVA, AVB) (Gendrel et al., 2016), but
that are not supported by other expression studies (Taylor et al., 2021) (Table 4).

In-vivo visualization of co-transmitter synapses in the ADF chemosensory neuron

We next used our toolkit to investigate the subcellular localization of vesicular transporters in a
co-transmitting neuron, ADF. The ADF neurons are a bilaterally symmetric pair of sensory
neurons in *C. elegans* known to regulate food exploration, chemotaxis and entry into the lethargic-
like dauer state (Bargmann & Horvitz, 1991). While ADF has long been known to use serotonergic
neurotransmission, our findings indicate that it is also capable of acetylcholine synthesis and
packaging (Figure 6H, S7C and S10A-B). Our findings are consistent with recent transcriptomic
and reporter-based studies (Pereira et al., 2015; Taylor et al., 2021; Wang et al., 2024).

To examine the subcellular distribution of serotonin or acetylcholine vesicular transporters in ADF,
we used previously developed tools to endogenously label the serotonin vesicular transporter
CAT-1/VMAT (Huang et al., 2023), and the acetylcholine vesicular transporter UNC-17/VAChT
with GFP (Figures 4 and 5). To label these vesicular transporters specifically in ADF, we drove
Flippase and GFP1-10 expression using the AFD-specific promoter, *srh-142* promoter (Maicas et
al., 2021). Both UNC-17/VAChT and CAT-1/VMAT displayed punctate fluorescence along the
ADF axon, consistent with the location of presynaptic sites expected from electron microscopy-
based 3D reconstructions (Figure S10C) (Koonce et al., 2024; White et al., 1986).

To then visualize the spatial relationship between these two vesicular transporters, we generated
a strain in which UNC-17/VAChT was tagged with mRuby and CAT-1/VMAT with GFP at their
respective endogenous loci, cell-specifically in ADF neurons (Figure 7A). *In vivo i*maging using a
spinning disk confocal microscope revealed that these transporters frequently co-localize within
the same synaptic boutons (Figure 7B and 7B'). Interestingly, we also found instances in which
the UNC-17/VAChT::mRuby and CAT-1/VMAT::GFP signals partially segregate into separate
boutons along the same axon (Figure 7C and 7C'), suggesting that these vesicular transporters
can be sorted into distinct vesicle populations. Consistent with the idea that UNC-17 and CAT-1
localizes to synaptic vesicles along the axon, we observed that UNC-17 and CAT-1 localization
was disrupted in a Kif1A/*unc-104* (*e1265*) mutant animal (Hall & Hedgecock, 1991), absent in the
axon and accumulated in the cell body (Figure 7D-D').
412

To test whether both transporters are present at all ADF synapses, we endogenously tagged the
active zone protein UNC-13/Munc13 with mScarlet and examined its spatial relationship to UNC-
17/VAChT::GFP and CAT-1/VMAT::GFP (Figure 7E). We observed active zone UNC-

13::mScarlet puncta that lacked either CAT-1/VMAT::GFP (Figure 7F) or UNC-17/VAChT::GFP

labeling (Figure 7G). These findings are consistent with the idea that, while these two vesicular

transporters display co-localization in synaptic varicosities, they also independently localize to

distinct subcellular compartments.

Synaptic varicosities in *C. elegans* can be within the diffraction limits of light microscopy,

preventing differentiation of co-localizing vesicular populations. To better understand the relative

distribution of these vesicular transporters, we next visualized ADF synapses with increased

resolution, using AiryScan imaging, which can achieve differentiation of fluorophores up to 120

nm apart (Wu & Hammer, 2021). We observed that, even in synaptic boutons in which the

vesicular transporters were observed to co-localize with traditional light microscopy methods,

UNC-17/VAChT::mRuby and CAT-1/VMAT::GFP differentially segregated when imaged using

AiryScan microscopy (Figure 7H).

Together these results suggest that acetylcholine and serotonin co-localize to synapses, but might

be packaged into distinct vesicles with specific synaptic subcellular localization that is detectable

upon super-resolution microscopy. Our findings underscore the importance of endogenous

labeling in determining the specific localization of these vesicular transporters and their use with

higher-resolution imaging methods, highlighting the value of the tools developed in this study to

understand the cell-biological organization of synapses *in vivo*, particularly for neurons using more

than one neurotransmitter.

DISCUSSION

The integration of anatomical connectivity, molecular identity, neural activity, and transmitter

usage provides a powerful framework for building models of neural circuit function. The *C.*

elegans community has access to a complete connectome (White et al., 1986); the cellular identity

of all neurons (Sulston & Horvitz, 1977; Sulston et al., 1983); whole-brain calcium imaging

(Nguyen et al., 2016; Prevedel et al., 2014; Schrödel et al., 2013); single-cell transcriptomic

profiles (Taylor et al., 2021); and a full neurotransmitter identity map for all neurons (Wang et al.,

2024). These datasets have inspired models describing how specific circuits may give rise to

behavior. However, validating these models *in vivo* requires tools that can precisely manipulate

the molecular components of individual synaptic connections (Dag et al., 2023; Hawk et al., 2018;

Kumar et al., 2024). SynaptoTagMe provides that missing capability (Table 1) for the

neurotransmitter systems that cover approximately 90% of the *C. elegans* nervous system
(GABA, glutamate, acetylcholine, and the monoamines). By enabling cell-specific labeling and
conditional knockout of vesicular transporters, we can now directly test the contribution of
individual neurotransmitters within defined circuits and link those changes to behavioral
outcomes. Moreover, due to the evolutionary conservation of vesicular transporters, the *in vivo*
validation of tagging strategies will help identify suitable labeling strategies for other organisms,
providing a path toward comparative and cross-species studies of synaptic dynamics based on
neurotransmitter identity.

Co-transmission is a conserved feature of neural systems across the animal kingdom (Granger
et al., 2017; Lacin et al., 2019; Vaaga et al., 2014), but its preponderance *in vivo*, its regulation
and its functional significance is still an area of active research. Using *in-vivo* reporters subject to
endogenous regulation, and contrasting our results with previous studies, we determine that more
than 10% of *C. elegans* neurons have co-transmission potential (Figure S9) (Tables 2 and 3). Our
in vivo characterization of co-transmitting neurons confirm and expand findings reported for the
latest neurotransmitter atlas of *C. elegans* (Wang et al., 2024) and yield three key insights. First,
co-transmission occurs throughout the nervous system of *C. elegans*, including both the
pharyngeal (enteric-like) and more central nervous systems, like the nerve ring and nerve cords
(Table 2). Second, neurons can co-transmit multiple neurotransmitters in specific combinations
that are conserved from nematodes to mammals (Figure 6 and S9) (Granger et al., 2017; Lacin
et al., 2019; Trudeau & El Mestikawy., 2018; Vaaga et al., 2014; Wang et al., 2024). Importantly,
the same neurons consistently exhibit co-transmission of the same neurotransmitter identities
across individual animals, consistent with co-transmitter identity mapping to neuronal identity
(Figure 6D-H). Third, co-transmission is part of every layer of a circuit, from sensory neurons to
interneurons and motor neurons (Table 2). This is especially interesting in light of recent studies
showing that co-transmission in sensory and motor circuits can be modulated by environmental
cues such as stress (Bertuzzi et al., 2018; Li et al., 2024; Pocock & Hobert, 2010) or light-dark
cycles (Chen et al., 2023; Maddaloni et al., 2024). With the tools developed here - based on
endogenously tagged vesicular transporters – it is now possible to monitor the dynamic
expression and subcellular distribution of specific vesicle populations *in vivo* and what molecular
mechanisms drive those changes.

Our conclusions are supported by independent and convergent lines of evidence, and the toolkit
developed here enables direct empirical interrogation of both the existence and functional

relevance of co-transmission *in vivo*. While expression of vesicular transporters represents one
line of evidence for co-transmission potential, it does not by itself establish functional co-release,
which additionally depends on neurotransmitter biogenesis, release competence, and activity-
dependent regulatory mechanisms. SynaptoTagMe now make it possible to systematically test
these requirements.
489
We note that the current characterization of co-transmitting neurons might be an under-estimate
of the total number of neurons which use co-transmission. For example, it has been proposed
that additional neurotransmitters, like betaine, may function in the *C. elegans* nervous system
(Wang et al., 2024). Accounting for neurons that express proteins capable of synthesizing or
packaging betaine, the proportion of potential co-transmitter neurons may exceed 20% of the
whole nervous system of *C. elegans*. Our characterization of co-transmission focused on the *C.*
elegans adult hermaphrodite, and co-transmitting neuron identities could be developmentally
regulated, or modulated based on prior experience. Consistent with this, it has been observed
that the identity of co-transmitting neurons is different between males and hermaphrodites
(Serrano-Saiz et al., 2017), underscoring the importance of future examination of the plasticity
and developmental regulation (Pereira et al., 2019; Pereira et al., 2015) of co-transmitting capacity
for individual neurons.
502
Expression of a vesicular transporter, while consistent with co-transmitting capacity, is not
conclusive for the existence of co-transmission for that neuron. For example, we identified co-
expression of the GABA and Glutamate vesicular transporters in the pharyngeal neurons I2
(Figure S8A). Notably, I2 does not express the GABA synthesis enzyme, *unc-25*/GAD (McIntire,
Jorgensen, & Horvitz, 1993) or the GABA re-uptake transporter, *snf-11* (Mullen et al., 2006). Thus,
it is unlikely that it produces GABA or uptakes it from the extracellular space. VGAT/UNC-47 has
also been reported to transport neurotransmitters such as glycine (Aubrey et al., 2007) and beta-
alanine (Juge et al., 2013), raising the possibility that I2 could co-transmit glutamate with an
unconventional neurotransmitter. Additionally, it is important to mention that the identification of
eat-4-positive neurons through the replacement of the *eat-4* coding sequence and introns (Figure
S7A) could result in the elimination of regulatory sequences. Thus, we conceptualize the list of
co-transmitting neurons as a hypothesis-generating framework to be further examined with the
tools developed in this study.
516

Our observations of the identity of co-transmitting neurons, and the specific combinations
represented in the neurons, suggest that there may be transmitter-specific rules of synaptic
biology important for circuit function (Silm et al., 2019; Trudeau & El Mestikawy., 2018). This is
consistent with findings in vertebrates, in which specific neurotransmitter combinations and their
distributions could underpin specific features of circuit function. For example, in Starburst
Amacrine cells in the mammalian retina, acetylcholine and GABA (O'Malley et al., 1992) are
packaged into distinct vesicle pools that exhibit different calcium sensitivities for release (Lee et
al., 2010). The distribution of specific vesicular populations and their release probabilities might
constitute an architecture that helps encode the sensory signals processed by Starburst Amacrine
cells. We similarly hypothesize that the specific distribution of co-transmitting synapses across
the *C. elegans* connectome, and the identities of the neurotransmitters used, might help encode
features important for circuit function and animal behavior. By allowing longitudinal, cell-specific
monitoring of the expression, regulation, and subsynaptic distribution of distinct vesicle
populations in living animals, this toolkit provides a platform for probing when, where, and how
co-transmission is deployed, and for defining the molecular and circuit-level mechanisms that
govern its use under different physiological and environmental conditions.
533
534   **METHODS**
535
Strains
Worms were maintained at 20°C using standard techniques (Brenner, 1973). Strains were
maintained on NGM plates seeded with *E. coli* (OP-50). The wild type (WT) is N2, and only
hermaphrodite worms were used for this study. A complete list of strains appears below. Strains
that were developed in this study as part of the SynaptoTagMe toolkit and appear in Table 1 can
be requested through CGC. The necessary sequencing information can be found at:
https://www.intralab.app/research-papers/cuentas-condori_etal-2026.
543
Generation of new alleles
For the strains engineered by Sunybiotech, as described below, strain design was
performed in the Colón-Ramos lab by Andrea Cuentas-Condori.
Sunybiotech used CRISPR/Cas9 to insert GFP FLP-on cassettes (Schwartz & Jorgensen,
2016) at either the N-termini of the *unc-17* locus (*syb7251*) or C-termini end of the *unc-17*
(ola503), or *eat-4 (syb8568)* locus, according to sequence design. mRuby FLP-on cassettes were
inserted similarly at the C-termini end of *eat-4* (*syb9193*) and *unc-17* (*syb7882*) locus. To visualize

the fluorescent signal tagged to the protein of interest, Sunybiotech generated single-copy MosCl
strains.

Sunybiotech used CRISPR/Cas9 to add full length GFP (*syb6990*), full length mKate
(*syb7358*) or split GFP (one (*syb7313*) or three (*syb7849*) copies of GFP11) to the *unc-47* locus
at the +893bp position. To visualize the reconstituted GFP signal in RIB neurons, complementary
GFP1-10 was driven with the P*sto-3b* promoter; and to visualize the GFP-reconstituted signal in
DD neurons, GFP1-10 was driven with the P*flp-13* promoter.

Sunybiotech used CRISPR/Cas9 to introduce an FRT site before the +1bp in the *unc-25*
gene locus (*syb5949*). In a second round of CRISPR editing, they introduced *let-858* 3' UTR
followed by a second FRT and nuclear mCherry (*syb6275)*.

cat-1 (*ky1101 ky1118*) cell-specific knock out strain was created using CRISPR/Cas9 to
introduce an FRT site immediately before the ATG in the *cat-1* gene locus (*ky1101*). A second
round of CRISPR editing in that strain introduced the *let-858* 3' UTR followed by a second FRT
and the mCherry coding region immediately after the stop codon of *cat-1* to generate *ky1118*.

Sunybiotech used CRISPR/Cas9 to introduce a T2A::Flippase sequence before the STOP
codon of *unc-47(syb8125)* and *unc-17(syb8059)*. All strains generated using CRISPR/Cas9 were
outcrossed twice before use.

Molecular Biology

Plasmids were constructed using Gibson cloning. First, Snapgene (Version 7.0.3)
software was used to design primers targeting the desired DNA vector backbone and DNA insert.
The vector backbone and DNA insert were PCR linearized and amplified using "CloneAmp HiFi
PCR Premix." To assemble the desired plasmid, the purified vector backbone DNA and insert
DNA were combined and incubated in solution with "2x Gibson Assembly Enzyme Premix."
Following incubation, the reaction mixture was used to transform Stellar Competent Cells, which
were subsequently plated and grown overnight on LB-Amp plates. All plasmids were verified with
Sanger sequencing.

Protein alignment and structure visualization

For each gene under study, NCBI BLAST was used to generate a protein sequence
alignment of the *C. elegans* gene with the closest orthologs from the other model organisms *M.*
musculus, *D. rerio*, *D. melanogaster*, and *H. sapiens*. Protein structure models for the C. elegans
genes were downloaded from the AlphaFold database (Jumper et al., 2021) and predicted models

for the CRISPR-Cas9 modified genes including fluorophores were generated by the Alphafold3
online server (Abramson et al., 2024). Visualization and image generation of protein structures
was done using the ChimeraX software (Pettersen et al., 2021). To color the structures by
sequence conservation, the alignments per gene were overlaid onto the structures with ChimeraX
and colored by resulting sequence conservation Z-scores as calculated by the AL2CO algorithm
(Pei & Grishin, 2001) within the software.

Table 5 – List of all molecular locations tested for labeling each synaptic vesicle
transporter

Gene	AA tested for fluorescent protein insertion	Functional tag?
eat-4	N-terminus (After START codon)	yes
unc-47	Between E145 – N146	yes
unc-17	Between AA P6-V7	yes
unc-17	Between AA N242 - P243	no
unc-17	C-terminus (Before STOP codon)	yes

Microscopy

Larval or young adult animals were immobilized on 2-10% agarose pads with 10mM
levamisole. A Nikon Ti2 microscope equipped with a CSU-W1 spinning disk head, ORCA-Fusion
BT SCMOS camera, high-speed piezo stage motor, 60X/1.40 Apo Lambda oil objective lens was
used for live imaging. Z-stack images were collected (0.3-0.5 µm/step), spanning the focal depth
of the nerve cord and nerve ring synapses. A Zeiss LSM880 microscope equipped with an
AiryScan detector and 63X NA 1.4 oil objective was used for AiryScan imaging. FIJI (Schindelin
et al., 2012) or NIS Elements AR analysis software (version 6.10.01) were used to create
maximum intensity projections and 2D renderings.

Thrashing Assay

C. elegans were raised at 20°C under standard laboratory conditions on agar plates
seeded with a lawn of *E. coli* (OP50). Worms were synchronously grown to L4-stage and placed
in individual wells of a Corning™ PYREX™ Spot Plate (Catalogue #722085) containing 1000 µl
of M9 buffer, ensuring the buffer remained within the well's borders. After a 30-second acclimation
period to M9, thrashes were manually counted for 1 min. A single thrash was defined as a change
in the direction of the worm's midbody bending, counting each time the worm's body flexed to one
side. Following each trial, the worm was removed using a pipette and disposed of, and the M9
buffer was absorbed and discarded. The well was then cleaned with 70% ethanol and wiped dry.
To avoid bias, the counter was blinded to each genotype. Each worm was tested only once, with

assays conducted on 10 worms per genotype per day, and repeated over 2–3 days to account
for potential day-to-day environmental variations.
616
Chemotaxis Assay
Worms were maintained at 20°C for at least two generations on Nematode Growth
Medium (NGM) seeded with OP50 *Escherichia coli* bacteria. The concentration of our attractant
(NaCl) is approximately 50 mM in NGM plates. "Training" plates were produced using NGM with
the further addition of 50 mM NaCl to a total concentration of 100 mM, then also seeded with
OP50.
Chemotaxis assay was modified from standard procedure (Ward, 1973). All assays were
performed on 50 mm diameter plates. Unseeded NGM plates were marked at the center and one
point 12.5 mm away from the center. A ~60-85 mM gradient of NaCl was created between the
center and outer point by adding 5 M NaCl at the outer point as drops of 4μL (20-24 hours before
the assay), 4μL (5 hours before), and 1.6μL (2 hours before); a sham gradient was created using
only water. Gradient prediction was determined as previously described (Crank, 1956; Pierce-
Shimomura et al., 1999); briefly, for every point some distance r in cm from the salt peak, the
concentration C in mM at any point in time was calculated as:
$$C(t_i, r) = C_o + \sum_{i=1}^{n} c(t_i, r) \qquad (1)$$
where C_o is the initial concentration of NaCl in the agar (50 mM), n is the drop number, and t_i is
the time in seconds since the drop had been applied; the contribution from each drop, in turn,
was calculated as:

$$c(t_i, r) = 10^6 \frac{N_i}{\sqrt{4\pi dD(t_i)}} e^{-\left(\frac{r^2}{4D(t_i)}\right)} \qquad (2)$$

where N_i is the moles of NaCl added per drop; d is the depth (cm) of the agar; and D is $1.590 \times$
10^{-5} cm^2/s, the diffusion coefficient for 5 M NaCl through an aqueous medium (Robinson RA,
1959). The resulting gradients were validated by electrical conductivity measurements using an
Oakton CON 6+ Handheld Conductivity Meter with a custom conductivity probe with 1 mm
insertion depth (Micro-electrodes, Inc, Bedford, NH). The conductivity readings from 50 mM and
100 mM NaCl NGM plates were used for calibration at specific room temperatures.
The day before experiments, L4 animals were transferred to a seeded NGM plate to
synchronize worms by developmental stage. At 5 hours before each assay, worms were
transferred to a training plate using standard NGM plate recipe adjusted to 100 mM NaCl. After
training, 8 worms were picked, with preference to those on the bacterial lawn, washed sequentially
in two 100 μL drops of liquid NGM buffer (25 mM potassium phosphate pH 6, 1 mM CaCl2, 1 mM

MgSO4, 50 mM NaCl) to remove adherent OP50, and transferred to a single 2 µL drop of NGM buffer at the center of the assay plate with prepared NaCl gradient as described. Data collection began when the water drop was fully absorbed into the assay plate and the first worm began to migrate from its starting point. Six assay plates were imaged for each strain across two separate days, yielding a total of 48 worms imaged per strain.

Images of chemotaxis behavior were acquired at 3.75 fps for 7 minutes using a Basler acA2440-35mm monochromatic sensor with an infrared filter on a commercially available WormLab imaging system and computer running WormLab 2023.1.1 software (MBF Bioscience LLC, Williston, VT USA). Individual worm position data was obtained by constructing tracks in WormLab software, then analyzed using custom scripts in R 4.4.1 (can be accessed through GitHub (https://github.com/colonramoslab/Cuentas-Condori-et-al.-2025-Toolkit-). The assay outcome was defined as the mean distance from the peak of the salt gradient for each worm, averaged over every available frame in the last minute of the assay. When a worm track was interrupted, e.g. by a worm exiting the camera field of view or by two worms intersecting, the last available position for the worm was repeated until the worm was re-detected.

Roaming Assay

Roaming assay plates were prepared 3-5 days prior to the experiment by seeding NGM agar plates with *E. coli* (OP50) culture no older than 2 days. Plates were seeded using a sterile glass rod to spread the bacteria evenly across each plate. Plates were left to dry completely between 3-5 days at room temperature to ensure the bacteria layer was fully dry, thereby allowing for visible worm tracks during the assay. *C. elegans* were raised at 20°C under standard laboratory conditions on agar plates seeded with a lawn of *E. coli* (OP50). On the day before the assay, worms synchronously grown to L4-stage were transferred to regular seeded plates and stored at 20°C. After approximately 10 hours, worms were transferred to individual assay plates and incubated at 20°C for 16 hours. After this time, worms were removed and a grid overlay (3 mm x 3 mm squares) covering the assay plate was used to count the number of squares the worms had traversed during the incubation period. The number of squares crossed provided a quantifiable measure of roaming activity. To avoid bias, the counter was blinded to each genotype. Each worm was tested only once, with assays conducted on 10 worms per genotype per day and repeated over 2–3 days to account for potential day-to-day environmental variations.

Aversion behavior Assay

Aversion behavior assay were performed as previously described (Feng et al.,
2025). Animals were fed on *E. coli* BW25113 for at least three generations before the behavioral
assay. 12.5 µL of overnight BW25113 cultures were seeded onto standard NGM agar plates,
grown at 37˚C incubator for 24 hours and then left at room temperature for another 24
hours. 15~20 animals at L4 stages from each genotype were transferred onto behavioral assay
plate and recorded at 21˚C for 20 hours, at a recording rate of 1 frame per minute. Biological
replicates across two different days were conducted. Videos were cropped and analyzed using
standard MatLab codes (Marquina-Solis et al., 2024). Aversion ratio was defined by the number
of worms outside the bacterial lawn over total number of worms on assay plates.
689
Statistical Analysis
We used the Shapiro-Wilk test to determine sample distribution. For comparisons between
2 normally distributed groups, Student's T-test was used and $p<0.05$ was considered significant.
ANOVA was used to compare between 3 or more normally distributed groups followed by
Dunnett's multiple-comparison test. If the samples were not normally distributed, we used a Mann-
Whitney test to compare two groups and a Kruskal-Wallis test to compare three or more groups.
Specific post-hoc statistical tests are listed in the figure legend of each experiment. Prism 10.4.2
was used to graph the data and for all statistical analysis.
698
699　**List of strains**

Strain name	Genotype
PHX8568	*eat-4::gfp FLP-on(syb8568)* III
DCR9575	*eat-4::gfp FLP-on(syb8568)* III 2X outcrossed
MT6308	*eat-4(ky5)* III
DCR9690	*Pflp-6::FLP(sybIs9606)* II
DCR9814	*Pflp-6::FLP(sybIs9606)* II 2X outcrossed
DCR9872	*eat-4::gfp FLP-on(syb8568)* III*; Peft-3::FLP (sybIs9614)* II
DCR9816	*eat-4::gfp FLP-on(syb8568)* III*; Pflp-6::FLP (sybIs9606)* II
DCR9681	*eat-4::mRuby FLP-on(syb9193)* III 2X outcrossed
DCR9963	*GFP::RAB-3 FLP-on(ox699)* III; *Pgcy-5::FLP (sybIs8828)*
DCR9965	*GFP::RAB-3 FLP-on(ox699)* III; *olaEx5706[*P*gcy-7*::Flippase::T2A::BFP*;* Punc-122::RFP]*

DCR9210	*unc-47::gfp (syb6990)* III 2X outcrossed
DCR9453	*unc-47::mKate2 (syb7358)* III 2X outcrossed
CB307	*unc-47(e307)* III
DCR9269	*unc-47::gfp (syb6990)* III*; olaEx5490[Psto-3b::BFP; Punc-122::RFP]*
PHX7313	*unc-47::gfp11 (syb7313)* III
DCR9280	*unc-47::gfp11 (syb7313)* III 1X outcrossed
DCR9738	*unc-47::gfp11 (syb7313)* III*; olaEx5686[Pflp-13::GFP1-10; Pmyo-2::mCherry]*
PHX7849	*unc-47::gfp11x3 (syb7849)* III
DCR9739	*unc-47::gfp11x3 (syb7849)* III 2X outcrossed
DCR9740	*unc-47::gfp11x3 (syb7849)* III*; olaEx5686[Pflp-13::GFP1-10; Pmyo-2::mCherry]*
DCR9741	*unc-47::gfp11x3 (syb7849)* III*; olaEx5687[Psto-3b::GFP1-10; Punc-122::RFP]*
CF4587	*muIs253 [Peft-3*::GFP1-10::*unc-54* 3'UTR + *Cbr-unc-119*(+)] II; *unc-119(ed3)* III
DCR9838	*muIs253 [Peft-3*::GFP1-10::*unc-54* 3'UTR + *Cbr-unc-119*(+)] II; *unc-119(ed3)* III *unc-47::gfp11x3 (syb7849)* III recombinant
PHX6275	*unc-25 (syb5949 syb6275)* III
DCR9898	*unc-25 (syb5949 syb6275)* III; bqSi506[P*rgef-1*::Flippase D5 + unc-119(+)] IV
CB933	*unc-17(e245)* IV
DCR9891	*unc-17(e245)* IV*; olaEx5703[Punc-17::*UNC-17cDNA]
DCR9074	*unc-17(e245)* IV*; olaEx5400[Punc-17::*GFP::UNC-17 (N-terminal tag)]
DCR9892	*unc-17(e245)* IV*; olaEx5704[Punc-17::*UNC-17::GFP (between TM6-7)]
OH15568	*unc-17::mKate2 (ot907)* IV
DCR9011	*unc-17::GFP FLP-on(ola503)* IV
DCR9211	*unc-17::GFP FLP-on(ola503)* IV 1X outcrossed
PHX7251	*GFP FLP-on::unc-17 (syb7251)* IV
DCR9265	*GFP FLP-on::unc-17 (syb7251)* IV 1X outcrossed
DCR9720	*Peft-3::FLP (sybIs9614)* II
DCR9733	*Peft-3::FLP (sybIs9614)* II *2X outcrossed*

DCR9742	*unc-17::GFP FLP-on(ola503)* IV*; Peft-3::FLP (sybIs9614)* II
DCR9374	*unc-17::mRuby FLP-on(syb7882)* IV 2X outcrossed
DCR9989	*unc-17::mRuby FLP-on(syb7882)* IV*; Peft-3::FLP (sybIs9614)* II
PHX7239	*cat-1::gfp11x3(syb7239)* X
DCR9370	*cat-1::gfp11x3(syb7239)* X 2X outcrossed
DCR9338	*cat-1::gfp11x3(syb7239)* X*; olaEx5516[Psrh-142::GFP1-10; Psrh-142::BFP; Punc-122::RFP]*
RB681	*cat-1(ok411)*
DCR9837	*muIs253* [P*eft-3*::GFP1-10::*unc-54* 3'UTR + *Cbr-unc-119*(+)] II; *unc-119(ed3)* III *cat-1::GFP11x3(syb7239)* X
DCR9414	*cat-1(ky1101 ky1118)* X 2X outcrossed
DCR9736	*Psrh-142::FLP (syb9159)* II 2X outcrossed
DCR9912	*Psrh-142::FLP (syb9159)* II; *cat-1(ky1101 ky1118)* X
DCR9342	*unc-17::GFP FLP-on(ola503)* IV*; unc-13::mScarlet FLP-on (wy1322)* I*; vlcSi1[unc-119(+); Psrh-142::Flippase]* III
DCR9574	*unc-17::mRuby FLP-on(syb7882)* IV*; cat-1::gfp11x3(syb7239)* X*; olaIs153 [Psrh-142::GFP1-10; Psrh-142::FLP; Punc-122::RFP]*
DCR9588	*unc-104(e1265)* II*; unc-17::mRuby FLP-on(syb7882)* IV*; cat-1::gfp11x3(syb7239)* X*; olaEx5548[Psrh-142::Flippase; Psrh142::GFP1-10; Punc-122::RFP]*
DCR9448	*cat-1::gfp11x3(syb7239)* X*; unc-13::mScarlet FLP-on (wy1322)* I*; olaEx5550[Psrh-142::Flippase; Psrh142::GFP1-10; Punc-122::RFP]*
DCR9583	*bas-1(syb5923[bas-1*::SL2::GFP::H2B]) III; *olaex5513* [P*srh-142*::BFP; P*unc-122*::RFP]
DCR9590	*mod-5(vlc47[mod-5*::T2A::mNeonGreen]); *olaex5513* [P*srh-142*::BFP; P*unc-122*::RFP]
DCR9577	*eat-4(kySi76 kySi77)* III [*eat-4* cell-specific KO]; *unc-17(syb8059)* IV [Flippase expression in *unc-17* locus]
DCR9584	*eat-4(kySi76 kySi77)* III [*eat-4* cell-specific KO]; *unc-47(syb8125)* III [Flippase expression in *unc-47* locus]

DCR9371	*eat-4(kySi76 kySi77)* III [*eat-4* cell-specific KO]; *bqSi614* IV [P*dat-1*::Flippase]
DCR9334	*eat-4(kySi76 kySi77)* III [*eat-4* cell-specific KO]; *bqSi488* IV [P*tph-1*::Flippase]
DCR9576	*unc-17::GFP FLP-on(ola503)* IV [*unc-17* cell-specific GFP knock-in]; *unc-47(syb8125)* III [Flippase expression in *unc-47* locus]
DCR9344	*unc-17::GFP FLP-on(ola503)* IV [*unc-17* cell-specific GFP knock-in]; *bqSi488* IV [P*tph-1*::Flippase] recombinant
DCR9897	*unc-17::GFP FLP-on(ola503)* IV [*unc-17* cell-specific GFP knock-in]; *bqSi614* IV [P*dat-1*::Flippase]

List of plasmids

Plasmid name	Genotype
DACR218	P*unc-122::RFP*
DACR704	P*myo-2*::mCherry
DACR4016	P*sto-3b::BFP*
DACR4027	P*unc-17*::UNC-17cDNA
DACR4033	P*unc-17*::UNC-17cDNA::GFP in TM6-7
DACR4038	P*unc-17*::GFP::UNC-17cDNA
DACR4064	P*sto-3b::GFP1-10*
DACR4078	P*srh-142::*BFP
DACR4083	P*srh-142::*Flippase
DCR4092	P*srh-142*::GFP1-10
pSH87	P*flp-13::*GFP1-10

ACKNOWLEDGEMENTS

We thank Nuria Flames Bonilla (Instituto de Biomedicina de Valencia, Spain), Rafa Alis (Instituto de Biomedicina de Valencia, Spain), Steven Flavell (MIT), and Yung-Chi Huang (MIT) for sharing reagents and constructs. We thank James Rand, Oliver Hobert (Columbia University) and Chen Wang (Columbia University) for sharing unpublished observations. We thank Ian Gonzalez for the creation of the name for this toolkit. We thank members of the Colón-Ramos Lab for feedback on figures and the manuscript. We also thank Stacy Wilson for technical support and training using

the AiryScan imaging set-up as part of the Yale Neuroscience Imaging Core Facility. We also thank Emerson Santiago, member of the Koelle Lab (Yale University), for expert advice on assays of serotonin function. Some strains were provided by the CGC, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440). Some figures were created in https://BioRender.com. This work was supported by National Institutes of Health grants to DC-R (R35NS132156 and R01NS076558) and to AW (K99AG083129), to EMJ (R01 NS034307), and to MLS (F32GM133139). The Pew Latin American Fellowship to ACC (AWD0006561), the Jane Coffin Childs Fellowship to ACC (AWD0006564), the HHMI Hanna Gray Fellowship to ACC (GT15993), and the Chan Zuckerberg Initiative to CIB.

FUNDING

ACC was supported by The Pew Foundation, the Jane Coffins Child, HHMI-HGF

AW was supported by K99AG083129

DC-R, PC-L and MT were supported by R35NS132156 and R01NS076558

MLS was supported by F32GM133139

EMJ was supported by R01 NS034307, R01 GM095817, and HHMI

MB, MSE, LF and CB were supported by a grant from the Chan Zuckerberg Initiative.

ORCID

Cuentas-Condori: 0000-0002-4847-0031

Feng: 0000-0003-4601-4224

Wolfe: 0000-0002-4926-5064

Schwartz: 0000-0002-5542-9109

Brown: 0000-0003-0547-6545

Ebert: 0000-0002-9122-7854

Jorgensen: 0000-0002-2978-8028

Bargmann: 0000-0002-8484-0618

Colón-Ramos: 0000-0003-0223-7717

REFERENCES

Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A.,...Jumper, J. M. (2024).
Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature*,
630(8016), 493-500. https://doi.org/10.1038/s41586-024-07487-w
Ahier, A., & Jarriault, S. (2014). Simultaneous expression of multiple proteins under a single
promoter in Caenorhabditis elegans via a versatile 2A-based toolkit. *Genetics*,
196(3), 605-613. https://doi.org/10.1534/genetics.113.160846
Albertson, D. G., & Thomson, J. N. (1976). The pharynx of Caenorhabditis elegans. *Philos*
Trans   R   Soc   Lond   B   Biol   Sci,   *275*(938),   299-325.
https://doi.org/10.1098/rstb.1976.0085
Alfonso, A., Grundahl, K., Duerr, J. S., Han, H.-P., & Rand, J. B. (1993). The Caenorhabditis
unc-17 gene: A putative vesicular acetylcholine transporter. *Science*, *261*.
Aubrey, K. R., Rossi, F. M., Ruivo, R., Alboni, S., Bellenchi, G. C., Le Goff, A.,...Supplisson, S.
(2007). The transporters GlyT2 and VIAAT cooperate to determine the vesicular
glycinergic   phenotype.   *J   Neurosci*,   *27*(23),   6273-6281.
https://doi.org/10.1523/JNEUROSCI.1024-07.2007
Bajar, B. T., Wang, E. S., Lam, A. J., Kim, B. B., Jacobs, C. L., Howe, E. S.,...Chu, J. (2016).
Improving brightness and photostability of green and red fluorescent proteins for live
cell   imaging   and   FRET   reporting.   *Sci   Rep*,   6,   20889.
https://doi.org/10.1038/srep20889
Bargmann, C. I., & Horvitz, R. H. (1991). Chemosensory neurons with overlapping functions
direc chemotaxis to multiple chemicals in C. elegans. *Neuron*, *7*.
Bellocchio, E. E., Hu, H., Pohorille, A., Chan, J., Pickel, V. M., & Edwards, R. H. (1998). The
localization of the brain-specific inorganic phosphate transporter suggests a specific
presynaptic role in glutamatergic transmission. *The journal of neuroscience*, *18*.
Bertuzzi, M., Chang, W., & Ampatzis, K. (2018). Adult spinal motoneurons change their
neurotransmitter phenotype to control locomotion. *Proc Natl Acad Sci U S A*,
115(42), E9926-E9933. https://doi.org/10.1073/pnas.1809050115
Brenner, S. (1973). The genetics of Caenorhabidits elegans. *Genetics*, *77*.
Chaudhry, F. A., Reimer, R. J., Bellocchio, E. E., Danbolt, N. C., Osen, K. K., Edwards, R. H.,
& Storm-Mathisen, J. (1998). The vesicular GABA transporter, VGAT, localizes to
synaptic vesicles in sets of glycinergic as well as GABAergic neurons. *The journal of*
neuroscience, *18*.
Chen, N., Zhang, Y., Rivera-Rodriguez, E. J., Yu, A. D., Hobin, M., Rosbash, M., & Griffith, L.
C. (2023). Widespread posttranscriptional regulation of cotransmission. *Sci Adv*,
9(22), eadg9836. https://doi.org/10.1126/sciadv.adg9836
Crank, J. (1956). *The Mathematics of Diffusion*.
Crawford, D. C., & Kavalali, E. T. (2015). Molecular underpinnings of synaptic vesicle pool
heterogeneity. *Traffic*, *16*(4), 338-364. https://doi.org/10.1111/tra.12262
Dag, U., Nwabudike, I., Kang, D., Gomes, M. A., Kim, J., Atanas, A. A.,...Flavell, S. W. (2023).
Dissecting the functional organization of the C. elegans serotonergic system at
whole-brain   scale.   *Cell*,   *186*(12),   2574-2592.e2520.
https://doi.org/10.1016/j.cell.2023.04.023
Duerr, J. S., Frisby, D. L., Gaskin, J., Duke, A., Asermely, K., Huddleston, D.,...Rand, J. B.
(1999). The cat-1 gene of Caenorhabditis elegans encodes a vesicular monoamine

transporter required for specific monoamine-dependent behaviors. *The journal of*
neuroscience, 19.

Duerr, J. S., Han, H. P., Fields, S. D., & Rand, J. B. (2008). Identification of major classes of
cholinergic neurons in the nematode Caenorhabditis elegans. *J Comp Neurol*,
506(3), 398-408. https://doi.org/10.1002/cne.21551

Edwards, R. H. (2007). The neurotransmitter cycle and quantal size. *Neuron, 55*(6), 835-858.
https://doi.org/10.1016/j.neuron.2007.09.001

Eiden, L. E. (1998). The cholinergic gene locus. *J Neurochem*, *70*(6), 2227-2240.
https://doi.org/10.1046/j.1471-4159.1998.70062227.x

Erickson, J., Eiden, L. E., & Hoffman, B. J. (1992). Expression cloning of a reserpine-sensitive
vesicular monoamine transporter. *Proceedings of the national academy of sciences*
of the United States of America, *89*.

Fei, H., Grygoruk, A., Brooks, E. S., Chen, A., & Krantz, D. E. (2008). Trafficking of vesicular
neurotransmitter transporters. *Traffic*, *9*.

Feng, L., Marquina-Solis, J., Yue, L., Harnagel, A., Greenfeld, Y., & Bargmann, C. I. (2025). *C.*
elegans interprets dietary quality through context-dependent serotonergic
modulation. *bioRxiv*. https://doi.org/https://doi.org/10.1101/2025.01.05.631367

Flavell, S. W., Pokala, N., Macosko, E. Z., Albrecht, D. R., Larsch, J., & Bargmann, C. I. (2013).
Serotonin and the neuropeptide PDF initiates and extend opposing behavioral states
in C. elegans. *Cell*, *154*.

Gendrel, M., Howell, E. G., & Hobert, O. (2016). A cellular and regulatory map of the
GABAergic nervous system of C. elegans. *eLIFE*, *5*.

Gouwens, N. W., Sorensen, S. A., Baftizadeh, F., Budzillo, A., Lee, B. R., Jarsky, T.,...Zeng, H.
(2020). Integrated Morphoelectric and Transcriptomic Classification of Cortical
GABAergic Cells. *Cell*, *183*(4), 935-953.e919.
https://doi.org/10.1016/j.cell.2020.09.057

Granger, A. J., Wallace, M. L., & Sabatini, B. L. (2017). Multi-transmitter neurons in the
mammalian central nervous system. *Curr Opin Neurobiol*, *45*, 85-91.
https://doi.org/10.1016/j.conb.2017.04.007

Hall, D. H., & Hedgecock, E. M. (1991). Kinesin-related gene unc-104 is required for axonal
transport of synaptic vesicles in C. elegans. *Cell*, *65*.

Hawk, J. D., Calvo, A. C., Liu, P., Almoril-Porras, A., Aljobeh, A., Torruella-Suárez, M.
L.,...Colón-Ramos, D. A. (2018). Integration of Plasticity Mechanisms within a Single
Sensory Neuron of C. elegans Actuates a Memory. *Neuron*, *97*(2), 356-367.e354.
https://doi.org/10.1016/j.neuron.2017.12.027

He, S., Cuentas-Condori, A., & Miller, D. M. (2019). NATF (Native and Tissue-specific
fluorescence: A strategy for bright, tissue-specific GFP labeling of native proteins in
Caenorhabditis elegans. *Genetics*, *212*.

Huang, Y.-C., Luo, J., Huang, W., Baker, C. M., Gomez, M. A., Meng, B.,...Flavell, S. W. (2023).
A single neuron in C. elegans orchestrates multiple motor outputs through parallel
modes of transmission. *Current Biology*, *33*.

Juge, N., Omote, H., & Moriyama, Y. (2013). Vesicular GABA transporter (VGAT) transports
β-alanine. *J Neurochem, 127*(4), 482-486. https://doi.org/10.1111/jnc.12393

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,...Hassabis, D.
(2021). Highly accurate protein structure prediction with AlphaFold. *Nature*,
596(7873), 583-589. https://doi.org/10.1038/s41586-021-03819-2

Kamalova, A., & Nakagawa, T. (2021). AMPA receptor structure and auxiliary subunits. *J*
Physiol, *599*(2), 453-469. https://doi.org/10.1113/jp278701

Koonce, N. L., Emerson, S. E., Bhaskar, D., Kuchroo, M., Moyle, M. W., Arroyo-Morales,
P.,...Colón-Ramos, D. (2025). NeuroSC: Exploring Neurodevelopment via
Spatiotemporal Collation of Anatomical Networks. *bioRxiv*.
https://doi.org/10.1101/2024.08.27.609993

Koonce, N. L., Emerson, S. E., Dhananjay, B., Kuchroo, M., Moyle, M. W., Arroyo-Morales,
P.,...Colon-Ramos, D. (2024). NeuroSCAN: Exploring neurodevelopment via
spatiotemporal collation of anatomical networks. *bioRxiv*.

Kumar, S., Sharma, A. K., & Leifer, A. M. (2024). An inhibitory acetylcholine receptor gates
context-dependent mechanosensory processing in. *iScience*, *27*(10), 110776.
https://doi.org/10.1016/j.isci.2024.110776

Lacin, H., Chen, H. M., Long, X., Singer, R. H., Lee, T., & Truman, J. W. (2019).
Neurotransmitter identity is acquired in a lineage-restricted manner in the. *eLIFE*, *8*.
https://doi.org/10.7554/eLife.43701

Lee, R. Y., Sawin, E. R., Chalfie, M., Horvitz, R. H., & Avery, L. (1999). EAT-4, homolog of a
mammalian sodium-dependent inorganic phosphate co-transporter, is necessary
for glutamatergic neurotransmission in Caenorhabditis elegans. *The journal of*
neuroscience, *19*.

Lee, S., Kim, K., & Zhou, Z. J. (2010). Role of ACh-GABA cotransmission in detecting image
motion and motion direction. *Neuron*, *68*(6), 1159-1172.
https://doi.org/10.1016/j.neuron.2010.11.031

Li, H. Q., Jiang, W., Ling, L., Pratelli, M., Chen, C., Gupta, V.,...Spitzer, N. C. (2024).
Generalized fear after acute stress is caused by change in neuronal cotransmitter
identity. *Science*, *383*(6688), 1252-1259. https://doi.org/10.1126/science.adj5996

Li, H. Q., Pratelli, M., Godavarthi, S., Zambetti, S., & Spitzer, N. C. (2020). Decoding
Neurotransmitter Switching: The Road Forward. *J Neurosci*, *40*(21), 4078-4089.
https://doi.org/10.1523/JNEUROSCI.0005-20.2020

Liu, C., Goel, P., & Kaeser, P. S. (2021). Spatial and temporal scales of dopamine
transmission. *Nat Rev Neurosci*, *22*(6), 345-358. https://doi.org/10.1038/s41583-
021-00455-7

Lopez-Cruz, A., Sordillo, A., Pokala, N., Liu, Q., McGrath, P. T., & Bargmann, C. I. (2019).
Parallel multimodal circuits control an innate foraging behavior. *Neuron*, *102*.

Maddaloni, G., Chang, Y. J., Senft, R. A., & Dymecki, S. M. (2024). Adaptation to photoperiod
via dynamic neurotransmitter segregation. *Nature*, *632*(8023), 147-156.
https://doi.org/10.1038/s41586-024-07692-7

Maicas, M., Jimeno-Martin, A., Millan-Trejo, A., Alkema, M. J., & Flames, N. (2021). The
transcription factor LAG-1/CSL plays a Notch-independent role in controlling
terminal differentiation, fate maintenance, and plasticity of serotonergic
chemosensory neurons. *PLoS Biology*, *19*.

Marquina-Solis, J., Feng, L., Vandewyer, E., Beets, I., Hawk, J., Colon-Ramos,
D.,...Bargmann, C. I. (2024). Antagosnism between neuropeptides and monoamines
in a distributed circuit for pathogen avoidance. *Cell Reports*, *43*.

Martens, H., Weston, M. C., Boulland, J.-L., Gronborg, M., Grosche, J., Kacza, J.,...Hartig, W.
(2008). Unique luminal localization of VGAT-C terminus allows for selective labeling
of active cortical GABAergic synapses. *The journal of neuroscience*, *28*.

Mathews, E. A., Mullen, G. P., Manjarrez, J. R., & Rand, J. B. (2015). Unusual regulation of
splicing of the cholinergic locus in Caenorhabditis elegans. *Genetics*, *199*.

McIntire, S. L., Jorgensen, E., & Horvitz, R. H. (1993). Genes required for GABA function in
Caenorhabditis elegans. *Nature*, *364*.

McIntire, S. L., Jorgensen, E., Kaplan, J., & Horvitz, R. H. (1993). The GABAergic nervous
system of Caenorhabditis elegans. *Nature*, *364*.

McIntire, S. L., Reimer, R. J., Schuske, K., Edwards, R. H., & Jorgensen, E. (1997).
Identification and characterization of the vesicular GABA transporter. *Nature*, *389*.

Morrie, R. D., & Feller, M. B. (2015). An Asymmetric Increase in Inhibitory Synapse Number
Underlies the Development of a Direction Selective Circuit in the Retina. *J Neurosci*,
35(25), 9281-9286. https://doi.org/10.1523/JNEUROSCI.0670-15.2015

Mullen, G. P., Mathews, E. A., Saxena, P., Fields, S. D., McManus, J. R., Moulder, G.,...Rand,
J. B. (2006). The Caenorhabditis elegans snf-11 gene encodes a sodium-dependent
GABA transporter required for clearance of synaptic GABA. *Mol Biol Cell*, *17*(7), 3021-
3030. https://doi.org/10.1091/mbc.e06-02-0155

Muñoz-Jiménez, C., Ayuso, C., Dobrzynska, A., Torres-Mendéz, A., Ruiz, P. C., & Askjaer, P.
(2017). An Efficient FLP-Based Toolkit for Spatiotemporal Control of Gene Expression
in. *Genetics*, *206*(4), 1763-1778. https://doi.org/10.1534/genetics.117.201012

Nguyen, J. P., Shipley, F. B., Linder, A. N., Plummer, G. S., Liu, M., Setru, S. U.,...Leifer, A. M.
(2016). Whole-brain calcium imaging with cellular resolution in freely behaving
Caenorhabditis elegans. *Proc Natl Acad Sci U S A*, *113*(8), E1074-1081.
https://doi.org/10.1073/pnas.1507110112

O'Malley, D. M., Sandell, J. H., & Masland, R. H. (1992). Co-release of acetylcholine and
GABA by the starburst amacrine cells. *J Neurosci*, *12*(4), 1394-1408.
https://doi.org/10.1523/JNEUROSCI.12-04-01394.1992

Pei, J., & Grishin, N. V. (2001). AL2CO: calculation of positional conservation in a protein
sequence alignment. *Bioinformatics*, *17*(8), 700-712.
https://doi.org/10.1093/bioinformatics/17.8.700

Pereira, L., Aeschimann, F., Wang, C., Lawson, H., Serrano-Saiz, E., Portman, D.
S.,...Hobert, O. (2019). Timing mechanism of sexually dimorphic nervous system
differentiation. *Elife*, *8*. https://doi.org/10.7554/eLife.42078

Pereira, L., Kratsios, P., Serrano-Saiz, E., Sheftel, H., Mayo, A. E., H., H. D.,...Hobert, O.
(2015). A cellular and regulatory map of the cholinergic nervous system of C. elegans.
eLIFE, *4*.

Pettersen, E. F., Goddard, T. D., Huang, C. C., Meng, E. C., Couch, G. S., Croll, T. I.,...Ferrin,
T. E. (2021). UCSF ChimeraX: Structure visualization for researchers, educators, and
developers. *Protein Sci*, *30*(1), 70-82. https://doi.org/10.1002/pro.3943

Pierce-Shimomura, J. T., Morse, T. M., & Lockery, S. R. (1999). The fundamental role of
pirouettes in Caenorhabditis elegans chemotaxis. *J Neurosci*, *19*(21), 9557-9569.
https://doi.org/10.1523/JNEUROSCI.19-21-09557.1999

Pocock, R., & Hobert, O. (2010). Hypoxia activates a latent circuit for processing gustatory
information in C. elegans. *Nat Neurosci*, *13*(5), 610-614.
https://doi.org/10.1038/nn.2537

Prevedel, R., Yoon, Y. G., Hoffmann, M., Pak, N., Wetzstein, G., Kato, S.,...Vaziri, A. (2014).
Simultaneous whole-animal 3D imaging of neuronal activity using light-field
microscopy. *Nat Methods*, *11*(7), 727-730. https://doi.org/10.1038/nmeth.2964

Robinson RA, S. R. (1959). *Electrolyte Solutions*.

Roghani, A., Feldman, J., Kohan, S. A., Shirzadi, A., Gundersen, C. B., Brecha, N., & Edwards,
R. H. (1994). Molecular cloning of a putative vesicular transporter for acetylcholine.
Proceedings of the national academy of sciences of the United States of America, *91*.

Santos, M., Park, K. C., Foss, S. M., & Voglmaier, S. M. (2013). Sorting of the vesicular GABA
transporter to functional vesicle pools by an atypical dileucine-like motif. *The journal*
of neuroscience, *33*.

Sato, H., Kunitomo, H., Fei, X., Hashimoto, K., & Lino, Y. (2021). Glutamate signaling from a
single sensory neuron mediates experience-dependent bidirectional behavior in
Caenorhabditis elegans. *Cell Reports*, *35*.

Scheffer, L. K., Xu, C. S., Januszewski, M., Lu, Z., Takemura, S. Y., Hayworth, K. J.,...Plaza, S.
M. (2020). A connectome and analysis of the adult Drosophila central brain. *Elife*, *9*.
https://doi.org/10.7554/eLife.57443

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,...Cardona,
A. (2012). Fiji: an open-source platform for biological-image analysis. *Nat Methods*,
9(7), 676-682. https://doi.org/10.1038/nmeth.2019

Schrödel, T., Prevedel, R., Aumayr, K., Zimmer, M., & Vaziri, A. (2013). Brain-wide 3D imaging
of neuronal activity in Caenorhabditis elegans with sculpted light. *Nat Methods*,
10(10), 1013-1020. https://doi.org/10.1038/nmeth.2637

Schwartz, M. L., & Jorgensen, E. (2016). SapTrap, a toolkit for high-throughput CRISPR/Cas9
gene modification in Caenorhabditis elegans. *Genetics*, *202*.

Seggewisse, A., & Winding, M. (2024). Mapping the fly nerve cord. *Elife*, *13*.
https://doi.org/10.7554/eLife.99804

Serrano-Saiz, E., Pereira, L., Gendrel, M., Aghayeva, U., Bhattacharya, A., Howell,
K.,...Hobert, O. (2017). A neurotransmitter atlas of the Caenorhabditis elegans male
nervous system reveals sexually dimorphic neurotransmitter usage. *Genetics*, *206*.

Serrano-Saiz, E., Poole, R. J., Felton, T., Zhang, F., De la Cruz, E. D., & Hobert, O. (2013).
Modula control of glutamateric neuronal identity in C. elegans by distinct
homeodomain proteins. *Cell*, *155*.

Seydoux, G., & Fire, A. (1994). Soma-germline asymmetry in the distributions of embryonic
RNAs in Caenorhabditis elegans. *Development*, *120*(10), 2823-2834.
https://doi.org/10.1242/dev.120.10.2823

Silm, K., Yang, J., Marcott, P. F., Asensio, C. S., Eriksen, J., Guthrie, D. A.,...Edwards, R. H.
(2019). Synaptic Vesicle Recycling Pathway Determines Neurotransmitter Content

and Release Properties. *Neuron*, *102*(4), 786-800.e785. https://doi.org/10.1016/j.neuron.2019.03.031

Sitko, A. A., Frank, M. M., Romero, G. E., Hunt, M., & Goodrich, L. V. (2025). Lateral olivocochlear neurons modulate cochlear responses to noise exposure. *Proc Natl Acad Sci U S A*, *122*(4), e2404558122. https://doi.org/10.1073/pnas.2404558122

Sulston, J. E., & Horvitz, H. R. (1977). Post-embryonic cell lineages of the nematode, Caenorhabditis elegans. *Dev Biol*, *56*(1), 110-156. https://doi.org/10.1016/0012-1606(77)90158-0

Sulston, J. E., Schierenberg, E., White, J. G., & Thomson, J. N. (1983). The embryonic cell lineage of the nematode Caenorhabditis elegans. *Dev Biol*, *100*(1), 64-119. https://doi.org/10.1016/0012-1606(83)90201-4

Südhof, T. C. (2021). The cell biology of synapse formation. *J Cell Biol*, *220*(7). https://doi.org/10.1083/jcb.202103052

Taylor, S. R., Santpere, G., Weinreb, A., Barrett, A., Reilly, M. B., Xu, C.,...Miller, D. M., 3rd. (2021). Molecular topography of an entire nervous system. *Cell*, *184*(16), 4329-4347.e4323. https://doi.org/10.1016/j.cell.2021.06.023

Trudeau, L.-E., & El Mestikawy., S. (2018). Glutamate cotransmission in cholinergic, GABAergic and monoamine systems: Contrasts and commonalities. *Frontiers in neural circuits*, *12*.

Uchida, O., Nakano, H., Koga, M., & Ohshima, Y. (2003). The C. elegans che-1 gene encondes a zinc finger trnascription factor required for specification of the ASE chemosensory neurons. *Development*, *130*.

Vaaga, C. E., Borisovska, M., & Westbrook, G. L. (2014). Dual-transmitter neurons: functional implications of co-release and co-transmission. *Curr Opin Neurobiol*, *29*, 25-32. https://doi.org/10.1016/j.conb.2014.04.010

Wang, C., Vidal, B., Sural, S., Loer, R. G., Merritt, D. M., Toker, I. A.,...Hobert, O. (2024). A neurotransmitter atlas of C. elegans males and hermaphrodites. *eLIFE*.

Ward, S. (1973). Chemotaxis by the nematode Caenorhabditis elegans: identification of attractants and analysis of the response by use of mutants. *Proc Natl Acad Sci U S A*, *70*(3), 817-821. https://doi.org/10.1073/pnas.70.3.817

White, J. G., Southgate, E., Thomson, J. N., & Brenner, S. (1986). The structure of the nervous system of the nematode Caenorhabditis elegans. *philosophical Transactions of the Royal Society of London*, *314*.

Wu, J. S., Yi, E., Manca, M., Javaid, H., Lauer, A. M., & Glowatzki, E. (2020). Sound exposure dynamically induces dopamine synthesis in cholinergic LOC efferents for feedback to auditory nerve fibers. *eLIFE*, *9*. https://doi.org/10.7554/eLife.52419

Wu, X., & Hammer, J. A. (2021). ZEISS Airyscan: Optimizing Usage for Fast, Gentle, Super-Resolution Imaging. *Methods Mol Biol*, *2304*, 111-130. https://doi.org/10.1007/978-1-0716-1402-0_5

Yi, Y., Li, Y., Zhang, S., Men, Y., Wang, Y., Jing, D.,...Zhao, H. (2024). Mapping of individual sensory nerve axons from digits to spinal cord with the transparent embedding solvent system. *Cell Res*, *34*(2), 124-139. https://doi.org/10.1038/s41422-023-00867-3

1003   Yu, C. Y., & Chang, H. C. (2022). Glutamate signaling mediates C. elegans behavioral
1004   plasticity to pathogens. *iScience*, *25*(3), 103919.
1005   https://doi.org/10.1016/j.isci.2022.103919
1006

1007   **Table 2 – List of potential co-transmitter neurons by neuronal type**

pharyngeal neurons	sensory neurons	interneurons	motor neurons
I2L	ADFL	AIML	HSNL
I5	ADFR	AIMR	HSNR
M4	AFDL	RIH	SMDDL
M5	AFDR	PVNL	SMDDR
MI	URXL	PVNR	SMDVL
	URXR	RIML	SMDVR
	SDQR	RIMR	VC04
	DVA	RICL	VC05
	PDEL	RICM	
	PDER		

1008

1009   **Table 3 – List of potential co-transmitter neurons reported by at least more than one study**

Co-transmitter neurons	Suggested by CenGEN expression data?*	Consistent with previous reports:	Combination suggested
ADF	Yes	Pereira et al. (2015), Wang et al. (2024) & this study	Serotonin + acetylcholine
AFD	Yes	Wang et al. (2024) & this study	Acetylcholine + glutamate
AIM	No	Jafari et al. (2011), Serrano-Saiz et al. (2013), Wang et al. (2024)	Serotonin + glutamate
DVA	Yes	Wang et al. (2024) & this study	Acetylcholine + glutamate
HSN	Yes	Duerr et al. (2001), Wang et al. (2024) & this study	Serotonin + acetylcholine
I2L	No	Wang et al. (2024) & this study	glutamate + UNC-47-dependent NT
I5	No	Jafari et al. (2011), Serrano-Saiz et al. (2013), Wang et al. (2024)	Serotonin + glutamate
M4	No	Wang et al. (2024) & this study	acetylcholine + UNC-47-dependent NT
M5	Yes	Wang et al. (2024) & this study	Acetylcholine + glutamate
MI	Yes	Serrano-Saiz et al. (2013), Wang et al. (2024)	Serotonin + glutamate
PDE	No	Wang et al. (2024) & this study	dopamine + glutamate
PVN	Yes	Wang et al. (2024) & this study	Acetylcholine + glutamate

RIC	Yes	Reilly et al. (2022), Wang et al. (2024)	glutamate + octopamine
RIH	Yes	Jafari et al. (2011), Wang et al. (2024)	Serotonin + acetylcholine
RIM	Yes	Alkema et al. (2005), Serrano-Saiz et al. (2013), Wang et al. (2024)	Glutamate + tyramine
SDQR	Yes	Wang et al. (2024) & this study	acetylcholine + UNC-47-dependent NT
SMD	Yes	Gendrel et al. (2016) & this study	GABA + acetylcholine
URX	No	Jafari et al. (2011), Wang et al. (2024)	Serotonin + acetylcholine
VC04	Yes	Duerr et al. (2001), Wang et al. (2024) & this study	Serotonin + acetylcholine
VC05	Yes	Duerr et al. (2001), Wang et al. (2024) & this study	Serotonin + acetylcholine

*Determined with threshold of 1

Table 4 – List of neurons inconsistent across studies to suggest co-transmission capacity

Previosly suggested:	Co-transmitter combination suggested	Reported by:	*cha-1*	*unc-17*	*unc-25*	*unc-47*	*unc-46*
RIB	GABA + acetylcholine	Gendrel et al. (2016)			X	X	X
AVB	GABA + acetylcholine	Gendrel et al. (2016)	X	X			X
AVA	GABA + acetylcholine	Gendrel et al. (2016)	X	X			

	Maximum CenGEN threshold that shows expression of each gene
4	X
3	X
2	X
1	X

1015 **FIGURE LEGENDS**

1016

Figure 1 - Endogenous fluorescence labeling of synaptic vesicle transporters

(Left) FLP-on and (Right) split GFP strategies to endogenously label synaptic vesicle transporters.

(A) Cartoon representation of the CRISPR-engineered worm in the endogenous locus of the

target synaptic vesicle gene. **(B)** Cell-specific driver that expresses (Left) Flippase or (Right)

GFP1-10. **(C)** Resulting tagged synaptic vesicle proteins with a (Left) full-length GFP or mRuby3

or (Right) by reconstituting GFP in the cell of interest. **(D)** Schematic of the resulting toolkit to label

and eliminate the endogenous machinery that packages or synthesizes glutamate, GABA,

acetylcholine and monoamines.

1025

Figure 2 – Probing glutamatergic transmission in *C. elegans*

(A) Predicted protein structure (using AlphaFold Protein Structure Database) of EAT-4 (cyan)

tagged with GFP (green) (Abramson et al., 2024). The last amino acid in the C-terminus end

(W576) corresponds to the tagged residue. **(B)** Schematic of the Vesicular Glutamate Transporter

(VGLUT/*eat-4*) gene structure, loss of function allele (*ky5*) and endogenously tagged versions

built for this study and by others. *eat-4* (*ky5*) mutants lack the first three exons. For cell-specific

knock-in (KI) tools, GFP (*syb8568*) and mRuby (*syb9193*) FLP-ON cassettes (Schwartz &

Jorgensen, 2016) were inserted before the STOP codon. To cell-specifically knock out (KO) *eat-*

4, two FRT sites flank the coding sequence of the *eat-4* gene (*kySi76 kySi77*) (Lopez-Cruz et al.,

2019). When recombination takes place, the *eat-4* coding sequence is removed and cytosolic

mCherry is inserted in-frame to be expressed as a proxy for *eat-4* sequence removal. **(C)**

Mutations in the *che-1* (prevents ASE development (Uchida et al., 2003)) (17.42 ± 7.7 mm) or *eat-*

4 gene (13.15 ± 4 mm) results in disrupted migration across the salt gradient. Wild-type animals

(7.76 ± 5.2 mm) migrate across the salt gradient similarly to EAT-4::GFP FLP-on (*syb8568*)

animals that express (7.6 ± 3.6 mm) or not (4.9 ± 4.3 mm) flippase pan-cellularly (*sybIs9614*).

Animals that express flippase in all cells (5.61 ± 3.9 mm) migrate across the salt gradient as a

wild-type animals. Results represent the mean distance of each worm from the salt peak,

averaged across the final minute of the assay, with each dot representing a single animal. Plots

are overlaid with Mean $\pm$ Standard Deviation. Kruskal-Wallis test with Dunn's multiple comparison

post hoc test. **** represents $p<0.0001$; *** represents $p<0.001$ and NS means "not significant".

(D) (Top) Fluorescent image of an adult worm expressing endogenously labeled EAT-4 with GFP

in all cells (P*eft-3*::Flippase). Scale Bar = 50μm. (Bottom) Zoom-in area of the head shows EAT-

4 expressed predominantly in the nerve ring. Scale Bar = 10μm. **(E)** (Left) Electron microscopy

rendering of ASE synapses in an L4 wild-type animal (White et al., 1986) (image generated with

NeuroSC (Koonce et al., 2025). (Right) Fluorescent image of endogenously tagged EAT-4 protein

cell-specifically in the ASE axons (P*flp-6*::Flippase). Scale Bar = 10μm.

1052

Figure S1

(A) Graphical representation of conservation of the amino acid predicted structure for the

Vesicular Glutamate Transporter across common model organisms (named in Figure S1B). Dark

colors denote the least conserved region, while clear colors denote the highest conservation. Note

the C-terminus end, where we introduced the GFP, is one of the darkest (least conserved) regions

in the structure. **(B)** Sequence alignment of the amino acid sequences for the Vesicular Glutamate

Transporter across common model organisms. Blue letters indicate conservation of the amino

acid properties (basic, acid, neutral) but not identities. Gray represents columns with gaps and no

conservation.

(C) Schematic of chemotaxis assay. (Top) A concentration gradient of NaCl (~60-85 mM) is

established on assay plates (blue rim) by adding high concentration NaCl drops to the outer point

of the assay plate at pre-determined times (See Methods). (Bottom) Larva stage 4 animals are

picked, and twenty-four hours later, the young adult animals are transferred to NaCl 100mM

training plates (red rim) for five hours. Animals are transferred to NGM buffer drops to wash off

bacteria, before being placed on the assay plates (blue rim, placement at blue dot in schematic),

where animal movement away from the highest [NaCl] is recorded (towards red dot in schematic).

(D) Displacement of trained wild type animals on a sham gradient (13.92 ± 5.3 mm) or untrained

wild-type animals on a gradient of NaCl (13.84 ± 5.4 mm). Wild-type animals (7.76 ± 5.2 mm)

migrate across the salt gradient like EAT-4::GFP FLP-on (*syb8568*) animals that express (8.36 ±

5.8 mm) flippase in ASE neurons (*sybIs9606*). Animals that only express flippase in ASE neurons

(7.61 ± 5.3 mm) migrate across the salt gradient similar to wild-type animals. Results represent

the mean distance of each worm from the salt peak, averaged across the final minute of the assay,

with each dot representing an individual animal. Plots are overlaid with Mean ± Standard

Deviation. Kruskal-Wallis test with Dunn's multiple comparison post hoc test. **** represents

$p < 0.0001$; *** represents $p < 0.001$; * represents $p < 0.05$; and NS means "not significant".

1078

1079   **Figure S2**

1080   Endogenous RAB-3::GFP synaptic vesicle clusters in the axonal process of ASER and ASEL.

1081

Figure 3 – Probing GABAergic transmission in *C. elegans*

(A) UNC-47 predicted protein structure from AlphaFold Protein Structure Database. GFP11 tag (red) was added between amino acids E145 and N146. Complementary GFP1-10 (green) was modeled as bound to GFP11. **(B)** Schematics of the Vesicular GABA Transporter (VGAT/*unc-47*) loss of function allele and endogenously tagged versions built for this study. *unc-47(e307)* mutant animals have a single base pair substitution (G to A in the first nucleotide of exon 6) that results in a splicing acceptor mutant. Fluorescent tags GFP (*syb6990*) and mKate2 (*syb7358*) were inserted between amino acids E145 and N146. For cell-specific endogenous labeling, UNC-47 was tagged with one (*syb7313*) or three copies (*syb7849*) of GFP11. To silence GABA transmission, we flanked the Glutamic Acid Decarboxylase/*unc-25* coding sequence with two FRT sites (*syb5949 syb6275*). Upon recombination, the *unc-25* coding sequence is removed and nuclear (black) mCherry (purple) is designed to be in-frame and expressed as a proxy for *unc-25* sequence removal. **(C)** (Top) Fluorescent image of an adult worm expressing endogenously labeled UNC-47 with reconstituted split-GFP in all cells (P*eft-3*::GFP1-10). Scale Bar = 50μm. (Bottom) Zoom-in area of the head shows UNC-47 expressed in the nerve ring and nerve cords. Scale Bar = 10μm. **(D)** (Left) Electron microscopy renderings of RIB synapses in an L4 wild-type animal (White et al., 1986) (image generated with NeuroSC (Koonce et al., 2025)). (Right) RIB-specific UNC-47 puncta (green) *in-vivo* using the UNC-47:GFP11 with reconstituted split-GFP in RIB (P*sto-3b*::GFP1-10) Scale Bar = 10μm. **(E)** *unc-47(e307)* mutant animals thrash significantly less per minute (30.8 ± 17) than wild-type animals (94.5 ± 10). Animals with non-reconstituted UNC-47::GFP11X3 (82.2 ± 9) (*syb7849*) thrash slightly less than wild-type animals (94.5 ± 10), while animals that express pan-cellular GFP1-10 (100.5 ± 9) (*muIs253*) are no different than wild-type animals. UNC-47::GFP11x3 animals that express Pan-cellular::GFP1-10 (with reconstituted GFP) (88.6 ± 7) thrash similarly to wild-type animals. FRT-flanked *unc-25* animals (*syb5949 syb6275*) that do not express Flippase (95.8 ± 12) thrash similarly to wild-type animals (91 ± 8). Cell-specific *unc-25* knockout animals (*syb5949 syb6275*) that express Flippase in every neuron (*bqSi506*) (49.36 ± 19) thrash to the same extent as *unc-25* (*e156*) mutant animals (40.2 ± 19). Plots are overlaid with Mean ± Standard Deviation. Kruskal-Wallis test with Dunn's multiple comparison post hoc test. **** represents $p < 0.0001$; *** represents $p < 0.001$; and NS means "not significant".

Figure S3

(A) Graphical representation of conservation of the Vesicular GABA Transporter amino acid predicted structure across common model organisms (named in Figure S2B). Dark colors denote the least conserved region, while clear colors denote the highest conservation. **(B)** Sequence alignment of the cytosolic loop between transmembrane domains 2 and 3 of the vesicular GABA transporter across common model organisms. Red letters indicate highly conserved columns (conservation of amino acid identity), and blue letters indicate conservation of the amino acid properties (basic, acid, neutral) but not identities. Gray represents columns with gaps. **(C)** Schematic of thrashing assay. "One thrash cycle" is scored when the animal bends as indicated in the schematic (from elongated-to-bent-to-elongated). We measured the number of thrashes per minute. **(D)** *unc-47(e307)* mutant animals thrash significantly less (41.5 ± 21.35) than wild-type animals (109.2 ± 23), while over-expression of UNC-47::GFP (100.2 ± 16), CRISPR knock-in UNC-47::GFP (107.9 ± 29) and UNC-47::mKate2 (103.1 ± 18) CRISPR-tagged animals thrash similarly to wild-type animals. Plots are overlaid with Mean $\pm$ Standard Deviation. Kruskal-Wallis test with Dunn's multiple comparison post hoc test. **** represents $p<0.0001$; *** represents $p<0.001$; ** represents $p<0.01$; * represents $p<0.05$; and NS means "not significant". **(E-F)** Fluorescent image of endogenously tagged **(E)** UNC-47::GFP (*syb6990*) and **(F)** UNC-47::mKate2 (*syb7358*). DNC = Dorsal Nerve Cord, VNC = Ventral Nerve Cord. Scale Bar = 10 μm.

Figure S4

(A) All presynaptic sites from known GABAergic neurons in the *C. elegans* nerve ring (RIS, RME, and RIB), according to 3D electron microscopy of a Larva stage 4 (L4) wild-type animal (White et al., 1986) (image generated with NeuroSC (Koonce et al., 2025)). **(B)** 3D-rendering of UNC-47::GFP (green) puncta in the *C. elegans* nerve ring. A-P denotes anterior-posterior axis. Scale bar = 10 μm. **(C)** Addition of RIB::BFP into Figure S3B. Green arrowheads point to UNC-47::GFP puncta that overlap with RIB::BFP. Yellow arrowheads point to the synapses we interpret, based on our labeling and distribution, to belong to the RME neurons. **(D)** Endogenous mScarlet::RAB-3 in RIB highlight the characteristic shape of the RIB axon. **(E)** *In-vivo* reconstitution of UNC-47::GFP11x3 in DD neurons when tagged with (Top) one or (Bottom) three copies of GFP11 (P*flp-13*::GFP1-10) (He et al., 2019). **(F-G) (F)** Line scans and **(G)** quantification of reconstituted GFP fluorescence intensity with one (11.3 ± 6) or three copies (38.3 ± 28) of GFP11 in DD neurons. Scale bar = 10 μm. Plots are overlaid with Mean $\pm$ Standard Deviation. Mann-Whitney test. **** represents $p<0.0001$.

Figure 4 – Probing cholinergic transmission in *C. elegans*

(A) Predicted UNC-17 protein structure (magenta) tagged with GFP (green) on the last amino

acid in the C-terminus end (W532). **(B)** Schematics of the Vesicular Acetylcholine Transporter

(VACHT/*unc-17*) loss-of-function allele and endogenously tagged versions tested in this study

(and built by others). *unc-17(e245)* mutant animals have a single base pair substitution in the third

exon which leads to an amino acid change (arrowhead). Full-body knock-in animal labels UNC-

17 with mKate (*ot907*). For cell-specific labeling, the N-terminus GFP FLP-ON cassette (*syb7251*)

was inserted between amino acids P6 and V7 (See Methods). Similarly, the C-terminus GFP FLP-

ON cassette (*ola503*) and mRuby FLP-on cassette (*syb7882*) were inserted before the STOP

codon. **(C)** *unc-17(e245)* mutant animals (1.9 ± 2) thrash significantly less than wild-type animals

(93 ± 10). UNC-17::GFP FLP-on (*ola503*) animals that express pan-cellular::Flippase (88 ± 1)

(*sybIs9614*) or animals that only express pan-cellular::Flippase (94.8 ± 12) are indistinguishable

from wild-type animals in their thrashing behavior. UNC-17::GFP FLP-on animals (98.7 ± 8) thrash

significantly more than wild-type animals. Mean $\pm$ Standard Deviation. Brown-Forsythe ANOVA

test with Dunnett's T3 multiple comparisons post hoc test. **** represents $p < 0.0001$; * represents

$p < 0.05$; and NS means "not significant". **(D)** (Top) Fluorescent image of an adult worm expressing

endogenously labeled UNC-17::GFP in all cells (P*eft-3*::Flippase). Scale Bar = 50μm. (Bottom)

Zoom-in area of the head shows UNC-17 expressed in the nerve ring, nerve cords and sub-lateral

cords. DNC = Dorsal Nerve Cord, VNC = Ventral Nerve Cord. Scale Bar = 10μm. **(E)** (Left)

Electron microscopy rendering of ADF synapses in an L4 wild-type animal (White et al., 1986)

(image generated with NeuroSC (Koonce et al., 2025). (Right) Fluorescence image of

endogenously tagged UNC-17::GFP protein specifically in the ADF neuron. . Scale Bar = 10μm.

1169

Figure S5

(A) Graphical representation of the amino acid conservation of the Vesicular Acetylcholine

Transporter (VAChT) across common model organisms. Dark colors denote the least conserved

region, while clear colors denote the highest conservation. Note the C-terminus end is one of the

darkest regions in the structure, where the fluorophore was added. **(B-C)** Amino acid sequence

alignment of VAChT across model organisms. Black arrowhead points to insertion site of

fluorescent tag. **(B)** Sequences between transmembrane domains 6 and 7, which was tagged in

olaEx5704 array (See panel E). **(C)** Sequence at the C-terminus end. Red letters indicate highly

conserved columns (exact conservation of the amino acid identity), and blue indicates

conservation of the amino acid properties (basic, acid, neutral) but not identities. Gray represents

areas of no conservation. **(D)** Fluorescence image of CRISPR-tagged UNC-17::mKate2 (*ot907*)

animal. Scale Bar = 10μm. **(E)** Schematic of UNC-17 predicted topology (magenta) along the

membrane (gray). Green represents the locations of fluorescent tags tested for thrashing assays.

Alleles, either extrachromosomal arrays expressed in *unc-17* (*e245*) mutants or CRISPR knock-

in strains in the *unc-17* locus, are listed based on where the fluorescent tag was inserted. **(F)** *unc-*

17(e245) mutant animals barely swim (1.7 $\pm$ 2). Over-expression of untagged P*unc-17*::UNC-17

(*olaEx5703*) (114.8 $\pm$ 18) and N-terminus-tagged UNC-17::GFP (*olaEx5400*) (91.1 $\pm$ 22) swim like

wild-type animals (99.4 $\pm$ 20). GFP-tag between TM6-7 (*olaEx5704*) (34.3 $\pm$ 19) or CRISPR-

insertion of C-terminus mKate2 tag (*ot907*) (77.7 $\pm$ 17) leads to reduced swimming behavior when

compared to wild-type animals. Insertion of GFP-FLP-on cassette at the N-terminus end

(s*yb7251*) (82.9 $\pm$ 16) or insertion of the GFP-FLP-on cassette at the C-terminus (*ola503*) (101.9

$\pm$ 16) thrash as well as wild-type animals (109 $\pm$ 25). We are unsure as to the phenotypic

differences of *ola503* allele and *ot907* in our thrashing assays, but note that in addition to

the use of different fluorescent proteins, each allele also employs distinct linker sequences

between UNC-17 and the fluorescent protein (new Figure S6). *syb7882* allele, which

employes mRuby3 with a similar linker to *ola503*, also does not display defects in the thrashing

assay (Figure S6). Mean $\pm$ Standard Deviation. Kruskal-Wallis test with Dunn's multiple

comparison post hoc test. **** represents $p<0.0001$; ** represents $p<0.01$; * represents $p<0.05$;

and NS means "not significant".

1199

Figure S6

(A) (Top) Fluorescent image of an adult worm expressing endogenously labeled UNC-

17::mRuby3 in all cells (P*eft-3*::Flippase). Scale Bar = 50μm. (Bottom) Zoom-in area of the head

shows UNC-17 expressed in the nerve ring. DNC = Dorsal Nerve Cord, VNC = Ventral Nerve

Cord. Scale Bar = 25μm. **(B)** *unc-17(e245)* mutant animals (3 $\pm$ 4.1) thrash significantly less than

wild-type animals (94 $\pm$ 7.8). UNC-17::mRuby3 FLP-on (*syb7882*) animals that express pan-

cellular::Flippase (99 $\pm$ 6.6) (*sybIs9614*) or animals that only express pan-cellular::Flippase (95 $\pm$

6.2) are indistinguishable from wild-type animals in their thrashing behavior. UNC-17::mRuby

FLP-on animals (100 $\pm$ 6.1) thrash significantly more than wild-type animals. Mean $\pm$ Standard

Deviation. Kruskal-Wallis ANOVA test with Dunn's multiple comparisons post hoc test. ****

represents $p<0.0001$; * represents $p<0.05$; and NS means "not significant".

1211

Figure 5 – Probing Monoaminergic transmission in *C. elegans*

(A) Predicted CAT-1 protein structure (blue) labeled with GFP11 (red) at C-terminus end (E145). Complementary GFP1-10 (green) was modeled with bound GFP11. **(B)** Schematic of the Vesicular Monoamine Transporter (VMAT/*cat-1*) loss-of-function allele and endogenously tagged versions used to monitor monoaminergic transmission. *cat-1(ok411)* mutant animals have a deletion that spans exons 7 and 8 (black line) of the *cat-1* gene. For cell-specific labeling, three copies of GFP11 (*syb7239*) were inserted before the STOP codon. For cell-specific silencing of *cat-1* activity, two FRT sites (*ky1101 ky1118*) flank the coding sequence of the *cat-1* gene. Upon recombination, expression of cytosolic mCherry (magenta) is used as a proxy for deletion of the *cat-1* coding sequence. **(C)** (Top) Fluorescence image of reconstituted CAT-1::GFP11x3 with expression of complementary GFP1-10 in the whole animal (P*eft-3*::GFP1-10). Scale Bar = 50μm. (Bottom) Zoom-in of the (Left) head and (Right) vulva area of an adult worm shows CAT-1 expressed in the nerve ring and pharyngeal neurons as well as in the vulva. Scale Bar = 10μm. **(D)** (Top) A well-fed day-1 adult animal is placed on NGM plates covered with a thin layer of bacteria, allowed to roam for 16 hours and the number of squares traveled was recorded. (Bottom) Wild-type animals (43.6 $\pm$ 10) roam less than *cat-1(ok411)* mutants (87.5 $\pm$ 12). CAT-1::GFP11x3 (*syb7239*) animals with GFP reconstituted (43.3 $\pm$ 20) (or not (56.8 $\pm$ 14)) roam similar to wild-type animals. Animals with pan-cellular expression of GFP1-10 (*muIs253*) (53.3 $\pm$ 20) also roam like wild-type animals. **(E)** (Top) A well-fed day-1 adult animal is placed on NGM plates covered with a thin layer of *E. coli*, allowed to roam for 20 hours and the aversion ratio was calculated. (Bottom) Wild-type animals (0.12 $\pm$ 0.07) display less aversion to *E. coli* lawns than ADF-specific *cat-1* conditional KO animals (*ky1101 ky1118; syb9159*) (0.4 $\pm$ 0.1), consistent with previous reports using serotonin-depletion mutants (Feng et al., 2025). *cat-1* conditional KO animals that do not express Flippase in ADF neurons (0.07 $\pm$ 0.03) are indistinguishable from wild-type. Mean $\pm$ Standard Deviation. Kruskal-Wallis test with Dunn's multiple comparison post hoc test. **** represents $p<0.0001$; ** represents $p<0.01$; * represents $p<0.05$; and NS means "not significant".

Figure 6 – Mapping of co-transmitter neurons in the *C. elegans* nervous system

(A) Schematic of the approach used to find co-expression of two vesicular transporters in the same cells. Flippase drivers (Element #1, blue) and Flippase-dependent cassettes that result in fluorescence (Element #2, orange) are used. Magenta cells are seen when both elements are co-expressed (in green box). **(B)** Strategy to track the co-expression of the vesicular glutamate and acetylcholine transporter in combination with the 4 most common neurotransmitters in *C. elegans*:

Acetylcholine, GABA, Serotonin and Dopamine. "Elements #1" or "Element #2" refers to the
genetic elements in the schematic in Fig 6A. **(C-D)** Specific example of the genetic strategy
outlined in Fig 6A, for the vesicular transporters of glutamate (EAT-4) and acetylcholine (UNC-
17). **(C)** We repurposed the *eat-4* conditional KO strain (*kySi76 kySi77*) (Figure 1B) (Lopez-Cruz
et al., 2019) in which the *eat-4* gene coding sequence is flanked by two FRT sites and followed
by cytosolic mCherry (Element #1). We crossed this line with a strain that has an endogenously
inserted self-cleaving peptide sequence (T2A) followed by Flippase before the STOP codon in
the *unc-17* gene locus (Table 1). **(D)** (Top) Co-expression of EAT-4 and UNC-17 results in
cytosolic mCherry. (Bottom) Cytosolic mCherry was detected in the head of three neurons: (i)
AFDR, AFDL and M5; and in two neurons in the tail region: (ii) PVN, and DVA. **(E)** Fluorescence
microscopy shows neurons with co-expression of **(E)** VGLUT/EAT-4 and VGAT/UNC-47; **(F)**
VGLUT/EAT-4 and dopamine synthesis gene DAT-1; **(G)** VAChT/UNC-17 and VGAT/UNC-47;
and **(H)** VAChT/UNC-17 and the serotonin synthesis gene TPH-1. All scale bars = 10μm.

1258

Figure S7
(A) To track glutamate-involved co-transmission, we repurposed the *eat-4* conditional KO strain
(*kySi76 kySi77*) (Lopez-Cruz et al., 2019) where the *eat-4* gene coding sequence is flanked by
two FRT sites and followed by cytosolic mCherry (Element #2, orange box). Crossing this line
with a panel of flippase drivers (Element #1, blue box) results in activation of cytosolic mCherry
only in cells were both elements were co-expressed (Readout, green box). P*dat-1*::Flippase and
P*tph-1*::Flippase driver strains were made available from (Muñoz-Jiménez et al., 2017). **(B)** To
track acetylcholine-involved co-transmission, we repurposed the unc-17::GFP conditional KI
strain (*ola503*) (Figure 3B) for which, after the *unc-17* gene coding sequence, there are FRT sites
flanking the STOP codon and 3' UT, followed by GFP (Element #2, orange box). Crossing this
line with a panel of flippase drivers (Element #1, blue box) results in the protein fusion of UNC-17
with GFP in cells where both elements were co-expressed (Readout, green box). **(C-D)**
Systematic mapping shows neurons with co-transmission of **(C)** glutamate in combination with
acetylcholine, GABA, and dopamine. **(D)** Co-transmission of acetylcholine is found in combination
with GABA and serotonin.

1274

Figure S8
Live-imaging of co-expression strategy of the *eat-4* conditional knockout strain (*kySi76 kySi77*)
with **(A)** endogenous P*unc-47*::UNC-47::T2A::Flippase and **(B)** P*dat-1*:: Flippase driver. Live-
imaging of co-expression strategy of the *unc-17* conditional GFP knock-in strain (*kySi76 kySi77*)

with **(C)** P*tph-1*:: Flippase driver, and with **(D)** endogenous P*unc-47*::UNC-47::T2A::Flippase.

(Top) DIC images. (Bottom) Fluorescence imaging. All scale bars = 10μm.

1281

Figure S9

Our results (bold) are consistent and extend findings from previous studies in the field (See Table

3 and numbered references below). Here, we present an atlas of co-transmission in the *C.*

elegans nervous system. Co-transmitter neurons are present in the (i) head, mid-body region and

(ii) tail of the animal.

*Note that I2L, M4 and SDQR express UNC-47/VGAT but do not stain positive for GABA staining

(Gendrel et al., 2016) nor express genes required for GABA synthesis or uptake.

[1] Jafari et al. (2011), Serrano-Saiz et al. (2013), and Wang et al. (2024)

[2] Serrano-Saiz et al. (2013), and Wang et al. (2024)

[3] Reilly et al. (2022), and Wang et al. (2024)

[4] Jafari et al. (2011), and Wang et al. (2024)

[5] Alkema et al. (2005), Serrano-Saiz et al. (2013), and Wang et al. (2024)

1294

Figure 7 – Visualizing the distribution of cholinergic and serotonergic vesicles in ADF

neurons.

(A) Dual-labeling of the endogenous acetylcholine (UNC-17, magenta) and serotonin (CAT-1,

green) vesicular transporters in ADF neurons. **(B)** UNC-17::mRuby and CAT-1::GFP overlap

along the ADF axon. White arrowheads denote overlap of both signals. **(B')** UNC-17::mRuby and

CAT-1::GFP intensity profile. **(C)** Example of UNC-17::mRuby and CAT-1::GFP when they

partially do not overlap along the ADF axon, green arrowheads point to CAT-1-only puncta. **(C')**

UNC-17::mRuby and CAT-1::GFP intensity profile. Scale bar = 10μm. **(D-D')** Endogenous UNC-

17::mRuby and CAT-1:::GFP are absent from the axon of *unc-104* mutant animals. Line scan of

axon **(D)** and fluorescent images **(D')**. **(E)** Schematic of the dual-labeling of endogenous CAT-

1::GFP or UNC-17::GFP with endogenous active zone protein UNC-13::mScarlet along the ADF

axon. **(F)** Live imaging of the ADF axon reveals UNC-13::mScarlet puncta that lack CAT-1::GFP.

Scale bar = 10 μm. **(F'- F'')** Zoom-in region of E. White arrowheads show overlapping vesicular

transporter tagged (green) with the active zone protein UNC-13 (magenta) in individual puncta.

Magenta arrowhead points to UNC-13::mScarlet puncta that do not overlap with CAT-1::GFP. **(G)**

Live imaging of the ADF axon reveals UNC-13::mScarlet puncta that lack UNC-17::GFP. Scale

bar = 10 μm. **(G'- G'')** Zoom-in region of F. White arrowheads show overlapping vesicular

transporter tagged (green) with the active zone protein UNC-13 (magenta) in individual puncta.

Magenta arrowhead points to UNC-13::mScarlet puncta that do not overlap with CAT-1::GFP.

Scale Bar = 2 μm. **(H)** AiryScan imaging of dual-labeled UNC-17::mRuby and CAT-1::GFP along

the ADF axon show localization in the same synaptic bouton but with distinct enrichment areas.

Green arrowhead head points to CAT-1 enrichment and magenta arrowhead points to UNC-17

enrichment. Scale Bar = 2 μm**.**

1318

Figure S10

(A) Schematic of genes required for serotonergic identity. Mammalian (bold) and *C. elegans*

(italics) homologue genes are listed. Detectable expression of **(A')** *tph-1,* **(A'')** *bas-1* and **(A''')**

mod-5 in ADF neuron (blue). Scale Bar = 5 μm**.**

(B) Schematic of genes required for cholinergic identity. Mammalian (bold) and *C. elegans* (italics)

homologue genes are listed. Detectable expression of **(B')** *cho-1* in ADF neuron (blue). Scale Bar

= 10 μm**.**

(C) Presynaptic densities (green) on ADF axons as determined by serial electron microscopy

reconstructions (White et al., 1986) (image generated with NeuroSC (Koonce et al., 2025). **(C')**

3D rendering of live fluorescence imaging of endogenous UNC-17::GFP and **(C'')** CAT-1::GFP

specifically in ADF neurons. Scale bar = 10 μm**.**

1330

Table 1 – List of alleles for neurotransmitter examination in vivo

Table 2 – List of potential co-transmitter neurons by neuronal type

Table 3 – List of potential co-transmitter neurons reported by at least more than one study

Table 4 – List of neurons inconsistent across studies to suggest co-transmission capacity

Table 5 – List of all molecular locations tested for labeling each synaptic vesicle

transporter

A FLP-on cassettes | Split GFP

CRISPR-engineered worm

FRT cassette

synaptic vesicle transporter gene
FRT STOP FRT fluorophore

GFP11

synaptic vesicle transporter gene
GFP11

B Cell-specific

Flippase

Cell-specific promoter → Flippase

GFP1-10

Cell-specific promoter → GFP1-10

C Resulting tag

Tagged protein

fluorophore
FRT

GFP tagged OR mRuby tagged

Reconstituted GFP tagged protein

+

D Toolkit for *in* vivo mapping and modulating of neurotransmission at single-cell resolution

knock-in strains

Glutamate | GABA | Acetylcholine | Monoamines

knock-out strains

Figure 2

A Tagged Vesicular Glutamate Transporter/EAT-4

N-terminus

C-terminus
GFP tag

B Resources for the study of glutamatergic transmission

Tool	Allele	Sequence representation	Reference
eat-4 loss of function	*ky5*		Mori et al. (1993)
eat-4::GFP FLP-on knock in	*syb8568*	FRT STOP FRT	This study
eat-4::mRuby FLP-on knock in	*syb9193*	FRT STOP FRT	This study
FRT-flanked *eat-4*	*kySi76* *kySi77*	FRT STOP FRT	Lopez-Cruz (2019)

C Chemotaxis Assay

Mean distance from [NaCl] peak (mm)

**** NS NS NS
**** **** ***

Wild type
che-1 (*luc174*)
eat-4 (*ky5*)
eat-4::GFP FLP-on
Pan-cellular Flippase driver
Pan-cellular EAT-4::GFP

D EAT-4::GFP FLP-on (*syb8568*) + Pan-cellular::Flippase (*sybIs9614*)

50μm

50μm

Nerve ring

E ASE-specific Flippase

ASE synapses

Electron microscopy rendering

10μm

EAT-4::GFP (*syb8568*)

Figure S1

A Vesicular Glutamate Transporter conservation

Z-score

1.50
0.88
0.25
-0.38
-1.00

C-terminal

B Vesicular Glutamate Transporter C-terminus sequence

Tag insertion site ▼

C. elegans	549	------------APGTNP[5]----------	DEHGSSGVVENPHYQQW--- 576
M. musculus	534	PPAPPPSYGATHSTVQPP RPPPPVRDY	-------------------- 560
H. sapiens	534	PPAPPPSYGATHSTFQPP RPPPPVRDY	-------------------- 560
D. rerio	541	LATRQKTYGTTDNSSGRK QGWKKKRGV[4]EDDHESNHYENGEYQTQyq-	590
D. melanogaster	571	--STAISYGATGHVANNP[13]DAPPTYGDV[4]QYGYTQGQMPSYDPQGYqqq	632

C Chemotaxis Assay set up

1. Preparation of NaCl gradient

12.5 mm

Placement of high concentration NaCl drops

~50 ~60 ~85
[NaCl] (mM)

2. Training of animals and set-up of chemotaxis assay

+ 24 hrs.

5 hrs.

Pick L4 animals

Transfer animals to 100mM plate

Wash food off animals

Wash 1 Wash 2

Record movement

D Chemotaxis Assay Controls

**** **** NS NS NS

NS **** **** ***

Mean distance from [NaCl] peak (mm)

30
20
10
0

N2 Sham gradient No training eat-4::GFP FLP-on ASE-specific Flippase ASE-specific EAT-4::GFP

Figure S2

ASEL

ASER

10μm

endogenous
RAB-3

endogenous
RAB-3

Figure 3

A

Tagged Vesicular GABA Transporter/UNC-47

C-terminal

GFP11 tag

Reconstituted GFP

B Resources for the study of GABAergic transmission

Tool	Allele	Sequence representation	Reference
unc-47 loss of function	*e307*	Splicing acceptor mutant	Brenner (1974)
unc-47::GFP full-body knock in	*syb6990*		This study
unc-47::mKate full-body knock in	*syb7358*		This study
unc-47::GFP11 cell-specific labeling	*syb7313*		This study
unc-47::GFP11x3 cell-specific labeling	*syb7849*		This study
FRT-flanked *unc-25*	*syb5949* *syb6275*	FRT · STOP · FRT	This study

C

UNC-47::GFP11x3 (*syb7849*) + Pan-cellular::GFP1-10 (*muIs253*)

50 µm

DNC

VNC

10 µm

DNC

Nerve ring

VNC

Head

D

RIB synapses

RIB-specific GFP1-10

Electron microscopy rendering

UNC-47:: GFP11x3 (*syb7849*)

E Thrashing Assay

Thrashes per minute

**** *** NS NS *** NS ***

**** **** **** **** NS

WT | *unc-47* (*e307*) | UNC-47::GFP11x3 | Pan-cellular GFP1-10 | UNC-47::GFP11x3 +Pan-cellular GFP1-10 | *unc-25* (*e156*) | FRT-flanked *unc-25* | FRT-flanked *unc-25* + Pan-cellular FLP

Figure S3

A Vesicular GABA Transporter conservation

Z-score
1.50
0.88
0.25
-0.38
-1.00

first loop   N-term

B Vesicular GABA Transporter sequence

Tag insertion site

	TM2		TM3	
C. elegans	120 WSIGAMVGVAYVCYWTGVLLIECLYENG------VKKRKTYREIADFY-KPGF	165
D. melanogaster	172 WAIVAMVGIAHICCYTGKVLVQCLYEPDPaTGQMVRVRDSYVAIAKVCFGPKL	224
M. musculus	147 LGLFLIIFAAVVCCYTGKILIACLYEENE-DGEVVRVRDSYVAIANACCAPRF	198
D. rerio	152 LGLFLIIFAAVVCCYTGKILIACLYEENE-DGQLVRVRDSYVDIANACCAPRF	203
H. sapiens	147 LGLFLIIFAAVVCCYTGKILIACLYEENE-DGEVVRVRDSYVAIANACCAPRF	198

C Thrashing assay

One thrash cycle

D

Thrashes per minute

****   NS   NS   NS
*   ***   **

WT   *unc-47(e307)*   *unc-47(e307)* + Ex[UNC-47::GFP]   UNC-47::GFP (*syb6990*)   UNC-47::mKate2 (*syb7358*)

E UNC-47::GFP

DNC   Nerve ring   VNC

F UNC-47::mKate2

DNC   Nerve ring   VNC

Figure S4

Figure 4

A
Tagged Vesicular Acetylcholine Transporter/UNC-47

C-terminus
GFP tag

B Resources for the study of cholinergic transmission

Tool	Allele	Sequence representation	Reference
unc-17 loss of function	*e245*	G347R	Brenner (1974)
unc-17::mKate full-body knock in	*ot907*		Pereira et al. (2019)
GFP::*unc-17* FLP-on knock in	*syb7251*	SL2 — FRT FRT	This study
unc-17::GFP FLP-on knock in	*ola503*	FRT STOP FRT	This study
unc-17::mRuby FLP-on knock in	*syb7882*	FRT STOP FRT	This study
LoxP-flanked *unc-17*	*syb5779* *syb5987*	LoxP LoxP	Huang et al. (2023)

C Thrashing Assay

**** NS * NS
**** **** ****

Thrashes per minute

WT · unc-17 (e245) · Pan-cellular Flippase · UNC-17::GFP FLP-on · UNC-17::GFP FLP-on + Flippase

D UNC-17::GFP FLP-on (*ola503*) + Pan-cellular::Flippase (*sybIs9614*)

50μm

DNC

VNC

Sub-lateral cords

10μm

Pharynx

Nerve ring

E ADF-specific Flippase

ADF synapses

Electron microscopy rendering

UNC-17::GF (*ola503*)

Figure S5

A Vesicular Acetylcholine Transporter conservation

B Vesicular Acetylcholine Transporter sequence between TM6-TM7

Tag insertion site

TM6 ▼ TM7

C. elegans	227	SFVCLADAIAVFMVINPHRRGTDSHGEKVQGT-PMWRLFMDPFIACCS	273
D. melanogaster	222	ALVCLLDGLMLLLVMKPVKEAMKQSKDVQDQViPIWRLLMDPYIAVCA	269
M. musculus	222	ACVCLADGILCLTVLKPFSS--RTRENMPVGT-PIYKLMIDPYIAVVA	266
D. rerio	249	AAVSLFDALLLLAVAKPFSAAARARANLPVGT-PIHRLMDPYIAVVA	295
H. sapiens	249	AAVSLFDALLLLAVAKPFSAAARARANLPVGT-PIHRLMDPYIAVVA	295

C Vesicular Acetylcholine Transporter sequence on C-termini end

Tag insertion site ▼

C. elegans	501	NQAQIPNHAVSFQDSRPQAEFPAG-YDPLNPQW	532
D. melanogaster	546	NPFQQQQQQQQQQQQQVQSRGPAApANPFRQGF	578
M. musculus	493	GTFAGQSKSFSEEETSEPEYI-------------	513
D. rerio	515	GPFDGCEDDYNYY-SRS----------------	530
H. sapiens	516	GPFDACEDDYNYYTRS----------------	532

D *unc*-17::mKate2 *(ot907)*

DNC

Nerve ring

VNC

E UNC-17 predicted topology

untagged = *olaEx5703*

1 2 3 4 5 6 7 8 9 10 11 12

olaEx5704

olaEx5400
syb7251

ot907
ola503

F Thrashing Assay

Figure S6

A. UNC-17::mRuby3 (*syb7882*) +
Pan-cellular::Flippase (*sybIs9614*)

50μm

Sub-lateral cords

DNC

VNC

25μm

Pharynx   Nerve ring

B. Thrashing Assay

Figure 5

A
Tagged Vesicular Monoamine Transporter/CAT-1

GFP11 tag

Reconstituted GFP

C
CAT-1::GFP11x3 (*syb7239*) + Pan-cellular::GFP1-10 (*muIs253*)

50 µm

NSM

10 µm

B Resources for the study of monoaminergic transmission

Tool	Allele	Sequence representation	Reference
cat-1 gene	*ok411*		Hukema et al. 2008
cat-1::GFP11x3 knock in	*syb7239*		Huang et al. 2023
FRT-flanked *cat-1*	*ky1101* *ky1118*	FRT STOP FRT	This study

D On-food exploration assay

E. coli OP50 → +16 hrs. → quantify

E Aversion assay

Camera

E. coli BW25113

Calculate aversion ratio

$$\frac{N_{out}}{N_{total}}$$

Figure 6

A

B

		Element #1			
		Acetylcholine *unc-17*	GABA *unc-47*	Serotonin *tph-1*	Dopamine *dat-1*
Element #2	Glutamate *eat-4*	X	X	X	X
	Acetylcholine *unc-17*		X	X	X

C *eat-4*-based co-transmission

Element #1

Element #2

D Readout: Endogenous *eat-4* promoter

E *eat-4* FLP-on + P*unc-47*::Flippase

G *unc-17* FLP-on + P*unc-47*::Flippase

F *eat-4* FLP-on + P*dat-1*::Flippase

H *unc-17* FLP-on + P*tph-1*::Flippase

Figure S7

A Glutamate-involved co-transmission

Element #1

unc-17 endogenous promoter — Cholinergic transporter — T2A — Flippase

unc-47 endogenous promoter — GABAergic transporter — T2A — Flippase

Promoter of serotonergic gene (*tph-1*) — Flippase

Promoter of dopaminergic gene (*dat-1*) — Flippase

Element #2

eat-4 endogenous promoter — X FRT — glutamatergic transporter — X FRT — mCherry

Readout: Endogenous *eat-4* promoter — FRT X — mCherry — cytosolic mCherry

B Acetylcholine-involved co-transmission

Element #1

unc-47 endogenous promoter — GABAergic transporter — T2A — Flippase

Promoter of serotonergic gene (*tph-1*) — Flippase

Promoter of dopaminergic gene (*dat-1*) — Flippase

Element #2

unc-17 endogenous promoter — cholinergic transporter — X FRT — STOP — X FRT — GFP

Readout: Endogenous *unc-17* promoter — cholinergic transporter — FRT X — GFP — UNC-17::GFP

C VGLUT/EAT-4-involved co-transmission

Glutamate + Acetylcholine: AFD, M5, DVA, PVN

Glutamate + GABA: I2L

Glutamate + Serotonin: No cell detected

Glutamate + Dopamine: PDE

D VAChT/UNC-17-involved co-transmission

Acetylcholine + GABA: M4, SDQR, SMD

Acetylcholine + Serotonin: ADF, HSN, VC4, VC5

Acetylcholine + Dopamine: No cell detected

Figure S8

A *eat-4* FLP-on + P*unc-47*::Flippase

I2L

B *eat-4* FLP-on + P*dat-1*::Flippase

PDE

C *unc-17* FLP-on + P*tph-1*::Flippase

ADF

D *unc-17* FLP-on + P*unc-47*::Flippase

M4
SMDV
SMDD

SDQR

HSN
VC4
VC5

Figure S9

Co-transmission atlas of the *C. elegans* nervous system

glutamate

acetylcholine

GABA

serotonin

dopamine

tyramine

octopamine

VGAT/UNC-47-dependent co-transmission*

Figure 7

A ADF presynaptic site
VAChT/UNC-17   VMAT/CAT-1

B VAChT/UNC-17 VMAT/CAT-1

C VAChT/UNC-17 VMAT/CAT-1

B' Fluorescence Intensity (A.U.) vs Distance (μm)

C' Fluorescence Intensity (A.U.) vs Distance (μm)

D KIF1A/*unc-104*
Fluorescence Intensity (A.U.) vs Distance (μm)

D' MERGE   VMAT/CAT-1   VAChT/UNC-17
soma   10 μm

E ADF presynaptic site   VAChT/UNC-17
Munc13/UNC-13   VMAT/CAT-1

F Munc13/UNC-13 VMAT/CAT-1
F'   **F"**   2 μm
10 μm

G Munc13/UNC-13 VAChT/UNC-17
G'   **G"**   2 μm
10 μm

H AiryScan imaging
VAChT/UNC-17   VMAT/CAT-1
2 μm

Figure S10