Identifying and targeting the Mg-Fe-PmrAB regulatory circuit reverses phosphate starvation-driven polymyxin resistance in *Enterobacteriaceae*

Guangming Zhang,[1] Ziqing Deng,[1,6,8] Jiezhang Jiang,[2,3,8] Minji Wang,[1,7] Xiaoyuan Wang,[4,5,*] Xiaoyun Liu,[2,3,*] Aixin Yan[1,9,*]

[1]School of Biological Sciences, Faculty of Science, The University of Hong Kong, Hong Kong SAR, China

[2]Department of Microbiology and Infectious Disease Center, NHC Key Laboratory of Medical Immunology, School of Basic Medical Sciences, Peking University Health Science Center, Beijing, China

[3]Department of Infectious Diseases, Peking University Third Hospital, Beijing, China

[4]State Key Laboratory of Food Science and Resources, Jiangnan University, Wuxi, China

[5]School of Biotechnology, Jiangnan University, Wuxi, China

[6]Present address: BGI Research, Beijing, China.

[7]Present address: School of Chemistry and Material Sciences, University of Shanghai for Science and Technology, Shanghai, China.

[8]These authors contributed equally

[9]Lead contact

* Corresponding author

E-mail: xwang@jiangnan.edu.cn (X.W.); xiaoyun.liu@bjmu.edu.cn (X.L.); ayan8@hku.hk (A.Y.).

Abstract

Phosphate (Pi) scarcity, a pervasive stressor prevalent in bacterial infections and dysbiotic host environments, potently drives antimicrobial resistance (AMR) against cationic antibiotics like polymyxins across diverse bacterial taxa. While established Pi depletion-induced AMR mechanisms often involve membrane phospholipid remodelling, *Enterobacteriaceae* largely lack these pathways, leaving their Pi scarcity-driven AMR mechanism unresolved. Here, we reveal that Pi limitation robustly induces polymyxin resistance in *Enterobacteriaceae* through 4-amino-4-deoxy-L-arabinose (L-Ara4N) modification of the Gram-negative outer membrane component lipid A, reflecting a distinct adaptive strategy. This modification is driven by strong *ugd-arn* operon induction, mediated by the PmrAB two-component system. We discover that Pi depletion triggers cellular Mg^{2+} release, prompting compensatory Fe^{3+} mobilization to the cell envelope that directly activates PmrAB. Crucially, this metal-dependent signaling axis offers a directly targetable mechanism for reversing polymyxin resistance, a significant deviation from the less accessible, PhoBR-regulated phospholipid remodelling strategies in other bacteria. We demonstrate that Mg^{2+} supplementation or Fe^{3+} chelation effectively suppresses PmrAB activation and restores polymyxin susceptibility, thereby establishing a novel, metal-centric paradigm for understanding and pharmacological intervention in stress-induced AMR.

Introduction

Within the dynamic and often hostile environment of the host, nutrient scarcity presents a pervasive challenge for invading bacteria, promoting the evolution of survival strategies and shaping pathogen persistence.[1,2] Among the macronutrients essential for bacteria, phosphorus (P) serves as a critical element for life. It is a fundamental component of key biomolecules such as nucleic acids, phospholipids, and ATP, and plays crucial roles in the essential processes of genetic inheritance, energy metabolism, membrane integrity, and signal transduction.[3,4] The primary assimilable form of phosphorus for bacteria is the orthophosphate anion (PO_4^{3-}), commonly known as inorganic phosphate (Pi). However, Pi is typically found at very low concentrations (<4µM) in diverse environments, from oceans (0.1-3µM),[5,6] and soil (0.0105 to 10.5µM, <1µM near plant roots),[7,8] to critically host-associated niches including macrophages,[9] Salmonella-containing vacuoles,[10] and cystic fibrosis sputum.[11] This localized Pi scarcity is a direct consequence of host nutritional immunity and intense microbial competition.[12,13] Evidence for widespread bacterial adaptation to Pi limitation is abundant, with metagenomic and metatranscriptomic studies consistently revealing robust transcriptional upregulation of high-affinity phosphate acquisition (Pho) regulon genes across various chronic infections and dysbiosis conditions.[11,14,15] Bacteria cope with these pervasive conditions via the conserved PhoBR two-component system, which activates genes for Pi acquisition, transport, and storage when extracellular Pi falls below 4 µM.[3,16,17]

Beyond direct Pi assimilation, Pi depletion triggers broader adaptive responses crucial for

pathogen survival, including enhanced virulence and, critically, antimicrobial resistance

(AMR).[18-20] For example, Pi limitation upregulates virulence factors in *Vibrio cholerae*,

EHEC, and *Pseudomonas aeruginosa*, and conversely, Pi supplementation can reduce

infection severity.[21-24] More importantly, Pi depletion is increasingly recognized as a potent

environmental cue driving AMR across diverse bacterial taxa, particularly against cationic

antibiotics like polymyxins. This is evidenced by increased vancomycin resistance in

Streptomyces coelicolor,[25] and enhanced polymyxin B/E resistance in *Pseudomonas*

fluorescens, *P. aeruginosa*, and *Acinetobacter baumannii* under Pi-limited conditions.[11,26-

28] Given that Pi limitation is a prevalent and dynamic stressor in host niches,

understanding its role in promoting AMR is vital for developing effective counter-strategies.

Mechanistically, Pi depletion-induced AMR is often linked to the remodelling of bacterial

membranes, where phosphorus-containing phospholipids are substituted with diverse

non-phosphorous lipids. This strategy spares intracellular phosphate and often modifies

membrane charge, which is critical for resistance to cationic antimicrobials. Examples

include the accumulation of ornithine lipids (OLs) in *Vibrio cholerae,*[29] diacylglyceryl-

N,N,N-trimethylhomoserine (DGTS) in *Rhizobium meliloti*,[30] and *Cryptococcus*

neoformans,[31] glycolipids in *Agrobacterium tumefaciens*,[32] and extensive lipidome

remodelling in *Mycobacterium tuberculosis*.[33] Specifically for polymyxin resistance, *P.*

aeruginosa synthesizes neutral monoglucosyl diacylglycerol (MGDG) and glucuronic acid

diacylglycerol (GADG),[11] while *A. baumannii* produces positively charged ornithine- and

lysine-containing aminolipids (OL and LL) under Pi limitation.[27] These diverse lipid

modifications effectively reduce the net negative charge of the bacterial membrane,

decreasing polymyxin binding and improving resistance.

However, these established lipid-remodelling strategies are not universally conserved.

Critically, genes for OL biosynthesis are absent in most *Enterobacteriaceae*, a family of

paramount clinical importance, and MGDG/GADG synthesis is not a predominant

mechanism for polymyxin resistance in this group.[34-36] Given the widespread prevalence

of Pi limitation in host environments and the escalating threat of polymyxin resistance in

Enterobacteriaceae, a fundamental question with significant clinical implications arises:

what alternative, yet unidentified, mechanisms underpin Pi depletion-induced polymyxin

resistance in this critical bacterial family, especially in the absence of the established OL

and MGDG/GADG pathways?

In this study, we address this critical gap by reporting that Pi limitation also induces

polymyxin resistance in *Enterobacteriaceae*, but through a distinct mechanism. Using *E.*

coli K-12 MG1655 as a model, we demonstrate that this process involves the upregulation

of conserved Ugd and ArnBCADTEF proteins, which catalyse the addition of the positively

charged 4-amino-4-deoxy-L-arabinose (L-Ara4N) to lipid A. This adaptive lipid A

modification is driven by a novel Pi-Mg-Fe-PmrAB regulatory circuit: loss of membrane-

stabilizing magnesium (Mg^{2+}) and compensatory iron (Fe^{3+}) mobilization to the cell

membrane activate the PmrAB two-component system, leading to upregulation of the *ugd-*

arn operon. Crucially, interfering with this circuit, via Mg^{2+} supplementation or iron

chelation, effectively reverses resistance in representative *Enterobacteriaceae* species,

but not in non-enteric species like *P. aeruginosa*. Phylogenetic analyses further reveal

distinct clades for *E. coli* PmrB (EcoPmrB) and *P. aeruginosa* PmrB (PaePmrB), with

PaePmrB lacking the conserved "ExxE" motif required for Fe^{3+} sensing. Our findings

uncover a conserved, targetable stress-responsive circuit underlying Pi depletion-induced

polymyxin resistance within *Enterobacteriaceae*, offering new avenues for managing

stress-induced antibiotic resistance.

Results

Pi depletion induces polymyxin resistance in *Enterobacteriaceae*

To investigate whether Pi limitation induces antibiotic resistance in *Enterobacteriaceae*,

we assessed the susceptibility of two representative species, *E. coli* MG1655 and

Salmonella enterica serovar Typhimurium ATCC14028. Bacteria were initially cultured in

standard MOPS synthetic minimal medium containing replete Pi (1.32 mM). Subsequently,

cultures were transferred to either Pi-replete (Pi+, 1.32 mM) or acute Pi-depleted (Pi-, 0

mM) conditions for 3 hours, a protocol adapted from R G Groat et. al.[37] Following this,

bacterial survival was determined via 3-hour polymyxin B killing assays (Figure S1).

Polymyxin B concentrations for these assays were established based on the minimum

inhibitory concentrations (MICs) of strains in MOPS minimal medium (Table S1) and a

preliminary dose-dependent killing assay (Figure S2). Results showed that Pi depletion

enhanced the survival of both species against polymyxin B (Figure 1A). Similar Pi

depletion-induced polymyxin resistance was also observed in *E. coli* O157 and four

uropathogenic *E. coli* strains isolated from urinary tract infection patients in the Queen

Mary Hospital, Hong Kong (Figure 1B)*.* Consistent with previous reports, *P. aeruginosa*

PAO1 subjected to Pi depletion also display enhanced resistance against polymyxin B

(Figure 1B). These results suggest that Pi depletion-induced polymyxin resistance is a

widespread phenomenon among Gram-negative bacteria.

Pi depletion induces 4-amino-4-deoxy-L-arabinose (L-Ara-4N)

modification of lipid A in *Enterobacteriaceae*

Since these species lack the MGDG/GADG and OL pathways described in *P. aeruginosa*

and *A. baumannii*, we employed *E. coli* K-12 MG1655 as a model to comprehensively

investigate the regulatory mechanisms underlying this stress-induced polymyxin

resistance. We performed quantitative proteomic analysis on *E. coli* MG1655 cells

subjected to the identical growth and treatment conditions as for the polymyxin B

challenging assay (Figure S1). Among the 2990 proteins detected which cover 69% of

the 4358 proteins annotated in the UniProt *E. coli* MG1655 database, abundance of 225

proteins was found to be altered following acute Pi depletion, with 142 proteins

upregulated and 83 downregulated (Figure S3) (Table S2). In addition to proteins

belonging to the Pho regulon, such as PhoB, PstB, PhoU, and PhnB (Figure 2A) in the

COG functional group of transportation and metabolism of inorganic ions (13% of

upregulated proteins) (Figure S3), the most abundant functional groups among the

significantly upregulated proteins include those involved in cell wall/membrane/envelope

biogenesis (13%) and energy production and conversion (10%) (Figure S3). Conversely, proteins involved in translation, ribosomal structure and biogenesis (33%) and amino acid transport and metabolism (17%) were significantly downregulated (Figure S3).

Specifically, among the most upregulated proteins ($\log_2$FC>10) induced by Pi depletion (Table S2), two proteins encoded in the *arnBCADTEF* operon: ArnB and ArnC, were identified (Figures 2A and 2B). Additionally, two other proteins in the Arn pathway, ArnA and Ugd, were also up-regulated with more than five-fold in response to Pi depletion treatment (Figures 2A and 2B). This finding is particularly intriguing, because Ugd and Arn proteins are known to catalyze the biosynthesis of 4-amino-4-deoxy-L-arabinose (L-Ara4N) and its modification of lipid A (Figure 2B), which confers resistance to the cationic antibiotic polymyxins due to simultaneous neutralization of a negative charge and introduction of a positive charge to lipid A.[38] To verify the upregulation of these proteins, we first examined the mRNA level of *ugd* and *arnB* genes by RT-qPCR and found their levels were 16- and 27-fold higher, respectively, under Pi- than under Pi+ conditions (Figure 2C). Next, we constructed ArnB-3×FLAG in MG1655 chromosome and observed a significant production of the ArnB protein under Pi- conditions, while its level was undetectable under Pi+ conditions (Figure 2D). We also constructed promoter-*lacZ* fusions and observed more than 10-fold higher promoter activities for both *ugd* and *arn* genes under Pi- conditions than Pi+ conditions (Figure 2E), confirming the upregulation of the Ugd-Arn system under Pi depletion.

We then extracted lipid A molecules from *E. coli* MG1655 cells cultured under identical

conditions as in the proteomic analysis, and analyzed the extract using ESI-MS. Native

lipid A, a hexacetylated glucosamine disaccharide with two phosphate groups at position

1 and 4' (m/z 1796.98), was detected in *E. coli* cells cultured under Pi sufficient conditions

(Figure 2F). In constant, the predominant lipid A species in *E. coli* cells subjected to Pi

depletion were found to be modified with L-Ara4N (m/z 1928.67) (Figure 2F). Next, we

deleted *arnT*, *arnB*, and *ugd* and found that deletion of these genes abolished Pi

depletion-induced polymyxin B resistance in *E. coli* (Figure 2G).

The induction of L-Ara4N modified lipid A, along with decrease in the unmodified form,

was also observed in *E. coli* O157 (Figure S4A), a representative UTI strain UTI-3 (Figure

S4B), and *S.* Typhimurium ATCC14028 (Figure S4C), each exhibiting varying ratios.

These fundings indicate that L-Ara4N modification of lipid A is widespread in

Enterobacteriaceae and correlates with increased polymyxin B resistance. We also

analyzed lipid A species in *P. aeruginosa* PAO1 and found that instead of the L-Ara4N

modification, a 2-hydroxymyristate modification of lipid A, which is catalyzed by the

aspartyl beta-hydroxylase LpxO, was detected in Pi-depletion treated cells (Figure S4D).

Altogether, these results indicated that unlike the strategy observed in PAO1 where Pi

stress induces resistance by replacing the negatively charged phospholipids with

MGDG/GADG, in *E. coli*, resistance is mediated by lipid A modification with L-Ara4N

catalyzed by Ugd and Arn proteins.

To delineate the sensitivity of the Ugd-Arn pathway to varying Pi levels, we measured the

promoter activities of P$_{ugd}$-*lacZ* and P$_{arn}$-*lacZ* across a gradient of phosphate

concentrations. We observed that the activities of both P_{ugd}-*lacZ* and P_{arn}-*lacZ* remained

robustly activated, at levels comparable to complete Pi depletion (0 mM), even when

0.040 mM or 0.132 mM Pi was supplemented to the medium (Figure S5). When Pi

supplementation reached 0.4 mM, their upregulation was attenuated by 60%. These

findings collectively suggest that moderate Pi limitation (below 0.4mM) is effective in

sustaining the activation of *ugd-arn* genes (Figure S5).

The PmrAB two component system is activated to up-regulate the *arn*

operon during Pi depletion

To elucidate the regulatory mechanisms underlying *ugd-arn* upregulation during Pi

limitation, we first investigated the involvement of the global phosphate regulators PhoB

and PhoR. We found that deleting either PhoB or PhoR did not affect the enhanced

polymyxin B resistance or the upregulation of the *arn* operon observed in Pi-depleted *E.*

coli (Figures 3A and S6). This observation suggested that other regulatory systems were

primarily responsible for upregulating the *arn* gene expression under Pi depletion.

The *arn* operon and *ugd* are known targets of the PmrAB two-component system (TCS)

in *E. coli.* [39] (Note the *E. coli* homolog of this system is BasSR; we adopt the PmrAB

nomenclature in this study to reflect its direct role in polymyxin resistance and its wider

usage in the literature.)[40] To determine if this regulation also extended to Pi depletion, we

constructed Δ*pmrB* and Δ*pmrA* mutants. Both the enhanced polymyxin B resistance and

the upregulation of the *arn* operon under Pi depletion were abolished in these mutants

(Figures 3A and 3B), indicating a critical role for PmrAB in *arn* regulation and polymyxin

B resistance during Pi depletion. The promoter activity of *ugd* remained unaffected in

Δ*pmrB* and Δ*pmrA* mutants (Figure S7), suggesting a distinct regulatory input for *ugd*.

Given that the PhoPQ TCS can indirectly activate the PmrAB regulon in *S.* Typhimurium

via PmrD (though this indirect regulation appears absent in *E. coli*), we also examined the

contribution of PhoPQ.[41] Deletion of *phoP* or *phoQ* did not alter the enhanced polymyxin

B resistance or *arn* operon upregulation under Pi depletion (Figures 3A and S6). These

findings suggest that the *arn* system is activated by PmrAB in response to an as-yet-

unidentified secondary signal induced by Pi depletion, rather than by direct Pi depletion

sensing or through the PhoPQ system.

Given the distinct regulation we observed for *ugd* promoter activity, we further investigated

its response to the PhoBR system. We measured *ugd* promoter activity in Δ*phoB*, Δ*phoR*,

Δ*phoP*, and Δ*phoQ* mutants and found P$_{ugd}$-*lacZ* activity decreased by approximately 30%

in both Δ*phoB* and Δ*phoR* mutants, but not in Δ*phoP* and Δ*phoQ* mutants (Figure S7).

This indicates that *ugd* is regulated by the PhoBR system, potentially alongside other

pathways, during Pi depletion.

Given the central role of the PmrAB system in mediating Pi depletion-induced polymyxin

resistance, we next investigated whether *eptA*, another PmrAB-regulated gene,

contributes to this phenotype. We constructed a Δ*eptA* mutant and assessed bacterial

survival in a polymyxin B killing assay. Deletion of *eptA* had no impact on the enhanced

polymyxin B resistance (Figure S8), indicating that *eptA* is not required for the resistance

phenotype induced by Pi depletion. This observation was further supported by ESI-MS

analysis (Figure 2F), which did not detect phosphoethanolamine (pEtN) modification of

lipid A, a hallmark of EptA activity, in Pi-depleted cells. Collectively, these results strongly

suggest that the activation of the Arn system via PmrAB is the primary mechanism

responsible for Pi depletion-induced polymyxin resistance.

To assess whether PmrAB activation occurs under Pi depletion, we examined the

phosphorylation status of the response regulator PmrA using Phos-tag SDS-PAGE, which

reliably detects phosphorylated PmrA in cells treated with known activator Fe^{3+}.[42] Results

showed that phosphorylated PmrA was detected in cells subjected to Pi depletion (Figure

3C), demonstrating that PmrAB TCS is indeed activated under Pi depletion condition. To

further confirm this in vivo, we performed a FRET-based whole-cell transcription factor-

promoter binding assay.[43] We incorporated a bright fluorescent unnatural amino acid, L-

(7-hydroxycoumarin-4-yl) ethylglycine (CouA)[44] at position G137 within the DNA-binding

domain of PmrA through genetic encoding, allowing it to serve as the FRET donor (Figure

3D).[43] The nucleic acid dye SYTO9 was utilized to stain promoter DNA of PmrA regulon

(Figure 3D),[43] acting as the FRET acceptor. The whole-cell FRET assay revealed that,

similar to the treatment with excessive $FeSO_4$ (500µM),[43] a FRET signal indicative of

PmrA binding to its target promoters was observed in cells subjected to Pi depletion

without excessive $FeSO_4$ (Figure 3E). These results confirmed that Pi depletion induced

activation of the PmrAB signal transduction system in *E. coli.*

Pi depletion drives Fe^{3+} elevation in *E. coli* cells to activate the PmrAB

TCS

PmrAB has been previously reported to be activated by ferric ion (Fe^{3+}), zinc ions (Zn^{2+}),

aluminum ions (Al^{3+}), or acidic pH.[45,46] Since no Al^{3+} was added in the culture, and the pH

remained largely unchanged (7.2-7.0) across the Pi depletion treatments (Figure S9), we

focused on the two metal ions and examined cellular metal profiles including iron (Fe),

zinc (Zn), magnesium (Mg), copper (Cu), and calcium (Ca). ICP-MS analysis revealed a

significant, greater than 5-fold increase in cellular Fe content under Pi depletion,

compared to Pi-sufficient conditions (Figure 4A). Conversely, cellular Mg levels

decreased, while Cu, Zn, and Ca levels remained largely unchanged (Figure 4A).

Corroborating this cellular Fe accumulation, the Fe content in the spent medium

decreased after 3 hours of *E. coli* incubation under Pi depletion (Figure 4B), indicating

active Fe mobilization from the medium into *E. coli* cells. These findings strongly

implicated Fe^{3+} as the activating signal for PmrAB under Pi depletion.

Given that *S.* Typhimurium PmrB (StPmrB) senses Fe^{3+} through conserved glutamic acid

residues (E36, E39, E61, E64) in its periplasmic domain,[42] we hypothesized a similar

mechanism for *E. coli* PmrB (EcoPmrB). To test this, we generated chromosomal

*pmrB*E36A and *pmrB*E39A mutations in *E. coli* MG1655 (Figure 4C). Consistent with their

proposed role in Fe^{3+} sensing, these mutations abolished the P*arn-lacZ* reporter response

to exogenous Fe^{3+} (100 μM) (Figure S10). Crucially, both mutations also abolished *arn*

operon upregulation under Pi depletion (Figure 4D), demonstrating that Pi depletion-

induced PmrAB activation requires the Fe^{3+} sensing capability EcoPmrB.

To confirm the necessity of Fe mobilization, we observed that *arn* operon upregulation

was abolished when cells were grown in Fe-free, Pi-depleted medium (Figure 4E).

Although Pi depletion significantly impaired bacterial growth (doubling time increased

from ~1 hr to ~6 hrs), co-depletion of Fe did not further exacerbate this effect, confirming

that the observed PmrAB activation specifically responds to Pi-related Fe mobilization

rather than general growth inhibition (data not shown). Investigating the sensitivity of this

response, we found that reintroduction of Fe to Pi-depleted cells restored P*arn-lacZ*

activity in a dose-dependent manner (Figure 4F), further confirming the role Fe

mobilization in activating PmrAB and *arn* genes during Pi depletion. Importantly, *arn*

activation in Pi-starved cells occurs at significantly lower exogenous Fe concentrations

(~10-fold less) than in Pi-sufficient conditions (Figure 4F), suggesting an enhanced

cellular capacity for Fe mobilization or sensing under Pi depletion.

Pi depletion triggers Mg^{2+} loss, leading to Fe^{3+} mobilization to

membrane and PmrAB activation

To elucidate the mechanism of Fe mobilization during Pi depletion, we first investigated

the role of canonical iron uptake pathways. Surprisingly, deletion of genes encoding outer

membrane iron-chelator complex receptors and enterobactin synthetase (*fecA*, *fepA*, *fiu*,

cirA, *entE*), individually or as a penta-deletion mutant (Δ5Fe), did not abolish P*arn-lacZ*

induction during Pi depletion (Figure S11A). ICP-MS analysis further confirmed that cell-

associated Fe still increased by over 5-fold in Δ5Fe cells during Pi depletion (Figure S11B).

These results indicated that Fe-mediated PmrAB activation during Pi depletion was

independent of enhanced intracellular Fe uptake and suggested an alternative

mechanism, likely involving Fe association with the cell surface, consistent with the known

ability of PmrAB to sense periplasmic and surface-bound Fe^{3+}.[42]

The outer leaflet of the Gram-negative outer membrane (OM) primarily consists of lipid A,

whose negatively charged phosphate groups are typically neutralized by cell-associated

Mg^{2+} and Ca^{2+} to maintain membrane stability.[47] Our ICP-MS analysis revealed a 44%

reduction in cellular Mg content, while Ca^{2+} levels remained unchanged, in *E. coli* cells

following 3 hours of Pi depletion treatment (Figure 4A). Monitoring the dynamics revealed

that cellular Mg progressively decreased upon transferring *E. coli* cells to Pi-depleted

medium, continuing over a 5-hour period (Figure 5A). This Mg^{2+} loss was independent of

Fe, as a similar degree of loss occurred in Fe-free, Pi-depleted medium (Figure S12).

Conversely, cellular Fe increased, peaking after two hours of Pi depletion before slightly

declining (Figure 5A). The temporal correlation between Fe accumulation and P*arn-lacZ*

activity (Figure 5B) suggested active Fe mobilization to the cell surface during Pi depletion,

potentially to compensate for Mg^{2+} loss and stabilize the membrane. The subsequent

activation of the PmrAB-Arn system that remodel lipid A by decreasing negative charges,

might thus represent an adaptive response to physiological changes, such as OM

instability, caused by Mg^{2+} dissociation. Supporting this, deletion of *pmrA* led to an even

greater and sustained increase in Fe mobilization during Pi depletion (Figure 5C), which

could be toxic to *E. coli*.[42]

To directly assess Fe association with the cell surface, we washed cell pellets with

deferoxamine (DFOM)-containing buffer, a membrane-impermeable Fe^{3+}-specific

chelator.[48] ICP-MS analysis showed that washing with DFOM-supplemented buffer

significantly reduced Fe content in cells treated with Pi depletion but had no significant

effect on cells harvested from Pi-sufficient conditions (Figure 5D). Quantification revealed

that Pi-starved cells exhibited a two-fold greater loss of Fe upon DFOM washing

compared to Pi-sufficient conditions (Figure 5E), demonstrating that Pi depletion

promoted Fe mobilization to cell surfaces, where it remained susceptible to removal by a

membrane-impermeable chelator.

To uncover the driving force behind this enhanced Fe mobilization, we supplemented the

culture medium with excess Mg (10 mM) to counteract Mg loss. This supplementation

significantly reduced Fe mobilization to MG1655 cells during Pi depletion (Figure 6A);

correspondingly, P_{arn}-*lacZ* activity declined as Mg concentration increased in the Pi-

depleted medium (Figure 6B). These results strongly suggest that Fe mobilization to the

cell surface during Pi depletion is dependent on Mg^{2+} loss. Further, monitoring Mg^{2+}

dynamics in the spent medium revealed a continuous increase in Mg levels throughout

the Pi depletion treatment (Figure 6C), confirming active Mg^{2+} dissociation from the cells.

Similar Mg^{2+} release was observed even during combined Mg and Pi depletion (Figure

S13). These results suggested that while Mg^{2+} is essential for bacterial growth, an

imbalance with phosphorus during Pi depletion appears to trigger active Mg^{2+} dissociation

from the cells. A comparable 45% Mg^{2+} loss was also observed in *P. aeruginosa* PAO1

under similar conditions (Figure S14). However, Fe mobilization was less pronounced

(2.4-fold vs. 5.4-fold in *E. coli*), reflecting species-specific adaptation strategies.[11]

Given that membrane-associated Mg^{2+} primarily neutralizes the negative charges of lipid

A to stabilize the OM, we hypothesized that Mg^{2+} loss during Pi depletion alters OM

properties, and Fe mobilization may serve to stabilize the OM. To test this, we employed

an OM-sensitive probe.[49] The probe reports membrane surface properties through two

orthogonal readouts: an overall fluorescence increase, which reflects probe disassembly

upon binding to the anionic bacterial surface, and the pyrene excimer-to-monomer

emission ratio (I_{482}/I_{375}), which captures differences in local surface interactions. The 2D

sensor output revealed distinct signals between cells grown under Pi-sufficient and Pi-

depleted conditions (Figure 6D), indicating altered OM characteristics. Notably, cells

grown in Fe-free, Pi-depleted medium exhibited even greater shifts (Figure 6D),

suggesting that Fe mobilization and subsequent activation of the PmrAB-Arn pathway

served as an adaptation strategy to maintain OM integrity during Pi stress. This strategy

likely prevents uncontrolled Fe accumulation and toxicity,[42] ultimately resulting in stress-

induced polymyxin resistance. A schematic model illustrating this process is presented

(Figure 7).

Disrupting the Mg-Fe-PmrAB-Arn circuit counters Pi depletion-

induced polymyxin resistance in *Enterobacteriaceae*

Our findings above identified a Mg-Fe-PmrAB-Arn circuit underpinning the polymyxin

resistance induced by stress adaptation to Pi depletion. To evaluate whether interference

with this pathway could mitigate the stress-induced polymyxin resistance, we

supplemented Pi depletion medium with excess Mg^{2+} (10mM) or applied a membrane

impermeable chelator DFOM (10μM). Both approaches effectively suppressed Pi

depletion-induced polymyxin resistance in *E. coli* and *S.* Typhimurium (Figure 8A). In

contrast, neither intervention reversed polymyxin resistance in *P. aeruginosa* despite

various combinations of Mg^{2+} supplementation and Fe chelation (Figure S15). This

highlights species-specific adaptative strategies and aligns with several observations

from our study as well as several previous research.[11,50] Specifically, in *P. aeruginosa,*

despite a 45% Mg loss during Pi depletion, Fe levels increased only two-fold compared

to over five-fold in *E. coli*, consistent with its reliance on MGDG/GADG lipids synthesis

regulated by the PhoBR system (Figure 8B).[11] Additionally, *P. aeruginosa* exhibits a

distinct lipid A modification, 2-hydroxy myristate (Figure S4D), that does not confer

polymyxin resistance.[51] Phylogenetic analysis of 250 PmrB homologs revealed that

EcoPmrB and PaePmrB cluster in separate clades, with PaePmrB lacking the conserved

"ExxE" motif essential for Fe^{3+} sensing (Figure 8C). Together, these findings underscore

the diverse stress response strategies employed by different classes of Gram-negative

bacteria and highlight the importance of understanding species-specific regulatory

mechanisms for developing targeted interventions against stress-induced antibiotic

resistance.

Discussion

Bacterial adaptation to environmental stressors is crucial for survival and pathogenesis. Phosphate (Pi) scarcity, a pervasive condition in diverse environments including host-associated niches such as macrophage phagosomes, salmonella-containing vacuoles, and cystic fibrosis sputum, is a potent cue driving antimicrobial resistance (AMR) against cationic antibiotics such as polymyxins.[9-11] Adaptive mechanisms previously characterized for Pi depletion-induced AMR often involve remodeling bacterial membranes by substituting phosphorus-containing phospholipids with Pi-free, neutral, or positively-charged lipids.[11,26-28] However, the specific mechanisms in clinically important Enterobacteriaceae, which largely lack these canonical phospholipid remodeling pathways, have remained largely unknown. In this study, we unveil a conserved, yet distinct, polymyxin B resistance mechanism in *Enterobacteriaceae*: Pi limitation triggers a cascade of cationic imbalance and Fe^{3+} signaling, leading to L-Ara4N modification of lipid A and outer membrane remodeling, a process distinct from phospholipid-based resistance pathways (Figure 7). This intricate Mg-Fe-PmrAB regulatory circuit represents a previously unappreciated ecophysiological vulnerability, offering a novel strategy to mitigate stress-induced polymyxin resistance.

Maintaining a balanced P/Mg ratio is paramount for bacterial cellular physiology, with ATP and rRNA serving as primary cytoplasmic reservoirs for both Pi and Mg^{2+}.[52-55] Previous work by Bruna et al. demonstrated that bacteria suppress Pi uptake under Mg^{2+} limitation

to preserve cellular function.[53] Herein, our study extends this understanding by showing that Pi scarcity itself actively triggers Mg^{2+} release from *E. coli* cells. This observation underscores the tightly coupled nature of Pi and Mg^{2+} homeostasis. Our comprehensive proteomic analysis further supports this, revealing that proteins involved in translation, ribosomal structure, and biogenesis were among the most significantly downregulated groups during Pi depletion (Figures 1A and S3). Concurrently, we observed a steady decline in cellular ATP levels under Pi-limited conditions (Figure S16), suggesting a profound arrest in ribosome biogenesis and protein synthesis that likely contributes to the observed Mg^{2+} release and cellular loss.

In *Enterobacteriaceae*, Mg^{2+} homeostasis is primarily governed by the PhoPQ two-component system, which is well-characterized in *S.* Typhimurium to be activated under low extracellular Mg^{2+} conditions.[56,57] Given the observed Mg^{2+} release during Pi-depletion, we investigated the involvement of this classical Mg^{2+}-sensing PhoPQ system. Our results showed that despite cellular Mg^{2+} loss during Pi depletion, the transcription of key PhoPQ regulon genes, such as *phoQ* and *mgtA*, remained unchanged (Figure S17). This indicates that while cellular Mg^{2+} levels are perturbed, they do not drop sufficiently to activate the canonical PhoPQ response. Instead, our data compellingly demonstrate that under acute Pi depletion, *Enterobacteriaceae* preferentially engage an alternative, Fe^{3+}-mediated membrane remodeling pathway. This pathway activates the *arn* operon and subsequent L-Ara4N modification independently of the classical Mg^{2+}-sensing PhoPQ system, highlighting a distinct adaptive strategy.

This Fe^{3+}-mediated lipid A modification strategy in *Enterobacteriaceae* offers a stark

contrast to Pi depletion-induced polymyxin resistance mechanisms described in other

Gram-negative bacteria, notably *P. aeruginosa*. In *P. aeruginosa*, Pi depletion primarily

activates the PhoBR regulon, orchestrating the biosynthesis of phosphorus-free

glycolipids (MGDG and GADG) to replace negatively charged phospholipids and maintain

membrane integrity (Figure 8B).[11] While *P. aeruginosa* possesses a PmrAB homolog, its

PmrB sensor kinase lacks the critical Fe^{3+}-sensing "ExxE" motifs found in

Enterobacteriaceae PmrB, precluding direct Fe^{3+} response. Instead, PmrAB activation in

P. aeruginosa is typically coupled to the PhoPQ system, which senses low extracellular

Mg^{2+}. Thus, while *P. aeruginosa* employs PhoBR-regulated phospholipid remodeling as

its primary Pi depletion response, *Enterobacteriaceae*, largely lacking these glycolipid

pathways, have evolved to utilize a secondary, Fe^{3+}-sensing mechanism. Here, the

PmrAB system, via its specific Fe^{3+}-sensing "ExxE" motifs in PmrB, orchestrates

membrane remodeling through L-Ara4N modification of lipid A as a key physiological

adaptation to Pi depletion.[50] While *P. aeruginosa* also exhibits lipid A modification (e.g.,

2-hydroxymyristate by LpxO) during Pi depletion, this modification does not contribute to

polymyxin resistance and is regulated independently of PhoBR, PhoPQ, or PmrAB,

highlighting distinct physiological roles.[51]

The biosynthesis of L-Ara4N and its transfer to lipid A are mediated by Ugd and

ArnABCDEFT proteins, encoded by the *ugd* gene and the *arn* operon, respectively[58-61].

Our data reveal a sophisticated regulatory separation: during Pi depletion, *arn*

transcription is tightly controlled by the PmrAB system, while ugd expression is not

PmrAB-dependent and exhibits partial regulation by the PhoBR system (Figure S6). This

bifunctional control is not unexpected, as *ugd* encodes UDP-glucose dehydrogenase, an

enzyme with dual roles in L-Ara4N biosynthesis and in the production of

exopolysaccharides like colanic acid.[62,63] Therefore, while PmrAB activation of *arn* is

absolutely critical for L-Ara4N modification of lipid A during Pi limitation, *ugd* regulation

reflects its broader involvement in diverse cellular metabolic and stress responses.

While preparing this manuscript, a recent study by Baijal et al. (2024) reported that the

ppk gene modulates *arn* operon upregulation under combined nutrient limitations (transfer

from rich LB to low-Pi, no-amino-acid MOPS medium) in *E. coli.*[64] Inspired by this, we

investigated the role of *ppk* in polymyxin resistance specifically under Pi depletion, our

primary focus. We found that *ppk* deletion did not affect *arn* or *ugd* promoter activity under

sole Pi-depleted conditions (Figure S18), indicating its dispensability when Pi is the only

limiting factor. To reconcile these findings, we replicated the Baijal et al. experimental

conditions, confirming *ppk*-dependent *arn* upregulation under combined Pi and amino acid

scarcity (Figure S19A). Our subsequent experiments, systematically disentangling these

stresses, revealed that while both Pi and amino acid depletion induce *arn* upregulation, Pi

depletion is the dominant driver, and *ppk*-dependent regulation emerges only under

combined Pi and amino acid scarcity (Figure S19B). These results clarify that while *ppk*

influences *arn* expression under specific multifactorial stresses, our identified PmrAB-arn

pathway is independently and dominantly activated by Pi depletion alone, providing

precise mechanistic insight into this fundamental environmental stressor.

In both natural habitats and host-associated niches, Pi scarcity imposes dual challenges:

nutrient acquisition and defense against antimicrobial threats. The Pi-Mg-Fe-PmrAB

regulatory circuit we have uncovered likely represents a sophisticated adaptive strategy

for *Enterobacteriaceae*, facilitating their survival in highly heterogeneous environments

with fluctuating Pi levels, such as during host colonization, transmission, or environmental

persistence.[65] This PmrAB-mediated membrane remodeling not only confers resistance

to polymyxins but may also enable bacteria to preserve membrane integrity and withstand

other cationic antimicrobial peptides (CAMPs) produced by competing microbes or host

defenses.

In summary, our findings elucidate a conserved yet distinct regulatory circuit in

Enterobacteriaceae that intricately links Pi and Mg^{2+} homeostasis to Fe^{3+}-mediated lipid

A modification, driving stress-induced antibiotic resistance. This mechanism is notably

distinct from those characterized in *P. aeruginosa* or *A. baumannii*. Crucially, we

demonstrate that targeting this circuit effectively mitigates stress-induced resistance.

Given that polymyxins remain indispensable as last-line antibiotics for treating multidrug-

resistant Gram-negative bacterial infections, our study not only offers critical insights into

the fundamental molecular mechanisms underlying stress-induced resistance but also

highlights a promising avenue for informing novel resistance management strategies.

Materials and methods

Bacterial strains and growth conditions

Strains, plasmids, and primers used in this study are listed in the S2 and S3 Tables. Bacteria were cultured in lysogeny broth (LB) broth/plates or MOPS minimal medium. MOPS medium was freshly prepared from 1× MOPS stock by supplementing 0.1 % (w/v) glucose and 1.32 mM K_2HPO_4. Acute Pi depletion was applied by transferring bacteria culture into the non-Pi MOPS medium. 1× MOPS consists of 40 mM MOPS, 4 mM Tricine, 10 µM $FeSO_4 \cdot 7H_2O$, 9.5 mM NH_4Cl, 0.276 mM K_2SO_4, 0.5 µM $CaCl_2 \cdot 2H_2O$, 525 µM $MgCl_2$, 50 mM NaCl, 0.003 µM $(NH_4)_6Mo_7O_{24}$, 0.4 µM H_3BO_3, 0.03 µM $CoCl_2$, 0.01 µM $CuSO_4$, 0.08 µM $MnCl_2$, and 0.01 µM $ZnSO_4$. Antibiotics applied to select and maintain the construct were chloramphenicol (25 µg/ml), kanamycin (20 µg/ml) and ampicillin (100 µg/ml).

Lipid A extraction and ESI-MS analysis

Lipid A was extracted from designated *E. coli* cells using Bligh-Dyer method as described with minor modifications.[66] *E. coli* MG1655 cultured in MOPS medium to OD_{600} of 0.5 was harvested as cells under Pi sufficient condition. The same volume of cells was washed with Pi depletion medium for 3 times and resuspended in Pi depletion medium and shake for 3 hrs at 37°C as cells subjected to Pi depletion. 200 ml of each type of cell pellets with three biological replicates were harvested by centrifugation at 4500 rpm for 30 min and

washed with H_2O. The cell pellets were suspended in 76 ml of a single-phase Bligh-Dyer

mixture (chloroform: methanol: water=1:2:0.8; v/v/v) and stirred at room temperature for

1 h. Insoluble debris was collected by centrifugation at 4500 rpm for 30 min, washed with

60 ml single-phase Bligh-Dyer mixture, and suspended in 27 ml of 12.5 mM sodium

acetate (pH 4.5). The mixture was heated at 100°C for 30 min and cooled to room

temperature. The suspension was mixed with 30 ml chloroform and 30 ml methanol.

Following vigorously shaken for extraction of lipid A and centrifugation for 30 min, the

lower phase containing lipid A was collected and dried with a rotary evaporator.

All the mass spectra of lipid A samples were acquired using Waters SYNAPT quadrupole

time-of-flight mass spectrometer (MS) equipped with an electrospray ionization source[67].

Lipid A samples were dissolved in a mixture of chloroform and methanol (4 : 1; v/v) and

subjected to ESI/MS in the negative ion mode. Data acquisition and analysis were

performed using MASS LYNX V4.1 software, Waters, Milford, MA, USA.

ICP-MS and DFOM treatment

Total cellular metal content was quantified using inductively coupled plasma mass

spectrometry (ICP-MS) following previous description with minor modifications[68]. *E. coli*

cells equivalent to OD_{600} as 5 were harvested by centrifugation at 4500 rpm for 15 mins

and washed three times with ice-cold PBS buffer. Cell pellets were subjected to acid

digestion with 100 µl 65.0 % HNO_3 (Fluka) at 65°C for 16hrs. Samples were diluted to a

final volume of 15 ml with 1% HNO_3 and subjected to ICP-MS detection on Agilent 7900

system (Agilent Technologies, CA). Samples were further diluted if the signals exceeded

the liner range of standard curve. The standard curves of tested metals were generated

from multi-element calibration standard-2A for ICP-MS (Agilent). Metal contents in

bacteria were calculated based on the standard curve and results were normalized to

original cell number.

Deferoxamine mesylate salt (sigma) were applied to examine the degree of iron

association to *E. coli* cell surface. *E. coli* cells equivalent to OD_{600} as 5 were harvested by

centrifugation at 4500 rpm for 15 mins. Cell pellets were resuspended in HEPES buffered

saline with and without 50µM DFOM and incubated in 37°C for 30 mins with 220rpm

agitation. Total cellular metal content of these treated cells was quantified using

inductively coupled plasma mass spectrometry (ICP-MS) as described above.

OM feature monitoring using PI-BactD

OM feature of *E. coli* was monitored by an OM-sensitive probe PI-BactD following previous

description with minor modifications.[49] *E. coli* cells grown under designated conditions

were harvested by centrifugation at 13000 rpm for 2 mins and washed with PBS buffer

(10 mM, pH 7.2) twice. Cell pellets were resuspended in PBS buffer to an OD_{600} of 0.5.

Fluorescence probe PI-BactD (gift from prof. Xu Z, Dalian) was added into 1mL cell

suspension to a final concentration of 2 mM and incubated at room temperature for 1 min.

Output signals of two channels: fluorescence increase ($\Delta S/S_0$) and fluorescence ratio

metric changes (I_{482}/I_{375}) were recorded to plot the 2D graph (Thermo Scientific

Varioskan® Flash spectral scanning multimode reader). Six replicates were conducted for

each strain and treatment.

Construction of P$_{promoter}$-*lacZ* (P$_{arn}$-*lacZ* as example) fusion in the single-copy plasmid pNN387

A DNA fragment corresponding to the -200 to -1 region relative to the ATG start codon of

arnB was amplified using *E. coli* MG1655 genomic DNA as the template and primers

specific to this fragment flanked with NotI and HindIII restriction sites at the 5'- and 3'-end,

respectively. Following digestion with NotI and HindIII and purification using the MiniBEST

agarose gel DNA extraction kit (TaKaRa), the fragment was ligated into the plasmid

pNN387 which was treated with the same restriction enzymes using the quick ligation kit

(NEB). The resulting construct was selected on LB agar plates supplemented with

chloramphenicol (25 µg/ml) and was verified by DNA sequencing (BGI, Shenzhen).

Beta-galactosidase activity assay

β-gal activity assay was performed based on a previous description with modifications.[68]

Cells were grown under designated conditions until OD$_{600}$ reached 0.4. 10 µg/ml

tetracycline was added to stop cell growth and protein synthesis, and cells were left on

ice until being assayed. 100µl of cell culture and 400 µl of Z buffer (60 mM Na$_2$HPO$_4$, 40

mM NaH$_2$PO$_4$, 10 mM KCl, 1 mM MgCl$_2$, 50 mM β-mercaptoethanol, pH 7.0, 4 °C) were

mixed. One drop of 0.1% SDS and two drops of chloroform were added to lyse the cells.

The reactions were initiated by adding 100 µl of 4 mg/ml ortho-nitrophenyl-β-galactoside

into cell suspension and were incubated at 28 °C. Color development in the mixtures was

monitored. Finally, 250µl of 1M Na_2CO_3 was added to stop the reaction. Color of the

mixtures were measured using spectrophotometer at OD_{420} and OD_{550}. The expression

value in Miller units was calculated, and results are presented as the mean of three

biological replicates.

RNA extraction and RT-qPCR

Bacterial cells grown under designated conditions was subjected to RNA extraction when

the OD_{600} reached 0.4. Cells were harvested by centrifugation at 4,500 rpm for 10 min at

4 °C. After removing the supernatant, the cell pellet was stored at -80 °C to aid cell lysis.

Total RNA was extracted using the MiniBEST Universal RNA Extraction Kit (Takara,

Japan) according to the manufacturer's instruction. Reverse transcription (RT) was

performed using PrimeScript^{TM} RT reagent Kit (Takara, Japan). Quantitative PCR was

performed using specific primers and the TB Green® Premix Ex Taq™ (Takara, Japan)

in a 20 µL reaction system. The reaction was performed in the ABI StepOnePlus real time

PCR system with *rrsB* as reference gene to normalize the relative expression of the target

genes. The results were presented as fold change expression of the target genes, and

results were presented as the mean of three biological isolates.

Construction of clean gene deletion, point mutation and tag insertion

in chromosome of *E. coli* MG1655

Gene deletion, mutation, and tag insertion in chromosomal were achieved employing a

CRISPR/Cas9-mediated approach developed by Zhao et al. with modifications.[69] For

gene deletion (deletion of *lacZ* as example), *E. coli* MG1655 was electrotransformated

with pCAGO plasmid which contains both SpCas9 and λ-RED to obtain MG1655-pCAGO.

A single colony of the resulting transformant was inoculated into LB medium

supplemented with ampicillin (100 mg/ml) and grew overnight at 30°C. 100 μl of the

overnight culture was inoculated into 10 ml fresh culture medium and cells were grown at

30°C with IPTG (0.1 mM) induction to OD600 as 0.6. The cells were collected and washed

with cold ddH$_2$O for three times, generating competent cells for electroporation. An aliquot

comprising 100 μl competent cells was mixed with 800 ng of the editing-cassette which

encompasses 40 bp of sequence among the CDS of *lacZ* gene followed by the

chloramphenicol resistant gene, the N20PAM sequence

(GTCCATCGAACCGAAGTAAGG), 40 bp upstream-homologous arm of *lacZ* gene, and

40bp downstream-homologous arm of *lacZ* gene, in a 2 mm Gene Pulser cuvette (Bio-

Rad) and was subjected to electroporation at 2.3 kV. Following recovering the

transformants by cultivation in LB medium for 2 hrs at 30°C, the mixture was spread onto

LB agar plate supplemented with ampicillin (100mg/ml), chloramphenicol (25 mg/ml), and

1% glucose (to avoid the background expression of the Cas9 protein). Single colonies

were collected and verified by colony PCR to obtain an authentic clone in which the

editing-cassette was embedded in chromosomal. The confirmed colony was inoculated

into LB medium supplemented with ampicillin (100 mg/ml), IPTG (0.1 mM), and L-

arabinose (0.2%) at 30°C overnight to enable expression of SpCas9 and the λ-RED

recombinase. The cells were then streaked on LB agar plates supplemented with

ampicillin (100mg/ml). Following colony PCR, desired colonies containing deletion of the

lacZ gene were verified by DNA sequencing (BGI, Shenzhen). For point mutation,

(mutation of *pmrB* E36 to A as example) *E. coli* MG1655 was conducted following the

same procedures except for the editing cassette sequence which encompasses 40 bp of

upstream-homologous arm of GAA (Glu36) followed by the chloramphenicol resistant

gene, the N20PAM sequence, the 40 bp upstream-homologous arm of GAA (Glu36), GCA

(Ala) and 40 bp downstream-homologous arm of GAA (Glu36). For tag insertion, (insertion

3xFLAG to C-terminal of *arnB* gene as example) *E. coli* MG1655 was conducted following

the same procedures except for the editing cassette sequence which encompasses 40 bp

of upstream-homologous arm of stop codon of *arnB* gene followed by the chloramphenicol

resistant gene, the N20PAM sequence, the 40 bp upstream-homologous arm of stop

codon of *arnB* gene, tag sequence and 40 bp downstream-homologous arm of stop codon

of *arnB* gene (include stop codon). After obtaining all desired genome modifications, the

editing plasmid was cured by growing the resulting cells at 42°C for 48 hrs.

Construction of pET28A-pBAD-*pmrA* for PmrA protein expression

A DNA fragment encoding the PmrA protein was amplified using genomic DNA of *E. coli*

MG1655 as the template and primers specific to this fragment flanked with NcoI and XhoI

restriction sites at the 5'- and 3'-end, respectively. After digestion with NcoI and XhoI, and

purification using the MiniBEST agarose gel DNA extraction kit (TaKaRa), the purified

fragment was ligated into the plasmid pET28a-pBAD and pET28a-pBAD-3xFLAG [43] which

was treated with the same restriction enzymes using the quick ligation kit (NEB). The

resulting construct was selected on LB agar plates supplemented with kanamycin (20

µg/ml) and was verified by DNA sequencing (BGI, Shenzhen). All plasmids constructed

are listed in the supplementary S3 Table.

Immunoblot analysis

Protein levels of ArnB and PmrA were analyzed by SDS-PAGE following Western Blot.

1ml of cells with OD_{600} of 0.5 was collected and resuspended in 200 µl sample buffer (13

mM Tris, 0.2% SDS, 5% glycerol, 0.004% bromophenol blue, pH 6.8) before being

subjected to cell lysis by sonication. Following centrifugation at 13000 rpm for 10 min at

4°C, 20 µl of the sample was separated on SDS-PAGE gels (TGX FastCast Acrylamide

Kit, 10%, BIO-RAD). Proteins were then transferred onto a polyvinylidene difluoride

(PVDF) membrane (BIO-RAD) using a Trans-Blot SD Semi-Dry Transfer Cell (BioRad) at

constant voltage (25 V) for 7 min. Membranes were blocked for 1 h with 5% non-fat milk

solution in TBST (20 mM Tris, 150 mM NaCl, 0.1% Tween-20) at room temperature

followed by incubation with 1:10,000 primary antibody (monoclonal or polyclonal anti-His

antibody) in 1% non-fat milk in TBST at room temperature with shaking for 1 hour. After

washing with TBST, the membrane was incubated with the secondary antibody (Goat anti-

mouse IgG HRP conjugate) in 1% non-fat milk in TBST at room temperature with shaking

for 1 hour. Chemiluminescence of protein bands was detected by ECL Plus Western

Blotting Detection Reagents (GE Healthcare, USA), and imaged using a X-ray film

development by Alliance Q9 (UVITEC, United Kingdom).

Phos-tag SDS-PAGE

In vivo phosphorylation of PmrA was examined using Phos-Tag SDS-PAGE following our previous description with minor modifications.[43] *E. coli* MG1655Δ*pmrA* cell harboring pET28a-pBAD-*pmrA*-3×FLAG (AY7068) was inoculated into MOPS minimum medium supplemented with kanamycin (20 μg/ml) and grew overnight at 37°C. 100 μl of the overnight culture was inoculated into 5 ml fresh MOPS medium and grew at 37°C for 4 hrs. Expression of PmrA was induced with 1mM L-arabinose for 1 hr. Cells were collected by centrifugation at 13000 rpm and were washed with MOPS medium supplemented with 0.1, 0.5, or 1 mM $FeSO_4$ or Pi- MOPS medium twice and shake for 1 hr at 37°C in the corresponding medium. 1ml of cells with OD_{600} of 0.5 was collected for the subsequent analysis. Cell pellet was washed with cold PBS twice and was resuspended in 200 μl sample buffer (13 mM Tris, 0.2% SDS, 5% glycerol, 0.004% bromophenol blue, pH 6.8) before being subjected to cell lysis by sonication. Following centrifugation at 13000 rpm for 10 min at 4°C, 20 μl of the sample was loaded onto a 10% Tris-Gly polyacrylamide gel containing 25 mM Phos-tag Acrylamide (Wako) and 50 mM $MnCl_2$. Samples were separated by electrophoresis for 2 hrs at 4°C, following washing with 50 ml transfer buffer containing 10 mM EDTA for 20 min to remove residual Mn^{2+}. Proteins on the gel were transferred onto a PVDF membrane. The membrane was blocked with 5% non-fat milk in TBST buffer and was incubated subsequently with monoclonal ANTI-FLAG® M2 antibody (Sigma) and goat anti-mouse IgG (H+L)-HRP conjugate (Bio-rad). To detect FLAG-tagged protein bands, the membrane was submerged in the ECL Plus Western Blotting Detection

656   Reagents (GE Healthcare) for 2 min and then subjected to the imaging system (UVITEC

657   Cambridge) to record signals.

658   Amber codon mutation of selected CouA incorporation sites in

659   pET28A-pBAD-*pmrA*

Selective amino acid located in the DNA binding domain of PmrA was subjected to

replacement by CouA, each of these CouA incorporation sites were mutated to a TAG

amber codon by site-directed mutagenesis PCR. PCR was conducted using a pair of

primers for the desired mutagenesis and pET28A-pBAD-*pmrA* plasmid as the template.

PCR product was purified using MiniBEST agarose gel DNA extraction kit (TaKaRa).

Following DpnI digestion of the template for 16 hrs, the PCR product was transformed into

E. coli DH5α. Transformants were recovered on LB agar plates supplemented with

kanamycin (20 µg/ml) at 37°C overnight. Colonies recovered were verified by DNA

sequencing (BGI, Shenzhen).

669   In vivo PmrA$_{CouA}$ - promoter$_{SYTO\ 9}$ binding assay

Binding of PmrA$_{CouA}$ and promoter$_{SYTO\ 9}$ was monitored by FRET assay following our

previous description with minor modifications[43]. Plasmid pET28a-pBAD-*pmrA*-G137TAG

was co-transformed with pEVOL-CouRS into *E. coli* MG1655 *ΔpmrA* strain, resulting in

AY7065 constructs. A fresh single colony of AY7065 was inoculated into MOPS medium

supplemented with kanamycin (20 µg/ml) and chloramphenicol (25 µg/ml) and grew

overnight at 37°C. 75 µl of the overnight culture was inoculated to 7.5 ml fresh medium to

subculture the bacteria. Following culturing at 37°C for 4 hrs, CouA incorporation and

expression of PmrA-G137CouA was induced mildly by supplementing 0.9% arabinose

and 1mM CouA at 37°C for 5 hrs. Approximately 3.5×10^{10} cells were harvested and

washed twice with fresh MOPS medium. Cells were resuspended in MOPS medium as

control, or MOPS medium supplemented with 0.5 mM $FeSO_4$, or Pi- MOPS medium and

shake for 1 hr at 37°C. After centrifugation at 4500 rpm for 15 min at 4°C, cell pellet was

resuspended in 600 µl of PBS. 85 µl of the suspension was added in each of 6 wells (3

wells for control and 3 wells for experimental group) on a 96-well black plate preloaded

with 15 µl of 50 µM SYTO 9 and was incubated for 15 min at room temperature in the dark

before being subjected to fluorescence recording (Thermo Scientific Varioskan® Flash

spectral scanning multimode reader).

Comparative proteomic analysis

E. coli MG1655 cultured in MOPS medium to OD_{600} of 0.5 was harvested as cells under

Pi sufficient condition. The same volume of cells was washed with Pi depletion medium

for 3times and resuspended in Pi depletion medium and shake for 3 hrs at 37°C as cells

subjected to Pi depletion. Each type of cell pellets with three biological replicates were

lysed by resuspending in 20 µl BugBuster reagent (Novagen, USA) (containing 25 mg/ml

lysozyme, 1 unit DNase I) at room temperature for 30 mins following centrifugation at

13000 rpm at 4°C for 20 min. Supernatant was transferred to a clean microcentrifuge tube

and mixed with 4×SDS sample buffer (40% glycerol, 8% SDS, 240 mM Tris/HCl pH 6.8,

0.04% bromophenol blue) supplemented with 5% β-mercaptoethanol. Sample were

incubated at 55 °C for 25 min. 20 μL of each sample were loaded onto a polyacrylamide

gel composed of a 10% resolving (lower) gel and a 6% stacking (upper) gel following

staining with Coomassie blue staining and fractionated into eight fractions. Gel slices were

cut into 1 mm^3 cubes and destained with 50% acetonitrile (ACN) in 50 mm triethyl

ammonium bicarbonate (TEAB) and dehydrated with pure CAN and dried in a vacuum

concentrator. Sequencing grade modified trypsin (Promega) diluted by an ice-cold buffer

containing 50 mM NH4HCO3 and 10% (v/v) ACN was added to cover the dehydrated gel

pieces on ice. After 30 min, more trypsin-containing buffer was added to cover the

expanded gel pieces followed by overnight incubation at 37 °C. Finally, the resulting tryptic

peptides were extracted from the gel by equilibrating the samples with 50% ACN and 5%

formic acid (FA) for 20 min in a 37 °C shaker. The extraction was repeated once, and all

fractions were combined and vacuum dried. The protein digest was resuspended in 100

μl of 100 mm TEAB, and then 4 μl of 0.6 m sodium cyanoborohydride (NaBH$_3$CN) were

added. Peptides from PI+ and Pi- were labeled with 4 μl of 4% formaldehyde (CH$_2$O) and

deuterated formaldehyde (CD$_2$O), respectively. The reaction mixture was vortexed and

incubated at room temperature for 1 h. The reaction was quenched by sequential addition

of 16 μl of 1% (v/v) ammonia solution and 8 μl of formic acid. Finally, light- and heavy-

labeled peptides were mixed, and vacuum dried immediately before further mass

spectrometric analyses. Extracted peptides were analyzed by Nanoflow LC-MS/MS

analyses using the EASY-nLC 1000 System (Thermo Scientific, USA) and LTQ Velos

(Thermo Scientific, USA).

Data processing was performed following a previous description[70]. For protein

identification, MS/MS spectra were extracted from RAW files using MaxQuant

(http://maxquant.org/, version 1.5.3.30). MASCOT (Matrix Science, version 2.3.02) was

used to perform database search against an *E. coli* database (strain MG1655, with

sequence files downloaded from UniProt in FASTA format, 4,358 entries. Average mass

was selected with precursor mass tolerance of 1.5 Da and fragment mass tolerance of 0.8

Da. Methionine oxidation was set as a dynamic modification. The resulting assignments

were filtered to achieve a false-discovery rate (FDR) of <1% at both the peptide and

protein levels using the target-decoy method. Relative abundances of protein were

represented by spectral counts, which are the number of repeated identification of

peptides during the entire analysis.

To identify proteins that were differentially expressed in the sets of samples, the spectral

counts data were subjected to statistic by DESeq package (Anders and Huber 2010) in R

(Team 2014). Briefly, to eliminate the variance caused by sample loading and machine

reading depth, the total counts data of two biological conditions was normalized to the

same size. The following statistics were performed: *foldChange* (fold change from

condition A to B), *pval* (p value for the statistical significance of this change), and *padj* (p

value adjusted for multiple testing with the Benjamini-Hochberg procedure, which controls

false discovery rate). Data set with *padj* <0.05 were considered to be significant (Anders

and Huber 2010), and corresponding proteins were listed in S1 Table.

Polymyxin challenging assay

Challenging *E .coli* cells with designated concentration of polymyxin B (Sigma. USA) were

performed as described with modifications.[71] *E. coli* cells grown under designated

conditions were harvested by centrifugation at 13000 rpm for 1min. Cell pellets were re-

suspended in MOPS medium to a final concentration of 10^8 cells/ml. The cell culture was

treated with polymyxin B with designated concentration for 1-3 hours at 37 °C without

agitation. Cell culture prior to and following polymyxin treatment was plated on LB plate,

CFU was counted after 12-16 hours incubation at 37°C. Survival of polymyxin challenging

was determined by the survival rate of *E. coli* cells after drug treatment. Results are

presented as the mean of three biological replicates.

Phylogenetic analysis of EcoPmrB homologs

The PmrB sequences from *E. coli* K-12 MG1655 were used as a query against γ-

proteobacteria via BLASTP searches in the UniProt database, with an E-threshold of 10

and the BLOSUM62 substitution matrix. The top 250 retrieved sequences were aligned

using the MUSCLE algorithm implemented in MEGA12 software. A phylogenetic tree of

EcoPmrB homologous was constructed using the Maximum Likelihood method, with

branch support assessed via adaptive bootstrap analysis using 562 bootstrap replicates

in MEGA12. The JTT model was applied as the evolutionary model for tree construction.

Statistical analysis

757   Statistical analysis was performed using GraphPad Prism. Group comparisons between

758   two groups were performed using two-tailed Student's *t* test. Group comparisons between

multiple groups were performed using one-way ANOVA or two-way ANOVA with post hoc tests.

Acknowledgments

We thank Ms. Yining Chen in Prof. A. Yan's lab for proofreading the manuscript. We thank Prof. Changhao Bi (Tianjin Institute of Industrial Biotechnology, Chinese Academy of Sciences) for the generous gift of pCAGO plasmid. We thank Prof. Zhaochao Xu and Prof. Lu Miao (CAS Key Laboratory of Separation Science for Analytical Chemistry, Dalian Institute of Chemical Physics, Chinese Academy of Sciences) for the generous gift of OM-sensitive probe PI-BactD. We thank Prof. Patrick C. Y. Woo (Department of Microbiology, University of Hong Kong) for the generous gift of clinical *E. coli* strains.

Funding

This work was supported by the

Research Grants Council of Hong Kong, Theme-based Research Scheme T21-705/20-N (A.Y. and T.Z.)

Natural Science Foundation of China 22193062 (T.Z.)

Research Grants Council of Hong Kong, Collaborative Research Fund C7033-20G (A.Y.)

National Key Research and Development Program of China 2022YFA1304500 (X.L.)

Natural Science Foundation of China 22174003 and 21974002 (X.L.)

Author contributions

Conceptualization: Guangming Zhang, Ziqing Deng, Aixin Yan.

Investigation Guangming Zhang, Ziqing Deng, Jiezhang Jiang, Minji Wang, Xiaoyuan Wang, Xiaoyun Liu, Aixin Yan.

Formal Analysis Guangming Zhang, Ziqing Deng, Xiaoyuan Wang, Xiaoyun Liu, Aixin Yan.

Funding Acquisition: Xiaoyun Liu, Aixin Yan.

Resources: Xiaoyuan Wang, Xiaoyun Liu, Aixin Yan.

Supervision: Xiaoyuan Wang, Xiaoyun Liu, Aixin Yan.

Visualization: Guangming Zhang.

Writing – original draft: Guangming Zhang, Aixin Yan.

Writing – review & editing: Guangming Zhang, Xiaoyun Liu, Aixin Yan.

Conflicts of interest

Authors declare that they have no competing interests.

791 **Data and materials availability**

792 All data are available in the main text or the supplementary materials.

References

1. Fang, F.C., Frawley, E.R., Tapscott, T., and Vázquez-Torres, A. (2016). Bacterial stress responses during host infection. Cell host & microbe 20, 133–143.

2. Poole, K. (2012). Bacterial stress responses as determinants of antimicrobial resistance. Journal of Antimicrobial Chemotherapy 67, 2069–2089.

3. Wanner, B.L. (1996). Phosphorus assimilation and control of the phosphate regulon. Escherichia coli and Salmonella: cellular and molecular biology 1, 1357–1381.

4. Fagerbakke, K.M., Heldal, M., and Norland, S. (1996). Content of carbon, nitrogen, oxygen, sulfur and phosphorus in native aquatic and cultured bacteria. Aquatic Microbial Ecology 10, 15–27.

5. Moore, C., Mills, M., Arrigo, K., Berman-Frank, I., Bopp, L., Boyd, P., Galbraith, E., Geider, R., Guieu, C., and Jaccard, S. (2013). Processes and patterns of oceanic nutrient limitation. Nature geoscience 6, 701–710.

6. Van Mooy, B.A., Fredricks, H.F., Pedler, B.E., Dyhrman, S.T., Karl, D.M., Koblížek, M., Lomas, M.W., Mincer, T.J., Moore, L.R., and Moutin, T. (2009). Phytoplankton in the ocean use non-phosphorus lipids in response to phosphorus scarcity. Nature 458, 69–72.

7. Vitousek, P.M., Porder, S., Houlton, B.Z., and Chadwick, O.A. (2010). Terrestrial phosphorus limitation: mechanisms, implications, and nitrogen–phosphorus interactions. Ecological applications 20, 5–15.

8. George, T.S., Hinsinger, P., and Turner, B.L. (2016). Phosphorus in soils and plants – facing phosphorus scarcity. Plant and Soil 401, 1–6. 10.1007/s11104-016-2846-9.

9. Nguyen, T.H., Wang, B.X., Diaz, O.R., Rajendram, M., McKenna, J.A., Butler, D.S.C., Hokamp, K., Hinton, J.C.D., Monack, D.M., and Huang, K.C. (2025). Profiling Salmonella transcriptional dynamics during macrophage infection using a comprehensive reporter library. Nature Microbiology 10, 1006–1023. 10.1038/s41564-025-01953-5.

10. Röder, J., Felgner, P., and Hensel, M. (2021). Single-cell analyses reveal phosphate availability as critical factor for nutrition of Salmonella enterica within mammalian host cells. Cell Microbiol 23, e13374. 10.1111/cmi.13374.

11. Jones, R.A., Shropshire, H., Zhao, C., Murphy, A., Lidbury, I., Wei, T., Scanlan, D.J., and Chen, Y. (2021). Phosphorus stress induces the synthesis of novel glycolipids in *Pseudomonas aeruginosa* that confer protection against a last-resort antibiotic. ISME J 15, 3303–3314. 10.1038/s41396-021-01008-7.

12. Ferreira, R.B.R., Antunes, L.C.M., and Sal-Man, N. (2025). Pathogen-pathogen interactions during co-infections. Isme j 19. 10.1093/ismejo/wraf104.

13. Murdoch, C.C., and Skaar, E.P. (2022). Nutritional immunity: the battle for nutrient metals at the host–pathogen interface. Nature Reviews Microbiology 20, 657–670. 10.1038/s41579-022-00745-6.

14. Rossi, E., Falcone, M., Molin, S., and Johansen, H.K. (2018). High-resolution in situ transcriptomics of Pseudomonas aeruginosa unveils genotype independent

patho-phenotypes in cystic fibrosis lungs. Nature Communications 9, 3459. 10.1038/s41467-018-05944-5.

15. Sun, N., Song, Y., Liu, C., Liu, M., Yu, L., and Wang, F. (2021). Analysis of In Vivo Transcriptome of Intracellular Bacterial Pathogen Salmonella enterica serovar Typhmurium Isolated from Mouse Spleen. Pathogens 10. 10.3390/pathogens10070823.

16. Makino, K., Shinagawa, H., Amemura, M., Kawamoto, T., Yamada, M., and Nakata, A. (1989). Signal transduction in the phosphate regulon of *Escherichia coli* involves phosphotransfer between PhoR and PhoB proteins. Journal of molecular biology 210, 551–559.

17. Hsieh, Y.-J., and Wanner, B.L. (2010). Global regulation by the seven-component Pi signaling system. Current opinion in microbiology 13, 198–203.

18. Crépin, S., Chekabab, S.-M., Le Bihan, G., Bertrand, N., Dozois, C.M., and Harel, J. (2011). The Pho regulon and the pathogenesis of *Escherichia coli*. Veterinary microbiology 153, 82–88.

19. Chekabab, S.M., Harel, J., and Dozois, C.M. (2014). Interplay between genetic regulation of phosphate homeostasis and bacterial virulence. Virulence 5, 786–793.

20. Romano, S., Schulz-Vogt, H.N., González, J.M., and Bondarev, V. (2015). Phosphate limitation induces drastic physiological changes, virulence-related gene expression, and secondary metabolite production in *Pseudovibrio* sp. strain FO-BEG1. Applied and environmental microbiology 81, 3518–3528.

21. Pratt, J.T., Ismail, A.M., and Camilli, A. (2010). PhoB regulates both environmental and virulence gene expression in *Vibrio cholerae*. Molecular microbiology 77, 1595–1605.

22. Jensen, V., Löns, D., Zaoui, C., Bredenbruch, F., Meissner, A., Dieterich, G., Münch, R., and Häussler, S. (2006). RhlR expression in *Pseudomonas aeruginosa* is modulated by the *Pseudomonas* quinolone signal via PhoB-dependent and-independent pathways. Journal of bacteriology 188, 8601–8606.

23. Ferreira, G.M., and Spira, B. (2008). The pst operon of enteropathogenic *Escherichia coli* enhances bacterial adherence to epithelial cells. Microbiology 154, 2025–2036.

24. Romanowski, K., Zaborin, A., Fernandez, H., Poroyko, V., Valuckaite, V., Gerdes, S., Liu, D.C., Zaborina, O.Y., and Alverdy, J.C. (2011). Prevention of siderophore-mediated gut-derived sepsis due to *P. aeruginosa* can be achieved without iron provision by maintaining local phosphate abundance: role of pH. BMC microbiology 11, 1–14.

25. Santos-Beneit, F., and Martin, J.F. (2013). Vancomycin resistance in *Streptomyces coelicolor* is phosphate-dependent but is not mediated by the PhoP regulator. J Glob Antimicrob Resist 1, 109–113. 10.1016/j.jgar.2013.03.003.

26. Dorrer, E., and Teuber, M. (1977). Induction of polymyxin resistance in *Pseudomonas fluorescens* by phosphate limitation. Archives of Microbiology 114, 87–89.

27. Olea-Ozuna, R.J., Campbell, M.J., Quintanilla, S.Y., Nandy, S., Brodbelt, J.S., and

Boll, J.M. (2025). Alternative lipid synthesis in response to phosphate limitation promotes antibiotic tolerance in Gram-negative ESKAPE pathogens. PLoS pathogens 21, e1012933.

28. Lewenza, S., Falsafi, R., Bains, M., Rohs, P., Stupak, J., Sprott, G.D., and Hancock, R.E. (2011). The *olsA* gene mediates the synthesis of an ornithine lipid in *Pseudomonas aeruginosa* during growth under phosphate-limiting conditions, but is not involved in antimicrobial peptide susceptibility. FEMS Microbiol Lett 320, 95–102. 10.1111/j.1574-6968.2011.02295.x.

29. Barbosa, L.C., Goulart, C.L., Avellar, M.M., Bisch, P.M., and von Kruger, W.M.A. (2018). Accumulation of ornithine lipids in Vibrio cholerae under phosphate deprivation is dependent on VC0489 (OlsF) and PhoBR system. Microbiology (Reading) 164, 395–399. 10.1099/mic.0.000607.

30. Geiger, O., Röhrs, V., Weissenmayer, B., Finan, T.M., and Thomas-Oates, J.E. (1999). The regulator gene phoB mediates phosphate stress-controlled synthesis of the membrane lipid diacylglyceryl-N,N,N-trimethylhomoserine in Rhizobium (Sinorhizobium) meliloti. Mol Microbiol 32, 63–73. 10.1046/j.1365-2958.1999.01325.x.

31. Lev, S., Rupasinghe, T., Desmarini, D., Kaufman-Francis, K., Sorrell, T.C., Roessner, U., and Djordjevic, J.T. (2019). The PHO signaling pathway directs lipid remodeling in Cryptococcus neoformans via DGTS synthase to recycle phosphate during phosphate deficiency. PLoS One 14, e0212651. 10.1371/journal.pone.0212651.

32. Geske, T., Vom Dorp, K., Dörmann, P., and Hölzl, G. (2013). Accumulation of glycolipids and other non-phosphorous lipids in Agrobacterium tumefaciens grown under phosphate deprivation. Glycobiology 23, 69–80. 10.1093/glycob/cws124.

33. Gray, R.M., Hunt, D.M., Silva dos Santos, M., Liu, J., Agapova, A., Rodgers, A., Fearns, A., Canseco, J.O., Garza-Garcia, A., MacRae, J.I., et al. (2025). Mycobacterium tuberculosis overcomes phosphate starvation by extensively remodelling its lipidome with phosphorus-free lipids. Nature Communications 16, 11317. 10.1038/s41467-025-66437-w.

34. Vences-Guzman, M.A., Geiger, O., and Sohlenkamp, C. (2012). Ornithine lipids and their structural modifications: from A to E and beyond. FEMS microbiology letters 335, 1–10.

35. Sohlenkamp, C., and Geiger, O. (2016). Bacterial membrane lipids: diversity in structures and pathways. FEMS microbiology reviews 40, 133–159.

36. Kiriukhin, M.Y., Debabov, D.V., Shinabarger, D.L., and Neuhaus, F.C. (2001). Biosynthesis of the glycolipid anchor in lipoteichoic acid of *Staphylococcus aureus* RN4220: role of YpfP, the diglucosyldiacylglycerol synthase. Journal of bacteriology 183, 3506–3514.

37. Groat, R.G., Schultz, J., Zychlinsky, E., Bockman, A., and Matin, A. (1986). Starvation proteins in Escherichia coli: kinetics of synthesis and role in starvation survival. Journal of Bacteriology 168, 486–493.

38. Raetz, C.R., Reynolds, C.M., Trent, M.S., and Bishop, R.E. (2007). Lipid A

921   modification systems in gram-negative bacteria. Annu. Rev. Biochem. 76, 295–329.

39.   Ogasawara, H., Shinohara, S., Yamamoto, K., and Ishihama, A. (2012). Novel regulation targets of the metal-response BasS-BasR two-component system of *Escherichia coli*. Microbiology (Reading) 158, 1482–1492. 10.1099/mic.0.057745-0.

40.   Fernandez-Ciruelos, B., Potmis, T., Solomin, V., and Wells, J.M. (2023). Cross-talk between QseBC and PmrAB two-component systems is crucial for regulation of motility and colistin resistance in Enteropathogenic Escherichia coli. PLoS Pathog 19, e1011345. 10.1371/journal.ppat.1011345.

41.   Winfield, M.D., and Groisman, E.A. (2004). Phenotypic differences between Salmonella and Escherichia coli resulting from the disparate regulation of homologous genes. Proceedings of the National Academy of Sciences 101, 17162–17167.

42.   Wösten, M.M., Kox, L.F., Chamnongpol, S., Soncini, F.C., and Groisman, E.A. (2000). A signal transduction system that responds to extracellular iron. Cell 103, 113–125.

43.   Wang, P., Zhang, G., Xu, Z., Chen, Z., Liu, X., Wang, C., Zheng, C., Wang, J., Zhang, H., and Yan, A. (2022). Whole-cell FRET monitoring of transcription factor activities enables functional annotation of signal transduction systems in living bacteria. J Biol Chem, 102258. 10.1016/j.jbc.2022.102258.

44.   Wang, J., Xie, J., and Schultz, P.G. (2006). A genetically encoded fluorescent amino acid. Journal of the American Chemical Society 128, 8738–8739.

45.   Chen, H.D., and Groisman, E.A. (2013). The biology of the PmrA/PmrB two-component system: the major regulator of lipopolysaccharide modifications. Annual review of microbiology 67, 83–112.

46.   Lee, L.J., Barrett, J.A., and Poole, R.K. (2005). Genome-wide transcriptional response of chemostat-cultured Escherichia coli to zinc. Journal of bacteriology 187, 1124–1134.

47.   Alves, E., Moreirinha, C., Faustino, M.A., Cunha, A., Delgadillo, I., Neves, M.G., and Almeida, A. (2016). Overall biochemical changes in bacteria photosensitized with cationic porphyrins monitored by infrared spectroscopy. Future Medicinal Chemistry 8, 613–628.

48.   Breuer, W., Epsztejn, S., and Cabantchik, Z.I. (1995). Iron Acquired from Transferrin by K562 Cells Is Delivered into a Cytoplasmic Pool of Chelatable Iron (II)(∗). Journal of Biological Chemistry 270, 24209–24215.

49.   Long, S., Miao, L., Li, R., Deng, F., Qiao, Q., Liu, X., Yan, A., and Xu, Z. (2019). Rapid Identification of Bacteria by Membrane-Responsive Aggregation of a Pyrene Derivative. ACS Sens 4, 281–285. 10.1021/acssensors.8b01466.

50.   Huang, J., Li, C., Song, J., Velkov, T., Wang, L., Zhu, Y., and Li, J. (2020). Regulating polymyxin resistance in Gram-negative bacteria: roles of two-component systems PhoPQ and PmrAB. Future microbiology 15, 445–459.

51.   Hofstaedter, C.E., Chandler, C.E., Met, C.M., Gillespie, J.J., Harro, J.M., Goodlett, D.R., Rasko, D.A., and Ernst, R.K. (2024). Divergent Pseudomonas aeruginosa

LpxO enzymes perform site-specific lipid A 2-hydroxylation. Mbio 15, e02823–02823.

52. Pontes, M.H., Sevostyanova, A., and Groisman, E.A. (2015). When too much ATP is bad for protein synthesis. Journal of molecular biology 427, 2586–2594.

53. Bruna, R.E., Kendra, C.G., Groisman, E.A., and Pontes, M.H. (2021). Limitation of phosphate assimilation maintains cytoplasmic magnesium homeostasis. Proc Natl Acad Sci U S A 118. 10.1073/pnas.2021370118.

54. Klein, D.J., Moore, P.B., and Steitz, T.A. (2004). The contribution of metal ions to the structural stability of the large ribosomal subunit. Rna 10, 1366–1379.

55. Pontes, M.H., and Groisman, E.A. (2018). Protein synthesis controls phosphate homeostasis. Genes & development 32, 79–92.

56. Perez, J.C., Shin, D., Zwir, I., Latifi, T., Hadley, T.J., and Groisman, E.A. (2009). Evolution of a bacterial regulon controlling virulence and Mg^{2+} homeostasis. PLoS genetics 5, e1000428.

57. Véscovi, E.G., Soncini, F.C., and Groisman, E.A. (1996). Mg^{2+} as an extracellular signal: environmental regulation of Salmonella virulence. Cell 84, 165–174.

58. Breazeale, S.D., Ribeiro, A.A., and Raetz, C.R. (2002). Oxidative decarboxylation of UDP-glucuronic acid in extracts of polymyxin-resistant Escherichia coli: Origin of lipid a species modified with 4-amino-4-deoxy-l-arabinose. Journal of Biological Chemistry 277, 2886–2896.

59. Breazeale, S.D., Ribeiro, A.A., and Raetz, C.R. (2003). Origin of Lipid A Species Modified with 4-Amino-4-deoxy-l-arabinose in Polymyxin-resistant Mutants of Escherichia coli: an aminotransferase (ArnB) that generates UDP-4-amino-4-deoxy-l-arabinose. Journal of Biological Chemistry 278, 24731–24739.

60. Yan, A., Guan, Z., and Raetz, C.R. (2007). An undecaprenyl phosphate-aminoarabinose flippase required for polymyxin resistance in *Escherichia coli*. J Biol Chem 282, 36077–36089. 10.1074/jbc.M706172200.

61. Breazeale, S.D., Ribeiro, A.A., McClerren, A.L., and Raetz, C.R. (2005). A formyltransferase required for polymyxin resistance in Escherichia coli and the modification of lipid A with 4-amino-4-deoxy-l-arabinose: identification and function of UDP-4-deoxy-4-formamido-l-arabinose. Journal of Biological Chemistry 280, 14154–14167.

62. Gunn, J.S., Lim, K.B., Krueger, J., Kim, K., Guo, L., Hackett, M., and Miller, S.I. (1998). PmrA–PmrB-regulated genes necessary for 4-aminoarabinose lipid A modification and polymyxin resistance. Molecular microbiology 27, 1171–1182.

63. Stevenson, G., Andrianopoulos, K., Hobbs, M., and Reeves, P.R. (1996). Organization of the *Escherichia coli* K-12 gene cluster responsible for production of the extracellular polysaccharide colanic acid. Journal of bacteriology 178, 4885–4893.

64. Baijal, K., Abramchuk, I., Herrera, C.M., Mah, T.-F., Trent, M.S., Lavallée-Adam, M., and Downey, M. (2024). Polyphosphate kinase regulates LPS structure and polymyxin resistance during starvation in E. coli. Plos Biology 22, e3002558.

65. Dahal, P. (2022). Enterobacteriaceae-definition, characteristics, identification.

Microbe Notes.

66.   Bligh, E.G., and Dyer, W.J. (1959). A rapid method of total lipid extraction and
purification. Canadian journal of biochemistry and physiology 37, 911–917.

67.   Sahonero-Canavesi, D.X., López-Lara, I.M., and Geiger, O. (2019). Membrane lipid
degradation and lipid cycles in microbes. Aerobic Utilization of Hydrocarbons, Oils,
and Lipids 10, 978–973.

68.   Xu, Z., Wang, P., Wang, H., Yu, Z.H., Au-Yeung, H.Y., Hirayama, T., Sun, H., and
Yan, A. (2019). Zinc excess increases cellular demand for iron and decreases
tolerance to copper in *Escherichia coli*. J Biol Chem 294, 16978–16991.
10.1074/jbc.RA119.010023.

69.   Zhao, D., Feng, X., Zhu, X., Wu, T., Zhang, X., and Bi, C. (2017). CRISPR/Cas9-
assisted gRNA-free one-step genome editing with no sequence limitations and
improved targeting efficiency. Sci Rep 7, 16624. 10.1038/s41598-017-16998-8.

70.   Liu, Y., Zhang, Q., Hu, M., Yu, K., Fu, J., Zhou, F., and Liu, X. (2015). Proteomic
analyses of intracellular *Salmonella enterica* serovar Typhimurium reveal extensive
bacterial adaptations to infected host epithelial cells. Infection and immunity 83,
2897–2906.

71.   Groisman, E.A., Kayser, J., and Soncini, F.C. (1997). Regulation of polymyxin
resistance and adaptation to low-Mg^{2+} environments. Journal of bacteriology 179,
7040–7045.

1027

Figure 1. Pi depletion induces polymyxin resistance in diverse bacterial species.

(**A**) Time-kill curves of *Escherichia coli* MG1655 (left panel) and *Salmonella*. Typhimurium

ATCC14028 (right panel) exposed to polymyxin B (PMXB) (4 µg/ml for *E. coli* strains, 20 µg/ml for

S. Typhimurium) following acute Pi-starvation treatment (Pi-, 0 mM Pi) compared to the group

without treatment (Pi+, 1.32 mM Pi).

(**B**) Survival rates for various *Enterobacteriaceae* strains (*E. coli* MG1655, O157, UTI-1, UTI-3,

UTI-4, UTI-5, *S.* Typhimurium ATCC14028) and *Pseudomonas aeruginosa* PAO1 exposed 1-hour

to polymyxin B (4 µg/ml for *E. coli* strains, 20 µg/ml for *S.* Typhimurium and 8 µg/ml for *P.*

aeruginosa). For all tested strains, phosphate depletion (Pi-, purple bars) significantly increases

polymyxin B resistance compared to phosphate-replete (Pi+, teal bars) conditions.

Error bars represent the standard deviation of at least three independent experiments. Statistical

1040   significance was determined using unpaired two-tailed Student's t-test. Significance levels are

1041   indicated as follows: $*p < 0.05$; $**p < 0.01$; $***p < 0.001$.

1042

Figure 2. Pi depletion induces the expression of *ugd-arn* system to modify lipid A with L-

Ara-4N, conferring stress-induced polymyxin resistance.

(**A**) Volcano plot depicting differentially expressed proteins in cells following acute Pi-starvation

treatment (Pi-, 0 mM) compared to the group without treatment (Pi+, 1.32 mM). Red (right) and

blue (left) dots represent significantly up-regulated and down-regulated proteins, respectively. The

horizontal dashed line depicts a p-value cutoff (0.05), and vertical dashed lines depict 1/2 and 2-

fold ratio cutoffs. Selected proteins are labeled.

(**B**) Genetic organization of the *arn* operon and *ugd* gene in MG1655. Ugd and ArnABCDEFT

catalyze the biosynthesis of 4-amino-4-deoxy-L-arabinose (L-Ara4N) and its modification of lipid

A.

(**C**) mRNA levels of *arnB* and *ugd* in cells following acute Pi-starvation treatment (Pi-, 0 mM)

compared to the group without treatment (Pi+, 1.32 mM).

(**D**) Production of ArnB-3×FLAG in cells following acute Pi-starvation treatment (Pi-, 0 mM)

compared to the group without treatment (Pi+, 1.32 mM). The protein band corresponding to ArnB-

3×FLAG is marked.

(**E**) β-galactosidase activities of P$_{arn}$-*lacZ* and P$_{ugd}$-*lacZ* in MG1655 Δ*lacZ* cells following acute Pi-

starvation treatment (Pi-, 0 mM) compared to the group without treatment (Pi+, 1.32 mM).

(**F**) ESI-MS spectra of lipid A molecules extracted from *E. coli* MG1655 cells following acute Pi-

starvation treatment (Pi-, 0 mM) compared to the group without treatment (Pi+, 1.32 mM). Peaks

corresponding to native lipid A and L-Ara-4N modified lipid A were shown.

(**G**) Survival rates for *E. coli* MG1655 parent and isogenic Δ*arnT*, Δ*arnB* and Δ*ugd* mutants

exposed 1-hour to polymyxin B (4 µg/ml) following acute Pi-starvation treatment (Pi-, 0 mM)

compared to the group without treatment (Pi+, 1.32 mM).

Error bars represent the standard deviation of at least three independent experiments. Statistical

significance was determined using two-tailed Student's t-test for **C** and **E**, and one-way ANOVA

followed by Dunnett's multiple comparisons test for **G**. Significance levels are indicated as follows:

*$p < 0.05$; **$p < 0.01$; ***$p < 0.001$.

Figure 3. The PmrAB two-component system is activated during Pi depletion and is

responsible for the up-regulation of *arn* genes under this condition.

(**A**) Survival rates for *E. coli* MG1655 parent and isogenic Δ*pmrB*, Δ*pmrA,* Δ*phoB,* and Δ*phoR*

mutants exposed 1-hour to polymyxin B (4 μg/ml) following acute Pi-starvation treatment (Pi-, 0

mM) compared to the group without treatment (Pi+, 1.32 mM).

(**B**) β-galactosidase activities of P$_{arn}$-*lacZ* in MG1655 Δ*lacZ* parent and isogenic Δ*pmrB*, Δ*pmrA*,

Δ*phoB*, and Δ*phoR* mutants following acute Pi-starvation treatment (Pi-, 0 mM) compared to the

group without treatment (Pi+, 1.32 mM).

(**C**) Phosphorylation of PmrA-3×FLAG detected by Phos-Tag SDS–PAGE following

immunoblotting with anti-FLAG antibody in cells following acute Pi-starvation treatment (Pi-, 0 mM)

compared to the group without treatment (Pi+, 1.32 mM). Protein bands representing PmrA-

3×FLAG and PmrA-3×FLAG-P are indicated.

(**D**) Schematic diagram of the whole-cell FRET measurement of PmrA-promoter binding assay. A

bright fluorescent unnatural amino acid donor CouA (red spheres) is incorporated into G137 of

PmrA in DNA-binding domain. SYTO9 (green spheres) is employed to stain chromosome DNA in

living *E. coli* MG1655.

(**E**) Quantification of the FRET effect between the CouA-SYTO9 pair in *E. coli* cells following acute

Pi-starvation treatment (Pi-, 0 mM Pi, 10 μM Fe) or iron excess treatment (Pi+, 1.32 mM Pi, 500

μM FeSO$_4$) compared to the group without treatment (Pi+, 1.32 mM Pi, 10 μM FeSO$_4$).

Error bars represent the standard deviation of at least three independent experiments. Statistical

significance was determined using one-way ANOVA followed by Dunnett's multiple comparisons

test for **A**, **B**, and **E**. Significance levels are indicated as follows: *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$.

1093

1094

Figure 4. Activation of PmrAB during Pi depletion is mediated by the secondary stress

signal iron (Fe^{3+}) induced under this condition

(**A**) Cellular metal contents determined by ICP-MS in *E. coli* MG1655 cells following acute Pi-

starvation treatment (Pi-, 0 mM) compared to the group without treatment (Pi+, 1.32 mM).

(**B**) Iron contents in the spent medium of *E. coli* MG1655 at the indicated time points during Pi

depletion treatment (Pi-, 0 mM).

(**C**) Schematic diagram of the sensor kinase PmrB. Two important residues for sensing Fe^{3+}, E36A

and E39A, are shown.

(**D**) β-galactosidase activities of P$_{arn}$-lacZ in MG1655 ΔlacZ parent and isogenic pmrB^{E36A} and

pmrB^{E36A} mutants following acute Pi-starvation treatment (Pi-, 0 mM) compared to the group

without treatment (Pi+, 1.32 mM).

(**E**) β-galactosidase activities of P$_{arn}$-lacZ in MG1655 ΔlacZ cells following Pi depletion treatment

(Pi-, 0 mM Pi, 10 µM Fe) or Fe, Pi depletion treatment (Pi- Fe-, 0 mM Pi, 0 µM Fe) compared to

the group without treatment (Pi+, 1.32 mM Pi, 10 µM $FeSO_4$).

(**F**) β-galactosidase activities of P$_{arn}$-lacZ in MG1655 ΔlacZ grown under Pi sufficient MOPS

medium in the presence of different concentrations of $FeSO_4$ and that subjected to Pi depletion

conditions in the presence of different concentrations of $FeSO_4$.

Error bars represent the standard deviation of at least three independent experiments. Statistical

significance was determined using one-way ANOVA followed by Dunnett's multiple comparisons

test for **B**, and two-tailed Student's t-test for **C**, **D**, and **E**. Significance levels are indicated as

follows: *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$.

1116

Figure 5. Response to Pi depletion induced iron mobilization to cell surfaces

1117

(**A**) Iron (Fe) and magnesium (Mg) contents in *E. coli* MG1655 cells following Pi depletion

1118

treatment (Pi-, 0 mM Pi) at the indicated time points.

1119

(**B**) β-galactosidase activities of P_{arn}-*lacZ* in MG1655 Δ*lacZ* cells and Fe contents in *E. coli*

1120

MG1655 cells following Pi depletion treatment at the indicated time points.

1121

(**C**) Fe contents in *E. coli* MG1655 parent and isogenic Δ*pmrA* mutant cells following Pi depletion

1122

treatment (Pi-, 0 mM Pi) at the indicated time points.

1123

(**D**) Fe contents in *E. coli* MG1655 cells with Pi depletion treatment (Pi-, 0 mM) and without Pi

1124

depletion treatment (Pi+, 1.32 mM) following washing by HEPES or DFOM-supplemented (50μM)

1125

HEPES buffer.

1126

(**E**) Reduction of cellular Fe content following washing with DFOM-supplemented (50μM) HEPES

1127

1128   buffer.

1129   Error bars represent the standard deviation of at least three independent experiments. Statistical

1130   significance was determined using two-tailed Student's t-test for **C** and **E**, and two-way ANOVA

1131   followed by Sidak's multiple comparisons test for **D**. Significance levels are indicated as follows:

1132   *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$.

1133

Figure 6. Iron mobilization is driven by Mg²⁺ dissociation during Pi depletion

1134

(**A**) Fe contents in *E. coli* MG1655 cells following acute Pi-starvation treatment (Pi-, 0 mM Pi, 0.53

1135

mM Mg) or Mg excess Pi depletion treatment (Pi-, 0 mM Pi, 10 mM Mg) compared to the group

1136

without treatment (Pi+, 1.32 mM, 0.53 mM Mg).

1137

(**B**) β-galactosidase activities of P_{arn}-*lacZ* in MG1655 Δ*lacZ* cells following acute Pi-starvation

1138

treatment (Pi-, 0 mM Pi, 0.53 mM Mg) or Mg excess Pi depletion treatment (Pi-, 0 mM Pi, 10 mM

1139

Mg) compared to the group without treatment (Pi+, 1.32 mM, 0.53-10 mM Mg).

1140

(**C**) Mg content in the spent medium during Pi depletion treatment at the indicated time points.

1141

(**D**) 2D graph plotted from output signals of two channels of the PI-BactD sensor: fluorescence

1142

increase (ΔS/S0) and fluorescence radiometric changes (I482/I375) in each of the treatments

1143

indicated. Data are presented as mean ± s.d. of three biological replicates for **A**, **B**, and **C**, and

six biological replicates for **D**.

Error bars represent the standard deviation of at least three independent experiments. Statistical

significance was determined using two-tailed Student's t-test for **a**, and one-way ANOVA followed

by Dunnett's multiple comparisons test for **B** and **C**. Significance levels are indicated as follows:

*$p < 0.05$; **$p < 0.01$; ***$p < 0.001$.

1150

Figure 7. A Mg-Fe-PmrAB regulatory circuit underpins the Pi-depletion induced polymyxin

resistance through activating the lipid A L-Ara4N modification system

In cell grown under Pi sufficient condition, ribosome and ATP serve as the major intracellular Mg^{2+}

and Pi reservoir. Mg^{2+} neutralizes negative charge of lipid A and stabilizes outer membrane (OM).

The PmrAB two-component system (TCS) is inactive, and the cell is sensitive to polymyxin. During

Pi stress, ATP biosynthesis and ribosome biogenesis are halted due to Pi depletion, disrupting the

balance between Pi and Mg^{2+}. Mg^{2+} disassociates from the cell which destabilizes the OM and

derives compensatory Fe^{3+} mobilization to cell surface. Mobilized Fe^{3+} activates the PmrAB TCS

which in turn upregulates the *arn* operon. Arn and Ugd proteins catalyze the biosynthesis of L-

Arn4N and modify lipid A with L-Arn4N to stabilize OM under stressed condition and prevent further

Fe mobilization. *E. coli* cells with L-Arn4N modified lipid A become resistant to polymyxin.

1162

1163

Figure 8. Disrupting the Mg-Fe-PmrAB signaling circuit abolished Pi-depletion induced

polymyxin resistance in *Enterobacteriaceae*

(**A**) Time-kill curves of *Escherichia coli* MG1655 (left panel) and *Salmonella*. Typhimurium

ATCC14028 (right panel) exposed to polymyxin B (PMXB) (4 µg/ml for *E. coli* strains, 20 µg/ml for

S. Typhimurium) following acute Pi-starvation treatment (Pi-, 0 mM Pi), Pi depletion treatment with

excess Mg (Pi-, 0 mM Pi, 10mM $MgCl_2$) or Pi depletion treatment with iron chelator (Pi-, 0 mM Pi,

10 µM DFOM) compared to the group without treatment (Pi+, 1.32 mM Pi).

(**B**) Distinct stress adaptation strategies in *E. coli* and *P. aeruginosa* under Pi depletion conditions.

Genes and regulatory systems activated in the two species are illustrated. Inactive regulatory

systems are depicted in faded effects. The regulatory mechanisms for *P. aeruginosa* are based

on findings from Jones et. al.[11]

(**C**) A phylogenetic tree of 250 PmrB homologous. A predominant clade is indicated by a blue circle,

comprising sequences from bacteria such as *Escherichia, Shigella, Citrobacter, Salmonella,*

Klebsiella, Yersinia, and *Enterobacter* species. A green circle marks several clades distinct from

the predominant one. The domain architecture and AlphaFold3-predicted structure of EcoPmrB

and PaePmrB are shown.

Error bars represent the standard deviation of at least three independent experiments. Statistical

significance was determined using one-way ANOVA followed by Dunnett's multiple comparisons

test. Significance levels are indicated as follows: *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$.

 Supporting information

1184

Figure S1. Workflow of acute Pi depletion treatment and following proteomic sample

preparation and polymyxin B challenging assay.

Bacteria were cultured to an OD_{600} of 0.4 in standard MOPS synthetic medium containing 1.32

mM inorganic phosphate (Pi), representing Pi-sufficient conditions. For Pi-depletion conditions,

bacteria were first grown to OD_{600} = 0.4 in the same medium, then transferred to Pi-depleted

MOPS medium (0 mM Pi). For comparative proteomic analysis, the collected bacterial cells were

subjected to sample preparation and proteomic analysis. For the polymyxin challenging assay, the

collected bacterial cells were resuspended in standard MOPS medium and treated with the

designated concentration of polymyxin B. Cell viability was assessed at the beginning and end of

treatment by serial dilution and plating.

Figure S2. Killing of *E. coli* MG1655 with different concentration of polymyxin B.

Survival rates for *E. coli* MG1655 cells exposed 1-hour to polymyxin B (1-16 μg/ml) following acute

Pi-starvation treatment (Pi-, 0 mM) compared to the group without treatment (Pi+, 1.32 mM).

Error bars represent the standard deviation of at least three independent experiments. Statistical

significance was determined using unpaired two-tailed Student's t-test. Significance levels are

indicated as follows: *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$.

1202

1203

Figure S3. Quantitative proteomic analysis of *E. coli* MG1655 in response to acute Pi

depletion.

Pie chart of total and differentially expressed proteins detected by LC-MS/MS analysis in *E. coli*

MG1655 cells subjected to Pi depletion for 3 hours, relative to cells under Pi sufficient conditions.

Red and blue colors represent significantly upregulated and downregulated proteins, respectively.

Clusters of Orthologous Groups (COG) functional categories of differentially expressed proteins

in *E. coli* MG1655 in response to Pi depletion. Red and blue colors represent significantly

upregulated and downregulated proteins, respectively.

1212

Figure S4. The ratio of L-Ara-4N modified Lipid A molecule increases in *E. coli* O157, *E. coli*

UTI-3, *S.* Typhimurium ATCC14028 and *P. aeruginosa* PAO1 cells subjected to Pi depletion.

(**A**) ESI-MS spectra of lipid A molecules extracted from *E. coli* O157 cells following acute Pi-

starvation treatment (Pi-, 0 mM) (right) compared to the group without treatment (Pi+, 1.32 mM)

(left). Peaks correspond to native lipid A lost a HPO_3 (1718), native lipid A (1798), L-Ara-4N

modified lipid A lost a HPO_3 (1848) and L-Ara-4N modified lipid A (1928) are shown. (**B**) ESI-MS

spectra of lipid A molecules extracted from *E. coli* UTI-3 cells following acute Pi-starvation

treatment (Pi-, 0 mM) (right) compared to the group without treatment (Pi+, 1.32 mM) (left). Peaks

correspond to native lipid A lost a HPO_3 (1718), native lipid A (1798), L-Ara-4N modified lipid A lost

a HPO_3 (1848) and L-Ara-4N modified lipid A (1928) are shown. (**C**) ESI-MS spectra of lipid A

molecules extracted from *S.* Typhimurium ATCC14028 cells following acute Pi-starvation

treatment (Pi-, 0 mM) (right) compared to the group without treatment (Pi+, 1.32 mM) (left). Peaks

correspond to native lipid A lost a HPO_3 (1718), native lipid A (1798), L-Ara-4N modified lipid A lost

a HPO_3 (1848), L-Ara-4N and -OH modified lipid A lost a HPO_3 (1863) and L-Ara-4N modified lipid

A (1928) are shown. (**D**) ESI-MS spectra of lipid A molecules extracted from PAO1 cells following

acute Pi-starvation treatment (Pi-, 0 mM) (right) compared to the group without treatment (Pi+,

1.32 mM) (left). Peaks correspond to hydroxylated lipid A lost a HPO_3 (1366), di-hydroxylated lipid

A lost a HPO_3 (1382) are shown.

1231

Figure S5. P*arn* and P*ugd* promoter activities in *E. coli* MG1655 during Pi depletion

supplemented with different concentration of Pi.

(A) β-galactosidase activities of P*arn*-*lacZ* in MG1655 Δ*lacZ* cells following Pi-starvation treatment

with different concentration of Pi (Pi-, 0-0.4 mM) compared to the group without treatment (Pi+,

1.32 mM).

(B) β-galactosidase activities of P*ugd*-*lacZ* in MG1655 Δ*lacZ* cells following Pi-starvation treatment

with different concentration of Pi (Pi-, 0-0.4 mM) compared to the group without treatment (Pi+,

1.32 mM).

1240   Error bars represent the standard deviation of at least three independent experiments. Statistical

1241   significance was determined using one-way ANOVA followed by Dunnett's multiple comparisons

1242   test. Significance levels are indicated as follows: *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$.

1243

Figure S6. Up-regulation of *arn* genes during Pi depletion is independent of PhoBR and

PhoPQ TCS in *E. coli* MG1655.

β-galactosidase activities of P_{ugd}-*lacZ* in MG1655 Δ*lacZ* parent and isogenic Δ*pmrB,* Δ*pmrA,*

Δ*phoB* and Δ*phoR* mutants following acute Pi-starvation treatment (Pi-, 0 mM) compared to the

group without treatment (Pi+, 1.32 mM).

Error bars represent the standard deviation of at least three independent experiments. Statistical

significance was determined using one-way ANOVA followed by Dunnett's multiple comparisons

test. Significance levels are indicated as follows: *p < 0.05; **p < 0.01; ***p < 0.001.

1252

1253

Figure S7. Up-regulation of *ugd* gene during Pi depletion is independent of PmrAB and

PhoPQ TCS, but partially dependent on PhoBR TCS in *E. coli* MG1655.

β-galactosidase activities of P_{arn}-*lacZ* in MG1655 Δ*lacZ* parent and isogenic Δ*phoR*, Δ*phoB*,

Δ*phoP* and Δ*phoQ* mutants following acute Pi-starvation treatment (Pi-, 0 mM) compared to the

group without treatment (Pi+, 1.32 mM).

Error bars represent the standard deviation of at least three independent experiments. Statistical

significance was determined using one-way ANOVA followed by Dunnett's multiple comparisons

test. Significance levels are indicated as follows: *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$.

1262

1263

Figure S8. Pi depletion induced polymyxin tolerance is not affected in *eptA* deletion strain.

Survival of *E. coli* MG1655 parent and isogenic ΔeptA mutant exposed 1-hour to polymyxin B (4

µg/ml) following acute Pi-starvation treatment (Pi-, 0 mM) compared to the group without treatment

(Pi+, 1.32 mM).

Error bars represent the standard deviation of at least three independent experiments. Statistical

significance was determined using two-tailed Student's t-test. Significance levels are indicated as

follows: *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$.

1271

Figure S9. pH of spent medium of _E. coli_ MG1655 after 5hrs Pi sufficient growth and after

1272

3hrs Pi depletion treatment.

1273

pH of the medium before bacteria inoculating and pH of the spent medium after 5hrs Pi sufficient

1274

growth or 3hrs Pi depletion treatment were measured.

1275

Error bars represent the standard deviation of at least three independent experiments.

1276

1277

1278

Figure S10. E36A and E39A mutation abolish the Fe^{3+} signaling capacity of PmrA.

β-galactosidase activities of P_{arn}-lacZ in MG1655 ΔlacZ parent and isogenic $pmrB^{E36A}$, $pmrB^{E39A}$

mutants growth under MOPS minimal medium supplemented with different concentration of

$FeSO_4$.

Error bars represent the standard deviation of at least three independent experiments. Statistical

significance was determined using two-way ANOVA followed by Dunnett's multiple comparisons

test. Significance levels are indicated as follows: *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$.

Figure S11. Deletion of canonical Fe^{3+} up-take genes does not affect the activation of PmrAB system and increase of Fe mobilization to _E. coli_ cell during Pi depletion.

(**A**) β-galactosidase activities of P$_{am}$-_lacZ_ in MG1655 Δ_lacZ_ parent and isogenic Δ_fecA_, Δ_fepA_, Δ_entE_, Δ_fiu_, Δ_cirA_, and Δ5Fe (Δ_fecA_Δ_fepA_Δ_entE_Δ_fiu_Δ_cirA_) mutants following acute Pi-starvation treatment (Pi-, 0 mM) compared to the group without treatment (Pi+, 1.32 mM).

(**B**) Fe contents in _E. coli_ MG1655 parent and isogenic Δ_fecA_ Δ_fepA_ Δ_entE_ Δ_fiu_ Δ_cirA_ mutant following acute Pi-starvation treatment (Pi-, 0 mM) compared to the group without treatment (Pi+, 1.32 mM).

Error bars represent the standard deviation of at least three independent experiments. Statistical significance was determined using one-way ANOVA followed by Dunnett's multiple comparisons test for **A** and two-way ANOVA followed by Sidak's multiple comparisons test for **B**. Significance levels are indicated as follows: *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$.

1299

Figure S12. Decrease of cellular magnesium during Pi depletion is independent of Fe

availability in medium.

Mg contents in *E. coli* MG1655 cell following Pi depletion treatment (0 Pi, 10 μM $FeSO_4$) or Fe-

free Pi depletion treatment (0 Pi, 0 $FeSO_4$) at the indicated time points.

Error bars represent the standard deviation of at least three independent experiments.

1305

Figure S13. Magnesium disassociates from *E. coli* cells when subjected to Mg-free Pi-

depleted medium.

Mg content in the spent medium during Mg-free Pi depletion treatment (0 Pi, 0 $MgCl_2$) at the

indicated time points.

Error bars represent the standard deviation of at least three independent experiments. Statistical

significance was determined using one-way ANOVA followed by Dunnett's multiple comparisons

test. Significance levels are indicated as follows: *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$.

1313

Figure S14. Cellular metal content in *P. aeruginosa* PAO1 cells with Pi depletion treatment

and without Pi depletion treatment.

Cellular metal contents determined by ICP-MS in *P. aeruginosa* PAO1 cells following acute Pi-

starvation treatment (Pi-, 0 mM) compared to the group without treatment (Pi+, 1.32 mM).

Error bars represent the standard deviation of at least three independent experiments. Statistical

significance was determined using two-tailed Student's t-test. Significance levels are indicated as

follows: *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$.

Figure S15. Disruption of Mg and Fe signaling showed no effect on Pi depletion induced polymyxin resistance in *P. aeruginosa*.

(A) Time-kill curves of *P. aeruginosa* PAO1 exposed to PMXB (8 µg/ml) following acute Pi-starvation treatment (Pi-, 0 mM Pi), Pi depletion treatment with excess Mg (Pi-, 0 mM Pi, 10mM MgCl$_2$) or Pi depletion treatment with iron chelator (Pi-, 0 mM Pi, 10 µM DFOM) compared to the group without treatment (Pi+, 1.32 mM Pi).

(B) Time-kill curves of *P. aeruginosa* PAO1 exposed to PMXB (8 µg/ml) following acute Pi-starvation treatment (Pi-, 0 mM Pi), Pi depletion treatment with Fe^{2+} chelator (Pi-, 0 mM Pi, 30 µM DIP) or Pi depletion treatment with both Fe^{2+} and Fe^{3+}chelator (Pi-, 0 mM Pi, 30 µM DIP, 10 µM

1331   DFOM) compared to the group without treatment (Pi+, 1.32 mM Pi).

1332   Error bars represent the standard deviation of at least three independent experiments. Statistical

1333   significance was determined using one-way ANOVA followed by Dunnett's multiple comparisons

1334   test. Significance levels are indicated as follows: $*p < 0.05$; $**p < 0.01$; $***p < 0.001$.

1335

Figure S16. Cellular ATP level in *E. coli* MG1655 cells continuously decrease during Pi

depletion.

Cellular ATP levels in *E. coli* MG1655 cells following Pi depletion treatment at the specified time

points. Error bars represent the standard deviation of at least three independent experiments.

1340

1341

Figure S17. Lack of upregulation of P*phoQ* and P*mgtA* promoter activities indicates that

the PhoPQ TCS is not activated during Pi depletion in *E. coli* MG1655.

β-galactosidase activities of P$_{phoQ}$-*lacZ* and P$_{mgtA}$-*lacZ* in MG1655 Δ*lacZ* parent following acute Pi-

starvation treatment (Pi-, 0 mM) compared to the group without treatment (Pi+, 1.32 mM).

Error bars represent the standard deviation of at least three independent experiments. Statistical

significance was determined using two-tailed Student's t-test. Significance levels are indicated as

follows: *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$.

1349

1350

Figure S18. Up-regulation of both P*arn* and P*ugd* promoter activities during Pi depletion

are not affected in *ppk* deletion strain.

(A) β-galactosidase activities of P$_{arn}$-*lacZ* in MG1655 Δ*lacZ* parent and isogenic Δ*ppk* mutant

following acute Pi-starvation treatment (Pi-, 0 mM) compared to the group without treatment (Pi+,

1.32 mM).

(B) β-galactosidase activities of P$_{ugd}$-*lacZ* in MG1655 Δ*lacZ* parent and isogenic Δ*ppk* mutant

following acute Pi-starvation treatment (Pi-, 0 mM) compared to the group without treatment (Pi+,

1.32 mM).

Error bars represent the standard deviation of at least three independent experiments. Statistical

significance was determined using two-way ANOVA followed by Tukey's multiple comparisons test.

Significance levels are indicated as follows: *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$.

1362

Figure S19. Pi depletion, amino acid depletion, and combined Pi–amino acid stress upregulate P*arn* promoter activity, while *ppk* deletion only attenuates upregulation under combined stress.

(A) β-galactosidase activities of P*arn*-*lacZ* in MG1655 Δ*lacZ* parent and isogenic Δ*ppk* mutant following acute nutrients downshift (MOPS 0.1 mM Pi) compared to the group without treatment (LB). Error bars represent the standard deviation of at least three independent experiments. Statistical significance was determined using two-way ANOVA followed by Tukey's multiple comparisons test.

(B) β-galactosidase activities of P*arn*-*lacZ* in MG1655 Δ*lacZ* parent and isogenic Δ*ppk* mutant following Pi depletion treatment (MOPS 0 Pi, 0.1% CAA), amino acid depletion treatment (MOPS 1.32 mM Pi, 0 CAA) and combined Pi–amino acid stress treatment (MOPS 0 Pi, 0 CAA) compared to the group without treatment (MOPS 1.32 mM Pi, 0.1% CAA).

Error bars represent the standard deviation of at least three independent experiments. Statistical significance was determined using two-way ANOVA followed by Tukey's multiple comparisons test.

Significance levels are indicated as follows: *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$.

1378   **Table S1.** Minimum inhibitory concentrations (MICs) of strains in MOPS minimal medium

Strain	PMXB (ug/ml)
MG1655	0.5
O157	1
UTI-1	0.5
UTI-3	1
UTI-4	1
UTI-5	1
PAO1	0.5
ATCC 14028	2

1379

Table S2. Proteins identified to be different expressed in *E. coli* MG1655 growing under Pi

sufficient and depletion conditions

prot_desc	id	log2FoldChange	padj
argF	sp|P06960|OTC2_ECOLI	-Inf	3.42E-18
gatZ	sp|P0C8J8|GATZ_ECOLI	-Inf	3.77E-09
fis	sp|P0A6R3|FIS_ECOLI	-Inf	2.78E-08
secY	sp|P0AGA2|SECY_ECOLI	-Inf	0.000159211
kgtP	sp|P0AEX3|KGTP_ECOLI	-Inf	0.001647533
gatC	sp|P69831|PTKC_ECOLI	-Inf	0.002435201
cysC	sp|P0A6J1|CYSC_ECOLI	-Inf	0.003341414
znuC	sp|P0A9X1|ZNUC_ECOLI	-Inf	0.025826417
rffH	sp|P61887|RMLA2_ECOLI	-Inf	0.036760697
cysD	sp|P21156|CYSD_ECOLI	-5.525585778	1.38E-11
mnmG	sp|P0A6U3|MNMG_ECOLI	-4.77187114	0.003253389
argA	sp|P0A6C5|ARGA_ECOLI	-4.649638457	2.18E-06
yhjE	sp|P37643|YHJE_ECOLI	-4.500028599	0.005992816
yegQ	sp|P76403|YEGQ_ECOLI	-4.482584177	0.015352456
cysN	sp|P23845|CYSN_ECOLI	-3.927062955	7.22E-14
pncB	sp|P18133|PNCB_ECOLI	-3.639270989	6.73E-05
fbp	sp|P0A993|F16PA_ECOLI	-3.600157407	0.023396164
moaA	sp|P30745|MOAA_ECOLI	-3.529794204	0.018342621

rpmC	sp|P0A7M6|RL29_ECOLI	-3.142935813	1.68E-07
ilvN	sp|P0ADF8|ILVN_ECOLI	-3.08821322	0.03765939
aroP	sp|P15993|AROP_ECOLI	-2.833642328	0.021648588
deaD	sp|P0A9P6|DEAD_ECOLI	-2.695068638	0.006647318
ilvB	sp|P08142|ILVB_ECOLI	-2.690257592	1.41E-07
asnA	sp|P00963|ASNA_ECOLI	-2.66746299	0.015909254
ptsG	sp|P69786|PTGCB_ECOLI	-2.510708646	0.000188954
rpmD	sp|P0AG51|RL30_ECOLI	-2.495454148	8.47E-09
ykgM	sp|P0A7N1|RL31B_ECOLI	-2.483011177	0.00644327
gatB	sp|P37188|PTKB_ECOLI	-2.418194429	0.035274735
cs	sp|P0A972|CSPE_ECOLI	-2.352007644	4.54E-08
metA	sp|P07623|META_ECOLI	-2.317452293	0.000139049
der	sp|P0A6P5|DER_ECOLI	-2.267771633	0.0102137
potA	sp|P69874|POTA_ECOLI	-2.183307337	0.03275735
leuA	sp|P09151|LEU1_ECOLI	-1.998105409	1.07E-08
glf	sp|P37747|GLF_ECOLI	-1.97691323	0.029751
tr	sp|P00895|TRPE_ECOLI	-1.943830992	0.005678046
miaB	sp|P0AEI1|MIAB_ECOLI	-1.92205685	0.004244044
pntB	sp|P0AB67|PNTB_ECOLI	-1.852085802	9.85E-06
purF	sp|P0AG16|PUR1_ECOLI	-1.826541229	4.76E-05
lepA	sp|P60785|LEPA_ECOLI	-1.791966706	0.01657399

gatY	sp|P0C8J6|GATY_ECOLI	-1.776699008	0.000182727
purT	sp|P33221|PURT_ECOLI	-1.764435997	0.000643699
prfC	sp|P0A7I4|RF3_ECOLI	-1.76312872	0.012506321
ybiC	sp|P30178|YBIC_ECOLI	-1.727397747	0.01941163
rpsH	sp|P0A7W7|RS8_ECOLI	-1.723379737	2.80E-10
purM	sp|P08178|PUR5_ECOLI	-1.678278829	0.003764
cyoA	sp|P0ABJ1|CYOA_ECOLI	-1.663437618	0.006838543
metL	sp|P00562|AK2H_ECOLI	-1.661017127	0.00181135
leuC	sp|P0A6A6|LEUC_ECOLI	-1.653771358	6.45E-06
pntA	sp|P07001|PNTA_ECOLI	-1.595004247	7.61E-08
trxB	sp|P0A9P4|TRXB_ECOLI	-1.571960722	0.002958769
metH	sp|P13009|METH_ECOLI	-1.565084962	0.000869601
metC	sp|P06721|METC_ECOLI	-1.510217214	0.025144896
rpmF	sp|P0A7N4|RL32_ECOLI	-1.498744569	0.000469522
gyrA	sp|P0AES4|GYRA_ECOLI	-1.448436432	7.72E-05
rpsR	sp|P0A7T7|RS18_ECOLI	-1.418043759	3.03E-05
rpmG	sp|P0A7N9|RL33_ECOLI	-1.381024188	0.006747605
purH	sp|P15639|PUR9_ECOLI	-1.376402854	0.000114658
argI	sp|P04391|OTC1_ECOLI	-1.37321011	0.000772873
rnb	sp|P30850|RNB_ECOLI	-1.364231063	0.003951903
rplT	sp|P0A7L3|RL20_ECOLI	-1.361561823	7.98781E-05

cysS	sp|P21888|SYC_ECOLI	-1.352510933	0.018153582
metK	sp|P0A817|METK_ECOLI	-1.28929395	0.000291382
sdhA	sp|P0AC41|SDHA_ECOLI	-1.284366661	8.12E-05
ydjN	sp|P77529|YDJN_ECOLI	-1.282130778	0.012801159
pyrD	sp|P0A7E1|PYRD_ECOLI	-1.24642368	0.017272295
rpsM	sp|P0A7S9|RS13_ECOLI	-1.231500265	4.82E-05
rpsC	sp|P0A7V3|RS3_ECOLI	-1.228519333	5.14E-06
rplU	sp|P0AG48|RL21_ECOLI	-1.200353979	0.000935979
pliG	sp|P76002|PLIG_ECOLI	-1.192616084	0.023769411
rplB	sp|P60422|RL2_ECOLI	-1.160209361	1.29E-05
rpsA	sp|P0AG67|RS1_ECOLI	-1.154439475	2.19E-06
rplS	sp|P0A7K6|RL19_ECOLI	-1.140490911	0.042048579
rpsT	sp|P0A7U7|RS20_ECOLI	-1.136468494	0.000373058
thrA	sp|P00561|AK1H_ECOLI	-1.131400346	7.55E-06
thrS	sp|P0A8M3|SYT_ECOLI	-1.130800143	0.017957548
rplR	sp|P0C018|RL18_ECOLI	-1.127973054	0.000234204
rplX	sp|P60624|RL24_ECOLI	-1.108686244	0.000297915
aceB	sp|P08997|MASY_ECOLI	-1.103226647	7.72E-05
aroA	sp|P0A6D3|AROA_ECOLI	-1.074448834	0.048313271
rplI	sp|P0A7R1|RL9_ECOLI	-1.074352178	3.31E-05
rplK	sp|P0A7J7|RL11_ECOLI	-1.030031069	0.000940436

argH	sp|P11447|ARLY_ECOLI	-1.029621386	0.010350803
cspC	sp|P0A9Y6|CSPC_ECOLI	-1.010776163	0.00524746
gpmA	sp|P62707|GPMA_ECOLI	1.007836346	0.000278112
gmhA	sp|P63224|GMHA_ECOLI	1.02030343	0.027607596
luxS	sp|P45578|LUXS_ECOLI	1.049588806	0.000506408
rpiA	sp|P0A7Z0|RPIA_ECOLI	1.052561101	0.01318169
pgi	sp|P0A6T1|G6PI_ECOLI	1.065800857	0.000188954
pykF	sp|P0AD61|KPYK1_ECOLI	1.081019653	2.73E-05
mlaC	sp|P0ADV7|MLAC_ECOLI	1.117643099	0.006345313
yfiF	sp|P0AGJ5|YFIF_ECOLI	1.128896995	0.03275735
uspG	sp|P39177|USPG_ECOLI	1.143746641	0.03765939
pgl	sp|P52697|6PGL_ECOLI	1.167828441	0.01318169
deoC	sp|P0A6L0|DEOC_ECOLI	1.174516518	0.018153582
grxB	sp|P0AC59|GLRX2_ECOLI	1.178432467	0.000732045
fbaB	sp|P0A991|ALF1_ECOLI	1.197658464	7.99E-05
adhE	sp|P0A9Q7|ADHE_ECOLI	1.201151692	3.88E-07
degP	sp|P0C0V0|DEGP_ECOLI	1.206061871	0.000278112
tas	sp|P0A9T4|TAS_ECOLI	1.219282681	0.036014767
katG	sp|P13029|KATG_ECOLI	1.220074858	7.01E-05
ribC	sp|P0AFU8|RISA_ECOLI	1.272415628	0.030285161
pmbA	sp|P0AFK0|PMBA_ECOLI	1.273270833	0.007784421

ftnA	sp|P0A998|FTNA_ECOLI	1.301974436	0.010044367
moaB	sp|P0AEZ9|MOAB_ECOLI	1.314241161	0.001746554
wrbA	sp|P0A8G6|NQOR_ECOLI	1.34050881	4.33E-06
hdhA	sp|P0AET8|HDHA_ECOLI	1.365397414	0.014224302
yahK	sp|P75691|YAHK_ECOLI	1.380128931	0.000137547
acnA	sp|P25516|ACNA_ECOLI	1.381383706	7.69E-05
sbp	sp|P0AG78|SUBI_ECOLI	1.410886097	0.001647533
degQ	sp|P39099|DEGQ_ECOLI	1.437968753	0.005431624
tktB	sp|P33570|TKT2_ECOLI	1.45986506	7.28E-07
yjgA	sp|P0A8X0|YJGA_ECOLI	1.470361111	0.0375636
glgP	sp|P0AC86|PHSG_ECOLI	1.489480956	0.004703968
yggE	sp|P0ADS6|YGGE_ECOLI	1.497996423	9.00E-05
mtlD	sp|P09424|MTLD_ECOLI	1.508297143	0.007422869
cbpA	sp|P36659|CBPA_ECOLI	1.520821408	0.040987772
deoB	sp|P0A6K6|DEOB_ECOLI	1.533048098	8.38E-05
ybjP	sp|P75818|YBJP_ECOLI	1.536131134	0.007427731
ybaY	sp|P77717|YBAY_ECOLI	1.541925932	0.00084693
tdcE	sp|P42632|TDCE_ECOLI	1.549713435	0.000108916
thiF	sp|P30138|THIF_ECOLI	1.58507647	0.00235383
hdeB	sp|P0AET2|HDEB_ECOLI	1.585194293	0.001902541
tpiA	sp|P0A858|TPIS_ECOLI	1.585411082	9.28E-09

uspE	sp|P0AAC0|USPE_ECOLI	1.605682754	0.000278112
rraA	sp|P0A8R0|RRAA_ECOLI	1.607904489	0.000417551
yeaY	sp|P0AA91|YEAY_ECOLI	1.609759679	0.001461256
nagD	sp|P0AF24|NAGD_ECOLI	1.651426571	0.020565443
slp	sp|P37194|SLP_ECOLI	1.661057589	0.000127411
osmC	sp|P0C0L2|OSMC_ECOLI	1.671087683	3.90E-06
ycfP	sp|P0A8E1|YCFP_ECOLI	1.676895218	0.000511934
acuI	sp|P26646|ACUI_ECOLI	1.680553338	0.001851227
pflB	sp|P09373|PFLB_ECOLI	1.700478434	9.87E-14
lpoB	sp|P0AB38|LPOB_ECOLI	1.721279546	0.000376072
clpB	sp|P63284|CLPB_ECOLI	1.727347175	1.23E-12
gstB	sp|P0ACA7|GSTB_ECOLI	1.748580068	0.001647533
slt	sp|P0AGC3|SLT_ECOLI	1.751322485	0.015251337
yggN	sp|P0ADS9|YGGN_ECOLI	1.768724665	0.049702565
loiP	sp|P25894|LOIP_ECOLI	1.795763405	0.012616436
pspA	sp|P0AFM6|PSPA_ECOLI	1.803047994	0.000834302
chrR	sp|P0AGE6|CHRR_ECOLI	1.829520642	0.00084693
gstA	sp|P0A9D2|GSTA_ECOLI	1.843282701	2.82E-06
msrA	sp|P0A744|MSRA_ECOLI	1.871059945	0.023254637
dps	sp|P0ABT2|DPS_ECOLI	1.8711205	1.92E-14
yghU	sp|Q46845|YGHU_ECOLI	1.947282377	0.000291495

lldD	sp|P33232|LLDD_ECOLI	1.949858867	0.035748827
yfcH	sp|P77775|YFCH_ECOLI	1.988861376	0.015909254
cdd	sp|P0ABF6|CDD_ECOLI	1.993269202	0.042932089
yeaG	sp|P0ACY3|YEAG_ECOLI	2.011516734	4.21E-11
pqiB	sp|P43671|PQIB_ECOLI	2.013507475	0.028681191
ygaU	sp|P0ADE6|YGAU_ECOLI	2.059465726	3.23E-14
blc	sp|P0A901|BLC_ECOLI	2.064974471	0.011100578
qorA	sp|P28304|QOR1_ECOLI	2.155671369	4.33E-07
curA	sp|P76113|CURA_ECOLI	2.175460559	0.010474873
bfr	sp|P0ABD3|BFR_ECOLI	2.180562547	1.56E-13
yegP	sp|P76402|YEGP_ECOLI	2.284730203	7.89E-10
ugd	sp|P76373|UDG_ECOLI	2.340441236	0.018153582
yceF	sp|P0A729|YCEF_ECOLI	2.382621181	0.040987772
yecA	sp|P0AD05|YECA_ECOLI	2.405409463	0.03275735
bepA	sp|P66948|BEPA_ECOLI	2.421711425	0.000774393
otsA	sp|P31677|OTSA_ECOLI	2.448868257	6.65E-08
katE	sp|P21179|CATE_ECOLI	2.50821371	2.35E-14
hchA	sp|P31658|HCHA_ECOLI	2.50879706	2.75E-12
dkgA	sp|Q46857|DKGA_ECOLI	2.649063918	1.64E-08
ybeL	sp|P0AAT9|YBEL_ECOLI	2.658797186	3.16E-08
ymdB	sp|P0A8D6|YMDB_ECOLI	2.671235984	0.024568779

fruB	sp|P69811|PTFAH_ECOLI	2.672541661	1.42E-05
ydbC	sp|P25906|YDBC_ECOLI	2.711914425	0.046758232
ynhG	sp|P76193|YNHG_ECOLI	2.733690087	3.31E-05
yciE	sp|P21363|YCIE_ECOLI	2.735728026	4.14E-05
ymbA	sp|P0AB10|YMBA_ECOLI	2.798163878	0.036198317
ydcL	sp|P64451|YDCL_ECOLI	2.82774542	2.04E-15
ldhA	sp|P52643|LDHD_ECOLI	2.837647013	2.28E-05
amn	sp|P0AE12|AMN_ECOLI	2.855927832	1.06E-13
yehZ	sp|P33362|YEHZ_ECOLI	2.946432256	0.017149234
mdtE	sp|P37636|MDTE_ECOLI	3.005567607	2.30E-05
gadB	sp|P69910|DCEB_ECOLI	3.026824738	1.75E-33
yhjG	sp|P37645|YHJG_ECOLI	3.114565954	0.004244044
gadC	sp|P63235|GADC_ECOLI	3.122545884	0.000234696
gadA	sp|P69908|DCEA_ECOLI	3.154673417	6.58E-36
ydhF	sp|P76187|YDHF_ECOLI	3.156625723	5.19E-05
yjgR	sp|P39342|YJGR_ECOLI	3.159652222	1.41E-12
yghA	sp|P0AG84|YGHA_ECOLI	3.175685027	1.40E-09
ycaC	sp|P21367|YCAC_ECOLI	3.185383669	2.44E-13
phoE	sp|P02932|PHOE_ECOLI	3.204721908	2.80E-10
ycgM	sp|P76004|YCGM_ECOLI	3.314835257	0.007920581
tam	sp|P76145|TAM_ECOLI	3.330887068	0.010474873

yhbO	sp|P45470|YHBO_ECOLI	3.335162899	0.000187361
phoR	sp|P08400|PHOR_ECOLI	3.473374236	0.001325039
mdaB	sp|P0AEY5|MDAB_ECOLI	3.504985139	1.26E-20
dkgB	sp|P30863|DKGB_ECOLI	3.638374711	5.12E-09
yniA	sp|P77739|YNIA_ECOLI	3.681925301	2.85E-07
gldA	sp|P0A9S5|GLDA_ECOLI	3.74947351	3.13E-05
phnD	sp|P16682|PHND_ECOLI	3.815420505	9.61E-10
ugpQ	sp|P10908|UGPQ_ECOLI	4.003964005	1.04E-06
ahr	sp|P27250|AHR_ECOLI	4.055942555	2.68E-05
pstS	sp|P0AG82|PSTS_ECOLI	4.113208109	3.15E-56
cysQ	sp|P22255|CYSQ_ECOLI	4.144617537	1.22E-07
glsA1	sp|P77454|GLSA1_ECOLI	4.234450197	0.024320745
yciF	sp|P21362|YCIF_ECOLI	4.316219533	0.000228764
ais	sp|P45565|AIS_ECOLI	4.363027397	0.02438333
kefG	sp|P0A756|KEFG_ECOLI	4.489497738	0.01318169
ydcS	sp|P76108|YDCS_ECOLI	4.581943058	0.009304996
phnB	sp|P16681|PHNB_ECOLI	4.606620186	4.21E-11
sodC	sp|P0AGD1|SODC_ECOLI	4.625310091	1.37E-05
arnA	sp|P77398|ARNA_ECOLI	4.698344251	1.36E-11
fic	sp|P20605|FIC_ECOLI	4.74532692	0.001647533
ugpB	sp|P0AG80|UGPB_ECOLI	4.789402823	1.62E-50

phoU	sp|P0A9K7|PHOU_ECOLI	4.829226009	7.34E-29
kdpC	sp|P03961|KDPC_ECOLI	4.927327983	0.001190624
pstB	sp|P0AAH0|PSTB_ECOLI	5.294764534	6.41E-25
yebE	sp|P33218|YEBE_ECOLI	5.774676565	1.18E-14
phoA	sp|P00634|PPB_ECOLI	5.790096911	7.32E-85
phoB	sp|P0AFJ5|PHOB_ECOLI	Inf	8.96E-35
arnB	sp|P77690|ARNB_ECOLI	Inf	8.99E-08
appA	sp|P07102|PPA_ECOLI	Inf	7.69E-05
yedP	sp|P76329|MPGP_ECOLI	Inf	7.72E-05
ugpC	sp|P10907|UGPC_ECOLI	Inf	0.000683993
arnC	sp|P77757|ARNC_ECOLI	Inf	0.008862011
yfcG	sp|P77526|YFCG_ECOLI	Inf	0.018153582
yraR	sp|P45469|YRAR_ECOLI	Inf	0.012911411
yhhA	sp|P0ADX7|YHHA_ECOLI	Inf	0.018724338
phnH	sp|P16686|PHNH_ECOLI	Inf	0.025826417
gspJ	sp|P45761|GSPJ_ECOLI	Inf	0.036463755
PmrA	sp|P30843|PMRA_ECOLI	Inf	0.037433165
alpA	sp|P33997|ALPA_ECOLI	Inf	0.037433165

1382

1383 **Table S3.** Bacteria strains used in this study.

Name	Description	Source
MG1655	*F- λ- ilvG- rfb-50 rph-1*	Lab collection
DH5α	*E. coli* F⁻ *endA1 glnV44 thi-1 recA1 relA1 gyrA96 deoR nupG purB20 φ80dlacZΔM15 Δ(lacZYA-argF)U169, hsdR17(r_K⁻m_K⁺), λ⁻*	Lab collection
Bl21 (DE3)	*E. coli* B F⁻ *ompT gal dcm lon hsdS_B(r_B⁻m_B⁻) [malB⁺]_{K-12}(λ^S)*	Lab collection
E. coli O157	EHEC, O157:H7	Clinical isolate from Queen Mary Hospital (HK)
UTI-1	UPEC isolated from UTI patients	Clinical isolate from Queen Mary Hospital (HK)
UTI-3	UPEC isolated from UTI patients	Clinical isolate from Queen Mary Hospital (HK)
UTI-4	UPEC isolated from UTI patients	Clinical isolate from Queen Mary Hospital (HK)
UTI-5	UPEC isolated from UTI patients	Clinical isolate from Queen Mary Hospital (HK)
Salmonella ATCC14028	*Salmonella* ATCC14028 wild type	Lab collection
P. aeruginosa	*P. aeruginosa* PAO1 wild type	Lab collection

PAO1

AY7010	MG1655 ArnB-3xFLAG	This study
AY0416	MG1655 $\Delta lacZ::Km^R$	This study
AY4875	MG1655 $\Delta arnT:: Km^R$	This study
AY4860	MG1655 $\Delta arnB:: Km^R$	This study
AY4862	MG1655 $\Delta ugd:: Km^R$	This study
AY4858	MG1655 $\Delta pmrB:: Km^R$	This study
AY4859	MG1655 $\Delta pmrA:: Km^R$	This study
AY2741	MG1655 $\Delta phoB:: Km^R$	This study
AY2742	MG1655 $\Delta phoR:: Km^R$	This study
AY7011	MG1655 $\Delta pmrB:: Km^R \Delta lacZ$	This study
AY7012	MG1655 $\Delta pmrA:: Km^R \Delta lacZ$	This study
AY7013	MG1655$\Delta phoB:: Km^R \Delta lacZ$	This study
AY7014	MG1655$\Delta phoR:: Km^R \Delta lacZ$	This study
AY7015	MG1655 $\Delta lacZ, pmrB^{E36A}$	This study
AY7016	MG1655 $\Delta lacZ, pmrB^{E39A}$	This study
AY7017	MG1655 $\Delta pmrA$	This study
AY7018	MG1655 $pmrB^{E36A}$	This study
AY7019	MG1655 $pmrB^{E39A}$	This study
AY7020	MG1655 $\Delta fecA$	This study
AY7021	MG1655 $\Delta fepA$	This study

AY7022	MG1655 Δ*entE*	This study
AY7023	MG1655 Δ*fiu*	This study
AY7024	MG1655 Δ*cirA*	This study
AY7025	MG1655 Δ*fecA* Δ*fepA* Δ*entE* Δ*fiu* Δ*cirA*	This study

1384

1385

 Table S4. Plasmids used in this study.

Name	Relevant characteristics	Source
pNN387	Single copy plasmid containing NotI and HindIII restriction sites before *lacZ*	Lab collection
pNN387(P*arn*-*lacZ*)	Ligate promoter of *arn* into NotI/HindIII linearized pNN387	This study
pNN387(P*ugd*-*lacZ*)	Ligate promoter of *ugd* into NotI/HindIII linearized pNN387	This study
pNN387(P*phoQ*-*lacZ*)	Ligate promoter of *phoQ* into NotI/HindIII linearized pNN387	This study
pNN387(P*mgtA*-*lacZ*)	Ligate promoter of *mgtA* into NotI/HindIII linearized pNN387	This study
pET28a-N	t7 promoter in pET-28a is replaced by araBad promoter	This study
pET28a-N-3xFLAG	C-terminal 6xHis tag in pET-28a-N is replaced by 3xFLAG tag	This study
pET-N-PmrA-6xHis	Ligate *pmrA* into NcoI/XhoI linearized pET-28a-N	This study
pET-N-PmrA-3xFLAG	Ligate *pmrA* into NcoI/XhoI linearized pET-28a-N-3xFLAG	This study
pET-N-PmrA137TAG-6xHis	Mutate G137 to TAG in pET28A-N-PmrA-6xHis	This study

| pEVOL-CouA-RS | Plasmid expression CouA tRNA synthetase | Lab collection |
| pCAGO | Editing plasmid containing CRISPR-Cas9 and λ-red | Gift from Prof. Changhao Bi |

1387