Social connections, group size and fine-scale manipulations of social stability shape learning, scrounging

and foraging rate in mixed flocks of wild songbirds

Michael S. Reichert[1,2], Josh A. Firth[3,4], Camille Troisi[2,5], John L. Quinn[2]

[1]Department of Biology, Oklahoma State University, Stillwater, Oklahoma, USA

[2]School of Biological, Earth and Environmental Sciences, University College Cork, Cork, Ireland

[3]Department of Biology, Edward Grey Institute, University of Oxford, Oxford, U.K.

[4]Faculty of Biological Sciences, University of Leeds, Leeds, U.K.

[5]Ethologie animale et humaine, Université de Rennes, Rennes, France

Correspondence: John Quinn, email: j.quinn@ucc.ie

ABSTRACT

The social environment provides both opportunities and challenges for foragers. How multiple

components of sociality and foraging interact simultaneously is poorly understood, while experimental

manipulation of sociality within wild foraging groups is rare. We tested whether production learning,

scrounging, and foraging rate are associated with individual social network metrics and manipulated

social stability in mixed species flocks of wild birds. First, individuals were randomly allowed access to,

and learned to forage primarily from, one of five feeders. Thereafter, we performed two reversals,

manipulating social connections by either assigning birds to a new feeder along with all others assigned

to their previous feeder (stable treatment) or reassigning each bird individually (unstable treatment).

Most of the effects observed were context dependent but in general we found that: i) Learning was

slower in the stable treatment, for individuals with higher weighted degree, and in smaller flocks; ii)

Scrounging was higher in the stable treatment, increased with weighted degree, but decreased with

flock size; and iii) Foraging rate was predicted only by weighted degree. Our findings demonstrate the

complex and dynamic nature of the relationship between sociality and foraging and suggest that

selection for group living is more nuanced than usually recognized.

INTRODUCTION

Foraging is the result of a complex set of decisions about when, where and how to obtain food (Stevens

et al. 2007). Because foraging provides the energy that is critical for fitness, a diverse set of strategies

have evolved to ensure foraging success (Waite and Field 2007). A key challenge for foragers is that this

behavior takes place in a constantly changing environment, and this environmental change can

drastically affect variables that determine the risks and rewards of any given foraging strategy such as

the availability and distribution of food, the technique required to extract it, and the necessary trade-

offs with other traits such as reproduction and vigilance against predators (Oom et al. 2004; Monk et al.

2018).

Additionally, for the many species that forage in social groups, sociality can offer both opportunities and

challenges to foraging individuals (Clark and Mangel 1986; Krause and Ruxton 2002). Social groups may

provide advantages to individual foragers such as increased discovery of food (Pitcher et al. 1982;

Snijders et al. 2021), increased efficiency in extracting it (Courchamp and Macdonald 2001; Beauchamp

2005; MacNulty et al. 2014), opportunities to socially learn from other individuals (Hoppitt and Laland

2013) and shared vigilance against predators (Cresswell 1994; Lima et al. 1999; Silk 2007; Sansom et al.

2008). However, foraging in a group also leads to increased competition (Cresswell 1997; Focardi and

Pecchioli 2005; Beltrão et al. 2022), as well as other general costs of sociality such as predator attraction

and disease spread (Godfrey et al. 2009; Drewe 2010; Weber et al. 2013). Previous studies show mixed

results for whether social group characteristics such as group size enhance or detract from foraging

success (Beauchamp 1998). In part this may be because foraging itself is a complex trait that is the

outcome of multiple functional variables, and individuals differ in how these combine to determine their

overall foraging strategy (Nachappa et al. 2010). For instance, some individuals are simply more active

foragers than others (DeLong et al. 2021), and for any given foraging effort some will spend more time

as producers and others as scroungers (Morand-Ferron et al. 2007). The interplay between sociality and

individual variation in multiple components of foraging is not well understood (Morand-Ferron et al.

2011; Aplin and Morand-Ferron 2017), but critical for a full understanding of the factors that determine

foraging success in natural populations.

Understanding the effects of sociality on foraging is further complicated because not all individuals in

the same social group experience the same social environment. Animal social network analyses have

shown that some individuals have stronger connections, are more central in the network or experience

specific types of connections more often than others (Whitehead 2008; Krause et al. 2015). The

temporal stability of social networks can also affect individual network position (Ansmann et al. 2012;

Murphy et al. 2020). Social network position in turn affects individual foraging behaviors (Giraldeau and

Caraco 2000). For instance, individuals that are more central in the network may experience greater

levels of foraging competition but also have more opportunities to scrounge from others (Aplin and

Morand-Ferron 2017; McMahon et al. 2024). Network density can also affect competition levels,

although the direction of this effect is highly variable (Beauchamp 1998; Vásquez and Kacelnik 2000;

Grueter et al. 2018; Sheppard et al. 2018). Individuals with close social ties may reduce their own

foraging effort in order to maintain those ties (Firth et al. 2015). And individuals that are closely

connected to others that act as sources of information—for example, individuals that discover a new

food patch or innovate a new foraging technique—may benefit from those connections by being an

early recipient of social information (Slagsvold and Wiebe 2011; Aplin et al. 2012; Kulahci et al. 2016;

Schakner et al. 2017; McMahon et al. 2024). Likewise, the overall structure of the social network may

impact on individual foraging behavior. For instance, networks that are more temporally stable (i.e. in

which connections between individuals remain relatively consistent over time) may allow for more

opportunities for social learning (Kulahci and Quinn 2019), but may also affect an individual's

competitiveness through the establishment of dominance hierarchies (McDonald and Shizuka 2013).

Different components of foraging success may be differentially impacted by social network structure

and position. First, foraging traits differ in how much they rely on or are affected by social interactions.

For example, scrounging requires the presence of other individuals, and is impacted by both social

network structure and social stability (Firth et al. 2015; Aplin and Morand-Ferron 2017). In contrast to

scrounging, other components of foraging behavior do not require social contact and may or may not be

affected by interactions with others in a social group. For instance, individual foraging rates could be a

product of individual hunger levels and extraction efficiency, but they can also be affected by social

group characteristics such as group size, for example through effects of competition (Rieucau and

Giraldeau 2009*a*) and the perception of predation risk driven by shared vigilance or the dilution effect

(Stevens et al. 2007). Foraging rates can also be improved by the strength of social interactions, which

may improve an individual's "sense of security" (Voelkl et al. 2016; Heathcote et al. 2017; Mady and

Blumstein 2017). Learning about beneficial sources of food and how these can be obtained is another

important component of foraging that is not necessarily a social process although it is often modulated

by social interactions. Second, the role of cognition, and the associated social environment effects on

cognitive performance, varies across foraging traits. Learning about where food can be found and how it

can be extracted is a common role for cognition in foraging, and is affected in multiple ways by the

social environment, not only via social learning but also through indirect effects of competition,

dominance and local enhancement (Pravosudov et al. 2003; Morand-Ferron and Quinn 2011; Avarguès-

Weber and Chittka 2014; Ashton et al. 2018, 2019; Langley et al. 2018), although there is variation in

these effects (Heinen et al. 2021*a*; Lambert and Guillette 2021). Learning is involved not only in

production but also scrounging (Reichert et al. 2021), but learning should have limited effects on

foraging performance when feeding on a known, readily accessible food source. The effects of sociality

on learning may be especially important when environmental conditions change, and previously learned

contingencies are no longer effective for finding food. When this happens, strong connections to other

individuals may be beneficial when there is social learning, such that when any individual accesses a new

food source others can quickly do the same (Aplin et al. 2012; Jones et al. 2017). However, strong

connections could also be costly when conditions change if social information is outdated or if foraging

opportunities available to some individuals are not available for others they are connected to

(Beauchamp et al. 1997; Kendal et al. 2005; Hall and Kramer 2008; Rieucau and Giraldeau 2009*b*; Firth et

al. 2016). In this case, individuals would be better off using their own private information to learn about

new food sources, but some may do so more readily than others (Thornton and Malapert 2009; Kurvers

et al. 2010; Pitera et al. 2025).

We tested social effects on foraging in wild mixed-species flocks of great tits (*Parus major*) and blue tits

(*Cyanistes caeruleus*) in the context of an experiment in which birds had to learn which feeder in an

array they could access to obtain food (Reichert et al. 2020). First, we tested whether an experimental

manipulation of social stability would affect three measures of foraging success: reversal learning of a

rewarded feeder location, propensity to scrounge, and overall foraging rate. On the one hand, social

instability could lead to disruptions in social relationships and opportunities for social learning, resulting

in slower reversal learning, and an increased tendency to scrounge. On the other hand, social instability

could facilitate reversal learning through disruptions to competitive hierarchies at feeders while

reducing scrounging opportunities. For either scenario, the effects of social instability should increase

with each successive reversal. To our knowledge, no previous study in a wild species has manipulated

the structure of the social network to determine the effects on multiple components of foraging

success. Second, we tested the relationship between two metrics of individual social network

positioning and foraging. The individual metrics used were the average size of the flocks birds were in

when visiting feeders and weighted degree, a measure of the strength of social connections with other

individuals in the network. Both of these metrics capture aspects of sociality that are likely to have

multiple impacts on foraging by influencing shared vigilance, levels of competition, opportunities for

social learning and expression of alternative foraging tactics (Dias 2006; Cantor et al. 2020). We

examined these relationships across three different learning tasks, an initial production learning and two

reversals. We predicted that the effects of sociality on foraging would vary across these learning tasks

because they involve different cognitive traits that have been shown differ in how they interact with the

social environment (Lambert and Guillette 2021): learning an initial association versus flexibility in

updating behavior in response to environmental change (Izquierdo et al. 2017).

MATERIALS AND METHODS

Study system

In winter, great tits and blue tits form flocks along with several other species and move about woodland

areas to forage (Ekman 1989). These flocks have a fission-fusion structure, in which individuals

frequently leave and rejoin the group, resulting in variation in the social bonds among individuals (Aplin

et al. 2015; Farine et al. 2015*b*). Multiple aspects of both their foraging behavior and their social

networks can be readily observed by tagging individuals with passive integrated transponders and then

monitoring their visits to RFID-equipped feeders (Firth and Sheldon 2015; Milligan et al. 2017; Cauchoix

et al. 2022). Furthermore, by programming feeders to allow access to the food inside for only certain

individuals, it is possible to both measure different components of foraging performance and to

manipulate the social structure (Firth and Sheldon 2015). Social structure and social network positioning

have important consequences for foraging performance (Aplin et al. 2012; Firth et al. 2015; Beck et al.

2024; McMahon et al. 2024) and for other important traits in these species (Farine and Sheldon 2015;

Firth and Sheldon 2016). Furthermore, these social network effects are not limited to conspecifics

because heterospecific social interactions also impact foraging (Aplin et al. 2012; Farine et al. 2012,

2014, 2015a).

We studied social behavior and foraging in the context of a discrimination learning experiment in a wild

mixed species foraging flock focused on great tits and blue tits at Wytham Woods, Oxfordshire, UK. The

general experimental design and methods are described in detail in our previous publications (Reichert

et al. 2020, 2021; Morand-Ferron et al. 2022). Briefly, we used arrays of sunflower-seed feeders with

radio frequency identification (RFID) antennas to monitor foraging of great tits and blue tits that had

been fitted with a passive integrative transponder (PIT) tag (IB Technology, Aylesbury, UK). Every bird

had a unique RFID code and we could therefore detect each visit by each individual to any of the feeders

in the array. Each array consisted of five feeders arranged linearly and spaced 1 m apart. Four arrays

were operative in different areas of the wood at any given time, and they were placed far enough apart

that birds rarely visited feeders at more than one array. One set of four feeders was monitored from

November-December 2017 and another set of four between January-February 2018. All procedures

were approved by the Animal Welfare Body at University College Cork (HPRA license number AE19130-

P017), and were in accordance with the Association for the Study of Animal Behaviour Guidelines for the

Treatment of Animals in Behavioural Research and Teaching. All research was conducted under British

Trust for Ornithology licenses as part of ongoing research in this population.

Prior to the learning experiments that contributed the data analyzed here, the experimental feeders

were placed in the area to attract birds and allow them to become accustomed to obtaining food from

them (for details see Reichert et al. 2020; Troisi et al. 2025). We initially placed two feeders 50 m apart

in an 'initial dispersed' stage to attract birds to the area. We then removed those feeders and placed five

feeders in a linear array, in the same position as they were used in the learning experiment. During this

'open clustered' stage, any tagged bird could feed from any feeder. During the learning experiments, the

focus of this study, each bird was assigned to one of the five feeders in the array, and it could only get

food from its assigned feeder. If it landed on the assigned feeder, a solenoid would activate, allowing the

bird to push open a door to gain access to the seeds inside. If it landed on any of the other feeders, its

visit would be recorded but the solenoid would not activate, so it would not be able to access the food.

The monitoring of the visits and activation of the solenoid was controlled by a custom program on a

'Darwin Board' (Stickman Technologies Inc., UK). The learning experiment was carried out in three

stages. The first was an initial learning stage in which each individual was randomly assigned to one of

the five feeders in the array and then we monitored their visits over eight days to determine if and when

they learned which feeder they were assigned to (see below). Second, in the first reversal stage, we

reassigned birds to a new feeder in the array and monitored their visits for another eight-ten days. In

contrast to the initial learning stage, in the reversal stage we applied two different 'social stability

treatments' that determined how birds were reassigned to new feeders. At four of the eight sites (two

in each year), we applied the 'stable' social treatment, in which the entire set of birds assigned to the

same feeder during the initial learning stage were reassigned to the same (randomly chosen) new

feeder. Thus, these birds would experience consistency in which other individuals in the flock would be

feeding from the same rewarded feeder. At the other four sites, we applied the 'unstable' social

treatment. In this treatment, each individual was randomly reassigned to a new feeder independently,

and thus was not always consistently feeding from the same rewarded feeder with the same other

individuals and may therefore have experienced disruptions in their social connections. Third, in the

second reversal stage, we again reassigned birds to new feeders, applied the stable and unstable

treatments as above, and monitored visits for eight to ten days.

In our previous analyses of this dataset, we provided an extensive description of demographic factors

affecting variation in participation, likelihood of learning, and learning speed in this system (Reichert et

al. 2020). We found both species- and individual-level variation in learning speed (Reichert et al. 2020),

and in the propensity to scrounge (Reichert et al. 2021). In an analysis focused on great tits, we also

found that the social stability treatment had lasting effects on the social network that carried over

across contexts (Troisi et al. 2025). Here, we extend this work by asking whether an experimental

manipulation of flock stability, and individuals' social network positions, affected individual foraging

behavior (learning, scrounging, foraging rate), and if so, which components were affected.

Social network and foraging measures

In order to capture various aspects of foraging behavior, we calculated variables corresponding to three

components of individual foraging strategies. Each of these variables was calculated separately for each

individual in each of the three experimental stages. First, we characterized production learning behavior

by calculating learning speed as the number of visits until an individual first met a criterion (following

previous research) of 16 correct visits within a window of 20 consecutive visits (Reichert et al. 2020).

Second, we characterized the propensity to scrounge as the proportion of rewarded visits (i.e. visits in

which the bird obtained food, which could happen either by visiting the correct feeder or scrounging)

that were scrounges. We defined scrounging as a visit by an individual to a feeder it was not assigned to

(and therefore where it could not obtain food by producing) that took place 1 s or less after the

departure of a bird that was assigned to that feeder. In such cases, the feeder door had not yet shut and

the scrounger could obtain food from that feeder (Reichert et al. 2021). Third, we characterized foraging

intake rate as a measure of the rate at which birds obtained food once they had learned the task,

quantified as the number of visits to their rewarded feeder on the day after it had met the learning

criterion.

We note that individuals of both great tits and blue tits foraged in the same mixed species flocks, and so

the social networks were constructed and social network metrics calculated using data from all detected

individuals, regardless of species. Monitoring at the study population also involves giving RFID tags to

any captured marsh tit (*Poecile palustris*) and nuthatch (*Sitta europaea*), thus social networks also

include these individuals when detected. We restrict our analyses to great tits and blue tits because only

these two species interacted with the learning apparatus in large numbers. The number of individuals of

each species included in the social network varied slightly between experimental stages (Table S1).

We built social networks by assigning birds' visits to discrete flocking events using a Gaussian mixture

model (Psorakis et al. 2012, 2015). We defined edges in the social network between each individual that

was detected in a flocking event (Troisi et al. 2025). We then quantified the association strength

between each possible pair of individuals using a simple ratio index calculated as the number of times

both individuals were seen in the same flocking event divided by the total number of flocking events in

which either (or both) individual was present (Cairns and Schwager 1987; Whitehead 2008). From these

data we calculated the two social metrics used in analyses for each individual: 1) its average flock size

(i.e. the mean number of individuals it was observed with in flocking events) and 2) weighted degree as

the weighted sum of its association strengths with all other individuals in the network. These metrics

were calculated separately for each of the three experimental stages.

Statistical analyses

All statistical analyses were conducted in R version 4.5.0 (R Development Core Team 2024). Our dataset

included 68 individual great tits and 115 individual blue tits that met the learning criterion in all three

experimental stages. Other individuals that were detected during the experiments but did not meet the

learning criteria were excluded, because the majority of these individuals made very few visits to the

feeders (Reichert et al. 2020), and because we were interested in the effects of the social stability

treatment on reversal learning performance, which could not be assessed in a meaningful way if birds

did not meet the initial learning criterion prior to the reversal. We used separate statistical models for

each of the three foraging strategy variables to determine how these were affected by the social

stability treatment and an individual's social network metrics. All models were run with the lme4 version

1.1-35.5 package (Bates et al. 2015). Where significant interactions were identified we used Tukey post-

hoc tests to determine which groups were different from one another. For interactions involving

categorical variables (experimental stage, species and social stability), we did post-hoc testing using the

emmeans function from the emmeans version 1.10.4 package in R (Lenth 2022), which calculates

estimated marginal means for each combination of factors. For interactions involving a continuous

variable (average flock size or weighted degree), we did post-hoc testing using the emtrends function in

the emmeans package, which tests for differences in slopes of the relationship between the continuous

predictor and the foraging variable between the levels of the categorical variables (experimental stage,

species). We checked model assumptions of normality or uniformity of residuals and non-collinearity of

variables using the check_model function in the performance version 0.14.0 package (Lüdecke et al.

2021). Learning speed (natural log-transformed) was modeled as a Gaussian variable in a linear mixed

model. Proportion scrounged was modeled as a binomial variable in a generalized linear mixed model.

Foraging rate was modeled as a Gaussian variable in a linear mixed model. Raw data and code are

available at https://figshare.com/s/8ddfb55d391138371d36.

For each of the three foraging variables, results for which we present in separate sections, we tested for

the effects of our social stability manipulation and social variables. First, we tested for effects of the

social stability treatment. For these models, we only included data from the two reversal learning

stages, because the social stability treatment was applied after the initial learning stage. We included

individual identity as a random effect and included the three-way interaction between species,

experimental stage and social stability treatment as the main factors of interest, expecting that the

effects of the social stability treatment could vary by species or between the two reversal stages.

Specifically, for the reversal stages, we expected that relationships between social metrics and foraging

should become stronger by the second reversal as the relationships become more established in the

stable group and the difference with the unstable group more pronounced. Without specifying any

specific a priori predictions, we expected that relationships between sociality and foraging could vary

among species because of differences in factors like body size, dominance, and predation risk, all of

which influence foraging behavior. We included three additional variables as fixed effects to control for

factors that were identified in previous analyses to affect learning performance: whether or not the

feeder was located at the edge of the array, and the amounts of time that an individual's assigned and

non-assigned feeders were inoperable due to equipment malfunction (Reichert et al. 2020). In our

previous study we found that birds assigned to edge feeders learned more quickly in all three

experimental stages (Reichert et al. 2020). Some feeders malfunctioned during the course of the

experiment because of power loss or antenna failure and did not open for any of the birds or record any

visits until they were repaired (further details and data on these occurrences are given in Reichert et al.

2020). We still include data from birds in such arrays, and our statistical approach aims to minimize

these disruptions; we acknowledge that feeder malfunctions could have affected individual learning

performance and are a source of noise for the other variables, which we assume was random with

respect to the hypotheses tested.

Second, we tested for effects of the two social network position variables, average flock size and

weighted degree. These models included data from all three experimental stages. Each model included

individual identity as a random effect and two three-way interaction terms: species, experimental stage

and average flock size, and species, experimental stage and weighted degree. Social stability treatment

was not included as a factor in this model because it was not applied during the initial learning stage and

because for these tests we were interested in the effects of the social network position itself, which

would have been obscured by including a factor that affects social network position, as we

demonstrated in a previous analysis of this dataset (Troisi et al. 2025). We again included additional

fixed effects of whether or not the feeder was located at the edge of the array, and the amounts of time

that an individual's assigned and non-assigned feeders were inoperable due to equipment malfunction.

RESULTS

There were 262530 records of the 183 focal individuals across the three experimental stages (great tits:

95818 visits, N = 68 individuals; blue tits: 166712 visits, N = 115 individuals). Across these stages, the

average flock size was (mean ± SE) 13.74 ± 0.26 for great tits and 12.59 ± 0.21 for blue tits, and the

average weighted degree was 6.65 ± 0.14 for great tits and 6.13 ± 0.12 for blue tits. Learning speed

averaged 62.4 ± 6.9 and 63.8 ± 4.62 visits until the learning criterion was met for great tits and blue tits,

respectively. For scrounging, great tits had 2347 total scrounging events (mean ± SE = 11.5 ± 1.8), and

blue tits had 6552 scrounges (mean ± SE = 19.0 ± 1.3). Foraging rate averaged 51.9 ± 1.4 and 46.0 ± 0.9

visits on the day after the learning criterion was met for great tits and blue tits, respectively.

Effects of sociality on learning speed

There was no significant three-way interaction between social stability treatment, experimental stage

and species (Table 1). However, there was a significant interaction between social stability treatment

and experimental stage (Table 1, Figure 1A, 1B). For both species, there was no difference between the

unstable and stable groups in learning speeds in the first reversal, but in the second reversal, individuals

in the unstable treatment learned more quickly (Table S2, Figure 1A). Learning speeds were not

significantly different in the two social stability treatment groups in the initial learning stage, confirming

that there were no differences prior to the treatment being applied (Table S3).

There were significant three-way interactions between species, experimental stage and both average

flock size and weighted degree on learning speed (Table 2). For blue tits, individuals in larger flocks

learned faster in the initial and first reversal stages, but this pattern was reversed in the second reversal

stage where individuals in smaller flocks learned faster (Table S4, Figure 1B). For great tits, individuals

generally learned faster in larger flocks, but, unlike in blue tits, the relationship between flock size and

learning speed did not differ across experimental stages (Table S4, Figure 1B). Blue tits were relatively

unaffected by weighted degree and showed no significant variation in the relationship between

weighted degree and learning speed across the three experimental stages, while great tits showed a

significantly different relationship between these two variables in the second reversal compared to

initial learning (Table S5, Figure 1C). Specifically, great tits' learning speeds were relatively independent

of weighted degree in the initial learning stage, but by the second reversal individuals with lower

weighted degrees learned more quickly than individuals with higher weighted degrees (Table S5, Figure

1C).

Effects of sociality on scrounging

There was a significant three-way interaction between species, experimental stage and social stability

treatment on the proportion scrounged (Table 1). Post-hoc tests indicated that scrounging levels were

significantly higher in the stable treatment compared to the unstable treatment for both species in the

second reversal (Table S6, Figure 2A). However, in the first reversal there was no effect of social stability

treatment on scrounging for great tits, and for blue tits there was a marginal effect, where again

scrounging levels were higher in the stable compared to the unstable treatment groups (Table S6, Figure

2A). However, we noticed that in the initial learning stage, there was a main effect of social stability

treatment: individuals at the sites that would be assigned to the stable treatment had higher

significantly scrounging levels than individuals in the unstable treatment, indicating that there may have

been a pre-existing difference between these groups for this trait, and this difference applied to both

species (Table S3). However, because we found no significant effect of social stability treatment in the

first reversal and then significant differences in the second reversal, the treatment had an effect over

and above any differences in the populations prior to treatment.

359 There were significant three-way interactions between species, experimental stage and both average

360 flock size and weighted degree on proportion scrounged (Table 2). For great tits, individuals in larger

361 flocks scrounged less often but this effect was much stronger in the reversal stages than in the initial

362 learning stage (Table S7, Figure 2B). There was also a significant difference between the two reversal

363 stages, with a stronger negative relationship in the first reversal than in the second reversal (Table S7,

364 Figure 2B). For blue tits, the relationship between flock size and proportion scrounged was overall much

365 weaker, but both reversal stages showed a more negative relationship between flock size and

366 proportion scrounged compared to the initial learning stage (Table S7, Figure 2B). In both species there

367 was an increasingly positive relationship between weighted degree and proportion scrounged as the

368 experiment proceeded from the initial learning stage through the reversals (Figure 2C): individuals with

369 higher weighted degree scrounged more often. For blue tits, the relationship became significantly

370 stronger and more positive in comparisons between each successive experimental stage (Table S8,

371 Figure 2C). For great tits, both reversal stages showed significantly stronger positive relationships

372 between weighted degree and proportion scrounged compared to the initial learning stage, but there

373 was no difference between the two reversal stages (Table S8, Figure 2C).

375 *Effects of sociality on foraging rate*

376 There was no effect of the social stability treatment on foraging rate (Figure 3A). This was the case both

377 for the full model including interactions with species and experimental stage (Table 1) and for a reduced

378 model that did not include the interactions (Table S9). In the reduced model, there was a statistically

379 significant effect of species, with great tits having a higher foraging rate than blue tits, and an effect of

380 feeder location, with higher foraging rates for feeders on the edge of the array (Table S9). Foraging rates

were not significantly different in the two social stability treatment groups in the initial learning stage,

confirming that there were no differences prior to the treatment being applied (Table S3).

There were no interactions between species, experimental stage and either average flock size or

weighted degree (Table 2). There was, however, a main effect of weighted degree: individuals with a

higher weighted degree had a higher foraging rate (Figure 3C). This effect was present in both the full

model (Table 2) and in a reduced model that did not include interaction effects (Table S10). In the full

model there was a marginal effect of average flock size (Table 2), although this effect was reduced and

not statistically significant in the reduced model without interactions (Table S10). Foraging rates

generally remained relatively constant or decreased slightly with increasing flock sizes (Figure 3B).

Figure 1. Effects of sociality on learning speed. A. Effects of social stability treatment and experimental

stage on learning speed (ln-transformed) for blue tits (left) and great tits (right). Black points represent

estimated marginal means (± 95% CI) from the model. Small colored points represent partial residuals

from the model for each individual. B. Relationship between average flock size and learning speed for

each species. Lines represent predicted estimate (± 95% CI) from the model, with separate colors for

each experimental stage, and points represent partial residuals from the model for each individual. C.

Relationship between weighted degree and learning speed for each species. Interpretation as in B.

Figure 2. Effects of sociality on scrounging. A. Effects of social stability treatment and experimental stage

on the proportion of feeds that were scrounges for blue tits (left) and great tits (right). Black points

represent estimated marginal means (± 95% CI) from the model. Small colored points represent partial

residuals from the model for each individual. B. Relationship between average flock size and scrounging

for each species. Lines represent predicted estimate (± 95% CI) from the model, with separate colors for

each experimental stage, and points represent partial residuals from the model for each individual. C.

Relationship between weighted degree and scrounging for each species. Interpretation as in B. The y-

axis is cut off at 0.6 for visualization purposes, there are 1, 13, and 13 outlying points not depicted in A,

B, and C, respectively. Figure S1 presents the same graphs but with all data points included.

Figure 3. Effects of sociality on foraging rate. A. Effects of social stability treatment and experimental

stage on foraging rate (number of rewarded visits on the day after the learning criterion was met) for

blue tits (left) and great tits (right). Black points represent estimated marginal means (± 95% CI) from the

model. Small colored points represent partial residuals from the model for each individual. B.

Relationship between average flock size and foraging rate for each species. Lines represent predicted

estimate (± 95% CI) from the model, with separate colors for each experimental stage, and points

represent partial residuals from the model for each individual. C. Relationship between weighted degree

and foraging rate for each species. Interpretation as in B.

 Table 1. Effects of social stability on foraging traits

Foraging trait	Effect	Estimate	SE	t	P
Learning speed	Intercept	3.407	0.145	23.553	
	Social stability treatment (unstable)	0.111	0.226	0.491	0.623
	Experiment (second reversal)	0.244	0.175	1.395	0.165
	Species (great tit)	0.127	0.218	0.583	0.560
	Own feeder malfunction time (hrs)	0.107	0.019	5.532	< 0.001
	Other feeder malfunction time (hrs)	0.061	0.012	5.018	< 0.001
	Feeder location (edge)	−0.46	0.12	−3.827	< 0.001
	Social stability treatment (unstable) X Experiment (second reversal)	−0.8	0.284	−2.819	0.005
	Social stability treatment (unstable) X Species (great tit)	−0.484	0.377	−1.283	0.201
	Experiment (second reversal) X Species (great tit)	−0.062	0.277	−0.224	0.823
	Social stability treatment (unstable) X Experiment (second reversal) X Species (great tit)	0.068	0.483	0.141	0.888
Scrounging	Intercept	−3.469	0.169	−20.535	
	Social stability treatment (unstable)	−0.532	0.273	−1.951	0.051
	Experiment (second reversal)	−0.142	0.038	−3.768	< 0.001
	Species (great tit)	−0.973	0.269	−3.615	< 0.001

	Own feeder malfunction time (hrs)	0.113	0.007	16.235	< 0.001
	Other feeder malfunction time (hrs)	0.065	0.005	12.411	< 0.001
	Feeder location (edge)	−0.016	0.035	−0.446	0.655
	Social stability treatment (unstable) X Experiment (second reversal)	−0.007	0.070	−0.095	0.924
	Social stability treatment (unstable) X Species (great tit)	−0.011	0.470	−0.023	0.982
	Experiment (second reversal) X Species (great tit)	0.800	0.069	11.56	< 0.001
	Social stability treatment (unstable) X Experiment (second reversal) X Species (great tit)	−0.564	0.149	−3.777	< 0.001
Foraging rate	Intercept	46.079	2.03	22.701	
	Social stability treatment (unstable)	-1.457	3.172	-0.459	0.646
	Experiment (second reversal)	0.142	2.418	0.059	0.953
	Species (great tit)	4.149	3.061	1.355	0.176
	Own feeder malfunction time (hrs)	-0.219	0.271	-0.807	0.42
	Other feeder malfunction time (hrs)	-0.233	0.17	-1.375	0.17
	Feeder location (edge)	3.79	1.67	2.269	0.024
	Social stability treatment (unstable) X Experiment (second reversal)	1.05	3.918	0.268	0.789
	Social stability treatment (unstable) X Species (great tit)	1.391	5.295	0.263	0.793
	Experiment (second reversal) X Species (great tit)	-1.817	3.829	-0.475	0.636

Social stability treatment (unstable) X Experiment (second reversal) X Species (great tit)	9.248	6.666	1.387	0.167

422   Fixed effects from mixed models with individual identity as a random effect and ln–transformed learning speed (Gaussian), proportion

423   scrounged (binary) or foraging rate (Gaussian) as the dependent variable. The reference category for social stability treatment is the stable

424   treatment, for experimental stage is the first reversal stage, for species is blue tits, and for the feeder location is center.

 Table 2. Effects of social characteristics on foraging traits

Foraging trait	Effect	Estimate	SE	t	P
Learning speed	Intercept	3.53	0.429	8.234	
	Experiment (first reversal)	0.144	0.518	0.279	0.781
	Experiment (second reversal)	−1.432	0.549	−2.607	0.009
	Species (great tit)	−0.214	0.715	−0.300	0.764
	Own feeder malfunction time (hrs)	0.086	0.014	6.244	< 0.001
	Other feeder malfunction time (hrs)	0.065	0.009	7.073	< 0.001
	Feeder location (edge)	−0.831	0.100	−8.271	< 0.001
	Flock size	−0.058	0.047	−1.222	0.222
	Weighted degree	0.080	0.095	0.842	0.400
	Flock size X Experiment (first reversal)	−0.090	0.074	−1.216	0.225
	Flock size X Experiment (second reversal)	0.184	0.071	2.604	0.010
	Flock size X Species (great tit)	0.013	0.080	0.167	0.868
	Experiment (first reversal) X Species (great tit)	0.638	0.909	0.702	0.483
	Experiment (second reversal) X Species (great tit)	0.253	1.001	0.252	0.801
	Weighted degree X Experiment (first reversal)	0.206	0.143	1.437	0.151

	Weighted degree X Experiment (second reversal)	−0.113	0.128	−0.880	0.380
	Weighted degree X Species (great tit)	−0.114	0.153	−0.746	0.456
	Flock size X Experiment (first reversal) X Species (great tit)	−0.035	0.119	−0.293	0.770
	Flock size X Experiment (second reversal) X Species (great tit)	−0.280	0.115	−2.442	0.015
	Weighted degree X Experiment (first reversal) X Species (great tit)	0.093	0.224	0.417	0.677
	Weighted degree X Experiment (second reversal) X Species (great tit)	0.631	0.204	3.096	0.002
Scrounging	Intercept	−3.725	0.25	−14.891	
	Experiment (first reversal)	0.936	0.132	7.076	< 0.001
	Experiment (second reversal)	0.252	0.152	1.656	0.098
	Species (great tit)	−0.959	0.510	−1.878	0.060
	Own feeder malfunction time (hrs)	0.093	0.005	18.612	< 0.001
	Other feeder malfunction time (hrs)	0.054	0.003	17.47	< 0.001
	Feeder location (edge)	−0.214	0.029	−7.381	< 0.001
	Flock size	0.015	0.019	0.752	0.452
	Weighted degree	−0.048	0.033	−1.461	0.144
	Flock size X Experiment (first reversal)	−0.116	0.018	−6.483	< 0.001
	Flock size X Experiment (second reversal)	−0.140	0.02	−7.151	< 0.001

	Flock size X Species (great tit)	−0.098	0.047	−2.081	0.037
	Experiment (first reversal) X Species (great tit)	2.736	0.381	7.173	< 0.001
	Experiment (second reversal) X Species (great tit)	2.446	0.427	5.731	< 0.001
	Weighted degree X Experiment (first reversal)	0.122	0.037	3.283	0.001
	Weighted degree X Experiment (second reversal)	0.251	0.039	6.387	< 0.001
	Weighted degree X Species (great tit)	0.129	0.079	1.632	0.103
	Flock size X Experiment (first reversal) X Species (great tit)	−0.266	0.052	−5.121	< 0.001
	Flock size X Experiment (second reversal) X Species (great tit)	−0.089	0.044	−2.008	0.045
	Weighted degree X Experiment (first reversal) X Species (great tit)	0.206	0.098	2.112	0.035
	Weighted degree X Experiment (second reversal) X Species (great tit)	0.034	0.083	0.412	0.680
Foraging rate	Intercept	35.602	6.486	5.489	
	Experiment (first reversal)	1.722	7.4	0.233	0.816
	Experiment (second reversal)	-4.579	7.864	-0.582	0.561
	Species (great tit)	12.105	10.808	1.12	0.263
	Own feeder malfunction time (hrs)	-0.363	0.205	-1.766	0.078
	Other feeder malfunction time (hrs)	-0.112	0.137	-0.816	0.415
	Feeder location (edge)	4.239	1.448	2.927	0.004

Flock size	-1.349	0.711	-1.896	0.058
Weighted degree	4.285	1.418	3.021	0.003
Flock size X Experiment (first reversal)	1.438	1.068	1.347	0.179
Flock size X Experiment (second reversal)	1.483	1.024	1.448	0.148
Flock size X Species (great tit)	0.235	1.208	0.194	0.846
Experiment (first reversal) X Species (great tit)	-9.344	12.995	-0.719	0.473
Experiment (second reversal) X Species (great tit)	15.762	14.338	1.099	0.272
Weighted degree X Experiment (first reversal)	-3.196	2.065	-1.547	0.123
Weighted degree X Experiment (second reversal)	-2.258	1.872	-1.206	0.228
Weighted degree X Species (great tit)	-1.242	2.295	-0.541	0.589
Flock size X Experiment (first reversal) X Species (great tit)	-0.598	1.715	-0.348	0.728
Flock size X Experiment (second reversal) X Species (great tit)	-1.415	1.657	-0.854	0.394
Weighted degree X Experiment (first reversal) X Species (great tit)	2.096	3.245	0.646	0.519
Weighted degree X Experiment (second reversal) X Species (great tit)	0.281	2.963	0.095	0.925

428   Fixed effects from mixed models with individual identity as a random effect and ln–transformed learning speed (Gaussian), proportion

429   scrounged (binary) or foraging rate (Gaussian) as the dependent variable. The reference category for experimental stage is the initial learning

430   stage, for species is blue tits, and for the feeder location is center.

432   Table 3. Summary of results and potential mechanisms

Social trait	Foraging trait	Effect	Potential mechanism
Social stability	Learning speed	Negative	Social instability disrupts dominance hierarchies, facilitating learning
	Scrounging	Positive	More competition in stable groups, or trade-off with learning speed
	Foraging rate	None	Individual variation in hunger, motivation or competitive ability
Flock size	Learning speed	Mostly positive	Learning is facilitated by shared vigilance, reduced risk of predation
	Scrounging	Negative	Trade-off with learning speed
	Foraging rate	None	Individual variation in hunger, motivation or competitive ability
Weighted degree	Learning speed	Negative	Individuals with strong social ties visit non-rewarded feeders with social partners
	Scrounging	Positive	Stronger connections facilitate learning to scrounge, or trade-off with learning speed
	Foraging rate	Positive	Increased sense of security and shared vigilance with strong social connections

For each combination of social and foraging trait measured, we give the direction for the general trend in the relationship between the two

traits, but note that all of these effects showed additional variation across species and experimental stages that in some cases resulted in a

qualitatively different relationship. Potential mechanisms for the effect are given, but these are speculative and our experiments were not

designed to isolate specific mechanisms.

DISCUSSION

Sociality had clear effects on all aspects of foraging behavior, indicating that foraging in social groups

affects food intake, which could carry over to affect allocation of resources to life history traits. In a

previous analysis of this dataset focused on great tits, we demonstrated that metrics of an individual's

social network position in these experiments are repeatable, indicating consistent among-individual

variation in sociality (Troisi et al. 2025). Here we show that these same metrics affect foraging behavior

in both great tits and blue tits, demonstrating an important consequence of individual variation in

sociality and also indicating that these summary metrics of social position do indeed capture meaningful

biological variation. Although there were multiple differences in effects on the two study species, these

were often in the magnitude rather than the direction of the effect, which also underscores the

robustness of our findings. The links among these traits were context dependent, and suggest

interactions with both individual factors like variation in cognitive ability and the positive and negative

effects of social grouping (reviewed in Table 3). Below, we explore and highlight the novelty in our

results, and generate hypotheses for future experimental work to understand the mechanisms by which

complex social structure affects different aspects of foraging success simultaneously.

Effects of sociality on learning speed

We had predicted that reversal learning would be faster in the stable treatment because of increased

opportunities for social learning, but we found no effect of treatment for the first reversal and a

significant effect in the opposite direction for the second reversal. For both species, individuals in the

unstable group were faster to learn in the second reversal. We note that a previous analysis on this

study system focused on the dynamics of the learning process, and therefore including many other

variables not analyzed here, found no effects of social stability treatment on reversal learning speed

(Reichert et al. 2020). The difference between our current study and these previous results can be

explained by our more focused analysis and including both experimental stages in the same model,

which increases power and allows the contrast between the first and second reversal to be identified.

Both species were affected similarly by the treatment, suggesting that the consequences of social

stability have general effects on learning in a foraging context despite species differences in competitive

ability and learning performance (Reichert et al. 2020, 2021). One explanation for our finding that

learning performance was not better in stable social groups is that most individuals learned the task

relatively quickly, potentially limiting the utility of social learning in this system (Reichert et al. 2020).

The somewhat improved learning performance in the unstable treatment in our study could have

occurred because this treatment disrupted social competition and dominance relationships that built up

among individuals attempting to access the same feeder. Our social network metrics do not capture

such relationships because they are based on flock membership, but they are plausible, because high

levels of competition and low dominance ranks are associated with reduced learning performance in

many species (Chalmeau and Gallo 1993; Pravosudov et al. 2003; Langley et al. 2018). Similarly, there is

evidence from multiple species, including great tits, that better problem solvers are poor competitors at

feeders (Cole and Quinn 2012), so the unstable treatment may have enabled those better at problem

solving to realize greater cognitive performance when otherwise negative hierarchical interactions with

known individuals were reduced. However, other studies of learning in wild systems found limited

effects of dominance and social stability (Heinen et al. 2021*a*, 2021*b*; Pitera et al. 2025), and other social

processes caused by our manipulation could also be responsible.

For great tits, we found limited evidence for effects of average flock size on learning speed, but in

general individuals with larger average flock sizes learned more quickly. For blue tits, individuals with

larger average flock size also learned more quickly in the first reversal, but more slowly in the second

reversal. In a previous study of great tits from the same population, rates of innovative problem solving

also increased with flock size (Morand-Ferron and Quinn 2011). Thus, the costs of being in larger groups,

such as increased competition, may be outweighed by other benefits, such as dilution of risk, in addition

to other more subtle benefits including better cognitive performance, at least for some contexts and

cognitive tasks. When we examined the effects of weighted degree, the strength of an individual's social

connections, we found that, at least for the second reversal, great tits with higher weighted degrees

were slower to learn the task. In a previous study, great tits reduced their own foraging to stay with

their mating partner (Firth et al. 2015). Particularly for the challenging second reversal, if similar

behavior occurred in our study where birds were similarly restricted in feeder access, then this tendency

to remain near strong social partners could have slowed down an individual's own learning. Blue tits

were less affected by this variable, perhaps because their cognitive performance is more strongly

affected by general social competition to access feeders.

Effects of sociality on scrounging

Scrounging rates were higher in the stable treatment during the second reversal stage for both species

and, for blue tits, also in the first reversal. These results ran counter to our expectation that individuals

in the unstable treatment would scrounge more because if individuals primarily scrounge from close

social associates (King et al. 2009; Harten et al. 2018), the unstable treatment would have broken up

those associations. However, we previously found that the across-stage correlation between the dyadic

associations between individuals in the social network did not differ between the stable and unstable

treatments (Troisi et al. 2025), so disruption of scrounging associations was probably not responsible for

our findings of more scrounging in the stable treatment. The most likely explanation for higher

scrounging in the stable treatment is that individuals in the unstable treatment were faster to learn to

produce, and therefore spent less time using the scrounging tactic, given the general trade-off between

producing and scrounging (Barnard and Sibly 1981).

For both great tits and blue tits, scrounging rates were lower for birds with larger average flock size in

both reversal stages. These results were surprising both because we had expected larger groups to

provide more opportunities to scrounge, and because individuals in larger groups generally learned

more quickly, which should have provided more opportunity for scroungers. However, in a previous

analysis of this dataset, we found that scrounging and production learning trade-off such that individuals

that learn quickly scrounge less often (Reichert et al. 2021). As we argued above for the effects of the

social stability treatment, the faster learning speed for birds in larger flocks therefore could have

corresponded with reduced benefits of scrounging relative to producing. Thus, the social factors

affecting production learning, in this case potentially beneficial effects of being in large groups, could

have indirectly negatively influenced the expression of an alternative foraging tactic. This explanation

could also apply to the positive relationship between scrounging and weighted degree in the reversals.

Particularly for the second reversal, great tits with higher weighted degree were slower to learn, which

could have induced more scrounging. However, determining the direction of causation is difficult.

Indeed, an alternative explanation is that because effective scrounging is a learned behavior in this

system (Reichert et al. 2021), birds with stronger connections may have had more opportunities to learn

effective scrounging techniques or could have developed more stable producer-scrounger relationships.

This would be consistent with a previous study of great tits in which birds with especially strong

connections were found to forgo production to stay with their mating partner, from whom they

obtained food by scrounging (Firth et al. 2015).

Effects of sociality on foraging rate

Individuals' foraging rates after learning the location of their rewarded feeder were not affected by the

social stability treatment for either species. Foraging rate at feeders is a repeatable trait in this system

(Crates et al. 2016). This individual consistency in behavior may have contributed to our findings that the

overall amount of foraging activity remained relatively stable despite any changes that the social

stability treatments caused to social relationships, overall social structure, and subsequent effects on

interactions at the feeders. Our findings suggest that the effects of living in groups on individual foraging

rates are not influenced by the ability to feed near specific individuals, which are unnecessary for

general benefits of social foraging like shared vigilance and the dilution effect, though we acknowledge

that these effects may have been limited by the relative ease with which food could be obtained post

learning.

Foraging rates were also unaffected by average flock size. A previous study found a similar pattern in

which overall visit lengths and inter-visit intervals were not affected by an experimental manipulation of

density at the feeder (Regan et al. 2022). However, it also found that individuals with stronger social ties

were more likely to visit feeders and that these effects interacted with social group density for some

foraging behaviors (Regan et al. 2022), pointing to the importance of weighted degree in foraging.

Indeed, we found that individuals had a higher foraging rate at our feeders when they had a higher

weighted degree. This effect was somewhat stronger for the initial learning stage, perhaps because at

this early stage scrounging levels were low and the social stability treatment had not yet affected social

connections. One important consideration is that our measure of foraging rate only includes visits to our

experimental feeders. Although feeders are important food sources, great tits have a broad diet in the

winter even when supplemental food is available (Coomes et al. 2025), and this is likely to be the case

for blue tits as well. Thus, individuals may vary in their tendency to visit feeders in the first place, and

this may be related to the strength of their social relationships, which could explain our findings.

Differences across stages and species

A clear outcome of our study is that effects of sociality on foraging varied across the experimental

stages, an initial associative learning task and two subsequent reversal learning tasks. However, this was

only the case for the two cognitively demanding foraging traits, learning speed and scrounging rate

(which is cognitively demanding because individuals learn to scrounge in our system; (Reichert et al.

2021)), but not for foraging rate. This implies that the effects of sociality may be especially pronounced

in cognitively challenging situations. The main difference between the experimental stages was in the

learning task required of the birds, from an initial association to two successive reversals. Therefore, one

explanation for variation across stages is that sociality interacted differently with the cognitive

challenges associated with each task. Initial learning is driven by an associative process between a

feeder and the reward, while reversal learning requires additional cognitive flexibility so that individuals

inhibit their response to the previously rewarded feeder and instead make an association with the new

rewarded feeder (Morand-Ferron et al. 2022). At the same time, some individuals were learning to

become increasingly proficient at scrounging and there were opportunities to socially learn from other

individuals visiting the feeders (Reichert et al. 2021). Some previous studies showed that the effects of

the social environment on cognition depend on the cognitive task (Kogan et al. 2000; Schrijver et al.

2004), and our results may indicate that these findings extend to foraging behaviors involving different

cognitive mechanisms in a wild population. Although further study is needed, our results point to an

effect of sociality on the opportunity to learn alternative foraging tactics, adding to the evidence that

cognition is shaped by social structure (Hoppitt and Laland 2013; Seyfarth and Cheney 2015; Kulahci and

Quinn 2019).

Another potential explanation for our finding of variation in the effects of sociality across stages is that

the overall structural properties of the global social network itself changed. Indeed, we previously

showed in this dataset that average flock size and weighted degree both increased from the initial

learning to the second reversal stages, and the increase in flock size was stronger for the stable

treatment (Troisi et al. 2025). Where we found significant interactions between experimental stage and

social network metrics on foraging behavior, the relationships between sociality and behavior were

generally stronger during the reversal stages, indicating that the increased strength of social

connections led to emergent impacts on foraging. At the same time, we found support for our

prediction that the effects of the social stability treatment on foraging would be stronger in the second

reversal than the first reversal, which indicates that multiple social disruptions have increasingly large

effects on foraging. The current study and the analyses in Troisi et al. (2025) clearly show that the

foraging and social processes were highly dynamic within groups during this experiment. We note that

some of these social processes were likely already in flux prior to the initial learning stage. Our

experimental design involved manipulating feeder placement and access to food to attract birds to the

area and acclimate them to the experimental devices before we began the learning experiments. Social

network traits and flock size varied even across these early stages, indicating that all individuals would

have been experiencing some social instability before our social stability manipulation (Troisi et al.

2025). Indeed, some baseline level of social instability is the norm in these foraging flocks, which are

characterized by a fission-fusion structure (Farine et al. 2015b). Our social stability manipulation

therefore aimed to add additional instability to the social group, and our results indicate that this did

have effects on foraging traits. Nevertheless, social instability can occur at different scales (e.g., at a fine

scale of accessing food at different areas in a plot to a large scale of flock membership), and further

work is needed to determine if social processes operating at different scales have different effects on

foraging.

We calculated the social network based on social ties across species, which is appropriate given the

strong tendency of great tits and blue tits to forage together. Although some social processes like mate

selection will be more strongly affected by within-species social ties (Firth et al. 2018), social foraging

tactics like scrounging and competition at feeders are likely to be affected by all network members, even

heterospecifics. Indeed, we found that the social stability treatment had qualitatively, if not

quantitatively, similar effects on both blue tits and great tits, and the impact of weighted degree and

flock size was also broadly similar on most measures of foraging in both species. These common effects

not only indicate real social ties that extend to heterospecifics but also give evidence for the

effectiveness of our experimental manipulation and the robustness of our results. Our findings also raise

important questions for future work, including whether social effects on foraging fundamentally differ

between single-species and mixed-species flocks and whether the nature of among-species interactions

plays a role.

Conclusions

Both social network position and foraging emerge from a complex interplay of factors including

competition, cognition, individual personality, levels of predation and other environmental

characteristics, yet disentangling these factors in the wild has been difficult due to the lack of

experiments. Our findings that sociality affects multiple aspects of foraging success in a wild system

expand on our knowledge of social foraging and demonstrate the utility of social network metrics for

capturing variation in key behavioral traits. The results demonstrate a series of costs (slower learning in

stable flocks and when weighted degree was higher) and benefits (greater scrounging opportunity with

stable and more connected flocks, and higher foraging rates in flocks with stronger relationships)

associated with greater sociality. However, there is still much progress to be made in identifying the

mechanisms behind these results. Many of our findings suggested a role of competition and so a key

next step will be to better characterize competitive interactions, which is challenging from RFID data but

increasingly possible with new monitoring technologies (Smith and Pinter-Wollman 2021). Our findings

also raise questions of causation: an individual's social position certainly influences its foraging behavior,

but it may also be the case that foraging behavior affects social position (Kulahci and Quinn 2019), for

example if some individuals serve as models for social learning or easy targets for scroungers (Harten et

al. 2018). Expanded efforts to monitor and manipulate the social dynamics of wild foraging flocks are

likely to yield exciting new insights into the factors that drive the expression of functional behaviors.

AUTHOR CONTRIBUTIONS

Conceptualization: M.R., J.F., J.Q.; Data curation: M.R., C.T., J.F.; Formal analysis: M.R., C.T., J.F.; Funding

acquisition: J.Q.; Investigation: M.R.; Methodology: M.R., J.F., J.Q.; Writing – original draft: M.R.; Writing

– review & editing: M.R., C.T., J.F., J.Q.

ACKNOWLEDGEMENTS

Sam Crofts and Keith McMahon assisted with bird ringing and fieldwork. Martin Whitaker helped design

the selective feeders, and Gabrielle Davidson, James Savage and Iván de la Hera Fernández helped with

their construction. Karen Cogan assisted with equipment sourcing and purchasing. Ben Sheldon

provided access to the study system at Wytham Woods and funding to maintain the long-term study.

This research was funded by the European Research Council under the European Union's Horizon 2020

Programme (FP7/2007-2013)/ERC Consolidator Grant 'EVOLECOCOG' Project No. 617509, awarded to

J.L.Q., and by a Science Foundation Ireland ERCSupport Grant 14/ERC/B3118 to J.L.Q.

REFERENCES

Ansmann, I. C., G. J. Parra, B. L. Chilvers, and J. M. Lanyon. 2012. Dolphins restructure social system after

reduction of commercial fisheries. Animal Behaviour 84:575–581.

Aplin, L. M., D. R. Farine, J. Morand-Ferron, and B. C. Sheldon. 2012. Social networks predict patch

discovery in a wild population of songbirds. Proceedings of the Royal Society B: Biological Sciences

279:4199–4205.

Aplin, L. M., J. A. Firth, D. R. Farine, B. Voelkl, R. A. Crates, A. Culina, C. J. Garroway, et al. 2015.

Consistent individual differences in the social phenotypes of wild great tits, *Parus major*. Animal

Behaviour 108:117–127.

Aplin, L. M., and J. Morand-Ferron. 2017. Stable producer–scrounger dynamics in wild birds: Sociability

and learning speed covary with scrounging behaviour. Proceedings of the Royal Society B: Biological

Sciences 284:20162872.

Ashton, B. J., A. R. Ridley, E. K. Edwards, and A. Thornton. 2018. Cognitive performance is linked to group

size and affects fitness in Australian magpies. Nature 554:364–367.

Ashton, B. J., A. Thornton, and A. R. Ridley. 2019. Larger group sizes facilitate the emergence and spread

of innovations in a group-living bird. Animal Behaviour 158:1–7.

Avarguès-Weber, A., and L. Chittka. 2014. Local enhancement or stimulus enhancement? Bumblebee

social learning results in a specific pattern of flower preference. Animal Behaviour 97:185–191.

Barnard, C. J., and R. M. Sibly. 1981. Producers and scroungers: A general model and its application to

captive flocks of house sparrows. Animal Behaviour 29:543–550.

Bates, D., M. Maechler, B. M. Bolker, and S. Walker. 2015. Fitting linear mixed-effects models using

lme4. Journal of Statistical Software 67:1–48.

Beauchamp, G. 1998. The effect of group size on mean food intake rate in birds. Biological Reviews

73:449–472.

———. 2005. Does group foraging promote efficient exploitation of resources? Oikos 111:403–407.

Beauchamp, G., M. Belisle, and L.-A. Giraldeau. 1997. Influence of conspecific attraction on the spatial

distribution of learning foragers in a patchy habitat. The Journal of Animal Ecology 66:671–682.

Beck, K. B., C. E. Regan, K. McMahon, S. Crofts, E. F. Cole, J. A. Firth, and B. C. Sheldon. 2024.

Experimental manipulation of population density in a wild bird alters social structure but not patch

discovery rate. Animal Behaviour 209:95–120.

Beltrão, P., A. C. R. Gomes, and G. C. Cardoso. 2022. Collective foraging: Experimentally increased

competition decreases group performance exploiting a permanent resource. Functional Ecology

36:1796–1805.

Cairns, S. J., and S. J. Schwager. 1987. A comparison of association indices. Animal Behaviour 35:1454–

1469.

Cantor, M., L. M. Aplin, and D. R. Farine. 2020. A primer on the relationship between group size and

group performance. Animal Behaviour 166:139–146.

Cauchoix, M., G. B. Jason, A. Biganzoli, J. Briot, V. Guiraud, N. El Ksabi, D. Lieuré, et al. 2022. The

OpenFeeder⍰: A flexible automated RFID feeder to measure interspecies and intraspecies differences in

cognitive and behavioural performance in wild birds. Methods in Ecology and Evolution 13:1955–1961.

Chalmeau, R., and A. Gallo. 1993. Social constraints determine what is learned in the chimpanzee.

Behavioural Processes 28:173–179.

Clark, C. W., and M. Mangel. 1986. The evolutionary advantages of group foraging. Theoretical

Population Biology 30:45–75.

Cole, E. F., and J. L. Quinn. 2012. Personality and problem-solving performance explain competitive

ability in the wild. Proceedings of the Royal Society B: Biological Sciences 279:1168–1175.

Coomes, J. R., J. Cuff, M. Reichert, G. L. Davidson, W. O. C. Symondson, and J. Quinn. 2025. Spatio-

temporal variation in diet among age and sex cohorts of a model generalist bird species, the great tit

Parus major: new insights revealed by DNA metabarcoding. Ecology and Evolution 15:e71565.

Courchamp, F., and D. W. Macdonald. 2001. Crucial importance of pack size in the African wild dog

Lycaon pictus. Animal Conservation 4:169–174.

Crates, R. A., J. A. Firth, D. R. Farine, C. J. Garroway, L. R. Kidd, L. M. Aplin, R. Radersma, et al. 2016.

Individual variation in winter supplementary food consumption and its consequences for reproduction

in wild birds. Journal of Avian Biology 47:678–689.

Cresswell, W. 1994. Flocking is an effective anti-predation strategy in redshanks, *Tringa totanus*. Animal

Behaviour 47:433–442.

———. 1997. Interference competition at low competitor densities in blackbirds *Turdus merula*. The

Journal of Animal Ecology 66:461–471.

DeLong, J. P., S. F. Uiterwaal, and A. I. Dell. 2021. Trait-based variation in the foraging performance of

individuals. Frontiers in Ecology and Evolution 9:649542.

Dias, R. I. 2006. Effects of position and flock size on vigilance and foraging behaviour of the scaled dove

Columbina squammata. Behavioural Processes 73:248–252.

Drewe, J. A. 2010. Who infects whom? Social networks and tuberculosis transmission in wild meerkats.

Proceedings of the Royal Society B: Biological Sciences 277:633–642.

Ekman, J. 1989. Ecology of non-breeding social systems of *Parus*. The Wilson Bulletin 101:263–288.

Farine, D. R., L. M. Aplin, C. J. Garroway, R. P. Mann, and B. C. Sheldon. 2014. Collective decision making

and social interaction rules in mixed-species flocks of songbirds. Animal Behaviour 95:173–182.

Farine, D. R., L. M. Aplin, B. C. Sheldon, and W. Hoppitt. 2015*a*. Interspecific social networks promote

information transmission in wild songbirds. Proceedings of the Royal Society B 282:20142804.

Farine, D. R., J. A. Firth, L. M. Aplin, R. A. Crates, A. Culina, C. J. Garroway, C. A. Hinde, et al. 2015*b*. The

role of social and ecological processes in structuring animal populations: a case study from automated

tracking of wild birds. Royal Society Open Science 2:150057.

Farine, D. R., C. J. Garroway, and B. C. Sheldon. 2012. Social network analysis of mixed-species flocks:

Exploring the structure and evolution of interspecific social behaviour. Animal Behaviour 84:1271–1277.

Farine, D. R., and B. C. Sheldon. 2015. Selection for territory acquisition is modulated by social network

structure in a wild songbird. Journal of Evolutionary Biology 28:547–556.

Firth, J. A., E. F. Cole, C. C. Ioannou, J. L. Quinn, L. M. Aplin, A. Culina, K. McMahon, et al. 2018.

Personality shapes pair bonding in a wild bird social system. Nature Ecology & Evolution 2:1696–1699.

Firth, J. A., and B. C. Sheldon. 2015. Experimental manipulation of avian social structure reveals

segregation is carried over across contexts. Proceedings of the Royal Society B 282:20142350.

———. 2016. Social carry-over effects underpin trans-seasonally linked structure in a wild bird

population. Ecology Letters 19:1324–1332.

Firth, J. A., B. C. Sheldon, and D. R. Farine. 2016. Pathways of information transmission among wild

songbirds follow experimentally imposed changes in social foraging structure. Biology Letters

12:20160144.

Firth, J. A., B. Voelkl, D. R. Farine, and B. C. Sheldon. 2015. Experimental evidence that social

relationships determine individual foraging behavior. Current Biology 25:3138–3143.

Focardi, S., and E. Pecchioli. 2005. Social cohesion and foraging decrease with group size in fallow deer

(*Dama dama*). Behavioral Ecology and Sociobiology 59:84–91.

Giraldeau, L. A., and T. Caraco. 2000. Social Foraging Theory. Princeton University Press, Princeton.

Godfrey, S. S., C. M. Bull, R. James, and K. Murray. 2009. Network structure and parasite transmission in

a group living lizard, the gidgee skink, *Egernia stokesii*. Behavioral Ecology and Sociobiology 63:1045–

1056.

Grueter, C. C., A. M. Robbins, D. Abavandimwe, V. Vecellio, F. Ndagijimana, T. S. Stoinski, and M. M.

Robbins. 2018. Quadratic relationships between group size and foraging efficiency in a herbivorous

primate. Scientific Reports 8:16718.

Hall, C. L., and D. L. Kramer. 2008. The economics of tracking a changing environment: competition and

social information. Animal Behaviour 76:1609–1619.

Harten, L., Y. Matalon, N. Galli, H. Navon, R. Dor, and Y. Yovel. 2018. Persistent producer-scrounger

relationships in bats. Science Advances 4:e1603293.

Heathcote, R. J. P., S. K. Darden, D. W. Franks, I. W. Ramnarine, and D. P. Croft. 2017. Fear of predation

drives stable and differentiated social relationships in guppies. Scientific Reports 7:41679.

Heinen, V. K., L. M. Benedict, A. M. Pitera, B. R. Sonnenberg, E. S. Bridge, and V. V. Pravosudov. 2021*a*.

Social dominance has limited effects on spatial cognition in a wild food-caching bird. Proceedings of the

Royal Society B: Biological Sciences 288:20211784.

Heinen, V. K., A. M. Pitera, B. R. Sonnenberg, L. M. Benedict, E. S. Bridge, D. R. Farine, and V. V.

Pravosudov. 2021*b*. Food discovery is associated with different reliance on social learning and lower

cognitive flexibility across environments in a food-caching bird. Proceedings of the Royal Society B:

Biological Sciences 288:20202843.

Hoppitt, W., and K. N. Laland. 2013. Social Learning. Princeton University Press, Princeton.

Izquierdo, A., J. L. Brigman, A. K. Radke, P. H. Rudebeck, and A. Holmes. 2017. The neural basis of

reversal learning: An updated perspective. Neuroscience 345:12–26.

Jones, T. B., L. M. Aplin, I. Devost, and J. Morand-Ferron. 2017. Individual and ecological determinants of

social information transmission in the wild. Animal Behaviour 129:93–101.

Kendal, R. L., I. Coolen, Y. van Bergen, and K. N. Laland. 2005. Trade-offs in the adaptive use of social and

asocial learning. Advances in the Study of Behavior 35:333–379.

King, A. J., N. J. B. Isaac, and G. Cowlishaw. 2009. Ecological, social, and reproductive factors shape

producer–scrounger dynamics in baboons. Behavioral Ecology 20:1039–1049.

Kogan, J. H., P. W. Franklandand, and A. J. Silva. 2000. Long-term memory underlying hippocampus-

dependent social recognition in mice. Hippocampus 10:47–56.

Krause, J., R. James, D. W. Franks, and D. P. Croft, eds. 2015. Animal Social Networks. Oxford University

Press, Oxford.

Krause, J., and G. Ruxton. 2002. Living in Groups. Oxford University Press, Oxford.

Kulahci, I. G., and J. L. Quinn. 2019. Dynamic relationships between information transmission and social

connections. Trends in Ecology and Evolution 34:545–554.

Kulahci, I. G., D. I. Rubenstein, T. Bugnyar, W. Hoppitt, N. Mikus, and C. Schwab. 2016. Social networks

predict selective observation and information spread in ravens. Royal Society Open Science 3:160256.

Kurvers, R. H. J. M., K. van Oers, B. a Nolet, R. M. Jonker, S. E. van Wieren, H. H. T. Prins, and R. C.

Ydenberg. 2010. Personality predicts the use of social information. Ecology letters 13:829–837.

Lambert, C. T., and L. M. Guillette. 2021. The impact of environmental and social factors on learning

abilities: a meta-analysis. Biological Reviews 96:2871–2889.

Langley, E. J. G., J. O. van Horik, M. A. Whiteside, and J. R. Madden. 2018. Group social rank is associated

with performance on a spatial learning task. Royal Society Open Science 5:171475.

Lenth, R. 2022. emmeans: estimated marginal means, aka least-squares means version 1.8.3.

Lima, S. L., P. A. Zollner, and P. A. Bednekoff. 1999. Predation, scramble competition, and the vigilance

group size effect in dark-eyed juncos (*Junco hyemalis*). Behavioral Ecology and Sociobiology 46:110–116.

Lüdecke, D., M. Ben-Shachar, I. Patil, P. Waggoner, and D. Makowski. 2021. performance: An R Package

for Assessment, Comparison and Testing of Statistical Models. Journal of Open Source Software 6:3139.

MacNulty, D. R., A. Tallian, D. R. Stahler, and D. W. Smith. 2014. Influence of group size on the success of

wolves hunting bison. PLoS ONE 9:e112884.

Mady, R. P., and D. T. Blumstein. 2017. Social security: are socially connected individuals less vigilant?

Animal Behaviour 134:79–85.

McDonald, D. B., and D. Shizuka. 2013. Comparative transitive and temporal orderliness in dominance

networks. Behavioral Ecology 24:511–520.

McMahon, K., N. M. Marples, L. G. Spurgin, H. M. Rowland, B. C. Sheldon, and J. A. Firth. 2024. Social

network centrality predicts dietary decisions in a wild bird population. iScience 27:109581.

Milligan, N. D., R. Radersma, E. F. Cole, and B. C. Sheldon. 2017. To graze or gorge: consistency and

flexibility of individual foraging tactics in tits. Journal of Animal Ecology 86:826–836.

Monk, C. T., M. Barbier, P. Romanczuk, J. R. Watson, J. Alós, S. Nakayama, D. I. Rubenstein, et al. 2018.

How ecology shapes exploitation: a framework to predict the behavioural response of human and

animal foragers along exploration–exploitation trade-offs. Ecology Letters 21:779–793.

Morand-Ferron, J., L.-A. Giraldeau, and L. Lefebvre. 2007. Wild Carib grackles play a producer scrounger

game. Behavioral Ecology 18:916–921.

Morand-Ferron, J., and J. L. Quinn. 2011. Larger groups of passerines are more efficient problem solvers

in the wild. Proceedings of the National Academy of Sciences 108:15898–15903.

Morand-Ferron, J., M. S. Reichert, and J. L. Quinn. 2022. Cognitive flexibility in the wild: Individual

differences in reversal learning are explained primarily by proactive interference, not by sampling

strategies, in two passerine bird species. Learning and Behavior 50:153–166.

Morand-Ferron, J., G. M. Wu, and L. A. Giraldeau. 2011. Persistent individual differences in tactic use in a

producer-scrounger game are group dependent. Animal Behaviour 82:811–816.

Murphy, D., H. S. Mumby, and M. D. Henley. 2020. Age differences in the temporal stability of a male

African elephant (*Loxodonta africana*) social network. Behavioral Ecology 31:21–31.

Nachappa, P., D. C. Margolies, J. R. Nechols, and T. J. Morgan. 2010. Response of a complex foraging

phenotype to artificial selection on its component traits. Evolutionary Ecology 24:631–655.

Oom, S. P., J. A. Beecham, C. J. Legg, and A. J. Hester. 2004. Foraging in a complex environment: from

foraging strategies to emergent spatial properties. Ecological Complexity 1:299–327.

Pitcher, T. J., A. E. Magurran, and I. J. Winfield. 1982. Fish in larger shoals find food faster. Behavioral

Ecology and Sociobiology 10:149–151.

Pitera, A. M., V. K. Heinen, B. R. Sonnenberg, L. M. Benedict, E. S. Bridge, and V. V. Pravosudov. 2025.

Social group membership does not facilitate spatial learning of fine-scale resource locations. Behavioral

Ecology and Sociobiology 79:104.

Pravosudov, V. V., S. P. Mendoza, and N. S. Clayton. 2003. The relationship between dominance,

corticosterone, memory, and food caching in mountain chickadees (*Poecile gambeli*). Hormones and

Behavior 44:93–102.

Psorakis, I., S. J. Roberts, I. Rezek, and B. C. Sheldon. 2012. Inferring social network structure in

ecological systems from spatio-temporal data streams. Journal of the Royal Society Interface 9:3055–

3066.

Psorakis, I., B. Voelkl, C. J. Garroway, R. Radersma, L. M. Aplin, R. A. Crates, A. Culina, et al. 2015.

Inferring social structure from temporal data. Behavioral Ecology and Sociobiology 69:857–866.

R Development Core Team. 2024. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria.

Regan, C. E., K. B. Beck, K. Mcmahon, S. Crofts, J. A. Firth, and B. C. Sheldon. 2022. Social phenotype-

dependent selection of social environment in wild great and blue tits : an experimental study.

Proceedings of the Royal Society B 289:20221602.

Reichert, M. S., S. J. Crofts, G. L. Davidson, J. A. Firth, I. G. Kulahci, and J. L. Quinn. 2020. Multiple factors

affect discrimination learning performance, but not between-individual variation, in wild mixed-species

flocks of birds. Royal Society Open Science 7:192107.

Reichert, M. S., J. Morand-Ferron, I. G. Kulahci, J. A. Firth, G. L. Davidson, S. J. Crofts, and J. L. Quinn.

2021. Cognition and covariance in the producer-scrounger game. Journal of Animal Ecology 90:2497–

2509.

Rieucau, G., and L.-A. Giraldeau. 2009*a*. Group size effect caused by food competition in nutmeg

mannikins (*Lonchura punctulata*). Behavioral Ecology 20:421–425.

———. 2009*b*. Persuasive companions can be wrong: the use of misleading social information in nutmeg

mannikins. Behavioral Ecology 20:1217–1222.

Sansom, A., W. Cresswell, J. Minderman, and J. Lind. 2008. Vigilance benefits and competition costs in

groups: do individual redshanks gain an overall foraging benefit? Animal Behaviour 75:1869–1875.

Schakner, Z. A., M. B. Petelle, M. J. Tennis, B. K. Van Der Leeuw, R. T. Stansell, and D. T. Blumstein. 2017.

Social associations between California sea lions influence the use of a novel foraging ground. Royal

Society Open Science 4:160820.

Schrijver, N. C. A., P. N. Pallier, V. J. Brown, and H. Würbel. 2004. Double dissociation of social and

environmental stimulation on spatial learning and reversal learning in rats. Behavioural Brain Research

152:307–314.

Seyfarth, R. M., and D. L. Cheney. 2015. Social cognition. Animal Behaviour 103:191–202.

Sheppard, C. E., R. Inger, R. A. McDonald, S. Barker, A. L. Jackson, F. J. Thompson, E. I. K. Vitikainen, et al.

2018. Intragroup competition predicts individual foraging specialisation in a group-living mammal.

Ecology Letters 21:665–673.

Silk, J. B. 2007. The adaptive value of sociality in mammalian groups. Philosophical Transactions of the

Royal Society B: Biological Sciences 362:539–559.

Slagsvold, T., and K. L. Wiebe. 2011. Social learning in birds and its role in shaping a foraging niche.

Philosophical transactions of the Royal Society of London. Series B, Biological sciences 366:969–977.

Smith, J. E., and N. Pinter-Wollman. 2021. Observing the unwatchable: Integrating automated sensing,

naturalistic observations and animal social network analysis in the age of big data. Journal of Animal

Ecology 90:62–75.

Snijders, L., S. Krause, A. N. Tump, M. Breuker, C. Ortiz, S. Rizzi, I. W. Ramnarine, et al. 2021. Causal

evidence for the adaptive benefits of social foraging in the wild. Communications Biology 4:94.

Stevens, D. W., J. S. Brown, and R. C. Ydenberg, eds. 2007. Foraging Behavior and Ecology. University of

Chicago Press, Chicago.

Thornton, A., and A. Malapert. 2009. Experimental evidence for social transmission of food acquisition

techniques in wild meerkats. Animal Behaviour 78:255–264.

Troisi, C. A., J. A. Firth, S. J. Crofts, G. L. Davidson, M. S. Reichert, and J. L. Quinn. 2025. Effects of fine-

scale changes in resource access and social stability on the sociality of foraging flocks of wild birds.

Animal Behaviour 221:123071.

Vásquez, R. A., and A. Kacelnik. 2000. Foraging rate versus sociality in the starling *Sturnus vulgaris*.

Proceedings of the Royal Society of London. Series B: Biological Sciences 267:157–164.

Voelkl, B., J. A. Firth, and B. C. Sheldon. 2016. Nonlethal predator effects on the turn-over of wild bird

flocks. Scientific Reports 6:33476.

Waite, T. A., and K. L. Field. 2007. Foraging with others: Games social foragers play. Pages 331–362 *in* D.

W. Stephens, J. S. Brown, and R. C. Ydenberg, eds. Foraging. Behavior and Ecology. University of Chicago

Press, Chicago.

Weber, N., S. P. Carter, S. R. X. Dall, R. J. Delahay, J. L. McDonald, S. Bearhop, and R. A. McDonald. 2013.

Badger social networks correlate with tuberculosis infection. Current Biology 23:R915–R916.

Whitehead, H. 2008. Analyzing animal societies: Quantitative methods for vertebrate social analysis. The

University of Chicago Press.