TITLE: Bridging the gap of late-gestation nephrogenesis using a non-human primate model

AUTHORS: Kairavee Thakkar[1,2], Sunitha Yarlagadda[3], Lyan Alkhudairy[3], Andrew Potter[4], Konrad Thorner[1], Praneet Chaturvedi[6], Kyle W. McCracken[3,5,6], Nathan Salomonis[1,5], Raphael Kopan[5,6], Meredith P Schuh[3,5,6]

[1]Division of Biomedical Informatics, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio, USA;

[2]Department of Pharmacology and Physiology, University of Cincinnati College of Medicine, Cincinnati, Ohio, USA

[3]Division of Nephrology and Hypertension, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio, USA;

[4]The Heart Institute, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio, USA

[5]Department of Pediatrics, University of Cincinnati College of Medicine, Cincinnati, Ohio, USA;

[6]Division of Developmental Biology, University of Cincinnati College of Medicine, Cincinnati, Ohio, USA.

Correspondence: Meredith P Schuh, MD,3333 Burnet Ave, MLC 7022 Cincinnati, OH 45229. E-mail: Meredith.Schuh@cchmc.org

Word count abstract: 259

Word count text: 4898

Number of figures: 6

Key points:

- Single-cell RNA sequencing analysis of late-gestation primate nephron progenitor cells (NPCs) identifies rhesus-specific self-renewing NPCs that are not observed in mid-gestation humans.

- Differential expression and ligand-receptor interactions suggest a compositional shift in WNT signaling within the naive NPC population during LBN.

- *FZD7* localized to the connecting segment and *SEMA3D* was identified in the ureteric bud (UB) tips of late-gestation primates.

Abstract:

Background: Prematurity is associated with low nephron endowment and an increased risk of chronic kidney disease. Human nephrogenesis is complete at 34-36 weeks gestation, with 60% of nephrons forming during the third trimester through lateral branch nephrogenesis (LBN). We hypothesized that a differentiated but dividing population of nephron progenitor cells (NPCs) would contribute to the amplification of nephrons in late gestation.

Methods: Single-cell RNA-sequencing (scRNA-Seq) was performed on cortically-enriched fetal rhesus kidneys (n=9) from late second trimester and third trimester during LBN. This data was integrated with publicly available human scRNA-seq datasets from 8-18 weeks gestation kidneys (n=8) using state-of-the-art bioinformatics pipelines. Differentially expressed genes and ligand-receptor interactions were assessed and validated using RNAScope™ on human and rhesus archival tissue.

Results: scRNA-Seq of 64,782 rhesus cells revealed 37 transcriptionally distinct cell populations, including 7,879 rhesus NPCs. Pseudotime analyses identified a late gestation-specific lineage branch of differentiated NPC in rhesus that was not observed in mid-gestation humans. Differential expression analyses identified increased *SFRP1, FZD4*, and *TLE2* and decreased *FZD7*, *SHISA2*, *SHISA3*, and *TLE4* within the late-gestation rhesus NPC compared to mid-gestation human NPC and increased SEMA3D within the rhesus UB tip, suggesting a compositional shift in WNT and SEMA signaling components within the naive NPC population during LBN.

Conclusion: The rhesus macaque uniquely enables molecular studies of late-gestation primate nephrogenesis. Our study suggests the hypothesis that a transitional state of self-renewing NPC supported by compositional shifts in key pathways may underlie the switch from branching phase nephrogenesis to lateral branch nephrogenesis and support ongoing nephron formation in late gestation.

INTRODUCTION

The period of new nephron formation, or nephrogenesis, in human kidney development typically extends through 34-36 weeks gestation (WG), with 60% of nephrons forming after 23 WG through a process termed lateral branch nephrogenesis (LBN).[1] Infants born prematurely must undergo LBN postnatally and are at the low end of nephron endowment and at high risk for chronic kidney disease (CKD) later in life.[2-10] Although preterm infants continue LBN postnatally, this process is associated with accelerated loss of SIX1 and RET expression in niche progenitors and earlier (by ~14 days) cessation of nephrogenesis.[11] In all mammals, nephrons are formed in a process associated with bifurcating branching of the ureteric bud (UB), which we will refer to as branching phase nephrogenesis (BpN) **(Fig 1A)**. In mice, BpN is the only mode of nephron formation and persists until postnatal day 2 (P2), at which point the progenitor pool differentiates en masse to amplify the nephron/tip ratio 2-5-fold and nephrogenesis is completed.[12, 13] In contrast, human nephrogenesis transitions through three stages, from BpN (5-15 WG) to the short arcading phase (15-22 WG) **(Fig 1B)** to LBN (23-36 WG), where an elongating, non-branching UB tip induces nephrons, each directly connecting to the ureteric stalk after induction **(Fig 1C)**.[14-17] BpN generates 33,000 human nephrons through 15 cycles of bifurcative branching,[14] while arcading[15, 16, 18] and LBN[14-17] combine to create additional nephrons at each of the 33,000 terminal branch tips to bring the final nephron count between 200,000 and 2.7 million.[14, 19] While mouse models have been instrumental in elucidating the fundamental paradigms that control nephron progenitor cell (NPC) behavior and nephron formation, which appear to be highly conserved in the earliest stages of human kidney organogenesis,[20] they cannot provide insight into this critical LBN period of development that produces the majority of nephrons in humans and is aberrantly regulated in babies born preterm, affecting approximately 10% of all births or over 13 million infants born annually[21].

The rhesus macaque (*Macaca mulatta*) gestational period is 165 days, and that in like humans, can be divided into three trimesters,[22] with LBN present in late gestation.[17] Given these similarities, we established the rhesus as an LBN model and identified age-dependent changes within the late gestation nephron progenitor cells[17] compared to early/mid-gestation human nephrogenesis **(Fig 1)**[23-25] and the fully differentiated kidney. However, the mechanisms of LBN, including UB elongation, maintenance of NPCs, and nephron induction, remain uncharacterized at the molecular level. Specifically, it is unclear how self-renewing NPCs are maintained in a

niche, comprised of a single non-branching UB tip, despite a known change in transcriptional signature[17] and an increase in WNT signaling over time[26].

In this study, we used integrative single-cell transcriptomics to perform the first comprehensive and high-resolution comparison of nephrogenic niche progenitors during LBN and BpN to investigate how nephron amplification occurs independent of branching in the primate-specific phases of nephron production. Our data support that a transitional self-renewing NPC, which exhibits features classically associated with an early induced state, contribute to ongoing nephron formation but also can also be maintained within the niche through changes in the signaling milieu. Potential compositional changes in the WNT and SEMA3 signaling pathways between the NPC and UB may retain these cells that would otherwise differentiate at earlier stages of kidney development.

METHODS

Animals

All primate procedures were performed at the California National Primate Research Center at University of California Davis according to the Institutional Animal Care and Use Committee approved protocol #20330 awarded to Claire Chougnet. Fresh kidneys from control animals euthanized in accordance with the protocol were removed in necropsy and shipped overnight on ice in histidine-tryptophan-ketoglutarate (HTK) solution. Kidney tissue was either prepared for single cell RNA sequencing as described below or fixed in 4% PFA or NBF and embedded in paraffin.

Rhesus single-cell RNA sequencing analyses

For all rhesus samples, count matrices from raw FASTQ files were generated using cellranger (v6.0.1) and rhesus reference genome (Mmul assembly 10). To remove ambient mRNA, we applied the SoupX protocol on the count matrices with automatic calculation of the contamination fraction. Post ambient mRNA removal, we used DoubletFinder to remove doublet cells using the parameters recommended by the authors of DoubletFinder. Rhesus genes were translated to human orthologs (using hg19, Ensmbl release 75). The intersection of genes observed between the human scRNA-seq datasets (see the next section in Methods) and the human orthologs observed in rhesus scRNA-seq datasets were considered for all downstream analyses.

Sample preparation and sequencing, cell type annotation, and subclustering can be found in the supplemental methods

Human single-cell RNA sequencing

Count matrices of the human scRNA-seq datasets were downloaded from GSE114530, GSE112570, and GSE102596. We filtered the count matrices for only the cells that had a cell type annotation in the original publication. To keep cross-species comparisons and analyses consistent, we also filtered the count matrices to only those genes that were also seen in the human ortholog genes from the rhesus samples. Additional information can be found in the supplement methods

Cross-species analyses of the early transitional NPCs

Differential gene expression analysis

Differentially expressed genes of NPCs and UB C6 in rhesus as compared to those in the human dataset were determined using the software cellHarmony, which calls a moderated t-test. Log-normalized gene expression matrices from human and rhesus were provided as the input. Genes with fold change greater than 1.2 and FDR-corrected p-value less than 0.05 were considered differentially expressed.

Lineage trajectory inference, pseudotime analysis, and Cell-cell signaling interactions

Pseudotime analysis was done separately on the rhesus and human datasets using Monocle2.

Cell-cell signaling interactions were determined individually for the human and rhesus datasets using the R package CellChat (v2.1.2). Additional information can be found within the supplemental methods.

RESULTS

Lateral branch nephrogenesis in the human and rhesus is characterized by elongation of a single active ureteric tip that induces nephrons without branching.

We previously evaluated the morphology of third-trimester (~130 days gestation (dg)) rhesus kidney, which is equivalent to 32 weeks gestation in the human.[17] However, despite identifying changes within the nephrogenic niche, we were unable to identify signaling pathways in the original dataset due to the sparsity of NPCs nearing nephrogenesis cessation at ~130dg (316 NPCs). In this study, we obtained rhesus fetal kidney tissue from three donors at 104-108dg (~25-26 WG in the human or late second trimester). Tissue was cleared and 3D rendering

of the nephrogenic niche was assembled as previously described (**Fig 1D-G**).[17] Like the ~130dg rhesus samples, the new samples demonstrated rosette-like organization of the nephrogenic niche as well as LBN morphology, but with higher numbers of NPCs (**Fig 1D, E**). LBN can be identified by the SIX2+ NPC cap associated with a single UB tip, with CDH1+ lateral branches extending from the ureteric stalk below (**Fig 1G blue arrow**). However, our previous study identified that rare bifurcating branching events still occurred at late gestation but occurred prior to 130dg. In this earlier LBN period, we identified rare ongoing BpN (**Fig 1G yellow arrow**), highlighting that this gestational age is suitable for capturing the LBN transition.

To investigate how nephron induction occurs during LBN in the absence of UB branching, we performed morphologic characterization of both human (25-26 WG) and rhesus (104-108 dg) fetal kidneys (**Fig 2B-C**). Our findings showed that the process is stereotypic and highly conserved between the two species, leading us to develop the model summarized in Fig 2A. At the leading distal edge of the epithelium, RET expression across the bulbous UB tip is spread to cover all contact with the asymmetrically located SIX1+ cells. (**Fig 2A-C; 1**). RET also extended below the UB tip, where its protein expression pattern followed and mirrored locations associated with induction and formation of the pre-tubular aggregate and renal vesicle (**Fig 2A-C; 2-3**). The site of distal nephron fusion occurred at the RET+ UB tip (**Fig 2A-C; 4**). The UB tip then extended past this fusion site with the new nephron as it continued to grow as an unbranched elongating tubule, and it continuously maintained close contact with the undifferentiated NPCs (**Fig 2 A-C; 5**).

Single-cell RNA-sequencing reveals late-gestation specific progenitor cell populations undergoing lateral branch nephrogenesis

Overall, the morphologic pattern of nephron induction and formation during LBN appeared similar to that observed during early stages of BpN, with the major exception being that nephrogenesis was uncoupled from UB branching. We hypothesized that comprehensive molecular characterization of the niche progenitors through transcriptomic studies would uncover the mechanisms underlying this distinct behavior. We supplemented our scRNA-seq dataset of four fetal rhesus kidneys at ~130dg[17] with five additional rhesus samples, including two samples at 105-108dg, for a total of nine cortically-enriched fetal kidney samples from distinct donors (**Supplemental Table 1**). We identified 37 transcriptionally distinct cell clusters from 64,782 cells using our previous cluster annotations as reference (**Supplemental Fig 1A**, **Supplemental Table 2, 3**).[17] We then

integrated these data with available human fetal kidney datasets (GSE114530[23], GSE112570[24], and GSE102596[25]) from seven samples (representing BpN and arcading), consisting of 29,484 cells, using Harmony (**Fig 3A, Supplemental Fig 1B, Supplemental Table 4**).[27] Four clusters (c25, c26, c28, c29) were marked with definitive NPC markers such as *SIX1*, *SIX2*, *CITED1* and *MEOX1*, with cluster 25 containing the naive NPCs (in rhesus, n = 7879 cells; in human, n=9434 cells) (**Fig 3B**). UB markers identifying clusters 11 and 13 included *WNT11*, *WNT9B*, *GATA3*, *ADAMTS18*, and *ELF5* (**Supplemental Fig 1C**). To specifically interrogate and resolve the niche progenitor states, we subclustered the four NPC and two UB progenitor clusters using ICGS2, an unsupervised non-negative matrix factorization-based clustering pipeline, which yielded 8 subclusters of NPCs and 6 subclusters of UB progenitors (**Fig 3C**, **Supplemental Table 5, 6**). NPC C4 represented the most naive subcluster, with high expression of the naive NPC markers, including *SIX1, SIX2, MEOX1, EYA1, CITED1, ITGA8, and TMEM100*. Clusters C8 and C3 represented proliferating NPC (*TYMS*, *PCNA*, *TOP2A*, *CENPF*, *CENPE*, *CDC20*), while cluster C7 presented primed NPC (*DLK1, FAM132A, AKAP12, SNAI2*) and cluster C2 contained further differentiated NPC markers (*CYSTM1*, *FXYD6*, *SYTL2*) (**Fig 3D, Supplemental Table 7**).

To reduce the likelihood that comparative analyses between rhesus and human would be confounded by committed NPC, we filtered the NPCs to include those that expressed at least two of the NPC markers: *SIX1*, *SIX2*, and *ITGA8 (*referred as enriched NPC, or eNPC, hereon*).* In rhesus, 80% of these 2,719 eNPC came from the 105-108dg samples (**Supplemental Fig 2, Supplemental Table 5, 6**). *CITED1* was not chosen as a marker for filtering due to its relatively broad expression throughout the rhesus cells (**Fig 3B**). After filtering for eNPC, 64.5% (previously 31.5%) and 93.9% (previously 55.8%) of the cells were from NPC C4 in rhesus and human, respectively (**Supplemental Fig 2**). To investigate the characteristics of the removed human C4 cells, we performed differential gene expression of eNPCs compared to the removed cells. We noted increased expression of *SIX1*, *SIX2,* and *ITGA8* in the eNPCs, supporting this subset as the most naïve progenitors while maintaining a sufficient number of cells in C4. Cells removed demonstrated upregulation in genes related to heat shock responses (HSPA1A, HSPA1B, HSPB1, HSP90AA1, TCIM), potentially representing differences in dissociation protocols. (**Supplemental Table 7**). Interestingly, we noted three subclusters (C1, C5, C6) in the unfiltered dataset that were rhesus predominant (**Supplemental Fig 2A**), and they all largely remained after filtering, supporting their identification as late-gestation specific progenitors. The marker genes characterizing

these three clusters post-filtering include *SPC25* (C1), *UTRN*, *SFRP2*, *DUSP6*, *EMX2*, *DLL1*, *ANXA4*, *FN1*, *CYSTM1*, *FXYD5* (C5), and *MAL* and *FBLN2* (C6) (**Fig 3D, Supplemental Table 7**).

Late-gestation NPC show a distinct trajectory from mid-gestation human NPC

To orthogonally confirm that C4 cluster represented the most naive NPC state and to delineate the differentiation trajectories using an unsupervised approach, we used Monocle[29] to perform pseudotime analysis of the eNPCs at mid gestation (human) and late gestation (rhesus) (**Fig 4**). In the mid-gestation human (**Fig 4A**), the first and only branch splits the NPC C4 into more differentiated NPC C4 (left branch) and NPC C8 (right branch), a proliferative NPC cluster (**Supplemental Table 7**). NPC C2 and C7 overlap with the left branch, while C1 and C3, the two most differentiated NPC, overlap with the right branch. Branch points along the constructed cell trajectory mark a suggested split into two or more distinct cell fates. Despite having a comparable number of C4 NPCs in rhesus and human, the pseudotime analysis identified two branch points in late-gestation rhesus NPCs compared to only one branch in the mid-gestation NPC (**Fig 4B**). In rhesus, the initial lineage branch point (Branch 1) creates two NPC populations: the right branch containing a subset of differentiated NPCs (NPC C4 and C8) found only in the late-gestation rhesus and the left "root" also containing C8 cells in addition to naïve progenitors (C4). Similar to branch 2 in the human dataset, the less differentiated left arm (Branch 2) either gives rise to NPC C7 (induced NPC), overlapping with NPC C4 and C8 (proliferating NPC), or to C1 and C3 (right arm post Branch 2), characterized by markers of cell cycle. Clusters C2, C5, and C6, found primarily in rhesus, represent a mixed population of both terminal branches, consisting of early differentiated NPC (C2), primed NPC (C6: *HEY1, PLOD2)*, markers of proximal (SFRP2, HES4, EMID1 for C5 and JAG1, CLDN1, GRIN2A for C6) and distal nephron (C5: SFRP2, HES4, EMID1)

To further understand this unexpected lineage bifurcation, we examined the transcriptional differences in the rhesus C4 cells using Monocle BEAM analysis and published literature.[24, 28, 30] Branch 1, found only in rhesus, distinguishes the differentiating eNPCs (left branch) vs self-renewing eNPCs (right branch). This rhesus-specific right branch of branch 1 includes NPC markers *EYA1*, *TMEM100, NRXN1, ITGA8*, and *RSPO3*, as well as what has previously been referred to as primed NPC[24], including *SFRP1*, *ROBO2*, and *TPM1*. However, this branch also contains markers of proliferation (*ATP1B1, CAMK2D)*, proximal (*MYC*), medial-distal (*TRIB2*), and distal segment (*LTBP4, PCDH18, COL6A3*). Additionally, these progenitors include markers typically associated with

extracellular matrix/interstitial progenitors including *COL1A2*, *DCN*, *VCAN*, and basement membrane markers *COL14A1* and *HMCN1*. (**Fig 4C, Supplemental Table 8**). In the left (more differentiated) branch of branch 1, genes included primed NPC markers *HEY1, ID1, ID3*, and *ID4*, proliferation markers *TOPS2A, NUSAP1, MKI67, CDC20*, and early renal vesicle/broadly proximal markers *JAG1, HES4, SFRP2* and distal markers *CCND1* and *BMP7*. Interestingly, both sides expressed early proximal marker CDH6 and podocyte marker *ADAMTS9*. These data indicate that the left-sided population would be the one giving rise to more differentiated NPC populations and derivatives via established developmental paradigms.

Branch 2 of rhesus and the single branch point of human demonstrate further differentiation into primed NPC (left branch) and proliferating cells (right branch). There is down-regulation of *RSPO3* in both left and right branches, but more prominent on the right. The left branch shows increased expression of ID1 and ID3 (primed NPC), medial/distal markers (*VIM, IGF2,* and *LTBP1*), proximal tubule markers (*FTL, CD81)*, and extracellular matrix COL1A1, as well as markers *MEIS3, TXNIP, HSBP1,* and *TSC22D1* (**Supplemental Fig 3**). The right branch has high expression of proliferation markers including *TOP2A, MKI67, NUSAP1, CCNA2, and CDK1.* In the human, we see a similar separation of differentiated NPC on the left, dominated by C4 but also C7 and C2, and cell cycle on the right, composed mostly of C8 with some C3. These findings suggest a transitional state of NPC in late-gestation that does not exist in mid-gestation and may provide the source of self-renewing NPC in late-gestation despite their induced state.

Enriched NPCs present age-dependent differences in *WNT* signaling

To evaluate the age-dependent changes in NPCs, we first performed differential expression of the NPC subclusters in rhesus vs human (**Fig 5A, Supplemental Table 9**). Genes prominent in the WNT pathway were identified within the NPC, including Frizzled (FZD) receptors, WNT inhibitors, and Transducin-Like Enhancer of Split (TLE) co-repressors (**Fig 5B**). Specifically, we identified increased *FZD4 (C7)* and *TLE2* and decreased *FZD7 (C3), SHISA2 (C4, C7), SHISA3 (C4), and TLE4 (C4, C7, C8)* (**Fig 5B**). We noted an increase in secreted Frizzled-related protein-1 (*SFRP1)* in the late-gestation rhesus relative to mid-gestation human, strongest within naïve NPC (C4) but present throughout the late-gestation NPC population, consistent with our findings of increased *SFRP1* within the self-renewing NPC branches of the Monocle pseudotime analyses. We also noted increased *SFRP2* in the more differentiated NPC clusters C2,3,5, and 6.

The mechanisms controlling nephrogenesis involve cross-talk of signaling pathways between the NPCs and the closely associated UB tips (**Fig 2**). Our initial subclustering (**Fig. 3C**) of UB cells (clusters c11 and c13) from the integrated dataset identified four stalk clusters (C1, C2, C3, C4) and two tip clusters (C5, C6), consisting of 3,507 total cells in rhesus and 391 cells[23] in human (**Fig 3C, Supplemental Table 10**). Marker genes of the UB tip of the late-gestation rhesus included *HIP1R, MYH10, STC1, AQP3 in C5* and *AKR1B1* and *EHF* in the UB C6, in addition to known UB markers *WNT9B*, *GATA3* and *GATA2* (**Fig 5C, Supplemental Figure 4A, Supplemental Table 11).** Over 95% of the UB cells from the human dataset were classified as UB C6, which limited the temporal comparative analyses to only UB C6 among the UB cells. Supporting a possible reduction in WNT response in the UB tip, we identified overall low expression of AXIN2 in the rhesus UB clusters (**Supplemental Figure 4A)**.

To further understand the signaling between eNPCs and UB, we interrogated the possible secretory receptor/ligand interactions between eNPC and UB in mid vs late gestation using CellChat.[31] Within the late-gestation rhesus, we identified distinct compositional changes within the WNT pathway (**Fig 5D-E, Supplemental Figure 4B, Supplemental Table 12**). While both human and rhesus exhibited WNT4 signaling to FZD4/6/8 from the more differentiated eNPC (C2), the rhesus NPC subcluster C5 demonstrated WNT signaling to UB tip and stalk through WNT5B ligand and FZD 4/6/8 interaction (**Fig 5D**). Additionally, WNT9B ligands within the UB tip C5 and C6 interacted with all of the NPC populations FZD2/3/4/6/8 and LRP5/6 receptors as expected (**Fig 5E**). Combined with the overall higher expression of the secreted antagonist SFRP1 in the naïve NPCs of rhesus, these data suggest an important role for differential modulation of WNT signaling activity as a mechanism governing LBN morphogenesis, consistent with the known role of the pathway as a pivotal regulator of kidney organogenesis.[26, 32-36] To further validate these findings, *in situ* hybridization was used to spatially localize the expression of differentially expressed components within the niche. We identified strong co-staining of *SFRP1* with *SIX1* in the cap NPC population (**Fig 6A**) in late-gestation rhesus and human, with extension surrounding the UB tip. This staining pattern confirmed our DEG analysis within the late-gestation rhesus and supports our pseudotime gene expression changes in the C4 rhesus progenitor population. *SFRP1* extended into the surrounding stroma in addition to the NPC. *SFRP2* was highly expressed in the early differentiated nephron structures, consistent with its high transcriptional level in the induced NPC *WNT4+*

population of cells.[26, 37] We next also noted a change in *FZD* composition within our DEG and highlighted in the cell chat analysis. Based on DEG, we anticipated high expression of *FZD4* and low expression of *FZD7* in late gestation compared to mid gestation. On tissue validation, we identified diffuse staining of FZD4 in both the UB and NPC (**Fig 6B**)*. FZD7*, while still present in the NPC, was most prominent in connecting segment of the distal nephron fusing to the UB tip (**Fig 6B**), which was not anticipated based on DEG. This unexpected localization was also seen at the protein level in the human archival sample, with increasing FZD7 protein expression at the distal fusion points with increasing gestational age, and present in examples for arcading (17wg) and LBN (27-30wg) (**Fig 6C**). Returning to the larger dataset, we identified increased *FZD7* in cluster 15 in late gestation. Although this cluster is labeled as Loop of Henle (with marker genes including *MAP1B*, *TFAP2B*, and *IRX2*), there is some overlap of these genes with collecting duct and urothelial expressions, including *FRAS1*, *TSPAN13*, *LLGL2*, *CPEP2* and *PTPN3*, suggesting broad expression of *FZD7* from Loop of Henle to collecting duct epithelium during the fusion of the nephron to the ureteric stalk.

Given that decreased *SHISA2* and *SHISA3* were identified on our differential expression analysis we next evaluated the expression of these two genes (**Supplemental Fig 4C**). Despite the low expression in the DEG analysis, we identified strong *SHISA2* within the NPC and early differentiating structures at late gestation. Interestingly, *SHISA3*, while still present in the NPC, appeared most prominent in the surrounding stroma (**Supplemental Fig 4C**). *TLE2* and *TLE4* were also evaluated using RNAScope™. *TLE4* was present within the NPC of late-gestation rhesus and human, while no distinct pattern was visible in *TLE2* expression (**Supplemental Fig 4D**).

Enriched UB cells present age-dependent differences in *SEMA3* signaling

SEMA3D expression was significantly upregulated in UB C6 of late-gestation rhesus compared to mid-gestation human samples (FDR p-value < 0.05) (**Supplemental Table 11**). Using CellChat, we identified SEMA3A and SEMA3B signaling from the differentiated eNPC C2, C3, and C5 clusters communicating to the UB stalk and tip clusters via receptors NRP1/2:PLXN1/2 in the late gestation rhesus samples as described above. We next interrogated the possible secretory receptor/ligand interactions from the UB to eNPC in the late gestation rhesus which suggested reciprocal semaphorin signaling in the niche (**Fig 5E**). SEMA3B ligand signaling to the NRP1/2 and PLXN1/2 receptors was observed ubiquitously across eNPCs, with the strongest

interactions noted in eNPC clusters C5 and C6. In contrast, SEMA3D signaling to these receptors originated specifically from the UB tip clusters. As *SEMA3D* was a prominent marker of the rhesus UB tip in both differential expression and UB to NPC secretory communication, we validated this marker using RNAScope™. We identified *SEMA3D* within the UB tip of late-gestation human and rhesus (**Fig 6D**). We also noted *SEMA3D* signal within the interstitium of the more mature deep cortex of the kidney.

DISCUSSION

Through our cross-time, cross-species investigation of the late-gestation rhesus macaque compared to mid-gestation human, we identified molecular differences in the nephron progenitor cells, including distinct NPC clusters in late gestation and distinct trajectories of the naïve NPC in late gestation. Our data suggests a transitional NPC state supported by a potential change in the signaling milieu between the NPC and UB within WNT and SEMA3 signaling pathways, allowing for this differentiated but self-renewing NPC to be maintained within the niche. While we acknowledge the limitations of RNA analyses, this hypothesis provides a potential mechanism for nephron amplification during LBN.

Distinct progenitor trajectories are seen within late-gestation rhesus

Distinct trajectory of the most naïve NPC in late gestation suggests there may be a population of late-gestation that does not exist mid-gestation NPC; a population of early differentiated NPC that contribute to self-renewal and remain in the niche. Rather than a single lineage branch dividing the self-renewing NPC from the differentiating NPC, the rhesus instead has two lineage branches, with a proliferating branch of NPC consisting of NPC C4 and C8. This right branch maintained strong cell cycle signals, markers of the naive NPC, as well as markers associated with the interstitial progenitors and early basement membrane markers, suggesting a shift in both location and early function despite their naive state that may be needed for LBN-nephrogenesis. This transitional proliferating population is distinct from those cells that go on to fully differentiate and become nephrons. The cells undergoing LBN may remain in a high cell cycle and early committed cell state (expressing early BM markers). Slowing of cell proliferation has been associated in the aging of the progenitor niche[38, 39]. However, changes in cell cycle expression at the RNA level are not the full story, as low translation in stem cell populations, independent of cell cycle, have been shown to maintain the progenitor state compared to high translation in differentiated cells.[40] Given that these populations of cells still maintain their progenitor signature,

they may represent the LBN and/or arcading phase of a self-renewing but early differentiated cell that can contribute to nephron formation in late gestation.

Changes in WNT signaling in late gestation may drive the transition from BpN to LBN

One of the potential pathways identified through this investigation included alterations within WNT signaling in late gestation compared to mid gestation. Jarmas et al found that the "tipping point" for nephron progenitor exit from the niche is controlled by the gradual increase in the translation of WNT agonists in individual nephron progenitor cells, enhancing the response to ureteric bud-derived WNT9b inputs.[26] It remains unknown how this shift in WNT responsiveness with age could impact ongoing nephrogenesis in a model with prolonged LBN prior to nephrogenesis cessation. SFRP1 is a secreted WNT antagonist that interacts directly with WNT ligand and can block its interaction with receptors.[41-44] Our data suggests that increased expression of SFRP1 within the naïve NPC may shield these nephron progenitor cells from the action of WNT agonists, as suggested by low expression of downstream target AXIN2, allowing for prolonged, ongoing nephrogenesis in late gestation without branching. Without early gestation rhesus samples for a within-species analyses this remains speculative. Future experiments including overexpression of *SFRP1* in a genetically tractable model of the NPC population may shed light on this question.

SHISA2 and *SHISA3* were downregulated in late gestation in the rhesus datasets, however our tissue validation still supports *SHISA2* was still present within the NPC in late gestation. SHISA2 has been shown to modulate both WNT and FGF signaling[45] and without an earlier time point we cannot conclude that there is decrease within the NPC at late gestation. Interestingly, SHISA3 had low presence in the NPC but had strong presence within the stroma. This supports our previous finding of interstitial progenitor markers within the LBN progenitor population. Similar to GNDF,[46] this may represent a shift in the spatial orientation of SHISA from the NPC to the stroma in late gestation, but its role in supporting LBN requires future investigation.

FZD7 identified the connecting segment/fusion of the distal nephron to the UB rather than the NPC population itself in late gestation. However, mid-gestation human data supports its expression throughout the NPC, as well as podocytes and loop of Henle, but only faintly expressed in the connecting tubule.[47] Our single cell data also supports expression within the Loop of Henle, despite tissue validated presence at the fusion of the developing nephron. FZD7 has also been identified as putative receptor for WNT6, which leads to de novo tubulogenesis in

kidney epithelial cells in culture,[48] but its role within the fusion of the nephron is not known. FZD7 can interact with LRP5/6 to transmit canonical Wnt signaling, as well as through ROR2 to transmit non-canonical Wnt signals.[49] WNT5A, with potential signaling highlighted within the rhesus-only clusters, may play a role in this distal fusion during LBN. WNT5A non-canonical signaling through ROR2 plays an important role in mediating cell proliferation and migration within the metanephric mesenchyme around the wolffian duct[50], further supporting this hypothesis.

The role of semaphorin 3D in UB elongation and induction of lateral branch nephrogenesis

Semaphorins are guidance proteins, mediating short and long-range repulsive and attractive guidance that influence organogenesis and disease.[51] Explant kidney cultures exposed to Sema3a demonstrate reduced ureteric branching, reduced GDNF expression, and reduced Ret phosphorylation. Conversely, Sema3a knockdown/knockout models resulted in increased bud branching.[52] Our study supports that *SEMA3D* is present within UB tips, as well as the surrounding interstitium, while our Cell chat data supports that *SEMA3B* signaling is prominent in the stalk. Our data suggests that increased *SEMA3D* signaling within the UB tip supports elongation rather than bifurcating branching, but future studies comparing the balance of *SEMA3B* to *SEMA3D* in the stalk vs tip require further investigation in early to late gestation. If early to mid-gestation tissue in the rhesus were available in the future, we would perform similar trajectory analyses of the UB overtime to better capture these changes.

The future of lateral branch nephrogenesis analyses

In summary, the gene expression approaches performed in this study help overcome current limitations in identifying pathways acting late in development, as the direct study of late-gestation human kidney development is not possible. The rhesus is the only resource that can be leveraged to define critical cellular and signaling changes that occur during human kidney development at single-cell resolution. Our study highlights a potential transitional, proliferating NPC that amplifies nephrons in late gestation LBN, which is the critical first step to maintaining these pathways otherwise impacted by preterm birth. However, the concepts proposed in this study remain hypothetical. Several limitations exist, resulting in unanswered questions. One of the greatest questions of this analysis is whether the identified changes within the late gestation NPC are time-dependent or species-dependent. As already discussed, no transcriptional data exists for the late-gestation human kidney, and

available human scRNA-seq datasets are limited to a developmental period before LBN begins. Additionally, human fetal archival material from autopsy can be challenging to obtain and can be subject to RNA degradation.[53, 54] Future work will seek to define signaling pathways in the rhesus at mid-gestation before LBN begins, with validation studies at both translational and protein levels to further test these hypotheses. Acquisition of these earlier gestation rhesus datasets will allow us to answer the many questions that remain when defining the unique maturational changes BpN to LBN transition and provide the high tissue samples for spatial validation of our computational findings.

DISCLOSURES

MPS receives research funding from Otsuka that is not relevant to this work

FUNDING

This work was supported by K08DK131259 (MPS), K08DK131259-03S1 (MPS), and ASN Discover Together Normal Seigel Research Scholar Grant (MPS).

ACKNOWLEDGEMENTS

The authors thank Dr. Claire Chougnet, Dr. Alan Jobe, and the California National Primate Center for donating the rhesus kidneys used for this study. This research was made possible, in part, using the Bio-Imaging and Analysis Facility (RRID:SCR_022628). and the Single Cell Genomics Facility (Kelly Rangel, Shawn Smith, RRID: SCR_022655). This study used samples from the Discover Together Biobank at Cincinnati Children's Research Foundation. We thank the Discover Together Biobank for supporting this study, as well as participants and their families, whose help and participation made this work possible. We thank Katrina Schlum for initial insights into the data.

AUTHOR CONTRIBUTIONS

R Kopan and MP Schuh conceptualized and planned the experiments, K Thakkar, R Kopan, and MP Schuh analyzed data. S Yarlagadda, L Alkhudairy, A Potter, and MP Schuh performed the experiments including single-cell dissociation. K Thakkar, N Salomonis, P Chaturvedi, and K Thorner performed bioinformatics analysis in collaboration with MP Schuh. K Thakkar and MP Schuh assembled figures. K Thakkar, KW McCracken, and MP Schuh wrote the manuscript. The final manuscript incorporated input from all authors.

DATA AVAILABILITY

The raw and processed files of scRNA-seq data from rhesus samples are deposited in GEO (Accession number: GSE305708). scRNA-seq data from human samples are available at GSE114530, GSE112570, and GSE102596. Code and scripts for reproducibility are available at https://github.com/kairaveet/integrative-nephrogenesis.

References

1. Hinchliffe SA, Sargent PH, Howard CV, Chan YF, van Velzen D: Human intrauterine renal growth expressed in absolute number of glomeruli assessed by the disector method and Cavalieri principle. *Laboratory investigation; a journal of technical methods and pathology,* 64**:** 777-784, 1991
2. Barker DJ: The intrauterine environment and adult cardiovascular disease. *Ciba Foundation symposium,* 156**:** 3-10; discussion 10-16, 1991
3. Barker D, Eriksson J, Forsen T, Osmond C: Fetal origins of adult disease: strength of effects and biological basis. *Int J Epidemiol,* 31**:** 1235-1239, 2002 10.1093/ije/31.6.1235
4. Barker DJP, Bagby SP: Developmental antecedents of cardiovascular disease: a historical perspective. *Journal of the American Society of Nephrology : JASN,* 16**:** 2537-2544, 2005 10.1681/ASN.2005020160
5. Barker DJ: The fetal and infant origins of disease. *European journal of clinical investigation,* 25**:** 457-463, 1995 10.1136/bmj.301.6761.1111
6. Barker DJP, Bagby SP, Hanson MA: Mechanisms of disease: in utero programming in the pathogenesis of hypertension. *Nature clinical practice Nephrology,* 2**:** 700-707, 2006 10.1038/ncpneph0344
7. Barker DJ, Bull AR, Osmond C, Simmonds SJ: Fetal and placental size and risk of hypertension in adult life. *BMJ (Clinical research ed),* 301**:** 259-262, 1990 10.1136/bmj.301.6751.552
8. Barker DJ, Osmond C, Golding J, Kuh D, Wadsworth ME: Growth in utero, blood pressure in childhood and adult life, and mortality from cardiovascular disease. *BMJ.* 1989 pp 564-567 10.1136/bmj.298.6673.564
9. Lackland DT, Bendall HE, Osmond C, Egan BM, Barker DJ: Low birth weights contribute to high rates of early-onset chronic renal failure in the Southeastern United States. *Archives of internal medicine,* 160**:** 1472-1476, 2000
10. Sayer AA, Cooper C, Evans JR, Rauf A, Wormald RPL, Osmond C, et al.: Are rates of ageing determined in utero? *Age and Ageing,* 27**:** 579-583, 1998 10.1093/ageing/27.5.579
11. Carpenter J, Yarlagadda S, VandenHeuvel KA, Ding L, Schuh MP: Human Nephrogenesis can Persist Beyond 40 Postnatal Days in Preterm Infants. *Kidney Int Rep,* 9**:** 436-450, 2024 10.1016/j.ekir.2023.10.032
12. Rumballe BA, Georgas KM, Combes AN, Ju AL, Gilbert T, Little MH: Nephron formation adopts a novel spatial topology at cessation of nephrogenesis. *Dev Biol,* 360**:** 110-122, 2011 10.1016/j.ydbio.2011.09.011
13. Hartman HA, Lai HL, Patterson LT: Cessation of renal morphogenesis in mice. *Dev Biol,* 310**:** 379-387, 2007 10.1016/j.ydbio.2007.08.021
14. al-Awqati Q, Goldberg MR: Architectural patterns in branching morphogenesis in the kidney. *Kidney Int,* 54**:** 1832-1842, 1998 10.1046/j.1523-1755.1998.00196.x
15. Osathanondh V, Potter EL: Development of human kidney as shown by microdissection. V. Development of vascular pattern of glomerulus. *Archives of pathology,* 82**:** 403-411, 1966
16. EL. P: Normal and abnormal development of the kidney. 305, 1972
17. Schuh MP, Alkhudairy L, Potter A, Potter SS, Chetal K, Thakkar K, et al.: The Rhesus Macaque Serves As a Model for Human Lateral Branch Nephrogenesis. *J Am Soc Nephrol,* 32**:** 1097-1112, 2021 10.1681/asn.2020101459
18. Schuh MP, Yarlagadda S, Alkhudairy L, Preusse K, Kopan R: Characterizing post-branching nephrogenesis in the neonatal rabbit. *Sci Rep,* 13**:** 19234, 2023 10.1038/s41598-023-46624-9

19. Moritz KM, Wintour EM, Black MJ, Bertram JF, Caruana G: Factors influencing mammalian kidney development: implications for health in adult life. *Advances in anatomy, embryology, and cell biology,* 196: 1-78, 2008

20. Lindstrom NO, McMahon JA, Guo J, Tran T, Guo Q, Rutledge E, et al.: Conserved and Divergent Features of Human and Mouse Kidney Organogenesis. *J Am Soc Nephrol,* 29: 785-805, 2018 10.1681/ASN.2017080887

21. Ohuma EO, Moller AB, Bradley E, Chakwera S, Hussain-Alkhateeb L, Lewin A, et al.: National, regional, and global estimates of preterm birth in 2020, with trends from 2010: a systematic analysis. *Lancet,* 402: 1261-1271, 2023 10.1016/s0140-6736(23)00878-4

22. Batchelder CA, Keyser JL, Lee CC, Tarantal AF: Characterization of growth, glomerular number, and tubular proteins in the developing rhesus monkey kidney. *Anat Rec (Hoboken),* 296: 1747-1757, 2013 10.1002/ar.22756

23. Hochane M, van den Berg PR, Fan X, Bérenger-Currias N, Adegeest E, Bialecka M, et al.: Single-cell transcriptomics reveals gene expression dynamics of human fetal kidney development. *PLoS Biol,* 17: e3000152, 2019 10.1371/journal.pbio.3000152

24. Lindstrom NO, De Sena Brandine G, Tran T, Ransick A, Suh G, Guo J, et al.: Progressive Recruitment of Mesenchymal Progenitors Reveals a Time-Dependent Process of Cell Fate Acquisition in Mouse and Human Nephrogenesis. *Dev Cell,* 45: 651-660 e654, 2018 10.1016/j.devcel.2018.05.010

25. Lindstrom NO, Guo J, Kim AD, Tran T, Guo Q, De Sena Brandine G, et al.: Conserved and Divergent Features of Mesenchymal Progenitor Cell Types within the Cortical Nephrogenic Niche of the Human and Mouse Kidney. *J Am Soc Nephrol,* 29: 806-824, 2018 10.1681/ASN.2017080890

26. Jarmas AE, Brunskill EW, Chaturvedi P, Salomonis N, Kopan R: Progenitor translatome changes coordinated by Tsc1 increase perception of Wnt signals to end nephrogenesis. *Nature Communications,* 12: 6332, 2021 10.1038/s41467-021-26626-9

27. Korsunsky I, Millard N, Fan J, Slowikowski K, Zhang F, Wei K, et al.: Fast, sensitive and accurate integration of single-cell data with Harmony. *Nature Methods,* 16: 1289-1296, 2019 10.1038/s41592-019-0619-0

28. Chen J, Bardes EE, Aronow BJ, Jegga AG: ToppGene Suite for gene list enrichment analysis and candidate gene prioritization. *Nucleic Acids Research,* 37: W305-W311, 2009 10.1093/nar/gkp427

29. Qiu X, Mao Q, Tang Y, Wang L, Chawla R, Pliner HA, et al.: Reversed graph embedding resolves complex single-cell trajectories. *Nat Methods,* 14: 979-982, 2017 10.1038/nmeth.4402

30. Qiu X, Hill A, Packer J, Lin D, Ma YA, Trapnell C: Single-cell mRNA quantification and differential analysis with Census. *Nat Methods,* 14: 309-315, 2017 10.1038/nmeth.4150

31. Jin S, Guerrero-Juarez CF, Zhang L, Chang I, Ramos R, Kuan CH, et al.: Inference and analysis of cell-cell communication using CellChat. *Nat Commun,* 12: 1088, 2021 10.1038/s41467-021-21246-9

32. Park J-S, Ma W, O'Brien Lori L, Chung E, Guo J-J, Cheng J-G, et al.: Six2 and Wnt Regulate Self-Renewal and Commitment of Nephron Progenitors through Shared Gene Regulatory Networks. *Developmental Cell,* 23: 637-651, 2012 https://doi.org/10.1016/j.devcel.2012.07.008

33. Stark K, Vainio S, Vassileva G, McMahon AP: Epithelial transformation of metanephric mesenchyme in the developing kidney regulated by Wnt-4. *Nature,* 372: 679-683, 1994 10.1038/372679a0

34. Merkel CE, Karner CM, Carroll TJ: Molecular regulation of kidney development: is the answer blowing in the Wnt? *Pediatr Nephrol,* 22: 1825-1838, 2007 10.1007/s00467-007-0504-4

35. Pulkkinen K, Murugan S, Vainio S: Wnt signaling in kidney development and disease. *Organogenesis,* 4: 55-59, 2008 10.4161/org.4.2.5849

36. Park J-S, Valerius MT, McMahon AP: Wnt/β-catenin signaling regulates nephron induction during mouse kidney development. *Development,* 134: 2533-2539, 2007 10.1242/dev.006155

37. Lawlor KT, Zappia L, Lefevre J, Park J-S, Hamilton NA, Oshlack A, et al.: Nephron progenitor commitment is a stochastic process influenced by cell migration. *eLife,* 8: e41156, 2019 10.7554/eLife.41156

38. Chen S, Brunskill Eric W, Potter SS, Dexheimer Phillip J, Salomonis N, Aronow Bruce J, et al.: Intrinsic Age-Dependent Changes and Cell-Cell Contacts Regulate Nephron Progenitor Lifespan. *Developmental Cell,* 35: 49-62, 2016 10.1016/j.devcel.2015.09.009

39. Short KM, Combes AN, Lefevre J, Ju AL, Georgas KM, Lamberton T, et al.: Global quantification of tissue dynamics in the developing mouse kidney. *Dev Cell,* 29: 188-202, 2014 10.1016/j.devcel.2014.02.017

40. Tahmasebi S, Alain T, Rajasekhar VK, Zhang JP, Prager-Khoutorsky M, Khoutorsky A, et al.: Multifaceted regulation of somatic cell reprogramming by mRNA translational control. *Cell Stem Cell,* 14**:** 606-616, 2014 10.1016/j.stem.2014.02.005

41. Finch PW, He X, Kelley MJ, Üren A, Schaudies RP, Popescu NC, et al.: Purification and molecular cloning of a secreted, Frizzled-related antagonist of Wnt action. *Proceedings of the National Academy of Sciences,* 94**:** 6770-6775, 1997

42. Cruciat C-M, Niehrs C: Secreted and transmembrane wnt inhibitors and activators. *Cold Spring Harbor perspectives in biology,* 5**:** a015081, 2013

43. Kawano Y, Kypta R: Secreted antagonists of the Wnt signalling pathway. *Journal of cell science,* 116**:** 2627-2634, 2003

44. Mii Y, Taira M: Secreted Wnt "inhibitors" are not just inhibitors: Regulation of extracellular Wnt by secreted Frizzled-related proteins. *Development, growth & differentiation,* 53**:** 911-923, 2011

45. Furushima K, Yamamoto A, Nagano T, Shibata M, Miyachi H, Abe T, et al.: Mouse homologues of Shisa antagonistic to Wnt and Fgf signalings. *Developmental Biology,* 306**:** 480-492, 2007 https://doi.org/10.1016/j.ydbio.2007.03.028

46. Magella B, Adam M, Potter AS, Venkatasubramanian M, Chetal K, Hay SB, et al.: Cross-platform single cell analysis of kidney development shows stromal cells express Gdnf. *Dev Biol,* 434**:** 36-47, 2018 10.1016/j.ydbio.2017.11.006

47. Lindström NO, Sealfon R, Chen X, Parvez RK, Ransick A, De Sena Brandine G, et al.: Spatial transcriptional mapping of the human nephrogenic program. *Developmental Cell,* 56**:** 2381-2398.e2386, 2021 https://doi.org/10.1016/j.devcel.2021.07.017

48. Beaton H, Andrews D, Parsons M, Murphy M, Gaffney A, Kavanagh D, et al.: Wnt6 regulates epithelial cell differentiation and is dysregulated in renal fibrosis. *Am J Physiol Renal Physiol,* 311**:** F35-45, 2016 10.1152/ajprenal.00136.2016

49. Phesse T, Flanagan D, Vincan E: Frizzled7: A Promising Achilles' Heel for Targeting the Wnt Receptor Complex to Treat Cancer. *Cancers (Basel),* 8, 2016 10.3390/cancers8050050

50. Nishita M, Qiao S, Miyamoto M, Okinaka Y, Yamada M, Hashimoto R, et al.: Role of Wnt5a-Ror2 signaling in morphogenesis of the metanephric mesenchyme during ureteric budding. *Mol Cell Biol,* 34**:** 3096-3105, 2014 10.1128/mcb.00491-14

51. Tran TS, Kolodkin AL, Bharadwaj R: Semaphorin regulation of cellular morphology. *Annu Rev Cell Dev Biol,* 23**:** 263-292, 2007 10.1146/annurev.cellbio.22.010605.093554

52. Tufro A, Teichman J, Woda C, Villegas G: Semaphorin3a inhibits ureteric bud branching morphogenesis. *Mechanisms of Development,* 125**:** 558-568, 2008 https://doi.org/10.1016/j.mod.2007.12.003

53. von Ahlfen S, Missel A, Bendrat K, Schlumpberger M: Determinants of RNA Quality from FFPE Samples. *PLOS ONE,* 2**:** e1261, 2007 10.1371/journal.pone.0001261

54. Groelz D, Sobin L, Branton P, Compton C, Wyrich R, Rainen L: Non-formalin fixative versus formalin-fixed tissue: a comparison of histology and RNA quality. *Exp Mol Pathol,* 94**:** 188-194, 2013 10.1016/j.yexmp.2012.07.002

FIGURES

Fig 1. Stages of nephrogenesis in human and comparative rhesus morphology. A) During branch phase nephrogenesis (BpN), nephrons are formed in a process associated with bifurcating branching of the ureteric bud (UB). **B)** 15 weeks gestation (WG), human nephrogenesis transitions to arcading phase followed by **C)** LBN at 23-36 WG, where an elongating, non-branching UB tip induces nephrons. Rhesus gestation in days included for reference. **D-F)** Early-LBN (108dg) rhesus displays a rosette-like organization of the NPC niche, similar to both 129dg rhesus and 32 wg human. with evidence of both LBN and rare BpN. G) the early-LBN rhesus (108dg) represents a critical transition period from BpN to LBN. Although lateral branches (blue arrow) are seen along the ureteric stalk, there is also evidence of recent BpN branching events (yellow arrow).

Fig 2. Schematic of the lateral branch nephrogenesis: nephron induction without UB branching. A) the schematic representation of nephron induction and fusion through LBN with representative immunofluorescent images from **B)** human and **C)** rhesus in late gestation. There is broad RET (yellow) expression across the UB tip and in contact with nephron progenitor cells (NPC, green) (panel 1). RET expression is visualized along the side of the ureteric tip with contact to the formation of renal vesicles (panels 2-3). Distal CDH1+ (red) fusion of the nephron occurs near the base of the UB tip in close proximity to the RET+ cells, with NPC just above this fusion point (panel 4). The bulbous UB tip then outpouches or grows above this fusion point, with the NPC traveling with it (panel 5), leaving the recently formed "lateral branch" below.

Fig 3. Cross-species integrative single-cell transcriptomics characterizes heterogeneity and temporal differences in NPCs. A) Integrated UMAP of 94,266 cells from cortically-enriched fetal kidneys from 9 rhesus (late second and third trimesters) samples and 5 human (8-13 weeks old) samples. Cells are colored by cellHarmony cluster labels (left panel) and by species (right panel). **B)** UMAP from panel A colored by gene expression of key NPC progenitor marker genes. **C)** Integrated UMAP of NPC and UB cells from rhesus and human samples. Cells are colored by cell subclusters within NPCs and UB (left panel) and by species (right panel). **D)** Dot plot indicating average expression of major marker genes of NPC subclusters in human (top panel) and rhesus (bottom panel) datasets. Dots are sized by the percent of cells that expressed the marker

gene. **E)** UMAP from Panel C highlighting the enriched NPCs (eNPCs) defined by expression for: *SIX1*, *SIX2*, *ITGA8* (at least 2 of the three genes expressed). The highlighted cells are colored by species, similar to right panel C.

Fig 4. Developmental trajectory of NPCs. Lineage trajectory of NPCs constructed by Monocle2 for **A)** human and **B)** rhesus. Pseudotime increases from top to the bottom of the tree. To clearly illustrate where clusters are located, the lineage tree is reproduced to highlight each rhesus cell population. **C)** Heatmap of normalized gene expression showing branch-dependent genes (Monocle BEAM test; FDR < 10%) as rows along the two fates of Branch 1 in rhesus. Left and right segments show the gene expression change along the left and right branches, respectively, after Branch 1 in the Monocle tree in panel B. Columns are points in pseudotime, and the dividing middle segment of the heatmap indicates beginning of pseudotime. Selected genes are highlighted on the right. Rows are clustered using hierarchical clustering.

Fig 5. Age-dependent changes in WNT signaling. A) Heatmap indicating the log2 fold changes of differentially expressed genes (FDR p-value < 0.001) of each NPC subcluster in rhesus as compared to human cells. Top 150 differentially expressed genes are considered from each comparison. **B, C)** Dot plot indicating average expression of **B)** selected genes associated with WNT signaling in NPC subclusters in human (top panel) and rhesus (bottom panel) datasets, and **C)** marker genes of UB subclusters in rhesus. Dots are sized by the percent of cells that expressed the gene. **D, E)** Bubble plots showing the WNT-associated ligand-receptor pairs (rows) that contribute to secreted signaling **D)** from NPC to UB and **E)** UB to NPC in rhesus. Dots are colored by communication probability and sized by p-value.

Fig 6. RNAScope™ validation of WNT pathway and SEMA3D. A) Representative late-gestation rhesus and human kidney with co-localization of SFRP1 and SIX1, with additional SFRP1 expression within the surrounding stroma. SFRP2 was noted further away from the cortical edge, likely representing the WNT4+ population of NPC. **B)** FZD 4 and 7 expression in the late-gestation rhesus and human. While FZD4 was broadly expressed in the NPC and developing nephron, FZD7 localized to the connecting segment or the distal nephron and fusion to the UB. **C)** FZD7 protein expression in the developing human kidney. FZD7 is seen at the fusion of the nephron segments in examples of arcading (17 wg) and lateral branch nephrogenesis (27-30 wg).

D) Late-gestation rhesus and human demonstrate strong *SEMA3D* signal in the outer UB and UB tip, with additional SEMA3D staining identified within the maturing interstitium.

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6