Title: Bivalent mRNA booster encoding virus-like particles elicits potent polyclass

RBD antibodies in pre-vaccinated mice

Authors: Chengcheng Fan[1,7], Alexander A. Cohen[1,7], Kim-Marie A. Dam[2,7], Annie V.

Rorick[1,3], Ange-Célia I. Priso Fils[1], Zhi Yang[1,4], Priyanthi N. P. Gnanapragasam[1], Luisa

N. Segovia[1], Kathryn E. Huey-Tubman[1], Woohyun J. Moon[5], Paulo J.C. Lin[5], Pamela J.

Bjorkman[1*], Magnus A. G. Hoffmann[2,6*]

Affiliations:

[1]Division of Biology and Biological Engineering, California Institute of Technology,

Pasadena, CA 91125, USA

[2]Gladstone Institutes, San Francisco, CA 94158, USA

[3]Present address: Department of Biochemistry, University of Washington, Seattle, WA

98109, USA

[4]Present address: Department of Molecular and Cell Biology, University of California,

Berkeley, Berkeley, CA 94720, USA

[5]Acuitas Therapeutics, Vancouver, BC, V6T 1Z3, CANADA

[6]Department of Medicine, University of California, San Francisco, CA 94143, USA

[7]These authors contributed equally

[*]To whom correspondence should be addressed: Email: bjorkman@caltech.edu,

magnus.hoffmann@gladstone.ucsf.edu

Abstract

mRNA vaccines emerged as a leading vaccine technology during the COVID-19 pandemic. However, their sustained protective efficacies were limited by relatively short-lived antibody responses and the emergence of SARS-CoV-2 variants, necessitating frequent and variant-updated boosters. We recently developed the ESCRT- and ALIX-binding region (EABR) mRNA vaccine platform, which encodes engineered immunogens that induce budding of enveloped virus-like particles (eVLPs) from the plasma membrane, thereby resulting in presentation of immunogens on cell surfaces and eVLPs. Prior studies showed that spike (S)-EABR mRNA-LNP immunizations elicited enhanced neutralizing antibody responses against ancestral and variant SARS-CoV-2 compared with conventional S mRNA-LNP in naïve mice, but the effectiveness of S-EABR mRNA-LNP boosters in the context of pre-existing immunity has not been investigated. Here, we evaluated monovalent Wuhan-Hu-1 (Wu1) and bivalent (Wu1/BA.5) S-EABR mRNA-LNP boosters in mice pre-vaccinated with conventional Wu1 S mRNA-LNP. Compared to conventional S mRNA-LNP boosters, the EABR approach enhanced monovalent and bivalent mRNA-LNP booster-induced neutralizing responses against Omicron subvariants BA.1, BA.5, BQ.1.1, and XBB.1, with bivalent S-EABR mRNA-LNP consistently eliciting the highest titers. Epitope mapping of polyclonal antisera by deep mutational scanning revealed that bivalent S-EABR mRNA-LNP boosted diverse "polyclass" anti-RBD responses, suggesting balanced targeting of multiple RBD epitope classes. In contrast, monovalent S, bivalent S, and monovalent S-EABR mRNA-LNP boosters elicited less diverse polyclonal serum responses primarily targeting immunodominant RBD epitopes. Cryo-EM structures demonstrated that bivalent mRNA

immunizations promote S heterotrimer formation, potentially enhancing bivalent S-EABR

mRNA-LNP booster-induced antibody breadth and polyclass epitope targeting by

activating cross-reactive B cells through intra-S crosslinking. These findings support the

future design of bivalent or multivalent S-EABR mRNA-LNP boosters as a promising

strategy to confer broader, and therefore potentially more durable, protection against

emerging SARS-CoV-2 variants and other rapidly evolving viruses.

Introduction

mRNA vaccines emerged as a leading platform for the rapid and scalable development of effective vaccines during the COVID-19 pandemic[1]. Currently approved mRNA vaccines encode the SARS-CoV-2 spike (S) trimer[2], the main target for neutralizing antibodies (Abs) produced during natural infection[3]. S-encoding mRNAs formulated in lipid nanoparticles (LNP) are taken up by cells, which promotes translation and cell surface expression of S trimers to activate B cells[4]. Translation of mRNA-encoded S protein within host cells also provides viral peptides for presentation on MHC class I molecules, which are recognized by cytotoxic T cells[5,6].

Initial two-shot COVID-19 mRNA vaccination series became available in December 2020 after clinical studies demonstrated protection from severe disease after SARS-CoV-2 infection[7,8]. However, the durability of mRNA vaccine-induced immune protection was limited by two factors. First, serum Ab levels contracted over several months[9]. Second, SARS-CoV-2 variants acquired mutations in the S protein, which reduced their sensitivity to vaccine-induced neutralizing Abs[10]. As a result, a third dose of the original mRNA vaccine encoding the ancestral Wuhan-Hu-1 (Wu1) S protein was offered starting in fall 2021[11].

The emergence of the Omicron BA.1 variant in November 2021[12] sparked concern because of its levels of substitution in S, especially in the receptor-binding domain (RBD), resulting in effective evasion of Ab-mediated immunity[13,14] and increased rates of symptomatic infections in previously vaccinated individuals[15,16]. Although undetectable

after the initial mRNA vaccine series, a third dose of the original mRNA vaccine elicited

neutralizing Abs against BA.1, but BA.1 neutralizing titers were lower compared to titers

against the ancestral Wu1 variant[17]. The subsequent emergence of Omicron subvariants

such as BA.5, BQ.1.1, and XBB.1, which exhibited progressively reduced sensitivities to

vaccine-induced neutralizing Abs[18], prompted the development of bivalent mRNA

boosters, which included mRNAs encoding both the Wu1 and the BA.5 S proteins[19].

Preclinical and clinical studies found that bivalent boosters elicited modestly improved

neutralizing Ab responses against Omicron subvariants compared to a fourth dose of the

original monovalent vaccine[19-23], although some studies found no differences between

monovalent and bivalent boosters following initial vaccination with the original Wu1 S-

encoding mRNA vaccine[24,25]. Several studies[21,22,24,26-28] showed that the effectiveness of

bivalent boosters was limited by immune imprinting, which occurs when prior exposure to

an antigen shapes immunological responses to related antigens during subsequent

exposures[29]. Instead of generating *de novo* Omicron-specific immune responses, bivalent

boosters primarily activated pre-existing cross-reactive memory B cells from previous

vaccinations and/or infections[26-28], leading to lower neutralizing activity against Omicron

subvariants compared to the ancestral variant[21,22,24].

Continued development of updated boosters in response to emerging variants was

slowed by the need for additional clinical testing, regulatory approvals, and adjustments

to the manufacturing process. As a result, bivalent boosters became available in fall 2022,

almost one year after the emergence of Omicron[30]. These challenges highlight the urgent

need for innovative vaccine strategies that achieve more lasting immune protection

against rapidly-evolving viruses such as SARS-CoV-2, thereby reducing the need for

frequent and updated boosters.

We recently developed the ESCRT- and ALIX-binding region (EABR) mRNA vaccine

platform that combines features of mRNA and protein nanoparticle (NP) vaccines through

delivery of mRNA-encoded self-assembling enveloped virus-like particles (eVLPs)[31].

eVLP assembly is achieved by fusing a short EABR sequence to the cytoplasmic tail of

viral surface proteins, which recruits proteins from the Endosomal Sorting Complex

Required for Transport (ESCRT) pathway. Thus, EABR mRNA vaccines promote dual

presentation of S trimers on cell surfaces and eVLPs that bud from the plasma membrane.

Immunizations in mice showed that an EABR mRNA vaccine encoding a Wu1 SARS-

CoV-2 S elicited superior neutralizing Ab potency and breadth against ancestral and

Omicron SARS-CoV-2 variants compared to conventional mRNA and protein NP

vaccines[31]. Similar to conventional mRNA vaccines, the EABR mRNA vaccine also

induced potent T cell responses, consistent with the presence of S-specific cytotoxic T

cells[31]. These findings suggest that the EABR mRNA vaccine approach has the potential

to enhance monovalent or bivalent booster-induced Ab responses against Omicron

subvariants in the context of pre-existing immunity induced by prior vaccination, infection,

or both.

Here, we evaluated the ability of EABR mRNA boost immunogens to elicit potent and

broad humoral immune responses against SARS-CoV-2 variants in pre-vaccinated mice.

We demonstrate that the EABR mRNA vaccine approach improves monovalent and

bivalent booster-mediated neutralizing Ab responses against Omicron subvariants

compared to conventional mRNA boosters. Epitope mapping of polyclonal antisera further

showed that a bivalent EABR mRNA booster elicited a more diversified anti-RBD IgG

response than other tested boosting immunogens. These findings support the use of the

EABR mRNA vaccine platform for the design of effective boosters that could confer more

lasting protection against SARS-CoV-2 variants and potentially other rapidly-evolving

viruses. In addition, our work provides structural evidence that bivalent mRNA

immunizations promote heterotrimer formation, which could enhance activation of cross-

reactive B cells through intra-S crosslinking. These insights inform the future design of

bivalent or multivalent COVID-19 boosters and highlight the need for further investigation

into potentially beneficial effects of S heterotrimer formation on vaccine-induced immune

responses against trimeric viral Ss.

Results

To expand upon our previous characterizations of a monovalent SARS-CoV-2 S-EABR

mRNA vaccine[31], we evaluated the potential of the EABR mRNA vaccine approach for

monovalent and bivalent SARS-CoV-2 immunizations in pre-vaccinated mice. Four

cohorts of 10 BALB/c mice each initially received prime-boost intramuscular (IM)

vaccinations on days 0 and 28 with 2 µg of a conventional mRNA vaccine encoding the

ancestral S protein from the Wu1 variant, similar to the immunogens used in initially

approved COVID-19 mRNA vaccines (Wu1 S mRNA-LNP) (Fig. 1A-C). To simulate the

timing of the third available COVID-19 mRNA vaccine dose in humans, booster

immunizations were administered on day 230, nearly 7 months after the primary

vaccination series, with each cohort receiving a different boost immunogen (Fig. 1A-C).

Consistent with the COVID-19 booster that became available in fall 2021[11], cohort 1

received a third dose of 2 µg Wu1 S mRNA-LNP (Fig. 1B-C). To evaluate whether

formation of mRNA-encoded S-EABR eVLPs enhances booster immunogenicity in pre-

vaccinated mice, cohort 2 received 2 µg Wu1 S-EABR mRNA-LNP (Fig. 1B-C). Cohort 3

received a bivalent Wu1/BA.5 S mRNA-LNP booster (1 µg Wu1 S mRNA-LNP combined

with 1 µg BA.5 S mRNA-LNP), which resembled the COVID-19 booster that became

available in fall 2022[30] (Fig. 1B-C). Lastly, cohort 4 received a bivalent Wu1/BA.5 S-EABR

mRNA-LNP booster to promote the formation of eVLPs that co-display Wu1 and BA.5 S

trimers (Fig. 1B-C). The Wu1 and BA.5 S-EABR constructs were designed as previously

described by fusing an endocytosis-preventing motif (EPM), a short Gly-Ser linker, and

an EABR sequence to the end of the cytoplasmic domain of Wu1 and BA.5 S trimers[31]

(Fig. S1A). The Wu1 and BA.5 S-EABR constructs induced similar levels of eVLP budding

in transiently transfected HEK293T cells (Fig. S1B).

Serum binding Ab responses were measured by enzyme-linked immunosorbent assay

(ELISA) four and 18 weeks after the primary vaccination series (days 56 and 154) and

two weeks after the booster immunizations (day 244). As previously observed, the initial

Wu1 S mRNA-LNP vaccinations elicited high binding Ab titers against soluble Wu1 S-6P

trimers[32], and as expected, these titers were similar for all cohorts on day 56 (Fig. 1D).

Robust binding Ab responses were also detected for all cohorts against Omicron BA.5,

BQ.1.1, and XBB.1 S-6P trimers on day 56 (Fig. 1E-G). On day 154, binding Ab levels

against all variants dropped slightly, which was consistent between cohorts (Fig. 1D-G).

On day 244, all boost immunogens increased binding Ab titers against Wu1, BA.5,

BQ.1.1, and XBB.1 S-6P trimers compared to titers measured on days 56 and 154, but

no differences were observed between cohorts (Fig. 1D-G).

Bivalent S-EABR mRNA-LNP booster elicits potent Ab responses targeting

Omicron RBDs.

We next evaluated the impact of the different boost immunizations on Ab responses

against the RBD, the primary target of neutralizing Abs[3]. The initial Wu1 S mRNA-LNP

prime-boost vaccinations elicited robust and similar Ab responses against the Wu1 RBD

for all cohorts on day 56 (Fig. 2A). The BA.5, BQ.1.1, and XBB.1 variants harbor multiple

RBD substitutions, particularly within the class 1 and 2 epitopes at the tip of the RBD[33]

(Fig. S2), promoting effective escape from neutralizing Abs[18]. Compared to Wu1 RBD

binding, Ab binding titers were weaker against the BA.5, BQ.1.1, and XBB.1 RBDs on day

56 (Fig. 2B-D). The robust Ab binding against the BA.5, BQ.1.1, and XBB.1 S trimers (Fig.

1E-G) in the absence of strong RBD responses was at least in part driven by potent

responses against the S2 region of S (Fig. S3), which is relatively conserved in SARS-

CoV-2 variants[34] (Fig. S2). Similar to binding titers against S-6P trimers (Fig. 1D-G), Wu1,

BA.5, BQ.1.1, and XBB.1 RBD binding slightly dropped on day 154 without statistically

significant differences between cohorts (Fig. 2A-D). The booster immunizations increased

Wu1 RBD binding on day 244 compared to day 56 and 154 levels, but no differences

were detected between the four boosting immunogens (Fig. 2A). In contrast, the bivalent

Wu1/BA.5 S-EABR mRNA-LNP booster elicited the highest BA.5, BQ.1.1, and XBB.1

RBD responses on day 244 (Fig. 2B-D), which were significantly higher than Wu1 S

mRNA-LNP booster-induced responses against BQ.1.1 RBD ($p = 0.0499$) (Fig. 2C) and

narrowly failed to reach statistical significance compared to Wu1 S mRNA-LNP booster-

induced responses against BA.5 RBD ($p = 0.07$) (Fig. 2B). Compared to baseline titers

measured prior to boost immunizations on day 154, the bivalent Wu1/BA.5 S-EABR

mRNA-LNP immunization elicited significantly larger increases in BA.5 RBD binding than

the monovalent Wu1 S mRNA-LNP ($p = 0.02$) and the bivalent Wu1/BA.5 S mRNA-LNP

boosters ($p = 0.01$) (Fig. 2B), as well as significantly larger increases in BQ.1.1 RBD

binding than the Wu1 S mRNA-LNP booster ($p = 0.002$) (Fig. 2C). While overall titers

were similar on day 244, the monovalent Wu1 S-EABR mRNA-LNP booster consistently

induced larger increases in Ab binding to the BA.5, BQ.1.1, and XBB.1 RBDs than the

monovalent Wu1 S mRNA-LNP booster (Fig. 2B-D), although these differences failed to

reach statistical significance. The monovalent Wu1 and bivalent Wu1/BA.5 S mRNA-LNP

boosters elicited similar binding titers against the BA.5 and XBB.1 RBDs (Fig. 2B and D),

but the bivalent booster induced slightly higher responses and a larger increase

compared to day 154 titers against BQ.1.1 RBD (not statistically significant) (Fig. 2C).

Overall, these results demonstrate that booster immunizations with the bivalent Wu1/BA.5

S-EABR mRNA-LNP elicited the highest Ab responses against Omicron RBDs in pre-

vaccinated mice. Notably, post-boost Ab binding against Omicron RBDs remained lower

than binding titers against Wu1 RBD in all cohorts, suggesting that booster responses

were affected by immune imprinting induced by the primary vaccination series.

Bivalent S-EABR mRNA-LNP booster elicits higher neutralizing responses against Omicron variants.

The ability of boost immunogens to elicit neutralizing Ab responses in pre-vaccinated mice was evaluated by in vitro pseudovirus neutralization assays[35,36]. The primary vaccination series with Wu1 S mRNA-LNP elicited potent and consistent autologous geometric mean neutralizing titers as assessed by half-maximal serum inhibitory dilution (ID_{50}) values against Wu1 pseudovirus in all cohorts on day 56 (Fig. 3A). Neutralizing activity was >100-fold lower against the BA.5 variant and undetectable in most animals against the BQ.1.1 and XBB.1 variants on day 56 for all cohorts (Fig. 3B-D). On day 154, neutralizing titers against the Wu1 and BA.5 variants were consistent in all cohorts and slightly lower compared to day 56 titers (Fig. 3A-B). Neutralizing activity against the BQ.1.1 and XBB.1 variants was undetectable in all but one animal on day 154 (Fig. 3C-D). Booster immunizations on day 230 increased autologous neutralizing titers against Wu1 pseudovirus on day 244 compared to day 154 in all cohorts (Fig. 3A). Monovalent Wu1 S mRNA-LNP and S-EABR mRNA-LNP immunizations elicited the smallest and largest increases in Wu1 neutralization (4.2-fold versus 6.6-fold), respectively. All boost immunogens elicited increased BA.5 neutralizing activity on day 244 compared to day 56 and 154 titers (Fig. 3B). Boosting with monovalent Wu1 S-EABR mRNA-LNP induced 4.8-fold higher BA.5 neutralizing titers than a third Wu1 S mRNA-LNP immunization, but this difference did not reach statistical significance. Using the respective day 154 titers as baseline, the Wu1 S-EABR mRNA-LNP booster induced a 23.8-fold increase in BA.5 neutralization on day 244 compared to only a 6.3-fold increase detected after boosting with Wu1 S mRNA-LNP (not statistically significant) (Fig. 3B). Compared to day 56

baseline titers, the Wu1 S-EABR mRNA-LNP booster induced a significantly higher

increase in BA.5 neutralization on day 244 than the conventional Wu1 S mRNA-LNP

booster (16.2-fold versus 3.1-fold; p = 0.048) (Fig. S4). The Wu1 S-EABR mRNA-LNP

booster also elicited 3.4-fold higher neutralizing responses than the Wu1 S mRNA-LNP

booster against the Omicron BA.1 variant on day 244 (p = 0.03) (Fig. S5). Based on day

154 titers, boosting with monovalent Wu1 S-EABR mRNA-LNP induced significantly

higher increases in BA.1 neutralization compared to monovalent Wu1 S mRNA-LNP (8.8-

fold versus 2.5-fold; p = 0.009) (Fig. S5). Interestingly, the monovalent Wu1 S-EABR

mRNA-LNP and the bivalent Wu1/BA.5 S mRNA-LNP boosters elicited similar BA.5

neutralizing responses on day 244 (Fig. 3B), suggesting that the EABR mRNA vaccine

approach might reduce the need for updating boost immunogens to achieve robust Ab

responses against emerging SARS-CoV-2 variants. The highest BA.5 neutralizing titers

were elicited by the bivalent Wu1/BA.5 S-EABR mRNA-LNP booster, which were 12.3-

fold higher than titers induced by the monovalent Wu1 S mRNA-LNP booster (p = 0.02)

and 2.4-fold higher than titers induced by the bivalent Wu1/BA.5 S mRNA-LNP booster

(not statistically significant). Based on titers measured on day 154, the Wu1/BA.5 S-EABR

mRNA-LNP booster induced a 67-fold increase in BA.5 neutralization, which was

significantly higher than the increase elicited by boosting with monovalent Wu1 S mRNA-

LNP (p = 0.003).

While the BA.5 neutralizing activity induced by the monovalent Wu1 S and S-EABR

mRNA-LNP boosters can likely be attributed to cross-neutralizing Abs that neutralize both

the Wu1 and BA.5 variants, it is possible that the bivalent Wu1/BA.5 S and S-EABR

mRNA-LNP boosters also elicited *de novo* strain-specific Abs that neutralize BA.5 but not

Wu1. Day 244 serum samples from mice that received bivalent Wu1/BA.5 S or S-EABR

mRNA-LNP boosters were depleted of Abs that bind to Wu1 RBD. Sera were passaged

over a column presenting either a mock protein (HIV-1 gp120) as a control or the Wu1

RBD to remove cross-neutralizing anti-RBD Abs while retaining BA.5 RBD-specific Abs.

Following depletions, residual neutralization activity was measured against the BA.5

variant. Interestingly, Wu1 RBD depletion resulted in 17- and 36-fold lower geometric

mean neutralizing titers compared to mock depletion for the bivalent Wu1/BA.5 S and S-

EABR mRNA-LNP boosters, respectively (Fig. S6). In both cohorts, BA.5 neutralizing

titers in all but two mice dropped below ID_{50} values of 1:120 following Wu1 RBD depletion,

consistent with the neutralizing activity in these samples being primarily driven by cross-

neutralizing Abs targeting the RBD induced by the primary vaccination series. The potent

residual neutralizing activity observed in two mice in each cohort after Wu1 RBD depletion

implies either the presence of boost-induced *de novo* BA.5-specific RBD Abs or

neutralizing non-RBD Abs that target another region of S, e.g., the N-terminal domain[37]

or S2[34].

Booster immunizations had less pronounced effects on neutralization activity against the

BQ.1.1 and XBB.1 variants. Against BQ.1.1 pseudovirus, the bivalent Wu1/BA.5 S-EABR

mRNA-LNP booster elicited the most potent neutralizing activity on day 244, which was

significantly higher than responses induced by a third immunization with the monovalent

Wu1 S mRNA-LNP booster (p = 0.0495) (Fig. 3C). The monovalent Wu1 S-EABR mRNA-

LNP and bivalent Wu1/BA.5 S mRNA-LNP boosters elicited 1.8- and 2.5-fold higher

BQ.1.1 titers than the monovalent Wu1 S mRNA-LNP booster, respectively, the latter of

which narrowly failed to reach statistical significance (p = 0.07). Based on day 154 BQ.1.1

neutralizing titers, the monovalent Wu1 S mRNA-LNP and bivalent Wu1/BA.5 S-EABR

mRNA-LNP boosters induced the smallest and largest increases in neutralizing activity

on day 244, respectively (1.4-fold versus 4.7-fold; p = 0.0495). The monovalent Wu1 S-

EABR mRNA-LNP and bivalent Wu1/BA.5 S mRNA-LNP boosters elicited 2.4- and 3.4-

fold increases in neutralizing BQ.1.1 titers, respectively. Notably, day 244 neutralizing

titers against BQ.1.1 were substantially lower than BA.5 neutralizing titers for all boost

immunogens (Fig. 3B-C), consistent with previous reports[18,38]. Notably, Ab binding

responses against the BA.5 and BQ.1.1 RBDs were similar (Fig. 2B-C), implying that RBD

binding responses are not always predictive of neutralizing activity against Omicron

subvariants.

Neutralizing responses against the XBB.1 variant were slightly lower than titers against

BQ.1.1 pseudovirus (Fig. 3D). Once again, the bivalent Wu1/BA.5 S-EABR mRNA-LNP

booster elicited the highest XBB.1 neutralizing titers, which were 2.1-fold higher than titers

induced by the monovalent Wu1 S mRNA-LNP booster (not statistically significant).

Based on day 154 titers, the monovalent Wu1 and bivalent Wu1/BA.5 S-EABR mRNA-

LNP boosters increased XBB.1 neutralization by 2.5- and 3.2-fold, respectively, while

conventional monovalent and bivalent S mRNA-LNP boosters only induced 1.5- and 2.2-

fold increases, respectively.

Notable differences were also observed in the number of mice in each cohort that reached

robust neutralizing titers (defined as ID_{50}s exceeding 1:100) against Omicron subvariants

following booster immunizations, titers that are likely important for the prevention of

symptomatic infections[39]. The primary vaccination series with Wu1 S mRNA-LNP elicited

robust neutralizing titers against the Wu1 variant in all mice on day 56 (Fig. 3F). However,

the initial Wu1 S mRNA-LNP vaccinations elicited robust titers in only 6 to 8, 1 to 3, and

0 to 2 of 10 mice against the BA.5, BQ.1.1, and XBB.1 variants on day 56, respectively.

On day 154, ID_{50}s over 1:100 were measured for 3 to 5 mice in each cohort against BA.5,

but for none of the animals against BQ.1.1 and XBB.1. While a third immunization with

Wu1 S mRNA-LNP induced robust titers against BA.5 in only 8 of 10 mice on day 244,

the monovalent Wu1 S-EABR mRNA-LNP booster induced robust titers against BA.5 in

all 10 mice, including the 4 mice that did not have detectable BA.5 neutralizing activity on

day 56 (Fig. 3F). As expected, robust neutralizing titers against the BA.5 variant were also

observed in all mice that received the bivalent Wu1/BA.5 S or S-EABR mRNA-LNP

boosters. Against the BQ.1.1 and XBB.1 variants, a third immunization with Wu1 S

mRNA-LNP induced robust neutralizing titers in only 2 and 3 mice, respectively (Fig. 3F).

In comparison, the bivalent Wu1/BA.5 S-EABR mRNA-LNP booster induced ID_{50}s over

1:100 in 7 and 6 mice against the BQ.1.1 and XBB.1 variants, respectively.

Overall, these results demonstrate that the EABR mRNA vaccine approach improves the

neutralizing activity against Omicron subvariants induced by monovalent and bivalent

boost immunogens in pre-vaccinated mice.

Bivalent S-EABR mRNA-LNP booster induces polyclass Ab responses.

Several distinct Ab-binding epitopes have been defined on RBDs, including the variable class 1 and 2 RBD epitopes and the more conserved class 3, 4, 1/4, and 5 RBD epitopes[33,40,41] (Fig. S2). We determined which RBD epitopes were targeted by booster-induced Abs in pre-vaccinated mice using deep mutational scanning (DMS) of yeast display libraries derived from Wu1[42] and XBB.1.5 RBDs[43]. In previous studies, we showed that equimolar mixtures of monoclonal Abs targeting distinct RBD epitopes produced less pronounced DMS escape peaks, which we defined as a polyclass Ab response[44]. This suggests that polyclonal serum that includes roughly equal distributions of Abs targeting multiple classes of RBD epitopes would exhibit relatively flat DMS profiles, while more defined DMS escape peaks would be observed for less diverse polyclonal serum responses containing Abs that primarily target particular RBD epitopes[44]. For this study, we used DMS escape fractions to define polyclass responses. Escape fractions for each RBD mutation range from 0 (no cells with this mutation are sorted into the serum Ab escape bin) to 1 (all cells with this mutation are sorted into the serum Ab escape bin)[45]. Total escape peaks represent the sum of escape fractions for all mutations at a specific site[45]. DMS profiles with total escape peaks below 0.5 at all sites were classified as polyclass responses. In contrast, DMS profiles with total escape peaks of 0.5-1, 1-2, or above 2 at one or more sites were classified as weak, moderate, or strong escape profiles, respectively, indicating weak, moderate, or strong targeting of particular RBD epitopes (Table S1). DMS was performed for 6 serum samples for each cohort, which were selected based on XBB.1 neutralization activity (1st, 2nd, 3rd, 5th, 6th, and 10th best

neutralizer from each cohort). For the Wu1 RBD library, only 5 DMS profiles were obtained

due to poor sequencing quality for one sample in each cohort.

Against the Wu1 RBD library, immunizations with the monovalent Wu1 S and S-EABR

mRNA-LNP boosters or the bivalent Wu1/BA.5 S mRNA-LNP booster resulted in DMS

profiles consistent with moderate targeting of the variable class 2 epitope primarily

inducing escape at RBD residue 484 (Fig. 4A-B). The bivalent Wu1/BA.5 S mRNA-LNP

booster also elicited weak to moderate responses that mapped to the variable class 1

and class 3 RBD epitopes. In contrast, the bivalent Wu1/BA.5 S-EABR mRNA-LNP

booster DMS profiles were consistent with polyclass Ab induction, suggesting a more

diverse response targeting multiple RBD epitopes (Fig. 4A-B). Notably, samples from 4

out of 5 mice in this cohort exhibited polyclass DMS profiles, whereas moderate to strong

escape fractions for class 1, 2, or 3 epitopes were observed in 3-4 mice in each of the

other cohorts (Fig. S7; Table S1).

As expected, no class 1 and 2 DMS profiles were observed against the XBB.1.5 RBD

library[43] since these epitopes are highly substituted in Omicron subvariants (Fig. S2). All

booster immunizations induced strong class 5 escape peaks targeting RBD residues 352,

357, 396, and 468 (Fig. 4C-D). Importantly, the bivalent Wu1/BA.5 S-EABR mRNA-LNP

booster induced a less pronounced overall class 5 profile than the other boost

immunogens, which was primarily driven by a strong class 5 response observed in a

single animal in this cohort (Fig. S8; Table S1). The other mice that received the bivalent

Wu1/BA.5 S-EABR mRNA-LNP booster exhibited polyclass or weak to moderate

responses against class 4 and 5 epitopes. In contrast, strong class 5 profiles were

detected in 4-5 mice in cohorts that received the monovalent Wu1 S or S-EABR mRNA-

LNP boosters or the bivalent Wu1/BA.5 S mRNA-LNP booster (Fig. S8; Table S1).

Overall, these results suggest that the bivalent Wu1/BA.5 S-EABR mRNA-LNP booster

induced a more diversified polyclass Ab response that likely contributed to the higher

neutralizing activity observed against Omicron subvariants, which could result in more

effective recognition of future variants.

Co-expression of Wu1 and BA.5 S results in heterotrimer formation.

The results from this and previous studies[19-22] demonstrated that bivalent booster

immunizations improve the potency and breadth of Ab responses against Omicron

subvariants compared to monovalent boosters. Serum depletion assays also showed that

the BA.5 neutralizing activity elicited by the bivalent Wu1/BA.5 S and S-EABR mRNA-

LNP boosters was primarily driven by cross-reactive Abs that bind to the Wu1 and BA.5

RBDs (Fig. S6). Preferential activation of cross-reactive B cells could be achieved through

co-display of Wu1 and BA.5 S trimers on the surface of a cell and/or an eVLP (Fig. 1C),

which could result in inter-S crosslinking between cross-reactive B cell receptors (BCRs)

and neighboring homotrimeric Wu1 and BA.5 S trimers (Fig. S9). As an additional

mechanism for bivalent booster-induced activation of cross-reactive B cells, we

hypothesized that co-expression of ancestral Wu1 and Omicron S in the same cell could

result in the formation of S heterotrimers consisting of Wu1 and Omicron S protomers and

could promote intra-S crosslinking, i.e., simultaneous binding of cross-reactive BCRs to

two neighboring RBDs within the same S trimer (Fig. S9). We previously postulated that

intra-S crosslinking is possible for several anti-RBD IgGs based on modeling of intact IgG

coordinates into single-particle cryo-electron microcopy (cryo-EM) structures of fragment

antigen-binding (Fab)-S complexes[33,40], and intra-S crosslinking was observed in EM

studies of IgG-S trimer complexes[46]. However, it is currently unknown whether co-

expression of Wu1 and Omicron Ss induces heterotrimerization, and if so, whether

heterotrimer formation affects the overall structure of the S ectodomain and influences

bivalent booster-induced immune responses.

To investigate whether S heterotrimers could form with different S protomers, we

designed 6P versions[32] of Wu1/BA.1 and Wu1/BA.1/Delta soluble heterotrimer Ss (HT2

and HT3, respectively) by co-transfection with specific C-terminally-tagged constructs.

HT2 was generated by co-transfecting StrepII-tagged Wu1 S and His-tagged BA.1 S,

while HT3 was generated using StrepII-tagged Wu1 S, His-tagged BA.1 S, and D7324[47]-

tagged Delta S (Fig. 5A). Each heterotrimer was subsequently purified using sequential

affinity purification specific to each tag, confirming the inclusion of distinct protomers in

the purified heterotrimers (Fig. 5A).

HT2 and HT3 S proteins were then evaluated using single-particle cryo-electron

microscopy (cryo-EM) to determine whether heterotrimeric Ss resembling their

homotrimeric counterparts were formed. To ensure that C-terminal tags did not disrupt

the formation of trimers, we included the following controls: (1) three homotrimers with

single tags (StrepII-tagged Wu1 S, His-tagged BA.1 S, and D7324-tagged Delta S), and

(2) a homotrimer comprised of His- and StrepII-tagged protomers (Wu1-His/Wu1-StrepII).

Single-particle cryo-EM analysis confirmed that all control S proteins formed trimers,

indicating that C-terminal tags did not disrupt trimerization (Fig. S10). Because most of

the EM constructions for homotrimeric S controls exhibited preferred orientations, we

determined overall conformational states (Fig. 5) for the control structures but did not fit

atomic models. Structural analysis also revealed trimerized HT2 and HT3 Ss (Fig. 5B-C,

Fig. S10). Although atomic models could not be built for the HT2 and HT3 trimer

reconstructions because the identities of individual protomers (Wu1 or BA.1 for HT2, and

Wu1, BA.1, or Delta for HT3) could not be conclusively determined at relatively low

resolution, the EM densities of S heterotrimers and control homotrimers showed no major

conformational differences in domain organization (Fig. 5B-C, Fig. S10). As previously

seen in SARS-CoV-2 S trimer structures[48,49], we observed two trimer populations in all

datasets—one with all three RBDs in the 'down' conformation and one with a single RBD

in an 'up' conformation (Fig. 5D). These data demonstrate heterotrimeric S formation for

soluble forms of SARS-CoV-2 S proteins. This result suggests that the bivalent S or S-

EABR mRNA-LNP boost immunogens likely induce heterotrimerization, which could

promote activation of cross-reactive B cells through intra-S cross-linking, a potential

mechanism warranting further investigation.

Discussion

The EABR mRNA vaccine platform delivers mRNA encoding engineered immunogens

that induce budding of enveloped virus-like particles (eVLPs) from the plasma membrane

to promote presentation of immunogens on cell surfaces and eVLPs[31]. This approach

elicited superior neutralizing Ab responses in naïve mice compared to a conventional

mRNA vaccine against original and variant SARS-CoV-2[31]. However, the majority of the

human population has already mounted immune responses against SARS-CoV-2 through

prior vaccinations, natural infections, or a combination of both. Here, we demonstrate that

the EABR mRNA vaccine approach improves humoral immune responses elicited by

monovalent and bivalent mRNA-LNP booster immunizations against Omicron subvariants

in pre-vaccinated mice.

The possibility of receiving a third dose of the original COVID-19 mRNA vaccine in fall

2021[11] coincided with the emergence of the Omicron variants in November 2021[12]. These

variants had acquired many substitutions, especially in the RBD, which allowed efficient

evasion of Ab-mediated immunity[13,14] and rapid global spread. Similar to the effects of a

third vaccination in humans[17], a third immunization in pre-vaccinated mice with a

monovalent Wu1 S mRNA-LNP booster elicited modest neutralizing activity against the

early Omicron variants BA.1 and BA.5. However, addition of an EABR sequence[31] to the

boosting immunogen resulted in higher Ab neutralizing titers against BA.1 and BA.5,

suggesting that a monovalent Wu1 S-EABR booster could have promoted enhanced

protection against early Omicron variants. Importantly, the monovalent Wu1 S-EABR

mRNA-LNP booster elicited similar neutralizing responses against the BA.5, BQ.1.1, and

XBB.1 subvariants compared to the conventional bivalent S mRNA-LNP booster that

contained both Wu1 and BA.5 S immunogens.

While updating of mRNA vaccines can be done relatively quickly, the development,

clinical testing, approval, and large-scale manufacturing of a bivalent booster that

includes updated immunogens based on new circulating variants cannot be accomplished

fast enough to keep up with viral evolution during a global pandemic such as COVID-19.

Indeed, the Wu1/BA.5 bivalent booster became available in fall 2022, almost one year

after the emergence of Omicron variants[30]. The results reported here suggest that the

EABR mRNA vaccine approach presents a potential solution that could reduce the need

for frequent development of updated COVID-19 boosters and potentially future pandemic

viruses. Combined with our previous finding that an initial two-shot vaccination series with

a Wu1 S-EABR mRNA-LNP immunogen elicited substantially higher neutralizing activity

against Omicron variants than conventional Wu1 S mRNA-LNP immunizations in naïve

mice[31], our results demonstrate that the EABR mRNA vaccine approach has the potential

to induce more effective and lasting humoral immunity against emerging SARS-CoV-2

variants and represents a promising platform for future pandemic preparedness.

Clinical studies showed that the bivalent COVID-19 mRNA booster outperformed the

monovalent mRNA booster in terms of preventing symptomatic illness, hospitalizations,

and deaths when administered as a fourth dose in humans[30,50]. Consistent with other

studies[19-23], the bivalent conventional and EABR-based boosters evaluated in this study

elicited higher neutralizing responses against Omicron subvariants than their monovalent

counterparts in pre-vaccinated mice, although improvements were relatively modest.

However, the bivalent Wu1/BA.5 S-EABR mRNA-LNP booster consistently elicited the

highest RBD binding and neutralizing titers against all tested Omicron subvariants and

induced robust neutralizing responses in the highest number of animals against the

BQ.1.1 and XBB.1 variants that are particularly difficult to neutralize[18,38] These results

highlight that the EABR mRNA vaccine approach and the bivalent booster strategy work

synergistically to induce more effective humoral immune responses.

In addition to eliciting the most potent neutralizing responses against Omicron

subvariants, the bivalent Wu1/BA.5 S-EABR mRNA-LNP booster also induced distinct

DMS signatures compared to the other boost immunogens, consistent with differences in

epitope mapping of polyclonal anti-RBD Ab responses. While the monovalent Wu1 S and

S-EABR mRNA-LNP and the bivalent Wu1/BA.5 S mRNA-LNP boosters elicited

predominantly class 1 and class 2 responses against Wu1 RBD and strong class 5

responses against XBB.1.5 RBD, boost immunizations with bivalent Wu1/BA.5 S-EABR

mRNA-LNP elicited polyclass serum Ab responses against Wu1 RBD and mostly

polyclass or moderate polyclass class 4 and class 5 responses against XBB.1.5 RBD.

These results imply that combining the bivalent booster strategy with the EARB mRNA

vaccine approach elicited a more diversified Ab response, resulting in simultaneous and

balanced targeting of multiple RBD epitopes. A more diverse anti-RBD Ab response might

be desirable as viral immune evasion would require mutations in multiple sites, including

the more conserved class 4 and 5 RBD epitopes. As a potential mechanism for the more

frequent induction of polyclass-like Ab responses, we hypothesize that EABR eVLP

formation promotes close proximity of Wu1 and BA.5 S trimers on densely-coated eVLPs,

consistent with cryo-electron tomography imaging of purified eVLPs showing that the

density of S trimers was equivalent or higher than on SARS-CoV-2 virions (20-40 S

trimers on 40-60 nm diameter eVLPs)[31]. The high density of co-displayed Wu1 and BA.5

S trimers on eVLPs should facilitate inter-S crosslinking by B cell receptors that recognize

epitopes that are conserved between Wu1 and BA.5 S trimers, leading to preferential

activation of cross-reactive B cells. By comparison, conventional bivalent mRNA booster-

mediated co-expression of Wu1 and BA.5 S trimers could result in lower S trimer density

on cell surfaces that are crowded with endogenous membrane proteins, which could

prevent efficient inter-S crosslinking.

As an additional mechanism for bivalent booster-induced preferential activation of cross-

reactive B cells, we provide evidence that co-expression of ancestral and Omicron S

results in heterotrimer formation that could promote intra-S crosslinking, i.e.,

simultaneous binding of both Fabs of a cross-reactive BCR to neighboring RBDs within

the same S trimer. We previously postulated that intra-S crosslinking is possible for

several anti-RBD Abs that target various RBD epitopes based on modeling of intact IgG

structures onto cryo-EM structures of Fab-S complexes[33,40]. The conventional mRNA-

and the EABR mRNA-based bivalent boost immunogens should promote similar degrees

of heterotrimerization and intra-S crosslinking. While our RBD serum depletion assays

showed that both bivalent boosters primarily elicited cross-neutralizing Abs that bind to

the Wu1 and BA.5 RBDs, only the EABR mRNA-based bivalent booster induced

predominantly polyclass DMS signatures. Thus, it is possible that the diversification of the

anti-RBD Ab response observed for the bivalent EABR mRNA booster was primarily

driven by inter-S crosslinking possibly in conjunction with intra-S crosslinking. Future

studies are needed to investigate the impact of inter- and intra-S crosslinking on the

activation of cross-reactive B cells and to evaluate whether the EABR mRNA vaccine

approach can improve other types of bivalent and multivalent vaccines.

532 Although our results highlight the potential of the EABR mRNA vaccine approach in the

533 context of pre-existing immunity, we observed that monovalent and bivalent EABR mRNA

534 booster-induced humoral immune responses were still largely dictated by immunological

535 imprinting resulting from the initial vaccination series. Post-boost BA.5 neutralizing

536 responses were roughly an order of magnitude lower than Wu1 neutralization titers, even

537 for bivalent boosters that included BA.5 S immunogens. In addition, despite eliciting

538 consistent polyclass anti-RBD Ab responses, bivalent EABR mRNA booster-induced

539 neutralizing activity against the BQ.1.1 and XBB.1 variants dropped over 10-fold

540 compared to BA.5 neutralization. Notably, Ab binding responses to BQ.1.1 and XBB.1

541 RBDs were similar compared to BA.5 RBD binding for all boost immunogens. This

542 discrepancy between RBD binding responses and neutralization titers indicates that only

543 a relatively small fraction of binding Abs has neutralizing activity and/or the Abs still bind

544 to the BQ.1.1 and XBB.1 RBDs, but fail to effectively prevent interactions with ACE2 on

545 the surface of target cells. The initial two-shot COVID-19 mRNA vaccination series and

546 all booster immunizations elicited potent binding responses against the S2 region, which

547 is relatively conserved among SARS-CoV-2 variants[34]. Consistent with previous reports[51],

548 our depletion studies showed that anti-S2 Abs had little impact on virus neutralization,

549 which was instead predominantly driven by anti-RBD Abs. However, anti-S2 Abs could

550 still provide valuable immune protection against symptomatic Omicron infections through

551 Fc-mediated effector functions[52]. In addition, mRNA vaccines induce potent cytotoxic T

552 cell responses that cross-recognize Omicron variants[53] and reduce COVID-19 disease

553 severity[54].

Taken together, our results demonstrate that the EABR mRNA vaccine approach

improves monovalent and bivalent booster-mediated humoral immune responses against

Omicron subvariants in pre-vaccinated mice. DMS epitope mapping studies further

showed that the bivalent EABR mRNA booster elicited a more diversified anti-RBD Ab

response than all other tested boost immunogens. Our work also provides evidence that

bivalent mRNA booster immunizations promote heterotrimer formation, which could

activate cross-reactive B cells through intra-S crosslinking. Despite these advances,

future work is needed to develop booster strategies that overcome immunological

imprinting more effectively and confer lasting immunity against emerging variants.

Methods

Design of EABR constructs

SARS-CoV-2 S-EABR constructs were designed as described[31]. The EABR domain

(residues 160-217) of the human CEP55 protein was fused to the C-terminus of the

SARS-CoV-2 S protein obtained from the Wu1 or BA.5 variants separated by a 4-residue

(Gly)$_3$Ser (GS) linker (Fig. S1A). These constructs contained the native furin cleavage

site, 2P stabilizing mutations[55], and the C-terminal 21 residues were truncated to remove

an endoplasmic reticulum retention signal[56]. The D614G substitution[57], naturally present

in BA.5, was inserted into Wu1 S and S-EABR constructs. To prevent endocytosis and

enhance the cell surface expression of S-EABR constructs, an EPM (residues 243-290

of mouse FcgRII-B1)[31,58] was inserted upstream of the 4-residue Gly-Ser linker and the

EABR domain as described[31] (Fig. S1A).

In vitro transfection

To verify that the designed constructs induce eVLP budding, HEK293T were transiently

transfected with DNA plasmids encoding Wu1 or BA.5 S-EABR using Fugene HD

(Promega). After 48 hours, transfected cell supernatants were harvested, and eVLPs

were purified by ultracentrifugation on a 20% sucrose cushion as described[31]. The purified

eVLP fractions were 3-fold serially diluted in 0.1 M NaHCO$_3$ pH 9.6, directly coated onto

Costar high binding 96-well plates (Corning), and eVLP-associated S protein levels were

quantified by ELISA. After 1 hour blocking in Tris-buffered saline with 0.1% TWEEN 20

(TBS-T) and 3% bovine serum albumin (BSA) (TBS-T/3% BSA) at room temperature, the

monoclonal anti-S2 Ab S2P6[59] was diluted to a concentration of 5 µg/mL in TBS-T/3%

BSA and added to plates for 1 hour at room temperature. After washing with TBS-T, HRP-

conjugated goat anti-human IgG (2015-05; SouthernBiotech) was diluted 1:8,000 in TBS-

T/3% BSA and added to plates for 30 min at room temperature. After washing with TBS-

T, plates were developed using 1-Step™ Ultra TMB-ELISA Substrate Solution (Thermo

Fisher Scientific) and absorbance was measured at 450 nm.

mRNA synthesis

Codon-optimized mRNAs encoding Wu1 S, Wu1 S-EABR, BA.5 S, and BA.5 S-EABR

constructs were synthesized by RNAcore (https://www.houstonmethodist.org/research-

cores/rnacore/) using proprietary manufacturing protocols as described[31]. In brief,

mRNAs were synthesized by T7 RNA polymerase-mediated in vitro transcription using

DNA templates encoding the immunogen open reading frame flanked by 5′ and 3'

untranslated regions and terminated by an encoded polyA tail. CleanCap 5' cap structures

(TriLink) were co-transcriptionally incorporated into the 5′ end. Uridine was completely

substituted with N1-methyl-pseudouridine to lower immunogenicity[60]. mRNAs were

purified by oligo-dT affinity purification and high-performance liquid chromatography

(HPLC) to remove double-stranded RNA contaminants[61]. Purified mRNAs were stored

at –80 °C.

LNP encapsulation of mRNAs

Purified mRNAs were formulated in LNP as described[31,62]. In brief, 1,2-distearoyl-*sn*-

glycero-3-phosphocholine, cholesterol, a PEG lipid, and an ionizable cationic lipid

dissolved in ethanol were rapidly mixed with an aqueous acidic solution containing the

mRNAs using an in-line mixer. The ionizable lipid and LNP composition are described in

the international patent application WO2017075531(2017). The post in-line solution was

dialyzed with PBS to remove ethanol and displace the acidic solution. Subsequently,

mRNA-LNP were measured for size (60-65 nm) and polydispersity (PDI < 0.075) by

dynamic light scattering (Malvern Nano ZS Zetasizer). The encapsulation efficiencies

were >97% as measured by the Quant-iT Ribogreen Assay (Invitrogen).

Immunizations

All animal procedures in this study were performed in compliance with the U.S.

Department of Agriculture's (USDA) Animal Welfare Act (9 CFR Parts 1, 2, and 3); the

Guide for the Care and Use of Laboratory Animals (Institute of Laboratory Animal

Resources, National Academy Press, Washington, D.C., 2011); and the National

Institutes of Health, Office of Laboratory Animal Welfare. Whenever possible, procedures

in this study were designed to avoid or minimize discomfort, distress, and pain to animals.

Animals were observed for signs of morbidity and/or mortality at least once daily,

according to animal welfare act and the Association for Assessment and Accreditation of

Laboratory Animal Care International (AAALACi). All mouse procedures were approved

by the Labcorp Institutional Animal Care and Use Committee (IACUC).

7-8 week-old female BALB/c mice (Charles River Laboratories) were purchased and

housed at Labcorp Drug Development, Denver, PA for immunization studies. All animals

were healthy upon receipt and were weighed and monitored during a 7-day acclimation

period before the start of the study. Mice were randomly assigned to experimental cohorts

of 10 animals. Up to 10 mice were co-housed together, and cages were kept in a climate-

controlled room at 63 to 77ºC at 50±20% relative humidity. Mice were provided Rodent

Diet #5001 (Purina Lab Diet) ad libitum.

2 µg of Wu1 S mRNA-LNP were administered to all cohorts of mice by IM injections on

days 0 and 28. On day 230, pre-vaccinated mice received different boosters by IM

injection: i) 2 µg Wu1 S mRNA-LNP; ii) 2 µg Wu1 S-EABR mRNA-LNP; iii) 1 µg Wu1 S

mRNA-LNP + 1 µg BA.5 S mRNA-LNP; iv) 1 µg Wu1 S-EABR mRNA-LNP + 1 µg BA.5

S-EABR mRNA-LNP. For bivalent boosters (iii and iv), the Wu1 and BA.5 S or S-EABR

mRNA-LNP were mixed prior to the IM injections. Serum samples for ELISAs and

neutralization assays were obtained on indicated days by retro-orbital bleeding. One

small drop of either 0.5% proparacaine hydrochloride ophthalmic solution or 0.5%

tetracaine hydrochloride ophthalmic solution was applied in the eye of the mouse prior to

performing retro-orbital blood collection. Eyes were alternated for each blood collection.

Euthanasia was performed in accordance with American Veterinary Medical Association

(AVMA) guidelines for euthanasia. Mice were individually placed in a CO2 chamber and

rendered unconscious. Mice were verified as unconscious by lack of response when toe

was pinched. Unconscious mice were subjected to bilateral jugular vein severance for

terminal exsanguination. Mice were placed back in the CO2 chamber until death

occurred. A double thoracotomy was performed as a second confirmation of death.

Protein expression and purification

Soluble SARS-CoV-2 S trimers (Wu1/D614G, BA.5, BQ.1.1, XBB.1)[32], RBDs (Wu1, BA.5,

BQ.1.1, XBB.1), and a prefusion-stabilized S2 protein[63] were expressed as described[37,64].

Briefly, Avi/His-tagged S trimers and His-tagged RBD and S2 proteins were purified from

transiently-transfected Expi293F cells (Gibco) by nickel affinity chromatography (HisTrap,

Cytiva) and size-exclusion chromatography (SEC) (Superose 6 Increase 10/300,

Cytiva)[33,37,64]. Peak fractions corresponding to S, RBD, or S2 proteins were pooled,

concentrated, and stored at 4ºC. Biotinylated S trimers for ELISAs were generated by co-

transfection of Avi/His-tagged S constructs with a plasmid encoding an endoplasmic

reticulum-directed BirA enzyme (kind gift from Michael Anaya, Caltech).

For heterotrimeric constructs, Wu1/BA.1 (HT2) and Wu1/BA.1/Delta (HT3) heterotrimers

were expressed via the addition of C-terminal tags and co-transfection of Expi293F cells

(Gibco) with specific plasmid DNA ratios. HT2 heterotrimers were produced by co-

transfecting a 1:1 plasmid DNA ratio of StrepII-tagged Wu1 S and His-tagged BA.1 S

constructs. HT3 heterotrimers were produced by co-transfecting a 1:1:1 plasmid DNA

ratio of StrepII-tagged Wu1 S, His-tagged BA.1 S, and D7324[47]-tagged Delta S

constructs. HT2 was sequentially purified using StrepTrap (Cytiva), HisTrap (Cytiva), and

SEC, while HT3 also underwent immunoaffinity chromatography using the D7324-specific

monoclonal Ab JR-52[47] (kind gift from James Robinson, Tulane University) before SEC.

The Wu1-His/Wu1-StrepII S homotrimer was generated by co-transfecting Expi293F cells

with a 1:1 plasmid DNA ratio of His-tagged Wu1 S and StrepII-tagged Wu1 S, followed by

purification as described for the HT2 heterotrimer. Single-tagged S homotrimers included

StrepII-tagged Wu1 S, His-tagged BA.1 S, and D7324-tagged Delta S, each purified using

their respective affinity chromatography steps followed by SEC.

Single-particle cryo-EM sample preparation

Grids of SARS-CoV-2 S homotrimers and heterotrimers were prepared at a protein

concentration of 2 mg/mL. Prior to freezing, fluorinated octylmaltoside solution (Anatrace)

was added to the S proteins at a final concentration of 0.02%(w/v). Three µL of S proteins

were immediately applied to Quantifoil 300 mesh 1.2/1.3 grids (Electron Microscopy

Sciences) that were freshly glow discharged with the PELCO easiGLOW (Ted Pella) for

60 seconds at 20 mA. Grids were blotted for 3 seconds using Whatman No.1 filter paper

with 0 blot force, 100% humidity and room temperature, and vitrified in 100% liquid ethane

using a Vitrobot Mark IV system (Thermo Fisher Scientific).

Single-particle cryo-EM data collection and processing

Single-particle cryo-EM datasets for SARS-CoV-2 S homotrimers and heterotrimers were

collected using SerialEM automatic data collection software[65] on a 300 keV Titan Krios

(Thermo Fisher Scientific) cryo-electron microscope equipped with a K3 direct electron

detector camera (Gatan). Movies were acquired in 40 frames with a defocus range of -1

to -3 μm and a total dosage of 60 electrons per $Å^2$ using the 3 by 3 beam image shift

pattern with 3 exposures per hole in the super-resolution mode at a pixel size of 0.416 Å.

Data processing workflows are outlined in Figure S10. For all datasets, motion correction

was done with patch motion correction with a binning factor of 2 and contrast transfer

function (CTF) parameters were estimated with Patch CTF in cryoSPARC[66]. Particles with

a diameter of 100 to 200 Å were picked with blob picker, inspected, extracted, and finally

classified with 2D classification in cryoSPARC[66]. Ice particles, junk particles, and particles

with preferred orientations were discarded, and the remaining particles were used for *ab*

initio reconstruction of three to four volumes. These reconstructions were further refined

with heterogeneous refinement in cryoSPARC[66]. Depending on the state for the receptor-

binding domains (RBDs) in the reconstructions, particles were combined for

homogeneous and non-uniform refinements to obtain the final reconstructions in

cryoSPARC[66].

ELISAs

ELISAs were performed as described[31,44]. To analyze serum binding to S trimers, pre-

blocked streptavidin-coated Nunc® MaxiSorp™ 384-well plates (ThemoFisher Scientific)

were coated with 5 μg/mL biotinylated S trimers in TBS-T/3% BSA for 1 hour at room

temperature. To analyze serum binding to RBDs or S2, Nunc® MaxiSorp™ 384-well

plates (Sigma) were coated with 2.5 μg/mL RBD or S2 in 0.1 M NaHCO$_3$ pH 9.8 and

incubated overnight at 4°C. Plates were blocked with TBS-T/3% BSA for 1 hour at room

temperature followed by removal of the blocking buffer by aspiration. Serum samples from

immunized mice were diluted 1:100, 4-fold serially diluted in TBS-T/3% BSA, and added

to plates. After a 3-hour incubation at room temperature, plates were washed with TBS-

T using an automated plate washer. HRP-conjugated goat anti-mouse IgG (715-035-150;

Jackson ImmunoResearch) was diluted 1:100,000 in TBS-T/3% BSA and added to plates

for 1 hour at room temperature. After washing with TBS-T, plates were developed using

SuperSignal™ ELISA Femto Maximal Signal Substrate (Thermo Fisher Scientific) and

absorbance was measured at 425 nm. Area under the curve (AUC) calculations for

binding curves were performed using GraphPad Prism 9.3.1 assuming a one-site binding

model with a Hill coefficient as described[67].

Neutralization assays

Lentivirus-based SARS-CoV-2 pseudoviruses were generated as described[35,36] using S

proteins from the Wu1/D614G, BA.1, BA.5, BQ.1.1, and XBB.1 variants in which the C-

terminal 21 residues of the S cytoplasmic tails were removed[35]. Serum samples from

immunized mice were heat-inactivated for 30 min at 56°C. Three-fold serial dilutions of

heat-inactivated samples were incubated with pseudoviruses for 1 hour at 37°C, followed

by addition of the serum-virus mixtures to pre-seeded HEK293T-ACE2 target cells. After

48-hour incubation at 37°C, BriteLite Plus substrate (Perkin Elmer) was added and

luminescence was measured. Half-maximal inhibitory dilutions (ID$_{50}$s) were calculated

using 4-parameter non-linear regression analysis in AntibodyDatabase[68] and ID_{50} values

were rounded to three significant figures.

Ab depletions

HiTrap™ NHS-activated resin (Cytiva) was coupled to purified SARS-CoV-2 Wu1 RBD or

a mock protein (HIV-1 gp120) at a 10 mg/mL concentration and stored in PBS

supplemented with 0.02% sodium azide at 4°C. For Ab depletions, 50 μL dry worth of

RBD-coupled resin were aliquoted into microcentrifuge tubes and spun down at 13,000 x

g for 2 minutes. The supernatant was discarded. The resin was washed with 100 μL of

sterile PBS to remove excess PBS-azide solution. The supernatant was again discarded.

10 μL of mouse serum was then diluted in 90 μL PBS, and added to the resin. The tubes

were mixed and incubated at room temperature for 10 minutes. The resin-serum mixture

was spun down at 13,000 x g for 7 minutes to allow the resin to settle, and 90 μL of the

supernatant was transferred into a fresh microcentrifuge tube. To ensure no resin was

present in the final sample, the collection tube was spun down again, and supernatant

was transferred to a fresh microcentrifuge tube. Sterile PBS was added for a final volume

of 200 μL. Prior to neutralization assays, the Ab depleted serum samples were heat-

inactivated at 56°C for 30 minutes.

Deep mutational scanning

759   DMS studies to map RBD epitopes recognized by serum Abs were performed in biological

760   duplicates using independent mutant RBD yeast libraries (Wu1[42,69] and XBB.1.5 RBD[43]

761   libraries were generously provided by Tyler Starr, University of Utah) as

762 described[44,64,70,71]. Briefly, yeast libraries were stained with serum Abs, labeled with a

763 secondary Ab (115-605-008, Jackson ImmunoResearch), and then processed using a

764 Sony SH800 cell sorter. Cells were gated to capture RBD mutants that had reduced Ab

765 binding for a relatively high degree of RBD expression. Ab-escaped cells were expanded

766 for DNA extraction and Illumina sequencing was done as described[44,70,71]. Raw

767 sequencing data are available on the NCBI SRA under BioProject PRJNA1067836.

768 Escape fractions were computed using previously described processing steps[44,70,71] and

769 implemented using a Swift DMS program available from authors upon request.

771 Escape map visualizations shown as static line plots, logo plots and structural depictions

772 were all created using Swift DMS as previously described[41,44,71]. In some visualizations,

773 RBD sites were categorized based on epitope region, which were defined as class 1 (pink)

774 (RBD residues 403, 405, 406, 417, 420, 421, 453, 455-460, 473-478, 486, 487, 489, 503,

775 504), class 2 (purple) (residues 472, 479, 483-485, 490-495), class 3 (blue) (residues

776 341, 345, 346, 437-450, 496, 498-501), class 4 (orange) (residues 365-390, 408)[72], or

777 class 5 (green) (352-357, 396, 462-468)[41]. Structural visualizations were performed using

778 an RBD surface (PDB 6M0J) colored by the site-wise escape metric at each site as

779 described[44].

781 **Statistical analysis**

782 Statistically significant titer differences between immunized cohorts of BALB/c mice (10

783 mice per cohort) for ELISAs were determined using analysis of variance (ANOVA) test

784 followed by Tukey's multiple comparison post hoc test calculated using Graphpad Prism

9.3.1. Titer differences between immunized cohorts of BALB/c mice for Wu1, BA.5,

BQ.1.1, and XBB.1 neutralization assays were determined using the non-parametric

Kruskal-Wallis test followed by Dunn's multiple comparison post hoc test calculated using

Graphpad Prism 9.3.1. For BA.1 neutralization titers, statistically significant differences

were determined using the non-parametric two-tailed Mann-Whitney test calculated using

Graphpad Prism 9.3.1.

Figure 1. Monovalent and bivalent boosters elicit similar Ab binding responses against variant S trimers in pre-vaccinated mice. A) Immunization schedule in BALB/c mice. Blood samples were collected on days 14, 56, 154, and 244. **B)** Immunization regimens for each cohort. All cohorts were initially vaccinated with 2 µg of Wu1 S mRNA-

LNP on days 0 and 28 by IM injection. Pre-vaccinated mice received different boost

immunogens on day 230 by IM injection. Cohorts 1-4 received 2 µg of monovalent Wu1

S mRNA-LNP (cyan), monovalent Wu1 S-EABR mRNA-LNP (orange), bivalent Wu1/BA.5

S mRNA-LNP (blue), or bivalent Wu1/BA.5 S-EABR mRNA-LNP (maroon), respectively.

C) Schematic showing the different boost immunogens that were administered on day

230. Wu1 S mRNA-LNP immunization results in cell surface presentation of membrane-

anchored Wu1 S trimers. Wu1 S-EABR mRNA-LNP immunization results in dual

presentation of Wu1 S trimers on cell surfaces and on eVLPs that are secreted after

budding from the plasma membrane. eVLP formation is induced by EABR-mediated

recruitment of host cell proteins from the ESCRT pathway. Bivalent Wu1/BA.5 S mRNA-

LNP immunization results in cell surface presentation of membrane-anchored Wu1 and

BA.5 S trimers. Wu1/BA.5 S-EABR mRNA-LNP immunization results in dual presentation

of Wu1 and BA.5 S trimers on cell surfaces and eVLPs that bud from the plasma

membrane. S trimers are shown as homotrimers for both monovalent and bivalent boost

immunogens. **D-G)** ELISA data from indicated time points for antisera from individual mice

(colored circles) presented as the geometric mean (bars) and standard deviation

(horizontal lines) of area under the curve (AUC) values. ELISAs evaluated binding of

purified Ss from the **D)** Wu1, **E)** BA.5, **F)** BQ.1.1, and **G)** XBB.1 variants. The $\log_{10}$-fold

change in binding titers from day 154 to day 244 is also shown for each S (plotted as

mean (bars) and standard deviation (horizontal lines)).

Figure 2. Bivalent S-EABR mRNA-LNP booster elicits potent Ab responses targeting Omicron RBDs. A-D) RBD binding titers for antisera from pre-vaccinated mice that received booster immunizations with 2 µg of monovalent Wu1 S mRNA-LNP (cyan), 2 µg of monovalent Wu1 S-EABR mRNA-LNP (orange), 2 µg of bivalent Wu1/BA.5 S mRNA-LNP (blue), or 2 µg of bivalent Wu1/BA.5 S-EABR mRNA-LNP (maroon) (immunization schedule in Fig. 1A-B). ELISA data are shown for indicated time points for antisera from individual mice (colored circles) presented as the geometric mean (bars) and standard deviation (horizontal lines) of area under the curve (AUC) values. ELISAs evaluated binding of RBDs from the **A)** Wu1, **B)** BA.5, **C)** BQ.1.1, and **D)** XBB.1 SARS-CoV-2 variants. The $\log_{10}$-fold change in binding titers from day 154 to day 244 is also shown for each RBD (plotted as mean (bars) and standard deviation (horizontal lines)).

830   Significant differences between cohorts linked by horizontal lines are indicated by

831   asterisks: $p < 0.05$ = *, $p < 0.01$ = **.

Figure 3. Bivalent Wu1/BA.5 S-EABR mRNA-LNP booster elicits higher neutralizing responses against Omicron subvariants. A-D) Neutralization titers for antisera from pre-vaccinated mice that received booster immunizations with 2 µg of monovalent Wu1 S mRNA-LNP (cyan), 2 µg of monovalent Wu1 S-EABR mRNA-LNP (orange), 2 µg of bivalent Wu1/BA.5 S mRNA-LNP (blue), or 2 µg of bivalent Wu1/BA.5 S-EABR mRNA-LNP (maroon) (immunization schedule in Fig. 1A-B). Neutralization data are shown for indicated time points for antisera from individual mice (colored circles) presented as the geometric mean (bars) and standard deviation (horizontal lines) of ID_{50} values. Results are shown for **A)** Wu1, **B)** BA.5, **C)** BQ.1.1, and **D)** XBB.1 pseudoviruses. Dashed

horizontal lines correspond to the background values representing the limit of detection

for neutralization assays (ID_{50} = 1:50). The $\log_{10}$-fold change in neutralization titers from

day 154 to day 244 is also shown for each pseudovirus (plotted as mean (bars) and

standard deviation (horizontal lines)). Significant differences between cohorts linked by

horizontal lines are indicated by asterisks: $p<0.05$ = *, $p<0.01$ = **. **E)** Number of mice in

each cohort that reached an $ID_{50} \leq$ 1:100 against each variant on days 56 (left), 154

(middle), and 244 (right).

Figure 4. Bivalent S-EABR mRNA-LNP booster induces polyclass Ab responses.
Sera for DMS studies were derived from pre-vaccinated mice that received booster immunizations with 2 µg of monovalent Wu1 S mRNA-LNP, monovalent Wu1 S-EABR mRNA-LNP, bivalent Wu1/BA.5 S mRNA-LNP, or bivalent Wu1/BA.5 S-EABR mRNA-LNP (immunization schedule in Fig. 1A-B). DMS was performed for 6 serum samples for each cohort, which were selected based on XBB.1 neutralization activity (1st, 2nd, 3rd, 5th, 6th, and 10th best neutralizer from each cohort). For the Wu1 RBD library, only five DMS profiles were obtained for each cohort. **A)** Line plots for DMS results using Wu1 (top) or XBB.1.5 (bottom) RBD libraries for sera from pre-vaccinated mice that received the indicated boost immunogens. The x axes display the RBD residue numbers, and the y

axes show the sum of the Ab escape of all mutations at a site (larger numbers indicating

increased Ab escape). Each faint line shows one antiserum with heavy lines representing

the average across the n = 5 (top) or n = 6 (bottom) sera in each cohort. Lines are colored

differently for RBD epitopes within different RBD epitope classes[33,40] (color definitions

shown in legend above line plots; epitopes are shown in Fig. S2; gray color for residues

not assigned to an epitope). **B)** The average site-total Ab escape for the Wu1 (top) or

XBB.1.5 (bottom) RBD libraries for pre-vaccinated mice that received the indicated boost

immunogens mapped to the Wu1 RBD surface (PDB: 6M0J). Gray coloring indicates no

escape, and red coloring indicates escape. Lines with numbers colored according to

epitope class indicate RBD positions with the most escape.

A

Wu1/BA.1 S heterotrimer (HT2)

1:1 transfection ratio

Wu1 S | StrepII tag
BA.1 S | His tag

StrepII affinity purification → His affinity purification → SEC

Wu1/BA.1/Delta S heterotrimer (HT3)

1:1:1 transfection ratio

Wu1 S | StrepII tag
BA.1 S | His tag
Delta S | D7324 tag

StrepII affinity purification → His affinity purification → D7324 affinity purification → SEC

B Wu1/BA.1 S (HT2)　　　　**C** Wu1/BA.1/Delta S (HT3)

Class I: 3 'down' RBDs　　Class II: 1 'up' RBD　　Class I: 3 'down' RBDs　　Class II: 1 'up' RBD

D

Trimers/Heterotrimers	Conformational states	
	0 'up' RBD 3 'down' RBDs	1 'up' RBD 2 'down' RBDs
Wu1/BA.1 S (HT2)	✔	✔
Wu1/BA.1/Delta S (HT3)	✔	✔
Wu1 S Tag Control (StrepII/His)	✔	✔
Wu1 S	✔	✔
BA.1 S	✔	✔
Delta S	✔	✔

Figure 5. Heterotrimer S formation occurs for soluble constructs. A) Schematic for

the generation and purification of soluble SARS-CoV-2 Wu1/BA.1 (HT2) and

Wu1/BA.1/Delta (HT3) S heterotrimers. HT2 heterotrimers were generated by co-

transfection of StrepII-tagged Wu1 S and His-tagged BA.1 S constructs at a 1:1 DNA

ratio, while HT3 heterotrimers were produced by co-transfecting StrepII-tagged Wu1 S,

His-tagged BA.1 S, and D7324-tagged Delta S constructs at a 1:1:1 ratio. Each

heterotrimer was sequentially purified using affinity chromatography specific for each tag

(StrepII, His, or JR-52) to ensure incorporation of distinct protomers. **B)** Cryo-EM density

maps for Wu1/BA.1 HT2 S class I with 3 'down' RBDs (3.8 Å resolution) and class II with

1 'up' RBD (3.8 Å resolution). **C)** 4.0 Å and 3.9 Å resolution cryo-EM density maps for

Wu1/BA.1/Delta HT3 S with 3 'down' RBDs and 1 'up' RBD. **D)** Table summarizing S

heterotrimer and control homotrimer conformational states.

A

EPM   ALPGNPDHREMGETLPEEVGEYRQPSGGSVPVSPGPPSGLEPTSSSPY
EABR FNSSINNIHEMEIQLKDALEKNQQWLVYDQQREVYVKGLLAKIFELEKKTETAAHSLP

Figure S1. Wu1 and BA.5 S-EABR constructs induce efficient eVLP budding. A) Design and sequence information for SARS-CoV-2 S-EABR constructs. Top: The Wu1 and BA.5 S (including a furin cleavage site, 2P stabilizing substitutions, the D614G substitution, and ΔCT, a cytoplasmic tail deletion) were fused to an EPM sequence, a (Gly)$_3$Ser (GS) spacer, and an EABR sequence. EPM = Endocytosis prevention motif. GS = (Gly)$_3$Ser linker. EABR = ESCRT- and ALIX-binding region. Bottom: EPM and EABR sequence information. **B)** DNA plasmids encoding Wu1 S-EABR (red) or BA.5 S-EABR (orange) constructs were transfected into HEK293T cells. 48 hours post-transfection, transfected cell culture supernatants were harvested, and eVLPs were purified by

896 ultracentrifugation on a 20% sucrose cushion. Purified eVLP samples were serially

897 diluted, directly coated onto 96-well plates, and eVLP-associated S protein levels were

898 quantified by ELISA.

Figure S2. Sarbecovirus RBD sequence conservation. A) Sequence conservation of 16 sarbecovirus RBDs calculated with ConSurf[73] and plotted on a surface representation of a SARS-CoV-2 RBD (PDB 7BZ5). Anti-RBD class 1, 2, 3, 4, 1/4, and 5 Ab epitopes are outlined with dotted lines in different colors using structural information from representative structures of mAb bound to SARS-CoV-2 S or RBD (C102: PDB 7K8M; C002: PDB 7K8T, S309: PDB 7JX3; CR3022: PDB 7LOP; C118: PDB 7RKV; WRAIR-

907   2063: PDB 8EOO). SARS-CoV-2 RBD substitutions (colored in red) in Omicron **B)** BA.1,

908   **C)** BA.5, **D)** BQ.1.1, and **E)** XBB.1 variants plotted on the RBD sequence conservation

909   plot.

ELISA / S2

Figure S3. Primary vaccinations and booster immunizations elicit potent anti-S2 binding responses. S2 binding titers for antisera from pre-vaccinated mice that received booster immunizations with 2 µg of monovalent Wu1 S mRNA-LNP (cyan), 2 µg of monovalent Wu1 S-EABR mRNA-LNP (orange), 2 µg of bivalent Wu1/BA.5 S mRNA-LNP (blue), or 2 µg of bivalent Wu1/BA.5 S-EABR mRNA-LNP (maroon). ELISA data for binding of Wu1 S2 are shown for indicated time points for antisera from individual mice (colored circles) presented as the geometric mean (bars) and standard deviation (horizontal lines) of area under the curve (AUC) values. The $\log_{10}$-fold change in binding titers from day 154 to day 244 is also shown (plotted as mean (bars) and standard deviation (horizontal lines)).

Figure S4. Monovalent and bivalent S-EABR mRNA-LNP boosters elicit higher increases in BA.5 neutralization compared to conventional S mRNA-LNP boosters. Neutralization titers for antisera from pre-vaccinated mice that received booster immunizations with 2 µg of monovalent Wu1 S mRNA-LNP (cyan), 2 µg of monovalent Wu1 S-EABR mRNA-LNP (orange), 2 µg of bivalent Wu1/BA.5 S mRNA-LNP (blue), or 2 µg of bivalent Wu1/BA.5 S-EABR mRNA-LNP (maroon) (immunization schedule in Fig. 1A-B). BA.5 neutralization data are shown for indicated time points for antisera from individual mice (colored circles) presented as the geometric mean (bars) and standard deviation (horizontal lines) of ID_{50} values. Dashed horizontal lines correspond to the background values representing the limit of detection for neutralization assays ($ID_{50} = 1:50$). The $\log_{10}$-fold change in neutralization titers from day 56 to day 244 is also shown (plotted as mean (bars) and standard deviation (horizontal lines)). Significant differences

936   between cohorts linked by horizontal lines are indicated by asterisks: p<0.05 = *, p<0.01

937   = **.

Figure S5. Monovalent Wu1 S-EABR mRNA-LNP booster elicits improved neutralizing responses against BA.1 variant. Neutralization titers for antisera from pre-vaccinated mice that received booster immunizations with 2 µg of monovalent Wu1 S mRNA-LNP (cyan) or monovalent Wu1 S-EABR mRNA-LNP (orange). Neutralization data are shown for indicated time points for antisera from individual mice (colored circles) presented as the geometric mean (bars) and standard deviation (horizontal lines) of ID_{50} values against BA.1 pseudovirus. Dashed horizontal lines correspond to the limit of detection for neutralization assays (ID_{50} = 1:50). The log_{10}-fold change in neutralization titers from day 154 to day 244 is also shown (plotted as mean (bars) and standard deviation (horizontal lines)). Significant differences between cohorts linked by horizontal lines are indicated by asterisks: $p < 0.05$ = *.

Figure S6. Depleting bivalent booster-induced antisera with Wu1 RBD reduces BA.5 neutralizing activity. Day 244 serum samples obtained from pre-vaccinated mice that received booster immunizations with 2 µg of bivalent Wu1/BA.5 S mRNA-LNP (blue) or 2 µg of bivalent Wu1/BA.5 S-EABR mRNA-LNP (maroon) were depleted using a mock protein (HIV-1 gp120; left) or Wu1 RBD (right). The neutralizing activity of the depleted antisera was measured against BA.5 pseudovirus. Neutralization data are shown as ID_{50} values for individual mice (colored circles). Data points for individual mice following mock or Wu1 RBD depletions are connected by colored lines. Dashed horizontal lines correspond to the limit of detection for neutralization assays (ID_{50} = 1:50).

Figure S7. Bivalent S-EABR mRNA-LNP booster induces polyclass Ab responses

against Wu1 RBD. Sera for DMS studies were derived from pre-vaccinated mice that

received booster immunizations with 2 µg of monovalent Wu1 S mRNA-LNP, monovalent

Wu1 S-EABR mRNA-LNP, bivalent Wu1/BA.5 S mRNA-LNP, or bivalent Wu1/BA.5 S-

EABR mRNA-LNP (immunization schedule in Fig. 1A-B). Line (left) and logo plots (right)

are shown for DMS results for individual sera from pre-vaccinated mice that received the

indicated boost immunogens. DMS was performed using a Wu1 RBD library. The x axes

of the line and logo plots display the RBD residue numbers. The y axis of the line plots

shows the sum of the Ab escape of all mutations at a site (larger numbers indicating

increased Ab escape). Sites with the strongest Ab escape are shown in logo plots, with

tall letters representing the most frequent mutations at a site. Logo plots are colored

differently for RBD epitopes within different classes[33,40,41] (epitopes are shown in Fig. S2;

gray color for residues not assigned to an epitope).

■ class 1 ■ class 2 ■ class 3 ■ class 4 ■ class 5

Figure S8. Bivalent S-EABR mRNA-LNP booster induces less pronounced class 3

Ab responses against XBB.1.5 RBD. Sera for DMS studies were derived from pre-

vaccinated mice that received booster immunizations with 2 µg of monovalent Wu1 S

mRNA-LNP, monovalent Wu1 S-EABR mRNA-LNP, bivalent Wu1/BA.5 S mRNA-LNP, or

bivalent Wu1/BA.5 S-EABR mRNA-LNP (immunization schedule in Fig. 1A-B). Line (left)

and logo plots (right) are shown for DMS results for individual sera from pre-vaccinated

mice that received the indicated boost immunogens. DMS was performed using an

XBB.1.5 RBD library. The x axes of the line and logo plots display the RBD residue

numbers. The y axis of the line plots shows the sum of the Ab escape of all mutations at

a site (larger numbers indicating increased Ab escape). Sites with the strongest Ab

escape are shown in logo plots, with tall letters representing the most frequent mutations

at a site. Logo plots are colored differently for RBD epitopes within different

classes[33,40] (epitopes are shown in Fig. S2; gray color for residues not assigned to an

epitope).

993
Figure S9. Cartoon diagram illustrating inter- and intra-S crosslinking. Left: model

showing potential inter-S crosslinking through the interactions between both Fabs of a

single IgG and two RBDs of adjacent viral S proteins. Right: model showing intra-S

crosslinking through the interaction between both Fabs of a single IgG and two RBDs on

the same viral S trimer. The hinge regions between the Fabs and Fc are modeled with

flexible linkers, shown as dotted lines, as these regions were not resolved in the crystal

structure of an intact IgG[74]. Disulfide bonds in the hinge region are shown as black lines.

These diagrams are modeled based on the cryo-EM structure of SARS-CoV-2 S

complexed with human C118 Fab (PDB ID 7RKV) and the crystal structure of human Fc

(PDB ID 1FC1). IgGs shown in this figure represent either circulating Abs or membrane-

bound BCRs.

1005
1006

Figure S10. Cryo-EM data collection and processing for heterotrimeric and control SARS-CoV-2 S structures. Representative micrographs, workflow for single-particle data processing, and final reconstructions are shown for SARS-CoV-2 (**A**) Wu1/BA.1 S HT2 heterotrimer, (**B**) Wu1/BA.1/Delta S HT3 heterotrimer, and homotrimer S controls

1011   including (**C**) Wu1 His/StrepII double-tag control, (**D**) StrepII-tagged Wu1, (**E**) His-tagged

1012   BA.1, and (**F**) D7324-tagged Delta constructs.

Boosting Immunogen	Animal	Wu1 DMS Profile	XBB.1.5 DMS Profile
Wu1 S mRNA-LNP	1_4	Moderate class 5 and 2	Strong class 5
Wu1 S mRNA-LNP	1_9		Strong class 5
Wu1 S mRNA-LNP	1_10	Moderate class 2	Strong class 5
Wu1 S mRNA-LNP	1_1	Moderate class 2	Strong class 5
Wu1 S mRNA-LNP	1_6	Polyclass	Strong class 5
Wu1 S mRNA-LNP	1_7	Polyclass	Moderate class 5
Wu1 S-EABR mRNA-LNP	2_3		Strong class 5
Wu1 S-EABR mRNA-LNP	2_5	Moderate class 2	Strong class 5
Wu1 S-EABR mRNA-LNP	2_9	Moderate class 2	Strong class 5
Wu1 S-EABR mRNA-LNP	2_2	Moderate class 2	Moderate class 5 and 4
Wu1 S-EABR mRNA-LNP	2_4	Polyclass	Polyclass
Wu1 S-EABR mRNA-LNP	2_7	Moderate class 2	Strong class 5
Wu1/BA.5 S mRNA-LNP	3_1	Moderate class 2	Strong class 5
Wu1/BA.5 S mRNA-LNP	3_2	Moderate class 1	Strong class 5
Wu1/BA.5 S mRNA-LNP	3_4	Strong class 2	Strong class 5
Wu1/BA.5 S mRNA-LNP	3_5	Polyclass	Strong class 5
Wu1/BA.5 S mRNA-LNP	3_6	Weak class 3	Strong class 5
Wu1/BA.5 S mRNA-LNP	3_9		Polyclass
Wu1/BA.5 S-EABR mRNA-LNP	4_6		Moderate class 5
Wu1/BA.5 S-EABR mRNA-LNP	4_7	Polyclass	Weak class 4
Wu1/BA.5 S-EABR mRNA-LNP	4_9	Polyclass	Moderate class 4
Wu1/BA.5 S-EABR mRNA-LNP	4_2	Polyclass	Weak class 5
Wu1/BA.5 S-EABR mRNA-LNP	4_3	Weak class 2	Polyclass
Wu1/BA.5 S-EABR mRNA-LNP	4_10	Polyclass	Strong class 5

	Escape fraction
Polyclass	<0.5
Weak	0.5-1
Moderate	1-2
Strong	>2

1013

Table S1. Bivalent S-EABR mRNA-LNP booster induces consistent polyclass anti-

RBD Ab responses. The DMS profiles for antisera from individual mice against the Wu1

and XBB.1.5 RBD libraries shown in Fig. S6 and S7 were classified as polyclass (total

escape peaks <1 at all sites; green), moderate escape (total escape peaks between 1

and 2 at one or more sites; yellow), or strong escape (total escape peaks >2 at one or

more sites; red) profiles. Total escape peaks represent the sum of escape fractions for all

mutations at a specific site[45]. For moderate and strong escape profiles, the RBD epitope

classes with the highest total escape peaks are specified.

References

References

1   Corbett, K. S. *et al.* SARS-CoV-2 mRNA vaccine design enabled by prototype
pathogen preparedness. *Nature* **586**, 567-571 (2020).
https://doi.org/10.1038/s41586-020-2622-0

2   Zheng, C. *et al.* Real-world effectiveness of COVID-19 vaccines: a literature
review and meta-analysis. *Int J Infect Dis* **114**, 252-260 (2022).
https://doi.org/10.1016/j.ijid.2021.11.009

3   Chen, Y. *et al.* Broadly neutralizing antibodies to SARS-CoV-2 and other human
coronaviruses. *Nat Rev Immunol* **23**, 189-199 (2023).
https://doi.org/10.1038/s41577-022-00784-3

4   Hogan, M. J. & Pardi, N. mRNA Vaccines in the COVID-19 Pandemic and
Beyond. *Annu Rev Med* **73**, 17-39 (2022). https://doi.org/10.1146/annurev-med-
042420-112725

5   Painter, M. M. *et al.* Rapid induction of antigen-specific CD4(+) T cells is
associated with coordinated humoral and cellular immunity to SARS-CoV-2
mRNA vaccination. *Immunity* **54**, 2133-2142 e2133 (2021).
https://doi.org/10.1016/j.immuni.2021.08.001

6   Oberhardt, V. *et al.* Rapid and stable mobilization of CD8(+) T cells by SARS-
CoV-2 mRNA vaccine. *Nature* **597**, 268-273 (2021).
https://doi.org/10.1038/s41586-021-03841-4

7   Polack, F. P. *et al.* Safety and Efficacy of the BNT162b2 mRNA Covid-19
Vaccine. *N Engl J Med* **383**, 2603-2615 (2020).
https://doi.org/10.1056/NEJMoa2034577

8   Baden, L. R. *et al.* Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine.
N Engl J Med **384**, 403-416 (2021). https://doi.org/10.1056/NEJMoa2035389

9   Zhang, Z. *et al.* Humoral and cellular immune memory to four COVID-19
vaccines. *Cell* **185**, 2434-2451 e2417 (2022).
https://doi.org/10.1016/j.cell.2022.05.022

10   Planas, D. *et al.* Reduced sensitivity of SARS-CoV-2 variant Delta to antibody
neutralization. *Nature* **596**, 276-280 (2021). https://doi.org/10.1038/s41586-021-
03777-9

11   Andrews, N. *et al.* Effectiveness of COVID-19 booster vaccines against COVID-
19-related symptoms, hospitalization and death in England. *Nat Med* **28**, 831-837
(2022). https://doi.org/10.1038/s41591-022-01699-1

12   WHO. Classification of Omicron (B.1.1.529): SARS-CoV-2 variant of concern.,
(2021).

13   Liu, L. *et al.* Striking antibody evasion manifested by the Omicron variant of
SARS-CoV-2. *Nature* **602**, 676-681 (2022). https://doi.org/10.1038/s41586-021-
04388-0

14   Cele, S. *et al.* Omicron extensively but incompletely escapes Pfizer BNT162b2
neutralization. *Nature* **602**, 654-656 (2022). https://doi.org/10.1038/s41586-021-
04387-1

15   Andrews, N. *et al.* Covid-19 Vaccine Effectiveness against the Omicron
(B.1.1.529) Variant. *N Engl J Med* **386**, 1532-1546 (2022).
https://doi.org/10.1056/NEJMoa2119451

16 Tseng, H. F. *et al.* Effectiveness of mRNA-1273 against SARS-CoV-2 Omicron
and Delta variants. *Nat Med* **28**, 1063-1071 (2022).
https://doi.org/10.1038/s41591-022-01753-y
17 Gruell, H. *et al.* mRNA booster immunization elicits potent neutralizing serum
activity against the SARS-CoV-2 Omicron variant. *Nat Med* **28**, 477-480 (2022).
https://doi.org/10.1038/s41591-021-01676-0
18 Wang, Q. *et al.* Alarming antibody evasion properties of rising SARS-CoV-2 BQ
and XBB subvariants. *Cell* **186**, 279-286 e278 (2023).
https://doi.org/10.1016/j.cell.2022.12.018
19 Scheaffer, S. M. *et al.* Bivalent SARS-CoV-2 mRNA vaccines increase breadth of
neutralization and protect against the BA.5 Omicron variant in mice. *Nat Med* **29**,
247-257 (2023). https://doi.org/10.1038/s41591-022-02092-8
20 Khoury, D. S. *et al.* Predicting the efficacy of variant-modified COVID-19 vaccine
boosters. *Nat Med* **29**, 574-578 (2023). https://doi.org/10.1038/s41591-023-
02228-4
21 Chalkias, S. *et al.* A Bivalent Omicron-Containing Booster Vaccine against Covid-
19. *N Engl J Med* **387**, 1279-1291 (2022).
https://doi.org/10.1056/NEJMoa2208343
22 Collier, A. Y. *et al.* Immunogenicity of BA.5 Bivalent mRNA Vaccine Boosters. *N*
Engl J Med **388**, 565-567 (2023). https://doi.org/10.1056/NEJMc2213948
23 Davis-Gardner, M. E. *et al.* Neutralization against BA.2.75.2, BQ.1.1, and XBB
from mRNA Bivalent Booster. *N Engl J Med* **388**, 183-185 (2023).
https://doi.org/10.1056/NEJMc2214293
24 Wang, Q. *et al.* Deep immunological imprinting due to the ancestral spike in the
current bivalent COVID-19 vaccine. *Cell Rep Med* **4**, 101258 (2023).
https://doi.org/10.1016/j.xcrm.2023.101258
25 Wang, Q. *et al.* SARS-CoV-2 neutralising antibodies after bivalent versus
monovalent booster. *Lancet Infect Dis* **23**, 527-528 (2023).
https://doi.org/10.1016/S1473-3099(23)00181-0
26 Alsoussi, W. B. *et al.* SARS-CoV-2 Omicron boosting induces de novo B cell
response in humans. *Nature* **617**, 592-598 (2023).
https://doi.org/10.1038/s41586-023-06025-4
27 Park, Y. J. *et al.* Imprinted antibody responses against SARS-CoV-2 Omicron
sublineages. *Science* **378**, 619-627 (2022).
https://doi.org/10.1126/science.adc9127
28 Carreno, J. M., Singh, G., Simon, V., Krammer, F. & group, P. V. I. s. Bivalent
COVID-19 booster vaccines and the absence of BA.5-specific antibodies. *Lancet*
Microbe **4**, e569 (2023). https://doi.org/10.1016/S2666-5247(23)00118-0
29 Cobey, S. & Hensley, S. E. Immune history and influenza virus susceptibility. *Curr*
Opin Virol **22**, 105-111 (2017). https://doi.org/10.1016/j.coviro.2016.12.004
30 Song, S. *et al.* A systematic review and meta-analysis on the effectiveness of
bivalent mRNA booster vaccines against Omicron variants. *Vaccine* **42**, 3389-
3396 (2024). https://doi.org/10.1016/j.vaccine.2024.04.049
31 Hoffmann, M. A. G. *et al.* ESCRT recruitment to SARS-CoV-2 spike induces
virus-like particles that improve mRNA vaccines. *Cell* (2023).
https://doi.org/https://doi.org/10.1016/j.cell.2023.04.024

32   Hsieh, C. L. *et al.* Structure-based design of prefusion-stabilized SARS-CoV-2
spikes. *Science* **369**, 1501-1505 (2020). https://doi.org/10.1126/science.abd0826
33   Barnes, C. O. *et al.* SARS-CoV-2 neutralizing antibody structures inform
therapeutic strategies. *Nature* **588**, 682-687 (2020).
https://doi.org/10.1038/s41586-020-2852-1
34   Li, C. J. & Chang, S. C. SARS-CoV-2 spike S2-specific neutralizing antibodies.
Emerg Microbes Infect **12**, 2220582 (2023).
https://doi.org/10.1080/22221751.2023.2220582
35   Crawford, K. H. D. *et al.* Protocol and Reagents for Pseudotyping Lentiviral
Particles with SARS-CoV-2 Spike Protein for Neutralization Assays. *Viruses* **12**
(2020). https://doi.org/10.3390/v12050513
36   Robbiani, D. F. *et al.* Convergent antibody responses to SARS-CoV-2 in
convalescent individuals. *Nature* **584**, 437-442 (2020).
https://doi.org/10.1038/s41586-020-2456-9
37   Wang, Z. *et al.* Analysis of memory B cells identifies conserved neutralizing
epitopes on the N-terminal domain of variant SARS-Cov-2 spike proteins.
Immunity **55**, 998-1012 e1018 (2022).
https://doi.org/10.1016/j.immuni.2022.04.003
38   Miller, J. *et al.* Substantial Neutralization Escape by SARS-CoV-2 Omicron
Variants BQ.1.1 and XBB.1. *N Engl J Med* **388**, 662-664 (2023).
https://doi.org/10.1056/NEJMc2214314
39   Feng, S. *et al.* Correlates of protection against symptomatic and asymptomatic
SARS-CoV-2 infection. *Nat Med* **27**, 2032-2040 (2021).
https://doi.org/10.1038/s41591-021-01540-1
40   Fan, C. *et al.* Neutralizing monoclonal antibodies elicited by mosaic RBD
nanoparticles bind conserved sarbecovirus epitopes. *Immunity* **55**, 2419-2435
e2410 (2022). https://doi.org/10.1016/j.immuni.2022.10.019
41   Fan, C. *et al.* Cross-reactive sarbecovirus antibodies induced by mosaic RBD
nanoparticles. *Proc Natl Acad Sci U S A* **122**, e2501637122 (2025).
https://doi.org/10.1073/pnas.2501637122
42   Starr, T. N. *et al.* Deep Mutational Scanning of SARS-CoV-2 Receptor Binding
Domain Reveals Constraints on Folding and ACE2 Binding. *Cell* **182**, 1295-1310
e1220 (2020). https://doi.org/10.1016/j.cell.2020.08.012
43   Taylor, A. L. & Starr, T. N. Deep mutational scans of XBB.1.5 and BQ.1.1 reveal
ongoing epistatic drift during SARS-CoV-2 evolution. *PLoS Pathog* **19**, e1011901
(2023). https://doi.org/10.1371/journal.ppat.1011901
44   Cohen, A. A. *et al.* Mosaic sarbecovirus nanoparticles elicit cross-reactive
responses in pre-vaccinated animals. *Cell* **187**, 5554-5571 e5519 (2024).
https://doi.org/10.1016/j.cell.2024.07.052
45   Greaney, A. J. *et al.* Mapping mutations to the SARS-CoV-2 RBD that escape
binding by different classes of antibodies. *Nat Commun* **12**, 4196 (2021).
https://doi.org/10.1038/s41467-021-24435-8
46   Callaway, H. M. *et al.* Bivalent intra-spike binding provides durability against
emergent Omicron lineages: Results from a global consortium. *Cell Rep* **42**,
112014 (2023). https://doi.org/10.1016/j.celrep.2023.112014

47 Sanders, R. W. *et al.* A next-generation cleaved, soluble HIV-1 Env trimer, BG505
SOSIP.664 gp140, expresses multiple epitopes for broadly neutralizing but not
non-neutralizing antibodies. *PLoS Pathog* **9**, e1003618 (2013).
https://doi.org/10.1371/journal.ppat.1003618
48 Walls, A. C. *et al.* Structure, Function, and Antigenicity of the SARS-CoV-2 Spike
Glycoprotein. *Cell* **181**, 281-292 e286 (2020).
https://doi.org/10.1016/j.cell.2020.02.058
49 Wrapp, D. *et al.* Cryo-EM structure of the 2019-nCoV spike in the prefusion
conformation. *Science* **367**, 1260-1263 (2020).
https://doi.org/10.1126/science.abb2507
50 Tan, C. Y. *et al.* Effectiveness of bivalent mRNA vaccines against medically
attended symptomatic SARS-CoV-2 infection and COVID-19-related hospital
admission among SARS-CoV-2-naive and previously infected individuals: a
retrospective cohort study. *Lancet Infect Dis* **23**, 1343-1348 (2023).
https://doi.org/10.1016/S1473-3099(23)00373-0
51 Bowen, J. E. *et al.* SARS-CoV-2 spike conformation determines plasma
neutralizing activity elicited by a wide panel of human vaccines. *Sci Immunol* **7**,
eadf1421 (2022). https://doi.org/10.1126/sciimmunol.adf1421
52 Balinsky, C. A. *et al.* Antibodies to S2 domain of SARS-CoV-2 spike protein in
Moderna mRNA vaccinated subjects sustain antibody-dependent NK cell-
mediated cell cytotoxicity against Omicron BA.1. *Front Immunol* **14**, 1266829
(2023). https://doi.org/10.3389/fimmu.2023.1266829
53 Keeton, R. *et al.* T cell responses to SARS-CoV-2 spike cross-recognize
Omicron. *Nature* **603**, 488-492 (2022). https://doi.org/10.1038/s41586-022-
04460-3
54 Rydyznski Moderbacher, C. *et al.* Antigen-Specific Adaptive Immunity to SARS-
CoV-2 in Acute COVID-19 and Associations with Age and Disease Severity. *Cell*
183, 996-1012 e1019 (2020). https://doi.org/10.1016/j.cell.2020.09.038
55 Pallesen, J. *et al.* Immunogenicity and structures of a rationally designed
prefusion MERS-CoV spike antigen. *Proc Natl Acad Sci U S A* **114**, E7348-
E7357 (2017). https://doi.org/10.1073/pnas.1707304114
56 McBride, C. E., Li, J. & Machamer, C. E. The cytoplasmic tail of the severe acute
respiratory syndrome coronavirus spike protein contains a novel endoplasmic
reticulum retrieval signal that binds COPI and promotes interaction with
membrane protein. *J Virol* **81**, 2418-2428 (2007).
https://doi.org/10.1128/JVI.02146-06
57 Korber, B. *et al.* Tracking Changes in SARS-CoV-2 Spike: Evidence that D614G
Increases Infectivity of the COVID-19 Virus. *Cell* **182**, 812-827 e819 (2020).
https://doi.org/10.1016/j.cell.2020.06.043
58 Miettinen, H. M., Rose, J. K. & Mellman, I. Fc receptor isoforms exhibit distinct
abilities for coated pit localization as a result of cytoplasmic domain
heterogeneity. *Cell* **58**, 317-327 (1989). https://doi.org/10.1016/0092-
8674(89)90846-5
59 Pinto, D. *et al.* Broad betacoronavirus neutralization by a stem helix-specific
human antibody. *Science* (2021). https://doi.org/10.1126/science.abj3321

60   Kariko, K. *et al.* Incorporation of pseudouridine into mRNA yields superior
nonimmunogenic vector with increased translational capacity and biological
stability. *Mol Ther* **16**, 1833-1840 (2008). https://doi.org/10.1038/mt.2008.200
61   Kariko, K., Muramatsu, H., Ludwig, J. & Weissman, D. Generating the optimal
mRNA for therapy: HPLC purification eliminates immune activation and improves
translation of nucleoside-modified, protein-encoding mRNA. *Nucleic Acids Res*
39, e142 (2011). https://doi.org/10.1093/nar/gkr695
62   Pardi, N. *et al.* Expression kinetics of nucleoside-modified mRNA delivered in
lipid nanoparticles to mice by various routes. *J Control Release* **217**, 345-351
(2015). https://doi.org/10.1016/j.jconrel.2015.08.007
63   Hsieh, C. L. *et al.* Prefusion-stabilized SARS-CoV-2 S2-only antigen provides
protection against SARS-CoV-2 challenge. *Nat Commun* **15**, 1553 (2024).
https://doi.org/10.1038/s41467-024-45404-x
64   Cohen, A. A. *et al.* Mosaic RBD nanoparticles protect against challenge by
diverse sarbecoviruses in animal models. *Science* **377**, eabq0839 (2022).
https://doi.org/10.1126/science.abq0839
65   Mastronarde, D. N. Automated electron microscope tomography using robust
prediction of specimen movements. *J Struct Biol* **152**, 36-51 (2005).
https://doi.org/10.1016/j.jsb.2005.07.007
66   Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC:
algorithms for rapid unsupervised cryo-EM structure determination. *Nat Methods*
14, 290-296 (2017). https://doi.org/10.1038/nmeth.4169
67   Cohen, A. A. *et al.* Mosaic nanoparticles elicit cross-reactive immune responses
to zoonotic coronaviruses in mice. *Science* **371**, 735-741 (2021).
https://doi.org/10.1126/science.abf6840
68   West, A. P., Jr. *et al.* Computational analysis of anti-HIV-1 antibody neutralization
panel data to identify potential functional epitope residues. *Proc Natl Acad Sci U*
S A **110**, 10598-10603 (2013). https://doi.org/10.1073/pnas.1309215110
69   Starr, T. N. *et al.* Shifting mutational constraints in the SARS-CoV-2 receptor-
binding domain during viral evolution. *Science* **377**, 420-424 (2022).
https://doi.org/10.1126/science.abo7896
70   Greaney, A. J. *et al.* Comprehensive mapping of mutations in the SARS-CoV-2
receptor-binding domain that affect recognition by polyclonal human plasma
antibodies. *Cell Host Microbe* **29**, 463-476 e466 (2021).
https://doi.org/10.1016/j.chom.2021.02.003
71   Hills, R. A. *et al.* Proactive vaccination using multiviral Quartet Nanocages to
elicit broad anti-coronavirus responses. *Nat Nanotechnol* **19**, 1216-1223 (2024).
https://doi.org/10.1038/s41565-024-01655-9
72   Barnes, C. O. *et al.* SARS-CoV-2 neutralizing antibody structures inform
therapeutic strategies. *Nature* **588**, 682-687 (2020).
https://doi.org/10.1038/s41586-020-2852-1
73   Landau, M. *et al.* ConSurf 2005: the projection of evolutionary conservation
scores of residues on protein structures. *Nucleic Acids Res* **33**, W299-302
(2005). https://doi.org/10.1093/nar/gki370
74   Saphire, E. O. *et al.* Crystal structure of a neutralizing human IGG against HIV-1:
a template for vaccine design. *Science* **293**, 1155-1159 (2001).

Acknowledgements

We thank J. Vielmetter, A. Lam, and the Caltech Protein Expression Center for assistance with protein production, S. Chen and the Caltech Cryo-EM facility for cryo-EM data collection, M. Anaya (Caltech) for a BirA expression plasmid, J.E. Robinson (Tulane University) for the JR-52 antibody, J. Bloom (Fred Hutchinson Cancer Research Center) and T. Starr (University of Utah) for RBD libraries, and J. Bloom (Fred Hutchinson) and P. Bieniasz (Rockefeller University) for neutralization assay reagents. We thank J. Keeffe for guidance on ELISA method development, A. West for Swift DMS support, Y. Tam (Acuitas Therapeutics) for careful reading of the manuscript, B. Wold, G. Tolomiczenko, and the Caltech Merkin Institute for Translational Research for helpful discussions. Cryo-EM was performed in the Beckman Institute Resource Center for Transmission Electron Microscopy at Caltech. We thank Labcorp Drug Development–Antibody Reagents and Vaccines (Denver, PA) (formerly Covance, Inc.) for mouse immunization studies, and RNAcore (Houston Methodist Research Institute) for synthesis of mRNAs and helpful discussion. Figures 1A-B and 5A were created with Biorender.com. These studies were funded by Wellcome Leap (P.J.B.), the National Institutes of Health P01-AI165075 (P.J.B.) and DP5OD033362 (M.A.G.H), Gates Foundation INV-034638 (P.J.B.) and INV-056219 (M.A.G.H.), the Merkin Institute for Translational Research (Caltech) (P.J.B), and the George Mason University Fast Grants (P.J.B.). This manuscript is the result of funding in whole or in part by the National Institutes of Health (NIH). It is subject to the NIH Public Access Policy. Through acceptance of this federal funding, NIH has been given a right to make this manuscript publicly available in PubMed Central upon the Official Date of Publication, as defined by NIH.

Author contributions:

Conceptualization: C.F., A.A.C., K.A.D., P.J.B., and M.A.G.H.; Methodology: C.F., A.A.C., K.A.D., A.-C.P.F., P.J.B., and M.A.G.H.; Investigation: C.F., A.A.C., K.A.D., A.V.R., A.-C.P.F., Z.Y., P.N.P.G., K.E.H.T., W.J.M., P.J.C.L., and M.A.G.H.; Writing – original draft: C.F., A.A.C., K.A.D., P.J.B., and M.A.G.H.; Writing – review and editing: C.F., A.A.C., K.A.D., A.-C.P.F., P.J.C.L., P.J.B., and M.A.G.H.; Visualization: C.F., A.A.C., K.A.D.; Supervision: P.J.B., and M.A.G.H.; Funding: P.J.B., and M.A.G.H.

Competing interests

M.A.G.H. and P.J.B. are inventors on a US patent application filed by the California Institute of Technology that covers the EABR technology described in this work. P.J.B. is a scientific advisor for Vaccine Company, Inc. M.A.G.H. is a scientific consultant for Vaccine Company, Inc. W.J.M. and P.J.C.L. are employees of Acuitas Therapeutics, a company developing lipid nanoparticle delivery technology; P.J.C.L. holds equity in Acuitas Therapeutics.

Materials, data, and code availability

Materials generated in this study are available upon request through a Materials Transfer Agreement. Code used for data processing and visualization of DMS results is available upon request. Main text and supplementary data will be publicly available. DMS raw sequencing data will be available under an NCBI SRA. Cryo-EM density maps are deposited in the EMDB with accession codes: EMD-71513, EMD-71514, EMD-71515 and EMD-71516. Materials are available upon request to corresponding authors with a signed