Rapid CAR screening and circRNA-driven CAR-NK cells for persistent shed-resistant immunotherapy

Joo-Yoon Chung[1,2, †], Jisu Hong[3, †], Su-Min Yee[1], Hye Won Lim[4], Dahye Lee[5], Sangwon Yoon[6], Seung-Hwan Lee[7,8], Yujin Jung[3], Seung Hun Shin[1,9], Song Cheol Kim[10], Chang-Han Lee[3,4,5,11,12, *], Mihue Jang[1,2,13] *

[1]*Medicinal Materials Research Center, Biomedical Research Division, Korea Institute of Science and Technology (KIST), Seoul 02792, Republic of Korea*

[2]*KHU-KIST Department of Converging Science and Technology, Kyung Hee University, Seoul 02447, Republic of Korea*

[3]*Department of Biomedical Sciences, Seoul National University College of Medicine, Seoul 03080, Republic of Korea*

[4]*Cancer Research Institute, Seoul National University College of Medicine, Seoul 03080, Republic of Korea*

[5]*Department of Pharmacology, Seoul National University College of Medicine, Seoul 03080, Republic of Korea*

[6]*Corporate Research & Development Center, UCI Therapeutics, Seoul 04784, Republic of Korea*

[7]*Department of Biochemistry, Microbiology and Immunology, Faculty of Medicine, University of Ottawa, Ottawa, Ontario K1N 6N5, Canada*

[8]*Ottawa Institute of Systems Biology, Faculty of Medicine and Centre for Infection, Immunity, and Inflammation, Faculty of Medicine, University of Ottawa, Ontario K1N 6N5, Canada*

[9]*Department of Biotechnology, College of Life Sciences and Biotechnology, Korea University, Seoul 02841, Republic of Korea*

[10]*Division of Hepato-Biliary and Pancreatic Surgery, Department of Surgery, University of*

Ulsan College of Medicine, Asan Medical Center 05505, Seoul, Republic of Korea

[11]*BK21 FOUR Biomedical Science Project, Seoul National University College of Medicine,*

Seoul 03080, Republic of Korea

[12]*Wide River Institute of Immunology, Seoul National University, Hongcheon 25159, Republic*

of Korea

[13]*Division of Bio-Medical Science and Technology, KIST School, University of Science and*

Technology (UST), Seoul 02792, Republic of Korea

[†]These authors contributed equally to this work.

KEYWORDS

41 *Corresponding authors:

42 Mihue Jang, Ph.D.

43 Tel: +82-2-958-6618; fax: +82-2-958-5909, e-mail: mihue@kist.re.kr

45 Chang-Han Lee, Ph.D.

46 Tel: +82-2-740-8278; email: chlee-antibody@snu.ac.kr

ABSTRACT

Chimeric antigen receptor (CAR)-based immunotherapies against solid tumors face two major hurdles, the "decoy effect" of shedding antigens that sequester CARs, and the limited persistence of immune effectors within the immunosuppressive tumor microenvironment. Here, we present a mechanistic approach to overcome these barriers by integrating a physiologically relevant screening platform with circular RNA (circRNA) engineering. Unlike conventional screens using immortalized cell lines, we performed rapid functional screening directly in human primary natural killer (NK) cells to identify a novel scFv, CLMS10. Structural modeling revealed that CLMS10 targets a membrane-proximal epitope that overlaps the proteolytic cleavage site, thereby evading inhibition by soluble mesothelin (solMSLN). Furthermore, we demonstrated that circRNA-mediated CAR expression, when codelivered with interleukin-21 (IL-21), confers sufficient stability to withstand the continuous antigen shedding induced by cancer-associated fibroblasts (CAFs), resulting in reduced CAR downregulation. In an *in vivo* metastatic pancreatic cancer model, IL-21-augmented circCAR-MS10-NK cells exhibited potency comparable to that of lentivirally engineered CAR-NK cells while offering superior manufacturability. Overall, this study establishes a paradigm for generating shed-resistant CAR therapeutics through the strategic integration of epitope-specific functional selection and enhanced RNA stability.

Introduction

Natural killer (NK) cells are innate immune effectors that exert rapid and potent cytotoxicity against cancer and virus-infected cells without prior sensitization, making them promising platforms for chimeric antigen receptor (CAR)-based therapies.[1] Unlike T cells, NK cells utilize a diverse array of germline-encoded receptors to recognize transformed cells, providing a layer of redundancy that enables target elimination even when antigen presentation mechanisms are impaired.[1-3] This intrinsic versatility suggests that CAR-NK cells may be less vulnerable to immune escape mechanisms, such as MHC downregulation, that limit conventional T-cell therapies.[1-4] CAR-NK cells integrate the antigen specificity of CAR constructs with the intrinsic cytotoxic mechanisms of NK cells, enhancing overall antitumor efficacy.[2] Furthermore, activated NK cells can remodel the tumor microenvironment (TME) by secreting proinflammatory cytokines like IFN-γ, potentially promoting broader immune engagement in otherwise immunologically cold tumors.[3, 4]

  While CAR-NK cells offer significant advantages over CAR-T cells, including a lower risk of graft-versus-host disease (GVHD) and cytokine release syndrome (CRS), their clinical translation is hindered by manufacturing challenges.[5-8] Although CAR-NK cells are amenable to off-the-shelf allogeneic applications, they remain difficult to genetically modify and exhibit substantial donor-to-donor variability.[2, 9, 10] Traditional viral transduction methods are limited by their low efficiency in NK cells and high cost.[9, 10] More importantly, the conventional viral approach is ill-suited for the iterative, rapid functional optimization required to identify the best CAR candidates among a vast pool of antibodies. Viral workflows are slow and resource-intensive, limiting combinatorial screening of CAR design parameters (scFv epitope, hinge length and signaling domains) in relevant primary cells, and causing costly late-stage attrition when binders fail after CAR reformatting due to synapse-specific constraints.[9, 11, 12]   Consequently, there is a growing demand for a rapid, nonviral

gene delivery platform that enables physiologically relevant functional screening directly in

primary NK cells, ensuring that selected constructs retain potency in a clinical setting. A

platform that can support fast design–build–test cycles in donor-derived NK cells would

enable a more engineering-driven approach to CAR discovery, where functional output

becomes the primary criterion rather than binding metrics alone.[13-15]

Mesothelin (MSLN) is a promising target for cancer immunotherapy because of its high

expression in various malignancies, including mesothelioma, ovarian cancer, pancreatic

adenocarcinoma, gastric adenocarcinoma, and triple-negative breast cancer, while its

expression in normal tissues is limited.[16-18] MSLN has been extensively pursued for various

therapeutic modalities; however, elevated MSLN levels often correlate with aggressive

clinical features and poor prognosis, partly due to its role in immune evasion.[19-21] Despite

their promise, MSLN-targeted strategies are severely compromised by "antigen shedding,"

which releases soluble MSLN (solMSLN) that acts as a circulating sink, sequestering

therapeutic agents and limiting productive engagement of tumor-associated MSLN.[22-25] In the

context of CAR therapies, this decoy phenomenon is particularly detrimental because soluble

antigen binding can reduce effector activation without triggering productive cytotoxicity.[25, 26]

Specifically, first-generation therapeutic agents targeting membrane-distal epitopes (Region I),

such as SS1P, have demonstrated limited efficacy due to susceptibility to the decoy effect.[23,

27] To overcome this limitation, recent strategies have shifted toward membrane-proximal

epitopes (Region III), as exemplified by hYP218 and 15B6,[25, 28] which are retained on the cell

surface after cleavage. Conceptually, targeting the membrane-proximal region should reduce

susceptibility to soluble decoys.[25, 26, 28, 29] Although theoretically superior, targeting this

region is structurally challenging because antibodies must bind close to the cell membrane

without steric hindrance while maintaining high affinity.[30, 31] The steric constraints imposed

by the glycocalyx and the specific geometry required for an optimal immune synapse mean

that many proximal binders may fail to induce signaling.[12, 30-32]Therefore, identifying a

"shed-resistant" binder that functions optimally within the structural constraints of a CAR

synapse requires rigorous validation beyond simple binding assays. The ideal binder must

discriminate membrane-bound antigen from soluble forms while supporting stable synapse

formation.[12, 25]

The performance of CAR-T and CAR-NK cell-based therapies is closely linked to the

affinity, epitope specificity, and membrane proximity of the selected antibody or single-chain

variable fragment (scFv).[11, 30, 33] While high affinity is generally desired, excessive affinity

can impair serial killing or exacerbate the antigen sink effect.[13, 34] Antibodies reformatted into

CARs often lose efficacy or fail to trigger proper immune synapses, necessitating iterative

testing.[12, 35] This "reformatting gap" is particularly acute in NK cells, where optimal signaling

domains differ from those in T cells.[2, 35] However, standard screening methods utilizing

immortalized cell lines (e.g., NK-92) often fail to predict efficacy in donor-derived cells.[12, 35,

36] Immortalized lines often differ in metabolic fitness and activation thresholds, leading to

false-positive predictions for candidates that underperform in primary cells.[2, 12, 35] To

successfully exploit membrane-proximal targeting, an efficient screening method is essential

to rapidly identify binders that are functionally optimized for the primary CAR-NK context

under physiologically relevant conditions, including the presence of soluble antigen.[12, 13, 26, 29,

35]

Beyond antigen engagement, durability remains a critical hurdle. The TME imposes

metabolic and suppressive barriers that limit NK cell persistence.[2, 37-39] Furthermore, stromal

components such as cancer-associated fibroblasts (CAFs) actively remodel the TME and

modulate protease activity, thereby accelerating antigen shedding.[21, 40, 41] A robust therapeutic

strategy must therefore address not only the direct interaction with the tumor antigen but also

the CAF-driven intensification of the decoy effect and the metabolic restrictions of the

TME.[21, 37, 41]

In this study, we address a central bottleneck in MSLN-directed CAR-NK therapy by

identifying binders that remain functional under soluble antigen decoy pressure while

sustaining activity in solid-tumor microenvironments. We establish a rapid, nonviral

workflow for iterative functional screening of membrane-proximal CAR binders directly in

primary NK cells and pair this with circRNA-based CAR expression. Together, this

integrated strategy provides a scalable route to engineer CAR-NK candidates aligned with the

biological constraints of MSLN shedding and the suppressive solid-tumor milieu.

Results

Selective scFv screening against mature MSLN (matMSLN) via the yeast display library

system

To generate high-affinity antibodies against matMSLN (residues 296–606), 6-week-old

BALB/c mice were immunized intraperitoneally with recombinant matMSLN protein **(Figs.**

1 and 2a, and Supplementary Fig. 1). Upon reaching a serum antibody titer of 1:100,000

(Supplementary Fig. 2), variable heavy (VH) and light (VL) chain genes were amplified

from the spleen and bone marrow of sacrificed immunized mice, fused to their respective

constant domains, and transformed into *Saccharomyces cerevisiae* strains JAR200 (MATa)

and YVH10 (MATα) to construct heavy-chain (HC) and light-chain (LC) Fab libraries.

Mating the HC and LC yeast libraries[42, 43] yielded a combinatorial Fab repertoire ($>$

$4\,\times\,10^7$) **(Supplementary Fig. 3a, and Supplementary Table 1).** Flow cytometric

selection of clones with high antigen-binding activity and subsequent cell-based panning of

MSLN$^+$ and MSLN$^-$ AsPC-1 cells narrowed the candidates to 96 unique Fab clones

(Supplementary Figs. 3 and 4). Most antibody candidates exhibited strong matMSLN

binding but negligible affinity for solMSLN (residues 296–592), indicating a favorable

selectivity that may reduce interference from shed MSLN (**Fig. 2a, b**).[25, 44, 45] To identify the

most promising clones, we applied sequential selection criteria: retaining the top 90% of

expressers to maximize recovery, confirming robust Fab assembly on the basis of diagonal

distribution, and isolating the top 5% of matMSLN binders exhibiting preferential specificity

for matMSLN over solMSLN. This systematic approach enabled the identification of

antibody candidates with optimal expression and selectivity (**Fig. 2b, and Supplementary**

Fig. 4).

On the basis of these selection criteria, seven antibody clones—CLMS10, CLMS20,

CLMS32, CLMS36, CLMS76, CLMS88, and CLMS95—were selected for further analysis

(**Fig. 2b**). After the production of recombinant scFvs[46, 47], their binding activities to both

matMSLN and solMSLN were evaluated via ELISA (**Fig. 2c, d**). Except for CLMS32 and

CLMS36, all clones selectively bound to matMSLN (${EC}_{50}$ = 1.7 ± 0.1–151.1 ±☐33.3 nM;

Supplementary Table 2) over solMSLN. Notably, CLMS10 demonstrated high affinity for

matMSLN (${EC}_{50}$ = 18.6 ± 1.1 nM), but minimal binding to solMSLN (${EC}_{50}$ > 1 μM). Surface

plasmon resonance (SPR) confirmed that the affinity of CLMS10 for matMSLN was 8.22 ±

0.40 nM (**Supplementary Fig. 5**). All scFvs, except CLMS20 and CLMS88, specifically

bound to MSLN-positive K562 cells, with ${EC}_{50}$ values ranging from a few nM to several

hundred nM**,** while negligible binding to MSLN-negative K562 cells was detected (**Fig. 2e, f,**

and Supplementary Table 3). These results indicate that our anti-MSLN antibodies

selectively recognize membrane-bound MSLN on the cell surface.

To further define the epitopes of the antibody candidates, we performed competitive

ELISA using reference antibodies with well-characterized binding sites: SS1, which targets

domain I (residues E296–L398)[28], and 15B6, which binds the C-terminal region (residues

Y594–L605).[25] CLMS10, CLMS32, CLMS36, CLMS76, and CLMS95 competed for antigen

binding, suggesting that they either share overlapping epitopes or exhibit steric hindrance that

prevents simultaneous binding **(Fig. 2g)**. Notably, CLMS10 and CLMS76 exhibited weak

binding to solMSLN, implying that their epitopes are localized near the C-terminus of

domain III, which is known to be susceptible to proteolytic shedding **(Fig. 2d, h)**.

Collectively, these findings suggest that the epitopes of CLMS10, CLMS32, CLMS36,

CLMS76, and CLMS95 are clustered near the C-terminal region of domain III, supporting

their therapeutic potential as shed-resistant antibodies **(Fig. 2h)**.

Rapid and efficient live-cell screening of functional scFvs via a linRNA-based CAR-NK

(linCAR-NK) cell platform

Standard screening methods often rely on immortalized cell lines (e.g., NK-92), which fail to

recapitulate the complex biology of donor-derived cells. To ensure translational relevance, we

implemented a "direct-to-primary" screening strategy. This streamlined approach enables

rapid live functional screening directly in clinically relevant human primary NK cells,

yielding high-fidelity phenotypic data and mitigating the risk of candidate failure during

translational scale-up **(Figs. 1 and 3a)**. Each scFv-containing CAR construct was synthesized

as a linRNA via *in vitro* transcription (IVT). The quality of the synthesized linRNAs was

validated via agarose gel electrophoresis and purity assessment **(Supplementary Fig. 6a)**.

Human primary NK (pNK) cells were electroporated with each linRNA, and CAR expression

and cell viability were assessed in a time- and dose-dependent manner. For example, CAR-

SS1-NK cells presented over 87% CAR expression at 1 µg linRNA per 10^6 cells, with

detectable expression persisting for up to 5 days, albeit at reduced levels **(Supplementary**

Fig. 6b, c). Following linRNA electroporation, flow cytometry confirmed CAR and GFP

expression across all clones, with GFP coexpressed via a P2A linker. CAR expression levels

varied among constructs and were quantified by staining with PE- or APC-conjugated

streptavidin in combination with biotinylated MSLN extracellular domain proteins

(**Supplementary Figs. 6d and 7**). The cell viability was largely consistent across the clones

(**Supplementary Fig. 6e**).

We established a rapid live-cell screening platform to assess the cytotoxic function of

linCAR-NK cells against an MSLN-overexpressing pancreatic cancer cell line (Capan-2).

Real-time cancer cell death by linCAR-NK cells was monitored by tracking Caspase-3/7

activation during live coculture, with CAR-SS1-NK and CAR-15B6-NK cells used as

controls (**Fig. 3b, c, and Supplementary Fig. 8a**). Notably, CAR-MS10-NK cells caused a

significant increase in cancer cell death, as evidenced by a 1.5-fold increase in Caspase-3/7-

positive Capan-2 cells compared with mock-NK cells. A luciferase-based cytotoxicity assay

revealed that CAR-MS10-NK cells achieved the strongest killing efficacy against both

Capan-2-Luc and SK-OV-3-Luc cells, which express different MSLN levels (**Supplementary**

Fig. 8b, c). We assessed the sensitivity of each CAR-NK construct to solMSLN-mediated

inhibition. In coculture with solMSLN, CAR-SS1-NK, CAR-MS32-NK, and CAR-MS36-

NK cells exhibited a dose-dependent reduction in killing activity. In contrast, CAR-MS10-

NK, CAR-15B6-NK, and CAR-MS20-NK cells maintained their cytotoxicity, indicating

resistance to solMSLN-mediated inhibition (**Fig. 3d**). The expression of CD107a, a marker of

degranulation was unaffected by solMSLN treatment in CAR-MS10-NK, CAR-MS20-NK,

and CAR-15B6-NK cells activated with matMSLN, further supporting their shedding

resistance profiles (**Fig. 3e**). We examined the extent of trogocytosis associated with impaired

CAR function and NK-mediated fratricide in linCAR-NK cells.[48, 49] Compared with CAR-

SS1-NK cells, CAR-MS10-NK cells exhibited significantly reduced trogocytosis

(**Supplementary Fig. 8d**). Furthermore, CAR-MS10-NK cells presented the greatest degree

of cell avidity among all the tested CAR constructs, suggesting the formation of robust

immunological synapses (**Fig. 3f**).[50] To validate these findings in a more clinically relevant

system, we introduced the scFvs into lentivirus-based CAR constructs, which were

transduced into pNK cells. The results of functional assays revealed similar cytotoxicity

patterns as those observed with linCAR-NK cells, with CAR-MS10-NK cells showing an

approximately 22% higher killing rate against Capan-2 cells than CAR-SS1- and CAR-15B6-

transduced cells **(Fig. 3g and Supplementary Fig. 9a)**. We evaluated whether this functional

trend persisted across immune cell types such as linCAR-T cells. In cytotoxicity assays

against Capan-2-Luc cells, linCAR-15B6-T cells showed comparable efficacy to linCAR-

MS10-T and linCAR-SS1-T cells **(Supplementary Fig. 9b–f).** CAR-MS20-T cells showed

limited activity, which is consistent with the NK cell results, suggesting that the functional

impact of scFvs may vary depending on the immune effector cell type.[13, 14, 32, 51]

Finally, *in silico* docking analysis revealed a critical structure–function relationship.

Unlike SS1, which binds to distal Region I, CLMS10 interacts with a conformational epitope

that overlaps with or lies immediately adjacent to the proteolytic cleavage site (**Fig. 3h**). This

unique binding geometry likely imposes steric hindrance that prevents access by sheddases or

stabilizes the membrane-bound conformation of MSLN.

Consequently, CLMS10 selectively engages membrane-anchored MSLN while

exhibiting drastically reduced affinity for the shed soluble form, providing a structural basis

for its superior performance in shedding inhibition assays. Additionally, our linRNA-based

screening platform offers an efficient functional approach for discovering and optimizing

clinically relevant MSLN-targeting CARs.

linRNA-driven screening for optimized CAR architecture and cytokine coexpression

We selected CAR-MS10 for further analysis. We employed a linRNA-based system to screen

various combinations of transmembrane and costimulatory domains. Specifically, CD8 and

NKG2D were chosen as the transmembrane domains (TMD_{CD8} and TMD_{NKG2D}), and four

CAR variants were designed: TMD_{CD8}–4-1BB–CD3ζ (4-1BBζ), TMD_{CD8}–OX40–CD3ζ

(OX40ζ), TMD$_{NKG2D}$–DAP10–DAP12 (DAP10/12), and TMD$_{NKG2D}$–2B4–CD3ζ (2B4ζ) **(Fig.**

4a).

The cytotoxic potential of the CAR variants against Capan-2-Luc cells was evaluated via

a luciferase-based killing assay. Among the variants, the CAR-MS10-OX40ζ construct

exhibited the strongest cytotoxicity, followed by CAR-MS10-4-1BBζ **(Fig. 4b and**

Supplementary Fig. 10a). A detailed analysis of the functional activities across the 4-1BBζ

and OX40ζ constructs revealed that, consistent with the cytotoxicity data, CAR-MS10-

OX40ζ resulted in a significant increase in CD107a activation compared with the prototype

CAR-MS10-4-1BBζ, accompanied by increased secretion of IFN-γ **(Fig. 4c, d)**. In addition to

direct cytotoxicity, metabolic fitness is a key determinant of CAR-NK cell survival in the

nutrient-deprived TME.[37] The incorporation of the OX40 costimulatory domain significantly

reprogrammed NK cell metabolism, enhancing both basal and maximal mitochondrial

respiration upon antigen stimulation **(Fig. 4e, f)**. This elevated spare respiratory capacity

suggests that CAR-MS10-OX40ζ-NK cells are better equipped to sustain energy production

under metabolic stress, contributing to prolonged antitumor activity.

To further investigate the underlying signaling pathways involved in OX40ζ-mediated

CAR activation, CAR-MS10-OX40ζ-NK cells were stimulated with an MSLN recombinant

protein and analyzed via western blotting **(Fig. 4g).** Compared with those in the 4-1BBζ

group, the phosphorylation of phospholipase C gamma and extracellular signal-regulated

kinase, key downstream molecules of CD3ζ signaling, was greater in the OX40ζ group.[52,53]

Collectively, these results identify OX40ζ as the optimal intracellular domain configuration

for CAR-MS10, enhancing both effector function and signaling capacity.

We cotransfected cytokine-encoding linRNAs into CAR-MS10-OX40ζ cells to increase

their cytotoxicity and functionality. Cytokines play pivotal roles in modulating NK cell

activity, proliferation, and persistence.[54-56] On the basis of their well-characterized functions,

we selected interleukin-2 (IL-2), IL-15, and IL-21 to reinforce NK cell effector function and

longevity. The synergistic effects of the selected cytokines were evaluated in CAR-MS10-NK

cells, generated by coelectroporating CAR-MS10-OX40ζ linRNAs with linRNAs encoding

each cytokine **(Fig. 4h and Supplementary Fig. 10b, c).** Among the various cytokine

combinations, IL-21–CAR-MS10-OX40ζ-coexpressing NK cells presented the strongest

antitumor activity against Capan-2-Luc cells, along with significantly increased CD107a

expression **(Fig. 4i, j)**. Taken together, the results of the linRNA-based approach

demonstrated that the IL-21–CAR-MS10-OX40ζ configuration endows NK cells with

superior cytotoxicity and functional activity.

CircRNA-based CAR-NK cells exhibited superior cytotoxicity with prolonged CAR

expression

To address the transient nature of linRNA expression in mRNA-based cell therapies, we

explored the use of circRNAs for prolonging CAR expression. Compared with linRNAs,

circRNAs have previously been used to prolong transgene expression and offer improved

structural stability and translational efficiency.[57-59] We monitored CAR expression and cell

viability over time in CAR-NK cells generated from either linRNAs or circRNAs **(Fig. 5a)**.

Notably, circCAR-NK cells presented significantly prolonged and sustained CAR expression,

as demonstrated by longitudinal flow-cytometry analyses of both the percentage of CAR□

cells and the CAR mean fluorescence intensity (MFI) across multiple timepoints (Days 1, 3,

and 5) **(Fig. 5b, c)**. In contrast, there was no notable difference in viability between linRNA-

and circRNA-transfected cells over the same time period **(Fig. 5d)**. We evaluated the

persistence of cytotoxicity using serial killing assays, in which CAR-NK cells were

rechallenged with fresh cancer cells for five consecutive rounds **(Fig. 5e)**. Consistent with

prolonged CAR expression, circCAR-NK cells maintained superior activity, although a

gradual decline was observed over time **(Fig. 5f)**. circCAR-NK cells exhibited robust IFN-γ

secretion, supporting a direct association between prolonged CAR expression and sustained

effector function **(Fig. 5g)**. circCAR-MS10-NK cells retained strong cytotoxic activity even

under solMSLN-shedding conditions, indicating resistance to solMSLN-decoy effects **(Fig.**

5h). In contrast, circCAR-SS1-NK cells showed limited cytotoxicity and a marked reduction

in IFN-γ secretion in the presence of solMSLN, highlighting the importance of target epitope

selection and construct design **(Fig. 5i)**. Importantly, circRNA-driven CAR-NK cells

consistently demonstrated stronger cytotoxicity and NK-cell activation than linCAR- or

$_{LV}$CAR-NK cells did, further supporting the superior functional durability of the circRNA

platform **(Supplementary Fig. 11).** Together, these findings underscore the superior

performance of circCAR-NK cells over their linRNA counterparts, demonstrating enhanced

CAR expression, prolonged cytotoxicity, and greater functional resilience under repeated

tumor challenge and antigen-shedding stress.

CircCAR-MS10-NK cells show potent antitumor activity in solMSLN-shedding models

in vitro **and** *in vivo*

The pancreatic TME is highly immunosuppressive and is characterized by a dense

extracellular matrix and abundant TGF-β signaling, which primarily originates from CAFs.

This milieu impairs NK cell function and greatly challenges the efficacy of

immunotherapies.[38-40] Recent studies have demonstrated that anti-mesothelin antibodies not

only target tumor cells but also disrupt antigen-presenting CAFs, reshaping the TME and

enhancing immune responses.[60] However, the therapeutic efficacy of MSLN-targeted CAR-T-

therapies is undermined by the decoy effect of solMSLN-mediated antigen shedding,

impeding CAR-Tcell activity.[25] To address the suppression of CAR-Tcell function induced by

solMSLN under physiologically relevant conditions, we established an *in vitro* cancer–CAF

coculture model to mimic the TME. Interestingly, we found that the presence of human

primary pulmonary fibroblasts or CAFs derived from patients with pancreatic cancer

significantly accelerated the secretion of solMSLN compared with that of cancer cells alone

(Fig. 6a, b, and Supplementary Fig. 12a, b).

Based on the observed increase in solMSLN secretion in the presence of CAFs, we

further assessed the therapeutic efficacy of CAR-NK cells under CAF-mediated shedding

inhibition via real-time live-cell imaging **(Fig. 6c, and Supplementary Fig. 12c)**. The CAF-

induced increase in solMSLN levels significantly reduced cancer cell death in the circCAR-

SS1-NK cell-treated group. In contrast, the cytotoxic efficacy of circCAR-MS10-NK cells

was maintained despite MSLN shedding. Notably, strong MSLN expression was associated

with advanced cancer progression, as evidenced by CLMS10-IgG staining in pancreatic

tumor tissues, suggesting that MSLN is a therapeutically actionable target **(Fig. 6d)**. We

established an *in vivo* intraperitoneal metastatic tumor model that mimics the TME[5, 25],

including CAF-mediated MSLN shedding, by coinjecting AsPC-1-MSLN-Luc tumor cells

and CAFs into NOD-*Prkdc*[em1Baek]*Il2rg*[em1Baek] (NSG) mice and treating them with various

CAR-NK platforms: linCAR-NK cells expressing CAR-SS1, CAR-15B6, or CAR-MS10;

circCAR-MS10-NK cells; and $_{LV}$CAR-MS10-NK cells **(Fig. 6e)**. To enhance NK cell

function and persistence, IL-21 linRNA was coelectroporated into all the mRNA-based CAR-

NK groups, but not into the lentivirus-based groups. Compared with the mock group, both the

circRNA-based and lentivirus-based CAR-MS10-NK groups demonstrated significant tumor

regression **(Fig. 6f-i)**. circCAR-MS10-NK cells showed anticancer activity comparable to

that of $_{LV}$CAR-MS10-NK cells. In contrast, compared with mock cells, linCAR-NK cells

induced a modest reduction in tumor size, although the difference was not significant. These

findings demonstrate that optimized circCAR-MS10-NK cells retain antitumor efficacy under

TME-mimicking conditions and elevated solMSLN levels, supporting the potential

applicability of the nonviral CAR-NK platform for treating solid tumors.

Given that IL-21 was codelivered to increase NK activity, we also assessed potential

systemic IL-21-associated toxicity. Following three repeated infusions of CAR-NK cells, we

observed no significant changes in body weight, gross liver and kidney morphology, or

hepatic and renal function markers, indicating the absence of acute organ toxicity

(**Supplementary Fig. 13**). These findings suggest that cytokine-augmented circCAR-NK

cells do not elicit measurable systemic toxicity, further supporting their translational potential

as safe and effective nonviral CAR-NK therapies for solid tumors. To further characterize the

behavior of the different CAR-NK modalities *in vivo*, we next examined the distribution and

persistence of the infused NK cells **(Supplementary Fig. 14)**. Quantification of CAR□ NK

cells in the peripheral blood at 72 h confirmed successful engraftment in all the CAR-bearing

groups, whereas the mock-NK group presented only background-level events. By day 7,

distinct persistence patterns had emerged among the platforms. As expected from the

transient nature of linear mRNAs, linCAR-NK cells were largely undetectable in the

circulation and secondary lymphoid tissues, highlighting the short-lived *in vivo* expression

driven by linRNAs. In contrast, IL-21–armored circCAR-NK cells demonstrated the most

robust persistence among all groups, with clear detection in the spleen 7 days after

intravenous injection, which aligns with the known survival- and proliferation-supporting

functions of IL-21. Conversely, even $_{LV}$CAR-NK cells showed limited persistence on day 7.

To complement these *in vivo* persistence data and to directly assess the stability of CAR

surface expression under repeated antigen exposure, we next evaluated CAR stability *in vitro*

by repeatedly challenging CAR-NK cells with Capan-2-Luc tumor cells for up to four

consecutive rounds (**Supplementary Fig. 15**). Across all platforms, serial stimulation

ultimately resulted in progressive CAR downregulation. Notably, the surface CAR MFI of

linCAR-, $_{LV}$CAR-, and nonarmored circCAR-NK cells rapidly decreased. However, IL-21–

incorporated circCAR-NK cells presented a modest transient increase in CAR expression

during the first stimulation round, followed by a gradual reduction over subsequent rounds.

Overall, compared with nonarmored circCAR controls, IL-21 armoring substantially

improved the maintenance of CAR surface expression across stimulation rounds within the

CAR☐ NK-cell population **(Supplementary Fig. 15)**. These findings indicate that transient

IL-21 delivery helps to preserve CAR surface expression during repeated antigen

encounters—an essential feature for sustaining NK-cell activity in solid-tumor contexts.

Discussion

In this study, we identified and addressed a critical failure mode of solid tumor

immunotherapies, specifically the suppression of CAR-NK activity by CAF-mediated antigen

shedding.[61] A major finding is that successful targeting of shed-prone antigens such as MSLN

requires a dual mechanism involving (1) structural evasion of the decoy effect via membrane-

proximal binders (CLMS10) and (2) sustained effector function via IL-21–armored,

circRNA-driven expression. Our approach integrated rational antibody engineering, rapid and

efficient mRNA-based functional screening, and circRNA technology to overcome the

limitations of current solid tumor therapies. Our data revealed that in the presence of CAFs,

which actively accelerate shedding, standard CARs (SS1) and transient expression systems

fail. In contrast, our optimized circCAR-MS10-NK cells maintained robust cytotoxicity,

providing a unique contribution to the field by defining the design principles required for

next-generation solid tumor therapies.

A major obstacle in targeting MSLN is the shedding of the extracellular domain, which

acts as a decoy receptor, sequestering CARs and diminishing cytotoxic potential.[26, 62] To

address this issue, we generated a panel of MSLN-specific antibodies and demonstrated the

performance of CLMS10 for binding membrane-proximal epitopes while displaying minimal

affinity for solMSLN (**Fig. 2**). Unlike standard screens, our two-step yeast surface display

workflow ensures the identification of binders that retain structural integrity in the CAR

format. A two-step Fab-to-scFv workflow further minimizes candidate dropout by first

recovering correctly paired, stable VH/VL candidates in Fab format and then validating their

compatibility after scFv conversion. This staged process ensured that only robust shed-

resistant binders advanced into CAR screening. In contrast to SS1, the CLMS10-based CAR

maintained strong cytotoxic activity even at high solMSLN concentrations, emphasizing the

importance of epitope selection (**Figs. 2c–h and 3d–g**). Notably, these shed-resistant CARs

demonstrated efficacy across both NK and T cells, indicating broad applicability across

effector platforms. Mechanistically, the membrane-proximal binding of CLMS10 not only

evades the "sink effect" but also significantly reduces trogocytosis compared with that of the

distal-binding SS1 CAR (**Supplementary Fig. 8d**). Trogocytosis is known to drive CAR

internalization, fratricide, and effector exhaustion.[34, 49] The lower trogocytic activity of

CLMS10 is likely attributable to its membrane-proximal, shed-resistant binding mode, which

minimizes soluble antigen–mediated CAR clustering and contributes to more controlled

synapse engagement. This restrained interaction may underlie the enhanced functional

durability observed with CLMS10-based CAR-NK cells.

A key advantage of our workflow is the transition from cell-line-based screening to rapid

mRNA-based functional assessment directly in primary immune cells. Conventional screens

using immortalized lines (e.g., NK-92) often fail to predict translational efficacy due to

divergent signaling and metabolic profiles. By employing electroporation-based delivery in

primary NK cells, we rapidly phenotyped multiple CAR constructs, identifying the OX40ζ

signaling domain and IL-21 codelivery as the optimal configuration. Notably, this

combination enhanced cytotoxicity and significantly improved metabolic fitness, as

evidenced by increased mitochondrial respiration (**Fig. 4**). As IL-21 is known to increase

PI3K–Akt activation[55, 63] and trigger the expression of T-BET, EOMES, and CEBPD[55, 64], our

findings suggest that the synergy between OX40 signaling and IL-21-mediated metabolic

reprogramming enables NK cells to persist and function within the nutrient-deprived TME.

Another major advance is the application of circRNAs for CAR expression in NK cells.

Compared with linRNAs, circRNAs exhibit prolonged transcript stability and sustained CAR

expression, owing to their resistance to exonuclease degradation.[65, 66] circCAR-NK cells

remained functionally active during serial tumor challenges, with cytotoxicity maintained

despite repeated antigen exposure **(Fig. 5)**. This highlights the unique advantage of circRNAs

in bridging transient and durable CAR expression systems, combining the safety of RNA-

based approaches with the persistence of integrated vectors. Despite its advantages, circRNA-

based CAR expression remains transient, indicating that further RNA engineering strategies

could enhance durability. The incorporation of stabilizing elements or translation-enhancing

motifs around the IRES may prolong protein production from circular transcripts.[67, 68]

Additional approaches that increase RNA stability or translational persistence could likewise

be applied to future CAR designs. Our data also show that IL-21 codelivery improves

functional persistence, suggesting that cytokine armoring can complement RNA engineering

by supporting NK-cell survival and activity. Together, these strategies represent promising

avenues to strengthen the durability and therapeutic performance of nonviral CAR-NK

platforms.

In terms of translational relevance, we tested our platform in a CAF-rich tumor model

characterized by dense stroma, high solMSLN levels, and immunosuppressive factors. While

CAFs compromised the cytotoxicity of circCAR-SS1-NK cells, circCAR-MS10-NK cells

retained potent cytotoxicity (**Fig. 6a–c**). Human pancreatic tumor tissues presented high

MSLN expression (**Fig. 6d**). In AsPC-1-MSLN-Luc/CAF xenografts, circCAR-MS10-NK

cells achieved tumor regression, exhibiting activity comparable to that of lentiviral CAR-NK

cells (**Fig. 6e–i**). These findings underscore the platform's shed-resistance and persistence

under clinically relevant TME conditions.

Our RNA engineering strategy supports the development of off-the-shelf, allogeneic

CAR-NK therapies. Compared with T cells, NK cells are inherently less prone to GVHD and

CRS, providing a potential universal donor base.[3, 4, 51] Previous trials using HLA-mismatched

cord blood-derived CAR-NK cells expressing IL-15 have demonstrated safe and durable

responses in patients.[69] By avoiding viral vectors, our mRNA and circRNA systems reduce

manufacturing complexity, shorten production timelines, and enable rapid adaptation to new

targets.[15] The transient expression profile of RNA further permits repeat dosing and modular

CAR redesign, offering unmatched flexibility. Furthermore, recent advances in circRNA

synthesis, particularly self-splicing intron-based circularization, have improved yield and

scalability, facilitating broader application of this approach in cell therapy

manufacturing.[68,70,71] While our shed-resistant circCAR-MS10-NK cells demonstrate strong

potential in addressing antigen shedding and mitigating elements of the immunosuppressive

TME, solid tumors such as pancreatic cancer remain highly refractory due to additional

resistance mechanisms. These include antigen downregulation and the secretion of

immunosuppressive factors such as TGF-β within the dense stromal matrix.[72-74] Consequently,

further studies are warranted to assess the long-term persistence, tumor trafficking, and safety

of circRNA-engineered CAR-NK cells *in vivo*. Future therapeutic strategies may benefit from

combining these cells with agents that target complementary immune escape pathways, such

as immune checkpoint inhibitors or TGF-β blockade, to increase antitumor efficacy within

the complex TME.

In conclusion, we propose a comprehensive framework for generating shed-resistant,

long-lasting CAR-NK therapies with enhanced cell–cell avidity against solid tumors. By

strategically integrating antibody engineering, rapid mRNA-based CAR discovery, and

circRNA-mediated expression combined with IL-21, we address key limitations of CAR-

based therapies for solid tumors. The IL-21–armored circCAR-MS10-NK system offers a

potent, scalable, and clinically translatable solution for treating malignancies such as

pancreatic cancer, effectively overcoming antigen shedding and immunosuppressive TMEs

that have previously impeded therapeutic success.

Materials and methods

Isolation and *ex vivo* expansion of human pNK cells.

PBMCs were purchased from Lonza (Walkersville, MD, USA). NK cells were cultured in

medium from Miltenyi Biotec (Bergisch Gladbach, Germany). pNK cells were cultured and

maintained in the presence of IL-2 (200 IU/mL; PeproTech, Rocky Hill, NJ, USA) and IL-15

(10 ng/mL; PeproTech), along with K562 feeder cells obtained from the American Type

Culture Collection (Manassas, VA, USA). Following a 2-week expansion period, pNK cells

were isolated via an NK cell isolation kit (Miltenyi Biotec) according to the manufacturer's

guidelines.

Animals

NSG mice and BALB/c nude mice were purchased from JABIO (Gyeonggi-do, Korea).

BALB/c mice were purchased from KOATECH (Gyeonggi-do, Korea). Male mice aged 6–8

weeks were randomly assigned to experimental groups for the study.

Yeast strains and culture conditions

S. cerevisiae JAR200 (MATa, GAL1-AGA1::KanM×4ura3D45, ura3–52 trp1 leu2D1

his3D200 pep4::HIS3 prb1D1.6R can1) and YVH10 (MATα, PDI::ADHII-PDI-Leu2 ura3-52

trp1 leu2D1 his3D200 pep4::HIS3 prb1D1.6p can1 GAL) strains were used to construct and

screen the yeast surface-display Fab library. Both strains were kindly provided by Prof. Dane

Wittrup (Massachusetts Institute of Technology, Cambridge, MA, USA). The haploid yeast

library was maintained in SDCAA (6.7 g/L yeast nitrogen base, 5 g/L casamino acids, 14.7

g/L sodium citrate, 4.29 g/L citric acid monohydrate, and 20 g/L dextrose in deionized water)

supplemented with either uracil (0.0002 g/L) for JAR200 or tryptophan (0.0004 g/L) for

YVH10, while the diploid Fab library was maintained in SDCAA without additional

supplements. The Fab surface display library was induced in 2× SGCAA medium (13.4 g/L

yeast nitrogen base, 10 g/L casamino acids, 5.4 g/L Na_2HPO_4, 8.56 g/L $NaH_2PO_4·H_2O$, and

20 g/L galactose in deionized water) at a starting OD_{600} of 0.5, with incubation at 20 °C and

160 rpm for 48 h.

Construction of the immune Fab library

The immune Fab library was constructed via a yeast surface display system to facilitate

proper eukaryotic protein folding, overcoming the structural limitations often associated with

prokaryotic phage display. A Fab-based format was employed to maximize combinatorial

library diversity by mating, thereby achieving a library size and screening throughput

significantly superior to those typically attainable in mammalian display systems. Following

the final immunization, the mice were sacrificed, and mRNA was isolated from the spleen

and bone marrow samples. The VH and VL genes were amplified from cDNA via a primer

mixture **(Supplementary Table 1)**. Through overlapping PCR, each amplified VH and VL

gene was fused to the CH1 and CL regions, respectively. The resulting VH-CH1 fragment (4

µg) was cotransformed with the linearized pYDS-H vector into *S. cerevisiae* strain JAR200,

while the VL-CL fragment was cotransformed with the linearized pYDS-K vector into *S.*

cerevisiae strain YVH10. Homologous recombination facilitated by electroporation was used

for both transformations. The resulting haploid HC library in JAR200 (MATa) and the LC

library in YVH10 (MATα) were cultured for 15 h at 30 °C and 160 rpm in 200 mL of

SDCAA medium supplemented with uracil (for JAR200) or tryptophan (for YVH10),

reaching an OD□□□ of ~5 in the late exponential phase. For mating, equal volumes

(OD□□□ = 100) of the HC and LC library cells were mixed thoroughly by vortexing and

inversion. The mixture was plated onto YPD agar (pH 4.5) and incubated at 30 °C for 8 h.

After incubation, the cells were washed twice with sterile distilled water, resuspended in

SDCAA medium at an initial OD□□□ of 0.1, and then grown overnight at 30 °C with

shaking at 160 rpm. To evaluate mating efficiency, serial dilutions of the mated cultures were

plated on agar with either SDCAA or SDCAA supplemented with tryptophan. Mating

efficiency was calculated as the ratio of colony counts on SDCAA to those on SDCAA with

tryptophan.

***In vitro* transcription (IVT) for linCAR**

To construct the CAR plasmid, the newly generated scFv containing the second-generation

CAR construct and GFP was cloned and inserted into the pmREX plasmid vector via the EZ-

Fusion HT Cloning Kit (Cat. EZ015TL, Enzynomics, Daejeon, Republic of Korea) along

with the SS1 (US Patent: US7081518) and 15B6 (US Patent: US7081518B1) scFv DNA

fragments. The sequences of these scFvs were subsequently cloned and inserted into template

DNA for mRNA synthesis. IVT of the mRNA was performed via the use of a linearized

plasmid vector as the template and a MEGAscript T7 transcription kit (Cat. AMB13345;

Invitrogen, Waltham, MA, USA). The synthesized mRNA was capped with CleanCap

Reagent AG (Cat. N-7113, Trilink, San Diego, CA, USA). mRNA was precipitated via

lithium chloride and purified, and the yield was measured via a NanoDrop 1000 UV visible

spectrophotometer (Thermo Fisher Scientific, Waltham, MA, USA).

Generation of CAR-NK cells via electroporation.

The circRNA construct, designed as a second-generation CAR incorporating either the SS1 or

CLMS10 scFv, was provided by GenScript (Nanjing, China). Expanded pNK cells were

collected by centrifugation at 300 ×g for 5 min, and the culture medium was removed. The

cells were washed twice with 10 mL of DPBS and resuspended in Resuspension Buffer T

from a Neon Transfection System 100 μL Kit (4.0 × 10□ cells/mL; Invitrogen). CAR-NK

cells were generated via IVT using 1 μg of mRNA per 10□ cells. The mRNA was mixed

thoroughly with 100 μL of cells suspended in Resuspension Buffer T. Electroporation was

performed via a Neon Transfection System (Invitrogen) to facilitate intracellular mRNA

delivery. For circCAR-NK generation, 0.5 μg of circRNA per 10□ cells was used for

electroporation. In the case of multiple transfections, the concentration of each mRNA was

adjusted on the basis of its corresponding molecular weight (bp size) to achieve the

appropriate molarity, and the final amount per 10□ cells was used for electroporation.

In vivo **efficacy of CAR-NK cells**

All animal procedures were conducted in accordance with institutional guidelines and

approved by the Institutional Animal Care and Use Committees of KIST (KIST–2024–079).

A metastatic tumor model reflecting the TME with high solMSLN secretion was established

via intraperitoneal injection of either 3 × 10□ AsPC-1-MSLN-Luc cells alone or a mixture of

AsPC-1-MSLN-Luc and CAF6 cells at a 3:1 ratio. After 11 days, the mice were randomly

divided into seven groups and treated under various conditions. Each tumor-bearing mouse

received three intravenous administrations of 1 × 10□ linCAR-NK, circCAR-NK, or $_{LV}$CAR-

NK cells. Tumor progression was monitored via *in vivo* luciferase imaging on an IVIS

Spectrum imaging system (Revvity, Waltham, MA, USA) following intravenous injection of

a luciferin substrate solution (Promega, Madison, WI, USA) under isoflurane anesthesia. The

tumor burden was quantified by measuring the average radiance (photons/s/cm²/sr).

Further methodological details are provided in the Supplementary Materials.

Statistical analysis

All the data are expressed as the means $\pm$ s.d. Statistical analyses were performed via

GraphPad Prism 10. Two-tailed Student's *t*-test or two-way ANOVA was performed,

depending on the experimental design, with Tukey's post hoc analysis. $p < 0.05$ was used to

identify statistically significant differences.

Ethics statement

All animal experiments were performed in accordance with guidelines approved by the

Institutional Animal Care and Use Committee of Seoul National University (protocol no.

SNU-221108-1-1) and KIST (protocol no. KIST–2024–079).

Acknowledgments

This work was supported by the National Research Foundation (NRF) of Korea, funded by

the Korean Government (grant nos. RS-2024-00463774, RS-2024-00400042, RS-2024-

00509114, RS-2023-NR076986, RS-2025-02216943, and RS-2023-00278980). Additional

support was provided by the KIST Institutional Program (grant no. 2E33761) and the Seoul

National University Hospital (SNUH) Research Fund (grant no. 03-2024-3030).

Conflict of interest

C-H.L. and J.S.H. (KR10-2024-0059322) and M.J., C.-H.L., J.Y.J., and J.S.H. are listed as inventors on provisional patents (KR10-2025-0013774). The authors declare that they have no other competing interests.

Author contributions

M.J. and C.L. conceived and designed the study; J.C., J.H., S.Yee., H.L., D.L., S.Yoon., S.L., Y.J., S.S., and S.K. performed the experiments and data analysis; J.C. and J.H. wrote the original draft of the manuscript; M.J. and C.L. revised the manuscript; and M.J. and C.L. supervised the project and validated the final version. All authors have read and approved the final manuscript.

Data availability statement

The original raw microscopy images underlying the figures, including uncropped images and ROI information, have now been deposited in a public repository (DOI: 10.6084/m9.figshare.31109158).

References

1. Vivier, E. et al. Innate or adaptive immunity? The example of natural killer cells. *Science* **331**, 44-49 (2011).

2. Laskowski, T.J., Biederstadt, A. & Rezvani, K. Natural killer cells in antitumour adoptive cell immunotherapy. *Nat Rev Cancer* **22**, 557-575 (2022).

3. Huntington, N.D., Cursons, J. & Rautela, J. The cancer-natural killer cell immunity cycle. *Nat Rev Cancer* **20**, 437-454 (2020).

4. Moscarelli, J., Zahavi, D., Maynard, R. & Weiner, L.M. The Next Generation of Cellular Immunotherapy: Chimeric Antigen Receptor-Natural Killer Cells. *Transplant*

Cell Ther **28**, 650-656 (2022).

5.   Li, Y., Hermanson, D.L., Moriarity, B.S. & Kaufman, D.S. Human iPSC-Derived Natural Killer Cells Engineered with Chimeric Antigen Receptors Enhance Anti-tumor Activity. *Cell Stem Cell* **23**, 181-192 e185 (2018).

6.   Marin, D. et al. Safety, efficacy and determinants of response of allogeneic CD19-specific CAR-NK cells in CD19(+) B cell tumors: a phase 1/2 trial. *Nat Med* **30**, 772-784 (2024).

7.   Ghobadi, A. et al. Induced pluripotent stem-cell-derived CD19-directed chimeric antigen receptor natural killer cells in B-cell lymphoma: a phase 1, first-in-human trial. *Lancet* **405**, 127-136 (2025).

8.   Lee, M.J., Cichocki, F. & Miller, J.S. CAR therapies: development, design, and implementation. *J Allergy Clin Immunol* (2025).

9.   Daher, M. & Rezvani, K. Next generation natural killer cells for cancer immunotherapy: the promise of genetic engineering. *Curr Opin Immunol* **51**, 146-153 (2018).

10.   Lamers-Kok, N. et al. Natural killer cells in clinical development as non-engineered, engineered, and combination therapies. *J Hematol Oncol* **15**, 164 (2022).

11.   Srivastava, S. & Riddell, S.R. Engineering CAR-T cells: Design concepts. *Trends Immunol* **36**, 494-502 (2015).

12.   Liu, D. et al. The Role of Immunological Synapse in Predicting the Efficacy of Chimeric Antigen Receptor (CAR) Immunotherapy. *Cell Commun Signal* **18**, 134 (2020).

13.   Liao, Y. & Cairo, M.S. CAR-NK's balancing act: when scFv affinity is not too tight, not too loose... but just right? *J Immunother Cancer* **13** (2025).

14.   Rahnama, R. et al. Single-chain variable fragment affinity tuning can optimize anti-

670   AML CAR-NK cell functionality. *J Immunother Cancer* **13** (2025).

15. McErlean, E.M. & McCarthy, H.O. Non-viral approaches in CAR-NK cell

engineering: connecting natural killer cell biology and gene delivery. *J*

Nanobiotechnology **22**, 552 (2024).

16. Hassan, R. et al. Mesothelin Immunotherapy for Cancer: Ready for Prime Time? *J*

Clin Oncol **34**, 4171-4179 (2016).

17. Faust, J.R., Hamill, D., Kolb, E.A., Gopalakrishnapillai, A. & Barwe, S.P. Mesothelin:

An Immunotherapeutic Target beyond Solid Tumors. *Cancers (Basel)* **14** (2022).

18. Morello, A., Sadelain, M. & Adusumilli, P.S. Mesothelin-Targeted CARs: Driving T

Cells to Solid Tumors. *Cancer Discov* **6**, 133-146 (2016).

19. Kachala, S.S. et al. Mesothelin overexpression is a marker of tumor aggressiveness

and is associated with reduced recurrence-free and overall survival in early-stage lung

adenocarcinoma. *Clin Cancer Res* **20**, 1020-1028 (2014).

20. Malla, M. et al. Mesothelin expression correlates with elevated inhibitory immune

activity in patients with colorectal cancer. *Cancer Gene Ther* **31**, 1547-1558 (2024).

21. Lee, Y.E. et al. Synergistic therapeutic combination with a CAF inhibitor enhances

CAR-NK-mediated cytotoxicity via reduction of CAF-released IL-6. *J Immunother*

Cancer **11** (2023).

22. Le, Q. et al. Therapeutic Targeting of Mesothelin with Chimeric Antigen Receptor T

Cells in Acute Myeloid Leukemia. *Clin Cancer Res* **27**, 5718-5730 (2021).

23. Awuah, P., Bera, T.K., Folivi, M., Chertov, O. & Pastan, I. Reduced Shedding of

Surface Mesothelin Improves Efficacy of Mesothelin-Targeting Recombinant

Immunotoxins. *Mol Cancer Ther* **15**, 1648-1655 (2016).

24. Chakraborty, A. et al. A Bispecific Antibody That Targets the Membrane-Proximal

Region of Mesothelin and Retains High Anticancer Activity in the Presence of Shed

Mesothelin. *Mol Cancer Ther* **23**, 1021-1030 (2024).

25.   Liu, X. et al. Highly active CAR T cells that bind to a juxtamembrane region of

mesothelin and are not blocked by shed mesothelin. *Proc Natl Acad Sci U S A* **119**,

e2202439119 (2022).

26.   Liu, X.F. et al. Tumor resistance to anti-mesothelin CAR-T cells caused by binding to

shed mesothelin is overcome by targeting a juxtamembrane epitope. *Proc Natl Acad*

Sci U S A **121**, e2317283121 (2024).

27.   Kelly, R.J., Sharon, E., Pastan, I. & Hassan, R. Mesothelin-targeted agents in clinical

trials and in preclinical development. *Mol Cancer Ther* **11**, 517-525 (2012).

28.   Tomar, S. et al. Development of Highly Effective Anti-Mesothelin hYP218 Chimeric

Antigen Receptor T Cells With Increased Tumor Infiltration and Persistence for

Treating Solid Tumors. *Mol Cancer Ther* **21**, 1195-1206 (2022).

29.   Chakraborty, A. et al. A Bispecific Antibody That Targets the Membrane-Proximal

Region of Mesothelin and Retains High Anticancer Activity in the Presence of Shed

Mesothelin. *Mol Cancer Ther*, OF1-OF10 (2024).

30.   Xiao, Q. et al. Size-dependent activation of CAR-T cells. *Sci Immunol* **7**, eabl3995

(2022).

31.   Zhang, Z. et al. Modified CAR T cells targeting membrane-proximal epitope of

mesothelin enhances the antitumor function against large solid tumor. *Cell Death Dis*

10, 476 (2019).

32.   Chao, Z. et al. Immunological synapse: structures, molecular mechanisms and

therapeutic implications in disease. *Signal Transduct Target Ther* **10**, 254 (2025).

33.   Sterner, R.C. & Sterner, R.M. CAR-T cell therapy: current limitations and potential

strategies. *Blood Cancer J* **11**, 69 (2021).

34.   Olson, M.L. et al. Low-affinity CAR T cells exhibit reduced trogocytosis, preventing

rapid antigen loss, and increasing CAR T cell expansion. *Leukemia* **36**, 1943-1946 (2022).

35. Celichowski, P. et al. Tuning CARs: recent advances in modulating chimeric antigen receptor (CAR) T cell activity for improved safety, efficacy, and flexibility. *J Transl Med* **21**, 197 (2023).

36. Mazinani, M. & Rahbarizadeh, F. CAR-T cell potency: from structural elements to vector backbone components. *Biomark Res* **10**, 70 (2022).

37. Li, L. et al. Loss of metabolic fitness drives tumor resistance after CAR-NK cell therapy and can be overcome by cytokine engineering. *Sci Adv* **9**, eadd6997 (2023).

38. Regis, S., Dondero, A., Caliendo, F., Bottino, C. & Castriconi, R. NK Cell Function Regulation by TGF-beta-Induced Epigenetic Mechanisms. *Front Immunol* **11**, 311 (2020).

39. Shaim, H. et al. Targeting the alphav integrin/TGF-beta axis improves natural killer cell function against glioblastoma stem cells. *J Clin Invest* **131** (2021).

40. Karamitopoulou, E. Tumour microenvironment of pancreatic cancer: immune landscape is dictated by molecular and histopathological features. *Br J Cancer* **121**, 5-14 (2019).

41. Wang, Z. et al. Cancer-associated fibroblast-derived MMP11 promotes tumor progression in pancreatic cancer. *Cancer Sci* **116**, 643-655 (2025).

42. Baek, D.S., Park, S.W., Adams, C., Dimitrov, D.S. & Kim, Y.S. Yeast Mating as a Tool for Highly Effective Discovery and Engineering of Antibodies via Display Methodologies. *Methods Mol Biol* **2491**, 313-333 (2022).

43. Baek, D.S. & Kim, Y.S. Construction of a large synthetic human Fab antibody library on yeast cell surface by optimized yeast mating. *J Microbiol Biotechnol* **24**, 408-420 (2014).

44.   Zhang, J. et al. Loss of mesothelin expression by mesothelioma cells grown in vitro

determines sensitivity to anti-mesothelin immunotoxin SS1P. *Anticancer Res* **32**,

5151-5158 (2012).

45.   Zhang, Y., Xiang, L., Hassan, R. & Pastan, I. Immunotoxin and Taxol synergy results

from a decrease in shed mesothelin levels in the extracellular space of tumors. *Proc*

Natl Acad Sci U S A **104**, 17099-17104 (2007).

46.   Lee, C.H. et al. IgG Fc domains that bind C1q but not effector Fcgamma receptors

delineate the importance of complement-mediated effector functions. *Nat Immunol* **18**,

889-898 (2017).

47.   Kim, M. et al. Construction and validation of a synthetic phage-displayed nanobody

library. *Korean J Physiol Pharmacol* **28**, 457-467 (2024).

48.   Hamieh, M. et al. CAR T cell trogocytosis and cooperative killing regulate tumour

antigen escape. *Nature* **568**, 112-116 (2019).

49.   Li, Y. et al. KIR-based inhibitory CARs overcome CAR-NK cell trogocytosis-

mediated fratricide and tumor escape. *Nat Med* **28**, 2133-2144 (2022).

50.   Wang, Y. et al. Affinity fine-tuning anti-CAIX CAR-T cells mitigate on-target off-

tumor side effects. *Mol Cancer* **23**, 56 (2024).

51.   Peng, L., Sferruzza, G., Yang, L., Zhou, L. & Chen, S. CAR-T and CAR-NK as

cellular cancer immunotherapy for solid tumors. *Cell Mol Immunol* **21**, 1089-1108

(2024).

52.   Shah, K., Al-Haidari, A., Sun, J. & Kazi, J.U. T cell receptor (TCR) signaling in

health and disease. *Signal Transduct Target Ther* **6**, 412 (2021).

53.   Peng, Y. et al. Engineering c-Met-CAR NK-92 cells as a promising therapeutic

candidate for lung adenocarcinoma. *Pharmacol Res* **188**, 106656 (2023).

54.   Issekutz, A.C., Derfalvi, B., Kasermann, F. & Rowter, D. Potentiation of cytokine-

induced proliferation of human Natural Killer cells by intravenous immunoglobulin G.

Clin Immunol **161**, 373-383 (2015).

55.   Shanley, M. et al. Interleukin-21 engineering enhances NK cell activity against

glioblastoma via CEBPD. *Cancer Cell* **42**, 1450-1466 e1411 (2024).

56.   Sun, J.C. & Lanier, L.L. NK cell development, homeostasis and function: parallels

with CD8(+) T cells. *Nat Rev Immunol* **11**, 645-657 (2011).

57.   Lasda, E. & Parker, R. Circular RNAs: diversity of form and function. *RNA* **20**, 1829-

1842 (2014).

58.   Wesselhoeft, R.A. et al. RNA Circularization Diminishes Immunogenicity and Can

Extend Translation Duration In Vivo. *Mol Cell* **74**, 508-520 e504 (2019).

59.   Chen, R. et al. Engineering circular RNA for enhanced protein production. *Nat

Biotechnol* **41**, 262-272 (2023).

60.   Huang, H. et al. Mesothelial cell-derived antigen-presenting cancer-associated

fibroblasts induce expansion of regulatory T cells in pancreatic cancer. *Cancer Cell* **40**,

656-673 e657 (2022).

61.   Wu, J., Wu, W., Zhou, B. & Li, B. Chimeric antigen receptor therapy meets mRNA

technology. *Trends Biotechnol* **42**, 228-240 (2024).

62.   Zhai, X., Mao, L., Wu, M., Liu, J. & Yu, S. Challenges of Anti-Mesothelin CAR-T-

Cell Therapy. *Cancers (Basel)* **15** (2023).

63.   Zhang, Y. et al. Co-expression of IL-21-Enhanced NKG2D CAR-NK cell therapy for

lung cancer. *BMC Cancer* **24**, 119 (2024).

64.   Wong, P. et al. T-BET and EOMES sustain mature human NK cell identity and

antitumor function. *J Clin Invest* **133** (2023).

65.   Hwang, H.J. & Kim, Y.K. Molecular mechanisms of circular RNA translation. *Exp

Mol Med* **56**, 1272-1280 (2024).

66.   Liu, X. et al. Circular RNA: An emerging frontier in RNA therapeutic targets, RNA

796   therapeutics, and mRNA vaccines. *J Control Release* **348**, 84-94 (2022).

797   67.   Pelletier, J. & Sonenberg, N. Internal initiation of translation of eukaryotic mRNA

798   directed by a sequence derived from poliovirus RNA. *Nature* **334**, 320-325 (1988).

799   68.   Tong, M. et al. Robust genome and cell engineering via in vitro and in situ

800   circularized RNAs. *Nat Biomed Eng* **9**, 109-126 (2025).

801   69.   Li, Y.R. et al. Managing allorejection in off-the-shelf CAR-engineered cell therapies.

802   *Mol Ther* (2024).

803   70.   Choi, S.W. & Nam, J.W. Optimal design of synthetic circular RNAs. *Exp Mol Med* **56**,

804   1281-1292 (2024).

805   71.   Du, Y. et al. Efficient circular RNA synthesis for potent rolling circle translation. *Nat*

806   *Biomed Eng* (2024).

807   72.   Lefler, D.S., Manobianco, S.A. & Bashir, B. Immunotherapy resistance in solid

808   tumors: mechanisms and potential solutions. *Cancer Biol Ther* **25**, 2315655 (2024).

809   73.   Zhang, M. et al. TGF-beta Signaling and Resistance to Cancer Therapy. *Front Cell*

810   *Dev Biol* **9**, 786728 (2021).

811   74.   Pan, Y. et al. Regulatory T cells in solid tumor immunotherapy: effect, mechanism and

812   clinical application. *Cell Death Dis* **16**, 277 (2025).

Figure Legends

Fig. 1. Streamlined chimeric antigen receptor (CAR) screening and circRNA-based

CAR (circCAR)-natural killer (NK) cell development to overcome antigen shedding

Schematic workflow illustrating (1) antibody discovery via a yeast surface-display platform

to isolate high-affinity clones targeting membrane-bound mesothelin (MSLN); (2) rapid CAR

optimization via mRNA-engineering of NK cells for functional screening; and (3)

development of persistent shed-resistant CAR-NK cells via circRNAs.

Fig. 2. Screening and characterization of mature mesothelin (matMSLN)-specific scFvs

(a) Schematic representation of matMSLN (residues 296–606) and its soluble counterpart

(solMSLN, residues 296–592). The top bar depicts the full-length matMSLN ending in a GPI

anchor at residue 606; the bottom bar represents solMSLN, which lacks a C-terminal GPI-

anchoring region. The corresponding amino acid sequence is aligned with the annotated

residue positions. **(b)** Heatmap showing the levels and antigen-binding affinities of individual

clones. Yeast-displayed clones were analyzed by flow cytometry using an anti-FLAG

antibody (expression) and 10 nM biotinylated matMSLN or solMSLN (binding). SS1 and

15B6 were included as positive controls, and an isotype antibody served as a negative control.

Clones are ranked by mean fluorescence intensity (MFI). **(c, d)** ELISA-based binding of

recombinant anti-MSLN scFvs to matMSLN **(c)** and solMSLN **(d)**. Binding in immobilized

scFvs incubated with serial dilutions of biotinylated antigen and detected via streptavidin-

HRP. Note that the yeast-display binding values in **(b)** represent semiquantitative MFI-based

screening at a fixed antigen concentration, whereas the ELISA results in **(c, d)** reflect the

quantitative binding of the purified scFvs. Apparent differences between these assays arise

from their distinct experimental formats, with ELISA providing a more accurate measure of

solMSLN binding. **(e, f)** Flow cytometry of cell surface binding by scFvs to MSLN$^+$ **(e)** and

MSLN⁻ K562 cells (**f**). Serially diluted antibodies were incubated with the cells, and the

bound scFvs were detected via an anti-His-FITC secondary antibody. (**g**) Epitope binning of

anti-MSLN scFvs (CLMS10, CLMS32, CLMS36, CLMS76, CLMS95, SS1, and 15B6) by

competitive ELISA. The immobilized scFvs were challenged with biotinylated matMSLN

preincubated with competing antibodies. The heatmap intensity reflects the degree of

competitive binding, indicating epitope overlap. (**h**) Predicted epitope mapping of anti-MSLN

scFvs on the basis of binning data. According to UniProt annotations, the extracellular

domain of MSLN is divided into Region I (residues 296–398), Region II (399–494), and

Region III (495–606). Three independent experiments were performed; the data are presented

as the means ± standard deviations (s.d.).

Fig. 3. mRNA-based screening identifies CLMS10 as a potent shedding-resistant CAR

construct in NK cells.

(**a**) Generation of linear mRNA-based CAR (linCAR)-NK cells via electroporation (EP) of

human primary NK cells. (**b, c**) Establishment of a rapid and cost-effective screening

platform for evaluating CAR-NK cytotoxicity against Capan-2 cells. (**b**) Visualization of

CAR-NK-mediated cancer cell killing. Capan-2 cells were stained with CellTrace Violet Cell

(blue) and live-cell Caspase-3/7 dye (red), and cocultured with CAR-NK cells (green). (**c**)

Quantification of apoptotic cancer cells based on Caspase-3/7 activation ($n = 3$). (**d**)

Shedding inhibition assay for determining CAR-NK cytotoxicity in the presence of

increasing concentrations of solMSLN (residues 296–592); dashed lines indicate the baseline

cytotoxicity of allogeneic NK cells. (**e**) CD107a activation in CAR-NK cells stimulated with

matMSLN-coated plates (20 µg/mL) in the presence or absence of solMSLN (50 µg/mL) was

analyzed by flow cytometry ($n = 3$). (**f**) Cell–cell avidity between linCAR-NK and Capan-2 cells

was quantified via acoustic force-based microfluidic microscopy ($n = 5$). (**g**) Luciferase-based

cytotoxicity of lentivirus (LV)-generated CAR($_{LV}$CAR)-NK cells against Capan-2-Luc and

SK-OV-3-Luc cells ($n = 3$). **(h)** Predicted docking model of CLMS10 scFv in complex with

matMSLN using Schrödinger software. Residues with side chains within 2.5 Å of the binding

interface are highlighted, revealing key antigen-antibody interactions. The soluble MSLN

regions are indicated in marine blue, and the membrane-bound matMSLN regions are

indicated in dark blue. The data are presented as the means ± s.d. Statistical analyses: one-

way ANOVA in **(c)** and **(g)** and two-way ANOVA in **(f)**; two-tailed Student's t tests in **(d)**

and **(e)**; *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.

Fig. 4. CAR-MS10 structural optimization and cytokine coexpression enhance CAR-NK

cytotoxicity and functionality

(a) Four CAR constructs were designed to evaluate combinations of transmembrane (CD8,

NKG2D), costimulatory (4-1BB, OX40, DAP10, and 2B4), and signaling (CD3ζ, DAP12)

domains. **(b)** Cytotoxicity of linCAR-NK cells at 12 h and 24 h post coculture with Capan-2

cells determined via a luciferase-based assay ($n = 3$). **(c)** CD107a degranulation in CD56$^+$

linCAR-NK cells after coculture with Capan-2 cells was analyzed by flow cytometry ($n = 4$).

(d) IFN-γ secretion after 24 h of coculture ($n = 3$). **(e, f)** Metabolic activity [oxygen

consumption rate (OCR)] after 2 h of stimulation on matMSLN-coated plates (20 µg/mL) (n

$= 3$). **(g)** Intracellular signaling pathway activation by the lead construct (linCAR-MS10-

OX40ζ), quantified by normalizing the levels of phosphorylated PLCγ and ERK to their

respective total forms ($n = 3$). **(h–j)** Functional assessment of primary NK cells cotransfected

with linCAR-MS10-OX40ζ and cytokine-encoding linRNAs (IL-2, IL-15, or IL-21).

Luciferase-based cytotoxicity **(i)** and CD107a degranulation **(j)** were measured in NK cells

cotransfected with single or combined cytokines ($n = 3$). The data are shown as the means ±

s.d. One-way ANOVA was used in panels **(b–d)**, **(f)**, **(i)**, and **(j)**; *$p < 0.05$, **$p < 0.01$, ***p

< 0.001, ****$p < 0.0001$.

Fig. 5. circCAR-MS10-CAR-NK cells exhibit prolonged CAR expression, sustained

cytotoxicity, and resistance to antigen shedding

(a) Generation of circCAR-NK cells via EP. **(b–d)** CAR expression and viability of linCAR-

and circCAR-NK cells over time. **(c)** CAR levels were measured and compared to those of

the EP-negative control ($n = 5$). **(d)** NK viability was assessed by live/dead staining following

EP ($n = 3$). **(e)** Serial killing assay of CAR-NK cells sequentially exposed to fresh Capan-2-

Luc cancer cells at 12 h-intervals. Serial cytotoxicity **(f)** and IFN-γ secretion **(g)** were

evaluated to determine the sustained antitumor activity of circCAR-NK cells ($n = 3$). **(h, i)**

Impact of solMSLN (50 µg/mL) on CAR-NK cell function during repeated tumor challenges.

Serial cancer killing **(h)** and IFN-γ secretion **(i)** were assessed following solMSLN treatment

(50 µg/mL) at each round ($n = 3$). The data are presented as the means ± s.d. Statistical

analyses: one-way ANOVA for **(c), (d), (f),** and **(g)**, and two-tailed Student's t tests for **(h)**

and **(i)**; *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.

Fig. 6. circCAR-MS10-NK cells maintain superior antitumor activity under CAF-

induced shedding conditions *in vitro* and *in vivo*

(a, b) Soluble MSLN (solMSLN) levels were elevated in Capan-2 cells cocultured with

human pancreatic primary fibroblasts (HPPFs) or CAFs (CAF5, CAF6, CAF16, and CAF22),

as shown by western blotting and quantified relative to those in cancer-only controls ($n = 3$).

(c) *In vitro* real-time cytotoxicity of circCAR-NK cells cocultured with mCherry-expressing

Capan-2 cancer cells, in the absence or presence of CAFs. CAFs were coincubated to mimic

the tumor microenvironment characterized by shed MSLN secretion. Cancer cell survival was

monitored by quantifying mCherry/RFP fluorescence, presented as relative cancer viability,

which was defined as the mCherry signal normalized to the 0 h value (set to 1.0) ($n = 3$).

Statistical comparisons were performed between the circCAR-SS1 and circCAR-MS10

groups at each indicated time point. **(d)** IHC of MSLN expression in human pancreatic tumor

tissues using a CLMS10-derived antibody across stages 1A, 2A, and 4. **(e, f)** *In vivo*

therapeutic effects of CAR-NK cells were evaluated in a metastatic mouse model with

intraperitoneal AsPC-1-MSLN-Luc tumors containing CAF6 cells. **(f)** Representative IVIS

images of mice treated with mock, linCAR-NK (SS1, 15B6, MS10), circCAR-MS10, or

$_{LV}$CAR-MS10 cells ($n = 4$ per group); all mRNA-CAR groups were coelectroporated with IL-

21 mRNA. **(g)** Body weight monitoring during treatment. **(h, i)** Tumor burden quantified via

IVIS imaging over time **(h)** and on day 21 **(i)** ($n = 4$). The data are presented as the means ±

s.d. Statistical analyses: one-way ANOVA for **(b)** and **(h)** and two-way ANOVA for **(c), (g),**

and **(i)**; $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$.

Part I
Antibody Discovery
Shed-resistant anti-MSLN antibody

Part II
CAR-NK Generation
mRNA-based CAR-NK platform

Part III
CAR Optimization & Validation
Fast-screening for therapeutic circCAR-NK

a

296 · matMSLN · 580 · 585 · 590 · 595 · 600 · 606 · GPI anchor

Q D D L D T L G L G L Q G G I P N G Y L V L D L S M Q E A L S

296 · solMSLN · 592

b

Expression

Binding to matMSLN

Binding to solMSLN

CLMS10 CLMS20 CLMS32 CLMS36 CLMS76 CLMS88 CLMS95

SS1 15B6 Isotype

30 — 300

30 — 2000

c Absorbance at 450 nm vs Concentration of matMSLN (nM)

d Absorbance at 450 nm vs Concentration of solMSLN (nM)

- SS1
- 15B6
- CLMS10
- CLMS20
- CLMS32
- CLMS36
- CLMS76
- CLMS88
- CLMS95

e MSLN-positive K562 — MFI (FITC-A, x1000) vs Concentration of anti-MSLN scFv (nM)

f MSLN-negative K562 — MFI (FITC-A, x1000) vs Concentration of anti-MSLN scFv (nM)

g

	SS1	15B6	CLMS10	CLMS32	CLMS36	CLMS76	CLMS95
SS1							
15B6							
CLMS10							
CLMS32							
CLMS36							
CLMS76							
CLMS95							

Strong — Weak

h

Distal

Domain I (296-398) — SS1

Domain II (399-494)

Strong competition

Weak competition

Cleavage site

Proximal

Domain III (495-606)

10 76 32 15B6 95 36

a

5' Cap (m7G) Kozak CD8a hinge & TMD P2A Poly A tail (110nt)

5'UTR — CD8SP — scFv — 4-1BB — CD3ζ — GFP — 3'UTR — AAA···

pNK Cells → IVT mRNA (mRNA 1, mRNA 2, mRNA 3, mRNA 4 ...) → Electroporation → mRNA-based CAR-NK

b linCAR-NK : Capan-2 (2.5:1)

Mock GFP | CAR SS1 | CAR 15B6 | CAR MS10 | CAR MS20 | CAR MS32 | CAR MS36 | CAR MS76 | CAR MS88 | CAR MS95

0h

12h

🟩 CAR-NK 🟦 Cancer 🟥 Caspase-3/7   Scale bar = 300 μm

c

Caspase-3/7 fluorescence [MFI]

Mock-G, CAR-SS1-G, CAR-15B6-G, 10-G, 20-G, 32-G, 36-G, 76-G, 88-G, 95-G
CAR-MS

** ; **** ; **

d linCAR-NK : Capan-2-Luc (1:1)

Mock-G | CAR-SS1-G | CAR-15B6-G | CAR-MS10-G

Cytotoxicity (%) vs solMSLN (μg/mL) 0 10 20 50 100

CAR-SS1-G: *, ***, ****, ****
CAR-15B6-G: ns, ns, ns, ns
CAR-MS10-G: ns, ns, ns, ns

CAR-MS20-G | CAR-MS32-G | CAR-MS36-G

CAR-MS20-G: ns, ns, ns, ns
CAR-MS32-G: ns, ns, *, *
CAR-MS36-G: ns, ns, **, **

e

CD107a⁺ % (fold)

☐ matMSLN ☐ matMSLN + solMSLN

Mock-GFP, CAR-SS1-G, CAR-15B6-G, CAR-MS10-G, CAR-MS20-G, CAR-MS32-G, CAR-MS36-G

*** ; ns ; ns ; ns ; *** ; ***

f linCAR-NK : Capan-2-Luc (1:5)

% CAR-NK Cells Bound vs Force (pN)

— EP only
— CAR-SS1
— CAR-15B6
— CAR-MS10
— CAR-MS20

n=5   ****

g

Anti-MSLN-CAR-GFP Lentivirus → pNK Cells → Virus transduction → Anti-MSLN CAR generation

LV CAR-NK:Capan-2-Luc (1:1)

Cytotoxicity (%)

Mock-GFP, CAR-SS1-G, CAR-15B6-G, CAR-MS10-G, CAR-MS20-G

*** ; * ; * ; * ; *

LV CAR-NK:SK-OV-3-Luc (1:1)

Cytotoxicity (%)

Mock-GFP, CAR-SS1-G, CAR-15B6-G, CAR-MS10-G, CAR-MS20-G

**** ; ** ; ** ; * ; ns

h

MSLN
CLMS10
GPI

D580
D577 2.3 Å
Y594 1.7 Å 2.3 Å 2.4 Å R563
1.8 Å 2.2 Å 2.6 Å
L595

▲ Cleavage site

a

Cancer cell only

Cancer with CAF

Soluble MSLN secretion

b

Band intensity (fold)

Capan-2 + | - + | - + | - + | - +

HPPF   CAF5   CAF6   CAF16   CAF22

c

circCAR-NK : Cancer (1:1)

circCAR-NK : Cancer/CAF (1:1)

Relative cancer viability

Time (h)

Time (h)

● Control ○ Mock ● circCAR-SS1 ● circCAR-MS10

d

CLMS10 antibody

Normal | Stage 1A | Stage 2A | Stage 4

Scale bar = 50 μm

e

AsPC-1-MSLN-Luc(3e6) CAF6 (1e6) (i.p.)

🟡 Bioluminescent imaging
🔴 CAR-NK administration

CAR-NK (1e7) (i.v.)

D0   6 8   11   14   17 18   21

Days after tumor injection

f

+ IL-21 linRNA

Control | Mock | CAR-SS1 | CAR-15B6 | linCAR-MS10 | circCAR-MS10 | _LV_CAR-MS10

D6
D8
D11
D18
D21

x10⁶ 1.5 / 1.0 / 0.5

Radiance (p/sec/cm²/sr)
Min: 6.50e4
Max: 1.50e6

x10⁶ 3.0 / 2.0 / 1.0

Radiance (p/sec/cm²/sr)
Min: 2.00e5
Max: 3.50e6

g

Body weight (g)

Days

ns

h

Radiance (p/sec/cm²/sr)

Days

○ Control
● Mock
● linCAR-SS1
● linCAR-15B6   + IL-21 linRNA
● linCAR-MS10
● circCAR-MS10
● _LV_CAR-MS10

i

Day 21

Radiance (p/sec/cm²/sr)

Control | Mock | linCAR-SS1 | linCAR-15B6 | linRNA | circRNA | LV

CAR-MS10