Microbes use transporters to regulate the release of metabolites based on value

Snorre Sulheim[1,2*], Gunn Broli[3], Peter F. Doubleday[4], Alisson Gillon[1], Julien S. Luneau[1], Prajwal Padmanabha[1], Andrew Quinn[1], Eric Ulrich[1], Margaret A. Vogel[1], Nicola Zamboni[4], Philipp Engel[1] & Sara Mitri[1*]

Affiliations

1. Department of Fundamental Microbiology, University of Lausanne

2. Department of Biological Sciences, University of Bergen

3. Department of Biotechnology and Nanomedicine, SINTEF Industry

4. Institute of Molecular Systems Biology, ETH Zurich

*Corresponding author. Email: snorre.sulheim@unil.ch; sara.mitri@unil.ch

Abstract

Microbial interactions are shaped by the exchange of metabolites, ranging from intermediates in central carbon metabolism to amino acids, vitamins and fermentation by-products[1]. Yet, the underlying reasons for metabolite release, and the mechanisms and factors influencing release rates are not well understood. Here, using a combination of computational and experimental approaches on seven different microbes we show that release rates are negatively correlated with the value of metabolites, and that this relationship explains more variability in release rates than other frequently associated factors. By measuring metabolite release from 66 *E. coli* mutants lacking individual metabolite-transport genes, we find that transporters both cause and counteract net metabolite release. These findings show that metabolite release constitutes a fitness cost that microbes have evolved to reduce and reveal the underlying mechanism. This new conceptual explanation for metabolite release reshapes our understanding of how microbial interactions emerge and persist.

Main

Introduction

Metabolites are frequently exchanged between microbes in natural communities[1]. Such cross-feeding plays a key role in maintaining community diversity and function[2–6]. While certain occurrences of cross-feeding arise from extracellular metabolism such as polymer degradation by extracellular enzymes[6,7], many observations of cross-feeding stem from the release of intracellular metabolites into the environment by microbial community members[8–10]. Recent *in vitro* quantifications of microbial cultures have demonstrated that this release includes high-energy glycolytic intermediates, amino acids, vitamins and organic acids[11–15]. The ubiquity of metabolite release may help explain the high frequency of auxotrophs isolated from natural environments that may be supported by co-cultivated partners[4,15–17]. However, why so many diverse intracellular metabolites are being released and what explains differences in release rates remains unanswered.

Previous efforts to answer why microbes release intracellular metabolites have revealed fundamental principles of metabolism, including rate-yield trade-offs and constraints on proteome allocation[18–22], mechanisms of pathway regulation and maintenance of homeostasis[23], optimal catabolic pathway lengths in different environments[24,25] and redox balancing[26,27]. These explanations are context-, metabolite- or species-specific. However, to fully understand metabolite release, we need to consider broader principles that apply across diverse microbial taxa and environments. Cell lysis and passive diffusion across the cell membrane are two such commonly proposed broad mechanisms[28–31].

Here, we approach the problem by asking: does metabolite release generally confer a benefit or impose a cost on the producer? While recent work has explored how metabolite release can be beneficial, e.g. by alleviating costs associated with molecular noise[31,32], we here propose and test an alternative hypothesis that metabolite release comes with a cost that microbes have evolved to reduce.

Metabolite release rates are negatively correlated with metabolite value

If metabolite release generally imposes a fitness cost, and natural selection acts to minimize such costs, then the release rates of specific metabolites should be negatively correlated with their relative importance, or value, to the microbe (Fig. 1A). Release of more valuable metabolites – that require more energy to produce – should have a more severe negative impact on fitness and be more strongly selected against.

To quantify metabolite release rates across different species, we first analysed an existing time-series exometabolome dataset with high temporal resolution and absolute quantification of 19 to 37 metabolites collected from batch cultures of *E. coli*, *B. licheniformis, S. cerevisiae* and *C. glutamicum* in high glucose concentrations[12]. We estimated net uptake and release rates for all metabolites using simple linear regression, modelling extracellular metabolite concentration as a linear function of the area under the growth curve (AUC), until the point where the extracellular concentration saturated or the exponential growth phase ended (Methods, Fig. 1B, and Figs. S1-S4). The initial dynamics of 44-95% of metabolites were well described by this simple linear model (two-sided Wald test, FDR < 0.05, R^2 > 0.5, Fig. S1-S4).

To estimate metabolite value, we used previously reconstructed enzyme-constrained genome-scale metabolic models (ecGEMs)[33–35]. Genome-scale metabolic models (GEMs) are mathematical representations of a species' metabolic capacity that can for example be used to predict optimal metabolic phenotypes across various environments[36,37]. The additional enzyme constraints account for the total enzyme pool limitation in cells[19], enabling prediction of relevant trade-offs and temporal dynamics, e.g. acetate overflow in *E. coli*[34]. Various measures of metabolite value have been proposed in the past[38–40]. Here we define metabolite value as the negative change in growth rate on metabolite release close to optimal growth as predicted by enzyme-constrained GEMs (Fig. 1C). This measure explicitly reflects the fitness penalty on growth rate associated with the marginal release of a metabolite, and it can easily be used for different species across different environmental contexts.

With these two measures, we could now test our hypothesis that release rate should be

negatively correlated with metabolite value. Indeed, this holds for *E. coli, S. cerevisiae, B.*

licheniformis in glucose medium (Pearson ρ, P < 0.001, Fig. 1D)*.* This did not hold for *C.*

glutamicum (Fig. 1D), possibly because the engineering of its central carbon metabolism for

enhanced lysine production has also affected the release of other metabolites[41].

To test whether our hypothesis held beyond simple glucose environments, we conducted new

well-controlled batch cultivations of *E. coli* in minimal media with galactose, L-malate or L-alanine

as the single carbon source. These carbon sources were chosen to span the variability in *E. coli*

metabolism to avoid biasing the results towards a particular class of compounds or metabolic flux

patterns (Methods, Fig. S5), since this has been shown to produce similar exometabolome

patterns[42]. Triplicate batch bioreactor experiments were performed under strictly controlled

aerobic conditions with active regulation of pH and dissolved oxygen (Fig. S6). Relative

extracellular concentrations of 126 metabolites were then quantified at three exponential-phase

samples and one sample from stationary phase. Of these metabolites, 41-56% were released at

significant rates during exponential phase (two-sided Wald test, FDR < 0.05, Figs. S7-S10),

consistent with previous findings showing that microbes generate rich exometabolomes[12,13].

From this set, we were able to estimate absolute release rates for 34 metabolites that we

compared with estimated metabolite values. Again, we found significant negative correlations for

all three conditions (Pearson ρ, P < 0.001, Fig. 1E).

Finally, we analysed a third dataset that in addition to *E. coli* and *Pseudomonas putida* contained

environmental isolates from the genera *Enterobacter* and *Pseudomonas*[42]. Unlike previous

datasets, these experiments were conducted without pH and oxygen control, offering an

opportunity to test the hypothesis under more variable conditions. Despite these differences, we

again found consistent negative correlations (Fig. 1F), suggesting that the relationship between

metabolite value and release rate is robust across species, nutrient conditions, and experimental

platforms. Importantly, the ecGEMs used here were reconstructed by different research groups

using different pipelines, suggesting that our results are not dependent on the choices made

during model reconstruction. This conclusion was also supported by consistent results when

metabolite values were estimated using four other *E. coli* GEMs (Fig. S11).

Metabolite value is the most important factor explaining variability in metabolite release

rates

The consistent negative correlation between metabolite release rates and metabolite values

encouraged us to ask how well this factor explains release variability compared to other metabolic

or physiochemical properties previously associated to metabolite release[32,43]. We focused on the

data from bioreactor batch cultures of *E. coli* with glucose, galactose, L-malate or L-alanine as the

carbon source, where conditions were well-defined, the measured metabolites were diverse and

reference values for intracellular metabolite concentrations were available[44–47]. We first built

univariate linear models to quantify the importance of these factors: metabolite value,

intracellular concentration, compound class, solubility in oil vs water (log P), molecular weight,

topological polar surface area, charge, hydrogen bond acceptor and donor count, rotatable bond

count, turnover and carbon source (Fig. 2A). Metabolite value explained 43% of the variability,

more than any other factor (Fig. 2B). The structure-based classification of compound class also

had good predictive power (29% explained), primarily because carboxylic and keto/hydroxy acids

were released at significantly higher rates than amino acids and other compounds (Fig. 2A).

However, these two factors are clearly confounded as class-level differences mirrored significant

differences in metabolite values (Fig. 2C). The number of rotatable bonds, solubility in oil vs water

(log P) and the number of hydrogen bonds, each related to membrane permeability[48,49], were

also predictive of metabolite release, suggesting passive diffusion as a relevant mechanism.

Metabolite turnover is predicted to correlate negatively with release rates, according to the noise-

averaging cooperation hypothesis (low turnover metabolites are more susceptible to noise and

therefore more useful to share within the population)[32], but our dataset shows little support for

this.

When we asked how well the best linear model could predict metabolite release rates, we found

that 63% of the variability could be explained by metabolite value, compound class, intracellular

concentration and charge (Fig. 2B). However, predictive accuracy declined when applied to out-
of-sample conditions (45%, P = 0.04, 52.9 ± 3.8% expected from random model, Fig. S12) or out-
of-sample metabolites (50%, P < 0.001, 55.2 ± 0.7% expected from random model, Fig. S13), likely
due to correlations among the factors, which warrants caution when interpreting factors' relative
importance (Fig. S14). Nevertheless, excluding metabolite value as a factor reduced model
performance on out-of-sample conditions (10% decline in median R^2, P = 0.003, one-sided Mann-
Whitney U, N = 20, Fig. S15), and model quality was largely dependent on including metabolite
value or compound class (Fig. S16). Furthermore, we found differences in metabolite value
between conditions to be negatively correlated with differences in release rate (Pearson ρ, P =
2e-3, Fig. S17). This not only supports the relevance of this factor in predicting out-of-sample rates
but also raises the question of whether microbes adapt their release rates to context-dependent
costs. Finally, when we estimated each factor's importance separately for each microbe in each
dataset (except *C. glutamicum*), metabolite value still emerged as the most robust explanatory
factor (Fig. 2D).
156
Cell lysis is commonly considered a key mechanism for metabolite release[30,31]. To measure its
importance in our experiments, we used *E. coli* batch cultures with paired sampling of intra- and
extracellular metabolites, as well as flow cytometry with live/dead staining to quantify the
fraction of lysed cells. In general, less than 10% of extracellular concentrations could be explained
by the release of intracellular metabolites following cell lysis, and often much less (Fig. 3A). We
then extended this analysis by comparing our collection of *E. coli* extracellular metabolite
concentrations to published intracellular concentrations[44–47]. To avoid asserting equal lysis rates
across conditions, we asked how large a fraction of the cell population would need to be lysed to
account for the extracellular levels, considering only the contribution from intracellular
metabolite pools (Fig. 3B). Only 10.4% (157/1508) of the data points fall within the range of
measured lysis fraction (0.2-2.3%, mean ± std across conditions = 0.6 ± 0.6%) or below, and more
than 56.3% (849/1508) are above the limit where more than the whole cell population is required.
Glutamate, glutamine and NAD are the only metabolites where most data points can be

accounted for by cell lysis, due to high intracellular concentrations and moderate to low

extracellular levels (Fig. S18).

Degradation of cell debris may also contribute to extracellular metabolite pools[50]. In *E. coli,* this

degradation is dependent on the presence of Lon protease in the cell lysate[51]. However, the

benefit of cell debris degradation by Lon was only significant well beyond exponential phase, and

only 19 ± 6% of oligopeptides (6-10 amino acids) in the lysate were catabolised by Lon over 20

hours[51]. Thus, in our dataset where 90% of the samples were collected before 26 hours (67% in

exponential phase), we expect only a fraction of the proteome from lysed cells to contribute to

the extracellular concentrations of amino acids. When we included 10% proteome degradation

in our analysis, only the extracellular levels of tryptophan and threonine additionally became fully

explained by cell lysis (Fig. S19). However, apart from amino acids, the metabolites in Fig. 3B are

not typical degradation products. Overall, these results suggest that cell lysis and proteome

degradation are insufficient to explain extracellular levels of most metabolites – consistent with

previous work[12,30].

Metabolic disruption and evolution change metabolite release rates

Given the effect of gene deletions on intracellular metabolite levels[52], we assumed that the lack

of negative correlation between metabolite value and release rate in *C. glutamicum* (Fig. 1D)

could be a result of its engineered metabolism[41]. To test the sensitivity of metabolite release

patterns to modifications in core metabolism, we measured the exometabolome of eight *E. coli*

knockout (KO) strains in late exponential phase in galactose medium (Fig. 4A). These gene

knockouts target different parts of core metabolism, were previously shown to affect intracellular

metabolite pools[52], and were predicted to change intracellular flux patterns (Fig. S20). In line with

our expectations, 14-38% (16-42 of 111) of the extracellular metabolite concentrations differed

significantly from the wild type across the KO strains (Welch's t-test, FDR < 0.05, Fig. 4B). In

contrast, only 2.7% of metabolite concentrations were different in the negative control strain

Δ*lacA*. Each KO strain displayed distinct exometabolome patterns (Fig. 4C), and we observed the

most pronounced differences for metabolites in proximity of the deleted reaction (Figs. S21-S23),

consistent with intracellular effects of enzyme-deletions[52]. The WT reference for these KO strains

from the KEIO collection[53] - *E. coli* BW25113 - had largely comparable rates as *E. coli* MG1655

used in the experiments presented in Fig. 1 (Fig. S24).

Our central hypothesis for the negative correlation between metabolite value and release rate is

that bacteria face a trade-off: the cost of preventing metabolite release versus the energy lost

through that release. If this trade-off is under strong selection, then disrupting the balance (e.g.

via a gene knockout) should trigger rapid evolutionary compensation to restore it. To test this, we

evolved the Δ*aceE* and Δ*sucB* mutants – chosen for their severe growth defects and distinct

exometabolome characteristics – for ~100 (replicates M2-M4) and ~200 generations (M5-M7) in

galactose medium, respectively (Fig. S25). The strains evolved in chemostats, where byproducts

are continuously removed to avoid accumulation and later uptake of metabolites. From each

chemostat endpoint, we isolated three strains and selected from each replicate the fastest

growing clone (Fig. 4, D and E). These clones showed 94-146% (Δ*aceE*) and 50-82% (Δ*sucB*) growth

rate increases relative to their ancestors (Fig. 4, F and G). To assess changes in metabolite release,

we measured extracellular concentrations of 5 metabolites at the end of exponential growth (Fig.

4, D and E). These metabolites were selected because their release was strongly disturbed in the

ancestors (Figs. S21-S23), and either near the deleted reaction (pyruvate and lactate for the

Δ*aceE*; citrate, isocitrate and *cis*-aconitate for Δ*sucB*) where we expected the strongest effects, or

located elsewhere in the metabolic network and of different chemical classes to probe more

widespread effects. Lactate levels were significantly reduced in 2/3 Δ*aceE* isolates (Fig. 4F), which

also exhibited the highest growth rates. The reductions were, however, not sufficient to reach WT

concentrations. In contrast, all three Δ*aceE* isolates showed increased pyruvate release, contrary

to expectations. These results were qualitatively invariant to growth rate normalization (Fig. S26).

For the Δ*sucB* isolates, we found in general small reductions in extracellular concentrations (Fig.

4G), but some of these differences change qualitatively upon growth rate normalization (Fig. S27).

For the metabolites elsewhere in the metabolic network, the outcomes were variable (Figs. S28

and S29).

To estimate the fitness contribution of reducing metabolite release, we used ecGEMs to predict

relative growth rate penalties associated with the observed release rates, using the ancestors'

mean growth rates as references. In the Δ*aceE* ancestor, lactate release is predicted to reduce

fitness by 23% whereas pyruvate release has negligible impact (0.02%). The reduced release of

lactate in the isolates from chemostat M2 and M4 is predicted to reduce the negative fitness

effect to only 16% and 7%, respectively. These contributions are small compared to the observed

increase (94-146%). The pattern is similar for the Δ*sucB* isolates: the estimated net fitness

reduction of citrate, isocitrate and cis-aconitate in the ancestor is only 0.37%, far from sufficient

to explain the improved growth rates.

To better understand the reasons for increased growth and changed release rates, we identified

sequence variants in the selected fast-growing strains and mutations that had fixed in the

chemostat populations (Table S1-S4). Aside from the Δ*aceE*-M4 isolate, which likely

hypermutated due to a *mutT* mutation, the evolved Δ*aceE* isolates have few sequence variants,

and either related to galactose uptake (e.g. *galS* in M2, M4 and M7, regulating galactose uptake)

or modulating fluxes near the deleted reaction (e.g. *poxB* in M2, *ldhA* in M4, and *icd, aceK* and

sdhA in M5, M6 & M7 and M7, respectively). Fixed non-synonymous variants in the chemostat

populations support these observations, e.g. *ldhA* and *galS* were fixed in M2 and *ilvG* was fixed in

M4. The Δ*sucB* populations have many more fixed mutations, but among the ones shared across

replicates we find again *galS* and *aceK*. Together, these results show that metabolite release rates

change as microbes evolve, but in our experiments, these changes were likely secondary to other,

more dominant drivers of fitness – leading to inconsistent changes in metabolite release across

replicates.

Transporters cause and counteract metabolite release

Our findings so far suggest that microbes can modulate metabolite release rates. To assess the

underlying mechanism, we profiled the exometabolomes of 66 *E. coli* KO mutants from the KEIO

collection, each strain devoid of a gene associated with active or passive transport across the

inner or outer membrane (Table S5). These genes were selected to span the diversity of

transporters, covering major functional classes (ABC, PTS, symporter, antiporter, porin, and

channel), a variety of substrates (amino acids, sugars, nucleosides, carboxylic acids, and others),

and expected direction (import, export, or both). We cultivated each KO mutant and the WT in

glucose minimal medium and quantified changes in extracellular metabolite concentrations over

the course of batch growth using flow-injection analysis time-of-flight mass spectrometry (FIA-

TOF MS)[54]. The final dataset comprises 2184 unique samples collected at 10 timepoints, yielding

more than 8700 mass spectrometry measurements. We could annotate m/z ion peaks to 333

metabolites in the *E. coli* metabolome. Three strains (Δ*tolC*, Δ*dppB,* and Δ*ptsG*) showed severely

reduced growth and were excluded from downstream analyses. We quantified the *effect* of each

transporter on each metabolite as the ratio between the mean KO-WT difference and the within-

strain variability (Fig. 5A). Statistical significance for all 15,848 strain-metabolite *effects* was

assessed using a permutation test (Methods). The dataset is made available for interactive

exploration at keio.unil.ch.

A positive effect implies that removing a transporter increases the net release of the respective

metabolite in the KO strain compared to the WT, while a negative effect implies the opposite. We

found a strong positive effect in 2.2% (343/15848) of the strain-metabolite pairs (*Effect* > 3*, P* <

0.005). In many cases, the strongest effect matched the annotated function of the transporter

(Fig. 5B). For example, deletion of the high-affinity proline/sodium symporter gene *putP* strongly

increased proline release, suggesting that PutP counteracts proline release in the WT by re-

importing proline lost via other mechanisms. This mechanism has already been described for *E.*

coli strains selected for proline secretion[55]. Similarly, deletion of the oxoglutarate importer *kgtP*

most strongly increased oxoglutarate release. For glutamate, three of the four annotated

transporters linked to positive effects, suggesting that they collectively counteract its release (*P* <

0.005, Fig. 5C). However, we also observed strong, and sometimes diverse effects extending

beyond the metabolites annotated to the respective transporters in EcoCyc[56]. For example,

deletion of the glutamine importer *glnP* strongly increased the extracellular concentration of 20

metabolites (*Effect* > 3, *P* < 0.005), including valine, hydroxybutanoic acid, aminobutanoic acid,

isoleucine/leucine, alanine, and glutamine (Fig. S30).

Strong negative effects were found in 2.4% (379/15848) of the strain-metabolite pairs (*Effect* < -

3, *P* < 0.005), showing that transporters frequently cause metabolite release. Some of these

effects were linked to transporters primarily associated with metabolite export, such as the valine

exporter YgaZ or RhtC, which mediates threonine efflux (Fig. 5D)[57,58]. However, we observed

equally strong effects for transporters that are important for metabolite uptake such as the purine

and adenosine transporters PunC and AdeP, respectively (Fig. 5E)[59–61]. Similarly, the "exporters"

YgaZ and RhtC showed positive effects comparable in magnitude to their negative ones (Fig. S30).

To test whether the measured effects of transporter deletion on metabolite release are

ecologically relevant, we co-cultivated the WT and 2-4 KO strains with a proline, methionine or

isoleucine auxotroph. With one exception, the Δ*ygaZ* strain which supported the growth of the

Δ*ilvE* isoleucine auxotroph better than the WT despite the measured negative effect on

isoleucine/leucine, strains with a measured positive effect on amino acid release supported the

growth of the corresponding auxotrophs better than the WT (Fig. S31).

Of the 63 KO strains with growth comparable to the WT, 55 strains strongly influenced one or

more metabolite (|*Effect*| > 3, *P* < 0.005, Fig. 5F). Twenty-nine strains strongly influenced ten or

more metabolites, and 37 showed both positive and negative effects. While the balance between

positive and negative effects differed between strains, the overall dataset showed a striking

symmetry with almost equally strong and numerous positive and negative effects (Fig. 5G). From

the metabolite perspective, 122 metabolites were affected by one or more strains, and 57 of

these were affected in both directions (|*Effect*| > 3, *P* < 0.005, Fig. 5F). We found no significant

differences in the distribution of effect sizes across metabolite classes, nor any correlation

between the value of a metabolite and how frequently it was affected by the deletion of a

transporter (Fig. S31). Likewise, effect sizes were similarly distributed across transporter classes,

and neither the number of affected metabolites nor the maximum effect size correlated with

average transporter expression levels (Fig. S31)[62]. Taken together, our results demonstrate that

transporters expressed in *E. coli* are both a cause of metabolite release and a mechanism for *E.*

coli to counteract it. The complexity of strain-metabolite connections suggests that trade-offs are

common.

Discussion

To better understand metabolic interactions between microbes, it is necessary to develop and

test hypotheses for the fundamental principles that govern them. Here we propose that the broad

release of intracellular metabolites is in general a loss of energy and that for each metabolite

there is a trade-off between the cost of its release and the cost of preventing it. This hypothesis

connects to previous work linking fitness costs to microbial traits, including amino acid uptake

rates[14], proteome composition[10,38], properties of extracellular enzymes[63] and metabolic

strategies[64].

We find consistent support for our hypothesis: (i) metabolite values are negatively correlated with

metabolite release rates across different species and nutrient environments, (ii) metabolite value

explains the most variability in release rates among all tested factors, (iii) release rates cannot be

explained by cell lysis alone, (iv) knock-out of metabolite-transport genes frequently change net

metabolite release, providing a mechanistic explanation for the observed release and for how

microbes can act to reduce the net release of valuable metabolites. The complexity of KO strain–

metabolite interactions (Fig. 5F) suggests why eliminating metabolite release entirely may be

challenging, given the apparent trade-off between release and recapture, and the need to express

transporters at basal levels to sense and adapt to environmental changes[65–67]. The high frequency

of metabolites with both significant positive and negative effects suggests that futile cycles might

be common and raises the question of whether such cycles contribute to intracellular

homeostasis, extending beyond their previously demonstrated role in metal ion regulation[68].

Furthermore, we find that release rates are sensitive to changes in intracellular metabolism, both

caused by gene knockouts and subsequent evolution. Yet on the timescale of our evolutionary

experiments, changes in release rates were inconsistent across replicates and did not always

match our expectations. This may reflect alternative evolutionary paths that improved fitness

independently of metabolite release, or selection favouring higher growth rate over higher yield.

The large-scale exometabolomics screen of *E. coli* transporter knockout strains provides a unique

resource for refining transporter annotations in databases and metabolic models. In parallel, our

finding that simple linear models based on metabolite value can predict order-of-magnitude

release rates offers a new avenue for such models to forecast microbial interactions. More

broadly, our findings reshape how we think about metabolite release and the evolution of

microbial cross-feeding. While it is commonly assumed that cross-feeding either emerges from

extracellular enzymes or from the release of waste products[2,69,70], our results suggest another

alternative: Microbes release metabolites because using, retaining or immediately recapturing

them is not worth the investment. This increases the scope for commensal cross-feeding to occur,

which in this scenario can arise without the need to invoke the evolution of cooperation[71] or being

vulnerable to cheaters[72].

A Hypothesis

C Estimation of metabolite values using ecGEMs

$$\text{Metabolite value} = -\frac{\Delta y}{\Delta x}$$

B Estimation of release rates from batch cultures and time-series exometabolomics

Poor fit: $R^2 < 0.5$ or $P > 0.05$

AUC of biomass [gDW/L · h]

D Paczia et al.

E This work

○ Carboxylic acids ○ Amino acids ○ Organooxygen compounds ○ Keto / hydroxy acids ○ Other

E. coli
$\rho = -0.53$ ($P = 7e\text{-}04$)

B. licheniformis
$\rho = -0.70$ ($P = 2e\text{-}06$)

S. cerevisiae
$\rho = -0.69$ ($P = 2e\text{-}04$)

C. glutamicum
$\rho = 0.10$ (ns)

Galactose
$\rho = -0.75$ ($P = 3e\text{-}04$)

L-alanine
$\rho = -0.66$ ($P = 6e\text{-}04$)

L-malate
$\rho = -0.76$ ($P = 2e\text{-}04$)

F Vila et al.

Release rate [mmol/gDW/h]

Metabolite value [gDW/mmol]

Figure 1: Correlation between microbial metabolite release rates and metabolite values. *A) Hypothesis: if metabolite release generally constitutes a fitness cost that microbes have evolved to reduce, metabolite release rates should be negatively correlated with metabolite values. Three metabolites are used to conceptually illustrate their relative position based on their estimated release rate and metabolite value, see panels B and C. B) Metabolite release rates were estimated from time-series exometabolomics data from batch cultures by linear regression of extracellular concentrations vs area under the curve (AUC) of biomass. This panel shows extracellular concentrations of 37 metabolites measured during exponential phase of E. coli cultivated in glucose medium by Paczia et al. (2012), and the linear regressions used to estimate net release rates. Linear regressions with an $R^2 < 0.5$ or $P > 0.05$ (two-sided Wald test) are drawn in red. Full time-series, including stationary phase and linear regression for E. coli, B. subtilis, C. glutamicum and S. cerevisiae are shown in Figs. S1-S4. Metabolite abbreviations: α-KG: alpha-ketoglutarate, DHAP: dihydroxyacetone phosphate, FBP: fructose 1,6-bisphosphate, F6P: fructose 6-phosphate, G6P: glucose 6-phosphate, 2/3PG: 2/3-phosphoglycerate, GA3P: glyceraldehyde 3-phosphate, R5P: ribose 5-phosphate, PEP: phosphoenolpyruvate, RU5P/X5P: ribulose 5-phosphate/xylulose 5-phosphate, E4P: erythrose-4-phosphate. C) We used enzyme-constrained genome-scale metabolic models to estimate metabolite values, quantified as the negative change in growth rate (Δy) on metabolite release (Δx) close to optimal growth. D-F) Three different datasets were used to test our hypothesis: D) Paczia et al. (2012), including S. cerevisiae, B. licheniformis, C. glutamicum and E. coli on glucose[12], E) E. coli on three other carbon sources (this work), and F) Vila et al. (2023) covering a smaller number of metabolites on E. coli, P. putida and two environmental strains of the genera Enterobacter and Pseudomonas on up to 8 different carbon sources[42]. The scatter plots show mean estimated rates +- standard errors vs estimated metabolite values. Black-filled circles are datapoints where the lower error bar is omitted because mean minus standard error is less than 0 and cannot be included on a log-scale. Axis labels and colour legend are shared for panel D-F. Annotations in panel F show the Pearson correlation and associated P value in parentheses.*

Figure 2: The importance of different factors in explaining metabolite release rates. *A) Among the tested chemical and metabolic factors, log-scaled metabolite value explains the most variability in log-scaled* E. coli *release rates. The tested factors are shown in order of how much variability (R^2) in log-scaled metabolite value they explain as a univariate linear model, shown in panel B. Compound class abbreviations: CA: Carboxylic acids, AA: Amino acids, KA: Keto/hydroxy acids, OC: Organooxygen compounds, O: Other. Statistical test for compound classes and carbon source: Kruskal-Wallis H-test followed by Conover's test with BH correction. For the other factors the reported P values are from Pearson correlations. B) The best linear model, ranked by BIC score, is a four-factor model that explains 63.4% of the variability in log-scaled release rates. C) Differences in metabolite values between compound classes. Abbreviations and statistical test as in Fig. 2A. D) Log-scaled metabolite value consistently ranks among the most important factors when evaluated separately for each species in each dataset. Factor abbreviations: RBC: Rotatable Bond Count, HBDC: Hydrogen Bond Donor Count, TPSA: Topological Polar Surface Area, HBAC: Hydrogen Bond Acceptor Count.*

Figure 3: Contribution from cell lysis and associated release of intracellular metabolites to extracellular concentrations. *A) Using paired intra- and extracellular metabolite samples from the exponential phase of* E. coli *batch cultures we quantify what fraction of extracellular metabolite concentrations can be explained by cell lysis. Point size reflects order-of-magnitude increase in extracellular concentration from the 2-hour sample to the late-exponential phase sample (OD$_{600}$ ≈ 1.1), and all values except leucine in the L-malate condition were in the µM range. B) We expanded this analysis to all metabolite measurements in our* E. coli *datasets by using literature values for intracellular concentrations (Methods). For each metabolite, we estimated the fraction of cell population that would need to be lysed for the intracellular concentrations released by cell lysis to explain the measured extracellular concentrations. The green region covers the range of measured fractions of lysed cells from A), and the red region is where more than the whole cell population is required to explain extracellular concentrations. Abbreviations: 2PG: 2-phosphoglycerate, 3PG: 3-phosphoglycerate, α-KG: Alpha-ketoglutarate, DHAP: Dihydroxyacetone phosphate, E4P: Erythrose 4-phosphate, FBP: Fructose 1,6-bisphosphate, F6P: Fructose 6-phosphate, G6P: Glucose 6-phosphate, PEP: Phosphoenolpyruvate, R5P: Ribose 5-phosphate, RU5P: Ribulose 5-phosphate, XU5P: Xylulose 5-phosphate.*

Figure 4: Effect of gene knockouts on metabolite release rates and the effect of subsequent evolution to higher fitness. *A)*
Growth curves of selected KO mutants, with sampling time-points indicated by open circles. B) Comparison of extracellular
metabolite concentrations in late exponential (OD_{600}~1.2-1.3) between KO mutants and WT. Significant differences are marked by
solid black outlines (Welch's t-test, FDR < 0.05). C) A PCA plot of exometabolome profiles, showing that each strain has a distinct
pattern. D & E) Batch culture growth curves and sampling timepoints for isolates from the endpoint of the evolution experiment
for ΔaceE and ΔsucB, respectively. F & G) Growth rates and extracellular metabolite concentrations for metabolites near the
deleted reaction, for ΔaceE and ΔsucB, respectively. Statistical significance was assessed using Welch's t-test.

Figure 5: Transporters cause and counteract metabolite release. *A) Approach: We screened the temporal dynamics of extracellular metabolites in batch cultures for 66 different E. coli KO strains using untargeted metabolomics. We quantified the effect of each strain on each metabolite by comparing for each metabolite the ratio between the mean KO-WT difference and the within-strain variability (Methods). B-E) Each panel shows the extracellular concentration of one metabolite (Z-score, median of four technical replicates) for E. coli BW25113 (WT) and 1-4 KO mutants across a batch culture, with the area-under-the-curve of the optical density (AUC of OD) on the x-axis. Small inset panels compare the growth curves of the respective strains (mean OD, three technical replicates). F) Overview of significant, large effects (|Effect| > 3, P < 0.005, permutation test). The left bar chart displays the number of positive (green) and negative (pink) large effects across all 66 KO strains, and the color label indicates the class of the deleted transporter. The right bar chart displays the number of positive (green) and negative (pink) effects on each of the metabolites affected in one or more of the KO strains (122 in total, |Effect| > 3, P < 0.005, permutation test). The color label indicates the classification of all metabolites in E. coli mapping to the corresponding m/z value, which in some cases maps to two different classes. The chord diagram displays which metabolites that are affected (and the strength) in each KO strain. Metabolites / strains displayed in panel B-E annotated / highlighted in bold, and the strongest effects are also annotated on the links in the chord diagram. G) Distribution of all effects across all KO strains and metabolites.*

References

1. Kost, C., Patil, K. R., Friedman, J., Garcia, S. L. & Ralser, M. Metabolic exchanges are ubiquitous in natural microbial communities. *Nat. Microbiol.* 1–9 (2023).

2. Culp, E. J. & Goodman, A. L. Cross-feeding in the gut microbiome: Ecology and mechanisms. *Cell Host Microbe* **31**, 485–499 (2023).

3. Douglas, A. E. The microbial exometabolome: ecological resource and architect of microbial communities. *Philos. Trans. R. Soc. B Biol. Sci.* **375**, 20190250 (2020).

4. D'Souza, G. *et al.* Ecology and evolution of metabolic cross-feeding interactions in bacteria. *Nat. Prod. Rep.* **35**, 455–488 (2018).

5. Solden, L. M. *et al.* Interspecies cross-feeding orchestrates carbon degradation in the rumen ecosystem. *Nat. Microbiol.* **3**, 1274–1284 (2018).

6. Pontrelli, S. *et al.* Metabolic cross-feeding structures the assembly of polysaccharide degrading communities. *Sci. Adv.* **8**, eabk3076 (2022).

7. Gralka, M., Szabo, R., Stocker, R. & Cordero, O. X. Trophic Interactions and the Drivers of Microbial Community Assembly. *Curr. Biol.* **30**, R1176–R1188 (2020).

8. Daniels, M., van Vliet, S. & Ackermann, M. Changes in interactions over ecological time scales influence single-cell growth dynamics in a metabolically coupled marine microbial community. *ISME J.* **17**, 406–416 (2023).

9. Chuang, Y.-C. *et al.* Bacterial adenine cross-feeding stems from a purine salvage bottleneck. *ISME J.* **18**, wrae034 (2024).

10. Mee, M. T., Collins, J. J., Church, G. M. & Wang, H. H. Syntrophic exchange in synthetic microbial communities. *Proc. Natl. Acad. Sci.* **111**, E2149–E2156 (2014).

11.   Campbell, K. *et al.* Self-establishing communities enable cooperative metabolite exchange

in a eukaryote. *eLife* **4**, e09943 (2015).

12.   Paczia, N. *et al.* Extensive exometabolome analysis reveals extended overflow metabolism

in various microorganisms. *Microb. Cell Factories* **11**, 122 (2012).

13.   Yu, J. S. L. *et al.* Microbial communities form rich extracellular metabolomes that foster

metabolic interactions and promote drug tolerance. *Nat. Microbiol.* **7**, 542–555 (2022).

14.   Zampieri, M., Hörl, M., Hotz, F., Müller, N. F. & Sauer, U. Regulatory mechanisms underlying

coordination of amino acid and glucose catabolism in Escherichia coli. *Nat. Commun.* **10**,

3354 (2019).

15.   Gregor, R. *et al.* Vitamin auxotrophies shape microbial community assembly on model

marine particles. *ISME J.* **19**, wraf184 (2025).

16.   Ryback, B., Bortfeld-Miller, M. & Vorholt, J. A. Metabolic adaptation to vitamin auxotrophy

by leaf-associated bacteria. *ISME J.* **16**, 2712–2724 (2022).

17.   Yousif, G. *et al.* Obligate cross-feeding of metabolites is common in soil microbial

communities. 2025.01.29.635426 Preprint at https://doi.org/10.1101/2025.01.29.635426

(2025).

18.   Baldazzi, V., Ropers, D., Gouzé, J.-L., Gedeon, T. & de Jong, H. Resource allocation accounts

for the large variability of rate-yield phenotypes across bacterial strains. *eLife* **12**, e79815

(2023).

19.   Basan, M. *et al.* Overflow metabolism in Escherichia coli results from efficient proteome

allocation. *Nature* **528**, 99–104 (2015).

20.   Huelsmann, M., Schubert, O. T. & Ackermann, M. A framework for understanding collective

microbiome metabolism. *Nat. Microbiol.* **9**, 3097–3109 (2024).

21.   Malina, C., Yu, R., Björkeroth, J., Kerkhoven, E. J. & Nielsen, J. Adaptations in metabolism and

protein translation give rise to the Crabtree effect in yeast. *Proc. Natl. Acad. Sci.* **118**,

e2112836118 (2021).

22.   Pfeiffer, T., Schuster, S. & Bonhoeffer, S. Cooperation and Competition in the Evolution of

ATP-Producing Pathways. *Science* **292**, 504–507 (2001).

23.   Reaves, M. L., Young, B. D., Hosios, A. M., Xu, Y.-F. & Rabinowitz, J. D. Pyrimidine homeostasis

is accomplished by directed overflow metabolism. *Nature* **500**, 237–241 (2013).

24.   Heinrich, R., Schuster, S. & Holzhütter, H.-G. Mathematical analysis of enzymic reaction

systems using optimization principles. *Eur. J. Biochem.* **201**, 1–21 (1991).

25.   Costa, E., Pérez, J. & Kreft, J.-U. Why is metabolic labour divided in nitrification? *Trends*

Microbiol. **14**, 213–219 (2006).

26.   van Hoek, M. J. & Merks, R. M. Redox balance is key to explaining full vs. partial switching to

low-yield metabolism. *BMC Syst. Biol.* **6**, 22 (2012).

27.   Vemuri, G. N., Eiteman, M. A., McEwen, J. E., Olsson, L. & Nielsen, J. Increasing NADH

oxidation reduces overflow metabolism in Saccharomyces cerevisiae. *Proc. Natl. Acad. Sci.*

104, 2402–2407 (2007).

28.   Bar-Even, A., Noor, E., Flamholz, A., Buescher, J. M. & Milo, R. Hydrophobicity and Charge

Shape Cellular Metabolite Concentrations. *PLoS Comput. Biol.* **7**, (2011).

29. Campbell, K., Herrera-Dominguez, L., Correia-Melo, C., Zelezniak, A. & Ralser, M.

Biochemical principles enabling metabolic cooperativity and phenotypic heterogeneity at

the single cell level. *Curr. Opin. Syst. Biol.* **8**, 97–108 (2018).

30. McKinlay, J. B. Are Bacteria Leaky? Mechanisms of Metabolite Externalization in Bacterial

Cross-Feeding. *Annu. Rev. Microbiol.* **77**, 277–297 (2023).

31. Gude, S., Pherribo, G. J. & Taga, M. E. Emergence of Metabolite Provisioning as a By-Product

of Evolved Biological Functions. *mSystems* **5**, 10.1128/msystems.00259-20 (2020).

32. Lopez, J. G. & Wingreen, N. S. Noisy metabolism can promote microbial cross-feeding. *eLife*

11, e70694 (2022).

33. Sánchez, B. J. *et al.* Improving the phenotype predictions of a yeast genome-scale metabolic

model by incorporating enzymatic constraints. *Mol. Syst. Biol.* **13**, 935 (2017).

34. Bekiaris, P. S. & Klamt, S. Automatic construction of metabolic models with enzyme

constraints. *BMC Bioinformatics* **21**, 19 (2020).

35. Mao, Z. *et al.* ECMpy 2.0: A Python package for automated construction and analysis of

enzyme-constrained models. *Synth. Syst. Biotechnol.* **9**, 494–502 (2024).

36. Orth, J. D., Thiele, I. & Palsson, B. Ø. O. What is flux balance analysis? *Nat Biotech* **28**, 245–

248 (2010).

37. Gu, C., Kim, G. B., Kim, W. J., Kim, H. U. & Lee, S. Y. Current status and applications of genome-

scale metabolic models. *Genome Biol.* **20**, 121 (2019).

38. Akashi, H. & Gojobori, T. Metabolic efficiency and amino acid composition in the proteomes

of Escherichia coli and Bacillus subtilis. *Proc. Natl. Acad. Sci.* **99**, 3695–3700 (2002).

39.   Mori, M., Cheng, C., Taylor, B. R., Okano, H. & Hwa, T. Functional decomposition of
metabolism allows a system-level quantification of fluxes and protein allocation towards
specific metabolic functions. *Nat. Commun.* **14**, 4161 (2023).

40.   Pacheco, A. R., Moel, M. & Segrè, D. Costless metabolic secretions as drivers of interspecies
interactions in microbial ecosystems. *Nat. Commun.* **10**, 103 (2019).

41.   Georgi, T., Rittmann, D. & Wendisch, V. F. Lysine and glutamate production by
Corynebacterium glutamicum on glucose, fructose and sucrose: Roles of malic enzyme and
fructose-1,6-bisphosphatase. *Metab. Eng.* **7**, 291–301 (2005).

42.   Vila, J. C. C. *et al.* Metabolic similarity and the predictability of microbial community
assembly. 2023.10.25.564019 Preprint at https://doi.org/10.1101/2023.10.25.564019
(2023).

43.   Wendering, P. & Nikoloski, Z. COMMIT: Consideration of metabolite leakage and community
composition improves microbial community reconstructions. *PLOS Comput. Biol.* **18**,
e1009906 (2022).

44.   Bennett, B. D., Yuan, J., Kimball, E. H. & Rabinowitz, J. D. Absolute quantitation of
intracellular metabolite concentrations by an isotope ratio-based approach. *Nat. Protoc.* **3**,
1299–1311 (2008).

45.   Park, J. O. *et al.* Metabolite concentrations, fluxes and free energies imply efficient enzyme
usage. *Nat. Chem. Biol.* **12**, 482–489 (2016).

46.   Sajed, T. *et al.* ECMDB 2.0: A richer resource for understanding the biochemistry of E. coli.
Nucleic Acids Res. **44**, D495–D501 (2016).

47. Thorfinnsdottir, L. B., García-Calvo, L., Bø, G. H., Bruheim, P. & Røst, L. M. Optimized Fast

Filtration-Based Sampling and Extraction Enables Precise and Absolute Quantification of the

Escherichia coli Central Carbon Metabolome. *Metabolites* **13**, 150 (2023).

48. Falcón-Cano, G., Molina, C. & Cabrera-Pérez, M. Á. Reliable Prediction of Caco-2 Permeability

by Supervised Recursive Machine Learning Approaches. *Pharmaceutics* **14**, 1998 (2022).

49. Liu, X., Testa, B. & Fahr, A. Lipophilicity and Its Relationship with Passive Drug Permeation.

Pharm. Res. **28**, 962–977 (2011).

50. Pontrelli, S. *et al.* Degradation of extracellular polymeric substances shapes microbial

community diversity. *PLOS Biol.* **23**, e3003287 (2025).

51. Gibson, S. E. R. *et al.* Bacteria encode post-mortem protein catabolism that enables altruistic

nutrient recycling. *Nat. Commun.* **16**, 1400 (2025).

52. Fuhrer, T., Zampieri, M., Sévin, D. C., Sauer, U. & Zamboni, N. Genomewide landscape of

gene–metabolome associations in Escherichia coli. *Mol. Syst. Biol.* **13**, 907 (2017).

53. Baba, T. *et al.* Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants:

the Keio collection. *Mol. Syst. Biol.* **2**, 2006.0008 (2006).

54. Fuhrer, T., Heer, D., Begemann, B. & Zamboni, N. High-throughput, accurate mass

metabolome profiling of cellular extracts by flow injection-time-of-flight mass spectrometry.

Anal. Chem. **83**, 7074–7080 (2011).

55. Rancourt, D. E., Stephenson, J. T., Vickell, G. A. & Wood, J. M. Proline excretion by Escherichia

coli K12. *Biotechnol. Bioeng.* **26**, 74–80 (1984).

56. Karp, P. D. *et al.* The EcoCyc database (2025). *EcoSal Plus* **0**, eesp-0019-2024 (2025).

57.   Park, J. H., Lee, K. H., Kim, T. Y. & Lee, S. Y. Metabolic engineering of Escherichia coli for the

560   production of l-valine based on transcriptome analysis and in silico gene knockout

561   simulation. *Proc. Natl. Acad. Sci.* **104**, 7797–7802 (2007).

562   58.   Zakataeva, N. P., Aleshin, V. V., Tokmakova, I. L., Troshin, P. V. & Livshits, V. A. The novel

563   transmembrane Escherichia coli proteins involved in the amino acid efflux. *FEBS Lett.* **452**,

564   228–232 (1999).

565   59.   Burton, K. Adenine transport in Escherichia coli. *Proc. R. Soc. Lond. B Biol. Sci.* **255**, 153–157

566   (1997).

567   60.   Papakostas,   K.,   Botou,   M.   &   Frillingos,   S.   Functional   Identification   of   the

568   Hypoxanthine/Guanine Transporters YjcD and YgfQ and the Adenine Transporters PurP and

569   YicO of Escherichia coli K-12. *J. Biol. Chem.* **288**, 36827–36840 (2013).

570   61.   Rodionova, I. A. *et al.* Identification of a transcription factor, PunR, that regulates the purine

571   and purine nucleoside transporter punC in E. coli. *Commun. Biol.* **4**, 991 (2021).

572   62.   Mori, M. *et al.* From coarse to fine: the absolute Escherichia coli proteome under diverse

573   growth conditions. *Mol. Syst. Biol.* **17**, e9536 (2021).

574   63.   Smith, D. R. & Chapman, M. R. Economical Evolution: Microbes Reduce the Synthetic Cost

575   of Extracellular Proteins. *mBio* **1**, e00131-10 (2010).

576   64.   Flamholz, A., Noor, E., Bar-Even, A., Liebermeister, W. & Milo, R. Glycolytic strategy as a

577   tradeoff between energy yield and protein cost. *Proc. Natl. Acad. Sci.* **110**, 10039–10044

578   (2013).

579   65.   Jacob, F. & Monod, J. Genetic regulatory mechanisms in the synthesis of proteins. *J. Mol.*

580   *Biol.* **3**, 318–356 (1961).

66.   Giuliani, S. E. *et al.* Environment sensing and response mediated by ABC transporters. *BMC*

Genomics **12**, S8 (2011).

67.   Martínez-Antonio, A., Janga, S. C., Salgado, H. & Collado-Vides, J. Internal-sensing machinery

directs the activity of the regulatory network in Escherichia coli. *Trends Microbiol.* **14**, 22–

27 (2006).

68.   Nies, D. H., Schleuder, G., Galea, D. & Herzberg, M. A flow equilibrium of zinc in cells of

Cupriavidus metallidurans. *J. Bacteriol.* **206**, e00080-24 (2024).

69.   Fritts, R. K., McCully, A. L. & McKinlay, J. B. Extracellular Metabolism Sets the Table for

Microbial Cross-Feeding. *Microbiol. Mol. Biol. Rev.* **85**, 10.1128/mmbr.00135-20 (2021).

70.   Lopez, J. A., Liu, B., Li, Z., Donia, M. S. & Wingreen, N. S. When is microbial cross-feeding

evolutionarily   stable?   2025.05.16.654511   Preprint   at

https://doi.org/10.1101/2025.05.16.654511 (2025).

71.   Preussger, D., Giri, S., Muhsal, L. K., Oña, L. & Kost, C. Reciprocal Fitness Feedbacks Promote

the Evolution of Mutualistic Cooperation. *Curr. Biol.* **30**, 3580-3590.e7 (2020).

72.   Zomorrodi, A. R. & Segrè, D. Genome-driven evolutionary game theory helps understand the

rise of metabolic interdependencies in microbial communities. *Nat. Commun.* **8**, 1563

(2017).

73.   Datsenko, K. A. & Wanner, B. L. One-step inactivation of chromosomal genes in Escherichia

coli K-12 using PCR products. *Proc. Natl. Acad. Sci.* **97**, 6640–6645 (2000).

74.   Lewis, N. E. *et al.* Omic data from evolved E. coli are consistent with computed optimal

growth from genome-scale models. *Mol. Syst. Biol.* **6**, 390 (2010).

75. Campello, R. J. G. B., Moulavi, D. & Sander, J. Density-Based Clustering Based on Hierarchical

Density Estimates. in *Advances in Knowledge Discovery and Data Mining* (eds Pei, J., Tseng,

V. S., Cao, L., Motoda, H. & Xu, G.) 160–172 (Springer Berlin Heidelberg, Berlin, Heidelberg,

2013).

76. Gallart-Ayala, H. *et al.* A global HILIC-MS approach to measure polar human cerebrospinal

fluid metabolome: Exploring gender-associated variation in a cohort of elderly cognitively

healthy subjects. *Anal. Chim. Acta* **1037**, 327–337 (2018).

77. van der Velpen, V. *et al.* Systemic and central nervous system metabolic alterations in

Alzheimer's disease. *Alzheimers Res. Ther.* **11**, 93 (2019).

78. Myers, J. A., Curtis, B. S. & Curtis, W. R. Improving accuracy of cell and chromophore

concentration measurements using optical density. *BMC Biophys.* **6**, 4 (2013).

79. Ren, Q., Henes, B., Fairhead, M. & Thöny-Meyer, L. High level production of tyrosinase in

recombinant Escherichia coli. *BMC Biotechnol.* **13**, 18 (2013).

80. Goelzer, A. *et al.* Quantitative prediction of genome-wide resource allocation in bacteria.

Metab. Eng. **32**, 232–243 (2015).

81. Zhang, Y. *et al.* A new genome-scale metabolic model of Corynebacterium glutamicum and

its application. *Biotechnol. Biofuels* **10**, 169 (2017).

82. McGill, S. L. *et al.* Pseudomonas aeruginosa reverse diauxie is a multidimensional, optimized,

resource utilization strategy. *Sci. Rep.* **11**, 1457 (2021).

83. Kuepper, J. *et al.* Metabolic Engineering of Pseudomonas putida KT2440 to Produce

Anthranilate from Glucose. *Front. Microbiol.* **6**, 1310 (2015).

84.   Niu, J. *et al.* Construction and Analysis of an Enzyme-Constrained Metabolic Model of

Corynebacterium glutamicum. *Biomolecules* **12**, 1499 (2022).

85.   Wu, K. *et al.* ecBSU1: A Genome-Scale Enzyme-Constrained Model of Bacillus subtilis Based

on the ECMpy Workflow. *Microorganisms* **11**, 178 (2023).

86.   Guo, J., Zhang, H., Wang, C., Chang, J.-W. & Chen, L.-L. Construction and analysis of a

genome-scale metabolic network for *Bacillus licheniformis* WX-02. *Res. Microbiol.* **167**, 282–

289 (2016).

87.   Nogales, J. *et al.* High-quality genome-scale metabolic modelling of Pseudomonas putida

highlights its broad metabolic capabilities. *Environ. Microbiol.* **22**, 255–269 (2020).

88.   Ebrahim, A., Lerman, J. A., Palsson, B. O. & Hyduke, D. R. COBRApy: COnstraints-Based

Reconstruction and Analysis for Python. *BMC Syst. Biol.* **7**, 74 (2013).

89.   Varma, A., Boesch, B. W. & Palsson, B. O. Stoichiometric interpretation of Escherichia coli

glucose catabolism under various oxygenation rates. *Appl. Environ. Microbiol.* **59**, 2465–

2473 (1993).

90.   Reznik, E., Mehta, P. & Segrè, D. Flux Imbalance Analysis and the Sensitivity of Cellular

Growth to Changes in Metabolite Pools. *PLOS Comput. Biol.* **9**, e1003195 (2013).

91.   Ram, Y. *et al.* Predicting microbial growth in a mixed culture from growth curve data. *Proc.*

Natl. Acad. Sci. **116**, 14698–14707 (2019).

92.   Reed, J. L., Vo, T. D., Schilling, C. H. & Palsson, B. O. An expanded genome-scale model of

Escherichia coli K-12 (iJR904 GSM/GPR). *Genome Biol.* **4**, R54 (2003).

93.   Feist, A. M. *et al.* A genome-scale metabolic reconstruction for Escherichia coli K-12 MG1655

that accounts for 1260 ORFs and thermodynamic information. *Mol. Syst. Biol.* **3**, 121 (2007).

94. Orth, J. D. *et al.* A comprehensive genome-scale reconstruction of Escherichia coli

metabolism—2011. *Mol. Syst. Biol.* **7**, 535 (2011).

95. Monk, J. M. *et al.* iML1515, a knowledgebase that computes Escherichia coli traits. *Nat.*

Biotechnol. **35**, 904–908 (2017).

96. Djoumbou Feunang, Y. *et al.* ClassyFire: automated chemical classification with a

comprehensive, computable taxonomy. *J. Cheminformatics* **8**, 61 (2016).

97. Seabold, S. & Perktold, J. Statsmodels: Econometric and Statistical Modeling with Python. in

92–96 (Austin, Texas, 2010). doi:10.25080/Majora-92bf1922-011.

98. Lukačišinová, M., Fernando, B. & Bollenbach, T. Highly parallel lab evolution reveals that

epistasis can curb the evolution of antibiotic resistance. *Nat. Commun.* **11**, 3105 (2020).

99. Chen, S. Ultrafast one-pass FASTQ data preprocessing, quality control, and deduplication

using fastp. *iMeta* **2**, e107 (2023).

100. Simon Andrews. FastQC: A Quality Control Tool for High Throughput Sequence Data.

Babraham Institute (2010).

101. Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–

3100 (2018).

102. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *GigaScience* **10**, giab008 (2021).

103. Cingolani, P. *et al.* A program for annotating and predicting the effects of single nucleotide

polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-

2; iso-3. *Fly (Austin)* **6**, 80–92 (2012).

104. Garrison, E. & Marth, G. Haplotype-based variant detection from short-read sequencing.

Preprint at https://doi.org/10.48550/arXiv.1207.3907 (2012).

105. Burz, S. D. *et al.* A Guide for Ex Vivo Handling and Storage of Stool Samples Intended for

Fecal Microbiota Transplantation. *Sci. Rep.* **9**, 8897 (2019).

106. Bennett, B. D. *et al.* Absolute metabolite concentrations and implied enzyme active site

occupancy in *Escherichia coli*. *Nat. Chem. Biol.* **5**, 593–599 (2009).

107. Mori, M., Schink, S., Erickson, D. W., Gerland, U. & Hwa, T. Quantifying the benefit of a

proteome reserve in fluctuating environments. *Nat. Commun.* **8**, 1–8 (2017).

108. Volkmer, B. & Heinemann, M. Condition-Dependent Cell Volume and Concentration of

Escherichia coli to Facilitate Data Conversion for Systems Biology Modeling. *PLoS ONE* **6**,

e23126 (2011).

109. Simensen, V. *et al.* Experimental determination of Escherichia coli biomass composition for

constraint-based metabolic modeling. *PLOS ONE* **17**, e0262450 (2022).

110. Saier, M. H. *et al.* The Transporter Classification Database (TCDB): 2021 update. *Nucleic Acids*

Res. **49**, D461–D467 (2020).

111. Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat.*

Methods **17**, 261–272 (2020).

112. Pignon, E., Holló, G., Steiner, T., van Vliet, S. & Schaerli, Y. Uptake and leakage rates

differentially shape community arrangement and composition of microbial consortia. *ISME*

J. **19**, wraf122 (2025).

113. Pignon, E. Engineering metabolic interactions to control spatial patterns and composition in

auxotrophic microbial communities. https://iris.unil.ch/handle/iris/264404 (2025).

Acknowledgements

We thank all members of the Mitri lab at the University of Lausanne for discussions and feedback. We also thank people at the Department of Biotechnology and Nanomedicine at SINTEF for technical support and advice on the experimental design, and Vassily Hatzimanikatis and the Laboratory of Computational Systems Biotechnology at EPFL for feedback on the computational analyses. Furthermore, we would like to thank Simon van Vliet, Olga Schubert, Guilhem Panneau, Jean C. C. Vila, Milton Saier and Uwe Sauer for providing feedback, Olga Schubert and Kim Schlegel for providing the KEIO strains, Sebastian Burz, You Zheng Teo and Clément Vulin for support with the live/dead staining protocol, Estelle Pignon for providing *E. coli* auxotrophs and Vladimir Sentchilo for support with the flow cytometry and sampling for metabolomics analyses. We also thank the Metabolomics unit at UNIL, and specifically Hector G. Ayala and Julijana Ivanisevic, not only for conducting the LC-MS analyses, but also for providing guidance and recommendations on related matters. Special thanks to Florent Mazel and Daniel Segrè for feedback on the draft manuscript. Finally, we thank Lilja B. Thorfinnsdottir, Jean C. C. Vila and Stephan Noack for sharing of experimental data.

Funding

Swiss National Science Foundation Swiss Postdoctoral Fellowship TMPFP3_217172 (SS)

National Center of Competence in Research Microbiomes grant SNF 51NF40_180575 (SM, SS, AG, MAV, JL, PP)

Faculty of Biology and Medicine, University of Lausanne (EU)

Swiss National Science Foundation Eccellenza grant PCEGP3_181272 (SM, MAV)

SINTEF Industry (GB)

Research Council of Norway, SFI Industrial Biotechnology, grant no 309558 (GB)

Author contributions

Conceptualization: SS, SM; Methodology: SS, PD, AQ, JSL, GB, MAV, EU; Investigation: SS, GB, PD, AG, EU, PP; Visualization: SS, PP, EU; Funding acquisition: SS, SM, PE; Project administration: SS, SM; Supervision: SS, SM; Writing – original draft: SS; Writing – review & editing: All authors

Competing interests

720 Authors declare that they have no competing interests.

Data and materials availability

723 Strains and isolates used in this study are available on request. Formatted data and code for data 724 analyses and visualization are available on GitHub at https://github.com/Mitri-lab/metabolite-725 release. A permanent archive of this repository will be deposited to Zenodo upon publication. 726 The raw sequencing data and metadata have been deposited to the NCBI SRA database under the 727 BioProject ID PRJNA1270783. The raw FIA-TOF MS data is available on MassIVE under the dataset 728 identifier MSV000097105.

Extended Data and Supplementary Information

730 Supplementary Figures S1-S32
731 Supplementary Tables S1-S7

Methods

Strains

735 All experiments were conducted with either *E. coli* K-12 MG1655, *E. coli* K-12 BW25113 (wild-type 736 ancestor for KEIO KO strains) or one of the 75 KO mutants selected from the KEIO collection[53] 737 (Table S6). The KEIO KO strains were verified by PCR with one primer binding inside the Kan^R 738 cassette inserted during gene knock-out (Keio_K1_Rev for genes on the (-) strand; 739 Keio_KT_RvComp_Fw for genes on the (+) strand) and one gene-specific primer binding 740 downstream of the knocked-out gene (primers are listed in Table S6)[53,73]. For all experiments, the 741 KEIO KO strains were used with the kanamycin resistance cassette in place.

Precultures and growth media

For all experiments detailed below, the strains were precultured as described here unless otherwise stated: From glycerol stocks stored at -70 °C each strain was streaked out onto LB agar plates and incubated overnight (37 °C). Then, one colony of each strain was picked and used to inoculate liquid precultures (10 mL LB in 50 mL Erlenmeyer flasks) that were incubated overnight (37 °C, 200 rpm). The kanamycin-resistant KEIO KO strains and strains evolved from these were always precultured in liquid LB and streaked out onto LB agar plates with 25 µg/mL kanamycin, while *E. coli* K-12 MG1655 and *E. coli* K12 BW25113 were always precultured in LB and streaked onto LB agar plates without selection. Unless otherwise noted, precultures were washed three times by centrifugation, with the supernatant removed after each step, and the cells resuspended in M9 medium with no carbon source to eliminate residual LB before inoculation. Unless otherwise stated, M9 medium was always prepared by mixing (per litre): 200 mL 5X M9 salts (M6030, Sigma-Aldrich, St. Louis, MO, USA), 2 mL 1M $MgSO_4$, 0.1 mL 1M $CaCl_2$, 1 mL 0.5 g/L $FeSO_4 \cdot 7H_2O$, MQ water and appropriate amounts of a carbon source stock solution. The final M9 medium was then pH-adjusted to pH = 7.0, filter sterilized (0.22 µm) and stored at 4°C until use.

Selecting carbon sources for *E. coli* K-12 MG1655 batch cultures

To complement the existing exometabolome data from batch cultures of *E. coli* on glucose[12], we aimed to select three carbon sources that differed in chemical class and caused variation in metabolic flux patterns. We used the ecGEM eciJO1366 of *E. coli* K-12 MG1655 without any modifications and parsimonious FBA (pFBA) to simulate optimal flux patterns across 104 different growth supporting metabolites[34,74]. We then performed a PCA analysis of the predicted fluxes (as Boolean values) to compare flux patterns between the carbon sources (Fig. S5). We used HDBSCAN[75] to identify clusters and selected from the different clusters in total 10 compounds that also differed in chemical class (sugars, amino acids and organic acids) and their entry point into central carbon metabolism (Fig. S5). We then screened the growth of the selected carbon sources in triplicates over 48 hours in 96-well plates (Fig. S5). For this experiment, we prepared M9 medium with each of the 10 different carbon sources to a concentration scaled to 120 mM carbon atoms. From washed LB precultures, *E. coli* K-12 MG16155 was inoculated to OD_{600} = 0.05

in 200 µL of media and cultivated with continuous shaking (double orbital, 425 cpm) in a BioTek

Synergy H1 (Agilent Technologies, Winooski, VT, USA) plate reader at 37 °C for 48 hours. We

ultimately chose galactose, L-alanine and L-malate, as they supported rapid growth, differed in

chemical class, entered central metabolism at different locations and were grouped into different

clusters based on flux patterns (Fig. S5).

E. coli K-12 MG1655 batch cultures in bioreactors with different carbon sources

E. coli K-12 MG1655 was cultivated in lab-scale bioreactors using three select carbon sources.

Their concentrations were normalized by the number of carbon atoms to a carbon atom

concentration of 120 mM, corresponding to 20 mM galactose (G0625, Sigma-Aldrich), 30 mM L-

malic acid (02290, Sigma-Aldrich) and 40 mM L-alanine (A7627, Sigma-Aldrich). M9 medium with

these three carbon sources were prepared as described above using 25X carbon source stock

solutions.

The strain was streaked from a frozen glycerol vial onto an LB agar plate and incubated at 37 °C

overnight. One colony was transferred to a 250 mL baffled shake flask containing 40 mL LB

medium. After 7 hours, a culture volume corresponding to a start OD_{600} = 0.05 was inoculated to

a 500 mL baffled shake flask containing 100 mL M9 + 40 mM L-alanine. This flask was incubated

for 36 h (200 rpm, 37 °C). For the two other carbon sources, the strain was incubated in LB for 17

h before inoculating a culture volume corresponding to OD_{600} = 0.05 in a 500 mL baffled shake

flask containing 100 mL M9 + 20 mM L-galactose or M9 + 30 mM L-malic acid, respectively. The

flasks were incubated for 24 h (200 rpm, 37 °C). Before inoculation to bioreactors, M9 precultures

were centrifuged (3220 rcf, 10 min, 4 °C), supernatant removed, and pellet resuspended in M9

medium without carbon.

Batch cultivations were performed in 1 L DASGIP bioreactors (Eppendorf DASGIP, Jülich, Germany)

with an initial volume of 600 mL M9 medium supplemented with either 20 mM galactose, 30 mM

L-malic acid or 40 mM L-alanine, three replicates per condition. The reactors were equipped with

two Rushton impellers and probes for measurements of dissolved oxygen (DO) and pH.

Submerged aeration was maintained at 0.5 vvm, and agitation was cascaded between 400 and

800, controlled by a DO set point of 30 %. pH set point was 7.0 and controlled using 2 M NaOH

and 2 M HCl. Start OD_{600} was 0.05.

Sampling was performed regularly during the exponential growth phase with additional samples

collected during stationary phase. OD_{600} was measured at all time points and supernatant stored:

2 mL culture was centrifuged (3220 rcf, 10 min, 4 °C), and supernatant frozen in -20 °C. At the end

of exponential phase and at the end of the experiment, samples were collected for cell dry weight

(CDW). 50 mL culture was centrifuged (3220 rcf, 10 min, 4 °C), supernatant removed, cell pellet

resuspended in 50 mL MQ water, centrifuged, supernatant removed, cell pellet resuspended in

5-10 mL MQ water and transferred to a pre-weighed aluminium beaker and dried at 105 °C for 24

hours. Beakers were then weighed again and CDW calculated.

Exometabolome analyses of time-series samples from *E. coli* K-12 MG1655 batch cultures

with different carbon sources

Four samples from each of the nine batch cultures (three replicates of three different media) and

media samples were analysed with LC-MS and GC-MS to measure the extracellular concentration

of metabolites at different time points throughout the experiment. The time points were selected

based on time-series cultivation data to cover the exponential growth phase (three samples) and

one stationary phase sample (Fig. S6). Samples from the different replicates were aligned based

on the timing of the end of the exponential growth phase (Fig. S6).

LC-MS of selected bioreactor samples was conducted by the Metabolomics facility at UNIL using

their method "high-coverage targeted analysis of polar metabolites", following this procedure:

Media samples were thawed on ice. An aliquot (20 µL) was extracted with the addition of ice-cold

methanol (80 µL), vortexed (30 sec) and centrifuged (15000 rpm, 15 min, 4°C). The supernatants

were transferred to vials for liquid chromatography – tandem mass spectrometry (LC-MS/MS)

analyses. The extracts were analyzed by hydrophilic interaction chromatography coupled to

tandem mass spectrometer (6496 iFunnel, Agilent Technologies) using multiple reaction

monitoring - MRM approach in both, positive and negative ionization modes, to maximize the
polar metabolome coverage. The analytical conditions have been described in detail
elsewhere[76,77]. Raw LC-MS/MS metabolome data were processed using the Mass Hunter
Quantitative analysis software (Agilent Technologies). The peak areas (or extracted ion
chromatograms (EICs) for the monitored MRM transitions) were used for relative comparison of
metabolite levels between different conditions or groups of samples. Data quality assessment,
including signal drift correction, was performed using pooled quality control (QC) samples
analyzed periodically throughout the entire batch.

Metabolite quantification was performed using stable isotope labeled internal standards and
calibration curves following the signal drift correction. Data processing was done using
MassHunter Quantitative analysis. Peak area integration was manually curated, and
concentrations were reported by selecting a 6-point portion of the linear standard curves relevant
for the observed concentrations.

The relative LC-MS data was standardized before subsequent analyses and data processing. Both
the relative and absolute LC-MS data were cleaned of obvious outliers. For the absolute data the
outliers were caused by carry-over from previous samples leading to too high malate (six samples)
and succinate (one sample) values. For the relative LC-MS data there were additional outliers of
unknown reasons (39 values in total) related to in total 14 metabolites. For several metabolites,
there were more missing values (not detected/not quantified) in the absolute data than in the
relative data. For a few metabolites (phenylalanine, proline, creatine), we leveraged good linear
relationships between relative and absolute data (R^2>0.8) to estimate the missing absolute values
using a simple linear transformation.

GC-MS analysis of the same samples was conducted in the following way: Analytical standards
were prepared as M9 medium with increasing concentrations of acetate, lactate, pyruvate,
formate and propionate and stored at -70°C until use. Samples from the *E. coli* bioreactor batch
cultures and analytical standards were thawed on iced. Then, 100 µL was transferred to an

Eppendorf tube and kept on a cold block (-20 °C) while adding 5 μL 11% HCl and 500 μL diethyl

ether. Tubes were capped, vortexed on a thermoblock (2000 rpm, 10 min, 1°C) and centrifuged

(13 000 rcf, 5 min, 4°C). The top organic phase layer was pipetted to a glass vial and derivatized

with 20 μL N-(t-butyldimethylsilyl)-N-methyltrifluoroacetamide (MTBSTFA) and vortexed briefly.

Samples were then placed on a heat block (90 min, 35°C) and kept at 16°C until analysis. The

samples were injected (1 μL) by a Pal3 autosampler onto an Agilent 8890-5977B GC-MSD (Agilent

Technologies) with a VF-5MS (30 m x 0.25 mm x 0.25 mm) column. The samples were injected

with a split ratio of 15:1, helium flow rate of 1 mL/min and inlet temperature of 230 °C. The

temperature was held for 2 min at 50 °C, raised at 25 °C/min to 175 °C, 30 °C/min to 280 °C and

held for 3.5 min. The MSD was run in scan mode from 40-500 Da. Analyte abundances were

calculated using the MassHunter Quantitative Analysis software (Agilent Technologies). Absolute

concentrations were calculated using standard curves.

Galactose concentrations were measured using the L-Arabinose/D-Galactose assay kit (K-ARGA,

Megazyme, Bray, Ireland), following their rapid protocol for analyses in 96-well plates.

Estimation of metabolite release rates of *E. coli*, *B. licheniformis*, *C. glutamicum* and *S.*

cerevisiae in glucose medium

Time-series data on biomass (in OD_{600}) and absolute extracellular metabolite concentrations from

batch cultivations of *E. coli, B. licheniformis, C. glutamicum* and *S. cerevisiae* in high glucose

concentrations (10-20 g/L)[12] was kindly provided by the authors (mean and standard deviations).

For metabolites reported as a sum of two metabolites, we attributed equal amounts to those

metabolites to enable direct comparison with metabolite values estimated for individual

metabolites (described below). This includes 2-phosphoglycerate/3-phosphoglycerate and

ribulose 5-phosphate/xylulose 5-phosphate. To estimate the metabolite release and uptake rates

of each species, we first converted the OD_{600} values to gDW/L. For *E. coli* we used the conversion

factor we estimated from our own cell dry weight measurements in M9 galactose medium (0.346

gDW/L/OD_{600}), which is comparable, but on the lower end of values found in the literature (0.36-

0.515 gDW/L/OD_{600})[78,79]. For the three other species, we used literature values: B. licheniformis

(data from B. subtilis) 0.48 gDW/L/OD$_{600}$[80]; *C. glutamicum* 0.27 gDW/L/OD$_{600}$[81]; *S. cerevisiae*

(mean of literature values) 0.746 gDW/L/OD$_{600}$[78]. We then integrated the translated biomass

time-series data to obtain biomass AUC data that we leveraged together with the mean absolute

exometabolome data to estimate the uptake or release rates for each measured metabolite for

each species using linear regression (Figs. S1-S4). Rates were only estimated for the exponential

growth phase, and only until each metabolite either saturated or was clearly being re-consumed.

Theoretical justification of estimating uptake or release rates from extracellular concentrations

We start out from a simple assumption that the change in extracellular concentration C_i of

metabolite *i* can be explained by the net release or uptake $a_i(t)$ (by any mechanism) by the

microbes with biomass density $X(t)$ in the batch culture:

899
$$\frac{dC_i}{dt} = X(t) \cdot a_i(t)$$

900 The concentration $C_i(t)$ at a given timepoint *t* is thus:

901
$$C_i(t) = \int_{t0}^{t} X(t) \cdot a_i(t)dt + C_i(t_0)$$

902 If we assume that the uptake or release rate $a_i(t)$ is constant, the concentration $C_i(t)$ can be

903 described by a linear function of the constant release rate and the integrated biomass (area under

904 the curve, AUC):

905
$$C_i(t) = a_i \int_{t0}^{t} X(t)dt + C_i(t_0) = a_i \cdot X_{AUC}(t) + C_0$$

Hence, when metabolites are released at constant rates the dynamics should be well explained

by a linear regression of extracellular metabolite concentrations on biomass AUC, and the specific

net rate (uptake or release) is the slope of the curve.

Estimation of metabolite release rates of *E. coli* in galactose, L-malate and L-alanine

For our bioreactor batch cultivations of *E. coli* in galactose, L-malate and L-alanine, we used paired

cell dry weight measurements and OD$_{600}$ readings to calculate conversion factors used to translate

OD$_{600}$ values for all time points (galactose: 0.346 ± 0.016; L-malate: 0.279 ± 0.017; L-alanine:

0.296 ± 0.011; mean $\pm$ std in gDW/L/OD$_{600}$). The translated time-series data was used to calculate

the biomass area under the curve used to estimate metabolite release and uptake rates further

described below.

We estimated the metabolite uptake or release rate by performing for each metabolite and each

carbon source a linear regression of measured concentration (relative or absolute) vs biomass

AUC using the three samples (timepoints T1-T3) acquired from the three replicate batch cultures

during exponential phase plus the initial M9 medium (T0; Figs. S7-S9). To avoid underestimating

release rates for metabolites that potentially were being re-consumed at the end of the

exponential phase, we discarded the last timepoint from the exponential phase in cases where

the rate was positive (indicating metabolite release) but there was a significant reduction in

extracellular concentration from T2-T3 ($P < 0.05$, one-sided t-test). For 17 (of 126) metabolites,

we chose to not include T0 for at least one of the three conditions in the linear regression, either

because of very large variability of T0 or because the trend from T0 to T1 was clearly different

from the general trend from T1-T3. We only estimated the rate for metabolites where we had at

least three data points from at least two different time points. For the metabolites where we had

both relative and absolute data, we used the absolute data if there were enough absolute data

points to get adequate slope estimates. For seven metabolites (glutamine, alpha-aminoadiapate,

serine, lysine, lactate, NAD and isocitrate) where both absolute and relative quantification was

performed, we could estimate the rate more accurately on the relative data set (because of more

missing data points in the absolute data set). However, there were sufficient absolute

measurements to estimate the absolute spread (standard deviation) of these data points,

enabling translation of the slope value from Z-score to μmol/gDW/h.

Estimation of metabolite release rates of *E. coli*, *P. putida,* an *Enterobacter* sp. and a

Pseudomonas sp. in various carbon source environments

The dataset from Vila et al. (2023) was kindly provided by the authors[42]. The dataset contains

exometabolome data from three timepoints (16, 28, and 48 hours) of *E. coli* MG1655, *P. putida*

KT2440, and two environmental isolates of the genera *Enterobacter* and *Pseudomonas,*

respectively[42]. All species were cultivated in M9 medium with one of five carbon sources (D-

glucose, D-fructose, glycerol, pyruvate or L-malate). Additionally, the dataset includes
exometabolome data from two timepoints (28 and 48 hours) of *E. coli* and the *Enterobacter* sp.
on D-ribose, L-arabinose and D-galactose, and of *P. putida* and the *Pseudomonas* sp. on acetate,
fumarate and succinate. To estimate metabolite release rates, we first used the plate reader
growth curves to calculate biomass AUC after 16, 28 and 48 hours and to identify the end of
exponential growth of each organism in each condition. The end of exponential phase was
automatically identified as the first peak in the second derivative of the smoothed growth curve
with an OD_{600} above 60% of maximum. We converted the AUC from OD_{600} to gDW/L using the
same conversion factor for all the different carbon source environments: 0.36 gDW/L/ OD_{600} for
E. coli and the *Enterobacter* sp., mean of measured values and literature values[78,79]; 0.476 gDW/L/
OD_{600} for *P. putida* and the *Pseudomonas* sp, mean of literature values[82,83]. Finally, we performed
a linear regression between extracellular metabolite concentrations and AUC on both duplicates
of each strain/carbon source combination including all datapoints before the end of exponential
growth + 2 hours (a buffer to account for some inaccuracy in the automatic detection). We only
included rate estimates for metabolites that were quantified at least twice for each strain. As the
dataset did not include an early timepoint or media measurement, we assumed the initial
concentration of each metabolite to be 0. If a metabolite was detected more at more than one
timepoint, we discarded the estimated rate if there was a qualitative discrepancy in slope value
when the slope was estimated with or without the T0 concentration assumed to be 0. If the
carbon source was among the measured metabolites, this metabolite was excluded from rate
estimation in those specific conditions.

Genome-scale metabolic models

For *E. coli, S. cerevisiae and C. glutamicum* we used the previously developed enzyme-constrained
genome-scaled metabolic models (GEMs) eciJO1366[34], ecCGL1[84] and ecYeastGEM 8.3.4[33],
respectively. For *B. licheniformis* there was no available enzyme-constrained GEM, and we
therefore opted to use the enzyme-constrained GEM ecBSU1 of the closely related species *B.*
subtilis[85]*,* which was available at github.com/tibbdc/ecBSU1. A normal GEM for *B. licheniformis* is
available[86], but with this model we could not achieve any feasible flux balance solution. To

simulate the environmental isolate of the genus *Enterobacter* we used the *E. coli* GEM eciJO1366.

For the environmental isolate of the genus *Pseudomonas* and for *P. putida* we used the enzyme-

constrained version of the *P. putida* GEM iJN1463[87] automatically generated by the ECMpy2

method[35], which we here refer to as eciJN1463.

We made a few minor changes to the published GEMs before predicting metabolite values. In the

E. coli GEM eciJO1366 the proteome weight of the forward reaction *GALKr* (galactokinase) was

extremely high (0.016), limiting the growth rate on galactose to much less than the growth rate

we measured in M9 galactose medium. To enable eciJO1366 to reach the measured growth rate

with galactose as the only carbon source, we replaced the proteome weight of the forward *GALKr*

reaction with the proteome weight of the reverse *GALKr* reaction which was much smaller

(0.00037). For the ecYeastGEM we only merged exchange reactions that were previously split to

simplify its use. Both ecBSU1 and eciJN1463 were translated from json to sMOMENT format in

xml[34], modified by merging split exchange reactions and fixing metabolite compartment names

and annotations. For ecBSU1 we also removed the enzyme constraint on the reaction

"PPAm_num1" (inorganic diphosphatase) as it caused unrealistic uptake and release rates of

phosphate and diphosphate, respectively. The *C. glutamicum* GEM ecCGL1 was also converted

from the ECMpy format[35] to the sMOMENT format[34] but otherwise kept unchanged. All GEMs

were used with their default biomass equations as objective when maximizing growth in all FBA

or pFBA analyses. Reframed 1.5.3 (github.com/cdanielmachado/reframed) or COBRApy 0.29.1[88]

were used to load, manipulate and run constraint-based analyses with Gurobi 10 (Gurobi

Optimization, LLC) as the solver.

Estimating metabolite value and turnover using genome-scale metabolic models

In constraint-based modelling a shadow price is an estimate of how sensitive the objective

function in a flux balance analysis (FBA) is to increased influx or efflux of a metabolite[89,90]. Hence,

one can interpret the shadow price as *metabolite value* as it quantifies the impact on growth

caused by the release of a metabolite (to have positive values we here define metabolite value

as the shadow price multiplied by -1). A similar approach to measure the cost of metabolite

1002 release has been used to identify costless secretions[40]. The metabolite value can be predicted for

1003 a specific organism in a specific context by using a GEM of the corresponding organism (or closely

1004 related) and then constraining this GEM to the specific context (i.e. primarily defining metabolite

1005 uptake rates). While shadow prices are usually estimated directly by the solver used to run FBA[36],

1006 the robustness of these values is often questioned because of numerical issues. To get more

1007 robust shadow price estimates we solved for each metabolite in each context a separate FBA,

1008 where we constrained the GEM to have an (additional) release of 0.01 mmol/gDW/h of that

1009 metabolite and quantified the change in the objective function, i.e. the change Δy in growth rate

1010 caused by the forced metabolite release. The metabolite value was then quantified as $\Delta y/0.01$. If

1011 the metabolite was predicted to be released by FBA, we added 0.01 mmol/gDW/h to the FBA-

1012 predicted value to estimate the rate. This was only relevant for acetate for *E. coli* in the L-malate

1013 condition, for ethanol, acetate, formate and pyruvate for *S. cerevisiae*, and for pyruvate for *P.*

1014 *putida* in the L-malate condition.

1015

1016 Using this approach, we estimated metabolite values for *E. coli* in batch cultures with glucose,

1017 galactose, L-malate or L-alanine as the carbon source by constraining the uptake of the

1018 corresponding exchange reaction in eciJO1366 to the uptake rate of the carbon source as

1019 estimated from the experimental data. The same approach was used for *C. glutamicum, B.*

1020 *licheniformis* and *S. cerevisiae* using the corresponding models. For *E. coli*, *P. putida* and the two

1021 environmental isolates of the genera *Enterobacter* and *Pseudomonas* we first extracted maximum

1022 growth rates for each organism in each environment from growth curves (using curveball[91]).

1023 Then, we used the respective GEMs to calculate the uptake rates of the corresponding carbon

1024 sources required to sustain the maximum growth rates. These uptake rates were then used to

1025 constrain the GEMs when estimating the metabolite values as described above. The turnover (i.e.

1026 the sum of positive fluxes producing an intracellular metabolite) was estimated by running

1027 parsimonious FBA[74], using the same GEMs constrained in the same way as described for above

1028 for metabolite values.

1029

To validate that the negative correlation between log-transformed metabolite values and release
rates, we compared estimates from eciJO1366 with estimates from the 4 benchmark *E. coli* K-12
MG1655 GEMs iJR904[92], iAF1260[93], iJO1366[94] and iML1515[95] that are of increasing complexity.
We then compared estimated metabolite values and estimated release rates using data from *E.*
coli in bioreactors with glucose, galactose, L-malate or L-alanine as the sole carbon source, the
same data used in Fig. 2A (Fig. S11). In the L-malate condition for the iJR904 model, we discarded
the metabolite value for formate as an outlier because it was extremely small ($<10^{-12}$), below the
solver tolerance. Note that no metabolite values were negative and hence no other values were
discarded upon log-transformation.

1039

Metabolite classification, chemical properties and intracellular concentrations
The charge and molecular weight of metabolites were obtained from eciJO1366[34]. All other
chemical properties and InChIKeys were obtained from PubChem using the python interface
PubChemPy (github.com/mcs07/PubChemPy). The InChIKeys were then used to classify
metabolites into a chemical taxonomy using ClassyFire[96]. To get an appropriate level of detail of
metabolic classification, we used the "Class" level category. However, we separated the class
"Carboxylic acids and derivatives" into "Amino acids" and "Carboxylic acids" based on the
"Subclass" level. We also merged the classes "Hydroxy acids and derivatives" and "Keto acids and
derivatives" into the joint class "Keto / hydroxy acids". Finally, metabolites that did not fall into
either of the abovementioned categories nor the class "Organooxygen compounds" were joined
into the category "Other".

1051

Linear models for evaluating the importance of different factors for predicting metabolite
release rates
Linear models were created and analysed using statsmodels[97]. Compound class and carbon
source was incorporated as categorical variables. Metabolites predicted to have zero turnover
was given a value of 10^{-4}, less than the minimum predicted turnover. Release rates where
corresponding values were missing in any of the compared factors were discarded before model
fitting. To evaluate the statistical significance of the out-of-sample predictions, we performed 10^4

permutations of model fitting and out-of-sample estimations where the labels (either metabolite

or condition) was shuffled on each permutation. We then calculated the fraction of out-of-sample

R^2 values from these permutations below the observed R^2 value to estimate the P value.

1062

Cultivation and exometabolome analyses of KEIO knockout strains

One objective was to study how disruption of key metabolic fluxes would affect extracellular

metabolite concentrations and how this would change if these strains were allowed to evolve in

a simple nutrient environment. The KEIO collection is a collection of non-essential *E. coli* KO

strains that we could use for this purpose[53]. To identify relevant KEIO KO strains, we used

eciJO1366 and parsimonious FBA[74] to predict the effect of a gene knockout on optimal metabolic

fluxes, and previous metabolomics data[52] of these strains to see the effect on intracellular

concentrations (Fig. S20A). We eventually chose seven strains (Δ*aceE*, Δ*cyoD*, Δ*nuoA*, Δ*pgi*, Δ*rpe*,

Δ*sdhB*, Δ*sucB*) that were predicted to have different optimal flux distributions, had many

significantly changed intracellular metabolite concentrations, and targeted different part of key

metabolic pathways (Fig. S20). We included Δ*lacA* as a negative control expected to behave like

the WT as done previously[98].

1075

1076 The selected strains from the KEIO collection were precultured on LB agar and liquid LB as

1077 previously described, and glycerol stocks used for experiments described below were prepared

1078 by adding glycerol to a final concentration of 25 %.

1079

For exometabolome sampling, the KEIO strains and *E. coli* BW25113 were precultured as detailed

above. They were then re-inoculated to OD_{600} = 0.05 in 10 mL M9 + 40 mM galactose (+ 25 µg/mL

kanamycin for KEIO strains) in 50 mL Erlenmeyer flasks (200 rpm, 37 °C, 31 h). 1 mL culture was

sampled into Eppendorf tubes and washed twice with M9 without carbon source by

centrifugation (6000 rcf, 6 min, 4°C). Ultimately, samples were resuspended in 0.5 mL M9 without

carbon source and used to inoculate 200 µL M9 + 40 mM galactose (+25 µg/mL kanamycin for

KEIO strains) in a flat bottom 96-well plate to OD_{600} = 0.05 and 6 mL M9 + 40 mM galactose (+25

µg/mL kanamycin for KEIO strains) to straight glass tubes (15.8 cm tall, 1.5 cm diameter) to OD_{600}

= 0.005. The 96-well plate was incubated in a Synergy H1 plate reader with continuous shaking

(double orbital, 425 cpm, 37 °C), and the glass tubes in a shaking incubator (200 rpm, 37 °C, 45°

angle), both for 108 hours. The experiment in the well plate was used to monitor more closely

the growth (as OD_{600}) of these strains and to help with the timing of the exometabolome

sampling. All strains were cultivated in three replicates. Two samples were collected during mid-

exponential growth phase for all cultures, mostly to ensure that at least one sample from each

replicate was obtained before the end of exponential phase. 1 mL culture volume was centrifuged

(14 000 rpm, 10 min, 4 °C), and supernatant stored in -70 °C for exometabolome analysis. The

second sample for each replicate (Fig. 4A) was analysed using LC-MS at the UNIL Metabolomics

facility, following their protocol described above. Additionally, we also analysed three samples of

pooled inoculums to verify that any carry-over with the inoculum was much lower (or below

detection limit) of than the sample metabolite concentrations. Three replicate pooled samples

were obtained by inoculating 6 mL M9 + 20 mM galactose medium with all the strains, each strain

to an OD_{600} = 0.005, and subsequently handled as described above for the culture samples.

1102

Chemostat evolution experiment with Δ*aceE* and Δ*sucB* in bioreactors

Two of the KEIO KO strains, Δ*aceE* and Δ*sucB*, were selected for the evolution experiment with

continuous cultivation (chemostats) in bioreactors based on their reduced growth compared to

their ancestor and difference in exometabolome patterns (Figs. 4A and S21-S23). The strains were

streaked from glycerol vials onto LB + 25 µg/mL kanamycin agar plates and incubated at 37 °C

overnight. 5-6 colonies were transferred to 500 mL baffled shake flasks containing 75 mL LB + 25

µg/mL kanamycin and incubated for 22 h (200 rpm, 37 °C).

1110

Chemostat cultivations were performed in 1 L DASGIP bioreactors with a constant volume of 300

mL M9 + 20 mM galactose medium + 25 µg/mL kanamycin (prepared as previously described, but

without pH adjustment because of practical challenges with the large volumes required, so pH ≈

7.1). Cultivation was performed at 37 °C with a submerged airflow maintained at 0.5 vvm, and a

cascaded agitation of 400-700 rpm, controlled by a DO set point of 30 %. pH control was started

at day 6 (Δ*aceE*) or day 20 (Δ*sucB*) with a set point of 7.1, controlled using 2 M NaOH (Fig. S25A).

1117

The reactors (three replicates per strain) were inoculated to OD_{600} = 0.1. During the late stage of

exponential growth, pumps were started to feed fresh culture medium and remove used medium.

Medium was fed using a pre-calibrated peristaltic pump and used medium was removed by a

steel pipe placed right above the culture surface. The steel pipe was coupled to a peristaltic pump

operating at a flow rate approximately 1.5x of the medium in flow, to avoid volume accumulation.

The set dilution rate was adjusted several times during the experiment to compensate for

(sometimes surprising) changes in growth dynamics (from 0-0.1 h^{-1} for Δ*aceE*, 0-0.3 for Δ*sucB*,

Fig. S25B). The actual dilution rate was monitored by keeping the flasks with fresh culture medium

on scales. OD_{600} was measured daily, and weekly, larger samples were collected. Culture was

centrifuged (3220 rcf, 10 min, 4 °C), and supernatant and pellet stored at -70 °C. In addition,

cultures were preserved at -70 °C as glycerol stocks. The number of generations were estimated

from the measured dilution rates across the experiment plus the generations during the initial

batch culture phase before the culture dilution was started.

1131

Cultivation of single colonies from evolution experiment and sampling of exometabolome

Culture samples from different bioreactors and timepoints collected in the chemostat experiment

were streaked out onto LB + 25 μg/mL kanamycin agar plates and incubated overnight at 37 °C.

Three colonies were picked and individually inoculated in 20 mL LB + 25 μg/mL kanamycin in 100

mL Erlenmeyer flasks and incubated (37 °C, 200 rpm, 16 h). 12 mL of the culture was then

centrifuged (3220 rcf, 6 min, RT), and the pellets stored at −20 °C for sequencing. The isolates

were stored as glycerol stocks at -70 °C made from these cultures.

1139

2 mL samples from the precultures were washed as previously described, resuspended in M9

without carbon source, and used to inoculate a flat bottom 96-well plate to OD_{600} = 0.05 with 200

μL M9 + 25 mM galactose + 25 μg/mL kanamycin. The M9 medium was prepared as previously

described, but to a pH of 7.1 as in the chemostats. The well plate was incubated in a Synergy H1

plate reader with continuous shaking (double orbital, 425 cpm, 37 °C, 120 h) and the OD_{600} was

read every 20 minutes.

1146

Based on these growth measurements we selected the fastest growing isolates from the last

timepoint of each of the six bioreactors (three bioreactors per strain). These isolates, the initial

KEIO KO strains Δ*aceE* and Δ*sucB*, and the KEIO ancestor *E. coli* BW25113 were precultured and

subsequently washed as described above. The washed preculture were used to inoculate 10 mL

M9 + 20 mM galactose to OD_{600} = 0.01 in straight glass tubes (15.8 cm tall, 1.5 cm diameter) in

triplicates and incubated for 20-84 hours (200 rpm, 37 °C, 45° angle). 1 mL samples for

exometabolome analysis were collected at OD_{600} ≈ 1 (Fig. 4, D and E), filtered using a 0.22 μm

Millex-GV PVDF syringe filter (SLGVR33RS, MilliporeSigma, Darmstadt, Germany) into a 1.5 mL

Eppendorf tube stacked in a cold block (-20 °C), and then stored at -70 °C until it was analysed.

The exometabolome LC-MS analyses were conducted by the UNIL Metabolomics facility, following

the procedure described above.

1158

DNA extraction of samples from selected isolates and chemostat culture samples

The DNA extraction of 14 different samples from the chemostat cultures (#37A D44 M2 Pellet,

#38A D44 M3 Pellet, #39A D44 M4 Pellet, #28B D30 M5 Pellet, #29B D30 M6 Pellet, #30B D30 M7

Pellet) and the selected isolates (sucB-M5-D30-4, sucB-M6-D30-6, sucB-M7-D30-4, sucB-

Ancestor, aceE-M2-D44-2, aceE-M3-D44-3, aceE-M4-D44-1, aceE-Ancestor) was performed by

adapting the protocol from the FastPure Bacteria DNA Isolation Mini Kit (DC103-01, Vazyme,

Nanjing, China). 1 mL GA buffer was added to the thawed pellet samples, and 230 μL of these

resuspensions were kept in Eppendorf tubes for DNA extraction. The rest was centrifuged (4000

rpm, 6 min, RT) and the pellet stored at –20 °C. Genomic DNA (gDNA) was extracted from the 230

μL sample following the manufacturer's protocol. In the final step, elution of the DNA was

performed twice with fresh elution buffer, eluting a total of 100 μL DNA. Extracted gDNA was

stored at –70 °C until sequencing.

1171

Genomic sequencing processing

Genomic DNA from the selected isolates and chemostat samples was sequenced by Novogene

(Planegg, Germany) using the Illumina NovaSeqX PE 150. The isolates and the bioreactor samples

were sequenced to 1 Gb and 10 Gb sequencing depth, respectively. Good quality of both datasets

was ensured using fastqc v. 0.12.1 before trimming the reads with fastp v. 0.24.0 (--

qualified_quality_phred 20 --cut_front --cut_tail --average_qual 20 --length_required 50)[99,100].

Both read sets were mapped against the *E. coli* BW25113 reference genome (NCBI RefSeq

assembly GCF_000750555.1) with minimap2's v. 2.28 default "sr" parameters[101]. The resulting

alignments were filtered with samtools view v. 1.20 (-b -f 3 -q 60) to filter out reads not properly

mapped in pairs or mapped to the reference with a MAPQ lower than 60[102]. Average coverage

was assessed for both datasets showing a minimum average coverage for the chemostat culture

samples of 2017 and 190 for the sequenced isolates. For both datasets, variants were identified

with freebayes v. 1.3.6 (--min-alternate-count 3 -p 1 --min-alternate-fraction 0.05 --pooled-

continuous --haplotype-length 0) and annotated with snpEff v. 5.0 using the BW25113 reference

genome[103,104]. A custom python script was used to filter the annotated variants. For the isolates,

variants with a frequency lower than 0.95 and a coverage lower than 100 were filtered out. For

the chemostat culture samples, variants with a frequency lower than 0.05 and coverage lower

than 100 were filtered out. A variant was considered fixed if its alle frequency was at least 0.9.

Paired intracellular and extracellular metabolomics and live/dead staining

M9 medium with 20 mM galactose, 30 mM malate or 40 mM L-alanine was prepared as detailed

above. *E. coli* K-12 MG1655 was precultured on LB agar plates and in liquid LB as detailed above.

1 mL preculture was washed as described above and used to start a second preculture in 20 mL

M9 + 40 mM L-alanine, inoculated to OD_{600} = 0.05. Another LB preculture of *E. coli* K-12 MG1655

was used to inoculate a second M9 + 20 mM galactose, and a M9 + 30 mM malate preculture in

the same way 12 hours later. After 36/24 hours, 1 mL was collected from each of the three

precultures and washed twice as detailed above. These washed precultures were used to

inoculate 10 mL of the same media (M9 + either 20 mM galactose, 30 mM malate or 40 mM L-

alanine) in straight glass tubes (15.8 cm tall, 1.5 cm diameter). These glass tubes were incubated

for two days (200 rpm, 37 °C, 45° angle). Growth was monitored by measuring OD_{600} directly in

the glass tubes (NANOCOLOR VIS II, MACHEREY-NAGEL, Düren, Germany) at 21 timepoints. The

glass tubes were vortexed briefly before any OD reading or sampling.

1203

A reference exometabolome 0.5 mL sample was collected after 2 hours following the procedure

detailed below. Then, 2 mL samples for paired intra- and extracellular metabolome and live/dead

analyses were collected in the late exponential phase (at $OD_{600} \approx 1.1$, Fig. S32). Exometabolome

samples were obtained by filtering 0.5 mL culture using a 0.22 μm Millex-GV PVDF syringe filter

(SLGVR33RS, MilliporeSigma) into a 1.5 mL Eppendorf tube stacked in a cold block (-20 °C). The

samples for intracellular metabolomics, i.e. cells, were obtained immediately after following this

procedure: A 25 mm 0.22 μm Durapore PVDF filter (GVWP02500, MilliporeSigma) was placed in

a new filter holder. A 5 mL syringe was connected to the inlet of the filter holder, and a vacuum

pump (250 mbar) was connected to the outlet of the filter holder. Then, 2 mL of 37 °C MQ water

was added to pre-wet the filter. Immediately after, 1 mL of culture sample was added to the

syringe and allowed to pass through the filter. If the culture did not pass through solely based on

the vacuum pump, light additional pressure was added by using the syringe itself to push the

culture through the filter. This process took 10-30 seconds. Still using the same set-up, the filter

was finally rinsed with 5 mL of 37 °C MQ water as previously recommended[47]. Two tweezers were

quickly rinsed in MQ water and used to carefully transfer the filter from the filter holder to 2 mL

lysis tubes filled with 1.4 mL of 80:20 methanol:water (v/v), pre-chilled to -20 °C and kept on a

cold block also pre-chilled to -20 °C. All samples were then quickly transferred to -70 °C for

storage. Live/dead staining was performed using a similar approach as previous work[105]: Briefly,

the *live* sample was made by diluting 10 μL culture in 990 μL PBS and the *dead* sample was made

by mixing 100 μL culture with 1 mL 70% isopropanol. Both samples were vortexed and incubated

at room temperature for 1 h. Then, the isopropanol was removed from the dead sample by

centrifugation (8000 rcf, 4 min, RT), pipetting off the supernatant, resuspending the pellet in 1 mL

PBS and vortexing. This procedure was repeated once. Then, both the live and the dead sample

were stained with propidium iodide (PI; P4170, Sigma-Aldrich) and SYBR green (Invitrogen S7563,

Thermo Fisher Scientific, Carlsbad, CA, USA) with the following procedure: 48 μL sample (live or

dead) was mixed in a 96-well plate with 49 μL PBS, 2 μL 0.5 mg/mL PI and 1 μL 100X SYBR green

and incubated for 15 min in the dark. Then, a 0.1X samples of both the stained live and the stained

dead samples were created by diluting 10 μL of each sample in 90 μL of PBS. The stained live/dead

samples were then analyzed immediately on a CytoFLEX S flow cytometer (Beckman Coulter,

Brea, CA, USA) and the CytExpert software (v2.4.0.28). The dead samples were used to make the
gatings required to quantify the fraction of dead cells in the live samples on the PE-H and FITC-h
signals (Table S7, Fig. S32). The extra- and intracellular metabolomics samples were analysed by
the metabolomics facility at UNIL to quantify the five metabolites leucine, glutamate, aspartate,
fructose 6-phosphate and cis-aconitate. These metabolites were chosen because they were
chemically and metabolically diverse and likely to be quantifiable both intracellularly and
extracellularly based on previous experience with similar samples. Also, as glutamate is the most
abundant intracellular metabolite[47,106], it serves as a best-case scenario in our attempt to test
how much of the extracellular metabolite levels that can be explained by cell lysis and the
associated release of intracellular metabolites. Intracellular metabolite samples were first
normalized by the amount of protein detected in the same sample and subsequently converted
to absolute intracellular concentrations using the estimated protein density in *E. coli* ($13.5 \cdot$
$10^{-8} \ \mu g/\mu m^3$)[107]. The fructose 6-phosphate value for sample 1B, timepoint 1, was discarded as
an outlier as it was one order of magnitude larger than any other fructose 6-phosphate
measurements.

1248

Contribution from intracellular metabolites and cell lysis to extracellular concentrations
Approximate cell volume was estimated from measured OD_{600} and a previous estimate of OD to
cell volume for *E. coli* of 3.6 $\mu L/OD_{600}/mL$[108]. From approximate cell volume, measured absolute
intracellular concentrations and cell lysis fractions we then calculated the corresponding change
in extracellular concentrations. This potential contribution was then compared to the net change
in extracellular concentrations between the T0 reference sample (after 2-hours) and the late
exponential phase sample. If all T0 values were below the detection limit, we assumed the initial
concentration of this metabolite to be 0. If only some values at T0 were below the detection limit
we imputed these values to the mean of the detected values, giving conservative estimates for
the net change in extracellular concentrations.
To identify typical biomass degradation products, we used the metabolites consumed by the
biomass equation of the E. coli model iML1515[95], although excluding the soluble pool (as
categorized in[93]) since this should correspond to the intracellular concentrations already

accounted for. Of the metabolites in our dataset (Fig. 3B), only the amino acids were among the

biomass components and only protein depolymerization is therefore covering the contribution

from biomass degradation in this case. To estimate a potential contribution from protein

depolymerization we estimated cell density in gDW/L from OD_{600} (0.346 gDW/L/OD_{600}) and from

this specific amino acid density (in g/L) from recent estimates of *E. coli* amino acid mass

fractions[109]. When mass fractions were attributed to pairs of amino acids (glutamate/glutamine

and aspartate/asparagine), we assumed equal contributions. Amino acid density was converted

from g/L to μM by dividing by the amino acids' molecular weight subtracted the weight of a water

molecule to account for the condensation reaction in protein polymerization. A similar procedure

as described above was followed when we extended this analysis by using literature values for

intracellular concentrations. To map extracellular concentrations in our dataset to literature value

for intracellular concentrations, we used the following procedure: We first tried to map to

literature values from cultivations in minimal medium with the same growth phase

(exponential/stationary) and carbon source. If this wasn't possible, we prioritized mapping to

data from minimal medium conditions of the correct growth phase. If concentrations from

minimal medium conditions were not available, we mapped to data from complex growth media.

When multiple intracellular concentrations mapped to our extracellular data point, we used the

median of the intracellular concentrations. When extracellular concentrations were reported as

two indistinguishable molecules, we attributed equal amounts to the two metabolites. This

includes 2-phosphoglycerate/3-phosphoglycerate and ribulose 5-phosphate/xylulose 5-

phosphate.

1283

Large-scale screening of metabolite release from transporter KO mutants

We selected sixty-seven strains from the KEIO collection[53] to assess if transporters (or

porins/channels) play a role in metabolite release (Table S5). Each of these strains lacks a gene

associated with metabolite transport. Functional annotations of these genes were obtained from

EcoCyc[56] and TCDB[110]. All strains were precultured and washed as described above and then

inoculated to OD_{600}=0.01 in 10 mL of 20 mM M9 glucose medium (pH=7.4) in straight glass tubes

(15.8 cm tall, 1.5 cm diameter) for 20 hours (37 °C and 200 rpm). The glass tubes were angled 45°

in the shaking incubator to improve mixing and aeration. The whole experiment was conducted

in 5 separate batches (conducted on separate days). The KEIO reference strain *E. coli* BW25113

was included in 6 technical replicates in each batch. All other strains cultivated in three technical

replicates.

1295

During the experiment, growth was monitored by measuring OD_{600} directly in the glass tubes on

1- or 2-hour intervals (NANOCOLOR VIS II, MACHEREY-NAGEL, Düren, Germany). The

exometabolomes were sampled at 4,8,10,11,12,13,14,15,16 and 20 hours (+ at 23 and 24 hours

for Δ*tolC*). These timepoints were selected to cover the whole exponential and stationary phase,

with highest temporal resolution in the last part of the exponential phase and during the

transition to stationary phase where expected the largest changes in extracellular metabolite

concentrations to occur. The exometabolome sampling was carried out by first vortexing each

glass tube and then immediately collecting 150 μL into a 96-well filter plate (Millipore MultiScreen

HTS GV Filter Plate, 0.22 μm Hydrophilic PVDF membrane, MSGVN22). The filter plate was stacked

on top of a receiver plate and briefly centrifuged (1 min, 1000 rcf) to filter the cultures. The

receiver plate and the filter plates were both immediately placed on an aluminum plate pre-

chilled and kept on dry ice to cool down the samples quickly before the samples were moved to

a -70°C freezer.

1309

The samples were measured by flow-injection analysis time-of-flight mass spectrometry (FIA-TOF

MS) using an Agilent 6550 quadrupole time of flight mass spectrometer as previously described[54].

Samples were injected online utilizing an Agilent 1100 Series HPLC system (Agilent) coupled to a

Gerstel MPS 3 autosampler (Gerstel). The mobile phase flow rate was set at 0.15 mL/min, with

the isocratic phase composed of 60:40 (v/v) isopropanol and water buffered to a pH of 9 with 4

mM ammonium fluoride. The instrument was run in 4 GHz full-scan mode, collecting spectra

between 50 and 1000 *m/z*. Online mass axis correction was performed with taurocholic acid and

Hexakis (1H, 1H, 3H-tetrafluoropropoxy–phosphazene). For these samples, masses were

annotated by exact mass to within 1.0 ppm of the theoretical monoisotopic mass against

Escherichia coli KEGG metabolites via in-house software as previously described[54]. Processed

mzML files for this dataset can be found at MASSIVE: MSV000097105. Samples from all batches

were initially analyzed in one continuous run (randomized within sampling plates) with four

injections (technical replicates) of each sample. A standard was analyzed at regular intervals

between samples. However, because of the large number of injections (>10,000) instrument

performance declined considerably during the analysis of batch 5, which was therefore re-

analyzed later in a separate run. FIA-MS data were annotated against KEGG identifiers to within 1ppm

of each metabolite's theoretical mass. In some cases, the complexity of MS data (i.e. close neighboring

peaks) yielded improperly centroided mass values which subsequently failed annotation to this stringent

and highly conservative mass accuracy. This led to some differences in annotated metabolites between

batch 1-4 and 5.

1330

TIC-normalized data was used for the analyses described below. First, we filtered the dataset by

removing *m/z* scores where none of the KEGG annotations mapped to the *E. coli* metabolite

database (ECMDB)[46]. SMILES from ECMDB were then used to classify metabolites with

ClassyFire[96]. We used primarily the superclass level to group metabolites, but we separated the

class "Organic acids and derivatives" into "Amino acids", "Carboxylic acids", "Keto / hydroxy acids"

and "Other organic acids". To reduce complexity, we further merged the classes that mapped to

few metabolites into a class named "Other", i.e. "Benzenoids", "Other organic acids",

"Phenylpropanoids and polyketides" and "Organic nitrogen compounds". We then removed

severe outliers, focusing on occurrences where multiple consecutive injections were affected, as

these errors could propagate despite using the median rather than the mean of 4 injections in

further data analyses. We standardized the data from batch 1-4 and batch 5 separately after

removing outliers and before calculating the median value across the 4 injections from each

sample.

1344

To quantify the effect of each KO on each metabolite, we compared the time-series median

standardized Z-score between the KO mutant and the WT, only using the data from the same

batch. To calculate the spread of the data for the WT and KO mutant we first fitted a smoothing

spline (using make_smoothing_spline in Scipy[111], with the regularization parameter = 10) to all

median Z-score values of each strain with the area-under-the curve of the OD600 readings on the

x-axis (Fig. 5A), with weights inversely proportional to the standard deviation of each data point.

We then calculated the spread as the mean absolute deviation from this smoothing spline. We

calculated the difference between the KO strain and the WT as the mean distance between the

two smoothing splines across x-axis locations corresponding to the underlying datapoints and

within the x-range of both datasets. We then calculated the *Effect* as the ratio between the mean

difference and the mean spread. To assess the significance of the estimated effects, we repeated

this procedure over 1000 permutations where we shuffled the KO strain / WT label of all data

points in each permutation. We then counted the number of permutations where we observed a

higher / lower effect than in the actual data (for positive and negative effects).

1359

Amino acid auxotroph co-cultures

E. coli strains auxotrophic for proline (*E. coli* K-12 MG1655 Δ*proC* mCherry), isoleucine (*E. coli* K-

12 MG1655 Δ*ilvE*) and methionine (*E. coli* K-12 MG1655 Δ*metB* GFP) were kindly provided by

Estelle Pignon[112,113]. The strains were transformed with the plasmid pEP28 providing

chloramphenicol resistance[112]. We then cultivated each auxotroph in monoculture, with *E. coli*

BW25113 and with 2-4 relevant transporter KO strains (proline: Δ*proP*, Δ*putP*; methionine:

Δ*brnQ, ΔcstA, ΔmetI, Δmtr*; isoleucine: Δ*glnP*, Δ*tsx*, Δ*yeaS*, Δ*ygaZ*). Strains were precultured (with

25 μg/mL chloramphenicol / kanamycin for auxotrophs / KEIO KO strains) and washed as

previously described, inoculated to OD_{600}=0.001 in 20 mM glucose M9 medium (200 μL) and

cultured in BioTek Synergy H1 platereader (24 hours, 37 °C) as previously described. After 24

hours, the auxotrophs abundances were quantified as CFUs/mL on LB+25 μg/mL chloramphenicol

agar plates. Because methionine was not quantified in the batch 5 data, we had no information

about the effect of Δ*metI* on methionine from the main dataset. However, not all metabolites

were affected by the deteriorating mass-spec quality in the first run of batch 5. We therefore went

back to the first run of batch 5, verified that the distribution of methionine peak intensities was

similar in batch 5 as in batch 1-4, and then estimated the effect of Δ*metI* on methionine from this

data (Effect = 2.56, *P < 0.005*).

1377