Antimicrobial and Biofilm-Inhibitory Potential of Green Synthesized Silver Nanoparticles from *Lantana camara* against *Pseudomonas aeruginosa*

Achal Dharmalal Rajratna, Saurav K. Saha, Tapas K. Sengupta*

Department of Biological Sciences

Indian Institute of Science Education and Research Kolkata

Mohanpur, Nadia- 741 246, West Bengal, India

***Email: senguptk@iiserkol.ac.in**

Abstract

Background:

Emerging antimicrobial resistance poses serious threat to human well-being. Current therapeutics often fail to treat bacterial infections as pathogenic bacteria often attach to various surfaces and form biofilm. One such pathogen, *Pseudomonas aeruginosa*, can form biofilm for its survival in the presence of antibacterial agents. This study aimed to evaluate the antibacterial, antibiofilm, and mechanistic activities of *Lantana camara* leaf extract mediated silver nanoparticles (LCLE-Ag NPs) against *Pseudomonas aeruginosa*, where the extract contributes for the reduction of Ag^+ to Ag^0.

Methods:

LCLE-Ag NPs were synthesized and characterized by using UV visible spectroscopy, dynamic light scattering, and electron microscopy. Antibacterial and antibiofilm activities of the synthesized NPs were assessed against *P. aeruginosa* isolates KPW.1-S1 and HRW.1-S3. Extracellular polysaccharides (EPS), and extracellular DNA (eDNA) and reactive oxygen species (ROS) generation were visualized using epifluorescence microscopy. The ability of the nanoparticles to eradicate preformed biofilm was also evaluated. The expressions of quorum-

26   sensing genes were analyzed using semi quantitative polymerase chain reaction. Cytotoxicity was

27   tested on human kidney epithelial-like cells.

Results:

The synthesized LCLE-Ag NPs displayed an absorption peak at 416 nm and a size range of 30 to

35 nm. They exhibited significant antibacterial and antibiofilm activities correlating with

decreased EPS and eDNA levels. LCLE-Ag NPs effectively disrupted both preformed and

antibiotic-induced biofilms and such activities were found to be associated with increased ROS

levels and decrease in *lasI* and *pqsA* expression. Cytotoxicity assays indicated no significant

toxicity at effective concentrations.

Conclusions:

36   *LCLE*-Ag NPs displayed promising antibacterial and antibiofilm activities against *P. aeruginosa*,

37   acting through biofilm matrix disruption, ROS generation, and quorum-sensing inhibition.

38   **Keywords:** Antibacterial activity; Biofilm; Green synthesized nanoparticles; *Lantana camara*;

39   *Pseudomonas aeruginosa,* Quorum sensing inhibition, Reactive oxygen species.

1. Introduction

Pseudomonas aeruginosa (PA), Gram-negative, rod-shaped bacteria, are well-known

opportunistic pathogens widely found in water, soil, plants, and animals, including humans. These

pathogens are essential to study due to their inherent ability to form biofilm on different surfaces

under adverse conditions as a defence mechanism and for its survival, making it clinically relevant.

Pseudomonas aeruginosa causes secondary infections in immunosuppressed individuals as well

as patients who have cystic fibrosis (1,2), chronic wounds (3), mastitis (4), endometritis (5), and

thermal wounds (6). As stated in reports by the Centers for Disease Control (CDC) and Prevention

experts, it is estimated that about 65% of human bacterial infections are linked to biofilms,

highlighting their impact on health and the challenges they pose for treatment (7,8).

Pseudomonas aeruginosa isolates are generally used as model organisms for biofilm-related

studies as they are avid biofilm formers, causing chronic infections (8). Besides the site of

infections, PA can grow as biofilm on a variety of surfaces, including medical implants and

catheters, making it clinically more vital. Biofilm also helps bacteria avoid phagocytosis and

defend against different constituents of the host's adaptive and innate immune systems, making it

resistant to antimicrobial treatments (8). It is reported that antimicrobial resistance (AMR) was

heightened in biofilm cells compared to planktonic cells (9,10). Thus, the increasing mortality and

antimicrobial resistance of *Pseudomonas aeruginosa* highlight the need to develop novel

therapeutics against the bacteria for combating these infections.

Different novel strategies are being developed against *Pseudomonas* infections in recent times due

to their ability to develop new resistance mechanisms to existing strategies. Newer strategies such

as anti-quorum sensing molecules, synthetic small organic molecules (11), antibiofilm peptides

(12), matrix-degrading enzymes (13), phage therapy (14), and CRISPR-Cas9 (15), photodynamic

therapies (16), vaccines (17), nanoparticles and biomaterials (18) are in use to enhance treatment

effectiveness against infections associated with biofilms. Nanoparticles, in particular, have been

an emerging tool in the recent field of therapeutics.

Nanotechnology-based tools have several advantages over different novel therapeutics, such as a

high surface area to volume ratio, reactivity, and stability (19). These nanotechnologies have

shown a positive impact in the field of biomedicine and pharmaceuticals, due to their effectiveness

in reducing infections with conventional drug administration methods (20). Synthesis of

nanoparticles through chemical synthesis involves an energy-intensive process with harsh

chemical involvement, which can be overridden by green synthesis of nanoparticles, as this method

offers non-toxic plant extract involvement, which acts as a reducing agent (21). The green

synthesized nanoparticles can be produced on a large scale with an eco-friendly approach. The

involvement of biological components in a reducing and capping agent avoids high energy and

pressure needs and, simultaneously less hazardous than chemically synthesized counterparts

(22,23). Among different types of nanoparticles, green synthesized metal-based nanoparticles have

exhibited good potential as antimicrobial agents (24), antifungal agents (25), and also as anticancer

agents (25,26).

Silver nanoparticles have been recognized for their potent antibacterial properties, effectively

targeting both Gram-negative and Gram-positive bacterial strains (27,28). Silver nanotechnology

can potentially inhibit bacterial infections by following the mode of action as cell membrane

damage, producing different types of reactive oxygen species (ROS), and inhibiting enzymes
(29,30). Such green synthesized silver nanoparticles are known to be effective, and they have many
potential applications (31).

Lantana camara, an invasive, adaptable weed native to the American tropics, is a flowering plant
of the Verbenaceae family. It has spread across the world, including India. *L. camara* has been
used in traditional medicine to treat various diseases (32). Different studies have found that *L.*
camara leaves display bactericidal, insecticidal, fungicidal, nematocidal, antioxidant, and
anticancer properties (33,34). Traditionally, various parts of this plant, like flowers, leaves, and
roots, have been utilized in herbal medicine (35). Bioactive compounds present in the plant have
been found to have antibacterial activity. As only a tiny fraction of plant species has been
thoroughly investigated for their therapeutic benefits, deep research into the potential of *L. camara*
as an antimicrobial agent may lead to the discovery of novel treatments that are both effective and
less toxic than current chemotherapeutic agents.

Metallic nanoparticles have been investigated over the past few years, and it has demonstrated the
potential against clinically relevant pathogens, including *P. aeruginosa* (36,37). Yadav et al.
showed that silver nanoparticles are quite effective in inhibiting *PA* biofilms (38). Though metal-
based nanoparticles have promise, their toxicity limits their uses (39). Thus, green synthesis
approaches have shown a promising alternative for PA infection due to their eco-friendly, less
toxic and biocompatible nature (21). Various biosynthesized nanoparticles were also found to be
effective against biofilm formation, as well as the virulence genes of PA (40). *Lantana camara*
conjugated silver nanoparticles were also found to have an antibacterial effect on various bacteria
(41). Although these studies confirm the antibacterial potential of the *Lantana camara* conjugated
silver nanoparticles, their antibiofilm potential has not yet been explored in depth.

Hence, in the current study, *Lantana camara* leaf extract conjugated silver nanoparticles (LCLE-
Ag NPs) were green synthesized and characterized. The synthesized nanoparticles were evaluated
for their antimicrobial and antibiofilm properties against two strains of *Pseudomonas aeruginosa,*
KPW.1-S1 and HRW.1-S3, which were isolated from contaminated water sources and found to
express various virulent genes (42). As biofilm formation depends on the extracellular matrix
components such as exopolysaccharides and extracellular DNA, the alterations of their abundance

due to nanoparticle were examined. Nanoparticle-mediated antibacterial activities are often

associated with oxidative imbalance. Therefore, reactive oxygen species (ROS) generation was

analyzed to find any probable contribution of oxidative stress associated with the antibacterial

activity of LCLE-Ag NPs. Moreover, because sub-MIC concentrations of antibiotics, including

gentamicin (42), are known to induce biofilm formation, the efficacy of the nanoparticles to reduce

or mitigate preformed biofilms induced by gentamicin was evaluated.

Quorum sensing plays a pivotal role in biofilm formation of *Pseudomonas aeruginosa*. The *lasI*

(gene name for acyl-homoserine-lactone synthase) and *pqsA* (gene name for anthranilyl-CoA

synthetase) are two important regulator genes for *P. aeruginosa* quorum sensing, which result in

the enhanced expression of virulence factors and biofilm formation. In the Las quorum sensing

system, *lasI* synthesizes 3-oxoC12-HSL which in turn activates other biofilm related genes.

Consequently, *pqsA* is part of the PQS (*Pseudomonas* Quinolone Signal) system, working

alongside the Las system during biofilm formation. It regulates biofilm physiology and

development and is responsible for eDNA production (43,44). Given the central role of these two

genes in the regulating biofilm, the expression of *lasI* and *pqsA* in the presence of nanoparticles

was analyzed to gain molecular insights into the working mechanism of LCLE-Ag NPs.

2. Materials and Methods

2.1. Bacterial strains of *Pseudomonas aeruginosa*

The *Pseudomonas aeruginosa* strains KPW.1-S1 and HRW.1-S3 were isolated from Kolkata Port

water and Haldi River water, respectively (45) and further identified by 16S rRNA sequencing

(NCBI Accession numbers FJ897721 and FJ897723, respectively). The strains were deposited at

the Microbial Type Culture Collection and Gene Bank (MTCC), Chandigarh, India, under the

catalogue numbers MTCC 10087 and MTCC 10088 (45).

For growth, *P. aeruginosa* KPW.1-S1 and HRW.1-S3 were inoculated and grown in a modified

Bushnell-Haas (BH) medium supplemented with 2% glucose as the carbon source (46).

2.2 Cell Culture of HEK293T cells

The HEK293T (human kidney epithelial-like) cell line was obtained from the National Centre for Cell Science, Pune, India. The cell line was maintained in Dulbecco's Modified Eagle Medium (DMEM) High Glucose (HiMedia, India), supplemented with 10% Fetal Bovine Serum (FBS) (HiMedia, India), 10000 U/mL Penicillin, and 10 mg/mL Streptomycin antibiotic solution (HiMedia, India). Cells were maintained at 37°C in a humidified environment with 5% CO_2 and 95% air to ensure optimal growth conditions (35).

2.3. *Lantana camara* leaf extract (LCLE) preparation

The leaf extract was prepared following the protocol described by Pal et al. (35), with slight modifications. Healthy *L. camara* leaves were collected from the Indian Institute of Science Education and Research, Kolkata campus (22°57′50″N, 88°31′28″E). The leaves were taxonomically verified by the Central National Herbarium, Botanical Survey of India (West Bengal, India), with a voucher reference number IISER/AP/01. Then they were thoroughly washed with water and then shade-dried for 48 hours. The dried leaves were powdered using a mortar and pestle for subsequent processing.

Leaf powder was extracted with absolute ethanol (Merck) at a 1:10 w/v ratio at 37 °C for 18 hours in a shaker. The extract was filtered, concentrated by a rotary evaporator, reconstituted in molecular-grade ethanol (Merck, Germany), sterilized through a syringe filter (HiMedia) of pore size 0.22-μm, and stored at 4 °C.

2.4. Synthesis of silver nanoparticles

A 2 mg/mL solution of ethanol-based leaf extract of *L. camara* dispersed in water and a 3 mM silver nitrate solution were mixed in a conical flask in a 2:1 ratio. The silver nitrate solution was added drop wise over the course of one hour. It was then incubated in a hot water bath at 60 °C for three hours until the solution turned brownish-yellow. The resulting mixture was centrifuged (13,000 rpm, 10 minutes), and the pellet was resuspended in water and stored at 4 °C for further use (41).

Based on the 2:1 ratio, the maximal concentration of plant extract of *L. camara* in synthesized nanoparticles would be 1.33 mg/mL, assuming complete utilization of the extract while nanoparticle formation.

2.5. Characterization of Nanoparticles

2.5.1. Ultraviolet-visible spectroscopy of LCLE-Ag NPs

The synthesized nanoparticles were analyzed using a UV-Vis spectrophotometer (Beckman Coulter DU 730). The analysis was conducted in the wavelength range of 200–800 nm, utilizing continuous scanning mode to capture detailed spectral data. Distilled water was employed as a blank.

2.5.2. Dynamic Light Scattering and Zeta Potential of LCLE-Ag NPs

The polydispersity index (PDI), average particle size, and zeta potential were measured using a Zetasizer (Nano ZS, Malvern) using DLS. The samples were prepared by dispersing them with sonication and diluting them with MiliQ water as required.

2.5.3 Scanning Electron Microscopy and Energy-Dispersive X-ray spectroscopy of LCLE-Ag NPs

The morphological characteristics and elemental composition of the synthesized LCLE-Ag NPs were studied using field emission scanning electron microscopy (FESEM) integrated with energy-dispersive X-ray spectroscopy (EDX) (Supra 55 VP, Carl Zeiss). The nanoparticle was drop-cast on cover slips, then mounted on aluminium stubs for SEM analysis, and *L. camara* powdered leaf extract was directly mounted on stubs for EDX analysis. The samples were subjected to sputter coating and subsequently analyzed under the electron microscope.

2.5.4. Transmission Electron Microscopy of LCLE-Ag NPs

Transmission electron microscopy (TEM) (JEM-2100 Plus, JEOL) was performed to determine the size of the nanoparticles. Samples were drop-cast on a carbon-coated copper grid, which was then left to air-dry overnight. The samples were desiccated and visualized under TEM at an accelerating voltage of 300 kV.

2.5.5. Atomic Force Microscopy of LCLE-Ag NPs

To prepare the LCLE-Ag NP sample for examination using Atomic Force Microscopy (AFM) (Cypher asylum Research AFM, Oxford Instruments), the nanoparticle was drop-cast on a silicon wafer and was placed on a tissue paper-lined petri dish, allowed to air dry in a laminar flow hood. Afterwards, the air-dried silicon wafer was desiccated and then examined under the microscope.

2.5.6. Attenuated Total Reflectance (ATR) FTIR of LCLE-Ag NPs

ATR was used to examine the presence of functional groups in the nanoparticles and *L. camara* leaf extract powdered samples. ATR was performed using a PerkinElmer Spectrum RX1 spectrophotometer in the scanning range of 500-4500 cm^{-1}.

2.6. Minimum Inhibitory Concentration (MIC) assay

The MIC assay was used to evaluate the minimal inhibitory concentration of LCLE-Ag NPs against two different strains of *Pseudomonas aeruginosa*, namely KPW.1-S1 and HRW.1-S3. Various concentrations of LCLE-Ag NPs (5- 10- 20- 50- 100 µg/mL) along with the maximal concentration of 1.33 mg/mL of LCLE were prepared and added to bacterial cultures. The individual bacterial cultures treated with nanoparticles and LCLE were incubated in a shaker (37 °C, 150 rpm) along with their respective blanks. Following 18 hours of incubation, the optical density (OD) was measured using a UV-Vis spectrophotometer (Schimadzu) at 600 nm.

2.7. Quantification of *Pseudomonas aeruginosa* Biofilm

The biofilm load assay was performed on 24-well plates coated with polystyrene (NUNC). 3 x 10^6 cells/mL of *Pseudomonas aeruginosa* KPW.1-S1 and HRW.1-S3 were individually cultured along with the different concentrations of LCLE-Ag NPs and also in the presence of 1.33 mg/ml LCLE in an orbital shaker (37 °C, 150 rpm). After 24 hours of incubation, the optical density of the planktonic cells was measured at 600nm using a UV-Vis spectrophotometer. The wells were rinsed twice with Phosphate Buffered Saline (PBS) to remove any excess planktonic cells, followed by staining the wells with 0.1% Crystal violet (CV) for 45 minutes. Post incubation, the CV was discarded and the wells were washed twice with PBS again. Then, 1 mL of 30% acetic acid was added to each well, and the plates were kept in the shaker (37 °C, 150 rpm) for 30 minutes to solubilize the stain. The optical density of the solubilized crystal violet was measured using a plate reader at 600 nm (BioTek ELx800).

2.8. Confocal Laser Scanning Microscopy for Biofilm Visualization

Pseudomonas aeruginosa KPW.1-S1 and HRW.1-S3 biofilm formation was further studied using confocal laser scanning microscopy (CLSM) to evaluate the impact of maximal concentration of LCLE and different doses of LCLE-Ag NPs. Biofilms were grown on coverslips in 24-well plates with respective treatment doses and incubated at 37 °C (150 rpm). Following 24 hours of incubation, the coverslips were carefully washed twice with PBS and stained with 0.005% (w/v) acridine orange for 5 minutes. Following staining, the coverslips were rinsed twice with PBS and mounted onto clean glass slides using 5% glycerol. Transparent nail polish was used to seal the edges of the coverslip. The samples were examined using a Nikon confocal microscope with an excitation wavelength of 488 nm, and emission was detected in the 293–586 nm range (42).

2.9. Scanning Electron Microscopy of *Pseudomonas aeruginosa* biofilm

The *Pseudomonas aeruginosa* KPW.1-S1 and HRW.1-S3 biofilm was grown on a glass coverslip in 24-well plates. After the 24-hour incubation period, each coverslip was washed twice with PBS to remove any planktonic cells. The coverslips were then immersed in 2.5% glutaraldehyde for 2 hours at 4 °C in the dark to fix the biofilm structure. Then they were washed thrice with PBS. Next, the coverslips were treated with 0.1% osmium tetroxide for 45 minutes at 4 °C in the dark. After another series of three PBS washes, the coverslips were subjected to a dehydration process using an alcohol gradient from 30% to 100%, with each concentration applied for 10 minutes. Finally, the coverslips were dried under a vacuum in a desiccator and examined in Scanning Electron Microscopy using Carl Zeiss smart software for detailed analysis of the biofilm structure (42).

2.10. Visualization and estimation of Extracellular polysaccharide and DNA (eDNA)

The two important extracellular matrix components, exopolysaccharides and extracellular DNA of *Pseudomonas aeruginosa* KPW.1-S1 and HRW.1-S3 biofilms grown absence and presence of different doses of nanoparticles were visualized and analyzed. The exo-polysaccharide and eDNA were visualized by dual staining of Concanavalin A (Con A) (25 µg/mL in PBS) and Propidium Iodide (PI) (0.2 µg/mL in PBS) using Olympus- IX83 fluorescence microscope. The coverslips with biofilm were washed twice with PBS and stained with Con A for 15 minutes, followed by staining with PI for 10 minutes. The coverslips were then washed with PBS and mounted on a

246   glass slide with 5% glycerol and imaged under the fluorescence microscope. The Con A and PI

247   were excited at 495 nm and 493 nm with emission peaks at 519 nm and 636 nm, respectively. The

248   fluorescence intensities of the images were measured using ImageJ software and graphs were

249   plotted accordingly.

2.11. Biofilm eradication assay

Similarly, a pre-formed biofilm load assay was performed on *Pseudomonas aeruginosa* KPW.1-

S1 and HRW.1-S3, to estimate the biofilm eradication potential of the LCLE-Ag NPs on the

formed biofilm. The biofilm was first grown for 24 hours. After this initial growth period of 24

hours, the media with planktonic cells were carefully removed and replaced with fresh BH 2%

glucose media with different concentrations of LCLE-Ag NPs. The rest of the steps were followed

according to the biofilm load assay protocol (Section 2.7).

2.12. Morphological study of the effect of LCLE-Ag NPs on gentamicin induced preformed biofilm

To check the effect of the nanoparticle on this gentamicin induced biofilm, CLSM was performed.

Biofilm was grown on coverslips in the presence of 1.5 μg/mL of gentamicin in BH 2% glucose

media in a shaker for 24 hours. Post-incubation, the media was replaced with fresh media

containing different doses of nanoparticles. Post incubation the coverslips were washed twice with

PBS, followed by a 5-minute staining in 0.005% acridine orange (w/v) solution in the dark. The

stained coverslips were washed with PBS and then mounted on a glass slide with 5% glycerol, and

sealed with nail polish. The samples were visualized using a Nikon confocal microscope with an

excitation wavelength of 488 nm, and emission was detected in the 293–586 nm range.

2.13. Reactive Oxygen Species (ROS) production assay

ROS generation was measured for both the strains using an oxidation-sensitive fluorescent

probe, dichlorodihydrofluorescein diacetate (DCFH-DA) (40). *P. aeruginosa* KPW.1-S1 and

HRW.1-S3 Biofilms were grown on coverslips in a 24-well plate and incubated at 37 °C (150

rpm) for 24 hours. After incubation, the coverslips were gently washed twice with PBS and

stained with 10μM DCFH-DA for 30 min. The prepared coverslips were mounted on a glass

slide using 5% glycerol and sealed with transparent nail polish. The samples were visualized

using an Olympus -IX81 fluorescence microscope, with a blue excitation filter (460-495nm) and

emission filter (510-550nm). The intensity was analyzed using ImageJ software.

2.14. Determination of mRNA level expression of *lasI* and *pqsA* in PA KPW.1-S1 biofilm cells

Semi quantitative polymerase chain reactions were performed to check the expression of quorum-

sensing genes, *lasI* and *pqsA* in absence and presence of different doses of nanoparticles. The

bacterial biofilm was grown in different conditions and harvested for RNA isolation. The RNA

was isolated following the protocol described in Pal et al**.** (35) with minor modifications. The RNA

was quantified and further used for the cDNA synthesis using PrimeScript cDNA synthesis kit

(TaKara, Japan) according to the manufacturer's protocol. The synthesized cDNA was then used

as a template for carrying out PCR of the genes (Table 1). The PCR reaction was carried 25 cycles

for 16s, 38 cycles for both *lasI* and *pqsA*. The amplified PCR products were run on agarose gel

electrophoresis and band quantification was done using GeneTools software. The intensity of the

bands were normalized using their respective housekeeping bands of the same experimental

condition and fold change was plotted.

Table 1: List of primers used in this study

Gene name	Primer sequence	Size	Reference
16S rRNA	FP- 5' ACACTGAGGTGCGAAAGCGTGG 3' RP-5' TCCCCAGCGGGTCGACTTATCG 3'	137 bp	Designed in this study
lasI	FP- 5' CAAAGGCTGGGACGTTAGT 3' RP-5' ATCTGGGTCTTGGCATTG 3'	404 bp	(47)
pqsA	FP- 5' CATGTGTTCCTCGCCAACCG 3' RP- 5' GCCCGCTTCCTCGATAGTGT 3'	117 bp	Designed in this study

2.11. Cell Viability Assay

The cytotoxicity of LCLE-Ag NPs, along with their control (equivalent Ag concentration), was tested on HEK293T cells using a standardized 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) colorimetric assay protocol (35). Briefly, human epithelial kidney like (HEK293T) cells were seeded in a 96-well culture plate with 5000 cells/well and allowed to adhere overnight under standard conditions (37 °C, 5% CO_2). Cells were then treated with 10- 20- 50- 100 μg/mL LCLE-Ag NPs and their respective silver concentrations, along with an untreated control, for 24 hours. Following 24-hour treatment, the MTT assay was performed by replacing spent media with fresh medium containing 0.5 mg/mL of MTT solution (SRL, India). After 4-hour incubation in darkness at (37 °C, 5% CO_2), the formazan conversion reaction was terminated by carefully removing the MTT solution and dissolving the crystalline products with 100 μL dimethyl sulfoxide per well and then incubated for 30 minutes in the dark at 37 °C. Then, the absorbance was taken in a microplate reader at 595 nm. The cytotoxic effect of the LCLE-Ag NPs and $AgNO_3$ was expressed as cell viability percentage compared to the untreated control.

2.12. Statistical analysis

The statistical analysis was carried out using GraphPad Prism 9.5.1, with results expressed as mean ± standard error. A two-tailed Student's t-test was employed to evaluate differences between group means, with statistically significant differences denoted by asterisks (*$P<0.05$; **$P<0.01$; ***$P<0.001$).

3. Results

3.1. Characterization of green synthesized silver nanoparticles

3.1.1. UV-Visible spectral analysis

UV-Visible spectroscopy was performed to check the absorbance band of the prepared LCLE-Ag NP. The distinct peak between 300-500 nm of the prepared nanoparticles revealed the Surface Plasmon Resonance (SPR) peak, indicating the formation of the nanoparticle, with the maximum absorbance at 416 nm (Figure 1).

3.1.2. Dynamic Light Scattering analysis

318 Dynamic Light Scattering was studied to examine the average size and PDI of LCLE-Ag NPs. The
319 average size of the nanoparticles determined by the size analyzer was 35 nm (Figure 2a). The
320 synthesized nanoparticles showed a PDI of 0.3, signifying a homogenous population of
321 nanoparticles. Overall, the result suggests that the synthesized nanoparticle is homogeneous, with
322 a size of around 20-40 nm.

3.1.3. Zeta potential analysis

324 The surface charge of the synthesized LCLE-Ag NPs was determined using a zeta sizer. The graph
325 displays the corresponding average zeta potential of -9.53 mV (Figure 2b). This zeta potential
326 value implies the stability of the synthesized nanoparticles, as the negatively charged nanoparticles
327 will exhibit a repulsive force, resulting in enhanced stability.

3.1.4. Atomic Force Microscopy analysis

329 AFM observations corroborate the obtained morphological results of synthesized nanoparticles.
330 The AFM images revealed that LCLE-Ag NPs exhibited a nearly spherical shape with some degree
331 of agglomeration from lateral and horizontal views (Figure 3a,b). The height and latitude studied
332 in the analysis support the data, suggesting the spherical shape of the prepared nanoparticle.

3.1.5. Scanning Electron Microscopy and Energy Dispersive X-ray Spectroscopy analysis

Scanning Electron Microscopy was performed to depict the morphology of the synthesized
nanoparticles (Figure 3c). The morphology of the prepared nanoparticles was observed to be nearly
spherical with fair agglomeration. The topographical analysis revealed that the average particle
size is 26 nm.

Energy Dispersive X-ray Spectroscopy (EDX) analysis was used to analyze the elemental
composition of the synthesized LCLE-Ag NPs (Supplementary Figure S1a). The graph indicates
a significant peak for Silver with a weight percentage of nearly 70%. The EDX analysis also
confirmed the formation of LCLE-Ag NPs.

The EDX of *L. camara* leaf extract was done to analyze elemental composition (Supplementary
Figure S1 b). Quantitative analysis elucidates that Cl, O, Ca, K, Si, and Ag elements are present in

the leaf extract. Out of these, O and K carried out the majority weight percentages of 78.53% and

11.26%, respectively, with traces of other elements.

3.1.5. Transmission Electron Microscopy

347   The Transmission Electron Microscopy images indicated the formation of approximately spherical

348   nanoparticles with slight agglomeration (Figure 3d).

3.1.7. Attenuated Total Reflectance -Fourier Transform Infrared spectroscopy

ATR-FTIR (attenuated total reflectance – Fourier transform infrared spectroscopy) was conducted

to identify functional groups in *L. camara* leaf extract as well as in the synthesized LCLE-Ag NPs.

The major intense peaks in the leaf extract were observed approximately around 1000 cm^{-1}, 1200

cm^{-1}, 1700 cm^{-1}, 2900 cm^{-1} and 3000 cm^{-1}. The firm, intense broad bands at 2921.62 cm^{-1} and

2853.67 cm^{-1} suggest the presence of amine salts and alkane groups in LCLE-Ag NPs, respectively

(Figure 4a). The small peak around 2500 cm^{-1} is the characteristic peak of the nitrile group and

alkyne in the nanoparticles. Another intense band between 1500 cm^{-1}-1750 cm^{-1} is attributed to a

carbonyl group (41). The peak signifying carbonyl group was found to be weaker in synthesized

LCLE-Ag NPs when compared with *L. camara* leaf extract (Figure 4b). This difference suggests

participation of the carbonyl groups present in the *L. camara* leaf extract along with the silver

during the nanoparticle formation. The phenol groups present in the leaf extract, around 1300 cm^{-1}

^{1}, are likely to be responsible for the bio-reduction of silver ions (48). The medium stretching of

alkane and aldehyde, along with minor stretching of nitrile group and alcohols or ethers, indicates

the presence of different phytoconstituents of *L. camara* in the nanoparticles, confirming the green

synthesis.

3.2. Antibacterial effect of LCLE-Ag NPs against *Pseudomonas aeruginosa*

366   The minimum inhibitory concentration of prepared LCLE-Ag Np was checked at 5-10-20-50-100

367   μg/mL concentrations against *Pseudomonas aeruginosa* KPW.1-S1 and HRW.1-S3 (Figure 5a).

368   For KPW.1-S1, at lower concentrations (5 μg/mL and 10 μg/mL), the nanoparticles exhibited non-

369   significant effects, with only slight reductions in bacterial growth. Further, at 20 μg/mL, the

370   nanoparticles achieved a significant reduction by 0.25-fold, whereas 50 μg/mL concentration of

nanoparticles resulted in a significant decrease by 0.996-fold. In the presence of 100 µg/mL, bacterial growth was entirely inhibited, confirming the effectiveness of LCLE-Ag NPs at this dose.

Similarly, when the minimum inhibitory concentration assay was performed on *Pseudomonas aeruginosa* HRW.1-S3 (Figure 5b), it was found that at the lower doses of nanoparticles (5- 10- 20 µg/mL), the LCLE-Ag NPs showed a 0.27-fold, 0.32-fold, and 0.44-fold effect on bacterial growth, respectively, which further increased to 0.85-fold in the presence of 50 µg/mL LCLE-Ag NPs. At the highest dose of 100 µg/mL LCLE-Ag NPs, the bacterial growth was inhibited entirely, confirming the efficacy of LCLE-Ag NPs at this dose.

Therefore, the MIC concentration of the LCLE-Ag NPs for both the strains was determined to be 100 µg/mL.

For both the strains, 50-100 µg/mL concentrations of LCLE-Ag NPs showed more effect as compared to 1.33 mg/mL of LCLE, which is the maximal possible concentration of LCLE in the prepared nanoparticle (Fig. 5a,b).

3.3. Antibiofilm effect of LCLE-Ag NPs against *Pseudomonas aeruginosa*

The Biofilm load assay was performed to study the effectiveness of the prepared LCLE-Ag NPs against *Pseudomonas aeruginosa* KPW.1-S1 and HRW.1-S3 biofilm. Interestingly, for KPW.1-S1, it was observed that when maximal possible concentration of LCLE (1.33 mg/mL) was used, it encouraged biofilm formation significantly by a 4.85-fold compared to the untreated control (Figure 6a). Importantly, it was observed that the prepared nanoparticles could reduce the biofilm with increasing doses. In the presence of 10 µg/mL LCLE-Ag NP, a non-significant decrease was seen but at 20 µg/mL concentration of LCLE-Ag NPs, the biofilm load decreased by 0.75-fold compared to the control. Higher concentration (50 µg/mL) of LCLE-Ag NP further reduced the biofilm by 0.67-fold and at the concentration of 100 µg/mL, further reduction in biofilm formation was observed, with a 0.97-fold decrease relative to the control (Figure 6a).

The biofilm load assay was also performed against *Pseudomonas aeruginosa* HRW.1-S3 (Figure 6b). Relative to control, the maximal possible concentration of 1.33 mg/mL LCLE showed a significant 4.73-fold increase in biofilm load. At the lower doses of 10 µg/mL and 20 µg/mL, the nanoparticles had a non-significant effect on the biofilm, but at higher doses of 50 µg/mL and 100

µg/mL, the biofilm load was significantly decreased to 0.65-fold and 0.73-fold, respectively
(Figure 6b). The images of the crystal violet-stained biofilms are given in Supplementary Figure
S2 as a supporting document.

Thus, in both the strains the leaf extract at its maximal concentration can induce biofilm in
comparison to control but the nanoparticle were effectively reducing biofilm at concentration of
50 µg/mL and 100 µg/mL.

3.4. Effect of LCLE-Ag NPs on the architecture of *Pseudomonas aeruginosa* biofilms

406 To visualize the reduction in the biofilm load and alterations in biofilm structures in the presence
407 of LCLE-Ag NPs, confocal laser scanning microscopy (CLSM) and scanning electron microscopy
408 (SEM) were performed.

3.4.1. Confocal Laser Scanning Microscopy analysis

410 CLSM analysis showed the inhibitory effect of LCLE-Ag NPs on biofilm formation by PA strains
411 (Figure 7). In comparison to the control samples, thick biofilm formation was observed in the
412 presence of 1.33 mg/mL *L. camara* leaf extract. Interestingly, the biofilm formation was found to
413 be decreased with the increasing concentrations of prepared LCLE-Ag NPs for both of the PA
414 strains (Figure 7a, b).

3.4.2. Scanning Electron Microscopy analysis

The SEM was performed to study the structure of the PA biofilms. For KPW.1-S1 (Figure 8a), it
was observed that most of the cells in control samples adhered to the surface and with each other
to form a dense biofilm. Treatment with 1.33 mg/mL *L. camara* leaf extract induced the bacterial
cells to form a thick and robust biofilm with increased presence of extracellular polymeric
substances and increased number of canopies. However, with the introduction of increasing
concentrations of prepared nanoparticles, it was observed that the biofilm thickness decreased
significantly. In the presence of 50 µg/mL of nanoparticles, mostly individually scattered cells
were present with no substantial EPS production, whereas in the presence of 100 µg/mL of
nanoparticles, very few individual cells, along with some cell debris, were visible (Figure 8a).

In the case of HRW.1-S3 (Figure 8b), the control condition showed the formation of a thick biofilm. The presence of LCLE alone further increased the biofilm formation, with a more thickened biofilm structure with an increased number of canopies. The introduction of low doses of nanoparticles decreased the overall biofilm formation in comparison to the control and LCLE alone. The presence of 50 µg/mL showed a few clumped cells, and in the highest dose (100 µg/mL), very few cells, along with cell debris, were visible (Figure 8b).

3.5. Visualization of Extracellular polysaccharide and extracellular DNA analysis

The alterations in the abundance of extracellular polysaccharides and extracellular DNA (eDNA) due to nanoparticles treatment were examined using dual staining of biofilms with Propidium iodide (PI) and Concanavalin A (Con A). PI binds to extracellular DNA contributed by secretion or membrane-compromised cells present in the matrix, resulting in characteristic red fluorescence, and Con A binds to mannose residues of the extracellular polysaccharides in the biofilm matrix, indicated by blue fluorescence, thus differentiating both the major components of the biofilm matrix. In case of PA KPW.1-S1 biofilm, Con A intensity decreased by 0.10-fold and 0.24-fold in comparison to control at lower doses (10 µg/mL and 20 µg/mL) of LCLE-Ag NPs respectively. At 50 µg/mL dose of the nanoparticles, the Con A intensity decreased by 0.52-fold, suggesting very less secretion of polysaccharides at the higher dose of LCLE-Ag NPs. Correspondingly, at 10 µg/mL and 20 µg/mL, PI intensity reduced by respective 0.27-fold and 0.68-fold, which further decreased significantly at higher doses (50 µg/mL) by 0.88-fold (Figure 9c).

Analyses of both the exopolysaccharide and eDNA components of the PA KPW.1-S1 biofilm matrix showed a significant decrease with the treatment of increasing concentrations of nanoparticle (Figure 9a). In the case of HRW.1-S3 strain, a similar trend was found in eDNA and extracellular polysaccharides (Figure 9c). The reduction in the extracellular components supports the SEM observations, where substantial decrease of EPS was visible. This suggests that the nanoparticle significantly reduces the secretion of biofilm matrix components resulting in an overall decrease in biofilm formation.

Similarly, for the PA HRW.1-S3, Con A fluorescence intensity decreased significantly, showing a 0.27-fold reduction at 10 µg/mL, 0.47-fold reduction at 20 µg/mL, and a 0.77-fold reduction at 50 µg/mL of nanoparticles. PI intensity also declined by 0.23-fold at 10 µg/mL, with further

reductions of 0.54-fold and 0.82-fold observed at 20 µg/mL and 50 µg/mL, respectively (Figure 9d). These observations indicate that LCLE-Ag nanoparticles effectively weaken the biofilm matrix by diminishing the secretion of major EPS components.

3.6. Effect of LCLE-Ag NPs on the preformed biofilm of *Pseudomonas aeruginosa*

Biofilm eradication assays were performed against *Pseudomonas aeruginosa* KPW.1-S1 and HRW.1-S3 to study the effectiveness of LCLE-Ag NPs on the preformed biofilm. For preformed biofilm in KPW.1-S1, addition of LCLE further increased the biofilm by 5.41-fold, whereas the LCLE-Ag NPs showed a decrease in the formed biofilm. Although there was no significant decrease in the presence of lower doses (10 µg/mL and 20 µg/mL) of nanoparticle, higher doses of nanoparticles, (50 µg/mL and 100 µg/mL) could reduced the formed biofilm by 0.71-fold and 0.82-fold, respectively, showcasing its ability to eradicate biofilm (Figure 9a).

In the case of HRW.1-S3, the 1.33 mg/ml LCLE showed a 5.62-fold increase in biofilm load. The lower doses of nanoparticles (10 µg/mL and 20 µg/mL) resulted in a non-significant decrease in the formed biofilm, whereas the higher doses of nanoparticles (50 µg/mL and 100 µg/mL) reduced the formed biofilm significantly by 0.61-fold and 0.67-fold (Figure 9b).

The images of the crystal violet-stained biofilm are given in Supplementary Figure S3 as a supporting document.

3.7. Morphological study on the effect of LCLE-Ag NPs on Gentamicin induced preformed biofilm

The confocal microscopy was performed to study the efficacy of the nanoparticles against preformed biofilms induced by gentamicin in PA strains KPW.1-S1 and HRW.1-S3 (Figure 11 a,b). In PA KPW.1-1, the lower doses of 10 µg/mL and 20 µg/mL LCLE-Ag NPs showed reduction in the biofilm thickness compared to the control biofilm. The higher doses of nanoparticles, 50,100-µg/mL, reduced the biofilm mass as well as the thickness decreased significantly (Figure 11a).

The PA strain HRW.1-S3 also showed significant reduction in biofilm in the presence of higher

doses of the nanoparticle indicating that the nanoparticle can effectively reduce biofilm formed

due to the sub-MIC dose of gentamicin (Figure 11b).

3.8. Quantitative analysis of ROS production in P. aeruginosa biofilms in the presence of

nanoparticle

The ROS production was estimated in *Pseudomonas aeruginosa* KPW.1-S1 and HRW.1-S3

biofilms in the presence and absence of LCLE-Ag NPs using fluorescence microscopy. It was

observed that the lower doses (10 and 20-µg/mL) of LCLE-Ag NPs could increase the ROS

production by 5.21-fold and 6.88-fold, respectively, as compared to control condition. However,

at a higher dose (50 µg/mL) of LCLE-Ag NPs, the ROS production increased only by 3.85-fold,

possibly due to high cell death in the given condition (Figure 12a,c).

In the case of the HRW.1-S3 strain, at the lower doses of LCLE-Ag NPs (10 µg/mL and 20 µg/mL),

the ROS production was increased by 4.49 and 6.24 folds, respectively. Subsequently, at the higher

dose of 50 µg/mL LCLE-Ag NPs, the ROS production increased only till 2.62-fold (Figure 12b,d)

may be due to the same reason as mentioned in case of KPW.1-S1 strain.

3.9. Effect of LCLE-Ag NPs on *lasI* and *pqsA* expression in PA KPW.1-S1

As LCLE-Ag NPs showed a significant effect on formation and eradication of *Pseudomonas*

aeruginosa biofilms, the expression of two important quorum sensing genes namely, *lasI* and *pqsA*

were analyzed.

Both the *lasI* and *pqsA* expressions were found to be decreasing with increasing doses of LCLE-

Ag NPs (Figure 13a). In case of *lasI*, the lower dose (10 µg/mL) of nanoparticles showed a non-

significant decrease, while the higher doses of 20 µg/mL and 50 µg/mL LCLE-Ag NPs showed

0.52-fold and 0.81-fold decreases, respectively (Figure 13b). Similarly, *pqsA* expression showed

a significant decrease of 0.37-fold, 0.61-fold, and 0.86-fold in the presence of 10 µg/mL, 20

µg/mL, and 50 µg/mL LCLE-Ag NPs, respectively (Figure 13c).

Overall, gene expression analysis of *lasI* and *pqsA* showed a decrease in expression pattern with
increasing LCLE-Ag NP doses, indicating disruption of quorum-sensing pathways which are
essential for biofilm formation.

3.10. Effect of LCLE-Ag NPs on the viability of HEK293T cells
The cytotoxic effect of different doses of LCLE-Ag NPs and their equivalent doses of silver ions
was studied in HEK293T cells. For all the doses, LCLE-Ag NPs could not induce any significant
cytotoxicity in HEK293T cells as compared to the untreated cells (Figure 10). Whereas the
equivalent doses of silver showed significant cytotoxicity in the HEK293T cells. These results
clearly indicate that normal silver ions are toxic to cells, but the green synthesized LCLE-Ag NP
has no significant toxicity on the HEK293T cells, making it as a potent candidate for medicinal
uses.
515

4. Discussion

In the present study, silver nanoparticles were successfully green-synthesized using *Lantana*
camara leaf extract, which has emerged as an eco-friendly and sustainable approach worldwide.
The phytochemical constituents of the plant extracts reduced silver ions (Ag^+ to Ag^0) and stabilized
the nanoparticles by acting as a capping agent. Previous studies have shown that bioactive
compounds present in the plant extract acts both as a reducing and a capping agent (49,50).
UV-visible spectroscopy serves as a key analytical tool for determining the formation and
assessing the stability of metal nanoparticles in aqueous solution with a distinct spectrum. In case
of our study, the successful formation of LCLE-Ag NPs was characterized by a surface plasmon
resonance (SPR) peak observed between 350-470 nm (Figure 1). Such SPR signatures are
consistent with previously reported phytogenic silver nanoparticles (51). Zeta potential and DLS
results indicated the good stability of the nanoparticle, with values indicating strong electrostatic
repulsion (-9.54 mV) and a particle size range of 30-50 nm (Figure 2). This suggests that the
phytochemicals present in the leaf extract are enhancing the repulsion between negatively charged
particles and preventing agglomeration as stated in previous reports (52,53).
While evaluating the morphology of LCLE-Ag NPs, the SEM analysis revealed the size of
nanoparticles to be 25-35 nm, having a spherical morphology with slight clustering, which was

further confirmed by TEM and AFM analysis (Figure 3). This clustering is normal due to some
factors, such as concentration and environmental conditions, despite their stabilizing effects
(44,45). EDX analysis confirmed the elemental composition of the synthesized LCLE-Ag NPs.
The EDX spectrum for *L. camara* leaf extract showed a weight percentage of 0% for Ag, indicating
that the amount of silver in nanoparticles is not from *L. camara* but due to the successful formation
of nanoparticles. The ATIR-FTIR study on chemical composition further confirmed the presence
of several functional groups and suggests their involvement in the formation of nanoparticles
(Figure 4). The spectra showed a strong signal for phytoconstituents, employing the successful
formation of green synthesized silver nanoparticles. The FTIR spectra revealed prominent peaks
corresponding to hydroxyl (-OH), carbonyl (C=O), and amine (N-H) groups, which are indicative
of the phytochemical constituents from *L. camara* acting as reducing and stabilizing agents. These
functional groups likely facilitate the capping of nanoparticles, enhancing their stability and
bioactivity as similarly shown in other green nanoparticle studies (46,47).
The LCLE-Ag NPs exhibited strong antibacterial, antibiofilm, and biofilm-eradicating activity
against PA strains KPW.1-S1 and HRW.1-S3 in a dose-dependent manner.
For the KPW.1-S1 strain, the lower concentrations of the nanoparticles did not result in any
significant reduction in growth. In contrast, the higher nanoparticle doses (50 µg/mL and 100
µg/mL) caused a marked and significant inhibition of bacterial growth. For the HRW.1-S3 strain,
all tested concentrations of LCLE-Ag NPs exhibited significant antibacterial activity, with the
higher doses exerting a more antibacterial effect than LCLE alone (Figure 5). Such differences in
inhibitory potentials between the nanoparticle and leaf extract at elevated NP doses strongly
support the superior efficiency of the nanoparticles over the crude extract and the probable reason
for this antibacterial activity can be attributed to the disruption of the cell membranes, ROS
generation, and impact on overall cell functions as seen in various other studies (57–59).
The biofilm load assay for both the strains further demonstrated that the synthesized nanoparticle
had an inhibitory effect on the biofilm load in a dose-dependent manner, whereas the maximal
possible concentration of LCLE rather promoted biofilm formation (Figure 6). Earlier reports
confirmed the biofilm-disruptive properties of AgNPs synthesized using *A. cobbe* leaf extract but
did not explore the biofilm-inducing effects of the corresponding plant extracts (60). Our findings,
therefore, significantly extend the existing knowledge by uncovering that *L. camara* leaf extract

alone may induce PA biofilm formation, thus highlighting the importance of LCLE-Ag NPs for both prevention and eradication of resilient bacterial communities.

This effect of LCLE-Ag NPs on the biofilm was further confirmed when we visualized the biofilm using CLSM and SEM. The CLSM imaging supported the observation that LCLE alone induced 3D biofilm formation, but LCLE-Ag NPs could inhibit the biofilm (Figure 7). Consistent with this, SEM analysis showed a substantial reduction in biofilm biomass and visibly diminished EPS in nanoparticle-treated samples of both PA strains, while the leaf extract alone enhanced biofilm structure, producing distinct canopy-like formations (Figure 8). Con A-PI staining confirmed the significant reduction of two extracellular polymeric components: exopolysaccharides and extracellular DNA, which are important for biofilm formation (44,61) in the presence of the nanoparticles (Figure 9). These results in the decrease in EPS components correlated with the observed SEM results (Figure 8). Siddique and his coworkers primarily demonstrated that silver nanoparticles suppress EPS production (62); however, the study did not investigate EPS subcomponents or 3D biofilm morphology in depth. Here, we not only confirmed biomass loss but also quantified matrix-specific disruption, offering mechanistic evidence for nanoparticle-induced biofilm reduction. Further, the biofilm eradication potential of the nanoparticles were assessed on preformed biofilm as well as on gentamicin induced preformed biofilm. Gentamicin is a commonly prescribed drug against PA, and previous findings showed that it can induce biofilm formation by PA in the presence of sub-MIC doses (42,63). The LCLE-Ag NPs were found to be able to reduce preformed biofilms, whereas leaf extract alone further enhanced the biofilm, as evident by the biofilm load assay (Figure 10). The effect of nanoparticles on gentamicin-induced biofilm was further studied using CLSM to understand how effectively nanoparticles can reduce it. The results of the CLSM studies showed that the higher doses of nanoparticles were effectively able to minimize the biofilm thickness and biomass compared to the control, suggesting the strong potential of nanoparticles in the therapeutic world (Figure 11). Tabassum et al. demonstrated the antibiofilm and biofilm eradication potential of Pyoverdine-silver nanoparticles (64). In continuation of these efforts, our study further evaluates the efficacy of nanoparticle against antibiotic-induced biofilms by demonstrating that LCLE-Ag NPs strongly disperse both conventional and gentamicin-induced biofilms, thereby reinforcing their therapeutic superiority.

Previous studies have shown that ROS affects biofilms by degrading the biofilm matrix (65,66), and thereby inhibits biofilm formation (67). In the present study, the ROS intensity was found to

be increased significantly in the presence of nanoparticles (Figure 12), however, the increase was

slightly less in the higher dose of 50 µg/ml, probably due to early cell death as reported earlier

(68). Hamida et al. demonstrated that silver nanoparticles synthesised using *Desertifilum sp.* exert

their antimicrobial activity mainly through oxidative stress induced by ROS generation, leading to

cell death (69). Notably, the effective concentrations of LCLE-AgNPs used in the present study

was comparatively lower, as compared to the nanoparticles reported by Hamida et al., showcasing

the better efficacy of the silver nanoparticle synthesized using LCLE.

A dose-dependent decrease in the expression of two major quorum-sensing genes*, lasI and pqsA,*

in PA was observed upon treatment with LCLE-Ag NPs (Figure 13). Down-regulation of these

genes indicates that the nanoparticles likely interfere with autoinducer synthesis in both the Las

and PQS systems, resulting in decreased biofilm formation. Gurunathan et al. highlighted the

phenotypic aspects of the antibacterial and antibiofilm effects of *A. cobbe* leaf extract mediated

silver nanoparticles (60). Our findings further complement these findings by demonstrating that

LCLE-Ag NPs modulate QS at the molecular level by directly downregulating QS-regulatory

genes, thereby inhibiting biofilm formation.

Having addressed the bactericidal and antibiofilm properties of LCLE-Ag NPs, it was essential to

assess the toxicity of LCLE-Ag NP on the mammalian cell line. The MTT assay of Human Kidney

Epithelial-like cells with different doses of LCLE-Ag NPs exhibited minimal toxicity, with cell

viability ranging from 90% to 100% (Figure 14). In contrast, equivalent doses of silver (with

respect to LCLE-Ag NP) showed high toxicity. Thus, the comparative toxicity profiling reinforces

that the green-synthesized nanoparticles offer a safer and favorable alternative over conventional

silver ion.

Several studies have demonstrated the successful green-synthesis of silver nanoparticles and their

efficacy against PA (70,71). Ajitha et al. reported the antibacterial potential of *L. camara* leaf,

which exhibited positive effects against both Gram-negative and Gram-positive bacteria (72). In

another study, Momani et al. demonstrated the antibiofilm activity of biosynthesized silver

nanoparticles using *P. harmala* seed extract against PA isolates obtained from cystic fibrosis

patients, highlighting the potential of plant-mediated nanoparticles (22), thus encouraged

exploration of *L. camara* leaf extract as a biocompatible agent in this study. Our present work

further advances this line of research by showing that L. camara-mediated silver nanoparticles are

capable of exerting antibacterial and antibiofilm effects, including the inhibition of biofilm

formation, as well as the eradication of conventional and antibiotic-induced biofilms against

Pseudomonas aeruginosa. At the cellular level, this study demonstrated nanoparticle effects

leading to cell damage/death by ROS production and molecular emphasis on quorum-sensing

genes. Additionally, *L. camara*-mediated silver nanoparticles could not exert any significant

cytotoxicity against Human Kidney Epithelial-like cells.

In conclusion, the enhanced efficacy and reduced toxicity of LCLE-Ag NPs synthesized using *L.*

camara may be attributed to their nanoscale size, high surface area-to-volume ratio, ROS

generation and their inhibitory effect on the quorum sensing systems. Future studies on elucidating

the detailed molecular mechanism underlying the antibiofilm activity of the LCLE-Ag NPs will

help to find relevant targets in *Pseudomonas* biofilms for remedial purposes.

Acknowledgment

All authors would like to acknowledge the imaging facilities of the Indian Institute of Science

Education and Research, Kolkata. The authors would also acknowledge Prof. Rahul Banerjee,

DCS IISER Kolkata, for helping with the ATR-FTIR analysis. Authors also thank Arundhaty Pal

and Sourav Sanyal, DBS IISER Kolkata for helping in MTT assays. SKS acknowledges IISER

Kolkata for providing fellowship.

CRediT authorship contribution statement

Conceptualization, **ADR, SKS and TKS;** Methodology**, ADR and SKS;** Investigation**, ADR and**

SKS; Formal analysis**, ADR, SKS and TKS;** Data curation**, ADR and SKS;** Resources, **TKS;**

writing—original draft preparation**, ADR and SKS;** writing—review and editing, **ADR, SKS and**

TKS; supervision**, TKS;** funding acquisition**, TKS; All authors have read and agreed to the**

published version of the manuscript.

Funding

The study was funded by the Indian Institute of Science Education and Research, Kolkata. The funding source has no involvement in the study design; collection, analysis and interpretation of data and writing and preparation of the article.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

All data generated or analyzed during this study are included in this present article.

Declaration

Conflict of interest

The authors declare that there is no conflict of interest.

Ethical Approval

Not applicable.

Consent to Participate

All authors had consented to participate in the study.

Consent for Publication

All authors have given consent for publication.

References

References

1.   Mirzaei R, Mohammadzadeh R, Alikhani MY, Shokri Moghadam M, Karampoor S, Kazemi S, et
al. The biofilm-associated bacterial infections unrelated to indwelling devices. IUBMB Life
[Internet]. 2020;72(7):1271–85. Available from:
https://iubmb.onlinelibrary.wiley.com/doi/abs/10.1002/iub.2266

2.   Pressler T, Bohmova C, Conway S, Dumcius S, Hjelte L, Høiby N, et al. Chronic Pseudomonas
aeruginosa infection definition: EuroCareCF Working Group report. J Cyst Fibros. 2011;10 Suppl
2:S75-8.

3.   Gondil VS, Subhadra B. Biofilms and their role on diseases. BMC Microbiol [Internet].
2023;23(1):203. Available from: https://doi.org/10.1186/s12866-023-02954-2

4.   Vestby LK, Grønseth T, Simm R, Nesse LL. Bacterial Biofilm and its Role in the Pathogenesis of
Disease. Antibiotics [Internet]. 2020;9(2):59. Available from: https://www.mdpi.com/2079-
6382/9/2/59

5.   Ferris RA, McCue PM, Borlee GI, Glapa KE, Martin KH, Mangalea MR, et al. Model of Chronic
Equine Endometritis Involving a Pseudomonas aeruginosa Biofilm. Infect Immun [Internet].
2017;85(12):10.1128/iai.00332-17. Available from:
https://journals.asm.org/doi/abs/10.1128/iai.00332-17

6.   Tuon FF, Dantas LR, Suss PH, Tasca Ribeiro VS. Pathogenesis of the Pseudomonas aeruginosa
Biofilm: A Review. Pathogens. 20220227th ed. 2022;11(3).

7.   Potera C. Forging a link between biofilms and disease. Science (1979). 1999;283(5409):1837,
1839.

8.   Thi MTT, Wibowo D, Rehm BHA. Pseudomonas aeruginosa Biofilms. Int J Mol Sci [Internet].
2020;21(22):8671. Available from: https://www.mdpi.com/1422-0067/21/22/8671

9.   Sharma D, Misba L, Khan AU. Antibiotics versus biofilm: an emerging battleground in microbial
communities. Antimicrob Resist Infect Control. 20190516th ed. 2019;8:76.

10.   Tabassum N, Khan F, Jeong GJ, Jo DM, Kim YM. Silver nanoparticles synthesized from
Pseudomonas aeruginosa pyoverdine: Antibiofilm and antivirulence agents. Biofilm [Internet].
2024;7:100192. Available from:
https://www.sciencedirect.com/science/article/pii/S2590207524000170

11.   Parrino B, Schillaci D, Carnevale I, Giovannetti E, Diana P, Cirrincione G, et al. Synthetic small
molecules as anti-biofilm agents in the struggle against antibiotic resistance. Eur J Med Chem.
20181017th ed. 2019;161:154–78.

12.   de la Fuente-Núñez C, Reffuveille F, Haney EF, Straus SK, Hancock RE. Broad-spectrum anti-
biofilm peptide that targets a cellular stress response. PLoS Pathog. 20140522nd ed.
2014;10(5):e1004152.

13.   Kaplan JB. Biofilm Matrix-Degrading Enzymes. In: Donelli G, editor. Microbial Biofilms:
Methods and Protocols [Internet]. New York, NY: Springer New York; 2014. p. 203–13.
Available from: https://doi.org/10.1007/978-1-4939-0467-9_14

14.   Furfaro LL, Payne MS, Chang BJ. Bacteriophage Therapy: Clinical Trials and Regulatory
Hurdles. Front Cell Infect Microbiol. 20181023rd ed. 2018;8:376.

15.   Hegde S, Nilyanimit P, Kozlova E, Anderson ER, Narra HP, Sahni SK, et al. CRISPR/Cas9-
mediated gene deletion of the ompA gene in symbiotic Cedecea neteri impairs biofilm formation

711		and reduces gut colonization of Aedes aegypti mosquitoes. PLoS Negl Trop Dis. 20191202nd ed.
712		2019;13(12):e0007883.
713	16.	Carpenter BL, Situ X, Scholle F, Bartelmess J, Weare WW, Ghiladi RA. Antiviral, Antifungal and
714		Antibacterial Activities of a BODIPY-Based Photosensitizer. Molecules [Internet].
715		2015;20(6):10604–21. Available from: https://www.mdpi.com/1420-3049/20/6/10604
716	17.	Koo H, Allan RN, Howlin RP, Stoodley P, Hall-Stoodley L. Targeting microbial biofilms: current
717		and prospective therapeutic strategies. Nat Rev Microbiol. 20170925th ed. 2017;15(12):740–55.
718	18.	Huang D ni, Wang J, Ren K feng, Ji J. Functionalized biomaterials to combat biofilms. Biomater
719		Sci [Internet]. 2020;8(15):4052–66. Available from: http://dx.doi.org/10.1039/D0BM00526F
720	19.	Öztürk K, Kaplan M, Çalış S. Effects of nanoparticle size, shape, and zeta potential on drug
721		delivery. Int J Pharm [Internet]. 2024;666:124799. Available from:
722		https://www.sciencedirect.com/science/article/pii/S0378517324010330
723	20.	Klębowski B, Depciuch J, Parlińska-Wojtan M, Baran J. Applications of Noble Metal-Based
724		Nanoparticles in Medicine. Int J Mol Sci. 20181213th ed. 2018;19(12).
725	21.	Ying S, Guan Z, Ofoegbu PC, Clubb P, Rico C, He F, et al. Green synthesis of nanoparticles:
726		Current developments and limitations. Environ Technol Innov [Internet]. 2022;26:102336.
727		Available from: https://www.sciencedirect.com/science/article/pii/S2352186422000359
728	22.	Al-Momani H, Almasri M, Al Balawi D, Hamed S, Albiss BA, Aldabaibeh N, et al. The efficacy
729		of biosynthesized silver nanoparticles against Pseudomonas aeruginosa isolates from cystic
730		fibrosis patients. Sci Rep. 20230601st ed. 2023;13(1):8876.
731	23.	Han C, Romero N, Fischer S, Dookran J, Berger A, Doiron AL. Recent developments in the use of
732		nanoparticles for treatment of biofilms. Nanotechnol Rev [Internet]. 2017;6(5):383–404. Available
733		from: https://doi.org/10.1515/ntrev-2016-0054
734	24.	Ali SG, Jalal M, Ahmad H, Sharma D, Ahmad A, Umar K, et al. Green Synthesis of Silver
735		Nanoparticles from Camellia sinensis and Its Antimicrobial and Antibiofilm Effect against
736		Clinical Isolates. Materials [Internet]. 2022;15(19):6978. Available from:
737		https://www.mdpi.com/1996-1944/15/19/6978
738	25.	Khedr WE, Shaheen MNF, Elmahdy EM, El-Bendary MA, Hamed AA, Mohamedin AH. Silver
739		and gold nanoparticles: Eco-friendly synthesis, antibiofilm, antiviral, and anticancer bioactivities.
740		Prep Biochem Biotechnol. 20230823rd ed. 2024;54(4):470–82.
741	26.	Wypij M, Jędrzejewski T, Trzcińska-Wencel J, Ostrowski M, Rai M, Golińska P. Green
742		Synthesized Silver Nanoparticles: Antibacterial and Anticancer Activities, Biocompatibility, and
743		Analyses of Surface-Attached Proteins. Front Microbiol. 20210422nd ed. 2021;12:632505.
744	27.	Ansari MA, Khan HM, Khan AA, Cameotra SS, Pal R. Antibiofilm efficacy of silver
745		nanoparticles against biofilm of extended spectrum β-lactamase isolates of Escherichia coli and
746		Klebsiella pneumoniae. Appl Nanosci [Internet]. 2014;4(7):859–68. Available from:
747		https://doi.org/10.1007/s13204-013-0266-1
748	28.	Salomoni R, Léo P, Montemor AF, Rinaldi BG, Rodrigues M. Antibacterial effect of silver
749		nanoparticles in Pseudomonas aeruginosa. Nanotechnol Sci Appl. 20170629th ed. 2017;10:115–
750		21.
751	29.	Augustine R, Hasan A, Primavera R, Wilson RJ, Thakor AS, Kevadiya BD. Cellular uptake and
752		retention of nanoparticles: Insights on particle properties and interaction with cellular components.
753		Mater Today Commun [Internet]. 2020;25:101692. Available from:
754		https://www.sciencedirect.com/science/article/pii/S2352492820327033

30. de la Fuente-Núñez C, Reffuveille F, Haney EF, Straus SK, Hancock RE. Broad-spectrum anti-
biofilm peptide that targets a cellular stress response. PLoS Pathog. 20140522nd ed.
2014;10(5):e1004152.

31. Alomar TS, AlMasoud N, Awad MA, AlOmar RS, Merghani NM, El-Zaidy M, et al. Designing
Green Synthesis-Based Silver Nanoparticles for Antimicrobial Theranostics and Cancer Invasion
Prevention. Int J Nanomedicine. 20240521st ed. 2024;19:4451–64.

32. Barreto F, Sousa E, Campos A, Costa J, Rodrigues F. Antibacterial Activity of Lantana camara
Linn and Lantana montevidensis Brig Extracts from Cariri-Ceará, Brazil. J Young Pharm.
2010;2(1):42–4.

33. Mansoori A, Singh NK, Dubey SK, Thakur TK, Alkan N, Das SN, et al. Phytochemical
Characterization and Assessment of Crude Extracts From Lantana camara L. for Antioxidant and
Antimicrobial Activity. In: Frontiers in Agronomy. 2020.

34. Vishal H, Firoj A, Asha S, Rohankumar C. Phytochemical analysis and antimicrobial activity of
Lantana camara. International Journal of Pharmaceutical Chemistry and Analysis [Internet].
2022;2022:15736. Available from: https://www.ijpca.org/article-details/15736

35. Pal A, Sanyal S, Das S, Sengupta TK. Effect of Lantana camara Ethanolic Leaf Extract on
Survival and Migration of MDA-MB-231 Triple-Negative Breast Cancer Cell Line. J Herb Med
[Internet]. 2024;43:100837. Available from:
https://www.sciencedirect.com/science/article/pii/S2210803323002154

36. Bruna T, Maldonado-Bravo F, Jara P, Caro N. Silver Nanoparticles and Their Antibacterial
Applications. Int J Mol Sci. 20210704th ed. 2021;22(13).

37. Sakthi Devi R, Girigoswami A, Siddharth M, Girigoswami K. Applications of Gold and Silver
Nanoparticles in Theranostics. Appl Biochem Biotechnol. 20220513th ed. 2022;194(9):4187–219.

38. Yadav S, Pawar S, Patil S. Inhibition of Pseudomonas aeruginosa Biofilm Formation Using Silver
Nanoparticles. Cureus. 20250122nd ed. 2025;17(1):e77848.

39. Omran BA, Tseng BS, Baek KH. Nanocomposites against Pseudomonas aeruginosa biofilms:
Recent advances, challenges, and future prospects. Microbiol Res [Internet]. 2024;282:127656.
Available from: https://www.sciencedirect.com/science/article/pii/S0944501324000570

40. Abdel-Fatah SS, Mohammad NH, Elshimy R, Mosallam FM. Impeding microbial biofilm
formation and Pseudomonas aeruginosa virulence genes using biologically synthesized silver
Carthamus nanoparticles. Microb Cell Fact [Internet]. 2024;23(1):240. Available from:
https://doi.org/10.1186/s12934-024-02508-9

41. Ajitha B, Ashok Kumar Reddy Y, Sreedhara Reddy P. Green synthesis and characterization of
silver nanoparticles using Lantana camara leaf extract. Mater Sci Eng C Mater Biol Appl.
20150109th ed. 2015;49:373–81.

42. Kumar A, Saha SK, Banerjee P, Prasad K, Sengupta TK. Antibiotic-Induced Biofilm Formations
in Pseudomonas aeruginosa Strains KPW.1-S1 and HRW.1-S3 are Associated with Increased
Production of eDNA and Exoproteins, Increased ROS Generation, and Increased Cell Surface
Hydrophobicity. Curr Microbiol. 20231117th ed. 2023;81(1):11.

43. Miranda SW, Asfahl KL, Dandekar AA, Greenberg EP. Pseudomonas aeruginosa Quorum
Sensing. Adv Exp Med Biol [Internet]. 2022 [cited 2025 Nov 18];1386:95–115. Available from:
https://link.springer.com/chapter/10.1007/978-3-031-08491-1_4

44. Tahrioui A, Duchesne R, Bouffartigues E, Rodrigues S, Maillot O, Tortuel D, et al. Extracellular
DNA release, quorum sensing, and PrrF1/F2 small RNAs are key players in Pseudomonas

aeruginosa tobramycin-enhanced biofilm formation. npj Biofilms and Microbiomes 2019 5:1
[Internet]. 2019 May 23 [cited 2025 Nov 18];5(1):15-. Available from:
https://www.nature.com/articles/s41522-019-0088-3

45.   Dasgupta D, Ghosh R, Sengupta TK. Biofilm-Mediated Enhanced Crude Oil Degradation by
Newly Isolated Pseudomonas Species . ISRN Biotechnol. 2013 Mar 5;2013:1–13.

46.   Bushnell LD, Haas HF. The Utilization of Certain Hydrocarbons by Microorganisms. J Bacteriol.
1941;41(5):653–73.

47.   Kumar A, Saha SK, Banerjee P, Prasad K, Sengupta TK. Antibiotic-Induced Biofilm Formations
in Pseudomonas aeruginosa Strains KPW.1-S1 and HRW.1-S3 are Associated with Increased
Production of eDNA and Exoproteins, Increased ROS Generation, and Increased Cell Surface
Hydrophobicity. Curr Microbiol. 20231117th ed. 2023;81(1):11.

48.   Manjamadha VP, Muthukumar K. Ultrasound assisted green synthesis of silver nanoparticles
using weed plant. Bioprocess Biosyst Eng. 20160111th ed. 2016;39(3):401–11.

49.   Roy A, Bulut O, Some S, Mandal AK, Yilmaz MD. Green synthesis of silver nanoparticles:
biomolecule-nanoparticle organizations targeting antimicrobial activity. RSC Adv [Internet]. 2019
Jan 18 [cited 2025 Dec 8];9(5):2673–702. Available from:
https://pubs.rsc.org/en/content/articlehtml/2019/ra/c8ra08982e

50.   Singh P, Kim YJ, Zhang D, Yang DC. Biological Synthesis of Nanoparticles from Plants and
Microorganisms. Trends Biotechnol [Internet]. 2016 Jul 1 [cited 2025 Dec 8];34(7):588–99.
Available from: https://www.sciencedirect.com/science/article/pii/S0167779916000408

51.   Khalil MMH, Ismail EH, El-Baghdady KZ, Mohamed D. Green synthesis of silver nanoparticles
using olive leaf extract and its antibacterial activity. Arabian Journal of Chemistry [Internet]. 2014
Dec 1 [cited 2025 Dec 8];7(6):1131–9. Available from:
https://www.sciencedirect.com/science/article/pii/S1878535213000968

52.   Mohammadi A, Hashemi N, Ghassabzadeh M, Sharafi A, Yazdinezhad A, Danafar H. Green
synthesis and toxicological evaluation of zinc oxide nanoparticles utilizing Punica granatum fruit
Peel extract: an eco-friendly approach. Scientific Reports 2025 15:1 [Internet]. 2025 Jul 1 [cited
2025 Dec 8];15(1):20853-. Available from: https://www.nature.com/articles/s41598-025-05544-6

53.   Min KH, Shin JW, Ki MR, Pack SP. Green synthesis of silver nanoparticles on biosilica diatomite:
Well-dispersed particle formation and reusability. Process Biochemistry [Internet]. 2023 Feb 1
[cited 2025 Dec 8];125:232–8. Available from:
https://www.sciencedirect.com/science/article/pii/S1359511322004585#sec0050

54.   Giri AK, Jena B, Biswal B, Pradhan AK, Arakha M, Acharya S, et al. Green synthesis and
characterization of silver nanoparticles using Eugenia roxburghii DC. extract and activity against
biofilm-producing bacteria. Sci Rep [Internet]. 2022;12(1):8383. Available from:
https://doi.org/10.1038/s41598-022-12484-y

55.   Jaast S, Grewal A. Green synthesis of silver nanoparticles, characterization and evaluation of their
photocatalytic dye degradation activity. Current Research in Green and Sustainable Chemistry
[Internet]. 2021;4:100195. Available from:
https://www.sciencedirect.com/science/article/pii/S2666086521001429

56.   Younis NS, Mohamed ME, El Semary NA. Green Synthesis of Silver Nanoparticles by the
Cyanobacteria Synechocystis sp.: Characterization, Antimicrobial and Diabetic Wound-Healing
Actions. Mar Drugs. 20220106th ed. 2022;20(1).

57. Dakal TC, Kumar A, Majumdar RS, Yadav V. Mechanistic Basis of Antimicrobial Actions of
Silver Nanoparticles. Front Microbiol [Internet]. 2016 Nov 16 [cited 2025 Dec 8];7(NOV):1831.
Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC5110546/

58. Morones JR, Elechiguerra JL, Camacho A, Holt K, Kouri JB, Ramírez JT, et al. The bactericidal
effect of silver nanoparticles. Nanotechnology [Internet]. 2005 Aug 26 [cited 2025 Dec
8];16(10):2346. Available from: https://iopscience.iop.org/article/10.1088/0957-4484/16/10/059

59. Tiwari H, Samal K, Geed SR, Bera S, Das C, Mohanty K. Green synthesis of silver nanoparticles
for ultrafiltration membrane surface modification and antimicrobial activity. Sustainable
Chemistry for Climate Action [Internet]. 2023;3:100031. Available from:
https://www.sciencedirect.com/science/article/pii/S2772826923000202

60. Gurunathan S, Han JW, Kwon DN, Kim JH. Enhanced antibacterial and anti-biofilm activities of
silver nanoparticles against Gram-negative and Gram-positive bacteria. Nanoscale Res Lett
[Internet]. 2014 [cited 2025 Dec 8];9(1):373. Available from:
https://pmc.ncbi.nlm.nih.gov/articles/PMC4127560/

61. Lewenza S. Extracellular DNA-induced antimicrobial peptide resistance mechanisms in
Pseudomonas aeruginosa. Front Microbiol [Internet]. 2013 Feb 14 [cited 2025 Nov
18];4(FEB):41081. Available from: www.frontiersin.org

62. Siddique MH, Aslam B, Imran M, Ashraf A, Nadeem H, Hayat S, et al. Effect of Silver
Nanoparticles on Biofilm Formation and EPS Production of Multidrug-Resistant Klebsiella
pneumoniae. Biomed Res Int [Internet]. 2020 [cited 2025 Dec 8];2020:6398165. Available from:
https://pmc.ncbi.nlm.nih.gov/articles/PMC7189323/

63. Song T, Duperthuy M, Wai SN. Sub-Optimal Treatment of Bacterial Biofilms. Antibiotics 2016,
Vol 5, Page 23 [Internet]. 2016 Jun 22 [cited 2025 Nov 18];5(2):23. Available from:
https://www.mdpi.com/2079-6382/5/2/23/htm

64. Tabassum N, Khan F, Jeong GJ, Jo DM, Kim YM. Silver nanoparticles synthesized from
Pseudomonas aeruginosa pyoverdine: Antibiofilm and antivirulence agents. Biofilm. 20240315th
ed. 2024;7:100192.

65. Kromer C, Schwibbert K, Radunz S, Thiele D, Laux P, Luch A, et al. ROS generating BODIPY
loaded nanoparticles for photodynamic eradication of biofilms. Front Microbiol. 20231012th ed.
2023;14:1274715.

66. Benoit DSW, Sims KR, Fraser D. Nanoparticles for Oral Biofilm Treatments. ACS Nano
[Internet]. 2019 May 28 [cited 2025 Nov 18];13(5):4869–75. Available from:
/doi/pdf/10.1021/acsnano.9b02816?ref=article_openPDF

67. Qayyum S, Oves M, Khan AU. Obliteration of bacterial growth and biofilm through ROS
generation by facilely synthesized green silver nanoparticles. PLoS One [Internet].
2017;12(8):e0181363. Available from: https://doi.org/10.1371/journal.pone.0181363

68. Ali HM, Karam K, Khan T, Wahab S, Ullah S, Sadiq M. Reactive oxygen species induced
oxidative damage to DNA, lipids, and proteins of antibiotic-resistant bacteria by plant-based silver
nanoparticles. 3 Biotech [Internet]. 2023 Dec 1 [cited 2025 Nov 18];13(12):414. Available from:
https://pmc.ncbi.nlm.nih.gov/articles/PMC10665289/

69. Hamida RS, Abdelmeguid NE, Ali MA, Bin-Meferij MM, Khalil MI. Synthesis of Silver
Nanoparticles Using a Novel Cyanobacteria Desertifilum sp. extract: Their Antibacterial and
Cytotoxicity Effects. Int J Nanomedicine [Internet]. 2020 [cited 2025 Dec 8];15:49. Available
from: https://pmc.ncbi.nlm.nih.gov/articles/PMC6956868/

70. Kemala P, Idroes R, Khairan K, Ramli M, Ginting B, Helwani Z, et al. Eco-friendly synthesis of
silver nanoparticles: Enhancing optimization reaction, characterization, and antimicrobial
properties with Lantana camara from geothermal area. S Afr J Chem Eng [Internet]. 2025;51:57–
67. Available from: https://www.sciencedirect.com/science/article/pii/S1026918524001306
71. Patil SP, Sahu KG, Andhale GS. Lantana camara mediated green synthesis of metallic
nanoparticles: A mini review. Inorg Chem Commun [Internet]. 2024;160:111853. Available from:
https://www.sciencedirect.com/science/article/pii/S138770032301465X
72. Ajitha B, Ashok Kumar Reddy Y, Shameer S, Rajesh KM, Suneetha Y, Sreedhara Reddy P.
Lantana camara leaf extract mediated silver nanoparticles: Antibacterial, green catalyst. J
Photochem Photobiol B [Internet]. 2015;149:84–92. Available from:
https://www.sciencedirect.com/science/article/pii/S1011134415001839

Figure 1.

Fig. 1 UV–vis absorbance spectra of AgNO₃ solution (blue line) and *L. camara* leaf extract
(Orange line), and LCLE-Ag NPs (grey line)

Figure 2.

Fig 2. Particle size and zeta potential analysis of green synthesized LCLE-Ag NPs (a) Particle size

analysis, (b) Zeta potential analysis of LCLE-Ag NPs

Figure 3.

Fig 3. Morphological and characteristic analysis of green synthesized LCLE-Ag NPs (a) AFM

height, (b) AFM amplitude, (c) SEM, (d) TEM

Figure 4.

Fig 4. FTIR spectrum of (a) *Lantana camara* leaf extract, (b) Green synthesized LCLE-Ag NPs

Figure 5.

Fig 5. Minimum inhibitory concentration assay of green synthesized LCLE-Ag NPs against *P.*
aeruginosa (a) KPW.1-S1 and (b) HRW.1-S3; Data presented as the mean ± SEM; n=9; N=3; *,
P ≤ 0.05, **P ≤0.01, ***P ≤0.001, ****P ≤0.0001, while 'ns' indicates non-significance

Figure 6.

Fig 6. Biofilm load assay of *P. aeruginosa* KPW.1-S1 and HRW.1-S3 in the presence of LCLE-
Ag NP (a) O.D. of biofilm cells of KPW.1-S1 (b) O.D. of biofilm cells of HRW.1-S3; Data
represent mean ± SEM of n=9; N=3; *, P ≤ 0.05, **P ≤0.01, ***P ≤0.001, ****P ≤0.0001, while
'ns' indicates non-significance

Figure 7.

Fig 7. Representative confocal laser scanning microscopy images of *P. aeruginosa* biofilm in the
presence of LCLE-Ag NP (a) 100X images of KPW.1-S1 biofilm (b) 100X images of HRW.1-S3
biofilm; Data represent mean ± SEM of n=9; N=3; *, P ≤ 0.05, **P ≤0.01, ***P ≤0.001, ****P
≤0.0001

Figure 8.

Fig 8. Representative Scanning Electron Microscope images of *P. aeruginosa* biofilm in the
presence of LCLE-Ag NP (a) 5.0KX images of KPW.1-S1 biofilm (b) 5.0KX images of HRW.1-
S3 biofilm; Data represent mean ± SEM of n=6; N=2; *, P ≤ 0.05, **P ≤0.01, ***P ≤0.001, ****P
≤0.0001

Figure 9.

Fig 9. Representative epifluorescence images of *P. aeruginosa* biofilm stained with Con A (blue)

and PI (red) in the presence of LCLE-Ag NP (a) 60X images of KPW.1-S1 biofilm (b) fold change

intensity for the strain KPW.1-S1 (c) 60X images HRW.1-S3 biofilm (d) fold change intensity for

the strain HRW.1-S3; Data represent mean ± SEM of n=6; N=2; *, P ≤ 0.05, **P ≤0.01, ***P

≤0.001, ****P ≤0.0001

Figure 10.

Fig 10. Biofilm eradication assay of *P. aeruginosa* KPW.1-S1 and HRW.1-S3 in the presence of
LCLE-Ag NP (a) O.D. of biofilm cells of KPW.1-S1 (b) O.D. of biofilm cells of HRW.1-S3; Data
represent mean ± SEM of n=9; N=3; *, $P \leq 0.05$, **$P \leq 0.01$, ***$P \leq 0.001$, ****$P \leq 0.0001$, while
'ns' indicates non-significance

Figure 11.

Fig 11. Representative confocal laser scanning microscopy images of gentamicin induced *P.*
aeruginosa KPW.1-S1 and HRW.1-S3 biofilm in the presence of LCLE-Ag NP (a) 100X images
of KPW.1-S1 biofilm (b) 100X images of HRW.1-S3 biofilm; Data represent mean ± SEM of n=6;
N=2; *, P ≤ 0.05, **P ≤0.01, ***P ≤0.001, ****P ≤0.0001

Figure 12.

Fig 12. Representative epifluorescence images ROS generation in *P. aeruginosa* KPW.1-S1 and

HRW.1-S3 biofilm cells in the presence of LCLE-Ag NP (a) 40X images of KPW.1-S1 biofilm

(b) 40X images of HRW.1-S3 biofilm (c) relative fold change for the strain KPW.1-S1 (d) relative

fold change for the strain HRW.1-S3; Data represent mean ± SEM of n=6; N=2; *, P ≤ 0.05, **P

≤0.01, ***P ≤0.001, ****P ≤0.0001

Figure 13.

Fig 13. Representative images of semi-quantitative PCR amplified DNA of respective genes in

agarose gel (a) agarose gel images of *lasI, pqsA* and 16s (b) fold change in *lasI* expression (c) fold

change in *pqsA* expression; Data represent mean ± SEM of n=6; N=2; *, P ≤ 0.05, **P ≤0.01, ***P

≤0.001, ****P ≤0.0001

Figure 14.

Fig 14. Quantification of viable cells by MTT assay in HEK293T cells in the presence of LCLE-Ag NPs and its equivalent silver concentration; Data represent mean $\pm$ SEM of n=9; N=3; *, P $\leq$ 0.05, **P $\leq$0.01, ***P $\leq$0.001, ****P $\leq$0.0001