# A phylogenomic and metagenomic meta-analysis of bacterial diversity in the phyllosphere lifts a veil on Hyphomicrobiales dark matter

**Authors:** Jean-Baptiste Leducq[1]*, Louis-Philippe St-Amand[1], David Ross[2] and Steven W. Kembel[2]

**Affiliations:** 1 - Département de Phytologie (FSAA), IBIS, CRIV - Université Laval - Québec (QC) G1V 0A6 – Canada
2 - Département des sciences biologiques – Université du Québec à Montréal – Canada

**Email address: (*corresponding author):** jean-baptiste.leducq.1@fsaa.ulaval.ca

**Abstract -** The phyllosphere, or above-ground part of plants, hosts diverse bacterial communities that play critical ecological roles and provide beneficial functions for the plant. Yet, the diversity of major plant-associated bacteria groups like Hyphomicrobiales (Alphaproteobacteria) remain largely uncharacterized in the phyllosphere, due to limitations of 16S rRNA metabarcoding and biases in reference databases toward model taxa and environments. Using a meta-analysis combining metabarcoding, metagenomics and phylogenomics, we explored the worldwide diversity of leaf-associated Hyphomicrobiales. We confirmed *Methylobacterium* was ubiquitous in the phyllosphere and revealed the dominance of two under-characterized Hyphomicrobiales taxa: *Lichenibacterium*, a recently described genus of lichen-associated bacteria previously identified as "1174-901-12" in taxonomic databases, and RH-AL1, an undescribed genus of bacteria related to Beijerinckiaceae, previously isolated from coal slag. Despite their abundance in the phyllosphere, *Lichenibacterium* and RH_AL1 could not be properly detected by classical 16S rRNA gene barcoding, underlining biases in reference databases and a lack of taxonomic resolution for this marker gene. As for *Methylobacterium,* a

significant proportion of *Lichenibacterium* and RH-AL1 were also detected in association with lichens and in environments with harsh conditions like exposed surfaces, air and snow, suggesting airborne or waterborne dispersal and high resilience in harsh environments. Overall our study stresses the need to move toward metagenomics and culturomics to increase the representativeness of leaf-associated bacterial taxa in reference databases, and to improve our understanding of the evolutionary and functional mechanisms underpinning bacteria adaptations to living on plants.

**Keywords:** Phyllosphere, Hyphomicrobiales, *Methylobacterium*, *Lichenibacterium*, metagenomics, phylogenomics.

## Highlights

- Global meta-analysis reveals core Hyphomicrobiales in the phyllosphere
- *Lichenibacterium* and *Methylobacterium* dominate the phyllosphere and exposed surfaces
- Novel phyllosphere clades RH-AL1 and JAJXWB01 were identified by phylogenomics
- 16S rRNA gene limits taxonomic resolution in the phyllosphere; metagenomics refines taxonomy

## INTRODUCTION

The phyllosphere, or the aerial parts of plants including leaves, has been intensively studied due to its crucial role in global carbon and nitrogen cycles. Recent advances in microbial ecology also revealed that, despite exposure to UV radiation, drought, and rapid environmental fluctuations, plant leaves host a large microbial diversity providing essential functions for plant health including nutrient acquisition, deterring pathogens, and facilitating resilience to stresses [40]. The phyllosphere microbiota can also modulate leaf surface chemical composition by

releasing plant growth promoters such as vitamins and hormones, and by degrading toxic compounds such as pesticides [47]. Our knowledge about beneficial plant-associated bacterial diversity has been dominated by studies of the rhizosphere, and the phyllosphere was less well studied until recently, first because of technical biases, as classical microbiology often relied on unrealistic lab conditions developed for the optimal growth of a few model microbes, and second for biological reasons, because of the inherent characteristics of different plants tissues. Roots are less exposed to environmental variation and are more nutrient-rich habitats than leaves. Therefore, root-associated microbes appear to be more abundant, and microbial diversity and culture conditions are more predictable in the rhizosphere than in the phyllosphere. For instance, Hyphomicrobiales (previously Rhizobiales) is a major plant-associated bacterial order that includes the intensively studied nitrogen-fixing bacteria in root nodules [12,42], while the phyllosphere-associated Hyphomicrobiales are mostly known for *Methylobacterium* (Methylobacteriaceae), a ubiquitous group of bacteria metabolizing one-carbon compounds such as methanol that are released by plants during cell-wall synthesis [38]. As a result, only a tiny fraction of leaf-associated microbes have been studied and cultivated so far, stressing the need for more studies that consider the phyllosphere, and for culture conditions that maximize their isolation success, such as those developed for the *Arabidopsis* microbiome [6]. Since the advent of molecular biology, microbial taxonomy has been based on universal marker genes (i.e. the 16S rRNA gene in bacteria), allowing identification of major taxonomic groups and their phylogenetic relationships [28]. Unfortunately, those markers do not give the resolution required to confidently define lower taxonomic units (genus, species), introducing unwanted artefacts in the phylogeny and classification of major groups of plant-associated microbes including *Methylobacterium* [2]. Alternatively, the *rpoB* gene has higher resolution [27] and has been used as a molecular marker to infer *Methylobacterium* diversity in the phyllosphere, with potential use at a larger taxonomic scale in the Hyphomicrobiales [19]. Overall, metabarcoding has allowed the quantification of microbial diversity without using cultivation, but is still limited by using a

single marker gene. Alternatively, shotgun metagenomics is a promising approach to infer the taxonomy, function and diversity of phyllosphere bacterial communities. This approach is however limited by the overall low bacterial abundance on the surface of plant leaves, and by the presence of high concentrations of plant host DNA for endophytic organisms. Finally, whole genome sequencing (WGS) of isolates, including plant-associated bacteria, coupled with phylogenomics (whole-genome-based phylogeny) substantially improved the taxonomy of Hyphomicrobiales in recent years [12,16]. Regardless of the method, taxonomic databases are still biased toward model taxa like *Methylobacterium*, and we argue that major phyllosphere-associated Hyphomicrobiales taxa are still omitted, underrepresented or misclassified in these databases. For instance, *1174-901-12* is an uncultured Hyphomicrobiales taxon identified in the 16S rRNA gene SILVA database [32] that has been consistently detected worldwide on the surface of plant leaves [3,15,18,19,22,35,43,45], where it typically represents 3 to 22% of total bacteria abundance. Little is known about the taxonomy and the ecological role of *1174-901-12*, although its detection in lichens [34], in aerosols [24] and on various exposed surfaces [4,25,33,39] suggests a broader ecological spectrum than the phyllosphere. According to the SILVA database (v138.1), *1174-901-12* is member of the Beijerinckiaceae, a catch-all family of phyllosphere-associated Hyphomicrobiales also including *Methyllocella*, *Methylocystis*, *Methylobacterium* and *Roseiarcus*, for instance. According to the List of Prokaryotic names with Standing in Nomenclature (LPSN), these genera belong to distinct families, with only *Methyllocella* being a "true" Beijerinckiaceae member. The underrepresentation and misclassification of major phyllosphere-associated Hyphomicrobiales such as *Methylobacterium* and *1174-901-12* illustrates the limits of the 16S rRNA gene in describing the composition of microbial communities in the phyllosphere. It also stresses the need to move towards a metagenomic approach to enrich databases with genomes from uncultured taxa, and to a phylogenomic-based approach to clarify their taxonomy.

Using a meta-analysis combining metabarcoding, metagenomics and phylogenomics, we explored worldwide the diversity of leaf-associated Hyphomicrobiales. We confirmed the ubiquity of *Methylobacterium* in the phyllosphere and revealed the dominance of two under-characterized Hyphomicrobiales taxa: *Lichenibacterium*, a recently described genus of lichen-associated bacteria formerly identified as *1174-901-12*, and RH-AL1, an undescribed genus of bacteria related to Beijerinckiaceae, previously isolated from coal slag. The fact that *Lichenibacterium* and RH_AL1 were not properly detected by classical 16S rRNA gene barcoding despite their abundance in the phyllosphere illustrates biases in reference databases and the lack of resolution for this marker gene. Like for *Methylobacterium,* a significant proportion of *Lichenibacterium* and RH-AL1 were also detected in lichen and in environments with harsh conditions like exposed surfaces, air and snow, suggesting airborne or waterborne dispersal and high resilience in stressful environments. Overall, our study stresses the biases of classical tools and reference database, as well as the power of metagenomics and phylogenomic to reveal uncultured leaf-associated bacteria. Future works should also use such tools, as well as culturomics, to increase the representation of these taxa in reference databases, and to address evolutionary and functional mechanisms underpinning bacterial adaptation to life on plants.

**METHODS**

***Metagenome and genome collection of representative Hyphomicrobiales taxa from the phyllosphere, moss and lichen***

We assembled a collection of Hyphomicrobiales (syn. Rhizobiales) genomes and metagenome-assembled genomes (MAGs) covering the diversity observed at the surface of tree leaves. We used 22 MAGs from the leaves of 25 tree species occurring in the Canadian temperate forest, that were previously identified as Rhizobiales with the GTDB database (unpublished data;

bioproject PRJNA1291777; Dataset 1). According to the GTDB-based taxonomic classification, these 22 leaf-associated MAGs were assigned to family Beijerinckiaceae, in genera *Methylobacterium* (n= 4), *Enterovirga* (n=1), *Lichenihabitans* (n= 8), RH-AL1 (n= 5) JAJXWB01 (n= 2) or unidentified at the genus level (n=2; Dataset 1). From here, we use exclusively the LPSN as a reference for the taxonomy of Hyphomicrobiales. According to the LPSN, *Methylobacterium* and *Enterovirga* are part of Methylobacteriaceae, a well-characterized phyllosphere-associated family [2]. *Lichenihabitans* [26] is part of Lichenihabitantaceae, a recently described lichen-associated family of bacteria, also including genera *Lichenifustis* [30], *and Lichenibacterium* [29]. RH_AL1 is represented by two strains isolated from coal slag [44]. JAJXWB01 was identified by metagenomics in mine drainage sediments (unpublished data; bioproject PRJNA666025; Dataset 1) and has no reference genome in NCBI.

Accordingly, we retrieved 97 reference genomes from NCBI, spanning the taxonomic and phylogenic range of Hyphomicrobiales taxa identified in leaf-associated MAGs (Methylobacteriaceae, Lichenihabitantaceae). For Methylobacteriaceae, we maximized the genomic diversity of reference genomes by selecting the best assembly from each lineage previously identified in *Methylobacterium* (n=23; [2]), *Enterovirga* (n=3), *Microvirga* (n=2) and in a newly described genera, *Hohaiivirga* (n=1; [41]; Dataset 1). Because other leaf-associated taxa had fewer representatives in genomic databases, we retrieved all available genomes for Lichenihabitantaceae (n=8), and RH-AL1 [44] (n=2; Dataset 1). According to the most recent phylogenomic-based study on Hyphomicrobiales [12], the aforementioned taxa form a monophyletic group within Hyphomicrobiales, which we will refer as the phyllosphere-associated Hyphomicrobiales, using Rhabdaerophilaceae as the closest known outgroup, for which we retrieved the reference genome (Dataset 1). To mitigate biases in phylogenomic reconstruction of the whole group, we also included NCBI RefSeq genomes from families that were not identified in leaf-associated MAGs, but were also members of this monophyletic group: Methylocystaceae (n=12), Beijerinckiaceae (n=10), Roseiarcaceae (n=1), Rhodoblastaceae

(n=3), Chelatococcaceae (n=11), Alsobacteraceae (n=3), Boseaceae (n=14) and

Salinarimonadaceae (n=3; Dataset 1).

We also retrieved any available MAGs in NCBI databases that could be assigned to the

phyllosphere-associated Hyphomicrobiales families. Most identified MAGs came from a

worldwide survey of lichen-associated microbiota (n=43; [37]) and from a study on moss-

associated microbiota from the Canadian boreal forest (n=5; [17]). Three additional

phyllosphere-associated Hyphomicrobiales MAGs were identified in glacier sediments [9], in

mining soil [23] and in mine drainage sediments (JAJXWB01; unpublished data; Bioproject

PRJNA666025; Dataset 1).

### Phylogenomic reconstruction of plant-associated Hyphomicrobiales

We reconstructed the phylogenetic tree of phyllosphere-associated Hyphomicrobiales as

described previously for Methylobacteriaceae [20] with the following modifications. Briefly, we

annotated *de novo* the 97 genomes and 73 MAGs using RAST [5] and identified core genes (i.e.

genes present in a single copy in all genomes), excluding hypothetical proteins. Because our

analysis included highly fragmented MAGs, we used more relaxed criteria to define the core

genome. First, we considered any gene present on average in 2 copies or less in at least 80% of

genome assemblies as candidate core genes (MAGs excluded). Second, we estimated the real

copy number of each candidate core gene in each assembly. As the proxy for the copy number,

we normalized the total size of gene copies observed for a given gene in each assembly by the

average size of all copies observed across all assemblies. We considered any normalized value

lower than 0.7 (missing or too fragmented copy) or above 1.3 (likely duplication) as missing data.

Third, we only considered candidate core genes with missing data in less than 30% of

assemblies as core genes. We performed sequence alignment for each core gene individually,

with the R package *msa* [8], using nucleotide sequence for rna-encoding genes and amino-acid

sequence for protein-encoding genes. We computed a maximum-likelihood phylogenetic tree for each core gene independently using RAxML (GTRGAMMA model, 1,000 replicated trees [36]). We reconstructed the phyllosphere-associated Hyphomicrobiales consensus tree using ASTRAL-III, a coalescent-based method combining ML trees determined for each core gene independently, accounting for incomplete lineage sorting and horizontal gene transfers [46].

**Identification of *1174-901-12* as *Lichenihabitantaceae***

We performed a preliminary phylogenetic analysis of 16S rRNA sequences used to define *1174-901-12* in the SILVA database, version 138.1, in order to validate that this taxon corresponded to *Lichenibacterium*, as identified in the SILVA database version 138.2. This included the 24 SILVA database sequences (that we numbered from 1 to 24), all complete 16S rRNA gene sequences available for the family Lichenihabitantaceae in NCBI databases, and two sequences from *Roseiarcus* as outgroup. The Maximum-Likelihood tree was inferred with RAxML (1,000 bootstraps, GTRGAMMA).

***Meta-analysis of barcoding-based phyllosphere studies***

We selected previously published metabarcoding data based on the 16S rRNA gene from the tree phyllosphere and potentially related microbial environments and sources (other plants, mosses, lichen, exposed surfaces, air, snow). We restricted our survey to 13 recent studies (Dataset 2) for which BioProjects and related metadata were publicly available. For each BioProject, we processed 16S rRNA gene reads to obtain an ASV (Amplicon Sequence Variants) abundance table per sample, using the package *dada2* in R [10] as described previously [19], with the following modifications. According to sequence quality profiles specific to each BioProject, 3' ends of forward and reverse reads from each sequence were trimmed

(option truncLen in function *filterAndTrim*) at specific lengths (Dataset 2). Additionally, 5' ends of both reads were trimmed 20bp to remove the primer part (option trimLeft in function *filterAndTrim*). To correct for the great variation in read number among samples and BioProjects, we performed rarefaction curves to determine a conservative number of sequences to conserve per sample. Accordingly, we randomly selected 3,000 sequences per sample, hence excluding samples with fewer than 3,000 sequences (Dataset 2). We pooled together unique ASV sequences found among all BioProjects after rarefaction (N=57,100 unique sequences) and inferred their taxonomy in two steps. We first classified all ASVs to the genus level with the SILVA 138.2 database [32] using the *assignTaxonomy* function (R package dada2). To verify SILVA v 138.2 accuracy in predicting phyllosphere-associated Hyphomicrobiales ASVs, we extracted 16S rRNA gene sequences from complete phyllosphere-associated Hyphomicrobiales reference genomes (see section *Phylogenomic reconstruction of phyllosphere-associated Hyphomicrobiales*) and classified them as we did for ASVs. We also performed a phylogenetic analysis of 16S rRNA sequences extracted from reference genomes for a comparison with our phylogenomic reconstruction. The Maximum-Likelihood tree was inferred with RAxML (1,000 bootstraps, GTRGAMMA). For the second step, we refined ASV classification restricted to phyllosphere-associated Hyphomicrobiales sequences (i.e. identified as Beijerinckiaceae, *Alsobacter* or *Phreatobacter* with SILVA v138.2 database) using the *assignTaxonomy* function with 16s rRNA gene sequences we extracted from complete phyllosphere-associated Hyphomicrobiales reference genomes as references. We subdivided Bioprojects from different climatic regions, or with samples from different substrates, in different categories. We subdivided Bioproject PRJNA100102 in three categories: Temperate forest, Subtropical forest and Tropical forest [43]. We subdivided Bioproject PRJNA290145 in three categories: Bark samples, Lichen samples and Moss Samples [4]. For each BioProject or category within BioProjects (n=17), we estimated the diversity of bacteria taxon as the relative abundance of sequences assigned to each taxon, focusing on Hyphomicrobiales dominant families and nested genera.

**Comparison of 16s rRNA and *rpoB*–based barcoding for the identification of phyllosphere-associated Hyphomicrobiales taxa.**

To assess the performance of the *rpoB* gene as a phylogenetic and taxonomic marker for phyllosphere-associated *Hyphomicrobiales*, we first retrieved complete *rpoB* gene sequences from reference genomes and MAGs with known taxonomy. *rpoB* sequences were extracted from RAST output files, aligned, and their phylogeny was assessed with RAxML as previously described for core genes. To compare *rpoB*- and 16S rRNA gene–based taxonomic resolution, we used amplicon data we previously generated [19], including 184 phyllosphere samples collected from two temperate forests in Quebec, Canada: *Station Biologique des Laurentides* (SBL, n = 99) and *Mont St-Hilaire* (MSH, n = 85). All samples were previously amplified with primers optimized for Methylobacteriaceae and targeting a hypervariable region of the *rpoB* gene [19]. We used *rpoB* ASV abundance tables and taxonomy from our previous study as a starting point. We reclassified ASVs using *rpoB* sequences from genomes and MAGs as references, excluding sequences that did not completely overlapped ASV sequences. Because reference sequences only included phyllosphere-associated Hyphomicrobiales, we only performed reclassification for ASVs previously identified as Rhizobiales (former Hyphomicrobiales), hence excluding Caulobacterales, Sphingomonadales and Rhodospirillales. Within Rhizobiales, we excluded ASVs previously assigned to other families (Aurantimonadaceae, Bradyrhizobiaceae, Brucellaceae, Hyphomicrobiaceae, Phyllobacteriaceae and Rhizobiaceae). Relative abundance per sample was thereafter recalculated for major Hyphomicrobiales taxa. We evaluated the congruence in relative abundance estimations between *rpoB*- and 16S rRNA gene–based barcoding. We compared the relative abundance of major phyllosphere-associated Hyphomicrobiales genera in 41 leaf samples for which metabarcoding was performed with both 16S rRNA and *rpoB* genes. To

evaluate the congruence, we used a linear model on relative abundances after a Hellinger

transformation to correct for rare taxa [13] and compared the observed estimate (slope) of the

model to its expected distribution by realizing 10,000 random permutations of relative

abundance values among leaf samples. Because relative abundances observed for a particular

taxa could be site-dependent in this dataset [19], we constrained permutations within sites (SBL

and MSH). We calculated p-values using the following formula: $p=(b+1)/(m+1)$ where b was the

number of expected values higher than the observed value and m, the number of permutations

[31].

## RESULTS

### *Phyllosphere-associated Hyphomicrobiales MAGs mostly belong to Methylobacteriaceae, Lichenihabitantaceae and two undescribed taxa*

Phyllosphere-associated Hyphomicrobiales (syn. Rhizobiales) MAGs were previously identified

in the phyllosphere of Canadian temperate forests (n=22; unpublished data; bioproject

PRJNA1291777), in mosses from the Canadian tundra (n=5; [17]), in lichens from a worldwide

survey (n=43; [37]), in glacier sediments (n=1; [9]) and in mining soils (n=2; [9,23]; Dataset 1).

We refined the taxonomic classification of phyllosphere-associated Hyphomicrobiales MAGs

using a phylogenomic-based approach. Combining the 73 MAGs and 97 reference

Hyphomicrobiales genomes covering the known phylogenomic diversity of phyllosphere-

associated Hyphomicrobiales [12] (Figure 1A; Figure S2; Dataset 1), we identified 10,726 unique

genes (hypothetical protein, repeat and mobile elements excluded). We considered 1,046 genes

that were present in at least 80% of reference genomes (MAGs excluded) as candidate core

genes. After excluding genes that were duplicated or missing data in more than 30% of

assemblies (genomes and MAGs combined), we identified 824 core genes (37 rna and 787

protein-encoding genes; Dataset 3) for which we could retrieve the DNA sequence in at least

70% of genomes and MAGs, and thus infer a phylogenetic tree (RAxML, 1,000 replicated tree, GTRGAMMA model). Our consensus phyllosphere-associated Hyphomicrobiales coalescent tree combining the 824 core gene phylogenetic trees (with ASTRAL-III) was largely congruent with a previous phylogenomic-based classification of Hyphomicrobiales [12], except for the genus *Salinarimonas*, previously classified in Salinarimonadaceae, but embedded within the family Methylobacteriaceae according to our consensus tree. Our tree also provided a few additions for the genera *Lichenifustis*, *Methylocella* and *Hohaiirga*, confirming they belong to the families Lichenihabitantaceae, Beijerinckiaceae and Methylobacteriaceae, respectively. Our consensus tree supported the classification of most phyllosphere-associated Hyphomicrobiales MAGs into five monophyletic groups (branch support =1; Figure 1B, Figure S2). These groups corresponded to the families Methylobacteriaceae (n=18), Lichenihabitantaceae (n=27), Rhabdaerophilaceae (n=1) and to two undescribed taxa that we named according to the previous GTDB-based classification of phyllosphere-associated Hyphomicrobiales MAGs (unpublished data; bioproject PRJNA1291777); namely: *RH-AL1* (n=21) and *JAJXWB01* (n=3). *RH-AL1* included only two reference genomes obtained from strains previously isolated from coal slag [44]. According to our consensus tree, it was a sister but distinct phylogenomic group from Beijerinckiaceae. *JAJXWB01* included no reference genome and was named according to a MAG identified in mine drainage sediments (unpublished data). According to our consensus tree, it was a sister but distinct phylogenomic group from Methylocystaceae and Rhodoblastaceae. No phyllosphere-associated Hyphomicrobiales MAG from related Hyphomicrobiales families could be identified (Alsobacteraceae, Boseaceae, Chelatococcaceae, Roseiarcaceae, Rhodoblastaceae, Methylocystaceae and Beijerinckiaceae). In Methylobacteriaceae, phyllosphere-associated Hyphomicrobiales MAGs from the genera *Enterovirga* (n=7) and *Methylobacterium* (n=11) were identified, but not from *Salinarimonas*, *Microvirga* or *Hohaiivirga*. In Lichenihabitantaceae, phyllosphere-associated Hyphomicrobiales MAGs were previously all assigned to Lichenihabitans according to the GTDB-based

classification, but spanned  all described genera according to our consensus tree, namely

*Lichenihabitans* (n=8), *Lichenifustis* (n=6) and *Lichenibacterium* (n=13).

**Limits and improvements in the classification of phyllosphere-associated**

**Hyphomicrobiales taxa with 16S rRNA metabarcoding**

A maximum-likelihood tree inferred from 24 16S rRNA sequences used to define *1174-901-12* in

the SILVA database, version 138.1, and identified as *Lichenibacterium* in version 138.2,

confirmed that this taxon corresponded to *Lichenibacterium* and to *Lichenifustis* (Figure S1). In

the tree, *1174-901-12* formed a monophyletic and strongly supported group with 16S rRNA

genera *Lichenibacterium* and *Lichenifustis*, but not with their related counterpart *Lichenihabitans*.

We evaluated the power of the 16S rRNA gene as a marker to classify phyllosphere-associated

Hyphomicrobiales taxa. We extracted 92 complete 16S rRNA sequences from reference

phyllosphere-associated Hyphomicrobiales genomes (for which we validated the taxonomy

using a phylogenomic approach; Figure 1A) that could be further used as a reference to refine

phyllosphere-associated Hyphomicrobiales ASV taxonomy. Due to incomplete assemblies, no

complete 16S rRNA gene sequence could be retrieved from most MAGs, but was recovered for

JAJXWB01. A maximum-likelihood tree built from complete 16S rRNA gene sequences shows

the limited power of this gene as a marker to infer the evolutionary history and taxonomy of

phyllosphere-associated Hyphomicrobiales taxa (Figure 2A). In this tree, the families

Methylobacteriaceae and Chelatococcaceae were not monophyletic, but deeply intermingled

together. Within Beijerinckiaceae, the genera *Methylocystis* and *Methylocapsa* were not

mophophyletic. According to the 16S rRNA gene tree, RH_AL1 and JAJXWB01 deeply

branched within families Beijerinckiaceae and Methylocystaceae, respectively (Figure 1A), while

these two undescribed groups formed sister but distinct clades from the two families in our

phylogenomic tree (Figure 2A). We classified the complete 16S rRNA gene sequences using the

SILVA 138.2 database and *assignTaxonomy* R function (Dataset 4). All phyllosphere-associated Hyphomicrobiales families and relatives (Boseaceae, Methylobacteriaceae, Chelatococcaceae, Roseiarcaceae, Rhodoblastaceae, Methylocystaceae, Beijerinckiaceae and Lichenihabitantaceae) were wrongly grouped by SILVA 138.2 in the Beijerinckaceae family, except for the two most basal families Alsobacteraceae and Rhabdaerophillaceae, which were identified as "Hyphomicrobiales *Incertae Sedis*" (Dataset 4). Most phyllosphere-associated Hyphomicrobiales genera were correctly assigned by SILVA 138.2, with few exceptions, mostly regarding genera for which taxonomy was updated after the database was released. RH_AL1 was classified as *Methylocella* and JAJXWB01 as *Methylocystis. Lichenifustis* and *Lichenihabitans* were both identified as *Lichenibacterium. Hohaiivirga* was not identified at the genus level and some *Methylobacterium* sequences were identified as *Methylorubrum* (Dataset 4).

We further explored the worldwide distribution and abundance of phyllosphere-associated Hyphomicrobiales taxa in the phyllosphere and related environments, using metabarcoding data from 13 studies based on the 16S rRNA gene (Dataset 2; Figure 2B). We identified 57,100 unique amplicon sequence variants (ASV) across 1998 samples. We first used SILVA 138.2 as a reference database to classify those ASVs at the order level. We identified 4,637 ASVs classified as Hyphomicrobiales, among which 3,089 were classified in phyllosphere-associated taxa, mostly Beijerinckiaceae (n=3,073). Because of SILVA databases limitations (Dataset 4), we used 92 complete 16 rRNA gene sequences extracted from complete phyllosphere-associated Hyphomicrobiales genomes (n=91) and MAG (n=1) as a database to reclassify the 3,089 ASVs at the family and genus level, according to the phylogenomic consensus tree (Figure 1A), with two exceptions. For family prediction, and according to the 16S rRNA gene phylogeny (Figure 2A), we classified JAJXWB01 within Methylocystaceae and RH_AL1 within Beijerinckiaceae (Dataset 4). Classification with SILVA, version 138.2 and with sequences

extracted from reference genomes resulted in the same genus for 52.22% of ASVs (Dataset 4). Because most genus were classified within Beijerinckiaceae in SILVA database, we could not directly compare both methods for family classification. We thus deduced families in SILVA classification from the genus (e.g. by classifying *Methylobacterium* ASVs in Methylobacteriaceae instead of Beijerinckiaceae) and found that it matched family classification with sequences extracted from reference genomes for 81.57% of ASVs. Family could not be assigned with at least one method for 17.03% of ASVs. Among 1,338 ASVs identified as *Lichenibacterium* with SILVA, v 138.2 (43.31% of ASVs), we confirmed that 1,213 belonged to Lichenihabitantaceae using sequences from reference genomes as a database, and refined their classification in *Lichenibacterium* (701 ASVs), *Lichenifustis* (312), *Lichenihabitans* (121) and unknown Lichenihabitantaceae (79; Dataset 4).

**Lichenihabitantaceae and Methylobacteriaceae are dominant phyllosphere-associated Hyphomicrobiales taxa worldwide according to 16S rRNA metabarcoding**

Worldwide, Pseudomonadota was the most abundant bacteria phylum in the phyllosphere of vascular, root-forming plants (excluding mosses and epiphytic plants), representing on average 56.1% of sequences identified as bacteria, or bacteria diversity [range: 25.1-87.1%] among Bioproject (Dataset 5; Figure 2C) and mostly consisted of Alphaproteobacteria (42.3 [16.7-78.7] % of Bacteria diversity). On average, Hyphomicrobiales dominated Alphaproteobacteria diversity in the phyllosphere (21.6 [6.8-40.6]% of total bacteria diversity) followed by Sphingomonadales (10.6 [2.2-21.0]%) and Acetobacterales (6.8[0.6-15.9%]). Hyphomicrobiales diversity in the phyllosphere was dominated by Lichenihabitantaceae (10.0 [2.0-19.7]% of total bacteria diversity) and by Methylobacteriaceae (5.8[0.7-11.8]% of total bacteria diversity). Other Hyphomicrobiales families, including Rhizobiaceae (2.2 [0.1-14.0]% of total bacteria abundance) and Beijerinckiaceae+RH_AL1 (1.7 [0.0-5.0]% of total bacteria diversity), could be occasionally detected at the surface of tree leaves, but barely Methylocystaceae+JAJXWB01 (0.3 [0.0-0.9]%

of total bacteria diversity). In the phyllosphere, ASVs identified as Methylobacteriaceae mostly corresponded to *Methylobacterium* (5.3 [0.6-11.0]% of total bacteria diversity) and ASVs identified as Lichenihabitantaceae mostly corresponded to *Lichenibacterium* (8.7 [2.0-17.5]% of total bacteria diversity) and *Lichenifustis* (1.1[0.0-3.7]%). ASVs classified as RH_AL1 were barely observed (0.4[0.0-2.0]% of total bacteria diversity) and JAJXWB01 was almost never detected (< 0.1%). On average, Hyphomicrobiales represented a smaller proportion of bacterial diversity in lichen habitats (8.1[7.8-8.4]% of total bacteria diversity) and moss habitats (2.9[2.4-3.5]% of total bacteria diversity) compared to the phyllosphere of vascular plants (21.6 [6.8-40.6]% of total bacteria diversity). In lichen, the Hyphomicrobiales were still dominated by Lichenihabitantaceae lichen (3.6[2.8-4.4]% of total bacteria diversity). Notable proportions of Lichenihabitantaceae were also detected in snow surface (10.6% of total bacteria abundance), the surface of tree bark (1.6%), in epiphytic plants (1.9%), and in aerosols (1.0%). Methylobacteriaceae was one of the dominant bacterial families detected at the surface of photovoltaic panels (30.2% of total bacteria abundance) and was also detected in aerosols (3.4%), on epiphyte plants (2.1%) and on snow surface (1.6%).

**Comparison of 16s rRNA and *rpoB*–based barcoding for the identification of phyllosphere-associated Hyphomicrobiales taxa.**

We explored the potential of the *rpoB* gene, which we previously adapted to metabarcoding to evaluate the diversity of *Methylobacterium* in the phyllosphere, as a broader taxonomic marker for Hyphomicrobiales [19]. We extracted 102 complete *rpoB* sequences from reference phyllosphere-associated Hyphomicrobiales genomes (n=95) and MAGs (n=7), which could be further used as a reference to refine the taxonomy of phyllosphere-associated Hyphomicrobiales ASVs. A maximum-likelihood phylogenetic tree built from complete *rpoB* gene nucleotide sequences shows that this gene is a reliable marker to infer the evolutionary history and

taxonomy of phyllosphere-associated Hyphomicrobiales taxa (Figure S2A). In this tree, all families predicted by the phylogenomic-based consensus tree (Figure 1A) were monophyletic (branch support > 0.90). Within phyllosphere-associated Hyphomicrobiales families (Lichenihabitantaceae, Methylobacteriaceae), all genera predicted by the phylogenomic-based consensus tree (Figure 1A) were monophyletic (branch support > 0.81).

We evaluated the abundance of Hyphomicrobiales taxa from *rpoB* metabarcoding previously developed to target Methylobacteriaceae and performed on 184 leaf samples from two temperate forests in Quebec [19]. ASVs were mostly assigned to Hyphomicrobiales (average per sample; 95.6%; range per forest: 93.8-97.7%) and Caulobacterales (3.9[1.7-5.8]%; Dataset 6). We refined previous taxonomic assignments only for Hyphomicrobiales ASVs, using *rpoB* sequences extracted from reference genomes and MAGs as a reference. In Hyphomicrobiales, ASVs were mostly classified in three families: Methylobacteriaceae (24.6[12.2-39.0%]) - dominated by *Methylobacterium* (23.4[11.1-37.7]%), RH_AL1 (19.8[13.1-25.5]%) and Lichenihabitantaceae (36.7[31.1-41.5]%) - dominated by *Lichenibacterium* (31.9[23.0-39.6]%) and *Lichenifustis* (4.3[1.1-8.0]%). *Enterovirga* (Methylobacteriaceae) and JAJXWB01 could only be detected at very low abundances (0.2[0.1-0.3]% and 0.1[0.0-0.3]%, respectively) (Figure S2B). We compared the relative abundance of Phyllosphere-associated Hyphomicrobiales genera in 41 leaf samples for which metabarcoding was performed with both 16S rRNA and *rpoB* genes, namely *Methylobacterium*, *Enterovirga*, *Lichenifustis*, *Lichenibacterium,* RH_AL1 and JAJXWB01 (Figure S3C). For these taxa, we observed a significant correlation between relative abundances estimated by 16S rRNA gene and *rpoB* gene barcoding ($R^2$= 0.529, p<0.001; Figure S3C). This correlation was not significantly different than expected after 10,000 permutations of relative abundances among leaf samples, within or among sites ($p > 0.05$; Figure S3D), indicating that it was mostly explained by the contrast between highly frequent and

rare taxa (*Lichenibacterium* and *Methylobacterium* vs *Lichenifustis* and *Enterovirga*), consistently observed among leaf samples, and by differences between forests (SBL vs MSH).

**DISCUSSION**

To clarify the distribution and composition of phyllosphere-associated Hyphomicrobiales, we examined over 1,900 phyllosphere 16S rRNA gene datasets from 13 studies spanning multiple continents. This meta-analysis confirmed that *Hyphomicrobiales* constitute a significant fraction of phyllosphere bacterial communities, with the families *Methylobacteriaceae* and *Lichenihabitantaceae* being dominant across environments. Notably, these groups also co-occurred in related aerial habitats such as lichen and moss surfaces, snow, and aerosols, suggesting they may form part of a larger aeroterrestrial biome. The identification of Lichenihabitantaceae, which was solely isolated and characterized in lichen [29], as a dominant taxon of the phyllosphere of trees, has profound implications for understanding microbial colonization dynamics in both the phyllosphere and extreme environments. The ecological distribution of Lichenihabitantaceae on exposed surfaces, leaves and potentially in snow, suggests its resilience to environmental stresses, marking it as a pioneer organism capable of colonizing newly exposed surfaces. This resilience aligns with potential pioneering capacities of *Lichenibacterium*, enabling it to establish communities on surfaces exposed to UV and nutrient scarcity [29]. For instance, metagenomics from lichens confirmed the abundance of *Lichenibacterium* and suggested that these bacteria may play a key role in providing essential nutrients to their fungal symbiotic partners in harsh environments [1]. Furthermore, *Lichenibacterium*'s dominance as an early leaf colonizer implies that it could facilitate subsequent microbial community development, potentially influencing plant health and ecosystem functioning.

Our analyses also highlighted the limitations of the 16S rRNA gene for resolving phyllosphere-associated Hyphomicrobiales diversity. Key families such as *Methylobacteriaceae* appeared polyphyletic based on 16S sequences, and clade-level patterns were often ambiguous or inconsistent across classification schemes, illustrating the low resolution of the 16S rRNA gene in inferring taxonomy and evolutionary history in some bacterial taxa [2,7,14]. We also determined that Lichenihabitantaceae, especially the genera *Lichenifustis* and *Lichenibacterium*, corresponded to *1174-901-12*, an undescribed Hyphomicrobiales taxon frequently associated with the phyllosphere [3,15,19,22,35,43,45]. Despite its abundance, the potential ecological role of *1174-901-12* on the surface of plant leaves remains poorly understood, likely due to the lack of data about this taxon in many reference databases and limitations of marker gene resolution. These issues can lead to overgeneralization or misinterpretation of community structure and ecological patterns in environmental studies.

To overcome these limitations, we applied a genome-resolved phylogenomic approach using core single-copy genes to analyze a representative set of 73 phyllosphere-associated Hyphomicrobiales MAGs alongside 97 reference *Hyphomicrobiales* genomes. This strategy provided a clearer taxonomic framework, revealing that the majority of phyllosphere-associated Hyphomicrobiales MAGs clustered into four robust clades: *Methylobacteriaceae*, *Lichenihabitantaceae*, and two previously undescribed but well-supported lineages—RH-AL1 and JAJXWB01.

RH-AL1 is represented by two reference isolates from coal slag and is related to *Beijerinckiaceae* [44], but forms a distinct and phylogenetically diverse lineage, including MAGs from the phyllosphere, lichens and from mosses. JAJXWB01 appears completely uncultured and forms a related but distinct lineage from *Methylocystaceae*, including MAGs form lichen and from the phyllosphere. Interestingly, RH-AL1 reference isolates, like many members of the Beijerinckiaceae and Methylocystaceae families, are methylotrophic bacteria, a trait shared with

*Methylobacterium* and allowing leaf-associated bacteria to use methanol released by plant as carbon source [38]. However, no MAG-assembled genome was placed within Methylocystaceae or Beijerinckiaceae, suggesting a relatively narrow and structured set of phyllosphere-associated Hyphomicrobiales lineages adapted to phyllosphere environments. A similar narrow ecological distribution was observed for Methylobacteriaceae, with phyllosphere-associated Hyphomicrobiales MAGs only identified within *Enterovirga* and *Methylobacterium*, but not in other genera. *Enterovirga* was traditionally associated with animal feces or aerosols [11,21] , but its detection in the phyllosphere and in lichens suggest a larger ecological niche. In *Methylobacterium*, MAGs were only detected in traditionally leaf-associated phylogroups A, B and D, but not in the most ancestral group C, confirming secondary leaf-association during the evolution of this genus [2,20]. The occurrence of phyllosphere-associated traits should be further investigated in phyllosphere-associated Hyphomicrobiales genera, for instance by screening the occurrence of methylotrophic pathways in phyllosphere-associated Hyphomicrobiales genomes and MAGs. The ability for these bacteria to effectively metabolize one-carbon compounds like methanol should be next validated by culturomics and phenotypic screening, in particular within taxa traditionally not associated with the phyllosphere like Lichenihabitantaceae and *Enterovirga*, or poorly represented in genomic databases and strain collections like RH_AL1 and JAJXWB01.

To complement the phylogenomic analysis of phyllosphere-associated Hyphomicrobiales, we evaluated the utility of the alternative marker gene *rpo*B that we previously used to improve resolution in *Methylobacterium* diversity characterization in the phyllosphere [19]. Unlike the 16S rRNA gene, *rpoB*-based phylogenies recovered phyllosphere-associated Hyphomicrobiales family- and genus-level relationships with much greater accuracy and congruence with genome-based placements. In particular, using *rpoB* amplicon sequencing revealed the strong prevalence of *Methylobacterium* and *Lichenibacterium*, supporting their dominance inferred from metagenomic data. Furthermore, the RH-AL1 clade, which was underrepresented in 16S

datasets, accounted for a large fraction of *Hyphomicrobiales* reads in *rpoB*-based profiles, of roughly the same abundance as *Methylobacterium* and *Lichenibacterium*, highlighting the rpoB marker's superior sensitivity and specificity for uncovering cryptic diversity in phyllosphere-associated Hyphomicrobiales.

Our study provides a revised and integrative view of phyllosphere-associated *Hyphomicrobiales*, revealing both the global dominance of known families such as *Methylobacteriaceae* and *Lichenihabitantaceae*, and the existence of undescribed but recurrent clades. We show that relying solely on the 16S rRNA gene obscures key phylogenetic and ecological patterns of bacterial communities in the phyllosphere, particularly due to its limited resolution and lack of data in reference databases. By integrating *rpoB* markers and genome-resolved phylogenetics, we propose a more robust framework to track phyllosphere-associated Hyphomicrobiales diversity and guide future ecological and evolutionary studies. Looking forward, the cultivation and characterization of the novel RH-AL1 and JAJXWB01 lineages, as well as a better delineation of the genera within *Lichenihabitantaceae*, are important research priorities. Combining long-read metagenomics, transcriptomics, and targeted functional assays will further allow us to link genome evolution to ecological adaptation, helping to uncover the roles these widespread Hyphomicrobiales play in the phyllosphere and beyond.

**Funding information –** JBL and SWK were supported by NSERC Discovery Grants RGPIN-2024-05825 and RGPIN-2025-05608 respectively.

**Acknowledgments**: NIngeborg Klarenberg and Wisnu Adi Wicaksono for providing Metadata; Marta Alonso Garcia and Zihui Wang for discussion.

**Data availability:** Code and data, including Datasets 1-6, are publicly available on Github repository: JBLED/Hyphomicrobiales-phyllosphere

**Figure Legend**

**Figure 1: Phyllosphere-associated Hyphomicrobiales MAGs mostly belong to
Methylobacteriaceae, Lichenihabitantaceae and to RH_AL1, an undescribed taxon related
to Beijerinckiaceae.** A) Phylogenomic tree of 96 reference genomes (NCBI ref_seq genomes)
and 73 MAGs (tree leaves, moss and lichen). The consensus evolutionary tree was inferred from
824 core gene phylogenies combined with ASTRAL-III (See Figure S2 for detailed tree). The
tree was rooted on *Rhabdaerophilum*. Scale represents coalescent time. Branch support
represents local posterior probability. B) Distribution of tree leaf-, moss- and lichen-associated
MAGs per Hyphomicrobiales genus. Each oval represents a MAG.

**Figure 2: Phyllosphere is dominated worldwide by Methylobacteriaceae and
Lichenihabitantaceae according to 16S rRNA gene metabarcoding.** A) Maximum-Likelyhood
tree (1,000 bootstrap, GTRGAMMA model) from complete 16S rRNA gene sequences extracted
from 96 phyllosphere-associated Hyphomicrobiales reference genomes (Fig 1). RH_AL1 is
embedded within Beijerinckiaceae and JAJXWBA01 in Methylocystaceae. Methylobacteriaceae,
Chelatococcaceae as well as some genera (*Chelatococcus*, *Methylocystis* and *Methylocapsa*)
appear polyphyletic. B) World map with BioProject origin (see **Dataset 2** for details). C) Bacterial
abundance according to 16S rRNA barcoding, focusing on phyllosphere-associated
Hyphomicrobiales families, in 13 BioProjects. Taxonomy from SILVA, refined with sequences
from phyllosphere-associated Hyphomicrobiales reference genomes (see Fig. 1). Bioprojects
were divided according to different climatic areas or substrates when applicable. For each
category, abundances are shown for main taxonomic subdivisions other than Hyphomicrobiales
(other Bacteria, Pseudomonadota and Alphaproteobacteria; left pie charts) and detailed for
major families within Hyphomicrobiales and for all genera within Methylobacteriaceae and

Lichenihabitantaceae (taxa with maximum relative abundance lower than 1% not showed, including Methylocystaceae and JAJXWBA01; right pie charts).

**References**

[1]      Ahmad, N., Ritz, M., Calchera, A., Otte, J., Schmitt, I., Brueck, T., Mehlmer, N. (2023) Biosynthetic Potential of Hypogymnia Holobionts: Insights into Secondary Metabolite Pathways. Journal of Fungi 9(5), 546, Doi: 10.3390/jof9050546.

[2]      Alleman, A.B., Stolyar, S., Marx, C.J., Leducq, J.-B. (2025) Led astray by 16S rRNA: phylogenomics reaffirms the monophyly of Methylobacterium and lack of support for Methylorubrum as a genus. ISME J 19(1), wraf011, Doi: 10.1093/ismejo/wraf011.

[3]      Ares, A., Pereira, J., Garcia, E., Costa, J., Tiago, I. (2021) The Leaf Bacterial Microbiota of Female and Male Kiwifruit Plants in Distinct Seasons: Assessing the Impact of Pseudomonas syringae pv. actinidiae. Phytobiomes Journal 5(3), 275–87, Doi: 10.1094/PBIOMES-09-20-0070-R.

[4]      Aschenbrenner, I.A., Cernava, T., Erlacher, A., Berg, G., Grube, M. (2017) Differential sharing and distinct co-occurrence networks among spatially close bacterial microbiota of bark, mosses and lichens. Molecular Ecology 26(10), 2826–38, Doi: 10.1111/mec.14070.

[5]      Aziz, R.K., Bartels, D., Best, A.A., DeJongh, M., Disz, T., Edwards, R.A., Formsma, K., Gerdes, S., Glass, E.M., Kubal, M., Meyer, F., Olsen, G.J., Olson, R., Osterman, A.L., Overbeek, R.A., McNeil, L.K., Paarmann, D., Paczian, T., Parrello, B., Pusch, G.D., Reich, C., Stevens, R., Vassieva, O., Vonstein, V., Wilke, A., Zagnitko, O. (2008) The RAST Server: Rapid Annotations using Subsystems Technology. BMC Genomics 9(1), 75, Doi: 10.1186/1471-2164-9-75.

[6]      Bai, Y., Müller, D.B., Srinivas, G., Garrido-Oter, R., Potthoff, E., Rott, M., Dombrowski, N., Münch, P.C., Spaepen, S., Remus-Emsermann, M., Hüttel, B., McHardy, A.C., Vorholt, J.A., Schulze-Lefert, P. (2015) Functional overlap of the Arabidopsis leaf and root microbiota. Nature

528(7582), 364–9, Doi: 10.1038/nature16192.

[7]     Bartoš, O., Chmel, M., Swierczková, I. (2024) The overlooked evolutionary dynamics of 16S rRNA revises its role as the "gold standard" for bacterial species identification. Sci Rep 14(1), 9067, Doi: 10.1038/s41598-024-59667-3.

[8]     Bodenhofer, U., Bonatesta, E., Horejš-Kainrath, C., Hochreiter, S. (2015) msa: an R package for multiple sequence alignment. Bioinformatics 31(24), 3997–9, Doi: 10.1093/bioinformatics/btv494.

[9]     Busi, S.B., Bourquin, M., Fodelianakis, S., Michoud, G., Kohler, T.J., Peter, H., Pramateftaki, P., Styllas, M., Tolosano, M., De Staercke, V., Schön, M., de Nies, L., Marasco, R., Daffonchio, D., Ezzat, L., Wilmes, P., Battin, T.J. (2022) Genomic and metabolic adaptations of biofilms to ecological windows of opportunity in glacier-fed streams. Nat Commun 13(1), 2168, Doi: 10.1038/s41467-022-29914-0.

[10]    Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., Holmes, S.P. (2016) DADA2: High resolution sample inference from Illumina amplicon data. Nat Methods 13(7), 581–3, Doi: 10.1038/nmeth.3869.

[11]    Chen, X., Li, Q.-Y., Li, G.-D., Lei, H., Jiang, Y., Han, L., Huang, X.-S., Jiang, C.-L. (2017) Enterovirga rhinocerotis gen. nov., sp. nov., isolated from Rhinoceros unicornis faeces. Antonie Van Leeuwenhoek 110(4), 553–62, Doi: 10.1007/s10482-016-0823-1.

[12]    diCenzo, G.C., Yang, Y., Young, J.P.W., Kuzmanović, N. (2024) Refining the taxonomy of the order Hyphomicrobiales (Rhizobiales) based on whole genome comparisons of over 130 type strains. Int J Syst Evol Microbiol 74(4), 006328, Doi: 10.1099/ijsem.0.006328.

[13]    Dixon, P. (2003) VEGAN, a package of R functions for community ecology. Journal of Vegetation Science 14(6), 927–30, Doi: 10.1111/j.1654-1103.2003.tb02228.x.

[14]    Hassler, H.B., Probert, B., Moore, C., Lawson, E., Jackson, R.W., Russell, B.T., Richards, V.P. (2022) Phylogenies of the 16S rRNA gene and its hypervariable regions lack concordance with core genome phylogenies. Microbiome 10(1), 104, Doi: 10.1186/s40168-022-

01295-y.

[15]     Herrmann, M., Geesink, P., Richter, R., Küsel, K. (2021) Canopy Position Has a Stronger Effect than Tree Species Identity on Phyllosphere Bacterial Diversity in a Floodplain Hardwood Forest. Microb Ecol 81(1), 157–68, Doi: 10.1007/s00248-020-01565-y.

[16]     Hördt, A., López, M.G., Meier-Kolthoff, J.P., Schleuning, M., Weinhold, L.-M., Tindall, B.J., Gronow, S., Kyrpides, N.C., Woyke, T., Göker, M. (2020) Analysis of 1,000+ Type-Strain Genomes Substantially Improves Taxonomic Classification of Alphaproteobacteria. Frontiers in Microbiology 11, 468, Doi: 10.3389/fmicb.2020.00468.

[17]     Ishak, S., Rondeau-Leclaire, J., Faticov, M., Roy, S., Laforest-Lapointe, I. (2024) Boreal moss-microbe interactions are revealed through metagenome assembly of novel bacterial species. Sci Rep 14(1), 22168, Doi: 10.1038/s41598-024-73045-z.

[18]     Klarenberg, I.J., Keuschnig, C., Russi Colmenares, A.J., Warshan, D., Jungblut, A.D., Jónsdóttir, I.S., Vilhelmsson, O. (2022) Long-term warming effects on the microbiome and nifH gene abundance of a common moss species in sub-Arctic tundra. New Phytologist 234(6), 2044–56, Doi: 10.1111/nph.17837.

[19]     Leducq, J.-B., Seyer-Lamontagne, É., Condrain-Morel, D., Bourret, G., Sneddon, D., Foster, J.A., Marx, C.J., Sullivan, J.M., Shapiro, B.J., Kembel, S.W. (2022) Fine-Scale Adaptations to Environmental Variation and Growth Strategies Drive Phyllosphere Methylobacterium Diversity. mBio 13(1), e03175-21, Doi: 10.1128/mbio.03175-21.

[20]     Leducq, J.-B., Sneddon, D., Santos, M., Condrain-Morel, D., Bourret, G., Martinez-Gomez, N.C., Lee, J.A., Foster, J.A., Stolyar, S., Shapiro, B.J., Kembel, S.W., Sullivan, J.M., Marx, C.J. (2022) Comprehensive Phylogenomics of Methylobacterium Reveals Four Evolutionary Distinct Groups and Underappreciated Phyllosphere Diversity. Genome Biology and Evolution 14(8), evac123, Doi: 10.1093/gbe/evac123.

[21]     Lee, H., Chaudhary, D.K., Kim, D.-U. (2024) Enterovirga aerilata sp. nov. and Knoellia koreensis sp. nov., isolated from an automobile air conditioning system. International Journal of

Systematic and Evolutionary Microbiology 74(5), 006383, Doi: 10.1099/ijsem.0.006383.

[22]    Long, J., Luo, W., Xie, J., Yuan, Y., Wang, J., Kang, L., Li, Y., Zhang, Z., Hong, M. (2021) Environmental Factors Influencing Phyllosphere Bacterial Communities in Giant Pandas' Staple Food Bamboos. Front. Microbiol. 12, Doi: 10.3389/fmicb.2021.748141.

[23]    Ma, B., Lu, C., Wang, Y., Yu, J., Zhao, K., Xue, R., Ren, H., Lv, X., Pan, R., Zhang, J., Zhu, Y., Xu, J. (2023) A genomic catalogue of soil microbiomes boosts mining of biodiversity and genetic resources. Nat Commun 14(1), 7318, Doi: 10.1038/s41467-023-43000-z.

[24]    Maki, T., Hosaka, K., Lee, K.C., Kawabata, Y., Kajino, M., Uto, M., Kita, K., Igarashi, Y. (2023) Vertical distribution of airborne microorganisms over forest environments: A potential source of ice-nucleating bioaerosols. Atmospheric Environment 302, 119726, Doi: 10.1016/j.atmosenv.2023.119726.

[25]    Moura, J.B., Delforno, T.P., do Prado, P.F., Duarte, I.C. (2021) Extremophilic taxa predominate in a microbial community of photovoltaic panels in a tropical region. FEMS Microbiology Letters 368(16), fnab105, Doi: 10.1093/femsle/fnab105.

[26]    Noh, H.-J., Baek, K., Hwang, C.Y., Shin, S.C., Hong, S.G., Lee, Y.M. (2019) Lichenihabitans psoromatis gen. nov., sp. nov., a member of a novel lineage (Lichenihabitantaceae fam. nov.) within the order of Rhizobiales isolated from Antarctic lichen. Int J Syst Evol Microbiol 69(12), 3837–42, Doi: 10.1099/ijsem.0.003695.

[27]    Ogier, J.-C., Pagès, S., Galan, M., Barret, M., Gaudriault, S. (2019) rpoB, a promising marker for analyzing the diversity of bacterial communities by amplicon sequencing. BMC Microbiol 19(1), 171, Doi: 10.1186/s12866-019-1546-z.

[28]    Pace, N.R. (1997) A molecular view of microbial diversity and the biosphere. Science 276(5313), 734–40, Doi: 10.1126/science.276.5313.734.

[29]    Pankratov, T.A., Grouzdev, D.S., Patutina, E.O., Kolganova, T.V., Suzina, N.E., Berestovskaya, J.J. (2020) Lichenibacterium ramalinae gen. nov, sp. nov., Lichenibacterium minor sp. nov., the first endophytic, beta-carotene producing bacterial representatives from

lichen thalli and the proposal of the new family Lichenibacteriaceae within the order Rhizobiales. Antonie van Leeuwenhoek 113(4), 477–89, Doi: 10.1007/s10482-019-01357-6.

[30]    Pankratov, T.A., Samylina, O.S., Tikhonova, E.N., Ianutsevich, E.A., Avtukh, A.N., Lee, Y.M. (2023) A novel bacteriobiont of the Arctic lichen Flavocetraria nivalis, Lichenifustis flavocetrariae gen. nov, sp. nov. demonstrating hydrolytic properties and containing a full set of the Calvin-Benson-Bassham cycle genes. Arch Microbiol 205(6), 232, Doi: 10.1007/s00203-023-03577-x.

[31]    Phipson, B., Smyth, G.K. (2010) Permutation p-values should never be zero: calculating exact p-values when permutations are randomly drawn. Statistical Applications in Genetics and Molecular Biology 9(1), Doi: 10.2202/1544-6115.1585.

[32]    Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., Glöckner, F.O. (2013) The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. Nucleic Acids Res 41(D1), D590–6, Doi: 10.1093/nar/gks1219.

[33]    Romani, M., Carrion, C., Fernandez, F., Intertaglia, L., Pecqueur, D., Lebaron, P., Lami, R. (2019) High bacterial diversity in pioneer biofilms colonizing ceramic roof tiles. International Biodeterioration & Biodegradation 144, 104745, Doi: 10.1016/j.ibiod.2019.104745.

[34]    Sierra, M.A., Danko, D.C., Sandoval, T.A., Pishchany, G., Moncada, B., Kolter, R., Mason, C.E., Zambrano, M.M. (2020) The Microbiomes of Seven Lichen Genera Reveal Host Specificity, a Reduced Core Community and Potential as Source of Antimicrobials. Front. Microbiol. 11, Doi: 10.3389/fmicb.2020.00398.

[35]    Siqueira, J.S., de Carvalho, L.A.L., Santos, C.H.B., Frezarin, E.T., Pinheiro, D.G., Nicodemo, D., Desoignies, N., Rigobelo, E.C. (2024) Influence of Growth Support on the Diversity, Composition, and Functionality of Microbial Communities Associated with Tillandsia recurvata. Microb Ecol 87(1), 129, Doi: 10.1007/s00248-024-02448-2.

[36]    Stamatakis, A. (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. Bioinformatics 30(9), 1312–3, Doi: 10.1093/bioinformatics/btu033.

[37]     Tagirdzhanova, G., Saary, P., Cameron, E.S., Allen, C.C.G., Garber, A.I., Escandón, D.D., Cook, A.T., Goyette, S., Nogerius, V.T., Passo, A., Mayrhofer, H., Holien, H., Tønsberg, T., Stein, L.Y., Finn, R.D., Spribille, T. (2024) Microbial occurrence and symbiont detection in a global sample of lichen metagenomes. PLoS Biol 22(11), e3002862, Doi: 10.1371/journal.pbio.3002862.

[38]     Tani, A., Masuda, S., Fujitani, Y., Iga, T., Haruna, Y., Kikuchi, S., Shuaile, W., Lv, H., Katayama, S., Yurimoto, H., Sakai, Y., Kato, J. (2023) Metabolism-linked methylotaxis sensors responsible for plant colonization in Methylobacterium aquaticum strain 22A. Front. Microbiol. 14, Doi: 10.3389/fmicb.2023.1258452.

[39]     Touchette, D., Gostinčar, C., Whyte, L.G., Altshuler, I. (2023) Lichen-associated microbial members are prevalent in the snow microbiome of a sub-arctic alpine tundra. FEMS Microbiology Ecology 99(12), fiad151, Doi: 10.1093/femsec/fiad151.

[40]     Vorholt, J.A. (2012) Microbial life in the phyllosphere. Nature Reviews Microbiology 10(12), 828–40, Doi: 10.1038/nrmicro2910.

[41]     Wang, H.-C., Huang, M.-H., Guo, D.-Y., He, W., Wang, L., Fu, Z.-Y., Li, W.-J., Zhang, A.H., Zhang, D.-F. (2024) Hohaiivirga grylli gen. nov., sp. nov., a New Member of the Family Methylobacteriaceae, Isolated from Cricket (Gryllus chinensis). Curr Microbiol 81(11), 392, Doi: 10.1007/s00284-024-03922-3.

[42]     Wang, S., Meade, A., Lam, H.-M., Luo, H. (2020) Evolutionary Timeline and Genomic Plasticity Underlying the Lifestyle Diversity in Rhizobiales. mSystems 5(4), e00438-20, /msystems/5/4/msys.00438-20.atom, Doi: 10.1128/mSystems.00438-20.

[43]     Wang, Z., Jiang, Y., Zhang, M., Chu, C., Chen, Y., Fang, S., Jin, G., Jiang, M., Lian, J.-Y., Li, Y., Liu, Y., Ma, K., Mi, X., Qiao, X., Wang, X., Wang, X., Xu, H., Ye, W., Zhu, L., Zhu, Y., He, F., Kembel, S.W. (2023) Diversity and biogeography of plant phyllosphere bacteria are governed by latitude-dependent mechanisms. New Phytologist 240(4), 1534–47, Doi: 10.1111/nph.19235.

[44]    Wegner, C.-E., Gorniak, L., Riedel, S., Westermann, M., Küsel, K. (2019) Lanthanide-Dependent Methylotrophs of the Family Beijerinckiaceae: Physiological and Genomic Insights. Appl Environ Microbiol 86(1), e01830-19, Doi: 10.1128/AEM.01830-19.

[45]    Wicaksono, W.A., Morauf, C., Müller, H., Abdelfattah, A., Donat, C., Berg, G. (2023) The mature phyllosphere microbiome of grapevine is associated with resistance against Plasmopara viticola. Front. Microbiol. 14, Doi: 10.3389/fmicb.2023.1149307.

[46]    Zhang, C., Rabiee, M., Sayyari, E., Mirarab, S. (2018) ASTRAL-III: polynomial time species tree reconstruction from partially resolved gene trees. BMC Bioinformatics 19(6), 153, Doi: 10.1186/s12859-018-2129-y.

[47]    Zhao, Q., Liu, W., Li, Y., Ke, M., Qu, Q., Yuan, W., Pan, X., Qian, H. (2020) Enantioselective effects of imazethapyr residues on *Arabidopsis thaliana* metabolic profile and phyllosphere microbial communities. Journal of Environmental Sciences 93, 57–65, Doi: 10.1016/j.jes.2020.03.009.

# Phylogenomic resolution of phyllosphere-associated Hyphomicrobiales
824 core genes from 97 genomes + 73 metagenome-assembled genomes (MAGs)

**Methylobacteriaceae**
- ***Methylobacterium*** *
- *Enterovirga*
- *Salinarimonas*
- *Microvirga*
- *Hohaiivirga*

**Model taxon***

Chelatococcaceae
Boseaceae
Alsobacteraceae

### Source of MAGs

Tree leaves (22)   Lichen (43)   Moss (5)   Other (3)

**Lichenihabitantaceae**
- *Lichenibacterium*
- *Lichenihabitans*
- *Lichenifustis*

- **Recently described from lichen** (few reference genomes)
- Identified as «1174-901-12» in SILVA database (abundant in the phyllosphere)

Methylocystaceae
- JAJXWB01
Rhodoblastaceae

- **Undescribed**
- Metagenomics only (sediments)

Beijerinckiaceae
- RH AL1
Roseiarcaceae

- **Undescribed**, only 2 genomes (coal slag)
- Barely detected by 16S rRNA barcoding

**Non-model taxa**

Other Hyphomicrobiales
Other Alphaproteobacteria

## The diversity we capture
16S rRNA gene barcoding
+ current databases (SILVA)
(13 BioProjects; >1900 samples)

## The diversity we expect
metagenomics + refined databases
(culturomics + phylogenomics)