# Reexamining the evolutionary history of the mammalian medial pterygoid muscle

Julia A. Schultz[1,*], Lucas N. Weaver[2,3], Kai R. K. Jäger[1], David M. Grossnickle[4,*]

[1]Section Palaeontology, Bonner Institute of Organismic Biology, Rheinische Friedrich-Wilhelms-Universität Bonn, Nussallee 8, D-53115 Bonn, Germany

[2]Museum of Paleontology, University of Michigan, Ann Arbor, MI 48109, USA

[3]Department of Earth and Environmental Sciences, University of Michigan, Ann Arbor, MI 48109, USA

[4]Natural Sciences Department, Oregon Institute of Technology, Klamath Falls, OR 97601, USA

*Correspondence: jaschultz@uni-bonn.de, david.grossnickle@oit.edu

26                                        ABSTRACT

In non-mammalian synapsids, feeding and hearing are closely linked because some jaw bones are

28     involved in both functions. The evolutionary decoupling of these two systems in early mammals

likely catalyzed greater specialization of feeding and hearing. Although fossil osteological changes

30     during this process are well documented, the corresponding evolutionary changes to soft tissue

anatomy are less certain. The medial pterygoid muscle is a jaw adductor that is central to this

32     evolutionary transition because in many fossil lineages it inserted near or possibly on jaw bones

involved in both feeding and hearing. In therians (placentals and marsupials), the medial pterygoid

34     muscle develops medial to Meckel's cartilage and inserts on the mandibular angular process.

Similarly, non-mammalian cynodonts are often reconstructed with a medial pterygoid muscle

36     passing medial to the ossified Meckel's cartilage, inserting on the dentary 'angular' (i.e.,

pseudangular) process. Thus, the traditional interpretation is that the medial pterygoid remained

38     medial to Meckel's cartilage through the evolutionary transitions from early cynodonts to therians.

Here we highlight issues with that interpretation: the medial pterygoid muscle inserts lateral (not

40     medial) to Meckel's cartilage in monotremes and, presumably, early mammal groups (e.g.,

spalacotherioids) that lacked an angular process. This suggests at least two possible explanatory

42     hypotheses: 1) the medial pterygoid muscle is evolutionarily labile, shifting in position relative to

Meckel's cartilage multiple times or 2) the medial pterygoid muscle did not insert on the

44     pseudangular process of non-mammalian cynodonts and instead inserted on the mandibular medial

ridge, dorsal to Meckel's cartilage. We advocate for the latter hypothesis, proposed by Patterson

46     (1956), which suggests that the medial pterygoid did not shift medial to Meckel's cartilage until the

complete separation of the ear and jaw in cladotherians (therians and close relatives), with the shift

48     in position possibly triggering the evolution of the therian angular process as an insertion site.

Patterson's hypothesis is in line with a growing body of evidence that indicate concomitant

50     evolutionary changes of muscles, ears, and jaws at the cladotherian node were important catalysts

for the evolution of hearing and feeding specializations in extant mammals.

**Key words** (5 to 10): Mastication, Jaw adductor muscles, Pterygoideus muscle, Mammaliaformes,

54    Mammalia, Cynodontia, Synapsida, Meckel's cartilage, Macroevolution, Mandibular angular process.

INTRODUCTION

Transformation of the jaw bones of non-mammalian synapsids to middle-ear ossicles of mammals is

60    among the best-documented evolutionary transitions in the fossil record, and it likely catalyzed

improved feeding and hearing abilities in early Mammalia (Patterson 1956, Rosowski 1992, Luo

62    2011, Anthwal et al. 2013, Maier and Ruf 2016, Luo et al. 2016, Grossnickle 2017, Bhullar et al. 2019,

Luo and Manley 2020, Mao et al. 2020, Mao and Meng 2020, Grossnickle et al. 2022, Schultz 2025).

64    Concomitant with those osteological changes—shrinking, repositioning, and ultimately detaching of

bony elements in the jaw into the middle-ear cavity of the skull—were changes in size, position, and

66    direction of contraction of the muscles that facilitated precise tooth-to-tooth occlusion and complex

chewing (Fig. 1; e.g., Crompton 1995, Luo and Manley 2020, Grossnickle et al. 2022). The jaw

68    adductor (or elevator) muscles facilitate mammalian chewing, and they include the temporalis,

masseter, and pterygoid muscles. Changes to the temporalis and masseter muscles are well

70    traceable through synapsid (mammalian total group) evolution because they are relatively large and

their positions are often associated with unambiguous bony features of the synapsid skull, such as

72    the sagittal crest, zygomatic arch, and masseteric fossa (Crompton 1963, DeMar and Barghusen

1972, Bramble 1978, Lautenschlager et al. 2017). The evolutionary history of the medial pterygoid

74    muscle (musculus pterygoideus internus or internal pterygoid), however, is less clear. This includes

uncertainty about when in the evolutionary history of synapsids the pterygoideus muscle gave rise

76    to the medial and lateral pterygoid muscles.

**Figure 1.** *A*, Summary of the anatomical relationship of the medial pteryoid muscle (MP) and Meckel's cartilage (MC) during the early evolution of mammals. Medial pterygoid muscle positions are approximate reconstructions. Meckel's cartilage of cladotherians is illustrated with a dashed line because it either does not ossify or, if ossified, is not medially displaced like those of eutriconodontans and spalacotherioids (e.g., Wang et al. 2022). Early crown mammal such as spalacotherioids and eutriconodontans have medially displaced ossified Meckel's cartilages (or 'prearticulars'; Kermack et al. 1973, Wang et al. 2001, Meng et al. 2003, Martin et al. 2015, Mao et al. 2020). Jaw images are of the left hemimandible in medial and posterior view. *B*, Ventral view of the skull of *Repenomamus* (IVPP V12549), modified from Wang et al. (2003), showing the medial separation of ossified Meckel's cartilage from the dentary. *C*, Reconstruction of the ventral portions of the medial and lateral pterygoid muscles in *Priacodon*.

The pterygoideus musculature is linked to the evolution of both the mammalian masticatory and auditory systems because, ancestrally, it likely attached to the postdentary bones of early cynodonts that were involved in both feeding and hearing prior to the separation of the mammalian middle ear (Fig. 1) (e.g., Parrington 1946; Crompton 1963). In extant therians, the medial pterygoid

muscle contributes to several types of jaw movements, including yaw (rotation around a

98    dorsoventral axis), roll (rotation around an anteroposterior axis), and lateral translation (Herring

and Scapino 1973, Hiiemae and Crompton 1985, Weijs 1980, Weijs 1994, Crompton 1995, Herring

100    2003, Crompton et al. 2010, Grossnickle 2017, Bhullar et al. 2019, Grossnickle et al. 2022). Prior to

the evolution of Cladotheria (therians and their close relatives), however, the pterygoid muscles

102    may have primarily assisted with stabilization of the jaw during mastication rather than facilitating

rotational movements (Crompton 1995) because most synapsids likely relied on relatively simple

104    orthal or palinal jaw motions (i.e., with very limited or no yaw, roll, or transverse movements;

Grossnickle et al. 2022). In early cladotherians, the medial pterygoid muscle was likely coopted to

106    facilitate more complex anterior and transverse occlusal movements (e.g., Grossnickle et al. 2022

and citations therein), concomitant with the evolution of a posteriorly positioned angular process

108    (Grossnickle 2017) on which the medial surface served as an insertion site for the medial pterygoid

muscle.

110    The medial and lateral pterygoid muscles likely derived from the sauropsid musculus

pterygoideus (termed pterygoideus muscle hereafter; Romer and Price 1940, Watson 1953);

112    however, others have proposed that the pterygoideus muscle is homoplastic among tetrapods

(Iordansky 2010), and it may have also given rise to additional mammalian muscles beyond the

114    medial and lateral pterygoids, such as the tensor veli palatini and tensor tympani (Barghusen 1986,

Crompton et al. 2018). Further uncertainties about the evolutionary history of the pterygoid

116    muscles are related to the phylogenetic appearance of the split of the single pterygoideus muscle

into multiple pterygoid muscles, and where those muscles inserted on the jaw. Previous studies

118    alternatively proposed that the medial and lateral pterygoids: (i) arose in early synapsids from a

muscle complex (anterior and posterior pterygoideus of Squamata) attaching to the angular bone

120    (Barghusen 1973, King 1981), (ii) evolved in early therapsids from the pterygoideus muscle and

attached to the articular/ossified Meckel's cartilage (e.g., synonymous with the term 'prearticular'

122    following Luo and Manley 2020) (Parrington 1955, Crompton 1963, Lautenschlager et al. 2017), or

(iii) evolved relatively later in synapsid history (likely in late 'cynodonts' but possibly as late as

124      Cladotheria) from the pterygoideus muscle and attached to the dentary (Patterson 1956, Bramble

1978, Barghusen 1986, Crompton 1995). Just like with the insertion sites, there is uncertainty and

126      debate on the cranial origin sites of the pterygoid muscles of synapsids (e.g., Barghusen 1973,

Lautenschlager et al. 2017, Crompton 2018). However, compared to jaw elements, the dearth of

128      well-preserved basicranial material in the fossil record makes questions concerning the origin sites

even more challenging to examine and unlikely to be resolved at this time.

130          Most relevant to the evolution of the mammalian chewing and hearing systems is the

evolution of pterygoid musculature among cynodonts. There is a general consensus that the

132      pterygoideus musculature shifted from inserting on the postdentary bones to the dentary (see

Lautenschlager et al. 2017 for a review) as the postdentary bones reduce in size in the lineage

134      leading from early cynodonts (e.g., *Thrinaxodon*) to mammaliaforms (Crompton 1963A, Allin and

Hopson 1992, Sidor 2001). Nonetheless, it is unclear whether that shift occurred early in the

136      evolutionary history of cynodonts (e.g., Crompton 1963B, Kemp 1972), later in the evolutionary

history of mammaliamorphans (Bramble 1978, Crompton 1995, Lautenschlager et al. 2017), or

138      among crown mammals (Patterson 1956).

         The posteroventral process of the non-therian cynodont dentary—alternatively referred to

140      as the 'angular process' or 'pseudangular process' by different authors (and hereafter we refer to it

as the pseudangular process)—is central to these debates. The pseudangular process first appears

142      among eucynodontian cynodonts and is also present on the dentaries of non-mammalian

mammaliaforms. Because the pseudangular process is roughly similar in position and shape to the

144      therian angular process (with the only major difference being the more anterior position of the

pseudangular process; Grossnickle 2017), it is often hypothesized that the mammaliaform and

146      therian angular processes are homologous (e.g., Rougier et al. 2015, Abdala and Damiani 2004).

Because the medial pterygoid attaches to the medial side of the angular process in extant therians,

148      the insertion site of the medial pterygoid in that evolutionary scenario remained on the 'angular'

process from early eucynodontians through therians (Adams 1918, Simpson 1928, Crompton 1963,

150     Kemp 1972, Lautenschlager et al. 2017). That the mammal-like division of the pterygoideus muscle

into the medial and lateral pterygoid muscles, with the medial pterygoid muscle inserting on the

152     dentary, is generally assumed to have occurred by the origin of Mammaliaformes (e.g., Kermack and

Mussett 1981) or potentially earlier among Eucynodontia (Lautenschlager et al. 2017) supports that

154     interpretation. Consequently, the early mammaliaform medial pterygoid muscle, originating on the

basicranium, is commonly reconstructed as passing medial to Meckel's cartilage and the

156     postdentary bones to insert on the 'angular' process (e.g., see the *Morganucodon* reconstruction in

Figure 1 and Kermack and Mussett 1981). Developmental data from extant therians further support

158     that view—the medial pterygoid muscle also inserts medial to Meckel's cartilage early in the

ontogenetic development of extant therians (Brocklehurst et al. 2015, Sánchez-Villagra and

160     Forasiepi 2017).

Nonetheless, the evolutionary scenario that the medial pterygoid passes medial to Meckel's

162     from early mammaliaforms to therians is complicated by recent fossil discoveries and phylogenetic

analyses demonstrating that several major early mammal clades phylogenetically intermediate to

164     early mammaliaforms and cladotherians (e.g., eutriconodontans, spalacotherioids,

multituberculates) lack an angular process (Fig. 1A; Krause et al. 2014, Krause et al. 2020, Rougier

166     et al. 2011, Rougier et al. 2012, Huttenlocker et al. 2017, Luo et al. 2017). These findings suggest that

there is an evolutionarily intermediate jaw morphology that lacks the classically invoked insertion

168     location for the medial pterygoid muscle (Fig. 1; Patterson 1956, Patterson and Olson 1961,

Grossnickle 2017, Grossnickle 2020, Grossnickle et al. 2022). Thus, we support the interpretation

170     that the pseudangular process is unlikely to be homologous with the therian angular process that

evolved at Cladotheria (Patterson 1956, Jenkins et al. 1983, Gow 1986, Grossnickle 2017).

172     Because some early mammal groups lack an angular process, it is worth reconsidering the

evolution of the medial pterygoid and its position relative to Meckel's cartilage. Although the medial

174     pterygoid muscle is traditionally reconstructed as passing medial to Meckel's cartilage in non-

mammalian synapsids (Fig. 1A), consistent with the early ontogeny of therians (Brocklehurst et al.

176   2015, Sánchez-Villagra and Forasiepi 2017), Meckel's cartilage runs very close to the dentary bone

in both of those cases (Fig. 1; Fig. 2G-I). Early mammal groups like eutriconodonts and

178   spalacotherioids, however, exhibit a medially inflected Meckel's cartilage that tethers otherwise-

detached middle-ear ossicles to the body of the dentary (Fig. 1B; Wang et al. 2003, Luo et al. 2007,

180   Mao et al. 2020). For the medial pterygoid muscle to insert medial to Meckel's cartilage in these

early mammals, it must have draped around the medially detached middle-ear bones to insert in the

182   deep pterygoid fossa and/or on the 'pterygoid shelf' (Sànchez-Villagra and Smith 1997; or

'pterygoid crest' of Simpson 1926); such an arrangement is very unlikely because it would have both

184   hindered the vibratory capabilities of the middle-ear complex and left an anomalous wide space

empty between the middle-ear bones and the pterygoid fossa. Thus, the medial pterygoid in early

186   mammals was likely positioned lateral to Meckel's cartilage (Fig. 1B), providing space for the

suspended middle-ear elements.

188        In addition to early mammals, monotremes also exhibit a medial pterygoid muscle (or

pterygoideus muscle) that passes lateral to Meckel's cartilage (i.e., the opposite of the therian

190   condition). Although it has been suggested that monotremes lack a 'true' pterygoid muscle

(Patterson 1956), the presence of the pterygoid (both medial and lateral parts) is reported for

192   *Ornithorhynchus* in multiple studies (e.g., Zeller 1989, Diogo and Powell 2019). Zeller (1989) traced

the medial pterygoid in different ontogenetic stages of *Ornithorhynchus* specimens, showing that

194   *Ornithorhynchus* exhibits a medial pterygoid muscle that, early in ontogeny, inserts lateral to

Meckel's cartilage, nestled within the pterygoid fossa with a significant gap between the Meckel's

196   cartilage and the dentary bone (Fig. 2B, 2D-F); such an arrangement is likely analogous to the

muscle and middle-ear arrangement of eutriconodontans and spalacotherioids (Fig. 1). In sum,

198   monotremes and therians appear to have conflicting arrangements of the medial pterygoid muscle

relative to the Meckel's cartilage and middle-ear complex (Fig. 2), mirroring the conflicts raised by

200   the fossil record (Fig. 1).

**Figure 2.** *A* Chondrocranium reconstruction of a platypus nestling (*Ornithorhynchus anatinus*, 70 mm crown-rump-length) modified from Zeller (1989). The vertical line indicates the orientation of coronal sections in C–F. *B* The 3D model of a juvenile platypus lower jaw based on illustrations in Zeller (1989), highlighting the position of the medial pterygoid muscle (pink) between the dentary (brown) and Meckel's cartilage (yellow). *C* Line-drawing of coronal histological section modified from Zeller (1989), with the red box indicating the area shown in D-F. *D-F* Histological sections modified from Zeller (1989), illustrating the monotreme condition in which the medial pterygoid muscle (mp) passes lateral to

210   Meckel's cartilage (mc), nested between the Meckel's cartilage and the dentary bone (d). *G*
      Chondrocranium and dermal bone skeleton of a water shrew fetus (*Potamogale velox*, 105 mm crown-
212   rump-length) modified from Ruf et al. (2020). The vertical line indicates the position of coronal section in
      H. *H* Histological section of a water shrew fetus (*Potamogale velox*, 48 mm crown-rump-length) modified
214   from Brocklehurst et al. (2016), illustrating the eutherian condition in which the medial pterygoid
      muscle passes medial to Meckel's cartilage (mc). *I* Histological section of a pouch young bandicoot
216   (*Perameles* sp., head length 17.5 mm) modified from Sánchez-Villagra and Forasiepi (2017), illustrating
      the metatherian condition in which the medial pterygoid muscle passes medial to Meckel's cartilage. For
218   orientation in *D-F*, *H*, and *I*, the positions of the CNV (fifth cranial nerve or trigeminal), $CNV_2$ (maxillary
      nerve) and $CNV_3$ (mandibular nerve), and CNVg (trigeminal ganglion) are indicated. Abbreviations: *ant*
220   anterior *dor* dorsal, *lat* lateral. Scale bars are approximately 1 mm.

224        If the traditional reconstruction for non-mammalian cynodonts is correct—that is, a medial
      pterygoid muscle inserting on the pseudangular process—then the evolutionary lineage to therians
226   must have included two evolutionary shifts of the medial pterygoid muscle relative to the position of
      Meckel's cartilage: one in early mammals, which lack an angular process, and one at Cladotheria
228   when the 'true' angular process evolved. These shifts are illustrated in Figure 1A. This raises
      questions about the accuracy of previous muscle reconstructions and how soft tissues like the
230   medial pterygoid muscle interacted with the detaching middle-ear complex. Answering those
      questions is crucial for understanding the chewing and hearing mechanisms of early mammals,
232   which lack extant analogs.
           Here we review hypotheses about the evolution of the mammalian medial pterygoid muscle
234   in Cynodontia and discuss their implications for homology and function. We propose a novel
      interpretation of medial pterygoid evolution that builds on a hypothesis proposed by Patterson
236   (1956). Nonetheless, we emphasize that each hypothesis has its merits and limitations, and our aim

is to ultimately reignite discussions and provide suggestions for future studies to help resolve this

238     evolutionary conundrum.

240          HYPOTHESES ON THE EVOLUTION OF THE MAMMALIAN MEDIAL PTERYGOID MUSCLE

242     Although there are numerous open questions on the evolution of the medial pterygoid muscle,

        hereafter we primarily focus on the issue of the medial pterygoid insertion site in the cynodont

244     lineage leading to mammals (Fig. 1). Specifically, we review three alternative hypotheses on the

        evolution of the mammalian medial pterygoid muscle (Fig. 3) and highlight conflicting lines of

246     evidence derived from what is known from the fossil record and ontogenetic studies of extant

        mammals, including monotremes. Two of the hypotheses (Simpson Hypothesis and Patterson

248     Hypothesis) were previously proposed but are updated here based on our current understanding of

        the fossil record, and we introduce a new hypothesis based on phylogenetic parsimony (Fig. 3). We

250     discuss the merits and shortcomings of each hypothesis.

**Figure 3**. Left hemimandible and medial pterygoid (or pterygoideus) muscle reconstructions in medial and posterior views. The Simpson Hypothesis is the 'traditional' reconstruction of the medial pterygoid muscle on the pseudangular process of non-cladotherian cynodonts (Fig. 1; Simpson 1928, Lautenschlager et al. 2017). Following the Patterson Hypothesis, the pterygoideus musculature (pink) may represent an undifferentiated pterygoideus muscle (Patterson 1956), or it may represent separate but adjacent medial pterygoid (MP) and lateral pterygoid (LP) muscles (see the alternative Patterson Hypothesis in the text). The highlighted phylogeny branches in the Phylogenetic Parsimony Hypothesis represent the clade (with spalacotherioids and multituberculates) that would need to be recovered in phylogenetic analyses to support the hypothesis. Blue stars and dashed maroon arrows indicate phylogenetic nodes at which the relative position of the medial pterygoid to Meckel's cartilage changes. Meckel's cartilages in early cladotherians (based in part on reconstructions in Allin and Hopson 1992) and early monotremes (represented by *Teinolophos*; Rich et al. 2016) are drawn with dashed lines because those elements are not presented in the fossil record, and in cladotherians it is likely only present early in ontogenetic development. The Meckel's cartilage position in *Teinolophos* is based on the position of Meckel's groove (Rich et al. 2016), and additional ear elements are not shown due to uncertainty of their positions.

1) Simpson Hypothesis

This hypothesis represents the traditional interpretation of medial pterygoid muscle

evolution, which we summarized in the *Introduction*. Its central tenet is that the medial

pterygoid muscle inserts on the pseudangular process of the dentary in non-mammalian

cynodonts (Figs. 1 and 3; Simpson 1928, Crompton 1963, Lautenschlager 2017). We title this

the 'Simpson Hypothesis' because, to our knowledge, Simpson (1928) was the first to argue

that the mammaliaform pseudangular process is homologous to the therian angular process.

In descriptions of the lower jaws of *Morganucodon* (e.g., Mussett and Kermack 1981) and

docodontans (e.g., Simpson 1928, Rougier et al. 2015), the pseudangular process is often

homologized with the angular process of therian mammals and assumed to be the

attachment site of the medial pterygoid muscle in all non-mammalian cynodonts with a

developed pseudangular/angular process. In this scenario, the medial pterygoid muscle in

non-mammalian cynodonts passes medial to the middle ear bones in the postdentary trough

(Fig. 3A). Assuming the pterygoideus muscle (or multiple pterygoid muscles) ancestrally

inserted on the angular-articular region of cynodonts (e.g., Lautenschlager et al. 2017), the

lateral pterygoid muscle insertion site would have shifted dorsally to the dentary condyle,

whereas the medial pterygoid muscle insertion site would have shifted in the opposite

direction, ventrally onto the dentary pseudangular process.

*Conflicting evidence:* Some early mammalian groups (i.e., eutriconodontans,

spalacotherioids, and multituberculates) lack a pseudangular or angular process and instead

exhibit a deep pterygoid fossa on the posteromedial portion of the dentary (Fig. 1).

Following the Simpson Hypothesis configuration, the medial pterygoid muscle would

therefore have to evolutionarily shift laterally 'over' the Meckel's cartilage and postdentary

trough to get from the pseudangular process (medial to the middle ear ossicles and Meckel's

cartilage) to occupy the pterygoid fossa (lateral to the middle ear ossicles and the Meckel's

298     cartilage), as shown in Figures 1 and 3. From its position lateral to Meckel's cartilage within

the pterygoid fossa, the medial pterygoid muscle would then have to shift again to pass

300     medial to Meckel's cartilage as is seen among extant therians (and presumably among their

close cladotherian relatives; Fig. 3). Thus, the Simpson Hypothesis requires the medial

302     pterygoid muscle to make two evolutionary shifts in position relative to Meckel's cartilage.

304          2)   Patterson Hypothesis

Patterson (1956) posited that the cynodont pterygoideus muscle did not differentiate into

306     the medial and lateral pterygoid muscles until Cladotheria (Fig. 3). He further suggested that

the undifferentiated pterygoideus attached to the 'medial ridge' (or 'dentary shelf'; Sidor and

308     Hopson 2017), which is dorsal to Meckel's cartilage and the postdentary bones of

mammaliaforms (Fig. 4), not to the pseudangular process as proposed for the medial

310     pterygoid muscle in the Simpson Hypothesis. In accordance with the Patterson Hypothesis, if

a muscle attached to the medial surface of the pseudangular process in non-mammalian

312     mammaliaforms, it was not the medial pterygoid muscle; instead, it may have been jaw-

depressor muscles such as the digastric (Bramble 1978) or detrahens mandibulae, the latter

314     of which is unique (as far as we know) to extant monotremes and inserts on the

pseudangular process of echidnas (Schulman 1906; referred to as the 'echidna angle' by

316     Patterson 1956 and Patterson and Olson 1961) or on the ventral surface of the mandibular

body in platypuses (Hopson 1966). As the postdentary bones detached and were medially

318     displaced from the jaw during their transition to mammalian middle-ear ossicles,

attachment of the pterygoideus muscles remained along the medial ridge, which is posited to

320     be homologous to the pterygoid shelf or fossa (Fig. 4) in early mammal clades that lack a

pseudangular or angular process (i.e., eutricondontans, spalacotherioids, and

322     multituberculates). In this evolutionary scenario, the medial pterygoid muscle shifted from a

position dorsal to the postdentary bones to a position lateral to Meckel's cartilage as the

324     pterygoid shelf shifted ventrally (Figs. 3 and 4), which is similar to the monotreme condition

        early in ontogeny (Fig. 2B, D-F).

Lateral pterygoid
Medial pterygoid
Pterygoideus m. (or MP&LP)
Meckel's cartilage
Surangular
Ectotympanic
Malleus (articular)
Incus
328     Dentary

**Figure 4.** *A*, Posterior part of the lower jaw of the eutriconodontan *Priacodon* with virtual middle ear
330     bones in place to illustrate the lateral position of the pterygoideus musculature (light red) in relation to
        the middle ear bones. *B*, Lower jaw of an *Ornithorhynchus* (monotreme) hatchling with the laterally
332     positioned medial pterygoid muscle (dark red) in relation to the middle ear bones. Only the medial
        pterygoid is reconstructed for *Ornithorhynchus* (see Fig. 2) and not the lateral pterygoid, which is visible
334     in Figure 2F. *C*, Following the Patterson Hypothesis, we reconstructed the insertion of the pterygoideus
        musculature (light red), which represents either a single muscle or adjacent medial and lateral pterygoid
336     muscles (because it is unclear if a separation pterygoid muscles existed in those taxa), in lower jaws of
        the morganucodontan *Morganucodon*, docodontan *Haldanodon*, and eutriconodontan *Priacodon*, and the
338     resulting cladotherian split of both pterygoids (medial [dark red] and lateral [pink]) in *Dryolestes*. Jaws
        are not to scale.

A central aspect of the Patterson Hypothesis is that the pterygoideus musculature inserted on the medial ridge of non-mammalian synapsids, and not on the pseudangular process (Fig. 4). In this scenario, the medial ridge of non-mammalian synapsids is homologous to the 'pterygoid shelf' (medioventral to the pterygoid fossa) of early mammals like eutriconodontans (Fig. 1C). It is worth mentioning that extant mammals often exhibit a 'endocondylar ridge,' a structure similar to the medial ridge that likely serves a bite force distributing function (Wilken et al. 2024). However, the endocondylar ridge and medial ridge are likely analogous, not homologous, because many early mammals (e.g., eutriconodontans, multituberculates, spalacotherioids) lack either structure, just like those groups lack an angular process.

An alternative version of the Patterson Hypothesis is that the medial and lateral pterygoid muscles differentiated earlier in synapsid evolution (at least by Mammaliaformes), which is a common view (e.g., Barghusen 1973, Romer and Price 1940, Watson 1953, Crompton 1963, Lautenschlager et al. 2017), and both muscles inserted on the medial ridge (or elsewhere dorsal to Meckel's cartilage) in non-mammalian mammaliaforms, not on the pseudangular process. In this scenario, the medial pterygoid muscle (rather than the entire pterygoideus muscle like in the original version of the Patterson Hypothesis) evolutionarily drops ventrally between Meckel's cartilage and the dentary bone in early mammals. This alternative hypothesis is supported by the presence of both a medial pterygoid and lateral pterygoid in monotremes (Fig. 2F; Zeller 1989, Diogo and Powell 2019), which is why we illustrate the medial pterygoid separately from the lateral pterygoid in monotremes (Figs. 2 and 4), except for *Teinolophos* in Figure 3.

*Conflicting evidence*: The original version of Patterson's hypothesis posits that the pterygoideus does not differentiate into the medial and lateral pterygoid muscles until Cladotheria, which conflicts with interpretations of most other studies on the evolution of

366      mammalian jaw adductor musculature (Simpson 1928, Romer and Price 1940, Watson 1953,

Crompton 1963, Barghusen 1973, Kermack and Mussett 1981, Lautenschlager et al. 2017)

368      and the observation of both a medial and lateral pterygoid in monotremes (Fig 2F: Zeller

1989, Diogo and Powell 2019). (Note that the alternative version of the Patterson

370      Hypothesis discussed in the previous paragraph helps to address this conflict.) Further, the

Patterson Hypothesis suggests that when the medial pterygoid evolved at Cladotheria it

372      passed medial to Meckel's cartilage (which is supported by ontogenetic evidence from

therians; Fig. 2H, I), whereas the pterygoideus muscle of early mammals passed lateral to

374      Meckel's cartilage (Fig. 3). In other words, some pterygoideus muscle fibers probably shifted

in position relative to Meckel's cartilage, which would also occur in the scenario presented

376      by the Simpson Hypothesis (Figs. 1 and 3). Nonetheless, adult cladotherians do not maintain

the ear-jaw connection via Meckel's cartilage; the connection is lost early in ontogeny.  Thus,

378      there would not have been an ossified Meckel's cartilage limiting jaw masticatory

movements in adults – this may be linked to the evolutionary shift in muscle position

380      (Grossnickle 2017, Grossnickle et al. 2022).

An additional issue with the Patterson Hypothesis is that the superficial masseter

382      muscle is commonly reconstructed to insert on the lateral surface of the pseudangular and

angular processes (e.g., Abdala and Damiani 2004, Lautenschlager et al. 2017; but see

384      Patterson and Olson 1961 for an opposing view). If these reconstructions are correct, it

supports the argument that pseudangular and angular processes are homologous (e.g.,

386      Rougier et al. 2015), conflicting with Patterson's assertion of non-homology of these

structures.

     3)   Phylogenetic Parsimony Hypothesis

390      This hypothesis, like the Simpson Hypothesis, reconstructs the medial pterygoid muscle as

inserting on the pseudangular process of the dentary in non-mammalian cynodonts (Fig. 3;

392        Simpson 1928, Crompton 1963, Lautenschlager et al. 2017). However, it minimizes the

number of evolutionary shifts in the position of the medial pterygoid muscle relative to

394        Meckel's cartilage by collapsing early crown lineages (monotremes, spalacotherioids,

eutriconodonts, and multituberculates) into a single clade; thus, it includes only a single shift

396        in medial pterygoid position (or evolutionary loss of the medial pterygoid muscle) at the

base of that clade (Fig. 3). If the mammaliaform pseudangular process and cladotherian

398        angular process are homologous (e.g., Rougier et al. 2015), then this hypothesis offers a

possible explanation for the lack of an angular process in multiple early mammal lineages;

400        those lineages may form a monophyletic clade and the loss of an angular process (in all

groups besides monotremes) may be a derived trait of the clade.

402                *Conflicting evidence:* The major conflicting line of evidence with the Phylogenetic

Parsimony Hypothesis is that it requires a phylogenetic topology that contradicts the

404        phylogenetic hypotheses of multiple independent studies (Meng et al. 2011, Rougier et al.

2012, Krause et al. 2014, Krause et al. 2020, Luo et al. 2015, Luo et al. 2017, Huttenlocker

406        2018, Hoffmann et al. 2020). To the best of our knowledge, no published phylogenetic

analysis has recovered monotremes, spalacotherioids, eutriconodonts, and

408        multituberculates as a clade. Due to the lack of phylogenetic evidence for this hypothesis, we

hereafter primarily focus on evaluating the Simpson and Patterson hypotheses.

410        Nevertheless, future studies could test this hypothesis by examining the likelihood of the

required tree topology.

PTERYGOIDEUS MUSCULATURE OF MONOTREMES AND THERIANS

A central observation of this study, which to the best of our knowledge has not been discussed in

416    previous literature, is that therians and monotremes have different positions of the medial

pterygoid muscle relative to Meckel's cartilage during ontogenetic development (Figs. 2 and 3;

418    Zeller 1989, Brocklehurst et al. 2015, Sánchez-Villagra and Forasiepi 2017). Thus, even if fossil

       evidence is ignored, there is still an issue with the homology of the medial pterygoid muscle among

420    Mammalia. We contend that the Patterson Hypothesis, not the Simpson Hypothesis, aligns more

       parsimoniously with those embryological studies (Fig. 3). That is, the Patterson Hypothesis does not

422    require the insertion site of the pterygoideus musculature to make an evolutionary shift relative to

       Meckel's cartilage in early mammals, whereas the Simpson Hypothesis does (Fig. 3). Following the

424    Patterson Hypothesis, if the pterygoideus musculature inserted on the pterygoid shelf and passed

       lateral to Meckel's cartilage in early mammals, then the pterygoideus musculature in monotremes

426    may have simply maintained that ancestral state (Fig. 2; Zeller 1989). (Patterson [1956] argued that

       monotremes lack a medial pterygoid or pterygoideus muscle, but other authors have shown that the

428    muscle is present in monotremes [Toldt 1905, Zeller 1989, Diogo and Powell 2019].) That the

       therian medial pterygoid muscle, in contrast, inserts medial to Meckel's cartilage early in ontogeny

430    may be linked to the evolution of the cladotherian angular process, which might in turn have

       initiated the reposition of the muscle insertion site. In further support of the Patterson Hypothesis, a

432    human embryo diagnosed with Treacher Collins syndrome (mandibulofacial dysostosis) was

       observed to have a medial pterygoid that passed *lateral* to Meckel's cartilage and does not insert on

434    the angular process, instead inserting dorsal to ear elements at or near the mylohyoid line (Herring

       et al. 1979) – this configuration is consistent with the ancestral mammalian condition following the

436    Patterson Hypothesis.

              The jaw morphology of the Early Cretaceous fossil monotreme, *Teinolophos* (Rich et al.

438    2016), also provides support for the Patterson Hypothesis. In *Teinolophos*, Meckel's groove is

       described as ventral to a trough for the mandibular branch of the trigeminal nerve ($V_3$) (Rich et al.

440    2016), suggesting that Meckel's cartilage passed very close to the ventral margin of the mandibular

       body (Fig. 3). Thus, there would have been very little space on the medial surface of the

442    pseudangular process for insertion of the medial pterygoid muscle, which would have likely

       interfered with middle ear elements occupying the same space. More likely is that the pterygoideus

444 musculature inserted dorsal to both Meckel's cartilage and the trough for $V_3$, which is how we

reconstruct it in Figure 3. That anatomical configuration for *Teinolophos* (and spalacotherioids and

446 eutriconodonts) is consistent with that of extant monotremes (Fig. 2A–F): the insertion site of the

medial pterygoid muscle in *Ornithorhynchus* is primarily dorsal to Meckel's cartilage, with the

448 muscle fibers passing lateral to Meckel's cartilage (Figs. 1 and 3).

450 THE PSEUDANGULAR PROCESS AND JAW DEPRESSOR MUSCLES

452 We follow the Patterson Hypothesis (Patterson 1956, Patterson and Olson 1961) in contending that

the medial face of the cynodont pseudangular process was not the insertion site for the medial

454 pterygoid, but rather jaw-depressor musculature. (Similarly, Patterson and Olson [1961] argued

that no part of the masseter muscle inserted on the pseudangular process.) In sauropsids and early

456 synapsids, the depressor mandibulae was the primary jaw-depressor muscle, and it inserted on the

articular, the postdentary bone forming the ancestral jaw joint with the quadrate of the cranium. As

458 the secondary—i.e., dentary-squamosal—jaw joint was established during the evolution of

cynodonts, however, the articular became increasingly reduced in size (e.g., Sidor 2003). Hopson

460 (1966) hypothesized that reduction of the articular bone resulted in the ancestral depressor

mandibulae losing its function, providing an evolutionary impetus for new jaw-depressor

462 musculature to emerge. He hypothesized that both the therian digastric muscle and the monotreme

detrahens mandibulae muscle therefore emerged as novel jaw depressors as a consequence of the

464 establishment of the dentary-squamosal jaw joint in cynodonts.

Indeed, consistent with Hopson's (1966) hypothesis, the pseudangular process first appears

466 among cynodonts in which the articular-quadrate jaw joint gives way to the dentary-squamosal jaw

joint (e.g., *Probelesodon*, *Probainognathus*) (Lautenschlager et al. 2017). Hopson (1966) further

468 hypothesized that reduction of the articular resulted in the loss of function of the sauropsid

depressor mandibulae muscle, which attached to the retroarticular process and new jaw-depressor

470　(mouth-opening) muscles like the digastric (of therians) and detrahens mandibulae (of

monotremes) could emerge. Following Hopson's (1966) hypothesis, the sauropsid depressor

472　mandibulae muscles became the "pterygoideus complex" in lineages leading to mammals, with the

depressor mandibulae itself likely becoming the tensor tympani muscle of mammals. Thus,

474　reduction of the articular bone may have driven the emergence of both novel mammalian jaw-

depressors muscles and a new site for their attachment along the posteroventral and medial portion

476　of the dentary (i.e., the pseudangular process). The view that a muscle other than the medial

pterygoid inserted on the pseudangular process in non-mammaliaform cynodonts is shared by

478　Bramble (1978) and seemingly supported by additional studies, such as Crompton and Hylander

(1986) and Crompton (1995).

480　　　　The evolutionary emergence of the pseudangular process also had implications for the

superficial masseter, which may have inserted on the pseudangular process's lateral surface (e.g.,

482　Abdala and Damiani 2004, Lautenschlager et al. 2017; but see Patterson and Olson 1961 for an

opposing view)—whether changes to the masseter musculature of cynodonts also contributed to

484　the emergence of the pseudangular process, or whether that shift occurred after the structure was

in place, remains unclear. Lastly, it is worth noting that some researchers suggest that the

486　pterygoideus also originally attached to the articular in early therapsids (Parrington 1955;

Crompton 1963; Lautenschlager et al. 2017). If that was the case, this suggests a rather elegant

488　evolutionary scenario in which the arrangement of the mammalian jaw musculature resulted from a

cascade of evolutionary changes brought on by the early establishment of the dentary-squamosal

490　jaw joint in cynodonts.

　　　　That the pseudangular process was an attachment site for jaw depressor muscles rather

492　than the medial pterygoid muscle (Patterson 1956, Patterson and Olson 1961) is also supported by

comparative anatomy and functional morphology. Relative to the cladotherian angular processes,

494　pseudangular processes of early cynodonts and mammaliaforms are anteriorly positioned relative

to the jaw joint, rather than posteriorly positioned (Fig. 5; Patterson 1956, Grossnickle 2017),

496    suggesting that they are not homologous and might be associated with different biomechanical

functions. Indeed, the pseudangular process may be functionally analogous to a therian 'second'

498    angular process, termed the 'marginal process' (Toldt 1905, Patterson 1956) that evolved in some

therian lineages, including bat-eared foxes (*Otocyon*), sloth bears (*Melursus*), and *Solenodon* (Fig. 5).

500    In these taxa, the marginal process serves as the insertion site for the jaw-depressing digastric

muscle, whereas the medial pterygoid inserts on the 'true' angular process (Toldt 1905, Allen 1910,

502    Ewer 1973, Wible 2008). Similarly, the pseudangular process of monotremes, including that of the

echnida and *Teinolophos*, is the insertion site of the detrahens mandibulae muscle, a jaw depressor

504    analogous to the therian digastric muscle (Fig. 4; Schulman 1906, Hopson 1966). The relatively

anterior position of the pseudangular process compared to the angular process may optimize

506    leverage for jaw depression by the posteriorly oriented jaw depressor muscles. Conversely, the

more posteriorly positioned cladotherian angular process is posited to be associated with

508    generating rotational jaw movements such as yaw (Grossnickle 2017, Grossnickle 2020) or roll

(Bhullar et al. 2019), which both result in transverse movements of molars during occlusion.

510    Further, the jaw pseudangular process is often described as bent laterally ('effected angle'; see

Schultz et al. 2017 for a summary), which may be due to the pull of the laterally attached superficial

512    masseter lacking a corresponding pull by a medially attached muscle such as the medial pterygoid.

In contrast, the angular process of therians, which have a definitive medial pterygoid muscle, is

514    inflected medially in many taxa, such as in most marsupials (e.g., Maier 1987, Sánchez-Villagra and

Smith 1997), likely enhancing roll rotation (Grossnickle 2020).

**Figure 5**. Comparative jaw anatomy of example eutherians that exhibit a 'marginal process' (red arrows;
520    Patterson 1956), which is the insertion site for the digastric muscle, and fossil cynodonts. (Note that the
eutherian jaws are in lateral view, but the medial pterygoid muscle inserts primarily on the medial
522    surface.) In terms of anatomical position along the length of the jaw, the pseudangular process (pAP) of
non-cladotherian cynodonts is more in-line with the therian 'marginal process' than the therian angular
524    process (AP; blue arrows). Jaw sizes are scaled roughly based on the jaw-joint-to-molars distance.

In sum, these observations on functional jaw anatomies support the argument by Patterson

528    (1956) and Patterson and Olson (1961) that (1) the pseudangular and angular processes are not

homologous and (2) that the pseudangular process served as an insertion site for a jaw depressor

530    muscle, not the medial pterygoid (Fig. 5), and thus serves a different functional purpose than the

angular process.

EXPANDING ON THE PATTERSON HYPOTHESIS

The Patterson Hypothesis makes two arguments that contradict traditional muscle reconstructions

536    (i.e., the Simpson Hypothesis): (1) the pterygoideus muscle does not give rise to medial and lateral

pterygoid muscles until Cladotheria and (2) the pterygoideus muscle inserts on the dentary medial

538    ridge (not the pseudangular process as commonly reconstructed) in non-mammalian

mammaliaforms (and possibly earlier taxa), suggesting homology of the medial ridge and medial

540    surface of the therian angular process. In this section, we examine these arguments in greater detail.

The first argument of the Patterson Hypothesis is challenging to assess due to the small size

542    of the lateral pterygoid muscle (making it unlikely for researchers to observe bone scarring or

osteological correlates) and the considerable uncertainty on the insertion sites of jaw musculature

544    in non-mammalian synapsids (see the *Introduction*). The therian lateral pterygoid muscle is

generally oriented to generate medially or anteriorly directed jaw movements and is often

546    associated with fine-scale jaw movements, such as slight protrusion and/or lateral movement of the

jaw in primates (Hylander 2006). But most non-cladotherian synapsid lineages lacked complex

548    chewing cycles that involved transverse movements, instead relying primarily on orthally or

posteriorly directed jaw movements (Grossnickle et al. 2022 and sources within), with taxa such as

550    docodontans (with pseudo-tribosphenic molars analogous to therian molars) and the dicynodont

*Sangusaurus* (Angielcyzk et al. 2017) being possible exceptions.

552    We also reiterate that an alternative version of the Patterson Hypothesis is that the medial

and lateral pterygoid muscles evolved prior to Cladotheria but that the medial pterygoid never

554    inserted on the pseudangular process. If the ancestral cynodont condition is that both lateral and

medial pterygoid muscles are present and lie next to each other in close proximity inserting on the

556    angular-articular region (like has been reconstructed in *Thrinaxodon* and *Probainognathus*; see

Lautenschlager et al. 2017 and sources within, Parrington 1955, Barghusen 1968, 1972, Crompton

558    1963), then the muscles may have remained in close association and both shifted dorsally to the

medial ridge. In other words, rather than a single pterygoideus muscle migrating dorsally to the

560    dentary as in the Patterson Hypothesis, it is possible that both the medial and lateral pterygoid

muscles shifted dorsally. This contradicts the Simpson Hypothesis in which there is a divided state

562    of the lateral and medial pterygoid muscles in early mammaliaforms, with the lateral pterygoid

muscle attaching close to the articular condyle (dorsal of postdentary bones) and the medial

564    pterygoid muscle inserting ventral to the postdentary bones on the pseudangular process.

The second argument of the Patterson Hypothesis—that the pterygoideus muscle inserts on

566    the dentary medial ridge in non-mammalian mammaliaforms—can be assessed by examining

osteological evidence and comparing this scenario to that of the Simpson Hypothesis (e.g., Crompton

568    1963, Lautenschlager 2017). For instance, in favor of the pterygoideus muscle inserting on the

dentary medial ridge is that non-mammalian mammaliaforms exhibit a well-developed medial ridge,

570    a bony crest above the postdentary trough (Fig. 4). The medial ridge includes significant surface

area (likely greater than the surface area of the medial side of the pseudangular process in most

572    taxa) for muscle attachment. However, it is also worth noting that beyond serving as a muscle

insertion site, the medial ridge might serve as the primary load path for distributing bite forces like

574    has been suggested for the endocondylar ridge of therian mammals (Panagiotopoulou et al. 2020;

Wilken et al. 2024). Non-mammalian mammaliaforms have complex middle ear elements inside the

576    grooves of the postdentary trough for sound transmission; a pterygoideus that inserts dorsal to the

middle ear elements is unlikely to have interfered with sound transmission. In contrast, a medial

578 pterygoid muscle inserting on the pseudangular process (i.e., the Simpson Hypothesis condition; Fig.

3) would pass adjacent to the ear elements, possibly limiting the vibration and thus function of the

580 ear elements.

The Patterson Hypothesis is further supported by our current knowledge of the evolutionary

582 detachment of the middle ear from the dentary. The middle ear ossicles detach posteriorly first

while they keep a connection to the lower jaw anterior to the middle ear ossicles via the Meckel's

584 cartilage (Luo et al. 2007, Mao et al. 2020, Mao and Meng 2020). Following the Patterson

Hypothesis, when ear elements are still anteriorly connected to the lower jaw via the Meckel's

586 cartilage the pterygoideus remains lateral to Meckel's cartilage (Figs. 1 and 3), even as the pterygoid

fossa expands ventrally. For example, in eutriconodonts such as *Yanoconodon*, the pterygoid fossa

588 and the jaw joint are confluent (Luo et al. 2007, Luo 2007) and the pterygoideus may have spread

along the entire region, keeping posterior fibers close to the jaw joint, which is where the lateral

590 pterygoid muscle is positioned in extant mammals. In contrast to the Simpson Hypothesis, this

evolutionary scenario is possible without a 'jumping' of the medial pterygoid muscle over middle

592 ear elements at/near Mammalia (Fig. 3). Luo and Manley (2020) suggest the pterygoid fossa of

mammals has 'replaced' the non-mammalian mammaliaform postdentary trough, but our

594 interpretation of evidence from monotreme ontogeny and the preserved ossified Meckel's cartilages

of eutriconodontans and spalacotherioids goes further, implying a homology of the non-mammalian

596 mammaliaform attachment area dorsal to the medial ridge and the mammalian pterygoid fossa.

598                                THE ALLOTHERIAN PERSPECTIVE

600 One major early mammalian group—Allotheria—has not yet been addressed, due in part to

uncertainties surrounding their phylogenetic relationships. Allotheria, if a valid monophyletic

602 group, include haramiyidans (including the Late Triassic-Early Jurassic *Haramiyavia* and *Thomasia*

and the Middle Jurassic Euharamiyida), gondwanatherians, and multituberculates (e.g., Butler 2000;

604 Zheng et al. 2013; Mao et al., 2023). However, many authors view 'Allotheria' as polyphyletic, with many specific variations in terms of which groups are included/excluded in Allotheria (Butler 2000,

606 Luo et al. 2002, Kielan-Jaworowska et al. 2004, Butler and Hooker 2005, Zhou et al. 2013, Luo et al. 2015b, Krause et al. 2020a, Hoffmann et al. 2020, King and Beck 2020). Disputes surrounding the

608 putative monophyly of Allotheria are centered on interpretations of the middle-ear configuration in haramiyidans. Multituberculata have a fully detached mammalian middle ear (Lillegraven and Hahn

610 1993, Szalay et al. 1993, Hurum et al. 1996, Rougier et al. 1996). In comparison, some authors contend that all haramiyidans had middle-ear bones still closely attached to the jaw (e.g., Zhou et al.

612 2013; Luo et al. 2015b), but some euharamiyidans may have exhibited partly attached middle ears still tethered to the jaw by Meckel's cartilage (i.e., analogous to eutriconodontans and

614 spalacotherioids) and some may have had fully detached definitive mammalian middle ear (e.g., Zheng et al. 2013; Mao et al. 2023). Until the phylogenetic relationships are resolved, interpreting

616 allotherian medial pterygoid muscle evolution remains challenging. Here, we briefly summarize the implications (following the Patterson Hypothesis) under three competing evolutionary scenarios:

618 (1) Allotheria are monophyletic and outside of crown Mammalia, (2) Allotheria are monophyletic and within crown Mammalia, and (3) Allotheria are polyphyletic.

620   If Allotheria are monophyletic and outside Crown Mammalia, the evolution of the middle ear and medial pterygoid muscle would largely mirror what we have proposed above for non-

622 allotherian mammaliaforms (Fig. 6). Like *Morganucodon*, the coeval *Haramiyavia* had middle-ear bones closely affixed to the jaw in a post-dentary trough (Luo et al. 2015b), whereas, many

624 euharamiyidans had a detached middle ear (e.g., Meng 2014; Mao et al., 2023) or a middle-ear that was tethered to the dentary by Meckel's cartilage like eutriconodontans and spalacotherioids. Like

626 cladotherians, multituberculates exhibit a fully detached middle ear (e.g., Hurum et al. 1996; Rougier et al. 1996). Following the Patterson Hypothesis in this scenario, the pterygoideus muscle would

628 have attached to the 'medial ridge' or 'pterygoid shelf' overhanging the postdentary trough in early allotherians like *Haramiyavia* (Luo et al. 2015b; Fig. 6), and it would have then shifted ventrally (and

630    lateral to Meckel's cartilage) to continue attaching along the prominent pterygoid shelf in

euharamiyidans, gondwanatherians, and multituberculates (e.g., Gambaryan and Kielan-

632    Jaworowska 1995, Meng et al. 2014, Krause et al. 2020b). Allotherians did not develop an angular

process like that seen in Cladotheria and the medial pterygoid (or pterygoideus) muscle presumably

634    remained lateral to Meckel's cartilage throughout allotherian evolution. If Allotheria are

monophyletic but within Crown Mammalia, the story largely remains the same, with the primary

636    difference that the full sequence of middle ear detachment (from fully attached, to partly detached,

to completely detached) occurred independently in allotherians within Crown Mammalia.

642    **Figure 6**. Evolution of the allotherian pterygoid musculature following the Patterson Hypothesis.

Whether haramiyidans are non-mammalian mammaliaforms (A; e.g., Luo et al. 2015b) or crown

644    mammals closely related to multituberculates in a monophyletic Allotheria (B; e.g., Zheng et al. 2013),

the undifferentiated pterygoid muscle likely attached to the medial ridge or pterygoid shelf above the

646    postdentary elements, either dorsal (as in *Haramiyavia* and, in scenario A, *Vilevolodon*) or lateral (as in

*Jeholbaatar* and, in scenario B, *Vilevolodon*) to Meckel's cartilage.

650        If Allotheria are polyphyletic and multituberculates are the only crown mammalian group,

multituberculates evolved a detached middle ear independently with no obvious evolutionary

652    precursor known from the fossil record. The earliest multituberculate specimens for which the

morphology of the posteromedial dentaries can be observed (e.g., Paulchoffatiidae like

654    *Pseudobolodon* Hahn 1969 or *Rugosodon* Yuan et al. 2013), that show no evidence of Meckel's

cartilage (or other post-dentary bones) attaching to the jaw. The medial pterygoid (or pterygoideus)

656    muscle is reconstructed to attach to the pterygoid shelf along the ventral margin of the

posteromedial dentary (e.g., Simpson, 1926; Wall and Krause, 1992; Gambaryan and Kielan-

658    Jaworowska, 1995; Fig. 6), but its origin (i.e., medial ridge or pseudangular process) is not clear due

to missing evidence in the fossil record leading to Multituberculata.

660        Relevant to the Patterson Hypothesis, the medial lower jaws of multituberculates do not

exhibit obviously different putative insertion locations for a split medial and lateral pterygoid.

662    Instead, the ventromedial portion of the jaw posterior to the molars is simply characterized by an

expansive fossa (e.g., Clemens and Kielan-Jaworowska, 1979; Kielan-Jaworowska et al., 2004).

664    Indeed, in previous muscle reconstructions of the multituberculate masticatory apparatus, the

medial and lateral pterygoids are either reconstructed as being right next to each other in the

666    pterygoid fossa (with the lateral pterygoid slight posterodorsal to the medial pterygoid; Gambaryan

and Kielan-Jaworowska, 1995) or undifferentiated as a single 'pterygoid' (Simpson, 1926; Wall and

668    Krause, 1992). Thus, the anatomy of multituberculate jaws in consistent with the Patterson

Hypothesis, which assumes the pterygoideus remained a single muscle until the evolution of

670    Cladotheria (Fig. 6).

        Unlike nearly all other non-mammalian mammaliaforms, allotherians are characterized by

672    bilateral mastication and exhibit mostly orthal and palinal (posteriorly oriented) chewing strokes

(Krause 1982, Jenkins et al. 1997, Butler 2000, Butler and Hooker 2005, Lazzari et al. 2010, Weaver

674    and Wilson 2021, Grossnickle et al. 2022). Thus, whereas the medial pterygoid muscle is essential

for facilitating yaw, roll, and translation during the unilateral occlusion of cladotherians, it was likely

676    primarily used for stabilizing the jaws during bilateral occlusion in allotherians (e.g., Butler 2000;

Butler and Hooker 2005; Gambaryan and Kielan-Jaworowska 1995). That the pterygoid shelves and

678    fossae of allotherians are very well-developed points to the importance of the pterygoid

musculature in general; yet, instead of enhancing transverse movements like cladotherians (e.g.,

680    Grossnickle 2017), the medial pterygoid (or pterygoideus) muscle probably helped prevent

deviations from the horizontal plane, maintaining the precise occlusion of interlocking longitudinal

682    cusp rows during the palinally directed power stroke that characterizes allotherian mastication.

684                        IMPLICATIONS FOR HEARING AND FEEDING

686    At the time of Patterson's (1956) publication, well-preserved fossils of eutriconodontans and

spalacotherioids were unknown, and our understanding of early mammalian phylogenetic

688    relationships were more poorly resolved, so it is unclear how he would have interpreted these new

revelations. Patterson (1956) did posit, however, that the medial pterygoid muscle evolved

690    concurrently with the evolutionary disconnection of the middle ear bones of the jaw, talonid shelf of

molars, and cladotherian angular process. Following the Patterson Hypothesis, this suggests that

692    specializations of the masticatory and hearing apparatuses evolved in concert with evolution of the

pterygoideus musculature. See Grossnickle et al. (2022) for an expanded discussion on how these

694    evolutionary changes may be linked to functional changes of the masticatory apparatus.

        The postdentary trough of early mammaliaforms, containing bony and cartilaginous

696    elements and responsible for sound transmission to the inner ear, naturally requires freedom of

movement for transmitting vibrations from one element to the other. Any jaw movement (not

698    necessarily chewing) probably interfered with such a fine-tuned system. In particular, soft tissue

structures, such as muscles and tendons, as reconstructed with the medial pterygoid muscle

700    attached to the pseudangular process (e.g., see the Simpson Hypothesis in Figure 3), may have

       hindered transmission process by wrapping closely around the elements preventing mobility. The

702    alternative placement advocated here (Patterson Hypothesis; Fig. 3) of muscle and tendon above the

       postdentary trough allows for more freedom of movement.

704         Following the Patterson Hypothesis, once in the pterygoid fossa, the pterygoideus muscle

       passes dorsal to (in early mammaliaforms) or lateral to (in early mammals) the middle ear elements

706    (Fig. 4). A pterygoideus muscle dorsal or lateral to the middle ear elements prevents possible close

       contact between the contracting muscle and ear elements. In contrast, following the Simpson

708    Hypothesis scenario in non-mammalian synapsids (Fig. 3), the medial pterygoid muscle would

       drape medio-ventrally around Meckel's cartilage, and the contracting muscle would likely interfere

710    with the mobility of the ossicles and limit the transmission of vibrations. Later, in early

       cladotherians, the ear completely separates from the jaw, which may have allowed the jaw joint to

712    shift dorsally (e.g., Grossnickle 2017) and provide space for the medial pterygoid insertion site to

       shift to a more posteroventral position on the jaw, likely triggering the evolution of the angular

714    process. As a result, cladotherians reflect the modern mammalian configuration of the medial and

       lateral pterygoid muscles (Fig. 4). The concurrent evolutionary change to cladotherian molars,

716    namely the origin of a talonid shelf that suggests the evolution of novel medially directed occlusal

       movements (Schultz and Martin 2014, Grossnickle 2017, Grossnickle et al. 2022), further suggests

718    that the new jaw and muscle morphologies were associated with a fundamental shift in chewing

       mechanics in cladotherians.

CONCLUSIONS

       There is considerable uncertainty about the position of the medial pterygoid muscle insertion site

724    throughout the evolution of synapsids. For instance, there is a lack of a consensus on the cynodont

       pterygoideus configuration (Crompton 1993, Bramble 1978, Barghusen 1986, Allin and Hopson

726    1992, Crompton 1995, Lautenschlager et al. 2017), which is necessary to understand the

evolutionary steps that lead to the mammalian condition of the pterygoid muscles. Newly

728    interpreted evidence from fossils and developmental investigations, in addition to the vetting of

three competing hypotheses (Fig. 3), highlight the difficulties of understanding the muscular

730    construction of jaw adductors in the history of mammals, but also allow for new perspectives in

regard of the evolution of the mammalian hearing and feeding system.

732          We favor Patterson's (1956) hypothesis that medial pterygoid muscle did not insert on the

pseudangular process in mammaliaforms, based on information derived from recent fossil findings

734    (Figs. 1, 3–6), monotreme ontogeny (Fig. 2), and comparative anatomy of mammalian jaws (Fig. 5).

Mammalian evolution may have involved migration of the pterygoideus muscle insertion site from

736    the articular and/or ossified Meckel's cartilage (i.e., prearticular) in early cynodonts to the dentary's

medial ridge above the postdentary trough in early mammaliaforms (Fig. 4), potentially in response

738    to articular reduction and the establishment of the dentary-squamosal jaw joint (Hopson, 1966).

Patterson (1956) argued that the pterygoideus does not give rise to the medial and lateral pterygoid

740    muscles until Cladotheria, but it is also possible that the muscles were present in earlier lineages but

both maintained insertion sites dorsal and/or lateral to Meckel's cartilage. The Patterson

742    Hypothesis indicates homology of the non-mammalian mammaliaform medial ridge and mammalian

pterygoid fossa due to the continued insertion of the pterygoideus on this structure during early

744    mammalian evolution. During the evolutionary separation of the middle ear elements from the jaw,

the medial pterygoid muscle likely 'dropped' ventrally between the middle ear and dentary bone,

746    occupying space lateral to the ossified Meckel's cartilage. The monotreme condition is a pterygoid

muscle that passes lateral to Meckel's cartilage (Fig. 2), which is the same configuration

748    reconstructed in early crown mammals (Fig. 1), suggesting that monotremes have maintained the

ancestral mammalian state. In further support of the Patterson Hypothesis, we present evidence

750    that the medial or posterior surface of the pseudangular process of non-mammalian

mammaliaforms was the insertion site of a jaw depressor muscle (possibly the monotreme

752     detrahens mandibulae or the therian digastric; Fig. 5) rather than the medial pterygoid muscle (as

posited by the Simpson Hypothesis and Phylogenetic Parsimony Hypothesis). We believe that the

754     evolutionary scenario of the Patterson Hypothesis is more parsimonious than that suggested by the

Simpson Hypothesis, which would require multiple changes in the position of the medial pterygoid

756     muscle relative to Meckel's cartilage (Figs. 1 and 3).

        Nonetheless, we emphasize that none of the hypotheses presented in this study (Fig. 3),

758     including the Patterson Hypothesis, are fully satisfactory because they each require seemingly

unlikely evolutionary scenarios. Each hypothesis requires either evolutionary shifts in the position

760     of medial pterygoid muscle relative to Meckel's cartilage or suggest timings of the origin or loss of

the medial pterygoid muscle that are inconsistent with various lines of evidence (e.g., fossil record,

762     ontogeny, phylogenetic parsimony). Thus, we recommend future studies further test competing

hypotheses on the evolution of medial pterygoid muscle, making use of various lines of evidence

764     from paleontology, developmental biology, phylogenetics, and biomechanics. In particular, the

strengths of competing hypotheses are sensitive to the ancestral position of the medial pterygoid

766     muscle on the postdentary elements in non-mammaliaform cynodonts—a ventral position on the

pseudangular process would support the Simpson Hypothesis and a dorsal position on postdentary

768     bones or the medial ridge would support the Patterson Hypothesis. This topic could be investigated

in mammaliamorphan cynodonts such as brasilodontids (Rawson et al. 2024), which possess a mix

770     of mammalian and non-mammalian traits. In addition, we emphasize that future studies should not

only utilize the ever-growing fossil record but should also examine the ontogeny of soft tissue parts

772     surrounding the developing middle ear in extant mammals, especially monotremes. Finally, we see

potential for utilizing biomechanical studies to test evolutionary hypotheses on the medial

774     pterygoid (and additional muscles like the superficial masseter), such as investigating load-bearing

patterns on non-mammalian mammaliaform lower jaws by comparing the two different attachment

776     sites (pseudangular process of the Simpson Hypothesis versus medial ridge of the Patterson

Hypothesis). Our goal here is to reinvigorate interest and debate on the topic of soft tissue evolution

778    in early mammals, and we hope to spur future research on the topic.

780                                                    MATERIALS & METHODS

782    We produced 3D digital models of fossil jaws of the mammaliaforms *Morganucodon*, *Haldanodon*,

the euticonodont *Priacodon*, and the cladotherian *Dryolestes*, which were then used to virtually

784    reconstruct origin and insertion areas of the pterygoides (or medial pterygoid and lateral pterygoid)

muscles. The 3D models were produced from micro-computed tomography (micro-CT) scans. The

786    lower jaws of *Morganucodon* (NHM M84028, Natural History Museum London) and *Priacodon*

(LACM 120451, Natural History Museum of Los Angeles) were scanned at the University of Texas

788    High-Resolution X-ray CT facility using the NSI scanner. Scan parameters for *Morganucodon*: Xradia,

60 kV, 10 W, 1 s acquisition time, 1 mm SiO2 filter, voxel size 19.64 μm. Scan parameters for

790    *Priacodon*: Fein focus high power source, 110 kV, 0.15 mA, no filter, voxel size 10.4 μm. For further

scan information, see Jäger et al. (2020). The lower jaws of *Haldanodon* (GuiMam 3-80) and

792    *Dryolestes* (GuiMam 3-78) were scanned at the Bonner Insitute of Organismic Biology (BIOB,

formerly Institute of Geosciences, Universität Bonn; section Palaeontology) using the v|tome|x s μCT

794    device (GE Sensing & Inspection Technologies GmbH phoenix|x-ray). Scan parameters for

*Haldanodon*: 240 tube, 90 kV, 180 μA, 333 ms exposure time, no filter, voxel size 16.49 μm. Scan

796    parameters for *Dryolestes*: 240 tube, 90 kV, 180 μA, 333ms exposure time, no filter, voxel size 24.66

μm.

798         The 3D model of the *Ornithorhynchus* hatchling lower jaw is a virtually edited polygonal

surface model based on the adult surface model of *Ornithorhynchus anatinus* downloaded from

800    MorphoSource (imnh:r:2125; OBJ mesh file, Idaho Museum of Natural History specimen S32167). In

a manual procedure the surface model was modified and edited in MAYA 16 (Autodesk) to create

802    the hatchling jaw morphology and middle ear bones based on illustrations of hatchling MO 39 from

Zeller (1989) and to fit the histological sections within (Fig. 2).

804        All 3D models (except *Ornithorhynchus*) were segmented from micro-CT-image stacks and

virtual 3D models were created using Avizo 8.1 (Visualization Sciences Group, France). Editing and

806    virtual alignment was performed with Polyworks (2014, InnovMetric Software Inc., Canada)

including Gouraud filters for smoothing surfaces, adding and removing triangles, reworking mesh to

808    reduce data size. The 3D models representing different evolutionary stages of the middle ear bones

and ossified Meckel's cartilage were manually created using MAYA 16 (Autodesk) based on

810    illustrations in Kermack et al. (1973), Kermack et al. (1981), Luo (2011), Luo et al. (2007), Luo et al.

(2016), Ji et al. (2009), and Mao et al. (2020).

## ACKNOWLEDGEMENTS

We are grateful to Thomas Martin, Rich Cifelli, Brian Davis and Pip Brewer for providing micro-CT

816    scans of the lower jaws of *Morganucodon*, *Priacodon*, *Haldanodon* and *Dryolestes*. Sue Herring, Alec

Wilken, Savannah Olroyd, James Rawson, and Abigail Tucker provided valuable feedback on earlier

818    drafts of the manuscript. Further, we benefitted from discussions with Zhe-Xi Luo, Thomas Martin,

James A. Hopson, Rebecca German, Paulina Jiménez Huidobro, Bastian Mähler, Simone Hoffman, Elsa

820    Panciroli, Elis Newham, Pamela Gill, Callum Ross, Dallas Krentzel, Allison K. Bormet, and Kenneth-

Dieter Benton. For technical support, we thank Peter Göddertz. Funding to LNW was provided by

822    National Science Foundation (NSF) Graduate Research Fellowship and EAR-PF 2052992, and DMG

was supported by an NSF Postdoctoral Research Fellowship in Biology (DBI-1812126). JAS profited

824    from funding by the Deutsche Forschungsgemeinschaft (DFG) during the two funding periods of

research unit FOR 771.

828    **Conflict of interest statement**: the authors declare that there is no conflict of interest.

       **Data availability statement**: there are no data to be archived.

830    **Author contributions**: JAS, DMG, and LNW wrote the manuscript and created the figures. JAS

       contributed *Haldanodon* and *Dryolestes* 3D data, creating and editing of 3D digital models. KRKJ

832    contributed *Morganucodon* and *Priacodon* 3D data and edited the manuscript.

REFERENCES

       Abdala F, Damiani R. 2004. Early development of the mammalian superficial masseter muscle in

838    cynodonts. *Palaeontologica Africana*, 40, 23-29.

       Allen GM. 1910. *Solenodon paradoxus. Memoirs of the Museum of Comparative Zoology at Harvard

840    College*, 40,1-54.

       Allin EF, Hopson JA. 1992. Evolution of the auditory system in Synapsida ('mammal-like reptiles' and

842    primitive mammals) as seen in the fossil record. In The evolutionary biology of hearing (eds. DB

       Webster, RR Fay, AN Popper), pp. 587-614. Springer.

844    Angielczyk KD, Hancox PJ, Nabavizadeh A. 2017. A redescription of the Triassic kannemeyeriiform

       dicynodont *Sangusaurus* (Therapsida, Anomodontia), with an analysis of its feeding system.

846    *Journal of Vertebrate Paleontology*, 37(sup1), 189-227.

       Anthwal N, Joshi L, Tucker AS. 2013. Evolution of the mammalian middle ear and jaw: adaptations

848    and novel structures. *Journal of Anatomy*, 222, 147-160.

       Barghusen HR. 1968. The lower jaw of cynodonts (Reptilia, Therapsida) and the evolutionary origin

850    of mammal-like adductor jaw musculature. *Postilla*, 116, 1-49.

       Barghusen HR. 1972. The origin of the mammalian jaw apparatus. In Morphology of the

852    Maxillomandibular Apparatus (ed. Schumacher GH), pp. 26–32. VEB Georg Thieme.

854  Barghusen HR. 1973. The adductor jaw musculature of *Dimetrodon* (Reptilia, Pelycosauria). *Journal of Paleontology*, 47, 823-834.

Barghusen HR. 1986. On the evolutionary origin of the therian tensor veli palatini and tensor

856  tympani muscles. In The Ecology and Biology of Mammal-like Reptiles, ed. N. Hotton Ill, P.D. MacLean, J.J. Roth, E.C. Roth, pp. 263-282. Smithsonian Press.

858  Barry TH. 1965. On the epipterygoid-alisphenoid transition in Therapsida. *Annals of the South African Museum*, 48, 399-426.

860  Bhullar BAS, Manafzadeh AR, Miyamae JA, Hoffman EA, Brainerd EL, Musinsky C. Crompton AW.

2019. Rolling of the jaw is essential for mammalian chewing and tribosphenic molar function.

862  *Nature*, 566(7745), 528-532.

Bramble DM. 1978. Origin of the mammalian feeding complex: models and mechanisms.

864  *Paleobiology*, 4, 271-301.

Brocklehurst RJ, Crumpton N, Button E, Asher RJ. 2016. Jaw anatomy of *Potamogale velox*

866  (Tenrecidae, Afrotheria) with a focus on cranial arteries and the coronoid canal in mammals. *PeerJ*, 4, e1906.

868  Butler PM. 2000. Review of the early allotherian mammals. *Acta Palaeontologica Polonica*, 45(4).

Butler PM, Hooker JJ. 2005. New teeth of allotherian mammals from the English Bathonian, including

870  the earliest multituberculates. *Acta Palaeontologica Polonica*, 50(2).

Clemens JR, Kielan-Jaworowska Z. 1979. Multituberculata. In Mesozoic Mammals: The First Two-

872  Thirds of Mammalian History (eds. Lillegraven JA, Kielan-Jaworowska Z, Clemens WA), pp. 99-149. University of California Press.

874  Crompton AW. 1995. Masticatory function in nonmammalian cynodonts and early mammals. In

Functional morphology in vertebrate Paleontology (ed. Thomason J), pp. 55-75. Cambridge

876  University Press.

Crompton AW. 1963. The evolution of the mammalian jaw. *Evolution*, 431-439.

878    Crompton AW, Parkyn DG. 1963. On the lower jaw of *Diarthrognathus* and the origin of the

       mammalian lower jaw. *Proceedings of the Zoological Society of London*, 140(4), 697-749.

880    Crompton AW, Hylander, WL. 1986. Changes in mandibular function following the acquisition of a

       dentary-squamosal jaw articulation. In The ecology and biology of mammal-like reptiles (eds.

882    Hotton N, MacLean PD, Roth JJ, Roth EC), pp. 263–282. Smithsonian Institution Press.

       Crompton AW, Owerkowicz T, Skinner J. 2010. Masticatory motor pattern in the koala

884    (*Phascolarctos cinereus*): a comparison of jaw movements in marsupial and placental herbivores.

       *Journal of Experimental Zoology A - Ecology Genetics Physiology*, 313, 564-578.

886    Crompton AW, Musinsky C, Bonaparte J, Bhullar BA, Owerkowicz T. 2018. Evolution of the

       mammalian fauces region and the origin of suckling. https://nrs.harvard.edu/URN-

888    3:HUL.INSTREPOS:37364482 (permanent link)

       De Beer GR. 1937. The development of the vertebrate skull. Clarendon Press.

890    DeMar R, Barghusen HR. 1972. Mechanics and the evolution of the synapsid jaw. *Evolution*, 26, 622-

       637.

892    Diogo R, Powell V. 2019. The origin and evolution of mammalian head muscles with special

       emphasis on the facial myology of primates and modern humans. In Heads, Jaws, and Muscles:

894    Anatomical, Functional, and Developmental Diversity in Chordate Evolution (eds. Ziermann JM,

       Diaz Jr. RE, Diogo R), pp. 253-293. Springer.

896    Ewer RF. 1973. The carnivores. Cornell University Press.

       Gambaryan PP, Kielan-Jaworowska Z. 1995. Masticatory musculature of Asian taeniolabidoid

898    multituberculate mammals. *Acta Palaeontologica Polonica*, 40(1).

       Gow CE (1986). A new skull of *Megazostrodon* (Mammalia, Triconodonta) from the Elliot Formation

900    (Lower Jurassic) of Southern Africa. *Palaeontologia Africana*, 26, 13-23.

       Grossnickle DM. 2017. The evolutionary origin of jaw yaw in mammals. Scientific Reports 7, 45094.

902    Grossnickle DM. 2020. Jaw roll and jaw yaw in early mammals. *Nature* 582, E6-E8.

Grossnickle DM, Weaver LN, Jäger KRK, Schultz JA. 2022. The evolution of anteriorly directed molar

904      occlusion in mammals. *Zoological Journal of the Linnean Society*, 194, 349-365.

Hahn G. 1969. Beiträge zur Fauna der Grube Guimarota Nr. 3. Die Multituberculata.

906      *Palaeontographica* A, 133, 1-100

Herring SW. 2003. TMJ anatomy and animal models. *Journal of musculoskeletal & neuronal*

908      *interactions*, 3, 391-394.

Herring SW, Rowlatt UF, Pruzansky S, Optiz JM. 1979. Anatomical abnormalities in mandibulofacial

910      dysostosis. *American Journal of Medical Genetics*, 3(3), 225-259.

Herring SW, Scapino RP. 1973. Physiology of feeding in miniature pigs. *Journal of Morphology*,

912      141(4), 427-460.

Hiiemae KM, Crompton AW. 1985. Mastication, food transport, and swallowing. In Functional

914      vertebrate morphology (eds. Hildebrand M, Bramble DM, Liem KF, Wake DB), pp. 262-290.

         Harvard University Press.

916  Hoffmann S, Beck RMD, Wible JR, Rougier GW, Krause DW. 2020. Phylogenetic placement of

         *Adalatherium hui* (Mammalia, Gondwanatheria) from the Late Cretaceous of Madagascar:

918      implications for allotherian relationships. *Journal of Vertebrate Paleontology*, 40 (Suppl. 1), 213-

920  Hopson JA. 1966. The origin of the mammalian middle ear. *American Zoologist*, 6(3), 437-450.

     Hurum JH, Presley R, Kielan-Jaworowsk, Z. 1996. The middle ear in multituberculate mammals. *Acta*

922      *Palaeontologica Polonica*, 41(3), 253-275.

     Huttenlocker AK, Grossnickle DM, Kirkland JI, Schultz JA, Luo ZX. 2018. Late-surviving stem mammal

924      links the lowermost Cretaceous of North America and Gondwana. *Nature*, 558(7708), 108-112.

     Hylander WL. 2006. Functional anatomy and biomechanics of the masticatory apparatus. In

926      Temporomandibular disorders: an evidenced approach to diagnosis and treatment (eds Laskin

         DM, Greene CS, Hylander WL), pp. 3-34. Quintessence Pub Co.

Iordansky NN. 2010. Pterygoideus muscles and other jaw adductors in amphibians and reptiles. *Biology Bulletin*, 37(9), 905-914.

Jäger KRK, Cifelli RL, Martin T. 2020. Molar occlusion and jaw roll in early crown mammals. *Scientific Reports*, 10(1), 22378.

Jenkins FA Jr, Crompton AW, Downs WR. 1983. Mesozoic mammals from Arizona: new evidence on mammalian evolution. *Science*, 222, 1233-1235

Ji Q, Luo ZX, Zhang X, Yuan CX, Xu L. 2009. Evolutionary development of the middle ear in Mesozoic therian mammals. *Science*, 326(5950), 278-281.

Kemp TS. 1972. The jaw articulation and musculature of the whaitsiid Therocephalia. In Studies in Vertebrate Evolution (eds Joysey KA and TS Kemp), pp. 213-230. Oliver & Boyd.

Kermack KA, Mussett F. 1958. The jaw articulation of the Docodonta and the classification of Mesozoic mammals. *Proceedings of the Royal Society of London, Series B*, 149, 204-215.

Kermack KA, Mussett F, Rigney HW. 1981. The skull of *Morganucodon*. *Zoological Journal of the Linnean Society*, 71, 1-158.

Kermack KA, Mussett F, Rigney HW. 1973. The lower jaw of *Morganucodon*. *Zoological Journal of the Linnean Society*, 53, 87-175.

Kielan-Jaworowska Z, Cifelli RL, Luo ZX. 2004. Mammals from the age of dinosaurs: origins, evolution, and structure. Columbia University Press.

King GM. 1981. The functional anatomy of a Permian dicynodont. Philosophical *Transactions of the Royal Society of London. B, Biological Sciences*, 291(1050), 243-322.

King B, Beck RM. 2020. Tip dating supports novel resolutions of controversial relationships among early mammals. *Proceedings of the Royal Society B*, 287(1928), 20200943.

Krause DW, Hoffmann S, Wible JR, Kirk EC, Schultz JA, von Koenigswald W, Groenke JR, Rossie JB, O'Connor PM, Seiffert ER, Dumont ER, Holloway WL, Rogers RR, Rahantarisoa LJ, Kemp AD,

Andriamialison H. 2014. First cranial remains of a gondwanatherian mammal reveal remarkable mosaicism. *Nature*, 515(7528), 512-517.

Krause DW, Hoffmann S, Hu Y, Wible JR, Rougier GW, Kirk EC, Groenke JR, Rogers RR, Rossie JB,

Schultz JA, Evans AR, von Koenigswald W, Rahantarisoa LJ. 2020a. Skeleton of a Cretaceous

956    mammal from Madagascar reflects long-term insularity. *Nature*, 581(7809), 421-427.

Krause DW, Hoffmann S, Wible JR, Rougier GW. 2020b. Lower jaw morphology of *Adalatherium hui*

958    (Mammalia, Gondwanatheria) from the Late Cretaceous of Madagascar. *Journal of Vertebrate*

*Paleontology*, 40(sup1), 81-96.

960    Krause DW. 1982. Jaw movement, dental function, and diet in the Paleocene multituberculate

*Ptilodus*. *Paleobiology*, 8(3), 265-281.

962    Lautenschlager S, Gill PG, Luo ZX, Fagan MJ, Rayfield EJ. 2017. Morphological evolution of the

mammalian jaw adductor complex. *Biological Reviews*, 92, 1910-1940.

964    Lazzari V, Schultz JA, Tafforeau P, Martin T. 2010. Occlusal pattern in paulchoffatiid

multituberculates and the evolution of cusp morphology in mammaliamorphs with rodent-like

966    dentitions. *Journal of Mammalian Evolution*, 17(3), 177-192.

Lillegraven JA, Hahn G. 1993. Evolutionary analysis of the middle and inner ear of Late Jurassic

968    multituberculates. *Journal of Mammalian Evolution*, 1(1), 47-74.

Luo ZX. 2011. Developmental patterns in Mesozoic evolution of mammal ears. *Annual Review of*

970    *Ecology, Evolution, and Systematics*, 42, 355-380.

Luo ZX. 2007. Transformation and diversification in early mammal evolution. *Nature*, 450, 1011-

972    1019.

Luo ZX, Manley GA. 2020. Origins and early evolution of mammalian ears and hearing function. In

974    The Senses: A comprehensive Reference (2nd Edition) (ed Fritzsch B), pp. 207-252. Academic

Press.

976    Luo ZX, Crompton AW. 1994. Transformation of the quadrate (incus) through the transition from

nonmammalian cynodonts to mammals. *Journal Vertebrate Paleontology*, 14, 341-374.

978    Luo ZX, Chen P, Li G, Chen M. 2007. A new eutriconodont mammal and evolutionary development in

early mammals. *Nature*, 446, 288-293.

980    Luo ZX, Meng QJ, Ji Q, Liu D, Zhang YG, Neander AI. 2015a. Evolutionary development in basal

        mammaliaforms as revealed by a docodontan. *Science*, 347(6223), 760-764.

982    Luo ZX, Gatesy SM, Jenkins Jr FA, Amaral WW, Shubin NH. (2015b). Mandibular and dental

        characteristics of Late Triassic mammaliaform *Haramiyavia* and their ramifications for basal

984    mammal evolution. *Proceedings of the National Academy of Sciences*, 112(51), E7101-E7109.

        Luo ZX, Schultz JA, Ekdale EG. 2016. Evolution of the middle and inner ears of mammaliaforms: The

986    approach to mammals. In Evolution of the vertebrate ear: evidence from the fossil record (eds JA

        Clack, RR Fay, AN Popper), pp. 139-174. Springer.

988    Luo ZX, Meng QJ, Grossnickle DM, Liu D, Neander AI, Zhang YG, Ji Q. 2017. New evidence for

        mammaliaform ear evolution and feeding adaptation in a Jurassic ecosystem. *Nature*,

990    548(7667), 326-329.

        Maier W. 1987. The angular process of *Monodelphis domestica* (Didelphidae, Marsupialia) and its

992    relation to the middle ear: an ontogenetic and evolutionary morphologic study. *Gegenbaurs*

        *Morphologisches Jahrbuch*, 133, 123-161.

994    Maier W, Ruf I. 2016. Evolution of the mammalian middle ear: a historical review. *Journal of*

        *Anatomy*, 228, 270-283.

996    Manley GA, Sienknecht UJ. 2013. The evolution and development of middle ears in land vertebrates.

        In The middle Ear: science, otosurgery, and technology (eds. Puria S, Fay RR, Popper AN), pp. 7-

998    30. Springer.

        Mao F, Li Z, Hooker JJ, Meng J. 2023. A new euharamiyidan, *Mirusodens caii* (Mammalia:

1000    Euharamiyida), from the Jurassic Yanliao Biota and evolution of allotherian mammals. *Zoological*

        *Journal of the Linnean Society*, 199(3), 832-859.

1002    Mao F, Hu Y, Li C, Wang Y, Chase MH, Smith AK, Meng J. 2020. Integrated hearing and chewing

        modules decoupled in a Cretaceous stem therian mammal. *Science*, 367, 305-308.

1004    Mao F, Meng J. 2020. Making a mammalian ear. Modular decoupling of the mammalian middle ear

and jaw discovered in a new species of Cretaceous stem therian mammals. *Zoology*, 140,

1006    125767.

Martin T, Marugán-Lobón J, Vullo R, Martín-Abad H, Luo ZX, Buscalioni AD. 2015. A Cretaceous

1008    eutriconodont and integument evolution in early mammals. *Nature*, 526, 380-384.

Meng J, Hu Y, Wang Y, Li C. 2003. The ossified Meckel's cartilage and internal groove in Mesozoic

1010    mammaliaforms: implications to origin of the definitive mammalian middle ear. *Zoological*

*Journal of the Linnean Society*, 138, 431-448.

1012    Meng J, Wang Y, Li C. 2011. Transitional mammalian middle ear from a new Cretaceous Jehol

eutriconodont. *Nature*, 472, 181-185.

1014    Meng J, Bi S, Wang Y, Zheng X, Wang X. 2014. Dental and mandibular morphologies of

*Arboroharamiya* (Haramiyida, Mammalia): a comparison with other haramiyidans and

1016    *Megaconus* and implications for mammalian evolution. *PloS One*, 9(12), e113847.

Panagiotopoulou O, Iriarte-Diaz J, Abraha HM, Taylor AB, Wilshin S, Dechow PC, Ross CF. 2020.

1018    Biomechanics of the mandible of *Macaca mulatta* during the power stroke of mastication:

loading, deformation, and strain regimes and the impact of food type. *Journal of Human*

1020    *Evolution*, 147, 102865.

Patterson B. 1956. Early Cretaceous mammals and the evolution of mammalian molar teeth.

1022    *Fieldiana Geology*, 13, 1-105.

Patterson B, Olson EC. 1961. A triconodontid mammal from the Triassic of Yunnan. In International

1024    colloquium on the evolution of lower and non-specialized mammals (ed. Vandebroek G.), pp

129-191. Koninklijke Vlaamse Academie voor Wetenschappen, Letteren en Schone Kunsten van

1026    Belgie.

Pfaff C, Schultz JA, Schellhorn R. 2019. The vertebrate middle and inner ear: A short overview.

1028    *Journal of morphology*, 280, 1098-1105.

Rawson JR, Martinelli AG, Gill PG, Soares MB, Schultz CL, Rayfield EJ. 2024. Brazilian fossils reveal homoplasy in the oldest mammalian jaw joint. *Nature*, 634(8033), 381-388.

Rosowski JJ. 1992. Hearing in transitional mammals: predictions from the middle-ear anatomy and hearing capabilities of extant mammals. In The evolutionary biology of hearing (eds DB Webster, AN Popper, RR Fay), pp. 615-631. Springer.

Rougier GW, Wible JR, Novacek MJ. (1996). Middle-ear ossicles of the multituberculate *Kryptobaatar* from the Mongolian late Cretaceous: implications for mammaliamorph relationships and the evolution of the auditory apparatus. *American Museum Novitates*, 3187, 1-45.

Rougier GW, Sheth AS, Carpenter K, Appella-Guiscafre L, Davis BM. 2015. A new species of *Docodon* (Mammaliaformes: Docodonta) from the Upper Jurassic Morrison Formation and a reassessment of selected craniodental characters in basal mammaliaforms. *Journal of Mammalian Evolution*, 22, 1-16.

Rougier GW, Wible JR, Beck RM, Apesteguía S. 2012. The Miocene mammal *Necrolestes demonstr*ates the survival of a Mesozoic nontherian lineage into the late Cenozoic of South America. *Proceedings of the National Academy of Sciences*, 109(49), 20053-20058.

Rougier GW, Apesteguía S, Gaetano LC. 2011. Highly specialized mammalian skulls from the Late Cretaceous of South America. *Nature*, 479(7371), 98-102.

Ruf I, Behrens H, Zeller U. 2020. Ontogeny and morphofunctional implications of the ethmoidal region in the semiaquatic *Potamogale velox* (Afrotheria, Mammalia). *Vertebrate Zoology*, 70, 679-697.

Sánchez-Villagra MR, Forasiepi AM. 2017. On the development of the chondrocranium and the histological anatomy of the head in perinatal stages of marsupial mammals. *Zoological Letters*, 3, 1-33.

Sánchez-Villagra MR, Smith KK. 1997. Diversity and evolution of the marsupial mandibular angular process. *Journal of Mammalian Evolution*, 4, 119-144.

1054 Schulman H. 1906. Vergleichende Untersuchungen über die Trigeminus-Musculatur der

Monotremen, sowie dabei in Betracht kommenden Nerven und Knochen. *Zoologische*

1056 *Forschungsreisen in Australien und dem Malayischen Archipel*, 3, 303-400.

Schultz JA. 2020. Eat and listen—how chewing and hearing evolved. *Science*, 367(6475), 244-246.

1058 Schultz JA. 2025. A perspective from the Mesozoic: Evolutionary changes of the mammalian skull

and their influence on feeding efficiency and high-frequency hearing. *The Anatomical Record*,

1060 DOI: 10.1002/ar.25652

Schultz JA, Bhullar BAS, Luo ZX. 2019. Re-examination of the Jurassic mammaliaform *Docodon victor*

1062 by computed tomography and occlusal functional analysis. Journal of Mammalian *Evolution*, 26,

9-38.

1064 Schultz JA, Martin T. 2014. Function of pretribosphenic and tribosphenic mammalian molars

inferred from 3D animation. *Naturwissenschaften*, 101, 771-781.

1066 Sidor CA. 2001. Simplification as a trend in synapsid cranial evolution. *Evolution* 55, 1419-1442.

Sidor CA, Hopson JA. 2017. *Cricodon metabolus* (Cynodontia: Gomphodontia) from the Triassic

1068 Ntawere Formation of northeastern Zambia: patterns of tooth replacement and a systematic

review of the Trirachodontidae. *Journal of Vertebrate Paleontology*, 37(Suppl. 1), 39-64.

1070 Simpson GG. 1928. Mesozoic Mammalia, XII - the internal mandibular groove of Jurassic mammals.

*American Journal of Science*, 5, 461-470.

1072 Szalay FS, Novacek MJ, McKenna MC. 1993. Mammal phylogeny: Mesozoic differentiation,

multituberculates, monotremes, early therians, and marsupials. Springer.

1074 Toldt C. 1905. Der Winkelfortsatz der Unterkiefer beim Menschen und bei den Säugetieren und die

Beziehungen der Kaumuskulatur zu demselben. *Sitzungsberichte der Kaiserliche Akademie der*

1076 *Wissenschaften in Wien*, 114, 315-476.

Wall CE, Krause DW. 1992. A biomechanical analysis of the masticatory apparatus of *Ptilodus*

1078 (Multituberculata). *Journal of Vertebrate Paleontology*, 12(2), 172-187.

Wang HB, Hoffmann S, Wang DC, Wang YQ. 2022. A new mammal from the Lower Cretaceous Jehol

1080    Biota and implications for eutherian evolution. *Philosophical Transactions of the Royal Society B*, 377(1847), 20210042.

1082    Wang Y, Hu Y, Meng J, Li C. 2001. An ossified Meckel's cartilage in two Cretaceous mammals and origin of the mammalian middle ear. *Science*, 294(5541), 357-361.

1084    Watson DMS. 1953. The evolution of the mammalian ear. *Evolution*, 7, 159-177.

Weaver LN, Wilson GP. 2021. Shape disparity in the blade-like premolars of multituberculate

mammals: functional constraints and the evolution of herbivory. *Journal of Mammalogy*, 102(4), 967-985.

Weijs WA. 1980. Biomechanical models and the analysis of form: a study of the mammalian masticatory apparatus. *American Zoologist*, 20, 707-719.

Weijs WA. 1994. Evolutionary approach of masticatory motor patterns. In Biomechanics of feeding in vertebrates (eds. Bels V, Chardon M, Vandewalle P), pp. 281-320. Springer.

Wible, JR. 2008. On the cranial osteology of the Hispaniolan solenodon, *Solenodon paradoxus* Brandt, 1833 (Mammalia, Lipotyphla, Solenodontidae). *Annals of Carnegie Museum*, 77 (3), 321-402.

Wilken AT, Schultz JA, Luo ZX, Ross CF. 2024. A new biomechanical model of the mammal jaw based on load path analysis. *Journal of Experimental Biology*, 227(18), jeb247030.

Yuan CX, Ji Q, Meng QJ, Tabrum AR, Luo ZX. 2013. Earliest evolution of multituberculate mammals revealed by a new Jurassic fossil. *Science*, 341(6147), 779-783.

Zeller U. 1989. Die Entwicklung und Morphologie des Schädels von *Onithorhynchus anatinus* (Mammalia: Prototheria: Monotremata). *Abhandlungen der Senckenbergischen Naturforschenden*

*Gesellschaft*, 545, 1-188.

Zheng X., Bi S., Wang X., Meng J. 2013. A new arboreal haramiyid shows the diversity of crown

1102    mammals in the Jurassic period. *Nature*, 500(7461), 199-202.

Zhou CF, Wu S, Martin T, Luo ZX. 2013. A Jurassic mammaliaform and the earliest mammalian

1104    evolutionary adaptations. *Nature*, 500(7461), 163-167.