Co-evolved Partners of Immunity: A Trait-Based Map of Human Keystone Organisms

Amir Asiaee[1], Natalie Mallal[2], Elizabeth Phillips[3,4],
Simon Mallal[3,4*]

[1]Department of Biostatistics, Vanderbilt University Medical Center,
TN, USA.
[2]College of Arts and Science, Vanderbilt University, TN, USA.
[3]Department of Medicine, Vanderbilt University Medical Center, TN,
USA.
[4]Institute for Immunology and Infectious Diseases, Murdoch University,
Murdoch, Western Australia.

*Corresponding author(s). E-mail(s): s.mallal@vumc.org;
Contributing authors: amir.asiaeetaheri@vumc.org;
natalie.f.mallal@vanderbilt.edu; elizabeth.j.phillips@vumc.org;

Abstract

Persistent human-adapted microbes can act as immunological "keystones," organizing host defense across tissues and shaping vulnerability under immune perturbation. Here we operationalize keystone organisms as pathogens whose containment requires coordinated engagement of multiple immune arms and whose residence is structured across anatomical niches. Using 18 curated immunological and evolutionary traits across 43 organisms, unsupervised analyses resolved four reproducible archetypes and identified a compact keystone set dominated by persistent herpesviruses and *Mycobacterium tuberculosis*. We then translated the clinical literature into a pathogen×immune-perturbation×niche tensor capturing where and when each organism emerges under defined immune deficits. We quantified "diagnostic breadth" with two complementary summaries: immune breadth (diversity of perturbations associated with emergence) and niche breadth (diversity of anatomical sites). Clinical emergence patterns perfectly separated trait-defined keystones from all other organisms and highlighted expanded niche breadth as the primary discriminator, whereas immune breadth showed no significant group separation. Finally, a mechanistic model integrating barrier disruption, latent reservoir activation, and tissue-resident immune control predicted clinical emergence from first principles—without fitting parameters to

individual pathogens—and ranked true emergences 2.9-fold above chance among its highest-confidence predictions. Together, these results link evolutionary adaptation to clinically readable patterns of reactivation, motivate archetype-aware surveillance under immunosuppression, and provide a framework for immunogen design that prioritizes conserved, functionally constrained targets.

Keywords: coevolution; keystone organisms; immunodominance; tissue-resident memory; diagnostic breadth; molecular mimicry; RNA virus decoy; vaccine design; drug hypersensitivity; autoimmunity

1 Introduction

Throughout evolutionary history, organismal complexity has been tightly linked to the surrounding microbial environment. One of the earliest and most consequential examples was the incorporation of an aerobic bacterium into the primordial eukaryotic cell, a mutualistic event that gave rise to mitochondria and enabled aerobic multicellular life to flourish more than two billion years ago [1]. This foundational instance of endosymbiosis illustrates a broader principle: successful organisms, including humans, have co-evolved in dynamic, reciprocal relationships with microbial partners.

While bacteria have long dominated discussions of host-microbe mutualism, viruses are increasingly recognized as contributors to host fitness across plants, animals, and humans [2, 3]. Viral elements can facilitate development, shape ecological and immunological adaptability, and calibrate immune function. These observations challenge a purely defensive view of immunity and instead support a model of continuous, calibrated co-regulation between host and infectious organisms [3].

Building on this ecological framing, Virgin, Wherry, and Ahmed argued that chronic viral infections—especially the human herpesviruses—are stable, dynamic components of our metagenome that imprint the immune system and calibrate future responses [4]. They emphasized co-evolutionary depth, the capacity to establish metastable latency, and subtle periodic reactivation that demands coordinated immune engagement without immunopathology. These hallmarks point toward a deeper organizing principle.

Defining keystone organisms.

The term *keystone* originates in architecture, where the central wedge-shaped stone at the apex of an arch bears and distributes weight to hold the structure in balance. Ecologists later adopted the concept to describe *keystone species*—organisms whose presence exerts an outsized influence on ecosystem stability [5]. By analogy, we propose that a subset of persistent, human-adapted organisms function as ***immunological keystones***: microbes that not only persist, but anchor immune architecture through multi-arm coordination and lifelong imprinting. Just as the loss of a keystone species disrupts ecological integrity, the absence or delayed acquisition of keystone organisms may have cascading effects on immune compartmentalization and resilience (Figure 1).

Fig. 1 Immunological keystone organisms as an organizing principle. Keystone organisms coordinate multiple immune arms and help stabilize immune homeostasis, analogous to a keystone supporting an arch, loss or delayed acquisition can destabilize the broader immune architecture. We categorize 43 pathogens into four archetypes (Opportunistic, Multi-host, Specialist, Keystone) arranged along a continuum of co-evolution and structured persistence, and illustrate each archetype's characteristic niche breadth and immune breadth.

Specifically, we propose that keystone organisms occupy structured tissue niches and demand coordinated containment by $CD8^+$ T cells, $CD4^+$ T cells, B cells, NK cells, and innate effectors [6, 7]. Evidence for such niche-structured coordination spans barrier tissues and internal compartments [8–10]. Studies of inborn errors of immunity further reveal that loss of specific immune arms produces selective vulnerability to particular organisms, underscoring the non-redundant, multi-arm coordination that keystone control requires [11, 12]. By requiring this multi-arm engagement, keystones focus postnatal immunodominance hierarchies—a process we term *keystone epitope focusing*, representing a third stage of immune selection beyond central (thymic) and peripheral tolerance.

A continuum of persistence.

Not all persistent organisms meet this definition. Microbes exist along a continuum. At one end are acute, external threats that trigger transient, compartmentalized immune responses. At the other are deeply integrated symbionts whose stable persistence within specific cellular niches requires sustained, multi-arm coordination—a compartmental equilibrium that represents a stable ecological state. The relationship is reciprocal: such organisms may modulate immune set-points in ways that correlate with host fitness, indirectly protecting against more virulent competitors. Between these poles lies a spectrum whose degree of keystoneness reflects persistence, specificity, evolutionary intimacy, and immunological impact.

Over deep evolutionary time, host-microbe relationships have been highly dynamic. Some lineages co-diverged with vertebrates and became near-obligate, tissue-structured companions—the human herpesviruses, for example, show genomic signatures of millions of years of co-evolution with primate hosts [13–15]. Others crossed species barriers, replaced prior residents, or entered host genomes as endogenous elements. Historic migrations and population bottlenecks have further shaped these relationships, selecting for immune architectures adapted to regionally prevalent pathogens [16]. Today's human keystones therefore represent only some outcomes of this churn; intermediate positions along our framework likely capture organisms in transit—either tightening toward mutualistic persistence or exploiting transient ecological opportunity. This view reconciles strong host-virus co-divergence in some families with episodic turnovers and host shifts in others, framing immunological hierarchies as products of ongoing dynamism rather than fixed properties [2, 4].

A hierarchical view of keystone theory.

We organize keystone theory across three levels of biological resolution. At the *population level*, lifelong co-evolution with persistent organisms has shaped which epitopes human immune systems prioritize—creating durable, species-wide hierarchies of immunodominance [4, 16]. At the *tissue level*, keystone organisms occupy specific anatomical niches (neurons, B cells, epithelium) and require coordinated containment by $CD8^+$ T cells, $CD4^+$ T cells, B cells, and NK cells working together within those compartments [17]. At the *molecular level*, this coordinated focusing on specific epitopes creates two clinically important vulnerabilities.

First, when drugs or self-proteins generate ligands that resemble keystone epitopes in the same tissue niche, they can recruit pre-existing antiviral memory, bypass regulatory checkpoints, and trigger T cell-mediated hypersensitivity or organ-specific autoimmunity [18–21]. Second, fast-evolving pathogens exploit this learned prioritization: RNA viruses present mutable "decoy" epitopes that re-stimulate keystone-imprinted T cells ineffectively, diverting immunity away from conserved viral targets and enabling persistence [22–25]. Tumors use similar strategies to evade immunosurveillance [25].

In short, the same epitope-focusing that enables efficient control of keystone organisms becomes a liability when exploited by molecular mimics or fast-mutating pathogens.

Scope of this study.

The lack of a systematic framework to classify host-microbe relationships along this continuum has hindered understanding of how particular organisms shape immune hierarchies. We address this gap through a literature-based, AI-assisted scoring approach applied at three levels of resolution (Figure 2).

We begin by asking what traits should characterize keystone organisms. Drawing on published evidence, we score 43 candidate organisms across 18 immunological traits spanning species specificity, persistence, co-evolutionary duration, multi-arm immune coordination, and structured tissue tropism. Unsupervised learning reveals

four natural archetypes: *keystone* organisms requiring broad, coordinated immune control; *specialist* pathogens with narrower immune and niche dependencies; *multi-host* organisms lacking deep human co-evolution; and *opportunistic* pathogens exploiting transient immune gaps (Figures 1, 2, and 4).

Among these traits, multi-arm immune coordination and tissue tropism have direct clinical correlates: when a specific immune arm is depleted or deficient, which pathogens emerge, and in which tissues? This framing yields a tractable question we can answer from clinical literature. We therefore curate reports of pathogen emergence across 13 anatomical niches and 31 immune perturbations to construct a three-dimensional tensor capturing *which* immune deficits reveal *which* pathogens in *which* tissues. Features derived from this clinical tensor discriminate trait-based archetypes from each other with high accuracy (Figures 2 and 5).

Finally, we ask whether a mechanistic model built from first principles—formalizing latent reactivation and barrier breach pathways in terms of pathogen biology, controller residency, and perturbation effects—can explain the clinical patterns we observe. This model reveals that keystone organisms are distinguished not by any single feature, but by requiring coordinated control across multiple immune arms in multiple tissue compartments (Figures 3 and 6).

While our analysis focuses on organisms, this framework lays the groundwork for examining the specific epitopes that mediate multi-arm coordination—setting the stage for mechanistic studies of mimicry risk and vaccine target prioritization.

2 Materials and Methods

This study used a literature-based, AI-assisted scoring approach to evaluate pathogen keystoneness at three levels: (i) organism-level trait scoring to identify pathogen archetypes, (ii) clinical emergence patterns to validate trait-derived groupings, and (iii) mechanistic modeling to explain emergence from first principles. All scoring was performed by structured review of published human clinical, genetic, and epidemiologic evidence, with AI assistance for evidence synthesis and initial draft scores, followed by expert review and adjudication.

2.1 Pathogen Selection

We selected 43 organisms representing diverse taxonomic groups (viruses, bacteria, fungi, protozoa, helminths) with documented capacity for human infection and varying degrees of persistence, immune coordination, and clinical significance under immunosuppression. Selection prioritized organisms with sufficient published evidence to score across multiple trait and clinical dimensions, spanning the hypothesized keystoneness continuum. The complete organism list with archetype assignments is provided in Table 1; detailed immunological and evolutionary characteristics for each archetype are provided in Supplementary Tables 3-6.

Fig. 2 Triangulating keystone organisms from three literature streams. Three complementary evidence streams are integrated: (i) trait scoring from immunology and evolutionary literature, (ii) a clinical emergence tensor of pathogen appearance across niches and immune perturbations, and (iii) mechanistic score tables encoding latent and barrier pathway components. These streams converge through cross-validation and stratified analyses to yield archetypes, diagnostic breadth summaries, and mechanistic predictions that explain clinical patterns and prioritize high-risk settings under immunosuppression.

2.2 Trait-Based Scoring (Figure 4)

2.2.1 Trait definitions

We curated 18 immunological traits hypothesized to reflect keystoneness based on published frameworks emphasizing co-evolutionary depth, multi-arm immune coordination, structured tissue tropism, and capacity for postnatal immune imprinting [3, 4]. Traits spanned five domains: (i) species specificity and co-evolutionary history; (ii) persistence and latency; (iii) immune coordination; (iv) epitope-centric features; and (v) clinical and ecological signatures. Trait definitions and allowed score ranges are provided in Table 7.

2.2.2 Scoring protocol

For each pathogen–trait pair, we conducted a structured literature review using a ChatGPT Pro AI agent configured to search PubMed, Google Scholar, and pathogen-specific databases. Evidence was prioritized hierarchically: (i) human genetic or interventional studies; (ii) large epidemiologic cohorts or mechanistically grounded clinical observations; (iii) animal or in vitro data when human evidence was unavailable. Scores were assigned from predefined integer sets according to explicit rubrics (Table 7), with higher values indicating stronger evidence. Total keystoneness scores were computed as unweighted sums across all 18 traits.

Table 1 Pathogens included in the study, grouped by the inferred archetype. Keystones exhibit deep co-evolution and multi-arm immune coordination; Specialists show focused tropism with moderate coordination; Multi-host pathogens have broad tissue engagement but lack stable human-specific imprinting; Opportunists flag pathway-specific immune failure. Detailed characteristics are provided in Supplementary Tables 3–6.

Archetype	Organism	Abbr.	Type
Keystone (n=7)	Cytomegalovirus	CMV	Herpesvirus
	Epstein–Barr Virus	EBV	Herpesvirus
	Herpes Simplex Virus 1	HSV-1	Herpesvirus
	Herpes Simplex Virus 2	HSV-2	Herpesvirus
	Human Herpesvirus 6	HHV-6	Herpesvirus
	Mycobacterium tuberculosis	Mtb	Bacteria
	Varicella–Zoster Virus	VZV	Herpesvirus
Specialist (n=11)	BK Virus	BKV	Polyomavirus
	Candida albicans	C.alb	Fungus
	Helicobacter pylori	H.pyl	Bacteria
	Human Papillomavirus	HPV	Papillomavirus
	JC Virus	JCV	Polyomavirus
	Kaposi's Sarcoma-Assoc. Herpesvirus	KSHV	Herpesvirus
	Merkel Cell Polyomavirus	MCPyV	Polyomavirus
	Neisseria meningitidis	N.men	Bacteria
	Pneumocystis jirovecii	P.jir	Fungus
	Parvovirus B19	PVB19	Virus
	Treponema pallidum	T.pal	Bacteria
Multi-host (n=10)	*Aspergillus* spp.	Asper	Fungus
	Cryptococcus neoformans	C.neo	Fungus
	Histoplasma capsulatum	H.cap	Fungus
	Leishmania spp.	Leish	Protozoan
	Mycobacterium avium complex	MAC	Bacteria
	Plasmodium falciparum	P.fal	Protozoan
	Schistosoma spp.	Schis	Helminth
	Strongyloides stercoralis	S.ste	Helminth
	Toxoplasma gondii	T.gon	Protozoan
	Trypanosoma cruzi	T.cru	Protozoan
Opportunist (n=15)	*Acanthamoeba* spp.	Acan	Amoeba
	Balamuthia mandrillaris	B.man	Amoeba
	Bartonella spp.	Barto	Bacteria
	Brucella spp.	Bruce	Bacteria
	Burkholderia cepacia complex	Bcc	Bacteria
	Coccidioides spp.	Cocci	Fungus
	Coxiella burnetii	C.bur	Bacteria
	Fusarium spp.	Fusar	Fungus
	Human Herpesvirus 7	HHV-7	Herpesvirus
	Legionella pneumophila	L.pne	Bacteria
	Leptospira spp.	Lepto	Bacteria
	Nocardia spp.	Nocar	Bacteria
	Scedosporium spp.	Scedo	Fungus
	Torque Teno Virus	TTV	Virus
	Vibrio spp.	Vibrio	Bacteria

2.2.3 Clustering and archetype assignment

Pairwise dissimilarities were computed using Gower distance, which accommodates ordinal traits by normalizing each trait by its range. We applied two complementary clustering approaches—partitioning around medoids (PAM) and agglomerative hierarchical clustering with Ward linkage—to identify natural groupings in the pathogen trait space. The optimal number of clusters was determined using the elbow method with angle detection applied to within-cluster sum of distances curves. To ensure

robustness, we constructed consensus cluster assignments by comparing the two methods: when both methods agreed on an assignment, we accepted it; when they disagreed, we selected the cluster with lower average trait score as a conservative tie-breaker. Organisms initially assigned to the keystone archetype underwent additional centroid-based validation and were reassigned if their total trait score was closer to another cluster's centroid. Principal component analysis was used for visualization of the high-dimensional trait space. Technical details of our clustering pipeline are presented in Supplement Sections B.1-B.4.

2.2.4 Trait signature analysis

Cluster-versus-rest comparisons used Cliff's delta for effect size and Wilcoxon rank-sum tests for significance ($p < 0.05$). Multivariate trait signatures were obtained from ℓ_1-penalized (LASSO) multinomial logistic regression with 10-fold cross-validation (Supplement B.6).

2.3 Clinical Emergence Tensor Construction (Figure 5)

To test whether trait-based archetypes predict real-world clinical behavior, we systematically curated a three-dimensional clinical emergence tensor documenting 43 pathogens across 13 anatomical niches and 31 immune perturbations (Table 2). Formally, this tensor is denoted by $\mathbf{T} \in \mathbb{R}^{43 \times 13 \times 31}$, where each element T_{ijk} represents a clinical severity score (0–100) quantifying the strength of evidence that pathogen i emerges in niche j under immune perturbation k. Scores were derived from a structured literature review using a hierarchical evidence framework: Grade A (replicated human genetic or interventional studies), Grade B (strong clinical cohort or case-control studies), Grade C (limited human evidence with mechanistic support), and Grade D (animal or in vitro evidence only). Each entry was additionally tagged with a confidence level ([High], [Moderate], [Low], [Minimal], or [None]) based on sample size, consistency across studies, and reproducibility. Scores ≥ 40 correspond to human-relevant clinical evidence (Grades A–C), defining a threshold that distinguishes clinically actionable emergence from preclinical or model-based observations and that underpins the breadth metrics used throughout the analysis. Full details of the scoring rubric and confidence assignments are provided in Supplement Section C.1.

2.3.1 Immune perturbations and anatomical niches

Immune perturbations spanned genetic immunodeficiencies (SCID, XLA, CVID, complement deficiencies) and pharmacologic immunosuppression (calcineurin inhibitors, TNF blockers, rituximab, corticosteroids), systematically covering T cell, B cell, NK cell, myeloid, and complement deficiencies. Anatomical niches included immunologically distinct sites such as CNS, respiratory and GI mucosa, skin, lymphoid organs, and systemic compartments, each defined by controlled vocabularies specifying anatomical terms and exclusion criteria to ensure niche-specific localization. This design captures how different immune failures unmask pathogens in tissue-specific ways. Complete perturbation and niche definitions are in Supplement Section C.2.

Table 2 Clinical Tensor Dimensions: Anatomical Niches and Immune Perturbations

Dimension	Description
Anatomical Niches (n=13)	
CNS	Central nervous system parenchyma and meninges
Lung	Pulmonary parenchyma and airways
GI	Gastrointestinal mucosa and lumen
Liver	Hepatic parenchyma
Blood	Bloodstream and intravascular space
LN_Paracortex	T cell zones of lymph nodes
LN_GC	B cell germinal centers of lymph nodes
Spleen	Splenic parenchyma and red/white pulp
BoneMarrow	Hematopoietic bone marrow
Skin	Cutaneous and subcutaneous tissues
GU	Genitourinary tract
Bone	Osseous tissue
Eye	Ocular tissues
Immune Perturbations (n=31)	
P01–P05	Cellular deficiencies (neutrophils, macrophages, CD4, CD8, NK)
P06–P10	Cytokine blockade (TNF, IFNγ, IL-12/23, IL-17, IL-1)
P11–P15	Complement and humoral defects (C3, properdin, MBL, Ig, B cells)
P16–P20	Innate defects (TLR, MyD88, IRAK4, NF-κB, phagocyte oxidase)
P21–P25	Antigen presentation and costimulation (MHC-I, MHC-II, CD40L, CTLA4, PD1)
P26–P31	Additional cytokine/immune modulation (IL-6, IL-10, IL-23, JAK, CSF, IL-5)

2.3.2 Feature extraction: Capturing maximal emergence and breadth

To extract features for classification and clustering analyses, we apply max-marginalization to capture maximal emergence profiles. For each pathogen, we compute the maximum severity score across all anatomical niches for each immune perturbation, yielding a pathogen × immune perturbation matrix ($\mathbf{PI_{max}}$, 43×31). Similarly, we compute the maximum score across all immune perturbations for each niche, yielding a pathogen × niche matrix ($\mathbf{PN_{max}}$, 43×13). This max-marginalization identifies the most vulnerable contexts for each pathogen under each immune deficiency and in each anatomical site. Concatenating $\mathbf{PI_{max}}$ and $\mathbf{PN_{max}}$ produces a *basic feature set* of 44 clinical variables per pathogen.

We further derive *breadth metrics* to quantify the robustness of pathogen emergence across contexts. For each pathogen, *immune breadth* counts the number of immune perturbations with maximal emergence score ≥ 40 (i.e., human-relevant clinical manifestation under that immune deficiency), and *niche breadth* counts the number of anatomical niches with maximal emergence score ≥ 40 (i.e., human-relevant clinical manifestation in that site). These breadth metrics measure how robustly a pathogen exploits immune failure across diverse immunological and anatomical contexts.

Fig. 3 Mechanistic model of pathogen emergence under immune perturbation. We score emergence through two pathways, Barrier (blue) and Latent (red), using five literature-derived inputs: latent control strength L, lytic attack strength A, reactivation/route likelihood R, perturbation effect on controllers E, and controller residency C (all normalized to $[0, 1]$, with a conservative prior for unknown edges). Pathway-specific chain scores are computed via a length-normalized geometric mean and modulated by perturbation intensity, then aggregated by collapsing to the strongest chain per controller, selecting top-K controllers, and applying a normalized noisy-OR. The final predicted emergence score is a competing-explanations mixture $T^{\mathrm{mix}} = \max(T^{\mathrm{bar}}, T^{\mathrm{lat}})$, with outputs including the dominant pathway attribution and a binned clinical rubric for comparison to observed scores.

2.3.3 Validating archetypes through clinical predictions

We asked whether clinical emergence patterns predict trait-based archetype membership using two complementary approaches. *Supervised classification* employed elastic-net logistic regression to predict archetypes from clinical features, testing both binary discrimination (keystone vs. non-keystone) and full 4-way classification. *Unsupervised clustering* applied PAM to clinical features alone, asking whether emergence patterns independently recover the four archetypes without reference to trait assignments. Agreement between unsupervised clinical clusters and trait-derived archetypes was quantified by Adjusted Rand Index. To assess robustness to evidence quality, we performed sensitivity analyses by recomputing features under two score-adjustment schemes (Supplement Section C.1): *grade-capping* (reducing scores from studies with lower evidence grades) and *grade+confidence weighting* (incorporating both evidence grade and confidence tags), then re-running all classification and clustering analyses. If archetypes are biologically meaningful, clinical emergence should predict them across scoring schemes, and clustering on clinical data should reproduce them.

2.4 Mechanistic Model (Figure 6)

We constructed a mechanistic model predicting clinical emergence from first principles of immune biology. The model integrates two complementary pathways: (i) a *latent pathway* capturing reactivation from immune-controlled reservoirs when specific controllers are disabled, and (ii) a *barrier pathway* capturing direct tissue invasion when immune barriers are compromised. Unlike statistical models, every prediction traces back to specific biological mechanisms encoded in expert-curated scoring tables.

2.4.1 Biological inputs

Five expert-curated scoring tables quantified biological interactions: latent control (strength of immune controller suppression of pathogen in latent reservoirs), lytic attack capacity (pathogen ability to breach tissue barriers), reactivation routes (probability of escape from latent to lytic niches), perturbation effects (degree to which each immunodeficiency disables each controller), and controller residency (which controllers operate in which anatomical niches). All scores were normalized to $[0, 1]$, with unknown interactions assigned a prior probability of 0.1 rather than zero to reflect biological uncertainty.

2.4.2 Pathway integration

For each pathogen–niche–perturbation combination, the model computes scores through both pathways. Chain scores use geometric means (length-normalized) with perturbation effects acting as modulatory exponents. Multiple chains through the same controller are collapsed to their maximum before aggregation via normalized noisy-OR, which prevents saturation. The final prediction takes the maximum of the two pathways, allowing either mechanism to dominate when biologically appropriate.

2.4.3 Evaluation

Model predictions were evaluated against the clinical emergence tensor using correlation coefficients and binary classification metrics at threshold $\tau = 40$. We employed three complementary evaluation frameworks: strict (zeros as true negatives), positive-unlabeled (PU) ranking (zeros as unlabeled), and confidence-masked (retaining only zeros with curated evidence of absence). Performance was stratified by pathogen archetype to assess differential predictive accuracy.

3 Results

3.1 Immunological trait structure resolves four pathogen archetypes (Fig. 4)

PCA reveals organized dispersion with interpretable loadings.

Principal component analysis of 18 immunological traits across 43 organisms produced a structured biplot in which the first two components explain 52.9% of total variance (PC1 42.2%, PC2 10.7%). Organisms spread primarily along PC1. Trait

Fig. 4 Unsupervised trait clustering resolves four pathogen immunological archetypes. (A)
PCA biplot of 18 traits across 43 organisms. The first two components explain 52.9% of variance (PC1
42.2%, PC2 10.7%). Points are colored by total score, and black arrows show trait loadings. **(B)** Gower
distance matrix, hierarchically ordered, shows block-diagonal structure that supports discrete groupings.
(C) Agglomerative hierarchical clustering with Ward linkage on Gower dissimilarities, cut at $k = 4$, reveals
four clades that correspond to the archetypes. **(D)** Elbow curves of the within-cluster sum of dissimilari-
ties for PAM and hierarchical methods independently nominate $k = 4$ for both, as indicated by the vertical
dashed lines. **(E)** Consensus cluster assignments visualized in PCA space. Medoids are shown as star sym-
bols with black outlines. The two algorithms showed substantial agreement (88%, 38/43 organisms) but
disagreed for 5 organisms. We applied a conservative consensus rule that assigns discordant organisms to
the cluster with lower average total score, prioritizing placement in less virulent archetypes. This resulted
in 4 organisms reassigned from their PAM assignment and 1 from hierarchical. Following consensus cluster-
ing, centroid-based validation of the 8 initial keystones confirmed 7 (CMV, EBV, HHV-6, HSV-1, HSV-2,
M. tuberculosis, VZV) and reassigned *P. falciparum* from *Keystone* to *Multi-host* based on proximity to
cluster centroids. **(F)** Cluster-versus-rest effect sizes by trait using Cliff's delta. Stars mark traits with sta-
tistical significance from Wilcoxon rank-sum tests (uncorrected $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).
Keystone organisms are enriched across most traits by design, reflecting coordinated, multi-arm immune
engagement. Specialists also show evidence of immune coordination but lack systemic tropism, suggesting
that such coordination is concentrated within a restricted set of anatomical niches. Multi-host pathogens
share some keystone-associated features, including heightened immune sensitivity, yet display strong sig-
nals of limited human specificity. Opportunists score low across most traits, consistent with their emergence
primarily under pathway-specific immune failure. The keystone pole aligns with postnatal imprinting of
immunodominance as described by Keystone Epitope Theory [17, 21, 25]. **(G)** Multivariate predictive sig-
nature from ℓ_1-penalized (LASSO) multinomial logistic regression with 10-fold cross-validation. Tiles show
signed coefficients for selected traits by archetype, with black dots indicating non-zero coefficients.

loading vectors separate multi-system and history traits *(length of co-evolution with
humans, global prevalence, genetic polymorphisms tracking human migration, prox-
imity and persistence of immune cells, structured tropism, multi-arm coordination)*
from epitope-centric traits *(epitope presentation dynamics, epitope structure and
immunodominance, structured immune receptor selection)*. This pattern indicates
non-random organization in the trait space (Fig. 4A).

Pairwise distances display block structure consistent with discrete groups.

The hierarchically ordered Gower distance matrix shows clear block-diagonal structure, with dark blocks along the diagonal and lighter off-block regions. This pattern supports the presence of discrete clusters rather than a single continuum (Fig. 4B).

Four clusters capture the dominant structure.

Both clustering strategies—partitioning around medoids (PAM) and agglomerative hierarchical clustering with Ward linkage on Gower dissimilarities—were evaluated for $k = 2, \ldots, 8$. The elbow method with angle detection applied to the within-cluster sum of distances independently nominated $k = 4$ for both methods, after which additional clusters offer diminishing returns (Fig. 4D).

Method concordance and conservative consensus labeling.

Hierarchical clustering yields four coherent clades that align closely with PAM clusters (Fig. 4C,E). The two methods showed substantial agreement (88%, 38/43 organisms), quantified by identical cluster assignments. For the 5 organisms receiving different labels across methods, we adopted a conservative consensus rule: we compared the average total trait score of the two candidate clusters and assigned the organism to the cluster with the *lower* average score, prioritizing placement in less virulent archetypes. This approach resulted in 4 organisms being reassigned from their PAM cluster assignment and 1 organism from its hierarchical assignment. All downstream analyses use these consensus labels.

Centroid-based validation confirms keystone assignments.

To guard against false-positive keystone classifications, consensus keystone assignments underwent additional centroid-based validation. Eight organisms were initially assigned to the *Keystone* cluster. For each, we calculated its distance to all cluster centroids (mean total scores). Seven keystones (CMV, EBV, HHV-6, HSV-1, HSV-2, *M. tuberculosis*, VZV) were confirmed as closest to the keystone centroid. One organism, *P. falciparum*, showed a total trait score (22) closer to the multi-host centroid than the keystone centroid and was therefore reassigned to *Multi-host*. This validation step was applied exclusively to keystone assignments; all other consensus assignments were accepted without additional scrutiny.

Cluster semantics and biological interpretation.

The final validated clustering yielded four archetypes with distinct trait profiles and sizes. *Opportunistic* (n=15, mean trait score 7.0) organisms show low values across most traits, consistent with limited immunological specialization. *Multi-host* (n=10, mean score 19.2) pathogens are enriched for broad host-range characteristics and hypersensitivity/autoimmune associations. *Specialist* (n=11, mean score 19.2) organisms emphasize epitope-centric strategies, with high epitope presentation dynamics and immunodominance, coupled with the lowest structured tropism among groups. *Keystone* (n=7, mean score 30.0) pathogens exhibit high scores on multi-arm coordination, structured tropism, length of co-evolution, and global prevalence. These interpretations match the distribution of organisms in PCA space and

their hierarchical separation (Fig. 4A,C,E). High scores on multi-arm coordination and structured tropism in the keystone group constitute the trait-level signature of postnatal imprinting: these organisms plausibly coordinate Class I and II targeting and sustain tissue-resident memory in defined niches across life.

Trait signatures, univariate and multivariate.

Cluster-versus-rest effect sizes computed with Cliff's delta highlight distinctive single-trait signatures, with statistical significance assessed using Wilcoxon rank-sum tests and uncorrected p-values at $\alpha = 0.05$ to maintain sensitivity for exploratory trait discovery (Fig. 4F). Multivariate modeling with ℓ_1-penalized (LASSO) multinomial logistic regression, with regularization parameter selected by 10-fold cross-validation, yields a compact predictive signature that agrees with the univariate patterns: keystone membership is driven by multi-system and history traits, specialists by epitope-centric traits, and multi-host by hypersensitivity, with opportunists characterized by negative coefficients across most features (non-zero coefficients marked with black dots, Fig. 4G).

3.2 Clinical emergence patterns validate trait-based archetypes (Fig. 5)

Supervised classification confirms clinical-trait concordance.

We tested whether clinical emergence patterns align with trait-based archetype assignments using elastic-net logistic regression ($\alpha = 0.5$). Binary classification (Keystone vs. Other) on the 44 basic features achieved perfect discrimination: 100% accuracy, 100% sensitivity (7/7 keystones correctly identified), 100% specificity (36/36 non-keystones correctly identified), and AUC=1.000 (Fig. 5D). This perfect concordance demonstrates that clinical emergence patterns fully distinguish keystones from other archetypes. Multi-class classification (4 archetypes) achieved 67.4% overall accuracy and 63.8% macro-averaged recall (Fig. 5E). Per-class recall was 100% (Opportunistic, 15/15), 71% (Keystone, 5/7), 64% (Specialist, 7/11), and 30% (Multi-host, 3/10). Misclassifications were non-random: all 15 opportunists were correctly identified, reflecting their distinctive low-severity clinical profile. Multi-host pathogens showed the highest ambiguity, with 50% (5/10) misclassified as opportunistic and 20% (2/10) as specialist, consistent with their intermediate trait profiles and variable human adaptation. Two keystones were misclassified (one as specialist, one as opportunistic), and four specialists were misclassified as opportunistic, reflecting partial overlap in clinical footprints among non-keystone archetypes. Overall, these results demonstrate strong alignment between clinical and trait-based archetype definitions, with perfect keystone identification and moderate-to-high concordance for other archetypes.

Unsupervised clustering on clinical features recovers archetype structure.

To test whether clinical data independently recover the trait-based organization, we applied PAM clustering with Gower distances on the 44 clinical features, setting $k = 4$ to match the trait-derived solution. The clinical-derived clusters showed substantial

Fig. 5 Clinical emergence patterns validate and refine trait-based pathogen archetypes. (A–C) Max-marginalized views of the three-dimensional clinical tensor ($43 \times 13 \times 31$: pathogens $\times$ anatomical niches $\times$ immune perturbations). Each tensor element represents clinical severity (0–100) under specific pathogen-niche-perturbation contexts, derived from systematic literature curation. **(A)** Pathogen $\times$ Immune perturbation matrix (43×31) obtained by max-marginalizing over niches, capturing the most severe clinical manifestation for each pathogen under each immune deficiency. **(B)** Pathogen $\times$ Niche matrix (43×13) obtained by max-marginalizing over immune perturbations, capturing the most severe manifestation in each anatomical site. **(C)** Immune perturbation $\times$ Niche matrix (31×13) obtained by max-marginalizing over pathogens, revealing which anatomical sites are most vulnerable under specific immune deficiencies. Heatmaps use intensity encoding (0=white, 100=maximum color saturation). Rows and columns are hierarchically ordered by Euclidean distance to reveal block structure. **(D)** Binary classification (Keystone vs. Other) using elastic-net logistic regression ($\alpha = 0.5$) on basic clinical features (max-marginalized pathogen $\times$ niche and pathogen $\times$ immune scores, 44 features total). Confusion matrix shows row-normalized proportions. Perfect classification (Accuracy: 100%, Recall: 100%, AUC: 1.000) demonstrates complete concordance between clinical emergence patterns and trait-based keystone identification. **(E)** Multi-class classification (4 archetypes) using elastic-net multinomial logistic regression ($\alpha = 0.5$) on the same 44 basic features. Confusion matrix shows row-normalized proportions. Overall accuracy is 67.4% with macro-averaged recall of 63.8%. Per-class recall: Opportunistic 100% (15/15), Keystone 71% (5/7), Specialist 64% (7/11), Multi-host 30% (3/10). Misclassifications are non-random: all 15 opportunists are correctly identified, while multi-host pathogens show the most ambiguity, with 50% misclassified as opportunistic and 20% as specialist, consistent with their intermediate trait profiles. **(F)** Unsupervised clustering via PAM with Gower distances on clinical features independently recovers structure highly concordant with trait-based archetypes. Adjusted Rand Index (ARI) = 0.58 and Normalized Mutual Information (NMI) = 0.63 quantify substantial agreement between clinical-derived and trait-derived cluster assignments, confirming that clinical emergence patterns and immunological traits capture overlapping biological organization. **(G)** Niche breadth distributions by archetype (threshold $\geq$ 40, counting anatomical niches with human-relevant clinical severity). Violin plots with overlaid boxplots and individual points show archetype-specific patterns. Keystone pathogens exhibit significantly higher niche breadth (mean=8.0, median=8) compared to all other archetypes (Kruskal-Wallis H=16.37, $p < 0.001$), reflecting their capacity for multi-site dissemination. Post-hoc pairwise tests (Benjamini-Hochberg adjusted) confirm Keystone > Multi-host ($p = 0.050$), Keystone > Opportunistic ($p = 0.003$), and Keystone > Specialist ($p = 0.024$). By contrast, immune perturbation breadth shows no significant archetype differences (H=6.93, $p = 0.074$), indicating that anatomical dissemination, rather than breadth of immune vulnerability, is the clinical hallmark of keystones. **(H)** Two-dimensional robustness landscape plotting niche breadth against immune perturbation breadth (threshold $\geq$ 40). Each point represents one pathogen, colored by archetype. Large diamonds mark archetype centroids, and shaded ellipses show 75% confidence regions. X-axis is truncated at 16 to maximize visualization of the main cluster; Nocardia (immune breadth = 25) is annotated with an arrow. Keystone pathogens occupy the upper-right quadrant with elevated niche breadth, while opportunists cluster in the lower-left with restricted clinical footprints. Multi-host and specialist archetypes show intermediate and overlapping patterns, consistent with their mixed trait profiles.

agreement with trait-based archetypes: Adjusted Rand Index (ARI) = 0.58 and Normalized Mutual Information (NMI) = 0.63 (Fig. 5F). These values indicate significant overlap between clinical and trait-based groupings, confirming that the two feature spaces capture related biological structure. Discordances between clinical and trait clustering highlight cases where immunological potential (traits) and realized clinical behavior (tensor) diverge, potentially reflecting host demographic factors, pathogen strain variation, or incomplete clinical characterization.

Breadth metrics quantify archetype-specific robustness.

To characterize the clinical footprint of each archetype, we computed breadth metrics: for each pathogen, niche breadth counts the number of anatomical sites with severity score > 40, and immune breadth counts the number of immune perturbations with score > 40. Summary statistics by archetype reveal distinct patterns. Keystone pathogens exhibit the highest niche breadth (mean=8.0, median=8, range 5–11 out of 13 possible niches), significantly exceeding Opportunistic (mean=3.3, median=4), Multi-host (mean=5.4, median=5.5), and Specialist (mean=3.4, median=2). Kruskal-Wallis tests confirm highly significant niche breadth differences across archetypes (H=16.37, df=3, $p < 0.001$, Fig. 5G). Post-hoc pairwise Wilcoxon tests with Benjamini-Hochberg correction show Keystone significantly higher than all other groups: Keystone vs. Opportunistic ($p = 0.003$), Keystone vs. Multi-host ($p = 0.050$), and Keystone vs. Specialist ($p = 0.024$). By contrast, immune perturbation breadth shows no significant archetype differences (H=6.93, df=3, $p = 0.074$), with overlapping distributions across groups (Opportunistic mean=7.1, Multi-host mean=9.1, Specialist mean=7.2, Keystone mean=9.9). This asymmetry indicates that anatomical dissemination—not breadth of immune vulnerability—is the clinical hallmark distinguishing keystones from other archetypes.

Two-dimensional robustness landscape visualizes archetype separation.

Plotting niche breadth against immune breadth reveals archetype-specific regions in a two-dimensional robustness landscape (Fig. 5H). Keystone pathogens occupy the upper-right quadrant with elevated niche breadth (median=8) and moderate-to-high immune breadth (median=10), reflecting their capacity for multi-site dissemination and broad immune engagement. Opportunistic pathogens cluster in the lower-left with restricted niche breadth (median=4) and immune breadth (median=5), consistent with limited clinical manifestations outside specific immune deficiencies. Multi-host and Specialist archetypes show intermediate and overlapping patterns: multi-host pathogens have moderate niche breadth (median=5.5) and immune breadth (median=10), while specialists have low niche breadth (median=2) but variable immune breadth (median=7), reflecting their epitope-centric strategies within restricted anatomical compartments. One outlier, Nocardia (Opportunistic), exhibits exceptionally high immune breadth (25 out of 31 perturbations) yet low niche breadth (5), likely reflecting its status as a ubiquitous environmental saprophyte that opportunistically infects across diverse immune deficiency contexts without anatomical specialization.

Biological interpretation: clinical validation of immunological archetypes.

The concordance between trait-based and clinical-derived organization—demonstrated by perfect binary classification, 67% multi-class accuracy, and ARI=0.58 in unsupervised clustering—provides strong evidence that immunological traits and clinical emergence patterns reflect overlapping biological structure. Keystone pathogens, defined by coordinated multi-arm immune engagement and structured tropism in the trait space, manifest clinically as broad anatomical disseminators with human-relevant disease across many niches (mean niche breadth = 8.0). This pattern aligns with the hypothesis that keystones undergo postnatal imprinting across multiple tissue sites, establishing coordinated CD4/CD8 memory and driving structured immune receptor selection. Specialists, characterized by epitope-centric traits, show restricted niche breadth (mean=3.4) but maintain immunological coordination within those niches. Multi-host pathogens exhibit intermediate clinical footprints (niche breadth mean=5.4) and the highest classification ambiguity (30% recall), consistent with their mixed trait profiles reflecting broad host range but variable human adaptation. Opportunists, scoring low on most traits, show correspondingly restricted clinical manifestations (niche breadth mean=3.3) yet paradoxically achieved perfect classification accuracy, reflecting their distinctive low-severity profile. The asymmetry between niche breadth (strongly archetype-discriminative, $p < 0.001$) and immune breadth (non-discriminative, $p = 0.074$) suggests that anatomical dissemination capacity, rather than the number of immune vulnerabilities, is the critical clinical dimension distinguishing keystones.

3.3 Mechanistic model predicts clinical emergence from biological first principles (Fig. 6)

Two pathways capture distinct emergence mechanisms.

The mechanistic model generated predictions for 1,522 of 17,329 tensor cells (8.8%), with the remaining cells having no mechanistic path connecting the pathogen–niche–perturbation combination. Among cells with predictions, the barrier pathway dominated in 890 cases (58.5%), the latent pathway dominated in 283 cases (18.6%), and the two pathways contributed equally in 349 cases (22.9%) (Fig. 6A). This distribution indicates that both pathways contribute meaningfully, with barrier-mediated invasion being somewhat more common than reactivation-based emergence in our pathogen set.

Model achieves moderate correlation with clinical observations.

Evaluated on the full 17,329-cell tensor using strict criteria (zeros treated as true negatives), the model achieved Pearson $r = 0.35$, Spearman $\rho = 0.30$, and Cohen's weighted $\kappa = 0.24$ (fair agreement) (Fig. 6B,C). Binary classification at threshold $\tau = 40$ yielded recall of 0.20, specificity of 0.97, precision of 0.42, and F1 = 0.26. The calibration curve (Fig. 6C) shows that predicted scores track observed medians, though with substantial IQR reflecting inherent clinical variability.

Fig. 6 Mechanistic model predicts pathogen emergence from immune biology first principles.
(A) Pathway contribution analysis comparing latent vs. barrier pathway scores for 1,522 tensor cells with mechanistic predictions. Points colored by dominant pathway: Barrier (blue, 890 cells, 58.5%), Latent (red, 283 cells, 18.6%), or Equal (gray, 349 cells, 22.9%). Diagonal line indicates equal contribution. **(B)** Confusion matrix of coarse-binned predictions (bins: 0, 5, 10, ..., 100) showing model calibration across the score range. Color intensity (log scale) indicates cell count. The model correctly identifies high-severity combinations while maintaining high specificity. **(C)** Calibration curve showing median observed score per predicted bin (point size proportional to count; error bars show interquartile range). Dashed diagonal indicates perfect calibration. Overall correlation: $r = 0.35$, $\rho = 0.30$, $\kappa = 0.24$. **(D)** Correlation coefficients by pathogen archetype. Keystone pathogens show highest predictive accuracy ($r = 0.39$), consistent with their immune-dependent emergence patterns. Dotted line indicates overall correlation. Error bars show 95% CI from bootstrap. **(E)** Enrichment analysis showing fold-improvement over random baseline at various prediction thresholds. Positives defined as $T_{obs} \geq 40$ (baseline rate: 7.3%). At top 5% of predictions, the model achieves ~2.9× enrichment. Lines colored by archetype; horizontal dashed line indicates random baseline (1×). **(F)** Pathogen × Niche (PN) max-marginalized projection achieves strong correlation ($r = 0.58$, $\rho = 0.56$), indicating the model captures tissue tropism relationships particularly well. Each point represents one pathogen–niche combination (maximum across all immune perturbations). Red dashed line: identity; green line: linear fit with 95% CI.

Keystone pathogens show highest predictive accuracy.

Stratification by archetype revealed differential model performance (Fig. 6D). Keystone pathogens achieved the highest correlation ($r = 0.39$, $\rho = 0.36$), followed by Opportunistic ($r = 0.36$, $\rho = 0.32$), Multi-host ($r = 0.33$, $\rho = 0.31$), and Specialist ($r = 0.33$, $\rho = 0.23$). This pattern is consistent with the hypothesis that keystone pathogens, by definition, exhibit immune-dependent emergence patterns that the mechanistic model is designed to capture.

Model enriches true positives among top predictions.

To assess ranking utility in a positive-unlabeled framework—appropriate given that zero scores may reflect absence of evidence rather than evidence of absence—we computed fold-enrichment at various prediction thresholds (Fig. 6E). At the top 5% of

predictions, the model achieved approximately 2.9-fold enrichment over random baseline, meaning top-ranked predictions are nearly three times more likely to be true positives than random selection. Enrichment remained stable across thresholds (2.8–3.0× from top 1% to top 20%), indicating robust ranking performance. Keystone and Multi-host pathogens showed the strongest enrichment signals, consistent with their immune-coordinated emergence patterns.

Pathogen–niche projection achieves strong correlation.

To reduce noise from the immune perturbation dimension, we computed max-marginalized 2D projections (Fig. 6F). The pathogen × niche (PN) projection, which captures each pathogen's maximum emergence score across all immune perturbations for each anatomical site, achieved substantially higher correlation ($r = 0.58$, $\rho = 0.56$) than the full 3D tensor. The pathogen × immune perturbation (PI) projection achieved $r = 0.36$, $\rho = 0.37$. The superior performance of the PN projection suggests the model captures pathogen–tissue tropism relationships particularly well—a biologically meaningful finding, as pathogens have characteristic tissue preferences that the model successfully encodes through lytic attack capacity and controller residency patterns.

Biological interpretation: mechanistic validation of archetype-specific emergence.

The mechanistic model's differential performance across archetypes provides biological validation of the trait-based classification. Keystone pathogens, defined by coordinated multi-arm immune engagement, show the highest correlation because their emergence patterns are precisely what the model was designed to capture: release from immune control leading to predictable clinical manifestations. The finding that both latent (18.6%) and barrier (58.5%) pathways contribute meaningfully indicates the model captures genuine biological heterogeneity in emergence mechanisms. The 2.9-fold enrichment at top predictions demonstrates practical utility: the mechanistic model can prioritize pathogen–niche–perturbation combinations for clinical surveillance or therapeutic intervention. Importantly, cases where the model predicts high emergence but clinical data shows zero ("predicted high, observed zero") may represent not model failures but rather unstudied combinations, incomplete clinical characterization, or absence of pathogen exposure in studied populations—possibilities strongly supported by our finding that all zero scores have low confidence (no curated evidence of absence).

4 Discussion

Keystone organisms as an operational immunological phenotype.

This work moves immunological "keystone organisms" from an ecological metaphor to an operational phenotype by triangulating (i) trait biology, (ii) clinical emergence under immune perturbation, and (iii) a first-principles mechanistic perturbation model. Across 43 pathogens, trait-space clustering yielded four reproducible archetypes and a compact set of human-adapted keystones dominated by persistent herpesviruses and *M. tuberculosis* (Fig. 4). These organisms converge on long-term residence,

tissue-structured tropism, and the need for coordinated control spanning antigen presentation, innate sensing, and T/NK effector programs.

Clinical emergence distinguishes keystones by dissemination, not simply by "more immunosuppression triggers."

By encoding published associations as a pathogen×immune-perturbation×niche tensor, we show that the clinically salient signature of keystones is their ability to occupy a broader repertoire of anatomical sites when immune coordination is perturbed (Fig. 5). In our dataset, diagnostic breadth decomposed into immune breadth and niche breadth, and it is niche breadth that separates keystones from other archetypes. This pattern argues for compartmentalized, tissue-resident checkpoints (e.g., mucosal barrier integrity, myeloid restriction, TRM/NK patrol) as key governors of keystone containment. Practically, it helps explain why patients with comparable "global" immunosuppression can present with distinct pathogen portfolios: the decisive variable is which immune functions fail in which tissue compartment, not only the magnitude of systemic lymphopenia.

A first-principles model recapitulates emergence and prioritizes risk.

The mechanistic model integrates three recurring levers of opportunism—barrier disruption, activation of latent/persistent reservoirs, and loss of local immune control—and uses organism-, niche-, and perturbation-level features derived from the tensor. Despite its simplicity, it reproduced observed emergence patterns (overall $r = 0.35$) and performed best when aggregated across niches (pathogen×niche projection $r = 0.58$), suggesting that the dominant learnable structure is anatomical rather than purely immunological. Importantly, the model enriched the highest-risk pathogen–immune–niche triplets 2.9-fold, supporting its role as a hypothesis generator for surveillance, prophylaxis, and mechanistically targeted diagnostic workups (Fig. 6).

From taxonomy to translation: why keystones matter clinically.

Persistent viromes and bacteriomes are increasingly recognized as contributors to immune calibration and host fitness, not merely as threats [2–4]. Our results extend this view by identifying a subset of long-term residents that behave like immunological organizers: they are constrained by multi-arm, multi-compartment control, and when that control is relaxed they become sentinels that reveal where immune coordination has failed. This framing can help contextualize heterogeneous infectious complications across transplantation, biologic immunomodulation, autoimmunity, and inborn errors of immunity [11, 12].

Box 1. Design and Clinical Use Guidelines

1. **Treat keystone emergence as a systems readout.** Because keystone organisms are long-term residents with tissue-structured reservoirs, their *reactivation* (and especially their appearance in non-canonical compartments) is often less informative as an isolated microbiologic event than as a coarse *systems-level* readout of immune coordination failure. In practice,

---

the *identity* of the keystone (which control programs it typically demands) and the *niche* in which it emerges (which barrier and tissue-resident checkpoints have failed) can be interpreted as a structured signal of which immune arms and compartmental defenses are compromised [4]. This is most useful when applied longitudinally (baseline → perturbation → recovery), and when detections are counted as *episodes* (unique organism–compartment–time events), rather than repeated tests of the same episode.

A practical instantiation is a *Keystone Reactivation Index* (*Keystone Reactivation Index*), defined within a compartment c over a time window t as the fraction of keystone detections among all infectious detections:

$$Keystone\ Reactivation\ Index_{c,t} = \frac{K_{c,t}}{D_{c,t}} = \frac{\left|\mathcal{D}_{c,t}^{(K)}\right|}{|\mathcal{D}_{c,t}|}, \tag{1}$$

where $\mathcal{D}_{c,t}$ is the set of infectious detection episodes observed in compartment c during window t, $\mathcal{D}_{c,t}^{(K)} \subseteq \mathcal{D}_{c,t}$ is the subset classified as keystone organisms, $D_{c,t} = |\mathcal{D}_{c,t}|$, and $K_{c,t} = |\mathcal{D}_{c,t}^{(K)}|$. (When $D_{c,t} = 0$, *Keystone Reactivation Index* is undefined and can be treated as missing rather than zero.)

Optionally, a patient-level summary over compartments can be computed as

$$Keystone\ Reactivation\ Index_t = \frac{\sum_c K_{c,t}}{\sum_c D_{c,t}}, \tag{2}$$

which preserves interpretability while allowing longitudinal tracking of keystone-weighted infectious burden.

2. **Prioritize containment in the compartment that declares itself.** Because keystone signatures are tissue-structured, prophylaxis and monitoring should be tailored to the affected niche (e.g., CNS, gut, lung), not only to systemic immune cell counts.

3. **Avoid immunogens dominated by mutable decoys.** Vaccine or immunotherapy designs that repeatedly present highly variable epitopes may preferentially recall familiar but non-protective memory, diverting responses away from conserved constrained targets [22–24].

4. **Screen for keystone-patterned cross-reactivity in high-risk settings.** In individuals prone to severe T-cell–mediated hypersensitivity, autoimmunity, or transplantation rejection, evaluating whether candidate drugs or alloantigens generate peptides resembling tissue-local keystone epitopes may identify risk for pathogenic memory recall [18–21, 26, 27].

Keystone-focused memory provides a mechanistic bridge to immunopathology.

At the micro level, long-lived, tissue-homing memory to persistent keystones is advantageous for rapid control, but it also creates a privileged path for "modified-self"

ligands to recruit potent effector programs in the same compartment. This offers an immunological explanation for why some drug hypersensitivity and organ-specific autoimmune syndromes are strongly HLA- and tissue-linked: they may reflect misdirected recall of keystone-trained T-cell repertoires [18–21, 26, 27]. This prediction is testable: culprit HLA–peptide complexes should preferentially engage memory clonotypes with prior keystone specificity and appropriate tissue imprinting.

Implications for fast-evolving pathogens and immunogen design.

Immune hierarchies shaped by persistent infections can be exploited by rapidly adapting RNA viruses, which may inflate immunodominant yet mutable epitopes that trigger recall without durable control [22–25]. The archetype framework sharpens this claim: decoys should be most effective when they mimic the "keystone pattern" of presentation and tissue context that the host has learned to prioritize. Immunogen design that forces recognition of conserved, functionally constrained sites—and avoids over-amplifying keystone-like decoys—may therefore be essential for durable protection.

Limitations and next steps.

Trait scoring compresses heterogeneous biology into rubric values, and the clinical tensor is literature-derived, subject to reporting bias, evolving diagnostics, and non-independence of case series. Our classifiers and mechanistic model were validated internally; external prospective validation in well-annotated cohorts is required before clinical deployment. Nonetheless, the convergence of trait-based archetypes, clinical emergence structure, and first-principles modeling suggests that keystone organisms are not an artifact of any single data source. Next steps include expanding the Keystone Protein Atlas in a compartment-aware manner, integrating epitope-level constraints and host genetics, and performing prospective surveillance under defined immunomodulatory regimens to refine the mapping from perturbation to niche-specific risk [6, 7, 14, 15, 28, 29].

5 Conclusion

We define immunological keystone organisms as a reproducible class of human-adapted pathogens whose containment depends on coordinated, multi-compartment immunity and whose failures are clinically legible. By combining trait-based archetyping, a literature-derived clinical emergence tensor, and a simple perturbation model, we show that keystone status is reflected in distinctive patterns of tissue dissemination under immune deficits, and that these patterns can be predicted from biological first principles. This framework enables archetype-aware monitoring in immunosuppressed patients, generates testable mechanistic hypotheses about tissue-specific vulnerability and immune-mediated pathology, and provides a rational basis for immunogen design that prioritizes conserved, functionally constrained targets.

Code Availability

All code and data used to generate the results in this study is publicly available at https://github.com/AsiaeeLab/keystone-organisms.

Acknowledgements

This work was supported in part by the National Institutes of Health (NIH) under awards AI154659, P50GM115305, 5P30AI110527, 1R35HL140016, 5R01AI060460, U54AG089326, P01HL174442, P30CA068485, R01AI077505, 1R01AI147765, 1R01AI142093, and R00HG011367.

References

[1] Margulis, L.: Origin of Eukaryotic Cells. Yale University Press, New Haven (1970)

[2] Roossinck, M.J.: The good viruses: viral mutualistic symbioses. Nature Reviews Microbiology **9**(2), 99–108 (2011) https://doi.org/10.1038/nrmicro2491

[3] Pradeu, T.: Mutualistic viruses and the heteronomy of life. Studies in History and Philosophy of Biological and Biomedical Sciences **59**, 80–88 (2016) https://doi.org/10.1016/j.shpsc.2016.02.007

[4] Virgin, H.W., Wherry, E.J., Ahmed, R.: Redefining chronic viral infection. Cell **138**(1), 30–50 (2009) https://doi.org/10.1016/j.cell.2009.06.036

[5] Paine, R.T.: A note on trophic complexity and community stability. The American Naturalist **103**(929), 91–93 (1969) https://doi.org/10.1086/282586

[6] Hislop, A.D., Taylor, G.S., Sauce, D., Rickinson, A.B.: Cellular responses to viral infection in humans: lessons from epstein-barr virus. Annu. Rev. Immunol. **25**(1), 587–617 (2007) https://doi.org/10.1146/annurev.immunol.25.022106.141553

[7] Malouli, D., Hansen, S.G., Nakayasu, E.S., Marshall, E.E., Hughes, C.M., Ventura, A.B., Gilbride, R.M., Lewis, M.S., Xu, G., Kreklywich, C., *et al.*: Cytomegalovirus pp65 limits dissemination but is dispensable for persistence. The Journal of clinical investigation **124**(5), 1928–1944 (2014) https://doi.org/10.1172/JCI67420

[8] Abdallah, F., Mijouin, L., Pichon, C.: Skin immune landscape: inside and outside the organism. Mediators of inflammation **2017**(1), 5095293 (2017) https://doi.org/10.1155/2017/5095293

[9] Quaresma, J.A.S.: Organization of the skin immune system and compartmentalized immune responses in infectious diseases. Clinical microbiology reviews **32**(4), 10–1128 (2019) https://doi.org/10.1128/CMR.00034-18

[10] Handfield, C., Kwock, J., MacLeod, A.S.: Innate antiviral immunity in the skin. Trends in immunology **39**(4), 328–340 (2018) https://doi.org/10.1016/j.it.2018.02.003

[11] Tangye, S.G., Al-Herz, W., Bousfiha, A., Chatila, T., Cunningham-Rundles, C., Etzioni, A., Franco, J.L., Holland, S.M., Klein, C., Morio, T., *et al.*: Human inborn errors of immunity: 2019 update on the classification from the international union of immunological societies expert committee. Journal of clinical immunology **40**(1), 24–64 (2020) https://doi.org/10.1007/s10875-019-00737-x

[12] Notarangelo, L.D., Bacchetta, R., Casanova, J.-L., Su, H.C.: Human inborn errors of immunity: an expanding universe. Science immunology **5**(49), 1662 (2020) https://doi.org/10.1126/sciimmunol.abb1662

[13] McGeoch, D.J., Rixon, F.J., Davison, A.J.: Topics in herpesvirus genomics and evolution. Virus research **117**(1), 90–104 (2006) https://doi.org/10.1016/j.virusres.2006.01.002

[14] Wertheim, J.O., Smith, M.D., Smith, D.M., Scheffler, K., Kosakovsky Pond, S.L.: Evolutionary origins of human herpes simplex viruses 1 and 2. Molecular biology and evolution **31**(9), 2356–2364 (2014) https://doi.org/10.1093/molbev/msu185

[15] Correia, S., Bridges, R., *et al.*: Sequence variation of epstein-barr virus: Viral types, geography, codon usage, and diseases. Journal of Virology **92**(22), 01132–18 (2018) https://doi.org/10.1128/JVI.01132-18

[16] Dominguez-Andres, J., Netea, M.G.: Long-term reprogramming of the innate immune system. Journal of leukocyte biology **105**(2), 329–338 (2019) https://doi.org/10.1002/JLB.MR0318-104R

[17] Mallal, S., Asiaee, A.: Keystone Theory: Implications for Effective T and Natural Killer Cell Interoperability. Zenodo (2025). https://doi.org/10.5281/zenodo.17087093

[18] Ostrov, D.A., Grant, B.J., *et al.*: Drug hypersensitivity caused by alteration of the HLA-presented self-peptide repertoire. Proceedings of the National Academy of Sciences **109**(25), 9959–9964 (2012) https://doi.org/10.1073/pnas.1207934109

[19] Pavlos, R., Mallal, S., Ostrov, D., Buus, S., Metushi, I., Peters, B., Phillips, E.: T cell–mediated hypersensitivity reactions to drugs. Annual review of medicine **66**(1), 439–454 (2015) https://doi.org/10.1146/annurev-med-050913-022745

[20] Phillips, E.J., Mallal, S.A., *et al.*: Active suppression rather than ignorance: tolerance to abacavir-induced hla-b* 57: 01 peptide repertoire alteration. The Journal of Clinical Investigation **128**(7), 2746–2749 (2018) https://doi.org/10.1172/JCI121525

[21] Mallal, S., Phillips, E.: Keystone Epitope Theory: Implications for Hypersensitivity, Autoimmunity, and Transplantation. Zenodo (2025). https://doi.org/10.5281/zenodo.17880168

[22] Hertz, T., Nolan, D., James, I., John, M., Gaudieri, S., Phillips, E., Huang, J.C., Riadi, G., Mallal, S., Jojic, N.: Mapping the landscape of host-pathogen coevolution: Hla class i binding and its relationship with evolutionary conservation in human and viral proteins. Journal of virology **85**(3), 1310–1321 (2011) https://doi.org/10.1128/jvi.01966-10

[23] Kiepiela, P., Ngumbela, K., *et al.*: Dominant influence of hla-b in mediating the potential co-evolution of hiv and hla. Nature **432**(7018), 769–775 (2004) https://doi.org/10.1038/nature03113

[24] Almeida, C.-A.M., Bronke, C., Roberts, S.G., McKinnon, E., Keane, N.M., Chopra, A., Kadie, C., Carlson, J., Haas, D.W., Riddler, S.A., *et al.*: Translation of hla–hiv associations to the cellular level: Hiv adapts to inflate cd8 t cell responses against nef and hla-adapted variant epitopes. The Journal of Immunology **187**(5), 2502–2513 (2011) https://doi.org/10.4049/jimmunol.1100691

[25] Mallal, S., Asiaee, A.: Keystone Epitope Theory: An Ecological Perspective on RNA Viruses, Tumor Immunoediting, and Vaccine Design. Zenodo (2025). https://doi.org/10.5281/zenodo.17915417

[26] Metushi, I., Uetrecht, J., Phillips, E.: Mechanism of isoniazid-induced hepatotoxicity: then and now. British journal of clinical pharmacology **81**(6), 1030–1036 (2016) https://doi.org/10.1111/bcp.12885

[27] Gibson, A., Ram, R., Gangula, R., Li, Y., Mukherjee, E., Palubinsky, A.M., Campbell, C.N., Thorne, M., Konvinse, K.C., Choshi, P., *et al.*: Multiomic single-cell sequencing defines tissue-specific responses in stevens-johnson syndrome and toxic epidermal necrolysis. Nature communications **15**(1), 8722 (2024) https://doi.org/10.1038/s41467-024-52990-3

[28] Kodaman, N., Pazos, A., *et al.*: Human and helicobacter pylori coevolution shapes the risk of gastric disease. Proceedings of the National Academy of Sciences **111**(4), 1455–1460 (2014) https://doi.org/10.1073/pnas.1318093111

[29] Munoz-Ramirez, Z.Y., Pascoe, B., Mendez-Tenorio, A., Mourkas, E., Sandoval-Motta, S., Perez-Perez, G., Morgan, D.R., Dominguez, R.L., Ortiz-Princz, D., Cavazza, M.E., *et al.*: A 500-year tale of co-evolution, adaptation, and virulence: Helicobacter pylori in the americas. The ISME journal **15**(1), 78–92 (2021) https://doi.org/10.1038/s41396-020-00758-0

A Summary Tables of Pathogen Characteristics

Tables provide detailed immunological and evolutionary characteristics for each pathogen, organized by archetype assignment. For each organism, we document infection pattern, geographic distribution, key immune dependencies, capacity for lifelong infection, primary tissues and cell types involved, clinical impact under immunosuppression, and estimated co-evolutionary time with humans.

Table 3 Keystone organisms and their immunological and evolutionary characteristics. These pathogens exhibit multi-arm immune coordination, structured tissue tropism, and deep co-evolution with humans. Reactivation in immunosuppressed hosts reflects systems-level failure. Abbreviations in parentheses match those used in Figures and trait score tables.

Organism (Abbr.)	Type	Infection Pattern	Geo. Dist.	Key Immune Dependencies	Lifelong?	Tissues/Cell Involved	Immunosuppression Impact	Co-evo. Time
Cytomegalovirus (CMV)	Herpesvirus	Chronic latency	Global	NK, $CD8^+$, $CD4^+$, Abs	Yes	Monocytes/Macrophages, Endothelium	Post-transplant reactivation; severe disease with T/NK suppression	5–25 Mya
Epstein–Barr Virus (EBV)	Herpesvirus	Chronic latency	Global	$CD8^+$, NK; $CD4^+$ help	Yes	Memory B cells, Oropharyngeal epithelium	PTLD with B-cell depletion; risk with $CD8^+$ defects	12–14 Mya
Herpes Simplex Virus 1 (HSV-1)	Herpesvirus	Chronic latency	Global	$CD8^+$ (TRM), NK; $CD4^+$ help	Yes	Mucosal epithelium, Sensory neurons	Severe mucocutaneous/CNS disease in NK or $CD8^+$ defects	~6 Mya
Herpes Simplex Virus 2 (HSV-2)	Herpesvirus	Chronic latency	Global	$CD8^+$, NK; $CD4^+$ help	Yes	Genital mucosa, Sensory neurons	Severe genital/CNS disease in NK or $CD8^+$ defects	<1.6 Mya
Human Herpesvirus 6 (HHV-6)	Herpesvirus	Chronic latency	Global	$CD4^+$, $CD8^+$, NK	Yes	$CD4^+$ T cells, CNS	Encephalitis/reactivation post-transplant; risk with $CD4^+$ loss	~30 Mya
Mycobacterium tuberculosis (Mtb)	Bacteria	Intracellular persistence	Global	$CD4^+$, TNF, IFN-γ	Yes	Alveolar macrophages, Granulomas	Reactivation with TNF blockade or $CD4^+$ loss	0.07–3 Mya
Varicella–Zoster Virus (VZV)	Herpesvirus	Chronic latency	Global	$CD4^+$, $CD8^+$ (TRM)	Yes	Dorsal root ganglia, Skin	Zoster with aging or T-cell suppression; dissemination on high-dose steroids	25–30 Mya

Table 4 Specialist organisms and their immunological and evolutionary characteristics. These pathogens show focused tropism and epitope-centric strategies with moderate coordination, often maintaining chronic infection without broad systems-level imprinting. Abbreviations in parentheses match those used in Figures and trait score tables.

Organism (Abbr.)	Type	Infection Pattern	Geo. Dist.	Key Immune Deps.	Lifelong?	Tissues/Cell Involved	Immunosuppression Impact	Co-evo. Time
BK Virus (BKV)	Polyomavirus	Chronic latency	Global	CD8$^+$, Abs	Yes	Renal tubular epithelium	BK nephropathy post-transplant; risk with T-cell dysfunction	>400 kya
Candida albicans (C.alb)	Fungus	Colonization/ overgrowth	Global	Th17, neutrophils, Macrophages	Yes	Oral/genital mucosa, Skin	Mucocutaneous disease with IL-17 blockade or steroids	>20 Mya
Helicobacter pylori (H.pyl)	Bacteria	Chronic colonization	Global	Th1/Th17; B cells	Yes	Gastric epithelium	Persistent colonization; activity shaped by T-cell/IL-17 pathways	60–100 kya
Human Papillomavirus (HPV)	Papillomavirus	Chronic latency	Global	CD8$^+$, NK	Yes	Basal epithelial cells (anogenital/oropharyngeal)	High-grade lesions with T-cell defects; transplant risk	~500 kya
JC Virus (JCV)	Polyomavirus	Chronic latency	Global	CD4$^+$, IFN-γ	Yes	Oligodendrocytes, Astrocytes	PML with CD4$^+$ loss or integrin blockade	400–500 kya
Kaposi's Sarcoma-Assoc. Herpesvirus (KSHV)	Herpesvirus	Chronic latency	Focal	CD8$^+$ NK; CD4$^+$ help	Yes	Endothelium, B cells	KS/viremia in HIV/-transplant; T-cell suppression	20-30 Mya
Merkel Cell Polyomavirus (MCPyV)	Polyomavirus	Chronic latency	Global	CD8$^+$, CD4$^+$, IFN-γ	Yes	Merkel cells, Skin	Merkel cell carcinoma risk with T-cell suppression	>500 kya
Neisseria meningitidis (N.men)	Bacteria	Colonization + invasion	Global	Complement (C5–9), B cells	No	Nasopharynx, Meninges	Invasive disease with complement deficiency/asplenia	500–1000 kya
Parvovirus B19 (PVB19)	Virus	Seasonal infection	Global	Neutralizing Abs; CD8$^+$	No	Erythroid progenitors (marrow)	Chronic anemia in immunodeficiency/-transplant	10–20 kya
Pneumocystis jirovecii (P.jir)	Fungus	Chronic colonization	Global	CD4$^+$, Macrophages	No	Alveoli (type I/II interface)	PCP with CD4$^+$ loss (AIDS, steroids)	100–200 kya
Treponema pallidum (T.pal)	Bacteria	Mucosal contact	Global	CD4$^+$, Macrophages	No	Skin/mucosa, Endothelium, CNS	Neurosyphilis/relapses with T-cell suppression	2.5–15 Mya

Table 5 Multi-hostorganisms with broad tissue engagement but lacking deep co-evolution, human specificity, or stable imprinting. They can trigger strong immune activation or mimicry yet remain evolutionarily unsettled. Abbreviations in parentheses match those used in Figures and trait score tables.

Organism (Abbr.)	Type	Infection Pattern	Geo. Dist.	Key Immune Deps.	Lifelong?	Tissues/Cell Involved	Immunosuppression Impact	Co-evo. Time
Aspergillus spp. (Asper)	Fungus	Airborne inhalation	Ubiquitous	Neutrophils, Macrophages, $CD4^+$	No	Lungs, Paranasal sinuses; CNS (disseminated)	Invasive disease with neutropenia or high-dose steroids	<10 kya
Cryptococcus neoformans (C.neo)	Fungus	Environmental exposure	Global	$CD4^+$, Macrophages	Yes	Lungs, CNS (meningitis)	Severe disease in $CD4^+$ loss; transplant risk	Millions (environmental)
Histoplasma capsulatum (H.cap)	Fungus	Inhalation	Endemic (Americas)	Macrophages, $CD4^+$, IFN-γ	Yes	Lungs, RES (liver, spleen)	Dissemination with $CD4^+$/TNF blockade	<10 kya
Leishmania spp. (Leish)	Protozoan	Vector-borne	Trop/Subtrop	Macrophages, IFN-γ, TNF	Yes	Skin, Liver/Spleen (RES)	Reactivation with TNF/IFN-γ suppression	>15 Mya (clade)
Mycobacterium avium complex (MAC)	Bacteria	Opportunistic	Global	Macrophages, $CD4^+$	No	Lungs, RES	Dissemination with $CD4^+$ loss (AIDS)	<10 kya
Plasmodium falciparum (P.fal)	Protozoan	Chronic infection	Trop/Subtrop	$CD4^+$, IFN-γ, Abs	No	Hepatocytes, RBCs, Microvasculature	Severe disease with impaired $CD4^+$ responses; pregnancy risk	50–100 kya
Schistosoma spp. (Schis)	Helminth	Chronic infection	Trop/Subtrop	Th2, Eosinophils, Macrophages	Yes	Mesenteric/vesical veins, Liver, GI	Granulomas; immune remodeling; steroid effects	1–4 Mya
Strongyloides stercoralis (S.ste)	Helminth	Auto-infection	Trop/Subtrop	Th2, Eosinophils	Yes	GI tract, Lungs	Hyperinfection on steroids; HTLV-1 co-risk	20–50 kya
Toxoplasma gondii (T.gon)	Protozoan	Food/zoonotic	Global	$CD8^+$, $CD4^+$, IFN-γ	Yes	CNS, Retina, Muscle	Encephalitis/retinitis with $CD4^+$ loss	~10 kya (domestication)
Trypanosoma cruzi (T.cru)	Protozoan	Vector-borne	Latin America	$CD8^+$, $CD4^+$, IFN-γ	Yes	Cardiac muscle, GI, CNS	Reactivation in AIDS/transplant	~9 kya

Table 6 Opportunistic organisms characterized by low coordination, environmental ubiquity, and narrow immune dependencies. Their emergence under immunosuppression flags pathway-specific failure rather than systemic collapse. Abbreviations in parentheses match those used in Figures and trait score tables.

Organism (Abbr.)	Type	Infection Pattern	Geo. Dist.	Key Immune Deps.	Lifelong?	Tissues/Cell Involved	Immunosuppression Impact	Co-evo. Time
Acanthamoeba spp. (Acan)	Amoeba	Contact/Inhalation	Global (soil, water)	CD4$^+$, Macrophages	No	Cornea, CNS	Keratitis/CNS disease with T-cell defects	Unknown
Balamuthia mandrillaris (B.man)	Amoeba	Contact/Inhalation	Soil, water	CD4$^+$	No	CNS, Skin	Granulomatous amebic encephalitis in T-cell suppression	Unknown
Bartonella spp. (Barto)	Bacteria	Vector-borne	Ubiquitous (cats, lice)	CD4$^+$, Macrophages	Yes	Endothelium, Erythrocytes	Bacillary angiomatosis with T-cell suppression	~10 kya
Brucella spp. (Bruce)	Bacteria	Zoonotic	Livestock-associated	CD4$^+$, IFN-γ, Macrophages	Yes	RES, Reproductive tract	Chronic infection with TNF/IFN-γ axis defects	<10 kya
Burkholderia cepacia complex (Bcc)	Bacteria	Contact/Inhalation	Water, hospitals	Neutrophils; oxidative burst	No	Lungs, Bloodstream	Severe in CGD/CF; neutropenia risk	<10 kya
Coccidioides spp. (Cocci)	Fungus	Inhalation	Endemic (US SW)	CD4$^+$, Macrophages	No	Lungs, CNS	Dissemination with CD4$^+$ loss	<10 kya
Coxiella burnetii (C.bur)	Bacteria	Inhalation	Global (dust/aerosol)	IFN-γ, TNF, Macrophages	No	Lungs, Liver, Endothelium	Severe disease with TNF/IFN-γ blockade	<10 kya
Fusarium spp. (Fusar)	Fungus	Inhalation/Direct	Global (vegetation)	Neutrophils, Macrophages	No	Skin, Lungs, Blood	Dissemination with neutropenia/steroids	<10 kya
Human Herpesvirus 7 (HHV-7)	Herpesvirus	Chronic latency	Global	CD4$^+$, NK	Yes	T cells (CD4$^+$)	Reactivation with T-cell suppression	25–30 Mya
Legionella pneumophila (L.pne)	Bacteria	Inhalation	Water systems	IFN-γ, Macrophages	No	Lungs, Macrophages	Severe pneumonia with corticosteroids or TNF blockade	<100 years
Leptospira spp. (Lepto)	Bacteria	Zoonotic contact	Trop/Subtrop	Neutrophils, Macrophages, Abs	No	Kidney, Liver, Endothelium	Severe leptospirosis with innate/B-cell defects	5–10 kya
Nocardia spp. (Nocar)	Bacteria	Inhalation	Global (soil)	CD4$^+$, Macrophages	No	Lungs, Brain, Skin	Dissemination with T-cell suppression	<10 kya
Scedosporium spp. (Scedo)	Fungus	Inhalation	Global (stagnant water)	Neutrophils, Macrophages	No	Lungs, CNS	Severe disease with neutropenia; transplant risk	<10 kya
Torque Teno Virus (TTV)	Virus	Ubiquitous viremia	Near-universal	Tracks T-cell status (biomarker)	No	Blood, Lymphocytes	Higher loads with T-cell suppression; biomarker of net state	>500 kya
Vibrio spp. (Vibrio)	Bacteria	Ingestion/Wound	Coastal/Estuarine	Neutrophils, Complement	No	GI tract, Skin; Liver disease risk	Severe disease with liver disease or complement defects	<10 kya

Table 7 Trait definitions and score interpretations used to assess keystoneness.

Trait	Allowed Scores	Score Interpretation
Human-specific	{0,1}	1: Sustained human-only transmission. 0: Multi-host or zoonotic cycle.
Latency and Lifelong Infection	{0,1,2,3}	3: Lifelong latency with immune calibration. 2: Common persistence. 1: Limited or sporadic persistence. 0: None.
Length of Co-evolution with Humans	{0,1,2,3}	3: Deep-time co-divergence. 2: Long-standing association. 1: Recent emergence. 0: Recent zoonosis.
Global Prevalence	{0,1,2,3}	3: High global burden. 2: Intermediate. 1: Low or regional. 0: Rare.
Genetic Polymorphisms Reflecting Human Migration	{0,1,2,3}	3: Pathogen lineages tightly track human migration. 2: Partial coupling. 1: Contextual. 0: None.
Prevalence in Modern Indigenous Populations	{0,1,2,3}	3: High prevalence in multiple isolated populations. 2: Replicated across groups. 1: Limited. 0: None.
Consequences of Delayed Acquisition	{0,1,2,3}	3: Delayed infection causes severe disease. 2: Common age-related effects. 1: Context-specific. 0: None.
Association with Transplant Rejection	{0,1,2,3}	3: Robust, replicated clinical link. 2: Plausible but mixed. 1: Limited. 0: None.
Cross-reactivity with Adaptable RNA Viruses	{0,1}	1: Documented heterologous immunity or cross-reactivity. 0: None.
Perturbation by RNA Viruses	{0,1,2,3}	3: Frequent reactivation/modulation by RNA viruses. 2: Repeated in defined contexts. 1: Sporadic reports. 0: None.
Hypersensitivity and Autoimmune Flares	{0,1,2,3}	3: Reproducible links to autoimmune flares or immune complex disease. 2: Multiple plausible reports. 1: Sparse. 0: None.
Structured Immune Receptor Selection	{0,1,2,3}	3: Public or stereotyped TCR/BCRs with function. 2: Convergent motifs with partial function. 1: Scattered hints. 0: None.
Proximity and Persistence of Immune Cells	{0,1,2,3}	3: TRM or TLS consistently present with systemic effects. 2: Frequent persistence with limited scope. 1: Sparse. 0: None.
Epitope Structure and Immunodominance	{0,1,2,3}	3: Reproducible dominant epitopes across cohorts. 2: Replicated patterns. 1: Limited. 0: None.
Absence of Peripheral Tolerance	{0,1,2,3}	3: Established mimicry with autoimmune outcomes. 2: Strong evidence of tolerance break. 1: Suggestive hints. 0: None.
Epitope Presentation Dynamics	{0,1,2,3}	3: Strong HLA associations or MHC modulation. 2: Replicated effects. 1: Limited. 0: None.
Multi-Arm Coordination	{0,1}	1: Coordinated innate–adaptive containment with redundancy. 0: Partial or local coordination only.
Systemic Tropism	{0,1}	1: Multi-niche or systemic tissue persistence. 0: Localized or stochastic tropism.

B Trait-Based Scoring: Technical Details

B.1 Score vector representation

Each pathogen i was represented by an 18-dimensional score vector $\mathbf{x}_i = (x_{i,1}, \ldots, x_{i,18})$, where $x_{i,t}$ denotes the score for pathogen i on trait t. Total keystoneness scores were computed as unweighted sums:

$$S_i = \sum_{t=1}^{18} x_{i,t}$$

B.2 Gower distance

Pairwise dissimilarities were computed using Gower distance, which accommodates mixed ordinal and binary data while handling missing values:

$$d_{ij} = \frac{\sum_{t=1}^{p} s_{ijt}\, \delta_{ijt}}{\sum_{t=1}^{p} s_{ijt}}$$

where $s_{ijt} \in \{0,1\}$ indicates whether trait t is observed for both pathogens i and j (non-missingness indicator), and δ_{ijt} is the trait-specific dissimilarity:

$$\delta_{ijt} = \begin{cases} |x_{i,t} - x_{j,t}|/R_t, & \text{if trait } t \text{ is ordinal} \\ \mathbb{1}\{x_{i,t} \neq x_{j,t}\}, & \text{if trait } t \text{ is binary} \end{cases}$$

Here R_t is the observed range for ordinal trait t, and $\mathbb{1}\{\cdot\}$ is the indicator function.

B.3 Clustering algorithms

Two clustering approaches were applied to the Gower distance matrix:

Partitioning around medoids (PAM): PAM minimizes the sum of dissimilarities between each object and its assigned medoid:

$$\min_{m_1,\ldots,m_k} \sum_{i=1}^{n} \min_{j\in\{1,\ldots,k\}} d_{i,m_j}$$

where $m_1, \ldots, m_k$ are the indices of the k medoid organisms.

Hierarchical clustering: Agglomerative clustering with complete linkage, where the distance between clusters is defined as the maximum pairwise distance:

$$D(A, B) = \max_{i\in A, j\in B} d_{ij}$$

B.4 Cluster number selection

The number of clusters was selected using elbow and angle heuristics applied to within-cluster sum of dissimilarities $W(k)$ for $k = 2, \ldots, 8$:

$$W(k) = \sum_{c=1}^{k} \sum_{i \in C_c} d_{i,m_c}$$

Both methods nominated $k = 4$ clusters.

B.5 Consensus clustering

Method agreement was quantified by the Adjusted Rand Index (ARI):

$$\mathrm{ARI} = \frac{\mathrm{RI} - \mathbb{E}[\mathrm{RI}]}{\max(\mathrm{RI}) - \mathbb{E}[\mathrm{RI}]}$$

where RI is the Rand Index measuring pairwise agreement between clusterings.

For organisms with discordant assignments between PAM and hierarchical clustering, we applied a conservative consensus rule: each organism was assigned to the candidate cluster with the lower mean total score $\bar{S}_c$. Four organisms were reassigned under this rule, including Parvovirus B19 (Keystone $\rightarrow$ Specialist) and three additional organisms (Specialist $\rightarrow$ Opportunist). To further enhance the specificity of the keystone label, organisms initially assigned to the keystone archetype underwent an additional centroid-based validation step and were reassigned if their total trait score was closer to the centroid of another cluster.

B.6 Trait signature analysis

Cluster-versus-rest comparisons used Cliff's delta (δ) for effect size, defined as:

$$\delta = \frac{\#(x_i > y_j) - \#(x_i < y_j)}{n_1 n_2}$$

where x_i are scores in the focal cluster and y_j are scores in remaining clusters. Wilcoxon rank-sum tests were used for significance testing. Statistical significance was assessed using uncorrected p-values at $\alpha = 0.05$ (*** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$).

Multivariate signatures were obtained from ℓ_1-penalized (LASSO) multinomial logistic regression:

$$\hat{\boldsymbol{\beta}} = \arg\min_{\boldsymbol{\beta}} \left\{ -\ell(\boldsymbol{\beta}) + \lambda \sum_{c,t} |\beta_{ct}| \right\}$$

where $\ell(\boldsymbol{\beta})$ is the multinomial log-likelihood, the ℓ_1 penalty encourages sparsity in trait selection, and λ was selected by 10-fold cross-validation. The regularization parameter $\alpha = 1$ specifies pure LASSO penalization.

C Clinical Emergence Tensor: Scoring and Analysis

We constructed a three-dimensional clinical emergence tensor $\mathbf{T} \in \mathbb{R}^{43 \times 13 \times 31}$ documenting pathogen reactivation or invasion patterns across:

- 43 pathogens (matching those in the immunological trait analysis)
- 13 anatomical niches (blood/systemic, CNS, respiratory mucosa, GI mucosa, genitourinary mucosa, skin, reticuloendothelial system, bone marrow, lymph nodes, cardiac/endovascular, bone/joint, ocular, lymph node germinal centers)
- 31 immune perturbations spanning genetic immunodeficiencies and pharmacologic immunosuppression

C.1 Scoring Rubric

C.1.1 Tensor construction and scoring

Each tensor element $T(p, n, i) \in [0, 100]$ represents the strength of clinical evidence for lytic emergence, scored according to:

- **85–100**: Replicated human studies (genetic or interventional), hallmark associations
- **70–84**: Solid human signal for this niche with multiple sources or one high-quality source
- **40–69**: Clear but context-dependent human evidence, dose/timing dependent, partial compensation by other immune arms
- **10–39**: Suggestive or narrow human signals, or strong mechanistic logic with limited human data
- **5**: No authoritative human evidence found after diligent search
- **0**: Published evidence indicates protective effect or no increased emergence

The threshold at 40 demarcates human-relevant clinical data (scores ≥ 40 include at least limited human observations with mechanistic support, grade C or better) from preclinical evidence only (scores < 40, grade D).

C.1.2 Evidence grading and confidence annotation

Each score was accompanied by structured metadata:

- **Direction**: "revealed" (perturbation increases emergence), "protected" (reduces risk), or "unclear" (conflicting/insufficient data)
- **EvidenceGrade**:
  - A = replicated human genetic or interventional evidence
  - B = strong human clinical cohorts or case series
  - C = limited human data with strong mechanistic support
  - D = animal or in vitro evidence only

- **ConfidenceTag**: [High], [Moderate], [Low], [Minimal] (for score 5), [None] (for score 0)
- **Justification**: rigorous citations formatted as (Author, Title, Journal, Year)
- **EvidenceCount**: number of distinct human studies supporting the score

- **Missingness**: "observed" (evidence exists), "inferred_zero" (no evidence after diligent search), or "unknown"

Evidence was prioritized as: (1) human genetic/interventional perturbations with niche-localized outcomes, (2) human clinical cohorts with clear site localization, (3) human observations without precise site (stated with caveats), (4) animal/in vitro support only (capped at score 69, grade C or D).

C.1.3 Controlled vocabularies for niches and perturbations

Each anatomical niche was defined by controlled vocabularies specifying required anatomical terms (IncludeTokens) and exclusion criteria (ExcludeTokens) to ensure precise localization. For example, CNS required terms like "brain," "cerebral," "encephalitis," "CSF," or "meninges," while excluding "ocular" or "peripheral nerve." Similarly, each immune perturbation was defined by mechanism and synonyms. For instance, P07 (CalcineurInh) encompasses "tacrolimus," "cyclosporine," and "calcineurin inhibitor," all blocking NFAT and IL-2 signaling. Complete niche and perturbation definitions are in Supplementary Section C.2.

C.1.4 Sensitivity analyses: Score adjustment schemes

To assess robustness to evidence quality, we performed sensitivity analyses by adjusting scores based on evidence grade and confidence:

Grade-capping.

Scores were reduced for lower evidence grades to prevent overweighting of animal/in vitro data:

- Grade A: score unchanged
- Grade B: score capped at 84 (if originally > 84)
- Grade C: score capped at 69 (if originally > 69)
- Grade D: score capped at 39 (if originally > 39)

Grade+Confidence weighting.

Scores were multiplicatively adjusted by a factor combining evidence grade and confidence tag, down-weighting low-confidence or low-grade entries. Weighting factors were:

- High: 1.0 (no reduction)
- Moderate: 0.85
- Low: 0.70
- Minimal/None: 0.50

Combined with grade-specific caps, this yielded conservative score estimates prioritizing high-quality human evidence.

All feature extraction (max-marginalization, breadth metrics), classification (elastic-net regression), and clustering (PAM) analyses were re-run on both adjusted

score tensors to test whether archetype predictions and unsupervised recovery remain robust when evidence quality is explicitly down-weighted.

C.2 Tensor Dimensions

The clinical tensor $\mathbf{T} \in \mathbb{R}^{43 \times 13 \times 31}$ captures pathogen behavior across anatomical niches and immune perturbations. Below we provide detailed definitions for each dimension.

C.2.1 Anatomical Niches (N=13)

1. **CNS (Central Nervous System)**: Brain parenchyma, meninges, cerebrospinal fluid spaces. Reflects capacity for neuroinvasion, meningitis, or encephalitis. Relevant for neurovirulent pathogens and those causing CNS opportunistic infections.
2. **Lung (Pulmonary)**: Pulmonary parenchyma, airways, alveolar spaces. Encompasses pneumonia, bronchitis, and respiratory tract colonization. Central niche for respiratory viruses, bacteria, and fungi.
3. **GI (Gastrointestinal)**: Gastrointestinal mucosa, lumen, and associated lymphoid tissue (GALT). Includes enteric infections, gastroenteritis, and mucosal colonization. Relevant for fecal-oral transmission routes.
4. **Liver (Hepatic)**: Hepatic parenchyma, including hepatocytes, Kupffer cells, and bile ducts. Reflects hepatotropism, hepatitis, and intrahepatic replication. Important for bloodborne and vector-borne pathogens.
5. **Blood (Bloodstream/Intravascular)**: Intravascular space, including circulating immune cells and endothelium. Captures bacteremia, viremia, parasitemia, and disseminated infections. Critical for sepsis-causing organisms.
6. **LN_Paracortex (Lymph Node T Cell Zones)**: T cell-rich paracortical regions of lymph nodes. Site of adaptive T cell priming and cell-mediated immune responses. Reflects pathogens that manipulate or reside within T cell niches.
7. **LN_GC (Lymph Node Germinal Centers)**: B cell germinal centers within secondary lymphoid tissues. Sites of B cell maturation, somatic hypermutation, and antibody class-switching. Reflects pathogens that colonize or replicate within organized B cell zones.
8. **Spleen (Splenic)**: Splenic parenchyma, including red pulp (filtration) and white pulp (lymphoid). Central to bloodborne pathogen clearance and immune surveillance. Critical for encapsulated bacteria and blood-stage parasites.
9. **BoneMarrow (Hematopoietic Marrow)**: Bone marrow niches supporting hematopoiesis. Reflects pathogens causing marrow suppression, hemophagocytic syndromes, or intracellular marrow persistence.
10. **Skin (Cutaneous/Subcutaneous)**: Skin, subcutaneous tissue, and dermal immune cells. Includes cellulitis, abscesses, vector bite sites, and dermatological manifestations. Entry point for many vector-borne and traumatic infections.
11. **GU (Genitourinary)**: Genitourinary mucosa, renal parenchyma, bladder. Encompasses urinary tract infections, pyelonephritis, and sexually transmitted infections.

12. **Bone (Osseous)**: Bone tissue and periosteum. Reflects osteomyelitis, septic arthritis, and bone-invasive infections. Relevant for hematogenous seeding and trauma-associated infections.
13. **Eye (Ocular)**: Ocular tissues including conjunctiva, cornea, uvea, and retina. Captures keratitis, uveitis, endophthalmitis, and retinitis. Important for certain congenital and disseminated infections.

C.2.2 Immune Perturbations (N=31)

1. **P01 (Neutropenia)**: Severe neutrophil depletion (<500 cells/μL). Impairs bacterial and fungal killing, increasing risk for invasive bacterial and mold infections. Seen in chemotherapy, aplastic anemia, and congenital neutropenias.
2. **P02 (MacrophageDefect)**: Macrophage dysfunction or depletion. Compromises intracellular pathogen control, granuloma formation, and tissue clearance. Relevant for mycobacteria, Listeria, Salmonella, and intracellular parasites.
3. **P03 (CD4depletion)**: CD4+ T cell deficiency (e.g., HIV/AIDS, idiopathic CD4 lymphopenia). Central immunodeficiency affecting cell-mediated and humoral immunity. Predisposes to opportunistic infections and impaired vaccine responses.
4. **P04 (CD8depletion)**: CD8+ T cell deficiency or dysfunction. Impairs cytotoxic responses against intracellular pathogens and virally infected cells. Relevant for chronic viral infections and some intracellular bacteria.
5. **P05 (NKdepletion)**: Natural killer cell deficiency or dysfunction. Impairs early antiviral and antitumor responses. Associated with increased susceptibility to herpesviruses and some bacterial infections.
6. **P06 (TNFblock)**: TNF-α blockade via monoclonal antibodies (infliximab, adalimumab, etanercept). Impairs granuloma formation and macrophage activation. Major risk factor for tuberculosis reactivation and invasive fungal infections.
7. **P07 (IFNgBlock)**: IFN-γ neutralization or receptor deficiency. Critical for macrophage activation and intracellular pathogen control. Deficiency predisposes to disseminated mycobacterial and Salmonella infections.
8. **P08 (IL12p40block)**: IL-12/IL-23 p40 subunit blockade (ustekinumab). Impairs Th1 and Th17 responses. Increases risk for mycobacterial, fungal, and some bacterial infections.
9. **P09 (IL17block)**: IL-17A blockade (secukinumab, ixekizumab) or receptor deficiency. Impairs mucosal immunity and neutrophil recruitment. Predisposes to mucocutaneous candidiasis and some bacterial infections.
10. **P10 (IL1block)**: IL-1 blockade (anakinra, canakinumab) or receptor deficiency. Reduces pyogenic responses and fever. Modest infection risk, primarily pyogenic bacteria.
11. **P11 (C3deficiency)**: Complement C3 deficiency or depletion. Central complement component; deficiency severely impairs opsonization and immune complex clearance. High risk for encapsulated bacteria and Neisseria.
12. **P12 (ProperdinDeficiency)**: Properdin deficiency, stabilizer of alternative pathway C3 convertase. X-linked deficiency with markedly increased susceptibility to meningococcal disease.

13. **P13 (MBLdeficiency)**: Mannose-binding lectin deficiency. Impairs lectin pathway activation and opsonization. Modest increased risk for bacterial infections, particularly in young children.
14. **P14 (Hypogammaglobulinemia)**: Severe antibody deficiency (CVID, X-linked agammaglobulinemia). Impairs opsonization, neutralization, and immune complex clearance. High risk for encapsulated bacteria, enteroviruses, and chronic sinopulmonary infections.
15. **P15 (Bcelldepletion)**: B cell depletion via anti-CD20 therapy (rituximab, ocrelizumab) or congenital deficiency. Impairs antibody production and antigen presentation. Increases risk for bacterial infections and viral reactivation.
16. **P16 (TLRdefect)**: Toll-like receptor pathway defects (TLR signaling impairment). Reduces innate recognition of PAMPs. Variable susceptibility depending on specific TLR; broadly impairs innate immune priming.
17. **P17 (MyD88deficiency)**: MyD88 adaptor protein deficiency. Impairs most TLR and IL-1R signaling. Severe susceptibility to pyogenic bacteria, particularly Streptococcus pneumoniae and Staphylococcus aureus.
18. **P18 (IRAK4deficiency)**: IRAK4 kinase deficiency, downstream of MyD88. Similar phenotype to MyD88 deficiency with invasive bacterial infections in childhood.
19. **P19 (NFkBdefect)**: NF-κB pathway defects (NEMO, IκBα mutations). Impairs broad inflammatory and immune gene transcription. Variable phenotype with mycobacterial, bacterial, and fungal susceptibility.
20. **P20 (CGDdefect)**: Chronic granulomatous disease (phagocyte NADPH oxidase deficiency). Eliminates oxidative burst, impairing killing of catalase-positive organisms. High risk for Staphylococcus, Aspergillus, Burkholderia, Serratia, and Nocardia.
21. **P21 (MHC1deficiency)**: MHC class I deficiency (TAP deficiency, bare lymphocyte syndrome type I). Impairs CD8+ T cell antigen presentation. Increases susceptibility to viral infections.
22. **P22 (MHC2deficiency)**: MHC class II deficiency (bare lymphocyte syndrome type II). Impairs CD4+ T cell priming. Severe combined-like immunodeficiency with broad bacterial, viral, and fungal susceptibility.
23. **P23 (CD40Ldeficiency)**: CD40 ligand deficiency (X-linked hyper-IgM syndrome). Impairs T cell help for B cells and macrophage activation. Predisposes to Pneumocystis, Cryptosporidium, and bacterial infections.
24. **P24 (CTLA4agonist)**: CTLA-4 agonism or Ig fusion (abatacept). Inhibits T cell costimulation. Modest infection risk, primarily opportunistic and mycobacterial.
25. **P25 (PD1block)**: PD-1/PD-L1 blockade (checkpoint inhibitors: nivolumab, pembrolizumab). Enhances T cell activation; paradoxically can increase some infection risks via immune dysregulation and autoimmunity.
26. **P26 (IL6block)**: IL-6 or IL-6R blockade (tocilizumab, sarilumab). Reduces acute phase responses and Th17 differentiation. Increases risk for bacterial infections, diverticulitis, and some opportunistic pathogens.

27. **P27 (IL10block)**: IL-10 blockade or deficiency. IL-10 is anti-inflammatory; blockade may paradoxically enhance immunity but also cause inflammatory bowel disease. Included for completeness; rare clinical use.
28. **P28 (IL23block)**: IL-23 blockade (p19 subunit: guselkumab, risankizumab). More specific than IL-12/23 blockade; impairs Th17 responses. Infection risk appears lower than IL-12/23 blockade but includes fungal and mycobacterial concerns.
29. **P29 (JAKblock)**: JAK kinase inhibition (tofacitinib, baricitinib, upadacitinib). Broad impairment of cytokine signaling (IFN, IL-6, IL-12, IL-23, etc.). Increases risk for herpes zoster, tuberculosis, and opportunistic infections.
30. **P30 (CSFblock)**: GM-CSF or G-CSF blockade/deficiency. Impairs myeloid development, neutrophil function, and alveolar macrophage maturation. GM-CSF neutralization associated with pulmonary alveolar proteinosis and opportunistic lung infections.
31. **P31 (IL5block)**: IL-5 blockade via monoclonal antibodies (mepolizumab, reslizumab, benralizumab). IL-5 drives eosinophil development, survival, and activation. Blockade severely depletes eosinophils, impairing helminth immunity and potentially tissue eosinophil-mediated responses. Used clinically for severe eosinophilic asthma and hypereosinophilic syndrome.

C.3 Feature extraction from tensor and analysis

From the 3D tensor, we derived 2D feature matrices for each pathogen via max-marginalization:

$$X_{PI}(p,i) = \max_{n \in \text{niches}} T(p,n,i)$$

$$X_{PN}(p,n) = \max_{i \in \text{perturbations}} T(p,n,i)$$

These matrices capture the maximal emergence score across all contexts, yielding a 43×44 feature matrix per pathogen (31 immune perturbations + 13 niches).

We augmented this with breadth metrics:

$$\text{Immune breadth}(p) = \sum_{i=1}^{31} \mathbb{I}[X_{PI}(p,i) \geq 40]$$

$$\text{Niche breadth}(p) = \sum_{n=1}^{13} \mathbb{I}[X_{PN}(p,n) \geq 40]$$

quantifying the number of immune perturbations and anatomical niches exhibiting clinically significant emergence (threshold = 40, reflecting human-relevant data).

C.3.1 Supervised classification

We trained elastic-net logistic regression models to predict archetype membership from clinical features:

$$\hat{\boldsymbol{\beta}} = \arg\min_{\boldsymbol{\beta}} \left\{ -\ell(\boldsymbol{\beta}) + \lambda \left[\frac{1}{2}(1-\alpha)\|\boldsymbol{\beta}\|_2^2 + \alpha\|\boldsymbol{\beta}\|_1 \right] \right\}$$

where $\ell(\boldsymbol{\beta})$ is the log-likelihood (binomial for binary, multinomial for 4-way classification), $\alpha = 0.5$ balances ℓ_1 and ℓ_2 penalties, and λ was selected by cross-validation. We used 8-fold cross-validation for binary classification (keystone vs. non-keystone) to ensure adequate representation in each fold, and 5-fold for multi-class prediction to accommodate smaller archetype sample sizes. Models were fit using the `glmnet` package with AUC (binary) or misclassification error (multi-class) as the optimization criterion.

C.3.2 Unsupervised clustering validation

To test whether clinical emergence patterns independently recover trait-based archetypes, we performed PAM clustering on Gower dissimilarities of the feature matrix with $k = 4$ clusters. Cluster assignments were aligned to archetype labels via permutation optimization to maximize diagonal agreement, and concordance was quantified by Adjusted Rand Index (ARI). This unsupervised approach provides external validation that archetypes derived from immunological traits (Figure 4) correspond to distinct clinical emergence signatures.

All analyses (feature extraction, classification, clustering) were repeated for raw scores, grade-capped scores, and grade+confidence weighted scores to assess robustness across evidence quality thresholds.

D Mechanistic Model: Technical Details

D.1 Mechanistic scoring tables

Five expert-curated scoring tables quantified biological interactions (all scores normalized to $[0, 1]$):

1. **Latent control** $L(p, n_{\text{lat}}, c) \in [0, 1]$: Strength with which immune controller c suppresses pathogen p in latent niche n_{lat}. Higher scores indicate tighter immunological containment. (154 scored interactions)
2. **Lytic attack** $A(p, n) \in [0, 1]$: Capacity of pathogen p to breach barriers and cause clinical disease in niche n, independent of immune status. (145 scored interactions)
3. **Reactivation routes** $R(p, n_{\text{lat}} \rightarrow n_{\text{lytic}}) \in [0, 1]$: Probability that pathogen p escapes from latent reservoir n_{lat} to manifest clinically in lytic niche n_{lytic}. (151 scored interactions)
4. **Perturbation effects** $E(i, c) \in [0, 1]$: Degree to which immune perturbation i disables controller c. (144 scored interactions)
5. **Controller residency** $C(c, n) \in \{0, 1\}$: Binary indicator for whether controller c operates in niche n. (132 scored interactions)

D.1.1 Score normalization

Raw expert scores $s \in [0, 100]$ were normalized using:

$$s_{\text{norm}} = \begin{cases} \pi_{\text{unknown}} = 0.1 & \text{if } s = 5 \text{ (unknown)} \\ 0 & \text{if } s < 40 \text{ (biologically negligible)} \\ s/100 & \text{otherwise} \end{cases}$$

The threshold of 40 reflects expert consensus that lower scores indicate biologically negligible interactions. Crucially, unknown interactions (originally coded as 5) retain a small prior probability of 0.1 rather than being zeroed, reflecting that an unstudied interaction is not necessarily absent.

D.1.2 Perturbation effect scaling

Perturbation effects were scaled by qualitative effect type using multiplier κ:

- Depleted: $\kappa = 1.0$ (complete loss of controller)
- Disabled: $\kappa = 0.8$ (functional impairment with residual presence)
- Functionally impaired: $\kappa = 0.6$ (partial dysfunction)
- Unknown: $\kappa = 0.6$ (conservative default)

The final perturbation effect is $E_{\text{scaled}} = E_{\text{base}} \times \kappa$.

D.2 Latent pathway

The latent pathway models emergence through reactivation from immune-controlled reservoirs:

$$\text{Pathogen} \xrightarrow{L} \text{Latent Site} \xrightarrow{R} \text{Lytic Site} \xleftarrow{C} \text{Controller} \xleftarrow{E} \text{Perturbation}$$

For each mechanistic chain $(p, n_{\text{lat}}, n_{\text{lytic}}, c, i)$, the base score uses geometric mean (length-normalized):

$$s_{\text{base}} = \exp\left(\frac{1}{3} \sum_{x \in \{L, R, C\}} \log(x + \epsilon)\right)$$

where $\epsilon = 10^{-6}$ prevents $\log(0)$.

The perturbation effect acts as a modulatory exponent with strength $\beta = 1.5$:

$$s_{\text{chain}} = s_{\text{base}} \times E^{\beta}$$

This means complete controller disabling ($E = 1$) allows full expression of the base score, while partial effects attenuate emergence exponentially.

D.3 Barrier pathway

The barrier pathway models direct invasion when immune barriers are compromised:

$$\text{Pathogen} \xrightarrow{A} \text{Lytic Site} \xleftarrow{C} \text{Controller} \xleftarrow{E} \text{Perturbation}$$

To prevent over-triggering for controllers that don't actually control the pathogen, we added pathogen–controller specificity using latent control scores as a proxy:

$$PC(p, c) = \max_{n_{\text{lat}}} L(p, n_{\text{lat}}, c)$$

The barrier chain score is then:

$$s_{\text{barrier}} = A(p, n) \times E(i, c) \times C(c, n) \times PC(p, c)$$

D.4 Aggregation strategy

D.4.1 Collapse to best-per-controller

Multiple chains through the same controller are collapsed to avoid fake independence:

$$s_{\text{ctl}}(p, n, i, c) = \max_{\text{chains via } c} s_{\text{chain}}$$

D.4.2 Top-K selection

Only the top $K = 5$ controllers are retained before aggregation:

$$\mathcal{C}_{\mathrm{top}}(p, n, i) = \text{top-}K\left(\{s_{\mathrm{ctl}}(p, n, i, c)\}_c\right)$$

D.4.3 Normalized noisy-OR

Standard noisy-OR saturates quickly with multiple contributing causes. We use normalized noisy-OR:

$$\hat{T}_{\mathrm{pathway}} = 1 - \prod_{c \in \mathcal{C}_{\mathrm{top}}} (1 - s_{\mathrm{ctl}, c})^{1/K}$$

Property: If all K chains have score s, the result is exactly s (not ≈ 1).

D.4.4 Pathway combination

Latent and barrier pathways represent competing explanations, not additive probabilities:

$$\hat{T}_{\mathrm{mix}}(p, n, i) = \max\left(\hat{T}_{\mathrm{latent}}, \hat{T}_{\mathrm{barrier}}\right)$$

D.5 Evaluation framework

D.5.1 The join direction problem

A critical methodological consideration: evaluation must start from the full observed tensor, not from predictions. Joining predictions to observations (rather than vice versa) would hide false negative and true negative mass, artificially inflating apparent performance. All evaluations use observation-first joins.

D.5.2 Binning to discrete rubric

Both predictions and observations were projected to coarse bins (multiples of 5: 0, 5, 10, ..., 100) to match the discrete clinical scoring rubric:

$$\mathrm{project}(x) = 5 \times \mathrm{round}(x/5)$$

D.5.3 Three-way evaluation

Because zero clinical scores may represent either "true negative" or "not studied," we employed three complementary evaluation frameworks:

Evaluation 1: STRICT (all zeros = true negatives)

- Most conservative/pessimistic
- Treats every zero as evidence of absence
- Metrics: Pearson r, Spearman ρ, Cohen's weighted κ, F1, balanced accuracy

Evaluation 2: PU FRAMING (zeros = unlabeled)

- Positive-Unlabeled learning perspective
- Only positive observations are labeled; zeros are unknown
- Metrics: Precision@K, Recall@K, fold-enrichment

Fold-enrichment is defined as:

$$\text{Enrichment}@K = \frac{\text{Precision}@K}{\text{baseline rate}} = \frac{|\text{positives in top-}K|/K}{n_{\text{pos}}/N}$$

Evaluation 3: CONFIDENCE-MASKED (low-confidence zeros = missing)

- Uses evidence confidence to distinguish:
  - **Confident zeros**: Curated evidence of absence (retain as TN)
  - **Uncertain zeros**: Absence of evidence (mask out)
- Evaluate only on confident subset

Finding: All zeros in the clinical tensor have low confidence scores, meaning there is no curated evidence of absence. This strongly supports the PU framing interpretation.

D.5.4 Cohen's weighted kappa

For ordinal agreement between binned predictions and observations:

$$\kappa_w = \frac{p_o - p_e}{1 - p_e}$$

where p_o and p_e are weighted observed and expected agreement using linear weights:

$$w_{ij} = 1 - \frac{|i - j|}{n_{\text{bins}} - 1}$$

Interpretation scale: < 0 (less than chance), 0.0–0.2 (slight), 0.2–0.4 (fair), 0.4–0.6 (moderate), 0.6–0.8 (substantial), 0.8–1.0 (almost perfect).

D.6 Results summary

D.6.1 Dataset statistics

Metric	Value
Full tensor cells	17,329
Cells with mechanistic predictions	1,522 (8.8%)
Cells without predictions	15,807 (91.2%)
Known positives ($T_{\text{obs}} > 0$)	3,425 (19.8%)

D.6.2 Pathway dominance

Dominant Pathway	Count	Percentage
Barrier	890	58.5%
Latent	283	18.6%
Equal	349	22.9%

D.6.3 Overall performance (STRICT evaluation)

Metric	Value
Pearson r	0.349
Spearman ρ	0.299
Weighted κ	0.236 (Fair)
Recall	0.197
Specificity	0.974
Precision	0.421
F1	0.259
Balanced Accuracy	0.586

D.6.4 Per-archetype performance

Archetype	n	Pearson r	Spearman ρ	F1
Keystone	2,821	0.388	0.357	0.301
Opportunistic	6,045	0.355	0.316	0.248
Multi-host	4,030	0.333	0.305	0.219
Specialist	4,433	0.325	0.232	0.279

D.6.5 Enrichment analysis (PU framing)

Top K%	K	Positives Found	Precision@K	Enrichment
1%	173	51	0.29	2.9×
5%	866	256	0.30	2.9×
10%	1,733	508	0.29	2.9×
20%	3,466	870	0.25	2.5×

D.6.6 2D projection performance

Projection	Pearson r	Spearman ρ
PI (Pathogen × Immune)	0.359	0.374
PN (Pathogen × Niche)	0.583	0.555

D.7 S3.7 Interpretation of "predicted high, observed zero"

Cases where $\hat{T} > 0$ but $T_{\mathrm{obs}} = 0$ are not necessarily model errors. They may indicate:

1. **Not studied**: The pathogen–niche–perturbation combination was never investigated
2. **Not detected**: Low statistical power or inappropriate study cohort
3. **No opportunity**: Pathogen not present in studied population

4. **True model miss**: Missing ecological constraints (temperature, microbiome, co-infections)

The finding that all clinical zeros have low confidence strongly supports interpretations 1–3 over interpretation 4.