Network Modeling Predicts How DYRK1A Inhibition Promotes Cardiomyocyte Cycling after Ischemic/Reperfusion Injury

Bryce C. Murillo[1]*, Alexander Young[2]*, Kaitlyn L. Wintruba[3], Alexander J. Eichert[4], Klara Siejda[1], Dennon Hoenig[1], Leigh A. Bradley[1], Bryana N. Harris[3], Catherine Zhao[3], Michelle Wu[3], Emmanuel Deau[5], Matthias F. Lindberg[5], Laurent Meijer[5], Jeffrey J. Saucerman[3]†, and Matthew J. Wolf[1]†

[1]Department of Pharmacology University of Virginia, Charlottesville VA, USA, [2]Department of Medicine, University of Virginia, Charlottesville VA, USA, [3]Department of Biomedical Engineering University of Virginia, USA, [4]Department of Molecular Physiology and Biological Physics, and [5]Perha Pharmaceuticals, Roscoff, France

*Contributed equally

One Sentence Summary: Inhibition of DYRK1A through LCTB-92 induces cardiomyocyte cycling and improved heart function in a mouse model of ischemic/reperfusion injury.

Co-Corresponding Authors' contact

†Matthew J. Wolf, M.D., Ph.D.
415 Lane Road
MR5, Room G213
Charlottesville, VA 22008
mjw5mc@virginia.edu

†Jeffery J. Saucerman, Ph.D.
415 Lane Road
MR5, Room 2207
Charlottesville, VA 22008
jjs3g@virginia.edu

ABSTRACT

The adult mammalian heart has a limited ability to regenerate lost myocardium following myocardial infarction (MI), largely due to the poor proliferative capacity of cardiomyocytes. Dual-specificity tyrosine phosphorylation-regulated kinase 1A (DYRK1A) is a known regulator of cell quiescence, though the mechanisms underlying its function remain unclear. Previous studies have shown that pharmacological inhibition of DYRK1A using harmine induces cardiomyocyte cell cycle re-entry after ischemia/reperfusion (I/R) MI. Here, we developed a computational network model of DYRK1A-mediated regulation of the cell cycle, which predicts how DYRK1A inhibition promotes cardiomyocyte re-entry. To validate these predictions, we tested selective DYRK1A inhibitors and observed robust induction of cell cycle activity in neonatal rat cardiomyocytes (NRCMs). Integrating our network model with bulk RNA-sequencing data from DYRK1A inhibitor-treated NRCMs, we identified E2F1 as a key transcriptional driver of cell cycle gene expression. Finally, we demonstrate that both pharmacological and post-developmental inhibition of DYRK1A enhances heart function and increases cardiomyocyte cycling following I/R MI. Our findings suggest that functional recovery induced by small molecule inhibitor of DYRK1A is mediated by the induction of cycling cardiomyocytes.

INTRODUCTION

Myocardial infarctions (MIs) affect ~800,000 individuals in the United States annually and cause significant morbidity and mortality[1]. The ischemia induced by the cessation of coronary blood flow during an MI and subsequent reperfusion leads to cardiomyocyte necrosis, infiltration of inflammatory cells, and activation of fibroblasts to produce fibrosis[2]. Unfortunately, the adult mammalian heart has no significant regenerative capacity to restore cardiomyocyte loss after MI[3,4]. Therefore, the current limitations in treating MIs and the need for new therapeutics to stimulate cardiomyocyte proliferation, in a controlled manner, represent an opportunity to improve cardiac function after injury.

A number of studies demonstrate that enhancing cardiomyocyte proliferation after MI improves heart function[5–7]. The genetic ablation or pharmacologic inhibition of proteins in the YAP (Yes-associated protein) pathway promotes cardiomyocyte cycling. For example, the genetic ablation of the Salvador scaffold protein or MST1 and MST2, two serine/threonine kinases that regulate YAP, promote robust cardiomyocyte proliferation[8,9]. Alternatively, the overexpression of cyclin-dependent kinase 1 (CDK1), CDK4, cyclin B1, and cyclin D1 or CDK4 and cyclin D1 in combination with Wee1 and TGFβ inhibition promotes cardiomyocyte proliferation in vivo[10]. We previously identified that genetic ablation or inhibition of dual specificity tyrosine phosphorylation regulated kinase 1A (DYRK1A) induced cardiomyocyte cell cycle entry and improved heart function after ischemic reperfusion (I/R) MI[11]. DYRK1A is an upstream mediator of the Myb-MuvB/dimerization partner, RB-like, E2F, and multi-vulval class B (DREAM) complex, which maintains cell cycle quiescence through suppression of G1/S and G2/M checkpoints[12]. DYRK1A promotes DREAM complex assembly through phosphorylation of LIN52 on residue serine 28[12]. DYRK1A also phosphorylates cyclin D2, targeting it for proteasomal degradation in cardiomyocytes[13]. In contrast, the overexpression of DYRK1A decreased cardiomyocyte proliferation during development in a mouse model of Down Syndrome[14]. This provides evidence

that DYRK1A regulates cell cycle quiescence, and its inhibition can promote re-entry of

cardiomyocytes into the cell cycle. However, the mechanisms through which DYRK1A influences

cell cycling remain enigmatic.

Our prior experiments have utilized Harmine, an inhibitor of DYRK1A that also inhibits

Monoamine Oxidase Inhibitor (MAOI)[11]. While Harmine has proven effective in promoting

cardiomyocyte cycling, it is not without its drawbacks. The drug's adverse effects, including

nausea, vomiting, drowsiness, and impaired concentration, significantly limit its potential clinical

use[15]. Therefore, our current efforts are focused on developing more specific, orally bioavailable

DYRK1A inhibitors that can effectively promote cardiomyocyte cycling without these adverse

effects, thereby paving the way for their potential clinical application.

Here, we employed computational network modeling to predict mechanisms by which

DYRK1A regulates cardiomyocyte cell cycling. These predictions were validated in neonatal rat

cardiomyocytes (NRCMs) using selective DYRK1A inhibitors, followed by RNA-sequencing.

Based on these findings, we tested the DYRK1A inhibitor Leucettinib-92 (LCTB-92) in a mouse

model of ischemia-reperfusion (I/R) myocardial infarction (MI), where it improved heart function

and increased cardiomyocyte cycling.

RESULTS

Network Model Predicts Mechanisms of DYRK1A-mediated Cell Cycle Activity

While we have previously found that both genetic and pharmacologic inhibition of DYRK1A

induce cardiomyocyte cycling[11], the mechanisms by which they do so have not been explored.

This led us to develop a network model by manual curation of literature that describes DYRK1A's

regulatory role of cell cycling molecules (**Table S1**). DYRK1A's role in cell cycling is believed to

be highly conserved across different tissue types, so we broadened our literature review beyond

papers that focused on cardiomyocytes (CMs) to leverage mechanistic knowledge from other cell

types[12,16–21]. We then used these relationships to create a logic-based network model using

Netflux[22,23], where we could mechanistically predict how DYRK1A knockdown increases cell cycle

activity.

The model predicts that in quiescent cardiomyocytes (baseline), high DYRK1A expression

induces DREAM complex formation and RB-1 activity through the direct and indirect degradation

of cyclin D and E respectively (**Fig. 1A**). Upon simulated DYRK1A knockdown, the model predicts

an increase in cell cycle entry (DNA replication) through the increase of cyclins D and E, which

promotes DREAM complex breakdown and inactivation of RB-1 (**Fig. 1A**). We then validated our

model's accuracy by comparing predictions with experimental results found in literature not used

to build the model (**Fig. 1B**). Overall, the model's predictions were highly accurate across multiple

experiments   performed   in   cardiomyocyte   (100%)   and   cancer   (85%)   literature[11,13,24–40].

Interestingly, the model's predicted effect of DYRK1A KO on DNA replication was consistent with

cardiomyocyte   literature   but   not   consistent   with   cancer   literature[31,38]. DYRK1a   may   play   an

alternative role in the context of some cancers[41] which may explain this discrepancy in literature.

To identify which network nodes are most sensitive to DYRK1A, we simulated a DYRK1A

concentration response by simulating knockdown of DYRK1A in 1% increments. Node sensitivity

was   assessed   based   on   predicted   EC50/IC50   values   (**Fig. 1C**).   Strikingly,   nodes   further

downstream targets were more sensitive to DYRK1A knockdown, indicating that even partial

DYRK1A inhibition may be sufficient to induce cell cycle entry. Finally, we performed a knockdown

screen in the context of a 100% DYRK1A knockdown to identify modulators of its response (**Fig.**

1D). We found that knockdown of E2F1,2, and 3 transcription factors attenuated the predicted

increase in DNA replication upon a DYRK1A knockdown. Additionally, knockdown of cyclin D and

E complexes did not attenuate the effect of DYRK1A knockdown on DNA replication, indicating

that they may play redundant roles in modulating cell cycle entry. All in all, our robust analysis

supports that we have built a reliable model of DYRK1A signaling.

123
Figure 1: Network model predicts mechanisms of DYRK1A-mediated cardiomyocyte cell
cycle activity. (A) Model predicts response to active DYRK1A (left) and when DYRK1A is
knocked out (right). **(B)** Model simulation of CDK2 overexpression's effect on DNA replication
(left) compared to experimental data found in literature (right). **(C)** Other simulations were
compared to experiments in both cardiomyocyte (n = 5) and cancer literature (n = 13) (n = # of
primary literature articles). Validation accuracy is measured by the # of predictions that match
literature results/simulations performed. **(D)** Concentration response showing resulting node
activity from increasing levels of DYRK1A knockdown. IC50 and EC50 values indicate the %
DYRK1A knockdown at which a node increases or decreases by 50% (determined by the change
in normalized activity by 50%). **(E)** Knockdown screen showing network model's response to each
node being 100% knocked down. Nodes were knocked out one at a time by setting their y-max
to 0. Simulations were performed under a 100% DYRK1A knockdown.

DYRK1A inhibition promotes cell cycling in neonatal rat cardiomyocytes

To experimentally validate the network model prediction that DYRK1A inhibition

upregulates cyclin D and promotes cell cycle entry, we tested recently reported DYRK1A

inhibitors, LCTB-21 and LCTB-92, in neonatal rat cardiomyocytes[42,43]. LCTB-92 and LCTB-21

work by binding to the active site of DYRK1A and preventing kinase activity. Inactive forms of

both compounds (ISO-92 and ISO-21, respectively) were used as negative controls along with

DMSO. High-content imaging and segmentation of α-actinin, Ki67, cyclin D1, and DAPI (**Fig. 2A**)

were used to analyze cardiomyocyte cell cycle activity at single-cell resolution. The network model

predicted that DYRK1A inhibition should increase DNA replication in a dose-dependent manner

(**Fig. 2B**). Consistent with these predictions, treatment with higher concentrations of LCTB-21 (1

μM) and LCTB-92 (0.1 and 1 μM) increased cell cycle activity as evidenced by a higher

percentage of Ki67-positive cardiomyocytes compared to DMSO and isomeric controls (**Fig. 2C**).

The model also predicted a dose-dependent increase in cyclin D levels following DYRK1A

inhibition (**Fig. 2D**). Supporting the model predictions, validation experiments showed an increase

in cyclin D1 expression in cardiomyocytes treated with LCTB-21 (1 μM) and LCTB-92 (0.1 and 1

μM) compared to DMSO and isomeric controls (**Fig. 2E**).

Given the substantial increase in both Ki67 and cyclin D1 expression in cardiomyocytes

treated with 1 μM LCTB-92, we further investigated the relationship between these markers and

the cell cycle phase under this condition. Analysis of the mean Ki67 and cyclin D1 intensities

revealed a shift from a unimodal distribution in cells treated with ISO-92 to a bimodal distribution

in LCTB-92-treated cardiomyocytes (**Fig. 2F**). The shift suggests a discrete bifurcation in Ki67

and cyclin D1 expression that creates distinct subpopulations of cardiomyocytes[44]. Analysis of

DNA content and Ki67 (**Fig. 2G**) revealed an increased proportion of cardiomyocytes in the G2/M

phase (**Fig. 2H**) and a decreased proportion in the G0/4c (**Fig. 2I**) and G0/2c phases (*c* refers to

DNA content, with 2c and 4c representing diploid and tetraploid levels, respectively) following

LCTB-92 treatment compared to ISO-92. This shift toward late cell cycle progression, along with

a reduction in multinucleation (**Fig. 2J**) suggests that LCTB-92 promotes true cell division rather

than incomplete cell cycle progression.

Figure 2: DYRK1A inhibitor LCTB-92 increases cell cycle activity of neonatal rat cardiomyocytes, validating network model. A) Representative images of cardiomyocytes treated with DYRK1A inhibitors (LCTB-21, LCTB-92) or controls (DMSO, ISO-21, ISO-92) and labeled with DAPI (blue), α-actinin (red), Ki67 (yellow), and cyclin D1 (magenta). Scale bar = 100 mm. **B)** Network model prediction of DNA replication and **C)** experimental quantification of the percent of cardiomyocytes expressing Ki67 in response to DYRK1A inhibition. **D)** Network model prediction of cyclin D expression and **E)** experimental quantification of the percentage of

cardiomyocytes expressing cyclin D1 in response to DYRK1A inhibition. **F)** Single-cell correlation
between Ki67 and cyclin D1 expression levels with histograms illustrating the distribution of
intensities for each marker. **G)** Bivariate analysis of DNA content and Ki67 to separate out cell
cycle phase. DNA content thresholds separating 2c, 3c, and 4c nuclei are shown as blue vertical
lines. Ki67 positive threshold is shown as a red horizontal line. Percent of cardiomyocytes in **H)**
G2/M and **I)** G0/4c phases from panel G. **J)** Percent of multinucleated cardiomyocytes. Error bars
represent mean ± SEM. Statistical significance for C and E was determined by one-way ANOVA
followed by Tukey's multiple comparisons test. ******$p < 0.05$ LCTB treatments compared to DMSO
(0 mM) control; *****$p < 0.05$ LCTB treatments compared to the respective isomeric control.
Statistical significance for H-J was determined by t-test. *****$p < 0.05$.

To uncover the mechanistic differences that explain the efficacy of LCTB-92 vs. LCTB-21

in promoting CM proliferation, we performed RNA-sequencing on drug-treated NRCMs. Principal

component analysis captured 91% of the variance across samples and revealed distinct clustering

of our LCTB-92 and -21 treated groups (**Fig. 3A)**. Differential gene expression was performed in

both drug-treated groups with respect to their inactive isomers (**Fig 3B-E).** Gene ontology (GO)

enrichment analysis revealed that LCTB-92 treatment was associated with multiple terms related

to cell cycle activity with cell division having the highest gene count (padj < 0.05) (**Fig 3B).** We

found a much more pronounced effect in LCTB-92 treated cells with LCTB-92 having 1800 unique

differentially expressed genes (DEGs), which could explain the more robust effect on cell cycling

(**Fig 3E**).

To predict drivers of these changes in gene expression, we performed transcription factor

enrichment using ChEA3 on the LCTB-92 DEGs[45]. We found that E2F1 was a top hit according

to mean library rank (**Fig. S1A**). We then used the ChEA3 results to map from the network model's

E2F1 node to DEGs associated with cell cycle related processes (**Fig. 3F**)**.** Several E2F1 targets

were also related to cardiac contraction, suggesting cardiomyocyte dedifferentiation which is a

critical component of cardiomyocyte proliferation[46]. We then visualized the expression levels of

genes involved in cell division and cardiac contraction, identified through our GO term enrichment

analysis, across all treatment groups relative to DMSO (**Fig. 3G–H**). Notably, LCTB-92 treatment

elicited a more pronounced transcriptional response, with increased expression of positive

regulators and decreased expression of negative regulators of cell cycling. Additionally, we

observed a greater decrease in cardiac contraction gene expression as well as reduced

sarcomere organization in the LCTB-92 treated samples, suggesting more robust cardiomyocyte

dedifferentiation. Taken together, the integration of model predictions and transcriptomic data

suggest LCTB-92 is inducing a more defined effect on cell cycling genes compared to LCTB-21,

possibly through higher activation of E2F1.

Figure 3: LCTB-92 treated NRCMs exhibit transcriptomic signatures associated with cell division. (A) Principal Component Analysis of drug-treated and respective control NRCM samples. **(B)** Gene Ontology Enrichment Analysis of LCT-92 treated NRCMs with respect to its inactive isomer (ISO-92). **(C-E)** Number of significantly differentiated expressed genes with ≥ 2-fold increase in LCTB-92 treated **(C)** or LCTB-21 treated **(D)** NRCMs compared to ISO-92 and ISO-21 respectively, and **(E)** the commonality between both treatment groups. **(F)** Network model simulation of LCTB-92 inhibition of DYRK1A and corresponding gene expression of cytokinesis, DNA replication, and cardiac contraction E2F1-target genes. **(G-H)** Z-score of differential gene expression in genes pulled from Cell Division **(G)** and Cardiac Contraction **(H)** Gene Ontology terms. Drug and isomeric controls groups were compared to DMSO for figures **G & H**. GO term enrichment analysis was performed with a p-value cutoff of 0.01 and q-value cutoff of 0.05.

LCTB-92 promotes cardiomyocyte cell cycle activity and improves heart function post-MI

Based on the in vitro experimental results, we investigated the effects of LCTB-92 in an *in vivo* model of ischemic/reperfusion myocardial infarction (I/R MI). Specifically, we used our previously described αDKRC::RLTG transgenic mouse that restricts Cre recombinase expression to adult cardiomyocytes that re-enter the cell cycle[47]. The reporter mouse relies on the activation of the Ki67 promoter, providing an in vivo validation of the Ki67 results in our NRCM experiments. Cardiomyocytes that re-enter the cell cycle, based on the activation of a Ki67 promoter, express GFP. αDKRC::RLTG mice underwent I/R MI surgeries and were treated with 10 mg/kg LCTB-92 or an inactive isomeric control (ISO-92) via daily oral gavage for ten days (**Fig. 4A**). Serial echocardiography revealed that mice treated with LCTB-92 had significantly improved left ventricular ejection fractions (LVEFs) compared to ISO-92 treatment at both 2- and 4-weeks post-MI (**Fig 4B**). However, LCTB-92 treatment did not improve infarct size compared to control 4-weeks post-MI, despite improvement in heart function, suggesting the mechanism of improved cardiac function was not through infarct size reduction (**Fig 4**. **C & D).** Additionally, the hearts of mice treated with LCTB-92 had significantly increased number of cycled (GFP+) cardiomyocytes compared to ISO-92 treatment (**Fig. 5A & B**). Interestingly, LCTB-92 did not induce a significant increase of CM cycling in uninjured hearts, suggesting an injury-dependent context for the effects of LCTB-92 on cardiomyocyte cycling.

We then investigated if the improvement in heart function was linked to the increase in cell

cycle activity induced by treatment with LCTB-92 using a previously described mouse model that

ablates cardiomyocytes upon entry into the cell cycle through the Cre-mediated expression of

Diphtheria toxin[48]. αDKRC::RLTG/DTA or +::RLTG/DTA mice underwent I/R MI and were treated

with 10 mg/kg LCTB-92 via daily oral gavage for ten days (**Fig. 6A**). Serial echocardiography

showed that the effects of LCTB-92 were attenuated when cycling cardiomyocytes were ablated

after MI (**Fig 6. B-D**). Infarct sizes were similar between the two groups (**Fig 6. E-F**). The results

suggest that the beneficial effects of LCTB-92 on cardiac function after MI require increases in

cycling cardiomyocytes.

Figure 4: LCTB-92 improves heart function after I/R MI. (A) Schematic of 12-week-old αDKRC::RLTG mice that were treated with LCTB-92 or isomeric control for 10 days (10 mg/kg) following I/R MI. **(B)** Cardiac function measurements of αDKRC::RLTG mice treated with LCTB-92 or isomeric control after I/R MI, showing left ventricular ejection fraction (LVEF) 2- and 4-weeks

post-MI. Error bars represent the mean±SD and open circles are measurements for individual
animals. **(C)** Average infarct size (%) across whole heart sections from hearts treated with LCTB-
92 or isomeric control 4 weeks after MI with **(D)** representative images. Error bars represent the
mean±SD and open circles are measurements for individual animals. P-values were calculated
using two-way ANOVA followed by Bonferroni's multiple comparisons test for panels B-D. Two
samples t-test was performed for panel C.

Figure 5: LCTB-92 promotes cardiomyocyte re-entry after I/R MI. (**A**) Quantification of the #
of GFP$^+$ cardiomyocytes collected from the midventricular section of hearts treated with LCTB-92
or isomeric control 4 weeks after MI with (**B**) representative images of Masson's trichrome stained
(left) and GFP$^+$ cardiomyocytes (right). P-values were calculated using two-way ANOVA followed
by Bonferroni's multiple comparisons test.

Figure 6: Ablation of cycling cardiomyocytes attenuates beneficial effects of LCTB-92 treatment. (A) Schematic of 12 week old αDKRC::RLTG/DTA or +::RLTG/DTA mice that were treated with LCTB-92 10 days (10 mg/kg) following I/R MI. **(B** and **C)** Cardiac function measurements of αDKRC::RLTG/DTA or control (+::RLTG/DTA) mice treated with LCTB-92 or isomeric control after I/R MI, showing absolute **(B)** absolute (%), and **(C)** percent change (% change from baseline) LVEF 2- and 4-weeks post-MI. **(D)** Representative echocardiography parasternal long-axis views of controls and DYRK1A k/o mice 4-weeks after I/R MI. **(E)** Average infarct size (%) across whole heart sections from hearts treated with LCTB-92 or isomeric control 4 weeks after MI with **(F)** representative images. P-values were calculated using two-way ANOVA followed by Bonferroni's multiple comparisons test for panels B and C. Two samples t-test was performed for panel E.

The post-developmental genetic ablation of DYRK1A recapitulates beneficial effects of

LCTB-92 treatment

Previously we observed aMHC-Cre::DYRK1A[flox/flox] mice in which cardiomyocyte-specific

deletion of DYRK1A starting in development had baseline cardiac hyperplasia and an increase in

the expression of cell cycle genes[11]. To complement LCTB-92 inhibitor experiments, we

investigated the effects of the post-developmental, cardiomyocyte-specific ablation on DYRK1A

in adult mice that underwent I/R MI. cTnnt2-Cre[Ert2]/+::Fucci2aR/Fucci2aR:DYRK1A[flox/flox]

(DYRK1A K/O) or cTnnt2-Cre[Ert2]/+::Fucci2aR/Fucci2aR (Control) mice were pulsed with

tamoxifen for 10 days followed by a 2-week recovery period and then underwent an I/R MI with

serial echocardiograms (**Fig. 7A**). DYRK1A K/O had improvements in LVEF at 2- and 4-weeks

after MI compared to controls (**Fig. 7B & C**). Similar to LCTB-92 treated mice that underwent I/R

MI, there were no significant differences in infarct size between our DYRK1A K/O and control

groups (**Fig. 7D & E**). Together, the genetic ablation of DYRK1A in adult cardiomyocytes

recapitulated the effects of LCTB-92 treatment after MI, further supporting the potential

therapeutic benefits DYRK1A inhibition after MI.

Figure 7: Post-developmental cardiomyocyte-specific deletion of DYRK1A recapitulates improved heart function induced by LCTB-92 treatment after I/R MI. (A) Schematic of 12 week old cTnnt2-Cre[Ert2]/+::Fucci2aR/Fucci2aR::DYRK1A[flox/flox] (Cardiomyocyte-specific DYRK1A K/O) or cTnnt2-Cre[Ert2]/+::Fucci2aR/Fucci2aR (Control) mice that undergo I/R MI. **B)** absolute (%), and **(C)** percent change (% change from baseline) LVEF 2- and 4-weeks post-MI. **(D)** Average infarct size (%) across whole heart sections from DYRK1A K/O or control hearts 4 weeks after MI with **(E)** representative images. P-values were calculated using two-way ANOVA followed by Bonferroni's multiple comparisons test for panel B. Two samples t-test was performed for panels C and D.

DISCUSSION

Here, we mechanistically propose how small molecule inhibition of DYRK1A leads to increased cell cycling of cardiomyocytes both *in vitro* and *in vivo*. Our network model accurately predicts DYRK1A-mediated cell cycle activity that we leveraged to inform our *in vitro* validation experiments in NRCMs. Treatment of NRCMs with LCTB-21 and LCTB-92 exhibited a dose-dependent response on both cyclin D1 and Ki67 expression validating our model predictions. RNA-sequencing of LCTB-21 and LCTB-92 treated NRCMs revealed distinct transcriptional regulation of cell cycling genes, which we mapped to our network model's predicted node activity via E2F1. LCTB-92 treatment induced CM cycling and improved heart function after I/R MI but did not alter infarct size. However, the beneficial effects of LCTB-92 treatment on heart function were lost when we genetically ablated cycling CMs, indicating that the functional benefits are mediated through cycling CMs. The post-developmental genetic ablation of DYRK1A attenuated the left ventricular dysfunction observed after MI, similar to the effects of LCTB-92. The two complementary approaches to prevent DYRK1A activity by pharmacological inhibition or genetic ablation specifically in adult CMs improved cardiac function after MI, supporting that DYRK1A knockdown after injury can lead to therapeutic benefits.

Our network model was built using literature in various cell types, as DYRK1A's regulation of cell cycling is believed to be highly conserved. The model predicted experimental outcomes reported in cardiomyocyte literature not used to develop the model, despite the use of predominately non-cardiomyocyte literature to construct the model. Interestingly, while the model retained high accuracy against experiments from most cancer literature, the model failed to recapitulate experimental results when knocking down or overexpressing DYRK1A[31,38]. The inconsistency of the model to predict DYRK1A perturbation in certain cancer cells could be due to the type of cancer cell studied, where DYRK1A has been reported to be an oncogene. For example, DYRK1A knockdown reduced tumor growth in both *in vitro* and *in vivo* models of colon, breast, and neck cancers (colon, breast, neck)[49].

We show here that while LCTB-92 promotes CM cycling NRCMs and after I/R MI *in vivo*,

it does not induce cycling in uninjured adult hearts. This could be due to the hypoxic environment

created after injury. Several reports have demonstrated hypoxic environments can induce CM

proliferation. Puente et al. (2014) showed that by exposing NRCMs to hypoxia, they could extend

their proliferative window by reducing oxidative stress and DNA damage signaling[50]. Similarly,

Johnson et al. (2023) showed that chronic systemic hypoxia in adult mice induced limited

cardiomyocyte proliferation in the right ventricle[51]. These findings suggest that the post-injury

hypoxic microenvironment may sensitize CMs to DYRK1A inhibition, thereby enhancing the

proliferative response observed with LCTB-92 after MI.

The gene dosage of DYRK1A has also been observed to play an important role in CM

cycling. Lana-Eloa et al. (2024) showed that Dp1Tyb mice, a line that has three copies of

chromosome 21, including DYRK1A, and phenocopies human Down Syndrome, have

proliferative defects in the embryonic heart[14]. Flow cytometry showed an increase in the

percentage of CMs in G1 phase, indicating a larger quiescent cell population. Additionally, they

observed reduced p-RB1 and transcription of E2F target genes, consistent with our model

predictions. Lastly, RNA-sequencing showed that LCTB-21 was able to partially rescue the

decreased expression of proliferative pathways in Dp1Tyb embryonic hearts. Treatment with

LCTB-92 might be able to induce a more robust recovery of these defects.

The αDKRC mouse model restricts Cre expression to adult CMs that express Ki67. One

limitation is that the reporter mouse cannot distinguish between CMs that enter the cell cycle and

undergo endoreplication to increase ploidy and CMs that undergo complete mitosis and

cytokinesis to promote new daughter cells. Our prior investigations demonstrated that ~10% of

adult CMs that enter the cell cycle undergo bona fide proliferation as determined by pairs of

clusters of Ki67-expressing CMs[48]. With this in mind, we have several hypotheses of how the

induction of CM cycling is responsible for the improvement of heart function we see with the

treatment of LCTB-92 after I/R. LCTB-92 could be inducing complete CM proliferation that can

restore some of the lost myocardium induced by I/R injury and improve the contractile function of

the heart. Alternatively, LCTB-92 could also result in the induction of more polyploid CMs which

have been shown to be more responsive to stress through enhanced mitophagy[52]. Lastly, cycling

CMs may produce paracrine factors that are cardioprotective to the surrounding myocardium.

Further work is necessary to understand the contributing role of LCTB-92 induced cycling CMs in

recovery following I/R MI.

The potential of the LCTB class of DYRK1A inhibitors to treat acute MI in patients is very

attractive. Currently, LCTB-21 is being investigated in a first-in-human Phase 1 trial in healthy

volunteers and subjects Down Syndrome and Alzheimer's disease (Clinical Trial NCT06206824).

While we show that both LCTB-21 and LCTB-92 induce cycling in CMs *in* vitro, LCTB-92 showed

stronger cell cycling effects which led to our investigation of LCTB-92 *in vivo.* Whether LCTB-21

has similar effects as LCTB-92 on cardiac function and CM cycling after MI remains to be seen.

Importantly, we did not observe tumorigenic effects of LCTB-92 in our studies. The results of this

trial, in conjunction with our results using LCTB-92 in a preclinical animal model of MI, are

encouraging in that the inhibition of DYRK1A may represent a new first-in-class therapy to treat

heart disease. Follow up studies that recapitulate the results here in more human relevant model

systems would be necessary to investigate the potential of these compounds in treating MI.

MATERIALS AND METHODS

Study Design: The aim of this study was to evaluate if a specific oral inhibitor of DYRK1a would improve left ventricular function after myocardial infarction and increase adult cardiomyocyte cycling. The study used cultured neonatal rat ventricular myocytes (NRVMs) to examine Ki67 expression in response to LCTB-92, LCTB-21, and their inactive isomers. **RNAseq was used to investigate changes in the gene expression of NRVMs in response to LCTB-92 and Iso-LCTB-92.** For in vivo studies, we used *αDKRC::RLTG* mice that restrict Cre expression to adult mouse cardiomyocytes that re-enter the cell cycle. *αDKRC::DTA* mice were used to express Diphtheria toxin and ablate in adult cardiomyocytes that re-enter the cell cycle. *cTnnt2-CreERt2::DYRK1a^flox/flox^* mice were used to investigate the genetic ablation of DYRK1a in adult cardiomyocytes. Ischemia-reperfusion myocardial infarctions were performed by ligating the proximal left anterior descending coronary artery for 60 minutes, followed by restoration for blood flow. This model recapitulated acute myocardial infarction and reperfusion in humans. Cycling cardiomyocytes were quantified by counting eGFP+ cells in 10-micron short-axis histological heart sections from *αDKRC::RLTG* mice. Infarct sizes were quantified by examining serial sections stained with Masson trichrome. Sample sizes were selected based on previous experience with similar methods. Mice were randomized to treatment groups based on genotypes for each experiment. Investigators were blinded to genotype when performing surgeries, echocardiography, quantification of eGFP+ cardiomyocytes, and analyses of infarct sizes. No samples were excluded from the study. Details on sample sizes representing biological replicates and statistical tests are detailed in figure legends and the Statistical Analysis section of the Materials and Methods.

Network Model Construction: A logic-based differential ordinary equations network model was used to predict mechanisms of DYRK1A-mediated cell cycle activity. The network model was built using Netflux[23], a software that auto-generates a logic-based differential equation model based

on a list of reaction rules and parameters[22]. This includes assigning the interactions (inhibitory or

stimulatory) between molecular components of interest (nodes) as well as the logic-gating

strategy ("OR" or "AND" gates). Further information on how this modeling approach works can be

found in Kraeutler et al. (2010) and Clark et al. (2024). A series of 6 primary and review literature

articles (**Table S3**) were used to infer the interactions and logic-gating strategy within the model

(**Table S1**). Literature was chosen based on its relevance to DYRK1A-mediated cell cycle control

and dimerization partner, RB-like, E2F and multi-vulval class B (DREAM) complex regulation.

While literature that used cardiomyocytes or other relevant experimental models were preferred,

we used literature across multiple experimental models (noted in **Table S3**) as this regulatory

process is believed to be highly conserved across tissue types. Literature search key terms

included: proliferation, cell cycling, DREAM complex, and DYRK1A.

Default reaction parameters included reaction weight (w=1), Hill coefficient (n=1.4) and

half-maximal effective concentration (EC_{50}=0.50) and default node parameters included initial

activation (Y_{init}=0), maximal activation (Y_{max}=1), and time constant (τ =1). Parameter optimization

was run on input nodes to allow the model to experience change with perturbations at baseline.

DYRK1A and B-Myb were then set to a Y_{max}=0.9 and 0.1 respectively to establish baseline activity.

As in past logic-based differential equation network models (Kraeutler et al., 2010; Tan et al.,

2017; Zeigler et al., 2016) species (nodes) refer to a small molecule, gene, protein, or process.

Reactions (or edges) are activating and inhibiting relationships between network species[22,53,54].

Network Model Validation: Model simulations were compared to experimental data from primary

literature that was separate from the literature that was used to build the model's interactions and

logic-gating strategy. Guided by the experimental design of each validation study, we simulated

knockdowns or overexpression of specific nodes to see how these perturbations could affect the

DNA replication node within the model. Knockdowns were performed by setting the Y_{max} of a

desired node to 0. Overexpression was performed by setting Y_{max}= 10 for input nodes or by setting

Y_{init}= 10 and tau= 10^9 for intermediate nodes. All perturbations were performed after the model

reached a steady state (<0.05% change in activity for all nodes). Model accuracy was determined

by if simulation results qualitatively matched the statistics from the corresponding experiment in

the primary literature (n = 19). Validation was performed across cancer and cardiomyocytes to

look at deviations in model predictions across different biological contexts. Experimental figures

were replotted with WebPlotDigitizer[55].

Network Model Simulations: We simulated a concentration response by knocking down

DYRK1A in increments of 1%. We then looked at the change in activity of the responding nodes

to characterize their sensitivity to a DYRK1A knockdown. EC50 and IC50 of affected nodes were

determined by the % DYRK1A knockdown at which their normalized activity was changed by 50%.

A knockdown screen was conducted through simulated 100% knockdowns on each node of the

model at baseline activity, where DYRK1A is highly active (Y_{max}=1). We then examined the change

in activity of the network in response to each individual knocked down node. We simulated "low,

medium, and high" DYRK1A inhibitor concentrations DYRK1A by increments of 15% to simulate

"low, medium, and high" drug concentrations. Our concentrations corresponded to these

simulated knockdown values; low: 15%, medium: 30%, and high: 45%. The results were then

compared to in vitro data of neonatal rat cardiomyocytes that received the LCTB-92 and LCTB-

21 DYRK1A inhibitors.

Neonatal rat cardiomyocyte isolation, immunostaining, and image analysis

Cardiomyocytes were isolated from 1-2-day-old Sprague-Dawley rats using a Neomyt

isolation kit (Cellutron). Each cell isolation is a mixture of cardiomyocytes from all 9-12 male and

female pups from that litter. The cells were cultured in plating media (low-glucose Dulbecco's

modified eagle media (DMEM), 17% M199, 10% horse serum, 5% fetal bovine serum (FBS), 1%

L-Glutamine, 10 U/mL penicillin, and 50 mg/mL streptomycin) at a density of 30,000 cells per well

in a 96-well Corning CellBIND plate coated with SureCoat (Cellutron). After being cultured for 48

hours, the cells were serum-starved in WE+B medium (William's E medium without phenol red

(Gibco) supplemented with Primary Hepatocyte Cell Maintenance Cocktail B (Gibco) containing

P/S, ITS+, GlutaMAX, and HEPES) for 4 hours before drug treatment. Cardiomyocytes were then

treated with the DYRK1A inhibitors LCTB92 or LCTB21 (0.01-1 mM) or respective isomeric and

DMSO (0.01%) controls in serum-free media for 48 hours.

Cells were then fixed in 4% paraformaldehyde and fluorescently labeled with anti α-actinin

(Sigma) or cardiac troponin T (Abcam), anti-Ki67 (Invitrogen), anti-cyclin D1 (Abcam), and DAPI

(Life Technologies). Stained cells were imaged with the Operetta CLS high-content imaging

system using a 10x 0.3 NA objective. Image analysis was performed using a CellProfiler pipeline

for cell segmentation to quantify the percentage of cardiomyocytes positive for Ki67 or cyclin D1[56].

Thresholds for mean nuclear Ki67 or cyclin D1 intensity were set manually to approximate the

local minimum of the bimodal distribution. Multinucleated cardiomyocytes were identified by

measuring the distance between adjacent nuclei, with a threshold of less than 2 pixels. Cell cycle

phase was assessed by combining DNA content analysis and Ki67 positivity as described in

previous studies[57,58].

Gene Expression Analysis: NRCMs were isolated and cultured as previously described, then

seeded at a density of 1,000,000 cells per well in 6-well Corning CellBIND plates coated with

SureCoat (Cellutron). Cells were treated with 1 μM of either LCTB-92 or LCTB-21, their respective

isomeric controls, or DMSO (n = 3 per condition), following the same treatment protocol as

described above. RNA extraction, cDNA library preparation (including adapter ligation), and

sequencing were performed by Genewiz, Inc. using an Illumina® HiSeq® system. Principal

component analysis (PCA) and differential gene expression were conducted using the **DESeq2**

package in R[59]. Gene Ontology (GO) enrichment analysis was performed with the **ClusterProfiler**

package in R[60]. Transcription factor enrichment was assessed using the **ChEA3** API platform,

with transcription factors ranked by average rank across all libraries[45]. To integrate RNA-seq data

with our network model, we linked the E2F1 node to its target genes identified via ChEA3 and

mapped these to gene sets associated with DNA replication, cytokinesis, and cardiac contraction

GO terms.

Mice. *αDKRC (αMHC-MerDreMer-Ki67p-RoxedCre::Rox-Lox-tdTomato-eGFP) mice were made*

and maintained as previously described. C57BL/6J (JAX stock #000664), *RC::RLTG* (*B6.Cg-*

Gt(ROSA)26Sor$^{tm1.2(CAG-tdTomato,-EGFP)Pjen}$/J),   and   *ROSA-DTA*   (*B6.129P2-*

Gt(ROSA)26Sor$^{tm1(DTA)Lky}$/J) mice were purchased from Jackson labs. *CAG-STOP-Fucci2aR* was

obtained from the European Mouse Mutant Archive. *cTnnt2-Cre^{ERt2}* was provided by Dr. Chenlen

Cai.   The   *αDKRC::RLTG*,   *αDKRC::DTA*,   *cTnnt2-Cre^{ERt2}::Fucci2aR*,   and   *cTnnt2-*

Cre^{ERt2}::Fucci2aR::DYRK1a$^{flox/flox}$ mice were created by standard breeding methods and all mice

were genotyped. All mice were housed and maintained in accordance with UVA Animal Care and

Use Committee approved protocols (UVA Animal Care and Use Committee (ACUC) Wolf Lab

protocol #4080).

Compounds: LCTB-92, Iso-LCTB-92, LCTB-21, and Iso-LCTB-21 were provided by Perha

Pharmaceuticals. For experiments using NRVMs, LCTB-92, Iso-LCTB-92, LCTB-21, and Iso-

LCTB-21 were dissolved in DMSO and added to cell media to give a final contraction of 0.01 uM,

0.1 uM, or 1 uM. For in vivo experiments, LCTB-92 or Iso-LCTB-92 was dissolve in 0.5%

carboxymethylcellulose in sterile water and 1 mg/kg was given daily by oral gavage.

Left Anterior Descending (LAD) I/R MI surgery: All animal care and surgeries were in

accordance with UVA ACUC Policy on Rodent Surgery and Perioperative Care under ACUC-

approved animal protocol (UVA ACUC Wolf Lab protocol #4080). The individual performing

surgeries was blinded to the mouse genotypes and treatments. Procedure: Male and female ten-

to twelve-week-old mice were randomized to MI surgeries, weighed and then anesthetized in an

induction chamber using the gas anesthetic Isoflurane (3% volume/weight, oxygen 500ml/min).

The animal was placed in the supine position on a face mask connected to the anesthesia system.

Isoflurane was adjusted to provide a maintenance level (1.8-2.2% volume/weight, oxygen

500ml/min) throughout the procedure. Anesthesia was monitored closely by the following

methods: (1) Depth and rate of respiration; (2) Heart rate by ECG; (3) Mucous membrane color;

(4) Body Temperature via Physio Suite Homoeothermic Temperature Monitoring System; (5)

Reflexes & toe pinch; (6) Overall appearance of muscle relaxation. All surgical procedures were

carried out with a stereo microscope exclusively for small animal surgery. Normal body

temperature was be maintained using an electrical heating pad on a feedback system via rectal

probe.

To prepare for surgery, the mouse was first given subcutaneous fluids (veterinary

Normosol), and Atropine as a pre-anesthetic to decrease mucous secretion and to prevent gag

reflex during intubation. The neck and chest wall were shaved, and then prepped three times with

alternating wipes of povidone-iodine and 70% ethanol. Mice were placed in the supine position

on a heating pad and an endotracheal intubation was performed under direct laryngoscopy with

a lighted fiber optic stylus to visualize the vocal cords and insert tracheal tube. The endotracheal

tube was then connected to the automated ventilator (Kent Scientific, Inc.) and the mouse was

ventilated (tidal volume = 1.0 mL, rate = 120 breaths/minute). Bupivacaine was infiltrated

subcutaneously into the area of the left 3rd and 4th intercostal space. A left anterior thoracotomy

was performed using sterile technique by making a small subcutaneous incision lateral from the

sternum with sharp scissors. The pectoralis muscle groups were identified and separated by blunt

dissection, and held open using fixed retractors. The thoracic cage was exposed and the 3rd

intercostal space was identified. Incision into the 3rd intercostal space was done by a combination

of blunt micro-scissors and micro-cauterizing tool in order to minimize bleeding. Retractors were

then repositioned onto the upper and lower ribs in order to visualize the heart. The pericardium

was then blunt dissected in order to expose the anterior view of the heart. The LAD on the surface

of the left ventricular wall was identified and an 8-0 prolene suture was placed through the

myocardium into the anterolateral LV wall underneath the LAD at the level of the lower atrium (1

mm below the left auricle), and the suture was tied. Occlusion of blood flow is observed by

blanching of the heart muscle and ST elevation confirmed by ECG. After 60 minutes the ligature

was removed. The chest wall, facial planes, and skin were sutured and the mouse was recovered

and treated with analgesics. In sham control mice, the entire procedure was identical except for

the ligation of the LAD. The mortality rate associated with surgeries was ~10%.

Histology and Immunohistochemistry: Hearts were excised and fixed in 10% Neutral Buffered

Formalin (Fisher, Inc.) for a minimum of four hours prior to embedding in paraffin. Ten-micron

sections were prepared in short axis orientation by microtome with 8 sections per glass slide.

Paraffin was removed and the tissue sections were rehydrated using Xylene and serial ethanol

wash steps, respectively. Antigen retrieval was performed by incubating tissue sections in boiling

1x Unmasking solution (Vector Labs H-3300) for 22 minutes. After cooling to room temperature,

the tissue sections were treated with Sudan Black to quench auto-fluorescence. Briefly, tissue

sections were incubated in 0.1% Sudan Black (Sigma, Cat# 199664) in 1x PBS and 70% ethanol

at room temperature for 20 minutes to quench auto fluorescence, followed by 3 x five-minute

washes in 1x PBS containing 0.02% Tween 20, and a final five-minute wash in 1x PBS.

Quantification of cardiomyocyte cycling: Images were obtained using a Leica DM2500

Fluorescence microscopy system with a Leica DFC7000 T fluorescence color camera or a Leica

THUNDER imaging system and Leica LAS X Multi Channel Acquisition software. For imaging,

fluorescence channels were calibrated to background of control sections stained with secondary

antibodies alone. eGFP+ CMs visualized using an YFP filter and quantified from 6-8 ten-micron

short axis sections of three slides per animal separated by ~400 microns per slide. The infarct

zone was identified by WGA. The non-infarct zone was divided into thirds with the two regions

adjacent to the infarct defined as the border zones and the middle region defined as the remote

zone. eGFP+ CMs present in the same location on sequential slides were counted once among

all sections to avoid over-representation of eGFP+ cells.

Echocardiography: Mice were anesthetized with Isoflurane and gently restrained on a Vevo

integrated rail system. The system included a physiological monitoring unit consisting of a heating

board with integrated ECG electrodes. The table temperature was maintained at ~38oC using a

rectal probe for temperature feedback. Hair was gently removed, and ultrasound contact gel

(warmed to 37oC) was applied to the chest. Echocardiography was performed using a high

frequency 30 MHz linear transducer and a Vevo 1100 (Visual Sonics) system similar to previously

described methods.[80,81] Initial B-mode images were obtained in the parasternal long axis with

the apex and aortic valves visualized. Next, B-mode short axis images were obtained by turning

the ultrasound probe ~90 degrees and identifying the mitral valve papillary muscles. The

parasternal long axis images were analyzed using VevoView software to calculate left ventricular

dimensions. The individuals performing and analyzing the echocardiography were blinded to the

treatment groups.

Statistics: GraphPad Prism 9 (GraphPad Software, Inc.) was used for statistical analyses. The

data was analyzed for normal distribution using Anderson-Darling, D'Agostino & Pearson,

Shapiro-Wilk, and Kolmogorov-Smirnov tests in GraphPad. One-way ANOVAs with Tukey test

corrections for multiple comparisons, two-way repeated measured ANOVA with Bonferroni test for

multiple comparisons, and student t-tests were used. We used a calculator from the IACUC at

Boston University, including an Excel template for calculation based on means/standard

deviations   and   proportions   (https://www.bu.edu/researchsupport/compliance/animal-

care/working-with-animals/research/sample-size-calculations-iacuc/).   For   experiments

quantifying eGFP+ CMs, we assumed a Type I (alpha) error of 0.05, a difference (delta) of 2-fold

(~2-3 eGFP+ cells in sham heart sections and 4-6 eGFP+ cells in MI heart sections), a standard

deviation of 2-3, and a Power of 0.9. We calculated that nine mice would be needed in each group.

For the echocardiography experiments, we assumed a Type I (alpha) error of 0.05, a difference

(delta) of ~10% (EF ~30% in the MI control group and ~40% in the drug treated group), a standard

deviation of 5%, and a Power of 0.95. We calculated that a minimum of ~seven mice would be

needed in each group.

ACKNOWLEDGEMENTS

Funding: This work was supported by the National Institutes of Health grants R01HL158718 to

MJW, R01HL162925 to JJS and MJW, R01HL160665 to JJS, and T32GM13661 to BCM.

Author contributions: J.J.S. and M.J.W. designed the study, obtained funding, supervised the

work, and edited the manuscript. B.C.M. performed the network modeling, RNA sequencing, and

experiments on tissue sections, and wrote the manuscript. K.L.W. performed the in vitro

experiments, image analysis, and wrote those sections of the manuscript. B.N.H., M.W., and C.Z.

performed network modeling. A.Y, L.A.B., K.S., D.H., L.M., and M.J.W. designed, performed, and

analyzed the mouse surgeries, echocardiography, histology, and immunohistochemistry. L.M.,

M.F.L., and E.D. provided the synthesized and purified LCTB-92, ISO-92, LCTB-21, and ISO-21

and expertise regarding the design of the mouse experiments.

Competing interests: All other authors declare they have no competing interests.

Data and materials availability: All data associated with this study are present in the paper or

Supplementary Materials. Bulk RNA sequencing data will be deposited in the Gene Expression

Omnibus.

REFERENCES

1.   Tsao, C. W. *et al.* Heart Disease and Stroke Statistics—2023 Update: A Report From the

American Heart Association. *Circulation* **147**, (2023).

2.   Prabhu, S. D. & Frangogiannis, N. G. The Biological Basis for Cardiac Repair After

Myocardial Infarction: From Inflammation to Fibrosis. *Circ Res* **119**, 91–112 (2016).

3.   Parmacek, M. S. & Epstein, J. A. Cardiomyocyte Renewal. *N Engl J Med* **361**, 86–88

(2009).

4.   Lázár, E., Sadek, H. A. & Bergmann, O. Cardiomyocyte renewal in the human heart:

insights from the fall-out. *Eur Heart J* **38**, 2333–2342 (2017).

5.   Hassink, R. J. *et al.* Cardiomyocyte cell cycle activation improves cardiac function after

myocardial infarction. *Cardiovascular Research* **78**, 18–25 (2008).

6.   Lin, Z. *et al.* Cardiac-Specific YAP Activation Improves Cardiac Function and Survival in

an Experimental Murine MI Model. *Circulation Research* **115**, 354–363 (2014).

7.   Xin, M. *et al.* Hippo pathway effector Yap promotes cardiac regeneration. *Proc. Natl. Acad.*

Sci. U.S.A. **110**, 13839–13844 (2013).

8.   Zhang, Y. & Del Re, D. P. A growing role for the Hippo signaling pathway in the heart. *J*

Mol Med **95**, 465–472 (2017).

9.   Zhou, Q., Li, L., Zhao, B. & Guan, K.-L. The Hippo Pathway in Heart Development,

Regeneration, and Diseases. *Circulation Research* **116**, 1431–1447 (2015).

10.   Mohamed, T. M. A. *et al.* Regulation of Cell Cycle to Stimulate Adult Cardiomyocyte

Proliferation and Cardiac Regeneration. *Cell* **173**, 104-116.e12 (2018).

11.   Young, A. *et al.* Inhibition of DYRK1a Enhances Cardiomyocyte Cycling After Myocardial

Infarction. *Circulation Research* **130**, 1345–1361 (2022).

12.   Litovchick, L., Florens, L. A., Swanson, S. K., Washburn, M. P. & DeCaprio, J. A. DYRK1A

protein kinase promotes quiescence and senescence through DREAM complex assembly. *Genes*

Dev **25**, 801–813 (2011).

13.   Hille, S. *et al.* Dyrk1a regulates the cardiomyocyte cell cycle via D-cyclin-dependent

Rb/E2f-signalling. *Cardiovasc Res* **110**, 381–394 (2016).

14.   Lana-Elola, E. *et al.* Increased dosage of DYRK1A leads to congenital heart defects in a

mouse model of Down syndrome. *Sci. Transl. Med.* **16**, eadd6883 (2024).

15.   Ables, J. L. *et al.* A Phase 1 single ascending dose study of pure oral harmine in healthy

volunteers. *J Psychopharmacol* **38**, 911–923 (2024).

16.   Najas, S. *et al.* DYRK1A-mediated Cyclin D1 Degradation in Neural Stem Cells

Contributes to the Neurogenic Cortical Defects in Down Syndrome. *EBioMedicine* **2**, 120–134

(2015).

17.   Guiley, K. Z. *et al.* Structural mechanisms of DREAM complex assembly and regulation.

Genes Dev **29**, 961–974 (2015).

18.   Fischer, M., Schade, A. E., Branigan, T. B., Müller, G. A. & DeCaprio, J. A. Coordinating

gene expression during the cell cycle. *Trends Biochem Sci* **47**, 1009–1022 (2022).

19.   Dyson, N. J. RB1: a prototype tumor suppressor and an enigma. *Genes Dev* **30**, 1492–

1502 (2016).

20.   Soppa, U. *et al.* The Down syndrome-related protein kinase DYRK1A phosphorylates

p27(Kip1) and Cyclin D1 and induces cell cycle exit and neuronal differentiation. *Cell Cycle* **13**,

2084–2100 (2014).

21.   Ren, B. *et al.* E2F integrates cell cycle progression with DNA repair, replication, and

G(2)/M checkpoints. *Genes Dev* **16**, 245–256 (2002).

22.   Kraeutler, M. J., Soltis, A. R. & Saucerman, J. J. Modeling cardiac β-adrenergic signaling

with normalized-Hill differential equations: comparison with a biochemical model. *BMC Syst Biol*

4, 157 (2010).

23.   Clark, A. P., Chowkwale, M., Paap, A., Dang, S. & Saucerman, J. J. Logic-based modeling

of biological networks with Netflux. (2024) doi:10.1101/2024.01.11.575227.

24.   Soonpaa, M. H. *et al.* Cyclin D1 overexpression promotes cardiomyocyte DNA synthesis

and multinucleation in transgenic mice. *J Clin Invest* **99**, 2644–2654 (1997).

25.   Liao, H. S. *et al.* Cardiac-specific overexpression of cyclin-dependent kinase 2 increases

smaller mononuclear cardiomyocytes. *Circ Res* **88**, 443–450 (2001).

26.   Sdek, P. *et al.* Rb and p130 control cell cycle gene silencing to maintain the postmitotic

phenotype in cardiac myocytes. *J Cell Biol* **194**, 407–423 (2011).

27.   Di Stefano, V., Giacca, M., Capogrossi, M. C., Crescenzi, M. & Martelli, F. Knockdown of

cyclin-dependent kinase inhibitors induces cardiomyocyte re-entry in the cell cycle. *J Biol Chem*

286, 8644–8654 (2011).

28.   Jin, Y. *et al.* B-Myb Is Up-Regulated and Promotes Cell Growth and Motility in Non-Small

Cell Lung Cancer. *Int J Mol Sci* **18**, 860 (2017).

29.   Biliran, H. *et al.* Overexpression of cyclin D1 promotes tumor cell growth and confers

resistance to cisplatin-mediated apoptosis in an elastase-myc transgene-expressing pancreatic

tumor cell line. *Clin Cancer Res* **11**, 6075–6086 (2005).

30.   Scaltriti, M. *et al.* Cyclin E amplification/overexpression is a mechanism of trastuzumab

resistance in HER2+ breast cancer patients. *Proc Natl Acad Sci U S A* **108**, 3761–3766 (2011).

31.   Li, Y.-L. *et al.* DYRK1A reinforces epithelial-mesenchymal transition and metastasis of

hepatocellular carcinoma via cooperatively activating STAT3 and SMAD. *J Biomed Sci* **29**, 34

(2022).

32.   Zhu, L. *et al.* Inhibition of cell proliferation by p107, a relative of the retinoblastoma protein.

Genes Dev **7**, 1111–1125 (1993).

33.   Claudio, P. P. *et al.* Mutations in the retinoblastoma-related gene RB2/p130 in lung tumors

and suppression of tumor growth in vivo by retrovirus-mediated gene transfer. *Cancer Res* **60**,

372–382 (2000).

34.   Jones, R. A. *et al.* RB1 deficiency in triple-negative breast cancer induces mitochondrial

protein translation. *J Clin Invest* **126**, 3739–3757 (2016).

35.   Liu, Q. *et al.* A MYBL2 complex for RRM2 transactivation and the synthetic effect of

MYBL2 knockdown with WEE1 inhibition against colorectal cancer. *Cell Death Dis* **12**, 683 (2021).

36.   Choi, Y. J. *et al.* The requirement for cyclin D function in tumor maintenance. *Cancer Cell*

22, 438–451 (2012).

37.   Geng, Y. *et al.* Kinase-independent function of E-type cyclins in liver cancer. *Proc Natl*

Acad Sci U S A **115**, 1015–1020 (2018).

38.   Radhakrishnan, A. *et al.* A dual specificity kinase, DYRK1A, as a potential therapeutic

target for head and neck squamous cell carcinoma. *Sci Rep* **6**, 36132 (2016).

39.   Ruiz, S. *et al.* Abnormal epidermal differentiation and impaired epithelial-mesenchymal

tissue interactions in mice lacking the retinoblastoma relatives p107 and p130. *Development* **130**,

2341–2353 (2003).

40.   Doan, A. *et al.* RB depletion is required for the continuous growth of tumors initiated by

loss of RB. *PLoS Genet* **17**, e1009941 (2021).

41.   Fernández-Martínez, P., Zahonero, C. & Sánchez-Gómez, P. DYRK1A: the double-edged

kinase as a protagonist in cell growth and tumorigenesis. *Molecular & Cellular Oncology* **2**,

e970048 (2015).

42.   Carey-Smith, S. L. *et al.* Efficacy of DYRK1A inhibitors in novel models of Down syndrome

acute lymphoblastic leukemia. *Haematologica* **109**, 2309–2315 (2024).

43.   Lindberg, M. F. *et al.* Chemical, Biochemical, Cellular, and Physiological Characterization

of Leucettinib-21, a Down Syndrome and Alzheimer's Disease Drug Candidate. *J. Med. Chem.*

66, 15648–15670 (2023).

44.   Chen, J.-Y., Lin, J.-R., Tsai, F.-C. & Meyer, T. Dosage of Dyrk1a Shifts Cells within a p21-

Cyclin D1 Signaling Map to Control the Decision to Enter the Cell Cycle. *Molecular Cell* **52**, 87–

100 (2013).

45.   Keenan, A. B. *et al.* ChEA3: transcription factor enrichment analysis by orthogonal omics

integration. *Nucleic Acids Research* **47**, W212–W224 (2019).

46.   Zhu, Y., Do, V. D., Richards, A. M. & Foo, R. What we know about cardiomyocyte

dedifferentiation. *Journal of Molecular and Cellular Cardiology* **152**, 80–91 (2021).

47.   Young, A., Bradley, L. A. & Wolf, M. J. In Vivo Methods to Monitor Cardiomyocyte

Proliferation. *JCDD* **9**, 73 (2022).

48.   Bradley, L. A., Young, A., Billcheck, H. O. & Wolf, M. J. Ablation of endogenously cycling

adult cardiomyocytes worsens myocardial function after injury. (2020)

doi:10.1101/2020.09.21.306852.

49.   Rammohan, M. *et al.* The chromosome 21 kinase DYRK1A: emerging roles in cancer

biology and potential as a therapeutic target. *Oncogene* **41**, 2003–2011 (2022).

50.   Puente, B. N. *et al.* The Oxygen-Rich Postnatal Environment Induces Cardiomyocyte Cell-

Cycle Arrest through DNA Damage Response. *Cell* **157**, 565–579 (2014).

51.   Johnson, J. *et al.* Systemic Hypoxemia Induces Cardiomyocyte Hypertrophy and Right

Ventricular Specific Induction of Proliferation. *Circulation Research* **132**, 723–740 (2023).

52.   Jiang, Y.-H. *et al.* Multinucleated polyploid cardiomyocytes undergo an enhanced

adaptability to hypoxia via mitophagy. *Journal of Molecular and Cellular Cardiology* **138**, 115–135

(2020).

53.   Zeigler, A. C., Richardson, W. J., Holmes, J. W. & Saucerman, J. J. A computational model

of cardiac fibroblast signaling predicts context-dependent drivers of myofibroblast differentiation.

Journal of Molecular and Cellular Cardiology **94**, 72–81 (2016).

54.   Tan, P. M., Buchholz, K. S., Omens, J. H., McCulloch, A. D. & Saucerman, J. J. Predictive

model identifies key network regulators of cardiomyocyte mechano-signaling. *PLoS Comput Biol*

13, e1005854 (2017).

55.   Rohatgi, A. *WebPlotDigitizer*.

56.   Stirling, D. R. *et al. CellProfiler 4: Improvements in Speed, Utility and Usability*.

http://biorxiv.org/lookup/doi/10.1101/2021.06.30.450416 (2021) doi:10.1101/2021.06.30.450416.

57.   Woo, L. A. *et al.* High-content phenotypic assay for proliferation of human iPSC-derived

cardiomyocytes identifies L-type calcium channels as targets. *Journal of Molecular and Cellular*

Cardiology **127**, 204–214 (2019).

58.   Woo, L. A. *et al.* Multi-omic analysis reveals VEGFR2, PI3K, and JNK mediate the small

molecule induction of human iPSC-derived cardiomyocyte proliferation. *iScience* **27**, 110485

(2024).

59.   Michael Love, S. A. *DESeq2*. ([object Object], 2017). doi:10.18129/B9.BIOC.DESEQ2.

60.   CompareCluster), G. Y. [Aut, Cre], Li-Gen Wang [Ctb], GiovanniDall'Olio [Ctb] (Formula

Interface   Of.   *clusterProfiler*.   ([object   Object],   2017).

doi:10.18129/B9.BIOC.CLUSTERPROFILER.