N-acetylcarnosine attenuates age-associated declines in multi-organ systems to improve survival

Edwin R. Miranda[1,2,3], Justin L. Shahtout[1,3,4], Shinya Watanabe[1,3], Jan Spaas[5], Norah Y. Milam[1,3], Grace Neiswanger[1,3], Benjamin Werbner[1,2,3], Deborah Stuart[6], Sohom Mookherjee[1,3], Jack Wilson[1], Michael Judge[1], Isabella Y. Goh[7], Tyler Slater[8], Molly R. Gallop[1,2,3], Guoli Hu[9], Takuya Karasawa[1,2,3], Jillian K. Landers[1,3], Bryann Rainbow[1,3], Kyle T. O'Connor[10], Nicholas J. Black[10], He Lan[10], Linda S. Nikolova[11], Ying Li[1,3,12], Crystal F. Davey[1,10,13], James E. Cox [3,12,14], Sihem Boudina[1,2,3,15], Courtney M. Karner[9,16], Natalia S. Harasymowicz[2,7], Nirupama Ramkumar[3,6], J. David Symons[1,2,3,16], Amandine Chaix[1,2,3], Jonathan Z. Long[5], Micah J. Drummond[1,2,3,4,15], Katsuhiko Funai[1,2,3,4,14,15].

Affiliations

[1]Department of Nutrition and Integrative Physiology, University of Utah, Salt Lake City, Utah, United States.

[2]Molecular Medicine Program, University of Utah, Salt Lake City, Utah, United States.

[3]Diabetes & Metabolism Research Center, University of Utah, Salt Lake City, Utah, United States.

[4]Department of Physical Therapy & Athletic Training, University of Utah, Salt Lake City, Utah, United States.

[5]Department of Pathology, Stanford University, Stanford, California, United States.

[6]Division of Nephrology, Department of Internal Medicine, University of Utah, Salt Lake City, Utah, United States.

[7]Department of Orthopaedic Surgery Operations, University of Utah, Salt Lake City, Utah, United States.

[8]Preclinical Imaging Core Facility, University of Utah, Salt Lake City, Utah, United States.

[9]Department of Internal Medicine, University of Texas Southwestern, Dallas, Texas, United

States.

[10]Transgenic Mouse Core Facility, University of Utah, Salt Lake City, Utah, United States.

[11]Electron Microscopy Facility, University of Utah, Salt Lake City, Utah, United States.

[12]Metabolomics Core Research Facility, University of Utah, Salt Lake City, United States.

[13]Mutation Generation & Detection Core Facility, University of Utah, Salt Lake City, United

States.

[14]Department of Biochemistry, University of Utah School of Medicine, Salt Lake City, United

States.

[15]Center on Aging, University of Utah, Salt Lake City, Utah, United States.

[16]Charles and Jane Pak Center for Mineral Metabolism and Clinical Research, University of

Texas Southwestern, Dallas, Texas, United States.

Corresponding Author:

Katsuhiko Funai

University of Utah

15N, 2030E, Salt Lake City, UT 84112, USA

Email: kfunai@health.utah.edu

Introductory Paragraph

Histidine containing dipeptides (HCDs) such as N-acetylcarnosine are endogenous metabolites that are ergogenic and mitigate metabolic dysfunction. We previously demonstrated that short-term N-acetylcarnosine treatment is highly efficacious in protecting muscle atrophy induced by disuse. Here we demonstrate that a 6-months treatment of N-acetylcarnosine attenuates a broad spectrum of age-associated maladies and improved survival by ~50% in female mice. A comprehensive survey of organ systems revealed that N-acetylcarnosine prevents decline in adiposity, diastolic function, vasodilation, muscle strength, and bone density. Together, N-acetylcarnosine substantially delays the onset of system-wide end-stage pathology to prolong lifespan. As an endogenously present metabolite, treatment with N-acetylcarnosine may be a safe and promising intervention to promote healthy aging in humans.

Main

With the aging global population, there remains an unmet need for therapies to mitigate age-related pathologies. While lifestyle modifications such as caloric restriction[1,2] and exercise[3] are effective strategies for prolonging health during aging, no equivocal pharmacologic strategies exist. Histidine containing dipeptides (HCDs) are endogenously produced metabolites primarily in excitable tissues[4] such as skeletal and cardiac muscle by conjugating histidine with beta-alanine or gamma-aminobutyric acid (GABA) in the case of homocarnosine. The supplementation of β-alanine, which is the rate limiting metabolite for HCD production, is ergogenic in humans, improving aerobic performance and muscle force production[5]. Conversely, decline in the HCD N-acetylcarnosine predicts mental and physical frailty in octogenarians[6]. The purported primary mechanism for these health benefits involves the scavenging of reactive carbonyl species, especially lipid-derived carbonyls such as 4-hydroxynonenal (4HNE) via the imidazole ring of histidine[7]. 4HNE and other reactive carbonyl species are generated via oxidation of their metabolic precursors. For example, 4HNE is produced primarily via peroxidation of ω6-polyunsaturated lipids. Formation of reactive carbonyls are accelerated in contexts of oxidative stress such as aging[8]. These reactive carbonyls non-enzymatically and irreversibly bind to proteins on nucleophilic residues incurring consequences such as impaired protein structure, function, and turnover[9,10].

HCD supplementation mitigates impairments in cardiovascular, musculoskeletal, neuronal and metabolic systems in contexts other than aging[11]. For example, supplementation with carnosine or its analog carnosinol improved glucose tolerance in mice and humans with obesity[12,13]. We recently demonstrated the ability of carnosine or N-acetylcarnosine supplementation to attenuate accumulation of 4HNE-modified proteins and preserve muscle mass and *ex vivo* force production in mice subjected to 7-days of inactivity[14]. Similarly, increasing muscle carnosine in older adults by supplementing with β-alanine improved muscle power and work

capacity[15]. Nonetheless, there has never been a study that examined the efficacy of long-term

HCD supplementation on aging *per se*. HCDs have excellent safety profiles with only mild side

effects[13,15,16], making them attractive to administer to humans over the course of aging. Herein,

we performed a 6-month preclinical N-acetylcarnosine trial in 18-24 months old mice and

performed systematic phenotyping of their organ systems to determine its effects on age-related

health decline. We chose N-acetylcarnosine primarily because we believed it would be more a

translatable therapy for humans due to its superior stability in human plasma compared to

carnosine. Unlike mice, humans ubiquitously express carnosinase (CN1) which hydrolyzes

carnosine and other HCDs limiting their stability in circulation. However, N-acetylcarnosine,

which is largely unique to humans, is not susceptible to CN1 hydrolysis.

C57Bl/6 mice from the aged rodent colony at the National Institute on Aging were acquired at 17

months of age. Following acclimation in our facility, the intervention was initiated at 18 months of

age and continued for 6 months. During the final month of the intervention, *in vivo* physiological

metrics spanning metabolic, neuro-endocrine, cardiovascular, renal, and musculoskeletal

systems were assessed. After the 6-month intervention, mice were anesthetized, and tissues

were collected for various assessments (Fig. 1a).

To confirm efficacy of our treatment, N-acetylcarnosine and other histidine-related metabolites

(Fig. 1b) were quantified in various tissues. As anticipated[4], N-acetylcarnosine was only

appreciably quantifiable in tissues from N-acetylcarnosine-treated mice (Fig. 1c, Supp. Fig. 2h).

Given the role of N-acetylcarnosine to scavenge free carbonyls and prevent protein

carbonylation, we also measured protein carbonyls in these tissues. Consistent with this notion,

abundances for protein carbonyls were lower in several tissues from N-acetylcarnosine-treated

mice compared to vehicle-treated mice (Fig. 1d, Supp. Fig. 2h). Carnosine, and the other HCDs

assayed were largely not enriched and some were even reduced by N-acetylcarnosine

treatment (Fig. 1e-h, Supp. Fig. 2h). In skeletal muscle, heart, and gonadal white adipose tissue

(gWAT), N-acetylcarnosine enrichment was greater in female mice than in male mice. This

observation is in accord with many of the sex-specific effects we observed on physiologic

functions assayed.

At the 24-month timepoint, N-acetylcarnosine increased survival by ~50% only in female mice

(Fig. 1i&j). In line with published female survival rates for this colony[17], approximately half of the

vehicle-treated female mice survived to approximately 24 months of age, while more than 75%

of N-acetylcarnosine-treated female mice survived to this age. N-acetylcarnosine did not

influence survival in male mice by the end of the study (~24-month of age) (Fig. 1j). It is

important to note that 75% survival for 24-month old male mice in this colony is in line with

published survival rates for male mice in this colony[17]. Concomitant with the increased survival,

N-acetylcarnosine promoted a retention of body weight that was also specific to female mice

(Fig. 1k&l). The effect of N-acetylcarnosine on maintaining body weight was exclusively

accounted for by its ability to retain fat mass without altering lean mass (Fig. 1m-p). Importantly,

these sex-specific effects were not explained by differences in food consumption, or exposure to

N-acetylcarnosine via water consumption (Supp. Fig 1a-f).

Short-term N-acetylcarnosine or other HCDs may mitigate metabolic dysfunction associated

with obesity[12,18-22]. Given the greater adiposity in N-acetylcarnosine-treated female mice, we first

interrogated the influence of N-acetylcarnosine on metabolic and neuroendocrine systems (Fig.

2a). Consistent with the body composition data, N-acetylcarnosine treatment promoted greater

gWAT mass without influencing inguinal white adipose tissue (iWAT) or brown adipose tissue

BAT mass in female (Fig. 2b) but not male (Supp. Fig. 1g) mice. It's noteworthy that N-

acetylcarnosine was robustly enriched in gWAT only in female mice (and not in male mice)

whereas it was equally enriched in iWAT between sexes (Fig. 1b, Supp. Fig. 2h). However,

protein carbonyls were equivocal in between groups in male and female mice (Fig. 1c, Supp.

Fig. 2h). Even though gWAT mass was greater in N-acetylcarnosine-treated female mice

compared to the vehicle, systemic glucose tolerance was not different between groups (Fig. 2d,

Supp. Fig. 1q). Liver, another metabolic organ influenced by adiposity, did not appear to be

affected in mass or histology by N-acetylcarnosine (Fig. 2c&d, Supp. Fig. 1g&h). Given the lack

of effect of N-acetylcarnosine on adipose and liver in males, it was unsurprising that there was

also no effect on their glucose tolerance (Supp. Fig. 1i&j).

We further examined the effect of N-acetylcarnosine on systemic metabolism by Indirect

calorimetry. Surprisingly, whole-body O_2 consumption and energy expenditure were consistently

reduced in the N-acetylcarnosine treatment group in both sexes (Fig. 2f, Supp. Fig. 1k&l, Supp.

Fig. 1r), despite greater body weight in N-acetylcarnosine-treated female mice (Fig. 1j).

Respiratory exchange ratio (RER), defined by CO_2 produced per O_2 consumed, is an indirect

parameter to estimate systemic substrate preference. A RER of 1.0 indicates exclusive

glycolytic oxidation, while a RER of 0.7 indicates exclusive fat oxidation. N-acetylcarnosine

treatment did not have any effect on RER in male mice (Supp. Fig. 1n). Somewhat surprisingly,

N-acetylcarnosine treatment reduced RER in female mice (Fig. 2g), normally a signature of

improved metabolic health[23]. These effects on metabolism are most likely explained by reduced

ambulatory activity that was observed in the female (Fig. 2h, Supp. Fig 1s&t) but not male

(Supp. Fig. 1o&p) N-acetylcarnosine-treated mice. Overall, N-acetylcarnosine preserved fat

mass in female mice without signs of metabolic dysfunction. In the context of aging, retention of

adiposity represents the ability of animals to maintain energy reserve which may delay frailty

and promote longevity.

For male mice, we also assessed the effect of N-acetylcarnosine on fertility by assessing sperm

count and sperm viability. Sperm count decreases dramatically between 18 and 25 months in

C57Bl/6 mice[24] and the HCD carnosine is able to preserve sperm morphology in an accelerated

aging senescent mouse model[25]. However, N-acetylcarnosine did not affect sperm count, or the

percentage of live sperm in 24-month-old mice (Supp. Fig2a).

Carnosine and other HCDs have also been implicated in improving neurologic function in aged

animal and human models[26-28]. Thus, we performed a wide variety of assessments to examine

the influence of N-acetylcarnosine treatment on the neurological system. N-acetylcarnosine

treatment did not have any effects on brain size in either sex (Fig. 2i). The behavioral

assessments also did not show differences in memory (fear conditioning and Y-maze) (Fig.

2j&k, Supp. Fig. 2b-d, Supp. Fig. 2f) or pain perception (mechanical or heat)[29] (Fig. 2l, Supp.

Fig. 2e&g) in male or female mice.

As one of the organ systems critical for survival we next profiled the influence of N-

acetylcarnosine on cardio-renal phenotypes (Fig. 3a). N-acetylcarnosine treatment did not

appear to alter relative heart weight (Fig. 3b, Supp. Fig. 3a) or histology (Fig. 3c, Supp. Fig. 3h).

There was a modest trend for lower heart rates in female mice (Fig. 3d), suggesting a potentially

positive effect of N-acetylcarnosine on heart or perhaps vagal tone. Fractional shortening (FS)

in female N-acetylcarnosine-treated hearts was lower compared to the vehicle-treated mice

(Fig. 3g) with no changes to cardiac output or ejection fraction (Fig. 3e&f). The lower FS was

driven by ~17% greater systolic left ventricular internal diameter compared to vehicle mice

(p=0.002, data not shown), which may have been related to the observed lower heart rate. N-

acetylcarnosine appeared to influence heart to a lesser extent in male mice (Supp. Fig. 3c-e),

except for improved E/e' indicative of late-diastolic ventricular filling (Supp. Fig. 3g). It is

important to point out that the N-acetylcarnosine treatment in our study started when mice were

18-months old, an age when some of the age-related decline in heart function had already

began[30]. Thus, it remains possible that the treatment may have had a more substantial effect if

initiated earlier. This is particularly the case as HCDs have been shown to improve excitation-

contraction coupling via regulating calcium handling[31]. Overall, N-acetylcarnosine

supplementation may improve some aspects of systolic function in aging female mice and

diastolic function in aging male mice.

Because female mice demonstrated substantial increase in survival, we further explored the

effect of N-acetylcarnosine on the vascular system in female mice. Acetylcholine-evoked,

endothelium-dependent vasodilation assessed using isometric myography was similar in vehicle

and N-acetylcarnosine-treated mice (Supp. Fig. 3i). However, addition of the endothelial nitric

oxide synthase (eNOS) inhibitor L-NMMA attenuated vasorelaxation to a greater extent in

vessels from N-acetylcarnosine compared to vehicle-treated mice (Fig. 3j&k), a result that was

substantiated by arteries from N-acetylcarnosine mice displaying a greater calculated nitric

oxide bioavailability compared to vehicle mice (Fig. 3l). Vascular smooth muscle responses

were similar between groups (not shown). These findings are consistent with previous work that

HCDs promote endothelium-dependent vasodilation[32,33], and might have implications

concerning systemic blood pressure.

The cardiovascular system intricately interacts with the kidney to control blood pressure and

filtration, a function that is known to decline drastically with aging. HCDs have been implicated

to be protective of the kidney in animal models of obesity[18]. Nonetheless, N-acetylcarnosine

treatment did not appear to influence kidney weight (Fig. 3m), histology (Fig. 3n), urine output

(Fig. 3o), creatinine clearance (Fig. 3p), or glomerular filtration rates (GFR) (Fig. 3q) in either

sex (Supp. Fig. 3j-p). These findings are particularly interesting given that other models of HCD

enrichment such as whole-body deletions of carnosinase 1 (CN1) demonstrate preferential HCD

enrichment in the kidney[34]. However, the GFR for both vehicle and N-acetylcarnosine treated

mice are similar to GFRs that of healthy young mice we reported recently[35] suggesting that

kidney dysfunction may not have manifested by 24 months of age.

Frailty is a hallmark of aging that very closely predicts mortality. Physical function, supported by

the musculoskeletal system (Fig. 4a), is known to drastically become impaired in the process of

aging to contribute to the decline in quality of life. Previously, we showed that N-acetylcarnosine

robustly attenuates muscle atrophy and weakness induced by 1-week of disuse in older mice[14].

However, sarcopenia occurs gradually over a longer period, so the current study provided a

perfect opportunity to determine the effects of N-acetylcarnosine treatment on loss of physical

function with aging. N-acetylcarnosine did not appear to strongly influence muscle mass or

myofiber cross-sectional area in fast-twitch extensor digitorum longus (EDL) or slow-twitch

soleus muscles in either sex (Fig. 4b&c, Supp. Fig. 4a). It is known that while C57Bl/6 mice

begin to experience atrophy at 24 months, loss of muscle mass at peaks at about 30 months of

age[36] so this perhaps was not too surprising. In contrast, muscle weakness precedes muscle

atrophy in sarcopenia[36]. Indeed, ex vivo assessment of skeletal muscle force-generating

capacity revealed consistent increase in both EDL and soleus muscle strength induced by N-

acetylcarnosine treatment in female mice (Fig. 4e&f), but not in male mice (Supp Fig. 4g&h). A

robust enrichment of N-acetylcarnosine in skeletal muscle in female mice, but not male mice

(Fig. 1c) is consistent with these observations. Nonetheless, these improvements were

insufficient to be reflected in the *in vivo* assessments of physical function (Fig. 4g-i, Supp. Fig.

4i-k). These data implicate N-acetylcarnosine as a potentially viable strategy to delay features of

sarcopenia in mice.

Aging also profoundly influences physical function by weakening bones, particularly in post-

menopausal women that experience osteoporosis. Micro-CT of femurs (Fig. 4j) revealed that N-

acetylcarnosine treatment increased trabecular number (Fig. 4k, Supp. Fig. 4l) but decreased

trabecular spacing (Fig. 4l, Supp. Fig. 4m). This in parallel with unaltered trabecular thickness

(Fig. 4m, Supp. Fig. 4n) led to increased bone volume fraction (Fig. 4n, Supp. Fig. 4o) in both

sexes in the distal femur. In contrast, neither tibial trabecular bone nor inflammation at the knee

joint were affected by N-acetylcarnosine treatment (Supp. Fig. 4p-r). These data indicate an

ability of N-acetylcarnosine to enhance bone density in the femurs of aging mice.

Discussion

Aging is associated with exacerbated oxidative stress, however strategies broadly targeting

reactive oxidative species (ROS) have been disappointing. This may be related to the presence

of many primary and secondary ROS-related species and the important roles some of these

species play in physiological signaling and hormetic response to stress. Carbonyls may be a

more promising target given they are secondary ROS species, their formation is proportional to

their metabolite precursors which may be altered in a context-dependent manner, and they are

canonically not known to possess physiological signaling roles. Supplementation with HCDs

such as N-acetylcarnosine, carnosine, their precursor β-alanine, and synthetic analogs have

established efficacy for mitigating age-related pathologies such as sarcopenia[14,15], insulin

resistance[12,21,22], cardiovascular[33,37,38] and renal dysfunction[39] primarily in contexts outside of

aging purportedly via scavenging of reactive carbonyls. Ours is the first study to simultaneously

investigate the effects of N-acetylcarnosine on these systems during aging *per se* in a

longitudinal study.

Supplementation with N-acetylcarnosine improved survival of female mice by approximately

50%. While the mechanism by which female mice preferentially benefited is unclear, certain

tissues from female N-acetylcarnosine-treated mice that displayed sex-specific effects, were

more enriched for N-acetylcarnosine (Fig. 1c). For example, N-acetylcarnosine was only

significantly elevated in gWAT from female N-acetylcarnosine-treated mice (Supp. Fig. 2h). In

parallel, we observed a disparate effect of N-acetylcarnosine to promote weight retention in

female mice which appeared to be driven by larger gWAT (Fig. 2b) without compromising

metabolic function. A finding in line with previous work demonstrating the ability of HCDs to

protect against obesity-related glucose intolerance[12,13,19]. While, the mechanism of this adipose

retention is unclear, our indirect calorimetry and activity data demonstrate decreased energy

expenditure and activity in female N-acetylcarnosine treated mice. Interestingly, recent research

suggests that decreased metabolic rate induced by a torpor-like state increases survivability and

delays aging in mice[40]. Carnosine treatment has also been demonstrated to mitigate adipocyte

senescence via scavenging of 4-HNE in aging mice[41] which could promote retention of fat mass

as we observed in the N-acetylcarnosine-treated mice.

The carbonyl scavenging ability of lipid carbonyls such as 4-HNE may have also contributed to

vascular, muscle, and bone function as well. 4-HNE uncouples NO production in endothelial

cells[42] leading to impaired NO bioavailability and carnosine has been shown to facilitate NO

production *in vitro[33]*. While this increased NO bioavailability we observed in the N-

acetylcarnosine treated mice did not enhance endothelium-dependent vasodilation, NO

bioavailability is closely associated with vascular health and appropriate blood pressure

regulation which we did not measure. Improved NO bioavailability may also protect old mice

from additional insults (e.g. ischemia reperfusion, dyslipidemia etc.) which should be tested in

future experiments. It is worth noting that N-acetylcarnosine drastically reduced protein carbonyl

burden in the heart, even though N-acetylcarnosine treated-mice did not exhibit overt

improvement in heart function discerned from echocardiography. However, heart functional

decline likely occurs earlier than 18 months which may have limited the therapeutic capacity of

N-acetylcarnosine in this study.

We previously found N-acetylcarnosine was effective at mitigating 4-HNE modified proteins in

muscle and preserved muscle force following 7-days of hindlimb unloading. Here, we

demonstrate that N- was effective at preserving *ex vivo* muscle force production during aging.

HCDs have been shown to chelate and alter the concentration of di-cations such as calcium

which may be an alternate mechanism to carbonyl sequestering that may have contributed to

this observation by altering cross bridge kinetics. This effect did not result in improved functional

metrics (i.e. treadmill, grip strength, rotarod) but these measurements mostly rely on

submaximal strength, do not necessarily reflect functional capacity relevant for trip and fall

prevention (important health factors for elderly individuals). Bone forming osteoblasts cells are

also negatively impacted by oxidative stress[43] which may have been attenuated by N-

acetylcarnosine treatment to increase femur trabecular bone density. Improved bone density is

also translationally relevant for older individuals with increased risk of bone fractures which can

dramatically increase morbidity and mortality especially in women. While the collection of these

physiologic improvements does not provide a clear explanation of why the female N-

acetylcarnosine mice improved survival by 50%, we speculate that the retention of adiposity

may have broadly delayed frailty onset which manifested with improved NO bioavailability,

muscle force capacity, and bone density.

In conclusion, N-acetylcarnosine treatment improves functions of multiple organ systems to

increase survival in female mice. As an endogenously present metabolite, N-acetylcarnosine

likely represents a safe, commercially scalable, and economically affordable treatment in

humans. The current intervention spanned 18-24 months of age, and it remains possible that N-

acetylcarnosine treatment has additional protective effects on age-related declines that occur

outside this window. Humans are known to have substantially different HCD metabolism and

pharmacokinetics than mice. Thus, N-acetylcarnosine is a strong candidate to be considered for

clinical testing to determine its efficacy in mitigating age-related health decline in humans.

Methods

Mice

All experiments were performed at the University of Utah and were approved by the Institutional

Animal Care and Use Committee. C57Bl/6 mice were ordered from the National Institutes on

Aging's aged mouse colony at 17-months of age and fed a normal chow diet, and housed with a

12-hour light/dark cycle in a temperature-controlled room for the duration of the study. Mice

were ordered in cohorts of 20 mice per months with three cohorts consisting of 10 males and 10

females, and five subsequent cohorts consisting of 20 female mice after observing certain

trends in female mice only. Upon arrival mice were weighed, counterbalanced by weight, and re-

caged to five mice per cage when possible and needed. Mice were monitored for the next three

weeks with minimum contact to allow them to acclimate to their new environment. After three

weeks of acclimation, baseline measures of body composition, grip strength, rotarod, and Y-

Maze were taken (described below).

N-acetylcarnosine Treatment

Cages of mice were randomized to receive normal drinking water (vehicle) or drinking water

supplemented with 80 mM N-acetylcarnosine prepared as previously described[14]. Briefly, tap

water was autoclaved and supplemented with N-acetylcarnosine to final concentration of 80

mM. Both vehicle and N-acetylcarnosine water were titrated to a final pH of 7.5. Water was

prepared fresh and replaced weekly. At the time water is replaced, the mice are weighed, fresh

bedding is provided, and food and water consumption are measured. The intervention begins 4

weeks after the mice have been acquired at 80 weeks of age and continued for 24 weeks. Over

the course of the intervention, mice are also monitored for general well-being and treated as

needed by the veterinarians. If deemed necessary by the veterinarians, mice were euthanized

prior to the end point. Reasons for early euthanasia included unresolving scruff wounds, anal

prolapse, enlarged tumors, and extreme weight loss. In addition to weekly routine monitoring,

mice were assessed every 8 weeks for body composition. Four weeks prior to the study

endpoint, physiological endpoint measures (body composition, grip strength, rotarod, treadmill

test, Y-maze, Comprehensive Laboratory Animal Monitoring System (CLAMS),

echocardiography, 24-hour urine collection, glomerular filtration rate, fear conditioning, Von Frey

test, hot plate test, and intraperitoneal glucose tolerance test). This timing was chosen to limit

the proximity of tests to one another to limit stressing the mice, and to prevent any resultant

stress from confounding terminal tissue collection. To limit measurement bias, the tester

performing these assessments was blinded and in the case of repeated measures, the same

tester performed the assessment at all time points.

Body Composition

Body composition was assessed at the beginning of the intervention and 8, 16, and 24 weeks of

treatment via Bruker Minispec NMR (Bruker, Germany).

Grip Strength

Whole body grip strength was assessed at baseline and between week 100 and 103 of the

intervention utilizing a grip strength meter with a mesh wire attachment (Columbus Instruments,

Columbus, OH, USA) with methods modified from[44]. Briefly mice were acclimated to the testing

room for 30 minutes. The tester then removed a mouse from the cage (blind from the group

allocation of that cage) and placed it onto the mesh wire attached to the dynamometer. Gripping

at the base of the tail, the mouse was pulled steadily, with its torso horizontal to the ground,

back towards the tester. This trial was repeated three times with 30 seconds between each bout

and the highest force being reported.

Rotarod

Rotarod assessment was performed at baseline and between week 100 and 103 of the

intervention as previously described[44]. Briefly, mice were trained on the Rotamex-5 (Columbus

Instruments, Columbus, OH) for three days prior to testing by placing mice on the apparatus at

a constant speed of 4 rpm for 5 minutes, replacing them if they were to fall. On the day of

testing, mice were acclimated to the testing room for 30 minutes prior to testing. After

acclimating, mice were placed on the apparatus and the instrument was run at 4 rpm,

accelerating 0.3 rpm/s. Mice were allowed to continue if they were able to grip the rotarod to

prevent a fall. Timer continued until the mice fell to the bottom. Once all of the mice were

assayed, the test was repeated starting with the first mouse assayed until all mice were assayed

three times with the best score being reported.

Treadmill Exercise Capacity

The treadmill test was performed between week 100 and 103 of the intervention as previously

described[30] prior to terminal experiments. Every day for the three days prior to the treadmill

exercise test, mice were acclimated to the treadmill by placing each mouse in the treadmill for 5

minutes with the belt being stationary followed by 5 minutes with the belt running at 5 meters

per minute. On the day of the test, mice were acclimated to the test room for 30 minutes. Four

mice at a time were placed on the treadmill with walls separating adjacent mice. Mice began by

running at 5 meters per minute at a 25% grade for 1 minute. The speed was increased by 1

meter per minute every minute until maximal exercise capacity was achieved. The mice were

encouraged to run by tapping the mice with a wire cleaner brush. Maximal capacity was

determined by a mouse's inability to run on the treadmill following 5 consecutive coercions with

the wire brush followed by 2 attempts to place the mouse in the middle of the treadmill by hand.

Maximal exercise capacity was calculated as follows:

$$Body\ Weight\ (kg) * Running\ Time\ (\text{minutes}) * Final\ Running\ Speed\ \left(\frac{m}{\text{minute}}\right)$$

$$* Treadmill\ Grade\ (25\%)$$

Y-Maze

Y-Maze was performed at baseline and between week 100 and 103 of the intervention. Mice

were acclimated to the testing room 30 minutes prior to the assessment. The Y-maze was

placed on the floor in the same orientation in the same room for repeated measures. A camera

was suspended above the Y-maze apparatus to record the test. The mouse was then placed in

the center for 20 seconds of the maze with three walls blocking the path to the maze arms to

acclimate to the apparatus. After acclimating to the apparatus, video recording was initiated and

the blockades to the maze arms were removed. The branches that the mouse traveled to was

recorded for 15 branches unless the mouse returned to the same arm twice in a row at which

point the test was terminated. Spontaneous alternations were calculated by counting the

number of unique groupings of three arms visited across the entire test.

CLAMS

Comprehensive laboratory animal metabolic system (CLAMS, Columbus Instruments,

Columbus, OH) was performed between week 100 and 103 of the study to measure energy

expenditure (VO_2), respiratory exchange ratio (RER; $VCO_2/\ VO_2$), food and water consumption,

and physical activity.

Echocardiography

Echocardiography was performed between week 100 and 103 of the study. Mice were induced

and maintained under light anesthesia by inhalation of isofluorane. Cardiac function was

measured using a VisualSonics Vevo 2100 ultrasound machine with MS400 probe. B- and M-

mode sequences were acquired in the parasternal long- and short-axis orientations and used to

assess systolic function and to quantify morphological features. Mitral inflow velocities (E and

A), and early diastolic mitral annulus tissue velocity (e') were measured in the apical, four-

chamber view using flow and tissue doppler respectively. These values were used to assess

diastolic function based on E/A and E/e' ratios. Echocardiographic data was analyzed using

VisualSonics VevoLab.

24-Hour Urine Collection and Creatinine Clearance

A 24-hour urine collection for determination of creatinine clearance was performed between

week 100 and 103 of the study. Mice were placed in an MMC100 Metabolic Cage for 24 h at

which point urine and cheek blood were collected. Body weight, urine volume, food, and water

intake were all observed throughout the experiment. Mice were fed with a standard chow gel

diet with free access to water. Urine samples were centrifuged at 21,000 x g for 15 min, then

aliquoted, and stored at -80 C. Urine and plasma creatinine were measured using QuantiChrom

Creatinine Assay Kit, BioAssay Systems, Hayward, CA (Cat. No. DIUR-500) according to the

manufacturer's instructions. Creatinine clearance was calculated as follows:

$$\frac{Urine\ Creatinine\ Concentration\ \left(\frac{mg}{dL}\right) * Urine\ Volume\ (mL)}{Plasma\ Creatinine\ Concentration\ (\frac{mg}{dL})}$$

Glomerular Filtration Rate

Glomerular filtration rate (GFR) was measured between week 100 and 103 of the study as

previously described[45]. Mice were injected with fluorescein isothiocyanate-senestrin

(Medibeacon, Mannheim, Germany) retro-orbitally (7.5 mg/100 g body weight). The NIC-Kidney

(Medibeacon) was used to detect fluorescence in the skin on the shaved back over 1 h. GFR

was calculated based on the kinetics of fluorescence decay as previously reported[46].

Fear Conditioning

Fear conditioning was conducted between week 100 and 103 of the study using a med-associates sound-attenuated chamber (29x25x26 cm) with no scent, plexiglass roof and front door, aluminum side walls, a plastic black wall and fitted with a top-down camera (*Context A*) (Med Associates; St. Albans, Vermont). Conditional and contextual memory were assessed on two subsequent days. On day 1 mice were acclimated to the test room for at least 1 hour. After shock calibration, an individual mouse was placed in the chamber for 3 minutes. At 120 seconds, a tone was delivered for 20 seconds. Two seconds prior to termination of the tone, a 0.3 mA foot shock was delivered lasting 2 seconds. The mouse remains in the chamber until the end of the 3 minutes. The mouse is then returned to its cage, and the chamber is cleaned with 70% ethanol before testing the next mouse. On day 2, contextual fear conditioning is tested by introducing mice to *Context A* for 3 minutes without tone or shock. The percent time freezing on day 1 and day 2 are reported as metrics of contextual memory. One hour later, mice are tested for conditional memory by being introduced to a new context which differs from *Context A* by placing curved plastic board inside the chamber, a plastic board underneath the grid floor, and the chamber is scented with vanilla extract (*Context B*). Mice are allowed to explore for 3 minutes without foot shocks but with the same tone as on the first day occurring during the last minute. The percent change (%Δ) of time freezing before and after the tone on day 1 compared to day 2 is reported as the metric for conditional memory.

Von Frey Test

The Von Frey test was performed between 100 and 103 weeks of the study. Mice were acclimated to the procedure room for 30 minutes prior to testing. Mice were placed in a small chamber with grid wire flooring. The Von Frey apparatus was fitted with three filaments of different tensions. These filaments were applied to the plantar region of the mouse's foot and

the force needed to elicit withdraw of the foot is recorded. Each test was repeated three times

with 15 minutes between each test.

Hot Plate Test

The hot plate test was performed between 100 and 103 weeks of the study. Mice were

acclimated to the procedure room for 30 minutes prior to testing. A glass enclosure was warmed

to 50 degrees Celsius and a camera was affixed to record the animal's behavior in response to

being placed in the enclosure. Change in behavior indicating pain (e.g. picking up paws, rearing

etc.) was scored and timed by 3 independent and blinded individuals.

Terminal Biological Sample Collection

On week 104, 4-hour fasted mice were anesthetized via intraperitoneal injection of 80 mg/kg

ketamine and 10 mg/kg xylazine. Hindlimb muscles (Tibialis anterior [TA], Extensor digitorus

longus [EDL], soleus, plantaris, and gastrocnemius) were quickly collected from live mice,

weighed, and either tied with sutures on distal and proximal tendons for *ex vivo force* (EDL and

soleus; described below), flash frozen in liquid nitrogen, or embedded in OCT and frozen in

liquid nitrogen-cooled isopentane for histological assessment (described below). After collecting

muscles, blood was collected via cardiac puncture followed by collecting, weighing, preservation

of the remainder of the tissues (heart, liver, kidney, inguinal white adipose tissue [iWAT],

gonadal [gWAT], brain, sperm, and bones.

Targeted Metabolomics

Plasma, urine, muscle (TA), whole brain, heart, iWAT, gWAT, kidney, and liver were processed

from frozen and assessed for N-acetylcarnosine, carnosine, histidine, anserine, and

homocarnosine concentrations. Sample preparation and metabolomics methods were based on

recently published protocols[4].

Frozen tissues were quickly weighed and immediately placed in HPLC-grade water with 10 mM

HCl (12 µL extraction solution per mg tissue) followed by homogenization in a bead mill

homogenizer at 4°C. Homogenates were centrifuged (20 min, 3000 × g, 4°C) and the resulting

supernatants were immediately diluted in a 3:1 ratio with ice-cold acetonitrile. The mixture was

vortexed for 15 s and kept on ice for 15 min. After a second centrifugation step (12 min, 20,000

× g, 4°C), samples were stored at –80°C until the day of analysis. Plasma and urine samples

(37.5% of total mixture) were combined with acetonitrile containing 1% formic acid (54%) and

HPLC-grade water (8.5%). Samples were vortexed for 15 s and centrifuged (15 min, 15,000 × g,

4°C), after which supernatant was collected and stored at –80°C until the day of analysis.

Targeted metabolite measurements were performed on a triple quadrupole LC-MS instrument

(Agilent 6470). Normal phase chromatography was used to separate polar metabolites using a

Luna 5µm NH_2 100 Å LC column (Phenomenex 00G-4378-E0) equipped with a SecurityGuard

cartridge (KJ0-4282, Phenomenex). Injection volume was 7 µL, except for urine (4 µL). Mobile

phases consisted of Buffer A (water with 10 mM ammonium formate) and Buffer B (95:5

acetonitrile:water with 10 mM ammonium formate). Each run started 100% B for 2 minutes,

followed by a gradient starting at 100% B changing linearly to 50% A and 50% B over the course

of 10 minutes, followed by 50% A and 50% B for 3 minutes, and re-equilibration at 100% B for 2

min (flow rate 0.7 mL/min). MS analysis was performed using AJS in positive mode with the

following parameters: dry gas temperature 250 °C, gas flow 12 L/min, nebulizer pressure 25 psi,

sheath gas temperature 300 °C, the sheath gas flow 12 L/min, capillary voltage 3,500 V.

Multiple reaction monitoring was performed to detect N-acetylcarnosine (269.1 → 110.1 *m/z*),

carnosine (227.1 → 110.1 *m/z*), anserine (241.1 → 109.1 *m/z*), homocarnosine (241.1 → 110.1

m/z), histidine (156.1 → 110.1 *m/z*). Secondary transitions[4] were used as confirmation of the

peak identity. Fragmentor voltage for the various metabolites was 80-100, collision energy 10-24

V. Endogenous metabolite concentrations were quantified using an external standard curve with

known concentrations analyzed alongside the biological samples in the same run, except for

homocarnosine. Total peak areas were used to calculate metabolite concentrations within each

sample.

Protein Carbonyl Assay

Protein carbonyls were measured via DNPH derivatization of proteins from tissues lysed in 50

mM MES buffer pH 7.4, 50 mM EDTA via protein carbonyl colorimetric assay kit (10005020,

Caymen Chemical, Ann Arbor, MI).

Ex Vivo **Muscle Force**

Force generating capacity of both the soleus and EDL were measured as previously

described[14,47-49]. Briefly, soleus and EDL were tied with suture at proximal and distal tendons

then attached to an anchor and force transducer in a tissue bath (Aurora Scientific, Model 801C)

while being submerged in oxygenated Krebs-Henseleit Buffer solution at 37° C. The optimal

length of the muscle was determined via maximum twitch force production. The buffer was then

replaced with a fresh buffer solution and equilibrated for 5 minutes. After equilibration, a force-

frequency sequence stimulated the muscle at increasing frequencies (10, 20, 30, 40, 60, 80,

100, 125, 150, and 200 Hz) with a 2-minute rest interval between each frequency. The rates of

contraction and relaxation were quantified as previously described[14,47,48]. The Aurora Scientific

DMAv5.321 software was utilized for analysis of force production data.

Muscle Histology and CSA

OCT embedded soleus and EDL samples were mounted on a precooled platform and sectioned

at 10 μm thickness with a cryostat (Microtome Plus). 6-8 serial sections from each sample were

transferred onto the same slide. Sections then underwent blocking for 1 hour at room

temperature with M.O.M. mouse IgG blocking reagent (Vector Laboratories, MKB-2213), after

which they were incubated overnight at 4° C with a concentrated primary antibody targeting

laminin (1:200, Sigma, L9393) in 2.5% horse serum in PBS. The next day, sections were

incubated for 1 hour at room temperature with anti-rabbit IgG conjugated to Alexa Fluor 647

(1:250, Invitrogen, A21245). Sections were then washed three times in PBS, fixed with methanol

for 5 minutes at room temperature and then washed 2 more times in PBS. Finally, slides were

mounted with mounting medium (Vector 136 Laboratories, H-1000). Slides were imaged on a

Zeiss Axioscan.Z1 at 20X magnification. Masks for fibers were generated via Cellpose (V3.0)[50]

and then imported into ImageJ software where CSA was quantified.

Tissue H&E and Trichrome Staining

Portions of tissues (e.g. liver, heart) or antiliteral tissues (e.g. kidney) that were not flash frozen

in liquid nitrogen, were taken from each mouse and immediately submerged in 4%

paraformaldehyde for 12 hours and then 70% ethanol for 48 hours. Tissues were sectioned

were then embedded in paraffin, sectioned at 10-µm thickness, stained for Masson's Trichrome

to assess fibrosis or hematoxylin and eosin (H&E) to determine fat droplet accumulation and

general morphological features. Samples were imaged on Axio Scan Z.1 (Zeiss).

Ex Vivo **Vascular Function**

Vasocontractile responses to potassium chloride (KCl, 20–100 mM) and phenylephrine (PE,

10^{-8}-10^{-5} M; not shown), and vasorelaxation responses to acetylcholine (10^{-8}-10^{-5} M] and

sodium nitroprusside (10^{-9}-10^{-4} M; not shown), were completed using isometric myography as

we have described[51]. To determine the contribution to vasorelaxation from endothelial nitric

oxide synthase (eNOS), a second Ach dose-response curve was completed (30-min later) in the

presence of the eNOS inhibitor NG methyl L arginine acetate salt (L-NMMA; 1mM]. To calculate

NO bioavailability for each group, maximal vasorelaxation to Ach in the presence of L-NMMA

was subtracted from maximal vasorelaxation to Ach in the absence of L-NMMA, and all values

were averaged, and compared between groups. For all responses, values from two femoral

artery segments per mouse were averaged.

Knee Joint Histology and Scoring

To evaluate the effects of N-acetylcarnosine on OA development in aged mice right hind limbs

were collected and fixed in 10% formalin. Bone microstructure analysis was performed as stated

below. Following bone microarchitecture analysis by micro-CT joints were decalcified,

dehydrated, and embedded in paraffin. The joint was sectioned at a thickness of 8 μm. To

determine OA severity, joint sections were stained with hematoxylin, fast green, and Safranin-O.

Three independent, blind graders scored joint sections for OA using a modified Mankin scoring

system (Glasson, 2007). Scores were averaged among graders for the whole joint.

Distal Femur Micro-Computed Tomography

Femur bone structure and morphological changes were measured using a Siemens Inveon®

Multi-Modality microPET/SPECT/CT machine using the following imaging parameters: an X-ray

tube potential of 80 kVp, an X-ray tube current of 100 μA, an integration time of 2600 ms, and

an isotropic voxel size of 10 μm. Image analysis was performed using Inveon® Research

Workplace (IRW) 4.0. A 100-slice volume of interest (VOI) was selected for the cortical bone,

centered about the mid-diaphysis. Cortical bone was delineated from the medullary cavity using

a fixed threshold that remained constant across specimens. A second 100-slice VOI was

selected for the trabecular bone, beginning just proximal to the distal epiphyseal plate and

extending into the metaphysis. Irregular anatomical contours were drawn slice-by-slice

throughout this VOI just within the endocortical boundary as described by Bouxsein and

colleagues[52]. Trabecular bone was delineated from the medullary cavity using a fixed threshold

that remained constant across specimens. IRW's *Bone Morphometry* tool was used to compute

bone volume fraction, trabecular spacing, trabecular thickness, and trabecular number. The

Annotation and Measurement tool was used to estimate the cortical thickness from four

measurements at the mid-diaphysis (corresponding to the major and minor axes of an ellipse

approximating the cross-section of the femur).

Proximal Tibia Micro Computed Tomography

Tibial bone structure and morphological changes were measured in the intact hind limbs of mice

by micro-CT (PerkinElmer Quantum GX2 microCT) with a 5.5-µm isotropic voxel resolution.

Subchondral/trabecular regions were segmented using CTAn automatic thresholding software.

The tibial and femoral epiphyses were selected using the subchondral plate and growth plate as

references. The tibial metaphysis was defined as the 300 µm region directly below the growth

plate. Micro-CT limb images were analyzed for bone volume/total volume (BV/TV), trabecular

bone number (Tb.N), trabecular bone thickness (Tb.Th), trabecular bone separation (Tb.Sp)[53].

Statistics

Significance for all statistical comparisons was set at $p<0.05$. Data were tested for normal

distribution via visual inspection of QQ plots for linearity. Student's T test or Wilcoxon Signed

Rank test were used for comparison of two groups depending on normality of data. Repeated

measurements, two-way ANOVAs or if some repeated measures were missing, mixed effects

two-way ANOVAs were utilized to test group by time interactions with Bonferroni Post Hoc

analyses applied where appropriate. Individual sample values as well as estimates of central

tendency (mean or media where indicated) are presented in all bar graphs. Standard error of

the mean is reported as the variance metric. Z-scores for radar plots in Fig. 1 were calculated by

considering the distribution of all samples regardless of sex or group. Statistical analyses

including two-way ANOVA, mean, and SEM for these data are presented in Supp. Fig. 2h.

Data Availability

Raw data will be available with reasonable request to the corresponding author.

Ethics Declaration

The authors declare no competing interests.

Author Contributions

ERM conceived of and designed the study, conducted experiments, performed data analysis, and wrote and edited the manuscript. KF conceived of and designed the study, analyzed data, and wrote and edited the manuscript. MJD and AC conceived of and designed the study, analyzed data, and edited the manuscript. JLS, SW, JS, SM, IYG, TS, MRG, SB, NSH, JDS, and JZL conducted experiments, analyzed data, and edited the manuscript. NYM, GN, BW, DS, JW, MJ, GH, TK, JKL, BR, KTO, NJB, HL, LSN, YL, CFD, JEC, CMK, NR conducted experiments and analyzed data.

Acknowledgements

This work was supported by NIH grants DK110858 (JEC); HL167866, DK128819 (SB); AR083904, AR071967 (CMK); DK133271 (NR); NS137105 (JDS); AG065993, CA272529, CA286584 (AC); DK130641, MH133313, DK136526, OD034455, DK124265 (JZL); AG074535, AG076075 (MJD); AG074535, DK127979, DK107397, GM144613 (KF); HL007576 (ERM); DK137475 (MRG); AHA predoctoral grant AHA1025910, University of Utah Graduate Research Fellowship (SM); and the grant-in-aid for Japan Society for Promotion of Science (JSPS) Fellows 24KJ2039 (TK). Figures were edited by Nikita Abraham. This work was also supported by the following University of Utah Core Facilities: Cell Imaging Core, Preclinical Imaging Core, and Metabolic Phenotyping Core.

Figure Legends

Figure 1 – N-acetylcarnosine supplementation preferentially improves female mouse survival.

a) Schematic of the preclinical trial design. C57Bl/6 mice from the NIA aging colony were acquired at 17 months of age. After acclimation, mice underwent baseline testing before being randomized to receive either vehicle or N-acetylcarnosine-supplemented water for 6 months. Mice were weekly monitored for body weight and food and water consumption. Various physiologic tests were conducted during the 5th month of the intervention. Mice were euthanized and tissues were harvested after 6 months of intervention period. **b)** Schematic representation of HCD metabolism. Radar plots of **c)** N-acetylcarnosine, **d)** protein carbonyls, **e)** Carnosine, **f)** Histidine, **g)** Anserine, and **h)** Homocarnosine demonstrate enrichment of N-acetylcarnosine. Values in radar plots are Z-scores. Corresponding concentrations and statistical analyses are in **Supplementary Figure 2h**. Vehicle treated female **(i)** and male **(j)** mice experienced typical survival rates (approximately 50% and 75% respectively) at 24 months. Female mice (starting n=45 vehicle, n=46 N-acetylcarnosine) treated with N-acetylcarnosine improved their survival probability by ~50% Body weights in the female (starting n=39 vehicle, n=40 N-acetylcarnosine) **(k)** and male (starting n=20 per group) **(n)** mice fell precipitously over the course of the trial regardless of treatment. Female mice treated with N-acetylcarnosine were able to delay the decrease in bodyweight. Bodyweights are reported as one-month averages and are presented as Mean ± SEM. N-acetylcarnosine did not affect lean mass in female **(l)** or male **(o)** mice. N-acetylcarnosine preserved fat mass in female mice **(m)** (n=57 per group) but not male **(p)** (n=20 per group) mice. Statistical testing for HCD enrichment was performed via two-way repeated measures ANOVA with Bonferroni post-hoc test where appropriate. Statistical testing for bodyweight and body composition comparisons was performed via one-way repeated measures ANOVA with Bonferroni post-hoc test where appropriate. Comparison of survival curves was performed via Log-rank (Mantel-Cox) test. Comparisons were considered statistically significant if $p<0.05$. HCD; Histidine Containing Dipeptide, GABA; gamma-aminobutyric acid, CN1; carnosinase 1, CARNS; carnosine synthase. Schematic images were generated via BioRender.

Figure 2 – Effects of 6-Months of N-acetylcarnosine supplementation on the Metabolic and Neuroendocrine system in female mice.

a) Visual representation of measures assessed related to adiposity, metabolism and neurophysiology. **b)** Preservation of fat mass in female mice was attributed to larger gWAT not iWAT or BAT fat pad (n=16 vehicle, n=20 N-acetylcarnosine). Despite increased adiposity, liver weight **(c)** was not altered by N-acetylcarnosine (n=15 vehicle, n=20 N-acetylcarnosine). **d)** H&E and trichrome stains did not indicate altered steatosis or fibrosis in livers. **e)** Glucose tolerance via intraperitoneal glucose tolerance test (IPGTT) was not different between treatment and vehicle mice (n=6 vehicle, n=9 N-acetylcarnosine). 72-hour indirect calorimetry revealed suppressed energy expenditure (VO$_2$) **(f)** and respiratory exchange ratio (RER) **(g)** during both light and dark phases (n=5 vehicle, n=8 N-acetylcarnosine). Respiratory exchange ratio is the volume of CO_2 consumed (VCO$_2$) proportional to the O_2 consumed (VO$_2$) and indicates substrate utilization whereby a value closer to 0.75 indicates complete reliance on fat oxidation and a value closer to 1 indicates complete reliance on glucose oxidation. Ambulatory frequency was also suppressed in N-acetylcarnosine mice **(h)** (n=5 vehicle, n=8 N-acetylcarnosine). **i)** Brain weights were not different between treatment groups (n=14 vehicle, n=19 N-

acetylcarnosine). **j)** Change in time spent freezing before and after an audible tone that proceeded with a shock on the first (condition day) but not the second (follow up) day was not different between groups indicating no effect of N-acetylcarnosine on conditional memory formation (n=5 vehicle, n=9 N-acetylcarnosine). **k)** Frequency of novel, non-consecutive exploration (alterations) between arms in the Y-maze was not different between groups indicating no effect of N-acetylcarnosine on short-term memory (n=20/14 pre/post vehicle, n=20/16 pre/post N-acetylcarnosine). **l)** Peripheral pain tolerance assessed via Von Frey hair filaments were not different between groups (n=5 vehicle, n=9 N-acetylcarnosine). Data are represented as Mean ± SEM. Statistical significance was assessed via two-way repeated measures ANOVA with Bonferroni correction, or unpaired t-test where appropriate. Comparisons were considered statistically significant if $p<0.05$. gonadal white adipose tissue; gWAT, inguinal white adipose tissue; iWAT, BAT; Brown Adipose Tissue, CLAMS; comprehensive lab animal monitoring system. Schematic images were generated via BioRender.

Figure 3 – Effects of 6-Months of N-acetylcarnosine supplementation on the cardiovascular and renal systems in female mice.

a) Visual representation of systems assessed related to the cardiovascular-renal axis including assessment of heart size, echocardiography, *ex vivo* vascular function, and kidney function. **b)** Heart weight normalized to tibial length was not different between groups of female mice. **c)** H&E staining of heart tissues was also indiscernible between groups. **d)** Heart rate measured via electrocardiogram was performed on anesthetized mice during echocardiography and was not different between groups. Metrics of systolic function including cardiac output **(e)** and ejection fraction **(f)** were not different between groups whereas fractional shortening was reduced by N-acetylcarnosine treatment **(g)**. Measures of diastolic function including E/A (ventricular filling) **(h)** and E/e' (marker of left ventricular filling pressures) **(i)** were not different between groups. Isolated femoral arteries (n=8 vehicle, n=15 N-acetylcarnosine, 2 segments per mouse) were pre-contracted to 65% of maximal developed tension, and vasorelaxation to cumulative doses of Ach was recorded as percent (%) vasorelaxation. No differences were observed between groups. A second Ach dose-response curve was completed (30-min later) in arteries from both groups that incubated with LNMMA for 30-min. While L-NMMA attenuated vasorelaxation in arteries from vehicle **(j)** and N-acetylcarnosine **(k)** -treated mice, the contribution from NO to vasorelaxation was greater in the latter group, an observation that was substantiated by calculating NO bioavailability **(l)**. Kidney weight was not different between groups **(m)**. H&E and Trichrome stains of kidney sections were unremarkable and indistinguishable between groups **(n)**. Mice were placed in single housed, urine collection chambers for 24 hours to collect 24-hour urine volume which was not different between groups **(o)**. 24-hour urine and plasma and urine creatinine concentration were used to determine creatinine clearance **(p)** which also was not affected by N-acetylcarnosine. Glomerular filtration determined by monitoring the filtration of a FIT-c labeled senestrin under anesthesia was also not different between groups in female mice (n=5 vehicle, n=7 N-acetylcarnosine) **(q)**. Data are represented as Mean ± SEM. Statistical significance was assessed via two-way repeated measures ANOVA with Bonferroni correction, or unpaired t-test where appropriate. Comparisons were considered statistically significant if $p<0.05$. Heart Rate; HR. Schematic images were generated via BioRender.

Figure 4 – Effects of N-acetylcarnosine on the musculoskeletal system in female mice.

a) Visual representation of systems assessed to probe physical function and musculoskeletal system. **b)** Lower hind limb weights were not different between groups (n=23-24 vehicle, n=35-37 N-acetylcarnosine). **c)** Histogram of the frequency of fibers sizes was not different between groups in EDL (n=6 vehicle, n=8 N-acetylcarnosine). **d)** Representative image of immunofluorescent stain for laminin used to determine muscle cross sectional area in the EDL. E*x vivo specific* force measured in EDL **(e)** (n=8 vehicle, n=8 N-acetylcarnosine) and SOL **(f)** (n=8 vehicle, n=9 N-acetylcarnosine) were elevated in N-acetylcarnosine-treated mice across stimulation frequencies ranging from 10-200 Hz. *In vivo* functional data as quantified by rotarod **(g),** grip strength **(h)**, and treadmill work capacity **(i)** were unaffected by N-acetylcarnosine treatment. **j)** Representative µCT image of femur analyzed for bone density in the distal femur. µCT of femurs revealed increased trabecular number **(k)** and decreased trabecular spacing **(l)** without a change in trabecular thickness **(m)** which contributed to a near-significant increase in the proportion of bone volume **(n)** in the ROI. Data are represented as Mean ± SEM. Statistical significance was assessed via two-way repeated measures ANOVA or two-way ANOVA with Bonferroni correction, or unpaired t-test where appropriate. Comparisons were considered statistically significant if $p<0.05$. Tibialis anterior; TA, plantaris; Plant, gastrocnemius; Gastroc, extensor digitorum longus; EDL, soleus; SOL, cross sectional area; CSA, micro computed tomography; µCT. Schematic images were generated via BioRender.

Supplementary Figure 1 – Effect of N-acetylcarnosine on male and female metabolism.

Free-living female food **(a)** and water **(b)** and male food **(c)** and water **(d)** were not different on a per cage basis. Data are represented as mean ± SEM of the grams of consumption per mouse per day. Food **(e)** and water **(f)** consumption was also determined during the CLAMS (n=8 vehicle, n=8 N-acetylcarnosine) assessment of single-housed mice. Food consumption was only significantly increased in male mice during the dark phase. **g)** Male iWAT, gWAT, and liver weights were not different between groups. **h)** H&E and Trichrome stains of liver sections in male mice were not remarkably different between groups demonstrating limited lipid infiltration and fibrosis in both groups. **i)** 2-hour IPGTT revealed no differences in plasma glucose in male mice at any time point or as represented as area under the curve (AUC) **(j)**. **k)** CLAMS revealed a trend for lower hourly energy expenditure (relative VO_2), and a significantly lower energy expenditure AUC in the N-acetylcarnosine treated male mice in the light and dark time points **(l)**. **m)** Hourly ambulation and RER **(n)** in male mice was not different between groups during CLAMS. Ambulation AUC was significantly higher in the male, N-acetylcarnosine treated mice **(o)** in light and dark time points. **p)** Motion in the Z-direction was not affected by N-acetylcarnosine in the male mice. **q)** Female glucose AUC during 2-hour IPGTT was not different between groups. **r)** Female energy expenditure (relative VO_2) AUC was lower in N-acetylcarnosine treated mice. **s)** Total ambulation was lower in light and dark time points in female N-acetylcarnosine treated mice. **t)** Z-motion frequency was not different between female vehicle and N-acetylcarnosine-treated mice. Two-way RMANOVA or T-tests were used for statistical comparison with Bonferroni post-hoc test where appropriate. Comparisons were considered statistically significant if $p<0.05$.

Supplementary Figure 2 – Effects of N-acetylcarnosine on male reproductive health and male and female neurophysiology.

Sperm collected from the epididymis of male mice were collected and a sperm count was performed at the time of euthanasia. **a)** N-acetylcarnosine did not affect total, dead, or alive sperm counts. **b)** Brain weights, and cognition as assessed by contextual and conditional fear

conditioning **(c and d)** were not affected by N-acetylcarnosine in male mice. **e)** Peripheral pain
perception as assess by Von Frey assessment were not affected by N-acetylcarnosine
treatment in male mice. Like male mice, contextual fear conditioning **(f)** was not affected by N-
acetylcarnosine. **g)** Hot plate assay like Von Frey indicated no effect of N-acetylcarnosine in
peripheral nociception in female mice. **h)** Table of HCD and carbonyl concentrations across
tissues. Details are provided in the table legend within the figure. Data are presented as mean ±
SEM and assessed by ttest or 2-way RMANOVA with Bonferroni correction applied for post hoc
testing where appropriate. Comparisons were considered statistically significant if *p*<0.05.

Supplementary Figure 3 – Effect of N-acetylcarnosine on cardiovascular-renal systems in
male mice and endothelium dependent vasorelaxation in female femoral arteries.

a) Heart weight relative to tibia length was not different between groups in male mice.
Echocardiographic assessment (n=12 vehicle, n=13 N-acetylcarnosine) of mouse systolic
function **(b-e)** was also not affected by N-acetylcarnosine in male mice. Early stage **(f)** left
ventricular filling was also found not to be different but late diastolic filling **(g)** was greater in N-
acetylcarnosine-treated male mice indicating some mild improvements in diastolic function in
male mice. **h)** Heart histology as assessed by H&E staining was morphologically similar
between groups. **i)** Acetylcholine-evoked, endothelium-dependent vasorelaxation of femoral
arteries was not impacted by N-acetylcarnosine in female mice. **j)** Kidney weight relative to body
mass was not different in male mice. 24-hour urine collection was conducted in a modified
single mouse cage. **k)** Urine volume trended higher but was not significant in N-acetylcarnosine-
treated male mice whereas plasma creatinine concentrations **(l)** trended lower and creatinine
clearance **(m)** was equivocal between groups in male mice. **n)** GFR was determined in a subset
of male mice (n=6 vehicle, n=6 N-acetylcarnosine) by measuring the excretion of FIT-c labeled
senestrin into the urine and was not different between groups. Kidney H&E **(o)** and trichrome **(p)**
did not reveal obvious differences in kidney morphology or fibrosis between groups. Data are
presented as mean ± SEM and assessed by unpaired t-test whereby comparisons were
considered statistically significant if *p*<0.05.

Supplementary Figure 4 – Effects of N-Acetylcarnosine on musculoskeletal system in
female and male mice.

a) CSA frequency and **(b)** representative image of soleus cross section from female mice
demonstrate the frequency of soleus muscle fibers trended smaller in N-acetylcarnosine treated
female mice (n=7 vehicle, n=6 N-acetylcarnosine). **c)** Hindlimb muscle weights relative to body
weight are not affected by N-acetylcarnosine in male mice. **d)** Representative image of soleus
and EDL muscles cryosectioned and stained for laminin for determination of muscle fiber cross
sectional area (CSA). **e)** Frequency of larger fibers is increased in EDL muscles from N-
acetylcarnosine treated male mice whereas no such effect was found in soleus **(f)**. *Ex vivo*
specific force assessed in EDL **(g)** and soleus **(h)** muscles were also not different between
groups in male mice. Despite greater CSA in EDL, rotarod **(i)**, grip strength of all four limbs **(j)**,
and treadmill work capacity **(k)** were not affected by N-acetylcarnosine treatment. Micro-CT
analysis of femur bones demonstrated increased trabecular number **(l)**, decreased trabecular
spacing **(m)**, no change in trabecular thickness **(n)**, and increased bone volume fraction **(o)**
collectively indicating greater bone density in male N-acetylcarnosine treated mice. **p)**
Representative images of safranin-O-stained knee joints subjected to Mankin scoring (n=10
vehicle, n=17 N-acetylcarnosine, not significant, data not shown). Representative µCT images
of the whole knee joint **(q)** and the tibial epiphysis **(r)**. Trabecular number, spacing, thickness**,**

and bone volume were not different between groups by µCT. Data are presented as mean ±
SEM and assessed by t-test or 2-way RMANOVA with Bonferroni correction applied for post hoc
testing where appropriate. Comparisons were considered statistically significant if $p<0.05$.

References

Mattison, J. A. *et al.* Caloric restriction improves health and survival of rhesus monkeys. *Nat Commun* **8**, 14063, doi:10.1038/ncomms14063 (2017).

Colman, R. J. *et al.* Caloric restriction delays disease onset and mortality in rhesus monkeys. *Science* **325**, 201-204, doi:10.1126/science.1173635 (2009).

Ruegsegger, G. N. & Booth, F. W. Health Benefits of Exercise. *Cold Spring Harb Perspect Med* **8**, doi:10.1101/cshperspect.a029694 (2018).

Van der Stede, T. *et al.* Extensive profiling of histidine-containing dipeptides reveals species- and tissue-specific distribution and metabolism in mice, rats, and humans. *Acta Physiol (Oxf)* **239**, e14020, doi:10.1111/apha.14020 (2023).

Saunders, B. *et al.* beta-alanine supplementation to improve exercise capacity and performance: a systematic review and meta-analysis. *Br J Sports Med* **51**, 658-669, doi:10.1136/bjsports-2016-096396 (2017).

Kameda, M., Teruya, T., Yanagida, M. & Kondoh, H. Frailty markers comprise blood metabolites involved in antioxidation, cognition, and mobility. *Proc Natl Acad Sci U S A* **117**, 9483-9489, doi:10.1073/pnas.1920795117 (2020).

Aldini, G., Carini, M., Beretta, G., Bradamante, S. & Facino, R. M. Carnosine is a quencher of 4-hydroxy-nonenal: through what mechanism of reaction? *Biochemical and Biophysical Research Communications* **298**, 699-706, doi:10.1016/S0006-291X(02)02545-7 (2002).

Czyzowska, A. *et al.* Elevated phospholipid hydroperoxide glutathione peroxidase (GPX4) expression modulates oxylipin formation and inhibits age-related skeletal muscle atrophy and weakness. *Redox Biol* **64**, 102761, doi:10.1016/j.redox.2023.102761 (2023).

Fritz, K. S. *et al.* 4-Hydroxynonenal Inhibits SIRT3 via Thiol-Specific Modification. *Chem Res Toxicol* **24**, 651-662, doi:10.1021/tx100355a (2011).

Coliva, G., Duarte, S., Perez-Sala, D. & Fedorova, M. Impact of inhibition of the autophagy-lysosomal pathway on biomolecules carbonylation and proteome regulation in rat cardiac cells. *Redox Biol* **23**, 101123, doi:10.1016/j.redox.2019.101123 (2019).

Wang, Q. *et al.* The impact of carnosine on biological ageing - A geroscience approach. *Maturitas* **189**, 108091, doi:10.1016/j.maturitas.2024.108091 (2024).

Anderson, E. J. *et al.* A carnosine analog mitigates metabolic disorders of obesity by reducing carbonyl stress. *J Clin Invest* **128**, 5280-5293, doi:10.1172/JCI94307 (2018).

de Courten, B. *et al.* Effects of carnosine supplementation on glucose metabolism: Pilot clinical trial. *Obesity (Silver Spring)* **24**, 1027-1034, doi:10.1002/oby.21434 (2016).

Eshima, H. *et al.* Lipid hydroperoxides promote sarcopenia through carbonyl stress. *eLife* **12**, e85289, doi:10.7554/eLife.85289 (2023).

del Favero, S. *et al.* Beta-alanine (Carnosyn™) supplementation in elderly subjects (60-80 years): effects on muscle carnosine content and physical capacity. *Amino Acids*, 49-56, doi:10.1007/s00726-011-1190-x (2012).

Baba, S. P. *et al.* Evaluation of supplementary carnosine accumulation and distribution: an initial analysis of participants in the Nucleophilic Defense Against PM Toxicity (NEAT) clinical trial. *Amino Acids* **56**, 55, doi:10.1007/s00726-024-03414-5 (2024).

Turturro, A. *et al.* Growth curves and survival characteristics of the animals used in the Biomarkers of Aging Program. *J Gerontol A Biol Sci Med Sci* **54**, B492-501, doi:10.1093/gerona/54.11.b492 (1999).

Aldini, G. *et al.* The carbonyl scavenger carnosine ameliorates dyslipidaemia and renal function in Zucker obese rats. *J Cell Mol Med* **15**, 1339-1354, doi:10.1111/j.1582-4934.2010.01101.x (2011).

Albrecht, T. *et al.* Carnosine Attenuates the Development of both Type 2 Diabetes and Diabetic Nephropathy in BTBR ob/ob Mice. *Sci Rep* **7**, doi:10.1038/srep44492 (2017).

Haus, J. M. & Thyfault, J. P. Therapeutic potential of carbonyl-scavenging carnosine derivative in metabolic disorders. *J Clin Invest* **128**, 5198-5200, doi:10.1172/JCI124304 (2018).

Matthews, J. J., Turner, M. D., Santos, L., Elliott-Sale, K. J. & Sale, C. Carnosine increases insulin-stimulated glucose uptake and reduces methylglyoxal-modified proteins in type-2 diabetic human skeletal muscle cells. *Amino Acids*, doi:10.1007/s00726-022-03230-9 (2023).

Hariharan, R. *et al.* Carnosine supplementation improves glucose control in adults with pre-diabetes and type 2 diabetes: A randomised controlled trial. *Nutr Metab Cardiovasc Dis* **34**, 485-496, doi:10.1016/j.numecd.2023.10.012 (2024).

Shoemaker, M. E. *et al.* Differences in muscle energy metabolism and metabolic flexibility between sarcopenic and nonsarcopenic older adults. *J Cachexia Sarcopenia Muscle* **13**, 1224-1237, doi:10.1002/jcsm.12932 (2022).

Fabricant, J. D. & Parkening, T. A. Sperm morphology and cytogenetic studies in ageing C57BL/6 mice. *J Reprod Fertil* **66**, 485-489, doi:10.1530/jrf.0.0660485 (1982).

Gopko, A. V., Kulibin, A. Y., Semenova, M. L. & Zakhidov, S. T. Carnosine-induced changes in the development of spermatogenic epithelial cells in senescence accelerated SAM mice. *Bull Exp Biol Med* **140**, 235-237, doi:10.1007/s10517-005-0454-4 (2005).

Rokicki, J. *et al.* Daily Carnosine and Anserine Supplementation Alters Verbal Episodic Memory and Resting State Network Connectivity in Healthy Elderly Adults. *Front Aging Neurosci* **7**, 219, doi:10.3389/fnagi.2015.00219 (2015).

Ding, Q. *et al.* Anserine/Carnosine Supplementation Preserves Blood Flow in the Prefrontal Brain of Elderly People Carrying APOE e4. *Aging Dis* **9**, 334-345, doi:10.14336/AD.2017.0809 (2018).

Banerjee, S., Mukherjee, B., Poddar, M. K. & Dunbar, G. L. Carnosine improves aging-induced cognitive impairment and brain regional neurodegeneration in relation to the neuropathological alterations in the secondary structure of amyloid beta (Abeta). *J Neurochem* **158**, 710-723, doi:10.1111/jnc.15357 (2021).

Novak, J. C. *et al.* Aging and neuropathic pain. *Brain Research* **833**, 308-310, doi:https://doi.org/10.1016/S0006-8993(99)01522-X (1999).

Cho, J. M. *et al.* Late-in-life treadmill training rejuvenates autophagy, protein aggregate clearance, and function in mouse hearts. *Aging Cell* **20**, e13467, doi:10.1111/acel.13467 (2021).

Creighton, J. V. *et al.* Physiological Roles of Carnosine in Myocardial Function and Health. *Adv Nutr* **13**, 1914-1929, doi:10.1093/advances/nmac059 (2022).

Ririe, D. G., Roberts, P. R., Shouse, M. N. & Zaloga, G. P. Vasodilatory actions of the dietary peptide carnosine. *Nutrition* **16**, 168-172, doi:https://doi.org/10.1016/S0899-9007(99)00268-3 (2000).

Takahashi, S., Nakashima, Y. & Toda, K. Carnosine facilitates nitric oxide production in endothelial f-2 cells. *Biol Pharm Bull* **32**, 1836-1839, doi:10.1248/bpb.32.1836 (2009).

Weigand, T. *et al.* A Global Cndp1-Knock-Out Selectively Increases Renal Carnosine and Anserine Concentrations in an Age- and Gender-Specific Manner in Mice. *Int J Mol Sci* **21**, doi:10.3390/ijms21144887 (2020).

Opurum, P. C. *et al.* Combined sedentarism and high-fat diet induce early signs of kidney injury in C57BL/6J mice. *Am J Physiol Renal Physiol* **328**, F850-F860, doi:10.1152/ajprenal.00259.2024 (2025).

Kerr, H. L. *et al.* Mouse sarcopenia model reveals sex- and age-specific differences in phenotypic and molecular characteristics. *J Clin Invest* **134**, doi:10.1172/JCI172890 (2024).

37   Hipkiss, A. R., Preston, J. E., Himswoth, D. T. M., Worthington, V. C. & Abbot, N. J.
Protective effects of carnosine against malondialdehyde-induced toxicity towards
cultured rat brain endothelial cells. *Neuroscience Letters* **238**, 135-138,
doi:10.1016/S0304-3940(97)00873-2 (1997).
38   Wetzel, C. *et al.* Histidine containing dipeptides protect epithelial and endothelial cell
barriers from methylglyoxal induced injury. *Sci Rep* **14**, 26640, doi:10.1038/s41598-024-
77891-9 (2024).
39   Elbarbary, N. S., Ismail, E. A. R., El-Naggar, A. R., Hamouda, M. H. & El-Hamamsy, M.
The effect of 12 weeks carnosine supplementation on renal functional integrity and
oxidative stress in pediatric patients with diabetic nephropathy: a randomized placebo-
controlled trial. *Pediatr Diabetes* **19**, 470-477, doi:10.1111/pedi.12564 (2018).
40   Jayne, L. *et al.* A torpor-like state in mice slows blood epigenetic aging and prolongs
healthspan. *Nat Aging* **5**, 437-449, doi:10.1038/s43587-025-00830-4 (2025).
41   Monroe, T. B. *et al.* Lipid peroxidation products induce carbonyl stress, mitochondrial
dysfunction, and cellular senescence in human and murine cells. *Aging Cell*, e14367,
doi:10.1111/acel.14367 (2024).
42   Whitsett, J., Picklo, M. J., Sr. & Vasquez-Vivar, J. 4-Hydroxy-2-nonenal increases
superoxide anion radical in endothelial cells via stimulated GTP cyclohydrolase
proteasomal degradation. *Arterioscler Thromb Vasc Biol* **27**, 2340-2347,
doi:10.1161/ATVBAHA.107.153742 (2007).
43   Hu, G. *et al.* Glutathione limits RUNX2 oxidation and degradation to regulate bone
formation. *JCI Insight* **8**, doi:10.1172/jci.insight.166888 (2023).
44   Bellantuono, I. *et al.* A toolbox for the longitudinal assessment of healthspan in aging
mice. *Nat Protoc* **15**, 540-574, doi:10.1038/s41596-019-0256-1 (2020).
45   Stuart, D. *et al.* Lack of renoprotective effects of targeting the endothelin A receptor and
(or) sodium glucose transporter 2 in a mouse model of Type 2 diabetic kidney disease.
Can J Physiol Pharmacol **100**, 763-771, doi:10.1139/cjpp-2022-0082 (2022).
46   Kittikulsuth, W., Stuart, D., Van Hoek, A. N. & Kohan, D. E. Lack of an effect of nephron-
specific deletion of adenylyl cyclase 3 on renal sodium and water excretion or arterial
pressure. *Physiol Rep* **3**, doi:10.14814/phy2.12316 (2015).
47   Eshima, H. *et al.* Neutralizing mitochondrial ROS does not rescue muscle atrophy
induced by hindlimb unloading in female mice. *J Appl Physiol (1985)* **129**, 124-132,
doi:10.1152/japplphysiol.00456.2019 (2020).
48   Shahtout, J. L. *et al.* Inhibition of the skeletal muscle Lands cycle ameliorates weakness
induced by physical inactivity. *J Cachexia Sarcopenia Muscle* **15**, 319-330,
doi:10.1002/jcsm.13406 (2024).
49   Ferrara, P. J. *et al.* Low lysophosphatidylcholine induces skeletal muscle myopathy that
is aggravated by high-fat diet feeding. *FASEB J* **35**, e21867, doi:10.1096/fj.202101104R
(2021).
50   Stringer, C., Wang, T., Michaelos, M. & Pachitariu, M. Cellpose: a generalist algorithm for
cellular segmentation. *Nat Methods* **18**, 100-106, doi:10.1038/s41592-020-01018-x
(2021).
51   Ramkumar, N. *et al.* Loss of Soluble (Pro)renin Receptor Attenuates Angiotensin-II
Induced Hypertension and Renal Injury. *Circ Res* **129**, 50-62,
doi:10.1161/CIRCRESAHA.120.317532 (2021).
52   Bouxsein, M. L. *et al.* Guidelines for assessment of bone microstructure in rodents using
micro-computed tomography. *J Bone Miner Res* **25**, 1468-1486, doi:10.1002/jbmr.141
(2010).
53   Harasymowicz, N. S. *et al.* Intergenerational Transmission of Diet-Induced Obesity,
Metabolic Imbalance, and Osteoarthritis in Mice. *Arthritis Rheumatol* **72**, 632-644,
doi:10.1002/art.41147 (2020).

Figure 1

Figure 2

Figure 3

Panels a–e: **Reproductive Health ♂** (Vehicle / N-acetylcarnosine); **Cognition/Nociception ♂** (Vehicle / N-acetylcarnosine): a, Sperm Count (Total, Dead, Alive); b, Brain Tissue Weight (mg); c, % Time Freezing (D1, D2); d, Δ% Freezing Time (D1, D2); e, Von Frey, Pressure (g).

Panels f–g: **Cognition/Nociception ♀** (Vehicle / N-acetylcarnosine): f, % Time Freezing (D1, D2); g, Hot Plate, Time (s).

Panel h:

Tissue	Metabolite	Female Vehicle	Female N-acetylcarnosine	Male Vehicle	Male N-acetylcarnosine	2-Way ANOVA p-value Treatment	Sex	Sex X Treatment
Plasma (µM)	N-acetylcarnosine	0.119 (0.038)	81.76 (5.095)	0.051 (0.010)	59.58 (9.438)	**<0.0001**	**0.0408**	**0.0417**
	Carnosine	0.938 (0.149)	1.173 (0.030)	0.816 (0.301)	0.775 (0.145)	0.723	0.353	0.617
	Anserine	1.085 (0.257)	2.292 (1.042)	2.737 (1.028)	2.681(0.134)	0.528	0.274	0.490
	Histidine	38.88 (3.174)	56.95 (8.564)	39.88 (12.48)	52.58 (8.255)	0.194	0.881	0.813
	Homocarnosine	ND	ND	ND	ND	N/A	N/A	N/A
nmol/µg	Protein Carbonyls	5.26 (.457)	4.535 (0.388)	3.790 (0.391)	3.061 (0.957)	0.2425	**0.0242**	0.9964
Urine (µM)	N-acetylcarnosine	29.26 (13.43)	16868 (1384)	1305 (806.2)	15375 (568.1)	**<0.0001**	0.891	0.0945
	Carnosine	23.57 (11.92)	28.95 (8.916)	23.31 (4.175)	34.91 (7.607)	0.343	0.747	0.724
	Anserine	9.948 (3.153)	13.44 (5.114)	35.90 (8.199)	32.15 (5.761)	0.983	**0.0020**	0.554
	Histidine	33.78 (8.553)	50.87 (9.297)	59.02 (10.33)	74.58 (4.587)	0.0728	**0.0112**	0.929
	Homocarnosine	2415 (416.7)	4282 (1373)	3388 (276.4)	3604 (271.6)	0.126	0.821	0.218
Muscle (µmol/kg WW)	N-acetylcarnosine	1.139 (0.726)	8.948 (2.076)	0.718 (0.141)	2.990 (0.424)	**0.0004**	**0.0118**	**0.0253**
	Carnosine	1291 (178.1)	943.8 (137.9)	2133 (57.26)	925.1 (228.1)	**0.0002**	**0.0223**	**0.0177**
	Anserine	4076 (481.3)	3120 (257.8)	5587 (153.5)	2757 (757.1)	**0.0010**	0.243	**0.0650**
	Histidine	394.3 (30.52)	197.7 (24.42)	315.5 (15.12)	124.4 (27.74)	**<0.0001**	**0.0080**	0.914
	Homocarnosine	ND	ND	ND	ND	N/A	N/A	N/A
nmol/µg	Protein Carbonyls	12.16 (3.26)	9.25 (1.35)	9.09 (1.054)	8.76 (0.701)	0.288	0.2436	0.3937
Brain (µmol/kg WW)	N-acetylcarnosine	ND	1.421 (0.154)	0.155 (0.010)	1.373 (0.336)	**<0.0001**	0.776	0.591
	Carnosine	51.74 (6.366)	61.42 (6.131)	81.34 (3.805)	59.800 (3.216)	0.260	**0.0140**	**0.0072**
	Anserine	6.800 (1.494)	3.680 (0.296)	6.220 (0.367)	4.520 (0.495)	**0.0097**	0.876	0.400
	Histidine	61.40 (6.421)	55.64 (4.457)	81.96 (10.94)	58.44 (6.315)	0.0662	0.135	0.249
	Homocarnosine	13417 (1376)	17284 (1766)	20427 (2019)	17889 (412)	0.668	**0.0235**	0.0514
nmol/µg	Protein Carbonyls	18.37 (6.44)	18.32 (3.36)	15.86 (2.47)	18.45 (2.79)	0.774	0.7873	0.766
Heart (µmol/kg WW)	N-acetylcarnosine	ND	6.094 (1.184)	ND	4.264 (0.732)	**<0.0001**	0.207	0.207
	Carnosine	1.776 (0.236)	2.875 (0.649)	7.032 (1.293)	5.172 (1.208)	0.694	**0.0011**	0.139
	Anserine	1.264 (0.134)	2.092 (0.435)	3.939 (0.857)	3.016 (0.478)	0.932	**0.0043**	0.125
	Histidine	56.26 (4.455)	46.26 (1.849)	72.05 (7.118)	49.02 (5.541)	**0.0052**	0.0885	0.221
	Homocarnosine	ND	ND	ND	ND	N/A	N/A	N/A
nmol/µg	Protein Carbonyls	74.51 (38.86)	9.19 (1.12)	34.38 (16.99)	10.18 (1.53)	**0.009**	0.228	0.206
Kidney (µmol/kg WW)	N-acetylcarnosine	0.139 (0.077)	111.2 (13.69)	0.216 (0.104)	117.0 (31.83)	**<0.0001**	0.868	0.871
	Carnosine	6.933 (1.798)	5.227 (1.665)	0.905 (0.066)	1.267 (0.148)	0.592	**0.0009**	0.412
	Anserine	6.059 (0.556)	5.594 (0.599)	2.903 (0.260)	4.907 (1.042)	0.271	**0.0116**	0.0860
	Histidine	21.90 (2.763)	27.01 (2.776)	34.08 (1.292)	35.05 (1.974)	0.203	**0.0004**	0.379
	Homocarnosine	ND	ND	ND	ND	N/A	N/A	N/A
nmol/µg	Protein Carbonyls	9.46 (3.27)	6.63 (1.23)	10.39 (2.45)	11.96 (4.05)	0.8489	0.347	0.507
Liver (µmol/kg WW)	N-acetylcarnosine	0.031 (0.031)	11.798 (1.381)	ND	18.535 (2.937)	**<0.0001**	**0.0318**	**0.0305**
	Carnosine	2.612 (0.293)	2.285 (0.317)	1.355 (0.242)	2.117 (0.360)	0.464	**0.0262**	0.0789
	Anserine	0.745 (0.094)	0.908 (0.149)	1.250 (0.087)	1.289 (0.106)	0.388	**0.0014**	0.592
	Histidine	150.7 (8.947)	155.6 (6.118)	150.2 (5.357)	206.4 (7.287)	**0.0006**	**0.0030**	**0.0025**
	Homocarnosine	ND	ND	ND	ND	N/A	N/A	N/A
nmol/µg	Protein Carbonyls	17.46 (3.87)	15.16 (1.76)	18.66 (5.22)	12.87 (2.78)	0.286	0.882	0.639
iWAT (µmol/kg WW)	N-acetylcarnosine	0.112 (0.022)	14.76 (4.174)	1.747 (0.778)	8.723 (5.318)	**0.0037**	0.496	0.242
	Carnosine	29.27 (13.43)	16867 (1385)	1306 (806.2)	15375 (568.1)	**<0.0001**	0.891	0.0945
	Anserine	50.53 (27.89)	19.32 (14.61)	10.08 (5.334)	212.3 (163.0)	0.236	0.289	0.113
	Histidine	22.93 (8.432)	9.142 (1.983)	9.954 (1.329)	19.34 (7.590)	0.700	0.808	0.0568
	Homocarnosine	ND	ND	ND	ND	N/A	N/A	N/A
nmol/µg	Protein Carbonyls	20.26 (3.24)	12.75 (2.87)	17.59 (4.36)	13.85 (3.20)	0.141	0.8313	0.611
gWAT (µmol/kg WW)	N-acetylcarnosine	0.0123 (0.071)	11.669 (7.148)	1.019 (0.446)	3.439 (0.381)	0.0872	0.352	0.250
	Carnosine	5.186 (4.892)	0.393 (0.247)	0.452 (0.054)	0.657 (0.110)	0.393	0.405	0.353
	Anserine	28.05 (25.17)	1.728 (1.119)	1.557 (0.155)	2.439 (0.348)	0.358	0.352	0.327
	Histidine	13.37 (7.114)	4.017 (0.736)	7.105 (0.802)	9.536 (2.703)	0.399	0.927	0.160
	Homocarnosine	ND	ND	ND	ND	N/A	N/A	N/A
nmol/µg	Protein Carbonyls	22.25 (9.83)	33.94 (4.02)	33.16 (12.96)	29.81 (8.42)	0.649	0.712	0.415

Histidine-Containing Dipeptide and Protein Carbonyl Concentrations Across Tissues. Targeted metabolomics was assessed via LC-MS to quantify N-acetylcarnosine, Carnosine, Anserine, Histidine, and Homocysteine in Plasma, Urine, Muscle (TA -HCDs, Gastroc -Carbonyls), Brain (Cortex), Heart (Ventricle), Kidney, Liver, iWAT, and gWAT. Data are presented as Mean ±SEM and were analyzed via 2-Way ANOVA. p-values for main effects of treatment, sex, and interaction are shown in the last three columns. Significance was set at p< 0.05, p-values that are below this threshold are bolded to indicate significance.