Buffer-dependent conformational dynamics of α-synuclein revealed by

nanopipette electrospray ionisation ion mobility mass spectrometry

Emily J. Byrd[1]•, Emma L. Norgate[1]•, Joel A. Crossley[1], Chalmers C. C. Chau[2,3], Bob

Schiffrin[1], Alexander Kulak[4], Sheena E. Radford[1], Paolo Actis[2,3*], Antonio N. Calabrese[1*],

Frank Sobott[1*]

[1]Astbury Centre for Structural Molecular Biology, School of Molecular and Cellular Biology,

Faculty of Biological Sciences, University of Leeds, Leeds, LS2 9JT, UK.

[2]School of Electronic and Electrical Engineering, University of Leeds, LS2 9JT, UK.

[3]Bragg Centre for Materials Research, University of Leeds, LS2 9JT, UK.

[4]School of Chemistry, University of Leeds, LS2 9JT, UK.

• These authors contributed equally to this work.

*Correspondence to: f.sobott@leeds.ac.uk, a.calabrese@leeds.ac.uk, p.actis@leeds.ac.uk

ABSTRACT

Electrolyte conditions *in vivo* and *in vitro* are known to influence protein structure and function. Intrinsically disordered proteins (IDPs) are particularly sensitive to their solution conditions such as ionic strength and molecular crowding, and their dynamic structural ensembles rapidly respond to the cellular environment. While structural mass spectrometry (MS) techniques are uniquely able to capture aspects of this structural diversity, technical limitations have largely precluded the use of native MS approaches to interrogate the conformational rearrangements of IDPs in response to high concentrations of non-volatile salts. Here, we overcome this challenge by employing sub 100-nm nanopipette electrospray emitters for more gentle and salt-tolerant analysis to study the conformational distribution of α-Synuclein (αS) using native MS and ion mobility-MS in varied solution conditions, including in phosphate buffered saline. We show using native MS that it is possible to capture salt and buffer induced changes in the αS conformational ensemble when using traditional biochemical buffers, which reflect structural changes from *in silico* predictions and in-solution measurements. This work demonstrates the power of nanopipette emitters for the study of IDPs, and establishes native MS as a method that can be routinely used to determine how solution conditions tune the conformational landscape of IDPs.

INTRODUCTION

Intrinsically disordered proteins (IDPs) and proteins containing intrinsically disordered regions (IDRs) partake in many protein-protein interactions, aided by their structural plasticity, which facilitates their functional roles in processes such as transcriptional regulation and cell signalling[1−4]. IDPs are known to explore a vast landscape of rapidly inter-converting and co-existing configurations which are determined by their sequence and the chemical composition of their environment[5]. Many IDPs and IDRs are implicated in disease pathogenesis and are therefore therapeutic targets. This includes diseases such as cancer[6,7] and neurodegenerative conditions, where many IDPs have been shown to undergo a disorder-to-order transition to form insoluble, β-sheet rich amyloid fibrils [8−10]. Deposits of amyloid fibrils are associated with a number of neurodegenerative diseases, including Parkinson's disease (PD), where α-synuclein (αS) fibrils are found in insoluble Lewy body deposits in the substantia nigra of the brain [11−17].

The structure and function of the IDP αS are affected by environmental conditions. For example, increasing NaCl concentration (from 50 mM up to 1000 mM) enhances liquid-liquid phase separation of αS in the presence of crowding agents (PEG400)[18], by exposing the non amyloid-β component (NAC) domain, a hydrophobic region of the sequence (residues 60-95) through the displacement of water molecules[19]. This ultimately results in modulation of the aggregation pathway, leading to different fibril morphologies[20−23]. Moderate physiological concentrations of NaCl (up to *ca.* 0.1-0.3 M) have been shown to weaken contacts between the N-terminal, C-terminal and the NAC domain of αS, resulting in fewer intramolecular contacts and enhancing aggregation into amyloid[24,25]. Additionally, monovalent and divalent ions such as Ca^{2+}, Cu^{2+}, Cu^+, Mn^{2+}, Mg^{2+}, Fe^{2+} and Zn^{2+}, many of which are shown to be present at elevated concentration in the brains of Parkinson's disease patients and are colocalised with αS in Lewy body assemblies[26,27], have been identified as regulators of the conformational landscape of αS and as accelerators of amyloid assembly[26,28−38]. Previous data from hydrogen-deuterium exchange-mass spectrometry identified that the conformational behaviour of αS differs between solution conditions that match the extracellular, cytosolic and lysosomal environments (which differ in pH and ionic strength), with discrete conformational changes identified in the central hydrophobic NAC domain of αS which forms the amyloid core[39,40]. These different solution conditions also result in altered αS amyloid assembly kinetics[39,41,42], highlighting the importance of replicating physiological conditions when studying IDPs *in vitro*.

Native mass spectrometry (native MS), especially when combined with ion mobility-MS (IM-

MS), is able to capture and characterise dynamic, transient, heterogeneous species within a

conformational ensemble and is a powerful technique for investigating IDP conformational

ensembles[43–50]. The first step in a native MS experiment is the gentle ionisation and transfer

of protein species which are kinetically trapped in their solution states into the gas phase.

This process is typically achieved using nano electrospray ionisation (nESI) under non-

denaturing solution conditions so that information about protein conformations adopted in

solution can be measured in the gas phase[51,52]. The desolvation process that occurs during

nESI requires the use of aqueous solutions comprising volatile salts such as ammonium

acetate (AmAc)[53]. The presence of salts that are commonly required during protein

preparation (such as Na^+, K^+, PO_4^{3+} and Tris) as well as buffer additives that may be critical

to maintain the structural integrity of proteins, can cause ion suppression which reduces

spectral quality, resolution and sensitivity [54–57]. Salt ions have also been shown to adduct to

protein ions[58–60], which can result in unresolvable mass spectra. Therefore, prior to native

MS analysis, proteins are typically buffer exchanged from their biochemical storage buffer (a

process which has been simplified recently with online buffer exchange[61,62]) into AmAc to

dilute/remove salt ions from solution to prevent adduction in mass spectra (ammonium ions

are volatile and do not adduct to proteins). However, many proteins require salts to remain

stable in solution, and produce poorly resolved native mass spectra in AmAc[60]. One way to

circumvent ion adduction is to generate smaller droplets during the nESI process so that salt

ions remain independent from the solvent droplets containing protein molecules[63–65]. This

approach has been applied successfully to the study of membrane proteins in detergents

and other proteins in biochemical buffers[59,66]. Using IM to measure the rotationally averaged

collision cross section (CCS) distributions of ion species coupled to MS is a powerful

technique for studying the conformational properties of IDPs [35,36,50,67–70] , and we, and others

have previously used this approach to probe the conformations of αS under different solution

conditions (e.g. in the presence of metal ions, dopamine and detergents)[35,36,38,50,71–74].

However, these experiments have been limited to using AmAc based solutions.

Here we demonstrate the application of quartz nanopipette nESI emitters with a tip diameter

of < 100 nm (standard borosilicate nESI emitters have diameters *ca.* 2-20 μm; Figure S1 and

S2) for native MS of an IDP, αS in its N-terminally acetylated form[53,75,76]. While previous

work has shown that submicron emitters can be used for nESI and that they enable the

acquisition of native mass spectra in diverse buffer conditions[57,63,65,66,77,78], these efforts

were limited to proof-of-concept studies of stably folded test proteins. Nanopipette emitters

have not been previously used to elucidate structural changes involving αS, or other disease

relevant IDPs. Moreover, the use of nanopipettes to characterise IDPs, a class of proteins

that are especially responsive to the solution conditions[67,79,80], and the conformational
changes that they undertake, has not been previously described. Given the role of solution
conditions in tuning the kinetics of αS aggregation[34,39,40,42] and the experimental evidence
demonstrating that different NaCl concentrations alter structural properties,[20,24,42] we chose
to interrogate how ionic strength and buffer composition, known modulators of αS
conformation and amyloid propensity[18–20,42,81], tune conformational distribution, as measured
by IM-MS. We show that native mass spectra of αS can be acquired in nonvolatile buffers
commonly used in biochemical/biophysical studies of protein structure and function *in vitro*,
including Dulbecco's phosphate buffered saline (PBS), Tris-HCl, potassium phosphate and
sodium phosphate, and in the presence of added NaCl. By combining data from IM-MS with
in silico and in solution measurements of protein conformation, we demonstrate that native
MS can reveal how both buffer composition and ionic strength tune the conformational
heterogeneity of αS. Combined, this work demonstrates that nanopipette nESI emitters can
expand the repertoire of possibilities for native MS and IM-MS studies of IDPs, to enable
new insights into how IDP conformational ensembles respond to changes in the solution
environment and understand how this impacts biological function or in the case of disease,
dysfunction.

RESULTS AND DISCUSION

Native MS in physiological buffers

To test the limits of nanopipette technology for nESI-MS analysis of αS, we first attempted to
use nanopipette emitters (quartz nanopipettes with a nanopore opening of *ca.* 40 nm, Figure
S1 and S2) to obtain native mass spectra of αS in PBS (Dulbeccos A), a buffer commonly
used in biochemical studies. Excitingly, we were able to measure a well-resolved spectrum
when using these nanopipette emitters, but (as expected) not when using standard µM-wide
emitters (Figure 1a,b). A similar αS charge state distribution was observed with both
emitters, suggesting that the smaller tip diameter of the nanopipette does not alter the
structural ensemble of monomeric αSas the charge states adopted in native MS relatively
reflects a proteins solvent exposed surface area[52,82,85–87].

In native MS analysis it is common to observe non-specific adducts and clusters of salts, e.g.
Na+ (+23 Da) and K+ (+39 Da)[83]. Reduced salt clustering and salt adducts can be observed
with nanopipette emitters (Figure 1b), compared to standard emitters (Figure 1a and Figure
S3), consistent with previous reports[54]. The native mass spectrum in Figure 1b shows an
overall decrease in salt adducts,as well as a marked reduction of salt clusters < 1000 *m/z*.

Overall, this shows that nanopipette emitters enable well-resolved nESI mass spectra of αS

in biochemical buffers, consistent with previous studies demonstrating an enhanced salt

tolerance when using sub-micron nESI emitters to analyse proteins from buffers with non-

volatile components [57,84]. By showing that it is possible to capture the conformational

diversity of αS in PBS using MS, we highlight the potential of nESI with nanopipette emitters

for conformational analysis of IDPs in complex solution environments which mimic more

biologically relevant conditions (cellular electrolytes, salts, crowding and liquid-liquid phase

separation promoting agents), and demonstrate that there is no need to avoid non-volatile

salts in native IM-MS.

Figure 1. Direct comparison of nanopipette and standard nESI emitters. *Representative*
native nESI mass spectra of 20 μM αS analysed in PBS, pH 7.2 using (a) standard emitters
and (b) nanopipette emitters using identical instrument acquisition parameters on Waters
Synapt HDMS (see Methods). I.D.: internal diameter.
161

When comparing native nESI mass spectra of αS acquired using nanopipettes in PBS

(Figure 1b) with those in AmAc buffer (Figure 2a), a shift in the charge state distribution (6+

to 14+ vs. 5+ to 18+) can be noticed as well as the emergence of dimer spectral peaks in

AmAc. The most intense charge state in PBS is 9+ whereas the 7+ charge state was most

intense in AmAc.

We next performed native MS and IM-MS to further understand how these changes in the

mass spectra are related to the conformational alterations of αS in a range of commonly

used biochemical buffers, by calculating collision cross section (CCS) distributions from the

IM-MS data (see Methods)[46]. Travelling wave (TW) IM measurements were performed in N_2

buffer gas for each charge state. Consistent with community standards[48] we refer to these

values as $^{TW}CCS_{N2}$. To provide a snapshot of the conformational ensemble captured by the

native IM mass specta, we measured $^{TW}CCS_{N2}$ values for each detected charge state and

plotted a cumulative distribution where the contributions from each individual charge state

are intensity-weighted and combined (using a scaling factor based on the intensity of each

charge state in the corresponding mass spectrum, see Methods). We acquired native mass

spectra of αS from the buffers Tris-HCl, potassium phosphate and sodium phosphate, Figure

2c,e,g and measured IM-MS $^{TW}CCS_{N2}$ distributions from these spectra (Figure 2b, d, f, h), for

comparison with data acquired in conventional AmAc (Figure 2a,b)[88,89].

Using nanopipette emitters, a resolvable charge state distribution was achieved for αS in

each of the tested buffer systems, and clear shifts in charge state intensities were observed

between buffers (Figure 2a,c,e,g). The native mass spectrum of αS in 20 mM AmAc (Figure

2a) appears as a multimodal distribution of charge states ranging from an extended

conformational family (+17 to +10) to a compact conformational family (+9 to +5). This is

reflected in the ion mobility $^{TW}CCS_{N2}$ plot of these data (Figure 2b) and has been shown

previously[36]. To enable a simple comparison between experimental conditions, we

calculated the average (mean) CCS value from the culmulative $^{TW}CCS_{N2}$ distributions

(vertical dashed lines in Figure 2). In 20 mM AmAc, this average $^{TW}CCS_{N2}$ value (2487 Å^2)

lies in between the extended and compact conformational families (Figure 2b). Interestingly,

the spectra and $^{TW}CCS_{N2}$ distributions in PBS (Figure 1b, Figure S4) and the other two

phosphate buffers tested (Figure 2e-h) are comparable. All three phosphate buffered

solutions resulted in $^{TW}CCS_{N2}$ distributions that were narrower and favoured more expanded

conformations compared to those observed in AmAc (average $^{TW}CCS_{N2}$ values of 2598 Å^2 ,

2772 Å^2 and 2723 Å^2 in PBS, potassium phosphate and sodium phosphate, respectively,

compared to 2487 Å^2 in AmAc). Contrary to this, the spectrum obtained in 20 mM Tris-HCl

buffer is shifted to populate lower charge states, and charge state 12+ and above are not

observed (Figure 2c). This is reflected in the $^{TW}CCS_{N2}$ distribution which demonstrates that

the compact conformational familes are favoured in Tris-HCl, with the average value of this

$^{TW}CCS_{N2}$ distribution being shifted to a lower value (2147 Å^2; Figure 2d). It appears that in

Tris-HCl, the conformational ensemble is less heterogeneous overall, populating two distinct

conformational families, compared with a broader, multimodal distribution detected in AmAc,

reflected by the decrease in the abundance of highly extended conformational families

(higher charge states 12+ to 18+), however the $^{TW}CCS_{N2}$ window in Å^2 is the same.

Conversely, the native mass spectra and $^{TW}CCS_{N2}$ distributions of αS in 10 mM potassium

phosphate, 20 mM sodium phosphate and PBS are consistent with more expanded αS

conformational families predominating than in AmAc, and the CCS fingerprint appears

distinctly different to αS in both AmAc and Tris-HCl.

Combined, these results show that the nm diameter orifice of the nanopipettes described in

this work can enable acquisition of resolvable mass spectra to obtain IM-MS fingerprints of

αS in a range of non-volatile biochemical buffers. The charge state distributions of αS

observed in the mass spectra and the $^{TW}CCS_{N2}$ distributions detected using IM-MS also differ in different buffer conditions. This suggests that modulation of the conformational landscape of the IDP αS in different buffers can be detected by native IM-MS using nanopipettes. Such a change has been reported previously for the folded protein bovine serum albumin (BSA) by analysis of native MS charge state distributions and gas phase thermal denaturation. AmAc was shown to stabilise compact conformational families of BSA and the presence of NaCl shifted the population of BSA towards more unfolded conformations[84].

Figure 2. Native MS and IM-MS of αS in biochemical buffers. *Native nESI mass spectra of 20 μM αS (left panel) and $^{TW}CCS_{N2}$ distributions (right panel) in (a,b) 20 mM AmAc, (c,d) 20 mM Tris-HCl, (e,f) 10 mM potassium phosphate, (g,h) 20 mM sodium phosphate. Selected charge states are labelled in the mass spectra. The key on the righthand side indicates the colour of the CCS distribution for each charge state, the black solid line is the cumulative fit and the black dotted line represents the average $^{TW}CCS_{N2}$ value from the distribution.*

Tip diameter reduces salt adduction enabling analysis of the effect of salt on CCS.

Next, we titrated NaCl into AmAc and Tris-HCl buffers to systematically investigate the effect of ionic strength on the conformational families populated by αS using IM-MS (Figure 3). In both AmAc and Tris-HCl, increasing the concentration of NaCl in the solution results in αS populating more highly charged species by enhancing the population of more expanded conformational families (Figure S5 and Figure S6, respectively with the 8+ charge state shown in Figure S7). This is supported by analysis of the $^{TW}CCS_{N2}$ distributions which also show a clear shift towards higher average CCS values with increasing salt concentrations (Figure 3a-h). We hypothesise that this shift towards extended conformations is likely due to a charge screening effect in which Na^+ ions interacts with the negatively charged C-terminal region and Cl^- ions interact with the overall positively charged N-terminal region[24], reducing intraprotein contacts. This is consistent with NaCl-dependent expansion of αS observed previously using small angle X-ray scattering (SAXS)[19].

As a control, we compared the effect of adding NaCl on the measured $^{TW}CCS_{N2}$ values of αS with those of natively folded and disulfide-stabilised hen egg white lysozyme (Figure S8), a well-studied globular protein of similar molecular weight to αS (14,350 Da compared with 14,502 Da for N-terminally acetylated αS used here)[90–92]. In both AmAc and Tris-HCl buffers, more compact species and narrower conformational distributions of lysozyme were observed upon addition of 25 mM NaCl, by contrast with the conformational extension observed with αS. This suggests that our observations for αS are a result of the protein structurally changing in response to its solution conditions, and that these structural features are captured in the gas-phase with IM-MS, and not a result of changes in the nESI process induced by the addition of salt. The results are consistent with a model in which the disordered and extended conformational families of αS are stabilised in elevated salt concetrations and that the protein is less driven to form intraprotein contacts which would stabilise more compact states (Figure 3i).

Figure 3. Salt titration of αS in AmAc and Tris-HCl by native ion mobility mass
spectrometry. $^{TW}CCS_{N2}$ distributions showing the effect of salt (NaCl) on αS. αS was
measured in (a) 20 mM AmAc with the addition of (b) 12.5, (c) 25 and (d) 125 mM NaCl and
also in (e) 20 mM Tris-HCl with the addition of (f) 12.5, (g) 25 and (h) 125 mM NaCl. (i) A
schematic of αS expansion in the presence of NaCl. The key on the righthand side indicates
the colour of the CCS distribution for each charge state, the black solid line is the cumulative
fit and the black dotted line represents the average $^{TW}CCS_{N2}$ value.
265

Course grained molecular dynamics (MD) simulations suggests expanded states of
267

αS are favoured at elevated NaCl concentrations
268

Previous coarse-grained MD simulations showed a salt dependent increase in the radius of
269

gyration (R_g) of αS up to a NaCl concentration of 0.1 M as a result of charge shielding[24]. We
270

sought to expand on these studies using coarse-grained simulations of αS in implicit solvent
271

using CALVADOS[93,94]. Simulations were carried out at ionic strengths ranging from 10 - 150

mM (these ionic strengths compare to the total ionic strength of 20 mM AmAc in the absence

and presence of NaCl used in IM-MS experiments here) and the R_g values of the resultant

model structures of αS were calculated. These data demonstrate a salt dependent increase

in the average R_g of αS monomers from the simulations over the range of ionic strength

values tested (Figure 4a). This change in average R_g is a result of shifts in the distribution of

the αS conformational ensemble towards structures with higher R_g values with increasing

ionic strength (Figure S9). We compared these computationally-generated R_g values with

the apparent R_g (R_{app}) derived from the weight-averaged mean of the $^{TW}CCS_{N2}$ distributions

(see Methods), as a proxy for an ensemble-averaged measurement of CCS (See Methods).

Similar to the computationally derived R_g values, we observe an increase in CCS as a

function of NaCl concentration in both AmAc and Tris-HCl infering that αS populates more

extended conformations (Figure 4b,c). Examination of representative structures from the

simulations at ionic strengths of either 10 mM or 150 mM demonstrates the role of

interactions between the N- and C-termini of αS in mediating chain collapse (Figure 4d),

consistent with previous reports[34,95].

Figure 4. The mean R_g of αS from coarse grained simulations and average (R_{app}) IM-
MS values both increase in response to increasing ionic strength. *(a) The mean radius*
of gyration (R_g) of αS as determined by coarse-grained implicit-solvent simulations at varying
ionic strengths (see Methods). The error shown is the standard error of the mean,
determined by considering the calculated R_g values for all αS models in the simulation after
a 5 ns equilibration (see Methods for details). (b, c) Average R_{app} calculated from the
average $^{TW}CCS_{N2}$ distributions of αS (See Methods, data from Figure 3) in (b) 20 mM Tris-

HCl or (c) 20 mM AmAc, with data recorded in the presence of different concentrations of
NaCl. (d) Representative coarse grained simulation frames of αS with calculated R_g values
of 31Å and 37Å.
299

Next, we sought to compare our gas phase $^{TW}CCS_{N2}$ measurements with solution phase

measurements using diffusion-ordered spectroscopy nuclear magnetic resonance (DOSY

NMR)[96]. We measured diffusion coefficients for αS in the absence or presence of NaCl (25

mM NaCl) in both AmAc and Tris-HCl buffers (Figure 5a,b,c and Figure S10) and the data

show that upon addition of NaCl, under both sets of buffer conditions, the measured diffusion

coefficient of αS decreases (~15%). The diffusion coefficient is inversely proportional to the

hydrodynamic radius (R_h) of the molecule, suggesting that addition of NaCl results in

expansion of αS under the buffer conditions tested[96,97]. This is consistent with previous

small angle X-ray scattering (SAXS) data that identified a NaCl-dependent increase in the R_g

of αS[19]. Taken together, these data provide experimental evidence that IM-MS $^{TW}CCS_{N2}$

gas-phase measurements are reflective of the solution-phase conformational distributions of

αS, and demonstrates that it is possible, using nanopipette IM-MS, to determine the

conformational properties of αS in different buffer conditions.

Figure 5. ^{1}H DOSY NMR indicates expansion of αS upon addition of NaCl. *Diffusion*
coeffici-ents for αS in solution are reduced in the presence of 25 mM NaCl. (a,b) Fitted ^{1}H
DOSY NMR data for αS in the presence or absence of 25 mM NaCl in (a) AmAc buffer, and
(b) Tris-HCl buffer. Errors were calculated from the signal-to-noise level in the ^{1}H spectra
(see Methods). (c) αS diffusion coefficients under different buffer conditions. Errors were
calculated using a Monte Carlo simulation approach and the spectral noise (see Methods).
Samples contained 50 µM αS, 0 mM or 25 mM NaCl, in 100 mM ammonium acetate, pH 7.2,
or 100 mM Tris-HCl, pH 7.2, at 24 °C.

To explore whether the conformational expansion of αS reflects a general sensitivity to ionic

strength, or an effect on conformation that is specific to adding NaCl, we performed IM-MS

of αS at elevated buffer concentrations (100 mM) of AmAc and Tris-HCl buffers. In the case

of AmAc, at 20 mM the average CCS of the $^{TW}CCS_{N2}$ distribution was 2487 Å^2 (Figure 1b),

and this increased to 2786 Å^2 in 100 mM AmAc (Figure 6). In the presence of an additional

25 mM NaCl we observed a CCS shift of αS towards higher, more expanded $^{TW}CCS_{N2}$

values, with the more compact conformational families significantly reducing in abundance

which is also reflected in the measured charge state distributions (Figure 6, S5). A similar

shift was observed in Tris-HCl buffer upon addition of NaCl (Figure 6, S6). This suggests

that the conformational changes that occur in elevated ionic strength conditions are not

specific to Na^+/Cl^- interactions but are instead driven by the overall ionic strength of the

solution. To summarise, we find that derived R_{app} from IM-MS measurements in the gas

phase, computationally calculated R_g values from CALVADOS simulations as well as in

solution determination of diffusion coefficient by DOSY NMR all show a clear

expansion/increase in size of αS upon addition of NaCl (Table S1).

Environmental factors including salt concentration are known to determine amyloid fibril

assembly conditions, leading to specific αS fibril morphologies[98]. Ions can stabilise both

intra- and intermolecular hydrophobic interactions through dehydration, promoting fibril

assembly[20,99–101]. Additionally, salt has been proposed to cause expansion of αS, exposing

the hydrophobic amyloid prone NAC domain of the protein leading to accelerated amyloid

assembly kinetics[19]. Here, in order to relate conformational changes to amyloid assembly

kinetics were monitored using Thioflavin T (ThT) fluorescence in the presence or absence of

NaCl (25 mM). The results showed that the rate of amyloid assembly increases with the

addition of 25 mM NaCl compared to 0 mM NaCl in both AmAc and Tris-HCl buffers (Figure

6e, f)[25,102]. This suggests that the change in structure we observe with IM-MS directly

impacts the amyloid assembly kinetics of αS, consistent with previous reports[36]. These

findings together exemplify the role that structural rearrangements of αS measured by IM-

MS correlate with its aggregation propensity and could help to elucidate disease relevant

conformational families for therapeutic intervention.

Figure 6. αS undergoes expansion at high ionic strength. $^{TW}CCS_{N2}$ *distributions showing the effect of NaCl on αS. αS was measured in (a) 100 mM AmAc with the addition of (b) 25 mM NaCl, and in (c) 100 mM Tris-HCl with the addition of (d) 25 mM NaCl. The key on the righthand side indicates the colour of the CCS distribution for each charge state, the black solid line is the cumulative fit and the black dotted linke represents the average* $^{TW}CCS_{N2}$ *value. (e) ThT fluorescence in 100 mM AmAc in the absence or presence of 25 mM NaCl and in (f) 100 mM Tris-HCl in the absence or presence of 25 mM NaCl. Three replicates are shown.*

CONCLUSION

This study demonstrates that there are remarkable shifts in the conformational ensembles of αS driven by changes in the ionic envirnoment, demonstrating that the structural landscape of αS is highly affected by salt concentration. The incorporation of nanopipette emitters has enabled us to show using native IM-MS that NaCl and the buffer composition modulate the conformational landscape of αS. We show that the conformational expansion of αS in presence of NaCl (Figure 3) determined using IM-MS correlates with in solution DOSY NMR measurements and computationally-derived R_g values (Figure 4, 5). IM-MS is well placed to capture snapshots of entire conformational ensembles and provide rich structural information for IDPs. Whereas a standard nESI set-up mostly limits analysis to volatile buffers which do

not represent physiological conditions, nanopipette emitters enable new capabilities for IM-

MS through the analysis of samples in common biochemical buffers and high-salt

environments.

Overall, a clear reduction in αS salt adducts is observed in native nESI mass spectra using

the nanopipette emitters described here (Figure 1, Figure 2 and S3) which we attribute to the

finer droplet spray[103,104]. The smaller droplets that emerge from the nanopipette partition the

solvent into smaller volumes that contain fewer salt ions on average which reduces

formation of adducts with proteins observed with standard ESI emitters. Our application of

nanopipette emitters opens the door to deploying native IM-MS to afford a new

understanding of how the conformations of IDPs relate to their function or dysfunction, such

as the role of αS in synucleopathies, and to analyse proteins in buffers closer to their native

cellular environment.

Our results pave the way for the structural analysis of proteins under physiological conditions

by native IM-MS. Future studies exploring the addition of other buffer components, and the

introduction of complex sample environments such as cellular compartments and cell

lysates, will further push this methodology forwards.

ACKNOWLEDGEMENTS

A.N.C. and E.J.B. acknowledge support through a Sir Henry Dale Fellowship jointly funded

by Wellcome and the Royal Society (220628/Z/20/Z). J.A.C acknowledges the support of the

University of Leeds and the BBSRC (BB/Y00034X/1). P.A., C.C.C.C. and A.N.C.

acknowledge funding from the BBSRC (BB/X003086/1) and F.S., P.A., E.L.N. from the MRC

(MR/W031515/1). S.E.R. is the grateful recipient of a Royal Society Research Professorship

(RSRP/R1/211057). Funding from the BBSRC enabled the purchase of MS equipment

(BB/E012558/1). The authors thank Dr Sri Ranjani Ganji and Dr Gemma Wildsmith from the

Biomolecular Mass Spectrometry Facility at the University of Leeds for their support and

assistance in this work. For access to the 600 MHz spectrometer, we acknowledge the

Astbury Biostructure Laboratory BioNMR facility which was funded by the University of

Leeds. We also thank Dr Arnout Kalverda for excellent technical support, and Dr Pijush

Chakraborty and Dr Alex Heyam for helpful discussions and advice.

METHODS

Standard nESI emitter fabrication

Borosilicate thin wall glass capillaries (0.78 mm) with filament (Harvard Apparatus, UK) were pulled using a P-97 Flaming/Brown micropipette puller (Sputter Instrument, CA, USA) to make two nanospray tips. The optimised procedure used three cycles as follows; heat 560, pull 250, velocity 10, time 50 s. Capillaries were placed into a glass petri dish and coated with palladium in a SC7620 mini sputter coater (Quorum Technologies, Sussex, UK) pressurised with argon at 2×10^{-2} mbar. When a current of 25-30 mA was applied the plasma enabled palladium atoms to deposit on the glass surface. The coating time lasted 75 s after which the petri dish was rotated by 90° followed by a second coating. Samples (10 µL) were loaded into capillaries which were then clipped to size using tweezers.

Nanopipette nESI emitter fabrication

The nanopipette nESI emitter tips were fabricated using 1.0 mm outer diameter and 0.5 mm inner diameter quartz capillaries (QF100-50-7.5; Sutter Instrument) with the SU-P2000 laser puller (World Precision Instruments). A two-line protocol was used: line 1 with HEAT 750/FIL 4/VEL 30/DEL 150/PUL 80, followed by line 2 with HEAT 850/FIL 3/VEL 40/DEL 135/PUL 225. For native IM-MS measurements, the emitters were loaded with analyte solution (8 µL) and fitted with a platinum wire (PT00-WR-000117; Goodfellow) prior to use.

Scanning electron microscopy (SEM) nESI emitter characterisation

The pore dimensions of the standard nESI and nanopipette nESI were characterised by field emission SEM with a FEI Nova 450 at an accelerating voltage of 3–5 kV. Images were prepared without coating, using a CBS detector (back-scattered electron detector).

Native ion mobility mass spectrometry

Native IM-MS experiments were performed on a Synapt HDMS mass spectrometer (Waters Corporation, Wilmslow, UK) with travelling (T-wave) ion mobility and a nano-ESI source. N-terminally acetylated αS, expressed and purified as described in[36], was analysed at a concentration of 20 µM in 20 mM AmAc, 20 mM Tris-HCl, 1 x PBS (Dulbeccos A; Thermo Fisher Scientific), 10 mM potassium phosphate, 20 mM sodium phosphate, 100 mM AmAc, and 100 mM Tris-HCl each at a pH of 7.2. For salt titrations, NaCl was added at 12.5 mM, 25 mM and 125 mM. Instrument parameters were set at: capillary voltage 1.0 kV, source temperature 30 °C, backing pressure 0.0 – 0.3 Bar, sampling cone 18 V, extraction cone 1.0 V, trap collision energy 5 V, transfer collision energy 2.0 V, trap DC bias 30 V, IM wave velocity 300 m/s, IM wave height 7.0 V. Gas pressures in the instrument were: trap cell

0.0258 mbar, IM cell 0.36 mbar. $^{TW}CCS_{N2}$ signifies CCS values calculated using traveling

wave ion mobility in N_2 buffer gas using calibrants acquired in N_2 buffer gas. Lysozyme was

analysed at 10 µM under identical conditions except for: sampling cone 10 V and trap DC

bias 12 V. MassLynx V4.1 (Waters Corporation, Wilmslow, UK) was used for data

processing. The IM spectra were calibrated according to the Bush database[46] using

denatured cytochrome C (charge states 13+ to 19+), myoglobin (charge states 15+ to 24+)

and ubiquitin (charge states 7+ to 13+) at 10 µM in 50 % (v/v) acetonitrile, 0.1 % (v/v) formic

acid. Calibrated intensity values were normalised using a scaling factor determined by the

signal intensity in the mass spectrum as well as the area under the arrival time distribution

peak. Hen egg white lysozyme (Sigma) IM spectra were also calibrated[46] using native

cytochrome C (charge states 6+ and 7+) and myoglobin (charge states 7+ to 9+) measured

in 100 mM AmAc. The apparent radius of gyration (R_{app}) was calculated using the following

equation: $R_{app} = \sqrt{\dfrac{Average\ CCS}{\pi}}$

^{1}H DOSY NMR

DOSY experiments were performed on a 600 MHz Bruker Avance III spectrometer equipped

with a 5 mm QCI-P CryoProbe. For each DOSY experiment under each sample condition, a

total of 24 1D acquisitions were performed at different gradient strengths between 2% and

98% where the maximum gradient strength was 48.148 G cm^{-1}. To improve data fitting the

gradient strength percentages were weighted quadratically towards both 2% and 98% as

follows: 2.00, 5.32, 8.76, 12.30, 16.10, 20.00, 24.10, 28.40, 33.00, 38.00, 43.40, 49.40,

50.60, 56.60, 62.00, 67.00, 71.60, 75.90, 80.00, 83.90, 87.70, 91.20, 94.70, 98.00. Following

processing, peaks were integrated in the methyl region between 0.6-0.9 ppm. To obtain the

error for each experiment, the noise at each gradient strength was obtained by integrating a

region of the baseline of the same size where no peaks were present (0.3-0.6 ppm). To

obtain the diffusion coefficient, following baseline correction and integration, the data were

fitted to the following equation:

$$\frac{I}{I_0} = e^{-D\,\gamma^2\,g^2\,\delta^2\left(\Delta-\frac{\delta}{3}\right)}$$

where I is the observed intensity, I_0 the reference intensity (the signal intensity of the first

point), D is the diffusion coefficient, γ is the gyromagnetic ratio of the observed nucleus, g is

the gradient strength, δ the length of the gradient, and Δ the diffusion time. The length of the

gradient (δ) and diffusion time (Δ) were 2 ms and 400 ms, respectively. Baseline adjustment,

integration, fitting and plotting was performed with in-house Python3 scripts which made use

of the NumPy (v1.26.4)[100], SciPy (v1.13.1)[101] and Matplotlib (v3.9.2)[102] libraries. The

integration was performed using the numpy.trapz function. To estimate the uncertainty in the

diffusion coefficient a Monte Carlo simulation approach, incorporating the errors associated

with each data point, was employed. To simulate the effect of the error in each data point on

the diffusion coefficient, we first generated 1000 synthetic datasets. In each dataset, every

original integrated signal intensity was perturbed by adding random noise drawn from a

Gaussian distribution with a mean of zero and a standard deviation equal to the error of that

data point. To determine the diffusion coefficient and associated error for each experiment

we used the mean and standard deviation of fits to these 1000 datasets. Samples contained

50 µM α-synuclein, 0 mM or 25 mM NaCl, in 100 mM ammonium acetate, pH 7.2, or 100

mM Tris-HCl, pH 7.2, at 24 °C.

ThT amyloid assembly kinetics

Kinetics of αS amyloid formation were monitored in a 96-well, non-binding, flat-bottom, half-

area microplate (Corning, USA; 10438082) containing one Teflon polyball (1/8" diameter;

Polysciences Europe, Eppelheim, Germany) per each well of sample. Samples of 100 µL

containing 100 µM aS with 20 µM Thioflavin T (ThT) in 100 mM ammonium acetate, pH 7.2

and 100 mM Tris-HCl with 0 mM NaCl and 25 mM NaCl were incubated at 37 °C shaking at

600 rpm in a FLUOstar omega plate reader (BMG Labtech, Ortenburg, Germany).

Fluorescence intensity was measured by exciting at 440 ± 10 nm and collecting emission at

482 ± 12 nm using a bandpass filter. Results were blank corrected and normalised to the

maximum fluorescence value of each curve.

Coarse-grained simulations

Coarse-grained implicit-solvent simulations of αS were performed in the in the OpenMM

framework (version 8.1.1)[108] using the CALVADOS python package and the CALVADOS2

force field[93,94]. Unless otherwise stated, simulation parameters used were kept as their

default values in the software. The N terminal positive charge was removed to match the N-

terminally acetylated sample used in the MS and NMR experiments. Simulations were

performed in a 30 nm^3 box at pH 7.5. Following energy minimization, simulations were run at

a temperature of 25 °C. Simulations were performed for a total of 500 ns each with

coordinates being saved every 0.5 ns. The simulation had a time step of 10 fs and a friction

coefficient of 10 fs^{-1}. The first 5 ns of each simulation was considered to be part of the

equilibration time and therefore discarded. In total, 10 simulations were carried out at ionic

strengths of 10, 15, 20, 25, 35, 45, 60, 80, 110 and 150 mM. The R_g was calculated using

the inbuilt function in the CALVADOS software for each of the 990 frames outside the

equilibration time, with the error bars representing the standard error of the mean.

REFERENCES

[1] S. E. Bondos, A. K. Dunker, V. N. Uversky, *Cell Commun. Signal.* **2022**, *20*, 20.

[2] E. L. Sipko, G. F. Chappell, R. B. Berlow, *Curr. Opin. Struct. Biol.* **2024**, *84*, 102742.

[3] J. Cornish, S. G. Chamberlain, D. Owen, H. R. Mott, *Biochem. Soc. Trans.* **2020**, *48*, 2669–2689.

[4] R. Chen, X. Li, Y. Yang, X. Song, C. Wang, D. Qiao, *Front Mol Biosci* **2022**, *9*, 985022.

[5] D. Moses, F. Yu, G. M. Ginell, N. M. Shamoon, P. S. Koenig, A. S. Holehouse, S. Sukenik, *J. Phys. Chem. Lett.* **2020**, *11*, 10131–10136.

[6] P. Santofimia-Castaño, B. Rizzuti, Y. Xia, O. Abian, L. Peng, A. Velázquez-Campoy, J. L. Neira, J. Iovanna, *Cell. Mol. Life Sci.* **2020**, *77*, 1695–1707.

[7] V. N. Uversky, *Int. J. Mol. Sci.* **2016**, *17*, DOI 10.3390/ijms17111874.

[8] W. T. Astbury, S. Dickinson, K. Bailey, *Biochem. J* **1935**, *29*, 2351-2360.1.

[9] C. M. Dobson, *Semin. Cell Dev. Biol.* **2004**, *15*, 3–16.

[10] S. Linse, T. Knowles, *Chem. Sci.* **2023**, *14*, 6491–6492.

[11] J. Parkinson, *J. Neuropsychiatry Clin. Neurosci.* **2002**, *14*, 223–236.

[12] M. H. Polymeropoulos, J. J. Higgins, L. I. Golbe, W. G. Johnson, S. E. Ide, G. Di Iorio, G. Sanges, E. S. Stenroos, L. T. Pho, A. A. Schaffer, A. M. Lazzarini, R. L. Nussbaum, R. C. Duvoisin, *Science* **1996**, *274*, 1197–1199.

[13] M. H. Polymeropoulos, C. Lavedan, E. Leroy, S. E. Ide, A. Dehejia, A. Dutra, B. Pike, H. Root, J. Rubenstein, R. Boyer, E. S. Stenroos, S. Chandrasekharappa, A. Athanassiadou, T. Papapetropoulos, W. G. Johnson, A. M. Lazzarini, R. C. Duvoisin, G. Di Iorio, L. I. Golbe, R. L. Nussbaum, *Science* **1997**, *276*, 2045–2047.

[14] L. I. Golbe, G. Di Iorio, V. Bonavita, D. C. Miller, R. C. Duvoisin, *Ann. Neurol.* **1990**, *27*, 276–282.

[15] M. G. Spillantini, M. L. Schmidt, V. M. Lee, J. Q. Trojanowski, R. Jakes, M. Goedert, *Nature* **1997**, *388*, 839–840.

[16] B. Holdorff, *J. Hist. Neurosci.* **2002**, *11*, 19–28.

[17] R. M. Meade, D. P. Fairlie, J. M. Mason, *Mol. Neurodegener.* **2019**, *14*, 29.

[18] R. Sternke-Hoffmann, X. Sun, A. Menzel, M. Dos Santos Pinto, U. Venclovaitė, M. Wördehoff, W. Hoyer, W. Zheng, J. Luo, *bioRxiv* **2024**, 2024.03.01.582895.

[19] S. Fujiwara, F. Kono, T. Matsuo, Y. Sugimoto, T. Matsumoto, A. Narita, K. Shibata, *J. Mol. Biol.* **2019**, *431*, 3229–3245.

[20] L. A. Munishkina, J. Henriques, V. N. Uversky, A. L. Fink, *Biochemistry* **2004**, *43*, 3289–3300.

[21] O. Ullman, C. K. Fisher, C. M. Stultz, *J. Am. Chem. Soc.* **2011**, *133*, 19536–19546.

[22] S. J. Roeters, A. Iyer, G. Pletikapić, V. Kogan, V. Subramaniam, S. Woutersen, *Sci. Rep.* **2017**, *7*, 41051.

[23] J. Gallardo, C. Escalona-Noguero, B. Sot, *ACS Chem. Neurosci.* **2020**, *11*, 872–879.

[24] R. Ramis, J. Ortega-Castro, B. Vilanova, M. Adrover, J. Frau, *Biomacromolecules* **2020**, *21*, 5200–5212.

[25] A. D. Stephens, J. Kölbel, R. Moons, C. W. Chung, M. T. Ruggiero, N. Mahmoudi, T. A. Shmool, T. M. McCoy, D. Nietlispach, A. F. Routh, F. Sobott, J. A. Zeitler, G. S. Kaminski Schierle, *Angew. Chem. Int. Ed Engl.* **2023**, *62*, e202212063.

[26] M. Bisaglia, I. Tessari, S. Mammi, L. Bubacco, *Neuromolecular Med.* **2009**, *11*, 239–251.

[27] B. Gardner, B. V. Dieriks, S. Cameron, L. H. S. Mendis, C. Turner, R. L. M. Faull, M. A. Curtis, *Sci. Rep.* **2017**, *7*, 10454.

[28] V. N. Uversky, J. Li, A. L. Fink, *J. Biol. Chem.* **2001**, *276*, 44284–44296.

[29] A. Binolfi, R. M. Rasia, C. W. Bertoncini, M. Ceolin, M. Zweckstetter, C. Griesinger, T. M. Jovin, C. O. Fernández, *J. Am. Chem. Soc.* **2006**, *128*, 9893–9901.

[30] A. Natalello, F. Benetti, S. M. Doglia, G. Legname, R. Grandori, *Proteins: Struct. Funct.*
Bioinf. **2011**, *79*, 611–621.
[31] M. C. Miotto, A. A. Valiente-Gabioud, G. Rossetti, M. Zweckstetter, P. Carloni, P.
Selenko, C. Griesinger, A. Binolfi, C. O. Fernández, *J. Am. Chem. Soc.* **2015**, *137*,
6444–6447.
[32] R. J. Mason, A. R. Paskins, C. F. Dalton, D. P. Smith, *Biochemistry* **2016**, *55*, 4737–
4741.
[33] P. Wongkongkathep, J. Y. Han, T. S. Choi, S. Yin, H. I. Kim, J. A. Loo, *J. Am. Soc.*
Mass Spectrom. **2018**, *29*, 1870–1880.
[34] A. D. Stephens, M. Zacharopoulou, G. S. Kaminski Schierle, *Trends Biochem. Sci.*
2019, *44*, 453–466.
[35] R. Moons, A. Konijnenberg, C. Mensch, R. Van Elzen, C. Johannessen, S. Maudsley,
A.-M. Lambeir, F. Sobott, *Sci. Rep.* **2020**, *10*, 16293.
[36] E. J. Byrd, M. Wilkinson, S. E. Radford, F. Sobott, *J. Am. Soc. Mass Spectrom.* **2023**,
34, 493–504.
[37] M. Gonzalez-Garcia, G. Fusco, A. De Simone, *Front Chem* **2023**, *11*, 1167766.
[38] C. Lantz, J. Lopez, A. K. Goring, M. A. Zenaidee, K. Biggs, J. P. Whitelegge, R. R.
Ogorzalek Loo, F.-G. Klärner, T. Schrader, G. Bitan, J. A. Loo, *J. Am. Soc. Mass*
Spectrom. **2023**, *34*, 2739–2747.
[39] N. Seetaloo, M. Zacharopoulou, A. D. Stephens, G. S. Kaminski Schierle, J. J. Phillips,
Anal. Chem. **2022**, *94*, 16711–16719.
[40] A. D. Stephens, M. Zacharopoulou, R. Moons, G. Fusco, N. Seetaloo, A. Chiki, P. J.
Woodhams, I. Mela, H. A. Lashuel, J. J. Phillips, A. De Simone, F. Sobott, G. S. K.
Schierle, *Nat. Commun.* **2020**, *11*, 2820.
[41] J. Santos, V. Iglesias, J. Santos-Suárez, M. Mangiagalli, S. Brocca, I. Pallarès, S.
Ventura, *Cells* **2020**, *9*, 145.
[42] W. Hoyer, T. Antony, D. Cherny, G. Heim, T. M. Jovin, V. Subramaniam, *J. Mol. Biol.*
2002, *322*, 383–393.
[43] F. W. Karasek, *Anal. Chem.* **1974**, *46*, 710A-720A.
[44] K. Giles, S. D. Pringle, K. R. Worthington, D. Little, J. L. Wildgoose, R. H. Bateman,
Rapid Commun. Mass Spectrom. **2004**, *18*, 2401–2414.
[45] D. P. Smith, T. W. Knapman, I. Campuzano, R. W. Malham, J. T. Berryman, S. E.
Radford, A. E. Ashcroft, *Eur. J. Mass Spectrom.* **2009**, *15*, 113–130.
[46] M. F. Bush, Z. Hall, K. Giles, J. Hoyes, C. V. Robinson, B. T. Ruotolo, *Anal. Chem.*
2010, *82*, 9557–9565.
[47] A. Marchand, S. Livet, F. Rosu, V. Gabelica, *Anal. Chem.* **2017**, *89*, 12674–12681.
[48] V. Gabelica, A. A. Shvartsburg, C. Afonso, P. Barran, J. L. P. Benesch, C. Bleiholder,
M. T. Bowers, A. Bilbao, M. F. Bush, J. L. Campbell, I. D. G. Campuzano, T. Causon, B.
H. Clowers, C. S. Creaser, E. De Pauw, J. Far, F. Fernandez-Lima, J. C. Fjeldsted, K.
Giles, M. Groessl, C. J. Hogan Jr, S. Hann, H. I. Kim, R. T. Kurulugama, J. C. May, J. A.
McLean, K. Pagel, K. Richardson, M. E. Ridgeway, F. Rosu, F. Sobott, K. Thalassinos,
S. J. Valentine, T. Wyttenbach, *Mass Spectrom. Rev.* **2019**, *38*, 291–320.
[49] R. Beveridge, A. N. Calabrese, *Front Chem* **2021**, *9*, 603639.
[50] M. C. S. Kit, I. Webb, *ChemRxiv* **2023**, DOI 10.26434/chemrxiv-2023-bb6p4.
[51] J. B. Fenn, *Angew. Chem. Int. Ed Engl.* **2003**, *42*, 3871–3894.
[52] R. Grandori, C. Santambrogio, S. Brocca, G. Invernizzi, M. Lotti, *Biotechnol. J.* **2009**, *4*,
73–87.
[53] S. Banerjee, S. Mazumdar, *Int. J. Anal. Chem.* **2012**, *2012*, 282574.
[54] H. J. Sterling, J. D. Batchelor, D. E. Wemmer, E. R. Williams, *J. Am. Soc. Mass*
Spectrom. **2010**, *21*, 1045–1049.
[55] H. Metwally, R. G. McAllister, L. Konermann, *Anal. Chem.* **2015**, *87*, 2434–2442.
[56] C. A. Cassou, E. R. Williams, *Analyst* **2014**, *139*, 4810–4819.
[57] A. C. Susa, Z. Xia, E. R. Williams, *Angew. Chem. Int. Ed Engl.* **2017**, *56*, 7912–7915.
[58] T. G. Flick, C. A. Cassou, T. M. Chang, E. R. Williams, *Anal. Chem.* **2012**, *84*, 7511–
7517.

[59] Z. Xia, J. B. DeGrandchamp, E. R. Williams, *Analyst* **2019**, *144*, 2565–2573.
[60] S. Tamara, M. A. den Boer, A. J. R. Heck, *Chem. Rev.* **2022**, *122*, 7269–7326.
[61] Z. L. VanAernum, F. Busch, B. J. Jones, M. Jia, Z. Chen, S. E. Boyken, A.
Sahasrabuddhe, D. Baker, V. H. Wysocki, *Nat. Protoc.* **2020**, *15*, 1132–1157.
[62] J. Eisert, E. V. S. Maciel, V. Dederer, A. Berwanger, H. J. Bailey, I. Đikić, S. Knapp, M.
Empting, S. Mathea, H. Jensen, F. Lermyte, *Small Methods* **2025**, e2500658.
[63] E. M. Panczyk, J. D. Gilbert, G. S. Jagdale, A. Q. Stiving, L. A. Baker, V. H. Wysocki,
Anal. Chem. **2020**, *92*, 2460–2467.
[64] J. S. Jordan, Z. Xia, E. R. Williams, *J. Am. Soc. Mass Spectrom.* **2022**, *33*, 607–611.
[65] N. Drachman, M. Lepoitevin, H. Szapary, B. Wiener, W. Maulbetsch, D. Stein, *Nat.*
Commun. **2024**, *15*, 7709.
[66] A. C. Susa, J. L. Lippens, Z. Xia, J. A. Loo, I. D. G. Campuzano, E. R. Williams, *J. Am.*
Soc. Mass Spectrom. **2018**, *29*, 203–206.
[67] D. Stuchfield, P. Barran, *Curr. Opin. Chem. Biol.* **2018**, *42*, 177–185.
[68] C. Eldrid, A. Ben-Younis, J. Ujma, H. Britt, T. Cragnolini, S. Kalfas, D. Cooper-
Shepherd, N. Tomczyk, K. Giles, M. Morris, R. Akter, D. Raleigh, K. Thalassinos, *J. Am.*
Soc. Mass Spectrom. **2021**, *32*, 1545–1552.
[69] J. Ujma, J. Jhingree, E. Norgate, R. Upton, X. Wang, F. Benoit, B. Bellina, P. Barran,
Anal. Chem. **2022**, *94*, 12248–12255.
[70] Z. Dai, A. Ben-Younis, A. Vlachaki, D. Raleigh, K. Thalassinos, *Biophys. Chem.* **2024**,
312, 107285.
[71] E. Illes-Toth, C. F. Dalton, D. P. Smith, *J. Am. Soc. Mass Spectrom.* **2013**, *24*, 1346–
1354.
[72] R. Moons, R. van der Wekken-de Bruijne, S. Maudsley, F. Lemière, A.-M. Lambeir, F.
Sobott, *Int. J. Mol. Sci.* **2020**, *21*, DOI 10.3390/ijms21217884.
[73] M. Cheung See Kit, T. C. Cropley, C. Bleiholder, C. D. Chouinard, F. Sobott, I. K. Webb,
Analyst **2023**, *149*, 125–136.
[74] A. Depraz Depland, S. Mikromanolis, I. Stroganova, A. Rijs, *ChemRxiv* **2025**, DOI
10.26434/chemrxiv-2025-n1zwv.
[75] M. Karas, U. Bahr, T. Dülcks, *Fresenius J. Anal. Chem.* **2000**, *366*, 669–676.
[76] A. Schmidt, M. Karas, T. Dülcks, *J. Am. Soc. Mass Spectrom.* **2003**, *14*, 492–500.
[77] E. M. Yuill, N. Sa, S. J. Ray, G. M. Hieftje, L. A. Baker, *Anal. Chem.* **2013**, *85*, 8498–
8502.
[78] G. T. H. Nguyen, T. N. Tran, M. N. Podgorski, S. G. Bell, C. T. Supuran, W. A. Donald,
ACS Cent Sci **2019**, *5*, 308–318.
[79] V. N. Uversky, *Curr. Alzheimer Res.* **2008**, *5*, 260–287.
[80] A. S. Holehouse, S. Sukenik, *J. Chem. Theory Comput.* **2020**, *16*, 1794–1805.
[81] G. Heesink, M. J. Marseille, M. A. A. Fakhree, M. D. Driver, K. A. van Leijenhorst-
Groener, P. R. Onck, C. Blum, M. M. A. E. Claessens, *Biomacromolecules* **2023**, *24*,
3680–3688.
[82] C. Santambrogio, A. Natalello, S. Brocca, E. Ponzini, R. Grandori, *Proteomics* **2019**, *19*,
e1800060.
[83] P. Lössl, J. Snijder, A. J. R. Heck, *J. Am. Soc. Mass Spectrom.* **2014**, *25*, 906–917.
[84] K. J. Lee, J. S. Jordan, E. R. Williams, *Anal. Chem.* **2024**, *96*, 17586–17593.
[85] I. A. Kaltashov, R. R. Abzalimov, *J. Am. Soc. Mass Spectrom.* **2008**, *19*, 1239–1246.
[86] M. L. Abramsson, C. Sahin, J. T. S. Hopper, R. M. M. Branca, J. Danielsson, M. Xu, S.
A. Chandler, N. Österlund, L. L. Ilag, A. Leppert, J. Costeira-Paulo, L. Lang, K. Teilum,
A. Laganowsky, J. L. P. Benesch, M. Oliveberg, C. V. Robinson, E. G. Marklund, T. M.
Allison, J. R. Winther, M. Landreh, *JACS Au* **2021**, *1*, 2385–2393.
[87] H. Osterholz, A. Stevens, M. L. Abramsson, D. Lama, K. Brackmann, A. Rising, A.
Elofsson, E. G. Marklund, S. Deindl, A. Leppert, M. Landreh, *JACS Au* **2025**, *5*, 281–
290.
[88] D. Brooke, N. Movahed, B. Bothner, *AIMS Biophys* **2015**, *2*, 336–342.
[89] G. J. Pielak, *Biochemistry* **2021**, *60*, 3436–3440.

[90] K. Araki, N. Yagi, R. Nakatani, H. Sekiguchi, M. So, H. Yagi, N. Ohta, Y. Nagai, Y. Goto, H. Mochizuki, *Sci. Rep.* **2016**, *6*, 30473.

[91] M. Runfola, A. De Simone, M. Vendruscolo, C. M. Dobson, G. Fusco, *Sci. Rep.* **2020**, *10*, 204.

[92] L. Kang, G. M. Moriarty, L. A. Woods, A. E. Ashcroft, S. E. Radford, J. Baum, *Protein Sci.* **2012**, *21*, 911–917.

[93] G. Tesei, T. K. Schulze, R. Crehuet, K. Lindorff-Larsen, *Proc. Natl. Acad. Sci. U. S. A.* **2021**, *118*, e2111696118.

[94] G. Tesei, K. Lindorff-Larsen, *Open Res. Eur.* **2022**, *2*, 94.

[95] A. D. Stephens, M. Zacharopoulou, R. Moons, G. Fusco, N. Seetaloo, A. Chiki, P. J. Hooper, I. Mela, H. A. Lashuel, J. J. Phillips, A. D. Simone, F. Sobott, G. S. K. Schierle, *bioRxiv* **2019**, 740241.

[96] C. S. Johnson Jr, *Prog. Nucl. Magn. Reson. Spectrosc.* **1999**, *34*, 203–256.

[97] P. S. Kandiyal, J. Y. Kim, D. L. Fortunati, K. H. Mok, *Methods Mol. Biol.* **2019**, *2039*, 173–183.

[98] F. Havemeister, M. Ghaeidamini, E. K. Esbjörner, *Biochem. Biophys. Res. Commun.* **2023**, *679*, 31–36.

[99] Y. R. Dahal, J. D. Schmit, *Biophys. J.* **2018**, *114*, 76–87.

[100] R. Gaspar, M. Lund, E. Sparr, S. Linse, *QRB Discov.* **2020**, *1*, e2.

[101] S. Al-Harthi, V. Kharchenko, P. Mandal, S. Gourdoupis, Ł. Jaremko, *Int. J. Biol. Macromol.* **2022**, *222*, 2878–2887.

[102] R. Guerrero-Ferreira, N. M. Taylor, A.-A. Arteni, P. Kumari, D. Mona, P. Ringler, M. Britschgi, M. E. Lauer, A. Makky, J. Verasdonck, R. Riek, R. Melki, B. H. Meier, A. Böckmann, L. Bousset, H. Stahlberg, *Elife* **2019**, *8*, DOI 10.7554/eLife.48907.

[103] C. Lübbert, W. Peukert, *Anal. Chem.* **2021**, *93*, 12862–12871.

[104] J. S. Jordan, E. R. Williams, *Anal. Chem.* **2021**, *93*, 1725–1731.

[105] C. R. Harris, K. J. Millman, S. J. van der Walt, R. Gommers, P. Virtanen, D. Cournapeau, E. Wieser, J. Taylor, S. Berg, N. J. Smith, R. Kern, M. Picus, S. Hoyer, M. H. van Kerkwijk, M. Brett, A. Haldane, J. F. Del Río, M. Wiebe, P. Peterson, P. Gérard-Marchant, K. Sheppard, T. Reddy, W. Weckesser, H. Abbasi, C. Gohlke, T. E. Oliphant, *Nature* **2020**, *585*, 357–362.

[106] P. Virtanen, R. Gommers, T. E. Oliphant, M. Haberland, T. Reddy, D. Cournapeau, E. Burovski, P. Peterson, W. Weckesser, J. Bright, S. J. van der Walt, M. Brett, J. Wilson, K. J. Millman, N. Mayorov, A. R. J. Nelson, E. Jones, R. Kern, E. Larson, C. J. Carey, İ. Polat, Y. Feng, E. W. Moore, J. VanderPlas, D. Laxalde, J. Perktold, R. Cimrman, I. Henriksen, E. A. Quintero, C. R. Harris, A. M. Archibald, A. H. Ribeiro, F. Pedregosa, P. van Mulbregt, SciPy 1.0 Contributors, *Nat. Methods* **2020**, *17*, 261–272.

[107] "Hunter, J.D. (2007) Matplotlib A 2D Graphics Environment. Computing in Science & Engineering, 9, 90-95. - References - Scientific Research Publishing," can be found under https://www.scirp.org/reference/referencespapers?referenceid=1560579, **n.d.**

[108] P. Eastman, J. Swails, J. D. Chodera, R. T. McGibbon, Y. Zhao, K. A. Beauchamp, L.-P. Wang, A. C. Simmonett, M. P. Harrigan, C. D. Stern, R. P. Wiewiora, B. R. Brooks, V. S. Pande, *PLoS Comput. Biol.* **2017**, *13*, e1005659.

<div align="center">**Supplementary information**</div>

Buffer-dependent conformational dynamics of α-synuclein revealed by

nanopipette electrospray ionisation ion mobility mass spectrometry

Emily J. Byrd[1]•, Emma L. Norgate[1]•, Joel A. Crossley[1], Chalmers C. C. Chau[2,3], Bob Schiffrin[1],

Alexander Kulak[4], Sheena E. Radford[1], Paolo Actis[2,3*], Antonio N. Calabrese[1*], Frank Sobott[1*]

[1]Astbury Centre for Structural Molecular Biology, School of Molecular and Cellular Biology, Faculty of

Biological Sciences, University of Leeds, Leeds, LS2 9JT, UK.

[2]School of Electronic and Electrical Engineering, University of Leeds, LS2 9JT, UK.

[3]Bragg Centre for Materials Research, University of Leeds, LS2 9JT, UK.

[4]School of Chemistry, University of Leeds, LS2 9JT, UK.

• These authors contributed equally to this work.

*Correspondence to: f.sobott@leeds.ac.uk, a.calabrese@leeds.ac.uk, p.actis@leeds.ac.uk

Figure S1. Fabrication of the nESI emitter tips. *The typical fabrication process for the standard nESI emitter tip (a) starts with a pre-pulled borosilicate glass pipette with a long taper. The long taper is clipped with tweezers, resulting in a tip diameter in the 2 - 20 micrometre range. For the fabrication of the nanopipette nESI emitter tip (b), a laser puller is used with quartz glass capillaries. This process results in the fabrication of a pair of nearly identical nanopipette nESI emitter tips, each with a nanometre range sized pore.*

Figure S2. Comparison of nanopipette nESI emitters with standard nESI emitters. *SEM images of (a) standard borosilicate nESI emitters clipped to size with tweezers (pore diameter ~ 16 µm) and (b) of nanopipette emitters (pore diameter ~ 43 nm).*

Figure S3. Zoomed in native nESI mass spectra of the 8+ charge state of αS. *Measured*
in PBS, pH 7.2 using (a) standard borosilicate emitters and (b) quartz nanopipettes. The
predicted m/z of the apo 8+ αS charge state is 1813. The full mass spectra are shown in
Figure 1.

Figure S4. **Representative** $^{TW}CCS_{N2}$ **distribution of αS in PBS buffer using a**
nanopipette nESI emitter. *The data are taken from a spectrum acquired of 10 μM αS in*
PBS buffer, pH 7.2. The key on the righthand side indicates the colour of the CCS
distribution for each charge state, the black solid line is the cumulative fit and the black
dotted line represents the average $^{TW}CCS_{N2}$ *value.*

Figure S5. Native nESI mass spectra of αS in 20 mM AmAc with NaCl titration. *αS (20 μM) was measured in (a) 20 mM AmAc, pH 7.2 with the addition of (b) 12.5 mM NaCl, (c) 25 mM NaCl, (d) 125 mM NaCl.*

Figure S6. Native nESI mass spectra of αS in 20 mM Tris-HCl with NaCl titration. *αS (20 μM) was measured in (a) 20 mM Tris-HCl, pH 7.2 with the addition of (b) 12.5 mM NaCl, (c) 25 mM NaCl, (d) 125 mM NaCl.*

Figure S7 Native nESI mass spectra of the 8+ charge state. *αS measured in (a) 20 mM*
AmAc, (b) 20 mM AmAc with 12.5 mM NaCl, (c) 20 mM Tris-HCl and (d) 20 mM Tris-HCl
with 12.5 mM NaCl. All measurements were taken at pH 7.2.

Figure S8. $^{TW}CCS_{N2}$ **Ion mobility distributions of lysozyme using nanopipette nESI**
emitters. *nESI mass spectra were acquired of 10 µM hen egg white lysozyme in (a) 100 mM*
AmAc, (b) 100 mM AmAc, pH 7.2 with 25 mM NaCl, (c) 100 mM Tris-HCl and (d) 100 mM
Tris with 25 mM NaCl.

Figure S9 Distributions of calculated Rg values for αS from coarse grained molecular dynamics simulations. *Each panel shows a different ionic strength value for the simulation. The mean R_g is shown as a vertical black dashed line. For details on the simulation and number of frames used, see Methods.*

793
794

Figure S10. Example 1D spectra of αS from DOSY NMR experiment.s *For each DOSY experiment under each sample condition, a total of 24 1D acquisitions were performed at different gradient strengths between 2% and 98%, where the maximum gradient strength was 48.148 G cm^{-1}. Data for the region used for integration (0.6-0.9 ppm) is shown. The sample contained 50 μM αS in 100 mM Tris-HCl, pH 7.2, at 24 °C.*

Ionic Strength	R_{app} (Å)	% change	R_g	% change	Diffusion coefficient × 10⁻□ (cm² s⁻¹)	% change
20 mM	28.14	8%	32.87	11%	8.99 ± 0.05	16.2%
125 mM	30.27		36.48		7.53 ± 0.04	

Table S1 A table comparing measurements of αS expansion at 20 mM ionic strength
and 125 mM ionic strength. *R_{app} in Å was calculated from the average $^{TW}CCS_{N2}$ values for*
αS in 20 mM AmAc, pH 7.2 and 100 mM AmAc with 25 mM NaCl, pH 7.2 (See methods for
calculation). R_g was calculated from coarse grained CALVADOS simulations of αS at 20 mM
and 125 mM ionic strength, pH 7.5. Diffusion coefficient (cm^2/s) were measured using DOSY
NMR. The percent change for each measurement represents an increase.