Herbivorous insects independently evolved salivary effectors to regulate plant immunity by destabilizing the malectin-LRR RLP NtRLP4

Xin Wang[1,2], Jia-Bao Lu[1], Yi-Zhe Wang[1], Xu-Hong Zhou[3], Jian-Ping Chen[1,2], Chuan-Xi Zhang[1], Jun-Min Li[1,#], Hai-Jian Huang[1,#]

[1] State Key Laboratory for Quality and Safety of Agro-Products, Key Laboratory of Biotechnology in Plant Protection of MARA, Zhejiang Key Laboratory of Green Plant Protection, Institute of Plant Virology, Ningbo University, Ningbo 315211, China

[2] Institute of Applied Ecology, Fujian Agriculture and Forestry University, Fuzhou 350002, China

[3] Office of Science and Technology, Yunnan University of Chinese Medicine, Kunming 650205, China

Abstract

Plants rely on receptor-like proteins and receptor-like kinases (RLPs/RLKs) to detect a variety of invading pathogens and insects. However, it remains unclear how these invaders are able to overcome receptor perception and disable the plant signaling pathways. Here, we show that whiteflies and planthoppers independently evolve salivary effectors to target defensive RLP4, therefore minimizing the plant immunity. RLP4 is a leucine-rich repeat RLP (LRR-RLP) that confers plant resistance against herbivorous insects by forming RLP4/SOBIR1 complexes. In whitefly *Bemisia tabaci*, BtRDP is an Aleyrodidae-restricted protein that attaches to salivary sheath. It interacts with RLP4 from different plants, causing the degradation of RLP4 in an ubiquitin-dependent manner. Transgenic plants overexpressing NtRLP4 exerts a detrimental impact on *B. tabaci* by exploiting the crosstalk between the salicylic acid (SA) and jasmonic acid (JA) pathways. Conversely, the overexpression of BtRDP or silencing of NtRLP4 effectively alleviates such negative effects. In planthopper *Nilaparvata lugens*, a Delphacidae-restricted salivary protein NlSP694 also targets and promotes degradation of OsRLP4 from rice plants. Our results reveal the

* Corresponding authors: Jun-Min Li, lijunmin@nbu.edu.cn; Hai-Jian Huang, huanghaijian@nbu.edu.cn

convergent evolution of salivary proteins in insects, and shed lights on the complex interactions

between plants and herbivorous insects.

Keywords

salivary effector, insect-plant interaction, receptor-like protein, convergent evolution

Introduction

The activation of innate immunity by membrane-localized receptors is a highly conserved

mechanism in eukaryotes [1]. In sessile plants, the first line of defense against potential pathogenic

microbes or insects relies on cell surface pattern-recognition receptors (PRRs) [2]. The plant genome

encodes hundreds of PRRs, including receptor-like kinases (RLKs) and receptor-like proteins (RLPs)

[2]. RLKs consist of different types of N-terminal ectodomains, a single transmembrane domain,

and a cytoplasmic protein kinase domain, while RLPs have a similar structure but lack the kinase

domain [3, 4]. As RLPs lack intracellular signaling domains, they are usually associated with

SOBIR1 or SOBIR1-like adaptors to form bimolecular receptor kinases [4]. Despite the large

number of PRRs encoded by plant genomes, only a few of them have been well characterized in

terms of their function in recognizing microbial-associated molecular patterns (MAMPs), damage-

associated molecular patterns (DAMPs), and herbivore-associated molecular patterns (HAMPs) [3,

5]. For instance, FLS2 is known to recognize bacterial flagellin [6], EFR can recognize bacterial

elongation factor Tu [7], and EIX2 is responsible for recognizing the fungal ethylene-inducing

xylanase [8].

Piercing-sucking insects, such as whiteflies, aphids, and planthoppers, damage plants by

feeding or transmitting viruses. They inject a mixture of saliva into plant tissues, which aids their

performance by binding to the leaked calcium, degrading extracellular DNA, and regulating plant

hormonal signals [9-12]. In response, plants have evolved the ability to detect herbivores by sensing

HAMPs present in insect saliva [5, 13]. This triggers pattern-triggered immunity (PTI), including

mitogen-activated protein kinase (MAPK) cascades, reactive oxygen species (ROS), and hormone

signaling [14-16]. Although a few salivary proteins have been reported to activate plant PTI, the

specific PRRs that recognize these proteins and trigger plant immunity are still not well understood

[17-19]. One exception is the inceptin receptor (INR), which is a legume-specific LRR-RLP

recognizing the Vu-In in insect oral secretions [5]. Moreover, plant PRRs have been widely reported

to be targeted by pathogen effectors [20-22]. It remains unknown how insects cope with the immune

recognition system.

The whitefly *Bemisia tabaci* (Hemiptera: Aleyrodidae) and brown planthopper *Nilaparvata*

lugens (Hemiptera: Delphacidae) are notorious pests globally. During the feeding process, both

insects secrete abundant species-specific salivary proteins into host plants [23, 24]. The plants have

been reported to initiate defense responses when they recognize insect feeding [17, 25]. However,

the specific PRRs responsible for this process are unknown. Both insects can successfully ingest

phloem saps with limited plant defenses. It remains unclear how insects attenuate plant defenses.

Given that many defense mechanisms are conserved across plants [1, 26], it is interesting to

investigate whether insect evolves a similar strategy to aid feeding in convergent evolution. In this

study, we characterize the function of RLPs in plant immunity responsive to piercing-sucking insects,

and demonstrate that *B. tabaci* and *N. lugens* independently evolve salivary proteins to regulate the

posttranslational modification of RLPs.

Results

BtRDP is most abundantly expressed in salivary glands and secreted into host plants

According to the transcriptomic results, BtRDP (*B. tabaci* RLP-degrading protein; GenBank

accession: MN738093) is the most abundantly expressed gene in salivary glands [24]. It is a

secretory protein containing a signal peptide at the N-terminal, and totally 6 unique peptides from

BtRDP are detected in *B. tabaci* watery saliva (Fig. S1). As there is no gene homolog identified in

the NCBI nr database, BtRDP is subjected to BLAST analysis against the transcriptomic or genomic

data of 30 insect species. The results show that RDP is distributed in all the analyzed Aleyrodidae

species, while no RDP homolog is detected in other species (Table S1). Based on phylogenetic

analysis, RDPs from five *B. tabaci* cryptic species are clustered in the same clade, showing a close

relationship to RDP from *Aleurocanthus spiniferus* (Fig. S2).

BtRDP can be secreted into host plants, with two bands between 25-35kDa being detected in

Nicotiana tabacum plants infested by *B. tabaci*, but not in the non-infested plants (Fig. 1a). The

smaller high intensity band is the same size as that of BtRDP in salivary glands, and always presents

in the *B. tabaci*-infested plants. For the larger lower intensity band, it presents in infested samples

with lower frequency. It is possible that the larger band might be the BtRDP that undergoes

modification after being secreted into plants, similar with another salivary effector previously

reported [27]. However, we cannot exclude the possibility that this larger band is the non-specific

N. tabacum protein induced by *B. tabaci* feeding. Western-blotting, qPCR and transcriptomic

analysis demonstrate that BtRDP is nearly exclusively expressed in the salivary glands (Fig. 1b, c;

Fig. S3a), which is significantly different from the actin or ribosomal 18S rRNA (Fig. S4a). In

addition, BtRDP is mainly expressed in the adult stages, but lowly expressed in the egg, nymph, and

pseudopupa stages (Fig. 1d; Fig. S3b), similar to another salivary protein BtFTSP [28] but different

from actin or ribosomal 18S rRNA (Fig. S4b). Immunohistochemical (IHC) staining shows that

BtRDP is specifically distributed in principle salivary gland, but not in the accessory salivary gland

or other tissues (Fig. 1e). A few salivary proteins have been reported to attach to salivary sheath

after secretion [29]. We also stain the salivary sheath secreted by *B. tabaci* using anti-BtRDP serum.

The results show that BtRDP signal is mainly detected in the protuberant structure of sheath (Fig.

S5), indicating that BtRDP attaches to salivary sheath after secretion.

BtRDP is important for *B. tabaci* feeding on tobacco plants

RNA interference is firstly employed to investigate the potential role of BtRDP in whitefly

feeding, and ds*BtRDP* treatment efficiently and specifically suppresses the target gene at

transcriptional and protein level (Fig. 2a; Fig. S6). We do not observe a significant difference in

salivary sheath morphology between ds*BtRDP*- and ds*GFP*-treated *B. tabaci* (Fig. S7a). However,

the salivary sheath length of ds*BtRDP*-treated *B. tabaci* is significantly shorter than that of the

ds*GFP*-treated control (Fig. S7b). Moreover, ds*BtRDP* treatment exerts no influence on the whitefly

survivorship (Fig. S7c). For reproduction analysis, the ds*BtRDP*-treated whiteflies oviposit less eggs

than the control (Fig. 2b). The whitefly feeding behavior, which can be classified into

nonpenetration (np), pathway duration (C), phloem salivation (E1), and phloem ingestion (E2) [30]

(Fig. S8), is recorded using the electrical penetration graph (EPG) technique. Compared with ds*GFP*

treatment, ds*BtRDP*-treated *B. tabaci* exhibits a dramatic decrease in phloem ingestion (Fig. 2c).

Moreover, ds*BtRDP*-treated whitefly exhibits a significantly increased pathway duration phase and

a slightly increased nonpenetration phase (Fig. 2c).

Salivary proteins are reported to function in intercellular space and/or extracellular space after

secreted into host plants [29, 31]. To investigate the role of BtRDP in different subcellular location
of host plants, we construct two transgenic *N. tabacum* overexpressing BtRDP, one with a complete
coding region (oeBtRDP) and the other without signal peptide (oeBtRDP[-sp]). The signal peptide of
BtRDP may enable protein secretion into the extracellular space. The flag tag is fused to the C-
terminal ends of recombinant proteins, respectively. The empty vector (EV) transgenic plants, which
exert no significant influence on insect performance compared with the WT (Fig. S9), is used as a
negative control. Fecundity assay demonstrates that both oeBtRDP and oeBtRDP[-sp] transgenic
plants are beneficial to whiteflies, with more eggs oviposited in these two overexpressing plants
compared with the EV controls (Fig. 2d-i; Fig. S10). Two-choice experiments exert similar results,
with more insects being settled and oviposited in oeBtRDP or oeBtRDP[-sp] transgenic plants than the
EV controls (Fig. 2d-i; Fig. S10). These results indicate that BtRDP is beneficial to whiteflies,
regardless of its subcellular location in host plants.

Transcriptomic sequencing is performed on EV, oeBtRDP#1, and oeBtRDP[-sp]#1 transgenic
plants. PCA analysis demonstrates that the biological replicates of each transgenic plant can be well
clustered (Fig. S11a). Totally 1253 and 1227 differentially expressed genes (DEGs) are identified
when comparing oeBtRDP#1 and oeBtRDP[-sp]#1 with EV plants, respectively (Table S2; Table S3).
Enrichment analysis of DEGs in these two comparison groups exerts similar results, with pathways
associated with MAPK signaling, environmental adaptation, signal transduction, and plant-
pathogen interaction significantly enriched (Fig. S10b, c). There are 464 DEGs overlap in these two
comparison groups (Fig. S11d). Interestingly, the majority of these DEGs exhibit similar expression
pattern in oeBtRDP#1 and oeBtRDP[-sp]#1 when compared with EV plants (Fig. S11e), suggesting
that BtRDP affect some plant pathways regardless of its subcellular location. There are 1084 DEGs
identified in comparison of oeBtRDP#1 and oeBtRDP[-sp]#1 plants (Table S4). Enrichment analysis
show that the DEGs associated with metabolism, diterpenoid biosynthesis, and plant-pathogen
interaction are significantly enriched (Fig. S12), suggesting that subcellular location of BtRDP also
has non-negligible influence on host plants.

BtRDP interacts with NtRLP4/NtSORBIR1 complex

To investigate the underlying mechanism of BtRDP in improving whitefly feeding, yeast two-
hybrid (Y2H) screening against a *N. benthamiana* cDNA library is conducted using BtRDP[-sp] as a

bait. Totally seven proteins are found to potentially interact with BtRDP^{-sp} (Table S5), including an

NbRLP4 (accession Niben261Chr07g1310001.1 in Niben261 genome). NbRLP4 contains a

predicted N-terminal signal peptide, a malectin-like domain, a LRR domain, followed by a

transmembrane domain. NbRLP4, as well as its closest homology in *N. tabacum* (NtRLP4;

GenBank accession: XM_016601109), does not contain the kinase domain (Fig. 3a).

The relative expression of *NtRLP4* in response to *B. tabaci* infestation is investigated. qRT-

RCR results demonstrate that the transcript level of *NtRLP4* is not significantly influenced by *B.*

tabaci (Fig. S13). However, a decrease in protein level of NtRLP4 is detected by western-blotting

assay (Fig. 3b). Point-to-point Y2H assays reveal that NtRLP4 without signal peptides and

transmembrane domains interacts with BtRDP^{-sp} (Fig. 3c). To investigate the specificity of NtRLP4-

BtRDP interaction, another RLP protein NtCf9 showing the highest sequence similarity to tomato

resistance protein Cf9 [32], as well as three randomly selected salivary proteins from *B. tabaci*, are

used as controls. The results show that BtRDP^{-sp} cannot interact with truncated NtCf9 without signal

peptides and transmembrane domains (Fig. 3c). Also, NtRLP4$_{(23-541)}$ cannot interact with the other

three salivary proteins without signal peptides (Fig. S14a, b). In addition, it is the LRR domain, but

not the malectin-like domain or the short intercellular region, that is required for the NtRLP4-BtRDP

interaction (Fig. 3c). To investigate the shortest LRR sequence responsible for BtRDP interaction,

LRR domain is divided into three segments. However, none of these segments interact with BtRDP^{-}

^{sp} (Fig. S14a). Co-immunoprecipitation (Co-IP) assays further confirm the specific interaction

between BtRDP and NtRLP4 (Fig. 3d). It is the C-terminal myc tag-fused NtRLP4 (NtRLP4-myc),

but not the NtCf9 (NtCf9-myc), that can be immunoprecipitated by C-terminal flag tag fused BtRDP

(BtRDP-flag). In bimolecular fluorescence complementation (BiFC) assay, YFP fluorescence is

observed when co-expressing the N-terminal nYFP tag fused NtRLP4 and N-terminal cYFP tag

fused BtRDP^{-sp} (Fig. S14c). Furthermore, oeBtRDP transgenic plants that overexpressing BtRDP-

flag are incubated with anti-flag beads. The results show that endogenous NtRLP4 can be

immunoprecipitated, which is significantly different from the control (Fig. S14d). These results

suggest that BtRDP interacts with NtRLP4.

The absence of kinase motifs in the C-terminal region of NtRLP4 suggests that it must function

in cooperation with other proteins to initiate the signaling. The receptor-like kinase SOBIR1, which

contained a kinase domain (Fig. S15a), is reported to be required for the function of several RLPs

involved in the innate immunity [4]. Through Co-IP assays, our study demonstrates that NtRLP4 is

capable of interacting with NtSOBIR1 (Fig. S15b).

NtRLP4 and NtSOBIR1 are resistant to *B. tabaci* infestation

The transgenic *N. tabacum* overexpressing NtRLP4 (oeRLP#1 and oeRLP#2, without any tags)

is constructed (Fig. 3e). Compared with the EV control, both oeRLP lines are less attractive to *B.*

tabaci, with less settling and ovipositing insects (Fig. 3f; Fig. S16a). Also, we observe the

significantly decreased fecundity of *B. tabaci* in oeRLP plants (Fig. 3g; Fig. S16b). Similar results

are observed after transient overexpression of NtRLP4-GFP in *N. tabacum* (Fig. S16c, d), indicating

that NtRLP4 is resistant to *B. tabaci* infestation.

Thereafter, the EV and oeRLP#1 plants are subjected to transcriptomic sequencing (Fig. 17a).

Totally 729 differentially expressed genes (DEGs) are identified, including 423 up-regulated and

306 down-regulated ones (Table S6). Enrichment analysis demonstrate that the majority of up-

regulated DEGs are associated with plant-pathogen interaction, environmental adaptation, MAPK

signaling pathway, and signal transduction (Fig. 17b). By contrast, the glutathione metabolism,

carbohydrate metabolism, and amino acid metabolism pathways are significantly enriched in down-

regulated DEGs (Fig. 17c). Noteworthily, there are many DEGs annotated as RLK/RLP or WRKY

transcription factor, with the majority of them being significantly upregulated (Fig. S17d, e). These

results suggest an increased defense in oeRLP plants. The crosstalk between salicylic acid (SA) and

jasmonic acid (JA) pathways is critical for plant defenses [12]. The increased defense of oeRLP

transgenic plant is partially caused by the altered hormonal signaling. The JA-associated genes

FAD7 is significantly induced, while the SA-associated genes *PAL* and *NPR1* are repressed in oeRLP

plants (Fig. 3h). Transient overexpression of NtRLP4-GFP in *N. tabacum* plants via *Agrobacterium*

infiltration exert similar results (Fig. S16e). Furthermore, NtRLP4-GFP significantly enhances the

H_2O_2 accumulation (Fig. S16f).

We also investigate the function of NtSOBIR1 by transient overexpression and hairpin RNAi (Fig.

S18). Overexpression of NtSOBIR1-GFP induces the cell death phenotype five days post agro-

injection (Fig. S18b). Insects prefer to settle and oviposit on GFP-expressed *N. tabacum* than on

NtSOBIR1-GFP-expressed control, even no obvious death phenotype is observed two days post

agro-injection (Fig. S18c, d). For NtSOBIR1 silencing, no cell death phenotype is observed, while

the NtSOBIR1-silenced leaves exhibit reduced growth (Fig. S18f). Compared with EV-silenced

control, NtSOBIR1-silenced *N. tabacum* is slightly attractive to *B. tabaci* (Fig. S18g, h).

Collectively, these results demonstrate that NtSOBIR1 is resistant to *B. tabaci* infestation.

BtRDP attenuates plant defenses by promoting the degradation of host RLP4

To investigate the interplay between NtRLP4 and BtRDP, we construct transgenic *N. tabacum*

silencing NtRLP4 (*RNAi-RLP*#1 and *RNAi-RLP*#2) (Fig. S19a). Compared with EV plants, *B.*

tabaci show a tendency to settle on NtRLP4-silenced plants and lay more eggs (Fig. S19b, c).

Thereafter, ds*GFP*- and ds*BtRDP*-treated *B. tabaci* are reared on EV, *RNAi-RLP*#1, and *RNAi-*

RLP#2, respectively. The ds*BtRDP*-treated *B. tabaci* produce less offspring than the control

(decreased by 28.3%; *P*=0.0007) on EV plants (Fig. 4a), consistent with the results on WT plants

(Fig. 2b). On *RNAi-RLP*#1 plant, the impair insect reproduction is still observed after ds*BtRDP*

treatment, but the extent is slightly reduced (decreased by 23.7%; *P*=0.002) (Fig. 4a). On *RNAi-*

RLP#2 plant, there is no statistical significance between ds*GFP* and ds*BtRDP* treatments (Fig. 4a).

These results suggest that *RNAi-RLP* plants are beneficial for insects and can partially, but not

completely, rescue the impaired feeding performance of ds*BtRDP*-treated insects.

To investigate the influence of BtRDP on NtRLP4, the GFP-tagged NtRLP4 (NtRLP4-GFP) is

transiently co-expressed with mCherry-tagged BtRDP (BtRDP-mCherry), red fluorescent protein

(RFP)-mCherry, and BtFTSP1-mCherry via agroinfiltration, respectively. All tags are fused to the

C-terminus. BtRDP-mCherry does not influence the location patterns of NtRLP4-GFP (Fig. S20).

However, a significant reduction in the fluorescent signal of NtRLP4-GFP is detected when co-

expressed with BtRDP-mCherry, but not the controls (Fig. S20). The co-expression of BtRDP-

mCherry/NtRLP4-GFP significantly reduces H_2O_2 accumulation (Fig. 4b) and suppresses the

expression of the JA-associated genes *FAD7* and *PDF.12*, while inducing the expression of the SA-

associated genes *PAL* and *NPR1* (Fig. 4c). Insect bioassays demonstrate that transient

overexpression of BtRDP-mCherry rescues the negative effects on whitefly performance caused by

NtRLP4-GFP overexpression (Fig. 4d, e). In host choice experiments, *B. tabaci* display a preference

for settling on *N. tabacum* leaves co-expressing BtRDP-mCherry/NtRLP4-GFP over those co-

expressing RFP-mCherry/NtRLP4-GFP (Fig. 4d). We observe almost no preference in *B. tabaci*

feeding on *N. tabacum* plants co-expressing BtRDP-mCherry/NtRLP4-GFP and plants expressing

GFP alone (Fig. S21).

The attenuated NtRLP4-GFP fluorescent signal (Fig. S20), along with the observed decrease

in the protein level of NtRLP4 during *B. tabaci* infestation (Fig. 3b), leads us to speculate on whether

BtRDP influences the accumulation of NtRLP4. To this end, the BtRDP^{-sp}-his and GFP-his proteins

are prokaryotic expressed in *Escherichia coli*. Different concentrations of purified BtRDP^{-sp}-his and

GFP-his proteins are infiltrated into *N. tabacum* plants overexpressing NtRLP4-myc. Interestingly,

the protein level of NtRLP4-myc shows a slight decrease with increasing amounts of BtRDP^{-sp}-his,

while no such effect is observed with GFP-his (Fig. S22). Moreover, the BtRDP^{-sp}-his protein does

not have significant effect on the accumulation of another RLP NtCf9-like (Fig. S22).

Then, we detect the NtRLP4 level in oeBtRDP and oeBtRDP^{-sp} plants. Compared with the EV

control, a reduced NtRLP4 abundance is observed in both oeBtRDP#1 and oeBtRDP#2 transgenic

lines, but not the oeBtRDP^{-sp} plants (Fig. 4f), suggesting the importance of BtRDP secretion in

extracellular space. BtRDP attaches to salivary sheath that localizes in plant apoplast (Fig. S5). The

presence of signal peptide promotes the secretion of BtRDP into extracellular space. Therefore, we

use the complete coding region of *BtRDP* in subsequent experiments. The NtRLP4-myc and

BtRDP-flag are transiently co-expressed in *N. benthamiana* plants through *Agrobacterium*

infiltration. The results show that BtRDP-flag significantly decreasing the protein level of NtRLP4-

myc, but not the NtCf9-myc (Fig. 4g). qRT-PCR analyses demonstrate that the transcript level of

NtRLP4 remains unchanged (Fig. S23). These findings indicate that the reduction in protein level

of NtRLP4 is not caused by transcriptional changes but rather to degradation.

The ubiquitin system and autophagy are two major pathways regulating protein degradation

[33]. At first, we employ MG132, a 26S proteasome inhibitor known to inhibit the degradation of

ubiquitin-conjugated proteins [34]. The results show that MG132 efficiently prevents the

degradation of NtRLP4-myc by BtRDP-flag (Fig. 4h). Then, we employ autophagy inhibitor 3-

Methyladenine (3-MA) and bafilomycin A1 (BAF) that can block autophagy in plants [35].

However, the decreased protein level of NtRLP4-myc is still observed in the presence of 3-MA or

BAF (Fig. S24). Additionally, NtRLP4-myc and HA-tagged ubiquitin (HA-UBQ) are co-expressed

in *N. benthamiana*. Total proteins are extracted from *N. benthamiana* leaves and then subjected to

immunoprecipitation with anti-myc beads. The ubiquitin attached to NtRLP4 is determined by

immunoblot using anti-HA antibody. Leaves expressing NtRLP4-myc or UBQ-HA along are used

as negative controls. The results show that a strong UBQ signal is detected in the

immunoprecipitated product only when NtRLP4-myc and UBQ-HA are co-expressed (Fig. 4i). The

molecular weight of UBQ-attached NtRLP4-myc is remarkably larger than NtRLP4-myc itself (Fig.

4i). These results suggest that BtRDP promotes NtRLP4 degradation through the ubiquitin system,

but not the autophagy pathway.

Degrading plant RLP4s by salivary effectors is a common strategy for insects

To investigate the ability of BtRDP to degrade RLPs from other host plants of *B. tabaci*,

Solanum lycopersicum RLP4 (SlRLP4; GenBank accession: XP_004232910) is selected for

analysis. Sequence analysis reveals a similar gene structure between SlRLP4 and NtRLP4 (Fig. S25;

Fig. S26a). The interaction between BtRDP and SlRLP4 is confirmed through Y2H and Co-IP

assays (Fig. S26b, c). Moreover, we transiently co-express SlRLP4-myc and SlCf9-like-myc with

varying concentrations of BtRDP-flag or GUS-flag, with the myc and flag tags fused to the C-

terminus. The results show that BtRDP-flag exerts no influence on the accumulation of SlCf9-like-

myc protein. By contrast, the protein level of SlRLP4-myc is significantly reduced in the presence

of BtRDP-flag (Fig. S26d), indicating the conserved function of BtRDP in *B. tabaci* adaptation to

different host plants.

BtRDP is an Aleyrodidae-restricted gene that does not have homologues in other species. To

explore whether RLP degradation by salivary effectors is conserved among different herbivorous

insects, a RLP4 homology from *Oryza sativa* (OsRLP4; GenBank accession: XP_015645303; Fig.

5a) and twenty salivary proteins from planthopper species without signal peptides are paired to

identify the potential interactions using Y2H assays. As a result, only the salivary protein NlSP694-

sp (GenBank accession: MF278694) interacts with OsRLP4$_{(29-551)}$ (OsRLP4 without signal peptides

and transmembrane domains), while the other salivary proteins analyzed do not (Fig. 5b, c; Fig.

S27). NlSP694 does not show any sequence or structure similarity to BtRDP (Fig. S28). Based on

truncation analysis, NlSP694-sp interacts with both the LRR domain and malectin-like domain of

OsRLP4 (Fig. 5c), slightly different from that of the BtRDP-NtRLP4 interaction (Fig. 3c).

Considering that both BtRDP and NlSP694 interact with the LRR domain of RLP4, the specific

region of LRR domain responsible for NlSP694 binding is further investigated. The LRR domain

of OsRLP4 is divided into three segments, in accordance with the LRR domain division in NtRLP4

(Fig. S14a). NlSP694^{-sp} interacts with the truncated LRR domain (OsRLP4$_{(480-526)}$) (Fig. 5c). By

contrast, none of the truncated LRR domain of NtRLP4 interact with BtRDP^{-sp} (Fig. S14a). These

results suggest that the binding sites of salivary proteins to RLP4 may be different between *N. lugens*

and *B. tabaci*.

NlSP694 is a planthopper-specific salivary protein that is nearly exclusively expressed in

salivary glands (Fig. 5d; Fig. S29). The function of NlSP694 or its gene homologs in other

planthopper species has not been well investigated previously. Silencing NlSP694 does not impact

insect survivorship (Fig. 5e). However, significant reductions in honeydew excretion and impaired

fecundity are observed in ds*NlSP694*-treated *N. lugens* (Fig. 5f, g), indicating the importance of

NlSP694 in insect feeding.

Subsequently, we transiently co-express OsRLP4-myc with a complete coding region of

NlSP694 (C-terminal flag tagged, NlSP694-flag). Another salivary protein NlSP7-flag and NtCf9-

myc are used as negative controls. The results demonstrate that NlSP694-flag specifically and

significantly decreases the protein level of OsRLP4-myc, which differs significantly from that of

the NlSP7-flag (Fig. 5h). Additionally, we demonstrate that NlSP694-flag fails to decrease the

protein level of NtCf9-myc (Fig. 5h). These findings suggest that *N. lugens* potentially evolves

salivary effectors to target rice RLPs.

Discussion

Plants have evolved a complex innate immune system to protect themselves against

pathogen/insect invasion, and the cell surface-localized PRRs function on the frontline [2]. To

counteract this response, the pathogens have been reported to develop various methods to avoid

plant PRR-mediated PTI for their own advantage [36]. However, it remains unknown whether

insects take the similar strategy to sabotage plant immunity. In this study, we demonstrate that the

RLP4/SOBIR1 complexes confer plant resistance against herbivorous insects. In response, both

planthoppers and whiteflies secrete salivary effectors that facilitate the degradation of defensive

RLP4, leading to attenuated plant immunity. Our results suggest that manipulation of PRRs may be

a conserved strategy for herbivorous insects.

PRRs recognize modified-self or non-self patterns and activate a series of immune signaling,

which subsequently regulate downstream defense responses and confer plant resistance against

invaders [2, 3]. While various insect salivary components have been reported to trigger plant

immune responses, the specific PRRs responsible for this process are elusive. Our study

demonstrates that RLP4 exerts a significant impact on insect performance by modulating plant

hormones and ROS (Fig. 3f-h; Fig. S16-S17), highlighting the critical role of this gene in plant

immunity. Nevertheless, the exact ligand for RLP4 remains unknown. It is possible that malectin-

LRR receptor RLP4 responds to the DAMPs or HAMPs that are released during insect feeding [37],

which deserved further investigation. RLP4 lacks the intracellular kinase domain, and forms

bimolecular receptor kinases by interacting with SOBIR1 (Fig. S15). Overexpression of NtRLP4 or

NtSOBIR1 enhances insect feeding, while silencing of either gene exerts the opposite effect (Fig.

Fig. S16-S19). The association between RLPs and SOBIR1 protein kinases is a structural and

functional equivalent to genuine receptor kinases [4]. Therefore, disrupting the stability of the

RLP4/SOBIR1 complex can potentially undermine the immune recognition process.

As pivotal regulators in immune signaling, PRRs are frequently targeted by multiple pathogens

through diverse mechanism. For example, *Pseudomonas syringae* hampered the functionality of

receptor complexes by disrupting the ligand-dependent association between receptor-like kinase

BAK1 and the flagellin receptor FLS2 [22], while *Phytophthora infestans* fostered alternative

splicing of the host RLPKs to enhance infection [21]. PRRs are usually subjected to degradation via

post-translational modifications [38]. In plants, the activity and stability of PRRs undergo stringent

regulation since excessive activation of PRRs can be detrimental [20, 39]. One prevalent regulatory

mechanism involves ubiquitination followed by subsequent degradation mediated by the 26S

proteasome [38]. Mounting evidence has demonstrated the exploitation of such mechanisms by

bacteria, fungi, viruses, and oomycetes to facilitate PRR degradation [40-43]. Nevertheless, the

occurrence of effector-mediated PRR degradation in insects is unreported. Our study reveals that

the salivary effectors BtRDP from *B. tabaci* and NlSP694 from *N. lugens* interact with plant RLP4s,

leading to their degradation in an ubiquitin-dependent manner (Fig. 4f-i). Notably, BtRDP and

NlSP694 share no sequence or structural similarity and lack resemblance to known eukaryotic

ubiquitin-ligase domains. Their interaction with RLP4s occurs in the extracellular space (Fig. 3d;

Fig. 5c), whereas the ubiquitin-proteasome system primarily functions in the cytosol and nucleus

[44]. Furthermore, NtRLP4 reduction is observed only in oeBtRDP transgenic plants, not in

oeBtRDP^{-sp} plants (Fig. 4f), suggesting that BtRDP exerts its influence on NtRLP4 in the

extracellular space. These findings might exclude the possibility that BtRDP/NlSP694 harbors E3

ligase activity that directly ubiquitinates RLP4s in plant cells. It is likely that salivary proteins

indirectly influence the post-translational modification of RLP4, increasing its degradation rate,

which warrants further investigation.

Herbivorous insects use a combination of salivary components to facilitate successful feeding

[29]. Previous studies have shown that *B. tabaci* suppresses plant defenses by leveraging SA-JA

crosstalk, with the salivary protein Bt56 playing a pivotal role during this process [12]. Bt56 triggers

the SA-signaling pathway by directly interacting with a KNOTTED 1-like homeobox transcription

factor [12]. This study investigates the effects of another salivary effector, BtRDP, on plant

hormones, and finds that it elicits similar effects to Bt56, albeit primarily impacting the signal

recognition process (Fig. 3h; Fig. S16e). This leads us to speculate that different salivary effectors

from the same insect species might simultaneously target the same signaling pathway to ensure

effective immune suppression. Our analysis reveals that BtRDP interacts with other host genes in

addition to NtRLP4 (Table S5), and silencing NtRLP4 cannot completely rescue the impaired

feeding performance of ds*BtRDP*-treated insects (Fig. 4a). Salivary proteins exert a crucial role in

mediating the interactions between herbivores and plants. The dynamic interplay between plants

and insects might be highly complex, one salivary protein can target multiple plant pathways, and

one plant pathway is influenced by multiple salivary proteins.

In conclusion, this study shows that suppressing PRR-mediated plant immunity may be the

common strategy for the successful feeding of herbivorous insects. Whiteflies and planthoppers

independently evolve salivary effectors that facilitate the degradation of to degrade defensive RLP4

of host plants in an ubiquitin-dependent manner, which can therefore disrupt the RLP4/SOBIR1

complexes for signal recognition (Fig. 6).

Material and methods

Insects and plants

A colony of *B. tabaci* (cryptic species MED) was originally collected from a soybean field in

Suzhou, Anhui Province, China. The insects were subsequently maintained on *N. tabacum* cv. K326

plants under controlled laboratory conditions (25 $\pm$ 1 °C, 50–70% relative humidity, 16:8 h

light:dark photoperiod). For experimental use, *N. benthamiana* and *N. tabacum* plants were grown

in a climate-controlled growth chamber (23 ± 1 °C, 16:8 h light: dark cycle).

Phylogenetic analysis of RDP

To identify potential RDP homologs across insect species, we performed a BLASTp search using

the BtRDP protein sequence as a query against the predicted proteomes of 30 insect species. Hits

were filtered using an E-value cutoff of 10^{-5}. The retrieved RDP homologs were aligned using

MAFFT (v7.310) with default parameters [45]. To improve alignment quality, we removed

ambiguously aligned regions using Gblocks with default settings. The optimal amino acid

substitution model (JTT+G) was determined using ModelTest-NG (v0.1.6). Phylogenetic

relationships were inferred using maximum likelihood analysis in RAxML (v0.9.0) with 1000

bootstrap replicates to assess branch support [46]. The final tree was visualized and annotated using

iTOL (https://itol.embl.de/).

Preparation of anti-BtRDP and anti-NtRLP4 serum

The coding sequence of BtRDP (excluding the signal peptide) was PCR-amplified and cloned

into the pET-28a vector (Novagen) to generate a C-terminal 6×His-tagged fusion protein. The

recombinant plasmid was transformed into *Escherichia coli* Transetta (DE3) competent cells

(TransGen Biotech, Beijing, China). Protein expression was induced with 0.5 mM isopropyl-β-D-

thiogalactoside (IPTG) at 28°C for 8 h. Bacterial cells were harvested by centrifugation and lysed

via ultrasonication for 30 min. The recombinant BtRDP^{-sp}-His protein was purified under native

conditions using Ni-NTA agarose (Qiagen, Hilden, Germany) according to the manufacturer's

protocol.

Purified BtRDP^{-sp}-His protein was used to immunize rabbits (Hua'an Biotechnology,

Hangzhou, China) for custom polyclonal antibody production. The anti-NtRLP4 serum, prepared

by immunizing rabbits with peptide GQNSDRPISTKNST, was produced via the custom service of

Genscript Biotechnology Company (Nanjing, China). The anti-BtRDP serum was subsequently

conjugated to Alexa Fluor™ 488 NHS Ester (#A20000, Thermo Fisher Scientific) following

standard labeling protocols.

The specificity of anti-BtRDP serum was tested by comparing ds*GFP*- and ds*BtRDP*-treated

Bemisia tabaci. The results showed that ds*BtRDP* significantly reduced the target gene. Similarly,

the specificity of anti-NtRLP4 serum was tested by comparing RNAi-EV and RNAi-NtRLP4 plants,

which showed a reduced NtRLP4 expression in *RNAi-NtRLP4* plants.

Immunohistochemistry (IHC) staining

Dissected *B. tabaci* heads with attached salivary glands and Parafilm-embedded salivary

sheaths were fixed in 4% paraformaldehyde (#E672002, Sango Biotechnology, Shanghai) for 30

min. Salivary sheaths adherent to Parafilm were similarly excised and fixed under identical

conditions. After fixation, samples were washed three times with PBST (PBS containing 0.1%

Tween-20) and blocked with 10% FBS for 2 h at room temperature. The tissues were then incubated

overnight at 4°C with Alexa Fluor 488-conjugated anti-BtRDP serum (1:200 dilution), followed by

co-staining with phalloidin-rhodamine (1:500, Thermo Fisher Scientific) for 30 min and 4',6-

diamidino-2-phenylindole (DAPI) solution (#ab104139, Abcam, Cambridge, USA) for nuclear

visualization. Fluorescent images were acquired using a Leica SP8 confocal microscope (Leica

Microsystems) to examine BtRDP localization patterns

Protein extraction and Western-blotting assays

To identify secreted salivary proteins in *N. tabacum* plants, approximately 200 *B. tabaci* adults

were confined in a 3-cm diameter feeding cage and allowed to feed on host plants for 24 h.

Following feeding, the infested tobacco leaves were collected, immediately frozen in liquid nitrogen,

and ground to a fine powder. The powdered tissue was then homogenized in RIPA Lysis Buffer

(#89900, Thermo Fisher Scientific) for protein extraction. To examine the effects of insect feeding

on NtRLP expression, groups of approximately 200 *B. tabaci* were allowed to feed on 4-5 true-leaf

stage tobacco plants for 12, 24, or 48 h, with uninfested plants serving as controls. For

silencing/overexpression efficiency tests, leaf samples were collected at 48 h post-*Agrobacterium*

infiltration, along with transgenic tobacco plants at the 4-5 true-leaf stage. For insect tissue analysis,

various *B. tabaci* organs were carefully dissected in 1×PBS buffer, including carcasses (20), fat

bodies (20), guts (40), salivary glands (60), and ovaries (10), with sample sizes indicated in

parentheses. All dissected tissues were transferred to lysis buffer for protein extraction. Total protein

concentrations in both insect and plant homogenates were quantified using a BCA Protein Assay

Kit (#CW0014S, CwBiotech, Taizhou, China). Equal protein amounts were separated by SDS-

PAGE and electro-transferred to PVDF membranes.

Membranes were probed with the following primary antibodies: anti-FLAG (1:10,000, #MA1-

91878, Thermo Fisher Scientific), anti-MYC (1:10,000, #MA1-21316, Thermo Fisher Scientific),

anti-GFP (1:10,000, #MA5-15256, Thermo Fisher Scientific), anti-NtRLP4 (1:5,000, GenScript

Biotechnology, Nanjing, China), and anti-BtRDP serum (1:5,000, Huaan Biotechnology, Hangzhou,

China). Following primary antibody incubation, membranes were treated with HRP-conjugated

secondary antibodies: goat anti-mouse IgG (1:10,000, #31430, Thermo Fisher Scientific) or goat

anti-rabbit IgG (1:10,000, #31460, Thermo Fisher Scientific). Protein bands were visualized using

an AI 680 image analyzer (Amersham Pharmacia Biotech, Buckinghamshire, UK), with Ponceau S

staining used to verify equal protein loading.

Quantitative real-time PCR (qRT-PCR) analysis

To prepare tissue-specific samples from *B. tabaci*, adult females were dissected to isolate

carcasses (10), fat bodies (10), guts (20), salivary glands (40), and ovaries (10). Additionally,

different developmental stages, including eggs (50), nymphs (20), pupae (20), and adult males (20)

and females (20), were collected for analysis. The number of insects in each sample is indicated in

parentheses. To extract RNA from *N. tabacum* plants, the samples were grinded using liquid

nitrogen after collection. Total RNA was extracted using the TRIzol Total RNA Isolation Kit (#9109,

Takara, Dalian, China) following the manufacturer's instructions. RNA purity and concentration

were assessed using a NanoDrop spectrophotometer (Thermo Fisher Scientific). First-strand cDNA

was synthesized from equal amounts of RNA (1 µg) using the HiScript II Q RT SuperMix (#R212-

01, Vazyme, Nanjing, China). Gene expression analysis was performed using a Roche LightCycler®

480 system (Roche Diagnostics, Mannheim, Germany) with SYBR Green Supermix (#11202ES08,

Yeasen, Shanghai, China). The thermal cycling protocol consisted of initial denaturation at 95°C for

5 min, and amplification for 40 cycles of 95°C for 10 s and 60°C for 30 s. Primers were designed

with Primer Premier v6.0 (Table S7). *B. tabaci Tubulin* and *N. tabacum Tubulin* served as internal

controls for normalization. Relative gene expression was calculated using the $2^{-\Delta\Delta Ct}$ method [47].

Three to six independent biological replicates, each repeated twice, were performed.

RNA interference

The target gene DNA sequences were amplified using primers listed in Table S7 and

subsequently cloned into the pClone007 Vector (#TSV-007, Tsingke Biotechnology, Beijing, China).

For dsRNA production, PCR-generated DNA templates containing T7 promoter sequences were

transcribed *in vitro* using the T7 High Yield RNA Transcription Kit (#TR101-01, Vazyme Biotech,

Nanjing, China). The RNAi experiments were performed according to established methods [48].

Briefly, newly emerged adult whiteflies were anesthetized with carbon dioxide for 5-10 seconds.

Approximately 50 nL of dsRNA solution (2 μg/μL) was microinjected into the mesothorax of each

insect using a FemtoJet microinjection system (Eppendorf, Hamburg, Germany). Following

injection, the whiteflies were maintained on healthy *N. tabacum* leaves under standard rearing

conditions. Only actively moving insects were selected for subsequent analyses. Gene silencing

efficiency was evaluated on day 4 post-injection using both qRT-PCR and Western-blotting

approaches.

Generation of transgenic plants

We generated two transgenic *N. tabacum* lines expressing different BtRDP variants: one

containing the full coding sequence (oeBtRDP) and another lacking the signal peptide (oeBtRDP⁻

ˢᵖ), which may affect protein secretion. The target sequences, along with the full-length NtRLP4

coding sequence and an NtRLP4 hairpin RNAi construct, were each cloned into the pBWA(V)HS

vector with C-terminal FLAG tags. All constructs were transformed into *A. tumefaciens* strain

GV3101 and used to generate transgenic tobacco plants (K326 background) through

Agrobacterium-mediated transformation performed by Wuhan Boyuan Biological Company.

Transgenic lines were verified by qRT-PCR and western blotting, with empty vector-transformed

plants serving as controls, and two independent lines for each construct were selected for further

experiments.

Target gene silencing in plants

The hairpin RNAi construct targeting *NtSOBIR1* was generated by cloning the specific

fragment into the pBWA(V)HS plasmid using BsaI and Eco31I restriction enzymes. The

recombinant vector was then transformed into *A. tumefaciens* strain GV3101 and subsequently

infiltrated into *N. tabacum* leaves using the agroinfiltration method. Following a 48-h incubation

period under standard growth conditions, silencing efficiency was quantitatively evaluated by qRT-
PCR analysis.

Transcriptomic sequencing

To investigate the transcriptional effects of NtRLP4, BtRDP, and BtRDP^{-sp} in *N. tabacum*, leaf
samples from EV, oeRLP4#1, oeBtRDP#1, and oeBtRDP^{-sp}#1 transgenic plants were homogenized
in TRIzol Reagent (#10296018, Invitrogen, Carlsbad, CA, USA) for total RNA extraction following
the manufacturer's protocol. Afterwards, total RNA was extracted according to the instructions of
the manufacturer, and the RNA samples were sent to Novogene Institute (Novogene, Beijing, China)
for transcriptomic sequencing. Briefly, poly(A)+ RNA was isolated from 20 μg pooled total RNA
using oligo(dT) magnetic beads, followed by fragmentation in divalent cation buffer at 94°C for 5
min. First-strand cDNA synthesis was performed using N6 random primers, with subsequent
second-strand synthesis to generate double-stranded cDNA. After end-repair and Illumina adapter
ligation, the products were PCR-amplified (15 cycles) and purified using a QIAquick PCR
Purification Kit (#28104, Qiagen, Hilden, Germany). The resulting cDNA libraries were quantified
and quality-checked before paired-end sequencing on an Illumina NovaSeq 6000 platform. All raw
sequencing data were deposited in the National Genomics Data Center under accession number
PRJCA025907.

Analysis of transcriptomic data

The raw sequencing reads were quality-filtered using Illumina's internal software, and the
resulting clean reads from each cDNA library were aligned to the *N. tabacum* reference genome in
Sol Genomics Network
(https://solgenomics.net/ftp/genomes/Nicotiana_tabacum/edwards_et_al_2017/) using HISAT
v2.1.0 [49]. Low-quality alignments were removed using SAMtools (v1.7) with default parameters
[50]. Transcript abundance was quantified as Transcripts Per Million (TPM) using Cufflinks (v2.2.1)
[51]. Differential expression analysis was performed with DESeq2 (v2.2.1) [52], identifying
significantly DEGs using thresholds of |log2(fold change)| > 1 and adjusted *p*-value < 0.05. PCA of
transcriptome profiles was conducted using a custom R script plotPCA (github.com/franco-
ye/TestRepository/blob/main/PCA_by_deseq2.R). For functional annotation, KEGG pathway

enrichment analysis was performed in TBtools (v2.083) using a one-sided hypergeometric test:

$P = 1 - \sum_{i=0}^{m-1} \left(\frac{\binom{M}{i}\binom{N-M}{n-i}}{\binom{N}{n}} \right)$, [53]. In this software, enriched *P*-values were calculated according to

one-sided hypergeometric test: with *N* representing the number of gene with KEGG annotation, *n*

standing for the number of DEGs in *N*, *M* indicating the number of genes in each KEGG term, and

m suggesting the number of DEGs in each KEGG term.

Insect bioassays

To survivorship analysis, newly emerged *B. tabaci* adults were microinjected with dsRNA and

placed on *N. tabacum* leaves. After 24 h, any dead individuals (potentially due to injection injury)

were removed. For each treatment, 20-30 surviving adults were transferred to leaf cages, and

mortality was recorded daily for 10 days. Three independent biological replications were performed.

For fecundity analysis, dsRNA-treated virgin females were paired with untreated males in

leaf cages. Five such pairs were maintained per cage (n=10-12 cages per treatment). Egg deposition

was quantified daily for 3 consecutive days under controlled conditions. Each male-female pair was

considered an experimental replicate.

For host preference test, detached *N. tabacum* leaves with moist cotton-wrapped petioles were

arranged in a 50-cm diameter choice arena with a central release chamber. Forty adult whiteflies

were released per chamber, and settling preference was recorded at 3, 6, 12, 24, 36, and 48 h post-

release. The experiment included 10-12 replicates per treatment, with leaf positions randomized

between replicates to eliminate positional bias.

EPG recording and analysis

Feeding behavior of *B. tabaci* was monitored using a GiGA-8d EPG amplifier (Wageningen

Agricultural University; 10 TΩ input resistance, <1 pA input bias current) with 50× gain and ±5 V

output range. Prior to recording, dsRNA-treated whiteflies were starved for 12 h on moist filter

paper, briefly anesthetized with CO_2 (10 s), and connected to the amplifier via a gold wire electrode

(20 μm diameter × 5 cm length) attached to the pronotum using conductive silver glue. Plant

electrodes consisted of copper wires (2 mm diameter × 10 cm length) inserted into the soil of potted

N. tabacum. Recordings were conducted for 8 h in a Faraday cage (120×75×67 cm; Dianjiang,

Shanghai), with only insects surviving the full duration retained for analysis.

Recorded waveforms were processed using PROBE 3.4 software (Wageningen Agricultural

University) and categorized into four distinct feeding phases: nonpenetration (np), pathway duration

(C), phloem salivation (E1), and phloem ingestion (E2) as defined in a previous study [30]. Each

treatment included ≥ 10 biological replicates, with ds*GFP*-injected and untreated whiteflies

serving as controls.

Agrobacterium*-mediated plant transformation in *N. tabacum* and *N. benthamiana

The recombinant expression vectors were introduced into *A. tumefaciens* strain GV3101 via

heat shock transformation. Transformed cells were selected on LB agar plates supplemented with

50 μg/mL kanamycin and 10 μg/mL rifampicin, followed by incubation at 28°C for 60 h. Positive

colonies were inoculated into liquid LB medium with the same antibiotics and grown to log phase.

Bacterial cells were then pelleted by centrifugation at 2,400 *g* for 2 min and resuspended in induction

buffer (10 mM MgCl$_2$, 10 mM MES [pH 5.6], 200 μM acetosyringone) adjusted to specific OD_{600}

values depending on the experiment.

For standard experiments, bacterial suspensions were normalized to $OD_{600} = 1.0$, while

degradation assays used three concentrations ($OD_{600} = 0.05, 0.3,$ and 1.0). Equal volumes of selected

bacterial suspensions were mixed and infiltrated into leaves of 4–5-week-old *N. tabacum* and *N.*

benthamiana plants using a needleless syringe. Infiltrated plants were maintained under controlled

conditions for subsequent analysis.

Diaminobenzidine (DAB) staining

To detect H_2O_2 accumulation in *N. tabacum* leaves, leaf samples were immersed in 1 mg/mL

3,3'-diaminobenzidine tetrahydrochloride (DAB-HCl, #D8001, Sigma-Aldrich, St. Louis, MO,

USA) solution (pH 3.8) for 6 hours at room temperature, allowing H_2O_2-dependent brown

polymerization to occur. Following incubation, the DAB solution was replaced with 100% ethanol

for chlorophyll removal, and samples were decolorized overnight at 65°C. The destained leaves

were then photographed under standardized conditions using a Canon EOS 80D digital camera

(Canon Inc., Tokyo, Japan).

Interaction assays between two proteins

In the Y2H screening assay, the coding sequence of BtRDP^{-sp} was cloned into the pGBKT7 bait vector (Clontech, USA), while a *N. benthamiana* cDNA library was constructed in the pGADT7 prey vector (Biogene Biotech, Shanghai, China). These recombinant vectors were co-transformed into yeast strain Y2HGold and plated on QDO medium (SD/-Ade/-His/-Leu/-Trp, #630428, Takara) for 3 days at 30°C. Positive colonies were cultured in QDO liquid medium, and plasmids were extracted using a yeast DNA kit (#DP112-02, TIANGEN, Beijing, China) before transformation into E. coli DH5α (#TSC-C14, Tsingke) for sequencing (YouKang Biotech, China) to identify interacting partners.

In the Y2H point-to-point verification assay, BtRDP^{-sp} and NlSP694^{-sp}, and mutant variants of NtRLP4/SlRLP4/ OsRLP4 were cloned into either pGBKT7 or pGADT7 vectors using primers from Table S7. Yeast co-transformants were first selected on DDO medium (SD/-Leu/-Trp) (#630417, Takara) for 3 days at 30°C, then spotted onto QDO medium to confirm interactions through growth after another 3 days of incubation.

In the Co-IP assays, Full-length sequences of *BtRDP*, *NlSP694*, *NtSOBIR1*, *NtRLP4*, *SlRLP4*, and *OsRLP4* were cloned into lic-myc or lic-flag vectors for C-terminal tagging, with BtFTSP-flag serving as a negative control. Five-week-old *N. benthamiana* leaves were agroinfiltrated with these constructs or collected from oeBtRDP transgenic plants. Total proteins were extracted from 1 g frozen tissue in IP lysis buffer (#87788, Thermo Scientific) with protease inhibitors (#56079200, Roche, Switzerland), followed by centrifugation at $1,000 \times g$ for 20 min. Cleared lysates were incubated with anti-flag beads (#L00432-1, GenScript, Nanjing, China) for 4 h at 4°C, washed with PBS, and eluted in 2× SDS-PAGE buffer (500 mM Tris-HCl, pH=6.8, 50% glycerin, 10% SDS, 1% bromophenol blue, and 2% β-mercaptoethanol) for Western-blotting assay.

In the BiFC assay, BtRDP^{-sp}, BtFTSP^{-sp}, NtRLP4, and NtCf9-like were cloned into the pCV-cYFP or pCV-nYFP vectors, respectively. After transforming *A. tumefaciens* GV3101, pairwise combinations were co-infiltrated into *N. benthamiana* leaves. Following 36-48 h incubation in a growth chamber, reconstituted YFP fluorescence was visualized using a Leica SP8 confocal microscope with standard filter sets for YFP detection.

RLP degradation by salivary effectors among different herbivorous insects

To examine salivary protein effects on RLPs, we co-infiltrated *A. tumefaciens* GV3101 strains

carrying various constructs into *N. benthamiana* leaves. The recombinant vectors expressed C-

terminally tagged proteins: BtRDP-flag, NlSP694-flag, GUS-flag, NlSP7-flag, and myc-tagged

RLPs (NtRLP4, SlRLP4, OsRLP4, SlCf9, NtCf9). Bacterial suspensions were mixed at

predetermined OD_{600} ratios before infiltration, with leaf samples collected 48 hours post-infiltration

for subsequent analysis.

For confocal microscopy studies, *N. benthamiana* leaves were co-infiltrated with *A.*

tumefaciens carrying: (1) NtRLP4-GFP plus BtRDP-mCherry, (2) NtRLP4-GFP plus RFP-mCherry,

or (3) NtRLP4-GFP plus BtFTSP-mCherry. The samples were imaged by confocal microscopy at

48 h post injection. The same parameters were used in photographing.

This study also conducted a treatment on *N. benthamiana* plants that were overexpressing

NtRLP4-myc using purified proteins. Briefly, the BtRDP^{-sp} and GFP coding sequences were cloned

into pET-28a for 6×His-tagged expression in *E. coli* Transetta (TransGen Biotech). Protein

expression was induced with 0.5 mM IPTG at 25°C for 8 h, followed by purification using Ni-NTA

agarose and buffer exchange into PBS using a 3-kDa molecular-weight cutoff Amicon Ultra-4

Centrifugal Filter Device (Millipore, MA, USA). Different concentrations of purified proteins were

infiltrated into leaves of *N. benthamiana* plants stably expressing NtRLP4-myc, with samples

collected 24 h post-treatment.

To investigate degradation pathways, we infiltrated *N. benthamiana* leaves with: 50 μM

MG132 (#M7449, Sigma-Aldrich, Steinheim, Germany), 5 μM 3-MA (#M9281, Sigma-Aldrich,),

or 10 nM BAF (#BML-CM110-0100, Enzo Life Sciences, Farmingdale, USA) at 24 h post-

agroinfiltration. The samples were incubated for additional24 h. Protein levels were detected using

Western-blotting assay.

***In planta* ubiquitination assay**

Ubiquitination detection in *N. benthamiana* was performed according to the methods

previously reported [54]. The recombinant vector for transient expression of the HA-fused ubiquitin

protein was kindly provided by Dr. Chao Zheng (Ningbo University). Myc-tagged proteins were co-

expressed with HA-UBQ by agroinfiltration, and 50 μM MG132 was injected into the *N.*

benthamiana leaves one day post *A. tumefaciens* infiltration. Immunoprecipitation was performed

using anti-myc beads (ytma-20, ChromoTek , Planegg-Martinsried, Germany). Ubiquitination

signals were detected by immunoblot with anti-HA antibody (HT301, TransGen Biotech, Beijing,

China).

Scanning electron microscopy (SEM)

B. tabaci adults were allowed to feed on artificial diet solutions for 24 h. Parafilm sections

containing salivary sheaths were carefully excised and rinsed with 1× PBS (pH 7.4) to remove

residual diet. The samples were mounted on aluminum stubs, desiccated under vacuum, and sputter-

coated with gold before imaging with a TM4000 II Plus SEM (Hitachi, Tokyo, Japan). Salivary

sheath length (from apex to base) was measured, with 20 sheaths analyzed per treatment to ensure

statistical reliability.

Statistical analysis

The log-rank test (SPSS Statistics 19, Chicago, IL, USA) was used to determine the statistical

significance of survival distributions. Two-tailed unpaired Student's *t*-test (comparisons between

two groups) or one-way ANOVA test followed by Tukey's multiple comparisons test (comparisons

among three groups) was used to analyze the results of qRT-PCR, fecundity analysis, and host choice

analysis. The EPG data were first checked for the normality and homogeneity of variance, and those

not fitting a normal distribution were $\log_{10}$ transformed before Student's *t* test, as described in

previous studies [30]. Data were graphed in GraphPad Prism 9.

Data availability

679 All data is included in the manuscript and/or supporting information. The transcriptomic data have

680 been submitted to the National Genomics Data Center under accession number PRJCA025907. The

681 original blot images have been displayed in Fig. S30-S33.

References

684 1.   Nürnberger T, Brunner F, Kemmerling B, & Piater L (2004) Innate immunity in plants and

685   animals: striking similarities and obvious differences. *Immunol Rev* 198(1):249-266.

686 2.   Macho AP & Zipfel C (2014) Plant PRRs and the activation of innate immune signaling. *Mol*

687   *Cell* 54(2):263-272.

688 3. Zipfel C (2014) Plant pattern-recognition receptors. *Trends Immunol* 35(7):345-351.

689 4. Gust AA & Felix G (2014) Receptor like proteins associate with SOBIR1-type of adaptors to

690   form bimolecular receptor kinases. *Curr Opin Plant Biol* 21:104-111.

691 5. Adam D. Steinbrennera, *et al.* (2020) A receptor-like protein mediates plant immune responses

692   to herbivore-associated molecular patterns. *Proc Natl Acad Sci USA* 117(49):31510-31518.

693 6. Chinchilla D, Bauer Z, Regenass M, Boller T, & Felix G (2005) The *Arabidopsis* receptor kinase

694   FLS2 binds flg22 and determines the specificity of flagellin perception. *Plant Cell* 18(2):465-

695   476.

696 7. Zipfel C, *et al.* (2006) Perception of the bacterial PAMP EF-Tu by the receptor EFR restricts

697   *Agrobacterium*-mediated transformation. *Cell* 125(4):749-760.

698 8. Ron M & Avni A (2004) The receptor for the fungal elicitor ethylene-inducing xylanase is a

699   member of a resistance-like gene family in tomato. *Plant Cell* 16(6):1604-1615.

700 9. Tian T, *et al.* (2021) A salivary calcium-binding protein from *Laodelphax striatellus* acts as an

701   effector that suppresses defense in rice. *Pest Manag Sci* 77(5): 2272-2281.

702 10. Huang HJ, *et al.* (2019) Salivary DNase II from *Laodelphax striatellus* acts as an effector that

703   suppresses plant defence. *New Phytol* 224(2):860-874.

704 11. Chen CY, *et al.* (2019) An effector from cotton bollworm oral secretion impairs host plant

705   defense signaling. *Proc Natl Acad Sci USA* 116(28):14331-14338.

706 12. Xu HX, *et al.* (2019) A salivary effector enables whitefly to feed on host plants by eliciting

707   salicylic acid-signaling pathway. *Proc Natl Acad Sci USA* 116(2):490-495.

708 13. Chen CY & Mao YB (2020) Research advances in plant-insect molecular interaction. *F1000Res*

709   9(F1000 Faculty Rev):198.

710 14. Guiguet A, *et al.* (2016) Shared weapons of blood- and plant-feeding insects: Surprising

711   commonalities for manipulating hosts. *J Insect Physiol* 84:4-21.

712 15. Wu J & Baldwin IT (2009) Herbivory-induced signalling in plants: perception and action. *Plant*

713   *Cell Environ* 32(9):1161-1174.

714 16. Jiang YJ, Zhang CX, Chen RZ, & He SY (2019) Challenging battles of plants with phloem-

715   feeding insects and prokaryotic pathogens. *Proc Natl Acad Sci USA* 116(47):23390-23397.

716 17. Gao H，*et al*. (2023) The salivary chaperone protein NlDNAJB9 of *Nilaparvata lugens* activated

plant immune responses. *J Exp Bot* 74(21):6874-6888.

18.   Guo H, *et al*. (2020) An aphid-secreted salivary protease activates plant defense in phloem. *Curr*

Biol 30(24): 4826-4836.

19.   Bos JI, *et al*. (2010) A functional genomics approach identifies candidate effectors from the

aphid species *Myzus persicae* (green peach aphid). *PLoS Genet* 6(11):e1001216.

20.   Zhang Y & Zeng L (2020) Crosstalk between ubiquitination and other post-translational protein

modifications in plant immunity. *Plant Commun* 1(4):100041.

21.   Huang J*, et al.* (2020) *Phytophthora* effectors modulate genome-wide alternative splicing of

host mRNAs to reprogram plant immunity. *Mol Plant* 13(10):1470-1484.

22.   Shan L*, et al.* (2008) Bacterial effectors target the common signaling partner BAK1 to disrupt

multiple MAMP receptor-signaling complexes and impede plant immunity. *Cell Host Microbe*

4(1):17-27.

23.   Huang HJ*, et al.* (2016) Screening and functional analyses of *Nilaparvata lugens* salivary

proteome. *J Proteome Res* 15(6):1883-1896.

24.   Huang HJ, *et al.* (2021) Identification of salivary proteins in the whitefly *Bemisia tabaci* by

transcriptomic and LC-MS/MS analyses. *Insect Sci* 28(5):1369-1381.

25.   Gao H, *et al*. (2022) *Nilaparvata lugens* salivary protein NlG14 triggered defense response in

plants. *J Exp Bot* 73(22):7477-7487.

26.   Mukhtar M, *et al*. (2011) Independently evolved virulence effectors converge onto hubs in a

plant immune system network. *Science* 333(6042):596-601.

27.   Huang J, *et al.* (2020) Salivary protein 1 of brown planthopper is required for survival and

induces immunity response in plants. *Front Plant Sci* 11:571280.

28.   Wang Y, *et al*. (2023) Horizontally transferred salivary protein promotes insect feeding by

suppressing ferredoxin-mediated plant defenses. *Mol Biol Evol* 40(10):msad221.

29.   Huang HJ*, et al.* (2023) Planthopper salivary sheath protein LsSP1 contributes to manipulation

of rice plant defenses. *Nat Commun* 14(1):737.

30.   Lu S, *et al*. (2021) EPG-recorded feeding behaviors reveal adaptability and competitiveness in

two species of *Bemisia tabaci* (Hemiptera: Aleyrodidae). *J Insect Behav* 34: 26-40.

31.   Yan ZW, *et al*. (2023) Endocytosis-mediated entry of a caterpillar effector into plants is

countered by Jasmonate. *Nat Commun* 14(1):6551.

32.   Jones DA, *et al*. (1994) Isolation of the tomato Cf-9 gene for resistance to *Cladosporium fulvum*

by transposon tagging. *Science* 266 (5186): 789-793 (1994).

33.   Varshavsky A, *et al*. (2017) The ubiquitin system, autophagy, and regulated protein degradation.

Annu Rev Biochem 86: 123-128.

34.   Li L*, et al.* (2022) Independently evolved viral effectors convergently suppress DELLA protein

SLR1-mediated broad-spectrum antiviral immunity in rice. *Nature Commun* 13(1):6920.

35.   Takatsuka C*, et al.* (2004) 3-Methyladenine inhibits autophagy in tobacco culture cells under

sucrose starvation conditions. *Plant Cell Physiol* 45: 265-274.

36.   Tariqjaveed M*, et al.* (2021) Versatile effectors of phytopathogenic fungi target host immunity.

J Integr Plant Biol 63(11):1856-1873.

37.   Doblas VG, *et al.* (2018) Cell wall integrity signaling in plants: Malectin-domain kinases and

lessons from other kingdoms. *Cell Surf* 3:1-11.

38.   Li B, Lu D, & Shan L (2014) Ubiquitination of pattern recognition receptors in plant innate

immunity. *Mol Plant Pathol* 15(7):737-746.

39.   Chen Y, Song Y, Liu J, Xu G, & Dou D (2022) Ubiquitination of receptorsomes, frontline of

plant immunity. *Int J Mol Sci* 23(6):2937.

40.   Langin G, González-Fuente M, & Üstün S (2023) The plant ubiquitin–proteasome system as a

target for microbial manipulation. *Annu Rev Phytopathol* 61:351-375.

41.   Rosas-Diaz T*, et al.* (2018) A virus-targeted plant receptor-like kinase promotes cell-to-cell

spread of RNAi. *Proc Natl Acad Sci USA* 115(6):1388-1393.

42.   Irieda H*, et al.* (2019) Conserved fungal effector suppresses PAMP-triggered immunity by

targeting plant immune kinases. *Proc Natl Acad Sci USA* 116(2):496-505.

43.   Qin J*, et al.* (2018) The Xanthomonas effector XopK harbours E3 ubiquitin-ligase activity that

is required for virulence. *New Phytol* 220(1):219-231.

44.   Xu FQ, Xue HW (2019) The ubiquitin-proteasome system in plant responses to environments.

Plant Cell Environ 42(10):2931-2944.

45.   Katoh K & Standley DM (2013) MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. *Mol Biol Evol* 30(4):772-780.

46.   Kozlov AM, Darriba D, Flouri T, Morel B, & Stamatakis A (2019) RAxML-NG: a fast, scalable

and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics*

35(21):4453-4455.

47.   Livak, K.J., and Schmittgen, T.D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ method. *Methods* 25, 402-408.

48.   Xu HJ*, et al.* (2015) Two insulin receptors determine alternative wing morphs in planthoppers. *Nature* 519(7544):464-467.

49.   Kim, D., Langmead, B., and Salzberg, S.L. (2015). HISAT: A fast spliced aligner with low memory requirements. *Nature Methods* 519(7544):464-467.

50.   Li, H. (2009). The Sequence Alignment/Map format and SAMtools. *BIOINFORMATICS - OXFORD*.

51.   Trapnell, C., Roberts, A., Goff, L., Pertea, G., Kim, D., Kelley, D.R., Pimentel, H., Salzberg, S.L., Rinn, J.L., and Pachter, L. (2012). Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. *Nature Protocols* 7, 562-578.

52.   Wang, L., Feng, Z., Wang, X., Wang, X., and Zhang, X. (2010). DEGseq: an R package for identifying differentially expressed genes from RNA-seq data. *Bioinformatics* 26, 136-138.

53.   Chen, C., Chen, H., Zhang, Y., Thomas, H.R., Frank, M.H., He, Y., and Xia, R. (2020). TBtools: an integrative toolkit developed for interactive analyses of big biological data. *Mol Plant* 13, 1194-1202.

54.   Wang, N., Yin, Z., Zhao, Y., Wang, J., Pei, Y., Ji, P., Daly, P., Li, Z., Dou, D., and Wei, L. (2022). An F-box protein attenuates fungal xylanase-triggered immunity by destabilizing LRR-RLP NbEIX2 in a SOBIR1-dependent manner. *New Phytol* 236(6):2202-2215.

Acknowledgements

This project has received funding from the National Key Research and Development Plan in the 14th five-year plan (2021YFD1401100: H.J.H, C.X.Z.), National Natural Science Foundation of China (32422075: H.J.H.; U23A6006: J.P.C and C.X.Z.), and the Natural Science Foundation of Zhejiang Province (LDQ24C140001: H.J.H).

Author Contributions Statement

H-JH, J-ML, C-XZ, and J-PC planned and designed the research. XW, J-BL, Y-ZW, H-JH, H-BL, Z-LZ, and Z-XY performed experiments and analyzed data. X-HZ provided the EV plants. H-JH

807   and XW drafted the manuscript.

809   **Competing Interests Statement**

810   The authors declare no competing interests.

 Figure Legends

 Fig. 1. BtRDP is a salivary protein and secreted into plants. (**a**) Detection of BtRDP in *Nicotiana*

 tabacum plants. The untreated and *Bemisia tabaci*-infested tobacco plants are collected for western-

 blotting assays. Six independent *B. tabaci*-infested plants are tested. The presence of lower intensity

 band (arrow) in each replicate is summarized in the bracket. Asterisk indicates non-specific binding.

 Coomassie brilliant blue (CBB) staining is conducted to visualize the sample loading. (**b**, **c**)

 Detection of BtRDP in different tissues by western-blotting and qRT-PCR assays. Ov, ovary; SG,

 salivary gland; Ca, carcass; FB, fat body. Silver staining and anti-actin serum are used to visualize

 the sample loading. (**d**) Expression patterns of BtRDP in different development stages. Ny, nymph;

 Ps, pseudopupa; Fe, female; Ma, male. *B. tabaci tubulin* is used as an internal control. The relative

 quantitative method ($2^{-\Delta\Delta Ct}$) is used to evaluate the quantitative variation. Data are presented as mean

 values ± SEM (n= 3 independent biological replicates). Different lowercase letters indicate

 statistically significant differences at the $P < 0.05$ level according to one-way ANOVA test followed

 by Tukey's multiple comparisons test. (**e**) Immunohistochemical staining of BtRDP. The salivary

 gland and its nearby tissues are dissected and incubate with anti-BtRDP serum or pre-immune serum

 conjugated with Alexa Fluor™ 488 NHS Ester (green) and actin dye phalloidinrhodamine (red).

 The nucleus is stained with DAPI (blue). PSG, principal salivary gland; ASG, accessory salivary

 gland. Experiments are repeated thrice for (**b**), while twice for (**e**). Similar results are observed and

 representative images are displayed.

 Fig. 2. Effects of BtRDP on *Bemisia tabaci*. (**a**) Treating *B. tabaci* with ds*BtRDP* significantly

 reduces the transcript and protein level of target gene. The ds*GFP*-treated *B. tabaci* is used as a

 control. qRT-PCR data are presented as mean values ± SEM (n= 6 independent biological replicates).

 (**b–c**) Effects of *BtRDP* knockdown on insect reproduction (**b**) and feeding behavior (**c**). Electrical

 penetration graph (EPG) is used to monitor the insect feeding behavior, which can be classified into

 nonpenetration (np), pathway duration (C), phloem salivation (E1) and phloem ingestion (E2)

 phases. All EPG recordings are performed for 8 h. Typical EPG waveforms are displayed in Fig. S8.

 (**d-f**) Insect performance on transgenic plants overexpressing BtRDP (oeBtRDP). (**d**) Detection of

 BtRDP in transgenic *Nicotiana tabacum* overexpressing a complete coding region of *BtRDP*. The

 empty vector (EV) plants are used as the control. The flag tag is fused to the C-terminal ends of

recombinant proteins. **(e)** Comparison of insect reproduction on oeBtRDP#1 and EV plants. Five *B.*

tabaci individuals are confined to indicated plants for 3 days, and the oviposited eggs are counted.

(f) Attraction of oeBtRDP#1 and EV leaves to *B. tabaci* in a two-choice equipment. A group of 40

female *B. tabaci* are released into a device containing oeBtRDP#1 and EV leaves. The number of

insects settling on each leaf is counted. After 48 h, the number of eggs on each leaf is counted. **(g-i)**

Insect performance on transgenic plants overexpressing BtRDP^{-sp} (oeBtRDP^{-sp}). **(g)** Detection of

BtRDP^{-sp} in transgenic *N. tabacum* overexpressing BtRDP without a signal peptide. The insect

reproduction **(h)** and settlement **(i)** on oeBtRDP^{-sp} transgenic plants are recorded. Data are presented

as mean values ± SEM. The EPG data are first checked for normality and homogeneity of variance,

and data not fitting a normal distribution are subjected to log10 transformation. *P*-values are

determined by two-tailed unpaired Student's *t* test. ***P < 0.001; **P < 0.01; *P < 0.05; ns, not

significant. Western-blotting assays are repeated thrice with the similar results.

Fig. 3. NtRLP4 interacts with BtRDP and confers plant resistance to *Bemisia tabaci*. (a)

Domain organization of NtRLP4. NtRLP4 contains a predicted N-terminal signal peptide (SP), a

malectin-like domain, a LRR domain, and a transmembrane domain (TM). **(b)** The protein level of

NtRLP4 in response to *B. tabaci* infestation. The untreated *Nicotiana tabacum* is used as a negative

control. Four independent biological replicates are performed, and the representative images are

displayed. The band density is measured using ImageJ. The density values from 4 biological

replicates are calculated and the mean value in the controls is set at 1.0. **(c)** Yeast two hybrid assays

showing the interaction between BtRDP and NtRLP4. The different combinations of constructs are

transformed into yeast cells, and are grown on the selective medium SD/-Trp/-Leu (DDO), and the

interactions are tested with SD/-Trp/-Leu/-His/-Ade (QDO). **(d)** Co-immunoprecipitation assay

showing the interaction between BtRDP and NtRLP4. Total proteins are extracted from *N.*

benthamiana leaves transiently co-expressing NtRLP4-myc/NtCf9-myc with BtFTSP-flag/BtRDP-

flag. All genes are expressed with a complete coding region, and the myc/flag tags are fused at the

C-terminus. Precipitation is performed using flag beads. The samples are probed with anti-flag and

anti-myc antibodies for immunoblotting analysis. **(e–h)** Analysis of transgenic *N. tabacum*

overexpressing NtRLP4 (oeRLP). **(e)** Detection of NtRLP4 level in oeRLP plants. The empty vector

(EV) plant is used as a control. Two independent oeRLP lines are selected. The samples are probed

with an anti-NtRLP4 antibody. Rubisco staining (RbcL) is used to visualize the amount of sample

loading. **(f)** Attraction of oeRLP#1 and EV leaves to *B. tabaci* in a two-choice equipment. A group

of 40 female *B. tabaci* are released into a device containing oeRLP#1 and EV leaves. The number

of insects settling on each leaf is counted at each time point. After 48 h, the number of eggs on each

leaf is counted. **(g)** Comparison of insect reproduction on oeRLP#1 and EV plants. Five *B. tabaci*

individuals are confined to indicated plants for 3 days, and the oviposited eggs are counted. **(h)**

Relative transcript levels of salicylic acid (SA)- and jasmonic acid (JA)-associated genes in

oeRLP#1 and EV plants. PAL, phenylalanine ammonia lyase; NPR1, nonexpressor of pathogenesis-

related protein 1; FAD7, fatty acid desaturase 7; PDF1.2, plant defensin 1.2. Two independent

biological replicates are performed in **(d)** and **(e)**. Data in **(f)**, **(g)**, and **(h)** are presented as mean

values ± SEM. For insect bioassays in **(f)** and **(g)**, n= 10 independent biological replicates. For qRT-

PCR in **(h)**, n= 3 independent biological replicates. *P*-values are determined by two-tailed unpaired

Student's *t* test. ***P < 0.001; **P < 0.01; *P < 0.05; ns, not significant.

Fig. 4. BtRDP suppress plant defenses by promoting NtRLP4 degradation. (a) Effects of

ds*BtRDP* suppression on insect reproduction (n= 10 independent biological replicates) when

feeding on empty vector (EV) and NtRLP4-silenced (*RNAi-RLP*) transgenic *Nicotiana tabacum*. **(b)**

Transient overexpressing BtRDP-mCherry attenuates H_2O_2 accumulation caused by NtRLP4-GFP

overexpression. *N. tabacum* co-expressing RFP-mCherry and NtRLP4-GFP is used as a control. All

genes are expressed with a complete coding region. The experiment is repeated 5 times with the

similar results. **(c)** Relative transcript level of salicylic acid (SA)- and jasmonic acid (JA)-associated

genes in BtRDP-mCherry/NtRLP4-GFP and RFP-mCherry/NtRLP4-GFP plants (n=3 independent

biological replicates). PAL, phenylalanine ammonia lyase; NPR1, nonexpressor of pathogenesis-

related protein 1; FAD7, fatty acid desaturase 7; PDF1.2, plant defensin 1.2. **(d)** Comparison of

insect reproduction (n= 16 independent biological replicates) on transient-expressed BtRDP-

mCherry/NtRLP4-GFP and RFP-mCherry/NtRLP4-GFP plants. **(e)** Attraction of BtRDP-

mCherry/NtRLP4-GFP and RFP-mCherry/NtRLP4-GFP leaves to *B. tabaci* in a two-choice

equipment (n= 16 independent biological replicates). The number of eggs on each leaf is counted at

48 h post insect release. **(f)** Effects of BtRDP and BtRDP^{-sp} overexpression on NtRLP4

accumulation. Transgenic oeBtRDP (left) and oeBtRDP^{-sp} (right) plants are probed with anti-

NtRLP4 and anti-flag antibodies for immunoblotting analysis. The experiments are repeated twice

with the similar results. **(g)** Degradation of NtRLP4 by BtRDP in *N. benthamiana* leaves. NtRLP4-

myc and BtRDP-flag are transiently co-expressed in *N. benthamiana* plants through *Agrobacterium*

infiltration. *Agrobacterium* carrying NtCf9-myc and GUS-flag are used as negative controls. **(h)**

Effects of 26S proteasome inhibitor (MG132) on NtRLP4 accumulation. Co-infiltrated leaves are

treated with MG132 at 24 h post injection. The samples are probed with anti-flag and anti-myc

antibodies for immunoblot analysis. Rubisco staining (RbcL) is conducted to visualize the amount

of sample loading. The small triangle indicates the different concentrations of *Agrobacterium*

(OD_{600} = 0.05, 0.3, and 1.0). **(i)** NtRLP4 is ubiquitinated *in planta*. NtRLP4-myc is co-expressed

transiently with HA-UBQ in *N. benthamiana* leaves. Extracted total proteins are

immunoprecipitated by anti-myc beads and immunoblotted with anti-myc or anti-HA antibody.

Experiments are repeated thrice for **(g)** and **(h)**, while twice for **(i)**. Band density is measured using

ImageJ. The density values from three biological replicates are calculated and the mean value in the

first lane is set at 1.0. Data are presented as mean values ± SEM. *P*-values are determined by two-

tailed unpaired Student's *t* test. ***P < 0.001; **P < 0.01; *P < 0.05; ns, not significant.

Fig. 5. Rice RLP4 are targeted by salivary protein NlSP694 from *Nilaparvata lugens*. (a)

Domain organization of *Oryza sativa* RLP4 (OsRLP4). **(b, c)** Yeast two-hybrid and Co-IP assays

showing the interaction between OsRLP4 and NlSP694. All genes are expressed with a complete

coding region in Co-IP assays, while NlSP694 without a signal peptide is used in yeast two-hybrid

assays. Experiments in **(b)** are repeated twice with the similar result. **(d)** The expression patterns of

NlSP694 in different *N. lugens* tissues. Te, testis; Ov, ovary; SG, salivary gland; Ca, carcass; FB, fat

body. Data are presented as mean values ± SEM (n= 3 independent biological replicates). Different

lowercase letters indicate statistically significant differences at *P* < 0.05 level according to one-way

ANOVA test followed by Tukey's multiple comparisons test. **(e-g)** Effects of dsRNA treatment on

insect survivorship **(e)**, reproduction **(f)**, and honeydew excretion **(g)**. For survivorship analysis, a

group of 30 *N. lugens* are reared in a cage. Three independent biological replications are performed.

Differences in survivorship between the two treatments are tested by log-rank test. ns, not significant.

For reproduction analysis, n= 10 and 13 individuals are tested in ds*GFP*- and ds*NlSP694*-treatment,

respectively. The sterile females are excluded from data analysis. For honeydew analysis, n= 10

independent biological replicates. *P*-values in **(f)** and **(g)** are determined by two-tailed unpaired

Student's *t* test. ****P* < 0.001; ***P* < 0.01. **(h)** Effect of NlSP694 on the accumulation of transient-

expressed OsRLP4. OsRLP4-myc is agro-injected into *Nicotiana benthamiana* together with

different concentrations of NlSP694-flag and NlSP7-flag. NtCf9-myc is used as a negative control.

The small triangle indicates the different concentrations (OD_{600} = 0.05, 0.3, and 1.0) of

Agrobacterium. Rubisco staining (RbcL) is conducted to visualize the amount of sample loading.

Experiments are repeated three times with the similar results. Band density is measured using

ImageJ. The density values from three biological replicates are calculated with the mean value in

the first lane being set at 1.0.

Fig. 6. The proposed model for the suppression of receptor-like proteins (RLPs)-mediated

plant defenses by salivary effectors. Host plants employ pattern-recognition receptors (PRRs) to

detect various damage-associated molecular patterns (DAMPs) and herbivore-associated molecular

patterns (HAMPs) triggered by insect feeding. The RLP4/SOBIR1 complex plays a vital role in

initiating pattern-triggered immunity (PTI), including H_2O_2 burst, upregulation of jasmonic acid

(JA), and downregulation of salicylic acid (SA), which hinders insect feeding. The whitefly *Bemisia*

tabaci and planthopper *Nilaparvata lugens* independently evolved salivary proteins that targeted

plant RLP4. *B. tabaci* salivary sheath protein BtRDP interacts with the leucine-rich repeat (LRR)

domain of RLP4 from *Nicotiana tabaci* and *Solanum lycopersicum*, while *N. lugens* NlSP694 targets

both the LRR domain and the malectin-like domain of *Oryza sativa* RLP4. These interactions

promote the ubiquitin-dependent degradation of RLP4, thereby disrupting the stability of the

RLP4/SOBIR1 complex. The presence of salivary effectors causes a hormonal shift and suppresses

the H_2O_2 burst, finally favoring insect feeding.

a Transcript level — $P=0.004$, ** ; dsGFP, ds*BtRDP*

α-BtRDP — 33kDa, 26kDa
α-Actin — 55kDa, 43kDa

b Eggs by 5 females — $P<2\times10^{-5}$, *** ; dsGFP, ds*BtRDP*

c Duration time (min) — np ($P=0.279$, ns), C ($P=0.024$, *), E1 ($P=0.987$, ns), E2 ($P=0.007$, **)

Plants overexpressing BtRDP

d α-flag — 43kDa, 33kDa ; RbcL — 55kDa, 43kDa ; EV, oeBtRDP #1 #2

e Eggs by 5 females — $P=0.012$, * ; EV, oeBtRDP#1

f Number of whiteflies — oeBtRDP#1, EV
3h ($P=0.093$, ns), 6 ($P=0.002$, **), 12 ($P=0.0006$, ***), 24 ($P=0.0001$, ***), 36 ($P=4\times10^{-5}$, ***), 48 ($P=0.0001$, ***)
Eggs per leaf — $P=0.0003$, *** ; EV, oeBtRDP#1

Plants overexpressing BtRDP^{-sp}

g α-flag — 43kDa, 33kDa ; RbcL — 55kDa, 43kDa ; EV, oeBtRDP^{-sp} #1 #2

h Eggs by 5 females — $P=0.002$, ** ; EV, oeBtRDP^{-sp}#1

i Number of whiteflies — oeBtRDP^{-sp}#1, EV
3h ($P=0.006$, **), 6 ($P=0.002$, **), 12 ($P=0.007$, **), 24 ($P=0.0001$, ***), 36 ($P=0.0001$, ***), 48 ($P=0.028$, *)
Eggs per leaf — $P=0.002$, ** ; EV, oeBtRDP^{-sp}#1

a

NtRLP4 SP Malectin-like LRR TM
1 22 31   366 375   541 573 625

b

	Untreat	12h	24h	48h
α-NtRLP4	1.00	0.70±0.14	0.42±0.08	0.40±0.10

55kDa — α-NtRLP4

55kDa — RbcL

c

BD-BtRDP^{BDP} ×

	DDO	×10	×10	×10²	×10³
AD-NtRLP4$_{(23-541)}$					
AAD-NtRLP4$_{(31-366)}$					
AD-NtRLP4$_{(375-541)}$					
AD-NtRLP4$_{(573-625)}$					
AD-NtCf9$_{(26-728)}$					

DDO   QDO

d

NtRLP4-myc	+	+	−
NtCf9-myc	−	−	+
BtFTSP-flag	+	−	−
BtRDP-flag	−	+	+

33kDa — α-flag
55kDa — α-myc

IP   Input

e

	EV	oeRLP #1 #2

55kDa — α-NtRLP4

55kDa / 43kDa — RbcL

f

oeRLP#1 ● / EV ■

Number of whiteflies — 3, 6, 12, 24, 36, 48(h)
*P=0.016, ***P=3×10⁻⁴, **P=0.002, ***, ***, ***P=3×10⁻⁴

Eggs per leaf — EV vs oeRLP#1 ***P=0.0005

g

Eggs by 5 females — EV vs oeRLP#1 ***P=2×10⁻⁴

h

Relative transcript level — EV ■ / oeRLP#1 ■

	PAL	NPR1	FAD7	PDF1.2
	P=0.007 **	P=0.009 **	P=0.014 *	P=0.50 ns

SA-associated   JA-associated