Herbivorous insects independently evolved salivary effectors to regulate plant immunity by destabilizing the malectin-LRR RLP NtRLP4

Xin Wang[1,2], Jia-Bao Lu[1], Yi-Zhe Wang[1], Xu-Hong Zhou[3], Jian-Ping Chen[1,2], Chuan-Xi Zhang[1], Jun-Min Li[1,#], Hai-Jian Huang[1,#]

[1] State Key Laboratory for Quality and Safety of Agro-Products, Key Laboratory of Biotechnology in Plant Protection of MARA, Zhejiang Key Laboratory of Green Plant Protection, Institute of Plant Virology, Ningbo University, Ningbo 315211, China

[2] Institute of Applied Ecology, Fujian Agriculture and Forestry University, Fuzhou 350002, China

[3] Office of Science and Technology, Yunnan University of Chinese Medicine, Kunming 650205, China

Abstract

Plants depend on receptor-like proteins and receptor-like kinases (RLPs/RLKs) to detect a variety of invading pathogens and insects. Although the mechanisms of microbial pathogens in suppressing plant immunity have been well investigated, it remains unclear how herbivorous insects overcome receptor-mediated defenses. Here, we show that salivary effectors evolve independently in whiteflies and planthoppers to target defensive RLP4, therefore minimizing the plant immunity. RLP4, as a leucine-rich repeat RLP (LRR-RLP), confers plant resistance against herbivorous insects by forming the RLP4/SOBIR1 complexes. In the whitefly *Bemisia tabaci*, BtRDP, the Aleyrodidae-specific salivary sheath protein, interacts with RLP4 from multiple plant species and promotes its ubiquitin-dependent degradation. Overexpression of NtRLP4 in transgenic plants exerts a detrimental impact on *B. tabaci* by exploiting the crosstalk between the salicylic acid (SA) and jasmonic acid (JA) pathways. Conversely, overexpression of BtRDP or silencing of NtRLP4 effectively alleviates such negative effects. In planthopper *Nilaparvata lugens*, the Delphacidae-

* Corresponding authors: Jun-Min Li, lijunmin@nbu.edu.cn; Hai-Jian Huang, huanghaijian@nbu.edu.cn

28   restricted salivary protein NlSP104 also targets and promotes the degradation of OsRLP4 from rice

29   plants. Our results reveal the convergent evolution of salivary proteins in insects, and highlight the

30   complex interactions between plants and herbivorous insects.

32   **Keywords**

33   salivary effector, insect-plant interaction, receptor-like protein, convergent evolution

Introduction

Plants have evolved sophisticated defense systems to survive the constant attacks of pathogens

and herbivorous insects. These defenses function at multiple levels, encompassing physical barriers

such as the cuticle and cell wall, chemical defenses that include toxic secondary metabolites and

anti-nutritive compounds, and indirect defenses that involve the emission of herbivore-induced

volatiles to attract natural enemies of the herbivores [1, 2]. Beyond these general strategies, plants

also rely on the highly specialized molecular immune responses to detect and rapidly respond to

invaders. The activation of innate immunity by membrane-localized receptors is a highly conserved

mechanism among eukaryotes [3]. In sessile plants, the first line of defense against potential

pathogenic microbes or insects relies on cell surface pattern-recognition receptors (PRRs) [4]. The

plant genome encodes hundreds of PRRs, including receptor-like kinases (RLKs) and receptor-like

proteins (RLPs) [4]. RLKs are consisted of different types of N-terminal ectodomains, a single

transmembrane domain, and a cytoplasmic protein kinase domain, while RLPs have a similar

structure but lack the kinase domain [5, 6]. As RLPs lack the intracellular signaling domains, they

usually bind to SOBIR1 or SOBIR1-like adaptors to form the bimolecular receptor kinases [6].

Despite the large number of PRRs encoded by plant genomes, only a few of them have been well

characterized regarding their functions in recognizing microbial-associated molecular patterns

(MAMPs), damage-associated molecular patterns (DAMPs), and herbivore-associated molecular

patterns (HAMPs) [5, 7]. For instance, FLS2 is known to recognize bacterial flagellin [8], EFR can

recognize the bacterial elongation factor Tu [9], and EIX2 is responsible for the recognition of the

fungal ethylene-inducing xylanase [10].

Piercing-sucking insects, such as whiteflies, aphids, and planthoppers, damage plants by

feeding or transmitting viruses. They inject a mixture of saliva into plant tissues, which enhances

their feeding efficiency by binding to leaked calcium, degrading extracellular DNA, and regulating plant hormonal signals [11-14]. In response, plants have evolved the ability to detect herbivores by sensing HAMPs present in insect saliva [7, 15]. This triggers pattern-triggered immunity (PTI), including mitogen-activated protein kinase (MAPK) cascades, reactive oxygen species (ROS), and hormone signaling [16-18]. A few salivary proteins have been reported to activate plant PTI, but the specific PRRs that recognize these proteins and trigger plant immunity are largely unclear [19-21]. One exception is the inceptin receptor (INR), which is a legume-specific LRR-RLP that can recognize the Vu-In in insect oral secretions [7]. Moreover, plant PRRs have been extensively reported to be targeted by pathogen effectors [22-24]. However, it remains unknown how insects cope with the immune recognition system.

The whitefly *Bemisia tabaci* (Hemiptera: Aleyrodidae) and brown planthopper *Nilaparvata lugens* (Hemiptera: Delphacidae) are notorious pests globally. During their feeding process, both insects secrete abundant species-specific salivary proteins into host plants [25, 26]. It is reported that plants initiate defense responses when they recognize insect feeding [19, 27]. However, the specific PRRs responsible for this process are still unknown. Both insects can successfully ingest phloem saps with limited plant defenses. At present, it remains unknown how insects attenuate plant defenses. Given that numerous defense mechanisms are conserved across plants [3, 28], it is interesting to investigate whether insect evolves a similar strategy to aid feeding during convergent evolution. In this study, the functions of RLPs in the plant immunity responsive to piercing-sucking insects were investigated, and *B. tabaci* and *N. lugens* were demonstrated to independently evolve salivary proteins to regulate the post-translational modification of RLPs.

Results

BtRDP is most abundantly expressed in salivary glands and secreted into host plants

Based on the transcriptomic results, BtRDP (*B. tabaci* RLP-degrading protein; GenBank accession: MN738093) was the most abundant gene in salivary glands [26]. It was a secretory protein that contained a signal peptide at the N-terminal, and totally 6 unique peptides from BtRDP were detected in the *B. tabaci* watery saliva (Fig. S1). As there was no gene homolog identified in the NCBI nr database, BtRDP was subjected to BLAST analysis against the transcriptomic or genomic data of 30 insect species. RDP was distributed in all the analyzed Aleyrodidae species,

while no RDP homolog was detected in other species (Table S1). Phylogenetic analysis indicated

that RDPs from five *B. tabaci* cryptic species were clustered in the same clade, exhibiting a close

correlation with RDP from *Aleurocanthus spiniferus* (Fig. S2).

A group of *B. tabaci* were allowed to feed on *Nicotiana tabacum* plants for 24 h, and the

presence of BtRDP in the infested and non-infested samples were assayed. One band between 26-

33 kDa was detected in the *N. tabacum* plants infested by *B. tabaci*, but not in the non-infested

plants (Fig. 1a). This band matched the molecular weight of BtRDP identified in the salivary glands.

The detection of BtRDP in the infested plants (Fig. 1a) and in watery saliva (Fig. S1) collectively

indicated that BtRDP was a salivary protein. According to Western-blotting, qPCR and

transcriptomic analysis, BtRDP was nearly exclusively expressed in the salivary glands (Fig. 1b, c;

Fig. S3a), which was significantly different from the actin or ribosomal 18S rRNA (Fig. S4a). In

addition, BtRDP was highly expressed in the adult stages, but lowly expressed in the egg, nymph,

and pseudopupa stages (Fig. 1d; Fig. S3b), similar to another salivary protein BtFTSP [29] but

different from actin and ribosomal 18S rRNA (Fig. S4b). According to immunohistochemical (IHC)

staining, BtRDP was specifically distributed in the principle salivary gland, but not in the accessory

salivary gland or other tissues (Fig. 1e). For most piercing-sucking insects, two types of saliva (gel

and watery saliva) are secreted into plant tissues. The salivary sheath, formed from gel saliva, serves

as an insoluble lining along the stylet path, potentially providing a scaffold for effector delivery [30].

The salivary sheath secreted by *B. tabaci* was also stained with the anti-BtRDP serum. The BtRDP

signal was mainly detected in the protuberant structure of sheath (Fig. 1f), suggesting that BtRDP

attaches to salivary sheath after secretion.

BtRDP is important for *B. tabaci* feeding on tobacco plants

RNA interference was conducted to investigate the role of BtRDP in whitefly feeding.

Treatment with ds*BtRDP* efficiently and specifically suppressed the target gene at both the transcript

and protein levels (Fig. 2a; Fig. S5). There was no significant difference in the morphology of the

salivary sheath between ds*BtRDP*-treated and ds*GFP*-treated *B. tabaci* (Fig. S6a). However, the

salivary sheath length of ds*BtRDP*-treated *B. tabaci* was significantly shorter than that of the

ds*GFP*-treated control (Fig. S6b). Furthermore, treatment with ds*BtRDP* did not affect the

survivorship of whiteflies (Fig. S6c). Upon conducting a reproduction analysis, it was found that

the ds*BtRDP*-treated whiteflies oviposited fewer eggs than the control (Fig. 2b). The feeding

behavior of whiteflies was monitored using the electrical penetration graph (EPG) technique, which

distinguishes nonpenetration (np), pathway (C), phloem salivation (E1), and phloem ingestion (E2)

phases [31] (Fig. S7). Compared to the ds*GFP* control, ds*BtRDP*-treated *B. tabaci* exhibited a

significant reduction in phloem ingestion and an extended pathway duration, indicating that BtRDP

is necessary for efficient feeding (Fig. 2c).

Salivary proteins are reported to function in intercellular space and/or extracellular space after

being secreted into host plants [30, 32]. To investigate the role of BtRDP in different subcellular

locations of host plants, we constructed two transgenic *N. tabacum* lines overexpressing BtRDP:

one carrying the full-length coding sequence that included the signal peptide (oeBtRDP), which is

expected to be secreted into the apoplast (extracellular space), and the other lacking the signal

peptide (oeBtRDP^{-sp}), which is likely to be retained in the cytoplasm. The flag tag was fused to the

C-terminal ends of the recombinant proteins, respectively. Transgenic plants containing the empty

vector (EV), which exerted no significant influence on insect performance compared with the WT

(Fig. S8), were used as negative controls. Fecundity and two-choice assays demonstrated that

BtRDP, whether localized in the apoplast (oeBtRDP) or cytoplasm (oeBtRDP^{-sp}), enhanced whitefly

settling and oviposition relative to the EV controls (Fig. 2d-i; Fig. S9), indicating that BtRDP

promotes whitefly feeding behavior irrespective of its subcellular location.

BtRDP interacts with the NtRLP4/NtSORBIR1 complex

To investigate the underlying mechanism of BtRDP in improving whitefly feeding, yeast two-

hybrid (Y2H) screening against a *N. benthamiana* cDNA library was conducted using BtRDP^{-sp} as

a bait. Totally seven proteins were found to potentially interact with BtRDP^{-sp} (Table S2), including

an NbRLP4 (accession Niben261Chr07g1310001.1 in Niben261 genome). NbRLP4 possesses a

predicted N-terminal signal peptide, a malectin-like domain, a LRR domain, and a transmembrane

domain. NbRLP4, as well as its closest homology in *N. tabacum* (NtRLP4; GenBank accession:

XM_016601109), did not contain the kinase domain (Fig. 3a).

The relative expression of *NtRLP4* in response to *B. tabaci* infestation was investigated. Based

on qRT-RCR results, the transcript level of *NtRLP4* was not significantly influenced by *B. tabaci*

(Fig. S10). However, a decrease in the protein level of NtRLP4 was detected by Western-blotting

assay (Fig. 3b). Point-to-point Y2H assays revealed that NtRLP4$_{(23-541)}$ (a truncated version lacking

the signal peptide and transmembrane domains) interacted with BtRDP^{-sp} (Fig. 3c). To investigate

the specificity of the NtRLP4-BtRDP interaction, another RLP protein NtCf9 with the highest

sequence similarity to tomato resistance protein Cf9 (Fig. S11) [33], as well as three randomly

selected salivary proteins from *B. tabaci* (BtFTSP, QHB15613; BtSP16.3, MN738005; BtSP37.4,

MN738158), were used as controls. As a result, BtRDP^{-sp} did not interact with the truncated NtCf9

construct, in which the signal peptide and transmembrane domains were removed (Fig. 3c). Also,

NtRLP4$_{(23-541)}$ could not interact with the other three salivary proteins without signal peptides (Fig.

S12a, b). These findings raised the question of which domain of NtRLP4 is responsible for binding

to BtRDP, as identifying the interacting domain can help to infer where the salivary protein contacts

the receptor in plants. We therefore dissected the NtRLP4 domains accordingly. The results showed

that it was the LRR domain, rather than the malectin-like domain or the short intercellular region,

that was required for the NtRLP4-BtRDP interaction (Fig. 3c). To investigate the shortest LRR

sequence responsible for BtRDP interaction, the LRR domain was divided into three segments.

However, none of these segments interacted with BtRDP^{-sp} (Fig. S12a). Co-immunoprecipitation

(Co-IP) assays further confirmed the specific interaction between BtRDP and NtRLP4 (Fig. 3d). It

was the C-terminal myc tag-fused NtRLP4 (NtRLP4-myc), but not the NtCf9 (NtCf9-myc), that

could be immunoprecipitated by the C-terminal flag tag fused BtRDP (BtRDP-flag). In the

bimolecular fluorescence complementation (BiFC) assay, YFP fluorescence was observed upon co-

expression of the N-terminal nYFP tag fused NtRLP4 and N-terminal cYFP tag fused BtRDP^{-sp} (Fig.

S12c). Furthermore, in oeBtRDP transgenic plants, endogenous NtRLP4 was specifically

immunoprecipitated with BtRDP-flag, which was significantly different from the control (Fig.

S12d). These results suggest that BtRDP interacts with NtRLP4.

The absence of kinase motifs in the C-terminal region of NtRLP4 suggests that it may function

in cooperation with other proteins to initiate signaling. The receptor-like kinase SOBIR1, which

contained a kinase domain (Fig. S13a), has been widely reported to be required for the function of

RLPs in the innate immunity [6]. Through Co-IP assays, our study demonstrated that NtSOBIR1

was capable of interacting with NtRLP4 but not BtRDP (Fig. S13b, c).

NtRLP4 and NtSOBIR1 are resistant to *B. tabaci* infestation

The transgenic *N. tabacum* overexpressing NtRLP4 (oeRLP#1 and oeRLP#2, without any tags)

was constructed (Fig. 3e). Compared with the EV control, both oeRLP lines were less attractive to

B. tabaci, alongside less settling and ovipositing insects (Fig. 3f; Fig. S14a). Also, *B. tabaci* had a

significantly decreased fecundity in oeRLP plants (Fig. 3g; Fig. S14b). Similar results were

observed after transient overexpression of NtRLP4-GFP in *N. tabacum* (Fig. S14c, d), indicating

that NtRLP4 is resistant to *B. tabaci* infestation.

Thereafter, the EV and oeRLP#1 plants underwent transcriptomic sequencing (Fig. S15a). A total

of 729 differentially expressed genes (DEGs) were identified, comprising 423 up-regulated and 306

down-regulated genes (Table S3). Enrichment analysis demonstrated that the majority of up-

regulated DEGs were associated with plant-pathogen interaction, environmental adaptation, MAPK

signaling pathway, and signal transduction (Fig. S15b). In contrast, the glutathione metabolism,

carbohydrate metabolism, and amino acid metabolism pathways were significantly enriched in

down-regulated DEGs (Fig. S15c). Noteworthily, numerous DEGs were annotated as RLK/RLP or

WRKY transcription factor, with the majority of them being significantly upregulated (Fig. S15d,

e). These results suggested an enhanced defense response in oeRLP plants. The crosstalk between

salicylic acid (SA) and jasmonic acid (JA) pathways was critical for plant defenses [14]. The

increased defense of oeRLP transgenic plants was partially caused by the altered hormonal signaling.

The JA-associated gene *FAD7* was significantly induced, while the SA-associated genes *PAL* and

NPR1 were repressed in oeRLP plants (Fig. 3h). Transient overexpression of NtRLP4-GFP in *N.*

tabacum plants via *Agrobacterium* infiltration exerted similar effects (Fig. S14e). Furthermore,

NtRLP4-GFP significantly enhanced H_2O_2 accumulation (Fig. S14f).

The function of NtSOBIR1 was also investigated by transient overexpression and hairpin RNAi

(Fig. S16). Overexpression of NtSOBIR1-GFP induced the cell death phenotype five days post agro-

injection (Fig. S16). Insects preferred to settle and oviposit on GFP-expressed *N. tabacum* than on

NtSOBIR1-GFP-expressed control, even though no obvious death phenotype was observed two

days post agro-injection (Fig. S16c, d). For NtSOBIR1 silencing, no cell death phenotype was

observed, while the NtSOBIR1-silenced leaves exhibited reduced growth (Fig. S16f). Compared

with the EV-silenced control, NtSOBIR1-silenced *N. tabacum* was slightly attractive to *B. tabaci*

(Fig. S16g, h). Collectively, these results demonstrate that NtSOBIR1 is resistant to *B. tabaci*

infestation.

BtRDP attenuates plant defenses by promoting the degradation of host RLP4

To investigate the interplay between NtRLP4 and BtRDP, transgenic *N. tabacum* silencing NtRLP4 (*RNAi-RLP*#1 and *RNAi-RLP*#2) plants were constructed. Both *RNAi-RLP* lines showed reduced NtRLP4 levels compared with EV plants, with *RNAi-RLP*#2 exhibiting a stronger silencing effect (Fig. S17a). Compared with EV plants, *B. tabaci* showed a tendency to settle on NtRLP4-silenced plants and lay more eggs (Fig. S17b, c). Thereafter, ds*GFP*- and ds*BtRDP*-treated *B. tabaci* were reared on EV, *RNAi-RLP*#1, and *RNAi-RLP*#2, respectively. The ds*BtRDP*-treated *B. tabaci* produced less offspring than the control (decreased by 28.3%; *P*=0.0007) on EV plants (Fig. 4a), consistent with the results obtained from WT plants (Fig. 2b). On the *RNAi-RLP*#1 plant, the impaired insect reproduction was still observed after ds*BtRDP* treatment, but the extent was slightly reduced (decreased by 23.7%; *P*=0.002) (Fig. 4a). On the *RNAi-RLP*#2 plant, there was no significant difference between the ds*GFP* and ds*BtRDP* treatments (Fig. 4a). The differential rescue effect between the two RNAi lines likely resulted from their different NtRLP4 silencing efficiencies, with the lower NtRLP4 level in *RNAi-RLP*#2 leading to a more complete rescue phenotype (Fig. S17a; Fig. 4a). Collectively, these results suggest that *RNAi-RLP* plants are beneficial for insects and can partially, but not completely, rescue the impaired feeding performance of ds*BtRDP*-treated insects.

To investigate the influence of BtRDP on NtRLP4, GFP-tagged NtRLP4 (NtRLP4-GFP) was transiently co-expressed with mCherry-tagged BtRDP (BtRDP-mCherry), red fluorescent protein (RFP)-mCherry, and BtFTSP1-mCherry through agroinfiltration, respectively. All tags were fused with the C-terminal region. As discovered, BtRDP-mCherry did not influence the location patterns of NtRLP4-GFP (Fig. S18). However, a significant reduction in the fluorescent signal of NtRLP4-GFP was detected after co-expression with BtRDP-mCherry, but not with the controls (Fig. S18). Besides, co-expression with BtRDP-mCherry/NtRLP4-GFP significantly reduced H_2O_2 accumulation (Fig. 4b), suppressed the expression of JA-associated genes *FAD7* and *PDF.12*, and induced that of the SA-associated genes *PAL* and *NPR1* (Fig. 4c). It was discovered by insect bioassays that, transient overexpression of BtRDP-mCherry rescued the negative effects on whitefly performance caused by NtRLP4-GFP overexpression (Fig. 4d, e). In the host choice experiments, *B. tabaci* displayed a preference for settling on *N. tabacum* leaves co-expressing BtRDP-

mCherry/NtRLP4-GFP over those co-expressing RFP-mCherry/NtRLP4-GFP (Fig. 4d). However, almost no preference was observed in *B. tabaci* feeding on *N. tabacum* plants co-expressing BtRDP-mCherry/NtRLP4-GFP and plants expressing GFP alone (Fig. S19).

The attenuated NtRLP4-GFP fluorescent signal (Fig. S18), along with the observed decrease in the protein level of NtRLP4 during *B. tabaci* infestation (Fig. 3b), led us to speculate on whether BtRDP influenced the accumulation of NtRLP4. To this end, the BtRDP^{-sp}-his and GFP-his proteins were prokaryotically expressed in *Escherichia coli*. Different concentrations of purified BtRDP^{-sp}-his and GFP-his proteins were infiltrated into *N. tabacum* plants overexpressing NtRLP4-myc. Interestingly, the protein level of NtRLP4-myc slightly decreased with the increasing amount of BtRDP^{-sp}-his, while no such effect was observed with GFP-his (Fig. S20). Moreover, the BtRDP^{-sp}-his protein did not have any significant effect on the accumulation of another RLP NtCf9-like (Fig. S20). We further assessed the impact of BtRDP on NtSOBIR1 following NtRLP4 destabilization. As the amount of BtRDP-flag increased, NtRLP4-myc accumulation was markedly reduced, while NtSOBIR1-flag levels remained unchanged (Fig. S21).

Then, the NtRLP4 levels in oeBtRDP and oeBtRDP^{-sp} plants were detected. Compared with the EV control, a reduced NtRLP4 abundance was observed in both oeBtRDP#1 and oeBtRDP#2 transgenic lines, but not the oeBtRDP^{-sp} plants (Fig. 4f), suggesting the importance of BtRDP secretion in extracellular space. BtRDP attached to salivary sheath that was localized in plant apoplast (Fig. 1f). The presence of signal peptide promoted the secretion of BtRDP into the extracellular space. Therefore, the complete coding region of *BtRDP* was used in subsequent experiments. The NtRLP4-myc and BtRDP-flag were transiently co-expressed in *N. benthamiana* plants through *Agrobacterium* infiltration. As a result, BtRDP-flag significantly decreased the protein level of NtRLP4-myc, but not NtCf9-myc (Fig. 4g). qRT-PCR analyses demonstrated that the transcript level of *NtRLP4* remained unchanged (Fig. S22). These findings indicate that the reduction in the protein level of NtRLP4 is not caused by transcriptional changes but rather by degradation.

The ubiquitin system and autophagy are two major pathways regulating protein degradation [34]. At first, MG132, a 26S proteasome inhibitor known to inhibit the degradation of ubiquitin-conjugated proteins, was employed [35]. The results showed that MG132 efficiently prevented the degradation of NtRLP4-myc by BtRDP-flag (Fig. 4h). Then, the autophagy inhibitor 3-

Methyladenine (3-MA) and bafilomycin A1 (BAF) that can block autophagy in plants were used

[36]. However, the decreased protein level of NtRLP4-myc was still observed in the presence of 3-

MA or BAF (Fig. S23). Additionally, NtRLP4-myc and HA-tagged ubiquitin (HA-UBQ) were co-

expressed in *N. benthamiana*. Total proteins were extracted from *N. benthamiana* leaves and then

subjected to immunoprecipitation with anti-myc beads. The ubiquitin attached to NtRLP4 was

determined by immunoblot using the anti-HA antibody. Leaves expressing NtRLP4-myc or UBQ-

HA along were used as negative controls. The results showed that a strong UBQ signal was detected

in the immunoprecipitated product only when NtRLP4-myc and UBQ-HA were co-expressed (Fig.

4i). The molecular weight of UBQ-attached NtRLP4-myc was remarkably larger than NtRLP4-myc

itself (Fig. 4i). These results suggest that BtRDP promotes NtRLP4 degradation through the

ubiquitin system, but not the autophagy pathway.

Degrading plant RLP4s by salivary effectors is a common strategy for insects

To investigate the ability of BtRDP to degrade RLPs from other host plants of *B. tabaci*,

Solanum lycopersicum RLP4 (SlRLP4; GenBank accession: XP_004232910) was selected for

further analysis. Sequence analysis revealed a similar gene structure between SlRLP4 and NtRLP4

(Fig. S24; Fig. S25a). Moreover, the interaction between BtRDP and SlRLP4 was confirmed

through Y2H and Co-IP assays (Fig. S25b, c). Additionally, SlRLP4-myc and SlCf9-like-myc were

transiently co-expressed with varying concentrations of BtRDP-flag or GUS-flag, with the myc and

flag tags fused with the C-terminus. The results showed that BtRDP-flag exerted no influence on

the accumulation of SlCf9-like-myc protein. In contrast, the protein level of SlRLP4-myc was

significantly reduced in the presence of BtRDP-flag (Fig. S25d), indicating the conserved function

of BtRDP in *B. tabaci* adaptation to different host plants.

BtRDP was a gene specific to the Aleyrodidae family, lacking homologues in other species.

To explore whether the degradation of RLPs by salivary effectors was a conserved mechanism

among different herbivorous insects, a RLP4 homology from *Oryza sativa* (OsRLP4; GenBank

accession: XP_015645303; Fig. 5a) and twenty salivary proteins from planthopper species lacking

signal peptides were paired to identify the potential interactions using Y2H assays. As a result, only

the salivary protein NlSP104^{-sp} (*N. lugens* salivary protein consisting of 104 amino acids, GenBank

accession: MF278694) interacted with OsRLP4$_{(29-551)}$ (OsRLP4 lacking signal peptides and

transmembrane domains), while the other salivary proteins tested did not show any interaction (Fig.

5b, c; Fig. S26). NlSP104 did not exhibit any sequence or structural similarity to BtRDP (Fig. S27).

Truncation analysis revealed that NlSP104^{-sp} interacted with both the LRR domain and the malectin-

like domain of OsRLP4 (Fig. 5c), which was slightly different from the BtRDP-NtRLP4 interaction

(Fig. 3c). Given that both BtRDP and NlSP104 interacted with the LRR domain of RLP4, further

investigation focused on the specific region of the LRR domain responsible for NlSP104 binding.

The LRR domain of OsRLP4 was divided into three segments, mirroring the division of the LRR

domain in NtRLP4 (Fig. S12a). NlSP104^{-sp} interacted with the truncated LRR domain (OsRLP4$_{(480-}$

$_{526)}$) (Fig. 5c). In contrast, none of the truncated LRR domain of NtRLP4 interacted with BtRDP^{-sp}

(Fig. S12a). These findings suggest that the binding sites for salivary proteins to RLP4 may differ

between *N. lugens* and *B. tabaci*.

NlSP104 was a planthopper-specific salivary protein nearly exclusively expressed in salivary

glands (Table S1; Fig. 5d; Fig. S28). The function of NlSP104 or its gene homologs in other

planthopper species has not been well investigated previously. Silencing NlSP104 did not affect

insect survivorship (Fig. 5e). However, significant reductions in honeydew excretion and impaired

fecundity were observed in ds*NlSP104*-treated *N. lugens* (Fig. 5f, g), indicating the importance of

NlSP104 in insect feeding.

Subsequently, OsRLP4-myc was transiently co-expressed with a complete coding region of

NlSP104 (C-terminal flag tagged, NlSP104-flag). Another salivary protein NlSP7-flag and NtCf9-

myc were used as negative controls. It was found that NlSP104-flag specifically and significantly

decreased the protein level of OsRLP4-myc, which differed significantly from that of the NlSP7-

flag (Fig. 5h). Additionally, NlSP104-flag failed to decrease the protein level of NtCf9-myc (Fig.

5h). These findings suggest that *N. lugens* potentially evolves salivary effectors to target rice RLPs.

Discussion

Plants have evolved a complex innate immune system to protect themselves against

pathogen/insect invasion, and the cell surface-localized PRRs function on the frontline [4]. To

counteract this response, the pathogens have developed various strategies to avoid plant PRR-

mediated PTI for their own advantage [37]. However, it remains unknown whether insects take the

similar strategy to sabotage plant immunity. In this study, we demonstrate that the RLP4/SOBIR1

complexes confer plant resistance against herbivorous insects. In response, both planthoppers and

whiteflies secrete salivary effectors that facilitate the degradation of defensive RLP4, leading to the

attenuated plant immunity. Our results suggest that manipulation of PRRs may be a conserved

strategy for herbivorous insects.

PRRs detect modified-self or non-self patterns and initiate a cascade of immune signaling

events, which subsequently regulate downstream defense responses, conferring plant resistance

against invaders [4, 5]. Although various insect salivary components have been identified as triggers

for plant immune responses, the specific PRRs involved in this process remain unclear. Our study

indicates that RLP4 significantly influences insect performance by modulating plant hormones and

ROS (Fig. 3f-h; Fig. S14), highlighting the critical role of RLP4 in plant immunity. However, the

precise ligand for RLP4 has yet to be identified. The malectin-LRR receptor RLP4 may respond to

the DAMPs or HAMPs released during insect feeding [38], a hypothesis that warrants further

investigation. RLP4 lacks an intracellular kinase domain, and instead forms bimolecular receptor

kinases by interacting with SOBIR1 (Fig. S13). Overexpression of NtRLP4 or NtSOBIR1 in *N.*

tabacum attracts *B. tabaci* and promotes insect reproduction, whereas silencing of either gene exerts

the opposite effect (Fig. S14-S17). The association between RLPs and SOBIR1 protein kinases is

structurally and functionally analogous to that of genuine receptor kinases [6]. Therefore,

destabilizing the RLP4/SOBIR1 complex can potentially disrupt the immune recognition processes.

It is noteworthily that although NtRLP4 interacts with SOBIR1, this alone does not confirm that it

operates strictly through this canonical module. Evidence from other RLPs shows that co-receptor

usage can be flexible, and some RLPs function partly or conditionally independent of SOBIR1 [39].

Therefore, a more definitive assessment of NtRLP4 signaling will therefore require genetic

dissection of its co-receptor dependencies, including but not limited to SOBIR1.

As pivotal regulators in immune signaling, PRRs are frequently targeted by multiple pathogens

through diverse mechanisms. For example, *Pseudomonas syringae* hampers the functionality of

receptor complexes by disrupting the ligand-dependent association between receptor-like kinase

BAK1 and the flagellin receptor FLS2 [24], while *Phytophthora infestans* promotes alternative

splicing of the host RLPKs to enhance infection [23]. PRRs are usually subjected to degradation via

post-translational modifications [40]. In plants, the activity and stability of PRRs are under stringent

regulation since excessive activation of PRRs could be detrimental [22, 41]. One prevalent

regulatory mechanism involves ubiquitination followed by subsequent degradation mediated by the

26S proteasome [40]. Mounting evidence has demonstrated that bacteria, fungi, viruses, and

oomycetes exploit such mechanisms to facilitate PRR degradation [42-45]. Nevertheless, the

occurrence of effector-mediated PRR degradation in insects is unreported. Our study reveals that

the salivary effectors BtRDP from *B. tabaci* and NlSP104 from *N. lugens* interact with plant RLP4s,

resulting in their degradation in a ubiquitin-dependent manner (Fig. 4f-i).

Notably, BtRDP and NlSP104 exhibit neither sequence nor structural similarity and do not

resemble the known eukaryotic ubiquitin-ligase domains. Their interaction with RLP4s occurs in

the extracellular space (Fig. 3d; Fig. 5c), whereas the ubiquitin-proteasome system primarily

functions within the cytosol and nucleus [46]. Furthermore, NtRLP4 reduction is observed solely in

oeBtRDP transgenic plants, not in oeBtRDP^{-sp} plants (Fig. 4f), indicating that BtRDP exerts its

influence on NtRLP4 in the extracellular space. These observations collectively refute the

possibility that BtRDP or NlSP104 possesses the intrinsic E3 ligase activity capable of directly

ubiquitinating RLP4s within plant cells. Importantly, the reduced NtRLP4 levels may not stem from

a direct physical interaction between BtRDP and NtRLP4. Instead, BtRDP may indirectly affect

RLP4 post-translational modification, thereby accelerating its degradation, which warrants further

investigation. Additionally, the independent evolution of RLP4-targeting effectors in various insect

lineages may expedite plant-insect co-evolution. Plants with robust RLP4-mediated defenses can

select for herbivores carrying effective effectors, driving an evolutionary "arms race". Conversely,

insects targeting similar host proteins may exert parallel selective pressures on unrelated plant

species, potentially shaping defense diversification across plant communities. Understanding these

dynamics provides insight into how molecular interactions scale up to influence herbivore

adaptation, host range, and the ecological outcomes of plant-insect interactions.

Herbivorous insects employ a mix of salivary components to ensure successful feeding [30].

Previous research has indicated that *B. tabaci* suppresses plant defenses by utilizing SA-JA crosstalk,

with the salivary protein Bt56 playing a crucial role in this process [14]. Bt56 activates the SA-

signaling pathway by directly interacting with a KNOTTED 1-like homeobox transcription factor

[14]. This study investigates the effects of another salivary effector, BtRDP, on plant hormones, and

finds that it elicits similar effects to Bt56, albeit primarily impacting the signal recognition process

(Fig. 3h; Fig. S14e). This leads us to speculate that different salivary effectors from the same insect

species may simultaneously target the same signaling pathway to ensure effective immune suppression. Our analysis reveals that BtRDP interacts with other host genes apart from NtRLP4 (Table S2), and silencing NtRLP4 cannot completely rescue the impaired feeding performance of ds*BtRDP*-treated insects (Fig. 4a). Salivary proteins exert a crucial effect on mediating the interactions between herbivores and plants. The dynamic interplay between plants and insects may be highly complex, one salivary protein can target multiple plant pathways, and one plant pathway is influenced by multiple salivary proteins.

In conclusion, this study shows that suppressing PRR-mediated plant immunity may be the common strategy for the successful feeding of herbivorous insects. Whiteflies and planthoppers independently evolve salivary effectors that facilitate the degradation of defensive RLP4 in host plants in an ubiquitin-dependent manner, therefore disrupting the RLP4/SOBIR1 complexes for signal recognition (Fig. 6).

Material and methods

Insects and plants

A colony of *B. tabaci* (cryptic species MED) was originally collected from a soybean field in Suzhou, Anhui Province, China. The insects were subsequently maintained on *N. tabacum* cv. K326 plants under controlled laboratory conditions (25 $\pm$ 1 °C, 50–70% relative humidity, 16:8 h light:dark photoperiod). For experimental use, *N. benthamiana* and *N. tabacum* plants were grown in a climate-controlled growth chamber (23 ± 1 °C, 16:8 h light: dark cycle).

Phylogenetic analysis of RDP

To identify potential RDP homologs across insect species, we performed a BLASTp search using the BtRDP protein sequence as a query against the predicted proteomes of 30 insect species. Hits were filtered using an E-value cutoff of 10^{-5}. The retrieved RDP homologs were aligned using MAFFT (v7.310) with default parameters [47]. To improve alignment quality, we removed ambiguously aligned regions using Gblocks with default settings. The optimal amino acid substitution model (JTT+G) was determined using ModelTest-NG (v0.1.6). Phylogenetic relationships were inferred using maximum likelihood analysis in RAxML (v0.9.0) with 1000 bootstrap replicates to assess branch support [48]. The final tree was visualized and annotated using

418   iTOL (https://itol.embl.de/).

Preparation of anti-BtRDP and anti-NtRLP4 serum

The coding sequence of BtRDP (excluding the signal peptide) was PCR-amplified and cloned

into the pET-28a vector (Novagen) to generate a C-terminal 6×His-tagged fusion protein. The

recombinant plasmid was transformed into *Escherichia coli* Transetta (DE3) competent cells

(TransGen Biotech, Beijing, China). Protein expression was induced with 0.5 mM isopropyl-β-D-

thiogalactoside (IPTG) at 28°C for 8 h. Bacterial cells were harvested by centrifugation and lysed

via ultrasonication for 30 min. The recombinant BtRDP^{-sp}-His protein was purified under native

conditions using Ni-NTA agarose (Qiagen, Hilden, Germany) according to the manufacturer's

protocol.

Purified BtRDP^{-sp}-His protein was used to immunize rabbits (Hua'an Biotechnology,

Hangzhou, China) for custom polyclonal antibody production. The anti-NtRLP4 serum, prepared

by immunizing rabbits with peptide GQNSDRPISTKNST, was produced via the custom service of

Genscript Biotechnology Company (Nanjing, China). The anti-BtRDP serum was subsequently

conjugated to Alexa Fluor™ 488 NHS Ester (#A20000, Thermo Fisher Scientific) following

standard labeling protocols.

The specificity of anti-BtRDP serum was tested by comparing ds*GFP*- and ds*BtRDP*-treated

Bemisia tabaci. The results showed that ds*BtRDP* significantly reduced the target gene. Similarly,

the specificity of anti-NtRLP4 serum was tested by comparing RNAi-EV and RNAi-NtRLP4 plants,

which showed a reduced NtRLP4 expression in *RNAi-NtRLP4* plants.

Immunohistochemistry (IHC) staining

Dissected *B. tabaci* heads with attached salivary glands and Parafilm-embedded salivary

sheaths were fixed in 4% paraformaldehyde (#E672002, Sango Biotechnology, Shanghai) for 30

min. Salivary sheaths adherent to Parafilm were similarly excised and fixed under identical

conditions. After fixation, samples were washed three times with PBST (PBS containing 0.1%

Tween-20) and blocked with 10% FBS for 2 h at room temperature. The tissues were then incubated

overnight at 4°C with Alexa Fluor 488-conjugated anti-BtRDP serum (1:200 dilution), followed by

co-staining with phalloidin-rhodamine (1:500, Thermo Fisher Scientific) for 30 min and 4',6-

diamidino-2-phenylindole (DAPI) solution (#ab104139, Abcam, Cambridge, USA) for nuclear

visualization. Fluorescent images were acquired using a Leica SP8 confocal microscope (Leica

Microsystems) to examine BtRDP localization patterns

Protein extraction and Western-blotting assays

To identify secreted salivary proteins in *N. tabacum* plants, approximately 200 *B. tabaci* adults

were confined in a 3-cm diameter feeding cage and allowed to feed on host plants for 24 h. After

feeding, the eggs deposited on the infested tobacco leaves were removed. The leaves showing no

visible insect contamination were immediately frozen in liquid nitrogen and ground to a fine powder.

The powdered tissue was then homogenized in RIPA Lysis Buffer (#89900, Thermo Fisher

Scientific) for protein extraction. To examine the effects of insect feeding on NtRLP expression,

groups of approximately 200 *B. tabaci* were allowed to feed on 4-5 true-leaf stage tobacco plants

for 12, 24, or 48 h, with uninfested plants serving as controls. For silencing/overexpression

efficiency tests, leaf samples were collected at 48 h post-*Agrobacterium* infiltration, along with

transgenic tobacco plants at the 4-5 true-leaf stage. For insect tissue analysis, various *B. tabaci*

organs were carefully dissected in 1×PBS buffer, including carcasses (20), fat bodies (20), guts (40),

salivary glands (60), and ovaries (10), with sample sizes indicated in parentheses. All dissected

tissues were transferred to lysis buffer for protein extraction. Total protein concentrations in both

insect and plant homogenates were quantified using a BCA Protein Assay Kit (#CW0014S,

CwBiotech, Taizhou, China). Equal protein amounts were separated by SDS-PAGE and electro-

transferred to PVDF membranes.

Membranes were probed with the following primary antibodies: anti-FLAG (1:10,000, #MA1-

91878, Thermo Fisher Scientific), anti-MYC (1:10,000, #MA1-21316, Thermo Fisher Scientific),

anti-GFP (1:10,000, #MA5-15256, Thermo Fisher Scientific), anti-NtRLP4 (1:5,000, GenScript

Biotechnology, Nanjing, China), and anti-BtRDP serum (1:5,000, Huaan Biotechnology, Hangzhou,

China). Following primary antibody incubation, membranes were treated with HRP-conjugated

secondary antibodies: goat anti-mouse IgG (1:10,000, #31430, Thermo Fisher Scientific) or goat

anti-rabbit IgG (1:10,000, #31460, Thermo Fisher Scientific). Protein bands were visualized using

an AI 680 image analyzer (Amersham Pharmacia Biotech, Buckinghamshire, UK), with Ponceau S

staining used to verify equal protein loading.

Quantitative real-time PCR (qRT-PCR) analysis

To prepare tissue-specific samples from *B. tabaci*, adult females were dissected to isolate carcasses (10), fat bodies (10), guts (20), salivary glands (40), and ovaries (10). Additionally, different developmental stages, including eggs (50), nymphs (20), pupae (20), and adult males (20) and females (20), were collected for analysis. The number of insects in each sample is indicated in parentheses. To extract RNA from *N. tabacum* plants, the samples were grinded using liquid nitrogen after collection. Total RNA was extracted using the TRIzol Total RNA Isolation Kit (#9109, Takara, Dalian, China) following the manufacturer's instructions. RNA purity and concentration were assessed using a NanoDrop spectrophotometer (Thermo Fisher Scientific). First-strand cDNA was synthesized from equal amounts of RNA (1 μg) using the HiScript II Q RT SuperMix (#R212-01, Vazyme, Nanjing, China). Gene expression analysis was performed using a Roche LightCycler® 480 system (Roche Diagnostics, Mannheim, Germany) with SYBR Green Supermix (#11202ES08, Yeasen, Shanghai, China). The thermal cycling protocol consisted of initial denaturation at 95°C for 5 min, and amplification for 40 cycles of 95°C for 10 s and 60°C for 30 s. Primers were designed with Primer Premier v6.0 (Table S4). *B. tabaci Tubulin* and *N. tabacum Tubulin* served as internal controls for normalization. Relative gene expression was calculated using the $2^{-\Delta\Delta Ct}$ method [49]. Three to six independent biological replicates, each repeated twice, were performed.

RNA interference

The target gene DNA sequences were amplified using primers listed in Table S4 and subsequently cloned into the pClone007 Vector (#TSV-007, Tsingke Biotechnology, Beijing, China). For dsRNA production, PCR-generated DNA templates containing T7 promoter sequences were transcribed *in vitro* using the T7 High Yield RNA Transcription Kit (#TR101-01, Vazyme Biotech, Nanjing, China). The RNAi experiments were performed according to established methods [50]. Briefly, newly emerged adult whiteflies were anesthetized with carbon dioxide for 5-10 seconds. Approximately 50 nL of dsRNA solution (2 μg/μL) was microinjected into the mesothorax of each insect using a FemtoJet microinjection system (Eppendorf, Hamburg, Germany). Following injection, the whiteflies were maintained on healthy *N. tabacum* leaves under standard rearing conditions. Only actively moving insects were selected for subsequent analyses. Gene silencing

efficiency was evaluated on day 4 post-injection using both qRT-PCR and Western-blotting

approaches.

Generation of transgenic plants

We generated two transgenic *N. tabacum* lines expressing different BtRDP variants: one

containing the full coding sequence (oeBtRDP) and another lacking the signal peptide (oeBtRDP-

sp), which may affect protein secretion. The target sequences, along with the full-length NtRLP4

coding sequence and an NtRLP4 hairpin RNAi construct, were each cloned into the pBWA(V)HS

vector. The oeBtRDP and oeBtRDP-sp were fused with C-terminal flag tags, while no tag was fused

to oeNtRLP4. All constructs were transformed into *A. tumefaciens* strain GV3101 and used to

generate transgenic tobacco plants (K326 background) through *Agrobacterium*-mediated

transformation performed by Wuhan Boyuan Biological Company. Transgenic lines were verified

by qRT-PCR and western blotting, with empty vector-transformed plants serving as controls, and

two independent lines for each construct were selected for further experiments.

Target gene silencing in plants

The hairpin RNAi construct targeting *NtSOBIR1* was generated by cloning the specific

fragment into the pBWA(V)HS plasmid using BsaI and Eco31I restriction enzymes. The

recombinant vector was then transformed into *A. tumefaciens* strain GV3101 and subsequently

infiltrated into *N. tabacum* leaves using the agroinfiltration method. Following a 48-h incubation

period under standard growth conditions, silencing efficiency was quantitatively evaluated by qRT-

PCR analysis.

Transcriptomic sequencing

To investigate the transcriptional effects of NtRLP4, BtRDP, and BtRDP-sp in *N. tabacum*, leaf

samples from EV, oeRLP4#1, oeBtRDP#1, and oeBtRDP-sp#1 transgenic plants were homogenized

in TRIzol Reagent (#10296018, Invitrogen, Carlsbad, CA, USA) for total RNA extraction following

the manufacturer's protocol. Afterwards, total RNA was extracted according to the instructions of

the manufacturer, and the RNA samples were sent to Novogene Institute (Novogene, Beijing, China)

for transcriptomic sequencing. Briefly, poly(A)+ RNA was isolated from 20 μg pooled total RNA

using oligo(dT) magnetic beads, followed by fragmentation in divalent cation buffer at 94°C for 5

min. First-strand cDNA synthesis was performed using N6 random primers, with subsequent

second-strand synthesis to generate double-stranded cDNA. After end-repair and Illumina adapter

ligation, the products were PCR-amplified (15 cycles) and purified using a QIAquick PCR

Purification Kit (#28104, Qiagen, Hilden, Germany). The resulting cDNA libraries were quantified

and quality-checked before paired-end sequencing on an Illumina NovaSeq 6000 platform. All raw

sequencing data were deposited in the National Genomics Data Center under accession number

PRJCA025907.

Analysis of transcriptomic data

The raw sequencing reads were quality-filtered using Illumina's internal software, and the

resulting clean reads from each cDNA library were aligned to the *N. tabacum* reference genome in

Sol   Genomics   Network

(https://solgenomics.net/ftp/genomes/Nicotiana_tabacum/edwards_et_al_2017/)   using   HISAT

v2.1.0 [51]. Low-quality alignments were removed using SAMtools (v1.7) with default parameters

[52]. Transcript abundance was quantified as Transcripts Per Million (TPM) using Cufflinks (v2.2.1)

[53]. Differential expression analysis was performed with DESeq2 (v2.2.1) [54], identifying

significantly DEGs using thresholds of |log2(fold change)| > 1 and adjusted *p*-value < 0.05. PCA of

transcriptome profiles was conducted using a custom R script plotPCA (github.com/franco-

ye/TestRepository/blob/main/PCA_by_deseq2.R). For functional annotation, KEGG pathway

enrichment analysis was performed in TBtools (v2.083) using a one-sided hypergeometric test:

$P = 1 - \sum_{i=0}^{m-1} \left(\frac{\binom{M}{i}\binom{N-M}{n-i}}{\binom{N}{n}} \right)$, [55]. In this software, enriched *P*-values were calculated according to

one-sided hypergeometric test: with *N* representing the number of gene with KEGG annotation, *n*

standing for the number of DEGs in *N*, *M* indicating the number of genes in each KEGG term, and

m suggesting the number of DEGs in each KEGG term.

Insect bioassays

To survivorship analysis, newly emerged *B. tabaci* adults were microinjected with dsRNA and

placed on *N. tabacum* leaves. After 24 h, any dead individuals (potentially due to injection injury)

were removed. For each treatment, 20-30 surviving adults were transferred to leaf cages, and

mortality was recorded daily for 10 days. Three independent biological replications were performed.

For fecundity analysis, dsRNA-treated virgin females were paired with untreated males   in

leaf cages. Five such pairs were maintained per cage (n=10-12 cages per treatment). Egg deposition

was quantified daily for 3 consecutive days under controlled conditions. Each male-female pair was

considered an experimental replicate.

For host preference test, detached *N. tabacum* leaves with moist cotton-wrapped petioles were

arranged in a 50-cm diameter choice arena with a central release chamber. Forty adult whiteflies

were released per chamber, and settling preference was recorded at 3, 6, 12, 24, 36, and 48 h post-

release. The experiment included 10-12 replicates per treatment, with leaf positions randomized

between replicates to eliminate positional bias.

EPG recording and analysis

Feeding behavior of *B. tabaci* was monitored using a GiGA-8d EPG amplifier (Wageningen

Agricultural University; 10 TΩ input resistance, <1 pA input bias current) with 50× gain and ±5 V

output range. Prior to recording, dsRNA-treated whiteflies were starved for 12 h on moist filter

paper, briefly anesthetized with CO_2 (10 s), and connected to the amplifier via a gold wire electrode

(20 μm diameter × 5 cm length) attached to the pronotum using conductive silver glue. Plant

electrodes consisted of copper wires (2 mm diameter × 10 cm length) inserted into the soil of potted

N. tabacum. Recordings were conducted for 8 h in a Faraday cage (120×75×67 cm; Dianjiang,

Shanghai), with only insects surviving the full duration retained for analysis.

Recorded waveforms were processed using PROBE 3.4 software (Wageningen Agricultural

University) and categorized into four distinct feeding phases: nonpenetration (np), pathway duration

(C), phloem salivation (E1), and phloem ingestion (E2) as defined in a previous study [31]. Each

treatment included ≥10 biological replicates, with ds*GFP*-injected and untreated whiteflies

serving as controls.

Agrobacterium-**mediated plant transformation in** *N. tabacum* **and** *N. benthamiana*

The recombinant expression vectors were introduced into *A. tumefaciens* strain GV3101 via

heat shock transformation. Transformed cells were selected on LB agar plates supplemented with

50 μg/mL kanamycin and 10 μg/mL rifampicin, followed by incubation at 28°C for 60 h. Positive

colonies were inoculated into liquid LB medium with the same antibiotics and grown to log phase.

Bacterial cells were then pelleted by centrifugation at 2,400 g for 2 min and resuspended in induction

buffer (10 mM $MgCl_2$, 10 mM MES [pH 5.6], 200 μM acetosyringone) adjusted to specific OD_{600}

values depending on the experiment.

For standard experiments, bacterial suspensions were normalized to $OD_{600} = 1.0$, while

degradation assays used three concentrations ($OD_{600} = 0.05$, 0.3, and 1.0). Equal volumes of selected

bacterial suspensions were mixed and infiltrated into leaves of 4–5-week-old *N. tabacum* and *N.

benthamiana* plants using a needleless syringe. Infiltrated plants were maintained under controlled

conditions for subsequent analysis.

Diaminobenzidine (DAB) staining

To detect H_2O_2 accumulation in *N. tabacum* leaves, leaf samples were immersed in 1 mg/mL

3,3'-diaminobenzidine tetrahydrochloride (DAB-HCl, #D8001, Sigma-Aldrich, St. Louis, MO,

USA) solution (pH 3.8) for 6 hours at room temperature, allowing H_2O_2-dependent brown

polymerization to occur. Following incubation, the DAB solution was replaced with 100% ethanol

for chlorophyll removal, and samples were decolorized overnight at 65°C. The destained leaves

were then photographed under standardized conditions using a Canon EOS 80D digital camera

(Canon Inc., Tokyo, Japan).

Interaction assays between two proteins

In the Y2H screening assay, the coding sequence of BtRDP^{-sp} was cloned into the pGBKT7

bait vector (Clontech, USA), while a *N. benthamiana* cDNA library was constructed in the pGADT7

prey vector (Biogene Biotech, Shanghai, China). These recombinant vectors were co-transformed

into yeast strain Y2HGold and plated on QDO medium (SD/-Ade/-His/-Leu/-Trp, #630428, Takara)

for 3 days at 30°C. Positive colonies were cultured in QDO liquid medium, and plasmids were

extracted using a yeast DNA kit (#DP112-02, TIANGEN, Beijing, China) before transformation

into E. coli DH5α (#TSC-C14, Tsingke) for sequencing (YouKang Biotech, China) to identify

interacting partners.

In the Y2H point-to-point verification assay, BtRDP^{-sp} and NlSP104^{-sp}, and mutant variants of

NtRLP4/SlRLP4/ OsRLP4 were cloned into either pGBKT7 or pGADT7 vectors using primers

from Table S4. Yeast co-transformants were first selected on DDO medium (SD/-Leu/-Trp)

(#630417, Takara) for 3 days at 30°C, then spotted onto QDO medium to confirm interactions

through growth after another 3 days of incubation.

In the Co-IP assays, Full-length sequences of *BtRDP*, *NlSP104*, *NtSOBIR1*, *NtRLP4*, *SlRLP4*,

and *OsRLP4* were cloned into lic-myc or lic-flag vectors for C-terminal tagging, with BtFTSP-flag

serving as a negative control. Five-week-old *N. benthamiana* leaves were agroinfiltrated with these

constructs or collected from oeBtRDP transgenic plants. Total proteins were extracted from 1 g

frozen tissue in IP lysis buffer (#87788, Thermo Scientific) with protease inhibitors (#56079200,

Roche, Switzerland), followed by centrifugation at $1,000 \times g$ for 20 min. Cleared lysates were

incubated with anti-flag beads (#L00432-1, GenScript, Nanjing, China) for 4 h at 4°C, washed with

PBS, and eluted in 2× SDS-PAGE buffer (500 mM Tris-HCl, pH=6.8, 50% glycerin, 10% SDS, 1%

bromophenol blue, and 2% β-mercaptoethanol) for Western-blotting assay.

In the BiFC assay, BtRDP^{-sp}, BtFTSP^{-sp}, NtRLP4, and NtCf9-like were cloned into the pCV-

cYFP or pCV-nYFP vectors, respectively. After transforming *A. tumefaciens* GV3101, pairwise

combinations were co-infiltrated into *N. benthamiana* leaves. Following 36-48 h incubation in a

growth chamber, reconstituted YFP fluorescence was visualized using a Leica SP8 confocal

microscope with standard filter sets for YFP detection.

RLP degradation by salivary effectors among different herbivorous insects

To examine salivary protein effects on RLPs, we co-infiltrated *A. tumefaciens* GV3101 strains

carrying various constructs into *N. benthamiana* leaves. The recombinant vectors expressed C-

terminally tagged proteins: BtRDP-flag, NlSP104-flag, GUS-flag, NlSP7-flag, and myc-tagged

RLPs (NtRLP4, SlRLP4, OsRLP4, SlCf9, NtCf9). Bacterial suspensions were mixed at

predetermined OD_{600} ratios before infiltration, with leaf samples collected 48 hours post-infiltration

for subsequent analysis.

For confocal microscopy studies, *N. benthamiana* leaves were co-infiltrated with *A.*

tumefaciens carrying: (1) NtRLP4-GFP plus BtRDP-mCherry, (2) NtRLP4-GFP plus RFP-mCherry,

or (3) NtRLP4-GFP plus BtFTSP-mCherry. The samples were imaged by confocal microscopy at

48 h post injection. The same parameters were used in photographing.

This study also conducted a treatment on *N. benthamiana* plants that were overexpressing

NtRLP4-myc using purified proteins. Briefly, the BtRDP^{-sp} and GFP coding sequences were cloned

into pET-28a for 6×His-tagged expression in *E. coli* Transetta (TransGen Biotech). Protein

expression was induced with 0.5 mM IPTG at 25°C for 8 h, followed by purification using Ni-NTA

agarose and buffer exchange into PBS using a 3-kDa molecular-weight cutoff Amicon Ultra-4

Centrifugal Filter Device (Millipore, MA, USA). Different concentrations of purified proteins were

infiltrated into leaves of *N. benthamiana* plants stably expressing NtRLP4-myc, with samples

collected 24 h post-treatment.

To investigate degradation pathways, we infiltrated *N. benthamiana* leaves with: 50 μM

MG132 (#M7449, Sigma-Aldrich, Steinheim, Germany), 5 μM 3-MA (#M9281, Sigma-Aldrich,),

or 10 nM BAF (#BML-CM110-0100, Enzo Life Sciences, Farmingdale, USA) at 24 h post-

agroinfiltration. The samples were incubated for additional24 h. Protein levels were detected using

Western-blotting assay.

***In planta* ubiquitination assay**

Ubiquitination detection in *N. benthamiana* was performed according to the methods

previously reported [56]. The recombinant vector for transient expression of the HA-fused ubiquitin

protein was kindly provided by Dr. Chao Zheng (Ningbo University). Myc-tagged proteins were co-

expressed with HA-UBQ by agroinfiltration, and 50 μM MG132 was injected into the *N.*

benthamiana leaves one day post *A. tumefaciens* infiltration. Immunoprecipitation was performed

using anti-myc beads (ytma-20, ChromoTek , Planegg-Martinsried, Germany). Ubiquitination

signals were detected by immunoblot with anti-HA antibody (HT301, TransGen Biotech, Beijing,

China).

Scanning electron microscopy (SEM)

B. tabaci adults were allowed to feed on artificial diet solutions for 24 h. Parafilm sections

containing salivary sheaths were carefully excised and rinsed with 1× PBS (pH 7.4) to remove

residual diet. The samples were mounted on aluminum stubs, desiccated under vacuum, and sputter-

coated with gold before imaging with a TM4000 II Plus SEM (Hitachi, Tokyo, Japan). Salivary

sheath length (from apex to base) was measured, with 20 sheaths analyzed per treatment to ensure

statistical reliability.

Statistical analysis

The log-rank test (SPSS Statistics 19, Chicago, IL, USA) was used to determine the statistical

significance of survival distributions. Two-tailed unpaired Student's t-test (comparisons between

two groups) or one-way ANOVA test followed by Tukey's multiple comparisons test (comparisons

among three groups) was used to analyze the results of qRT-PCR, fecundity analysis, and host choice

analysis. The EPG data were first checked for the normality and homogeneity of variance, and those

not fitting a normal distribution were $\log_{10}$ transformed before Student's t test, as described in

previous studies [31]. Statistical tests, including Student's t-test and one-way ANOVA, and data

visualization were conducted in GraphPad Prism 9.

Data availability

All data is included in the manuscript and/or supporting information. The transcriptomic data have

been submitted to the National Genomics Data Center under accession number PRJCA025907. The

original blot images have been displayed in Fig. S29-S32.

References

1.   Howe GA & Jander G (2008). Plant immunity to insect herbivores. *Annu Rev Plant Biol* 59, 41-
66.

2.   Wang H, *et al*. (2024), Release brake and set plant defense in motion. *New Plant Prot* 1: e11.

3.   Nürnberger T, Brunner F, Kemmerling B, & Piater L (2004) Innate immunity in plants and
animals: striking similarities and obvious differences. *Immunol Rev* 198(1):249-266.

4.   Macho AP & Zipfel C (2014) Plant PRRs and the activation of innate immune signaling. *Mol*
Cell 54(2):263-272.

5.   Zipfel C (2014) Plant pattern-recognition receptors. *Trends Immunol* 35(7):345-351.

6.   Gust AA & Felix G (2014) Receptor like proteins associate with SOBIR1-type of adaptors to
form bimolecular receptor kinases. *Curr Opin Plant Biol* 21:104-111.

7.   Adam D. Steinbrennera, *et al*. (2020) A receptor-like protein mediates plant immune responses
to herbivore-associated molecular patterns. *Proc Natl Acad Sci USA* 117(49):31510-31518.

8.   Chinchilla D, Bauer Z, Regenass M, Boller T, & Felix G (2005) The *Arabidopsis* receptor kinase
718   FLS2 binds flg22 and determines the specificity of flagellin perception. *Plant Cell* 18(2):465-
719   476.

720   9.   Zipfel C, *et al.* (2006) Perception of the bacterial PAMP EF-Tu by the receptor EFR restricts
721   *Agrobacterium*-mediated transformation. *Cell* 125(4):749-760.

722   10.   Ron M & Avni A (2004) The receptor for the fungal elicitor ethylene-inducing xylanase is a
723   member of a resistance-like gene family in tomato. *Plant Cell* 16(6):1604-1615.

724   11.   Tian T, *et al.* (2021) A salivary calcium-binding protein from *Laodelphax striatellus* acts as an
725   effector that suppresses defense in rice. *Pest Manag Sci* 77(5): 2272-2281.

726   12.   Huang HJ, *et al.* (2019) Salivary DNase II from *Laodelphax striatellus* acts as an effector that
727   suppresses plant defence. *New Phytol* 224(2):860-874.

728   13.   Chen CY, *et al.* (2019) An effector from cotton bollworm oral secretion impairs host plant
729   defense signaling. *Proc Natl Acad Sci USA* 116(28):14331-14338.

730   14.   Xu HX, *et al.* (2019) A salivary effector enables whitefly to feed on host plants by eliciting
731   salicylic acid-signaling pathway. *Proc Natl Acad Sci USA* 116(2):490-495.

732   15.   Chen CY & Mao YB (2020) Research advances in plant-insect molecular interaction. *F1000Res*
733   9(F1000 Faculty Rev):198.

734   16.   Guiguet A, *et al.* (2016) Shared weapons of blood- and plant-feeding insects: Surprising
735   commonalities for manipulating hosts. *J Insect Physiol* 84:4-21.

736   17.   Wu J & Baldwin IT (2009) Herbivory-induced signalling in plants: perception and action. *Plant*
737   *Cell Environ* 32(9):1161-1174.

738   18.   Jiang YJ, Zhang CX, Chen RZ, & He SY (2019) Challenging battles of plants with phloem-
739   feeding insects and prokaryotic pathogens. *Proc Natl Acad Sci USA* 116(47):23390-23397.

740   19.   Gao H，*et al*. (2023) The salivary chaperone protein NlDNAJB9 of *Nilaparvata lugens* activated
741   plant immune responses. *J Exp Bot* 74(21):6874-6888.

742   20.   Guo H, *et al*. (2020) An aphid-secreted salivary protease activates plant defense in phloem. *Curr*
743   *Biol* 30(24): 4826-4836.

744   21.   Bos JI, *et al*. (2010) A functional genomics approach identifies candidate effectors from the
745   aphid species *Myzus persicae* (green peach aphid). *PLoS Genet* 6(11):e1001216.

22.   Zhang Y & Zeng L (2020) Crosstalk between ubiquitination and other post-translational protein

modifications in plant immunity. *Plant Commun* 1(4):100041.

23.   Huang J*, et al.* (2020) *Phytophthora* effectors modulate genome-wide alternative splicing of

host mRNAs to reprogram plant immunity. *Mol Plant* 13(10):1470-1484.

24.   Shan L*, et al.* (2008) Bacterial effectors target the common signaling partner BAK1 to disrupt

multiple MAMP receptor-signaling complexes and impede plant immunity. *Cell Host Microbe*

4(1):17-27.

25.   Huang HJ*, et al.* (2016) Screening and functional analyses of *Nilaparvata lugens* salivary

proteome. *J Proteome Res* 15(6):1883-1896.

26.   Huang HJ*, et al.* (2021) Identification of salivary proteins in the whitefly *Bemisia tabaci* by

transcriptomic and LC-MS/MS analyses. *Insect Sci* 28(5):1369-1381.

27.   Gao H*, et al.* (2022) *Nilaparvata lugens* salivary protein NlG14 triggered defense response in

plants. *J Exp Bot* 73(22):7477-7487.

28.   Mukhtar M*, et al.* (2011) Independently evolved virulence effectors converge onto hubs in a

plant immune system network. *Science* 333(6042):596-601.

29.   Wang Y*, et al.* (2023) Horizontally transferred salivary protein promotes insect feeding by

suppressing ferredoxin-mediated plant defenses. *Mol Biol Evol* 40(10):msad221.

30.   Huang HJ*, et al.* (2023) Planthopper salivary sheath protein LsSP1 contributes to manipulation

of rice plant defenses. *Nat Commun* 14(1):737.

31.   Lu S*, et al.* (2021) EPG-recorded feeding behaviors reveal adaptability and competitiveness in

two species of *Bemisia tabaci* (Hemiptera: Aleyrodidae). *J Insect Behav* 34: 26-40.

32.   Yan ZW*, et al.* (2023) Endocytosis-mediated entry of a caterpillar effector into plants is

countered by Jasmonate. *Nat Commun* 14(1):6551.

33.   Jones DA*, et al.* (1994) Isolation of the tomato Cf-9 gene for resistance to *Cladosporium fulvum*

by transposon tagging. *Science* 266 (5186): 789-793 (1994).

34.   Varshavsky A*, et al.* (2017) The ubiquitin system, autophagy, and regulated protein degradation.

Annu Rev Biochem 86: 123-128.

35.   Li L*, et al.* (2022) Independently evolved viral effectors convergently suppress DELLA protein

SLR1-mediated broad-spectrum antiviral immunity in rice. *Nature Commun* 13(1):6920.

36.   Takatsuka C*, et al.* (2004) 3-Methyladenine inhibits autophagy in tobacco culture cells under

sucrose starvation conditions. *Plant Cell Physiol* 45: 265-274.

37. Tariqjaveed M*, et al.* (2021) Versatile effectors of phytopathogenic fungi target host immunity.

J Integr Plant Biol 63(11):1856-1873.

38. Doblas VG, *et al.* (2018) Cell wall integrity signaling in plants: Malectin-domain kinases and

lessons from other kingdoms. *Cell Surf* 3:1-11.

39. Albert I, *et al.* (2015). An RLP23–SOBIR1–BAK1 complex mediates NLP-triggered immunity.

Nat Plants 1: 15140.

40. Li B, Lu D, & Shan L (2014) Ubiquitination of pattern recognition receptors in plant innate

immunity. *Mol Plant Pathol* 15(7):737-746.

41. Chen Y, Song Y, Liu J, Xu G, & Dou D (2022) Ubiquitination of receptorsomes, frontline of

plant immunity. *Int J Mol Sci* 23(6):2937.

42. Langin G, González-Fuente M, & Üstün S (2023) The plant ubiquitin–proteasome system as a

target for microbial manipulation. *Annu Rev Phytopathol* 61:351-375.

43. Rosas-Diaz T*, et al.* (2018) A virus-targeted plant receptor-like kinase promotes cell-to-cell

spread of RNAi. *Proc Natl Acad Sci USA* 115(6):1388-1393.

44. Irieda H*, et al.* (2019) Conserved fungal effector suppresses PAMP-triggered immunity by

targeting plant immune kinases. *Proc Natl Acad Sci USA* 116(2):496-505.

45. Qin J*, et al.* (2018) The Xanthomonas effector XopK harbours E3 ubiquitin-ligase activity that

is required for virulence. *New Phytol* 220(1):219-231.

46. Xu FQ, Xue HW (2019) The ubiquitin-proteasome system in plant responses to environments.

Plant Cell Environ 42(10):2931-2944.

47. Katoh K & Standley DM (2013) MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. *Mol Biol Evol* 30(4):772-780.

48. Kozlov AM, Darriba D, Flouri T, Morel B, & Stamatakis A (2019) RAxML-NG: a fast, scalable

and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics*

35(21):4453-4455.

49. Livak, K.J., and Schmittgen, T.D. (2001). Analysis of relative gene expression data using real-

time quantitative PCR and the $2^{-\Delta\Delta CT}$ method. *Methods* 25, 402-408.

50. Xu HJ*, et al.* (2015) Two insulin receptors determine alternative wing morphs in planthoppers.

Nature 519(7544):464-467.

51. Kim, D., Langmead, B., and Salzberg, S.L. (2015). HISAT: A fast spliced aligner with low

memory requirements. *Nature Methods* 519(7544):464-467.

52.   Li, H. (2009). The Sequence Alignment/Map format and SAMtools. *BIOINFORMATICS - OXFORD*.

53.   Trapnell, C., Roberts, A., Goff, L., Pertea, G., Kim, D., Kelley, D.R., Pimentel, H., Salzberg, S.L., Rinn, J.L., and Pachter, L. (2012). Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. *Nature Protocols* 7, 562-578.

54.   Wang, L., Feng, Z., Wang, X., Wang, X., and Zhang, X. (2010). DEGseq: an R package for identifying differentially expressed genes from RNA-seq data. *Bioinformatics* 26, 136-138.

55.   Chen, C., Chen, H., Zhang, Y., Thomas, H.R., Frank, M.H., He, Y., and Xia, R. (2020). TBtools: an integrative toolkit developed for interactive analyses of big biological data. *Mol Plant* 13, 1194-1202.

56.   Wang, N., Yin, Z., Zhao, Y., Wang, J., Pei, Y., Ji, P., Daly, P., Li, Z., Dou, D., and Wei, L. (2022). An F-box protein attenuates fungal xylanase-triggered immunity by destabilizing LRR-RLP NbEIX2 in a SOBIR1-dependent manner. *New Phytol* 236(6):2202-2215.

Acknowledgements

This project has received funding from the National Key Research and Development Plan in the 14th five-year plan (2021YFD1401100: H.J.H, C.X.Z.), National Natural Science Foundation of China (32422075: H.J.H.; U23A6006: J.P.C and C.X.Z.), and the Natural Science Foundation of Zhejiang Province (LDQ24C140001: H.J.H).

Author Contributions Statement

H-JH, J-ML, C-XZ, and J-PC planned and designed the research. XW, J-BL, Y-ZW, and H-JH performed experiments and analyzed data. X-HZ provided the EV plants. H-JH and XW drafted the manuscript.

Competing Interests Statement

The authors declare no competing interests.

Figure Legends

 Fig. 1. BtRDP is a salivary protein and secreted into plants. (**a**) Detection of BtRDP in *Nicotiana*

tabacum plants. The untreated and *Bemisia tabaci*-infested tobacco plants, as well as the salivary

gland samples, are collected for western-blotting assays. Asterisk indicates non-specific binding.

Rubisco staining (RbcL) is conducted to visualize the amount of sample loading. (**b**, **c**) Detection

of BtRDP in different tissues by western-blotting and qRT-PCR assays. Ov, ovary; SG, salivary

gland; Ca, carcass; FB, fat body. Silver staining and anti-actin serum are used to visualize the sample

loading. (**d**) Expression patterns of BtRDP in different development stages. Ny, nymph; Ps,

pseudopupa; Fe, female; Ma, male. *B. tabaci tubulin* is used as an internal control. The relative

quantitative method ($2^{-\Delta\Delta Ct}$) is used to evaluate the quantitative variation. Data are presented as mean

values ± SEM (n= 3 independent biological replicates). Different lowercase letters indicate

statistically significant differences at the *P* < 0.05 level according to one-way ANOVA test followed

by Tukey's multiple comparisons test. (**e, f**) Immunohistochemical staining of BtRDP in salivary

glands (**e**) and salivary sheath secreted from *B. tabaci* (**f**). The salivary gland and its nearby tissues

are dissected and incubate with anti-BtRDP serum or pre-immune serum conjugated with Alexa

Fluor™ 488 NHS Ester (green) and actin dye phalloidinrhodamine (red). The nucleus is stained

with DAPI (blue). PSG, principal salivary gland; ASG, accessory salivary gland. The lower images

represent the enlarged images of the boxed area in the upper image. Experiments are repeated thrice

for (**a** and **b**), while twice for (**e** and **f**). Similar results are observed and representative images are

displayed.

Fig. 2. Effects of BtRDP on *Bemisia tabaci*. (**a**) Treating *B. tabaci* with ds*BtRDP* significantly

reduces the transcript and protein level of target gene. The ds*GFP*-treated *B. tabaci* is used as a

control. (**b–c**) Effects of *BtRDP* knockdown on insect reproduction (**b**) and feeding behavior (**c**).

Electrical penetration graph (EPG) is used to monitor the insect feeding behavior, which can be

classified into nonpenetration (np), pathway duration (C), phloem salivation (E1) and phloem

ingestion (E2) phases. All EPG recordings are performed for 8 h. Typical EPG waveforms are

displayed in Fig. S7. (**d-f**) Insect performance on transgenic plants overexpressing BtRDP

(oeBtRDP). (**d**) Detection of BtRDP in transgenic *Nicotiana tabacum* overexpressing a complete

coding region of *BtRDP*. The empty vector (EV) plants are used as the control. The flag tag is fused

to the C-terminal ends of recombinant proteins. **(e)** Comparison of insect reproduction on

oeBtRDP#1 and EV plants. Five *B. tabaci* individuals are confined to indicated plants for 3 days,

and the oviposited eggs are counted. **(f)** Attraction of oeBtRDP#1 and EV leaves to *B. tabaci* in a

two-choice equipment. A group of 40 female *B. tabaci* are released into a device containing

oeBtRDP#1 and EV leaves. The number of insects settling on each leaf is counted. After 48 h, the

number of eggs on each leaf is counted. **(g-i)** Insect performance on transgenic plants overexpressing

BtRDP^{-sp} (oeBtRDP^{-sp}). **(g)** Detection of BtRDP^{-sp} in transgenic *N. tabacum* overexpressing BtRDP

without a signal peptide. The insect reproduction **(h)** and settlement **(i)** on oeBtRDP^{-sp} transgenic

plants are recorded. The EPG data are first checked for normality and homogeneity of variance, and

data not fitting a normal distribution are subjected to log10 transformation. Data are presented as

mean ± SEM. Statistical significance in panels **(a)**, **(b)**, **(c)**, **(e)**, **(f)**, **(h)**, and **(i)** was assessed using

two-tailed unpaired Student's *t*-tests. ***P < 0.001; **P < 0.01; *P < 0.05; ns, not significant.

Western-blotting assays are repeated thrice with the similar results.

Fig. 3. NtRLP4 interacts with BtRDP and confers plant resistance to *Bemisia tabaci*. (a)

Domain organization of NtRLP4. NtRLP4 contains a predicted N-terminal signal peptide (SP), a

malectin-like domain, a LRR domain, and a transmembrane domain (TM). **(b)** The protein level of

NtRLP4 in response to *B. tabaci* infestation. The untreated *Nicotiana tabacum* is used as a negative

control. Four independent biological replicates are performed, and the representative images are

displayed. The band density is measured using ImageJ. The density values from 4 biological

replicates are calculated and the mean value in the controls is set at 1.0. **(c)** Yeast two hybrid assays

showing the interaction between BtRDP and NtRLP4. The different combinations of constructs are

transformed into yeast cells, and are grown on the selective medium SD/-Trp/-Leu (DDO), and the

interactions are tested with SD/-Trp/-Leu/-His/-Ade (QDO). **(d)** Co-immunoprecipitation assay

showing the interaction between BtRDP and NtRLP4. Total proteins are extracted from *N.*

benthamiana leaves transiently co-expressing NtRLP4-myc/NtCf9-myc with BtFTSP-flag/BtRDP-

flag. All genes are expressed with a complete coding region, and the myc/flag tags are fused at the

C-terminus. Precipitation is performed using flag beads. The samples are probed with anti-flag and

anti-myc antibodies for immunoblotting analysis. **(e–h)** Analysis of transgenic *N. tabacum*

overexpressing NtRLP4 (oeRLP). **(e)** Detection of NtRLP4 level in oeRLP plants. The empty vector

(EV) plant is used as a control. Two independent oeRLP lines are selected. The samples are probed

with an anti-NtRLP4 antibody. Rubisco staining (RbcL) is used to visualize the amount of sample

loading. **(f)** Attraction of oeRLP#1 and EV leaves to *B. tabaci* in a two-choice equipment. A group

of 40 female *B. tabaci* are released into a device containing oeRLP#1 and EV leaves. The number

of insects settling on each leaf is counted at each time point. After 48 h, the number of eggs on each

leaf is counted. **(g)** Comparison of insect reproduction on oeRLP#1 and EV plants. Five *B. tabaci*

individuals are confined to indicated plants for 3 days, and the oviposited eggs are counted. **(h)**

Relative transcript levels of salicylic acid (SA)- and jasmonic acid (JA)-associated genes in

oeRLP#1 and EV plants. PAL, phenylalanine ammonia lyase; NPR1, nonexpressor of pathogenesis-

related protein 1; FAD7, fatty acid desaturase 7; PDF1.2, plant defensin 1.2. Two independent

biological replicates are performed in **(d)** and **(e)**. Data in **(f)**, **(g)**, and **(h)** are presented as mean

values ± SEM. For insect bioassays in **(f)** and **(g)**, n= 10 independent biological replicates. For qRT-

PCR in **(h)**, n= 3 independent biological replicates. *P*-values are determined by two-tailed unpaired

Student's *t* test. ***P < 0.001; **P < 0.01; *P < 0.05; ns, not significant.

Fig. 4. BtRDP suppress plant defenses by promoting NtRLP4 degradation. (a) Effects of ds*BtRDP* suppression on insect reproduction (n= 10 independent biological replicates) when feeding on empty vector (EV) and NtRLP4-silenced (*RNAi-RLP*) transgenic *Nicotiana tabacum*. **(b)** Transient overexpressing BtRDP-mCherry attenuates H_2O_2 accumulation caused by NtRLP4-GFP overexpression. *N. tabacum* co-expressing RFP-mCherry and NtRLP4-GFP is used as a control. All genes are expressed with a complete coding region. The experiment is repeated 5 times with the similar results. **(c)** Relative transcript level of salicylic acid (SA)- and jasmonic acid (JA)-associated genes in BtRDP-mCherry/NtRLP4-GFP and RFP-mCherry/NtRLP4-GFP plants (n=3 independent biological replicates). PAL, phenylalanine ammonia lyase; NPR1, nonexpressor of pathogenesis-related protein 1; FAD7, fatty acid desaturase 7; PDF1.2, plant defensin 1.2. **(d)** Comparison of insect reproduction (n= 16 independent biological replicates) on transient-expressed BtRDP-mCherry/NtRLP4-GFP and RFP-mCherry/NtRLP4-GFP plants. **(e)** Attraction of BtRDP-mCherry/NtRLP4-GFP and RFP-mCherry/NtRLP4-GFP leaves to *B. tabaci* in a two-choice equipment (n= 16 independent biological replicates). The number of eggs on each leaf is counted at

48 h post insect release. (**f**) Effects of BtRDP and BtRDP^{-sp} overexpression on NtRLP4

accumulation. Transgenic oeBtRDP (left) and oeBtRDP^{-sp} (right) plants are probed with anti-

NtRLP4 and anti-flag antibodies for immunoblotting analysis. The experiments are repeated twice

with the similar results. (**g**) Degradation of NtRLP4 by BtRDP in *N. benthamiana* leaves. NtRLP4-

myc and BtRDP-flag are transiently co-expressed in *N. benthamiana* plants through *Agrobacterium*

infiltration. *Agrobacterium* carrying NtCf9-myc and GUS-flag are used as negative controls. (**h**)

Effects of 26S proteasome inhibitor (MG132) on NtRLP4 accumulation. Co-infiltrated leaves are

treated with MG132 at 24 h post injection. The samples are probed with anti-flag and anti-myc

antibodies for immunoblot analysis. Rubisco staining (RbcL) is conducted to visualize the amount

of sample loading. The small triangle indicates the different concentrations of *Agrobacterium*

(OD_{600} = 0.05, 0.3, and 1.0). (**i**) NtRLP4 is ubiquitinated *in planta*. NtRLP4-myc is co-expressed

transiently with HA-UBQ in *N. benthamiana* leaves. Extracted total proteins are

immunoprecipitated by anti-myc beads and immunoblotted with anti-myc or anti-HA antibody.

Experiments are repeated thrice for (**g**) and (**h**), while twice for (**i**). Band density is measured using

ImageJ. The density values from three biological replicates are calculated and the mean value in the

first lane is set at 1.0. Data are presented as mean values ± SEM. *P*-values are determined by two-

tailed unpaired Student's *t* test. ***P < 0.001; **P < 0.01; *P < 0.05; ns, not significant.

Fig. 5. Rice RLP4 are targeted by salivary protein NlSP104 from *Nilaparvata lugens*. (a)
Domain organization of *Oryza sativa* RLP4 (OsRLP4). **(b, c)** Yeast two-hybrid and Co-IP assays
showing the interaction between OsRLP4 and NlSP104. All genes are expressed with a complete
coding region in Co-IP assays, while NlSP104 without a signal peptide is used in yeast two-hybrid
assays. Experiments in **(b)** are repeated twice with the similar result. **(d)** The expression patterns of
NlSP104 in different *N. lugens* tissues. Te, testis; Ov, ovary; SG, salivary gland; Ca, carcass; FB, fat
body. Data are presented as mean values ± SEM (n= 3 independent biological replicates). Different
lowercase letters indicate statistically significant differences at *P* < 0.05 level according to one-way
ANOVA test followed by Tukey's multiple comparisons test. **(e-g)** Effects of dsRNA treatment on
insect survivorship **(e)**, reproduction **(f)**, and honeydew excretion **(g)**. For survivorship analysis, a
group of 30 *N. lugens* are reared in a cage. Three independent biological replications are performed.
Differences in survivorship between the two treatments are tested by log-rank test. ns, not significant.

For reproduction analysis, n= 10 and 13 individuals are tested in ds*GFP*- and ds*NlSP104*-treatment,

respectively. The sterile females are excluded from data analysis. For honeydew analysis, n= 10

independent biological replicates. *P*-values in **(f)** and **(g)** are determined by two-tailed unpaired

Student's *t* test. ***$P < 0.001$; **$P < 0.01$. **(h)** Effect of NlSP104 on the accumulation of transient-

expressed OsRLP4. OsRLP4-myc is agro-injected into *Nicotiana benthamiana* together with

different concentrations of NlSP104-flag and NlSP7-flag. NtCf9-myc is used as a negative control.

The small triangle indicates the different concentrations (OD_{600} = 0.05, 0.3, and 1.0) of

Agrobacterium. Rubisco staining (RbcL) is conducted to visualize the amount of sample loading.

Experiments are repeated three times with the similar results. Band density is measured using

ImageJ. The density values from three biological replicates are calculated with the mean value in

the first lane being set at 1.0.

Fig. 6. The proposed model for the suppression of receptor-like proteins (RLPs)-mediated

plant defenses by salivary effectors. Host plants employ pattern-recognition receptors (PRRs) to

detect various damage-associated molecular patterns (DAMPs) and herbivore-associated molecular

patterns (HAMPs) triggered by insect feeding. The RLP4/SOBIR1 complex plays a vital role in

initiating pattern-triggered immunity (PTI), including H_2O_2 burst, upregulation of jasmonic acid

(JA), and downregulation of salicylic acid (SA), which hinders insect feeding. The whitefly *Bemisia*

tabaci and planthopper *Nilaparvata lugens* independently evolved salivary proteins that targeted

977 plant RLP4. *B. tabaci* salivary sheath protein BtRDP interacts with the leucine-rich repeat (LRR)

978 domain of RLP4 from *Nicotiana tabaci* and *Solanum lycopersicum*, while *N. lugens* NlSP104 targets

979 both the LRR domain and the malectin-like domain of *Oryza sativa* RLP4. These interactions

980 promote the ubiquitin-dependent degradation of RLP4, thereby disrupting the stability of the

981 RLP4/SOBIR1 complex. The presence of salivary effectors causes a hormonal shift and suppresses

982 the H_2O_2 burst, finally favoring insect feeding.

1. Supplementary Figures

a

cleavage site

```
  1: MHKLL AFLVL AGVAL AVPTA TRRCA NYNHY NDWKG NYLVD KEWVV AASTA
 51: SGSVF GWRND YEFCD AYFTY GVPQC GSCTF IRFIP NSRYS GDFGG YTMVQ
101: YAWNS NRQDY DYTEN FIGRA YNGYN APVYR CYTRE PFDRT YFGYK DVMDP
151: DFVPY DAWSN DLTYS AGVLS YCPKN NIVIL VICNN FQYMN YVDNY PIVVV
201: LAQDY KLNAD QKNYI VSELE KNNIN PYGVI LRDNY NCKAR TLYNP SIYL
```

Fig. S1. Characteristic of BtRDP. (a) Deduced amino acid sequence of BtRDP. Arrow indicates the signal peptide cleavage site. **(b-g)** Mass spectrums of the identified unique peptides in watery saliva of *Bemisia tabaci*.

Fig. S2. Analysis of insect RDP. (**a**) Phylogenetic tree of RDP sequences from Aleyrodidae species. The trees are constructed by RAxML v0.9.0 using the maximum likelihood method with 1000 bootstrap replicates. Nodes with bootstrap values greater than 50 are displayed. (**b**) Sequence alignments of RDPs. The amino acid sequence of RDPs used for phylogenetic tree construction are aligned using ClusterX software. Black shades indicate the conserved regions.

Fig. S3. Expression patterns of *Bemisia tabaci BtRDP*. Transcripts per million (TPM) expression values of *BtRDP* in different tissues (**a**) and at different developmental stages (**b**) are determined based on the transcriptomic data. Data are presented as mean values ± SEM (n=3 independent biological replicates).

Fig. S4. Expression patterns of *BtActin* and *Bt18s rRNA*. The transcript level of each gene in different tissues (**a**) and at different developmental stages (**b**) is quantified by qRT-PCR. *Bemisia tabaci tubulin* is used as an internal control. The relative quantitative method ($2^{-\Delta\Delta Ct}$) is used to evaluate the quantitative variation. Ov, ovary; SG, salivary gland; Ca, carcass; FB, fat body; Ny, nymph; Ps, pseudopupa; Fe, female; Ma, male. Data are presented as mean values ± SEM (n=3 independent biological replicates). Different lowercase letters indicate statistically significant differences at *P* < 0.05 level according to one-way ANOVA test followed by Tukey's multiple comparisons test.

Fig. S5. BtRDP is efficiently and specifically suppressed by ds*BtRDP*. (**a**) Relative transcript level of *BtRDP*, *BtFTSP*, *BtE3*, and *Bt56* after dsRNA treatments. *Bemisia tabaci* are treated with ds*GFP* and ds*BtRDP*. The expression pattern of each gene is determined 4 days post treatment using qRT-PCR methods. *B. tabaci tubulin* is used as an internal control. The relative quantitative method ($2^{-\Delta\Delta Ct}$) is used to evaluate the quantitative variation. Data are presented as mean values ± SEM (n= 3 independent biological replicates) (**b**) Protein level of BtRDP after dsRNA treatments. Actin is used as an internal control. Two independent biological replicates are displayed. (**c**) Immunohistochemical staining of BtRDP after dsRNA treatment. The salivary gland (SG) and its nearby tissues are collected from ds*GFP*- and ds*BtRDP*-treated *Bemisia tabaci*. The samples are incubated with anti-BtRDP serum conjugated with Alexa Fluor™ 488 NHS Ester (green) and actin dye phalloidinrhodamine (red), and examined by Leica SP8. The nucleus is stained with DAPI (blue). The lower image in each treatment represents the enlarged images of the boxed area in the upper image. Experiments are repeated twice, with each inspecting more than 10 salivary glands.

Fig. S6. Effects of dsRNA treatment on insect survivorship and salivary sheath formation. (**a**, **b**) Effects of dsRNA treatment on salivary sheath formation. Newly emerged *Bemisia tabaci* adults are injected with ds*GFP* and ds*BtRDP*. The dsRNA-treated *B. tabaci* are fed on artificial diets. The salivary sheaths left on parafilm are inspected by scanning electron microscopy (**a**), and the length of salivary sheath is measured from the top to base of salivary sheath (**b**). Data are presented as mean values ± SEM. Twenty salivary sheaths from each treatment are measured. *P*-value is determined by two-tailed unpaired Student's *t* test. *$P < 0.05$. (**c**) Effects of dsRNA treatment on insect survivorship. A group of 20-30 *B. tabaci* are placed in a leaf cage, and their mortality is recorded for ten consecutive days. Three independent biological replications are performed. Differences in survivorship between the two treatments are tested by log-rank test. ns, not significant.

Fig. S7. Typical EPG waveforms for *Bemisia tabaci* feeding on *Nicotiana tabacum*.
The *B. tabaci* feeding behavior can be classified into nonpenetration (np), pathway
duration (C), phloem salivation (E1) and phloem ingestion (E2).

Fig. S8. Effects of empty vector (EV) transgenic plants on *Bemisia tabaci*. Two-choice experiment is performed to investigate the attraction of wild type (WT) and EV transgenic plants to *B. tabaci*. A group of 40 female *B. tabaci* are released into a device containing EV and WT leaves. The number of insects settling on each leaf is counted at 3, 6, 12, 24, 36, and 48 h (**a**). After 48 h, the number of eggs on each leaf is counted (**b**). Data are presented as mean values ± SEM. Ten independent biological replicates are performed. *P*-values are determined by two-tailed unpaired Student's *t* test. ns, not significant.

Transgenic plants overexpressing BtRDP

Transgenic plants overexpressing BtRDP⁻ˢᵖ

Fig. S9. Effects of BtRDP and BtRDP⁻ˢᵖ overexpression on *Bemisia tabaci*. Insect performance on oeBtRDP#2 (**a, b**) and oeBtRDP⁻ˢᵖ#2 (**c, d**) transgenic plants are tested. (**a, c**) Comparison of insect reproduction on empty vector (EV) and oeBtRDP#2/oeBtRDP⁻ˢᵖ#2 transgenic plants. Five *B. tabaci* individuals are confined to indicated plants for 3 days, and the oviposited eggs are counted. Twelve independent biological replicates are performed. (**b, d**) Attraction of EV and oeBtRDP#2/oeBtRDP⁻ˢᵖ#2 leaves to *B. tabaci* in a two-choice equipment. A group of 40 female *B. tabaci* are released into a device containing indicated leaves. The number of insects settling on each leaf is counted at 3, 6, 12, 24, 36, and 48 h. After 48 h, the number of eggs on each leaf is counted. Data are presented as mean values ± SEM. Ten independent biological replicates are performed. *P*-values are determined by two-tailed unpaired Student's *t* test. ****P* < 0.001.

Fig. S10. Expression patterns of *NtRLP4* in response to *Bemisia tabaci* infestation. Relative transcript level of *NtRLP4* in response to *B. tabaci* infestation are determined by qRT-PCR. Data are presented as mean values ± SEM. Different lowercase letters indicate statistically significant differences at *P* < 0.05 level according to one-way ANOVA test followed by Tukey's multiple comparisons test. Three independent biological replicates are performed.

Fig. S11. Sequence alignments of SlCf9 and NtCf9. The amino acid sequence of SlCf9 and NtCf9 are aligned using ClusterX software. Black shades indicate the conserved regions.

Fig. S12. BtRDP interacts with NtRLP4. (a) Yeast two hybrid assays showing the interaction between NtRLP4 and different whitefly salivary proteins (BtFTSP^{-sp}, BtSP16.3^{-sp} and BtSP37.4^{-sp}, without signal peptides), as well as BtRDP^{-sp} and truncated NtRLP4. The different combinations of constructs are transformed into yeast cells, and are grown on the selective medium SD/-Trp/-Leu (DDO), and the interactions are tested with SD/-Trp/-Leu/-His/-Ade (QDO). **(b)** Expression of three control salivary proteins at the protein level. The target bands are indicated by asterisks. **(c)** Bimolecular fluorescence complementation assay showing the specific interaction between BtRDP and NtRLP4. YFP fluorescence is observed when co-expressing the N-terminal nYFP tag fused NtRLP4 and N-terminal cYFP tag fused BtRDP^{-sp}. Bar =40 μm. **(d)** Co-immunoprecipitation (Co-IP) assays on oeBtRDP transgenic plants. oeBtRDP transgenic plants that overexpressing BtRDP-flag are incubated with anti-flag beads. Endogenous NtRLP4 can be immunoprecipitated by BtRDP-flag in oeBtRDP transgenic plants, while the flag tag alone in EV plants fails to immunoprecipitate endogenous NtRLP4. Rubisco staining (RbcL) is used to visualize the amount of sample loading.

Fig. S13. NtSOBIR1 interacts with NtRLP4 but not BtRDP. (a) Domain organization of NtSOBIR1. NtSOBIR1 contains a predicted N-terminal signal peptide (SP), a leucine-rich repeat (LRR) domain, a transmembrane (TM) domain, and a kinase domain. **(b)** Co-immunoprecipitation assay showing the interaction between NtSOBIR1 and NtRLP4. **(c)** Co-immunoprecipitation assay showing the interaction between NtRLP4 and BtRDP, but not NtSOBIR1 and BtRDP. The complete coding region of *NtSOBIR1* and *NtRLP4* are fused with flag and myc tags at the C-terminal ends. Total proteins are extracted from *N. benthamiana* leaves co-expressing NtRLP4-myc with GFP-flag or NtSOBIR1-flag. Precipitation is performed using flag beads. The samples are probed with anti-flag and anti-myc antibodies for immunoblot analysis. Experiments are repeated twice with the similar results.

Fig. S14. Effect of NtRLP4 overexpression on *Bemisia tabaci* performance. (a) Attraction of empty vector (EV) and oeNtRLP4#2 (oeRLP#2) transgenic *Nicotiana tabacum* to *B. tabaci* in a two-choice equipment. A group of 40 female *B. tabaci* are released into a device containing oeRLP#2 and EV leaves. The number of insects settling on each leaf is counted at 3, 6, 12, 24, 36, and 48 h. After 48 h, the number of eggs on each leaf is counted. **(b)** Comparison of insect reproduction on EV and *oeRLP*#2 transgenic plants. Five *B. tabaci* individuals are confined to indicated plants for 3 days, and the oviposited eggs are counted. **(c)** Attraction of leaves transiently overexpressing GFP and NtRLP4-GFP to *B. tabaci* in a two-choice equipment. **(d)** Comparison of insect reproduction on *N. tabacum* leaves transiently overexpressing GFP and NtRLP4-GFP. **(e)** Relative transcript level of salicylic acid (SA)- and jasmonic acid (JA)-associated genes in *N. tabacum* leaves transiently overexpressing GFP and NtRLP4-GFP. PAL, phenylalanine ammonia lyase; NPR1, nonexpressor of pathogenesis-related protein 1; FAD7, fatty acid desaturase 7; PDF1.2, plant defensin

1.2. **(f)** Transiently overexpressing NtRLP4-GFP induces H_2O_2 accumulation. For bioassays in (**a**), (**b**), (**c**), and (**d**), ten independent biological replicates are performed. For qPCR analysis in (**e**), three independent biological replicates are performed. Data are presented as mean values ± SEM. *P*-values are determined by two-tailed unpaired Student's *t* test. ***$P < 0.001$. The experiment in (**f**) is repeated five times with the similar results.

Fig. S15. Transcriptomic comparison of empty vector (EV) and oeNtRLP4#1 (oeRLP#1) transgenic plants. (**a**) Principal component analysis (PCA) of gene expression patterns in EV and oeRLP#1 transgenic *Nicotiana tabacum* plants. The first two principal components (PC1 and PC2) based on transcriptomic results are shown. Four independent biological replicates are performed. (**b**, **c**) Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analysis of differentially expressed genes (DEGs) that significantly up-regulated (**b**) and down-regulated (**c**). Enriched *P*-values are calculated according to one-sided hypergeometric test using TBtools software. (**d, e**) Expression pattens of DEGs that annotated as RLK/RLP (**d**) or WRKY transcription factor (**e**). Gene expression patterns are illustrated by a heatmap. The max transcripts per million value of each gene is set as 100.

Fig. S16. Effect of NtSOBIR1 overexpression and silencing on *Bemisia tabaci* performance. (**a**) Validation of NtSOBIR1 transient overexpression using qRT-PCR (n= 4 biological replicates). (**b**) NtSOBIR1 transient overexpression induces cell death phenotype five days post agro-injection. (**c**) Comparison of insect reproduction on *Nicotiana tabacum* transiently overexpressing GFP and NtSOBIR1-GFP. Five *B. tabaci* individuals are confined to indicated plants for 3 days, and the oviposited eggs are counted. (**d**) Attraction of GFP and NtSOBIR1-GFP leaves to *B. tabaci* in a two-choice equipment. A group of 40 female *B. tabaci* are released into a device containing

GFP and NtSOBIR1-GFP leaves. The number of insects settling on each leaf is counted at 3, 6, 12, 24, 36, and 48 h (left). After 48 h, the number of eggs on each leaf is counted (right). (**e**) Silencing efficiency of hairpin NtSOBIR1 using qRT-PCR (n= 3 independent biological replicates). (**f**) Transiently silencing NtSOBIR1 affect growth of infiltrated leaves. (**g**) Comparison of insect reproduction on *Nicotiana tabacum* leaves infiltrated with hairpin empty vector (EV) and NtSOBIR1. (**h**) Attraction of hairpin EV and NtSOBIR1-treated leaves to *B. tabaci* in a two-choice equipment. The experiments in (**b**) and (**f**) are repeated ten times with the similar results. In (**c**), (**d**), (**g**), and (**h**), ten independent biological replicates are performed. *P*-values are determined by two-tailed unpaired Student's *t* test. ***$P < 0.001$; **$P < 0.01$.

Fig. S17. Effects of *RLP* silencing on *Bemisia tabaci* performance. (**a**) Detection of NtRLP4 protein in empty vector (EV) and *NtRLP4*-silenced (*RNAi-RLP*) transgenic *Nicotiana tabacum*. Rubisco staining (RbcL) is conducted to visualize the amount of sample loading. (**b**, **c**) Attraction of EV and *RNAi-RLP* plants to *B. tabaci* in a two-choice equipment. Two NtRLP4-silenced plants (RNAi-RLP#1 in **b**; RNAi-RLP#2 in **c**) are assayed. A group of 40 female *B. tabaci* are released into a device containing two leaves. The number of insects settling on each leaf is counted at 3, 6, 12, 24, 36, and 48 h. After 48 h, the number of eggs on each leaf is counted. Ten independent biological replicates are performed. Data are presented as mean values ± SEM. *P*-values are determined by two-tailed unpaired Student's *t* test. ***$P < 0.001$.

Fig. S18. Influence of BtRDP on NtRLP4 accumulation by fluorescent analysis. The NtRLP4-GFP is transiently co-expressed with red fluorescent protein (RFP)-mCherry, BtFTSP-mCherry, or BtRDP-mCherry via agroinfiltration. The samples are imaged by confocal microscopy at 48 h post injection. RFP-mCherry and BtFTSP-mCherry are used as negative controls. Three independent biological replicates are performed, and three representative images are taken in each biological replicate. Fluorescence intensity is measured using ImageJ. The intensity values from three biological replicates are calculated and the mean value in the upper left image is set at 1.0. Representative fluorescence images are displayed. The small triangle indicates the different concentrations (OD_{600} = 0.1 and 1.0) of *Agrobacterium*. Bar= 40 μm.

Fig. S19. Effects of BtRDP on suppressing NtRLP4-associated plant defenses. (a) Comparison of insect reproduction on *Nicotiana tabacum* plants transiently overexpressing GFP along and BtRDP-mCherry/NtRLP4-GFP. Five *Bemisia tabaci* individuals are confined to indicated plants for 3 days, and the oviposited eggs are counted. **(b)** Attraction of GFP- and BtRDP-mCherry/NtRLP4-GFP-expressed leaves to *B. tabaci* in a two-choice equipment. A group of 40 female *B. tabaci* are released into a device containing two leaves. The number of insects settling on each leaf is counted at 3, 6, 12, 24, 36, and 48 h. After 48 h, the number of eggs on each leaf is counted. Data are presented as mean values ± SEM. Ten independent biological replicates are performed. *P*-values are determined by two-tailed unpaired Student's *t* test. *P < 0.05; ns, not significant.

Fig. S20. Degradation of NtRLP4 by purified BtRDP^{-sp} in *Nicotiana benthamiana* leaves. Recombinant BtRDP^{-sp}-his and GFP-his proteins are expressed in *Escherichia coli*, and purified by Ni-NTA. *N. benthamiana* plants overexpressing NtRLP4-myc or NtCf9-myc are then infiltrated with different concentrations of purified BtRDP^{-sp}-his and GFP-his. The samples are probed with anti-myc antibodies for immunoblot analysis. Rubisco staining (RbcL) is conducted to visualize the amount of sample loading. Coomassie brilliant blue (CBB) staining is conducted to visualize the amount of recombinant BtRDP^{-sp}-his and GFP-his proteins. Experiments are repeated three times with the similar results. Band density is measured using ImageJ. The density values from three biological replicates are calculated and the mean value in the first lane is set at 1.0. Data are presented as mean values ± SEM.

Fig. S21. Effect of BtRDP on NtRLP4 and NtSOBIR1. NtRLP4-myc and NtSOBIR1-flag were transiently co-expressed with different concentration of BtRDP-flag in *Nicotiana benthamiana* plants through *Agrobacterium* infiltration. The samples are probed with anti-flag and anti-myc antibodies for immunoblot analysis. Rubisco staining (RbcL) is conducted to visualize the amount of sample loading. The amount of BtRDP were controlled by infiltrating different concentrations of *Agrobacterium* (OD_{600} = 0.1, 0.3, and 1.0).

Fig. S22. Effect of BtRDP on the transcript level of *NtRLP4*. NtRLP4-myc is agroinfiltrated together with different concentration of BtRDP-flag or GFP-flag into *Nicotiana benthamiana* leaves. The transcript level of *NtRLP4* is determined by qRT-PCR. Data are presented as mean values ± SEM. Three independent biological replicates are performed. The same lowercase letters "a" indicate no statistically significant differences at *P* < 0.05 according to the one-way ANOVA test followed by Tukey's multiple comparisons test.

Fig. S23. Effects of autophagy inhibitor on NtRLP4 accumulation. NtRLP4-myc and BtRDP-flag were transiently co-expressed in *Nicotiana benthamiana* plants through *Agrobacterium* infiltration. Co-infiltrated leaves are treated with autophagy inhibitor BAF and 3-MA at 24 h post injection. The samples are probed with anti-flag and anti-myc antibodies for immunoblot analysis. Rubisco staining (RbcL) is conducted to visualize the amount of sample loading. The small triangle indicates the different concentrations of *Agrobacterium* (OD_{600} = 0.1 and 1.0). Experiments are repeated three times with the similar results. Band density is measured using ImageJ. The density values from three biological replicates are calculated and the mean value in the first lane is set at 1.0. Data are presented as mean values ± SEM. *P*-values are determined by two-tailed unpaired Student's *t* test. ***P < 0.001; **P < 0.01; *P < 0.05; ns, not significant.

Fig. S24. Sequence alignment and phylogenetic tree of RLP4 homologs. (a) The amino acid sequence of RLP4s from *Nicotiana tabacum*, *N. benthamiana*, *Solanum lycopersicum*, and *Oryza sativa* are aligned using ClusterX software. Black shades indicate the conserved regions. **(b)** Phylogenetic analysis of RLP4 and Cf9-associated genes used in this study. The unrooted phylogenetic trees are constructed by RAxML v0.9.0 using the maximum likelihood method with 1000 bootstrap replicates.

Fig. S25. Influence of BtRDP on RLP4 homology in *Solanum lycopersicum*. (a) Domain organization of *S. lycopersicum* RLP4 (SlRLP4). SlRLP4 contains a predicted N-terminal signal peptide (SP), a malectin-like domain, a LRR domain, and a transmembrane (TM) domain. **(b, c)** Yeast two-hybrid and co-immunoprecipitation (Co-IP) assays showing the interaction between BtRDP and SlRLP4. In **(b)**, BtRDP is expressed without a signal peptide (BtRDP[-sp]), while SlRLP4 and SlCf9 are expressed without a signal peptide and transmembrane domain (SlRLP4[(23-546)] and SlCf9[(24-524)]). **(d)** Effect of BtRDP on the accumulation of SlRLP4. SlRLP4-myc and SlCf9-myc are agro-injected together with different concentration of BtRDP-flag or GFP-flag. In **(c, d)**, the complete coding region of *BtRDP*, *SlRLP4*, *SfCf9* are fused with flag or myc tags at C-terminal ends, respectively. The small triangle indicates the different concentrations (OD_{600} = 0.05, 0.3, and 1.0) of *Agrobacterium*. Rubisco staining (RbcL) is conducted to visualize the amount of sample loading. Experiments are repeated three times with the similar results. Band density is measured using ImageJ. The density values from three biological replicates are calculated. For SlRLP4-myc bands (lane 1-6), the lane 1 is set at 1.0. For SlCf9-myc bands (lane 7-9), the lane 7 is set at 1.0.

Fig. S26. Yeast two hybrid assays showing the interaction between OsRLP4 and salivary proteins from planthopper species. OsRLP4$_{(29-551)}$ (OsRLP4 without signal peptides and transmembrane domains) is fused to pGADT7 vector, while salivary proteins without signal peptides are fused to pGBKT7 vector, respectively. The different combinations of constructs are transformed into yeast cells, and are grown on the selective medium SD/-Trp/-Leu (DDO), and the interactions are tested with SD/-Trp/-Leu/-His/-Ade (QDO). Twenty salivary proteins are selected. The accession number of each protein is listed as follow: (1) MF278694.1, (2) XP_039291719.1, (3) KU365967.1, (4) MF278706.1, (5) MF278711.1, (6) MF278714.1, (7) MF278715.1, (8) MF278720.1, (9) XP_022195702.1, (10) XP_022196818.2, (11) XP_039284081.1, (12) XP_022192221.2, (13) XP_022207944.2, (14) KT764973,15) RZF44823.1, (16) RZF42644.1, (17) RZF33006.1, (18) RZF48570.1, (19) RZF42817.1, (20) RZF33751.1.

Fig. S27. Three-dimensional structure of BtRDP and NlSP104. AlphaFold2 is used to predicate the protein structure. The structures are colored as a spectrum from N-terminus (blue) to C-terminus (red).

Fig. S28. Expression patterns of *Nilaparvata lugens NlSP104*. Transcripts per million (TPM) expression values of *NlSP104* in different tissues are determined based on the transcriptomic data. The head that contains salivary glands exhibits highest *NlSP104* expression in male and female *N. lugens*. Data are presented as mean values ± SEM (n=3 independent biological replicates).

Fig. S29. Original images for blots in Figures 1-3 and Supplementary Figure S13b

Fig. S30. Original images for blots in Figures 4-5

Fig. S31. Original images for blots in Figures 5 and Supplementary Figure 5-17.

Fig. S32. Original images for blots in Supplementary Figure 20-25.

2. Supplementary Tables

Table S1. Identification of RDP and SP101 homologs in insect species

Family	Species	Accession	Total Base	Number of RDP homologs	Number of SP101 homologs	Note
Aleyrodidae	*Bemisia tabaci* MED	GCA_918797505	609.6 MB	1	0	Assembled genome
	Bemisia tabaci MEAM1	GCF_001854935	600.6 MB	1	0	Assembled genome
	Bemisia tabaci India	SRR1159209	9.7 GB*	1	0	Assembled genome
	Bemisia tabaci SSA1	GCA_902825415	657.8 MB	1	0	Assembled genome
	Bemisia tabaci SSA2	GCA_903994125	625.3 MB	1	0	Assembled genome
	Aleurocanthus spiniferus	SRR17330024	9.2 GB*	1	0	RNA-seq, whole body
	Aleyrodes proletella	SRX14998536	1.9 GB*	1	0	RNA-Seq, whole body
	Trialeurodes vaporariorum	GCA_011764245	814.7 MB	1	0	Assembled genome
	Dialeurodes citri	SRR2980521	5.0 GB*	1	0	RNA-Seq, whole body
Delphacidae	*Nilaparvata lugens*	GCF_014356525	1,088 MB	0	1	Assembled genome
	Laodelphax striatellus	GCA_017141395	510.2 MB	0	1	Assembled genome
	Sogatella furcifera	GCA_017141385	563.8 MB	0	1	Assembled genome
Aphididae	*Acyrthosiphon pisum*	GCA_005508785	533.6 MB	0	0	Assembled genome

	Myzus persicae	GCA_001856785	347.3 MB	0	0	Assembled genome
	Sitobion miscanthi	GCA_008086715	397.9 MB	0	0	Assembled genome
Aphalaridae	Pachypsylla venusta	GCA_012654025	482.0 MB	0	0	Assembled genome
	Diaci psyllid	GCA_000475195	485.7 MB	0	0	Assembled genome
Alydidae	Riptortus pedestris	GCA_019009955	1.0 GB	0	0	Assembled genome
Lygaeidae	Oncopeltus fasciatus	GCA_000696205	1.1 GB	0	0	Assembled genome
Pentatomidae	Halyomorpha halys	GCA_000696795	998.2 MB	0	0	Assembled genome
Cimicidae	Cimex lectularius	GCA_000648675	510.8 MB	0	0	Assembled genome
Reduviidae	Rhodnius prolixus	GCA_000181055	706.8 MB	0	0	Assembled genome
Gerridae	Gerris buenoi	GCA_001010745	994.4 MB	0	0	Assembled genome
Drosophilidae	Drosophila melanogaster	GCA_002310755	120.4 MB	0	0	Assembled genome
Culicidae	Aedes albopictus	GCA_018104305	1.3 GB	0	0	Assembled genome
Tenebrionidae	Tribolium castaneum	GCA_000002335	165.9 MB	0	0	Assembled genome
Chrysomeloidea	Anoplophora glabripennis	GCA_000390285	679.8 MB	0	0	Assembled genome
Bombycidae	Bombyx mori	GCA_027366755	456.6 MB	0	0	Assembled genome
Noctuidae	Spodoptera frugiperda	GCA_023101765	383.9 MB	0	0	Assembled genome
Pediculidae	Pediculus humanus	GCA_000006295	110.8 MB	0	0	Assembled genome

[*] The total base of raw data retrieved from SRA database.

Table S2. Proteins from a *Nicotiana benthamiana* cDNA library screened by yeast two hybrid using BtRDP as a bait

N. benthamiana v2.6.1 Accession [1]	Annotation	Number of colonies	Region
Niben261Chr07g1310001.1	Putative leucine-rich repeat receptor-like serine/threonine-protein kinase	3	455-709
Niben261Chr13g1396009.1	Chloroplast stem-loop binding protein of 41 kDa b	1	162-374
Niben261Chr02g0086007.1	Sorbitol dehydrogenase-like	1	95-305
Niben261Chr17g0179002.1	Guanine nucleotide-binding protein subunit beta-like protein	1	92-326
Niben261Chr01g1673009.1	Ribulose bisphosphate carboxylase/oxygenase activase 2	2	228-420
Niben261Chr07g1128067.1	Ribulose-1,5-bisphosphate carboxylase/oxygenase small subunit	5	23-181
Niben261Chr14g0161005.1	Cytochrome b6-f complex iron-sulfur subunit 2	5	1-216

[1] The sequence can be downloaded from Sol Genomics Network (https://solgenomics.net/ftp/genomes/Nicotiana_benthamianaV261)