p63-GATA2 molecular switch mediates the tumor suppressor to oncogene transition of glucocorticoid receptor in the prostate

Johannes Hiltunen[1], Niina Aaltonen[1], Heini Sohlberg[1], Laura Kemppi[1], Ville Paakinaho[1, *]

[1]Institute of Biomedicine, University of Eastern Finland, Kuopio, Finland.

[*]Correspondence: ville.paakinaho@uef.fi

Abstract

Although glucocorticoids are widely used to alleviate side effects of prostate cancer (PCa) treatment, the glucocorticoid receptor (GR) exhibits a dual role exerting tumor-suppressive effects by inhibiting early-stage PCa cell proliferation, while also promoting oncogenic progression by mediating antiandrogen resistance. The mechanisms underlying this functional dichotomy have remained elusive and poorly characterized. Using genome-wide analyses and CRISPR-based genome editing, we identified the tumor protein p63 as a key mediator of GR's tumor-suppressive chromatin activity. Loss of p63 reprograms GR activity toward an oncogenic state, marked by enhanced cell migration, invasion, and altered morphology. This shift is driven by increased GATA2 expression, which alters GR's chromatin binding and transcriptional output. Together, our findings uncover a p63–GATA2 molecular switch that governs the dual role of GR in PCa, establishing transcription factor crosstalk as a critical regulator of GR-driven oncogenic reprogramming and cellular plasticity.

Keywords

chromatin, GATA2, glucocorticoid receptor, p63, prostate cancer

Introduction

Prostate cancer (PCa), one of the most prevalent cancers worldwide [1], is primarily driven by signaling through the androgen receptor (AR) [2,3]. AR belongs to the steroid receptor subfamily of nuclear receptors, which also includes estrogen receptor (ER) and glucocorticoid receptor (GR). These receptors are ligand-dependent transcription factors (TFs) that, upon hormone binding, translocate to the nucleus, bind chromatin, and regulate gene expression. This regulation is mediated through coordinated recruitment of TFs and coregulators with histone-modifying and chromatin-remodeling functions [4,5].

Because AR-driven gene regulatory programs are central to PCa development and progression, inhibition of AR signaling remains the cornerstone of PCa therapy. Depending on disease stage, treatment strategies include androgen deprivation therapy (ADT), antiandrogens such as enzalutamide and darolutamide, and androgen biosynthesis inhibitors like abiraterone [2,3]. Although these therapies are initially effective, many tumors eventually develop resistance to AR-targeted treatments. In this resistant state, cancer progression is often sustained by alternative TFs [6]. Notably, TFs such as FOXA1, GATA2, and SOX2 can reprogram chromatin accessibility and promote lineage plasticity, enabling continued tumor growth in the absence of AR activity [7–9]. Moreover, these TFs can drive antiandrogen resistance by modulating the activity of other steroid receptors, including GR [10,11].

Glucocorticoids—the activating ligands of GR—have traditionally been used as adjuvant therapy in PCa to alleviate treatment-related side effects [12]. However, GR is now recognized as a context-dependent oncogene in PCa, capable of mediating drug resistance by compensating for the loss of AR signaling under antiandrogen therapy [13]. While the pro-tumorigenic role of GR in therapy-resistant PCa has been well documented by us and others [10,14,15], its function during earlier stages of the disease remains elusive. Notably, early studies suggested a tumor suppressive role for GR in primary PCa, where it inhibited proliferation and oncogenic signaling pathways [16]. These findings point to a striking functional shift in GR activity—from

tumor suppressor to oncogene—throughout disease progression. However, the molecular mechanisms underlying this transition remain poorly understood. Given GR's ubiquitous expression, this switch is likely governed by changes in the transcriptional landscape and signaling networks driven by other TFs.

It is well established that the tumor suppressive or oncogenic activity of a TF can be modulated by TF crosstalk, as well as by lineage and epigenetic plasticity [7,17,18]. For steroid receptors, the clearest examples of this come from breast cancer. Both GR and AR exhibit tumor suppressive roles in ER-positive breast cancers, while exerting oncogenic functions in ER-negative subtypes [4,19,20]. This highlights how the presence or absence of ER signaling can dramatically reshape the functional consequences of other steroid receptors. Moreover, this context-dependent shift is not limited to interactions among steroid receptors. For example, AR's tumor-suppressive role in ER-positive breast cancer can be reprogrammed into an oncogenic function through activation of the JAK/STAT pathway [21]. These findings underscore the capacity of distinct TF networks and signaling pathways to serve as molecular switches, governing the tumor suppressor-to-oncogene transition of steroid receptors.

To investigate the transition of GR from a tumor suppressor to an oncogene in the prostate, we examined its transcriptional activity across various PCa cell lines and immortalized normal prostate epithelial cells. We found that both the GR-regulated transcriptome and its cistrome differ markedly between normal and cancerous prostate cells, regardless of PCa cells' AR status. In normal prostate cells, GR binding sites are uniquely co-occupied by the basal epithelial TF p63, a member of the p53 family. Loss of p63 diminishes GR's anti-proliferative effects and instead promotes GR-driven cell invasion. This phenotypic switch is associated with increased GATA2 activity. In the absence of p63, GATA2 reprograms GR's transcriptional output from tumor suppressive to oncogenic pathways. Together, these findings suggest that GR's oncogenic transition in prostate cancer is governed by dynamic changes in transcription factor crosstalk.

Results
Chromatin binding and GR regulated transcriptome separates prostate cells from each other
For comprehensive investigation of GR action in the prostate, we examined GR-regulated transcriptome with RNA-seq and chromatin binding (cistrome) of GR with ChIP-seq from two AR-positive (VCaP, 22Rv1) [10] and two AR-negative (PC3, DU145) PCa cell lines (Supplementary Figure S1A) [22]. In addition, we employed immortalized normal prostate epithelial cell line RWPE1 [23], which express similar levels of GR as DU145 cells (Supplementary Figure S1B) [15], as model for non-cancerous prostate cells. Intriguingly, principal component analysis (PCA) of GR's transcriptomes (Figure 1A) and cistromes (Figure 1B), when treated with synthetic GR agonist dexamethasone (Dex), display clear separation based on the subtype of prostate cells. VCaP and 22Rv1 formed one cluster while another was formed by PC3 and DU145 cells. Moreover, GR action in RWPE1 cells differentiated from both AR-positive and -negative PCa cells. As an example, GR binding sites (GRBs) at *PER1* locus are shared between all the cell lines, while *KRT16* locus shows RWPE1-specific GRBs and *PTPRJ* locus shows primarily PCa-specific GRBs (Figure 1C). The prostate cell subtype-specific GR activity could be explained by the distinct cooperative partners of the receptor, since the transcriptional activity of GR is modulated by other TFs in PCa [13].

To investigate such TF partners, we analyzed motif enrichment of the GRBs (Supplementary Table S4). Glucocorticoid response elements (GREs) were prominently enriched in all cell lines, with PCa cells exhibiting higher enrichment than RWPE1 cells (Figure 1D). While motifs for luminal epithelial TFs, such as GATA2, FOXA1 and HOXB13 were accumulated at GRBs in VCaP and 22Rv1 cells, AP-1 motif was considerably enriched at PC3 and DU145 cells. These distinct motif enrichments are concordant with the known role of the former TFs in AR-positive PCa and the latter TF in AR-negative PCa [6]. Most intriguingly, tumor suppressor p53-family motifs were prominently enriched only at the GRBs of RWPE1 cells. The motif enrichments matched with the TF expression levels; *GATA2*, *FOXA1*, and *HOXB13* are highly expressed in AR-positive PCa cells, *FOSL1*, subunit of AP-1, is prominently expressed in AR-negative PCa cells, and *TP53* and *TP63* are strongly expressed in RWPE1 cells (Figure 1E, Supplementary Figure S1C). These

results suggest that GR's cooperative TFs mediate its action in non-cancerous and cancerous prostate cells, likely influencing the tumor suppressive and oncogenic activities of the receptor.

Figure 1. GR has distinct chromatin binding profile in normal and malignant prostate cells. (**A-B**) PCA plot of (**A**) GR-regulated transcriptome and (**B**) GR cistrome from RWPE1, VCaP, 22Rv1, PC3, and DU145 cells. (**C**) Genome browser tracks of GR ChIP-seq at *PER1*, *KRT16*, and *PTPRJ* loci. Red arrows point specific peaks for each locus. (**D-E**) Heatmap of (**D**) GRB motif enrichment and (**E**) expression of corresponding TFs. Motif enrichment values calculated as fold change over background, and expression as TPM values. (**F**) Relapse-free survival of TCGA-PRAD patients expressing either high or low levels of (left) *NR3C1* or (right) *TP63*. (**G**) Progression-free survival of TCGA-PRAD patients expressing high levels of both *TP63* and *NR3C1*, or low levels of both *TP63* and *NR3C1*. Other patients were classified as unclassified. Survival p-values calculated with log-rank test. PRAD, prostate adenocarcinoma; TCGA, the cancer genome atlas.

To complement the RWPE1 data, we also generated GR cistrome from immortalized normal prostate epithelial cell line PNT1A [24]. PCA of GR cistromes indicated closer association of PNT1A cells with RWPE1 than DU145 cells (Supplementary Figure S1D-E). Moreover, p53-family motifs were prominently

enriched also at the GRBs of PNT1A cells (Supplementary Figure S1F), and both RWPE1 and PNT1A peaks were equally associated with p53 pathway (Supplementary Figure S1G). During the progression of PCa, the *NR3C1* expression shows minimal change, while *TP63* is the only p53-family member that is prominently decreased during cancer progression (Supplementary Figure S1H). Intriguingly, the high expression of *NR3C1* and *TP63* are associated with significantly improved relapse-free survival (Figure 1F), while no change is seen with *TP53* and *TP73* expression (Supplementary Figure S1I). Moreover, PCa patients with concurrent high expression of *NR3C1* and *TP63* have the best progression-free survival (Figure 1G). In sum, our results indicate that p53-family members, especially p63, could mediate the tumor suppressive activities of GR in the normal prostate.

GR cistrome exhibits substantial overlap with p63 binding sites

Considering that p53-family is well-characterized for its overall significance in cancer [25], and normal prostate GRBs seem to be linked with p53-family members, we further explored their role in modulating GR activity. We profiled the chromatin binding of both p53 and p63 in RWPE1 cells. While the epithelial isoform DNp63α is the most abundant p63 in prostate tissue [26], for simplicity we will utilize p63 term throughout the text. Both TFs overlapped with GRBs, but compared to p53, p63 showed a considerable (over 70%) overlap with GRBs (Figure 2A, Supplementary Figure S2A). Moreover, p53-family motifs were enriched only at GRBs that showed p63 binding regardless of p53 occupancy (Figure 2B, Supplementary Table S4). GRBs that overlapped only with p53 showed no p53-family motifs enrichment. The lack of motif enrichment at p53 binding sites can be partly explained by the transient nature of p53 stability in non-stressed cells [27]. Intriguingly, GRBs co-occupied with p63 showed minimal enrichment of GREs, while the center of the GRBs showed accumulation of p53 and p63 motifs (Figure 2C). Close inspection of the motif indicated that part of the GRE half site overlaps with the p63 motif (Supplementary Figure S2B). Around 11% of the p63-GR sites had either complete (5.7%) or partial (5.6%) overlap between the half GRE and p63 motif (Supplementary Figure S2C). In addition, the p63 binding was significantly stronger in sites with either complete or partial motif overlap compared to sites without overlap (Supplementary Figure S2D). These analyses highlight that p63 is anticipated to modulate the chromatin binding of GR to its binding sites in RWPE1 cells. Based on these observations, we focused on the relationship between GR and p63.

Clustering of GR and p63 binding sites indicated that GR activation had minimal effect on the chromatin occupancy of p63 (Figure 2D-E). This suggests that p63 chromatin occupancy precedes GR activation, likely facilitating GR recruitment to these sites. Intriguingly, the binding of both p63 and GR was significantly stronger at shared sites compared to sites where they bind on their own (Figure 2E). Moreover, the active enhancer histone mark, histone 3 lysine 27 acetylation (H3K27ac), was most prominently enriched at p63-GR shared binding sites (Figure 2F). These results highlight the biological relevance of GR and p63 co-occupying sites. This is supported by pathway analyses showing a significant enrichment of p63-GR binding sites with the p53 as well as apoptosis and epithelial-mesenchymal transition (EMT) pathways (Figure 2G).

While p63 motif was specifically enriched at RWPE1 GRBs, AP-1 motifs showed enrichment at GRBs in multiple cell lines (Figure 1D). To further assess the distinctive relationship between GR and p63, we profiled chromatin binding of FRA1 in RWPE1 cells. We selected FRA1 for its prominent role in PCa [6]. FRA1 had a prominent (over 60%) overlap with GR while its overlap with p63 was only 8% (Supplementary Figure S2E). Most p63 sites overlapped with GR without FRA1. The GRBs that harbored p63, with or without FRA1, had high enrichment whereas GRBs with FRA1 lacking p63 had low enrichment of H3K27ac (Supplementary Figure S2F). This suggests that GRBs with p63 are the most active enhancers in RWPE1 cells. Finally, single-cell RNA-seq (scRNA-seq) analyses indicate that over 75% of RWPE1 cells expressed both *NR3C1* and *TP63* in the same single cell (Figure 2H). This highlights the interconnected relationship between GR and p63. Interestingly, GR activation downregulated the transcription of both *NR3C1* and *TP63* (Supplementary Figure S2G), reducing the number of cells expressing both genes (Figure 2H). This supports the existence of negative regulatory feedback loop, as previously observed by Sevilla et al. (2024)[28]. Thus, GR and p63 crosstalk is likely a multifaceted phenomenon occurring not only at the chromatin level but also via transcriptional regulation.

Figure 2. GRBs have extensive overlap with p63 in RWPE1 cells. (**A**) Overlap of GR (blue), p53 (purple), and p63 (green) ChIP-seq peaks in Dex-treated RWPE1 cells. (**B**) Percentage of p53 (grey), p63 (orange) and GRE (blue) motif enrichment at GRBs shared with only p53, with both p53 and p63, only with p63, or neither p53 nor p63. (**C**) Enrichment of p53 (yellow) and p63 (blue) motifs surrounding GR and p63 shared binding sites. (**D**) p63 and GR ChIP-seq profiles at GR, p63-GR, and p63 sites in RWPE1 cells. GR sites bind only GR, p63-GR sites bind both GR and p63, and p63 sites bind only p63. For GR sites, only top 7000 sites are shown in the heatmap. Each heatmap represents ± 500 bp around the center of the peak, with binding intensity (tags per bp per site) scale is noted below on a linear scale. (**E**) Log_2 tag enrichment of (left) p63 and (right) GR at the indicated binding sites. (**F**) Log_2 tag enrichment of H3K27ac at the indicated binding sites. (**G**) Hallmark and CORUM Gene Sets pathway analysis of p63-GR sites. Color scale represents $-log_{10}$ p-value, and distinct pathways highlighted in red. Pathway names with asterisk are truncated. (**H**) Single cells expressing *NR3C1* (orange), *TP63* (grey), both (blue), or neither (yellow) in scRNA-seq data from RWPE1 cells. Expression is defined as a non-zero value. Statistical significance in box plots was calculated with One-way ANOVA with Bonferroni *post hoc* test. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

Loss of p63 leads to prominent changes in GR-regulated transcriptome

To investigate the degree of p63 influence on GR activity, we utilized CRISPR-Cas9 to knockout p63 from RWPE1 cells (p63KO) (see Methods for details). The utilized guide RNA (gRNA) generated a frameshift mutation resulting in premature STOP codon at exon 5 before critical amino acids for p63's DNA binding (Supplementary Figure S3A). Two monoclonal p63KO cell lines, clone 1 (CL#1) and 2 (CL#2), were utilized in the downstream analyses (Figure 3A). RWPE1 cell line exposed to non-targeting gRNA followed by monoclonal selection was used as control (CTRL) cell line. PCA plot indicated that compared to CTRL, the transcriptome of p63KO cells, especially CL#1, reshaped towards the AR-negative PCa cell transcriptome (Supplementary Figure S3B). Intriguingly, while the transcriptome of p63KO cells were close to p63-null PC3 cells, the CTRL transcriptome resembled the transcriptome of PC3 cells overexpressing p63 [29] (Supplementary Figure S3C). These results suggest that gain or loss of p63 leads to bidirectional transcriptomic changes.

AR-positive PCa cells exhibit a luminal epithelial (LE) phenotype, while the disappearance of basal epithelial (BE) cell layers is potentially associated with PCa development [30]. Since parental RWPE1 cells display LE-BE intermediate phenotype [23], and p63 is a known regulator of BE phenotype and inhibitor of EMT [31,32], we quantified the expression of LE, BE, and EMT marker genes in CTRL and p63KO cells. Interestingly, the loss of p63 led to phenotypic switch wherein p63KO cells expressed more LE and less BE marker genes compared to CTRL (Figure 3B, Supplementary Figure S3D-E). Despite the increase in LE marker genes, we observed no *AR* or *KLK3* expression upon p63KO. In parallel with LE marker gene induction, genes associated with EMT were prominently increased upon p63KO (Figure 3B, Supplementary Figure S3F). This was corroborated at pathway level with differentially regulated genes (DEGs) in p63KO displaying a prominent enrichment of EMT pathway (Figure 3C-D). Moreover, other cancer progression related pathways, such as VEGFA VEGFR2 signaling and embryonic stem cell pluripotency, were enriched with p63KO DEGs. Focusing on Dex-regulated DEGs, the loss of p63 lead to a clear transcriptomic change where several genes gained or lost their Dex-mediated regulation (Figure 3E). Intriguingly, while GR pathway was enriched with Dex-regulated genes in both CTRL and p63KO cells, only the latter showed enrichment of PCa malignancy related pathways, such as VEGFA VEGFR2 signaling, AR network in PCa, and cancer pathways (Figure 3F). These results imply that p63 loss alters GR to have a more oncogenic potential in prostate cells.

Figure 3. Depletion of p63 alters cell identity and EMT marker gene expression. (**A**) Immunoblotting of GR, p63, and tubulin protein levels in RWPE1 control (CTRL) and p63 depleted (p63KO) cells. Two clones (CL#1 and CL#2) of p63KO cells were utilized. (**B**) Average Z-scores for luminal epithelial (LE) (left; green), basal epithelial (BE) (middle; red), and epithelial-mesenchymal transition (EMT) (right; blue) marker gene expression in CTRL and p63KO cell lines. (**C**) Overlap of DEGs in CL#1 (green) and CL#2 (red) p63KO cells. (**D**) WIKI pathway analysis of CL#1 and CL#2 DEGs. (**E**) Overlap of Dex-regulated DEGs in CTRL (blue), CL#1 (green) and CL#2 (red) p63KO cells. (**F**) WIKI pathway analysis of shared or CL#1 and CL#2 (CL#1+2) specific Dex-regulated DEGs. In pathway heatmaps, color scales represent -log$_{10}$ p-value, and distinct pathways are highlighted in red. Pathway names with asterisk are truncated.

Absence of p63 leads to pronounced changes in cell morphology, proliferation, migration, and invasion
Given the p63-depletion induced expression of LE and EMT marker genes and GR's prominent role in PCa cell survival [15], we sought to investigate whether these changes corresponded with enhanced oncogenic potential. In agreement with phenotypic transcriptome changes, p63KO cells showed altered morphology with round shape converted to a more jagged and elongated appearance (Figure 4A-B). The opposite effect has been seen with PC3 cells where overexpression of p63 converts cells' elongated shape to a rounder appearance [33]. The p63KO-mediated change in morphology was associated with higher proliferation rates and attenuation of the anti-proliferative effects of Dex (Figure 4C, Supplementary Figure S4A-B). Moreover, a similar significant increase in migration and suppression of the anti-migrative effects of Dex was observed in p63KO compared to CTRL cells (Figure 4D, Supplementary Figure S4C-E). The concordant effect on migration is seen with overexpression of p63 in PC3 cells [33]. Thus, increased proliferation and migration are driven by the loss of p63 while GR's effect on these changes is diminished.

Since EMT pathway was prominently enriched with p63KO DEGs, including Dex-regulated genes, we measured the 3D invasiveness of the cells (see Methods for details). While proliferation and migration displayed significant changes, the cells' invasiveness showed only slight change upon p63KO (Figure 4E, Supplementary Figure S4F). However, Dex treatment led to significant increase in p63KO cells' invasiveness which was not observed in CTRL cells (Figure 4E, Supplementary Figure S4F). This indicates that Dex-regulated genes emerged upon p63 loss (Figure 3E-F) mediate the increased invasive potential of

the cells. Therefore, the loss of p63 shifts GR behavior from tumor suppressive towards more oncogenic activity.

Figure 4. Cells' morphology, proliferation, migration, and invasion are altered by p63 depletion. (**A**) Immunofluorescence (IF) images of RWPE1 CTRL and p63KO cells with nuclei (DAPI; blue) and cell membrane (actin; red) staining. Scale bar, 20 μm. (**B**) Aspect ratio from CTRL and p63KO cells, displaying the ratio of cell length to width (n=37-41). (**C**) Relative cell proliferation of CTRL and p63KO cells on day 5 (n=5) with indicated treatments. (**D**) Relative migration in wound healing assay of CTRL and p63KO cells on day 2 (n=8) with indicated treatments. (**E**) 3D gel invasion of CTRL and p63KO cells on day 6 with indicated treatments. Two of the highest invasion points were utilized (n=36-52). Statistical significance in box plots was calculated with One-way ANOVA with Bonferroni *post hoc* test. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

Reprogramming GR cistrome upon loss of p63

Due to the clear Dex-induced changes in the cells' invasiveness, we profiled GR cistrome from p63KO cells. Comparison of GR cistromes indicated that p63KO led to both loss and gain of GRBs (Supplementary Figure S5A). Similar to transcriptomes (Supplementary Figure S3B), PCA plot showed that the GR cistromes of p63KO cells, especially CL#1, advanced towards the GRB profile of PCa cell lines (Supplementary Figure S5B). As an example of GRBs, *MAOA* and *GNAO1* loci showed loss of p63 co-occupying GRBs, while *GCNT1* locus displayed gain of GRB (Figure 5A). Moreover, these changes were correlated at gene regulatory level, as *MAOA* and *GNAO1* lost while *GCNT1*, associated with extracapsular PCa occurrence [34], gained Dex-induction in p63KO cells (Figure 5B). To investigate what drives these changes, we performed motif enrichment analyses from the p63KO GRBs (Supplementary Table S4). In accordance with the depletion of p63, the p53-family motifs displayed the most prominent loss at GRBs (Figure 5C). At transcript level only *TP63* showed significant reduction, indicating that the loss of p63 drives

the changes. While GR itself, *i.e.*, GRE enrichment and *NR3C1* transcript levels, remained largely unaffected, *FOSL1* transcripts were significantly increased in p63KO cells. However, GRBs showed no prominent difference in AP-1 motif enrichment. In comparison to the mixed AP-1 differences, p63KO cells exhibited a distinct increase in GATA-family motif enrichment and elevated *GATA2* transcript levels (Figure 5C).

To better examine what mediates the GRB changes upon p63KO, we divided GR cistromes to p63KO-lost, -retained, and -gained GRBs (see Methods for details) (Figure 5D, Supplementary Figure S5C). The lost and retained GRBs were enriched at the p53 pathway, whereas only gained GRBs showed prominent enrichment of androgen response pathway (Supplementary Figure S5D). This suggests that both gained GRBs and Dex-regulated genes in p63KO cells (Figure 3F), mimic the PCa cells' regulatory profiles. In addition, the chromatin accessibility and H3K27ac enrichment were significantly decreased at lost GRBs whereas they were significantly increased at gained GRBs upon p63KO (Supplementary Figure S5E-F). These results highlight that p63 mediates the enhancer activity of lost GRBs, while other TFs likely have a reciprocal effect at gained GRBs. To identify such TFs, we performed motif analyses. While GREs were enriched at all GRBs, with the clearest enrichment at retained GRBs, p63 motif was enriched only at lost GRBs (Figure 5E). The latter was corroborated with p63 ChIP-seq data from parental cells, indicating that lost GRBs were the most p63 occupied sites (Figure 5D, Supplementary Figure S5C). This confirms that GR binding to these sites is determined by p63, and p63 loss leads to the concomitant loss of GR binding. Even though AP-1 is a known mediator of GR action [35] and AP-1 motifs showed some enrichment at gained GRBs, the most prevalent enrichment of AP-1 motifs were observed at lost and retained GRBs. FRA1 ChIP-seq data from parental cells indicated occupancy mainly at retained GRBs (Figure 5D, Supplementary Figure S5C). This suggests that although *FOSL1* transcript levels are increased, FRA1 alone is not mediating GR binding to gained GRBs in p63KO cells.

Intriguingly, gained GRBs presented prominent enrichment of GATA-family motifs that were not observed at lost or retained GRBs (Figure 5E). Of the GATA-family members, only *GATA2* and *GATA3* showed a prominent increase upon p63KO (Supplementary Figure S5G), and the promoters of *GATA2* and *GATA3* displayed significantly increased chromatin accessibility and H3K27ac enrichment in p63KO compared to CTRL cells (Supplementary Figure S5H). Interestingly, GATA2 has a prominent role in PCa [36], and its expression is high in early-stages of PCa (Supplementary Figure S5I). Moreover, LE marker expression was increased in p63KO cells and GATA2 is associated with luminal differentiation. Thus, we focused on the role of GATA2. First, we confirmed the prominent increase of GATA2 protein levels in p63KO cells (Supplementary Figure S5J), followed by GATA2 ChIP-seq analyses from CTRL and p63KO cells. In agreement with the increased protein levels, more GATA2 occupied sites were observed in p63KO compared to CTRL cells (Supplementary Figure S5K). More importantly, GATA2 binds to the gained GRBs in p63KO cells being absent from the lost GRBs (Figure 5D, Supplementary Figure S5C). Furthermore, the GATA2-GR binding sites were associated with EMT and androgen response pathways (Figure 5F). Our results suggest that both p63 and GATA2 function as pioneer factors, as previously indicated [37,38], mediating GR's occupancy at distinct sites in asynchronous fashion. The loss of p63 leads to increased GATA2 levels resulting in the reshaping of GR chromatin occupancy from p63 to GATA2 binding sites.

Figure 5. Depletion of p63 modulates the GR cistrome. (**A**) Genome browser tracks of p63 ChIP-seq from parental RWPE1 cells, and GR ChIP-seq from CTRL and p63KO cells at *MAOA*, *GNAO1*, and *GCNT1* loci. Red arrows point specific peaks for each locus. (**B**) RNA-seq TPM values of *MAOA*, *GNAO1*, and *GCNT1* from EtOH- and Dex-treated CTRL and p63KO cells. Bar graphs represent mean ±SD, n=2. (**C**) Heatmap of GRB motif enrichment and corresponding TF expression change in p63KO cells. Motif enrichment and expression values calculated as log2 fold change over CTRL cells. (**D**) ChIP-seq profiles of p63 and FRA1 from parental RWPE1 cells, GATA2 and GR from CTRL and p63KO cells at p63KO-lost, -retained, and -gained GRBs. Each heatmap represents ± 1 kb around the center of the peak, and binding intensity (tags per bp per site) is noted below on a linear scale. (**E**) Motif density of GRE, p63, AP-1, and GATA at indicated p63KO GRBs. Histograms represent ± 240 bp around the center of the peak, and enrichment intensity represents motif per bp per site. (**F**) Hallmark Gene Sets pathway analysis of GATA2-GR sites. Color scale represents -$\log_{10}$ *p*-value, and distinct pathways highlighted in red.

GR's harmful effects in p63-null cells can be mitigated with GATA2 depletion

Our results highlight that GR's chromatin action is reprogrammed through the crosstalk with p63 and GATA2 with the former mediating GR's tumor suppressive-like effects while the latter conveys GR's oncogenic activities. Intriguingly, scRNA-seq data from human and mouse prostate tissue as well as from PCa patients shows the general mutually exclusive cell type-specific expression of *TP63* and *GATA2*, wherein *TP63* is expressed mainly in basal cells while *GATA2* is expressed in luminal and PCa cells (Figure 6A). In comparison, *NR3C1* is expressed in most cells, suggesting that other TFs, such as p63 and GATA2, mediates its cell type-specific action. At the level of survival, the high expression of *GATA2* is associated with decreased relapse-free survival (Supplementary Figure S6A), which is opposite to the *NR3C1* and *TP63* expression (Figure 1F). More importantly, PCa patients with the high *TP63* and low *GATA2* expression have the best while high *GATA2* and low *TP63* expression have the worst progression-free survival (Figure 6B). This indicates that the differential expression of *TP63* and *GATA2* has clinical significance, possibly through altering GR activity.

To further evaluate GATA2's contribution, we depleted it from the p63KO cells using siRNA knockdown (Supplementary Figure S6B), and analyzed transcriptomic changes with RNA-seq. Depletion of GATA2 altered the expression of numerous transcripts including *AREG*, *ITGB1*, and *SGK1* (Supplementary Figure S6C). As an example, the last is a known GR target gene that shows increased GATA2 and GR binding upon p63KO (Figure 6C), which lost its p63KO-potentiated Dex-induction when GATA2 was depleted (Figure 6D). Pathway analyses indicated that siGATA2 downregulated genes were enriched at EMT pathways (Figure 6E). Similar enrichment was observed with Dex-regulated genes that showed significant attenuation of Dex response upon siGATA2 treatment. Importantly, only siGATA2-attenuated Dex-regulated genes were enriched at androgen response pathway (Figure 6E). Thus, while GATA2 regulates EMT in Dex-dependent and -independent manner, GR together with GATA2 is responsible for the regulation of androgen response pathway (Figure 3F, Figure 5F).

Since EMT and androgen response pathway were impacted by siGATA2, we analyzed the proliferation and invasion of the cells after GATA2 depletion. The proliferation of p63KO cells was significantly inhibited with siGATA2 treatment (Figure 6F, Supplementary Figure S6D-E). However, while anti-proliferative effects of Dex were prominent in p63-intact cells (Figure 4C), GATA2 depletion did not have any effect on the Dex responsiveness of p63KO cells (Figure 6F, Supplementary Figure S6D-E). In comparison, GATA2 depletion significantly impaired the GR-induced invasiveness of p63KO cells (Figure 6G). This suggests that the crosstalk between GATA2 and GR regulates the expression of EMT pathway manifesting in increased cell invasion.

Since depletion of GATA2 did not completely remove all the GR effects induced by p63KO, we investigated other potential contributing TFs using epigenetic landscape in silico deletion analysis (LISA) [39]. GRBs with GATA2 were associated with GATA-family members, as expected, whereas AP-1 subunit FRA2 (encoded by *FOSL2*) was highly associated with GRBs without GATA2 (Supplementary Figure S6F). The expression of *FOSL2* did not change, but the expression of *FOSL1* was significantly increased in p63KO compared to

CTRL cells (Supplementary Figure S6G). Moreover, siGATA2 significantly decreased *FOSL1* but not *FOSL2* expression (Supplementary Figure S6H). This suggests that FRA1 (encoded by *FOSL1*) could influence GR's transcriptional activity in p63KO cells. However, since GATA2 and FRA1 primarily occupy different GRBs (Figure 5D), they likely influence the action of GR in a distinct manner.

Our results suggest that GR's tumor suppressive and oncogenic action in prostate cells is largely dependent on its crosstalk partner. With p63, GR regulates basal and p53 pathway genes restricting the proliferation and migration of cells (Figure 6H). Upon loss of p63, GR switches crosstalk partner to GATA2. With GATA2, GR regulates luminal and EMT pathway genes inducing the invasiveness of cells (Figure 6H). Thus, GR's expression can remain unchanged during the initiation and progression of PCa, but its action can be drastically altered by other TFs mediating its chromatin activities.

Figure 6. GATA2 promotes GR-induced invasion. (**A**) Single cell expression of *Gata2*, *Trp63* and *Nr3c1* in mouse prostate, and *GATA2*, *TP63* and *NR3C1* in human prostate and metastatic (m)CRPC samples. Data obtained from Single Cell Portal, and intensity of the color depicts the scaled expression, and size of the circle depicts the percentage of cells that express the gene. (**B**) Progression-free survival of TCGA-PRAD patients expressing high levels of both *TP63* and *NR3C1*, or low levels of both *TP63* and *NR3C1*. Other patients were classified as unclassified. (**C**) Genome browser track of GR and GATA2 ChIP-seq from CTRL and p63KO cells at *SGK1* locus. (**D**) RNA-seq TPM values of *SGK1* from EtOH- and Dex-treated CTRL and p63KO cells, and siNON- and siGATA2-treated p63KO cells. Bar graphs represent mean ±SD, n=2. (**E**) Hallmark Gene Sets pathway analysis of siGATA2 downregulated (DN) genes where Dex-regulation is lost or without Dex-regulation. Color scale represents -$\log_{10}$ *p*-value, and distinct pathways highlighted in red. (**F**) Relative cell proliferation of p63KO cells on day 4 (n=5) with indicated treatments. (**G**) 3D gel invasion of p63KO cells on day 6 with indicated treatments. Two of the highest invasion points were utilized (n=50-54). (**H**) Model of the reprogramming of GR action through p63 and GATA2. (Upper) GR-p63 regulates basal identity, p53 pathway and has anti-proliferative and -migrative effects. Upon loss of p63 and increase of GATA2, (lower) GR-GATA2 regulates luminal identity, androgen response (in the absence of AR) and EMT. Survival *p*-values calculated with log-rank test. Statistical significance in box plots was calculated with One-way ANOVA with Bonferroni *post hoc* test. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$. PRAD, prostate adenocarcinoma; TCGA, the cancer genome atlas.

Discussion

Glucocorticoids are among the most widely prescribed drugs globally [40] and are extensively used to mitigate treatment-related side effects in cancer patients [12]. In PCa, supplemental glucocorticoids are particularly important to counteract the suppression of serum cortisol levels and prevent a compensatory rise in adrenocorticotropic hormone caused by abiraterone treatment. However, the role of the GR in PCa is complex. Studies have demonstrated both pro- and anti-tumorigenic effects of GR [10,14–16], underscoring the clinical importance of understanding when GR functions as a tumor suppressor versus as an oncogene [13]. Despite this significance, GR's context-dependent activity and the underlying molecular mediators have remained elusive. Here, we identify a molecular switch involving p63 and GATA2 that governs GR's transition from a tumor-suppressive to an oncogenic state.

Our genome-wide analyses revealed that GR's chromatin binding and transcriptional responses differ markedly between PCa cells and immortalized prostate epithelial cells. In PCa cells, GR activity was closely associated with the TFs FOXA1, GATA2, and AP-1, all well-established oncogenic drivers in PCa [6] with FOXA1 and GATA2 specifically mediating GR's pro-tumorigenic functions [10,41,42]. In contrast, our data indicated that in immortalized prostate cells, GR predominantly associated with the p53-family member p63. Our findings demonstrate that p63 plays a central role in directing GR's anti-tumorigenic activity. Given that GR expression remains relatively constant across normal and malignant prostate tissues [43,44], and that GR function shifts during PCa progression [13], it is likely that differential TF crosstalk, rather than GR abundance, underlies its context-dependent activity.

PCa typically exhibits a loss of BE cells, while its origin is most commonly traced to LE cells [30,45]. However, BE cells can also serve as a source of PCa following their differentiation into an LE-like state [46]. Our investigation reveals that p63-null prostate cells display a diminished BE phenotype and an enhanced LE-like identity, consistent with the well-established role of p63 as a key regulator of BE cell fate [31,32]. In addition to this shift in epithelial identity, p63-null cells exhibit notable morphological changes and elevated expression of EMT markers, supporting previous findings that p63 suppresses EMT in the prostate [32,47]. Intriguingly, despite the upregulation of EMT markers, GR activation was required to drive the invasive behavior of p63-null cells.

In p63-intact prostate cells, GR regulated p53-associated gene expression programs through active enhancers co-occupied by both GR and p63. At these shared regulatory sites, GR suppressed cell proliferation and migration. Patient data supported these tumor-suppressive effects, showing the most favorable survival outcomes in individuals with high expression of both GR and p63. However, the protective role of GR was markedly altered upon p63 depletion. In p63-null prostate cells, GR's chromatin binding profile was reprogrammed, losing occupancy at p63-bound enhancers and gaining access to GATA2-bound enhancers.

From these newly acquired sites, GR activated oncogenic transcriptional programs, including EMT and androgen response pathways, leading to diminished anti-proliferative effects and enhanced cell invasiveness. These molecular changes were mirrored in patient survival data, which revealed the best outcomes in tumors with high p63 and low GATA2 expression, and the poorest outcomes in those with high GATA2 and low p63 expression.

Although GR and p63 have not previously been linked in the prostate, their cooperative roles have been documented in other disease contexts. GR and p63 have been shown to co-occupy chromatin in basal-like pancreatic ductal adenocarcinoma cells [48]. In keratinocytes, GR and p63 exhibit substantial overlap in chromatin binding, and physically interact at shared regulatory sites [28]. Notably, GR loss in this context leads to atopic dermatitis, driven by a p63-mediated increase in inflammatory signaling, suggesting that GR counteracts the detrimental effects of p63 in skin. Similarly, in the prostate, our findings indicate that GR and p63 cooperate to suppress tumorigenic processes, highlighting a beneficial partnership in disease attenuation. However, this cooperative effect is likely not universally protective. While p63 restricts EMT in PCa, it has been shown to promote EMT in other malignancies, such as squamous cell carcinoma [26,49,50]. These findings underscore the context- and disease-specific nature of GR–p63 crosstalk, which may yield either protective or pathogenic outcomes depending on cellular and molecular environments.

GATA2 is a well-established oncogenic factor in PCa, known to promote disease progression through both AR-dependent and -independent mechanisms [9,36,37,51]. Most notably, GATA2 enhances AR signaling by modulating AR expression and facilitating its chromatin binding [9,37] and contributes to antiandrogen resistance by regulating GR expression [42]. Our findings extend GATA2's oncogenic role by demonstrating its involvement in reprogramming GR activity following p63 depletion. In p63-null prostate cells, GR's acquired regulation of EMT and androgen response pathways was abrogated upon GATA2 silencing. Furthermore, GATA2 depletion significantly reduced GR-driven cell invasion. These results highlight GATA2 as a critical mediator of GR's oncogenic reprogramming.

Although the interplay between AP-1 and GATA family members has not been extensively characterized in PCa, their cooperative activity at shared enhancers has been documented in other biological contexts. For example, GATA2 and AP-1 can jointly reprogram human fibroblasts into hematopoietic progenitor cells [52], and in breast cancer, GATA3 collaborates with AP-1 to establish functional enhancers [53]. Notably, p63 depletion leads to increased expression of AP-1 subunits, such as *FOSL1*, at levels comparable to GATA2. In p63-null prostate cells, GRBs that are not associated with GATA2 are also frequently linked to AP-1, suggesting that AP-1 may also contribute to GR's oncogenic activity in the absence of p63. Indeed, AP-1 is a well-established mediator of GR action [35] and has been implicated in advanced stages of PCa [6]. The involvement of AP-1 provides a plausible explanation for the persistence of some oncogenic effects of GR in p63-null cells even after GATA2 depletion, highlighting a broader network of TF crosstalk driving GR-mediated tumor progression.

Our data indicates that p63, in cooperation with GR, exerts anti-tumorigenic effects in prostate cells. However, a recent study suggests that p63 can adopt oncogenic role in advanced non-luminal PCa. Specifically, loss of the methyltransferase KMT2C leads to elevated p63 expression, driving the transdifferentiation of luminal adenocarcinoma cells into squamous-like, double-negative PCa that do not express AR or neuroendocrine markers [54]. This aligns with the established pro-tumorigenic role of p63 in squamous cell carcinoma [26]. Additionally, p63 has been shown to restrict transdifferentiation from squamous to neuroendocrine cells in esophageal cancer [55], suggesting that its oncogenic activity is preserved in squamous cell contexts. Intriguingly, the induction of p63 and associated transdifferentiation has been linked to the loss of NF-κB subunit p65–mediated repression of *TP63* [54]. Given that inflammation promotes basal-to-luminal differentiation in the prostate [56], it is plausible that p65 also represses p63 expression in this tissue. AP-1, another inflammation-associated TF, might also play a role in the differentiation process. Since GR is known to exert anti-inflammatory effects by repressing p65 and AP-1 activity [13], glucocorticoids may influence p63-mediated transdifferentiation. These findings suggest that GR–p63 cooperation may function

differently in double-negative PCa cells, a notion supported by the distinct transcriptional activity of GR observed between prostate cells and double-negative PCa cell lines such as DU145 and PC3.

Our findings suggest that TF crosstalk, specifically a p63–GATA2 molecular switch, governs the transformation of GR activity from tumor-suppressive to oncogenic in prostate cells. This mechanism highlights a broader principle of steroid receptor plasticity. A similar, though mechanistically distinct, phenomenon has been observed in breast cancer, where the tumor-suppressive functions of the AR can be reprogrammed into oncogenic activity via the JAK/STAT pathway [21]. Notably, our study demonstrates that such reprogramming of a steroid receptor can occur even in non-cancerous prostate cells. Furthermore, our results point to a potential therapeutic strategy to mitigate the detrimental effects of glucocorticoids in PCa. Given that GATA2 also regulates *NR3C1* expression in antiandrogen-treated PCa cells [42], targeting GATA2 may be an effective strategy to suppress GR-driven oncogenic activity across diverse PCa contexts. Small-molecule GATA2 inhibitors, such as K-7174 [42,57] and the repurposed vasodilator Dilazep [58], offer promising avenues for intervention, although they have yet to be evaluated in clinical trials.

Acknowledgements
We thank Siri Mäki-Ikola, Merja Räsänen, Saara Pirinen, and Kirsi Kainulainen for technical assistance. The EMBL GeneCore team (Heidelberg, Germany) is greatly acknowledged for deep sequencing services. This work was carried out with the support of UEF Single Cell Genomics Core and UEF Cell and Tissue Imaging Unit, University of Eastern Finland, Finland, Biocenter Kuopio and EuroBioimaging Finland. The computational analyses were performed on servers provided by UEF Bioinformatics Center, University of Eastern Finland, Finland. This work was supported by the Research Council of Finland; Cancer Foundation Finland; Sigrid Jusélius Foundation; University of Eastern Finland strategic funding; University of Eastern Finland Doctoral Programme of Molecular Medicine.

Author contributions
Johannes Hiltunen: Conceptualization; Formal Analysis; Investigation; Visualization; Writing – original draft; Writing – review & editing. **Niina Aaltonen**: Formal Analysis; Investigation; Visualization; Writing – review & editing. **Heini Sohlberg**: Investigation; Writing – review & editing. **Laura Kemppi**: Investigation; Writing – review & editing. **Ville Paakinaho**: Conceptualization; Formal Analysis; Funding acquisition; Investigation; Project administration; Supervision; Visualization; Writing – original draft; Writing – review & editing.

Methods
Cell culture and treatments
RWPE1 cells (ATCC, # CRL-3607, RRID:CVCL_3791) were cultured in Keratinocyte SFM (Thermo-Fischer, #17005042) supplemented with BPE (50 µg/ml) and EGF (5 ng/ml) and 1 U/µl penicillin and 1 µg/ml streptomycin (Gibco, #15140122). DU145 cells (DSMZ, #ACC261, RRID:CVCL_0105) were cultured in MEM (Thermo-Fisher, #10370047) supplemented with 10% FBS (Gibco, #11573397), 2 mM l-glutamine (Gibco, #A2916801), and 1 U/µl penicillin and 1 µg/ml streptomycin (Gibco, #15140122). PC3 cells (ATCC, # CRL-1435, RRID:CVCL_0035) were cultured in Ham's F12-K medium (Gibco, #21127022) supplemented with 10% FBS (Gibco, #11573397), 1 U/µl penicillin and 1 µg/ml streptomycin (Gibco,

#15140122). PNT1A cells (Sigma, #95012614, RRID:CVCL_2163) were cultured in RPMI-1640 medium (Gibco, #1187509) supplemented with 10% FBS (Gibco, #11573397), 2 mM l-glutamine (Gibco, #A2916801), and 1 U/µl penicillin and 1 µg/ml streptomycin (Gibco, #15140122). For experiments DU145, PC3, and PNT1A were cultured in experiment medium containing 5% charcoal stripped FBS (Gibco, #10270106), for minimum of 2 days before the start of hormone treatment. Dexamethasone (Dex) (Sigma-Aldrich, #D4902) treatments were conducted with 100 nM concentration. The cells were regularly tested for mycoplasma contamination.

Generation of knockout cell lines using CRISPR-Cas9
Predesigned Alt-R CRISPR-Cas9 crRNA (IDT) (Supplementary Table S1) recognized the DNA binding domain in exon 5 of *TP63* gene. Alt-R CRISPR-Cas9 negative control crRNA (IDT, # 1072544) was used as control. crRNA and tracrRNA (IDT, #1072533) were combined to create each guide RNA (gRNA). *TP63*-specific gRNA was predicted to induce breaks before the key amino acids critical for Zinc Finger, DNA binding, and Intradimer functionality [59]. Alt-R *S.p.* Cas9 Nuclease V3 (IDT, #1081058), was diluted to 1 µM concentration in Cas9 working buffer and was incubated with gRNA oligos to form ribonucleoprotein (RNP) complex. Generation of gRNAs, RNPs and cationic lipid delivery were performed according to manufacturer's instructions (IDT). RNP complexes were reverse-transfected into RWPE1 cells using Opti-MEM (Thermo Fisher, #31985070) and Lipofectamine RNAiMAX Transfection Reagent (Invitrogen, #13778150). Initial gRNAs functionality was tested with Alt-R Genome Editing Detection Kit (IDT, # 1075932) and Agilent Bioanalyzer 2100 DNA 1000 Analysis kit (Agilent, #5067–1504). After 48 h exposure of RWPE1 cells to RNPs, cells were collected and plated in 96-well plate with serial dilutions to obtain monoclonal cultures, with monoclonality verified by microscopy. After the expansion of monoclonal cultures, the absence of p63 was verified with immunoblotting. For verification of genome editing, genomic DNA was isolated using DNA Extraction Solution (Epicentre, #QE09050), and Sanger sequenced (Macrogen Europe) (Supplementary Table S1).

Antibodies
Primary antibodies: anti-GR (Cell Signaling Technology, #12041, RRID:AB_2631286), anti-α tubulin (Santa Cruz Biotechnology, #sc-5286, RRID:AB_628411), anti-p53 (Abcam, #ab1101, RRID: AB_297667), anti-p63α (Cell Signaling Technology, #13109, RRID:AB_2637091), anti-FRA1 (Cell Signaling Technology, #5281, RRID:AB_10557418), anti-GATA2 (Cell Signaling Technology, #79802, RRID:AB_3105938), anti-H3K27ac (Active Motif, #39133, RRID:AB_2561016). Secondary antibodies: goat anti-rabbit (Invitrogen, #G-21234, RRID:AB_1500696), goat anti-mouse (Invitrogen, #62-6520, RRID:AB_2533947).

Immunoblotting
Protein isolation and separation was done as before [60]. Primary antibodies: anti-GR (1:1000), anti-p63α (1:1000), anti-GATA2 (1:1000). Anti-tubulin antibody (1:3000) was used as a loading control. Secondary antibodies (1:10000): anti-mouse (tubulin), anti-rabbit (GR, p63, GATA2). The detection was done using Pierce ECL Western Blotting Substrate kit (Thermo Fisher, #32106) with Bio-Rad ChemiDoc Imager.

RNA extraction and RNA-seq
For RNA extraction, the cells were plated on 6-well plates (PC3, 200k cells per well; DU145, 300k cells per well; RWPE1, 250k cells per well). The cells were treated with vehicle (0.1% EtOH), or 100 nM Dex for 6 h before the start of RNA extraction. For RNA interferences, the cells were reverse siRNA transfected using Opti-MEM (Thermo-Fisher, #31985070) and Lipofectamine RNAiMAX Transfection Reagent (Invitrogen, #13778150) according to manufacturer's instructions. SiRNAs were used at 60 nM concentration: ON-TARGET plus Non-targeting Control SMARTpool (siNON, #D-001810–10), ON-TARGET plus Human GATA2 SMARTPOOL (siGATA2, L-009024-02-0020). Immunoblotting was used to verify siRNA effectiveness as above. RNA was extracted using TriPure Isolation Reagent (Roche, #1166715700), and Monarch Total RNA Miniprep Kit (New England BioLabs, #T2010S). RNA sequencing (RNA-seq) libraries

were generated with NEBNext Poly(A) mRNA Magnetic Isolation Module (New England BioLabs, #E7490) and NEBNext Ultra II Directional RNA Library Prep Kit (New England BioLabs, #E7765) according to manufacturer's instructions using 2 µg of purified RNA. Library quality was measured with Agilent 2100 Bioanalyzer using DNA 1000 analysis kit (Agilent, #5067-1504). Two to three replicates were then sequenced with Illumina NextSeq 500 (75SE) or Illumina NextSeq 2000 (100SE).

Human hg38 genome alignment was done using STAR2.7.9a (RRID:SCR_004463) [61] with default settings allowing max 10 mismatches and max 10 multi-mapped reads. Differential expression was assessed with DESeq2 (RRID:SCR_015687) [62] and HOMER (4.1.1) (RRID:SCR_010881) [63]. For visualization genes were normalized transcript per million (TPM). Differentially expressed genes were defined with log_2 fold change $\pm$ 0.5 and FDR < 0.05 between specific conditions. DESeq2 data and TPM values are presented in Supplementary Table S2. Z-score pathway average was calculated by taking an average from single pathway gene Z-scores. Metascape pathway enrichment analysis (RRID:SCR_016620) [64] was used for comparison of specific gene clusters. Hallmark, CORUM, and WIKI pathway gene sets were used with criteria $p < 0.01$, at least three gene overlap per gene set, and > 1.5 pathway enrichment. For genes downregulated by siGATA2 (siGATA2 DN), log_2 fold change >1 and FDR < 0.05 requirement were utilized. For Dex-regulated siGATA2 DN genes, siGATA2 led to loss of Dex-regulation wherein the target genes no longer met the above set criteria.

ChIP-seq and ATAC-seq
For chromatin immunoprecipitation (ChIP), the cells were plated on 10-cm plates (PC3, 1M cells per plate; DU145, 2M cells per plate; RWPE1, 2M cells per plate; PNT1A, 1.5M cells per plate). Cells were treated with vehicle (0.1% EtOH), or 100 nM Dex for 1 h before the start of ChIP. ChIP experiments were done as previously indicated [65]. Antibodies per IP: GR, 12,5 µl; p63, 3 µg; FRA1, 3 µg; GATA2, 5 µl; H3K27ac, 2 µg. Two to five IP samples were pooled for one sequencing sample. Libraries were made using NEBNext Ultra II DNA Library Prep Kit (New England BioLabs, #E7645L) according to manufacturer's instructions. Library quality control was done with Agilent Bioanalyzer 2100 DNA 1000 Analysis kit (Agilent, #5067–1504) or NGS Fragment Kit (1-6000bp) (Agilent, DNF-473-0500). In general, two replicates were then sequenced with Illumina NextSeq 500 (75SE) or Illumina NextSeq 2000 (100SE).

For assay for transposase-accessible chromatin with sequencing (ATAC-seq), RWPE1 cells were plated on 10-cm plates (2M cells per plate). Cells were treated with vehicle (0.1% EtOH), or 100 nM Dex for 1 h before the start of nuclei isolation. ATAC-seq was performed as previously described [66,67]. Quality analysis of ATAC-seq libraries was done using High Sensitivity DNA Analysis kit (Agilent, #5067-4626) or NGS Fragment Kit (1-6000bp) (Agilent, DNF-473-0500). Two biological replicates were sequenced using Illumina NextSeq 2000 (50PE).

ChIP-seq and ATAC-seq data analysis
ChIP-seq and ATAC-seq alignment to hg38 and read filtering was done with Bowtie2 (RRID:SCR_005476) [68]. HOMER (4.1.1) was used for data analysis [63] with findPeaks used to call peaks over ChIP input sample. Replicate consensus peaks were used with p53 and p63 ChIP-seq data. Differentially bound peaks were discovered using getDifferentialPeaksReplicates and getDifferentialPeaks. For overlapping peaks, mergePeaks using given size of the peaks were utilized. ChIP-seq peaks are presented in Supplementary Table S3. The GR ChIP-seq peaks after p63 ablation were categorized as: p63KO-lost, -gained, and -retained, based on the overlap with p63-positive control. GR peaks in Control, not present in either p63KO line were classified as p63KO-lost. GR peaks gained after p63KO, shared between p63KO lines were classified as p63KO-gained. GR peaks present in both Control and p63KO cells were classified as p63KO-retained sites. Downstream analyses were normalized to 10 million mapped reads. Heatmaps were generated with 10 or 20 bp bins with $\pm$ 0.5 kb or 1 kb surrounding the peak. Tag counts in box plots were normalized using log_2 or variance stabilized and transformed (rlog) options. Box plots were drawn with Tukey method. Motif enrichment was measured using findMotifsGenome.pl on default settings. Motif data is presented in Supplementary Table S4. Motif histograms were normalized to motif per bp per site and displayed $\pm$ 100-250

bp from peak center. H3K27ac histograms were normalized to tags per bp per site and displayed ± 1 kb from peak center. Motif enrichment $\log_2$ fold difference was calculated based on the relation of percentage target sequences with motif to the percentage of background sequences with motif. For the comparison of p63 motif and half GRE motif overlap, motif distances were called with HOMER annotatePeaks.pl and the motifs filtered based on the relative distance. The motifs were then aligned by sequence overlap. Half GRE motifs located inside p63 motifs were classified as complete overlap, while the rest of the motifs containing an overlap of two base pairs or more were classified as partial overlap. For pathway analysis, peaks were first associated with target genes using annotatePeaks.pl followed by Metascape pathway analysis like with RNA-seq data. For the prediction of transcriptional regulators from peaks, epigenetic landscape in silico deletion analysis (LISA2) [39] from Coverage command was used with default settings and hg38 genome. Gene output from annotatePeaks.pl was utilized as the gene list, and H3K27ac BigWig files as coverage over genome.

scRNA-seq
For single-cell RNA-seq (scRNAseq), RWPE1 cells were plated on 10-cm plates (2M cells per plate). The cells were treated with vehicle (0.1% EtOH) or 100 nM Dex for 6 hours before the start of extraction. Single cell suspensions were isolated according to published protocol and diluted to 700 cells per µl (10xGenomics, #CG00054). Library generation was performed using Single Cell 3' Kit v3.1 (10x Genomics, #PN-1000269, #PN-1000127, #PN-1000215) according to protocol (10x Genomics, #CG000315) with target cell recovery set to 2000 cells. Library quality was done using Agilent 2100 Bioanalyzer using High Sensitivity DNA Analysis kit (Agilent, #5067-4626), and libraries were sequenced with Illumina NextSeq 2000 (100PE). The scRNA-seq data analyses were done as previously indicated [69]. In total 4058 single cells were obtained from sequencing of EtOH (2169 cells) and Dex (1889 cells) samples. Cells with low (<750) or high (>7500) feature counts, and cells with high mitochondrial reads (>20 %) and low ribosomal reads (<5 %) were filtered out. After filtration 1468 EtOH- and 1592 Dex-treated single cells were qualified for downstream analyses. Downstream analysis was performed using Seurat 4.3.0 (RRID:SCR_007322) (Hao et al. 2021). Violin plots represent the average of the normalized expression values. Differential expression was determined with default algorithm; $\log_2$ fold change < 0.10, p-value < 0.05, min.pct =0.25). Cells with non-zero expression values were counted as expressing a transcript.

Confocal imaging
For confocal imaging, the cells were plated on 8-well chamber slides (Ibidi GmbH, #80826) (10k cells per well). The cells were fixed with 4% paraformaldehyde (PFA) for 20 min and washed with 0.1 M phosphate buffer. The fixed cells were permeabilized for 15 minutes with 0.1% Triton X-100 in 1% BSA and stained with Phalloidin-iFluor 594 reagent (Abcam, #ab176757) for 20 min at room temperature. Nuclei were labeled with 4',6-diamidino-2-phenylindole (DAPI) (Sigma-Aldrich, #D8417). The fluorescent images were obtained with LSM800 confocal microscope (Carl Zeiss Microscopy GmbH) using 40x objective. Image processing was performed using ZEN 3.10 software (Carl Zeiss Microscopy GmbH). For determining the cell length-to-width ratio (aspect ratio), the cell length and width in pixels were first measured with ImageJ software (National Institutes of Health, RRID:SCR_003070) and the aspect ratio was calculated by dividing the length with width (n=37-41).

Cell proliferation assay
For automated live cell proliferation experiments, the cells were seeded onto 96-well plate (10k cells per well). At the start of the experiment (day 0), the cells were treated with vehicle (0.1% EtOH) or 100 nM Dex and the nuclei were stained with Incucyte Nuclight Rapid Red reagent (Essen BioSciences, #4717). The cells were monitored every day for 5 days using Incucyte S3 Live-Cell Imaging System (Sartorius). Incucyte S3 2024A software (Sartorius) was used to count the number of cells. Results were normalized to day 0 and are shown as relative percentage in cell count, representing the mean ±SD of 8-10 images per day. Box plots were drawn with Tukey method. For RNA interference, siRNA treatments were conducted as described

above before plating the cells. Statistical significance was determined by Two-ANOVA with Bonferroni *post hoc* test.

Migration assay

For automated live cell migration experiments, the cells were seeded onto 96-well plate (Sartorius Imagelock, #BA-04855) (70k cells per well). The following day, scratch wounds were generated with Incucyte Woundmaker Tool (Sartorius) and cells were washed twice with PBS. Subsequently, the cells were treated with vehicle (0.1% EtOH) or 100 nM Dex and were monitored every 3 h for total of 48 h using Incucyte S3 Live-Cell Imaging System (Sartorius). Incucyte S3 2024A software (Sartorius) was used to analyze wound healing. Results are shown as relative wound density, representing the mean ±SD of 4-5 images per time point. Box plots were drawn with Tukey method. Statistical significance was determined by Two-ANOVA with Bonferroni *post hoc* test.

Invasion assay

For 3D invasion, the cells were plated onto 8-well chamber slides (Ibidi GmbH, #80826) (250k to 350k cells per well). The following day, Cultrex Basement Membrane Extract (R&D systems, #3432) and Type I collagen (R&D Systems, #3440) (1:1) gel mixture was added on each well and allowed to polymerize at 37°C for 2 h. Subsequently, media containing either vehicle (0.1% EtOH) or 100 nM Dex were added on the wells and cells were allowed to invade the gel for 6 days. Fresh media was added to the slides every 48 h. The wells were fixed with 4% PFA for 2 h at room temperature and washed with 0.1 M phosphate buffer. The wells were permeabilized with 0.1% Triton-X100 in 1% BSA for 30 min and stained with Phalloidin-iFluor 594 reagent (Abcam, #ab176757) for 1 h at room temperature. Nuclei were labeled with DAPI (Sigma-Aldrich, #D8417). Cell invasion was imaged with LSM800 confocal microscope (Carl Zeiss Microscopy GmbH) using 20x objective by taking 1 µm z-intervals. Images were processed and invasion distances measured with ZEN 3.10 software (Carl Zeiss Microscopy GmbH). Box plots were drawn with Tukey method. For RNA interference, siRNA treatments were conducted as described above before plating the cells. Statistical significance was determined by One-way ANOVA with Bonferroni *post hoc* test.

Public datasets and patient data analysis

The following publicly available sequencing datasets were used: GSE214757 for GR ChIP-seq and RNA-seq from VCaP and 22Rv1 cells [10], GSE204813 for RNA-seq from p63 overexpressed PC3 cells [29]. The data was processed as indicated above. PCa patient, survival, and single-cell expression data were obtained through Combined Transcriptome dataset of PCa Cell lines (CTPC) [22], Prostate cancer atlas [70], Xena (RRID:SCR_018938) [71], cBioPortal (RRID:SCR_014555) [72], Single Cell portal (RRID:SCR_014816) [73]. For survival, the patients were divided based on either median or top 25% and bottom 25% quartiles. Statistical significance was calculated with log-rank test for survival.

References

1. Bray F, Laversanne M, Sung H, Ferlay J, Siegel RL, Soerjomataram I, et al. Global cancer statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. CA Cancer J Clin. 2024;74(3):229–63.

2. Wang G, Zhao D, Spring DJ, DePinho RA. Genetics and biology of prostate cancer. Genes Dev. 2018 Sep 1;32(17–18):1105–40.

3. Feng Q, He B. Androgen Receptor Signaling in the Development of Castration-Resistant Prostate Cancer. Front Oncol. 2019;9:858.

4. Paakinaho V, Palvimo JJ. Genome-wide crosstalk between steroid receptors in breast and prostate cancers. Endocr Relat Cancer. 2021 Sep 1;28(9):R231–50.

5.   Burris TP, de Vera IMS, Cote I, Flaveny CA, Wanninayake US, Chatterjee A, et al. International Union of Basic and Clinical Pharmacology CXIII: Nuclear Receptor Superfamily-Update 2023. Pharmacol Rev. 2023 Nov;75(6):1233–318.

6.   Tang F, Xu D, Wang S, Wong CK, Martinez-Fundichely A, Lee CJ, et al. Chromatin profiles classify castration-resistant prostate cancers suggesting therapeutic targets. Science. 2022 May 27;376(6596):eabe1505.

7.   Mu P, Zhang Z, Benelli M, Karthaus WR, Hoover E, Chen CC, et al. SOX2 promotes lineage plasticity and antiandrogen resistance in TP53- and RB1-deficient prostate cancer. Science. 2017 Jan 6;355(6320):84–8.

8.   Linder S, Hoogstraat M, Stelloo S, Eickhoff N, Schuurman K, de Barros H, et al. Drug-Induced Epigenomic Plasticity Reprograms Circadian Rhythm Regulation to Drive Prostate Cancer toward Androgen Independence. Cancer Discov. 2022 Sep 2;12(9):2074–97.

9.   Zhou T, Yu C, Han Y, He B, Feng Q. GATA2 up-regulation restores androgen receptor chromatin association and advances darolutamide resistance in prostate cancer. Genes Dis. 2025 Jul;12(4):101508.

10. Helminen L, Huttunen J, Tulonen M, Aaltonen N, Niskanen EA, Palvimo JJ, et al. Chromatin accessibility and pioneer factor FOXA1 restrict glucocorticoid receptor action in prostate cancer. Nucleic Acids Res. 2024 Jan 25;52(2):625–42.

11. Williams A, Gutgesell L, de Wet L, Selman P, Dey A, Avineni M, et al. SOX2 expression in prostate cancer drives resistance to nuclear hormone receptor signaling inhibition through the WEE1/CDK1 signaling axis. Cancer Lett. 2023 Jul 1;565:216209.

12. Kalfeist L, Galland L, Ledys F, Ghiringhelli F, Limagne E, Ladoire S. Impact of Glucocorticoid Use in Oncology in the Immunotherapy Era. Cells. 2022 Feb 22;11(5):770.

13. Hiltunen J, Helminen L, Paakinaho V. Glucocorticoid receptor action in prostate cancer: the role of transcription factor crosstalk. Front Endocrinol. 2024 Jul 4;15:1437179.

14. Arora VK, Schenkein E, Murali R, Subudhi SK, Wongvipat J, Balbas MD, et al. Glucocorticoid receptor confers resistance to antiandrogens by bypassing androgen receptor blockade. Cell. 2013 Dec 5;155(6):1309–22.

15. Puhr M, Hoefer J, Eigentler A, Ploner C, Handle F, Schaefer G, et al. The Glucocorticoid Receptor Is a Key Player for Prostate Cancer Cell Survival and a Target for Improved Antiandrogen Therapy. Clin Cancer Res Off J Am Assoc Cancer Res. 2018 Feb 15;24(4):927–38.

16. Yemelyanov A, Czwornog J, Chebotaev D, Karseladze A, Kulevitch E, Yang X, et al. Tumor suppressor activity of glucocorticoid receptor in the prostate. Oncogene. 2007 Mar 22;26(13):1885–96.

17. Rogerson C, Britton E, Withey S, Hanley N, Ang YS, Sharrocks AD. Identification of a primitive intestinal transcription factor network shared between esophageal adenocarcinoma and its precancerous precursor state. Genome Res. 2019 May;29(5):723–36.

18. Parreno V, Loubiere V, Schuettengruber B, Fritsch L, Rawal CC, Erokhin M, et al. Transient loss of Polycomb components induces an epigenetic cancer fate. Nature. 2024 May;629(8012):688–96.

19. Hickey TE, Selth LA, Chia KM, Laven-Law G, Milioli HH, Roden D, et al. The androgen receptor is a tumor suppressor in estrogen receptor-positive breast cancer. Nat Med. 2021 Feb;27(2):310–20.

20. Prekovic S, Chalkiadakis T, Roest M, Roden D, Lutz C, Schuurman K, et al. Luminal breast cancer identity is determined by loss of glucocorticoid receptor activity. EMBO Mol Med. 2023 Dec 7;15(12):e17737.

21. Asemota S, Effah W, Holt J, Johnson D, Cripe L, Ponnusamy S, et al. A molecular switch from tumor suppressor to oncogene in ER+ve breast cancer: Role of androgen receptor, JAK-STAT, and lineage plasticity. Proc Natl Acad Sci U S A. 2024 Oct;121(40):e2406837121.

22. Cheng S, Yu X. CTPC, a combined transcriptome data set of human prostate cancer cell lines. The Prostate. 2023;83(2):158–61.

23. Bello D, Webber MM, Kleinman HK, Wartinger DD, Rhim JS. Androgen responsive adult human prostatic epithelial cell lines immortalized by human papillomavirus 18. Carcinogenesis. 1997 Jun;18(6):1215–23.

24. Degeorges A, Hoffschir F, Cussenot O, Gauville C, Le Duc A, Dutrillaux B, et al. Recurrent cytogenetic alterations of prostate carcinoma and amplification of c-myc or epidermal growth factor receptor in subclones of immortalized PNT1 human prostate epithelial cell line. Int J Cancer. 1995 Sep 15;62(6):724–31.

25. Pflaum J, Schlosser S, Müller M. p53 Family and Cellular Stress Responses in Cancer. Front Oncol [Internet]. 2014 Oct 21 [cited 2025 Jan 23];4. Available from: https://www.frontiersin.org/journals/oncology/articles/10.3389/fonc.2014.00285/full

26. Li Y, Giovannini S, Wang T, Fang J, Li P, Shao C, et al. p63: a crucial player in epithelial stemness regulation. Oncogene. 2023 Nov;42(46):3371–84.

27. Tsvetkov P, Reuven N, Shaul Y. Ubiquitin-independent p53 proteasomal degradation. Cell Death Differ. 2010 Jan;17(1):103–8.

28. Sevilla LM, Pons-Alonso O, Gallego A, Azkargorta M, Elortza F, Pérez P. Glucocorticoid receptor controls atopic dermatitis inflammation via functional interactions with P63 and autocrine signaling in epidermal keratinocytes. Cell Death Dis. 2024 Jul 28;15(7):535.

29. Sultanov R, Mulyukina A, Zubkova O, Fedoseeva A, Bogomazova A, Klimina K, et al. TP63–TRIM29 axis regulates enhancer methylation and chromosomal instability in prostate cancer. Epigenetics Chromatin. 2024 Mar 14;17:6.

30. Brawer MK, Peehl DM, Stamey TA, Bostwick DG. Keratin immunoreactivity in the benign and neoplastic human prostate. Cancer Res. 1985 Aug;45(8):3663–7.

31. Bergholz J, Xiao ZX. Role of p63 in Development, Tumorigenesis and Cancer Progression. Cancer Microenviron. 2012 Jul 31;5(3):311.

32. Olsen JR, Oyan AM, Rostad K, Hellem MR, Liu J, Li L, et al. p63 attenuates epithelial to mesenchymal potential in an experimental prostate cell model. PloS One. 2013;8(5):e62547.

33. Tucci P, Agostini M, Grespi F, Markert EK, Terrinoni A, Vousden KH, et al. Loss of p63 and its microRNA-205 target results in enhanced cell migration and metastasis in prostate cancer. Proc Natl Acad Sci. 2012 Sep 18;109(38):15312–7.

34. Kojima Y, Yoneyama T, Hatakeyama S, Mikami J, Sato T, Mori K, et al. Detection of Core2 β-1,6-N-Acetylglucosaminyltransferase in Post-Digital Rectal Examination Urine Is a Reliable Indicator for Extracapsular Extension of Prostate Cancer. PLoS ONE. 2015 Sep 21;10(9):e0138520.

35. Biddie SC, John S, Sabo PJ, Thurman RE, Johnson TA, Schiltz RL, et al. Transcription factor AP1 potentiates chromatin accessibility and glucocorticoid receptor binding. Mol Cell. 2011 Jul 8;43(1):145–55.

36. Rodriguez-Bravo V, Carceles-Cordon M, Hoshida Y, Cordon-Cardo C, Galsky MD, Domingo-Domenech J. The role of GATA2 in lethal prostate cancer aggressiveness. Nat Rev Urol. 2017 Jan;14(1):38–48.

37. Wu D, Sunkel B, Chen Z, Liu X, Ye Z, Li Q, et al. Three-tiered role of the pioneer factor GATA2 in promoting androgen-dependent gene expression in prostate cancer. Nucleic Acids Res. 2014 Apr 1;42(6):3607–22.

38. Yu X, Singh PK, Tabrejee S, Sinha S, Buck MJ. ΔNp63 is a pioneer factor that binds inaccessible chromatin and elicits chromatin remodeling. Epigenetics Chromatin. 2021 Apr 17;14(1):20.

39. Qin Q, Fan J, Zheng R, Wan C, Mei S, Wu Q, et al. Lisa: inferring transcriptional regulators through integrative modeling of public chromatin accessibility and ChIP-seq data. Genome Biol. 2020 Feb 7;21(1):32.

40. Pofi R, Caratti G, Ray DW, Tomlinson JW. Treating the Side Effects of Exogenous Glucocorticoids; Can We Separate the Good From the Bad? Endocr Rev. 2023 Nov 9;44(6):975–1011.

41. Huttunen J, Aaltonen N, Helminen L, Rilla K, Paakinaho V. EP300/CREBBP acetyltransferase inhibition limits steroid receptor and FOXA1 signaling in prostate cancer cells. Cell Mol Life Sci CMLS. 2024 Apr 2;81(1):160.

42. Yuan F, Hankey W, Wu D, Wang H, Somarelli J, Armstrong AJ, et al. Molecular determinants for enzalutamide-induced transcription in prostate cancer. Nucleic Acids Res. 2019 Nov 4;47(19):10104–14.

43. Karthaus WR, Hofree M, Choi D, Linton EL, Turkekul M, Bejnood A, et al. Regenerative potential of prostate luminal cells revealed by single-cell analysis. Science. 2020 May 1;368(6490):497–505.

44. He MX, Cuoco MS, Crowdis J, Bosma-Moody A, Zhang Z, Bi K, et al. Transcriptional mediators of treatment resistance in lethal prostate cancer. Nat Med. 2021 Mar;27(3):426–33.

45. Wang ZA, Toivanen R, Bergren SK, Chambon P, Shen MM. Luminal cells are favored as the cell of origin for prostate cancer. Cell Rep. 2014 Sep 11;8(5):1339–46.

46. Pitzen SP, Dehm SM. Basal epithelial cells in prostate development, tumorigenesis, and cancer progression. Cell Cycle. 2023;22(11):1303–18.

47. Tucci P, Agostini M, Grespi F, Markert EK, Terrinoni A, Vousden KH, et al. Loss of p63 and its microRNA-205 target results in enhanced cell migration and metastasis in prostate cancer. Proc Natl Acad Sci. 2012 Sep 18;109(38):15312–7.

48. Ekstrom TL, Rosok RM, Abdelrahman AM, Parassiadis C, Manjunath M, Dittrich MY, et al. Glucocorticoid receptor suppresses GATA6-mediated RNA polymerase II pause release to modulate classical subtype identity in pancreatic cancer. Gut. 2025 Jan 30;gutjnl-2024-334374.

49. Dang TT, Esparza MA, Maine EA, Westcott JM, Pearson GW. ΔNp63α Promotes Breast Cancer Cell Motility through the Selective Activation of Components of the Epithelial-to-Mesenchymal Transition Program. Cancer Res. 2015 Sep 14;75(18):3925–35.

50. Lambert AW, Fiore C, Chutake Y, Verhaar ER, Strasser PC, Chen MW, et al. ΔNp63/p73 drive metastatic colonization by controlling a regenerative epithelial stem cell program in quasi-mesenchymal cancer stem cells. Dev Cell. 2022 Dec 19;57(24):2714-2730.e8.

51. Jin Z, Wang H, Tang R, Pan B, Lee HJ, Liu S, et al. GATA2 promotes castration-resistant prostate cancer development by suppressing IFN-β axis-mediated antitumor immunity. Oncogene. 2024 Aug;43(34):2595–610.

52. Gomes AM, Kurochkin I, Chang B, Daniel M, Law K, Satija N, et al. Cooperative Transcription Factor Induction Mediates Hemogenic Reprogramming. Cell Rep. 2018 Dec 4;25(10):2821-2835.e7.

53. Orlando KA, Grimm SA, Wade PA. Pioneering new enhancers by GATA3: role of facilitating transcription factors and chromatin remodeling. Nucleic Acids Res. 2025 Jun 6;53(11):gkaf473.

54. Guo J, Li N, Liu Q, Hao Z, Zhu G, Wang X, et al. KMT2C deficiency drives transdifferentiation of double-negative prostate cancer and confer resistance to AR-targeted therapy. Cancer Cell [Internet]. 2025 Apr 24 [cited 2025 Apr 25];0(0). Available from: https://www.cell.com/cancer-cell/abstract/S1535-6108(25)00139-4

55. Zhang Y, Karagiannis D, Liu H, Lin M, Fang Y, Jiang M, et al. Epigenetic regulation of p63 blocks squamous-to-neuroendocrine transdifferentiation in esophageal development and malignancy. Sci Adv. 2024 Oct 11;10(41):eadq0479.

56. Kwon OJ, Zhang L, Ittmann MM, Xin L. Prostatic inflammation enhances basal-to-luminal differentiation and accelerates initiation of prostate cancer with a basal cell origin. Proc Natl Acad Sci U S A. 2014 Feb 4;111(5):E592–600.

57. Umetani M, Nakao H, Doi T, Iwasaki A, Ohtaka M, Nagoya T, et al. A novel cell adhesion inhibitor, K-7174, reduces the endothelial VCAM-1 induction by inflammatory cytokines, acting through the regulation of GATA. Biochem Biophys Res Commun. 2000 Jun 7;272(2):370–4.

58. Kaochar S, Rusin A, Foley C, Rajapakshe K, Robertson M, Skapura D, et al. Inhibition of GATA2 in prostate cancer by a clinically available small molecule. Endocr Relat Cancer. 2021 Oct 12;29(1):15–31.

59. Chen C, Gorlatova N, Kelman Z, Herzberg O. Structures of p63 DNA binding domain in complexes with half-site and with spacer-containing full response elements. Proc Natl Acad Sci. 2011 Apr 19;108(16):6456–61.

60. Launonen KM, Paakinaho V, Sigismondo G, Malinen M, Sironen R, Hartikainen JM, et al. Chromatin-directed proteomics-identified network of endogenous androgen receptor in prostate cancer cells. Oncogene. 2021 Jul;40(27):4567–79.

61. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-seq aligner. Bioinforma Oxf Engl. 2013 Jan 1;29(1):15–21.

62. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol. 2014;15(12):550.

63. Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, et al. Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. Mol Cell. 2010 May 28;38(4):576–89.

64. Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, et al. Metascape provides a biologist-oriented resource for the analysis of systems-level datasets. Nat Commun. 2019 Apr 3;10(1):1523.

65. Paakinaho V, Kaikkonen S, Makkonen H, Benes V, Palvimo JJ. SUMOylation regulates the chromatin occupancy and anti-proliferative gene programs of glucocorticoid receptor. Nucleic Acids Res. 2014 Feb;42(3):1575–92.

66. Buenrostro JD, Wu B, Chang HY, Greenleaf WJ. ATAC-seq: A Method for Assaying Chromatin Accessibility Genome-Wide. Curr Protoc Mol Biol. 2015 Jan 5;109:21.29.1-21.29.9.

67. Paakinaho V, Lempiäinen JK, Sigismondo G, Niskanen EA, Malinen M, Jääskeläinen T, et al. SUMOylation regulates the protein network and chromatin accessibility at glucocorticoid receptor-binding sites. Nucleic Acids Res. 2021 Feb 26;49(4):1951–71.

68. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods. 2012 Mar 4;9(4):357–9.

69. Hiltunen J, Helminen L, Aaltonen N, Launonen KM, Laakso H, Malinen M, et al. Androgen receptor-mediated assisted loading of the glucocorticoid receptor modulates transcriptional responses in prostate cancer cells. Genome Res. 2025 Jun 2;gr.280224.124.

70. Bolis M, Bossi D, Vallerga A, Ceserani V, Cavalli M, Impellizzieri D, et al. Dynamic prostate cancer transcriptome analysis delineates the trajectory to disease progression. Nat Commun. 2021 Dec 2;12(1):7033.

71. Goldman MJ, Craft B, Hastie M, Repečka K, McDade F, Kamath A, et al. Visualizing and interpreting cancer genomics data via the Xena platform. Nat Biotechnol. 2020 Jun;38(6):675–8.

72. Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. Cancer Discov. 2012 May;2(5):401–4.

73. Tarhan L, Bistline J, Chang J, Galloway B, Hanna E, Weitz E. Single Cell Portal: an interactive home for single-cell genomics data [Internet]. bioRxiv; 2023 [cited 2025 Jun 17]. p. 2023.07.13.548886. Available from: https://www.biorxiv.org/content/10.1101/2023.07.13.548886v1