Adaptive evolution of odorant receptors is associated with elaborations of social organization in ants

Yoann Pellen[1], Joel Vizueta[2], Lukas Schrader[3], Eyal Privman[1#]

[1] Institute of Evolution, Department of Evolutionary and Environmental Biology, University of Haifa, Israel
[2] Section for Ecology and Evolution, Department of Biology, University of Copenhagen, DK-2100, Copenhagen, Denmark
[3] Institute for Evolution & Biodiversity, University of Münster, Germany
[#] Corresponding author: eprivman@univ.haifa.ac.il

Abstract

Cooperation in social insect colonies depends on a complex vocabulary of chemical signals, which requires a correspondingly complex array of chemosensory receptors. In the genomes of ants, the odorant receptor (OR) gene family is dramatically expanded compared to other insects. This is most notable in the "9-exon" OR subfamily, which was implicated in responding to cuticular hydrocarbons, a major class of signalling compounds. The compilation of a large, high-quality ant genome dataset by the Global Ant Genomics Alliance allows investigating ant OR evolution in unprecedented detail. By analysing 55,068 ORs of 163 representative ant species, we tested for association between the development of sociobiological traits and adaptive evolution of ORs. We provide a detailed analysis of gene family expansions and adaptive sequence evolution at the OR subfamily level, allowing focused analysis of distinct ant lineages. We identified strong enrichment of positive selection on 9-exon ORs in the ancestor of the formicoid clade, the largest clade comprising 90% of the extant ants. Formicoid species are characterized by greater colony size and reproductive division of labour. Thus, our results support a key role of adaptive diversification of chemical communication systems for the evolution of more complex elaboration of social organization in ants. We also observed enrichment of positive selection on 9-exon ORs associated with the more recent evolution of continuous worker polymorphism in multiple lineages. Surprisingly, the recent evolution of other sociobiological traits was associated with depletion of positive selection on ORs. These results suggest that the evolution of worker polymorphism involves more extensive adaptation of chemical communication compared to other aspects of ant sociobiology. Our analysis of an unprecedented ant OR dataset offers new insights into the specific context in which ORs played a major role in the evolution of elaborations in social traits across the diverse phylogeny of ants.

Introduction

Chemical communication is a vital function of insect societies, mediating many aspects of colony organisation. The perception of chemical signals relies on a large repertoire of odorant receptors (ORs), as well as other smaller chemosensory receptor families (Bargmann 2006; Bear et al. 2016; Robertson 2019). Colonies of social insects such as ants are characterised by elaborate cooperation among nestmates. Communication is central to the efficient coordination of necessary activities, including foraging for food, caring for the queen and brood, nest construction and defence. One of the fundamental functions of chemical signalling is nestmate recognition. Cuticular hydrocarbons (CHCs) are a diverse group of long-chain hydrocarbons thought to be a major component of the chemical cues used for recognition, including sexual communication and colony discrimination, which requires classifying individuals based on their CHC profile (van Zweden and d'Ettorre 2010; Blomquist et al. 2020).

ORs are critical for social communications in ants (Trible et al. 2017). In particular, the 9-exon ("9e") subfamily of ORs was suggested to harbour many of the receptors that respond specifically to the diverse array of CHCs displayed by ants (Engsontia et al. 2015; Zhou et al. 2015; McKenzie et al. 2016; Pask et al. 2017; Slone et al. 2017). This clade of the OR gene tree is dramatically expanded in ants, accounting for about a third of all ORs (C.D. Smith et al. 2011; C.R. Smith et al. 2011; Zhou et al. 2012; Zhou et al. 2015; McKenzie et al. 2016). Pask et al. (2017) characterized 22 ORs from the 9e clade and showed that they are narrowly tuned to respond to specific CHCs. Most of the 9e ORs display worker-biased expression, in line with a role in social coordination in the colony (Zhou et al. 2012). These ORs are also mainly expressed in the ventral half of the antennal club that comes in contact with other ants when they antennate each other for nestmate recognition, in line with these being receptors of non-volatile contact pheromones such as CHCs (McKenzie et al. 2016). In social insects, discrimination against non-colony members is especially important to avoid any form of colony exploitation. Therefore, strong selection on these discrimination abilities is expected.

The transition from solitary to social organisation in ants was accompanied by a dramatic expansion of the OR gene family (C.D. Smith et al. 2011; Zhou et al. 2012). This may be limited to the ants, as recent work found no evidence of a general relationship between the evolution of eusociality in Hymenoptera and the expansion of the OR repertoire (Gautam et al. 2024). Ant genomes harbour hundreds of ORs, and up to 503 ORs reported for the clonal raider ant *Ooceraea biroi* (McKenzie and Kronauer 2018). Studies have also shown that the reductive social behaviour in social parasite species (e.g. *Formica selysi*, *Acromyrmex ameliae*, *Lasius cf. spathepus*, *Tetramorium atratulum*) was linked to a loss of OR genes (Schrader et al. 2021; Jongepier et al. 2022; Vizueta et al. 2025). Many ant ORs show evidence of positive selection on their protein sequence (Engsontia et al. 2015; Zhou et al. 2015; Saad et al. 2018) making them attractive candidate genes for the study of adaptive evolution of chemical communication in ants. In total, Saad et al. (2018) reported positive selection in 7.2% of the tested OR gene tree branches (252 out of 3507). Recent species-specific OR gene duplications were followed by a higher rate of positive selection compared to ORs that were not recently duplicated (Engsontia et al. 2015; Saad et al. 2018), as expected for adaptive evolution in the birth-

84 and-death model of gene family evolution (Ohno 1970; Demuth and Hahn 2009; Innan
85 and Kondrashov 2010; Mendivil Ramos and Ferrier 2012).
86
The study of ant ORs greatly benefits from the recent high-quality genomic dataset
produced by the Global Ant Genomics Alliance (GAGA; Vizueta et al., 2025). GAGA
sequenced genomes of 145 species of ants, which, in addition to 18 published high-
quality genomes, represent 27 tribes in 12 subfamilies. Most of these genomes were
produced using long-read sequencing that resulted in highly contiguous genome
assemblies. This is especially important for accurate and complete assembly and
annotation of the large tandem arrays of OR genes. Previous studies based on short-read
genome assemblies revealed only a partial picture, typically underestimating the number
of ORs. For example, the first genome assembly of *Camponotus floridanus* that was
based on Illumina short-read sequencing allowed the annotation of 352 complete OR
genes and 55 partial genes (Zhou et al. 2012), while the more recent PacBio long-read
assembly (Shields et al. 2018) resulted in annotations of 466 complete ORs and 61 partial
genes. The unprecedented genomic dataset produced by GAGA allows us to study the
dynamic evolution of the OR gene family in ants with greater precision.
101
The genome-wide analysis of single-copy gene families in the GAGA genomes (Vizueta
et al. 2025) revealed a high rate of positive selection on protein sequences in the branch
leading to the ancestor of formicoid ants, a clade containing most extant ant species,
which split from the poneroid ants ca. 125-145 Mya. The formicoid clade exhibits the most
elaborated forms of reproductive division of labour, large colony size (Dornhaus et al.
2012), worker polymorphism (Oster and Wilson 1978), and extended queen longevity
(Keller and Genoud 1997). This enrichment of positive selection in this branch is
consistent with previous work published by Romiguier et al. (2022) which identified this
branch as a hotspot for genes under positive selection. The GAGA analyses also revealed
expansions in multi-gene families in the ancestor of ants. Many of these expanded gene
families were implicated in chemoperception (22 families) and biosynthesis of CHCs (17
families). These results are in line with the hypothesis that the evolution of sociality in the
ant ancestor involved the evolution of sophisticated chemical communication.
115
Here, we focus on the evolutionary dynamics of ant ORs and further explore whether
positive selection on ORs is linked to the evolution of traits such as diet, colony size,
worker polymorphism, and worker reproduction. We tested hypotheses using sets of
representative species across the entire phylogeny, focusing on branches with inferred
trait shifts. We identified enrichment of positive selection on the 9e ORs subfamily in the
formicoid ancestor. Surprisingly, more recent branches with a transition to bigger colony
sizes were depleted for positive selection, and that depletion was mostly in the non 9e
OR subfamilies. We generally observed more depletion than enrichment in the recent
branches we tested. However, positive selection on 9e ORs was enriched in branches of
lineages that evolved continuous worker polymorphism. Our analyses leverage the
unprecedented large-scale data from the GAGA project to reveal a complex pattern of
adaptive evolution of ant ORs associated with some, but not all, elaborations of
sociobiology in the ant phylogeny.

Materials and methods

Traits of interest

The GAGA project included the compilation of a trait table with information on social and ecological traits for each species from literature and experts, including for example their habitat, diet, foraging stratum, colony size, worker polymorphism, etc. We focused on traits that we hypothesised to be related to ORs evolution: colony size, diet, worker polymorphism, and worker reproductive functions. These traits require sensory function such as recognition and discrimination, whether of nestmates or of food sources, as well as complex social interactions that require complex signalling. Therefore, ORs can be expected to play a major role in the evolution and elaboration of these traits. Ancestral state reconstruction for each trait across the ant phylogeny was conducted using the R package *ape* (version 5.7-1; Paradis and Schliep, 2019) evaluating the best fitting model by using the Akaike information criterion (AIC). Further details can be found in Vizueta et al. (2025). Trait reconstructions are plotted on the phylogeny in Supplementary Figures S1 to S4. We used these reconstructions to identify branches with inferred trait changes, which we refer to as "transition branches" (marked on Figure 1). We hypothesized that there was positive selection on ORs related to the evolution of the transitioning traits in these branches. For example, we identified branches with a transition from an omnivorous to a herbivorous diet (e.g. in the ancestor of *Messor* harvester ants) and branches where colony size increased (e.g. in the ancestor of *Lasius* and *Myrmecocystus*).

The number of species analysed here coupled with the large number of ORs annotated in each of them resulted in a dataset of 55,068 OR genes. Tests for positive selection using dN/dS models are not feasible for such a large gene tree. Therefore, we focused the positive selection analyses on nine clades with small subsets of species (four to nine species in each; 75 species in total), selected to include the transition branches to be tested. We also tested the three most ancestral branches of the formicoid clade (rectangles in Figure 1) using a selection of ten representative species across the entire ant phylogeny. The first of these branches is the ancestor of all formicoid ants, the second is the branch after the split of the *Dorylinae* subfamily, and the third is the ancestor of *Formicinae* and *Myrmicinae* after the split of the *Myrmeciinae*, *Pseudomyrmicinae*, *Aneuretinae* and *Dolichoderinae* subfamilies.

Genome sequences

Most of the species in the GAGA dataset have been sequenced using PacBio long-read sequencing (125 out of 163 species), which is advantageous relative to Illumina short-read sequencing for the assembly and annotation of repetitive regions such as the typically large arrays of tandem-duplicated OR genes. Twenty species, including some rare species like *Leptanilla*, had limited sample biomass and could only be sequenced with stLFR (single-tube long fragment read), a sequencing technique that requires fewer DNA amount than PacBio. Scaffolding of contigs into whole chromosomes was achieved for 12 species using HiC, a technique that captures genome-wide chromatin interactions (Belton et al., 2012). The published genomes of 18 additional species were retrieved from NCBI, giving a total of 163 genomes. Among the 75 genomes used in our analyses, 65

were sequenced by PacBio and seven sequenced by stLFR. Three of the non-GAGA
genomes were based on 454 sequencing, which is an obsolete short-read technology
that is expected to result in more partial and missing ORs. Further details about the
genome sequencing and assembly can be found in Vizueta et al. (2025).

Gene annotation

OR   gene   annotation   was   done   using   the   *HAPpy-ABCENTH*   pipeline
(https://github.com/biorover/HAPpy-ABCENTH). The *ABCENTH* (Annotation Based on
Conserved Exons Noticed Through Homology) program is an algorithm for building gene
models that was designed for multigene families like ORs, with high sequence divergence
but a conserved exon structure. *ABCENTH* builds HMM profiles separately for each exon
based on known protein coding genes, which increases sensitivity and precision.
ABCENTH then looks for splice sites of the candidate exons according to the expected
exon length and phase. The genes used as queries were taken from manually curated
ant ORs of the species: *Harpegnathos saltator*, *Stigmatomma sp*, *Ooceraea biroi*,
Camponotus floridanus and *Solenopsis invicta*. Subsequently, novel OR annotations
were evaluated by comparing them to OR sequences from *Drosophila melanogaster* and
the five abovementioned ants. We identified the OR co-receptor (ORCO) in each species.
Annotated ORs were classified in three categories: complete genes with the expected
exon structure, pseudogenes with in-frame stop codons, and partial or fragmented
annotations with incomplete exon structures, which can be caused by pseudogenization,
fragmented assembly or assembly and annotation errors.

Phylogenetic analysis

The large number of complete OR sequences – 55,068 – made it impossible to analyse
the full dataset jointly. Therefore, each of the 29 OR subfamilies defined by McKenzie et
al. (2016) was analysed separately, and the results were pooled together for statistical
analyses. ORs were assigned to a subfamily using *BLASTP* (version 2.11.0; Altschul et
al., 1990), based on the top hit to sequences classified in the published subfamilies. 98%
of the ORs (54,051 out of 55,068) had their top five hits within the same subfamily. The
9e subfamily comprises 19,528 sequences in our dataset, so it was further subdivided
into six monophyletic clades that had 100% bootstrap support (we also observed issues
in the alignment of these clades to each other, which would result in false positives in the
dN/dS tests).

204
We constructed a multiple sequence alignment and tree for each OR subfamily and
subset of species chosen for the positive selection analysis. Nucleotide coding sequences
were used to build codon alignments using *Guidance2* (version 2.02; Sela et al. 2015)
with the aligner *PRANK* (version 140110; Löytynoja and Goldman, 2005). Unreliably
aligned codons were masked at a cutoff on the *Guidance* confidence scores that was
chosen ad hoc for each alignment. We started by masking each alignment at a threshold
of 0.8, and if more than 20% of the codons were masked we lowered the threshold by 0.1
and tested the percentage of masked codons again. We repeated the operation until we
achieved a maximum of 20% masked codons, to avoid losing too much information for
the inference of positive selection. This procedure was demonstrated as good practice
for controlling the false positive rate of the branch-site test for positive selection (Privman

et al. 2012). The OR gene trees were built using *RAxML* (version 8.1.15; Stamatakis,
2014) with the translated sequence alignments as input, using the PROTCALG model,
the LG rate matrix, and 100 bootstraps repeats.

Positive selection inference

Positive selection was inferred using the branch-site test for positive selection (Zhang
2005) as implemented in *Godon* (Davydov et al. 2019), using the M0 model to estimate
branch length, four categories for codon rate variation, and testing all branches. A
likelihood ratio test was conducted for each gene tree branch contrasting between the
null model, not allowing positive selection, and the alternative model, allowing for positive
selection in that branch. The likelihood ratio test p-values were then corrected for multiple
testing using the Benjamini-Hochberg (1995) method.

Testing for association between trait evolution and OR sequence evolution

We developed a novel method for mapping positive selection results from genes trees of
multi-copy gene families to the species tree, which is described in detail in our
methodological paper (Pellen and Privman 2025). Briefly, we tested for enrichment of
positive selection on ORs in species tree branches where we inferred a change in our
traits of interest (the "transition" branches). The *Generax* (version 1.1.0; Morel et al.,
2020) algorithm for reconciliation between gene trees and species trees was used to map
branches of the gene trees onto branches of the species tree, with the UndatedDL
reconciliation model. This step allows mapping of gene tree branches that were tested for
positive selection to the species tree branch that we wish to test for enrichment. We then
used a hypergeometric test to test for enrichment of positive selection (i.e. gene tree
branches having *Godon* p-value < 0.05) in the species tree "transition" branches. Since
we expect it to be more likely that positive selection occurred in a long branch than in a
short branch, we normalized the number of gene tree branches by a factor defined as
their branch length relative to the average gene tree branch length of the studied clade.
This normalized number of branches was used as the size of the population that is tested
for enrichment in the hypergeometric test. The ratio of observed/expected frequency of
gene tree branches with positive selection were calculated to determine by what factor
was the transition branch enriched or depleted for positive selection.

OR gene family expansions and contractions

To analyse the evolution of OR genes numbers, we applied the BDI stochastic model
implemented in the *BadiRate* program (Librado et al. 2012). To avoid bias due to genome
incompleteness or fragmentation, we excluded the 20 low quality genomes (short-read
based with N50 < 500Kb), except for the *Leptanilla* outgroup, and added *Apis mellifera*
as an outgroup. We then applied a global rates model (BDI-GR-CSP in *BadiRate*; Sankoff
parsimony) to the numbers of complete OR genes. We note that the number of ORs
reconstructed for the ant ancestor may be an underestimate due to *Leptanilla sp.*, the
most basal ant species with only 151 complete ORs annotated. This genome has a lower
quality compared to the rest of the dataset, as it is sequenced with the stLFR method
only. It has an N50 size of 0.13 Mb, while the average for the dataset is 4.55 Mb. This low

assembly contiguity may result in many ORs appearing to be partial genes which are not
counted in the final dataset.
260
We also conducted a Phylogenetic Generalized Least Squares (PGLS) analysis using the
pgls function in the R package *caper* (version 1.0.3, https://CRAN.R-
project.org/package=caper) using the dated ant phylogeny. Again, we excluded the 20
species with lower quality genome assemblies by using *drop.tip* in the R package *treeio*
to prune the species tree. We used PGLS to fit regression models between traits and
ORs numbers (Supplementary Table S2). OR subfamilies with mostly single copy genes
were removed from the final results to reduce spurious correlations.
268

Results

The odorant receptor repertoire of ants

The total number of ORs found through the annotation of 163 ant genomes was 66,169.
This count includes partial ORs, which can be missing some terminal and/or internal parts
of the full-length sequence. The percentage of partial ORs in each species varied from
4% to 56% (Supplementary Table S1) and was related to the contiguity of the genome
assembly (scaffold N50 size), with fragmented assemblies having more partial ORs
(Pearson correlation r = -0.28 between N50 scaffold size and the percentage of partial
ORs, p-value = 0.00074). For example, *Gigantiops destructor* has the lowest N50 (8,895
bp) and the highest proportion of partial ORs (56%). Low quality genomes (N50 < 500Kb)
were therefore excluded from analyses regarding gene numbers as they are expected to
have biased under-estimates of OR numbers. We excluded partial ORs from further
analyses, which left a total of 55,068 complete ORs across all 163 genomes, and 50,657
ORs without the low-quality genomes, ranging from 90 (*Tetramorium "Anergates"*
atratulus) to 687 (*Pseudoneoponera rufipes*), with an average of 338 ORs per species
(Figure 1). The ancestral reconstruction of OR gene numbers estimated 378 ORs for the
ancestor of ants not including *Leptanilla*, which was excluded due to its low-quality
genome (see inferred ancestral numbers in Supplementary Figure S5). The number of
ORs expanded to 455 in the ancestor of the formicoid clade, while there was no further
significant expansion in the poneroids ancestor (with 375 ORs).

Figure 1: OR evolution along the ant phylogeny, including ancestral reconstruction of OR numbers and positive selection enrichment tests. Branch colour indicates the inferred ancestral number of ORs (see inferred ancestral numbers in Supplementary Figure S5). Coloured shapes mark branches with trait shifts that were tested for enrichment of positive selection on ORs. Shape colour indicates the statistical significance of enrichment or depletion based on the multiple testing corrected *p*-values (*q*-values). The species used in the positive selection analysis are marked with stars. The heatmap shows the number of complete ORs for each species partitioned to the 9-exon clade and the rest of the ORs, as well as partial genes.

We observed substantial variation in OR numbers along the phylogeny. Most species have fewer 9e ORs than non 9e ORs (Figure 1). The only exceptions being the small clade of army ants. *Ooceraea biroi* is the most extreme case with 315 9e ORs against 185 non 9e, as previously reported for this species (McKenzie et al. 2016; McKenzie and Kronauer 2018). Conversely, we note the *Myrmicocrypta* species as an extreme outlier for the number of non 9e ORs, with 458. Gene numbers of many OR subfamilies vary dramatically across ant species, most notably in the 9e subfamily and the other four largest subfamilies (E, L, U, V). Looking at the variation in the 9e ORs (Supplementary Figure S6), one species stands out, the workerless social parasite *Tetramorium atratulus*, which has the smallest number of 9e ORs. This species also shows drastic reductions in the other large OR subfamilies and is the only ant species that completely lost the medium-sized subfamilies P, T, M, and N. In subfamily P (Supplementary Figure S7), we noted a clade-specific expansion in the clade that includes *Camponotus*, *Polyrachis* and *Anoplolepis* (with smaller number in *Oecophylla*), and ponerine clade-specific expansions in *Neoponera* and in the clade of *Odontomachus* and *Leptogenys*. In subfamily H (Supplementary Figure S8), *Myrmicocrypta squomosa uncinata* stands out with 236 ORs, against an average of 12 over the rest of the ants. This seems to be due to extensive tandem duplications of subfamily H ORs that are restricted to this species (Supplementary Figure S9).

318
319 **Figure 2: Subfamily composition of the OR superfamily across the ant phylogeny.** The plotted data
320 include only complete ORs, and exclude the species with low-quality genomes.
321
322

Correlation of OR gene numbers with social and ecological traits

The elaboration of sociobiological traits may require an extended OR repertoire to enable more sophisticated chemical communication in ant colonies. Therefore, we predicted a correlation between the number of ORs and traits such as colony size or worker polymorphism. We also expected that adaptation for different diets may be associated with the OR gene numbers. Therefore, we tested for an association between the total number of ORs or the size of OR subfamilies and specific traits of interest using a PGLS analysis (Figure 3A and Supplementary Table S2). We observed an expansion of the 9e OR subfamily in species with larger colony size (*q*-value = 0.012), while another large subfamily, V, showed a significant contraction with bigger colony size (*q*-value = 0.012). Our study confirms previously described significant contractions of the OR gene family associated with the evolution of social parasitism (Schrader et al. 2021; Jongepier et al. 2022; Vizueta et al. 2025). This contraction was in the total number of ORs (*q*-value = 0.000015) and specifically in the 9e (*q*-value = 0.000027) as well as several other OR subfamilies. The only exception is subfamily N that was expanded in social parasites (*q*-value = 0.070). We further focused the analysis on inquilines, the more extreme form of social parasitic species that live in the nest of another ant species. Inquilines show an even stronger contraction (*q*-value = 0.000000027), which is particularly striking in *Tetramorium atratulum*, a workerless inquiline species which retained only 90 complete ORs, 26 of which are in the 9e subfamily compared to an average of 120 9e ORs in other ant species (Figure 3 and Supplementary Table S1). Contrary to our expectations, we found no significant correlation between OR gene numbers and worker polymorphism, worker reproduction, or diet.

A)

ORs subfamily	Average colony size	Social parasitism	Inquilinism	Worker polymorphism	Diet	Worker ovary	Worker spermatheca	Worker sterility
All ORs	0.145	1.49E-05	2.67E-08	0.878	0.852	0.935	1	0.997
9-exon ORs	0.0119	2.67E-05	2.67E-08	0.878	0.302	1	1	0.997

B)

Figure 3: A) Phylogenetic Generalized Least Squares (PGLS) tests for association of gene family expansion or contraction with chosen traits, for the whole OR superfamily and for the 9-exon subfamily. Results are presented after correction for multiple testing (*q*-values). B) Number of ORs for species that are either not socially parasitic, temporary parasites, slave makers or inquilines.

Positive selection on ORs

We used the branch-site test for positive selection to infer adaptive evolution of OR protein sequences in individual branches of the OR gene tree. We then evaluated whether species tree branches were enriched or depleted in positive selection on ORs (see Methods). We focused this analysis on subsets of the species phylogeny that allowed testing branches where we observed shifts in our chosen target traits ("transition branches") based on ancestral trait reconstruction: colony size, worker reproduction, worker polymorphism, and diet (marked by shapes in Figure 1). This analysis involved testing branches of the gene trees of each OR subfamily for each subset of species, 249 gene trees in total. Out of a total of 89,802 gene tree branches tested, positive selection was detected in 652 branches (0.7%), with a false discovery rate of 10% after correction for multiple testing.

We pooled the results for transition branches showing the same trait change. For example, we pooled the ten different branches that showed a convergent increase in colony size. We generally observed depletion in positive selection associated with most trait transitions (Figure 4). Despite being statistically significant, the depletion seen for increased colony size has a negligible effect size (0.998-fold depletion, *p*-value=0.036). The evolution of continuous worker polymorphism was the exception, with a significant enrichment of positive selection (1.18-fold enrichment, *p*-value=0.007). Also the ancestral branches of the formicoid clade were enriched for positive selection (1.075-fold enrichment, *p*-value=0.0476).

We were especially interested in positive selection in the 9e subfamily because it was suggested to be involved in social communication. We found that the enrichment in the formicoid ancestor was driven by the 9e subfamily (1.43-fold enrichment, *p*-value=0.016; Figure 3), while non 9e ORs were depleted for positive selection. A similar pattern was observed for the evolution of continuous worker polymorphism (1.36-fold enrichment in the 9e ORs, *p*-value=0.0022). In contrast, the shift to discrete polymorphism showed an opposite effect, with a significant depletion (0.863-fold depletion, *p*-value=0.014). In general, the enrichment or depletion in the 9e ORs mostly follows the pattern for total ORs but shows a stronger effect. The exception is the depletion associated with changes in diet, which was most pronounced in the non 9e ORs.

In addition to the pooled results, we also inspected individual transition branches. We found significant enrichment of positive selection in 11 branches and depletion in 23 branches (*q*-value < 0.10; Supplementary Table S3). For the three ancestral formicoid branches, the two more ancient branches were enriched and the most recent one was depleted for positive selection. Some branches display especially strong depletion or enrichment for positive selection. For example, the common ancestor of *Wasmannia* and *Pheidole*, where increased colony size was inferred (Supplementary Figure S1), was strongly depleted for positive selection (0.534-fold depletion, *q*-value=0.03262). Out of the 14 branches in which discrete polymorphism evolved (Supplementary Figure S3), four were enriched and ten depleted for positive selection. Among these, the common ancestor of the clade including *Oecophylla* and *Camponotus* was strongly depleted (0.477-fold depletion, *q*-value=0.03262) of positive selection. The common ancestor of

 Cardiocondyla and *Tetramorium*, showing a loss of functional ovaries in workers
398 (Supplementary Figure S4), was strongly depleted for positive selection (0.595-fold
399 depletion, *q*-value=0.02461).
400

401
402 **Figure 4: Enrichment for positive selection on ORs associated with trait evolution.** Results of the
403 hypergeometric test for enrichment of branches under positive selection (considering Godon's p-values
404 under 0.05 as positive) for the pooled branches exhibiting the same change in trait.
405 a: Length-normalized number of gene tree branches mapped on the species tree branch that is tested.
406

Discussion

The high quality of the GAGA genomes and wide coverage across the ant phylogeny,
coupled with an annotation pipeline specifically tailored for ant ORs, resulted in 50,657

ORs across 143 species, a more comprehensive and complete dataset for the study of
the evolution of ant ORs than any previously generated dataset. We used this
unprecedented gene set to infer ancestral OR repertoire sizes, reconstruct gene trees for
all ant OR subfamilies, and test for adaptive evolution of ORs that may be associated with
important sociobiological and ecological traits of ants.
415
Our tests found only 0.7% of branches under positive selection, which is low compared
to previous studies that tested for positive selection in ant ORs (Zhou et al. 2015; Saad
et al. 2018). This may be because we focused mainly on relatively recent events and
short branches in the ant phylogeny, whereas previous studies analysed only a few
species representing different ant subfamilies, making the tested branches much longer.
It could also be because we tested many more branches, forcing us to be more stringent
in our FDR correction.
423
Our results confirmed previous analyses, finding evidence of an ancestral expansion of
the OR gene family in ants (C.D. Smith et al. 2011; Engsontia et al. 2015; Mier et al.
2022), and especially the 9e subfamily. We also observed further expansion in the
formicoid ancestor, consistent with the further elaboration of social traits inferred in this
lineage. Genome-wide analyses also inferred significant expansions in the common
ancestor of ants in gene families implicated in chemical communication, including
chemoperception and synthesis of cuticular hydrocarbons (Vizueta et al. 2025). The
ancestral expansion of the OR gene family, likely playing a major role in the evolution of
sociality in ants, has been reversed in socially parasitic species (Schrader et al. 2021;
Jongepier et al. 2022; Vizueta et al. 2025), and this results also occurs in the 9e subfamily.
434
Previous studies identified an enrichment in positive selection on the branch of the
ancestor of all formicoid ants (Romiguier et al. 2022; Vizueta et al. 2025), but these were
genome-wide analyses limited to single-copy genes, and did not include large gene
families such as ORs. This is because the study of adaptive evolution in multi-copy gene
families requires complex reconciliation of gene trees with the species tree, to account
for gene duplications and losses (Page and Charleston 1998). Here we put in first
evidence for enrichment of positive selection on the OR gene family in the formicoid
ancestor. This pattern is driven by the 9e subfamily of ORs that is highly enriched for
positive selection, while non 9e ORs are actually depleted. This is in-line with the
hypothesis that 9e ORs are responsible for social communication functions in ants,
including their putative role in nestmate recognition, which is supported by their specificity
for cuticular hydrocarbons and their expression in workers (Engsontia et al. 2015; Zhou
et al. 2015; McKenzie et al. 2016; Pask et al. 2017; Slone et al. 2017). We propose that
as the ancestor of formicoid ants evolved various elaborate sociobiological traits such as
larger colonies and greater specialization of the non-reproductive castes, they evolved
additional complexity in their chemical communication, including new 9e ORs that
respond to chemical signals like CHCs shaped by positive selection.
452
Overall, we observed more depletions in positive selection than enrichments in the
transition branches we tested, especially in the most recent ones, and the 9e subfamily
seems to be the major driver of these changes. We observed depletion of positive

selection associated with increased colony size, complementing the findings by Vizueta
et al. (2025) of genome-wide relaxation of purifying selection in species with increased
colony size. Colony size was suggested to be a major driver in the evolution of ants, with
many other social traits depending on it, including worker polymorphism (Bell-Roberts et
al. 2024; Vizueta et al. 2025). The genome-wide analysis by Vizueta et al. revealed
intensified selection associated with the recent loss of worker ovaries in multiple formicoid
lineages, but predominantly relaxation of selection associated with the evolution of worker
polymorphism. Our analysis found depletion of positive selection on ORs associated with
the loss of worker reproductive functions. While we found that evolution of discrete
polymorphism shows a depletion in positive selection, we detected enrichment of positive
selection associated with the evolution of continuous polymorphism. This intriguing
pattern would be interesting to investigate further, as it could imply that the evolution of
different forms of worker polymorphism involved different mechanisms of chemical
communication. The differences between our results and the analyses by Vizueta et al.
may be attributed to the very different nature of the genes analysed: while we studied the
fast-evolving multi-gene OR family, their analyses were limited to conserved single-copy
gene families. Enrichment or depletion of positive selection on ORs is most strongly
represented in the 9e subfamily for all tested sociobiological traits. It is interesting to note
the exception of transition to herbivorous diet, which was depleted for positive selection.
This depletion was most pronounced in the non 9e ORs, a result that is in line with the
hypothesis that non-social olfactory functions are mediated by other OR subfamilies.
Overall, our results indicate that only the evolution of certain worker traits involved a
detectable signature of positive selection on ORs, whereas the elaboration of other
sociobiological traits was associated with less adaptive evolution of ORs.

480 Conclusion

ORs expanded in the ant ancestor, mainly driven by the 9e subfamily that is associated
with social communication. Our analyses detected enrichment of positive selection on
ORs in the ancestor of the formicoid ants, mostly in the 9e subfamily, which was
accompanied by a further expansion of OR gene numbers. These results support a
central role for 9e ORs in the evolution of this large and diverse ant clade. Enrichment of
positive selection in the 9e ORs was also associated with the evolution of continuous
worker polymorphism, providing additional evidence for the role of this subfamily in social
evolution. Our study demonstrates the potential of molecular evolutionary analyses of the
OR gene family for explaining the evolution of social traits in ants. Our analyses provide
new insights into the role of ORs in the social evolution and diversification of ants and
raise new questions about specific aspects of OR evolution in the context of specific
sociobiological traits and in specific lineages. It would be interesting to investigate the
underlying causes for the depletion in positive selection in more recent branches, and
what explains the differences observed between discrete and continuous worker
polymorphism. Future studies including more species as well as functional genomic data
could shed light on these evolutionary processes.
497

Acknowledgements

We thank Sean McKenzie for help with using the *HAPpy-ABCENTH* pipeline for OR annotations. This study was funded by Israel Science Foundation grant number 1845/22.

Data availability

All the data generated in this study was submitted to the zenodo repository (10.5281/zenodo.16753530) including the OR annotations in GFF and FASTA format, multiple sequence alignments, gene trees, and generax results.

References

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J Mol Biol* 215:403–410.

Bargmann CI. 2006. Comparative chemosensation from receptors to ecology. *Nature* 444:295–301.

Bear DM, Lassance J-M, Hoekstra HE, Datta SR. 2016. The Evolving Neural and Genetic Architecture of Vertebrate Olfaction. *Current Biology* 26:R1039–R1049.

Bell-Roberts L, Turner JFR, Werner GDA, Downing PA, Ross L, West SA. 2024. Larger colony sizes favoured the evolution of more worker castes in ants. *Nat Ecol Evol* 8:1959–1971.

Blomquist GJ, Tittiger C, Jurenka R. 2020. Cuticular Hydrocarbons and Pheromones of Arthropods. In: Wilkes H, editor. Hydrocarbons, Oils and Lipids: Diversity, Origin, Chemistry and Fate. Cham: Springer International Publishing. p. 213–244. Available from: http://link.springer.com/10.1007/978-3-319-90569-3_11

Davydov II, Salamin N, Robinson-Rechavi M. 2019. Large-Scale Comparative Analysis of Codon Models Accounting for Protein and Nucleotide Selection.Pupko T, editor. *Molecular Biology and Evolution* 36:1316–1332.

Demuth JP, Hahn MW. 2009. The life and death of gene families. *BioEssays* 31:29–39.

Dornhaus A, Powell S, Bengston S. 2012. Group Size and Its Effects on Collective Organization. *Annual Review of Entomology* 57:123–141.

Engsontia P, Sangket U, Robertson HM, Satasook C. 2015. Diversification of the ant odorant receptor gene family and positive selection on candidate cuticular hydrocarbon receptors. *BMC Res Notes* 8:380.

Gautam S, McKenzie S, Katzke J, Hita Garcia F, Yamamoto S, Economo EP. 2024. Evolution of odorant receptor repertoires across Hymenoptera is not linked to the

evolution of eusociality. *Proceedings of the Royal Society B: Biological Sciences* 291:20241280.

Innan H, Kondrashov F. 2010. The evolution of gene duplications: classifying and distinguishing between models. *Nat Rev Genet* 11:97–108.

Jongepier E, Séguret A, Labutin A, Feldmeyer B, Gstöttl C, Foitzik S, Heinze J, Bornberg-Bauer E. 2022. Convergent Loss of Chemoreceptors across Independent Origins of Slave-Making in Ants.Wittkopp P, editor. *Molecular Biology and Evolution* 39:msab305.

Keller L, Genoud M. 1997. Extraordinary lifespans in ants: a test of evolutionary theories of ageing. *Nature* 389:958–960.

Librado P, Vieira FG, Rozas J. 2012. BadiRate: estimating family turnover rates by likelihood-based methods. *Bioinformatics* 28:279–281.

Löytynoja A, Goldman N. 2005. An algorithm for progressive multiple alignment of sequences with insertions. *Proc Natl Acad Sci U S A* 102:10557–10562.

McKenzie SK, Fetter-Pruneda I, Ruta V, Kronauer DJC. 2016. Transcriptomics and neuroanatomy of the clonal raider ant implicate an expanded clade of odorant receptors in chemical communication. *Proceedings of the National Academy of Sciences* 113:14091–14096.

McKenzie SK, Kronauer DJC. 2018. The genomic architecture and molecular evolution of ant odorant receptors. *Genome Res.* 28:1757–1765.

Mendivil Ramos O, Ferrier DEK. 2012. Mechanisms of Gene Duplication and Translocation and Progress towards Understanding Their Relative Contributions to Animal Genome Evolution. *International Journal of Evolutionary Biology* 2012:846421.

Mier P, Fontaine J-F, Stoldt M, Libbrecht R, Martelli C, Foitzik S, Andrade-Navarro MA. 2022. Annotation and Analysis of 3902 Odorant Receptor Protein Sequences from 21 Insect Species Provide Insights into the Evolution of Odorant Receptor Gene Families in Solitary and Social Insects. *Genes (Basel)* 13:919.

Morel B, Kozlov AM, Stamatakis A, Szöllősi GJ. 2020. GeneRax: A Tool for Species-Tree-Aware Maximum Likelihood-Based Gene Family Tree Inference under Gene Duplication, Transfer, and Loss. *Molecular Biology and Evolution* 37:2763–2774.

Ohno S. 1970. Evolution by Gene Duplication. Berlin, Heidelberg: Springer Available from: http://link.springer.com/10.1007/978-3-642-86659-3

Oster GF, Wilson EO. 1978. Caste and Ecology in the Social Insects. Princeton University Press

Page RDM, Charleston MA. 1998. Trees within trees: phylogeny and historical associations. *Trends in Ecology & Evolution* 13:356–359.

Paradis E, Schliep K. 2019. ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics* 35:526–528.

Pask GM, Slone JD, Millar JG, Das P, Moreira JA, Zhou X, Bello J, Berger SL, Bonasio R, Desplan C, et al. 2017. Specialized odorant receptors in social insects that detect cuticular hydrocarbon cues and candidate pheromones. *Nat Commun* 8:297.

Pellen Y, Privman E. 2025. Testing for positive selection in multi-copy gene families using a reconciliation approach. *bioRxiv*.

Privman E, Penn O, Pupko T. 2012. Improving the Performance of Positive Selection Inference by Filtering Unreliable Alignment Regions. *Molecular Biology and Evolution* 29:1–5.

Robertson HM. 2019. Molecular Evolution of the Major Arthropod Chemoreceptor Gene Families. *Annu. Rev. Entomol.* 64:227–242.

Romiguier J, Borowiec ML, Weyna A, Helleu Q, Loire E, La Mendola C, Rabeling C, Fisher BL, Ward PS, Keller L. 2022. Ant phylogenomics reveals a natural selection hotspot preceding the origin of complex eusociality. *Current Biology* 32:2942-2947.e4.

Saad R, Cohanim AB, Kosloff M, Privman E. 2018. Neofunctionalization in Ligand Binding Sites of Ant Olfactory Receptors.Chang B, editor. *Genome Biology and Evolution* 10:2490–2500.

Schrader L, Pan H, Bollazzi M, Schiøtt M, Larabee FJ, Bi X, Deng Y, Zhang G, Boomsma JJ, Rabeling C. 2021. Relaxed selection underlies genome erosion in socially parasitic ant species. *Nat Commun* 12:2918.

Sela I, Ashkenazy H, Katoh K, Pupko T. 2015. GUIDANCE2: accurate detection of unreliable alignment regions accounting for the uncertainty of multiple parameters. *Nucleic Acids Res* 43:W7–W14.

Shields EJ, Sheng L, Weiner AK, Garcia BA, Bonasio R. 2018. High-Quality Genome Assemblies Reveal Long Non-coding RNAs Expressed in Ant Brains. *Cell Reports* 23:3078–3090.

Slone JD, Pask GM, Ferguson ST, Millar JG, Berger SL, Reinberg D, Liebig J, Ray A, Zwiebel LJ. 2017. Functional characterization of odorant receptors in the ponerine ant, *Harpegnathos saltator*. *Proc. Natl. Acad. Sci. U.S.A.* 114:8586–8591.

Smith CD, Zimin A, Holt C, Abouheif E, Benton R, Cash E, Croset V, Currie CR, Elhaik E, Elsik CG, et al. 2011. Draft genome of the globally widespread and invasive

Argentine ant (*Linepithema humile*). *Proc. Natl. Acad. Sci. U.S.A.* 108:5673–
5678.

Smith CR, Smith CD, Robertson HM, Helmkampf M, Zimin A, Yandell M, Holt C, Hu H,
Abouheif E, Benton R, et al. 2011. Draft genome of the red harvester ant
Pogonomyrmex barbatus. *Proc. Natl. Acad. Sci. U.S.A.* 108:5667–5672.

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis
of large phylogenies. *Bioinformatics* 30:1312–1313.

Trible W, Olivos-Cisneros L, McKenzie SK, Saragosti J, Chang N-C, Matthews BJ, Oxley
PR, Kronauer DJC. 2017. *orco* Mutagenesis Causes Loss of Antennal Lobe
Glomeruli and Impaired Social Behavior in Ants. *Cell* 170:727-735.e10.

Vizueta J, Xiong Z, Ding G, Larsen RS, Ran H, Gao Q, Stiller J, Dai W, Jiang W, Zhao J,
et al. 2025. Adaptive radiation and social evolution of the ants. *Cell* [Internet].
Available   from:
https://www.sciencedirect.com/science/article/pii/S0092867425006178

Zhang J. 2005. Evaluation of an Improved Branch-Site Likelihood Method for Detecting
Positive Selection at the Molecular Level. *Molecular Biology and Evolution*
22:2472–2479.

Zhou X, Rokas A, Berger SL, Liebig J, Ray A, Zwiebel LJ. 2015. Chemoreceptor Evolution
in Hymenoptera and Its Implications for the Evolution of Eusociality. *Genome*
Biology and Evolution 7:2407–2416.

Zhou X, Slone JD, Rokas A, Berger SL, Liebig J, Ray A, Reinberg D, Zwiebel LJ. 2012.
Phylogenetic and Transcriptomic Analysis of Chemosensory Receptors in a Pair
of Divergent Ant Species Reveals Sex-Specific Signatures of Odor Coding.Moran
NA, editor. *PLoS Genet* 8:e1002930.

van Zweden JS, d'Ettorre P. 2010. Nestmate recognition in social insects and the role of
hydrocarbons. In: Blomquist GJ, Bagnères A-G, editors. Insect Hydrocarbons. 1st
ed.   Cambridge   University   Press.   p.   222–243.   Available   from:
https://www.cambridge.org/core/product/identifier/CBO9780511711909A021/type
/book_part

631