LFA-1 Interaction with GBP-130 on *Plasmodium falciparum*-infected Red Blood

Cells mediates NK Cell Activation and Parasite Control

Osama Mukhtar[1], Ravi Dutt[2], Ashutosh Panda[1], Poonam Kumari[1, 5], Suneet Shekhar

Singh[1], Gourab Paul[1, 4], Neha Prakash[3], Madiha Abbas[3], Md. Muzahidul Islam[3], Priya

Arora[3], Asif Mohmmed[3], Dhiraj Kumar[2*], Pawan Malhotra[1*].

Affiliations:

[1]Malaria Biology Group, International Centre for Genetic Engineering and

Biotechnology (ICGEB), Aruna Asaf Ali Marg, New Delhi – 110067, India.

[2]Cellular Immunology Group, International Centre for Genetic Engineering and

Biotechnology, New Delhi 110067, India.

[3]Parasite Cell Biology Group, International Centre for Genetic Engineering and

Biotechnology, New Delhi 110067, India.

[4]Department of Bioinformatics, Amity School of Biological science, Amity

University Punjab.

[5]Amity Institute of Biotechnology (AIB), Amity University, Kant Kalwar, NH-11C,

Jaipur, Rajasthan 303002, India

* Corresponding authors

E-mails; pawanmal@gmail.com; dhiraj@icgeb.res.in

Abstract

Natural Killer (NK) cells contribute to early immunity against *Plasmodium falciparum* by recognizing and eliminating infected red blood cells (iRBCs), a process mediated in part by the integrin LFA-1. However, the cognate parasite ligand for LFA-1 has remained unknown. Here, we identify Glycophorin Binding Protein-130 (*Pf*GBP-130) as a surface-expressed ligand on iRBCs that binds the I-domain of LFA-1 (LFA-1 αI). Using an LFA-1 αI–Fc fusion protein, we demonstrate stage-specific binding to iRBCs, and LC-MS/MS analysis of immunoprecipitates of αI–Fc bound to iRBC revealed *Pf*GBP-130 as a high-confidence interactor. Recombinant *Pf*GBP-130 binds NK and THP-1 cells in an LFA-1–dependent manner. Co-culture assays show that *Pf*GBP-130 promotes NK cell activation, degranulation, and facilitates contact-dependent killing of iRBCs. Neutralizing antibodies against *Pf*GBP-130 significantly impair these responses. Our findings establish *Pf*GBP-130 as the LFA-1 ligand on iRBCs, providing new insight into NK cell–mediated immunity in malaria and identifying a potential target for host-directed interventions.

Introduction

NK cells are innate immune cells involved in early defence against microbial pathogens, parasitic infections and tumor cells (Burrack et al., 2019). NK cells are found in various tissues including liver, peritoneal cavity as well as placenta and constitute up to 15% of peripheral blood lymphocytes. NK cells were initially identified due to their ability to kill certain tumor cells *in vitro* (Cerwenka and Lanier, 2001). Subsequent studies using *in vivo* murine models demonstrated that depletion of NK cells leads to enhanced tumor formation (Karre, 2002). Activation of NK cells results from the concerted action/response of cytokine signalling, adhesion molecules and the interaction of activating receptors with their corresponding ligands expressed on the surface of pathogenic infected cells or tumors. Despite having understood the role(s) of NK cells in controlling the microbial infeuction and tumor surveillance, mechanisms of NK cells recognition by pathogenic cells are not well understood.

Malaria infection and pathogenesis involves a complex series of interplays between parasite and human host factors and a better understanding of host cells (factors) may be important in developing innovative host derived therapeutic approaches. NK cells provide first line of defence against malaria parasite infection. Studies in mouse malaria models have shown that various immune cells such as natural killer (NK) cells, dendritic cells, T cells and B cells contribute to antiparasitic immunity (Chen et al., 2014; Ye et al., 2018). *Plasmodium* development in humans occurs in two phases: the merozoite stage in the liver and the symptomatic blood stage. During the liver stage, immunity against *Plasmodium* is primarily mediated by IFN-gamma. NK cells secrete IFN-gamma, which eliminates infected hepatocytes either directly or by activating other immune cells, including cytotoxic T cells and macrophages (Artavanis-Tsakonas and Riley, 2002; Burrack et al., 2019).

NK cells contribute to blood stage immunity through three main mechanisms. First, they facilitate the clearance of infected red blood cells (iRBCs) via a bystander mechanism, involving the secretion of IL-12 and IL-18. These cytokines prompt macrophages and dendritic cells to clear iRBCs or stimulate adaptive immune cells. Second, NK cells engage in antibody-dependent cell-mediated cytotoxicity (ADCC), wherein antibody-opsonized iRBCs are recognized via the CD16 (FcγRIII) receptor. This leads to the release of cytotoxic granules, including perforin and granulysin, enabling targeted lysis of infected cells. Third, NK cells can directly kill infected or

transformed cells through activating receptor–ligand interactions, a mechanism well characterized in antiviral and tumor immunity. NK cells are critical in limiting acute malaria infection as depletion of NK cells in mouse malaria model has been associated with higher parasitemia and accelerated disease progression (Chen et al., 2014; Ye et al., 2018). NK cell protective effects are mediated through both cytotoxic activity and secretion of interferon-gamma (IFN-γ). Clinical studies in *P. falciparum*-infected children have revealed an inverse correlation between NK cell frequency or functional activity and parasite burden (Ojo-Amaize et al., 1981). Furthermore, experimental infections in malaria-naïve individuals have demonstrated that NK cells are among the earliest responders, capable of directly lysing iRBCs and releasing IFN-γ and soluble granzyme (Hermsen et al., 2003; Ye et al., 2018).

Plasmodium Infected RBCs (iRBCs) activates NK cell, enhancing IFN-γ production and cytotoxicity through perforin and granzyme release. This activation is contact-dependent, as demonstrated by the lack of upregulation of NK cell activation markers during transwell incubation. Co-incubation of iRBCs with NK cells promote *Plasmodium* parasite killing, and this contact-dependent interaction is mediated in part by the lymphocyte function-associated antigen-1 (LFA-1) on NK cells (Chen et al., 2014; Korbel et al., 2005). In this study, we demonstrate that iRBCs bind specifically to the "αI-domain" of LFA-1 and identified *Plasmodium* Glycophorin Binding Protein-130 (*Pf*GBP-130) as a major iRBC surface ligand interacting with LFA-1. Furthermore, we show that *Pf*GBP-130 expressed on the iRBC surface binds to THP-1 and NK cells, inducing activation and degranulation of these cells. Together, our findings identify *Pf*GBP-130 as a ligand for LFA-1 that facilitates direct contact-mediated killing of *Plasmodium*-infected erythrocytes by NK cells, highlighting its potential role in host–parasite immune interactions.

Results

LFA αI domain of Lymphocyte Function Associated Antigen-1 (LFA-1) is involved in ligand binding on iRBCs surface.

Contact-dependent killing of *Plasmodium*-infected red blood cells (iRBCs) by NK cells is essential for effective malaria control and is mediated by LFA-1 (CD11a/CD18), a β2 integrin composed of an αL (CD11a) and β (CD18) subunit. Notably, CD11a contains an inserted (I) domain of approximately 200 amino acids,

homologous to the von Willebrand factor type A (vWA) domain family (Suppl. Fig.

1), which plays a critical role in ligand binding, including interaction with ICAM-

1(Qu and Leahy, 1995). Here, we generated an LFA-1 αI-Fc fusion protein to assess

αI domain binding to iRBCs. Schematics of CD11a (a subunit of LFA-1) and it's I

domain fused to human IgG1 Fc domain is shown in Figure 1A. DNA sequences

corresponding to LFA-1 αI-Fc fusion protein were cloned in a plasmid pFUSE-

hIgG1-Fc2 (Suppl. Fig. 2A & 2B) and the protein was expressed in CHO K1 cells

under serum-free conditions, Fusion protein was purified via protein-A affinity beads

and purified protein was analysed by SDS-PAGE and western blot using secondary

anti-human IgG. A protein of a ~45 kDa was seen in SDS-PAGE and western blot

(Fig. 1B), which corresponded to the molecular weight of fusion construct. We next

analysed the binding of purified LFA αI-Fc protein with iRBCs and detected binding

using secondary human antibody conjugated to PE Texas red or anti-human antibody

conjugated to FITC. Binding was also analyzed for all the three asexual blood stages

of *Plasmodium* ring, trophozoites and schizonts by FACS. As shown in figure 1C,

high affinity binding was observed with all the three stages that was significantly

higher than that of observed with the human IgG (hIgG) isotype control. These results

showed that the αI domain of LFA-1 is involved in binding to infected RBC cells.

***P. falciparum* Glycophorin Binding Protein-130 protein on iRBC surface binds to**

LFA αI domain of LFA-1 protein.

To elucidate the molecular basis of LFA-1 mediated interactions with *Plasmodium*

falciparum infected erythrocytes, we undertook a comprehensive proteomic approach.

Briefly, recombinant LFA-1 αI-domain Fc fusion protein (LFA-1 αI-Fc) used as an

affinity reagent to capture iRBC surface-interacting proteins, following DTSSP cross-

linking. Human IgG1 (hIgG1) served as a control to distinguish specific from non-

specific background interactions. Immunoprecipitated proteins from both LFA-1 αI-

Fc and hIgG1 conditions were analyzed by LC-MS/MS, enabling a comparative

assessment of specifically interacting proteins. Comparative analysis revealed a

distinct set of proteins specifically enriched in the LFA-1 αI-Fc fraction (Fig. 2A,

Suppl. Fig. 2C), with *Pf*GBP-130 (PF3D7_1016300) identified as a top candidate

based on peptide coverage and spectral abundance across three biological replicates.

LC-MS/MS analysis thus strongly suggested a specific and robust interaction between

LFA-1 αI domain and *Pf*GBP-130.

To validate the observed interaction and provide orthogonal confirmation of our

proteomic findings, we expressed the N-terminal region of *Pf*GBP-130 (aa 69-270

encompassing one GBP repeat) in *Escherichia coli* and purified the protein to near

homogeneity (Fig. 2B. i). Polyclonal antibodies were raised against recombinant

GBP-130 N terminal protein fragment referred here as *Pf*GBP130-N in rabbit (Fig.

2B. ii). Antibodies raised against *Pf*GBP130-N were specific as they could detect the

native *Pf*GBP-130 on iRBCs surface and this staining showed partially overlap with

Band-3 staining, a RBC resident membrane protein (Fig. 2B. iii). To confirm the

specificity of anti-*Pf*GBP-130 N antibodies, we next performed a western blot

analysis of trophozoite stage parasite lysate and also for the immunoprecipitaed

fraction of LFA-1 αI-Fc protein bound iRBC lysate using anti-PfGBP-130 N

antibodies. As shown in figure 2B. iv, a distinct band at ~110kDa, consistent with the

reported molecular weight of *Pf*GBP-130, were observed exclusively within the LFA-

1 αI-Fc bound iRBCs eluate as well as in iRBCs lysate control. Critically, this band

was absent in the hIgG1 control eluate, confirming the specificity anti-*Pf*GBP-130

antibodies as well as of the observed interaction between *Pf*GBP-130 and

recombinant LFA-1 αI-Fc protein, thus ruling out non-specific binding artifacts.

Biophysical assessment of the interaction between *Pf*GBP130-N and LFA-1 αI-Fc

fragments.

To confirm the interaction between LFA-1 and *Pf*GBP-130, real-time interaction

studies were performed using recombinant *Pf*GBP-130 N and LFA-1 αI-domain

proteins by Bio-layer interferometry (BLI) on the Sartorius Octet K2 system. Briefly,

LFA-1 αI-domain was immobilized onto an AR2G sensor (Octate Amine Reactive

Second-Generation biosensor) at a concentration of 100ng and increasing

concentrations of *Pf*GBP-130 showed a sharp association curve with the bound LFA-

1 αI-domain, followed by a dissociation curve. The dissociation constant (KD) was

1.5 □ 10^{-8} M, indicating a significant binding affinity between LFA-1 αI-Fc and

*Pf*GBP130-N (Fig. 2C. i).

Since LFA-1 αI-domain displayed a high binding affinity for *Pf*GBP-130 N, we next

performed protein-protein docking studies to examine the molecular-level interactions

between these two proteins. The *Pf*GBP-130 model was docked against LFA-1 αI-

domain using the Cluspro2.0 protein-protein docking server, and the binding site for

the generated complex was analyzed with PyMOL. The best resulting structure was

submitted to PDBSum to identify the amino acid residues involved in the interactions.

Examination of the interaction surface showed that LFA1 binds within a curved

pocket of *Pf*GBP130 formed by an anti-parallel helix structure, and the interacting

residues for LFA1 were mainly localized in the N-terminal domain. A total of 13

hydrogen bonds and five salt bridges were observed for the *Pf*GBP130/LFA1

complex. The representative hydrogen bonds are shown in figure 2C. ii.

To assess the strength of the interactions, the buried surface area of the complex was

calculated using the PDBePISA web server, yielding a value of 1528.7 Å² per

molecule, which is in accordance with the buried area suggested for known protein-

protein complexes. Next, we evaluated the stability of the docked complex through

molecular dynamics (MD) simulations in an aqueous environment over 200

nanoseconds (ns). The Root Mean Square Deviation (RMSD) trajectory plot,

indicated that after a minor fluctuation during the first 30 ns, the complex remained

stable for the rest of the simulation (Fig. 2C. iii). In conclusion, both real-time BLI

and *in-silico* studies suggested a stable interaction between LFA-1 expressed on NK

cells with *Pf*GBP-130.

_Pf_GBP-130 ectodomain shows a specific LFA-1-dependent binding to human

THP-1 monocytic and primary NK cells.

*Pf*GBP-130 possesses an N-terminal cytosolic domain, a transmembrane domain, and

an extracellular domain characterized by repeats (Fig. 3A). Since, we identified

*Pf*GBP-130 as a putative ligand for LFA-1 on NK cells, we next investigated the

direct interaction of *Pf*GBP-130 with immune cells, specifically THP-1 monocytes

and NK cells. Briefly, recombinant protein construct comprising of the extracellular

domain of *Pf*GBP-130, encompassing the putative LFA-1 binding sites, fused to the

Fc region of human IgG1 (*Pf*GBP-Fc) was generated (Fig. 3A). The *Pf*GBP-Fc fusion

protein was engineered to ensure stability and solubility for binding assays, while the

human IgG1 Fc tag facilitated detection via secondary anti-human IgG antibodies.

The *Pf*GBP-Fc construct was transiently expressed in CHO K1 cells cultured in

serum-free medium to minimize potential interference from serum components. The

secreted fusion protein was subsequently purified from the culture supernatant using

protein-A affinity chromatography (Suppl. Fig. 3A & 3B). The interaction between

*Pf*GBP_-Fc and THP-1 was investigated, focusing on the role of LFA-1 as a potential

receptor using siRNAs corresponding to CD11a subunit of LFA-1 mRNA. Next the

cultured THP-1 monocytic cells with and without siRNA treatment were fixed and

incubated with recombinant *Pf*GBP-Fc or control hIgG in FACS staining buffer for 2

hours. After washing, cells were stained with fluorescent anti-human PE-Texas Red

antibodies (invitrogen) for binding analysis via flow cytometry to detect the binding

of *Pf*GBP-Fc. Flow cytometric analysis demonstrated a significant increase (3-fold) in

binding of *Pf*GBP -Fc compared to an hIgG isotype control to THP-1 cells, indicating

a specific interaction of *Pf*GBP to THP-1 cells (Fig. 3B, i). To know whether the

*Pf*GBP-Fc binding to THP-1 cells is LFA-1 dependent, we next performed Smartpool

Accel siRNA-mediated gene silencing of CD11a subunit of LFA-1 in THP-1 cells

(Suppl. Fig. 4B). As shown in figure 3B (ii), western blot analysis confirmed efficient

depletion of the CD11a subunit of LFA-1 in THP-1 cells. Subsequent flow cytometric

analysis revealed a substantial reduction in *Pf*GBP130-Fc binding to LFA-1-deficient

THP-1 cells compared to untreated controls (Fig. 3B. i). This reduction thus

definitively established LFA-1 as a crucial mediator of *Pf*GBP-Fc binding to THP-1

monocytic cells.

We further extended these findings to other immune cell populations, such as primary

NK cells. Consistent with the results obtained in THP-1 cells, incubation of NK cells

with *Pf*GBP-Fc resulted in a significant increase in binding intensity relative to the

hIgG1 isotype control (Fig. 3C i). Smartpool Accel siRNA-mediated downregulation

of CD11a subunit of LFA-1 in NK cell (Fig3C-ii &Suppl. Figure 4A), reduced

*Pf*GBP130-Fc binding to NK cells (Fig. 3C. ii), thereby confirming the role of LFA-1

as a critical receptor for *Pf*GBP-Fc binding to NK cell.

Engineered chimeric GBP expressing CHO cell line activates NK cell through

LFA-1 binding.

To elucidate the mechanism by which *P. falciparum* infected red blood cells (iRBCs)

activate natural killer (NK) cells through LFA-1 engagement, we hypothesized that

*Pf*GBP130 serves as a cognate ligand for LFA-1 on the iRBC surface. To test this

hypothesis, a Chinese Hamster Ovary (CHO) cell line was engineered for the stable

expression of *Pf*GBP-130 extracellular domain on its surface by leveraging lentiviral

transduction. The engineered CHO-K1 expressed the extracellular C-terminal domain

of *Pf*GBP-130 fused to the transmembrane domain of transferrin receptor (TfR-TM)

for the surface presentation (Suppl. Fig. 3C). The surface expression was verified

using anti-*Pf*GBP 130 antibodies by immunofluorescence (Fig. 4A. i). Flow

cytometry analysis demonstrated robust binding of recombinant LFA-1 αI-Fc to

*Pf*GBP-130–expressing CHO cells, while no significant binding was observed with

mock-transduced CHO cells, confirming the specificity of the interaction (Fig. 4A. ii).

To further investigate the role of *Pf*GBP-130 in NK cell activation and degranulation

via direct engagement with its cognate receptor LFA-1, we co-cultured CHO cells

stably expressing the extracellular domain of *Pf*GBP-130 (CHO–*Pf*GBP-ECD) with

primary human NK cells at a 2:1 ratio (CHO–*Pf*GBP-ECD: NK). Given that LFA-1

affinity is dynamically regulated through inside-out signaling, co-cultures were

stimulated with Poly I:C/ Lipofectamine 2000 (synthetic double-stranded RNA

analogue), to induce a high-affinity conformational state of LFA-1. Subsequently, NK

cell activation was evaluated by flow cytometric detection of early activation markers

CD69 and CD25.Compared to mock CHO cells, *Pf*GBP130 expressing CHO cells

elicited a significant ~50% increase in CD69 and ~40% increase in CD25 expression,

indicating enhanced NK cell activation by *Pf*GBP-130. This activation was specific as

addition of anti-CD11a antibodies in co-cultures reduced this activation significantly

(Fig. 4B).

Furthermore, we examined the impact of *Pf*GBP-130 expression on primary NK cells

degranulation, an indirect indicator of cytotoxicity. For this surface expression of

CD107a, a marker of lysosomal degranulation was measured using flow cytometry.

Co-culture of NK cell with CHO–*Pf*GBP-ECD resulted in a substantial increase in

CD107a on primary NK cells as compared to mock CHO control. This increase was

significantly reduced in the presence of anti-CD11a antibodies, indicating that the

enhanced degranulation was specifically mediated through LFA-1 engagement with

*Pf*GBP-130 on the CHO cell surface (Fig. 4B). Together, these results indicate that

*Pf*GBP-130 protein promotes LFA-1 mediated activation and cytotoxic degranulation

of NK cells.

296   Human NK cells eliminate iRBCs in a co-culture study and this elimination is

297   dependent on GBP-130 expression on iRBCs.

To gain direct evidence of NK cell mediated killing of iRBCs via *Pf*GBP-130

dependent interaction, we performed co-culture experiments using purified human

NK cells and iRBCs as described earlier by Chen at al 2014 (Chen et al., 2014), in the

presence or absence of anti-*Pf*GBP-130 antibodies. An isotype control antibody

served as a negative control for non-specific antibody effects. Briefly, isolated human

NK cell was treated with Fc receptor blocker using anti-CD16 (3G8 clone) prior to

addition of iRBC (0.5% parasitemia) in 10: 1 ratio (NK: iRBC). The co-cultures were

maintained for 48 hour and 96 hours, a time frame sufficient for NK cell activation

and parasitemia control respectively.

In a parallel set of experiments, we sought to specifically examine the effect of

blocking *Pf*GBP-130 on iRBC with anti-*Pf*GBP neutralizing antibody on NK cell

activation and parasitemia control. Isolated NK cells were co-cultured with iRBCs for

an extended period of 96 hours to allow for a more comprehensive assessment of

parasite growth, which was assayed with Hoechst staining and number of NK cells

were assessed by anti-CD56 NK antibody staining in a flow cytometry analysis. The

iRBCs were positive for Hoechst, but negative for CD56, while NK cells were

positive for both. The co-culture of NK cells with iRBC either alone or in presence of

rabbit IgG (rIgG) isotype control antibody, significantly reduced parasitemia

compared to iRBCs cultured in the absence of NK cells. In contrast, the addition of

anti-*Pf*GBP antibody to NK cell-iRBC co-cultures resulted in a marked and

statistically significant increase in parasitemia, indicating that blockade of *Pf*GBP-130

impairs NK cell–mediated parasite clearance (Fig. 5A). Conversely, parasitemia

levels in the co-cultures supplemented with the isotype control antibody were

observed to be comparable to those in the NK cell culture alone, indicating that the

observed reduction in parasitemia control was specifically attributable to the blockade

of *Pf*GBP-130 and not due to non-specific antibody interactions (Fig. 5A). This

finding confirms the specificity of the anti-*Pf*GBP antibody and reinforces the critical

role of *Pf*GBP in NK cell-mediated parasite clearance.

We further studied the levels of activation and degranulation markers in NK cell in

the absence and presence of anti-GBP-130 antibody (Fig. 5B). Briefly, NK cell and

iRBC were co-cultured in the presence to anti-GBP-130 antibody or rabbit IgG (rIgG)

isotype control antibody. Incubation of human NK cells with iRBC for 48 hrs showed

a significant percent change in MFI of activation markers; CD69 and CD25 as well as

degranulation marker; CD107a. This effect was reduced in the presence of anti-

*Pf*GBP antibody, which signified that *Pf*GBP-130 engagement is necessary for the

activation and degranulation of NK cell (Fig. 5B). These results were specific as

levels of activation and degranulation markers were same in NK cells incubated with

iRBC in the presence or absence of rIgG control. Taken together, the reduced NK cell

activation and increase in parasitemia following *Pf*GBP-130 blockade strongly support the conclusion that direct interaction between LFA-1 on NK cells and GBP-130 on iRBCs is a critical determinant of contact dependent NK cell-mediated immune responses. These findings further establish *Pf*GBP-130 as a key ligand in the LFA-1–dependent activation of NK cell during *Plasmodium falciparum* infection.

Discussion

NK cells have been well described for their ability to eliminate viral-infected cells, tumors and several pathogens including *P. falciparum* (Burrack et al., 2019). A fine-tuned panel of activating and inhibitory receptors regulates the activation of NK cells. Among these, LFA-1, an integrin family member, serves not only as an adhesion molecule but also as an important activating receptor on NK cells (Urlaub et al., 2017). Beyond mediating adhesion, LFA-1 signaling is essential for the polarization of lytic granules towards target cells, a prerequisite for efficient NK cell cytotoxicity(Bryceson et al., 2005; Kabanova et al., 2018).

There are mounting evidence to show that NK cells contribute to immune responses in the clearance of parasites, removal of infected hepatocytes and infected RBCs through cytotoxicity and ADCC (Burrack et al., 2019). Using an immunodeficient humanized RAG-IL2Rγc–deficient (RICH) mice, a study by Chen et. al., showed that NK cells, rather than macrophages are responsible for regulating the parasite growth *in vivo* by directly interacting with iRBCs (Chen et al., 2014). This study also implicated LFA-1 as key receptor involved in NK cell mediated killing of iRBCs. LFA-1 is a heterodimer composed of CD11a/CD18, also called α1/β2. LFA-1α subunit,αL has two prominent structural features; I domain of 200 aa residues and three EF hand-like domains, which are crucial for ligand binding. Two other integrins expressed on leukocytes; Mac-1 (CD11b/CD18) and p150.95 (CD11c/CD18) share the same CD18 or ß2 integrin subunit and have homologues α subunits (Kolanus et al., 1996). The I domain of these proteins are homologous to motifs in von Wliilibrand factor, cartilage matrix protein, collagen type VI, complement factor C2 and factor B (Huang and Springer, 1995). Structure-function studies on LFA-1 and Mac-1 have implicated their I domains in ligand binding (McDowall et al., 1998). Here, we set out experiments to know whether LFA-1 αI, a 200 aa domain of LFA-1 on NK cells alone recognizes the iRBCs. To do so, LFA-1α1 domain was expressed

as a fusion protein with Fc domain of hIgG and was allowed to bind *P. falciparum*

infected erythrocyte. LFA-1α1-Fc fusion protein bound significantly with iRBCs in

comparison with hIgG control protein.

Having established LFA-1α1 as a key binding domain to iRBCs, we next carried out

studies to identify the parasite derived ligand on iRBC surface that interacts with

LFA-1. Up till now, LFA-1 has been shown to bind human and mouse ICAM-1

(Huang and Springer, 1995). A unique property of LFA-1 is that its binding affinity

can be dynamically modulated by inside-out (intracellular signals triggered by

antigens and chemokines) and outside-in (ligand binding) signaling pathways (Gerard

et al., 2021; Kondo et al., 2022). Conformation changes and membrane clustering of

LFA-1 have been shown to significantly influence its binding affinity (Shi and Shao,

2023; Urlaub et al., 2017). To fish out the parasite ligand on the surface of infected

RBCs for LFA-1, here we cross-linked LFA-1α1-Fc fusion protein with iRBCs,

followed by immunoprecipitation and LC-MS/MS analysis. *Pf*GBP-130 protein was

identified as a prominent LFA-1interacting protein on the iRBC surface. As the name

suggests, *Pf*GBP-130 is a glycophorin binding protein consisting of 11 highly

conserved 50 amino acid repeats and a charged N-terminal region of 225 amino acids

(Kochan et al., 1986). A number of studies have shown that *Pf*GBP-130 is exported

out of iRBC surface and is involved in erythrocyte invasion (Soni et al., 2016). We

further confirmed the iRBCs surface expression of *Pf*GBP-130 protein using anti-

*Pf*GBP antibodies, thus supporting the observation that it probably is involved in

recognition of iRBCs by NK cells. Western blot analysis of LFA-1 αI -Fc bound

iRBCs lysate using anti-*Pf*GBP antibodies further strengthened the hypothesis that

*Pf*GBP-130 on iRBCs surface is the ligand for LFA-1 that helps NK cells to recognize

the iRBCs. To evaluate the specificity of LFA-1 interaction with *Pf*GBP-130 on iRBC

surface, we overexpressed external domain of *Pf*GBP130 as a Fc fusion protein and

evaluated its binding to THP-1 monocytes and primary NK cells. Recombinant

*Pf*GBP-Fc protein efficiently bound to the surface of both cell types. Importantly,

siRNA-mediated knockdown of CD11a subunit LFA-1 significantly reduced this

binding, confirming that *Pf*GBP-130 directly interacts with LFA-1 on immune cells,

thereby advocating GBP-130 as a ligand for LFA-1.

NK cells are predominantly cytolytic lymphocytes, and their primary effector function

relies on the degranulation of cytotoxic molecules. LFA-1 not only mediates tight

adhesion to target cells but also plays a crucial role in early NK cell activation and
degranulation (Barber et al., 2004). To assess whether *Pf*GBP-130 triggers NK cell
activation through LFA-1 engagement, we analyzed the expression of activation
markers CD25 and CD69, and the degranulation marker CD107a, in NK cells co-
cultured with CHO cells expressing the extracellular domain of *Pf*GBP-130.
Compared to mock controls, co-culture with *Pf*GBP-130–expressing CHO cells
significantly upregulated all three markers. Importantly, this activation was abrogated
by the addition of anti-CD11a antibodies, confirming that *Pf*GBP-130 mediates NK
cell activation and degranulation via LFA-1. CD107a (LAMP-1), a lysosomal
membrane protein, serves as a well-established marker of NK cell degranulation
(Urlaub et al., 2017; Ye et al., 2018).

Besides mediating the tight adhesion of NK cells to various target cells, LFA-1–
mediated signaling plays a pivotal role in the polarization of lytic granules towards
the immunological synapse, thereby enhancing NK cell–mediated cytotoxicity
(Barber et al., 2004; Shi and Shao, 2023). To know whether such a toxicity is also
generated as a result of interaction between NK cells and iRBCs, we analyzed the
expression of two activation markers; CD25 and CD 69 and a degranulation marker;
CD107a in NK cells co-cultured with CHO cells overexpressing *Pf*GBP-130 protein
on their surface as described by Ye, W et al. 2018 (Ye et al., 2018). All these three
markers showed significant upregulation in primed NK cells in comparison to naïve
NK cells. This upregulation was specific as the presence of anti-LFA-1 antibodies
reversed these upregulations. To determine whether the activation of primary NK
cells is specifically mediated by direct engagement between LFA-1 on NK cells and
*Pf*GBP-130 on iRBCs at the immunological synapse, we performed targeted co-
culture assays. we co-cultured purified human NK cells with iRBCs and measured
both parasite growth as well as the levels of activation and degranulation markers in
NK cells, in the presence or absences of anti-*Pf*GBP-130 antibodies. The results of
co-cultured study showed that the presence of iRBCs resulted in activation of NK
cells and this resulted in reduced parasitemia. This activation of NK cells was as a
result of *Pf*GBP-130 expression on iRBCs as anti-*Pf*GBP-130 antibodies reduced the
activation of NK cells. These results are in line with a recent study that showed
upregulation of these markers in NK cells co-cultured with iRBCs from human
population (Arora et al., 2018; Ye et al., 2018).

In conclusion, the results presented here identify *Pf*GBP-130 as a novel ligand for LFA-1, in particularly its LFA-1 αI domain on NK cells. This interaction mediates firm adhesion between human NK cells and *Plasmodium falciparum* infected red blood cells resulting in the activation of NK cells. The ensuing activation subsequently triggers NK cells cytotoxic degranulation resulting in targeted killing of iRBCs. Given that NK cells constitute first line of defense against malaria, this molecular mechanism appears to be important in early control of *P. falciparum* infection in particularly by implementing a host directed therapy.

Materials and Method:

Plasmodium culture of 3D7

Plasmodium falciparum 3D7 parasites were maintained in O+ human erythrocytes at 4% hematocrit in RPMI 1640 medium (pH 7.4) supplemented with 50 mg/l hypoxanthine, 5% Albumax II, 2 g/l sodium bicarbonate, and 20 µg/ml gentamycin according to the protocol described by Trager and Jensen (Trager and Jensen, 1976). Cultures were maintained at 37 °C in a gas mixture of 90% N2, 5% CO2, and 5% O2. Parasite synchronization was performed using 5% (w/v) sorbitol.

Cell lines and Primary Human NK cell culture.

THP-1 cells, CHO-K1, and HEK293T cells were cultured in RPMI 1640 (Invitrogen) supplemented with 10% fetal bovine serum (FBS), or in DMEM (for HEK293T) with 10% FBS. Primary human NK cells were isolated from peripheral blood mononuclear cells (PBMCs) by negative selection using the BioLegend NK Cell Isolation Kit, achieving >95% purity.

Fusion protein construct and its expression and purification.

The LFA-1 αI domain was PCR-amplified from THP-1 cDNA and cloned into the pFUSE-hIgG Fc2 vector (InvivoGen). The construct was transfected into CHO-K1 cells using JetPrime (Polyplus). Post-transfection, cells were maintained in IgG-depleted FBS-containing media and later switched to 10% Nuserum medium. After 72 h, the culture supernatant was collected, filtered, and the secreted fusion protein purified using Protein A agarose affinity chromatography (G-Biosciences). Proteins

467 were concentrated using 3 kDa Centricon filters (Merck) and quantified via BCA

468 assay.

Immunoprecipitation and Mass spectrometry.

Schizont-stage *P. falciparum* iRBCs were isolated via Percoll gradient centrifugation

and incubated with 100 µg LFA-1 αI-Fc or human IgG control for 2 h at 37 °C.

Crosslinking was performed using 5 mM DTSSP (Thermo Fisher) for 30 min. Cells

were lysed with IP Native Lysis Buffer and incubated with Protein A magnetic beads

(Thermo Fisher) for 2 h. Beads were washed and proteins eluted with 5 mM DTT.

Eluates were processed for LC-MS/MS analysis. Eluates were processed for LC-

MS/MS analysis. The eluates were first precipitated using 10% trichloroacetic acid

(TCA), 5% acetone, and 5 mM DTT. Samples were air-dried and resuspended in 100

mM ammonium bicarbonate (ABC) buffer containing 8 M urea, then diluted to 1 M

urea with 100 mM ABC buffer. Followed by reduction with 10 mM DTT for 1 h at

room temperature (RT), and alkylation with 40 mM iodoacetamide (Sigma-Aldrich)

for 1 h at RT in the dark. The alkylated eluates were further digested with trypsin

(1:50, wt/wt) at 37 °C for 12–16 h. Peptides were acidified with 0.1% formic acid and

analyzed on an Orbitrap Fusion Lumos mass spectrometer coupled to a Nano-LC

1200 (Thermo Fisher Scientific). Data were processed with Proteome Discoverer v2.4

using the SEQUEST and AMANDA algorithms.

iRBC and Immune cell binding assay.

P. falciparum 3D7 parasites were cultured to 5% parasitemia and synchronized. Ring,

trophozoite, and schizont stages were isolated using Percoll gradient. Approximately

10□ iRBCs from each stage were incubated with 250 nM purified LFA-1 αI-Fc

fusion protein or 250 nM human IgG (hIgG) control for 2 h. After washing, cells were

stained with PE-Texas Red or FITC-conjugated anti-human Fc antibodies (Invitrogen)

for 30 min and analyzed by flow cytometry. Similarly, THP-1 cells and primary

human NK cells, including siRNA-treated populations, were fixed and incubated with

GBP130-Fc (250 nM) or hIgG control (250 nM) in FACS buffer for 2 h. After

washing, cells were stained with PE-Texas Red–conjugated anti-human secondary

antibodies and analysed by flow cytometry.

 ***In vitro Pf*GBP-130/LFA-1 αI-Fc interactions analysis with BLI, *in silico* docking**

and Molecular Dynamic stimulation analysis

The real-time interactions between LFA-1 and its potential binding partner, were studied using Bio-layer interferometry (BLI) on the Sartorius Octet K2 system. In this setup, LFA1 was immobilized onto an AR2G sensor (Octate Amine Reactive Second-Generation biosensor) at a concentration of 100ng, and the unbound protein was washed off using I X Phosphate-buffered Saline (PBS) pH7.4 buffer. To determine the binding affinity, *Pf*GBP was tested at various concentrations. Throughout the experiment, IX PBS pH7.4 was used as both the running and the dissociation buffer, and all working dilutions of *Pf*GBP proteins were prepared in the same buffer. For data normalization, a control sensor with immobilized LFA1 was run in parallel, where only the running buffer (1X PBS) flowed over the sensor. The interaction kinetics, including the association and dissociation curves, were monitored for 800 seconds in a series of increasing concentrations of the *Pf*GBP protein. The data was acquired with Octet BLI discovery 12.2 software, and the KD value was calculated through the Octet Evaluation software, applying a 1:1 binding model for fitting the curve. The experiment was performed at 25°C**.**

In silico docking and simulation, approaches were employed to investigate the molecular interaction between LFA1 with *Pf*GBP. To achieve this, initially, a homology model for PfGBP was generated using Alphafold(Jumper et al., 2021), and the quality of the generated model was assessed with PROCHECK and PyMOL. While for LFA1, its crystal structure (PDB code: 1LFA) was employed in the interaction studies. Next, rigid body protein-protein docking was carried out using the Cluspro2.0 web server (Comeau et al., 2004), a top performer at CAPRI (Critical Assessment of Predicted Interactions) challenges. ClusPro ranked the docked models based on cluster size and energy. The default set of docking parameters was applied, and the top-ranked complexes were selected based on the largest cluster sizes and lowest energy values. The models were further analyzed to examine the interaction interface using PyMOL (PyMOL Molecular Graphics System, v2.1 by Schrödinger, LLC). The final docked model for the LFA1/*Pf*GBP complex was chosen based on significant interactions and the model with the heights buried surface area (BSA).

PDBsum software was utilized to analyze the interactions within the
complex(Laskowski et al., 2018).

To examine the stability, the LFA1/*Pf*GBP complex was subjected to molecular
dynamics (MD) simulations using the Desmond module of Schrödinger software.
Firstly, the Schrödinger protein preparation tool was used to prepare the protein; by
removing the non-bonded water (> 3Å from protein residues), optimizing the H-
bonds, and energy minimizing the final structures using OPLS3. Next, these
complexes were placed in an orthorhombic water box at a buffer distance of 10 Å,
and solvated using the TIP4PEW solvation model while maintaining a NaCl
concentration of 0.15 M to ensure a physiological ionic strength. After solvation, the
complex was subjected to energy minimization using OPLS4 force field parameters
followed by relaxation, and the simulation was performed for 200 ns at NPT
conditions with a frame captured at every 0.2 ns. For determining the trajectories, the
frames were collected and examined through a simulation interaction diagram, which
helped in determining fluctuations (RMSD). The final frame of the simulation was
saved in a PDB file format. Further, PDBePISA (Protein Interfaces, Surfaces, and
Assemblies (PISA) server was employed to analyze the intermolecular
interfaces(Krissinel and Henrick, 2007), including computing buried surface area
(BSA Å) for the complex. PyMol was used to represent these interactions.

Chimeric CHO stable cell line generation.

550
To generate CHO-K1 cells stably expressing *Pf*GBP130, the codon-optimized
synthetic gene encoding the extracellular domain of *Pf*GBP130 was cloned into the
lentiviral transfer plasmid pMSCV-puro. Lentiviral particles were produced by co-
transfecting HEK293T cells with pMSCV-puro-*Pf*GBP130, psPAX2 (packaging
plasmid), and pMD2.G (VSV-G envelope plasmid) using JetPrime transfection
reagent (Polyplus), following the manufacturer's instructions. After 72 h, viral
supernatants were harvested and concentrated using Lentivirus Concentrator Solution
(Takara). CHO-K1 cells were transduced with the concentrated lentivirus for 48 h to
facilitate viral entry and integration of *Pf*GBP 130 gene, and selected with puromycin
(5 µg/mL) for 7 days to enrich for stably transduced cells. Following antibiotic
selection, cells were expanded in fresh culture medium, and PfGBP130 surface
expression was confirmed by fluorescence microscopy.

NK cell activation assay.

*Pf*GBP130-expressing CHO stable cells were trypsinized and counted before being co-incubated with purified primary human NK cells at a 2:1 ratio (CHO: NK) f. Following this co-incubation, a Lipofectamine/PolyI:C complex (100 µg/mL) was added to each well to stimulate a response for 24hrs. To investigate the role of LFA-1, NK cells were pre-incubated with a neutralizing antibody against the LFA-1 receptor for 30 minutes at room temperature before being added to the *Pf*GBP130-CHO cells.

siRNA mediated knockdown of THP1 and NK cells.

siRNA mediated CD11a knockdown was performed in both THP-1 cells and primary human NK cells using Accell SmartPool siRNA (Dharmacon). THP-1 cells were seeded at 1×10 cells/well in 96-well plates in Accell siRNA transfection medium containing 2.5% FBS and 1µM CD11a-targeting siRNA. Primary NK cells (1×10) were suspended in Accell siRNA transfection medium supplemented with 10 ng/mL IL-15 and transfected with 2 µM CD11a siRNA. Cells were incubated for 72 h, after which knockdown efficiency was confirmed by immunoblotting with anti-CD11a antibodies.

Flowcytometry.

The following antibodies were used for staining: anti-human CD3 (biotin, OKT3; Elabsciences), anti CD16 (3G8; Biolegend), anti CD11a (HI111; BioLegend), CD56-PE (5.1H11; Elabsciences), Ultra-LEAF™ Purified Human IgG1 Isotype (Biolegend), Goat anti-Human IgG Alexa Fluor™ 488 (Invitrogen), Goat anti-Human IgG Secondary Antibody, Alexa Fluor™ 594 (Invitrogen), (CD107a (H4A3; Elabsciences), CD25-Alexa Fluor® 488 (BioLegend), and CD69 (FN50; Elabsciences). Purified NK cells, either alone or co-cultured with iRBCs, were stained in 100 µL PBS containing 0.2% BSA and 0.05% sodium azide for 30 min on ice. After washing, stained cells were analyzed on a BD LSR Fortessa X-20 flow cytometer, and data were processed using FlowJo software (BD Biosciences).

NK cell co-culture with RBC and iRBC.

Human NK cells were purified from PBMCs using the BioLegend Human NK Cell Isolation Kit, achieving >95% purity. NK cells were pre-incubated with Fcγ receptor

blocker (anti-human CD16, clone 3G8; 2 µg per 10 cells) for 30 min at room
temperature. Late-stage *P. falciparum* schizont-infected RBCs (iRBCs) were purified
at 0.5% parasitemia. For neutralization experiments, either anti-*Pf*GBP130 antibody
or isotype control IgG1 was added to the culture wells for 30 min prior to NK cell
addition. Pre-treated NK cells were then co-cultured with iRBCs or uninfected RBCs
at a 10:1 ratio (NK: iRBC or NK: RBC) for 48 h. For activation marker analysis,
cells were collected and stained with relevant antibodies. For parasitemia assessment,
co-cultures were maintained for 96 h. Cells were stained with Hoechst 33342
(20 µg/mL) and anti-human CD56-PE (BioLegend) for 15 min, washed, and
analyzed using a BD LSR Fortessa X-20 flow cytometer.
607

608 **Statistical Analysis**
609 Data are presented as mean ± SEM. Differences between groups were analyzed via
610 Student *t* test. A P value <0.05 was considered statistically significant. All
611 calculations were performed using GraphPad Prism 10 software package.
612

Declaration of competing interest
The authors have no competing interests to declare.

Ethical Approval
The study was conducted according to the guidelines of the Declaration of Helsinki
and approved by the International Centre for Genetic Engineering and Biotechnology
(ICGEB)'s Scientific Ethical Review Unit and the Institutional Animal Ethics
Committee.

Funding and acknowledgements
The research work in PMs and AMs is supported by the Flagship Project (BT/IC-
06/003/91), DBT-Wellcome; Team Science Grant (WTA/24/006); and JC Bose Grant
(DST/20/015) from the Department of Science and Technology, Govt. of India. The
funders had no role in the design of the study; the collection, analysis, and
interpretation of the data; or the writing of the manuscript. We also thank the Rotary
Blood Bank (India) for providing human red blood cells. We also acknowledge the
Immunobiology group at ICGEB for providing infrastructure and insightful
suggestion for the completion of this study.

Author Contributions

O.M., R.D., performed the research experiments. S.S.S., contributed expertise and
necessary tools for LC–MS studies. G.P., and A.P., performed the protein
purification, antibody generation and immunolocalization assays related to *Pf*GBP.
P.K., performed the BLI related experiments. N.P., M.A., M.M.I. and P.A.,
maintained *Plasmodium* Culture and provided assistance in immunolocalization
studies. P.M., D.K., and A.M., designed the study, interpreted the data and wrote the
manuscript; P.M., A.M., D.K., O.M., R.D., and A.P., contributed to the experimental
design, data interpretation, and manuscript preparation.

References

Arora, G., G.T. Hart, J. Manzella-Lapeira, J.Y. Doritchamou, D.L. Narum, L.M. Thomas, J. Brzostowski, S. Rajagopalan, O.K. Doumbo, B. Traore, L.H. Miller, S.K. Pierce, P.E. Duffy, P.D. Crompton, S.A. Desai, and E.O. Long. 2018. NK cells inhibit Plasmodium falciparum growth in red blood cells via antibody-dependent cellular cytotoxicity. *Elife* 7:

Artavanis-Tsakonas, K., and E.M. Riley. 2002. Innate immune response to malaria: rapid induction of IFN-gamma from human NK cells by live Plasmodium falciparum-infected erythrocytes. *J Immunol* 169:2956-2963.

Barber, D.F., M. Faure, and E.O. Long. 2004. LFA-1 contributes an early signal for NK cell cytotoxicity. *J Immunol* 173:3653-3659.

Bryceson, Y.T., M.E. March, D.F. Barber, H.G. Ljunggren, and E.O. Long. 2005. Cytolytic granule polarization and degranulation controlled by different receptors in resting NK cells. *J Exp Med* 202:1001-1012.

Burrack, K.S., G.T. Hart, and S.E. Hamilton. 2019. Contributions of natural killer cells to the immune response against Plasmodium. *Malar J* 18:321.

Cerwenka, A., and L.L. Lanier. 2001. Natural killer cells, viruses and cancer. *Nat Rev Immunol* 1:41-49.

Chen, Q., A. Amaladoss, W. Ye, M. Liu, S. Dummler, F. Kong, L.H. Wong, H.L. Loo, E. Loh, S.Q. Tan, T.C. Tan, K.T. Chang, M. Dao, S. Suresh, P.R. Preiser, and J. Chen. 2014. Human natural killer cells control Plasmodium falciparum infection by eliminating infected red blood cells. *Proc Natl Acad Sci U S A* 111:1479-1484.

Comeau, S.R., D.W. Gatchell, S. Vajda, and C.J. Camacho. 2004. ClusPro: a fully automated algorithm for protein-protein docking. *Nucleic Acids Res* 32:W96-99.

Gerard, A., A.P. Cope, C. Kemper, R. Alon, and R. Kochl. 2021. LFA-1 in T cell priming, differentiation, and effector functions. *Trends Immunol* 42:706-722.

Hermsen, C.C., Y. Konijnenberg, L. Mulder, C. Loe, M. van Deuren, J.W. van der Meer, G.J. van Mierlo, W.M. Eling, C.E. Hack, and R.W. Sauerwein. 2003. Circulating concentrations of soluble granzyme A and B increase during natural and experimental Plasmodium falciparum infections. *Clin Exp Immunol* 132:467-472.

Huang, C., and T.A. Springer. 1995. A binding interface on the I domain of lymphocyte function-associated antigen-1 (LFA-1) required for specific interaction with intercellular adhesion molecule 1 (ICAM-1). *J Biol Chem* 270:19008-19016.

Jumper, J., R. Evans, A. Pritzel, T. Green, M. Figurnov, O. Ronneberger, K.
Tunyasuvunakool, R. Bates, A. Zidek, A. Potapenko, A. Bridgland, C. Meyer,
S.A.A. Kohl, A.J. Ballard, A. Cowie, B. Romera-Paredes, S. Nikolov, R. Jain,
J. Adler, T. Back, S. Petersen, D. Reiman, E. Clancy, M. Zielinski, M.
Steinegger, M. Pacholska, T. Berghammer, S. Bodenstein, D. Silver, O.
Vinyals, A.W. Senior, K. Kavukcuoglu, P. Kohli, and D. Hassabis. 2021.
Highly accurate protein structure prediction with AlphaFold. *Nature* 596:583-
589.
Kabanova, A., V. Zurli, and C.T. Baldari. 2018. Signals Controlling Lytic Granule
Polarization at the Cytotoxic Immune Synapse. *Front Immunol* 9:307.
Karre, K. 2002. NK cells, MHC class I molecules and the missing self. *Scand J*
Immunol 55:221-228.
Kochan, J., M. Perkins, and J.V. Ravetch. 1986. A tandemly repeated sequence
determines the binding domain for an erythrocyte receptor binding protein of
P. falciparum. *Cell* 44:689-696.
Kolanus, W., W. Nagel, B. Schiller, L. Zeitlmann, S. Godar, H. Stockinger, and B.
Seed. 1996. Alpha L beta 2 integrin/LFA-1 binding to ICAM-1 induced by
cytohesin-1, a cytoplasmic regulatory molecule. *Cell* 86:233-242.
Kondo, N., Y. Ueda, and T. Kinashi. 2022. LFA1 Activation: Insights from a Single-
Molecule Approach. *Cells* 11:
Korbel, D.S., K.C. Newman, C.R. Almeida, D.M. Davis, and E.M. Riley. 2005.
Heterogeneous human NK cell responses to Plasmodium falciparum-infected
erythrocytes. *J Immunol* 175:7466-7473.
Krissinel, E., and K. Henrick. 2007. Inference of macromolecular assemblies from
crystalline state. *J Mol Biol* 372:774-797.
Laskowski, R.A., J. Jablonska, L. Pravda, R.S. Varekova, and J.M. Thornton. 2018.
PDBsum: Structural summaries of PDB entries. *Protein Sci* 27:129-134.
McDowall, A., B. Leitinger, P. Stanley, P.A. Bates, A.M. Randi, and N. Hogg. 1998.
The I domain of integrin leukocyte function-associated antigen-1 is involved
in a conformational change leading to high affinity binding to ligand
intercellular adhesion molecule 1 (ICAM-1). *J Biol Chem* 273:27396-27403.
Ojo-Amaize, E.A., L.S. Salimonu, A.I. Williams, O.A. Akinwolere, R. Shabo, G.V.
Alm, and H. Wigzell. 1981. Positive correlation between degree of
parasitemia, interferon titers, and natural killer cell activity in Plasmodium
falciparum-infected children. *J Immunol* 127:2296-2300.
Qu, A., and D.J. Leahy. 1995. Crystal structure of the I-domain from the
CD11a/CD18 (LFA-1, alpha L beta 2) integrin. *Proc Natl Acad Sci U S A*
92:10277-10281.
Shi, H., and B. Shao. 2023. LFA-1 Activation in T-Cell Migration and Immunological
Synapse Formation. *Cells* 12:
Soni, R., D. Sharma, and T.K. Bhatt. 2016. Plasmodium falciparum Secretome in
Erythrocyte and Beyond. *Front Microbiol* 7:194.
Trager, W., and J.B. Jensen. 1976. Human malaria parasites in continuous culture.
Science 193:673-675.
Urlaub, D., K. Hofer, M.L. Muller, and C. Watzl. 2017. LFA-1 Activation in NK
Cells and Their Subsets: Influence of Receptors, Maturation, and Cytokine
Stimulation. *J Immunol* 198:1944-1951.
Ye, W., M. Chew, J. Hou, F. Lai, S.J. Leopold, H.L. Loo, A. Ghose, A.K. Dutta, Q.
Chen, E.E. Ooi, N.J. White, A.M. Dondorp, P. Preiser, and J. Chen. 2018.

Microvesicles from malaria-infected red blood cells activate natural killer cells
via MDA5 pathway. *PLoS Pathog* 14:e1007298.

Figure legends.

Figure 1. The αI domain of Lymphocyte Function-Associated Antigen-1 (LFA-1) binds to *Plasmodium falciparum*–infected erythrocytes. (**A**) Schematic representation of full-length of CD11a, a subunit of LFA-1 and the recombinant construct encoding the C-terminal Fc-tagged αI domain of LFA-1. (**B**) Expression and purification of recombinant LFA-1 αI-Fc fusion protein. The αI domain of LFA-1 was cloned into the pFUSE-hIgG1-Fc2 vector and expressed in CHO-K1 cells. The fusion protein was purified from culture supernatants using Protein-A affinity chromatography. (**i**) SDS-PAGE analysis of the purified protein revealed a prominent Coomassie-stained band at ~45 kDa, corresponding to the expected molecular weight of the LFA-1 αI-Fc fusion protein. (**ii**) Western blot analysis using anti-human IgG antibody confirmed the identity of the fusion protein. (**C**) Binding of LFA-1 αI-Fc (250nM) fusion protein to *P. falciparum*–infected red blood cells. Flow cytometry analysis using PE-Texas Red–conjugated anti-human IgG antibody demonstrated specific binding of the LFA-1 αI-Fc protein to ring, trophozoite, and schizont stages of iRBCs, indicating interaction across all major asexual blood stages.

Figure 2. *Pf*GBP-130 on the surface of *Plasmodium falciparum*–infected erythrocytes bind the LFA-1 αI domain.

(**A**) Identification of *Pf*GBP-130 as an interacting partner of LFA-1 αI-Fc. LC-MS/MS analysis of immunoprecipitates from *P. falciparum*–infected erythrocyte lysates pulled down with LFA-1 αI-Fc fusion protein revealed *Pf*GBP-130 (PF3D7_1016300) as a major interacting protein. The table summarizes proteins specifically enriched in the LFA-1 αI-Fc pull-down compared to an hIgG1 isotype

control, with *Pf*GBP-130 showing high peptide coverage and spectral abundance, indicating a specific and robust interaction. (**B**) Characterization and localization of *Pf*GBP-130. (**i**) Schematic representation of the domain organization of *Pf*GBP-130 and the N-terminal fragment (amino acids 69–270) that was expressed in *E. coli* (termed *Pf*GBP-130-N). (**ii**) SDS-PAGE and western blot analysis of purified *Pf*GBP-130-N using anti-rabbit *Pf*GBP-130 antibodies. A prominent band at ~30 kDa corresponds to the expected molecular weight of the recombinant fragment. (**iii**) Immunofluorescence assay (IFA) demonstrating surface localization of *Pf*GBP-130 on trophozoite-stage iRBCs using anti-*Pf*GBP-130 antibodies. *Pf*GBP-130 (green) partially colocalizes with Band 3 (red), a known erythrocyte membrane marker. Nuclei were stained with DAPI (blue), confirming surface expression. (**iv**) Western blot analysis of iRBC lysate and of immunoprecipitate of LFA1 αI-Fc bound iRBC (figure 2A) using anti-rabbit GBP-130 antibody. Lane1, shows the presence of *Pf*GBP-130 in the immunoprecipitate. The ~130kDa band is notably absent in the control IP eluate where no LFA-1 αI-Fc, was bound to iRBC, demonstrating the specificity of the LFA-1 αI-Fc and *Pf*GBP-130 interaction. Lane 2 shows the detection of native *Pf*GBP-130 as a ~110 kDa protein in trophozoite stage *P. falciparum* lysate, consistent with its predicted molecular weight. (**C**) Biophysical and computational validation of *Pf*GBP-130 and LFA-1 αI interaction.

(**i**) Biolayer interferometry (BLI) was used to assess real-time binding between *Pf*GBP-130-N and LFA-1 αI-Fc. Sensorgrams at increasing concentrations of *Pf*GBP-130-N demonstrate dose-dependent binding kinetics. (**ii**) *In silico* docking analysis showing the energy-minimized complex of *Pf*GBP-130 (salmon) and LFA-1 αI domain (green) generated using ClusPro 2.0. Representative hydrogen bonds and interacting residues are shown as sticks. Visualizations were prepared using PyMOL. (**iii**) Molecular dynamics (MD) simulation of the *Pf*GBP-130/LFA-1 αI complex. The graph depicts the root mean square deviation (RMSD) over time, confirming structural stability of the protein–protein complex.

Figure 3: Specificity of interaction between LFA-1 on Primary NK cells and THP-1 cells with *Pf*GBP-130. Primary NK and THP-1 cells treated with LFA-1 siRNAs showed reduced binding to recombinant extracellular domain of *Pf*GBP-130 expressed in CHO K1 cells. (**A**) Schematics representation of *Pf*GBP-130 and its extracellular domain cloned in pFUSE-hIgG1-Fc2 vector for the expression in CHO

K1 cells. The construct comprises the extracellular domain of *Pf*GBP-130 (including

putative LFA-1 binding sites) fused to the Fc region of human IgG1. The Fc fusion

provides stability, solubility, and facilitates detection. **(B)** Interaction of *Pf*GBP-130

ECD-Fc with THP-1 cells. **(i)** Flow cytometric analysis shows strong binding of

*Pf*GBP-130 ECD-Fc to THP-1 cells, compared to an hIgG1 isotype control, indicating

specific interaction. LFA-1 knockdown in THP-1 cells via siRNA treatment

significantly reduced PfGBP-130 ECD-Fc binding, thus confirming the specificity of

interaction between *Pf*GBP130 ECD-Fc and LFA-1 on THP cells. **(ii)** Western blot

analysis using anti-CD11a antibody confirmed the siRNA-mediated knockdown of

CD11a subunit of LFA-1 protein on THP-1 cells. **(C)** Interaction of *Pf*GBP-130

ECD-Fc with primary human NK cells. **(i)** Flow cytometry revealed a marked

increase in *Pf*GBP-130 ECD-Fc binding to NK cells over the isotype control. siRNA-

mediated knockdown of LFA-1 in NK cells led to a notable reduction in *Pf*GBP-130

ECD-Fc binding, confirming that LFA-1 is essential for this interaction. (**ii**) Western

blot analysis using anti-CD11a antibody confirmed the siRNA-mediated knockdown

of CD11a subunit of LFA-1 protein in NK cells, confirming LFA-1 as a critical

receptor for *Pf*GBP130 ECD-Fc binding to both NK as well as THP-1 cells. *

Denotes $p < 0.05$, ** denotes $p < 0.01$, and *** denotes $p < 0.001$.

Figure 4. NK cells activated in the presence of *Pf*GBP-130. (A). (i) Expression of

*Pf*GBP-130 ECD fused with Transferrin membrane domain on the membrane of CHO

K1 cells by infecting the lentiviral vector; pMSCV Puro and its immunofluorescence

analysis using anti-rabbit *Pf*GBP antibody. **(ii)** CHO K1 cells expressing *Pf*GBP-130

ECD bind LFA-1 αI-Fc. Binding of purified LFA-1 αI-Fc to *Pf*GBP-130 ECD

expressing CHO cells was assessed by FACS using an PE-Texas red anti-human IgG

antibody. **(B)** NK cells activation in the presence of CHO K1 cells expressing *Pf*GBP

ECD. Human NK cells were purified (>95%) from fresh PBMC and co-cultured with

CHO K1 cells expressing *Pf*GBP ECD in 2: 1 ratio (20,000 CHO-K1 cells:10,000 NK

cells) and these cells were stimulated with (Poly I:C/lipofectamine 2000) for 24 h. NK

cells were separated from adherent CHO K1 cells and NK cells activation was

assessed by assaying the expression of activation markers (CD69, CD25) and a

degranulation marker; CD107a . NK cells co-cultured with CHO K1 cells expressing

*Pf*GBP-ECD protein showed significant increase in the expression of CD25 and

CD69, as well as CD107a in comparison to the NK cells co-cultured with mock CHO

cells. Addition of anti-CD11a (HI111 clone) antibodies reduced the expression of

both activation and degranulation markers. * Denotes p < 0.05, ** denotes p < 0.01,

and *** denotes p < 0.001.

Figure 5. NK cells activated in the presence of iRBCs, controls parasite infection

(A) Activated human NK cells eliminate iRBCs *in vitro*. Human NK cells when co-

cultured with iRBCs reduce parasitemia significantly after 96h and in the presence of

anti-*Pf*GBP-130 antibodies this reduction in parasitemia was blocked. Presence of

anti-GBP-130 abs resulted in parasitemia simialr to the control when NK cells were

incubated with iRBCs alone. **(B)** NK cells activation in the presence of iRBCs.

Human NK cells were were purified (>95%) from fresh PBMC and co-cultured with

synchronized schizont stage iRBCs at a parasitemia of 0.5% in 10:1 ratio (NK: iRBC)

for 48h. Quantification of activation and degranulation markers was performed after

48 hours. NK cells co-cultured with iRBCs showed significant increase in the

expression of CD25 and CD69, the two activation markers as well as for the

expression of CD107a, a degranulation marker in comparison to the NK cells co-

cultured with RBCs alone. Addition of anti-rabbit PfGBP-130 antibodies reduced the

expression of both activation and degranulation markers in these NK cells in

comparison to rabbit IgG isotype control. * Denotes p < 0.05, ** denotes p < 0.01,

and *** denotes p < 0.001.

Supplementary Figures Legends:

Suppl. Figure 1. Amino acid sequence alignments of "αI domain" of Lymphocyte Function Associated Antigen-1 (LFA-1) with similar Von Willebrand Factor (vWA) domains of human and Mac-1 I-domain showing the similarity and differences among these proteins.

Suppl. Figure 2. (A) Schematic showing Vector map of pFUSE-hIgG1-Fc2 and sites used for cloning the LFA αI domain gene used for the expression of LFA αI-Fc fusion protein. **(B)** Agarose gel electrophoresis showing PCR amplification of DNA corresponding to LFA αI domain and restriction analysis of cloned plasmid; pFUSE-hIgG1- LFA αI -Fc2. **(C)** LC MS/MS analysis of immunoprecipitates from iRBC lysate alone using an hIgG1 isotype control from *P. falciparum*-infected erythrocyte (iRBC) lysates. **(D) (i)** Agarose gels showing the PCR amplification of 600 bp fragment of *PfGBP-130* gene (Pf3D7_1016300). **ii. & iii.** Agarose gel electrophoresis showing restriction analysis of cloned *PfGBP-130* gene fragment in pGEM-T and pET-28b vectors respectively.

Suppl. Figure 3. (A) SDS-PAGE and **(B)** western blot analysis showing the purified *Pf*GBP130-Fc fusion protein expressed in CHO K1 line. Western blot analysis was performed using anti-IgG abs. **(C)** Schematics of cloned *Pf*GBP130-ECD fused to TfR-TM domain in pMSV puro plasmid to produce lentiviral vector. **(D)** Staining of purified primary NK cells using anti-CD3 and anti-CD56 showing purity of NK cells.

Suppl. Figure 4. (A) FACS analysis to show expression of LFA-1 subunit CD11a using anti CD11a antibody on primary NK cells and its expression inhibition on the same cells using Smart pool Accel siRNA corresponding to CD11a of LFA-1. **(B)** FACS analysis to show expression of CD11a subunit of LFA-1 on THP-1 cells and its expression inhibition on the same cells using Smartpool Accel siRNA corresponding to CD11a of LFA-1. Secondary anti-mouse PE-Texas Red was used for detection.

Figure 1

(A). Domain Organization of CD11a (a subunit of LFA-1)

(B).

(i).

Marker (kDa) LFA-1 αI-Fc

150
100
70
55 ← ~ 48kDa
35
25
15
10

(ii).

Marker (kDa) LFA-1 αI-Fc

150
100
70
55
35 ← ~ 48kDa
25
15
10

(C).

iRBC + hIgG1 (dot plot: SSC-A vs LFA-1 αI-Fc Binding (Texas Red), gate 2.59)

iRBC(mixed) + LFA-1 αI-Fc (dot plot, gate 51.3)

Histogram
- Schizont Stage
- Trophozoite Stage
- Ring Stage
- Control (hIgG)

LFA-1 αI-Fc Binding (Texas Red)

Pf–Ring Stage (dot plot, gate 33.1)

Pf–Trophozoite Stage (dot plot, gate 18.9)

Pf–Schizont Stage (dot plot, gate 75.6)

LFA-1 αI-Fc Binding (Texas Red)

Figure 2

(A).

Accession	Description	Coverage [%]	# Peptides	# PSMs	# Unique Peptides	MW [kDa]
PF3D7_1016300	GBP130 protein	61	10	48	10	95.8
PF3D7_1357000	elongation factor 1-alpha	33	13	19	13	48.9
PF3D7_0708400	heat shock protein 90	15	8	14	8	86.1
PF3D7_1462800	glyceraldehyde-3-phosphate dehydrogenase	26	6	8	6	36.6
PF3D7_1105000	histone H4	40	5	9	5	11.4
PF3D7_0818900	heat shock protein 70	9	5	8	5	73.9
PF3D7_0922200	S-adenosylmethionine synthetase	19	5	7	5	44.8
PF3D7_0917900	heat shock protein 70	11	5	6	5	72.3

(B).

Figure 3

(A).

(B).

(i).

(ii).

(C).

(i).

(ii).

Figure 4

(A).

(i).

(ii).

(B).

CD25 Activation

CD69 Activation

CD107a Degranulation

CD25 Activation

CD69 Activation

CD107a Degranulation

Purified NK cells Isolated from Human PBMC

LFA-1
*Pf*GBP ECD

CHO cells Expressing the extracellular domain (ECD) of *Pf*GBP

CHO+mock+NK
CHO+*Pf*GBP+NK
CHO+*Pf*GBP+NK+antiCD11a

Figure 5

(A).

(B).

CD25 Activation
MFI

CD69 Activation
MFI

CD107a Degranulation
MFI

Purified NK cells
Isolated from Human
PBMC

LFA-1
*Pf*GBP

*Plasmodium
falciparum* Infected
erythrocyte

CD25 Activation
Comp-FITC-A

CD69 Activation
PE-A

CD107a Degranulation
Comp-BV650-A

iRBC+NK+antiGBP
iRBC+NK+Isotype
iRBC+NK
RBC+NK