Constraints on the G1/S transition pathway may favor selection of

multicellularity as a passenger phenotype

Selection of multicellularity as a passenger phenotype

Tom Louis Ducrocq, Damien Laporte and Bertrand Daignan-Fornier*

Univ. Bordeaux, CNRS, IBGC, UMR 5095, F-33000 Bordeaux, France

ORCID : TLD (0009-0009-6148-4057); DL (0000-0002-1556-5253); BDF (0000-0003-2352-9700)

*Corresponding author

bertrand.daignan-fornier@u-bordeaux.fr

Abstract

Multicellularity has emerged in the three branches of the tree of life. The formation of simple multicellular entities can either result from cells aggregating or staying together after mitosis. However, it is not yet fully understood how, once formed, these simple multicellular entities could be maintained or even selected for. Here, using the *ace2* yeast snowflake model of simple multicellularity, we aimed at identifying genetic conditions favoring its maintenance. Growth-competition experiments revealed that, while the *ace2* mutation by itself does not provide any fitness advantage or disadvantage, the *ace2* snowflakes were strongly selected when combined to conditions affecting regulators of the G1/S transition of the cell cycle, such as Cln3 or Whi5. We show that this selection results from a faster exit from quiescence of the *ace2* snowflake cells. Importantly, this advantage is not dependent on the multicellular phenotype, but rather on the *ace2* genotype itself. We found that the *ace2* selective advantage in the *cln3* background fully depends on the *KSS1* gene, a target of the Ace2 transcription factor. Finally, we show that phenotypes observed for *ace2* mutants are phenocopied by the *AMN1*[368D] allelic form found in "non-laboratory" yeast strains, hence adding physiological relevance to these observations. Altogether, our results support the hypothesis that simple multicellularity could, in some cases, persist, not because it provides a direct selective advantage due to multicellularity itself, but rather as a "passenger" phenotype that is maintained alongside other selected traits.

Introduction

Multicellularity has evolved repeatedly and independently in distinct lineages within bacteria, archaea and eukaryotes (1). This evolutionary process firstly involves the selection of simple forms of multicellularity such as clusters of cells. Several selective drivers of simple multicellularity have been proposed (2) that are generally associated to changes in geometry allowing access to new resources (faster sedimentation, increased mobility, foraging, dispersal), as well as other advantages associated with size increase (predation avoidance, stress resistance, improved extracellular metabolism) and finally, chimerism: the capacity to combine different cellular phenotypes in a single entity, allowing the establishment of intra-organism division of labor.

Many forms of multicellularity are facultative and their formation is triggered by external signals. One of the best studied case is that of the slime mold *Dictyostelium discoideum*, which cells, in response to resource limitation, aggregate to form a fruiting body that favors dispersal of spores (3,4). Another example comes from *Saccharomyces cerevisiae* cells that, under poor nitrogen conditions, can become highly polarized and stay attached, thereby favoring foraging, a phenomenon known as pseudohyphal growth (5,6). In such cases, multicellularity provides alternative options to the non-proliferation observed in unicellular organisms in response to local nutritional starvation (7). These multicellularity-specific responses are triggered by nutritional cues and, some of them may have evolved by functional recruitment of unicellular pre-existing sensors and pathways. It is hence likely that some actors of nutritional control of cell cycle progression should modulate facultative multicellularity. Indeed, there are several experimental evidences connecting the regulators of G1/S transition in the cell cycle to multicellularity. As described below for several species, this mostly concerns two major regulators, retinoblastoma (RB) and Cyclin D. RB is a negative regulator of E2F, a transcription factor critical for G1/S progression, while cyclin D is a negative regulator of RB (8).

Several experimental evidences connect regulators of G1/S transition, and notably RB, to multicellularity. First, when RB from the simple multicellular volvocine *Gonium pectorale* was

expressed in its close unicellular relative *Chlamydomonas*, the C*hlamydomonas* cells behaved as a

simple multicellular colonial entity (9). This points to an important role for RB in triggering uni- to

multicellular transition. Second, in the aggregative multicellular slime mold, *Dictyostelium discoideum*,

the RB orthologue, RblA, was found to be involved in the stalk *vs* spore cell-fate preference (10). Finally,

another key regulator of G1/S transition, namely Cyclin D, has also been associated to uni- to

multicellular transition in two different yeast species, *Candida albicans* and *Saccharomyces cerevisiae*

(11,12).   Depending on environmental conditions, *Candida albicans* can grow either as a yeast

unicellular planktonic form, or as a multicellular hyphal form (13). The transition from one form to the

other is associated with invasion and virulence. Strikingly, the knock-down of the *C. albicans* cyclin D

orthologue Cln3 resulted in hyphal growth suggesting that the G1 phase of the cell cycle is important

in the transition from uni- to multicellular development in *C. albicans* (11). Similarly, in *S. cerevisiae*

deletion of *CLN3*, the budding yeast cyclin D homolog, enhanced pseudohyphal growth (12).

Interestingly, *CLN3* encodes an inhibitor of Whi5 which is the yeast functional equivalent of RB (13). It

hence seems that, during evolution, uni- to multicellularity transitions have repeatedly used pre-

existing regulators of the G1/S transition, such as RB and cyclin D. These intriguing observations, from

phylogenetically distant organisms, connecting G1/S transition regulators to multicellularity led us to

ask the following question. Could the evolution of the G1/S regulation pathway itself have somehow

favored multicellularity emergence and/or maintenance?

In this work we used *S. cerevisiae* to question the complex relationship between G1/S cell cycle

regulators and multicellularity. We took advantage of the snowflake model (14,15) in which

multicellularity is due to a genetic defect in cell separation caused by the deletion of Ace2, a

transcription factor that control the expression of chitinases and endoglucanases required for cell wall

degradation after cytokinesis completion (16,17). When planktonic *ACE2* and snowflake *ace2* cells

were co-cultured without applied selective pressure and their fitness compared, we observed that

CLN3 deletion, as well as *WHI5* overexpression, favored the *ace2* snowflake form. We showed that exit

from quiescence is the critical stage leading to the *ace2 cln3* snowflakes selection. Importantly we

demonstrated that the *ace2* mutation, but not the snowflake phenotype, was responsible for the *cln3*

faster quiescence exit phenotype. Finally, we show that both the multicellularity and quiescence exit

phenotypes observed for *ace2* mutants are phenocopied by the *AMN1³⁶⁸ᴰ* allele found in "non-

laboratory" yeast strains. Our results point to the possibility that evolution of cell cycle control could

promote the evolution of multicellularity as a passenger phenotype.

Results

The *ace2 cln3* snowflake strain is selected over the *ACE2 cln3* planktonic strain.

In order to investigate the relationship between the G1/S cell cycle regulators and multicellularity, we

used the yeast snowflake model in which the absence of Ace2, a transcription activator of several

chitinase and endoglucanases genes (18–20), results in the lack of separation of the daughter cell from

the mother cell after cytokinesis. It ensues that *ace2* mutant yeasts form cell clusters that grow and

end-up breaking most probably due to mechanical constraints (21). A major advantage of studying

snowflakes rather than inducible forms of multicellularity, such as pseudo-hyphae, is that unicellular

(planktonic, PK) and multicellular (snowflake, SF) yeast, differing only by one mutation in the *ACE2*

gene, can be co-cultivated and their fitness compared. We hence grew isogenic diploid *ACE2/ACE2* and

ace2/ace2 cells of the BY (S288c-derived) background in minimal enriched medium (SD casa WAU).

Cultures of both genotypes were mixed to obtain a 50/50 ratio of *ace2* SF and *ACE2* PK strains, either

at the phenotype level (number of entities observed by microscopy, a SF being an entity with 4 or more

cells, Figure 1-figure supplement 1) or at the cell genotype level (by monitoring the *ace2::kanMX*

disruption using qPCR) (Figure 1A). The mixtures were then repeatedly grown to saturation and diluted

every 2 days, resulting in a total of more than 100 generations. Interestingly, the *ace2* strain showed

no selective advantage or disadvantage over the *ACE2* strain, both at the SF phenotype level (Figure

1B, black bars) and the genotype level (Figure 1C, black bars). This indicates that, under these growth

conditions, the *ace2* mutation and its multicellular phenotype have no effect on fitness. Strikingly,

when similar experiments were performed with *ace2* or *ACE2* strains carrying a deletion of the *CLN3*

gene, a deletion known to lengthen the G1 phase and to significantly increase cell volume (22), the

ace2 snowflake strains rapidly took over the population (red bars, Figure 1B and 1C). A similar result

was obtained when *WHI5,* encoding the major downstream target of *CLN3*, was overexpressed (red

bars, Figure 1D). These results strongly suggest that the observed effect on population fitness takes

place through the *CLN3/WHI5* G1/S transition regulatory pathway. Together our results reveal a robust

phenotypic interaction between the G1/S transition major players, *CLN3* and *WHI5*, and the *ace2*

snowflakes.

ace2 cln3 cells exit faster from quiescence than *ACE2 cln3*

In our competition experimental setup (Figure 1A), cultures alternate phases of active cell division with

stationary phases during which cells are in a quiescent state (Figure 2-figure supplement 1A).

Theoretically, the fitness advantage of the *ace2 cln3* strain over its *ACE2 cln3* isogenic counterpart

could be due to a faster doubling time during exponential growth, to a higher biomass yield of the

population, to a better survival in stationary phase or to a faster exit from stationary phase, each of

these possibilities (schematized Figure 2-figure supplement 1A) acting either alone or in combination.

We tested these four possibilities separately. First, we found that the population doubling times of the

two sets of strains was not significantly different (Figure 2-figure supplement 1B). In addition, *ace2* and

ACE2 cell populations led to similar biomass accumulation indicating that the yield of carbon utilization

was not significantly modified by the *ace2* and/or the *CLN3* mutation (Figure 2-figure supplement 1C).

We then estimated cell survival in stationary phase by staining 2 days-old cells with methylene blue, a

dye that specifically accumulates in dead cells. No significant differences in dead cell percentages were

observed between *ace2* and *ACE2* cells should they be *cln3* or *CLN3* (Figure 2-figure supplement 1D).

From these experiments, we conclude that neither the doubling time, nor the biomass yield, or the cell

death rate could explain the selective advantage of the *ace2 cln3* cells over their *ACE2 cln3* counterpart.

We then compared the time required for *ace2 cln3* and *ACE2 cln3* quiescent-cell populations to exit

stationary phase. We found that the time required for population to resume proliferation after

refeeding, referred to as lag-time, was not significantly different between *ace2 CLN3* and *ACE2 CLN3*

cells (grey and black dots, Figure 2A). In addition, we confirmed that the lag-time was significantly

longer for the *ACE2 cln3* (light red dots) than for *ACE2 CLN3* cell population (black dots) (Figure 2A), as

previously described (23,24). Indeed, after refeeding of stationary phase cultures, *CLN3* transcription

is increased by 5- to 10-fold within five minutes (24). Cln3 is hence likely to play an important role for

quiescent cells to move out from G1 and reenter proliferation. Importantly, this difference was

partially but significantly suppressed in *ace2 cln3* (red dots) cell populations (Figure 2A) suggesting that

the *ace2* mutation genetically suppresses the *cln3* strain lag-time defect. We then re-fed quiescent

cells on microscope agarose pads containing new medium and addressed cell ability to emit a new bud

i.e. to exit quiescence. While the time required by cells to exit quiescence was not significantly different

between *ace2 CLN3* and *ACE2 CLN3* strains (Figure 2-figure supplement 1E), it was significantly lower

for the *ace2 cln3* cells than for the *ACE2 cln3* cells (Figure 2C). Importantly, similar observations were

made with haploid *ace2 cln3* cells (Figure 2D) for which snowflakes are much smaller (Figure 2-figure

supplement 1G-K). Indeed, not only haploid cells display a smaller cell volume than diploid cells, but

their proximal budding pattern make haploid snowflakes much more globular than diploid snowflakes

that are rather elongated (25). This observation confirmed the robustness of the *cln3* phenotype and

its suppression by *ace2*. It also suggests that this phenotype is not dependent on the size of the

snowflake entity. Altogether these results point to quiescence exit as the key step accounting for the

competition outcomes.

A simple prediction from these results is that a competition experiment in co-culture ran under

conditions where the cells are always maintained in proliferation should not result in any fitness

difference between for *ace2 cln3* and *ACE2 cln3* cells. This is what was observed when maintaining

these two strains in proliferation for 94hrs (corresponding to at least 70 generations, Figure 2E, left

panel), by contrast to an equivalent number of generations obtained by alternating proliferation and

stationary phases (Figure 2E, right panel). Reciprocally, as described previously (26), increasing the

duration of stationary phase caused an increase of the lag phase length at the population level (Figure

2F). Importantly, it also increased the fitness advantage of *ace2 cln3* over *ACE2 cln3* cells (red and pink

bars Figure 2G), while there was still no difference between *ace2 CLN3* and *ACE2 CLN3* cells (black and

grey bars Figure 2G). Altogether, our results pinpoint exit from quiescence as the critical stage in which

the *ace2 cln3* snowflakes are positively selected.

The advantage of *ace2 cln3* during quiescence exit is independent of the

snowflake phenotype

Why do *ace2 cln3* cells exit quiescence faster than *ACE2 cln3* cells? This could be due to some specificity

associated with *cln3* mutation that would modify the snowflake properties. Alternatively, it could be

dependent on the specific *ace2 cln3* mutation combination independently of the snowflake

phenotype. To answer this question, we dissociated the snowflake phenotype from the *ace2* mutation

and vice versa.

First, we asked whether the snowflake phenotype was sufficient, independently of the *ace2* mutation,

to rescue the quiescence-exit default associated with *cln3*. Ace2 is a transcription factor that induces

the expression of several chitinases and endoglucanases necessary for the septum digestion and cell

separation. To mimic the *ace2* snowflake phenotype, we constructed a quintuple chitinase and

endoglucanase mutant (*cts1, dse2 dse4, egt2, scw11*), hereafter "quintuple mutant", that results in a

snowflake phenotype comparable to that of the *ace2* mutant (Figure 3-figure supplement 1) while

carrying the wild-type *ACE2* allele. When combined with the *cln3* mutation, the five deletions have no

effect on quiescence exit efficiency (Figure 3A). Accordingly, the *cln3 ace2* strain exited quiescence

much more efficiently than the *cln3* quintuple mutant (Figure 3B). This indicates that the *ace2* mutation

itself, and not multicellularity, suppresses the quiescence exit defect of the *cln3* mutant*.* Finally,

competition experiments in co-cultures initially containing equal proportion of quintuple mutant and

WT cells confirmed that the snowflake phenotype is not sufficient to give the *cln3* mutant a

phenotypical advantage (Figure 3C).

Reciprocally, we asked whether the *ace2* mutation itself, independently of the snowflake phenotype,

could be sufficient to rescue the quiescence-exit default associated with *cln3*. We thus examined

whether planktonic cells obtained in an *ace2* mutant background would still behave as their snowflake

counterparts. To obtain planktonic *ace2* cells, we used sonication to physically break the snowflakes,

before triggering quiescence exit. Mild sonication resulted in an increased proportion of planktonic

cells in the *ace2 cln3* cell population (up to 50%, Figure 4-figure supplement 1A) with no defect in cell

survival (Figure 4-figure supplement 1B). After sonication, isolated cells, small and large cell clusters

displayed a similar quiescence exit efficiency (Figure 4A). Furthermore, the sonicated and non-

sonicated cells had the same quiescence exit efficiency, indicating that sonication *per se* had no

significant impact (Figure 4-figure supplement 1C). Still, after sonication, *cln3 ace2* isolated cells exited

quiescence faster than their *cln3 ACE2* counterpart (Figure 4B). These results show that the snowflake

phenotype is not necessary for the ability of the *ace2 cln3* mutant to exit quiescence faster than *ACE2*

cln3 cells.

Together, our results with the haploid strains, the quintuple mutant and sonication experiments

establish that the *ace2* mutation, but not the snowflake phenotype, is responsible for the *cln3* fast

quiescence exit phenotype.

The *ace2 cln3* fast quiescence exit phenotype is dependent on the Kss1 MAP kinase

As previously mentioned, a delay of *cln3* mutants to exit quiescence was already documented (24).

Why is this phenotype suppressed by the *ace2* deletion? Among Ace2 well-established targets , *AMN1*,

BUD9, CHS7, CTS1, DSE2, DSE4 and *SCW11* are activated, while *KSS1* is down regulated by Ace2 (18).

Four of those, namely *CTS1, DSE2, DSE4* and *SCW11*, are not involved in suppressing the delay

phenotype since they are mutated in the quintuple mutant that, in combination with *cln3*, has the

same delay as a single *cln3* mutant (Figure 3D). We focused on Kss1, a MAP kinase involved in

filamentous growth and mating (27,28), because the Kss1 pathway was previously found to positively

regulate *CLN1,* a G1 cyclin encoding gene that is partially redundant to *CLN3* (29). We hypothesize that

deletion of *ACE2* would result in higher *KSS1* expression (18) that in turn, could lead to higher *CLN1*

expression and thereby be responsible for the *cln3* mutation suppression. We hence tested whether

suppression of *cln3* by *ace2* was dependent on *KSS1* by comparing quiescence exit efficiency of *ace2*

cln3 KSS1 and *ace2 cln3 kss1* cells. Clearly, the knock-out of *KSS1* totally abolished the phenotypic

suppression of *cln3* by *ace2* (Figure 5A). Further, *CLN1* expression upon quiescence exit was

significantly higher in the *ace2 cln3* than in the *ACE2 cln3* strain (Figure 5B), this difference of

expression being entirely abolished in the absence of *KSS1* (Figure 5C). We conclude that the

competitive advantage of the *ace2* mutant over its *ACE2* wild-type counterpart is most likely due to

higher expression of *CLN1* that, at least partially, compensates for the absence of *CLN3*. Importantly,

this transcriptional effect of Ace2 on *CLN1*, *via* Kss1, is independent of the role of Ace2 on

multicellularity.

The *AMN1*[368D] allelic-form found in the wild, phenocopies the *ace2* deletion

In laboratory strains, the Ace2 protein level is constant throughout the cell cycle, while it is not always

the case in non-laboratory strains (25). This is due to the presence of the recessive *amn1[368V]* allelic

form in laboratory strains (hereafter *amn1-368V*), which was probably initially selected to avoid

"unwanted" cell clusters. In non-laboratory strains, the Amn1-368D protein induces the proteolysis of

Ace2 through the ubiquitin proteasome pathway (25). It thus appears that the Amn1/Ace2 tandem has

an important physiological role associated with optional post-mitotic cell separation. We first observed

that expression of *AMN1[368D]* in the BY background resulted in a snowflake phenotype comparable to

that of *ace2* mutant strains (Figure 6-figure supplement 1). We then asked whether *AMN1[368D]* allelic

form could suppress the quiescence exit delay of the *cln3* mutant. Indeed, the *cln3 AMN1[368D]* strain

exited quiescence much more rapidly than the *cln3 amn1-368V* strain (Figure 6A). Finally, we found

that the *AMN1[368D]* strain was strongly enriched in competition experiments when associated with a

cln3 mutation (Figure 6B) or with *WHI5* overexpression (Figure 6C). We conclude that, in haploid yeast,

the *AMN1[368D]* allelic-form phenocopies both the multicellularity and the suppression phenotypes of

the *ace2* mutant.

Discussion

In this work, we took advantage of the snowflake model of multicellularity that is genetically

determined, by *ace2* or *AMN1* and is hence not triggered by external cues, as it is often the case for

facultative multicellularity. This experimental design allows direct fitness measurements between uni-

and multicellular entities. Importantly, in a wild-type background, multicellularity was neither selected

nor counter-selected, even following a substantial period of co-culture. This establishes that simple

multicellular cluster formation should not always come at a cost, contrary to the general assumption

in the literature (30,31). Such a neutral effect of multicellularity on fitness could easily result in

phenotypical *aller-retours* between unicellularity and multicellularity during evolution. In fact, such

aller-retours are observed during the life cycle of non-laboratory strains in which diploids are

planktonic while haploid strains form clusters (32). Here we show that the evolution of the cell cycle

regulation may interact with the multicellular phenotype. We revealed a genetic connection between

major G1/S regulators and *AMN1* and *ACE2*, key actors modulating cell cohesion at the end of mitosis

(Figure 7). Interestingly, this connection operates differently in laboratory strains carrying the *amn1-*

368V allele (Figure 7A) and strains carrying the *AMN1^368D* allele (Figure 7B). We established that this

genetic interaction is due to the pleiotropy of the *ace2* mutation, affecting not only the chitinases and

endoglucanases that are critical for septum digestion and cell separation, but also the Kss1 MAP kinase

which contributes to the activation of Cln1, one of the G1 cyclins. This phenotypical link between *CLN1*

expression and multicellularity, achieved by Ace2 function, is particularly important during quiescence

exit, a situation that is likely to happen very often under ecological conditions, such as boom and bust

cycles when moving from rotting fruit to rotting fruit. We propose that in some instances,

multicellularity may have evolved as a side effect, named here "passenger phenotype", of another

phenotype that was the one selected (Figure 7B). Since transcription factors often have multiple

targets, an alteration in their expression or function leading to the selection of a phenotype due to one

of the targets could easily lead to the selection of additional phenotypes due to other targets. This

could be a common feature involved in evolutionary processes.

Connections between G1/S regulators and multicellularity have been repeatedly observed in

phylogenetically distant species (see introduction section), suggesting a propitious genetic interplay

for co-evolution. Olson and co-workers have shown that RB could have played a role in early steps of

multicellularity evolution *via* a cooption mechanism favoring the appearance of simple multicellular

entities (9). Our work further supports a role for G1/S regulators in evolution of multicellularity, but

the mechanism appears different and affects selection of multicellularity rather than its initial

appearance. We show that the *ace2* or *AMN1^368D* allelic forms are sufficient to counteract a negative

fitness effect of the overexpression of the RB functional orthologue Whi5 (Figure 1D, Figure 6F).

Importantly we have shown that this phenotypic suppression is independent of the multicellular

phenotype associated with the *ace2* or *AMN1^368D* allelic forms. This observation led us to propose that

some regulatory effects on the G1/S transition could be alleviated by Amn1 degradation of Ace2 and

thereby connected to multicellularity. In this hypothesis the selective pressure applies on G1/S

transition efficiency and not on multicellularity itself. It is hence different from the cooption hypothesis

in which the RB protein acquired new functionalities leading to multicellularity that would result in

some directly selectable fitness advantage. Both scenarios imply co-evolution of cell cycle regulation

and multicellularity, which could therefore be a much more general feature than first thought. It could

indeed be a case of convergent evolution as suggested by the fact that it may involve distinct

mechanisms.

More generally, our results prompted us to reconsider whether simple multicellularity, which

appeared multiple times in evolution, was always selected for the new physical and physiological

properties it provides (2). Alternatively, it is possible that multicellularity was initially selected as a

byproduct, allowing it to be maintained although not necessarily increasing fitness as such. This would

contribute to disconnect the initial selection of simple multicellularity from the selection of further

associated advantages, such as division of labor, that would require time and necessitate that the

multicellularity phenotype would not be counter-selected in the first place. Such a passenger selection

of multicellularity would make it independent of physical constraints such as size or shape that could

restrain its further evolution. The work presented here provides a proof of concept that such a scenario

is plausible and may have contributed to the multiple appearance of multicellularity in the tree of life.

Materials and methods

Yeast media and strains, plasmids and oligonucleotides

Yeast cells were grown in liquid or solid media at 30 °C. SD is a synthetic minimal medium containing

0.5% ammonium sulfate, 0.67% yeast nitrogen base (Difco), 2% glucose. SDcasaWAU is an SD medium

supplemented with 0.2% casamino acids (Difco) tryptophan (0.2 mM), adenine (0.3 mM) and uracil

(0.3 mM). Unless specified, cells were grown in SDCasaWAU medium. When indicated, medium was

depleted in uracil, resulting in a medium named SDCasaWA, allowing the selection of cells expressing

a vector carrying *URA3* marker. SC medium was prepared as described by Sherman (33) and was

supplemented by histidine (0.06 mM), leucine (0.5 mM), lysine (0.07 mM), tryptophan (0.2 mM),

adenine (0.3 mM), and uracil (0.3 mM). The medium obtained by the depletion of both uracil and

leucine, called SC-U-L medium, allows the selection of cells expressing both *URA3* and *LEU2* markers.

The YPD medium contained 1% yeast extract, 2% peptone, and 2% glucose.

Yeast strains (listed in Table 1) are derived from both FY4 and FY5 prototrophic strains and from

disrupted strains isogenic to BY4741 or BY4742 purchased from Euroscarf (Frankfurt, Germany). Strains

described in Table 1 were either obtained by crossing, sporulation, and micromanipulation of meiosis

progeny or by transformation of cells with a DNA fragment of interest (plasmid, PCR product). Yeast

strains were transformed using the lithium acetate method (34).

Table 1: Strains

Strain name	Genotype
FY4	
Y12580, Y12581, Y12582	*cln3::URA3/cln3::URA3 ura3Δ0/ura3Δ0 leu2Δ0/leu2Δ0*
Y12584, Y12585, Y12586	*ace2::KanMX4/ace2::KanMX4 cln3::URA3/cln3::URA3 ura3Δ0/ura3Δ0 leu2Δ0/leu2Δ0*
Y12606, Y12607, Y12608	*leu2Δ0/leu2Δ0*
Y12973, Y12974, Y12975	*ace2::KanMX4/ace2::KanMX4 leu2Δ0/leu2Δ0*
Y12575 + p5314	*ace2::KanMX4/ace2::KanMX4 ura3Δ0/ura3Δ0* carrying empty plasmid 2μ
(Y11418xY12599) + p5314	*ura3Δ0/ura3Δ0* carrying empty plasmid 2μ
Y12575 + p4726	*ace2::KanMX4/ace2::KanMX4 ura3Δ0/ura3Δ0* carrying plasmid 2μ *WHI5*
(Y11418xY12599) + p4726	*ura3Δ0/ura3Δ0* carrying plasmid 2μ *WHI5*
Y12512	*ace2::KanMX4/ace2::KanMX4 HTB1-3xCFP-LEU2/HTB1-3xCFP-LEU2 ura3Δ0/ura3Δ0 his3Δ1/HIS3*
Y12565, Y12566, Y12958	*ace2::KanMX4 cln3::URA3 ura3Δ0 leu2Δ0*
Y12567, Y12568, Y12959	*cln3::URA3 ura3Δ0 leu2Δ0*

Y12408, Y12595, Y12596	*ace2::KanMX4 leu2Δ0*
Y11417, Y12571	*leu2Δ0*
Y12709, Y12710, Y12711	*cln3::URA3/cln3::URA3 ura3Δ0/ura3Δ0 leu2Δ0/leu2Δ0*
Y12716, Y12717, Y12718	*dse2::KanMX4/dse2::KanMX4 cts1::KanMX4/cts1::KanMX4 scw11::KanMX4/scw11::KanMX4 egt2::KanMX4/egt2::KanMX4 dse4::KanMX4/dse4::KanMX4 cln3::URA3/cln3::URA3 ura3Δ0/ura3Δ0 leu2Δ0/leu2Δ0*
Y12976, Y12977, Y12978	*dse2::KanMX4/dse2::KanMX4 cts1::KanMX4/cts1::KanMX4 scw11::KanMX4/scw11::KanMX4 egt2::KanMX4/egt2::KanMX4 dse4::KanMX4/dse4::KanMX4 leu2Δ0/leu2Δ0*
Y12967, Y12968	*kss1::KanMX4 ace2::KanMX4 cln3::URA3 ura3Δ0 leu2Δ0*
Y12969, Y12970	*kss1::KanMX4 cln3::URA3 ura3Δ0 leu2Δ0*
Y12838, Y12839, Y12840	*AMN1368D-URA3-AMN1368V ura3Δ0 leu2Δ0*
Y12935, Y12936, Y12937	*AMN1368V-URA3-AMN1368V ura3Δ0 leu2Δ0*
Y12942, Y12943, Y12944	*AMN1368D-URA3-AMN1368V cln3::KanR ura3Δ0 leu2Δ0 his3Δ1*
Y12946, Y12947, Y12949	*AMN1368V-URA3-AMN1368V cln3::KanR ura3Δ0 leu2Δ0 his3Δ1*
Y12838 + p338	*AMN1368D-URA3-AMN1368V ura3Δ0 leu2Δ0* carrying empty plasmid 2µ
Y12935 + p6053	*AMN1368D-URA3-AMN1368V ura3Δ0 leu2Δ0* carrying plasmid 2µ *WHI5*
Y12838 + p338	*AMN1368D-URA3-AMN1368V ura3Δ0 leu2Δ0* carrying empty plasmid 2µ
Y12935 + p6053	*AMN1368D-URA3-AMN1368V ura3Δ0 leu2Δ0* carrying plasmid 2µ *WHI5*
GN-1C	*HO:NatR AMN1368D FLO8+*

All plasmids and oligonucleotides are listed in Table 2 and Table 3, respectively. To construct the

plasmids allowing the overexpression of *WHI5* in yeast strains, the open reading frame of the *WHI5*

gene, driven by its own promoter, was amplified using the genomic DNA of the FY4 strain as a template.

The PCR product was then cloned into two 2µ vectors, YEpLac195 and YEpLac181, carrying respectively

URA3 and *LEU2* markers. Two plasmids carrying respectively two different alleles of the *AMN1* gene

were also constructed. The open reading frame of the *AMN1* gene was amplified using either the FY4

strain genomic DNA (*amn1^{368V}* PCR product) or the GN-1C strain genomic DNA (*AMN1^{368D}* PCR product).

The GN-1C strain was a generous gift from Pr. Marullo. The PCR products were cloned into YIpLac211,

that allowed integration of the cloned *AMN1* allele at the yeast *AMN1* locus.

S2 Table: Plasmids

Plasmid Name	Description	
p5314	YEpLac195: 2µ *URA3* AmpR	(35)
p4726	*WHI5* in YEpLac195	lab collection
p338	YEpLac181 : 2µ *LEU2* AmpR	(35)
p6053	*WHI5* in YEpLac181	This study
p6164	YIpLac211 : integrative plasmid *URA3* AmpR	(35)
p6165	*AMN1-368D* in YIpLac211	This study
p6167	*AMN1-368V* in YIpLac211	This study

323 **S3 Table: Oligonucleotides**

Oligonucleotide name	Sequence 5'-3'
qPCR locus ACE2 -26 forward	GGACCAAAAACGGTGTTAATACAATC
qPCR KanMX4 specific reverse	CTGGCGCGCCTTAATTAACC
qPCR ACT1 forward (coding)	CCCCAGAAGAACACCCTGTTC
qPCR ACT1 reverse (coding)	CGTAGAAGGCTGGAACGTTG
qPCR CLN1 coding forward	ATCGATCAGCAACCGGAGAT
qPCR CLN1 coding reverse	AACCTGACAGCGTGGAAGAA
KanB reverse	CTGCAGCGAGGAGCCGTAAT
ACE2 promoter forward	CGTCACTCCATTAGAATCCC
CLN3 promoter forward	TCCTCATTCGGTTTAACTCC
CLN3 terminator reverse	TGACTAGAGGAAGTAAGGAG
DSE2 promoter forward	CAGTAGAGCTAACCACAGTC
CTS1 promoter forward	ACTGTCGCTCGTTTCACAAC
SCW11 promoter forward	CAGTTACGCAACAAAGACAG
DSE4 promoter forward	ACTACAAGCGAGGGTAAAGG
EGT2 promoter forward	GATGCTGGTTTGATGCTAAG
CLN3 promoter -500 forward	TGAGGAAAGAGGACTATACC
CLN3 terminator +500 reverse	TAGGTAGCGATGAAGATTGG
URA3 promoter -500 forward	TCATCATCTCATGGATCTGC
URA3 terminator +500 reverse	TACGCCAGTACACCTTATCG
CLN1 promoter -200 forward	CCAAGGAGTTCTTCGTTCGC
CLN1 terminator +200 reverse	TCGCGTCATCTTTTCCGTTC
AMN1 promoter -75 forward	GTTTAATATCCATCCATTCC
AMN1 +459 forward	CCTTTGACTGCTCAACATCAG
AMN1 +882 forward	CATCAAGTAACACAACCAG
AMN1 +1269 forward	GGTTGTGATGTTGATGATG
AMN1 terminator +169 reverse	GTTTCTTCGGCCCTTCTGGA
AMN1 +1269 reverse	CATCATCAACATCACAACC

AMN1 +882 reverse	CTGGTTGTGTTACTTGATG
AMN1 +376 reverse	GAAAGACTGGATGCAGAAAC
Mat a and alpha forward	AGTCACATCAAGATCGTTTATGG
Mat a specific reverse	CTCCACTTCAAGTAAGAGTTTG
Mat alpha specific reverse	GCACGGAATATGGGACTACTTCG
WHI5 BamHI forward	CGCGGATCCCAAATCGGATATGAGCAGCTG
WHI5 PstI reverse	ACGTCTGCAGGCTCGAGGCGATCTGTCGC
M13 reverse	CAGGAAACAGCTATGACC
AMN1 BamHI forward	GTTGGATCCATGAAACTAGAACGCG
AMN1 PstI reverse	AACCTGCAGCTAGTCCACATTATTCTCTATTTCG

Description of a Snowflake population

Snowflake populations or co-culture between Snowflake and Planktonic strains were observed in a

Kova Glasstic slide (Kova international, 87144F). The proportion of the Snowflake phenotype in these

populations was determined by counting entities formed by four or more cells. To characterize the size

of Snowflakes within a population, the population was observed between slide and coverslip separated

by a layer of double-sided adhesive tape. The longest segment of the Snowflake entity was measured,

as well as the longest segment perpendicular to the first segment and lying in the same Z plane.

Competitions experiment in co-culture

Snowflake strains (either *ace2* or quintuple mutant or *AMN1*^{368D}) and planktonic strains were grown in

monocultures to stationary phase, for 2 days unless specified. Snowflake and planktonic strains were

then mixed in indicated proportion and co-cultured. Unless specified, every 2 days, co-cultures, after

they reached stationary phase, were diluted by a factor of 1000 into fresh liquid medium (10µl in 10ml).

To set up the co-cultures for which the proportion of Snowflake phenotype was monitored, the entity

(Snowflake entity or single-cell entity) concentrations in each of the monocultures were determined.

The volumes of the 2 monocultures in the mix were accordingly determined and then adjusted to

obtain 50% of Snowflake entity in the mix. The proportion of Snowflakes in the co-culture was

determined every two days before dilution in fresh medium.

To set up the co-cultures for which the proportion of *ace2* genotype was monitored, the two strains

were mixed to obtain a 50/50 cell ratio. Every 2 days, the genomic DNA of the stationary phase co-

cultures was extracted by phenol-chloroform protocol. Quantitative PCR amplifications were then

performed with GoTaq® qPCR Master Mix (Promega, A6001), using 50ng of DNA extracted from co-

cultures. The first set of primers was designed to amplify the *ace2::KanMX4* locus with the first primer

targeting the region directly upstream of the ATG of the *ACE2* gene, and the second primer targeting

a region specific to the *KanMX4* marker (Supplemental Table 3). The second set of primers targeted a

region of *ACT1* gene (Supplemental Table 3).

For competitions in which the co-cultures were maintained in proliferation phase for 94 hours, the co-

cultures were diluted every 8 hours to ensure that the OD_{600nm} was always maintained below 0.5.

Population measurement

To determine the population doubling time of yeast strains, cells were kept in exponential proliferation

phase ($OD_{600nm} \leq 1.0$) for at least 24 hours by successive dilutions before any measurement. Population

doubling time was then determined by following the optical density at 600 nm during 5 hours on at

least three independent exponential growth culture. Population doubling time was then determined

by exponential growth regression of the proliferation curves using GraphPad Prism 8 (GraphPad

Software, Inc. La Jolla, USA).

To determine cell viability, the cells were stained with a solution containing 0.2% of methylene blue

(Sigma-Aldrich, Saint Louis, MI, USA) and 2% sodium citrate pH7. After 5 minutes of incubation, the

proportion of blue-stained dead cells was scored manually.

The biomass yield of cultures was determined by filtration of 2-day stationary phase cultures on 0.45

μm polyamide pre-weighted filters (25006–47-N; Sartorius). The filter was washed twice with 10 mL of

miliQ water and then dried with successive 30 seconds cycles in a microwave oven at 800 W until the

mass of the filter and cells was constant.

Cell volume was measured by microscopy using the spheroid formula: $4/3*\pi*a^2*c$, where a and c are

the equatorial and polar radius respectively.

Refeeding of stationary phase cultures

Unless the age of cultures is otherwise specified, 2-day stationary phase liquid monocultures were

refed either on liquid or solid fresh SDcasaWAU medium.

Stationary phase cultures were refed by dilution into liquid fresh SDcasaWAU medium at

approximately 0.05 OD_{600nm}. The OD_{600nm} was monitored upon refeeding for at least 7 hours covering

both 'lag' and exponential phases (until at least a 5-fold increase of the OD_{600nm} compared to the initial

point). To quantify the duration of the 'lag' phase, we normalized all the OD_{600nm} values of each

individual growth curve to their respective initial OD_{600nm} value and then applied the logarithmic

function to all the normalized OD_{600nm} values of the growth curve. We next performed a linear

regression on the points corresponding to the proliferation phase of the growth curve using GraphPad

Prism 8 (GraphPad Software, Inc. La Jolla, USA). The time value for which the linear regression

intersects the Time-X axis corresponds to the end of the 'lag' phase and the beginning of the

proliferation phase, and is referred as 'the duration of lag phase'.

Quiescent cells from stationary phase cultures were also refed onto solid medium. Cells were spread

onto half a 3% agarose microscope pad made of fresh SDcasaWAU. Bright field images of individual

cells were then captured every 15 minutes for up to 4 hours. The cells forming an unequivocal new

bud upon refeeding were scored.

When indicated, an aliquot of stationary phase culture can be sonicated before a refeeding experiment

on microscope agarose pad. Cycles of 30 seconds in a sonication bath were carried out until the

proportion of single-cell entities in the *ace2* SF culture increased to a desired proportion

(approximately 40%).

Determination of *CLN1* mRNA level upon refeeding by Reverse transcriptase-

PCR.

To determine the level of *CLN1* mRNA upon refeeding of 2-day stationary phase cultures, 15 mL of a

2-day culture was diluted in 85 mL of fresh medium. Fifteen minutes upon refeeding, 25 mL of culture

was sampled and centrifuged at 4000 rpm at 4°C for 3 min. The pellet was washed in 1 mL of RNase-

free H2O, and centrifuged at 12,000 g at 4°C for 30 seconds. A Tri-Reagent (Sigma, T9424) protocol

was then performed to extract total RNA. A reverse-transcription of 200ng of total RNA using the 'High

Capacity cDNA Reverse Transcription' Kit (ThermoFisher, 4368813) was then performed. The cDNA was

diluted to 1/25 and amplified with a Quantitative-PCR protocol, using the GoTaq® qPCR Master Mix

(Promega, A6001) and primers specific to the coding region of *CLN1* or the coding region of *ACT1*

(Supplemental Table 3).

Microscopy

The refeeding experiments on microscope pad, as well as the Snowflake size-measurement

experiments were carried out using a Zeiss 200M inverted microscope (Carl Zeiss, Thornwood, NY,

USA) with the additional equipment : a MS-2000 stage (Applied Scientific Instrumentation, Eugene,

OR, USA), a Lambda LS 300 W xenon light source (Sutter, Novato, CA, USA), a 5 positions filter turret, a

100X 1.4NA Plan-Apochromat objective and a CoolSnap HQ camera (Roper Scientific, Tucson, AZ, USA).

The filters are from Chroma Technology Corp. The microscope and additional equipment were controlled by SlideBook software 5.0. (Intelligent Imaging Innovations, Denver, CO, USA). The images were analyzed with Image J. All other microscopy experiments were carried out using Zeiss Primo Star microscope, equipped with a 40X/0.65 Plan-Achromat objective.

Statistical analysis

All experiments were carried out at least three times on independent biological samples. The exact number of independent samples is indicated in the figure legends. Data are presented as mean ± SD. All the statistical analyses were done using GraphPad Prism 8 (GraphPad Software, Inc. La Jolla, USA). The proportion of SF in competition experiments, the refeeding experiments on microscope pad and the proportion of dead cells were analyzed by Fisher's exact test on a contingency table. Other experiments are analyzed by Welch's unpaired t-test. p-values below 0.05 are represented by *; p values inferior to 0.005 are represented by **; and p values inferior to 0.001 are represented by ***. p-values superior to 0.05 were considered non-significant and are represented by 'ns'.

Acknowledgments and funding sources

The authors thank Benoît Pinson for technical advising, the members of the Sagot lab for discussions on the project and Isabelle Sagot for critical reading of the manuscript. The authors also thank Sabine Vaur for her help with the qPCR method. This work was supported by a grant from Ligue Régionale Contre le Cancer (2022 Dordogne) to BD-F. TD was supported by a grant from University of Bordeaux, NewMoon RRI Program on Cancer in Evolution.

Author contributions

434   T.L.D., D.L. and B.D.-F. designed research; T.L.D. performed research; T.L.D., D.L. and B.D.-F. analyzed

435   data; and T.L.D. and B.D.-F. wrote the paper.

Competing interests

438   The authors declare no competing interest.

References

1. Grosberg RK, Strathmann RR. The Evolution of Multicellularity: A Minor Major Transition? Annu Rev Ecol Evol Syst. 1 déc 2007;38(1):621-54. doi:10.1146/annurev.ecolsys.36.102403.114735

2. Tong K, Bozdag GO, Ratcliff WC. Selective drivers of simple multicellularity. Current Opinion in Microbiology. juin 2022;67:102141. doi:10.1016/j.mib.2022.102141

3. La Fortezza M, Schaal K, Velicer GJ. Why Aggregate? On the Evolution of Aggregative Multicellularity [Internet]. 2021 [cité 28 avr 2025]. Disponible sur: https://www.preprints.org/manuscript/202105.0451/v1 doi:10.20944/preprints202105.0451.v1

4. Newell PC, Telser A, Sussman M. Alternative Developmental Pathways Determined by Environmental Conditions in the Cellular Slime Mold *Dictyostelium discoideum*. J Bacteriol. nov 1969;100(2):763-8. doi:10.1128/jb.100.2.763-768.1969

5. Gimeno CJ, Ljungdahl PO, Styles CA, Fink GR. Unipolar cell divisions in the yeast S. cerevisiae lead to filamentous growth: Regulation by starvation and RAS. Cell. mars 1992;68(6):1077-90. doi:10.1016/0092-8674(92)90079-R

6. Kumar A. The Complex Genetic Basis and Multilayered Regulatory Control of Yeast Pseudohyphal Growth. Annu Rev Genet. 23 nov 2021;55(1):1-21. doi:10.1146/annurev-genet-071719-020249

7. Daignan-Fornier B, Laporte D, Sagot I. Quiescence Through the Prism of Evolution. Front Cell Dev Biol. 29 oct 2021;9:745069. doi:10.3389/fcell.2021.745069

8. Bertoli C, Skotheim JM, De Bruin RAM. Control of cell cycle transcription during G1 and S phases. Nat Rev Mol Cell Biol. août 2013;14(8):518-28. doi:10.1038/nrm3629

9. Hanschen ER, Marriage TN, Ferris PJ, Hamaji T, Toyoda A, Fujiyama A, et al. The Gonium pectorale genome demonstrates co-option of cell cycle regulation during the evolution of multicellularity. Nat Commun. 22 avr 2016;7(1):11370. doi:10.1038/ncomms11370

10. MacWilliams H, Doquang K, Pedrola R, Dollman G, Grassi D, Peis T, et al. A retinoblastoma ortholog controls stalk/spore preference in *Dictyostelium*. Development. 1 avr 2006;133(7):1287-97. doi:10.1242/dev.02287

11. Bachewich C, Whiteway M. Cyclin Cln3p Links G_1 Progression to Hyphal and Pseudohyphal Development in *Candida albicans*. Eukaryot Cell. janv 2005;4(1):95-102. doi:10.1128/EC.4.1.95-102.2005

12. Loeb JDJ, Kerentseva TA, Pan T, Sepulveda-Becerra M, Liu H. Saccharomyces cerevisiae G1 Cyclins Are Differentially Involved in Invasive and Pseudohyphal Growth Independent of the Filamentation Mitogen-Activated Protein Kinase Pathway. Genetics. 1 déc 1999;153(4):1535-46. doi:10.1093/genetics/153.4.1535

13. Cross FR, Buchler NE, Skotheim JM. Evolution of networks and sequences in eukaryotic cell cycle control. Phil Trans R Soc B. 27 déc 2011;366(1584):3532-44. doi:10.1098/rstb.2011.0078

14. Ratcliff WC, Pentz JT, Travisano M. TEMPO AND MODE OF MULTICELLULAR ADAPTATION IN
EXPERIMENTALLY EVOLVED *SACCHAROMYCES CEREVISIAE*. Evolution. juin 2013;67(6):1573-81.
doi:10.1111/evo.12101

15. Ratcliff WC, Denison RF, Borrello M, Travisano M. Experimental evolution of multicellularity. Proc
Natl Acad Sci U S A. 31 janv 2012;109(5):1595-600. doi:10.1073/pnas.1115323109 PubMed
PMID: 22307617; PubMed Central PMCID: PMC3277146.

16. Ratcliff WC, Fankhauser JD, Rogers DW, Greig D, Travisano M. Origins of multicellular evolvability
in snowflake yeast. Nat Commun. mai 2015;6(1):6102. doi:10.1038/ncomms7102

17. Koschwanez JH, Foster KR, Murray AW. Improved use of a public good selects for the evolution
of undifferentiated multicellularity. eLife. 2 avr 2013;2:e00367. doi:10.7554/eLife.00367

18. Mostovoy Y, Thiemicke A, Hsu TY, Brem RB. The Role of Transcription Factors at Antisense-
Expressing Gene Pairs in Yeast. Genome Biol Evol. juin 2016;8(6):1748-61.
doi:10.1093/gbe/evw104

19. Voth WP, Olsen AE, Sbia M, Freedman KH, Stillman DJ. *ACE2* , *CBK1* , and *BUD4* in Budding and
Cell Separation. Eukaryot Cell. juin 2005;4(6):1018-28. doi:10.1128/EC.4.6.1018-1028.2005

20. Doolin M, Johnson AL, Johnston LH, Butler G. Overlapping and distinct roles of the duplicated
yeast transcription factors Ace2p and Swi5p. Molecular Microbiology. avr 2001;40(2):422-32.
doi:10.1046/j.1365-2958.2001.02388.x

21. Jacobeen S, Pentz JT, Graba EC, Brandys CG, Ratcliff WC, Yunker PJ. Cellular packing, mechanical
stress and the evolution of multicellularity. Nature Phys. mars 2018;14(3):286-90.
doi:10.1038/s41567-017-0002-y

22. Nash R, Tokiwa G, Anand S, Erickson K, Futcher AB. The WHI1+ gene of Saccharomyces cerevisiae
tethers cell division to cell size and is a cyclin homolog. The EMBO Journal. déc
1988;7(13):4335-46. doi:10.1002/j.1460-2075.1988.tb03332.x

23. Newcomb LL, Diderich JA, Slattery MG, Heideman W. Glucose Regulation of *Saccharomyces
cerevisiae* Cell Cycle Genes. Eukaryot Cell. févr 2003;2(1):143-9. doi:10.1128/EC.2.1.143-
149.2003

24. Wu M, Newcomb L, Heideman W. Regulation of Gene Expression by Glucose inSaccharomyces
cerevisiae: a Role for ADA2and ADA3/NGG1. J Bacteriol. 15 août 1999;181(16):4755-60.
doi:10.1128/JB.181.16.4755-4760.1999

25. Fang O, Hu X, Wang L, Jiang N, Yang J, Li B, et al. Amn1 governs post-mitotic cell separation in
Saccharomyces cerevisiae. Barsh GS, éditeur. PLoS Genet. 1 oct 2018;14(10):e1007691.
doi:10.1371/journal.pgen.1007691

26. Laporte D, Jimenez L, Gouleme L, Sagot I. Yeast quiescence exit swiftness is influenced by cell
volume and chronological age. Microb Cell. 5 févr 2018;5(2):104-11.
doi:10.15698/mic2018.02.615

27. Courchesne WE, Kunisawa R, Thorner J. A putative protein kinase overcomes pheromone-
induced arrest of cell cycling in S. cerevisiae. Cell. 22 sept 1989;58(6):1107-19. doi:10.1016/0092-
8674(89)90509-6

28. Madhani HD, Styles CA, Fink GR. MAP Kinases with Distinct Inhibitory Functions Impart Signaling
Specificity during Yeast Differentiation. Cell. nov 1997;91(5):673-84. doi:10.1016/S0092-
8674(00)80454-7

29. Madhani HD, Galitski T, Lander ES, Fink GR. Effectors of a developmental mitogen-activated
protein kinase cascade revealed by expression signatures of signaling mutants. Proc Natl Acad Sci
USA. 26 oct 1999;96(22):12530-5. doi:10.1073/pnas.96.22.12530

30. Bozdag GO, Libby E, Pineau R, Reinhard CT, Ratcliff WC. Oxygen suppression of macroscopic
multicellularity. Nat Commun. 14 mai 2021;12(1):2838. doi:10.1038/s41467-021-23104-0

31. Kapsetaki SE, West SA. The costs and benefits of multicellular group formation in algae*.
Evolution. juin 2019;73(6):1296-308. doi:10.1111/evo.13712

32. Barrere J, Nanda P, Murray AW. Alternating selection for dispersal and multicellularity favors
regulated life cycles. Current Biology. mai 2023;33(9):1809-1817.e3.
doi:10.1016/j.cub.2023.03.031

33. Sherman F. [1] Getting started with yeast. In: Methods in Enzymology [Internet]. Elsevier; 1991
[cité 16 juin 2025]. p. 3-21. Disponible sur:
https://linkinghub.elsevier.com/retrieve/pii/007668799194004V doi:10.1016/0076-
6879(91)94004-V

34. Gietz D, Jean ASt, Woods RA, Schiestl RH. Improved method for high efficiency transformation of
intact yeast cells. Nucl Acids Res. 1992;20(6):1425-1425. doi:10.1093/nar/20.6.1425

35. Gietz RD, Akio S. New yeast-Escherichia coli shuttle vectors constructed with in vitro
mutagenized yeast genes lacking six-base pair restriction sites. Gene. déc 1988;74(2):527-34.
doi:10.1016/0378-1119(88)90185-0

Figures

Figure 1

A

B

SF/PK co-culture in:

■ *CLN3/CLN3* background ■ *cln3Δ/cln3Δ* background

C

ace2Δ/ace2Δ - ACE2/ACE2 co-culture in:

■ *CLN3/CLN3* background
■ *cln3Δ/cln3Δ* background

D

Overexpression vector expressed
in the SF/PK co-culture:

■ Empty ■ *WHI5*

Figure 1 - figure supplement 1

Figure 1. Competition experiments. (A) Experimental set-up of the co-culture competition

experiments between *ace2/ace2* and *ACE2/ACE2* strains. A 50/50 mix of the two strains based either

on phenotype (snowflake vs planktonic, microscopy) or on genotype (*ace2* vs *ACE2,* qPCR) was

inoculated in SDcasaWAU liquid medium. The resulting co-culture was grown for 2 days until stationary

phase was reached and then diluted in new SDcasaWAU medium. For each dilution step, the

proportion of the snowflake phenotype or *ace2* genotype was determined (by microscopy or qPCR

respectively). **(B)** Evolution of the percentage of snowflake entities during independent co-culture

competitions between *ace2/ace2* and *ACE2/ACE2* strains, either *cln3/cln3* (red bars) or *CLN3/CLN3*

(black bars) (N=4, n>250, mean ± SD, ns p>0.05, *p<0.05, ***p<0.001). The percentage of snowflakes

is statistically compared between *CLN3/CLN3* and *cln3/cln3* co-cultures at each dilution round, using a

Fisher's exact test. **(C)** The proportion of *ace2/ace2* genotype was monitored by qPCR in independent

competitions in *cln3/cln3* (red bars) and *CLN3/CLN3* (black bars) backgrounds. The *ACT1* locus,

amplified in all the cells of the population, was used to normalize the proportion of *ace2* cells in the

co-culture cell population (N=3, mean ± SD, unpaired t-test, Welch correction, ns p>0.05, ***p<0.001).

(D) Evolution of the percentage of snowflake entities during independent co-culture competitions

between *ace2/ace2* and *ACE2/ACE2* strains overexpressing *WHI5* (orange bars) or not (black bars)

(N=3, n>235, mean ± SD, Fisher's exact test, ns p>0.05, *p<0.05, **p<0.005, ***p<0.001).

Figure supplement 1. (A) Number of cells per Snowflake entity determined by counting the number

of *HTB1-CFP* positive nuclei per Snowflake entity. **(B-E)** Representative images of *CLN3/CLN3* strains,

either *ACE2/ACE2* **(B)** or *ace2/ace2* **(C)**, and cln3/cln3 strains, either *ACE2/ ACE2* **(D)** or *ace2/ace2* **(E)**.

Bars are 10 μm. **(F)** Cell volume measured at 2 days of culture (error bars ± SD, unpaired t-test, Welch

correction, ***p<0.001, ns p>0.05).

Figure 2

Figure 2 - figure supplement 1

Figure 2. Quiescence exit efficiency explains the results of competition experiments. (A) Lag phase

duration after cell 2-day old population refeeding with SDCasaWAU medium. The optical density at

600nm was followed after population refeeding for 4 independent experiments. The mean and the SD

are indicated together with an unpaired t-test, Welch correction, ns p>0.05, ***p<0.001. **(B)**

Representative image series of *cln3/cln3* cells either *ace2/ace2* or *ACE2/ACE2* after refeeding of 2 days

cultures on a SDCasaWAU medium containing microscope pad. Arrows point to emerging buds. Bars

are 10μm. **(C)** Proportion of *cln3/cln3* cells that have formed a new bud after refeeding on a

SDCasaWAU medium containing microscope pad of either *ace2/ace2* or *ACE2/ACE2* 2 days cultures

(N=3, n>101, Fisher's exact test, mean ± SD, ns p>0.05, *p<0.05, **p<0.005, ***p<0.001). **(D)** Same as

C in a *cln3* haploid background (N=3, n>100, mean ± SD, Fisher's exact test, **p<0.005, ***p<0.001).

(E) Percentage of snowflake entities observed in independent co-cultures of *ace2/ace2* and *ACE2/ACE2*

strains. Cultures either went through cycles of proliferation and stationary phase as described in Fig1A

(right panel), or maintained in constant exponential phase (left panel) (N>3, n>208, mean ± SD, Fisher's

exact test, ns p>0.05). **(F)** Same as A except that the refeeding was done after a culture of 5 days (N=3,

mean ± SD, unpaired t-test, Welch correction, **p<0.005). **(G)** Percentage of snowflake entities

observed in independent co-cultures of *ace2/ace2* and *ACE2/ACE2* strains diluted every two days (red

and black bars) or every five days (grey and pink bars) (N≥3, n>700, mean ± SD, Fisher's exact test, ns

p>0.05, **p<0.005, ***p<0.001).

Figure supplement 1. **(A)** Theoretical diagram showing the various phenomena that can explain, either

alone or combined, how one subpopulation can outcompete another subpopulation in a co-culture.

One subpopulation may divide more rapidly in exponential phase (a), may have a higher yield (b), may

undergo less cell death during stationary phase (c) or may re-enter proliferation phase faster after

refeeding (d). **(B)** Doubling time of populations in proliferation phase. Independent cultures (N>3) were

maintained in constant proliferation phase for more than 24 hours, then the optical density at 600 nm

was measured every 15 minutes for 8 hours (mean ± SD, unpaired t-test, Welch correction, ns p>0.05).

(C-D) The dry biomass determined by filtration (**C**, N=3, mean ± SD, unpaired t-test, Welch correction,

ns p>0.05) and the percentage of dead cells stained with ethylene blue (**D**, N=3, n>280, mean ± SD,

Fisher's exact test, ns p>0.05,) were also measured for 2 days stationary phase cultures. **(E)** Proportion

of *CLN3/CLN3* cells that have formed a new bud after refeeding on a SDCasaWAU medium containing

microscope pad of either *ace2/ace2* or *ACE2/ACE2* 2 days cultures (N=3, n>101, Fisher's exact test,

mean ± SD, ns p>0.05). **(F)** Proportion of *CLN3* cells that have formed a new bud after refeeding on a

SDCasaWAU medium containing microscope pad of either *ace2* or *ACE2* 2 days cultures (N=3, n>91,

Fisher's exact test, mean ± SD, ns p>0.05). **(G-J)** Representative images of diploid *ace2/ace2* snowflakes

either *cln3/cln3* (**G**) or *CLN3/CLN3* (**H**), and haploid *ace2* snowflakes either *cln3* (**I**) or *CLN3* (**J**). Bars are

10 μm. **(K).** Size distribution of snowflakes in haploid (empty circles) or diploids (plain circles)

backgrounds, either *CLN3* (black) or *cln3* (red). Size is estimated by measuring both the longest

segment of the snowflake entity and the longest segment perpendicular to the first one (N=3, n>72).

We compared the length of each segment independently with an unpaired t-test with a Welch

correction (error bars ± SD, unpaired t-test, Welch correction, ***p<0.001, account for both segment).

Figure 3

Figure 3 - figure supplement 1

A

Quintuple mutant strain

B

C

Figure 3. Quintuple *cts1, dse2, dse4, egt2, scw11* mutant strains do not rescue the quiescence-exit

default associated with *cln3* and are not selected in competition experiments. (A) Proportion of

cln3/cln3 cells, either quintuple mutant or WT, that formed a new bud after refeeding of 2 days

cultures on a SDCasaWAU medium containing microscope pad (N=3, n>102, mean ± SD, Fisher's exact

test, ns p>0.05, *p<0.05). **(B)** Proportion of *cln3/cln3* cells, either quintuple mutant or *ace2/ace2*, that

formed a new bud after refeeding of 2 days cultures on a SDCasaWAU medium containing microscope

pad (N=3, n>75, mean ± SD, Fisher's exact test, ns p>0.05, *p<0.05, **p<0.005, ***p<0.001). **(C)**

Evolution of the percentage of snowflake entities during independent co-culture competitions

between quintuple mutant and WT strains either *cln3/cln3* (red bars) or *CLN3/CLN3* (black bars) (N=3,

n>250, mean ± SD, Fisher's exact test).

Figure supplement 1. Snowflake characterization of quintuple mutant strains. (A) Representative

image of the quintuple mutant strain. Bar is 10µm. **(B)** Size distribution of quintuple mutant (triangles)

or *ace2/ace2* (circles) snowflakes. Size is estimated by measuring both the longest segment of the

snowflake entity and the longest segment perpendicular to the first one. We compared the length of

each segment independently with an unpaired t-test with a Welch correction (N=3, n>60, mean ± SD).

(C) Proportion of snowflake entities in a population of quintuple mutant or *ace2/ace2* strains (N=3,

n>201, mean ± SD, Fisher's exact test, ns p>0.05).

Figure 4

A

B

Figure 4 - figure supplement 1

Figure 4. Rescue of the quiescence-exit default associated with *cln3* depends on the *ace2* mutation

and not on the Snowflake phenotype. (A) A 2-day *ace2/ace2 cln3/cln3* culture was sonicated and refed

onto a fresh SDCasaWAU microscope pad. The proportion of cells that formed a new bud after

refeeding on a SDCasaWAU medium containing microscope pad was assessed within different

multicellular size ranges (N=3, n>184, mean ± SD, Fisher's exact test). **(B)** 2-day *cln3* cultures, either

ace2 or *ACE2*, were sonicated and refed onto a fresh SDCasaWAU microscope pad. Within entities

formed of 1 to 3 cells, the proportion of cells that formed a new bud after refeeding on a SDCasaWAU

medium containing microscope pad was assessed (N=3, n>63, mean ± SD, Fisher's exact test, ns p>0.05,

*p<0.05, **p<0.005, ***p<0.001).

Figure supplement 1. **Effect of sonication on SF phenotype, cell death and quiescence exit**. **(A-B)** The

effect of sonication bath on an *ace2* culture was assessed by measuring the proportion of single cell

entities (**A**, N=3, n>212) and the percentage of dead cells (**B**, N=3, n>207) in the population before and

after the culture was sonicated. **(C)** Proportion of *cln3* cells, either *ace2* or *ACE2*, and either sonicated

or not, that formed a new bud after refeeding of 2 days cultures on a SDCasaWAU medium containing

microscope pad (N=3, n>63, Fisher's exact test, mean ± SD).

Figure 5

Figure 5. Rescue of the quiescence-exit default associated with *cln3* is dependent on the Kss1 MAP kinase, which favors transcription of the *CLN1* cyclin gene. (A) Proportion of *cln3 kss1* cells, either *ace2* or *ACE2*, that formed a new bud after refeeding of 2-day cultures on a fresh SDCasaWAU medium containing microscope pad (N=3, n>86, mean ± SD, Fisher's exact test, ns p>0.05). **(B)** Ratio of *CLN1* mRNA level and *ACT1* mRNA level, determined by RT-qPCR, 15 minutes after refeeding of 2 days *cln3* cultures, either *ace2* or *ACE2* (N=3, mean ± SD, unpaired t-test, Welch correction, **p<0.005). **(C)** Ratio of *CLN1* mRNA level and *ACT1* mRNA level, determined by RT-qPCR, 15 minutes after refeeding of 2-day *kss1 cln3* cultures, either *ace2* or *ACE2* (N=3, mean ± SD, unpaired t-test, Welch correction, ns p>0.05).

Figure 6

Figure 6 - figure supplement 1

Fig 6. Quiescence exit efficiency and selection in competition experiments of Snowflake $AMN1^{368D}$ strains. (A) Proportion of *cln3* cells, either $AMN1^{368D}$ or *amn1-368V*, that formed a new bud after refeeding of 2 days cultures on a fresh SDCasaWAU medium containing microscope pad (N=3, n>77, mean ± SD, Fisher's exact test, ns $p>0.05$, *$p<0.05$, **$p<0.005$, ***$p<0.001$). **(B)** Evolution of the percentage of snowflake entities during independent co-culture competitions between $AMN1^{368D}$ and *amn1-368V* strains, either *cln3/cln3* (red bars) or *CLN3/CLN3* (black bars) (N=3, n>252, mean ± SD, Fisher's exact test, ns $p>0.05$, **$p<0.005$, ***$p<0.001$). **(C)** Evolution of the percentage of snowflake entities during independent co-culture competitions between $AMN1^{368D}$ and *amn1-368V* strains overexpressing *WHI5* (orange bars) or not (black bars) (N=3, n>302, Fisher's exact test, ns $p>0.05$, **$p<0.005$, ***$p<0.001$).

Figure supplement 1. Snowflake characterization of $AMN1^{368D}$ strains. (A) Representative image of the $AMN1^{368D}$ strain. Bar is 10µm. **(B)** Proportion of snowflake entities in populations of $AMN1^{368D}$ or *ace2* haploid strains (N=3, n>219, mean ± SD, Fisher's exact test, ns $p>0.05$). **(C)** Size distribution of $AMN1^{368D}$ (diamond) or *ace2* (circle) snowflakes. Size is estimated by measuring both the longest segment of the snowflake entity and the longest segment perpendicular to the first one (N=3, n>74, mean ± SD, unpaired t-test, Welch correction, ***$p<0.001$, account for both segment).

Figure 7

A

B

Fig 7. Model explaining the selection of the Snowflake phenotype as a passenger phenotype. In strains expressing the non-functional *amn1-368V* allele **(A)**, a default in quiescence exit after refeeding was observed in a *cln3* mutant. In non-laboratory strains **(B)**, the functional *AMN1^368D* allele leads to degradation of Ace2p, which can no longer repress the expression of the Kss1 Map kinase. Kss1p favors the transcription of *CLN1*, which would favor progression to S phase and partially suppress the default in quiescence exit associated with the *cln3* mutation. This observed selective advantage (a) leads to the selection of *AMN1^368D* genotype (b), and because of the pleiotropy of Ace2p transcription factor, to the selection of the Snowflake multicellular phenotype as a passenger phenotype (c).