Red List criteria underestimate climate-related extinction risk of
range-shifting species

Raya Keuth[1,*], Susanne A. Fritz[2,3], Damaris Zurell[1]

Affiliations:

[1] University of Potsdam, Institute of Biochemistry and Biology, Potsdam, Germany

[2] German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Germany

[3] Friedrich Schiller University Jena, Institute of Biodiversity, Ecology and Evolution, Jena,
Germany

*corresponding author: raya.keuth@uni-potsdam.de

ORCIDs: RK: 0000-0002-6694-8887; SF: 0000-0002-4085-636X; DZ: 0000-0002-4628-3558

Abstract

Climate change causes global species redistribution and elevates extinction risk, making
early identification of vulnerable species critical for timely conservation. The IUCN Red List
provides guidelines for assessing climate-related extinction risk using species distribution
models (SDMs) and spatially explicit population models (SEPMs). However, a systematic
evaluation of these guidelines is currently missing. Using simulations of virtual species with
diverse life-history traits and range dynamics, we found that SDMs consistently
underestimated extinction risk for range-shifting species. This was due to a concave
relationship between population size and habitat loss, which contradicts the linear
assumption in the Red List guidelines. For range-contracting species, SDMs provided
adequate warning times. Probabilistic extinction estimates from SEPMs provided delayed
warning for all species, particularly for highly threatened ones. Our results reveal key
limitations of current Red List guidelines under climate change. Based on our findings, we
provide tentative recommendations for updating the IUCN Red List guidelines.

Keywords: IUCN Red List, extinction risk, species distribution models, climate change,
virtual species, spatially explicit population models

Introduction

Climate change poses a major threat to biodiversity, affecting virtually all biological processes[1] and potentially elevating extinction risk for many species[2,3]. Species dynamically respond to climate change, for example by tracking suitable habitat[4], which has been projected to lead to substantial global redistribution of species[5]. To protect biodiversity before declines become irreversible, early identification of the most threatened species is essential. Conservation assessments should hence provide sufficient warning time, defined as the time between identification of a threatened species and its extinction without any interventions[6].

The IUCN Red List for Threatened Species (Red List hereafter) is the most widely used framework for assessing extinction risk and informing conservation and policy decisions[7,8]. It classifies species into one of three threatened categories (Vulnerable, Endangered, Critically endangered) based on five quantitative criteria (A-E) that are derived from the theory of extinction trajectories for single populations[7,9,10]. The five different criteria provide alternative ways of assessing a species in the Red List, with qualifying for a threatened category under one criterion being sufficient to classify species in the category. The assessment of the species is mostly based on population size and trends, as well as geographic range area[9]. Originally designed to identify species already in decline, the IUCN has since introduced guidelines for assessing future extinction risk under climate change. These recommend species distribution models (SDMs) to infer habitat loss, and spatially explicit population models (SEPMs) to infer population loss or estimate extinction probability[11,12].

Despite their growing use, the predictive performance of SDMs under climate change remains debated[13–15]. SDMs are correlative models that relate observed species occurrences to prevailing environmental conditions and use this species-environment relationship to predict the suitable habitat under future climatic scenarios[16]. However, SDMs omit key biological processes, such as dispersal limitations, which can lead to an overprediction of future range size[17,18]. Nevertheless, SDMs currently constitute the most pragmatic approach for projecting potential future climate impacts. The Red List guidelines suggest to use SDM-predicted changes in habitat suitability as a proxy for population size reduction under criterion A3[8(p.102)]. This approach assumes a linear relationship between population loss and SDM-predicted habitat loss, which is acknowledged in the Red List guidelines as a source of uncertainty[8(p.99)]. Beyond simply causing uncertainty, this assumption actually lacks empirical validation[19,20], and could be influenced by the non-uniform distribution of species in the

74 landscape[21], extinction debts[22], and extinction thresholds related to minimum viable
75 population sizes[23].
76
SEPMs, in contrast, simulate demographic and dispersal processes over space and time,
potentially offering more realistic predictions of species responses to climate change[24,25]. Yet,
wider application of these models has been hampered by technical and data challenges[26,27]
and they have thus been rarely used for assessing climate-related future extinction risk of
species[28]. A key open question is whether SEPMs provide more accurate or timely extinction
risk estimates than SDMs, particularly for species with different range dynamics under
climate change. So far, only a single case study has compared these modelling approaches
in the context of IUCN assessments[29] while a systematic evaluation of modelling approaches
across varying life histories and range dynamics is missing. This gap prevents evidence-
based improvements to current Red List guidelines.
87
In this study, we aim to test the Red List guidelines for assessing species threatened by
climate change. Specifically, we test (i) whether the assumed linear relationship between
habitat loss and population decline holds across species and range dynamics, and (ii)
whether model-based assessments differ in their ability to identify extinction risk early
enough for conservation action. Due to the lack of data regarding climate change-driven
extinctions, we chose a simulation approach[30,31]. We generated virtual species with
contrasting life-history and dispersal traits (niche position, niche breadth, growth rate and
dispersal distance)[32] and used a spatially explicit modelling framework[33] to project population
and range dynamics under climate change across three different artificial landscapes (Fig.
1). Finally, we compared the performance of SDMs and SEPMs for assessing extinction risk,
focusing on their ability to provide timely warnings for species with varying life history traits
and undergoing either range contraction or range shift.

Figure 1: Conceptual figure of the modelling workflow. We simulated virtual species under climate change using a spatially explicit individual-based model. Species varied in four traits, which resulted in a variety of range distributions and dynamics. Before climate change set in, distribution data were sampled from the simulations as input to train correlative species distribution models (SDMs). SDMs were then used to predict habitat loss under climate change. Finally, Red List classifications under climate change were compared for SDM-derived habitat loss and simulation-derived population size reduction and extinction probability.

Results

Population dynamics and SDM performance

The considered trait combinations led to distinct differences in range and population dynamics. Notably, species with marginal niche position mainly occurred at the "northern" edge of the landscape and their ranges contracted towards that edge under climate change, while species with central niche position had centrally located ranges and exhibited range shifting before going extinct under climate change (Fig. 2). As could be expected, species with slow growth, narrow niches and short dispersal went extinct earlier than species with fast growth, wide niches and long dispersal (Fig. S1). The differences in temporal dynamics between species with different trait values were more pronounced for range-shifting species than for range-contracting species.

We sampled species occurrence from the simulations before climate change set in, and fitted three different SDM algorithms and an ensemble model. The SDM ensembles showed high pre-climate change predictive accuracy for all species and landscapes (AUC: 0.95 ± 0.02; TSS: 0.78 ± 0.06). The predictive performance was slightly lower for species with

central compared to marginal niches and for species with short compared to long dispersal (Fig. S2 + S3). Under climate change, SDMs correctly predicted range loss of range-contracting species, but they underestimated the range loss of dispersal-limited range-shifting species (Fig. 2). This pattern was consistent across replicate landscapes (Fig. S1).

Figure 2: Distribution of virtual species and SDM-predicted habitat suitability for two example species. Top panels (a) show the maps of simulated „true" species abundance and maps of habitat suitability predicted by SDMs for one replicate run under equilibrium conditions (year 0) and under climate change (year 20 / 30) in one example landscape. Bottom panels (b) show the corresponding time trajectories of the relative change in habitat suitability sums predicted from SDMs and in simulated population size, both averaged over ten selected replicate runs, as well as simulated extinction probability. Results are shown for two example species differing in their range dynamics (i.e., in marginal vs. central niche position). Both species have a narrow niche, slow growth rate, and short dispersal.

Population size - habitat loss relationship

For all species and replicate landscapes, the relationship between relative population size and SDM-derived habitat loss deviated from the linear assumption proposed in the Red List guidelines (Fig. 3). For range-contracting species, the relationship was convex with an initial

smaller rate of decline of population size in comparison to habitat loss. Population size
steeply declined once habitat loss exceeded 50% across all replicate landscapes (Fig. S4 +
S5). In contrast, range-shifting species showed a concave relationship with considerable
declines in population size already happening at low rates of habitat loss, and with larger
differences between the different replicate landscapes (Fig. 3 + S5). We only found
comparatively small differences between the relationships for the other traits, with the
variability between the different landscapes often being larger than among the different trait
levels (Fig. S4 + S5). A Bayesian generalised mixed-model analysis of covariance showed
that the population size - habitat loss relationship significantly differed across all traits, but
the effect sizes for niche position (i.e. for range-shifting vs. range-contracting species) were
considerably larger than for all other traits (Table S1).

Figure 3: Population size - habitat loss relationship for the different range dynamics. Relative
population size was obtained from the simulation (observed true data for virtual species) and habitat
loss was predicted by the SDMs. The dot-dash grey line shows the assumption of a linear relationship
between population size and habitat loss as proposed by the Red List guidelines. The relationships
are displayed using an ordered beta regression model, in which landscape was included as a random
effect. Lines indicate the mean model predictions and the transparent bands the 95% confidence
interval. We only show results here for the trait niche position, resulting in range-contracting and
range-shifting species, while results for the other traits are displayed in Fig. S4.

Red List classification

An important question is to which extent SDM-derived habitat loss under climate change can
be used for assessment in the Red List. To answer this, we determined the time at which a
species would be classified into the three threatened categories VU, EN and CR
(Vulnerable, Endangered, and Critically endangered) of the Red List. Specifically, we
compared assessments against criterion A3 when i) applied to the true (simulated)
population size and ii) to the sum of habitat suitability derived from SDMs as recommended
by the Red List guidelines[8(pp.101-111)]. Additionally, we compared these to the assessment
under criterion E based on the quantitative extinction probability extracted from our
simulation. Since all species went extinct during the 90 years of climate change, all of them
qualified for all threatened categories throughout their life span. Again, we found the
strongest differences between range-contracting and range-shifting species reflecting the
differences in the population size - habitat loss relationship (Fig. 4 + Fig. S6).

Figure 4: Classification times for listing species in different IUCN Red List threatened
categories. Results are shown separately for range-contracting (left) vs. range-shifting species
(right). Classification time refers to the years when species would be classified into the three different
threatened categories (Vulnerable, Endangered, Critically endangered) using true population size or
SDM-derived habitat loss against criterion A3, and simulated extinction probability against criterion E.
Note that all species went extinct during the 90 years of climate change, thus qualifying for all
threatened categories throughout their life span. The grey line shows the mean year of extinction
across species and the grey bands the 95% confidence interval. Sampling size per box plot is n=240
(8 species x 3 replicate landscapes x 10 replicate runs).

For range-contracting species, the classification time did not differ between true population
size and SDM-derived habitat loss. By contrast, for range-shifting species SDM-based risk
classification was substantially later than for true population size because range loss was
underestimated by the SDMs. This resulted in a very short warning time for range-shifting
species when using SDM-based risk classification (Fig. 4 + Fig. S6). In a sensitivity analysis,
we tested whether SDM-based predictions of range-shifting species could be improved by
including simple dispersal assumptions but found no effect (Fig. S7). Other traits did not
result in pronounced differences in classification times (Fig. S6). Interestingly, we found that
for higher threatened categories (EN, CR) quantitative estimates of extinction probability and
listing under criterion E resulted in considerably later classification time, and thus very short
warning time, than information on population size and listing under criterion A3 (Fig. 4 + S6).
The Red List guidelines indicate that criteria A-D are more precautionary than criterion
E[8(p.17)]. Still, our results show that listing under these criteria can differ by up to 20
generations for our virtual species (Fig. 4).

Discussion

Based on idealised simulated species, we systematically evaluated the guidelines of the Red
List for assessing species-level extinction risk under climate change. Our results revealed
that the effectiveness of current Red List criteria strongly depend on expected range
dynamics. SDM-based assessment provided sufficient warning time only for range-
contracting species. For range-shifting species, warning time was substantially shorter
mainly because of the highly non-linear relationship between population size and SDM-
predicted habitat loss. Other life-history traits had comparatively little effect on warning
times. Further, model-based assessment based on probabilistic extinction estimates led to
considerably shorter warning times than when based on predicted population sizes.

The strong effect of range dynamics stems from fundamental differences in how population
size relates to predicted habitat loss. For range-contracting species we found a convex
relationship, meaning that species initially retained stable population sizes despite ongoing
habitat loss while population sizes dropped very suddenly for higher rates of habitat loss.
This is consistent with a non-uniform distribution of individuals in the landscape, where
sparsely populated edge habitats are lost first during range-contraction[21,34]. Additionally, the
curve is indicative of an extinction threshold, a critical amount of habitat beyond which
population size declines steeply[23]. Crossing the threshold can initiate the extinction vortex, in
which reinforcing processes such as inbreeding depression and genetic drift increase

population decline, ultimately leading to species extinction[35]. Even though we did not
consider species genetics or stage-structured populations in our simulation, we still observed
an extinction threshold. In our simulations, this threshold was around 50% habitat loss, but
empirical evidence shows that these thresholds are highly case specific and species
dependent[23]. A convex shape of population decline over time was already reported for
several species[36], but research on the exact relationship between population size, habitat
loss and extinction risk remains limited[20]. While the linear assumption in Red List guidelines
is violated in these cases, our results suggest that SDM-based assessment still offer
sufficient warning time for range-contracting species.

In contrast, range-shifting species exhibited a concave relationship indicating a strong
underestimation of range loss by SDMs. This likely reflects the inability of some species to
fully track shifting climates due to dispersal limitations[37,38] and time lags[39,40]. The
underestimation resulted in shorter warning time for SDM-based assessment of species in
comparison to true population size. Because effective conservation measures often require
several decades of advance warning, SDM-based assessment may not leave enough
warning time to protect range-shifting species from extinction[29]. Stronger dispersal ability is
expected to improve climate tracking and the survival of range-shifting species under climate
change[32,41]. Interestingly, we found no such influence of dispersal ability on the classification
time of range-shifting species but found delayed warning for all range-shifting species. As
inclusion of simple dispersal assumptions did not change these results, our study suggests
that SDMs will not be adequate for assessing extinction risk in range-shifting species
threatened with climate change. For these species, spatially explicit population models are
likely more adequate.

Our results further revealed that Red Listing based on probabilistic extinction estimates
under criterion E led to delayed classifications and insufficient warning times for all species,
particularly those most at risk. This aligns with previous findings that criterion E often results
in less precautionary extinction risk status in comparison to other criteria[42]. In our study, the
short warning time resulted from steep, sigmoidal extinction risk curves, which led to rapid
population collapse and abrupt extinction of species. Such curve shapes have been shown
to realistically describe species' survival probabilities under changing environmental
conditions[43,44]. In stochastic population models, sigmoidal extinction curves typically emerge
when full parameter knowledge during the model building process is available[45]. With higher
parameter uncertainty the extinction risk curves are expected to become more shallow[45],
which consequently influences the classification time in the Red List. Interestingly, a
previous study[29] reported that criterion A leaves less warning time in comparison to listing

against criterion E when using a coupled SDM-population model approach with a real
species. It is thus unclear whether a higher parameter uncertainty in the model building
process for criterion E might leave more warning time or may introduce additional biases. As
a more robust approach for species threatened by climate change, especially those
undergoing range shifts, we recommend estimating expected minimum abundance (EMA)
from SEPMs. Our results suggest that assessing extinction risk based on these projections
provides a more reliable basis for classification than relying on quantitative extinction
probabilities alone.

Our simulations were idealised, with full knowledge of the true population and range
dynamics, perfect data to fit SDMs, and simplified assumptions. In practice, missing data
and sampling biases as well as environmental and demographic stochasticity could
introduce additional uncertainties in the assessment process[45–47]. Here, we want to echo the
Red List guidelines in stressing transparency regarding uncertainties in the assessment
process to minimize their effect[8(pp.23-25)]. Additionally, simulating realistic climate change
scenarios and more detailed species characteristics, such as stage-structured populations,
stochastic movement dispersal and density-dependent traits could lead to more diverse
responses of species influencing the assessment process. Furthermore, climate change is
known to not only influence the distribution of species but also have diverse effects on
species[48], which we did not consider in our study. Nevertheless, our results revealed that
even under idealised conditions, current Red List criteria are highly sensitive to species
range dynamics under climate change.

To improve assessment, we recommend including a screening step into model-based Red
List assessments. In this initial screening, SDMs can serve to assess whether a species is
likely to experience exclusive range contraction or to shift its range. For range-shifting
species, we strongly recommend using spatially explicit population models and assess
extinction risk based on expected minimum abundance, rather than extinction probabilities.
While adjusting probability thresholds may improve assessments in some cases, further
research is needed to define thresholds that offer both realism and sufficient warning time.
For range-contracting species, SDMs may be appropriate to assess the extinction risk of
species. Still, the assumption of a linear relationship between population size and habitat
size should be treated with caution as our results show that sudden population reductions
can cause delayed warning by SDMs. Furthermore, the known limitations of SDMs should
be kept in mind[18] and the assessment based on SDMs should follow community
standards[49,50] to avoid wrong classifications. To further guide the novelization of the Red List

under climate change, we recommend empirically testing our results for a variety of species
for which adequate range loss and population loss data are available.

Methods

We used the RangeShifter framework[33,51,52] to simulate population dynamics of virtual
species under equilibrium conditions and under climate change in three different artificial
landscapes. Species differed in four traits: niche position (central vs. marginal), niche width
(narrow vs. wide), growth rate (slow vs. fast) and dispersal (short vs. long), leading to 16 (2^4)
different species to cover all trait combinations. Niches were defined by two environmental
variables that could be interpreted as temperature and precipitation. Species occurrence
data and climate data were extracted from the simulation at equilibrium before climate
change set in. Based on these data we fitted species distribution models (SDMs) and
predicted habitat suitability under climate change. From the simulations, we observed
population size across the entire time frame of 90 years of climate change and estimated
extinction probability per year. We ran simulations for 90 years to ensure the extinction of all
species. From the SDMs, we derived the predicted habitat suitability values for each year
and compared these to true population size. Last, we applied the Red List criteria to each of
these three metrics to classify species into different threatened categories.

Virtual species
In our simulation approach, we first modelled the population dynamics of the 16 virtual
species in the three artificial landscapes under climate change. The artificial landscape
consisted of two environmental variables (temperature and precipitation) with a spatial extent
of 512 x 512 cells at 1000 m resolution. The environmental layers were generated using the
package "NLMR"[53]. Temperature was represented by a latitudinal gradient with spatially
auto-correlated noise. For this the arithmetic mean of a landscape with a linear gradient and
a landscape built with midpoint displacement with a Hurst exponent of 0.96 was taken. To
create spatially auto-correlated precipitation patterns, we generated a landscape using
midpoint displacement with a Hurst exponent of 0.75. Landscape values ranged between 0
and 1. We derived three replicate landscapes using the same settings (Fig. S8).

The virtual species niche was modelled using the package "virtualspecies"[54]. First,
Gaussian-shaped response curves for both environmental variables were simulated. The
mean of the response curve for temperature was adapted to the respective niche position
(marginal: 0.27, central: 0.5), the mean of the precipitation response curve was set to 0.5. To
model different niche breadths, the standard deviation of both response curves was adapted

based on the respective niche breadth (narrow: 0.045, wide: 0.055). We then used the
product of both response curves to map the virtual species niche in the artificial landscapes.
To simulate climate change, temperature was linearly increased over 90 years, while
precipitation remained static. We selected a time frame of 90 years to model a continuous
increase of temperature that would result in no suitable habitat being left at the end of
climate change, ensuring the extinction of all species. To the linear temperature increase we
also included a temporal autocorrelated noise. For this we used an ARIMA model[55] of the
order 0,0,1 and added it to the linear model using the following equations:

$\eta_t = \theta * \varepsilon_t - 1 + \varepsilon_t$

$y_t = \alpha + \beta * t + \eta_t$

Where, η_t represents the time series obtained from the ARIMA model with θ set to 0.5 and ε
representing the residuals at the respective time points. This time series was then added to
a linear model with α set to 0.5, β set to 0.6, and t representing the respective year. The time
series was rescaled between 0 and 0.9 and added to the temperature value of every cell in
the artificial landscape. Values that were larger than 1 were reduced to 1 to stay within the
range of 0 to 1.

To model the population dynamics of the virtual species, we used the spatially explicit and
individual-based modelling platform RangeShifter[33,51], operated via the package
"RangeShiftR"[52]. We implemented a cell-based model using the previously created maps of
the virtual species niche with the resolution set to 1000 m and the carrying capacity set to
0.05 ind/ha. We modelled the virtual species as an asexual species with non-overlapping
generations meaning that each individual only lived for one year. The competition coefficient
was set to 1 (i.e. no over- or undercompensation) and the value for the maximum growth
rate was adapted according to the growth rate (slow: 3, fast: 5). After reproduction the adults
died and the offspring dispersed. Dispersal is modelled explicitly in three phases in
RangeShifter: emigration, transfer, and settlement. The emigration probability for each
individual was set to 0.4 and was density-independent. The transfer was modelled using a
negative exponential dispersal kernel. For short-distance dispersal, the mean dispersal
distance was set to 5000 m. For long-distance dispersal, a double exponential dispersal
kernel was used to include rare long-distance dispersal events. The double kernel is
composed of two negative exponential kernels with different means and probabilities of
occurrence. The first (shorter) kernel occurred with a probability of 95% and had a mean
dispersal distance of 15 000 m, the second (longer) kernel occurred with a probability of 5%
and had a mean dispersal distance of 250 000 m. If the arrival point of an individual went
beyond the boundaries of the landscape or within the natal cell the dispersal process was
repeated. The individual settled in the arrival cell if the cell contained suitable habitat. If the

arrival cell was not suitable, the individual was displaced to one of the eight neighbouring
cells, if one of them was suitable, or else died. If a cell became unsuitable due to climate
change, all individuals inhabiting that cell died immediately.
378
The simulation was initialised using the map with equilibrium climatic conditions and
individuals were distributed in all suitable cells at half-carrying capacity. The simulation ran
for 101 spin-up years to remove any initialisation effect. After the spin-up period, the
landscape was changed yearly to represent the virtual species niche shifting northwards in
response to climate change. In the following, year 100 will be referred to as year 0 and year
1-90 represent the years under climate change. Each of the 16 species was simulated with
100 replicate runs for each of the three artificial landscapes.
386
387 From the output of each simulation, we obtained the population size of the species for each
388 year and replicate run. To calculate the extinction probability for each year, we divided the
389 number of replicate runs without a viable population of that year by the total number of
390 replicate runs.
391

Species distribution models

At the end of the spin-up period, before climate change set in, we extracted species
presence and absence information as well as climate variables from the simulation to fit
SDMs. To account for variation between the replicate runs, we randomly selected ten
replicate runs from the simulation and fitted separate SDMs to each. This resulted in 480
SDMs (16 species x 3 landscape replicates x 10 replicate runs). For the species data we
assumed perfect sampling with no error in detection. Data points were partially thinned by
selecting every second cell to avoid spatial autocorrelation using the package "dismo"[56].
Based on the recommendations of the Red List[8(p.108)] we used three distinct modelling
algorithms (Generalised Linear Model (GLM), Random Forest and MaxEnt) and further
calculated an ensemble model based on the arithmetic mean. For the GLM, linear and
quadratic predictors were used with a binomial error distribution. For Random Forest, we
fitted 1000 trees with a minimum terminal node size of 20 to fit comparably smooth response
surfaces. In the MaxEnt algorithm, we used the feature classes linear, quadratic and hinge.
The probability of occurrence was predicted for each of the 90 years under climate change
using the ensemble model. The predictive performance of the model algorithms and the
ensemble model was evaluated using the 99 replicate runs that were not used for fitting the
SDM. For the performance measures we selected the AUC (area under the ROC curve) and
TSS (true skill statistics)[16]. The calculated performance measures were averaged over all 99
replicate runs using the arithmetic mean. For the calibration and evaluation of the SDMs we

used the following packages: "randomForest"[57], "maxnet"[58], "gbm"[59], "PresenceAbsence"[60]

and "lhs"[61].

For testing the effect of dispersal assumptions in SDMs on the assessment in the Red List,

we first binarized the predictions of each species and landscape replicate using the maxTSS

threshold[62]. Around the known presences we created a buffer with the size of the estimated

dispersal distance of an individual in the respective time frame and set the predicted habitat

suitability outside the buffer to 0 following the recommendation of the Red List

guidelines[8(p.110)]. Buffer sizes were determined by the empirical dispersal distances observed

from the RangeShifter simulations. For this, we extracted the dispersal distances of all

emigrated individuals for year 0 from the simulations, and calculated the arithmetic mean of

dispersal distance.

Red List classification

To assess Red Listing of species threatened by climate, we compared the assessment of

the species against two different Red List criteria using three metrics. For the criterion A3,

we used true population size observed from the RangeShifter simulations and the sum of the

SDM-derived habitat suitability values. For the criterion E, we used the extinction probability

for every species obtained from the RangeShifter simulations. For the population size, we

only used the population size from the ten replicate runs that were selected for fitting the

SDMs. To calculate the sum of the habitat suitability values from the SDM predictions we

followed the recommendations from the Red List guidelines[8(p.99)]. We first removed cells with

values below a certain threshold value, for which we used the maxTSS[62]. We then summed

up the predicted habitat suitability of the remaining cells and for every year. For the

classification into the different categories of the Red List, different thresholds and time

frames apply for the different criteria (see Tab. 1).

448 Table 1: Thresholds and time frames used to classify species into threatened categories, based on
449 the Red List Criteria[8]

Criterion	Red List Threatened Categories		
	Critically Endangered	Endangered	Vulnerable
A3: future population size reduction / future reduction of sum of SDM-derived habitat suitability	>= 80% in 10 years	>= 50% in 10 years	>= 30% in 10 years
E: probability of extinction	>= 50% in 10 years	>= 20% in 20 years	>= 10% in 100 years

For each species, we determined the time of classification into the three different threatened
categories using all three metrics (i.e., population size, habitat suitability and extinction
probability). For this, we used a moving window approach, testing for every year if the
species would fulfil the Red List criteria in the following years (see Table 1 for considered
time spans). This means that if a species was predicted to reach a habitat loss of 30%
between year 0 and year 10, the year 0 would be marked as classification time point for
listing the species as "Vulnerable" under criterion A3 when using the metric sum of SDM-
derived habitat suitability values. If the species was predicted to reach a habitat loss of 30%
between year 25 and 35, then the year 25 would be marked as classification time for the
"Vulnerable" category. Thereby, the loss of suitable habitat and the population loss was
always calculated relative to the starting year of the moving window. As all species
continuously experienced population and habitat loss under climate change until their
extinction, each species qualified for all threatened categories throughout their life span.
In a sensitivity analysis, the assessment of classification times based on SDM-derived
habitat suitability values was repeated for more realistic dispersal assumptions using the
buffer approach described earlier.

Statistical analysis

To investigate the relationship between population size and habitat loss, we fitted a
generalised mixed-model ANCOVA using an ordered beta regression implemented by the
package "ordbetareg"[63,64]. This package is a front-end to the package "brms"[65]. We selected

an ordered beta regression since our response variable (simulated "true" populated size)
was continuous with a zero and one inflation. As predictors, we considered habitat loss as
continuous variable, the four traits (niche position, niche width, dispersal distance and
growth rate) as categorical variable with two levels each, and their pairwise interactions with
habitat loss. We further controlled for the landscapes by including them as random
intercepts. The best model was selected based on the difference in expected log predictive
density. Posterior distributions were estimated using Markov chain Monte Carlo (MCMC)
sampling with 4 chains and a delta value of 0.95. Each chain had a length of 6000 iterations,
with the first 2000 being used as a warm-up. For the prior, the default was selected.
481
482 For the study R 4.2.2[66] was used. In addition to the already mentioned packages the
483 following were used for data preparation and visualization: "scales"[67], "dplyr"[68], "ggplot2"[69],
484 "terra"[70], "gridExtra"[71], "tibble"[72], "RColorBrewer"[73], "doParallel"[74], "foreach"[75], "data.table"[76].
485

Code availability:
487 The codes for this study are available at https://doi.org/10.5281/zenodo.16909451
488

Author Contributions:
490 RK and DZ conceptualised the study. RK ran all analyses and led the writing. All authors
491 contributed to revising the text.
492

Competing Interest Statement:
494 The authors declare that they have no conflict of interest to disclose.
495

Acknowledgments:
497 This study was supported by the German Research Foundation DFG (grant no. ZU 361/6-1).
498 S.A.F. gratefully acknowledges the support of iDiv funded by the German Research
499 Foundation (DFG– _FZT 118, 202548816) and of the Leibniz Competition by the Leibniz
500 Association (P52/2017).

509 **References:**

1. Scheffers, B. R. *et al.* The broad footprint of climate change from genes to biomes to people. *Science* **354**, aaf7671 (2016).

2. Freeman, B. G., Scholer, M. N., Ruiz-Gutierrez, V. & Fitzpatrick, J. W. Climate change causes upslope shifts and mountaintop extirpations in a tropical bird community. *Proceedings of the National Academy of Sciences* **115**, 11982–11987 (2018).

3. Urban, M. C. Climate change extinctions. *Science* **386**, 1123–1128 (2024).

4. Bellard, C., Bertelsmeier, C., Leadley, P., Thuiller, W. & Courchamp, F. Impacts of climate change on the future of biodiversity. *Ecology Letters* **15**, 365–377 (2012).

5. Pecl, G. T. *et al.* Biodiversity redistribution under climate change: Impacts on ecosystems and human well-being. *Science* **355**, eaai9214 (2017).

6. Stanton, J. C., Shoemaker, K. T., Pearson, R. G. & Akçakaya, H. R. Warning times for species extinctions due to climate change. *Global Change Biology* **21**, 1066–1077 (2015).

7. IUCN. *IUCN Red List Categories and Criteria: Version 3.1.* (2001).

8. IUCN Standards and Petitions Committee. *Guidelines for Using the IUCN Red List Categories and Criteria. Version 16*. (2024).

9. Mace, G. M. & Lande, R. Assessing Extinction Threats: Toward a Reevaluation of IUCN Threatened Species Categories. *Conservation Biology* **5**, 148–157 (1991).

10. Mace, G. M. *et al.* Quantification of extinction risk: IUCN's System for Classifying Threatened Species. *Conservation Biology: The Journal of the Society for Conservation Biology* **22**, 1424–1442 (2008).

11. Akçakaya, H. R., Butchart, S. H. M., Mace, G. M., Stuart, S. N. & Hilton-Taylor, C. Use and misuse of the IUCN Red List Criteria in projecting climate change impacts on biodiversity. *Global Change Biology* **12**, 2037–2043 (2006).

12. IUCN Standards and Petitions Committee. *Guidelines for Using the IUCN Red List Categories and Criteria. Version 8.0*. (2010).

13. Valavi, R., Guillera-Arroita, G., Lahoz-Monfort, J. J. & Elith, J. Predictive performance of presence-only species distribution models: a benchmark study with reproducible code. *Ecological Monographs* **92**, e01486 (2022).

14. Urban, M. C. Accelerating extinction risk from climate change. *Science* **348**, 571–573 (2015).

15. Wüest, R. O. *et al.* Macroecology in the age of Big Data – Where to go from here? *Journal of Biogeography* **47**, 1–12 (2020).

16. Guisan, A., Thuiller, W. & Zimmermann, N. E. *Habitat Suitability and Distribution Models: With Applications in R*. (Cambridge University Press, Cambridge, 2017).

17. Araújo, M. B. & Guisan, A. Five (or so) challenges for species distribution modelling. *Journal of Biogeography* **33**, 1677–1688 (2006).

18. Elith, J. & Leathwick, J. R. Species Distribution Models: Ecological Explanation and Prediction Across Space and Time. *Annual Review of Ecology, Evolution, and Systematics* **40**, 677–697 (2009).

19. Weber, M. M., Stevens, R. D., Diniz-Filho, J. A. F. & Grelle, C. E. V. Is there a correlation between abundance and environmental suitability derived from ecological niche modelling? A meta-analysis. *Ecography* **40**, 817–828 (2017).

20. Zurell, D., Fritz, S. A., Rönnfeldt, A. & Steinbauer, M. J. Predicting extinctions with species distribution models. *Cambridge Prisms: Extinction* **1**, e8 (2023).

21. Rodríguez, J. P. Range Contraction in Declining North American Bird Populations. *Ecological Applications* **12**, 238–248 (2002).

22. Kuussaari, M. *et al.* Extinction debt: a challenge for biodiversity conservation. *Trends Ecol Evol* **24**, 564–571 (2009).

23. Fahrig, L. How much habitat is enough? *Biological Conservation* **100**, 65–74 (2001).

24. Keith, D. A. *et al.* Predicting extinction risks under climate change: coupling stochastic population models with dynamic bioclimatic habitat models. *Biology Letters* **4**, 560–563 (2008).

25. Maschinski, J., Baggs, J. E., Quintana-Ascencio, P. F. & Menges, E. S. Using Population Viability Analysis to Predict the Effects of Climate Change on the Extinction Risk of an Endangered Limestone Endemic Shrub, Arizona Cliffrose. *Conservation Biology* **20**, 218–228 (2006).

26. Akcakaya, H. R. & Sjögren-Gulve, P. Population Viability Analyses in Conservation Planning: An Overview. *USA. — Per Sjögren-Gulve* **11733**, (2000).

27. Briscoe, N. J. *et al.* Forecasting species range dynamics with process-explicit models: matching methods to applications. *Ecology Letters* **22**, 1940–1956 (2019).

28. Cazalis, V. *et al.* Bridging the research-implementation gap in IUCN Red List assessments. *Trends in Ecology & Evolution* **37**, 359–370 (2022).

29. Keith, D. A. *et al.* Detecting Extinction Risk from Climate Change by IUCN Red List Criteria. *Conservation Biology* **28**, (2014).

30. Zurell, D. *et al.* The virtual ecologist approach: simulating data and observers. *Oikos* **119**, 622–635 (2010).

31. Zurell, D. *et al.* Benchmarking novel approaches for modelling species range dynamics. *Global Change Biology* **22**, 2651–2664 (2016).

32. Foden, W. B. *et al.* Identifying the World's Most Climate Change Vulnerable Species: A Systematic Trait-Based Assessment of all Birds, Amphibians and Corals. *PLOS ONE* **8**, e65427 (2013).

33. Bocedi, G. *et al.* RangeShifter 2.0: an extended and enhanced platform for modelling spatial eco-evo-lutionary dynamics and species' responses to environmental changes. *Ecography* **44**, 1453–1462 (2021).

34. Sagarin, R. D. & Gaines, S. D. The 'abundant centre' distribution: to what extent is it a biogeographical rule? *Ecology Letters* **5**, 137–147 (2002).

35. Gilpin, M. E. & Soulé, M. E. Minimum viable populations: processes of extinction. in *Conservation Biology: The Science of Scarcity and Diversity* 19–34 (Sinauer Associates, Sunderland, MA, 1986).

36. Fagan, W. F. & Holmes, E. E. Quantifying the extinction vortex. *Ecology Letters* **9**, 51–60 (2006).

37. Travis, J. M. J. Climate Change and Habitat Destruction: A Deadly Anthropogenic Cocktail. *Proceedings: Biological Sciences* **270**, 467–473 (2003).

38. Hof, C., Levinsky, I., Araújo, M. B. & Rahbek, C. Rethinking species' ability to cope with rapid climate change. *Global Change Biology* **17**, 2987–2990 (2011).

39. Lenoir, J. *et al.* Species better track climate warming in the oceans than on land. *Nat Ecol Evol* **4**, 1044–1059 (2020).

40. Chen, I.-C., Hill, J. K., Ohlemüller, R., Roy, D. B. & Thomas, C. D. Rapid Range Shifts of Species Associated with High Levels of Climate Warming. *Science* **333**, 1024–1026 (2011).

41. Mancini, G. *et al.* A standard approach for including climate change responses in IUCN Red List assessments. *Conservation Biology* **38**, e14227 (2024).

42. Gärdenfors, U. Population Viability Analysis in the Classification of Threatened Species: Problems and Potentials. *Ecological Bulletins* 181–190 (2000).

43. Ball, T. *et al.* A general relationship between carrying capacity and extinction risk. Preprint at https://doi.org/10.33774/coe-2024-m9dqp-v2 (2025).

44. Dushoff, J. Carrying Capacity and Demographic Stochasticity: Scaling Behavior of the Stochastic Logistic Model. *Theoretical Population Biology* **57**, 59–65 (2000).

45. Taylor, B. L. The Reliability of Using Population Viability Analysis for Risk Classification of Species. *Conservation Biology* **9**, 551–558 (1995).

46. Akcakaya, H. R. Population viability analyses with demographically structured models. *Ecological Bulletins* **48**, 23–38 (2000).

47. Possingham, H. P., McCarthy, M. A. & Lindenmayer, D. B. Population Viability Analysis. in *Encyclopedia of Biodiversity (Second Edition)* (ed. Levin, S. A.) 210–

219 (Academic Press, Waltham, 2013). doi:10.1016/B978-0-12-384719-5.00173-8.

48. Walther, G.-R. *et al.* Ecological responses to recent climate change. *Nature* **416**, 389–395 (2002).

49. Araújo, M. B. *et al.* Standards for distribution models in biodiversity assessments. *Sci Adv* **5**, eaat4858 (2019).

50. Sofaer, H. R. *et al.* Development and Delivery of Species Distribution Models to Inform Decision-Making. *BioScience* **69**, 544–557 (2019).

51. Bocedi, G. *et al.* RangeShifter: a platform for modelling spatial eco-evolutionary dynamics and species' responses to environmental changes. *Methods in Ecology and Evolution* **5**, 388–396 (2014).

52. Malchow, A.-K., Bocedi, G., Palmer, S. C. F., Travis, J. M. J. & Zurell, D. RangeShiftR: an R package for individual-based simulation of spatial eco-evolutionary dynamics and species' responses to environmental change. *Ecography* (2021) doi:10.1111/ecog.05689.

53. Sciaini, M., Fritsch, M., Scherer, C. & Simpkins, C. E. NLMR and landscapetools: An integrated environment for simulating and modifying neutral landscape models in R. *Methods in Ecololgy and Evolution* **00**, 1–9 (2018).

54. Leroy, B., Meynard, C. N., Bellard, C. & Courchamp, F. virtualspecies, an R package to generate virtual species distributions. *Ecography* (2015) doi:10.1111/ecog.01388.

55. Box, G. E. P. & Jenkins, G. M. *Time Series Analysis: Forecasting and Control*. (San Francisco: Holden-Day, 1970).

56. Hijmans, R. J., Phillips, S., Leathwick, J. & Elith, J. *Dismo: Species Distribution Modeling*. (2021).

57. Liaw, A. & Wiener, M. Classification and Regression by randomForest. *R News* **2**, 18–22 (2002).

58. Phillips, S. *Maxnet: Fitting 'Maxent' Species Distribution Models with 'Glmnet'*. (2021).

59. Greenwell, B., Boehmke, B., Cunningham, J. & Developers, G. B. M. *Gbm: Generalized Boosted Regression Models*. (2020).

60. Freeman, E. A. & Moisen, G. PresenceAbsence: An R Package for Presence Absence Analysis. *Journal of Statistical Software* **23**, 1–31 (2008).

61. Carnell, R. *Lhs: Latin Hypercube Samples*. (2024).

62. Liu, C., Berry, P. M., Dawson, T. P. & Pearson, R. G. Selecting thresholds of occurrence in the prediction of species distributions. *Ecography* **28**, 385–393 (2005).

63. Kubinec, R. Ordered Beta Regression: A Parsimonious, Well-Fitting Model for Continuous Data with Lower and Upper Bounds. *Political Analysis* **31**, 519–536 (2023).

64. Kubinec, R. *Ordbetareg: Ordered Beta Regression Models with 'Brms'*. (2025).

65. Bürkner, P.-C. brms: An R Package for Bayesian Multilevel Models Using Stan. *Journal of Statistical Software* **80**, 1–28 (2017).

66. R Core Team. *R: A Language and Environment for Statistical Computing*. (R Foundation for Statistical Computing, Vienna, Austria, 2022).

67. Wickham, H. & Seidel, D. *Scales: Scale Functions for Visualization*. (2022).

68. Wickham, H., François, R., Henry, L., Müller, K. & Vaughan, D. *Dplyr: A Grammar of Data Manipulation*. (2023).

69. Wickham, H. *Ggplot2: Elegant Graphics for Data Analysis*. (Springer-Verlag New York, 2016).

70. Hijmans, R. J. *Terra: Spatial Data Analysis*. (2023).

71. Auguie, B. *gridExtra: Miscellaneous Functions for 'Grid' Graphics*. (2017).

72. Müller, K. & Wickham, H. *Tibble: Simple Data Frames*. (2022).

73. Neuwirth, E. *RColorBrewer: ColorBrewer Palettes*. (2014).

74. Weston, S. & Microsoft. *doParallel: Foreach Parallel Adaptor for the 'parallel' Package*. (2022).

75. Microsoft & Weston, S. *Foreach: Provides Foreach Looping Construct*. (2022).

76. Barrett, T. *et al. Data.Table: Extension of `data.Frame`*. (2024).