Red List criteria underestimate climate-related extinction risk of
range-shifting species

Raya Keuth[1,*], Susanne A. Fritz[2,3], Damaris Zurell[1]

Affiliations:
[1] University of Potsdam, Institute of Biochemistry and Biology, Potsdam, Germany
[2] German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Germany
[3] Friedrich Schiller University Jena, Institute of Biodiversity, Ecology and Evolution, Jena,
Germany

*corresponding author: raya.keuth@uni-potsdam.de

ORCIDs: RK: 0000-0002-6694-8887; SF: 0000-0002-4085-636X; DZ: 0000-0002-4628-3558

Abstract
Climate change causes global species redistribution and elevates extinction risk, making
early identification of vulnerable species critical for timely conservation. The IUCN Red List
provides guidelines for assessing climate-related extinction risk using species distribution
models (SDMs) and spatially explicit population models (SEPMs). However, a systematic
evaluation of these guidelines is currently missing. Using simulations of virtual species with
diverse life-history traits and range dynamics, we found that SDMs consistently
underestimated extinction risk for range-shifting species. This was due to a concave
relationship between population size and habitat loss, which contradicts the linear
assumption in the Red List guidelines. For range-contracting species, SDMs provided
adequate warning times. Probabilistic extinction estimates from SEPMs provided delayed
warning for all species, particularly for highly threatened ones. Our results reveal key
limitations of current Red List guidelines under climate change. Based on our findings, we
provide tentative recommendations for updating the IUCN Red List guidelines.

Keywords: IUCN Red List, extinction risk, species distribution models, climate change,
virtual species, spatially explicit population models

Introduction

Climate change poses a major threat to biodiversity, affecting virtually all biological

processes[1] and potentially elevating extinction risk for many species[2,3]. Species dynamically

respond to climate change, for example by tracking suitable habitat[4], which has been

projected to lead to substantial global redistribution of species[5]. To protect biodiversity

before declines become irreversible, early identification of the most threatened species is

essential. Conservation assessments should hence provide sufficient warning time, defined

as the time between identification of a threatened species and its extinction without any

interventions[6].

The IUCN Red List for Threatened Species (Red List hereafter) is the most widely used

framework for assessing extinction risk and informing conservation and policy decisions[7,8]. It

classifies species into one of three threatened categories (Vulnerable, Endangered, Critically

endangered) based on five quantitative criteria (A-E) that are derived from the theory of

extinction trajectories for single populations[7,9,10]. The five different criteria provide alternative

ways of assessing a species in the Red List, with qualifying for a threatened category under

one criterion being sufficient to classify species in the category. The assessment of the

species is mostly based on population size and trends, as well as geographic range area[9].

Originally designed to identify species already in decline, the IUCN has since introduced

guidelines for assessing future extinction risk under climate change. These recommend

species distribution models (SDMs) to infer habitat loss, or spatially explicit population

models (SEPMs) to infer population loss or estimate extinction probability[11,12].

Despite their growing use, the predictive performance of SDMs under climate change

remains debated[13–15]. SDMs are correlative models that relate observed species occurrences

to prevailing environmental conditions and use this species-environment relationship to

predict the suitable habitat under future climatic scenarios[16]. However, SDMs omit key

biological processes, such as demography and dispersal[17,18], and simplistic dispersal

assumptions (e.g. no dispersal or full dispersal) can lead to an over- or underestimation of

the expected range size[14]. Nevertheless, SDMs currently constitute the most pragmatic

approach for projecting potential future climate impacts. The Red List guidelines suggest to

use SDM-predicted changes in habitat suitability as a proxy for population size reduction

under criterion A3[8(p.102)]. This approach assumes a linear relationship between population

loss and SDM-predicted habitat loss, which is acknowledged in the Red List guidelines as a

source of uncertainty[8(p.99)]. Beyond simply causing uncertainty, this assumption actually lacks

empirical validation[19,20], and could be influenced by the non-uniform distribution of species in

the landscape[21], extinction debts[22], and extinction thresholds related to minimum viable

population sizes[23].

SEPMs, in contrast, simulate demographic and dispersal processes over space and time,

potentially offering more realistic predictions of species responses to climate change[24,25]. Yet,

wider application of these models has been hampered by technical and data challenges[26,27]

and they have thus been rarely used for assessing climate-related future extinction risk of

species[28]. A key open question is whether SEPMs provide more accurate or timely extinction

risk estimates than SDMs, particularly for species with different range dynamics under

climate change. So far, only a single case study has compared these modelling approaches

in the context of IUCN assessments[29] while a systematic evaluation of modelling approaches

across varying life histories and range dynamics is missing. This gap prevents evidence-

based improvements to current Red List guidelines.

In this study, we aim to test the Red List guidelines for assessing species threatened by

climate change. Specifically, we test (i) whether the assumed linear relationship between

habitat loss and population decline holds across species and range dynamics, and (ii)

whether model-based assessments differ in their ability to identify extinction risk early

enough for conservation action. Due to the lack of data regarding climate change-driven

extinctions, we chose a simulation approach[30,31]. We generated sixteen virtual species with

contrasting life-history and dispersal traits (niche position, niche breadth, growth rate and

dispersal distance)[32] and used a spatially explicit modelling framework[33] to project population

and range dynamics under climate change across three different stochastically replicated

artificial landscapes obtained from a neutral landscape model (Fig. 1). Although all species

eventually went extinct due to habitat loss, half of the species (those with central niche

positions) first underwent a range shift before extinction, while the other half of the species

(those with marginal, cold-adapted niches) only experienced range contractions, also leading

to extinction. Finally, we compared the performance of SDMs and SEPMs for assessing

extinction risk, focusing on their ability to provide timely warnings for species with varying life

history traits and undergoing either range contraction or range shift.

Figure 1: Conceptual figure of the modelling workflow. We simulated virtual species under climate change using a spatially explicit individual-based model. Species varied in four traits, which resulted in a variety of range distributions and dynamics. Before climate change set in, distribution data were sampled from the simulations as input to train correlative species distribution models (SDMs). SDMs were then used to predict habitat loss under climate change. Finally, Red List classifications under climate change were compared for SDM-derived habitat loss (criterion A3) and simulation-derived population size reduction (criterion A3) and extinction probability (criterion E).

Results

Population dynamics and SDM performance

The considered niche traits led to distinct differences in range and population dynamics (here, population refers to the number of individuals across the entire landscape), while the traits growth rate and dispersal only showed minor effects (Fig. S1 + S2). Notably, species with marginal niche position mainly occurred at the "northern" edge of the landscape and their ranges contracted towards that edge under climate change. Species with central niche position had centrally located ranges and exhibited range shifting before going extinct under climate change (Fig. 2). Notably, not all range-shifting species (i.e., species with a central niche position) were able to fully shift their ranges and reach the northern boundary of the landscape. Some species became extinct earlier due to dispersal limitations, as suitable habitat was still available in the northern part of the landscape (Fig. 2 + Fig. S1). As could be expected, species with slow growth, narrow niches and short dispersal went extinct earlier than species with fast growth, wide niches and long dispersal (Fig. S1). The differences in temporal dynamics between species with different trait values were more pronounced for range-shifting species than for range-contracting species.

We sampled species occurrence from the simulations before climate change set in, and

fitted three different SDM algorithms and an ensemble model. The SDM ensembles showed

high pre-climate change predictive accuracy for all species and landscapes (AUC: 0.95 ±

0.02; TSS: 0.78 ± 0.06). The predictive performance was slightly lower for species with

central compared to marginal niches and for species with short compared to long dispersal

(Fig. S3 + S4). Under climate change, SDMs correctly predicted range loss of range-

contracting species (i.e., the distribution of simulated abundances aligned well with the SDM-

derived habitat predictions), but they underestimated the range loss of dispersal-limited

range-shifting species (i.e., the distribution of simulated abundances was smaller than the

SDM-derived habitat distribution; Fig. 2). This pattern was consistent across replicate

landscapes (Fig. S1).

Figure 2: Distribution of virtual species and SDM-predicted habitat suitability for two example

species. Top panels (a) show the maps of simulated „true" species abundance and maps of habitat

suitability predicted by SDMs for one replicate run under equilibrium conditions (year 0) and under

climate change (year 20 / 30) in one example landscape. Bottom panels (b) show the corresponding

time trajectories of the relative change in habitat suitability sums predicted from SDMs and in

simulated population size, both averaged over ten selected replicate runs, as well as simulated

extinction probability. Results are shown for two example species differing in their range dynamics

(i.e., in marginal vs. central niche position). Both species have a narrow niche, slow growth rate, and

short dispersal.

Population size - habitat loss relationship

For all species and replicate landscapes, the relationship between relative population size
and SDM-derived habitat loss observed across simulated species and landscapes deviated
from the linear assumption proposed in the Red List guidelines (Fig. 3, S5 + S6). For range-
contracting species (with marginal, cold-adapted niche positions), the relationship was
convex with an initial smaller rate of decline of population size in comparison to habitat loss.
Population size steeply declined once habitat loss exceeded 50% across all replicate
landscapes (Fig. S5 + S6). In contrast, range-shifting species (with central niche positions)
showed a concave relationship with considerable declines in population size already
happening at low rates of habitat loss (Fig. 3). Yet, we also found considerable differences in
the population size - habitat loss relationship for range-shifting species in different replicate
landscapes, with some being closer to a linear relationship (Fig. S5 + S6). We only found
comparatively small differences in the population size - habitat loss relationships for the
other traits, with the variability between the different landscapes often being larger than
among the different trait levels (Fig. S5 + S6). We quantified the relative importance of the
different traits using a Bayesian generalised mixed-model analysis of covariance. This
revealed that the population size - habitat loss relationship significantly differed across all
traits, but the effect sizes for niche position (i.e., for range-shifting vs. range-contracting
species) were considerably larger than for all other traits (Table S1).

Figure 3: Population size - habitat loss relationship for the different range dynamics. Relative
population size was obtained from the simulation (observed true data for virtual species) and habitat
loss was predicted by the SDMs. The dot-dash grey line shows the assumption of a linear relationship
between population size and habitat loss as proposed by the Red List guidelines. The relationships
are displayed using an ordered beta regression model, which was fitted using the "true" population
size values obtained from the simulations as response variable and habitat loss, and the 16 trait
combinations across the 10 selected replicate runs as predictor variables, and in which landscape
was included as a random effect. Lines indicate the mean model predictions and the transparent
bands the 95% confidence interval. We only show results here for the trait niche position, resulting in
range-contracting and range-shifting species, while results for the other traits are displayed in Fig. S5.

Red List classification

An important question is to which extent SDM-derived habitat loss under climate change can
be used for assessment in the Red List. To answer this, we determined the time at which a
species would first be classified into each threatened category of the Red List: VU, EN, and
CR (Vulnerable, Endangered, and Critically endangered). Specifically, we compared
assessments against criterion A3 when i) applied to the true (simulated) population size and
ii) to the sum of habitat suitability derived from SDMs as recommended by the Red List
guidelines[8(pp.101-111)]. Additionally, we compared these to the assessment under criterion E
based on the quantitative extinction probability extracted from our simulation. Since all
species went extinct during the 90 years of climate change in our simulations, all of them
qualified for all threatened categories throughout their life span. Again, we found the
strongest differences between range-contracting and range-shifting species, reflecting the

 differences in the population size - habitat loss relationship (Fig. 4 + Fig. S7).

Figure 4: Classification times for listing species in different IUCN Red List threatened categories, compared to their extinction times. Results are shown separately for range-contracting species with marginal, cold-adapted niche position (left) vs. range-shifting species with central niche positions (right). Classification time refers to the years when species would first be classified into each of the three different threatened categories (Vulnerable, Endangered, Critically endangered). The boxplots show the classification times across simulated species, landscapes, and replicate runs using three approaches: true population size extracted from the simulation or SDM-derived habitat loss against criterion A3, and extinction probability calculated from the simulations against criterion E. Note that all species went extinct during the 90 years of climate change, thus qualifying for all threatened categories throughout their life span. The grey horizontal line shows the mean year of the extinction across species and the grey bands the 95% confidence interval. Sampling size per box plot is n=240 (8 species x 3 replicate landscapes x 10 replicate runs).

For range-contracting species, the classification time did not differ between true population size and SDM-derived habitat loss (VU$_{pop}$: 0±0 years; VU$_{habitat}$: 0±0; EN$_{pop}$: 0±0; EN$_{habitat}$: 0±0; CR$_{pop}$: 1.40±1.84; CR$_{habitat}$: 1.74±0.44). By contrast, for range-shifting species SDM-based risk classification was substantially later than for true population size because range loss was underestimated by the SDMs (VU$_{pop}$: 1.88±3.58 years; VU$_{habitat}$: 10.71±1.39; EN$_{pop}$: 6.67±4.52; EN$_{habitat}$: 12.16±0.81; CR$_{pop}$: 14.2±5.34; CR$_{habitat}$: 25.07±1.21). This resulted in a very short warning time for range-shifting species when using SDM-based risk classification (Fig. 4 + Fig. S7). For range-shifting species that showed a full range shift (i.e., reaching the suitable habitat in the northern part of the landscape) due to high dispersal ability and fast growth, classification time between true population size and SDM-derived habitat loss did

also not differ (Fig. S8). In a sensitivity analysis, we tested whether SDM-based predictions

of range-shifting species could be improved by including simple dispersal assumptions but

found no effect (Fig. S9). Other traits did not result in pronounced differences in classification

times (Fig. S7). Interestingly, we found that for higher threatened categories (EN, CR)

quantitative estimates of extinction probability and listing under criterion E resulted in

considerably later classification time, and thus very short warning time, than information on

population size and listing under criterion A3 (Fig. 4 + Fig. S7). The Red List guidelines

indicate that criteria A-D are more precautionary than criterion E[8(p.17)]. Still, our results show

that listing under these criteria can differ by up to 20 generations for our virtual species (Fig.

4; species were simulated with one generation per year).

Discussion

Based on idealised simulated species and unbiased SDMs, we systematically evaluated the

guidelines of the Red List for assessing species-level extinction risk under climate change.

Our results revealed that the effectiveness of current Red List criteria strongly depend on

expected range dynamics. SDM-based assessment provided sufficient warning time only for

range-contracting species. For range-shifting species with time-lags in colonisation of newly

suitable habitat, warning time was substantially shorter mainly because of the highly non-

linear relationship between population size and SDM-predicted habitat loss. Other life-history

traits had comparatively little effect on warning times. Further, model-based assessment

based on probabilistic extinction estimates led to considerably shorter warning times than

when based on predicted population sizes.

The strong effect of range dynamics stems from fundamental differences in how population

size relates to predicted habitat loss. For range-contracting species, we found a convex

relationship between population loss observed in the simulations and SDM-derived

predictions of habitat loss. In these cases, species initially retained stable population sizes

despite ongoing habitat loss while population sizes dropped very suddenly for higher rates of

habitat loss. Such a convex relationship can be caused by a non-uniform distribution of

individuals in the landscape, where sparsely populated edge habitats are lost first during

range-contraction[21,34], and by an extinction threshold, a critical amount of habitat beyond

which population size declines steeply[23]. Crossing the threshold can initiate the extinction

vortex, in which reinforcing processes such as inbreeding depression increase population

decline, ultimately leading to species extinction[35]. Even though we did not consider species

genetics or stage-structured populations in our simulation, we still observed an extinction

threshold, which has been shown to be highly case specific and species dependent[23]. A

convex shape of population decline over time was already reported for several species[36], but

research on the exact relationship between population size, habitat loss and extinction risk

remains limited[20]. While the linear assumption in Red List guidelines is violated in these

cases, our results suggest that SDM-based assessment still offers sufficient warning time for

range-contracting species.

In contrast, range-shifting species exhibited a concave population size - habitat loss

relationship indicating a considerable underestimation of range loss by SDM predictions,

although the slope of the relationship differed between replicate landscapes. The concave

curve likely reflects the inability of some species to fully track shifting climates due to

dispersal limitations[37,38] and time lags[39,40]. Certain landscape characteristics such as

fragmentation, patch size and connectivity can enhance the dispersal ability of the species

causing them to better track their suitable habitat and thereby influence the form of the

curve[32]. The underestimation of range loss resulted in generally shorter warning time for

SDM-based assessment of species in comparison to true population size. Because effective

conservation measures often require several decades of advance warning, SDM-based

assessment may not leave enough warning time to protect range-shifting species from

extinction[29]. In contrast, SDM-based assessments did not lead to shorter warning time for

fully range-shifting species with high dispersal ability and fast growth, highlighting the

relevance of explicitly considering dispersal and demographic processes in climate impact

assessments[17]. Stronger dispersal ability is expected to improve climate tracking and the

survival of range-shifting species under climate change[32,41]. Interestingly, we only found such

influence of dispersal ability on the classification time for one range-shifting species and

found delayed warning for all other range-shifting species. As inclusion of simple dispersal

assumptions did not change these results our study suggests that SDMs will not be

adequate for assessing extinction risk in range-shifting species threatened with climate

change. For these species, spatially explicit population models are likely more adequate.

Our results further revealed that Red Listing based on probabilistic extinction estimates

under criterion E led to delayed classifications compared to criterion A3 and insufficient

warning times for all species, particularly those most at risk. This aligns with previous

findings that criterion E often results in less precautionary extinction risk status in

comparison to other criteria[42]. In our study, the short warning time resulted from steep,

sigmoidal extinction risk curves, which led to rapid population collapse and abrupt extinction

of species. Such curve shapes have been shown to realistically describe species' survival

probabilities under changing environmental conditions[43,44]. In stochastic population models,

sigmoidal extinction curves typically emerge when full parameter knowledge during the

model building process is available[45]. With higher parameter uncertainty the extinction risk

curves are expected to become more shallow[45], which consequently influences the

classification time in the Red List. Interestingly, a previous study[29] reported that criterion A

leaves less warning time in comparison to listing against criterion E when using a coupled

SDM-population model approach with a real species. It is thus unclear whether a higher

parameter uncertainty in the model building process for criterion E might leave more warning

time or may introduce additional biases. As a more robust approach for species threatened

by climate change, especially those undergoing range shifts, we recommend estimating

expected minimum abundance (EMA) from SEPMs. Our results suggest that assessing

extinction risk based on these projections provides a more reliable basis for classification

than relying on quantitative extinction probabilities alone.

Our simulations were idealised, with full knowledge of the true population and range

dynamics, perfect (i.e., error-free and unbiased) occurrence data to fit SDMs, and perfect

knowledge and information on the environmental drivers, which allowed the estimation of

unbiased SDMs. In practice, missing data, sampling biases, and missing knowledge about

species ecology and dispersal as well as environmental and demographic stochasticity could

introduce additional uncertainties in the models and thus in the assessment process[46–48].

Here, we want to echo the Red List guidelines in stressing transparency regarding

uncertainties in the assessment process to minimize their effect[8(pp.23-25)]. Additionally, our

climate change scenario was simplistic with only temperature changing over time. Simulating

more driver variables and more realistic climate change scenarios, which include extreme

events and more realistic fluctuations in temperature and precipitation, could lead to more

diverse responses of species. Furthermore, more detailed species characteristics, such as

stage-structured populations, stochastic movement dispersal and density-dependent traits

could also influence the responses of the species and thereby the assessment process.

Nevertheless, our results revealed that even under idealised conditions, current Red List

criteria are highly sensitive to species range dynamics under climate change.

To improve assessment, we recommend including a screening step into model-based Red

List assessments. In this initial screening, SDMs can serve to assess whether a species is

likely to experience exclusive range contraction or whether suitable habitat is likely to shift

geographically. If range-shifting is likely, we strongly recommend using spatially explicit

population models and assess extinction risk based on expected minimum abundance,

rather than extinction probabilities. While adjusting thresholds for extinction probabilities

under criterion E or the time frames in general may improve assessments in some cases,

further research is needed to define thresholds and time frames that offer both realism and

sufficient warning time. For range-contracting species, SDMs may be appropriate to assess

the extinction risk of species. Still, the assumption of a linear relationship between population

size and habitat size should be treated with caution as our results show that sudden

population reductions can cause delayed warning by SDMs. Furthermore, the known

limitations of SDMs should be kept in mind[18] and the assessment based on SDMs should

follow best practices[49,50] to avoid wrong classifications. Here, we want to stress again that

our SDMs represented the ideal case with perfectly known and unbiased data while in reality

species ecology, dispersal and observer behaviour could bias SDMs. To further guide the

novelization of the Red List under climate change, we recommend empirically testing our

results for a variety of species for which adequate range loss and population loss data are

available.

Methods

We used the RangeShifter framework[33,51,52] to simulate population dynamics of sixteen virtual

species under equilibrium conditions and under climate change in three different artificial

landscapes. Species differed in four traits: niche position (central vs. marginal), niche width

(narrow vs. wide), growth rate (slow vs. fast) and dispersal (short vs. long), leading to 16 (2^4)

different species to cover all trait combinations. Niches were defined by two environmental

variables that could be interpreted as temperature and precipitation. We aimed to simulate a

representative set of virtual species spanning a wide range of different traits and

characteristics as well as extinction dynamics of these species to assess their effect on the

classification in the IUCN Red List. Species occurrence data and climate data were

extracted from the simulation at equilibrium (after 100 years of spinup) before climate

change set in. Based on these data we fitted species distribution models (SDMs) and

predicted habitat suitability under climate change. From the simulations, we observed

population size across the entire time frame of 90 years of climate change and estimated

extinction probability per year. We ran simulations for 90 years of climate change to ensure

the extinction of all species. From the SDMs, we derived the predicted habitat suitability

values for each of these 90 years and compared these to true population size. Last, we

applied the Red List criteria to each of these three metrics to classify species into different

threatened categories.

Virtual species

In our simulation approach, we first modelled the population dynamics of the 16 virtual

species in the three artificial landscapes under climate change. The three artificial

landscapes are stochastic replicates obtained using a neutral landscape model[53]. The

artificial landscape consisted of two environmental variables (which could be interpreted as

temperature and precipitation) with a spatial extent of 512 x 512 cells at 1000 m resolution.

The environmental layers were generated using the package "NLMR"[54]. Temperature was

represented by a latitudinal gradient with spatially auto-correlated noise. For this, the

arithmetic mean of a landscape with a linear gradient and a landscape built with midpoint

displacement with a Hurst exponent of 0.96 was taken. To create spatially auto-correlated

precipitation patterns, we generated a landscape using midpoint displacement with a Hurst

exponent of 0.75. Landscape values ranged between 0 and 1. We derived three replicate

landscapes using the same settings (Fig. S10).

The virtual species niche was modelled using the package "virtualspecies"[55]. First,

Gaussian-shaped response curves for both environmental variables were simulated. The

mean of the response curve for temperature was adapted to the respective niche position

(marginal: 0.27, central: 0.5), the mean of the precipitation response curve was set to 0.5. To

model different niche breadths, the standard deviation of both response curves was adapted

based on the respective niche breadth (narrow: 0.045, wide: 0.055). The response curves of

the simulated niches are included in the appendix (Fig. S11). We then used the product of

both response curves to map the virtual species niche in the artificial landscapes. The

different niche positions (i.e., marginal and central) were explicitly selected to model different

range dynamics to test their influence on the extinction of the species and thus their

classification in the IUCN Red List. A marginal, cold-adapted niche position caused range

contraction under climate change and extinction, while a central niche position led to range-

shifting prior to range contraction and extinction. We decided to only model cold-adapted

marginal niches in the north, as a warm-adapted marginal niche position in the south would

have resulted in the same range-shifting pattern as the central niche position but with a

longer time period for the range-shift.

To simulate climate change, temperature was linearly increased over 90 years (Fig. S12),

while precipitation remained static. We selected a time frame of 90 years to model a

continuous increase of temperature that would result in no suitable habitat being left at the

end of climate change, ensuring the extinction of all species. To the linear temperature

increase we also included a temporal autocorrelated noise. For this we used an ARIMA

model[56] of the order 0,0,1 and added it to the linear model using the following equations:

$\eta_t = \theta * \varepsilon_t{-}1 + \varepsilon_t$

$y_t = \alpha + \beta * t + \eta_t$

Where, η_t represents the time series obtained from the ARIMA model with θ set to 0.5 and ε

representing the residuals at the respective time points. This time series was then added to

a linear model with α set to 0.5, β set to 0.6, and t representing the respective year. The time

series was rescaled between 0 and 0.9 and added to the temperature value of every cell in
the artificial landscape. Values that were larger than 1 were reduced to 1 to stay within the
range of 0 to 1. We did not aim to mimic realistic climate change projections but modeled
climate change in a way to force the species to extinction within a reasonable time frame.
Precipitation remained static under climate change and extreme events were not considered
to reduce complexity.
412
To model the population dynamics of the virtual species, we used the spatially explicit and
individual-based modelling platform RangeShifter[33,51], operated via the package
"RangeShiftR"[52]. We implemented a cell-based model using the previously created maps of
the virtual species niche with the resolution set to 1000 m and the carrying capacity set to
0.05 ind/ha. We modelled the virtual species as an asexual species with non-overlapping
generations meaning that each individual only lived for one year. The competition coefficient
was set to 1 (i.e., no over- or undercompensation) and the value for the maximum growth
rate was adapted according to the growth rate (slow: 3, fast: 5). After reproduction the adults
died and the offspring dispersed. Dispersal is modelled explicitly in three phases in
RangeShifter: emigration, transfer, and settlement. The emigration probability for each
individual was set to 0.4 and was density-independent. The transfer was modelled using a
negative exponential dispersal kernel. For short-distance dispersal, the mean dispersal
distance was set to 5000 m. For long-distance dispersal, a double exponential dispersal
kernel was used to include rare long-distance dispersal events. The double kernel is
composed of two negative exponential kernels with different means and probabilities of
occurrence. The first (shorter) kernel occurred with a probability of 95% and had a mean
dispersal distance of 15 000 m, the second (longer) kernel occurred with a probability of 5%
and had a mean dispersal distance of 250 000 m. If the arrival point of an individual went
beyond the boundaries of the landscape or within the natal cell the dispersal process was
repeated. The individual settled in the arrival cell if the cell contained suitable habitat. If the
arrival cell was not suitable, the individual was displaced to one of the eight neighbouring
cells, if one of them was suitable, or else died. If a cell became unsuitable due to climate
change, all individuals inhabiting that cell died immediately.
Trait values were selected based on two criteria: all species had to achieve a stable
population size after the spin-up period of simulation, while the simulation still had to be
computationally feasible and the differences between the trait values had to be large enough
to show different effects. We used realistic species (i.e., small rodents and birds) as
guidance for the determination of the growth rate and dispersal distances.
441
The simulation was initialised using the map with equilibrium climatic conditions and

individuals were distributed in all suitable cells at half-carrying capacity. The simulation ran

for 100 spin-up years to remove any initialisation effect. After the spin-up period, the

landscape was changed yearly to represent the virtual species niche shifting northwards in

response to climate change. In the following, we refer to year 0 as the last year of

equilibrium (after spinup) and year 1-90 represent the years under climate change. Each of

the 16 species was simulated with 100 replicate runs for each of the three replicate artificial

landscapes.

From the output of each simulation, we obtained the population size of the species for each

year and replicate run. To calculate the extinction probability for each year, we divided the

number of replicate runs without a viable population (i.e., no individual was left in the

landscape) of that year by the total number of replicate runs[57].

Species distribution models

At the end of the spin-up period, before climate change set in, we extracted species

presence and absence information (Fig. S2) as well as climate variables from the simulation

to fit SDMs. To account for variation between the replicate runs, we randomly selected ten

replicate runs from the simulation and fitted separate SDMs to each. This resulted in 480

SDMs (16 species x 3 landscape replicates x 10 replicate runs). For the species data we

assumed perfect sampling with no error in detection. Data points were partially thinned by

selecting every second cell to avoid spatial autocorrelation using the package "dismo"[58].

Based on the recommendations of the Red List[8(p.108)] we used three distinct modelling

algorithms (Generalised Linear Model (GLM), Random Forest and MaxEnt) and further

calculated an ensemble model based on the arithmetic mean. For the GLM, linear and

quadratic predictors were used with a binomial error distribution. For Random Forest, we

fitted 1000 trees with a minimum terminal node size of 20 to fit comparably smooth response

surfaces. In the MaxEnt algorithm, we used the feature classes linear, quadratic and hinge.

The probability of occurrence was predicted for each of the 90 years under climate change

using the ensemble model. The predictive performance of the model algorithms and the

ensemble model was evaluated using the 99 replicate runs that were not used for fitting the

SDM. As performance measures we selected the AUC (area under the ROC curve) and TSS

(true skill statistics)[16]. The calculated performance measures were averaged over all 99

replicate runs using the arithmetic mean. For the calibration and evaluation of the SDMs we

used the following packages: "randomForest"[59], "maxnet"[60], "gbm"[61], "PresenceAbsence"[62]

and "lhs"[63].

For testing the effect of dispersal assumptions in SDMs on the assessment in the Red List,

we first binarized the predictions of each species and landscape replicate using the maxTSS

threshold[64]. Around the known presences we created a buffer with the size of the estimated

dispersal distance of an individual in the respective time frame and set the predicted habitat

suitability outside the buffer to 0 following the recommendation of the Red List

guidelines[8(p.110)]. Buffer sizes were determined by the empirical dispersal distances observed

from the RangeShifter simulations. For this, we extracted the dispersal distances of all

emigrated individuals for year 0 from the simulations, and calculated the arithmetic mean of

dispersal distance.

Red List classification

To assess Red Listing of species threatened by climate, we compared the assessment of

the species against two different Red List criteria using three metrics. For the criterion A3,

we used true population size observed from the RangeShifter simulations and the sum of the

SDM-derived habitat suitability values. For the criterion E, we used the extinction probability

for every species obtained from the RangeShifter simulations. For the population size, we

only used the population size from the ten replicate runs that were selected for fitting the

SDMs. To calculate the sum of the habitat suitability values from the SDM predictions we

followed the recommendations from the Red List guidelines[8(p.99)]. We first removed cells with

values below a certain threshold value, for which we used the maxTSS[64]. We then summed

up the predicted habitat suitability of the remaining cells and for every year. For the

classification into the different categories of the Red List, different thresholds and time

frames apply for the different criteria (see Tab. 1).

Table 1: Thresholds and time frames used to classify species into threatened categories, based on

the Red List Criteria[8]

Criterion	Red List Threatened Categories		
	Critically Endangered	Endangered	Vulnerable
A3: future population size reduction / future reduction of sum of SDM-derived habitat suitability	>= 80% in 10 years	>= 50% in 10 years	>= 30% in 10 years
E: probability of extinction	>= 50% in 10 years	>= 20% in 20 years	>= 10% in 100 years

For each species, we determined the time of first classification into the three different

threatened categories using all three metrics (i.e., population size, habitat suitability and
extinction probability). As the Red List criteria are tied to specific time frames, we to tested
for every consecutive year if the species would fulfill the Red List criteria in the following
years (see Table 1 for considered time spans). If the criterion was fulfilled, then this
particular year was noted as the time of classification for that criterion. If the criterion was not
fulfilled, we repeated the analyses for the next consecutive year until we found the year in
which the species would first be classified according to the criterion under question. As all
species continuously experienced population and habitat loss under climate change until
their extinction, each species qualified for all threatened categories throughout their life span
(within the 90 years of simulated climate change).
In a sensitivity analysis, the assessment of classification times based on SDM-derived
habitat suitability values was repeated for more realistic dispersal assumptions using the
buffer approach described earlier.
520
Statistical analysis
To quantify the relative importance of the different traits for explaining differences in the
population size - habitat loss relationship, we fitted a generalised mixed-model ANCOVA
using an ordered beta regression implemented by the package "ordbetareg"[65,66]. This
package is a front-end to the package "brms"[67]. We selected an ordered beta regression
since our response variable (simulated "true" populated size) was continuous with a zero
and one inflation. As predictors, we considered habitat loss as continuous variable, the four
traits (niche position, niche width, dispersal distance and growth rate) as categorical
variables with two levels each, and their pairwise interactions with habitat loss. We further
controlled for the landscapes by including them as random intercepts. The best model was
selected based on the difference in expected log predictive density. Posterior distributions
were estimated using Markov chain Monte Carlo (MCMC) sampling with 4 chains and a delta
value of 0.95. Each chain had a length of 6000 iterations, with the first 2000 being used as a
warm-up. For the prior, the default was selected.
535
536   For the study R 4.2.2[68] was used. In addition to the already mentioned packages the
537   following were used for data preparation and visualization: "scales"[69], "dplyr"[70], "ggplot2"[71],
538   "terra"[72], "gridExtra"[73], "tibble"[74], "RColorBrewer"[75], "doParallel"[76], "foreach"[77], "data.table"[78],
539   "ggtext"[79].
540
541
542
543

Code availability:

The codes for this study are available at https://doi.org/10.5281/zenodo.18220071

Author Contributions:

RK and DZ conceptualised the study. RK ran all analyses and led the writing. All authors

contributed to revising the text.

Competing Interest Statement:

The authors declare that they have no conflict of interest to disclose.

Acknowledgments:

This study was supported by the German Research Foundation DFG (grant no. 518316503).

S.A.F. gratefully acknowledges the support of iDiv funded by the German Research

Foundation (DFG– _FZT 118, 202548816) and of the Leibniz Competition by the Leibniz

Association (P52/2017). We thank Cara Gallagher for helpful discussions during early

phases of the work.

References:

1. Scheffers, B. R. *et al.* The broad footprint of climate change from genes to biomes to people. *Science* **354**, aaf7671 (2016).

2. Freeman, B. G., Scholer, M. N., Ruiz-Gutierrez, V. & Fitzpatrick, J. W. Climate change causes upslope shifts and mountaintop extirpations in a tropical bird community. *Proceedings of the National Academy of Sciences* **115**, 11982–11987 (2018).

3. Urban, M. C. Climate change extinctions. *Science* **386**, 1123–1128 (2024).

4. Bellard, C., Bertelsmeier, C., Leadley, P., Thuiller, W. & Courchamp, F. Impacts of climate change on the future of biodiversity. *Ecology Letters* **15**, 365–377 (2012).

5. Pecl, G. T. *et al.* Biodiversity redistribution under climate change: Impacts on ecosystems and human well-being. *Science* **355**, eaai9214 (2017).

6. Stanton, J. C., Shoemaker, K. T., Pearson, R. G. & Akçakaya, H. R. Warning times for species extinctions due to climate change. *Global Change Biology* **21**, 1066–1077 (2015).

7. IUCN. *IUCN Red List Categories and Criteria: Version 3.1.* (2001).

8. IUCN Standards and Petitions Committee. *Guidelines for Using the IUCN Red List Categories and Criteria. Version 16*. (2024).

9. Mace, G. M. & Lande, R. Assessing Extinction Threats: Toward a Reevaluation of IUCN Threatened Species Categories. *Conservation Biology* **5**, 148–157 (1991).

10. Mace, G. M. *et al.* Quantification of extinction risk: IUCN's System for Classifying Threatened Species. *Conservation Biology: The Journal of the Society for Conservation Biology* **22**, 1424–1442 (2008).

11. Akçakaya, H. R., Butchart, S. H. M., Mace, G. M., Stuart, S. N. & Hilton-Taylor, C. Use and misuse of the IUCN Red List Criteria in projecting climate change impacts on biodiversity. *Global Change Biology* **12**, 2037–2043 (2006).

12. IUCN Standards and Petitions Committee. *Guidelines for Using the IUCN Red List Categories and Criteria. Version 8.0*. (2010).

13. Valavi, R., Guillera-Arroita, G., Lahoz-Monfort, J. J. & Elith, J. Predictive performance of presence-only species distribution models: a benchmark study with reproducible code. *Ecological Monographs* **92**, e01486 (2022).

14. Urban, M. C. Accelerating extinction risk from climate change. *Science* **348**, 571–573 (2015).

15. Wüest, R. O. *et al.* Macroecology in the age of Big Data – Where to go from here? *Journal of Biogeography* **47**, 1–12 (2020).

16. Guisan, A., Thuiller, W. & Zimmermann, N. E. *Habitat Suitability and Distribution Models: With Applications in R*. (Cambridge University Press, Cambridge, 2017).

17. Araújo, M. B. & Guisan, A. Five (or so) challenges for species distribution modelling. *Journal of Biogeography* **33**, 1677–1688 (2006).

18. Elith, J. & Leathwick, J. R. Species Distribution Models: Ecological Explanation and Prediction Across Space and Time. *Annual Review of Ecology, Evolution, and Systematics* **40**, 677–697 (2009).

19. Weber, M. M., Stevens, R. D., Diniz-Filho, J. A. F. & Grelle, C. E. V. Is there a correlation between abundance and environmental suitability derived from ecological niche modelling? A meta-analysis. *Ecography* **40**, 817–828 (2017).

20. Zurell, D., Fritz, S. A., Rönnfeldt, A. & Steinbauer, M. J. Predicting extinctions with species distribution models. *Cambridge Prisms: Extinction* **1**, e8 (2023).

21. Rodríguez, J. P. Range Contraction in Declining North American Bird Populations. *Ecological Applications* **12**, 238–248 (2002).

22. Kuussaari, M. *et al.* Extinction debt: a challenge for biodiversity conservation. *Trends Ecol Evol* **24**, 564–571 (2009).

23. Fahrig, L. How much habitat is enough? *Biological Conservation* **100**, 65–74 (2001).

24. Keith, D. A. *et al.* Predicting extinction risks under climate change: coupling stochastic population models with dynamic bioclimatic habitat models. *Biology Letters* **4**, 560–563 (2008).

25. Maschinski, J., Baggs, J. E., Quintana-Ascencio, P. F. & Menges, E. S. Using Population Viability Analysis to Predict the Effects of Climate Change on the

Extinction Risk of an Endangered Limestone Endemic Shrub, Arizona Cliffrose. *Conservation Biology* **20**, 218–228 (2006).

26. Akcakaya, H. R. & Sjögren-Gulve, P. Population Viability Analyses in Conservation Planning: An Overview. *USA. — Per Sjögren-Gulve* **11733**, (2000).

27. Briscoe, N. J. *et al.* Forecasting species range dynamics with process-explicit models: matching methods to applications. *Ecology Letters* **22**, 1940–1956 (2019).

28. Cazalis, V. *et al.* Bridging the research-implementation gap in IUCN Red List assessments. *Trends in Ecology & Evolution* **37**, 359–370 (2022).

29. Keith, D. A. *et al.* Detecting Extinction Risk from Climate Change by IUCN Red List Criteria. *Conservation Biology* **28**, (2014).

30. Zurell, D. *et al.* The virtual ecologist approach: simulating data and observers. *Oikos* **119**, 622–635 (2010).

31. Zurell, D. *et al.* Benchmarking novel approaches for modelling species range dynamics. *Global Change Biology* **22**, 2651–2664 (2016).

32. Foden, W. B. *et al.* Identifying the World's Most Climate Change Vulnerable Species: A Systematic Trait-Based Assessment of all Birds, Amphibians and Corals. *PLOS ONE* **8**, e65427 (2013).

33. Bocedi, G. *et al.* RangeShifter 2.0: an extended and enhanced platform for modelling spatial eco-evolutionary dynamics and species' responses to environmental changes. *Ecography* **44**, 1453–1462 (2021).

34. Sagarin, R. D. & Gaines, S. D. The 'abundant centre' distribution: to what extent is it a biogeographical rule? *Ecology Letters* **5**, 137–147 (2002).

35. Gilpin, M. Minimum viable populations : Processes of species extinction. in (1986).

36. Fagan, W. F. & Holmes, E. E. Quantifying the extinction vortex. *Ecology Letters* **9**, 51–60 (2006).

37. Travis, J. M. J. Climate Change and Habitat Destruction: A Deadly Anthropogenic Cocktail. *Proceedings: Biological Sciences* **270**, 467–473 (2003).

38. Hof, C., Levinsky, I., Araújo, M. B. & Rahbek, C. Rethinking species' ability to cope with rapid climate change. *Global Change Biology* **17**, 2987–2990 (2011).

39. Alexander, J. M. *et al.* Lags in the response of mountain plant communities to climate change. *Global Change Biology* **24**, 563–579 (2018).

40. Chen, I.-C., Hill, J. K., Ohlemüller, R., Roy, D. B. & Thomas, C. D. Rapid Range Shifts of Species Associated with High Levels of Climate Warming. *Science* **333**, 1024–1026 (2011).

41. Mancini, G. *et al.* A standard approach for including climate change responses in IUCN Red List assessments. *Conservation Biology* **38**, e14227 (2024).

42. Gärdenfors, U. Population Viability Analysis in the Classification of Threatened Species: Problems and Potentials. *Ecological Bulletins* 181–190 (2000).

43. Ball, T. *et al.* A general relationship between carrying capacity and extinction risk. Preprint at https://doi.org/10.33774/coe-2024-m9dqp-v3 (2025).

44. Dushoff, J. Carrying Capacity and Demographic Stochasticity: Scaling Behavior of the Stochastic Logistic Model. *Theoretical Population Biology* **57**, 59–65 (2000).

45. Taylor, B. L. The Reliability of Using Population Viability Analysis for Risk Classification of Species. *Conservation Biology* **9**, 551–558 (1995).

46. Akcakaya, H. R. Population viability analyses with demographically structured models. *Ecological Bulletins* **48**, 23–38 (2000).

47. Possingham, H. P., McCarthy, M. A. & Lindenmayer, D. B. Population Viability Analysis. in *Encyclopedia of Biodiversity (Second Edition)* (ed. Levin, S. A.) 210–219 (Academic Press, Waltham, 2013). doi:10.1016/B978-0-12-384719-5.00173-8.

48. Walther, G.-R. *et al.* Ecological responses to recent climate change. *Nature* **416**, 389–395 (2002).

49. Araújo, M. B. *et al.* Standards for distribution models in biodiversity assessments. *Sci Adv* **5**, eaat4858 (2019).

50. Sofaer, H. R. *et al.* Development and Delivery of Species Distribution Models to Inform Decision-Making. *BioScience* **69**, 544–557 (2019).

51. Bocedi, G. *et al.* RangeShifter: a platform for modelling spatial eco-evolutionary dynamics and species' responses to environmental changes. *Methods in Ecology and Evolution* **5**, 388–396 (2014).

52. Malchow, A.-K., Bocedi, G., Palmer, S. C. F., Travis, J. M. J. & Zurell, D. RangeShiftR: an R package for individual-based simulation of spatial eco-evolutionary dynamics and species' responses to environmental change. *Ecography* (2021) doi:10.1111/ecog.05689.

53. With, K. A. & King, A. W. The Use and Misuse of Neutral Landscape Models in Ecology. *Oikos* **79**, 219–229 (1997).

54. Sciaini, M., Fritsch, M., Scherer, C. & Simpkins, C. E. NLMR and landscapetools: An integrated environment for simulating and modifying neutral landscape models in R. *Methods in Ecololgy and Evolution* **00**, 1–9 (2018).

55. Leroy, B., Meynard, C. N., Bellard, C. & Courchamp, F. virtualspecies, an R package to generate virtual species distributions. *Ecography* (2015) doi:10.1111/ecog.01388.

56. Box, G. E. P. & Jenkins, G. M. *Time Series Analysis: Forecasting and Control*. (San Francisco: Holden-Day, 1970).

57. Grimm, V. & Wissel, C. The intrinsic mean time to extinction: a unifying approach to analysing persistence and viability of populations. *Oikos* **105**, 501–511 (2004).

58. Hijmans, R. J., Phillips, S., Leathwick, J. & Elith, J. *Dismo: Species Distribution Modeling*. (2021).

59. Liaw, A. & Wiener, M. Classification and Regression by randomForest. *R News* **2**, 18–22 (2002).

60. Phillips, S. *Maxnet: Fitting 'Maxent' Species Distribution Models with 'Glmnet'*. (2021).

61. Greenwell, B., Boehmke, B., Cunningham, J. & Developers, G. B. M. *Gbm: Generalized Boosted Regression Models*. (2020).

62. Freeman, E. A. & Moisen, G. PresenceAbsence: An R Package for Presence Absence Analysis. *Journal of Statistical Software* **23**, 1–31 (2008).

63. Carnell, R. *Lhs: Latin Hypercube Samples*. (2024).

64. Liu, C., Berry, P. M., Dawson, T. P. & Pearson, R. G. Selecting thresholds of occurrence in the prediction of species distributions. *Ecography* **28**, 385–393 (2005).

65. Kubinec, R. Ordered Beta Regression: A Parsimonious, Well-Fitting Model for Continuous Data with Lower and Upper Bounds. *Political Analysis* **31**, 519–536 (2023).

66. Kubinec, R. *Ordbetareg: Ordered Beta Regression Models with 'Brms'*. (2025).

67. Bürkner, P.-C. brms: An R Package for Bayesian Multilevel Models Using Stan. *Journal of Statistical Software* **80**, 1–28 (2017).

68. R Core Team. *R: A Language and Environment for Statistical Computing*. (R Foundation for Statistical Computing, Vienna, Austria, 2022).

69. Wickham, H. & Seidel, D. *Scales: Scale Functions for Visualization*. (2022).

70. Wickham, H., François, R., Henry, L., Müller, K. & Vaughan, D. *Dplyr: A Grammar of Data Manipulation*. (2023).

71. Wickham, H. *Ggplot2: Elegant Graphics for Data Analysis*. (Springer-Verlag New York, 2016).

72. Hijmans, R. J. *Terra: Spatial Data Analysis*. (2023).

73. Auguie, B. *gridExtra: Miscellaneous Functions for 'Grid' Graphics*. (2017).

74. Müller, K. & Wickham, H. *Tibble: Simple Data Frames*. (2022).

75. Neuwirth, E. *RColorBrewer: ColorBrewer Palettes*. (2014).

76. Weston, S. & Microsoft. *doParallel: Foreach Parallel Adaptor for the 'parallel' Package*. (2022).

77. Microsoft & Weston, S. *Foreach: Provides Foreach Looping Construct*. (2022).

78. Barrett, T. *et al. Data.Table: Extension of `data.Frame`*. (2024).

79. Wilke, C. O. & Wiernik, B. M. *Ggtext: Improved Text Rendering Support for 'Ggplot2'*. (2022).