Cryo-EM Structure of the Cyclase Domain and Evaluation of Substrate Channeling in a Bifunctional Class II Terpene Synthase

Matthew N. Gaynes,[a] Kollin Schultz,[b,c] Eliott S. Wenger,[a] Ronen Marmorstein,[b,d] and David W. Christianson[a,*]

[a]Roy and Diana Vagelos Laboratories, Department of Chemistry, University of Pennsylvania, Philadelphia, Pennsylvania, 19104-6323, USA

[b]Abramson Family Cancer Research Institute, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA

[c]Graduate Group in Biochemistry, Biophysics, and Chemical Biology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA

[d]Department of Biochemistry and Biophysics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA

*Corresponding author: David W. Christianson, email: chris@sas.upenn.edu, phone: (215) 898-5714

ABSTRACT

Copalyl diphosphate synthase from *Penicillium verruculosum* (PvCPS) is a bifunctional class II terpene synthase containing a prenyltransferase that produces geranylgeranyl diphosphate (GGPP) and a class II cyclase that utilizes GGPP as a substrate to generate the bicyclic diterpene copalyl diphosphate. The various stereoisomers of copalyl diphosphate establish the greater family of labdane natural products, many of which have environmental and medicinal impact. Understanding structure-function relationships in class II diterpene synthases is crucial for guiding protein engineering campaigns aimed at the generation of diverse bicyclic diterpene scaffolds. However, only a limited number of structures are available for class II cyclases from bacteria, plants, and humans, and no structures are available for a class II cyclase from a fungus. Further, bifunctional class II terpene synthases have not been investigated with regard to substrate channeling between the prenyltransferase and the cyclase. Here, we report the 2.9 Å-resolution cryo-EM structure of the 63-kD class II cyclase domain from PvCPS. Comparisons with bacterial and plant copalyl diphosphate synthases reveal conserved residues that likely guide the formation of the bicyclic labdane core, but divergent catalytic dyads that mediate the final deprotonation step of catalysis. Substrate competition experiments reveal preferential GGPP transit from the PvCPS prenyltransferase to the cyclase, even when prepared as separate constructs. These results are consistent with a model in which transient prenyltransferase-cyclase association facilitates substrate channeling due to active site proximity.

Introduction

Terpenes comprise a ubiquitous class of natural products with an extensive range of applications as flavors, fragrances, pharmaceuticals, insecticides, and next-generation biofuels.[1-6] Terpene biosynthesis is rooted in primary metabolism: a prenyltransferase catalyzes the condensation of two C_5 substrates, dimethylallyl diphosphate (DMAPP) and one or more equivalents of isopentenyl diphosphate (IPP), to produce C_{10} geranyl diphosphate (GPP), C_{15} farnesyl diphosphate (FPP), C_{20} geranylgeranyl diphosphate (GGPP), and even longer isoprenoid diphosphates.[7,8] These linear isoprenoids have diverse biological functions, e.g., in protein prenylation to direct cellular localization, or small-molecule prenylation to generate intermediates in alkaloid or polyketide biosynthesis.[9-12] However, the most intricate reactions of terpene biosynthesis are found in secondary metabolism, where isoprenoid cyclization reactions yield products containing multiple rings and stereocenters. These reactions are catalyzed by terpene cyclases, enzymes that direct astoundingly complex carbocation-mediated cyclization cascades.[13-15]

A terpene cyclase catalyzes the first committed step in the biosynthesis of a terpenoid natural product and usually belongs to one of two principal enzyme classes based on domain architecture and chemical strategy for initial carbocation formation. A class I cyclase contains an active site in the middle of an α helical bundle designated as the α domain, with two metal binding motifs at the mouth of the active site. The α domain fold of class I terpene cyclases is shared with prenyltransferases, and both utilize a trinuclear metal cluster[16] to initiate ionization of the substrate diphosphate group to generate an allylic carbocation. A class II cyclase adopts a different α helical fold, denoted as a β domain that is most often associated with a structurally homologous γ domain.[14,17] The γ domain is inserted between the first and second helices of the β domain and caps the active site in the β domain. The β domain contains an aspartic acid that protonates the terminal C=C bond (or epoxide) in the isoprenoid substrate to generate a tertiary

carbocation.[18] Of note, class II cyclization products retain the substrate diphosphate group, e.g., as observed for *ent*-copalyl diphosphate synthase.[19,20]

Prenyltransferases and terpene cyclases are found individually across all domains of life, but bifunctional terpene synthases containing both enzymes in a single polypeptide chain have been identified mainly in fungal species.[21] The first reported bifunctional terpene synthase, fusicoccadiene synthase from *Phomopsis amygdali* (PaFS), contains a prenyltransferase (GGPP synthase) and a class I diterpene cyclase.[22] The first bifunctional terpene synthases identified containing a prenyltransferase (GGPP synthase) and a class II cyclase were copalyl diphosphate synthases from *Penicillium verruculosum* (Figure 1) and *Penicillium fellutanum* (PvCPS and PfCPS, respectively).[23] In both class I and class II bifunctional terpene synthases, the prenyltransferase is connected to the cyclase through a flexible linker containing 40–125 residues, with domain architecture represented as α~α and α~βγ, respectively.

Figure 1. Reaction sequence catalyzed by PvCPS. Prenyltransferase substrates DMAPP and IPP are utilized by the prenyltransferase α domain to produce GGPP. The class II cyclase active site at the βγ domain interface then cyclizes GGPP to generate copalyl diphosphate. The primary structure of PvCPS illustrated at bottom shows that the γ domain is inserted between the first and second helices of the β domain. The γ domain is structurally homologous to the β domain and is thought to be the result of primordial gene duplication and fusion. A 125-residue flexible linker connects the prenyltransferase to the cyclase.

Bifunctional terpene synthases have been referred to as "assembly-line" synthases[24] because some (but not all) class I bifunctional synthases exhibit substrate channeling in the first

two reactions of terpene biosynthesis.[25-27] Cryo-EM studies of PaFS[28,29] and the related

bifunctional synthase variediene synthase from *Emericella variecolor* (EvVS)[27] reveal a central

prenyltransferase octamer or hexamer, respectively, surrounded by class I cyclase domains with

overall $(\alpha{\sim}\alpha)_8$ or $(\alpha{\sim}\alpha)_6$ architecture. In PaFS, cyclase domains are randomly splayed out

around the central prenyltransferase octamer but are capable of transient binding to the side of

the prenyltransferase oligomer, which is believed to facilitate intramolecular substrate

channeling.[26,28,29] Intramolecular substrate channeling is not observed in EvVS, and cyclase

domains are fixed in apical positions above and below the prenyltransferase hexamer.[27]

To date, substrate channeling has not been explored in any class II bifunctional terpene

synthase. Moreover, no high-resolution structure of a bifunctional prenyltransferase-class II

cyclase has been reported to date. Even so, the X-ray crystal structure of the prenyltransferase

hexamer of PvCPS and a molecular model of the class II cyclase domain were used to interpret

a low resolution molecular envelope of the full-length hexamer calculated from small-angle X-

ray scattering (SAXS) data, revealing a fascinating $(\alpha{\sim}\beta\gamma)_6$ "starburst"-like architecture.[30]

Interestingly, the cryo-EM structure of the prenyltransferase hexamer of PfCPS reveals a

partially-open hexamer conformation thought to represent a possible intermediate between

hexameric and octameric prenyltransferase assemblies.[31]

We previously reported the X-ray crystal structure of the prenyltransferase domain of

S723T-PvCPS, showing IPP bound to the prenyltransferase domain in a nonproductive

location.[32] We hypothesized that if the IPP binding site corresponded to a possible isoprenoid

diphosphate binding site as GGPP transits from the prenyltransferase domain to the cyclase

domain, then a productive prenyltransferase-cyclase complex might be stabilized for

visualization in cryo-EM studies of full-length S723T-PvCPS. However, while we do not observe

prenyltransferase-cyclase association, we do observe well-defined density for the class II

cyclase domain. Accordingly, we now present the 2.9 Å-resolution cryo-EM structure of the

cyclase domain in full-length S723T-PvCPS, confirming general features of the model of this domain previously fit to the molecular envelope derived from SAXS data.[30] Due to its relatively small size, the 63-kD βγ cyclase domain would ordinarily be considered a challenging target for cryo-EM structure determination. Even so, cryo-EM data were of sufficient quality to yield an atomic resolution reconstruction. Additionally, we report biochemical assays showing that GGPP channeling from the prenyltransferase to the cyclase is operative in the $(\alpha{\sim}\beta\gamma)_6$ hexamer of wild-type PvCPS.

Methods

Reagents

Unless overwise noted, chemicals and reagents used for protein expression and purification were purchased from Fisher Scientific, Millipore Sigma, or GoldBio and were used without further purification.

Protein Preparation and Purification

Wild-type PvCPS and S723T-PvCPS. Codon-optimized plasmids of wild-type PvCPS (Uniprot: A0A348FUE1) and S723T-PvCPS were purchased from Genscript. All PvCPS constructs utilize a pET28a(+) vector with an N-terminal His tag and a TEV cleavage site for purification. Detailed methods for the purification of PvCPS including wild-type and mutants have been previously reported and were followed without major modifications.[30] Briefly, BL21(DE3) competent *E. coli* cells (New England Biolabs) were transformed with PvCPS plasmids and grown on agar plates. Six 2-L flasks containing 1 L of terrific broth (TB) supplemented with 50 μg/mL kanamycin were inoculated using 15-mL aliquots of overnight starter cultures. Expansion cultures were shaken at 37°C and 250 RPM until the OD reached approximately 0.6, at which point isopropyl β-D-thiogalactopyranoside was added to a final concentration of 500 μM. After shaking at 16°C for 18 h, cells were pelleted by centrifugation at 7808g for 20 min. Cell pellets were collected and stored at -80°C until further use.

Upon thawing, the cell pellet was resuspended in 100 mL Buffer A [25 mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) (pH 7.5), 250 mM NaCl, 30 mM imidazole, 1 mM tris(2-carboxyethyl)phosphine (TCEP), and 10% glycerol] also containing 100 mg lysozyme, 2 EDTA-free cOmplete Mini Protease Inhibitor tablets, 250 units Benzonase, and approximately 17 mg of phenylmethylsulfonyl fluoride (PMSF; dissolved in 100% ethanol prior to addition to the lysate mixture). The lysate was stirred at room temperature for 45 min and at 4°C for 30 min followed by sonication (30 Hz, cycling between 1 s on/2 s off for 10 min). The lysate mixture was clarified by centrifugation at 41,646g for 30 min at 4°C. The supernatant was loaded onto a 5-mL HisTrap HP prepacked column (Cytiva) pre-equilibrated with Buffer A. Protein was eluted with an initial 10% step to Buffer B [Buffer A plus 500 mM imidazole], followed by a 50 mL gradient to 100% Buffer B. Fractions containing the largest quantities of PvCPS were pooled, filtered with a 0.22 µM PVDF syringe filter (Millipore), and loaded onto a 26/60 Superdex 200 size-exclusion column (Cytiva) pre-equilibrated with sizing buffer [25 mM HEPES (pH 7.5), 150 mM NaCl, 10% glycerol, 1 mM TCEP]. Fractions were analyzed using SDS-PAGE and pure fractions were pooled and supplemented with 100% glycerol to bring the final glycerol concentration to 30%. The sample was then concentrated using a 50 kD Amicon centrifugal filter unit to 10 mg/mL and stored at -80°C.

PvCPS$_{CY}$ and PvCPS$_{PT}$. Purification of wild-type PvCPS was achieved as described above with the addition of a 1-h incubation with Endoproteinase Glu-C (Hampton Research) at a 1:1000 mg ratio at room temperature. The mixture was then reloaded onto a 26/60 Superdex 200 size-exclusion column (Cytiva) pre-equilibrated with sizing buffer [25 mM HEPES (pH 7.5), 150 mM NaCl, 10% glycerol, 1 mM TCEP]. Two major peaks eluted, the first containing an approximately 37-kD protein and the second containing an approximately 63-kD protein based on SDS-PAGE analysis. Fractions associated with the first and second peak were pooled as

PvCPS$_{PT}$ and PvCPS$_{CY}$, respectively. PvCPS$_{PT}$ was concentrated to 10 mg/mL, PvCPS$_{CY}$ was concentrated to 20 mg/mL, and both were stored at -80°C.

PvCPS$_{SL}$. A construct of PvCPS was designed in which the 125-residue linker segment L567–T692 was replaced by the 20-residue segment AISDHTSRAIDLCRNPPLWG corresponding to an extension of helix A in the prenyltransferase domain of PfCPS. The codon-optimized plasmid was purchased from Genscript. The expression and purification of this variant containing a shortened linker, designated "PvCPS$_{SL}$", were carried out in the same manner as described above for full-length PvCPS.

CotB2. A pET28a(+) expression plasmid containing the gene encoding cyclooctatenol synthase (CotB2) was purchased from Genscript. Previously published protocols[33,34] were utilized to express and purify the single domain class I cyclase CotB2 as described without modification (Uniprot ID: C9K1X5).

PaFS$_{CY}$. The cyclase domain of PaFS, designated "PaFS$_{CY}$", consists of residues 1–344. The preparation of this construct was achieved using previously reported procedures without modification.[26]

Mass Photometry

Samples of wild-type PvCPS and PvCPS$_{SL}$ were initially diluted to approximately 50 nM in mass photometry (MP) buffer [25 mM HEPES (pH 7.5), 250 mM NaCl]. The mass photometer (Refeyn Ltd.) was calibrated using a standard curve based on measurements with β-amylase and thyroglobulin, which allowed for the conversion of ratiometric contrast to particle mass in kD. A typical measurement involved addition of 12 µL of mass photometry buffer to the glass slip receptacle which was first utilized to obtain optimal focus. Following this, 3 µL of the diluted protein sample was mixed with the drop to yield a final protein concentration of approximately

10 nM. Data were collected for 60 s (approximately 60,000 frames) and analyzed using RefeynMP software.

Cryo-EM Data Collection

S723T-PvCPS was utilized for cryo-EM structure determination. An 80-µL sample (3.0 mg/mL) was dialyzed overnight in cryo-EM buffer (25 mM HEPES pH 7.5, 50 mM NaCl, 1 mM TCEP) and then filtered using a 0.22 µM PVDF centrifugal filter (Millipore) to yield a concentration of approximately 2.0 mg/mL. A 10X ligand and detergent stock was made containing $MgCl_2$, IPP, and NP-40 detergent mixed in cryo-EM buffer and filtered with a 0.22 µM PVDF centrifugal filter (Millipore). Protein was diluted to 1.0 mg/mL (approximately 10 µM) and the substrate mix was added to final concentrations of 5 mM $MgCl_2$, 0.5 mM IPP, and 0.025% NP-40. Grid setup utilized R1.2/1.3 300-mesh copper grids (QUANTIFOIL™) which were glow discharged for 1 min (easiGlow, Pelco) prior to blotting. A Vitrobot Mark IV was utilized to blot 3 µL of protein sample with a blot force of 0 for 6 s prior to vitrification in ethane. Frozen grids were clipped and transferred to a Titan Krios G3i cryogenic transmission electron microscope (Thermo Fischer Scientific) operating at 300 keV at the Laboratory for Biomolecular Structure at Brookhaven National Laboratory (Upton, NY). Images were recorded with a K3 Summit detector at 81,000x magnification (0.53 Å/pixel) with a defocus of -0.8 to -2.5 µM. Data were collected utilizing EPU software with parameters of 40 frames taken at a dose of 15 e⁻/pixel/s (33 e⁻/$Å^2$). A total of 3,310 movies were recorded.

Cryo-EM Data Processing

Data processing was completed using cryoSPARC (version 4.6.2), summarized by the workflow shown in Figure S1. Movies were imported into cryoSPARC and aligned using patch motion correction with output Fourier cropping set to 1/2. Contrast transfer function (CTF) estimations were calculated with patch CTF estimation and resulting exposures were manually

curated resulting in the rejection of 87 micrographs. An initial selection of particles was obtained

by using the Blob Picker in cryoSPARC with a minimum and maximum particle diameter of 150

Å and 300 Å, respectively. The resulting particle picks were extracted and utilized to establish an

initial set of two-dimensional (2D) classes. The best classes were selected and included in a

template picker job in cryoSPARC set to identify particles with a diameter of 200 Å; following

inspection, this procedure yielded 1,686,323 particles. Extraction of particles with a large box

size (400 pixels), which would accommodate a hexameric particle, revealed low quality 2D

classes of almost exclusively βγ domain monomers. To generate higher quality 2D classes of βγ

domain monomers, particles were extracted with a smaller box size of 180 pixels Fourier-

cropped to 90 pixels and utilized to establish 200 2D classes. Optimal particles were selected

and utilized for *ab initio* three-dimensional (3D) reconstruction with two classes. The best

volume was lowpass filtered to 20 Å and utilized for several rounds of heterogeneous

refinement for both classes. The resulting highest resolution class was utilized for non-uniform

refinement. Despite promising 2D classes of the class II cyclase, the final 3D reconstruction in

cryoSPARC still suffered greatly from alignment issues, exemplifying the significant challenge of

working with a relatively small 63-kD particle. This particle size approached the lower limit of

current cryo-EM capabilities (Figure S1).

A new cryo-EM tool in RELION, Blush regularization, has recently been shown to yield

high-resolution 3D reconstructions of low molecular weight biomolecules using data with low

signal-to-noise ratio; for example, application of Blush regularization yielded a 2.5 Å-resolution

map of a 40-kD protein-RNA complex.[35] To obtain a high-resolution 3D reconstruction of the βγ

domains of S723T-PvCPS, the highest quality 2D classes were selected for a stack of

approximately 500,000 particles. CTF-corrected micrographs with these particle picks were

transferred from cryoSPARC to RELION, and particles were re-extracted with a box size of 180

pixels. 3D refinement was run with all 500,000 particles followed by 3D classification. Optimal

3D classes were used to select a subset of particles for a final stack of approximately 350,000 particles. This particle stack was subject to iterative rounds of 3D refinement, post-processing for B-factor sharpening, and CTF refinement to establish a set of polished particles. These particles yielded a final 3D reconstruction at 2.9 Å-resolution as indicated by the Gold Standard Fourier Shell Correlation plot (Figure S2).

A model of the S723T-PvCPS βγ cyclase domain was generated with AlphaFold3[36] and fit into the 2.9 Å-resolution map using ChimeraX.[37] The fit model and map were then transferred to PHENIX for real-space refinement using secondary structure restraints and simulated annealing to optimize the model fit to the experimental map.[38] Iterative cycles of model adjustment with Coot[39] and real-space refinement yielded the final model of the class II cyclase validated with the cryo-EM validation tool in PHENIX as well as with Molprobity.[40] Data collection and refinement statistics are recorded in Table S1.

Substrate Competition Experiments

Wild-type PvCPS and PvCPS$_{SL}$ were utilized for substrate competition experiments designed to probe GGPP channeling between prenyltransferase and cyclase domains. Assays were performed on a 200-µL scale in 2-mL glass screw-cap vials. Reactions were run in sizing buffer also containing 1 mM MgCl$_2$ in which the concentrations of PvCPS and diterpene cyclase competitor (CotB2 or PaFS$_{CY}$) were 5 µM (MgCl$_2$ concentrations higher than 1 mM were found to be significantly inhibitory for the generation of copalyl diphosphate/copalol). Experiments were also conducted with equimolar 5 µM mixtures of individual PvCPS prenyltransferase (PvCPS$_{PT}$), PvCPS cyclase (PvCPS$_{CY}$), and diterpene cyclase competitors mentioned above. A mixture of IPP and DMAPP was prepared by adding equal volumes of 10 mM DMAPP and 30 mM IPP. Reactions were initiated by the addition of 20 µL of either the DMAPP+IPP substrate mixture or a 5 mM solution of GGPP. All isoprenoids were purchased from isoprenoids.com and were resuspended in a 7:3 methanol:10 mM NH$_4$HCO$_3$ mixture at various concentrations.

Reactions were incubated at room temperature overnight. The following day, rCutSmart Buffer (NEB) was added to each reaction to a final concentration of 1X from a 10X stock, followed by the addition of 40 units of Quick CIP (NEB). Reaction mixtures were then incubated at 37°C overnight. The following day, reactions were quenched by addition of 200 µL ethyl acetate and then vortexed for 3 x 5-s intervals. The entire reaction volume was transferred to a 1.7-mL Eppendorf tube and spun at 2415g for 15 s. A portion of the organic phase was subsequently transferred to a screw cap vial with a 300-µL insert.

Gas chromatography-mass spectrometry (GC-MS) was employed for product identification and quantification using an Agilent 8890 GC / 5597C MSD system equipped with a J&W HP-5 ms Ultra Inert GC capillary column (30 m x 0.25 mm x 0.25 µm). While GC-MS can be utilized to detect and quantify cyclooctatenol, GC-MS cannot be utilized to detect and quantify copalyl diphosphate due to the presence of the diphosphate group. This necessitated a dephosphorylation step to generate copalol, which can then be detected and quantified using GC-MS. Accordingly, dephosphorylation of copalyl diphosphate in product mixtures was achieved using calf intestinal phosphatase (CIP) as previously described by Ma and colleagues.[41] To ensure complete dephosphorylation of copalyl diphosphate, CIP reactions were carried out for increasing time intervals until the copalol signal plateaued in control experiments. The optimal dephosphorylation time was determined to be 18 h.

Copalol-cyclooctatenol and copalol-fusicoccadiene product mixtures were analyzed using a GC program initiated with an oven temperature of 60°C held for 2 min followed by a ramp to 320°C at 10°C/min with a hold at 320°C for 2 min. Following a solvent delay of 3 min, EI+ mode was used to collect MS data. Diterpene products were identified and confirmed by comparison with spectra archived in the National Institute of Standards and Technology (NIST) database.

Results

Cryo-EM Structure

Cryo-EM grids of full-length S723T-PvCPS were prepared at a protein concentration of

1.0 mg/mL (approximately 10 µM), a concentration at which hexamers and monomers are

observed by size exclusion chromatography-multi-angle light scattering (SEC-MALS).[30] Mass

photometry experiments with a 10^3-fold lower protein concentration (10 nM PvCPS) indicate a

mixture of oligomeric species, including monomer, dimer, trimer, tetramer, pentamer, and

hexamer (Figure S3), consistent with the previously-reported concentration-dependence of

PvCPS oligomerization.[30] While oligomeric heterogeneity of PvCPS persists at the 10 µM

concentration used to prepare cryo-EM grids, prenyltransferase hexamers were visible in 2D

class averages (Figure 2A). Because a relatively small number of prenyltransferase hexamer

particles were observed, and because we previously reported high-resolution X-ray crystal

structures of wild-type and S723T-PvCPS prenyltransferase hexamers,[30,32] we did not pursue a

3D reconstruction of the prenyltransferase hexamer. Finally, mass photometry experiments

indicate that 10 nM PvCPS$_{SL}$ is predominantly hexameric (Figure S3), but this variant did not

yield suitable grids for cryo-EM analysis.

Associated cyclase domains were not visible in 2D class averages of S723T-PvCPS

prenyltransferase hexamers, consistent with a model in which cyclase domains are randomly

splayed out from the prenyltransferase domains regardless of the oligomeric state of the protein.

This is consistent with our previously reported cryo-EM studies of copalyl diphosphate synthase

from *Penicillium fellutanum* (PfCPS).[31] Despite the oligomeric heterogeneity of S723T-PvCPS in

cryo-EM grids, reconstructions of the class II cyclase domain would be insensitive to oligomeric

state. Since the S723T substitution occurs in the prenyltransferase domain, these cryo-EM data

provided an ideal opportunity for closer study of the wild-type cyclase domain. Indeed, these

cryo-EM data yielded 2D class averages of the 63-kD class II cyclase that clearly revealed

secondary structure and characteristic βγ domain architecture. Interestingly, some βγ didomain particles were paired in close proximity with others in various orientations, suggesting the possibility of weak βγ-βγ interdomain interactions (Figure 2B).

Although 2D classes of the class II cyclase appeared to be promising (Figure 2C), achieving a high-resolution 3D reconstruction remained a significant challenge in view of the relatively small size of the particle, which at 63-kD approaches the lower limit of current cryo-EM capabilities. However, deployment of a newly developed small-particle processing pipeline in RELION, Blush regularization, enabled a 3D reconstruction at 2.9 Å-resolution. This reconstruction thus serves as a noteworthy example of cryo-EM structure determination of a relatively small protein. The reconstruction clearly reveals the expected didomain architecture of βγ domain assembly. The final map reveals all 12 helices of the β domain; however, two helices of the γ domain are characterized by ambiguous and weak density, so only 9 helices are included in the final model (Figure 2D).

Figure 2. Representative 2D classes and final 3D reconstruction of the class II cyclase domain. (A) 2D class showing a hexameric assembly corresponding to the prenyltransferase domain. (B) 2D class averages showing βγ-βγ pairs. (C) High quality 2D classes of individual βγ domains. (D) Final 3D reconstruction of the class II cyclase with local resolution estimation as indicated.

The final cryo-EM reconstruction of the PvCPS class II cyclase domain allows for visualization of active site residues (Figure 3A, 3B) and can be compared with other class II cyclases that generate a variety of copalyl diphosphate stereoisomers or regioisomers (Figure 3C, 3D). To date, structures of the following copalyl diphosphate synthases are available in the Protein Data Bank (PDB): (+)-copalyl diphosphate synthase contained in the bifunctional

enzyme abietadiene synthase from *Abies grandis*,[42] (+)-copalyl diphosphate synthase contained in the bifunctional enzyme miltiradiene synthase from *Selaginella moellendorffii*,[43] *ent*-copalyl diphosphate synthase from *Arabidopsis thaliana*,[20] *ent*-copalyl diphosphate synthase from *Streptomyces platensis*,[44] *syn*-copalyl diphosphate synthase from *Oryza sativa*,[41] and 7,13-copalyl diphosphate synthase from *Grindelia robusta.*[45] The structure of the PvCPS class II cyclase presented here is the first example of a fungal class II cyclase and the first example of a class II cyclase contained in a bifunctional prenyltransferase-cyclase system. PvCPS is also the third example of a class II cyclase that generates the copalyl diphosphate synthase stereoisomer shown in Figure 1, also known as (+)- or "normal" copalyl diphosphate.

Figure 3. (A) Final model of the PvCPS class II cyclase (β domain = green, γ domain = yellow). (B) Active site of the class II cyclase domain showing key residues in the cryo-EM map, including general acid D313 in the DXDD motif along with several hydrophobic and aromatic residues. D313 is not characterized by well-defined density and side chain atoms are accordingly omitted from the final model shown. (C) Alignment of the βγ domains of PvCPS (cyan), a bacterial *ent*-copalyl diphosphate (magenta, PDB: 5BP8), copalyl diphosphate synthase from bifunctional abietadiene synthase (green, PDB: 3S9V), and a plant *ent*-copalyl diphosphate synthase (yellow, PDB: 4LIX) reveals generally similar overall protein folds. (D) Comparison of active site residues in aligned class II cyclases reveals generally similar active site structures, except for A256 and M214.

Comparisons of selected normal and *ent*-copalyl diphosphate synthases across different species reveal both conserved and subtly divergent structural features (Figure 3C, 3D).

Alignment of the PvCPS cyclase domain with a bacterial *ent*-copalyl diphosphate synthase, a plant *ent*-copalyl diphosphate synthase, and the copalyl diphosphate synthase from bifunctional abietadiene synthase yields root-mean-square deviation (RMSD) values of 2.1, 3.8, and 3.4 Å, respectively, indicating generally similar overall structures. Several aromatic residues defining the active site contour, F262, W517, W266, and Y523, are conserved in these copalyl diphosphate synthases despite the fact that these enzymes generate two different product stereoisomers (Figure 3D). F262 and Y523 are substituted by Y317 and H501, respectively, in *syn*-CPS.

Interestingly, certain active site residues differ slightly in their positioning or identity among the copalyl diphosphate synthases, including F304 and W400 in PvCPS (Figure 4). In PvCPS, W400 is positioned lower in the active site, potentially guiding the stereochemical outcome towards normal copalyl diphosphate. In *ent*- or *syn*-copalyl diphosphate synthases, the corresponding tryptophan is positioned higher in the active site (Figure 4), which may influence catalysis. Active site aromatic residues were recently subjected to mutagenesis in *ent*-CPS enzymes from *Erwinia tracheiphila* and *Arabidopsis thaliana*, revealing that the tryptophan equivalent to W400 in PvCPS is more important for catalytic efficiency rather than directing product outcome.[46] The structures of CPS domains from abietadiene synthase from *Abies grandis* (AgAS), multiradiene synthase from *Selaginella moellendorffii*, and CPS from *Grindelia robusta* are excluded from this comparison since the loop containing the corresponding tryptophan residue adopts an alternative conformation that does not substantially contribute to the active site contour (for example, PvCPS and AgAS are compared in Figure S4). Also of note, the diphosphate-binding region of copalyl diphosphate synthase active sites exhibits varying degrees of accessibility. In PvCPS and AtCPS this region is more constricted, whereas in *ent*-CPS and *syn*-CPS this region is much less constricted. The conserved glutamate in this

region (E139 in PvCPS) coordinates to Mg^{2+} ions that are also coordinated by the substrate diphosphate group.[18,47]

Figure 4. Product and active site structures of copalyl diphosphate synthases. (A) Molecular structures of copalyl diphosphate stereoisomers. (B) Active site contours of PvCPS, *ent*-CPS from *Streptomyces platensis*, *syn*-CPS from *Oryza sativa*, and *ent*-CPS from *Arabidopsis thaliana* reveal significant variations, some of which appear to be highly complementary to the particular product stereoisomer generated by the enzyme. The active site contour of PvCPS was generated from an AlphaFold3 model since some active site loops in the cryo-EM structure are disordered. However, residues shown for PvCPS derive from the cryo-EM structure.

A general base is required to quench the final carbocation intermediate. While this base is located in a conserved region of the active site, the identity of this general base varies depending on product stereochemistry and species of origin.[18,48,49] Alternative general bases appear to guide the generation of alternative bicyclic diterpene skeletons. For example, tuberculosinyl diphosphate

(TBP) synthase[50] from *Mycobacterium tuberculosis* and terpentedienyl diphosphate (TPP)

synthase[51] from *Kitasatospora sp.* CB02891 generate halimane and clerodane diterpene skeletons,

respectively, which differ from the labdane diterpene skeleton of copalyl diphosphate (Figure 5). The

catalytic base in TBP synthase is a tyrosine positioned properly for a final deprotonation event

accompanied by a series of 1,2-hydride and 1,2-methyl shifts to give the halimane diterpene

product.[50] In TPP synthase, the catalytic acid (D296) also serves as the catalytic base to yield the

clerodane core.[51]

Figure 5. Molecular strategies for quenching the final carbocation intermediate in the generation of bicyclic diterpenes. (A) Terpentedienyl diphosphate (TPP) synthase is thought to utilize the conjugate base of general acid D296 to abstract an axial proton in the A ring, which is accompanied by four 1,2-methyl and -hydride migrations to yield the clerodane skeleton. While these migrations are illustrated as concerted for the sake of brevity, additional carbocation intermediates enroute to the clerodane product are possible. (B) Tuberculosinyl diphosphate (TBP) synthase likely utilizes the phenolic side chain of Y479 as a general base to abstract a proton in the B ring, which triggers 1,2-hydride/1,2-methyl/1,2-hydride migrations to yield the halimane skeleton. Although shown as a concerted reaction, additional carbocation intermediates are possible enroute to the halimane product. (C) PvCPS may utilize the imidazole group of H190 or the carboxylate group of E519 as a general base to abstract a proton from the exo methyl group of the B ring to generate the labdane skeleton.

Examples of class II diterpene cyclases with product outcomes that differ only by stereochemistry, but which utilize distinct catalytic bases, have also been identified (Figure 6A–D). For example, it has been proposed that plant *ent*-CPS in gibberellin phytohormone biosynthetic pathways utilize a histidine-asparagine catalytic dyad contained in L**H**S and P**N**V motifs that positions a hydrogen bonded water molecule to serve as the general base that quenches the final carbocation intermediate.[18,48,49,52,53] However, these motifs are not strictly conserved and also include histidine-tyrosine or histidine-threonine pairs (the latter involving just the first motif as L**HT**).[18, 54-56] Mutagenesis studies of *ent*-CPS from the fungus *Gibberella fujikuroi* indicate a conserved function for the histidine in the L**H**S motif as well as the threonine in an SG**T** motif that replaces the P**N**V motif in this enzyme.[49] The histidine-threonine dyad is thought to mediate the final deprotonation step leading to *ent*-copalyl diphosphate formation. The crystal structure of *Streptomyces platensis ent*-CPS synthase reveals a unique aspartate-histidine dyad in which the aspartate is contained in a different loop compared to the asparagine in the P**N**V motif; even so, this dyad is proposed to activate a hydrogen bonded water molecule or utilize the aspartate-activated histidine to serve as a general base.[44,49] Sequence alignment of various CPS enzymes illustrates the diversity of catalytic dyads that appear to serve general base functions (Figure 6E).

Figure 6. Catalytic dyads thought to serve general base functions in the generation of normal, *ent-*, and *syn*-copalyl diphosphate stereoisomers. (A) The catalytic dyad in *ent*-CPS from *Arabidopsis thaliana* consists of an asparagine and histidine that hydrogen bond to a water molecule thought to mediate the final deprotonation. (B) The catalytic dyad in *ent*-CPS from *Streptomyces platensis* consists of an aspartate and a histidine. (C) The catalytic dyad in *Abies grandis* abietadiene synthase (AgAS) consists of a histidine-tyrosine pair. (D) The catalytic dyad of PvCPS consists of a glutamate and histidine, the latter of which may serve to mediate the final deprotonation step. (E) Sequence alignment of various copalyl diphosphate synthases, indicating product stereochemistry as (n) for normal and (e) for *ent*. Triangles indicate residues comprising the catalytic dyad in each enzyme: PvCPS, CPS from *Penicillium verruculosum*; SpCPS, *ent*-CPS from *Streptomyces platensis*; AtCPS, *ent*-CPS from *Arabidopsis thaliana*; AgAS, CPS contained in abietadiene synthase from *Abies grandis*; GfCPS, *ent*-CPS from *Gibberella fujikuroi*.

Substrate Channeling

Recent studies show that certain bifunctional class I terpene synthases are capable of channeling GGPP directly from the prenyltransferase to the cyclase.[25-27] It has not yet been determined whether a bifunctional class II terpene synthase can similarly engage in substrate channeling. Cryo-EM studies of bifunctional class I terpene synthases suggest that weak, transient association of a cyclase domain with the side of a prenyltransferase oligomer facilitates substrate channeling.[26-29] However, cryo-EM studies of PvCPS reported herein do not reveal any particles showing prenyltransferase hexamers with side-on binding of cyclase domains.

To ascertain the possibility of substrate channeling in PvCPS, we performed a competition assay in which an equimolar mixture of PvCPS and the diterpene cyclase cyclooctatenol synthase (CotB2) or fusicoccadiene synthase (PaFS) were incubated with GGPP. In a second experiment, the equimolar enzyme mixture was incubated with DMAPP and IPP such that the only source of GGPP in the reaction mixture was that generated by the PvCPS prenyltransferase. If substrate channeling were operative, the ratio of copalyl diphosphate-derived copalol/cyclooctatenol or copalol/fusicoccadiene would increase when the enzyme mixture is incubated with DMAPP and IPP, indicating that GGPP generated by the prenyltransferase remains on the enzyme for cyclization to form copalyl diphosphate.

Quantitation of copalyl diphosphate-derived copalol and cyclooctatenol using GC-MS reveals that the copalol/cyclooctatenol ratio increases, by more than five-fold, when the enzyme mixture is incubated with DMAPP and IPP compared to GGPP (Figure 7, Table 1). If the PvCPS cyclase and prenyltransferase are included as separate constructs along with the competitor cyclase, the ratio with DMAPP and IPP still increases, but to slightly less than five-fold. When an equimolar mixture of PvCPS$_{SL}$ and CotB2 is studied using the same competition assay, the copalol/cyclooctatenol ratio increases by more than three-fold when the enzyme mixture is

incubated with DMAPP and IPP compared to GGPP (Figure 7, Table 1). These observations suggest that much of the GGPP generated by the PvCPS prenyltransferase remains on the enzyme for cyclization to form copalyl diphosphate, i.e., there is some degree of substrate channeling from the prenyltransferase to the cyclase regardless of whether they are covalently linked or not. These results are consistent with a model in which transient prenyltransferase-cyclase association facilitates GGPP transfer, but the optimal positioning of prenyltransferase and cyclase domains is unknown. Channeling appears to be slightly less efficient in PvCPS$_{SL}$, perhaps because cyclase domains are held in a less favorable position relative to the prenyltransferase due to the shorter linker.

Figure 7. Products resulting from substrate competition experiments with PvCPS or CotB2 (A) and PvCPS and PaFS$_{CY}$ (B). GC traces reveal products resulting from incubation of the equimolar enzyme mixture with GGPP (front trace) or DMAPP and IPP (back trace). COT, cyclooctatenol; COP, copalol; FS, fusicoccadiene. Traces are normalized such that the largest peak is equivalent in size.

Table 1. Product ratios resulting from substrate competition experiments.

Equimolar Enzyme Mixture	Products Formed[a]	Product Ratio from GGPP	Product Ratio from DMAPP and IPP
PvCPS, CotB2	COP:COT	0.64 ± 0.08 : 1	3.33 ± 0.18 : 1
PvCPS, PaFS$_{CY}$	COP:FS	1.06 ± 0.07 : 1	1.55 ± 0.13 : 1
PvCPS$_{PT}$, PvCPS$_{CY}$, CotB2	COP:COT	0.29 ± 0.03 : 1	1.35 ± 0.03 : 1
PvCPS$_{PT}$, PvCPS$_{CY}$, PaFS$_{CY}$	COP:FS	1.98 ± 0.16 : 1	2.26 ± 0.23 : 1
PvCPS$_{SL}$, CotB2	COP:COT	0.43 ± 0.05 : 1	1.41 ± 0.10 : 1
PvCPS$_{SL}$, PaFS$_{CY}$	COP:FS	1.05 ± 0.10 : 1	0.52 ± 0.01 : 1

[a]COP, copalol; COT, cyclooctatenol; FS, fusicoccadiene

When the substrate competition experiment is performed with an equimolar mixture of PvCPS and PaFS$_{CY}$, or an equimolar mixture of PvCPS$_{PT}$, PvCPS$_{CY}$, and PaFS$_{CY}$, the copalol/fusicoccadiene ratio increases only slightly, by 1.46-fold, and 1.14-fold, respectively, when the enzyme mixture is incubated with DMAPP and IPP compared to GGPP (Figure 7, Table 1). Channeling in PvCPS appears to be marginal when the PvCPS cyclase competes with PaFS$_{CY}$ for GGPP generated by the PvCPS prenyltransferase. Curiously, when the cyclase domain of PvCPS$_{SL}$ competes with PaFS$_{CY}$ for GGPP generated by the PvCPS$_{SL}$ prenyltransferase domain, there is increased production of fusicoccadiene. This result suggests that GGPP preferentially transits to PaFS$_{CY}$ rather than the cyclase domain of PvCPS$_{SL}$. In other words, intermolecular channeling of GGPP to PaFS$_{CY}$ appears to be preferred over intramolecular channeling of GGPP to the cyclase domain of PvCPS$_{SL}$. Intermolecular GGPP channeling to PaFS$_{CY}$ was similarly observed with variediene synthase, where a transient side-on binding mode of PaFS$_{CY}$ to the prenyltransferase hexamer was proposed to account for intermolecular channeling.[27]

Discussion

In the past two decades, cryo-EM has come to rival X-ray crystallography for protein structure determination at high resolution; indeed, proteins that are refractory to crystallization are often amenable to structure determination by cryo-EM.[57] However, cryo-EM has its limitations as well, in that smaller proteins <100 kD often fail to yield satisfactory reconstructions due to alignment complications associated with a lower signal-to-noise ratio.[35] To facilitate the study of smaller proteins using cryo-EM, enabling tools such as nanobodies have been developed to bind to a target particle so as to increase the particle size.[58] Absent such approaches, high-resolution reconstructions of particles less than ~100 kD are relatively rare, especially in cases where molecular symmetry cannot be exploited to enhance the signal-to-noise ratio. A hexamer of full-length PvCPS is approximately 660 kD and therefore should be a

reasonable target for cryo-EM analysis. However, due to significant flexibility of the 125-residue linker connecting the α and βγ domains in each α~βγ protomer, such that the six βγ domains of the class II cyclase are randomly splayed-out around a central prenyltransferase α_6 hexamer, high-resolution structure determination of the full-length protein using either cryo-EM or X-ray crystallography is impossible.

Even so, structure determinations of the individual catalytic domains of PvCPS are feasible. For example, crystallization of the prenyltransferase α_6 hexamer yielded a 2.41 Å-resolution X-ray crystal structure,[30] crystal structures of site-specific prenyltransferase variants have been determined at resolutions ranging 2.00–3.15 Å,[32] and the cryo-EM structure of the related prenyltransferase domain of PfCPS has been determined at 2.81 Å resolution.[31] Although the βγ domain assembly of the PvCPS cyclase was not amenable to crystallization, we show here that it is amenable to atomic resolution structure determination by cryo-EM. Moreover, a separate construct was not required for structure determination, since cryo-EM grids prepared with full-length PvCPS yielded separate 2D reconstructions of the prenyltransferase hexamer and the class II cyclase domains. Since the structure of the prenyltransferase hexamer had already been reported, we focused our attention on the 63-kD class II cyclase domain. The use of the new Blush regularization tool in RELION[35] enabled reconstruction of this relatively small particle at 2.9 Å-resolution.

The structure of the PvCPS cyclase domain is useful for advancing our understanding of structure-function relationships in the greater family of class II terpene cyclases. There are no structures reported to date for a class II cyclase from a fungal species, so it is instructive to compare the PvCPS cyclase with plant and bacterial class II cyclases. In particular, both bacterial and plant copalyl diphosphate synthases share a number of conserved residues lining the active site pocket, although key differences appear to direct alternative mechanistic pathways for the generation of alternative stereoisomers of the labdane skeleton. For example,

plant enzymes involved in gibberellin biosynthesis have a histidine-asparagine catalytic dyad which interacts with a water molecule functional serving to facilitate the final deprotonation step leading to *ent*-CPS products.[18,48,49] The identify of the catalytic base in class II diterpene cyclases has been found to be highly variable though, with examples that include tyrosine-histidine or histidine-threonine dyads.[18, 54-56] Notably, in a bacterial *ent*-CPS the catalytic dyad consists of an aspartate-histidine pair hydrogen bonded to a water molecule that may be responsible for the final proton abstraction.[44] Interestingly, if G487 of a bacterial TPP synthase is substituted by aspartate, the product outcome is altered so as to generate a labdane skeleton instead of the native clerodane skeleton.[51] The cryo-EM structure of the PvCPS cyclase reveals that E519 corresponds to the glutamate or aspartate observed in bacterial class II cyclases that generate the labdane core, suggesting that this residue similarly serves as a general base in catalysis. Since fungal species are more closely related to plants, it is interesting that PvCPS seems to have adopted the mechanistic strategy of a bacterial cyclase.

The current study is the first to reveal GGPP channeling from the prenyltransferase to the cyclase in a class II bifunctional terpene synthase. Transient prenyltransferase-cyclase association is proposed to account for intramolecular as well as intermolecular channeling, as previously proposed for certain class I bifunctional terpene synthases,[26-29] but the mode of prenyltransferase-cyclase association in a class II system remains unknown. Of note, PvCPS oligomerization is sensitive to concentration, and oligomerization could conceivably influence GGPP channeling due to the increased number of enzyme active sites held in close proximity to each other. Previous biophysical studies of PvCPS suggest that this system does not form hexamers exclusively at 5 µM enzyme concentrations (the concentration utilized for channeling experiments), but instead is a mixture of hexamer and lower order oligomers.[30] This is consistent with the relatively low occurrence of prenyltransferase hexamers found in 2D class averages of cryo-EM samples prepared with 10 µM enzyme concentrations (Figure 2A).

Concluding Remarks

The 2.9 Å-resolution cryo-EM structure of the 63-kD class II cyclase domain of PvCPS is notable in that this relatively small protein yielded a reconstruction at atomic resolution. The structure of the PvCPS cyclase enables comparisons with other class II cyclases, including enzymes that generate alternative regioisomers or stereoisomers of copalyl diphosphate and related bicyclic diterpenes. Natural products derived from copalyl diphosphate, such as abietane and primane-type diterpenes, are found in the plant oleoresin secreted in response to insect attack or physical injuries.[59] Copalyl diphosphate is also associated with molecules which have a broad range of medicinal applications, including modified grindelic acids and native sclareols that exhibit anti-tumor, anti-inflammatory, and anti-pathogenic activities.[60-62] A greater understanding of the prenyltransferase-cyclase assembly line will advance synthetic biology approaches for the generation of valuable natural products. Future studies will probe the nature of transient prenyltransferase-cyclase association and the role of preferential GGPP transit in these systems.

ASSOCIATED CONTENT

Supporting Information

Supporting Information is available free of charge at:

Table S1, cryo-EM structure determination statistics; Figure S1, cryo-EM workflow; Figure S2, particle orientation and Gold-Standard Fourier Shell Correlation plot; Figure S3, mass photometry data; Figure S4, structural comparison of the W400 active site loop.

Accession Codes

The cryo-EM map of the PvCPS cyclase has been deposited in the Electron Microscopy Data Bank (EMDB, www.ebi.ac.uk/pdbe/emdb) with accession codes EMD-72194. Atomic coordinates of the PfCPS prenyltransferase hexamer have been deposited in the Protein Data Bank (www.rcsb.org) with accession code 9Q3I.

AUTHOR INFORMATION

Corresponding Author

David W. Christianson – Roy and Diana Vagelos Laboratories, Department of Chemistry, University of Pennsylvania, Philadelphia, Pennsylvania 19104-6323, United States; orcid.org/0000-0002-0194-5212; Email: chris@sas.upenn.edu

Authors

Matthew N. Gaynes – Roy and Diana Vagelos Laboratories, Department of Chemistry, University of Pennsylvania, Philadelphia, Pennsylvania 19104-6323, United States. orcid.org/0000-0001-9649-601X.

Kollin Shultz – Abramson Family Cancer Research Institute, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104

Graduate Group in Biochemistry and Molecular Biophysics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104. orcid.org/0000-0002-4456-8349

Eliott S. Wenger – Roy and Diana Vagelos Laboratories, Department of Chemistry, University of Pennsylvania, Philadelphia, Pennsylvania 19104-6323, United States. orcid.org/0000-0001-6391-8036

Ronen Marmorstein – Abramson Family Cancer Research Institute, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104

Department of Biochemistry and Biophysics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104. orcid.org/0000-0003-4373-4752

Funding

This research was supported by NIH grants R01 GM56838 to D.W.C. and R35 GM118090 to R.M. M.N.G. and K.S. were supported by Chemistry-Biology Interface NIH Training Grant T32 GM133398.

Conflict of Interest Statement

The authors declare no competing interests.

ACKNOWLEDGMENTS

This research utilized cryo-EM facilities at the Laboratory for BioMolecular Structure (LBMS), Brookhaven National Laboratory, which is supported by the DOE Office of Biological and Environmental Research (KP1607011).

References

1. Stepanyuk, A., Kirschning, A. (2019) Synthetic terpenoids in the world of fragrances: Iso E Super® is the showcase. *Beilstein J. Org. Chem. 15*, 2590–2602.

2. Masyita, A., Sari, R. M., Dwi, A. A., Yasir B, Rahma, R. N., Emran, T. B., Nainu, F., Simal-Gandara, J. (2022) Terpenes and terpenoids as main bioactive compounds of essential oils, their roles in human health and potential application as natural food preservatives. *Food Chem X 13*, 100217.

3. Gozari, M., Alborz, M., El-Seedi, H. R., Jassbi, A. R. (2021) Chemistry, biosynthesis and biological activity of terpenoids and meroterpenoids in bacteria and fungi isolated from different marine habitats. *Eur. J. Med. Chem. 210*, 112957.

4. Cox-Georgian, D., Ramadoss, N., Dona, C., Basu, C. (2019) Therapeutic and medicinal uses of terpenes. In: Joshee, N., Dhekney, S., Parajuli, P., Eds. *Medicinal Plants,* 333–359.

5. Peralta-Yahya, P., Ouellet, M., Chan, R., Mukhopadhyay, A., Keasling, J. D., Lee, T. S. (2011) Identification and microbial production of a terpene-based advanced biofuel. *Nat. Commun. 2*, 483.

6. Woodroffe, J. D., Harvey, B. G. (2020) High-performance, biobased, jet fuel blends containing hydrogenated monoterpenes and synthetic paraffinic kerosenes. *Energy Fuels 34*, 5929–5937.

7. Poulter, C. D., Rilling, H. C. (1978) The prenyl transfer reaction. enzymic and mechanistic studies of the 1'–4 coupling reaction in the terpene biosynthetic pathway. *Acc. Chem. Res. 11*, 307–313.

8. Kellogg, B.A., Poulter, C. D. (1997) Chain elongation in the isoprenoid biosynthetic pathway. *Curr. Opin. Chem. Biol. 1*, 570–578.

9. Zhang, F. L., Casey, P. J. (1996). Protein prenylation: molecular mechanisms and functional consequences. *Annu. Rev. Biochem. 65*, 241–269.

10. Wang, M., Casey, P. J. (2016) Protein prenylation: unique fats make their mark on biology. *Nat. Rev. Mol. Cell Biol. 17*, 110–122.

11. Wang, H., Yang, Y., Abe, I. (2024) Modifications of prenyl side chains in natural product biosynthesis. *Angew. Chem. Int. Ed. Engl. 63*, e202415279.

12. Kuzuyama, T., Noel, J., Richard, S. (2005) Structural basis for the promiscuous biosynthetic prenylation of aromatic natural products. *Nature 435*, 983–987.

13. Christianson, D. W. (2007) Roots of biosynthetic diversity. *Science 316*, 60–61.

14. Christianson, D. W. (2017) Structural and chemical biology of terpenoid cyclases. *Chem. Rev. 117*, 11570–11648.

15. Whitehead, J. N., Leferink, N. G. H., Johannissen, L. O., Hay, S., Scrutton, N. S. (2023) Decoding catalysis by terpene synthases. *ACS Catal. 13*, 12774–12802.

16. Aaron, J.A., & Christianson, D.W. (2010) Trinuclear metal clusters in catalysis by terpenoid synthases. *Pure Appl. Chem. 82*, 1585–1597.

17. Oldfield, E., Lin, F.-Y. (2012) Terpene biosynthesis: modularity rules. *Angew. Chem. Int. Ed. Engl. 51*, 1124–1137.

18. Pan, X., Rudolf, J. D., Liao-Bin, D. (2024) Class II terpene cyclases: structures, mechanisms, and engineering. *Nat. Prod. Rep. 41*, 402–433.

19. Prisic, S., Xu, J., Coates, R. M., Peters, R. J. (2007) Probing the role of the DXDD motif in class II diterpene synthases. *ChemBioChem 8*, 869–874.

20. Köksal, M., Hu, H., Coates, R. M., Peters, R. J., Christianson, D. W. (2011) Structure and mechanism of the diterpene cyclase *ent*-copalyl diphosphate synthase. *Nat. Chem. Biol. 7*, 431–433.

21. Chen, R., Jia, Q., Mu, X., Hu, B., Sun, X., Deng, Z., Chen, F., Bian, G., Liu, T. Systematic mining of fungal chimeric terpene synthases using an efficient precursor-providing yeast chassis. (2021) *Proc. Natl. Acad. Sci. USA 118,* e2023247118.

22. Toyomasu, T., Tsukahara, M., Kaneko, A., Niida, R., Mitsuhashi, W., Dairi, T., Kato, N., Sassa, T. (2007) Fusicoccins are biosynthesized by an unusual chimera diterpene synthase in fungi. *Proc. Natl. Acad. Sci. USA 104*, 3084–3088.

23. Mitsuhashi, T., Okada, M., Abe, I. (2017) Identification of chimeric αβγ diterpene synthases possessing both type II terpene cyclase and prenyltransferase activities. *ChemBioChem 18*, 2104–2109.

24. Faylo, J. L., Ronnebaum, T. A., Christianson, D. W. (2021) Assembly-line catalysis in bifunctional terpene synthases. *Acc. Chem. Res. 54*, 3780–3791.

25. Pemberton, T. A., Chen, M., Harris, G. G., Chou, W. K. W., Duan, L., Köksal, M., Genshaft, A. S., Cane, D. E., Christianson, D. W. (2017) Exploring the influence of domain architecture on the catalytic function of diterpene synthases. *Biochemistry 56*, 2010–2023.

26. Wenger, E. S., Schultz, K., Marmorstein, R., Christianson, D. W. (2024) Engineering substrate channeling in a bifunctional terpene synthase. *Proc. Natl. Acad. Sci. USA 121*, e2408064121.

27. Wenger, E. S., Christianson, D. W. (2025) Structure of bifunctional variediene synthase yields unique insight on biosynthetic diterpene assembly and cyclization. *Nat. Commun. 16*, 5089.

28. Faylo, J. L., van Eeuwen, T., Kim, H. J., Gorbea Colón J. J., Garcia, B. A., Murakami, K., Christianson, D. W. (2021) Structural insight on assembly-line catalysis in terpene biosynthesis. *Nat. Commun. 12*, 3487.

29. Faylo, J. L., van Eeuwen, T., Gupta, K., Murakami, K., Christianson, D. W. (2022) Transient prenyltransferase-cyclase association in fusicoccadiene synthase, an assembly-line terpene synthase. *Biochemistry 61*, 2417–2430.

30. Ronnebaum, T. A., Gupta, K., Christianson, D. W. (2020) Higher-order oligomerization of a chimeric αβγ bifunctional diterpene synthase with prenyltransferase and class II cyclase activities is concentration-dependent. *J. Struct. Biol. 210*, 107463.

31. Gaynes, M. N., Ronnebaum, T. A., Schultz, K., Faylo, J. L., Marmorstein, R., Christianson, D. W. (2024) Structure of the prenyltransferase in bifunctional copalyl diphosphate synthase from *Penicillium fellutanum* reveals an open hexamer conformation. *J. Struct. Biol. 16*, 108060.

32. Ronnebaum, T. A., Eaton, S. A., Brackhahn, E. A. E., Christianson, D. W. (2021) Engineering the prenyltransferase domain of a bifunctional assembly-line terpene synthase. *Biochemistry 60*, 3162–3172.

33. Kim, S. Y., Zhao, P., Igarashi, M., Sawa, R., Tomita, T., Nishiyama, M., Kuzuyama, T. (2009) Cloning and heterologous expression of the cyclooctatin biosynthetic gene cluster afford a diterpene cyclase and two P450 hydroxylases. *Chem. Biol. 16*, 736–743.

34. Janke, R., Görner, C., Hirte M., Brück, T., Loll, B. (2014) The first structure of a bacterial diterpene cyclase: CotB2. *Acta Cryst. D70*, 1528–1537.

35. Kimanius, D., Jamali, K., Wilkinson, M. E., Lövestam, S., Velazhahan, V., Nakane, T., Scheres, S. (2024) Data-driven regularization lowers the size barrier of cryo-EM structure determination. *Nat. Methods 21*, 1216–1221.

36. Abramson, J., et al. (2024) Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature 630*, 493–500.

37. Pettersen, E. F. T. D., Goddard, C. C., Huang, G. S., Couch, D. M., Greenblatt, E. C., Meng, T. E., Ferrin, T. E. (2004) UCSF Chimera: a visualization system for exploratory research and analysis. *J. Comput. Chem. 25*, 1605–1612.

38. Afonine, P. V., Poon, B. K., Read, R. J., Sobolev, O. V., Terwilliger, T. C., Urzhumtsev, A., Adams, P. D. (2018) Real-space refinement in *PHENIX* for cryo-EM and crystallography. *Acta Cryst. D74*, 531–544.

39. Emsley, P., Lohkamp, B., Scott, W. G., Cowtan K. (2010) Features and development of *Coot. Acta Cryst. D66*, 486–501.

40. Williams, C. J., et al. (2018) MolProbity: more and better reference data for improved all-atom structure validation. *Protein Sci. 27*, 293–315.

41. Ma, X., Xu, H., Tong, Y., Luo, Y., Dong, Q., Jiang, T. (2023) Structural and functional investigations of *syn*-copalyl diphosphate synthase from *Oryza sativa. Commun Chem 6*, 240.

42. Zhou, K., Gao, Y., Hoy, J. A., Mann, F. M., Honzatko, R. B., Peters, R. J. (2012) Insights into diterpene cyclization from structure of bifunctional abietadiene synthase from *Abies grandis. J. Biol. Chem. 287*, 6840–6850.

43. Tong, Y., Ma, X., Hu, T., Chen, K., Cui, G., Su, P., Xu, H., Gao, W., Jiang, T., Huang, L. (2023) Structural and mechanistic insights into the precise product synthesis by a bifunctional miltiradiene synthase. *Plant Biotechnol. J. 21*, 165–175.

44. Rudolf, J. D., Dong, L. B., Cao, H., Hatzos-Skintges, C., Osipiuk, J., Endres, M., Chang, C. Y., Ma, M., Babnigg, G., Joachimiak, A., Phillips, G. N., Shen, B. (2016) Structure of the *ent*-copalyl diphosphate synthase PtmT2 from *Streptomyces platensis* CB00739, a bacterial type II diterpene synthase. *J. Am. Chem. Soc. 138*, 10905–10915.

45. Cowie, A. E., Pereira, J. H., DeGiovanni, A., McAndrew, R. P., Palayam, M., Peek, J. O., Muchlinski, A. J., Yoshikuni, Y., Shabek, N., Adams, P. D., Zerbe, P. (2024) The crystal structure of *Grindelia robusta* 7,13-copalyl diphosphate synthase reveals active site features controlling catalytic specificity. *J. Biol. Chem. 300*, 107921.

46. Raslan, A. M. A. A., Lemke, C., Larsen, R., Peters, R. (2025) Investigating conserved aromatic residues in *ent*-copalyl pyrophosphate synthases required for gibberellin phytohormone biosynthesis. *Phytochem. 240*, 114635.

47. Pan, X., Du, W., Zhang, X., Lin, X., Li, F.-R., Yang, Q., Wang, H., Rudolf, J. D., Zhang, B. (2022) Discovery, structure, and mechanism of a class II sesquiterpene cyclase. *J. Am. Chem. Soc. 144*, 22067–22074.

48. Jia, Q., Brown, R., Köllner , T. G., Fu, J., Chen, X., Wong, G. K., Gershenzon, J., Peters, R. J., Chen, F. (2022) Origin and early evolution of the plant terpene synthase family. *Proc. Natl. Acad. Sci. 119*, e2100361119.

49. Lemke, C., Potter, K. C., Schulte, S., Peters, R. J. (2019) Conserved bases for the initial cyclase in gibberellin biosynthesis: from bacteria to plants. *Biochem. J. 476*, 2607–2621.

50. Zhang, Y., Prach, L. M., O'Brien, T. E., DiMaio, F., Prigozhin, D. M., Corn, J. E., Alber, T., Siegel, J. B., Tantillo, D. J. (2020) Crystal structure and mechanistic molecular modeling studies of *Mycobacterium tuberculosis* diterpene cyclase Rv3377c. *Biochemistry 59*, 4507–4515.

51. Stowell, E. A., Ehrenberger, M. A., Lin, Y. L., Chang, C. Y., Rudolf, J. D. (2022) Structure-guided product determination of the bacterial type II diterpene synthase Tpn2. *Commun. Chem. 5*, 146.

52. Potter, K., Criswell, J., Zi, J., Stubbs, A., Peters, R. J. (2014) Novel product chemistry from mechanistic analysis of *ent*-copalyl diphosphate synthases from plant hormone biosynthesis. *Angew. Chem. Int. Ed. 53*, 7198–7202.

53. Potter, K. C., Zi, J., Hong, Y. J., Schulte, S., Malchow, B., Tantillo, D. J., Peters, R. J. (2016) Blocking deprotonation with retention of aromaticity in a plant *ent*-copalyl diphosphate synthase leads to product rearrangement. *Angew Chem. Int. Ed. Engl. 55*, 634–638.

54. Mafu, S., Potter, K. C., Hillwig, M. L., Schulte, S., Criswell, J., Peters, R. J. (2015) Efficient heterocyclisation by (di)terpene synthases. *Chem. Commun. 51*, 13485–13487.

55. Criswell, J., Potter, K., Shephard, F., Beale, M. H., Peters, R. J. (2012) A single residue change leads to a hydroxylated product from the class II diterpene cyclization catalyzed by abietadiene synthase. *Org. Lett. 14*, 5828–5831.

56. Schulte, S., Potter, K. C., Lemke, C., Peters, R. J. (2018) Catalytic bases and stereocontrol in Lamiaceae class II diterpene cyclases. *Biochemistry 57*, 3473–3479.

57. Kühlbrandt, W. (2014) The resolution revolution. *Science 343*, 1443–1444.

58. Wu, X., Rapoport, T. A. (2021) Cryo-EM structure determination of small proteins by nanobody-binding scaffolds (Legobodies). *Proc. Natl. Acad. Sci. USA 118*, e2115001118.

59. Peters, R. J., Flory, J. E., Jetter, R., Ravn, M. M., Lee, H. J., Coates, R. M., Croteau, R. B. (2000) Abietadiene synthase from grand fir (*Abies grandis*): Characterization and mechanism of action of the ″pseudomature″ recombinant enzyme. *Biochemistry. 39*, 15592–15602.

60. Zhou, J., Xie, X., Tang, H., Peng, C., Peng, F. (2022) The bioactivities of sclareol: A mini review. *Front. Pharmacol. 13*, 1014105.

61. Gierlikowska, B., Filipek, A., Gierlikowski, W., Kania, D., Stefańska, J., Demkow, U., Kiss A. K. (2021) *Grindelia squarrosa* extract and grindelic acid modulate pro-inflammatory functions of respiratory epithelium and human macrophages. *Front. Pharmacol. 11*, 534111.

62. Reta, G. F., Chiaramello, A. I., Garcia, C., León, L. G., Martin, V. S., Padrón, J. M., Tonn, C. E., Donadel, O. J. (2013) Derivatives of grindelic acid: from a non-active natural diterpene to synthetic antitumor derivatives. *Eur. J. Med. Chem. 67*, 28–38.

TOC Graphic

Substrate channeling   Cryo-EM structure

PvCPS
Class II Cyclase